The Diffusive Nature of Housing Prices

Antoine-Cyrus Becharat^{1,2},* Michael Benzaquen^{1,2,3}, and Jean-Philippe Bouchaud^{1,3,4}

¹*Chair of Econophysics and Complex Systems, Ecole Polytechnique, 91128 Palaiseau Cedex, France ´*

2 *LadHyX UMR CNRS 7646, Ecole Polytechnique, 91128 Palaiseau Cedex, France ´*

³*Capital Fund Management, 23 Rue de l'Universite, 75007 Paris, France and ´*

⁴*Academie des Sciences, 23 Quai de Conti, 75006 Paris, France ´*

(Dated: December 20, 2024)

We analyze the French housing market prices in the period 1970-2022, with high-resolution data from 2018 to 2022. The spatial correlation of the observed price field exhibits logarithmic decay characteristic of the twodimensional random diffusion equation – *local* interactions may create *long-range* correlations. We introduce a stylized model, used in the past to model spatial regularities in voting patterns, that accounts for both spatial and temporal correlations with reasonable values of parameters. Our analysis reveals that price shocks are persistent in time and their amplitude is strongly heterogeneous in space. Our study confirms and quantifies the diffusive nature of housing prices that was anticipated long ago [\[1,](#page-4-1) [2\]](#page-4-2), albeit on much restricted, local data sets.

Complex spatial patterns often result from a subtle interplay between random forcing and diffusion, like for example surface growth [\[3\]](#page-4-3) or fluid turbulence [\[4\]](#page-4-4). One can also expect such competition between heterogeneities and diffusion to take place in socio-economic contexts. For example, word of mouth leads to spreading of information or of opinions. Provided the spreading mechanism is local enough (i.e. before the advent of social media), the large scale description of such phenomena is provided by the diffusion equation that leads to specific predictions for the long-range nature of spatial correlations of voting patterns, which seems to be validated by the analysis of empirical data [\[5–](#page-4-5)[7\]](#page-4-6).

One may argue that housing prices should display similar patterns. Indeed, it is intuitively clear that the price of real estate in a given district is affected, among many other factors, by the price of the surrounding districts, through a sheer proximity effect. This is enough to generate a diffusion term in any coarse-grained description of the spatio-temporal evolution of prices – see below and SI-1 for more precise statements. The aim of this work is to present such a phenomenological description of the *price field* in a given region of space, and to compare analytical prediction to empirical data using spatially resolved transaction prices in France for the period 1970 to 2022 – see Fig. [1](#page-0-0) for a visual representation of the price field that motivates our analysis. We will find what we consider to be rather remarkable agreement with theory, in view of the minimal amount of modeling ingredients. In particular, the logarithmic dependence of spatial correlations, characteristic of two-dimensional diffusion, is clearly visible in the data at all scales (see Fig. [3](#page-3-0) below).

Due to its potent macroeconomic and systemic risk implications, the housing market and its corresponding price field have long been studied by economists, see [\[9\]](#page-4-7). One of the most famous description of the housing market is through the Hedonic prices hypothesis (see e.g. [\[10\]](#page-4-8)), which states that *goods are valued for their utility-bearing attributes. Hedonic prices are defined as the implicit prices of attributes and are revealed from observed prices of differentiated products and the specific amounts of characteristics associated with them*. In essence, we shall argue that real-estate prices in the vicinity of a given

Figure 1. Spatial transaction log-prices p distribution in France in 1970 (left) and in 2022 (right). We use a sigmoid transformation of the log prices rescaled by their mean and divided by their standard deviation in order to highlight price differences. As seen in this plot, high prices are concentrated around France's principal cities and on the coasts and mountains, but the price pattern clearly displays spatial diffusion. Data from [\[8\]](#page-4-9).

location is one of these characteristics.

There is also a great body of empirical literature highlighting the links between the housing market prices and, for example, violence [\[11\]](#page-4-10) or school grades [\[12\]](#page-4-11). This has naturally led to models of the housing market using reasonable assumptions. In particular, recent agent-based models of the housing market have been designed to explain price dynamics [\[9\]](#page-4-7), or its link with social segregation. Ref. [\[13\]](#page-4-12) observed that segregation patterns can be observed even with the simplest parameter setting in an agent-based model of the housing market. Ref. [\[14\]](#page-4-13) showed how such models could be very helpful to test and apply effective policies to prevent social/racial segregation, in the same vein as Ref. [\[15\]](#page-4-14) where the effectiveness of macro-prudential policies is tested on an agent-based model of the UK housing market. Interestingly, [\[16\]](#page-4-15) showed that social segregation is also strongly linked with social influence.

Concerning spatial patterns, studies from the mid-1990's have suggested the potential importance of spatial diffusion effects. For example, Clapp & Tirtiroglu [\[1\]](#page-4-1) find evidence of local price diffusion from their empirical study of the metropolitan of Hartford, Connecticut. Pollakowski & Ray [\[2\]](#page-4-2) confirms these results at the local level, and conclude that housing prices are inefficient: *If housing markets were efficient, [...] shocks would either be confined to one area, in which case information transfer is irrelevant, or affect a number of areas, in which case the price changes should occur nearly simultaneously, not one after another.* These authors also note that price changes are auto-correlated in time (a feature that we will explicitly include in our theoretical model), which is a further sign of price inefficiency. Indeed, properly anticipated prices should not be predictable [\[17\]](#page-4-16).

As we argue below, such *local* diffusion of prices is expected to create *long-range correlations* in the price field both in space and in time, which we will indeed observe in the data. Although the presence of spatial correlations were noticed in [\[18\]](#page-4-17), no mention was made of their long-range nature, let alone their specific logarithmic dependence discussed below. Other socio-economic variables, on the other hand, are known to be long-range correlated [\[6\]](#page-4-18), with far-reaching consequences on the statistical significance of many results in spatial economics, as forcefully argued in [\[19\]](#page-4-19).

Our theoretical framework aims at modeling the dynamics of the housing price field in a similar spirit as for the dynamics of opinions or intentions [\[5,](#page-4-5) [20–](#page-4-20)[22\]](#page-4-21). We introduce a twodimensional field $\psi(\mathbf{r}, t)$ which represents the deviation from the (possibly time dependent) mean of the log-price of housing around point r at time t . We then posit that such a field evolves in time according to the following stochastic partial differential equation

$$
\frac{\partial \psi(\mathbf{r},t)}{\partial t} = D\Delta\psi(\mathbf{r},t) - \varkappa\psi(\mathbf{r},t) + \eta(\mathbf{r},t) + \xi(\mathbf{r}), \quad (1)
$$

where Δ is the Laplacian operator, D a diffusion coefficient, κ a mean-reversion coefficient, $\eta(\mathbf{r}, t)$ a Langevin noise with zero mean and short range time and space correlations, and $\xi(\mathbf{r})$ a static random field with zero mean and short range correlations. The correlators of these terms are assumed to be of the following type:

$$
\langle \eta(\mathbf{r},t)\eta(\mathbf{r}',t')\rangle = \frac{A^2}{Ta^2}e^{-|t-t'|/T}g_a(|\mathbf{r}-\mathbf{r}'|);
$$

$$
\langle \xi(\mathbf{r})\xi(\mathbf{r}')\rangle = \frac{\Sigma^2}{a^2}g_a(|\mathbf{r}-\mathbf{r}'|),
$$
 (2)

where $g_a(r)$ is a bell-shaped function that decays over length scale a, such that $2\pi \int_{r>0} g_a(r)r dr = a^2$. Note that in terms of dimensions, $[A^2] = [D] = [L^2T^{-1}], [\varkappa] = [T^{-1}]$ and $[\Sigma] = [LT^{-1}].$

The four different terms of Eq. [\(1\)](#page-1-0) capture the following features: (i) the diffusion term describes the proximity effect alluded to in the introduction and documented in Refs. [\[1,](#page-4-1) [2\]](#page-4-2): pricey districts tend to progressively gentrify; conversely, rundown districts lower the market value of their surroundings. (A more technical version of this argument is given in SI-1). (ii) The mean-reversion term can be seen as a coupling between local log-prices and the mean log-price, here set to zero, and can be thought of as the result of long-range economic forces that keep prices within a country more or less in sync through the effect of e.g. migrations, policies or wealth inequalities. (iii) The time-dependent noise term η models all idiosyncratic shocks affecting the "hedonic" variables determining the price of properties – for example the creation of a local metro or train station, of a pedestrian zone, or adverse shocks like increase in local crime, floods, etc. The impact of such shocks is often drawn out in time, so we assume η to be auto-correlated with a decay time T , in line with the observations reported in [\[2\]](#page-4-2). (iv) The time-independent stochastic term ξ is meant to represent persistent biases in the local quality of life in different regions, due to e.g. geographical features (close to the sea-shore, or to river banks, etc.). For simplicity, We have assumed that the spatial correlation lengths of both η and ξ are equal to the same value a.

Now, Eq. [\(1\)](#page-1-0) makes detailed predictions for the spatial and temporal correlations of the field $\psi(\mathbf{r}, t)$. To wit, the spatial variogram $\mathbb{V}(\ell,0) := \langle (\psi(\mathbf{r},t) - \psi(\mathbf{r}',t))^2 \rangle_{|\mathbf{r}-\mathbf{r}'|=\ell}$ can be explicitly computed in the range $\max(a, \sqrt{DT}) \ll l \ll l^*$ (where $\ell^* := \sqrt{D/\varkappa}$), and reads (see SI-2.2):

$$
\mathbb{V}(\ell,0) \approx \frac{A^2}{2\pi D} \log \ell - \frac{\Sigma^2}{4\pi D^2} \ell^2 \log \ell + C,\tag{3}
$$

where C is a constant. Note that the first term is the familiar logarithmic correlation of the Gaussian free-field in two di-mensions, see e.g. [\[23\]](#page-4-22). For $\ell \gtrsim \ell^*$, the variogram reaches a plateau value.

Similarly, the temporal variogram $\mathbb{V}(0, \tau) := \langle (\psi(\mathbf{r}, t) \psi(\mathbf{r}, t + \tau)$ ² can be computed, but the final expression is cumbersome and depends on the relative position of three time scales: x^{-1} , the correlation time T and the typical diffusion time $S = a^2/D$ over length scale a, see SI-2.3. There are typically four regimes, a short time regime where $\mathbb{V}(0, \tau) \propto$ τ^2 that reads

$$
\mathbb{V}(0,\tau) = \frac{A^2}{16\pi D} \log \left(\frac{1 + \frac{T}{S}}{1 + \varkappa T} \right) \frac{\tau^2}{T^2}, \quad \tau \ll T, S \quad (4)
$$

followed by two intermediate regimes where $\mathbb{V}(0, \tau) \propto \tau$ and $\log \tau$, and finally a saturated regime for $\varkappa \tau \gg 1$.

In the next sections, we will compare these predictions to empirical data, with good overall agreement. We will find that the spatial variogram is well described by a pure logarithm, i.e. the first term of Eq. (3) – this allows us to determine the ratio A^2/D . With the same value of A^2/D , we then fit the temporal variogram with reasonable values of T and S .

We conducted extensive empirical analyses based on two data sources. The first one is accessible online via the DVF (Demande de Valeur Foncière) website, and displays every housing market transaction in France between 2018 and 2022. This data include the price of the property, its surface and its spatial coordinates. This allows us to study both transaction prices and prices per square meter, up to the granularity of a given point in space. The second data source comes from [\[8\]](#page-4-9), where the authors compiled a wealth of socio-economic

indicators, spanning from [1](#page-2-0)970 to $2022¹$, including housing market prices, but the dataset only contains average transaction prices per *communes* in France up to 2022 and average prices per squared meter per *communes* from 2014 to 2022. [2](#page-2-1) Even though the second data source is less granular than the DVF dataset, its time span of 52 years allows us to investigate the temporal variogram of prices, see below. (The DVF data only span 5 years, which will turn out to be of the same order of magnitude as the correlation time T of the noise). For empirical findings on prices per square meters from DVF, see SI-4.

We first show a color map of transaction log-prices $p :=$ $log P$ across France in Figure [\(1\)](#page-0-0), sourced from [\[8\]](#page-4-9), to compare the spatial distribution of prices in France over the past five decades, a key aspect of our investigation. Indeed, one can see that the price distribution in France is far from uniform, and reveals spatial diffusion around big cities, coastal regions or ski resorts.

Then, it is interesting to study the distribution of *individual* transaction log-prices p, unconditionally over the whole of France. Using the DVF data base, we find that the distribution of prices has a double hump shape, probably reflecting the superposition of two different price distributions for cities and for the countryside, see Fig. [2.](#page-2-2) We show in SI-4, Fig. [6](#page-11-0) a comparison between the distribution of prices in the *département* of la Creuse (chosen to represent a typical countryside district) and in Paris, highlighting the mixture of two distributions seen in the global price distribution for the whole of France. The tail of the distribution of the transaction prices decays as $P^{-1-\mu}$ with $\mu \approx 1.5$, implying that the variance of the transaction prices is mathematically infinite. This should be compared to the Pareto tail of the wealth distribution in France, which decays with a similar exponent [\[24\]](#page-4-23). The distribution of prices per square meters does not have the same shape, but has again a similar power-law tail, as shown in SI-4, Fig. [7.](#page-12-0)

We now shift our focus to the spatial correlations of the logarithm of prices, which we characterize by the equal-time variogram $\mathbb{V}(\ell, 0)$ defined above. The square-root of this quantity measures how different the (log-)prices are when considering two properties a distance ℓ away.^{[3](#page-2-3)} We studied this quantity inside cities, *departements ´* , *regions ´* and the whole of France, with a different coarse-graining scale for the elementary cells over which we average the transaction prices P in order to define the log-price field $p(\mathbf{r})$. We choose hexagonal cells of area 0.73 km^2 for the 17 cities considered,^{[4](#page-2-4)} 5 km² for *départements*, 30 km² for *régions*, and 250 km² for France. The results are shown in Fig. [3.](#page-3-0) At all scales, we

Figure 2. Distribution of all transaction log-prices $p := \log P$, for the 5 years of DVF data. Note the double hump shape, reflecting a mixture of two distributions, corresponding to prices in cities and prices in the countryside. The right tail, for property prices above 500, 000 Euros, corresponds to a power-law tail for prices as $P^{-1-\mu}$ with $\mu \approx 1.5$.

observe a logarithmic dependence on ℓ , provided ℓ is smaller than the size of sector considered (see further down). Furthermore, the slope predicted by Eq. [\(3\)](#page-1-1) is the same at all scales and equal to $A^2/2\pi D \approx 0.19$. The measured (log-)slopes of the variograms are extremely stable over the period 2018-2022 spanned by the DVF data. The other data source [\[8\]](#page-4-9) allows one to measure the spatial variogram over a much longer history. However, the data collection and averaging procedures used in [\[8\]](#page-4-9) seem to induce distortions in the price variograms when compared to the raw DVF data, that we do not fully understand. Still, the analysis of these variograms reveals that the slope of the short-distance logarithmic behaviour is only weakly time dependent, before reaching a plateau value for $\ell \approx 70$ km in 1970 and 300 km nowadays, as seen in SI-4, Fig. [8.](#page-12-1) A possible interpretation is that this crossover length is set by $\ell^* = \sqrt{D/\varkappa}$ which has increased with time, either because D has increased (faster spatial propagation of price changes) or because \varkappa has decreased, reflecting larger wealth inequalities that allows for larger price dispersion, or both.

A reasonable value for D is $-$ say $-$ 50 km²/year, corresponding to prices adapting to a local shock on a scale of 7 km after a year. This leads to a value of $A^2 \approx 2\pi \times 0.19D \sim 60$ km²/year. We will comment on this value below, after having discovered that the noise amplitude A^2 is in fact space dependent.

The reader must have noticed that although the *slopes* of the variograms are the same at all scales, they are shifted up and down in the y-direction. This is expected if one accounts for measurement noise. Indeed, the "true" price field $p(\mathbf{r}, t)$ is approximated here by an empirical average over the chosen cells of transaction prices. The larger the cell size and the smaller the dispersion of prices within each cell, the smaller such idiosyncratic contributions to the difference of prices for

¹ For the specific case of the housing market. Other socio-economic indicators cover an even longer time span. We in fact found similar logarithmic correlations for, e.g., the alphabetization rate in France.

² The housing market data compiled by [\[8\]](#page-4-9) for the years 2014-2022 comes from the DVF database, and is averaged per *communes*.

³ The spatial structure of transaction prices per square meters is investigated in SI-4, Fig. [5.](#page-11-1)

⁴ This leads, for instance, to the division of Paris into 185 neighborhoods.

Figure 3. Spatial variogram for the log price field $p(\mathbf{r})$ averaged over the period 2018-2022 for France as a whole, its *régions*, *départements* and cities, with their respective cross-sectional variability highlighted in shaded colors and the averages for each scale as filled circles. The black dashed lines have a slope equal to 0.19 for all scales, corresponding to $A^2/D = 1.2$. The different off-sets in the y direction corresponds to the measurement noise contribution to the empirical field $p(\mathbf{r})$.

two neighbouring cells.

Finally, note that the spatial variograms do not seem to reveal any departure from the $\log \ell$ behaviour predicted by the first term of Eq. [\(3\)](#page-1-1), except at large distances where finite size and boundary effects start playing a role. Comparing the two terms of Eq. [\(3\)](#page-1-1), one concludes that the second term remains negligible provided $\ell \leq D/\Sigma$. Choosing $D = 50 \text{ km}^2/\text{year}$, and assuming that idiosyncratic effects lead to persistent differential of price variations of at most 10% /year over 1 km, one finds $D/\Sigma \sim 500$ km. This justifies why one may safely neglect the second term in Eq. [\(3\)](#page-1-1).

Turning to the temporal variogram of prices, there are two different empirical definitions for such an object, which should lead to similar results if the system is (statistically) spatially homogeneous. One $(\mathbb{V}_1(\tau))$ is to compute the temporal variance of local price changes $p(\mathbf{r}, t) - p(\mathbf{r}, t + \tau)$ over the full time period, which is then averaged over r. The second $(\mathbb{V}_2(\tau))$ is to remove from $p(\mathbf{r}, t)$ the spatial average of the log-price at time t, i.e. $\bar{p}(t) = \langle p(\mathbf{r}, t) \rangle_{\mathbf{r}}$, and then compute the average of $[p(\mathbf{r}, t) - \bar{p}(t) - (p(\mathbf{r}, t + \tau) - \bar{p}(t + \tau))]^2$ over both t and r . For a statistically homogeneous system, these two procedures lead to comparable results. However, as shown in Fig. [4,](#page-3-1) our data reveals strong differences between $\mathbb{V}_1(\tau)$ and $\mathbb{V}_2(\tau)$, which can be accounted for by assuming that the variance A^2 of the driving noise η is space dependent: $A^2 \rightarrow A^2(\mathbf{r})$. In this case, spatial correlations lose their translation invariance but if one insists on computing them as a function of $\ell = |\mathbf{r} - \mathbf{r}'|$, one recovers Eq. [\(3\)](#page-1-1) with A^2 replaced by its spatial average $\langle A^2 \rangle_r$, see SI-4, Fig. [9.](#page-13-0)

Now, it turns out that in the presence of spatial heterogeneities, the temporal variogram $\mathbb{V}_1(\tau)$ is also given by

Figure 4. Comparison between $\mathbb{V}_1(\tau)$ and $\mathbb{V}_2(\tau)$ for the empirical data, in a log-log representation. We also show (in red) the fit found for $\mathbb{V}_1(\tau)$ with our theoretical equation. Note that the short time behaviour of $\mathbb{V}_1(\tau)$ is in-between τ and τ^2 , indicating a non-zero correlation time T. We find $T = 3.5$ years, $S = 1$ year, $D = 50$ km² per year and $A^2/D \approx 1.2$. The observed shift between $\mathbb{V}_1(\tau)$ and $\mathbb{V}_2(\tau)$ is a consequence of strong spatial heterogeneities, see SI-4, Fig. [9.](#page-13-0) Note that with 50 years of data, only the first 10 years of lags are reliable.

Eq. [\(4\)](#page-1-2) with $A^2 \rightarrow \langle A^2 \rangle_r$, see SI-4, Fig. [9.](#page-13-0) Hence we focus our attention to $\mathbb{V}_1(\tau)$ and attempt to fit it with our theoretical formula (see SI-2.3) with T, S as adjustable parameters, with $\langle A^2 \rangle_{\mathbf{r}}/D$ fixed and set to 1.2, close to the value inferred from spatial variograms. $(D$ itself has negligible influence on the goodness-of-fit). The optimal values are then found to be $S = 1$ year, corresponding to a correlation length for shocks $a = \sqrt{DS} = 7$ km, and a correlation time of $T = 3.5$ years, such that $\sqrt{DT} = 13$ km. The order of magnitude of A^2 is expected to be $a^2/T \sim 30 \text{ km}^2/\text{year}$, a factor two times smaller than expected if $D = 50 \text{ km}^2/\text{year}$, but not unreasonable in view of the crudeness of our model and the possibility to change the value of parameters without substantially affecting the joint goodness-of-fit of spatial and temporal variograms. For example, choosing $\langle A^2 \rangle_{\mathbf{r}}/D = 1.3$ leads to $T = S = 2.5$ years and in this case $a^2/T \sim 50$ km²/year. Note that the short-time regime of $\mathbb{V}_1(\tau)$ is a sign that price changes are persistent, which is inconsistent with the hypothesis that the housing market is "efficient" [\[2\]](#page-4-2). In view of the large transaction costs incurred when buying a house, this is hardly surprising.

Finally, in order to account for the empirical difference between the two temporal variograms $\mathbb{V}_1(\tau)$ and $\mathbb{V}_2(\tau)$, one needs to introduce rather strong spatial heterogeneities in the noise amplitude A^2 , that must vary by a factor of 10 depending on the considered region, see SI-4, Fig. [9.](#page-13-0) This is not very surprising in view of the very different structure of the housing market in international cities like Paris or Nice and the remote, low density regions like Lozère. An generalized version of our model, Eq. [\(1\)](#page-1-0), that properly accounts for geographical heterogeneities that make both D and A^2 space dependent, would however require a different, much more granular calibration strategy.

In conclusion, we have shown that housing prices in France reveal clear, robust statistical regularities. Such regularities are expected if the dynamics of prices is diffusive, that is, the spatial variogram of prices has a logarithmic dependence on distance. Indeed this is a signature of two-dimensional diffusing fields driven by random noise, captured by our stylized model, Eq. [\(1\)](#page-1-0), which was already used in the past to model spatial regularities in voting patterns [\[5,](#page-4-5) [6\]](#page-4-18). Note that a model where prices propagate in a ballistic way ($r \sim t$) instead of diffusing $(r \sim \sqrt{t})$ would lead to completely different spatial correlations. The temporal fluctuations of prices can be accounted for within the same framework, provided the shocks are persistent over a time scale that we find to be around 3 years. The data also suggests, not surprisingly, that the amplitude of the price shocks is spatially heterogeneous, with a large variation span. All the dimensional parameters obtained from fitting the spatial and temporal correlations appear to be of reasonable order of magnitude.

Our study thus confirms and quantifies the diffusive nature of housing prices that was anticipated long ago [\[1,](#page-4-1) [2\]](#page-4-2), albeit on more restricted, local data sets. Case studies, like the opening of a TGV (*Train à Grande Vitesse*) railway station, or of a new metro line that are expected to boost nearby housing prices, would be quite interesting as independent validations of the model proposed in this paper. Future work should attempt couple the random diffusion equation for prices to the population field in order to describe social mobility, as a twofield extension of our previous work [\[25\]](#page-4-24). Extending our analysis to other spatial socio-economic variables would also shed light on the mechanisms underlying diffusion of socio-cultural traits, as suggested in [\[22\]](#page-4-21).

Acknowledgements We thank Xavier Gabaix, Swann Chelly, Nirbhay Patil and Max Sina Knicker for fruitful comments and discussions. We also thank Thomas Piketty for useful explanations about how the data published in [\[8\]](#page-4-9) was created. This research was conducted within the Econophysics & Complex Systems Research Chair, under the aegis of the Fondation du Risque, the Fondation de l'École polytechnique, the Ecole polytechnique and Capital Fund Management. ´

∗ antoine-cyrus.becharat@polytechnique.edu

- [1] J. M. Clapp and D. Tirtiroglu, Positive feedback trading and diffusion of asset price changes: Evidence from housing transactions, [Journal of Economic Behavior and Organization](https://doi.org/https://doi.org/10.1016/0167-2681(94)90040-X) **24**, 337 [\(1994\).](https://doi.org/https://doi.org/10.1016/0167-2681(94)90040-X)
- [2] H. O. Pollakowski and T. S. Ray, Housing price diffusion patterns at different aggregation levels: An examination of housing market efficiency, [Journal of Housing Research](http://www.jstor.org/stable/24833634) **8**, 107 (1997).
- [3] A.-L. Barabási and H. E. Stanley, *Fractal concepts in surface growth* (Cambridge university press, 1995).
- [4] U. Frisch, *Turbulence: the legacy of AN Kolmogorov* (Cambridge university press, 1995).
- [5] C. Borghesi and J.-P. Bouchaud, Spatial correlations in vote statistics: a diffusive field model for decision-making [10.1140/epjb/e2010-00151-1](https://doi.org/10.1140/epjb/e2010-00151-1) (2010).
- [6] C. Borghesi, J.-C. Raynal, and J.-P. Bouchaud, Election turnout statistics in many countries: similarities, differences, and a diffusive field model for decision-making, PloS one **7**, e36289 (2012).
- [7] J. Fernández-Gracia, K. Suchecki, J. J. Ramasco, M. San Miguel, and V. M. Eguíluz, Is the voter model a model for voters?, Physical review letters **112**, 158701 (2014).
- [8] J. Cagé and T. Piketty, *Une histoire du conflit politique*. *Elections et in ´ egalit ´ es sociales en France, 1789-2022. ´* (Le Seuil, 2023).
- [9] J. Geanakoplos, R. Axtell, D. J. Farmer, P. Howitt, B. Conlee, J. Goldstein, M. Hendrey, N. M. Palmer, and C.-Y. Yang, Getting at systemic risk via an agent-based model of the housing market, American Economic Review **102**, 53 (2012).
- [10] S. Rosen, Hedonic prices and implicit markets: Product differentiation in pure competition.
- [11] T. Besley and H. Mueller, Estimating the peace dividend: The impact of violence on house prices in northern ireland, [Ameri](https://doi.org/10.1257/aer.102.2.810)[can Economic Review](https://doi.org/10.1257/aer.102.2.810) **102**, 810–33 (2012).
- [12] D. N. Figlio and M. E. Lucas, What's in a grade? school report cards and the housing market, [American Economic Review](https://doi.org/10.1257/0002828041464489) **94**, [591–604 \(2004\).](https://doi.org/10.1257/0002828041464489)
- [13] F. Feitosa and W. Zesk, [Spatial patterns of residential segrega](https://www.researchgate.net/publication/220913299)[tion: A generative model](https://www.researchgate.net/publication/220913299) (2008).
- [14] M. Pangallo, J.-P. Nadal, and A. Vignes, Residential income segregation: A behavioral model of the housing market, [Jour](https://doi.org/https://doi.org/10.1016/j.jebo.2019.01.010)[nal of Economic Behavior and Organization](https://doi.org/https://doi.org/10.1016/j.jebo.2019.01.010) **159**, 15 (2019).
- [15] R. Baptista, J. D. Farmer, M. Hinterschweiger, K. Low, D. Tang, and A. Uluc, [Staff working paper no. 619 macroprudential pol](http://www.bankofengland.co.uk/pra/Documents/publications/ps/2014/ps914.pdf)[icy in an agent-based model of the uk housing market](http://www.bankofengland.co.uk/pra/Documents/publications/ps/2014/ps914.pdf) (2016).
- [16] L. Gauvin, A. Vignes, and J.-P. Nadal, Modeling urban housing market dynamics: Can the socio-spatial segregation preserve some social diversity?, [Journal of Economic Dynamics](https://doi.org/10.1016/j.jedc.2013.03.001) and Control **37**[, 1300 \(2013\).](https://doi.org/10.1016/j.jedc.2013.03.001)
- [17] P. A. Samuelson, Proof that properly anticipated prices fluctuate randomly, in *The world scientific handbook of futures markets* (World Scientific, 2016) pp. 25–38.
- [18] S. Basu and T. Thibodeau, Analysis of spatial autocorrelation in house prices, [The Journal of Real Estate Finance and Economics](https://EconPapers.repec.org/RePEc:kap:jrefec:v:17:y:1998:i:1:p:61-85) **17**[, 61 \(1998\).](https://EconPapers.repec.org/RePEc:kap:jrefec:v:17:y:1998:i:1:p:61-85)
- [19] M. Kelly, *The Standard Errors of Persistence*, Working Paper WP19/13 (University College Dublin, UCD School of Economics, Dublin, 2019).
- [20] F. Schweitzer and J. A. Hołyst, Modelling collective opinion formation by means of active brownian particles, The European Physical Journal B-Condensed Matter and Complex Systems **15**, 723 (2000).
- [21] F. Schweitzer, Coordination of decisions in a spatial model of brownian agents, in *The Complex Dynamics of Economic Interaction: Essays in Economics and Econophysics* (Springer, 2004) pp. 303–318.
- [22] J.-P. Bouchaud, C. Borghesi, and P. Jensen, On the emergence of an 'intention field'for socially cohesive agents, Journal of Statistical Mechanics: Theory and Experiment **2014**, P03010 (2014).
- [23] S. F. Edwards and D. Wilkinson, The surface statistics of a granular aggregate, Proceedings of the Royal Society of London. A. Mathematical and Physical Sciences **381**, 17 (1982).
- [24] S. Bach, A. Thiemann, and A. Zucco, The top tail of the wealth distribution in germany, france, spain, and greece, (2015).
- [25] R. Zakine, J. Garnier-Brun, A.-C. Becharat, and M. Benzaquen, Socioeconomic agents as active matter in nonequilibrium

sakoda-schelling models, Phys. Rev. E **109**[, 044310 \(2024\).](https://doi.org/10.1103/PhysRevE.109.044310)

SI-1: Analytical derivation of the diffusive term

We assume that the diffusive term in the price field evolves through a mechanism of supply and demand such that the time evolution of the field ψ depends on the difference of the field between two locations $\psi(R_{\alpha}) - \psi(R_{\beta})$ where R_{α} and R_{β} refer to the considered locations. We then propose the following generic equation to describe the propagation of the field with respect to its surrounding influences:

$$
\partial_t \psi(R_\alpha, t) = \sum_{\beta} \Gamma_{\alpha, \beta} \psi(R_\beta) - \sum_{\beta} \Gamma_{\beta, \alpha} \psi(R_\alpha), \tag{5}
$$

where Γ is a symmetric influence matrix such that:

$$
\Gamma_{\alpha,\beta} = \Gamma(R_{\alpha}|R_{\beta}) = t(R_{\alpha} - R_{\beta}|R_{\beta}).
$$
\n(6)

Hence, in the continuous limit and in one dimension for simplicity, it comes:

$$
\partial_t \psi(x,t) = \int t(x-x'|x')\psi(x',t)dx' - \int t(x'-x|x)\psi(x,t)dx',\tag{7}
$$

which we can re write as:

$$
\partial_t \psi(x,t) = \int t(y|x-y)\psi(x-y,t)dy - \int t(y|x)\psi(x,t)dy,
$$
\n(8)

changing variables to $y = x - x'$. The Kramers-Moyal expansion of [\(8\)](#page-6-0) up to the order 2 in y then gives:

$$
\partial_t \psi(x,t) = -\partial_x \left[R_1(x)\psi(x) \right] + \frac{1}{2}\partial_x^2 \left[R_2(x)\psi(x) \right],\tag{9}
$$

where:

$$
R_1(x) = \int yt(y, x)dy;
$$
\n(10)

$$
R_2(x) = \int y^2 t(y, x) dy.
$$
\n⁽¹¹⁾

Moreover, the influence matrix is symmetric, hence the drift term $R_1(x)$ is set to zero and we retrieve the one dimensional diffusion equation:

$$
\partial_t \psi(x,t) = \partial_x^2 \left[D(x)\psi(x) \right] \tag{12}
$$

with $D(x) = \frac{1}{2} \int y^2 t(y, x) dy$. Note that we retrieve here a non-uniform diffusion coefficient, but we assume in the rest of the study that we can take $D(x) = D$.

SI-2: Theoretical predictions for the variograms

SI-2.1: Computation of the generic space-time variogram

Let us consider the following stochastic partial differential equation:

$$
\frac{\partial \psi(\mathbf{r}, \mathbf{t})}{\partial t} = D \Delta \psi(\mathbf{r}, \mathbf{t}) - \varkappa \psi(\mathbf{r}, \mathbf{t}) + \eta(\mathbf{r}, \mathbf{t}) + \xi(\mathbf{r}), \tag{13}
$$

where Δ is the Laplacian operator, D a diffusion coefficient, \varkappa a mean-reversion coefficient, $\eta(\mathbf{r}, \mathbf{t})$ a Langevin noise with zero mean and short range time and space correlations, and $\xi(\mathbf{r})$ a static random field with zero mean and short range correlations. The correlators of these terms are assumed to be of the following type:

$$
\langle \eta(\mathbf{r}, \mathbf{t}) \eta(\mathbf{r}', \mathbf{t}') \rangle = \frac{A^2}{Ta^2} e^{-|t - t'|/T} g_a(|\mathbf{r} - \mathbf{r}'|);
$$

$$
\langle \xi(\mathbf{r}) \xi(\mathbf{r}') \rangle = \frac{\Sigma^2}{a^2} g_a(|\mathbf{r} - \mathbf{r}'|),
$$
 (14)

8

where $g_a(r)$ is a bell-shaped function that decays over length scale a, such that $2\pi \int_{r\geq 0} g_a(r) r dr = a^2$. For the rest of the calculations, we consider the regime where $|\mathbf{r} - \mathbf{r}'| = \ell \gg a$ which leads to $\frac{1}{a^2} g_a(|\mathbf{r} - \mathbf{r}'|) \approx \delta(|\mathbf{r} - \mathbf{r}'|)$. Moreover, the space time correlation function can be written as:

$$
\mathbb{C}(|\mathbf{r} - \mathbf{r}'|, |t - t'|) = \langle \psi(\mathbf{r}, t)\psi(\mathbf{r}', t')\rangle = \int \int e^{-i\mathbf{k}\mathbf{r} - i\mathbf{k}'\mathbf{r}'} \langle \psi_{\mathbf{k}}(t)\psi_{\mathbf{k}'}(t')\rangle \frac{d\mathbf{k}}{(2\pi)^2} \frac{d\mathbf{k}'}{(2\pi)^2},
$$
(15)

where $\psi_{\mathbf{k}}$ is the solution of the following equation in Fourier space:

$$
\frac{\partial \psi_{\mathbf{k}}(t)}{\partial t} = -D\mathbf{k}^{2}\psi_{\mathbf{k}}(t) - \varkappa\psi_{\mathbf{k}}(t) + \eta_{\mathbf{k}} + \xi_{\mathbf{k}}.
$$
\n(16)

Hence:

$$
\psi_{\mathbf{k}}(t) = \psi_{\mathbf{k}}(0)e^{-(D\mathbf{k}^2 + \varkappa)t} + \int_0^t e^{-(D\mathbf{k}^2 + \varkappa)(t-\tau)} (\eta_{\mathbf{k}}(\tau) + \xi_{\mathbf{k}})d\tau.
$$
 (17)

Because of the two fields η and ξ - assumed to be independent - we will separate the calculation for the correlation function into two contributions. In the long time limit, the first contribution in Fourier space, coming from field η , is:

$$
\int_0^t \int_0^{t'} dt_1 dt_2 e^{-(D\mathbf{k}^2 + \varkappa)(t - t_1) - (D\mathbf{k}'^2 + \varkappa)(t' - t_2)} \langle \eta_{\mathbf{k}}(t_1) \eta_{\mathbf{k}'}(t_2) \rangle,
$$
\n(18)

leading to:

$$
\frac{A^2(2\pi)^2}{T} \int_0^t \int_0^{t'} dt_1 dt_2 e^{-(D\mathbf{k}^2 + \varkappa)(t - t_1) - (D\mathbf{k}'^2 + \varkappa)(t' - t_2)} e^{-\frac{|t_1 - t_2|}{T}} \delta(\mathbf{k} + \mathbf{k}'). \tag{19}
$$

We find, in the long time limit, that the integral yields in Fourier space:

$$
\frac{A^2 (2\pi)^2}{2T} \left[\frac{e^{-(D\mathbf{k}^2 + \varkappa)|t'-t|}}{2(D\mathbf{k}^2 + \varkappa)(D\mathbf{k}^2 + \varkappa + \frac{1}{T})} + \frac{e^{-\frac{|\mathbf{k}-t'|}{T}}}{(D\mathbf{k}^2 + \varkappa)^2 - \frac{1}{T^2}} - \frac{e^{-(D\mathbf{k}^2 + \varkappa)|t'-t|}}{2(D\mathbf{k}^2 + \varkappa)(D\mathbf{k}^2 + \varkappa - \frac{1}{T})} \right].
$$
 (20)

This can be condensed as:

$$
\frac{A^2(2\pi)^2}{2T((D\mathbf{k}^2+\varkappa)^2-\frac{1}{T^2})}\left[e^{-\frac{|t-t'|}{T}}-\frac{e^{-(D\mathbf{k}^2+\varkappa)|t'-t|}}{T(D\mathbf{k}^2+\varkappa)}\right].
$$
\n(21)

Similarly, we can compute the contribution for the correlation function coming from field $\xi(\mathbf{r})$:

$$
(2\pi)^2 \Sigma^2 \int_0^t \int_0^{t'} dt_1 dt_2 e^{-(D\mathbf{k}^2 + \varkappa)(t - t_1) - (D\mathbf{k}'^2 + \varkappa)(t' - t_2)} \delta(\mathbf{k} + \mathbf{k}'). \tag{22}
$$

This yields, in the long time limit:

$$
\frac{(2\pi)^2 \Sigma^2}{(D\mathbf{k}^2 + \varkappa)^2}.
$$
\n(23)

In the next sections, we will show how, starting from what has just been shown, we compute both the spatial and the temporal variograms, defined as $\mathbb{V}(\ell,0) := \langle (\psi(\mathbf{r},t) - \psi(\mathbf{r}',t))^2 \rangle$ and $\mathbb{V}(0,\tau) := \langle (\psi(\mathbf{r},t) - \psi(\mathbf{r},t+\tau))^2 \rangle$.

SI-2.2: Computation of the spatial variogram

We come back to the first contribution (coming from field η) in Fourier space for the space time correlation function:

$$
\frac{A^2(2\pi)^2}{2T((Dk^2+\varkappa)^2-\frac{1}{T^2})}\left[e^{-\frac{|t-t'|}{T}}-\frac{e^{-(Dk^2+\varkappa)}|t'-t|}{T(Dk^2+\varkappa)}\right].
$$
\n(24)

We now focus on the static behavior of this term, hence imposing $t = t'$. This yields:

$$
\frac{A^2(2\pi)^2}{2T((D\mathbf{k}^2+\varkappa)^2-\frac{1}{T^2})}\left[1-\frac{1}{T(D\mathbf{k}^2+\varkappa)}\right].\tag{25}
$$

Using notations $|\mathbf{k}| = k$, $\mathbf{k} \cdot (\mathbf{r} - \mathbf{r}') = k\ell \cos(\theta)$ and notation \mathbb{C}_{η} to describe the contribution from η to the correlation function, it comes in polar coordinates:

$$
\mathbb{C}_{\eta}(\ell,0) = \frac{A^2}{2T(2\pi)^2} \int dk \int d\theta e^{-ik\ell\cos(\theta)} \frac{k}{((Dk^2 + \varkappa)^2 - \frac{1}{T^2})} \left[1 - \frac{1}{T(Dk^2 + \varkappa)}\right].
$$
 (26)

The integral is defined for $1/\ell^* \ll k \ll 1/a$, which ensures that $Dk^2 \gg \frac{D}{\ell^{*2}} = \varkappa$. We can hence neglect the mean-reversion term in the computation. Moreover, we can neglect $D^2 k^4$ in favor of $\frac{1}{T^2}$ if $Dk^2 < \frac{1}{T}$, hence if $\ell > \sqrt{DT}$. This is typically the regime that we consider for this study, since we estimate (see in the main text) $\sqrt{DT} \approx 13$ km, so we assume here that this term is negligible. Finally, we can identify the Bessel function

$$
\frac{1}{2\pi} \int_0^{2\pi} d\theta e^{ik\ell \cos(\theta)} = J_0(k\ell) = J_0(-k\ell),\tag{27}
$$

so:

$$
\mathbb{C}_{\eta}(\ell,0) \approx \frac{A^2}{4\pi} \int_{1/\ell^*}^{1/a} dk \frac{J_0(k\ell)}{Dk}.
$$
 (28)

The Bessel function can be expanded for $k\ell \to 0$, and yields $J_0(k\ell) \approx 1 - \ell^2 k^2/4 + o(k^4\ell^4)$. Moreover, the Bessel function decays to zero when $k\ell \gg 1$, concentrating the integral towards its lower bound. This gives, up to constant contributions:

$$
\mathbb{C}_{\eta}(\ell,0) \approx -\frac{A^2}{4\pi D}\log\ell + K(\ell)
$$
\n(29)

with correction term $K(\ell)$. Similarly, we can compute the contribution from field ξ :

$$
\mathbb{C}_{\xi}(\ell,0) = \frac{\Sigma^2}{2\pi D^2} \int_{1/\ell^*}^{1/a} dk \frac{J_0(k\ell)}{k^3} = \frac{\Sigma^2}{2\pi D^2} \ell^2 \int_{\ell/\ell^*}^{\ell/a} du \frac{J_0(u)}{u^3}.
$$
 (30)

In order to have a non-constant contribution here, we must go to the second order in the expansion of the Bessel function towards the lower bound of the integral. This yields:

$$
\mathbb{C}_{\xi}(\ell,0) \approx \frac{\Sigma^2}{2\pi D^2} \ell^2 \int_{\ell/\ell^*}^{\ell/a} du \frac{1 - \frac{u^2}{4}}{u^3},\tag{31}
$$

which finally yields, up to constant terms:

$$
\mathbb{C}_{\xi}(\ell,0) \approx \frac{\Sigma^2}{8\pi D^2} \ell^2 \log \ell + K'(\ell)
$$
\n(32)

with correction $K'(\ell)$. Furthermore, the variogram is defined as $\mathbb{V}(\ell,0) = 2\langle \psi(\mathbf{r},0)^2 \rangle - 2\mathbb{C}(\ell,0)$. Hence, summing both contributions yields:

$$
\mathbb{V}(\ell,0) \approx \frac{A^2}{2\pi D} \log \ell - \frac{\Sigma^2}{4\pi D^2} \ell^2 \log \ell + C,\tag{33}
$$

where C is a constant. This result is of course only valid in the range where $a \ll \ell \ll \ell^*$.

SI-2.3: Computation of the temporal variogram

As we are now interested in the temporal variation of the same point in space, we will neglect the random static field $\xi(\vec{r})$ in the computation which will only yield constant terms. Moreover, we will again neglect the contribution \varkappa in the calculations as the integration back to real space will impose $Dk^2 \gg \kappa$, as seen in the previous section. Our starting point is therefore the following:

$$
\frac{A^2(2\pi)^2}{2T(D^2\mathbf{k}^4 - \frac{1}{T^2})} \left[e^{-\frac{|\mathbf{t} - \mathbf{t}'|}{T}} - \frac{e^{-D\mathbf{k}^2|\mathbf{t}' - \mathbf{t}|}}{TD\mathbf{k}^2} \right].
$$
\n(34)

When
$$
\tau = |t - t'| \gg T
$$

When $\tau = |t - t'| \gg T$, we can set $e^{-\frac{|t - t'|}{T}}$ to zero. Coming back in real space yields:

$$
\mathbb{C}(0, |t - t'|) = -\frac{A^2}{T^2 (2\pi)^2} \int d\mathbf{k} \frac{e^{-D\mathbf{k}^2 |t - t'|}}{2D\mathbf{k}^2 (D^2 \mathbf{k}^4 - \frac{1}{T^2})},
$$
(35)

which gives in polar coordinates:

$$
\mathbb{C}(0,\tau) = -\frac{A^2}{T^2(2\pi)^2} \int dk \int d\theta \frac{k e^{-Dk^2\tau}}{2Dk^2(D^2k^4 - \frac{1}{T^2})}.
$$
\n(36)

It comes:

$$
\mathbb{C}(0,\tau) = -\frac{A^2}{8\pi DT^2} \int_{\frac{D\tau}{\ell^*2}}^{\frac{D\tau}{a^2}} du \frac{e^{-u}}{u(\frac{u^2}{\tau^2} - \frac{1}{T^2})}.
$$
\n(37)

Moreover, $\frac{u}{\tau} < \frac{1}{T}$ if $S = \frac{a^2}{D} > T$, which allows us to neglect this term, leading to:

$$
\mathbb{C}(0,\tau) \approx \frac{A^2}{8\pi D} \int_{\frac{D\tau}{\ell^{*2}}}^{\frac{D\tau}{a^2}} du \frac{e^{-u}}{u}.
$$
\n(38)

Hence, in the regime where $T < S \ll \tau \ll \varkappa^{-1} = \frac{\ell^{*2}}{D}$ $\frac{1}{D}$:

$$
\mathbb{C}(0,\tau) \approx -\frac{A^2}{8\pi D} \log \tau,\tag{39}
$$

up to constant terms. This finally yields:

$$
\mathbb{V}(0,\tau) \approx \frac{A^2}{4\pi D} \log \tau.
$$
\n(40)

When $S \ll T \ll \tau \ll \varkappa^{-1}$, logarithmic contributions can once again be obtained by performing a partial fraction decomposition in [\(37\)](#page-9-0) prior to integration. For completeness, in the regime where $\tau \gg \varkappa^{-1}$, S, T, the computation yields a constant value.

When
$$
\tau = |t - t'| \ll T
$$

We come back to:

$$
\frac{A^2(2\pi)^2}{2T(D^2\mathbf{k}^4 - \frac{1}{T^2})} \left[e^{-\frac{|t - t'|}{T}} - \frac{e^{-D\mathbf{k}^2|t'-t|}}{TD\mathbf{k}^2} \right].
$$
\n(41)

If $\tau \ll S$, we can expand up to the order two in the exponentials for $D\mathbf{k}^2\tau \longrightarrow 0$, in addition to the expansion for $\frac{\tau}{T} \longrightarrow 0$, leading to:

$$
\frac{A^2(2\pi)^2}{2T(D^2\mathbf{k}^4 - \frac{1}{T^2})} \left[\frac{TD\mathbf{k}^2 - 1}{TD\mathbf{k}^2} + \frac{1}{2}(1 - TD\mathbf{k}^2) \frac{\tau^2}{T^2} \right].
$$
\n(42)

Hence, the temporal contribution in the correlation function, coming back to real space, is:

$$
\mathbb{C}(0,\tau) = \frac{1}{2\pi} \int_{\frac{1}{\ell^*}}^{\frac{1}{\alpha}} dk \frac{A^2 k}{2T \left(D^2 k^4 - \frac{1}{T^2}\right)} \frac{1}{2} (1 - T D k^2) \frac{\tau^2}{T^2}.
$$
\n(43)

This yields, after integration and up to constant terms:

$$
\mathbb{C}(0,\tau) \approx \frac{A^2}{32\pi D} \log \left(\frac{\frac{TD}{\ell^{*2}} + 1}{TD/a^2 + 1}\right) \frac{\tau^2}{T^2},\tag{44}
$$

which we can re write as:

$$
\mathbb{C}(0,\tau) \approx -\frac{A^2}{32\pi D} \log \left(\frac{\frac{T}{S} + 1}{\varkappa T + 1}\right) \frac{\tau^2}{T^2}.
$$
\n(45)

This finally yields:

$$
\mathbb{V}(0,\tau) \approx \frac{A^2}{16\pi D} \log \left(\frac{\frac{T}{S} + 1}{\varkappa T + 1}\right) \frac{\tau^2}{T^2}.
$$
\n(46)

If $\tau \geq S$, we cannot expand in the second exponential term of [\(21\)](#page-7-0). This leads us to study separately both terms. The first one will give, after expanding up to the second order in $\frac{\tau}{T}$:

$$
\frac{1}{2\pi} \int kdk \frac{A^2}{2T(D^2k^4 - \frac{1}{T^2})} \left(1 - \frac{\tau}{T} + \frac{\tau^2}{2T^2}\right),\tag{47}
$$

which yields:

$$
\frac{A^2}{32\pi D}\log\left(\frac{\left|\frac{T}{S}-1\right|(\varkappa T+1)}{\left(\frac{T}{S}+1\right)|\varkappa T-1|}\right)\left(1-\frac{\tau}{T}+\frac{\tau^2}{2T^2}\right).
$$
\n(48)

The second term:

$$
-\frac{1}{2\pi} \frac{A^2}{2T(D^2\mathbf{k}^4 - \frac{1}{T^2})} \frac{e^{-D\mathbf{k}^2|t'-t|}}{TD\mathbf{k}^2}
$$
(49)

will give:

$$
-\frac{A^2}{8\pi T^2 D} \int_{\frac{1}{\ell^*}}^{\frac{1}{a}} dk \frac{e^{-Dk^2 \tau}}{k(Dk^2 - \frac{1}{T})(Dk^2 + \frac{1}{T})}.
$$
\n(50)

Changing variables to $u = Dk^2 \tau$ yields, after a few integration steps:

$$
-\frac{A^2}{32\pi D} \left[e^{\tau/T} \log \left(\frac{\left|\frac{T}{S} - 1\right|}{\left|\varkappa T - 1\right|} \right) + e^{-\tau/T} \log \left(\frac{\frac{T}{S} + 1}{\varkappa T + 1} \right) - 2 \log \left(\varkappa S \right) \right],\tag{51}
$$

which gives, after expanding the two exponentials $e^{\tau/T}$ and $e^{-\tau/T}$ up to the order two in $\frac{\tau}{T}$:

$$
-\frac{A^2}{32\pi D}\log\left(\frac{\left|\frac{T}{S}-1\right|(\varkappa T+1)}{\left(\frac{T}{S}+1\right)|\varkappa T-1|}\right)\frac{\tau}{T}-\frac{A^2}{64\pi D}\log\left(\frac{\left|\frac{T^2}{S^2}-1\right|}{\left|\varkappa^2T^2-1\right|}\right)\frac{\tau^2}{T^2}.\tag{52}
$$

This finally yields, after adding the first and second term contribution from [\(21\)](#page-7-0):

$$
\mathbb{V}(0,\tau) \approx \frac{A^2}{8\pi D} \log \left(\frac{\left|\frac{T}{S} - 1\right| (\varkappa T + 1)}{\left(\frac{T}{S} + 1\right) |\varkappa T - 1|} \right) \frac{\tau}{T} + \frac{A^2}{16\pi D} \log \left(\frac{\frac{T}{S} + 1}{\varkappa T + 1} \right) \frac{\tau^2}{T^2}.
$$
\n
$$
\tag{53}
$$

We hence lose the quadratic behavior for the variogram when $S \leq \tau \ll T$ and the dominant behavior becomes linear.

Figure 5. Spatial variogram for the log-price field per squared meter $\hat{p}(\mathbf{r}) := \log(\hat{P})$, where the notation \hat{P} indicates the prices per squared meter, averaged over the period 2018-2022 for France as a whole, its *régions*, *départements* and cities, with their respective cross-sectional variability highlighted in shaded colors and the averages for each scale as filled circles. The different off-sets in the y direction corresponds to the measurement noise contribution to the empirical field $\hat{p}(\mathbf{r})$. The observed empirical behavior is once again logarithmic.

Figure 6. Distribution of all transaction log-prices $p := \log P$, averaged for the 5 years of DVF data, both for the *département* of la Creuse and for Paris. These locations were chosen as typical examples of both the countryside and cities, showing clearly two different shapes. This explains the double-hump nature of the global log-price distribution for the whole of France, discussed in the main text.

Figure 7. Distribution of all transaction log-prices per squared meter \hat{p} , for the 5 years of DVF data. The right tail corresponds to a power-law tail for prices per squared meter as $\hat{P}^{-1-\mu}$ with $\mu \approx 1.4$, close to 1.5, as found for the log-prices above.

Figure 8. Spatial variograms for the log price field $p(r)$ for every year between 1970 to 2022, using the data from [\[8\]](#page-4-9). We see that the slope of these variograms is only weakly time-dependent, and that the logarithmic behavior is robust in time. The variogram reaches a plateau for $\ell \approx 70$ km in 1970 and for $\ell \approx 300$ km in 2022.

Figure 9. Theoretical predictions for the spatial variogram, computed when the noise amplitude is uniform, equal to A^2 , and when the noise amplitude $A^2(\mathbf{r})$ is strongly heterogeneous, with $\langle A^2(\mathbf{r}) \rangle = A^2 = 2\pi D \times 0.19$. We obtain a similar logarithmic behavior in both cases. The inset shows a comparison between $\mathbb{V}_1(\tau)$ and $\mathbb{V}_2(\tau)$ computed for data simulated on a lattice with the same strongly heterogeneous noise amplitude $A^2(\mathbf{r})$. We hence qualitatively retrieve the observed empirical temporal behavior.