

Stability of chiral magnon condensate in antiferromagnetic insulators

Therese Frostad,^{1,*} Anne Louise Kristoffersen,^{1,*} Verena Brehm,¹
Roberto E. Troncoso,² Arne Brataas,¹ and Alireza Qaiumzadeh¹

¹*Center for Quantum Spintronic, Department of Physics,
Norwegian University of Science and Technology, NO-7491 Trondheim, Norway*
²*School of Engineering and Sciences, Universidad Adolfo Ibáñez, Santiago, Chile*

(Dated: December 20, 2024)

Magnon Bose-Einstein condensates in ferromagnetic insulators [1] has been a field of much interest, while condensation in anti-ferromagnetic systems are yet to be explored in detail. We analyze the stability of condensed chiral magnons in two antiferromagnetic insulators: an easy-axis system and a biaxial system. We show that two-component magnon condensation and inter-magnon interactions are essential to create metastable magnon condensation. The uniaxial system with a Rashba-type Dzyaloshinskii–Moriya interaction supports two condensate populations at finite wavevectors. The condensation state is stable only when the distribution of magnons between the two populations is symmetric. On the other hand, in the biaxial system without Dzyaloshinskii–Moriya interaction we predict that the magnons condensate will not be stabilized. The results indicate that the lack of stability is a general feature of single-component quasiparticle condensates.

I. INTRODUCTION

Bose-Einstein condensation (BEC) is a celebrated phenomenon in quantum physics describing a phase transition where bosonic particles form an equilibrium macroscopic state and behave collectively as a single quantum entity characterized by a coherent wave function [2–10]. This phase transition often requires temperatures close to absolute zero. When the temperature decreases, the thermal wavelength of the particles increases, leading to a macroscopic occupation of the lowest quantum state. Although BEC was initially observed in dilute atomic gases, the phenomenon has been extended to incorporate condensation processes of emerging bosonic quasiparticles in various condensed matter systems. Exploring *quasi-equilibrium* BEC states of quasiparticles, such as excitons [11], phonons [12], and magnons [1], opens up new avenues for studying quantum phenomena in diverse physical systems, from superfluidity to novel states of matter with potential application in quantum technologies.

In 2006, Demokrotov et al. observed a room-temperature magnon BEC (mBEC) in a ferromagnetic (FM) film of yttrium iron garnet (YIG) at room temperature [1]. In their experimental setup, non-equilibrium magnons were excited under microwave parallel pumping, and the magnon condensate was investigated using Brillouin light scattering. This prompted a discussion of the formation, coherence, stabilization, and lifetime of mBEC states in FM systems [13–27]. Most of the experimental studies focus on mBEC in FM materials generated parametric pumping. However, recent experiments also successfully generated mBEC electrically via the spin-transfer torque mechanism [28], and thermally via a rapid cooling process [29].

The magnon spectra of the thin film YIG have double degenerate minima at finite momenta. This results in a nonuniform condensate ground state in real space, which as been confirmed by experimental measurements [19]. The phase and magnon distribution between the two condensates, along with the stability of the condensate, are influenced by nonlinear magnon interactions [20, 24, 30, 31].

The double condensate mBEC, formed in two different energy minima of the magnon system, can be characterized by the number of magnons in each of the two mBECs and their relative phase. These two parameters have been suggested as the basis for a classical analog of a qubit logic [32, 33]. The possibility of Josephson oscillations and persists spin currents in mBEC systems has also been proposed theoretically, but the phenomena are not yet confirmed experimentally [22, 34].

While magnon condensation in ferromagnets has been studied extensively, the counterpart phenomena in antiferromagnets (AFM) has been less investigated. AFMs have recently gained new interest in spintronics [35, 36]. The absence of parasitic stray fields and the ability to operate in the THz regime make AFM systems promising candidates for next-generation high-speed, nanoscale spintronic technologies. Non-equilibrium magnon excitation in AFM insulators have been recently realized via spin-transfer torque and laser excitation [37, 38]. The possibility of parallel pumping in orthorhombic AFM systems was theoretically and experimentally investigated long ago [39]. However, it has not been successfully demonstrated in conventional AFM systems within the context of modern AFM spintronics.

To the best of our knowledge, there are only a few theoretical studies on generation and stability analysis of quasi-equilibrium mBEC states in AFM systems [31, 40–42]. It is worthy to mention that the *equilibrium* magnon condensation at cryogenic temperatures has already been observed in *quantum* AFM systems in which the phase transition to the BEC state is controlled by

* These authors contributed equally to this work

an external DC magnetic field [41, 43]. In this article, we investigate the stability of quasi-equilibrium mBEC in collinear AFM systems with uniaxial and biaxial magnetic anisotropy.

In compensated and collinear AFM systems, dipolar interactions are negligible and their magnon spectra often have a minimum at the centre of the magnetic Brillouin zone, i.e. at the Γ symmetry point with zero momentum. Magnons in collinear AFM systems have two opposite chiral modes. While these two magnon modes are degenerate in uniaxial easy-axis AFM systems, the degeneracy can be broken by applying an external magnetic field or in the presence of a hard-axis magnetic anisotropy [36, 44, 45]. In these cases the minima of both magnon modes remain at the Γ symmetry point. On the other hand, in systems with broken inversion symmetry, an antisymmetric Dzyaloshinskii–Moriya (DM) interaction is permitted, which lifts the degeneracy of the two chiral modes, causing them to split along the momentum axis [46]. As a result, the two magnon modes have two degenerate minima at finite wavevector. Such momentum-dependent band splitting of AFM magnons has been measured in α - $\text{Cu}_2\text{V}_2\text{O}_7$ by inelastic neutron scattering [47]. These AFM systems may facilitate the presence of a nonzero wavevector double condensate state.

In this paper, we explore the stability of magnon condensation by analytical calculations on two distinct AFM systems: a uniaxial easy-axis AFM with DM interaction and a biaxial AFM without DM interaction. In the first case, the magnon bands exhibit Rashba-type splitting along the momentum axis, while in the second case, the magnon bands are split along the energy axis, akin to Zeeman-type splitting.

The rest of the paper is organized as follows: Section II introduces the spin Hamiltonian of our AFM systems. Section III is a short presentation of the Holstein-Primakoff and Dyson-Maleev transformations. Section IV introduces the first AFM system we study, which is a uniaxial AFM with DM interaction. The *zero-sound spectrum* of this system is calculated in section Section V. Section VI introduces the second AFM system; a biaxial AFM without DM interaction. In this system we study the condensate stability when the interband magnon interactions are suppressed by applying a strong magnetic field. We compare the two AFM systems and present our concluding remarks in Section VII. Finally, Appendix Section A contains a comparison of the Holstein-Primakoff and Dyson-Maleev bosonization techniques.

II. SPIN HAMILTONIAN

We consider a collinear two-sublattice AFM system, where the sublattice spins \mathbf{S}_A and \mathbf{S}_B align antiparallel to each other along the Néel vector, oriented in the \hat{z} direction, described by the following extended quantum

Heisenberg model,

$$\mathcal{H} = \mathcal{H}_{\text{ex}} + \mathcal{H}_{\text{ani}} + \mathcal{H}_{\text{D}} + \mathcal{H}_{\text{Z}}. \quad (1)$$

The Heisenberg exchange interaction \mathcal{H}_{ex} acts between nearest-neighbour (n.n.) spins at lattice sites i and j with an AFM exchange coupling $J > 0$,

$$\mathcal{H}_{\text{ex}} = J \sum_{\substack{i \in A, B \\ j \in n.n.}} \mathbf{S}_i \cdot \mathbf{S}_j. \quad (2)$$

The magnetic anisotropy Hamiltonian incorporates single-ion easy and hard-axes magnetic anisotropies [44, 48],

$$\mathcal{H}_{\text{ani}} = -K_z \sum i(S_i^z)^2 + K_x \sum i(S_i^x)^2. \quad (3)$$

Here, $K_z \geq 0$ parametrize the uniaxial easy-axis magnetic anisotropy along the \hat{z} direction, while $K_x \geq 0$ denotes the hard-axis magnetic anisotropy along the \hat{x} direction.

The DM Hamiltonian reads,

$$\mathcal{H}_{\text{D}} = \sum_{\substack{i \in A, B \\ j \in n.n.}} \mathbf{D}_{ij} \cdot (\mathbf{S}_i \cdot \mathbf{S}_j), \quad (4)$$

where the DM vector is antisymmetric, $\mathbf{D}_{ij} = -\mathbf{D}_{ji}$. The vector is oriented along the \hat{z} direction so that $\mathbf{D}_{ij} = D\nu_{ij}\hat{z}$, with $\nu_{ij} = \pm 1$ for clockwise and counterclockwise hopping directions.

Finally, the interaction between an external magnetic field along the \hat{z} direction and localized spins are modelled by the Zeeman coupling,

$$\mathcal{H}_{\text{Z}} = -\mu_{\text{B}}h \sum_i S_i^z, \quad (5)$$

where μ_{B} is the Bohr magneton and h is the magnetic field amplitude.

III. NONLINEAR BOSON REPRESENTATIONS IN THE THEORY OF ANTIFERROMAGNETS

At low temperatures, the elementary excitations of an ordered magnet are magnons. To describe these excitations, it is suitable to pass from spin representation to boson operators. There are several bosonic representations of spins [49, 50].

In this paper, we consider both the Holstein-Primakoff and Dyson-Maleev transformations, and we compare the two methods in Appendix A. In particular, the Dyson-Maleev transformation use spin raising and lowering operators which are not Hermitian conjugates of each other. For this reason, we investigate whether the choice of bosonization technique affects the final analytical results in Appendix A.

i. Holstein-Primakoff transformation: In this transformation we define raising and lowering spin operators

which are Hermitian conjugates of each other, $\hat{S}^+ = (\hat{S}^-)^\dagger = \hat{S}^x + i\hat{S}^y$. We denote the bosonic creation (annihilation) by \hat{a}_i^\dagger (\hat{a}_i) for spins at sublattice A, and \hat{b}_j^\dagger (\hat{b}_j) for spins at sublattice B. Given the quantization axis along the \hat{z} axis, we have the following transformation for sublattice A in terms of the spin S up to third order in the bosonic operators,

$$\hat{S}_i^+ = \sqrt{2S - \hat{a}_i^\dagger \hat{a}_i \hat{a}_i} \approx \sqrt{2S}(\hat{a}_i - \frac{\hat{a}_i^\dagger \hat{a}_i \hat{a}_i}{4S}), \quad (6a)$$

$$\hat{S}_i^z = S - \hat{a}_i^\dagger \hat{a}_i. \quad (6b)$$

For sublattice B the operators are,

$$\hat{S}_j^+ = \hat{b}_j^\dagger \sqrt{2S - \hat{b}_j^\dagger \hat{b}_j} \approx \sqrt{2S}(\hat{b}_j^\dagger - \frac{\hat{b}_j^\dagger \hat{b}_j^\dagger \hat{b}_j}{4S}), \quad (7a)$$

$$\hat{S}_j^z = -(S - \hat{b}_j^\dagger \hat{b}_j). \quad (7b)$$

Since we are interested in the lowest nonlinear interaction in the spin Hamiltonian, we expanded the square roots in the above expressions assuming S is large and a low number of bosons.

ii. Dyson-Maleev transformation: In this representation, the raising and lowering spin operators are not Hermitian conjugates of each other, $\hat{S}^+ \neq (\hat{S}^-)^\dagger$. The spin operators on sublattice A are given by,

$$\hat{S}_i^+ = \sqrt{2S}(\hat{a}_i - \frac{\hat{a}_i^\dagger \hat{a}_i \hat{a}_i}{2S}), \quad (8a)$$

$$\hat{S}_i^- = \sqrt{2S}\hat{a}_i^\dagger, \quad (8b)$$

$$\hat{S}_i^z = S - \hat{a}_i^\dagger \hat{a}_i, \quad (8c)$$

and for sublattice B they are,

$$\hat{S}_j^+ = \sqrt{2S}(\hat{b}_j^\dagger - \frac{\hat{b}_j^\dagger \hat{b}_j \hat{b}_j}{2S}), \quad (9a)$$

$$\hat{S}_j^- = \sqrt{2S}\hat{b}_j^\dagger, \quad (9b)$$

$$\hat{S}_j^z = -(S - \hat{b}_j^\dagger \hat{b}_j). \quad (9c)$$

In the next sections, we only use Holstein-Primakoff transformations. We present our results for Dyson-Maleev calculations in the Appendix A. There we show that both transformations results in the same stability analysis of the magnon condensate in the uniaxial AFM.

IV. MAGNON BEC IN UNIAXIAL EASY-AXIS AFM CASE

First we study a uniaxial AFM with DM interaction. In this case, we set $K_x = 0$. We use the Holstein-Primakoff transformation and the Fourier transform $\hat{a}_i = (N/2)^{-1/2} \sum_{\mathbf{k}} \hat{a}_{\mathbf{k}} e^{-i\mathbf{k}\cdot\mathbf{r}_i}$, where N denotes the number of sites, to express the Hamiltonian in terms of bosonic operators in momentum space.

The non-interaction Hamiltonian consisting of quadratic bosonic operators reads,

$$\begin{aligned} \mathcal{H}^{(2)} = & SJ \sum_{\mathbf{k}z} [\hat{a}_{\mathbf{k}}^\dagger \hat{a}_{\mathbf{k}} + \hat{b}_{\mathbf{k}}^\dagger \hat{b}_{\mathbf{k}}] \\ & + 2SJ \cos(k_y) [\hat{a}_{\mathbf{k}} \hat{b}_{-\mathbf{k}} + \hat{a}_{\mathbf{k}}^\dagger \hat{b}_{-\mathbf{k}}^\dagger] \\ & + 2SK_z \sum_{\mathbf{k}} [\hat{a}_{\mathbf{k}}^\dagger \hat{a}_{\mathbf{k}} + \hat{b}_{\mathbf{k}}^\dagger \hat{b}_{\mathbf{k}}] - \mu_B h \sum_{\mathbf{k}} [\hat{b}_{\mathbf{k}}^\dagger \hat{b}_{\mathbf{k}} - \hat{a}_{\mathbf{k}}^\dagger \hat{a}_{\mathbf{k}}] \\ & - 4SD \sum_{\mathbf{k}} \sin(k_y) [\hat{a}_{\mathbf{k}}^\dagger \hat{b}_{-\mathbf{k}b}^\dagger + \hat{a}_{\mathbf{k}} \hat{b}_{-\mathbf{k}}], \end{aligned} \quad (10)$$

where z is the coordination number and \mathbf{k} is the wavevector. To diagonalize this bosonic Hamiltonian, we perform a Bogoliubov transformation,

$$\hat{\alpha}_{\mathbf{k}} = u_{\mathbf{k}} \hat{a}_{\mathbf{k}} + v_{\mathbf{k}} \hat{b}_{-\mathbf{k}}^\dagger, \quad (11a)$$

$$\hat{\beta}_{\mathbf{k}} = v_{-\mathbf{k}} \hat{a}_{-\mathbf{k}}^\dagger + u_{-\mathbf{k}} \hat{b}_{\mathbf{k}}, \quad (11b)$$

where bosonic $\hat{\alpha}_{\mathbf{k}}$ and $\hat{\beta}_{\mathbf{k}}$ operators denote two species of magnons in the AFM system.

We define the parameters $\omega_{\text{ex}} = 2SJ$, $\omega_z = SK_z$, $\omega_{ez} = \omega_{\text{ex}} + \omega_z$, $\omega_D = 2SD$, $\omega_H = \frac{1}{2}\mu_B$. The Bogoliubov coefficients are then expressed as,

$$u_{\mathbf{k}}^2 = \frac{1}{2}(\omega_{ez}[\omega_{ez}^2 - (\omega_{\text{ex}} \cos(k_y) - \omega_D \sin(k_y))^2]^{-\frac{1}{2}} + 1), \quad (12a)$$

$$v_{\mathbf{k}}^2 = \frac{1}{2}(\omega_{ez}[\omega_{ez}^2 - (\omega_{\text{ex}} \cos(k_y) - \omega_D \sin(k_y))^2]^{-\frac{1}{2}} - 1) \quad (12b)$$

The non-interacting Hamiltonian in the diagonalized magnon basis reads,

$$\mathcal{H}^{(2)} = \sum_{\mathbf{k}} \mathbf{k} \left[\varepsilon_{\mathbf{k}}^\alpha \hat{\alpha}_{\mathbf{k}}^\dagger \hat{\alpha}_{\mathbf{k}} + \varepsilon_{\mathbf{k}}^\beta \hat{\beta}_{-\mathbf{k}} \hat{\beta}_{-\mathbf{k}}^\dagger \right]. \quad (13)$$

The dispersion relations for the left-handed $\hat{\alpha}_{\mathbf{k}}$ and right-handed $\hat{\beta}_{\mathbf{k}}$ magnons are given by,

$$\varepsilon_{\mathbf{k}}^\alpha = \sqrt{\omega_{ez}^2 - (\omega_{\text{ex}} \cos(k_y) - \omega_D \sin(k_y))^2} + \omega_H, \quad (14a)$$

$$\varepsilon_{\mathbf{k}}^\beta = \sqrt{\omega_{ez}^2 - (\omega_{\text{ex}} \cos(k_y) - \omega_D \sin(k_y))^2} - \omega_H \quad (14b)$$

From these dispersion relations, it is evident that the DM interaction term leads to momentum-dependent splitting of magnon bands, while a Zeeman field leads to an energy-dependent splitting of bands, see Fig. 1. In the assumption that $\omega_D < \omega_{\text{ex}}$, the minima of the two magnon bands happen at finite wavevector $Q = \tan^{-1}(\omega_D/\omega_{\text{ex}})$

We obtain the lower-order interaction Hamiltonian, which is quartic in boson operators within the Holstein-




FIG. 1: Dispersion relations for the right- and left-handed magnon populations α (blue) and β (orange). The parameters for the Dzyaloshinskii–Moriya interaction, anisotropy, and external magnetic field strength from Eq. (10) are $\omega_D/\omega_{\text{ex}} = 0.03$, $\omega_z/\omega_{\text{ex}} = 0.025$, $\omega_H S/\omega_{\text{ex}} = 0.0025$ and $S = 5/2$. The dashed vertical lines mark the boundaries of the first Brillouin zone.

Primakoff transformation,

$$\begin{aligned}
\mathcal{H}^{(4)} = & - \sum_{\mathbf{k}, \mathbf{q}, \mathbf{q}'} \frac{f_J(\mathbf{k})}{(N/2)} \left[\hat{a}_{\mathbf{k}} \hat{b}_{\mathbf{k}+\mathbf{q}+\mathbf{q}'}^\dagger \hat{b}_{\mathbf{q}} \hat{b}_{\mathbf{q}'} + \text{h.c.} \right. \\
& + \hat{a}_{\mathbf{k}+\mathbf{q}+\mathbf{q}'}^\dagger \hat{a}_{\mathbf{q}} \hat{a}_{\mathbf{q}'} \hat{b}_{\mathbf{k}} + \text{h.c.} \\
& \left. + 4 \hat{a}_{\mathbf{q}-\mathbf{k}}^\dagger \hat{a}_{\mathbf{q}} \hat{b}_{\mathbf{q}'+\mathbf{k}}^\dagger \hat{b}_{\mathbf{q}'} \right] \\
& - \sum_{\mathbf{k}, \mathbf{q}, \mathbf{q}'} \frac{K_z}{(N/2)} \left[\hat{a}_{\mathbf{q}-\mathbf{k}}^\dagger \hat{a}_{\mathbf{q}} \hat{a}_{\mathbf{q}'+\mathbf{k}}^\dagger \hat{a}_{\mathbf{q}'} + \hat{b}_{\mathbf{q}-\mathbf{k}}^\dagger \hat{b}_{\mathbf{q}} \hat{b}_{\mathbf{q}'+\mathbf{k}}^\dagger \hat{b}_{\mathbf{q}'} \right] \\
& + \sum_{\mathbf{k}, \mathbf{q}, \mathbf{q}'} \frac{f_D(\mathbf{k})}{(N/2)} \left[\hat{a}_{\mathbf{k}}^\dagger \hat{b}_{\mathbf{q}} \hat{b}_{\mathbf{q}'}^\dagger \hat{b}_{\mathbf{k}+\mathbf{q}+\mathbf{q}'} + \hat{a}_{\mathbf{k}} \hat{b}_{\mathbf{k}+\mathbf{q}+\mathbf{q}'}^\dagger \hat{b}_{\mathbf{q}} \hat{b}_{\mathbf{q}'} \right. \\
& \left. - \hat{a}_{\mathbf{q}}^\dagger \hat{a}_{\mathbf{q}'}^\dagger \hat{a}_{\mathbf{q}+\mathbf{q}'+\mathbf{k}} \hat{b}_{\mathbf{k}}^\dagger - \hat{a}_{\mathbf{q}+\mathbf{q}'+\mathbf{k}}^\dagger \hat{a}_{\mathbf{q}} \hat{a}_{\mathbf{q}'} \hat{b}_{\mathbf{k}} \right] \quad (15)
\end{aligned}$$

where the Hermitian conjugate terms are denoted by "h.c.", and we defined $f_J(\mathbf{k}) = J \cos(k_y)$ and $f_D(\mathbf{k}) = D \sin(k_y)$.

We can rewrite the nonlinear Hamiltonian of Eq. (15) in terms of magnon operators, using the Bogoliubov transformation. To investigate mBEC stability, we are only interested in magnons at the bottom of each respective band, i.e. $\hat{a}_{k=-Q}$ and $\hat{\beta}_{k=+Q}$. Therefore, the nonlinear Hamiltonian for condensed magnons can be rewritten as the sum of intra- and inter-band terms $\mathcal{H}_Q^{(4)} = \mathcal{H}_Q^{\text{intra}} + \mathcal{H}_Q^{\text{inter}}$. The intra- and inter-band terms

are,

$$\mathcal{H}_Q^{\text{intra}} = \mathcal{A}_Q (\hat{\alpha}_{-Q} \hat{\alpha}_{-Q}^\dagger + \hat{\beta}_Q \hat{\beta}_Q^\dagger), \quad (16a)$$

$$\begin{aligned}
\mathcal{H}_Q^{\text{inter}} = & \mathcal{B}_Q (\hat{\alpha}_{-Q} \hat{\beta}_Q) \\
& + \mathcal{C}_Q (\hat{\alpha}_{-Q} \hat{\beta}_Q + \hat{\alpha}_{-Q} \hat{\alpha}_{-Q}^\dagger \\
& + \hat{\alpha}_{-Q} \hat{\alpha}_{-Q}^\dagger + \hat{\alpha}_{-Q}^\dagger \hat{\beta}_Q) \\
& + \mathcal{D}_Q (\hat{\alpha}_{-Q} \hat{\beta}_Q + \hat{\alpha}_{-Q}^\dagger \hat{\beta}_Q^\dagger). \quad (16b)
\end{aligned}$$

The amplitudes of the different condensate magnon interaction terms are,

$$\begin{aligned}
\mathcal{A}_Q = & -\frac{2}{N} [2(f_J(Q) + f_D(Q))\eta_3 \\
& - 2J\eta_2 - K_z(\eta_1 - 2\eta_2)], \quad (17a)
\end{aligned}$$

$$\begin{aligned}
\mathcal{B}_Q = & -\frac{2}{N} [8(f_J(Q) + f_D(Q))\eta_3 \\
& - 4J(\eta_1 - \eta_2) - 4K_z\eta_2], \quad (17b)
\end{aligned}$$

$$\begin{aligned}
\mathcal{C}_Q = & -\frac{2}{N} [(f_J(Q) + f_D(Q))(\eta_1 + \eta_2) \\
& - 4J\eta_3 - 2K_z\eta_3], \quad (17c)
\end{aligned}$$

$$\begin{aligned}
\mathcal{D}_Q = & -\frac{2}{N} [2(f_J(Q) + f_D(Q))\eta_3 \\
& - 2J\eta_2 - K_z\eta_2], \quad (17d)
\end{aligned}$$

where we have defined the following coefficients,

$$\eta_1 = u_{-Q}^4 + v_{-Q}^4 + 4u_{-Q}^2 v_{-Q}^2, \quad (18)$$

$$\eta_2 = 2u_{-Q}^2 v_{-Q}^2, \quad (19)$$

$$\eta_3 = -u_{-Q} v_{-Q}^4 (u_{-Q}^2 v_{-Q}^2). \quad (20)$$

When the magnon density is high, the interaction terms become important. Our goal is to analyze the nonlinear potential energy of the condensate magnons. We express the two condensate populations as two macroscopic wave functions using the Madelung transformation [20],

$$\langle \hat{\alpha}_{-Q} \rangle = \sqrt{N_\alpha} e^{i\phi_\alpha}, \quad (21a)$$

$$\langle \hat{\beta}_Q \rangle = \sqrt{N_\beta} e^{i\phi_\beta}, \quad (21b)$$

where $N_{\alpha(\beta)}$ is the number of chiral $\alpha(\beta)$ -magnons at the bottom of the bands, and $\phi_{\alpha(\beta)}$ refers to the macroscopic phase. We can define the total number of condensed magnons as $N_c = N_\alpha + N_\beta$, the phase sum $\Phi = \phi_\alpha + \phi_\beta$, and the difference in population of chiral condensate magnons $\delta = N_\alpha - N_\beta$. Finally, we find the following

interaction potential for condensate magnons,

$$\mathcal{V}^{(4)} = \mathcal{V}_Q^{\text{intra}} + \mathcal{V}_Q^{\text{inter}}, \quad (22a)$$

$$\mathcal{V}_Q^{\text{intra}} = \frac{N_c^2}{2} \mathcal{A}_Q \left[1 + \left(\frac{\delta}{N_c} \right)^2 \right], \quad (22b)$$

$$\begin{aligned} \mathcal{V}_Q^{\text{inter}} = & \frac{N_c^2}{2} \left[2\mathcal{C}_Q \sqrt{1 - \left(\frac{\delta}{N_c} \right)^2} \cos(\Phi) \right. \\ & - \left(\mathcal{D}_Q \cos(2\Phi) + \frac{\mathcal{B}_Q}{2} \right) \left(\frac{\delta}{N_c} \right)^2 \\ & \left. + \frac{\mathcal{B}_Q}{2} + \mathcal{D}_Q \cos(2\Phi) \right]. \quad (22c) \end{aligned}$$

We consider the total number of condensate magnons N_c to be constant. In general, the number of bosonic quasiparticles is not conserved due to e.g. damping, so this assumption is only valid in a time-scale proportional to the magnon condensate lifetime. We note that the interaction Hamiltonian of Eqs. (16a) and (16b) in its generalized form is directly comparable to the interaction Hamiltonian of ferromagnetic systems with two-component condensates, as seen in [24]. To investigate whether the condensate magnons are stable, we search for the minima of $\mathcal{V}^{(4)}(\delta, \Phi)$. First, we compute the extremum of $\mathcal{V}^{(4)}(\delta, \Phi) = 0$. We find the following critical points,

$$\begin{aligned} \text{i)} & \left(\frac{\delta_1}{N_c} \right)^2 = 0, \Phi_1 = 0 \\ \text{ii)} & \left(\frac{\delta_2}{N_c} \right)^2 = 0, \Phi_2 = \pi \\ \text{iii)} & \left(\frac{\delta_3}{N_c} \right)^2 = 1 - \left[\frac{\mathcal{C}_Q \cos(\Phi_3)}{\mathcal{A}_Q - \frac{\mathcal{B}_Q}{2} - \mathcal{D}_Q \cos(2\Phi_3)} \right]^2, \Phi_3 = 0 \\ \text{iv)} & \left(\frac{\delta_4}{N_c} \right)^2 = 1 - \left[\frac{\mathcal{C}_Q \cos(\Phi_3)}{\mathcal{A}_Q - \frac{\mathcal{B}_Q}{2} - \mathcal{D}_Q \cos(2\Phi_3)} \right]^2, \Phi_4 = \pi \\ \text{v)} & \left(\frac{\delta_5}{N_c} \right)^2 = 0, \Phi_5 = \arccos\left(\frac{-\mathcal{C}_Q}{\mathcal{D}_Q}\right), \\ \text{vi)} & \left(\frac{\delta_6}{N_c} \right)^2 = 1 - \left[\frac{\mathcal{C}_Q \cos(\Phi_6)}{\mathcal{A}_Q - \frac{\mathcal{B}_Q}{2} - \mathcal{D}_Q \cos(2\Phi_6)} \right]^2, \\ & \Phi_6 = \arccos\left(\frac{-\mathcal{C}_Q}{\mathcal{D}_Q} \left[1 - \left(\frac{\delta_6}{N_c} \right)^2 \right]^{-1/2}\right). \quad (23) \end{aligned}$$

To find the minima of the potential, if any, we check the second derivative sign of the potential. The discriminant is defined as,

$$\text{Disc} = \left[\left(\frac{\partial^2 \mathcal{V}^{(4)}}{\partial \delta^2} \right) \left(\frac{\partial^2 \mathcal{V}^{(4)}}{\partial \Phi^2} \right) - \left(\frac{\partial^2 \mathcal{V}^{(4)}}{\partial \delta \partial \Phi} \right)^2 \right]. \quad (24)$$

The nonlinear condensed magnon potential has a minimum if $\text{Disc} > 0$, $\left(\frac{\partial^2 \mathcal{V}^{(4)}}{\partial \delta^2} \right) > 0$, and $\left(\frac{\partial^2 \mathcal{V}^{(4)}}{\partial \Phi^2} \right) > 0$. These requirements depend on the numerical value of the potential interactions \mathcal{A}_Q , \mathcal{B}_Q , \mathcal{C}_Q and \mathcal{D}_Q , which are functions of spin parameters.




FIG. 2: The value of the interaction strength $\mathcal{V}^{(4)}$ (blue solid line) from Eq. (22a) evaluated for the symmetric magnon distribution $\delta = 0$, $\Phi = \pi$, listed as minima (ii) in Eq. (23). The parameters are $\omega_D/\omega_{\text{ex}} = 0.3$, $S = 5/2$ and $Q = \tan^{-1}(\omega_D/\omega_{\text{ex}})$. The interaction can be separated into the intra-condensate component $\mathcal{V}_Q^{\text{intra}}$ (blue dashed) and the inter-condensate component $\mathcal{V}_Q^{\text{inter}}$ (blue dotted). We include the second derivatives $\left(\frac{\partial^2 \mathcal{V}^{(4)}}{\partial \delta^2} \right)$ (dark red) and $\left(\frac{\partial^2 \mathcal{V}^{(4)}}{\partial \Phi^2} \right)$ (bright red).

We inspect the potential $\mathcal{V}^{(4)}(\delta, \Phi)$ and its second derivatives for our parameter space. We find that extreme point ii) is the only extrema listed in Eq. (23) which fulfils the above criteria for stable condensation.

In Fig. 2, we plot the magnon condensate interaction potential $\mathcal{V}^{(4)}$ as a function of the easy-axis magnetic anisotropy strength K_z/J for extremum point (ii). We also plot the second derivative of the interaction potential with respect to the distribution difference δ and phase sum Φ . The plot shows the intra- and inter-band contributions to the potential. We find that the interactions are dominated by the inter-band scatterings. We predict that extremum point (ii) is the only stable condensate state. The state has a symmetric magnon distribution between the two minima. This is in contrast to what is observed in ferromagnetic systems, where previous theory indicate that magnon condensates in FM systems can be stabilized for both asymmetric and symmetric populations, depending on parameters such as the anisotropy strength and external field strength [24].

V. ZERO-SOUND-LIKE EXCITATIONS IN MAGNON BEC

In the two-component magnon condensate, the difference between the two condensate phases can be a new Goldstone mode. This out-of-phase mode in

the collision-less regime shows similarities to the zero-sound phenomena in Fermi liquid theory. We calculate the zero-sound spectrum by solving the related Gross-Pitaevskii equation. Our condensate wavefunction is $\Psi_{\pm} = \sqrt{N_{\pm Q}} e^{i\phi_{\pm Q}}$. We consider the Hamiltonian operator of Eq. (1) which gives the interaction terms listed in Eqs. (16a) and (16b). We follow a similar approach as in Ref. [20] and we obtain an expression for the energy per volume V ,

$$\begin{aligned} E/V = \int dr \left[\frac{\hbar^2}{2mV} (|\nabla\Psi_+|^2 + |\nabla\Psi_-|^2) \right. \\ + \mathcal{A}_Q (|\nabla\Psi_+|^4 + |\nabla\Psi_-|^4) \\ + \mathcal{B}_Q (|\nabla\Psi_+|^2 + |\nabla\Psi_-|^2) \\ + \mathcal{C}_Q (\Psi_+\Psi_- + \Psi_+^\dagger\Psi_-^\dagger) (|\Psi_+|^2 + |\Psi_-|^2) \\ \left. + \mathcal{D}_Q (\Psi_+\Psi_+\Psi_-\Psi_- + \Psi_+^\dagger\Psi_+^\dagger\Psi_-^\dagger\Psi_-^\dagger) \right]. \quad (25) \end{aligned}$$

Here, \hbar is Planck's constant and m is the boson mass. In the previous section, we showed that the condensed magnons are stabilized for a symmetric magnon distribution. We now proceed to consider small deviations from this symmetric state, where $N_{\pm} = \frac{1}{2}N_c \pm \delta n$ and $\Phi = \pi$. We use the commutation relation $[\delta\phi, \delta n] = i$ and the equation of motion in a similar approach as Ref. [20], and get the following coupled equations,

$$\begin{aligned} \hbar \frac{\partial \delta\phi}{\partial t} = -\frac{\hbar^2}{2m} \frac{1}{N_c} \nabla^2 \delta n \\ + (\mathcal{A}_Q - \mathcal{B}_Q - \mathcal{D}_Q + \mathcal{C}_Q) V \delta n, \quad (26) \end{aligned}$$

$$\hbar \frac{\partial \delta n}{\partial t} = \frac{\hbar^2}{2m} N_c \nabla^2 \delta\phi. \quad (27)$$

To solve these equation of motions, we use the following harmonic-oscillator ansatz for the density and phase,

$$\delta n(t) = \delta n_0 e^{i\omega t} e^{-ikr} \quad (28a)$$

$$\delta\phi(t) = \delta\phi_0 e^{i\omega t} e^{-ikr} \quad (28b)$$

Finally, we get the following dispersion relation for the zero-sound-like excitation,

$$\omega^2 = \frac{\hbar^2}{(2m)^2} k^4 + \frac{N_c}{2m} (\mathcal{A}_Q - \mathcal{B}_Q - \mathcal{D}_Q + \mathcal{C}_Q) k^2. \quad (29)$$

The spectra of Eq. (29) is similar to the symmetric spectra of the ferromagnetic system in Ref. [20]. In our uniaxial AFM system, the slope $(\mathcal{A}_Q - \mathcal{B}_Q - \mathcal{D}_Q + \mathcal{C}_Q)$ is negative. For ferromagnetic systems, Ref. [20] found that if omitting terms similar to \mathcal{D}_Q , the slope can be both positive and negative, depending on parameters such as the film thickness and strength of the external magnetic field.

VI. MAGNON BEC IN BIAxIAL AFM CASE

Now we consider a biaxial AFM system without DM interaction. The system is similar to the one studied in

Ref. [45]. For materials such as NiO we assume that $J > K_x > K_z$ [44, 45]. We introduce $\omega_x = SK_x$. The magnon eigenenergies in the dispersion relation minima from Ref. [45] are given by,

$$\begin{aligned} \omega_{\alpha,\beta}^2 = \frac{1}{4} (\omega_{\text{ex}} + \omega_x + 2\omega_z + \omega_H^2 - \frac{1}{4} (\omega_x^2 + \omega_{\text{ex}}^2)) \\ \pm \frac{1}{2} \sqrt{4\omega_H^2 ((\omega_{\text{ex}} + \omega_x + 2\omega_z)^2 - \omega_{\text{ex}}^2) + \omega_{\text{ex}}^2 \omega_x^2}. \quad (30) \end{aligned}$$

Here, β -magnons have the lowest energy. We see that in this case the degeneracy of chiral magnon bands is broken even in the absence of the magnetic field, $\omega_H = 0$, due to the presence of the hard-axis magnetic anisotropy. The magnon spectra have the minima at $k = 0$. We proceed to investigate the four-magnon population in the lowest energy state. In this sense, the inter-band magnon interactions are suppressed. We use the following Bogoliubov transformation,

$$\hat{a} = u_\alpha \hat{\alpha} - v_\beta \hat{\beta}^\dagger, \quad (31a)$$

$$\hat{b}^\dagger = -v_\alpha \hat{\alpha} + u_\beta \hat{\beta}^\dagger. \quad (31b)$$

We use the approximated Bogoliubov parameters from Ref. [45] evaluated at $k = 0$,

$$u_{\alpha,\beta} = \sqrt{\frac{\omega_{\text{ex}} + \omega_x + 2\omega_z + \omega_{\alpha,\beta}}{2\omega_{\alpha,\beta}(\omega_H = 0)}}, \quad (32a)$$

$$v_{\alpha,\beta} = \sqrt{\frac{\omega_{\text{ex}} + \omega_x + 2\omega_z - \omega_{\alpha,\beta}}{2\omega_{\alpha,\beta}(\omega_H = 0)}}. \quad (32b)$$

We analyze the interactions of the magnon population of lowest energy, $k = 0$, which means that we only consider the condensed β -magnon interactions,

$$\mathcal{H}_\beta^{(4)} = \mathcal{A}_2 (\hat{\beta}\hat{\beta}\hat{\beta}^\dagger\hat{\beta}^\dagger) + \mathcal{A}_3 (\hat{\beta}\hat{\beta}\hat{\beta}\hat{\beta}^\dagger + \text{h.c.}) + \mathcal{A}_4 (\hat{\beta}\hat{\beta}\hat{\beta}\hat{\beta} + \text{h.c.}) \quad (33)$$

As before, we perform the Madelung transformation, $\langle \beta \rangle \rightarrow \sqrt{N_\beta} e^{i\phi_\beta}$, where ϕ_β is the phase and N_β is the number of condensed magnons. We obtain a general expression for the condensed β -magnon potential,

$$\mathcal{V}_\beta^{(4)} N_\beta^2 [\mathcal{A}_2 + 2\mathcal{A}_3 \cos(2\phi_\beta) + 2\mathcal{A}_4 \cos(4\phi_\beta)], \quad (34)$$

where the interaction parameters are given by,

$$\begin{aligned} \mathcal{A}_2 = \frac{4}{N} [-2J(u_\beta^2 v_\beta^2) + Ju_\beta v_\beta (u_\beta^2 + v_\beta^2) \\ - \frac{1}{8} K_z (u_\beta^4 + v_\beta^4) - 4K_x (u_\beta^4 + v_\beta^4)], \quad (35) \end{aligned}$$

$$\mathcal{A}_3 = -\frac{8}{N} K_x (u_\beta^4 + v_\beta^4), \quad (36)$$

$$\mathcal{A}_4 = 0. \quad (37)$$

We note that \mathcal{A}_3 appears purely due to the presence of the hard-axis magnetic anisotropy.

The expression in Eq. (34) shows that the interaction potential $\mathcal{V}_\beta^{(4)}$ is proportional to N_β^2 . This means that the expression for $\mathcal{V}_\beta^{(4)}$ does not have a nontrivial minimum for some finite number of condensed magnons. This indicates that there is no stable magnon condensation state for the system. We conclude that a single component magnon condensate state cannot be stabilized in magnetic systems.

VII. SUMMARY AND CONCLUDING REMARKS

We have calculated the condensate interactions in a uniaxial and biaxial AFM system using both the Holstein-Primakoff and Dyson-Maleev bosonization technique. The uniaxial easy-axis system can host two magnon condensate populations, resembling the condensation observed in FM thin-film YIG. The interactions are a function of the phase sum Φ and magnon distribution difference δ . We analyze the interactions for a range of easy-axis magnetic anisotropy strengths and DM interaction. We find that only the symmetric condensed magnon distribution $\delta = 0$ leads to a stable magnon condensation. This indicates that even in the absence of dipolar interactions, which play a crucial role in the stability of two-component magnon condensates in FM thin-film YIG, it is possible to have a meta-stable magnon condensate state in AFM systems. We predict that the stable condensate state will have a symmetric distribution of magnons between the two condensate populations. We found that both the Holstein-Primakoff and Dyson-Maleev transformations lead to the same conclusions on condensate stability.

In the biaxial AFM without DM interaction, the degeneracy of the magnon minima energy is lifted, and there can be only one chiral magnon population at the bottom energy minimum. However, when we consider the condensate magnon potential, we find that since there is no intermagnon interaction, there is no stable condensate state. In reality, there can still be weak intermagnon interactions between magnons at the lowest β -magnon band and magnons at the minimum of the higher α -magnon band, but we expect that such interactions would be too weak to stabilize the condensation.

We propose that our theory can be tested experimentally in a uniaxial AFM material with Rashba-type DM interaction. Without an external magnetic field, this system has two chiral magnon bands minima with the same energy at finite wavevector. By applying a magnetic field, one can lift this energy degeneracy. Our theory shows that a metastable two-component magnon condensate in the absence of the magnetic field can be created, and that

applying an external magnetic field along the magnetic ground-state can destroy it because of a reduction in the interband interaction of the condensate magnons.

ACKNOWLEDGEMENT

This project has been supported by the Research Council of Norway through its Centers of Excellence funding scheme, Project No. 262633 "QuSpin".

Appendix A: Uniaxial AFM case: Dyson-Maleev transformation

We want to verify whether our results in the main text depends on the bosonization technique. We introduce the alternative Dyson-Maleev approach [51]. Then, we follow the same procedure as in the Holstein-Primakoff framework. The Dyson-Maleev transformation gives us similar results for the non interaction quadratic Hamiltonian of Eq. (10). When analyzing the four-magnon quartic Hamiltonian of Eqs. (16a) and (16b), we find the same prefactors for \mathcal{A}_Q , \mathcal{B}_Q and \mathcal{D}_Q . However, the prefactor \mathcal{C}_Q must be modified. We find that the new Hamiltonian terms in the Dyson-Maleev bosonization are,

$$\mathcal{H}_Q^{\text{intra}} = \mathcal{A}_Q(\hat{\alpha}_{-\mathbf{Q}}\hat{\alpha}_{-\mathbf{Q}}^\dagger + \hat{\beta}_{\mathbf{Q}}\hat{\beta}_{\mathbf{Q}}^\dagger), \quad (\text{A1a})$$

$$\begin{aligned} \mathcal{H}_Q^{\text{inter}} &= \mathcal{B}_Q(\hat{\alpha}_{-\mathbf{Q}}\hat{\beta}_{\mathbf{Q}}) \\ &+ \mathcal{C}_Q^{\text{var1}}(\hat{\alpha}_{-\mathbf{Q}}\hat{\beta}_{\mathbf{Q}} + \hat{\alpha}_{-\mathbf{Q}}\hat{\alpha}_{-\mathbf{Q}}^\dagger) \\ &+ \mathcal{C}_Q^{\text{var2}}(\hat{\alpha}_{-\mathbf{Q}}\hat{\alpha}_{-\mathbf{Q}}^\dagger + \hat{\alpha}_{-\mathbf{Q}}^\dagger\hat{\beta}_{\mathbf{Q}}) \\ &+ \mathcal{D}_Q(\hat{\alpha}_{-\mathbf{Q}}\hat{\beta}_{\mathbf{Q}} + \hat{\alpha}_{-\mathbf{Q}}^\dagger\hat{\beta}_{\mathbf{Q}}^\dagger). \end{aligned} \quad (\text{A1b})$$

Here we have defined $\mathcal{C}_Q^{\text{var1}} = \mathcal{C}_Q + \Delta_{\mathcal{C}_Q}$ and $\mathcal{C}_Q^{\text{var2}} = \mathcal{C}_Q - \Delta_{\mathcal{C}_Q}$. The prefactors \mathcal{A}_Q , \mathcal{B}_Q and \mathcal{D}_Q are listed in Eq. (17).

We perform a Madelung transform and obtain the following correction to the \mathcal{C} -terms of the condensed magnon potential in Eq. (22c),

$$\mathcal{V}_4^{\mathcal{C}} = \frac{N_c^2}{2} \sqrt{1 - \left(\frac{\delta}{N_c}\right)^2} (2\mathcal{C}_Q \cos(\Phi) + 2i\frac{\delta}{N_c} \sin(\Phi)\Delta_{\mathcal{C}_Q}). \quad (\text{A2})$$

The Holstein-Primakoff transformation results in a real potential, where the interaction of Eq. (A2) has $\Delta_{\mathcal{C}_Q} = 0$. However, in the Dyson-Maleev transformation, we find that $\Delta_{\mathcal{C}} = -\frac{2}{N}(3\eta_2 - \eta_1)(f_J(Q) + f_D(Q))$. This means that while the real part of Eq. (A2) is similar for the Holstein-Primakoff and Dyson-Maleev transformation, the latter transformation results in imaginary terms in the interaction potential. We note that the stable condensation state found has $\Phi = \pi$, and the interaction potential is fully real.

- [1] Bose–einstein condensation of quasi-equilibrium magnons at room temperature under pumping. *Nature*, 443(7110):430–433, 2006.
- [2] Pyotr Kapitza. Viscosity of liquid helium below the λ -point. *Nature*, 141(3558):74–74, 1938.
- [3] John F Allen and AD Misener. Flow of liquid helium ii. *Nature*, 141(3558):75–75, 1938.
- [4] M. H. Anderson, J. R. Ensher, M. R. Matthews, C. E. Wieman, and E. A. Cornell. Observation of bose-einstein condensation in a dilute atomic vapor. *Science*, 269(5221):198–201, 1995.
- [5] C. C. Bradley, C. A. Sackett, J. J. Tollett, and R. G. Hulet. Evidence of bose-einstein condensation in an atomic gas with attractive interactions. *Phys. Rev. Lett.*, 75:1687–1690, Aug 1995.
- [6] K. B. Davis, M. O. Mewes, M. R. Andrews, N. J. van Druten, D. S. Durfee, D. M. Kurn, and W. Ketterle. Bose-einstein condensation in a gas of sodium atoms. *Phys. Rev. Lett.*, 75:3969–3973, Nov 1995.
- [7] J. T. Mäkinen, S. Autti, and V. B. Eltsov. Magnon bose–einstein condensates: From time crystals and quantum chromodynamics to vortex sensing and cosmology. *Applied Physics Letters*, 124(10):100502, 03 2024.
- [8] AS Borovik-Romanov, Yu M Bunkov, VV Dmitriev, and Yu M Mukharskii. Long-lived induction signal in superfluid $^3\text{He-b}$. *JETP Lett*, 40(6):1033, 1984.
- [9] Yu. M. Bunkov, S. N. Fisher, A. M. Guénault, and G. R. Pickett. Persistent spin precession in ^3B in the regime of vanishing quasiparticle density. *Phys. Rev. Lett.*, 69:3092–3095, Nov 1992.
- [10] Iulia Georgescu. 25 years of bec. *Nature Reviews Physics*, 2(8):396–396, 2020.
- [11] Zefang Wang, Daniel A Rhodes, Kenji Watanabe, Takashi Taniguchi, James C Hone, Jie Shan, and Kin Fai Mak. Evidence of high-temperature exciton condensation in two-dimensional atomic double layers. *Nature*, 574(7776):76–80, 2019.
- [12] O.V. Misochko, Muneaki Hase, K. Ishioka, and M. Kitajima. Transient bose–einstein condensation of phonons. *Physics Letters A*, 321(5):381–387, 2004.
- [13] V. E. Demidov, O. Dzyapko, S. O. Demokritov, G. A. Melkov, and A. N. Slavin. Thermalization of a parametrically driven magnon gas leading to bose-einstein condensation. *Phys. Rev. Lett.*, 99:037205, Jul 2007.
- [14] O Dzyapko, V E Demidov, S O Demokritov, G A Melkov, and A N Slavin. Direct observation of bose–einstein condensation in a parametrically driven gas of magnons. *New Journal of Physics*, 9(3):64, mar 2007.
- [15] V. E. Demidov, O. Dzyapko, S. O. Demokritov, G. A. Melkov, and A. N. Slavin. Observation of spontaneous coherence in bose-einstein condensate of magnons. *Phys. Rev. Lett.*, 100:047205, Jan 2008.
- [16] V. E. Demidov, O. Dzyapko, M. Buchmeier, T. Stockhoff, G. Schmitz, G. A. Melkov, and S. O. Demokritov. Magnon kinetics and bose-einstein condensation studied in phase space. *Phys. Rev. Lett.*, 101:257201, Dec 2008.
- [17] O. Dzyapko, V. E. Demidov, S. O. Demokritov, G. A. Melkov, and V. L. Safonov. Monochromatic microwave radiation from the system of strongly excited magnons. *Applied Physics Letters*, 92(16):162510, 04 2008.
- [18] Roberto E Troncoso and Álvaro S Núñez. Dynamics and spontaneous coherence of magnons in ferromagnetic thin films. *Journal of Physics: Condensed Matter*, 24(3):036006, dec 2011.
- [19] P Nowik-Boltyk, O Dzyapko, VE Demidov, NG Berloff, and SO Demokritov. Spatially non-uniform ground state and quantized vortices in a two-component bose-einstein condensate of magnons. *Scientific reports*, 2(1):482, 2012.
- [20] Fuxiang Li, Wayne M Saslow, and Valery L Pokrovsky. Phase diagram for magnon condensate in yttrium iron garnet film. *Scientific reports*, 3(1):1372, 2013.
- [21] Alexander A Serga, Vasil S Tiberkevich, Christian W Sandweg, Vitaliy I Vasyuchka, Dmytro A Bozhko, Andrii V Chumak, Timo Neumann, Björn Obry, Gennadii A Melkov, Andrei N Slavin, et al. Bose–einstein condensation in an ultra-hot gas of pumped magnons. *Nature communications*, 5(1):3452, 2014.
- [22] Roberto E. Troncoso and Álvaro S. Núñez. Josephson effects in a bose–einstein condensate of magnons. *Annals of Physics*, 346:182–194, 2014.
- [23] P. O. Sukhachov, S. Banerjee, and A. V. Balatsky. Bose-einstein condensate of dirac magnons: Pumping and collective modes. *Phys. Rev. Res.*, 3:013002, Jan 2021.
- [24] Therese Frostad, Philipp Pirro, Alexander A. Serga, Burkard Hillebrands, Arne Brataas, and Alireza Qaiumzadeh. Anisotropy-assisted magnon condensation in ferromagnetic thin films. *Phys. Rev. Res.*, 6:L012011, Jan 2024.
- [25] Therese Frostad, Hans L. Skarsvåg, Alireza Qaiumzadeh, and Arne Brataas. Spin-transfer-assisted parametric pumping of magnons in yttrium iron garnet. *Phys. Rev. B*, 106:024423, Jul 2022.
- [26] Morteza Mohseni, Alireza Qaiumzadeh, Alexander A Serga, Arne Brataas, Burkard Hillebrands, and Philipp Pirro. Bose–einstein condensation of nonequilibrium magnons in confined systems. *New Journal of Physics*, 22(8):083080, aug 2020.
- [27] Wenhao Xu, Andrey A Bagrov, Farhan T Chowdhury, Luke D Smith, Daniel R Kattnig, Hilbert J Kappen, and Mikhail I Katsnelson. Friction versus bose-einstein condensation in pumped bosonic systems. *arXiv preprint arXiv:2411.00058*, 2024.
- [28] B Divinskiy, H Merbouche, VE Demidov, KO Nikolaev, L Soumah, D Gouéré, R Lebrun, V Cros, Jamal Ben Youssef, P Bortolotti, et al. Evidence for spin current driven bose-einstein condensation of magnons. *Nature communications*, 12(1):6541, 2021.
- [29] Michael Schneider, Thomas Brächer, David Breitbach, Viktor Lauer, Philipp Pirro, Dmytro A Bozhko, Halyna Yu Musiienko-Shmarova, Björn Heinz, Qi Wang, Thomas Meyer, et al. Bose–einstein condensation of quasiparticles by rapid cooling. *Nature Nanotechnology*, 15(6):457–461, 2020.
- [30] Nick P Proukakis, David W Snoke, and Peter B Littlewood. *Universal themes of Bose-Einstein condensation*. Cambridge university press, 2017.
- [31] Naoya Arakawa. Controlling stability of bose-einstein condensation of interacting magnons in an antiferromagnet by an external magnetic field. *Phys. Rev. B*, 99:014405, Jan 2019.
- [32] Morteza Mohseni, Vitaliy I Vasyuchka, Victor S L’vov, Alexander A Serga, and Burkard Hillebrands. Classical

- analog of qubit logic based on a magnon bose–einstein condensate. *Communications physics*, 5(1):196, 2022.
- [33] S. N. Andrianov and S. A. Moiseev. Magnon qubit and quantum computing on magnon bose-einstein condensates. *Phys. Rev. A*, 90:042303, Oct 2014.
- [34] Kouki Nakata, Kevin A. van Hoogdalem, Pascal Simon, and Daniel Loss. Josephson and persistent spin currents in bose-einstein condensates of magnons. *Phys. Rev. B*, 90:144419, Oct 2014.
- [35] Tomas Jungwirth, X Marti, P Wadley, and J Wunderlich. Antiferromagnetic spintronics. *Nature nanotechnology*, 11(3):231–241, 2016.
- [36] V. Baltz, A. Manchon, M. Tsoi, T. Moriyama, T. Ono, and Y. Tserkovnyak. Antiferromagnetic spintronics. *Rev. Mod. Phys.*, 90:015005, Feb 2018.
- [37] A. Simoncig, R. Mincigrucci, E. Principi, F. Bencivenga, A. Calvi, L. Foglia, G. Kurdi, A. Matruglio, S. Dal Zilio, V. Masciotti, M. Lazzarino, and C. Masciovecchio. Generation of coherent magnons in nio stimulated by ev pulses from a seeded free-electron laser. *Phys. Rev. Mater.*, 1:073802, Dec 2017.
- [38] Christian Tzschaschel, Kensuke Otani, Ryugo Iida, Tsutomu Shimura, Hiroaki Ueda, Stefan Günther, Manfred Fiebig, and Takuya Satoh. Ultrafast optical excitation of coherent magnons in antiferromagnetic nio. *Phys. Rev. B*, 95:174407, May 2017.
- [39] Hitoshi Yamazaki. Parallel pumping of spin-waves in an orthorhombic antiferromagnet. *Journal of the Physical Society of Japan*, 32(5):1227–1233, 1972.
- [40] Eirik Løhaugen Fjærbu, Niklas Rohling, and Arne Brataas. Electrically driven bose-einstein condensation of magnons in antiferromagnets. *Phys. Rev. B*, 95:144408, Apr 2017.
- [41] Thierry Giamarchi, Christian Rüegg, and Oleg Tchernyshyov. Bose–einstein condensation in magnetic insulators. *Nature Physics*, 4(3):198–204, 2008.
- [42] Yury M. Bunkov and Vladimir L. Safonov. Magnon condensation and spin superfluidity. *Journal of Magnetism and Magnetic Materials*, 452:30–34, 2018.
- [43] Yosuke Matsumoto, Simon Schnierer, Jan AN Bruin, Jürgen Nuss, Pascal Reiss, George Jackeli, Kentaro Kitagawa, and Hidenori Takagi. A quantum critical bose gas of magnons in the quasi-two-dimensional antiferromagnet ybcl₃ under magnetic fields. *Nature Physics*, pages 1–8, 2024.
- [44] Sergio M. Rezende, Antonio Azevedo, and Roberto L. Rodríguez-Suárez. Introduction to antiferromagnetic magnons. *Journal of Applied Physics*, 126(15):151101, 10 2019.
- [45] F. L. A. Machado, P. R. T. Ribeiro, J. Holanda, R. L. Rodríguez-Suárez, A. Azevedo, and S. M. Rezende. Spin-flop transition in the easy-plane antiferromagnet nickel oxide. *Phys. Rev. B*, 95:104418, Mar 2017.
- [46] Masataka Kawano, Yoshinori Onose, and Chisa Hotta. Designing rashba–dresselhaus effect in magnetic insulators. *Communications Physics*, 2(1):27, 2019.
- [47] G. Gitgeatpong, Y. Zhao, P. Piyawongwatthana, Y. Qiu, L. W. Harriger, N. P. Butch, T. J. Sato, and K. Matan. Nonreciprocal magnons and symmetry-breaking in the noncentrosymmetric antiferromagnet. *Phys. Rev. Lett.*, 119:047201, Jul 2017.
- [48] Mahroo Shiranzaei, Roberto E Troncoso, Jonas Fransson, Arne Brataas, and Alireza Qaiumzadeh. Thermal squeezing and nonlinear spectral shift of magnons in antiferromagnetic insulators. *New Journal of Physics*, 24(10):103009, oct 2022.
- [49] Andrei A Katanin and Valentin Yu Irkhin. Magnetic order and spin fluctuations in low-dimensional insulating systems. *Physics-Uspekhi*, 50(6):613, jun 2007.
- [50] Assa Auerbach. *Interacting electrons and quantum magnetism*. Springer Science & Business Media, 2012.
- [51] Mona Helen Alexandra Kalthoff. *Nonequilibrium materials engineering in correlated systems via light-matter coupling*. PhD thesis, Staats-und Universitätsbibliothek Hamburg Carl von Ossietzky, 2022.