

Tracer and current fluctuations in driven diffusive systems

Théotim Berlioz,¹ Olivier Bénichou,¹ and Aurélien Grabsch¹

¹*Sorbonne Université, CNRS, Laboratoire de Physique Théorique de la Matière Condensée (LPTMC), 4 Place Jussieu, 75005 Paris, France*

Interacting particles diffusing in single-file is a fundamental model of transport in narrow channels where particles cannot bypass each other. An important result has been obtained by Kollmann for the mean square displacement of a tracer for any such single-file model. However, since then, no general result has been obtained in the important case where the particles are driven by an external field. Here, we fill this gap and determine the fluctuations and the skewness of the tracer's position for any driven diffusive system. In addition, we also consider a variety of important observables such as the integrated current, the response of the system to the perturbation induced by the displacement of the tracer, and the correlations between several tracers. Furthermore, we also unveil fundamental relations underlying the out-of-equilibrium dynamics of driven diffusive systems. This work constitutes a first step toward a full description of driven one-dimensional systems of interacting particles.

Introduction.— The investigation of the dynamical properties of interacting particle systems in both equilibrium and non-equilibrium settings has been a prominent area of research in the last decades [1–5]. A paradigmatic example is the single-file model, which describes particles diffusing in narrow channels, so that they cannot bypass each other. In this context, two quantities have attracted a lot of attention: the position X_t of a tracer, and the integrated current (which measures the total particles flux through the origin). It was shown that at long times the mean squared displacement of a tracer particle grows sub-linearly with time, $\langle X_T^2 \rangle \propto T^{1/2}$ [6–8]. This is a general property of single-file systems, which has been observed experimentally in various systems [9–12]. However the precise determination of the prefactor, which contains all the dependence on the physical properties of the system, such as the nature of the interaction, the value of the density, or the initial condition, is subtle.

For a general single-file system, starting from an equilibrium distribution with mean density $\bar{\rho}$, the mean-squared displacement (MSD) of the tracer, including the prefactor, has first been determined for reflecting Brownian particles [6] [13], and later for the Simple Exclusion Process (SEP) [8] [14], which is a paradigmatic model of single-file system. A major progress has been accomplished by Kollman [15] who obtained the result for *any* single-file system,

$$\langle X_T^2 \rangle_{T \rightarrow \infty} \simeq \frac{\sigma(\bar{\rho})}{\bar{\rho}^2 \sqrt{\pi D(\bar{\rho})}} \sqrt{T}, \quad (1)$$

where $D(\rho)$ is the collective diffusion coefficient, and $\sigma(\rho)$ the mobility, which controls both the current fluctuations, and the response of the current to a small drive. The fluctuations of the integrated current Q_T through the origin, defined as the number of particles that have crossed the origin up to time T from left to right minus those from right to left, have also been determined, and




FIG. 1. A system of diffusive particles in one dimension, which interact with a given potential (represented in red) and driven by a field ν (green arrows). At large scales this system can be described by two transport coefficients: the collective diffusion coefficient $D(\rho)$ and the mobility $\sigma(\rho)$ [3, 5, 27, 28], which both depend on the local density of particles $\rho(x, t)$. We consider here two standard observables: the position X_t of a tagged particle (the tracer, in blue) at time t , and the integrated current Q_t through the origin up to time t .

take a similar form [16],

$$\langle Q_T^2 \rangle_{T \rightarrow \infty} \simeq \frac{\sigma(\bar{\rho})}{\sqrt{\pi D(\bar{\rho})}} \sqrt{T}. \quad (2)$$

The distribution of both X_T and Q_T has been shown to be non-Gaussian [17, 18]. Recently, there has been a lot of interest in characterizing these distributions beyond the variance, particularly through higher-order cumulants, which quantify deviations from Gaussianity. All the cumulants have first been determined for specific models, such as reflecting Brownian particles [18–21], and the SEP [17, 22–25]. For general single-file systems the question is more complex and, beyond the MSD, only the fourth cumulant has been determined recently [26].

A natural question is how X_T or Q_T behave when the full system is subjected to an external driving field. This is a ubiquitous situation, for instance encountered in traffic flow [29] or molecular motors [30]. Unlike the case of non-driven single-file diffusion described above, the presence of a driving field keeps the system out of equilibrium, leading to non-trivial steady states and complex dynamics. Despite their importance, the determination

of the fluctuations (and higher order cumulants) of X_T or Q_T for *any* driven diffusive systems, i.e. the extension of (1) and (2) to driven systems is still an open question. We stress that even in specific cases, such as the central model of the weakly asymmetric simple exclusion process (WASEP, see below for definition), the fluctuations have not been determined.

Here, we fill this gap and take the first step to characterize the fluctuation in *any* driven diffusive system. Using the macroscopic fluctuation theory (MFT) formalism, we obtain both the second and third cumulant (which characterizes the skewness of the distribution) of the integrated current Q_T , and the associated correlation profiles, which describe the correlations of Q_T with the medium. Using a duality mapping between general 1D driven diffusive systems (which extends the one found for non-driven systems in [31]), we deduce the first three cumulants of X_T , as well as the bath-mediated correlations between two tracers in these systems. We stress that our results for *any* $D(\rho)$ and $\sigma(\rho)$ allow one to characterize driven systems of diffusive particles interacting with a realistic potential (such as Lennard-Jones or Weeks-Chandler-Andersen involved in single-file experiments on colloids [11]) [32], as shown in Fig. 1.

Driven diffusive systems.— Driven systems typically display a ballistic behaviour, meaning that the cumulants of X_T or Q_T grow linearly with time [33]. This is for instance well-known for the asymmetric simple exclusion process (ASEP), in which particles hop (with exclusion) on a 1D lattice with different rates p_{\pm} towards the left and right [4, 28, 34, 35]. It is a minimal model of interacting particles driven by an external field $F = (p_+ - p_-)/2$. Such systems can be studied using ballistic macroscopic fluctuation theory [36–40]. However, for a *weak drive*, the system remains diffusive, i.e. the cumulants of Q_T or X_T scale as \sqrt{T} . A typical example is the weakly asymmetric simple exclusion process (WASEP), which has attracted a lot of attention [41–45], in particular due to its close relation with the Kardar-Parisi-Zhang equation [46]. In the WASEP, which is a specific case of the ASEP, the hopping rates are chosen as $p_{\pm} = 1 \pm \frac{\nu}{\sqrt{T}}$ when the observable (Q_T or X_T) is measured at a large time T . Note that this scaling of the drive with T is required to have a diffusive dynamics. Similarly, a general system of driven diffusive particles displays a diffusive behaviour if the drive F scales with the observation time T as

$$F = \nu/\sqrt{T}, \quad \text{with } \nu \text{ fixed.} \quad (3)$$

Importantly, this scaling of the drive with the observation time, which defines driven diffusive systems, allows one to obtain general results which interpolate between non-driven (equilibrium) and strongly driven (non-equilibrium) situations. Technically, we will rely on the MFT formalism.

Macroscopic fluctuation theory. The MFT is a general framework developed by Bertini et al. [5] to describe dif-

fusive systems far from equilibrium at large scales (long times and large distances). Rescaling space by \sqrt{T} and time by T allows to introduce a density $\rho(x, t)$ and a current $j(x, t)$ which characterise the system at the macroscopic scale. The main idea is that these fields are stochastic, and verify the constitutive equations

$$\partial_t \rho + \partial_x j = 0, \quad j = -D(\rho) \partial_x \rho + \nu \sigma(\rho) + \frac{\sqrt{\sigma(\rho)}}{T^{1/4}} \zeta, \quad (4)$$

together with the boundary conditions $\rho(\pm\infty, t) = \bar{\rho}$ and $j(\pm\infty, t) = \nu \sigma(\bar{\rho})$, and where ζ is a Gaussian white noise, $\langle \zeta(x, t) \zeta(x', t') \rangle = \delta(x - x') \delta(t - t')$. Note that the noise that arises from the stochastic dynamics scales with $T^{-1/4}$ due to the rescaling of space and time. The transport coefficients $D(\rho)$ and $\sigma(\rho)$ embed all the microscopic details (dynamics, interaction, etc) of the system. For a given system, these transport coefficients have to be calculated from the microscopic scale. Exact formulas are available only for a few specific models, such as the WASEP for which $D(\rho) = 1$ and $\sigma(\rho) = 2\rho(1 - \rho)$. Nevertheless, in the general case where $D(\rho)$ and $\sigma(\rho)$ cannot be computed exactly, there exists a variational formula [1] that yields approximate expressions [47, 48].

The microscopic integrated current through the origin can be expressed in terms of the macroscopic current as

$$Q_T = \sqrt{T} \int_0^1 j(0, t) dt, \quad (5)$$

where the factor \sqrt{T} emerges from the macroscopic description (see Supplemental Material (SM) [49] for details). The analysis of (5) in the MFT framework relies on a path integral formulation of the stochastic equations (4), which provides the probability of observing a trajectory $\{\rho(x, t), j(x, t)\}$. In the long times limit, only the optimal trajectory is relevant, and is determined by the Euler-Lagrange equations [5, 17],

$$\partial_t q = \partial_x [D(q) \partial_x q - \sigma(q) \partial_x p - \nu \sigma(q)], \quad (6a)$$

$$\partial_t p = -D(q) \partial_x^2 p - \frac{1}{2} \sigma'(q) (\partial_x p)^2 - \nu \sigma'(q) \partial_x p. \quad (6b)$$

Here $q(x, t)$ is the optimal realisation of $\rho(x, t)$, while $p(x, t)$ is a conjugate field that imposes the conservation relation (4). These equations are completed by boundary conditions, which depend on the observable (Q_T or X_T) under consideration. For Q_T (5) they take the form [17]

$$p(x, 1) = \lambda \Theta(x), \quad p(x, 0) = \lambda \Theta(x) + \int_{\bar{\rho}}^{q(x, 0)} \frac{2D(r)}{\sigma(r)} dr. \quad (7)$$

The statistical properties of Q_T , encoded in the cumulants $\hat{\kappa}_n$ (scaled by \sqrt{T}) defined by

$$\hat{\psi}(\lambda) \equiv \lim_{T \rightarrow \infty} \frac{1}{\sqrt{T}} \ln \langle e^{\lambda Q_T} \rangle \equiv \sum_{n=1}^{\infty} \hat{\kappa}_n \frac{\lambda^n}{n!}, \quad (8)$$

can be directly determined from the solution of the MFT equations (see SM [49]) as

$$\frac{d\hat{\psi}}{d\lambda} = \nu\sigma(\bar{\rho}) + \int_0^\infty [q(x, 1) - q(x, 0)]dx. \quad (9)$$

Note that, even in the non-driven case $\nu = 0$, no general solution of the MFT equations (6) is available. Exact solutions have recently been obtained [50–55], but only for specific models. Here, in the case of arbitrary $D(\rho)$, $\sigma(\rho)$ and drive $\nu \neq 0$, we determine the first cumulants $\hat{\kappa}_n$ exactly by solving the MFT equations (6,7) order by order in λ [16, 18, 20, 26].

Cumulants of the integrated current.— Calculations, provided in SM [49], finally yield explicit formulas for the first three cumulants of Q_T for general 1D driven diffusive systems,

$$\hat{\kappa}_1 = \nu\sigma(\bar{\rho}), \quad \hat{\kappa}_2 = \frac{\sigma(\bar{\rho})}{\sqrt{D(\bar{\rho})}} \mathcal{G}\left(y \equiv \frac{\nu\sigma'(\bar{\rho})}{2\sqrt{D(\bar{\rho})}}\right), \quad (10)$$

$$\begin{aligned} \hat{\kappa}_3 = & \nu \frac{\sigma(\bar{\rho})\sigma'(\bar{\rho})^2}{4D(\bar{\rho})^2} + \nu \frac{\sigma(\bar{\rho})^2\sigma''(\bar{\rho})}{4D(\bar{\rho})^2} [1 + 3(y^2 - \mathcal{G}(y)^2)] \\ & + \frac{D'(\bar{\rho})\sigma(\bar{\rho})^2}{4D(\bar{\rho})^{5/2}} \left[y(1 - 3\text{erf}(y)^2) - 3\text{erf}(y) \frac{e^{-y^2}}{\sqrt{\pi}} \right], \quad (11) \end{aligned}$$

where we denoted $\mathcal{G}(y) = \frac{e^{-y^2}}{\sqrt{\pi}} + y\text{erf}(y)$. These results constitute the first description of *any* driven diffusive system. In addition, they allow one to address important situations and go beyond the study of Q_T by considering: (i) other observables such as X_T , (ii) the ballistic limit of driven diffusive systems, (iii) the response of the system to the perturbation induced by the displacement of the tracer, (iv) the extension to the case of several tracers. Furthermore, they also enable to reveal fundamental relation underlying the out-of-equilibrium dynamics of driven diffusive systems.

Cumulants of the tracer's position.— First, we show that the cumulants of X_T can be directly deduced from those of Q_T (10,11). Indeed, the substitutions

$$D(\rho) \rightarrow \frac{1}{\rho^2}D\left(\frac{1}{\rho}\right), \quad \sigma(\rho) \rightarrow \rho\sigma\left(\frac{1}{\rho}\right), \quad (12)$$

$$\nu \rightarrow -\nu, \quad \lambda \rightarrow -\lambda, \quad \bar{\rho} \rightarrow \frac{1}{\bar{\rho}}. \quad (13)$$

map the position X_T of a tracer in a given driven diffusive system onto the opposite of the integrated current through the origin in a dual system (see SM [49]), generalising the duality relation derived in [31] for symmetric ($\nu = 0$) systems. For instance the variance takes the compact form,

$$\frac{\langle X_T^2 \rangle_c}{\sqrt{T}} \underset{T \rightarrow \infty}{\simeq} \frac{\sigma(\bar{\rho})}{\bar{\rho}^2\sqrt{D(\bar{\rho})}} \mathcal{G}\left(z \equiv \nu \frac{\sigma(\bar{\rho}) - \bar{\rho}\sigma'(\bar{\rho})}{2\bar{\rho}\sqrt{D(\bar{\rho})}}\right), \quad (14)$$

where the index c refers to the cumulants. This result constitutes the extension of the celebrated result (1) of Kollmann [15] to *any* driven diffusive system. Remarkably, the effect of the drive ν on the fluctuations of X_T vanishes if $\sigma(\rho) \propto \rho$. This is for instance the case in models of reflecting Brownian particles, for which the drive can be absorbed by a Galilean transform. In all other cases, the presence of a drive strictly increases the fluctuations of X_T .

Large drive limit: ballistic regime.— Next, we show that our results, obtained under the scaling (3), contain the ballistic regime as a limit. This is a priori not guaranteed, since the limits $\nu \rightarrow \infty$ and $T \rightarrow \infty$ do not necessarily commute. In fact, inserting $\nu = F\sqrt{T}$ into the cumulants (10,11) and performing the duality mapping (12,13), we obtain

$$\langle X_T \rangle \underset{T \rightarrow \infty}{\simeq} T F \frac{\sigma(\bar{\rho})}{\bar{\rho}}, \quad (15)$$

$$\langle X_T^2 \rangle_c \underset{T \rightarrow \infty}{\simeq} T |F| \frac{\sigma(\bar{\rho})}{2\bar{\rho}^3 D(\bar{\rho})} |\sigma(\bar{\rho}) - \bar{\rho}\sigma'(\bar{\rho})|, \quad (16)$$

$$\begin{aligned} \langle X_T^3 \rangle_c \underset{T \rightarrow \infty}{\simeq} T F \frac{\sigma(\bar{\rho})}{4\bar{\rho}^3 D(\bar{\rho})^3} & \left[\sigma''(\bar{\rho})\sigma(\bar{\rho}) \right. \\ & \left. + (\sigma(\bar{\rho}) - \bar{\rho}\sigma'(\bar{\rho})) \left(\frac{3\sigma(\bar{\rho}) - \bar{\rho}\sigma'(\bar{\rho})}{\bar{\rho}^2} + \frac{\sigma(\bar{\rho})D'(\bar{\rho})}{\bar{\rho}D(\bar{\rho})} \right) \right]. \quad (17) \end{aligned}$$

We argue that these results are the exact first cumulants of X_T in the ballistic regime. A strong indication supporting this claim is that, by setting $D(\rho) = 1$ and $\sigma(\rho) = 2\rho(1 - \rho)$, we recover from these formulas the first three cumulants of the ASEP [34, 35]. In fact, in this case, all the cumulants are equal, $\langle X_T^n \rangle_c \simeq F(1 - \bar{\rho})T$, indicating that the position of a tracer at long times in the ASEP is Poissonian [35]. This is expected for a larger class of systems, that can be determined by imposing that the cumulants (15-17) are all equal, leading to the necessary condition $D(\rho) = \frac{|\sigma(\rho) - \rho\sigma'(\rho)|}{2\rho^2}$. Note that, considering the dual system defined by (12), this condition simplifies into $2D(\rho) = |\sigma'(\rho)|$. In turn, this condition characterizes the celebrated zero-range processes (ZRP) [56] defined as lattice models in which the hopping rates of a particle depend only on the site it currently occupies (hence “zero range”). Our results finally show that the statistics of X_T is Poissonian only if the dual model is a ZRP.

Beyond the cumulants: correlation profiles.— Beyond the cumulants of Q_T and X_T , our approach also gives access to the correlations between these observables and the density of surrounding particles. These correlation profiles are both (i) physically important since they quantify the response of the bath to the displacement of a tracer, and (ii) technically relevant since they were shown to satisfy a simple closed equation in the case of the




FIG. 2. Correlation profile $\Phi_1(x)$ (19) between the macroscopic density $\rho(x, 1)$ (describing the microscopic system at $t = T$) and the integrated current Q_T , as a function of x for different values of the rescaled drive $f = \frac{\nu\sigma'(\bar{\rho})}{\sqrt{4D(\bar{\rho})}}$.

SEP [50]. These correlations between the macroscopic density $\rho(x, 1)$ describing the microscopic system at time T and the integrated current Q_T (or X_T) at the same time, are encoded in the generalised correlation profiles, which are given by [57]

$$\frac{\langle \rho(x, 1)e^{\lambda Q_T} \rangle}{\langle e^{\lambda Q_T} \rangle} \underset{T \rightarrow \infty}{\simeq} q(x, 1) \equiv \Phi(x) = \sum_{n=0}^{\infty} \frac{\lambda^n}{n!} \Phi_n(x). \quad (18)$$

The correlation profiles Φ_n up to $n = 2$ are determined in SM [49]. For instance,

$$\Phi_1(x) = \text{sign}(x) \frac{\sigma(\bar{\rho})}{4D(\bar{\rho})} \text{erfc} \left(\text{sign}(x) \frac{x - \nu\sigma'(\bar{\rho})}{2\sqrt{D(\bar{\rho})}} \right), \quad (19)$$

quantifies the correlations $\langle \rho(x, 1)Q_T \rangle_c$ between the density and Q_T , and is represented in Fig. 2. It is positive for $x > 0$ and negative for $x < 0$, indicating that an increase of Q_T is correlated with an accumulation of particles on $x > 0$, and a depletion on $x < 0$. Remarkably, when the rescaled drive $f = \frac{\nu\sigma'(\bar{\rho})}{\sqrt{4D(\bar{\rho})}}$ is increased to a value of the order of 2, Q_T completely decorrelates with the density on $x < 0$ (see Fig. 2). By symmetry, the same holds on $x > 0$ when f is decreased to the order of -2 . Importantly, the region of space which decorrelates is not necessarily related to the direction in which the particles are driven by the field ν , since $\sigma'(\bar{\rho})$ can be negative. This is for instance the case in the WASEP, for which $\sigma'(\bar{\rho}) < 0$ if $\bar{\rho} > \frac{1}{2}$. We finally note that correlations between the density and the tracer, are similarly obtained by the duality mapping, and present the same qualitative behavior with f replaced by $\nu \frac{\sigma(\bar{\rho}) - \bar{\rho}\sigma'(\bar{\rho})}{\sqrt{4\bar{\rho}^2 D(\bar{\rho})}}$.

Correlations between two tracers.— Importantly, our results also allow us to address the question of the correlation between the positions of two tracers, which are expected to be strong in 1D due to bath-mediated interactions [58–61]. The correlations between the positions




FIG. 3. Normalised covariance between the positions X_T and Y_T of two tracers, initially separated by a distance $\xi = (Y_0 - X_0)/\sqrt{T}$. The correlations are represented for different values of $z = \nu \frac{\sigma(\bar{\rho}) - \bar{\rho}\sigma'(\bar{\rho})}{\sqrt{4\bar{\rho}^2 D(\bar{\rho})}}$.

of two tracers X_T and Y_T , which initially start at $X_0 = 0$ and $Y_0 = \xi\sqrt{T}$ [62] are given by (see SM [49]),

$$\frac{\langle X_T Y_T \rangle_c}{\sqrt{T}} \underset{T \rightarrow +\infty}{\simeq} \frac{\sigma(\bar{\rho})}{\bar{\rho}^2 \sqrt{D(\bar{\rho})}} \left(\mathcal{G}(z + \xi) + \mathcal{G}(z - \xi) - \frac{\xi}{\bar{\rho}\sqrt{D(\bar{\rho})}} \right), \quad (20)$$

with z defined in Eq. (14). As expected, these correlations decrease when the initial distance between the tracer increases, and increase with the drive (see Fig. 3). Our Eq. (20) quantifies these trends for *any* driven diffusive system.

Physical boundary conditions for driven diffusive systems.— In addition to providing answers to the variety of questions considered above, our approach also enables us to reveal fundamental relations underlying the out-of-equilibrium dynamics of driven diffusive systems. Indeed, we claim that the profile Φ (18) satisfies the set of simple physical boundary conditions,

$$\mu(\Phi(0^+)) - \mu(\Phi(0^-)) = \lambda, \quad [\partial_z \mu(\Phi)]_{0^-}^+ = 0, \quad (21)$$

$$\hat{\psi} + \nu \partial_\nu \hat{\psi} = -2 (\partial_x \mu(\Phi)|_{0^+} - 2\nu) \int_{\Phi(0^-)}^{\Phi(0^+)} D(r) dr, \quad (22)$$

where $\mu(\rho) = \int^\rho \frac{2D(r)}{\sigma(r)} dr$ is the chemical potential and $[f]_a^b = f(b) - f(a)$. These relations also hold for the profiles associated to X_T , but with μ replaced by the pressure $P(\rho) = \int^\rho \frac{2rD(r)}{\sigma(r)} dr$ in the first relation of (21).

The conditions (21) are obtained from the MFT equation (6,7). Remarkably, these relations have the same form as in the non-driven case [55], which in turn demonstrates their generality. Equation (22) generalizes the boundary condition derived for symmetric ($\nu = 0$) systems [26] to general driven diffusive systems and is a conjecture, supported by the following points: (i) Based on

our determination of the first three cumulants (10,11), we verify that it holds for arbitrary $D(\rho)$, $\sigma(\rho)$ and ν up to order 3 in λ . (ii) For arbitrary λ , this relation is verified in the case of the WASEP (see SM [49]). (iii) Relation (22) is invariant under the duality mapping (12,13). Technically, this last relation provides a shortcut to the determination of the cumulants from the correlation profile Φ as it bypasses the computation of the integral (9).

We stress that the boundary relations (21,22) hold in out-of-equilibrium situations even if they involve thermodynamic functions, defined for equilibrium systems. The key point is that these thermodynamic functions are evaluated for the out-of-equilibrium quantities $\Phi(0^\pm)$ and $\Phi'(0^\pm)$.

Conclusion.— We have considered both the integrated current of particles Q_T and the displacement X_T of a tracer in *any* driven diffusive system. We have determined explicit expressions for the first three cumulants of both quantities. In addition to characterizing the fluctuations of these observables and the skewness of their distributions, we have characterized the response of the bath by determining (i) the full spatial structure of the bath-observable correlations (up to second order) and (ii) the correlations between the positions of two tracers, mediated by the bath. Importantly, we have also obtained general physical boundary conditions satisfied by the correlation profiles in these out-of-equilibrium systems. Altogether, this work constitutes a first step toward a full description of driven one-dimensional systems of interacting particles.

Acknowledgements.— The authors thank Kirone Mallick, Alexandre Krajenbrink and Pierre Le Doussal for inspiring discussions.

[1] H. Spohn, *Large scale dynamics of interacting particles* (Springer Berlin, Heidelberg, 1991).
 [2] M. R. Evans and T. Hanney, *J. Phys. A* **38**, R195 (2005).
 [3] B. Derrida, *J. Stat. Mech.* **2007**, P07023 (2007).
 [4] T. Chou, K. Mallick, and R. K. Zia, *Rep. Prog. Phys.* **74**, 116601 (2011).
 [5] L. Bertini, A. De Sole, D. Gabrielli, G. Jona-Lasinio, and C. Landim, *Rev. Mod. Phys.* **87**, 593 (2015).
 [6] T. E. Harris, *J. Appl. Probab.* **2**, 323 (1965).
 [7] D. G. Levitt, *Phys. Rev. A* **8**, 3050 (1973).
 [8] R. Arratia, *Ann. Probab.* **11**, 362 (1983).
 [9] T. T. Perkins, D. E. Smith, and S. Chu, *Science* **264**, 819 (1994).
 [10] K. Hahn, J. Kärger, and V. Kukla, *Phys. Rev. Lett.* **76**, 2762 (1996).
 [11] Q.-H. Wei, C. Bechinger, and P. Leiderer, *Science* **287**, 625 (2000).
 [12] B. Lin, M. Meron, B. Cui, S. A. Rice, and H. Diamant, *Phys. Rev. Lett.* **94**, 216001 (2005).
 [13] The model of reflecting Brownian particles refers to the 1D system of Brownian particles with a hardcore exclu-

sion constraint.
 [14] The SEP is a models of particles on a lattice which randomly hop with symmetric rates to neighbouring sites, with a hardcore exclusion constraint.
 [15] M. Kollmann, *Phys. Rev. Lett.* **90**, 180602 (2003).
 [16] P. Krapivsky and B. Meerson, *Phys. Rev. E* **86**, 031106 (2012).
 [17] B. Derrida and A. Gerschenfeld, *J. Stat. Phys.* **137**, 978 (2009).
 [18] P. L. Krapivsky, K. Mallick, and T. Sadhu, *J. Stat. Phys.* **160**, 885 (2015).
 [19] C. Hegde, S. Sabhapandit, and A. Dhar, *Phys. Rev. Lett.* **113**, 120601 (2014).
 [20] P. L. Krapivsky, K. Mallick, and T. Sadhu, *Phys. Rev. Lett.* **113**, 078101 (2014).
 [21] T. Sadhu and B. Derrida, *J. Stat. Mech: Theory Exp.* **2015**, P09008 (2015).
 [22] B. Derrida and A. Gerschenfeld, *J. Stat. Phys.* **136**, 1 (2009).
 [23] P. Illien, O. Bénichou, C. Mejía-Monasterio, G. Oshanin, and R. Voituriez, *Phys. Rev. Lett.* **111**, 038102 (2013).
 [24] T. Imamura, K. Mallick, and T. Sasamoto, *Phys. Rev. Lett.* **118**, 160601 (2017).
 [25] T. Imamura, K. Mallick, and T. Sasamoto, *Commun. Math. Phys.* **384**, 1409 (2021).
 [26] A. Grabsch and O. Bénichou, *Phys. Rev. Lett.* **132**, 217101 (2024).
 [27] H. Spohn, *J. Phys. A* **16**, 4275 (1983).
 [28] K. Mallick, *Physica A* **418**, 17 (2015), proceedings of the 13th International Summer School on Fundamental Problems in Statistical Physics.
 [29] A. S. Debashish Chowdhury, Ludger Santen, *Physics Reports* **329**, 199–329 (2000).
 [30] R. Lipowsky, S. Klumpp, and T. M. Nieuwenhuizen, *Phys. Rev. Lett.* **87**, 108101 (2001).
 [31] P. Rizkallah, A. Grabsch, P. Illien, and O. Bénichou, *J. Stat. Mech: Theory Exp.* **2023**, 013202 (2023).
 [32] If the interaction potential decays slower than $1/r$ for large distances r , the dynamics of the system will no longer be diffusive, but will instead depend on the precise exponent of the decay of the potential [63].
 [33] For the ASEP, under specific initial conditions one can observe a non-ballistic behaviour, such as the Kardar-Parisi-Zhang scaling $t^{1/3}$ [46].
 [34] A. De Masi and P. A. Ferrari, *J. Stat. Phys.* **38**, 603 (1985).
 [35] P. A. Ferrari and L. R. G. Fontes, *Journal of Applied Probability* **33**, 411 (1996).
 [36] B. Doyon and J. Myers, in *Ann. Henri Poincaré*, Vol. 21 (Springer, 2020) pp. 255–302.
 [37] J. Myers, J. Bhaseen, R. J. Harris, and B. Doyon, *SciPost Phys.* **8**, 007 (2020).
 [38] G. Peretto and B. Doyon, *SciPost Phys.* **10**, 116 (2021).
 [39] B. Doyon, G. Peretto, T. Sasamoto, and T. Yoshimura, *SciPost Phys.* **15**, 136 (2023).
 [40] B. Doyon, G. Peretto, T. Sasamoto, and T. Yoshimura, *Phys. Rev. Lett.* **131**, 027101 (2023).
 [41] C. Enaud and B. Derrida, *J. Stat. Phys.* **114**, 537 (2004).
 [42] T. Bodineau and B. Derrida, *J. Stat. Phys.* **123**, 277 (2006).
 [43] T. Bodineau and B. Derrida, *Phys. Rev. E* **72**, 066110 (2005).
 [44] S. Prohac and K. Mallick, *J. Phys. A* **42**, 175001 (2009).
 [45] M. Gorissen and C. Vanderzande, *Phys. Rev. E* **86**,

- 051114 (2012).
- [46] T. Imamura and T. Sasamoto, *Phys. Rev. Lett.* **108**, 190603 (2012).
 - [47] C. Arita, P. L. Krapivsky, and K. Mallick, *Phys. Rev. E* **95**, 032121 (2017).
 - [48] C. Arita, P. L. Krapivsky, and K. Mallick, *J. Phys. A* **51**, 125002 (2018).
 - [49] Supplemental material.
 - [50] A. Grabsch, A. Poncet, P. Rizkallah, P. Illien, and O. Bénichou, *Sci. Adv.* **8**, eabm5043 (2022).
 - [51] K. Mallick, H. Moriya, and T. Sasamoto, *Phys. Rev. Lett.* **129**, 040601 (2022).
 - [52] E. Bettelheim, N. R. Smith, and B. Meerson, *Phys. Rev. Lett.* **128**, 130602 (2022).
 - [53] E. Bettelheim, N. R. Smith, and B. Meerson, *J. Stat. Mech: Theory Exp.* **2022**, 093103 (2022).
 - [54] A. Krajenbrink and P. Le Doussal, *Phys. Rev. E* **107**, 014137 (2023).
 - [55] A. Grabsch, P. Rizkallah, and O. Bénichou, *SciPost Phys.* **16**, 016 (2024).
 - [56] T. Bodineau and B. Derrida, *Phys. Rev. Lett.* **92**, 180601 (2004).
 - [57] A. Poncet, A. Grabsch, P. Illien, and O. Bénichou, *Phys. Rev. Lett.* **127**, 220601 (2021).
 - [58] S. Majumdar and M. Barma, *Physica A* **177**, 366 (1991).
 - [59] A. Poncet, O. Bénichou, V. Démery, and G. Oshanin, *Phys. Rev. E* **97**, 062119 (2018).
 - [60] A. Poncet, O. Bénichou, V. Démery, and G. Oshanin, *Phys. Rev. Res.* **1**, 033089 (2019).
 - [61] A. Grabsch, T. Berlioz, P. Rizkallah, P. Illien, and O. Bénichou, *Phys. Rev. Lett.* **132**, 037102 (2024).
 - [62] If the two tracers are initially separated by a fixed (independent of T) distance, they will collapse at the macroscopic scale into a single tracer, leading to trivial correlations.
 - [63] R. Dandekar, P. L. Krapivsky, and K. Mallick, *Phys. Rev. E* **107**, 044129 (2023).

Supplemental Material for Tracer and current fluctuations in driven diffusive systems

Théotim Berlioz, Olivier Bénichou and Aurélien Grabsch

CONTENTS

I. Microscopic to macroscopic description	7
A. Lattice gases	8
B. Continuous models	8
II. Macroscopic fluctuation theory	9
A. The transport coefficients and the fluctuating hydrodynamics	9
B. The current through the origin	10
C. MFT for the integrated current	11
D. Duality mapping	11
III. Solving the MFT equations for the first cumulants	12
A. First Order	13
B. Second order	13
1. The term in σ''	15
2. The term in D'	16
3. Summary	17
IV. Cumulants and profiles for the integrated current	18
A. Cumulants	18
B. Correlation profile	18
V. Cumulants and profiles for the tracer	19
A. Cumulants	19
B. Correlation profiles	19
VI. Large asymmetry: ballistic limit	20
VII. Extension: correlations between two tracers or two currents	20
VIII. The weakly asymmetric simple exclusion process (WASEP)	21
A. Generalities on the ASEP	22
B. Microscopic equations for the ASEP	22
C. Microscopic equations for the WASEP	23
D. Macroscopic equations for the WASEP	23
E. Cumulants and correlations profiles	24
IX. Boundary conditions for general driven diffusive systems	24
A. Derivation from MFT	24
B. A conjecture for the cumulant generating function	25
References	25

I. MICROSCOPIC TO MACROSCOPIC DESCRIPTION

We first show how we can build a macroscopic description at a long time T from a microscopic model of interacting particles, and in particular how the integrated current at the microscopic scale is related to the macroscopic fields. We discuss the two cases of lattice gases (particles hop on discrete sites) and of particles on the real line.

A. Lattice gases

We first consider particles on a 1D infinite lattice, and denote $n_i(t)$ the number of particles at site i at time t . In exclusion models such as the SEP or the ASEP, n_i can only take values 0 or 1, but other models such as zero range processes can host an arbitrary number of particles per site [S1]. We also introduce $Q_t^{(i)}$ which counts the number of particles that have crossed the bond from i to $i+1$, minus those that have crossed from $i+1$ to i , up to time t . This is the integrated current up to t on each site i . These quantities satisfy a microscopic conservation relation,

$$n_i(t) - n_i(0) = Q_t^{(i-1)} - Q_t^{(i)}, \quad (\text{S1})$$

which states that the variation of the number of particle on site i comes from the particles that have arrived from site $i-1$ minus those which have left to site $i+1$.

We construct a macroscopic description of this problem by rescaling time by the large observation time T and space by \sqrt{T} (since we consider systems with diffusive dynamics). We introduce macroscopic density and current fields, defined as [S2, S3],

$$\rho(x, t) = \frac{1}{\sqrt{T}} \sum_i n_i(tT) \delta\left(x - \frac{i}{\sqrt{T}}\right), \quad j(x, t) = \sum_i \delta\left(x - \frac{i}{\sqrt{T}}\right) \frac{d}{dt} Q_t^{(i)} \Big|_{t \rightarrow tT}, \quad (\text{S2})$$

where we have denoted $\delta(x)$, the Dirac delta function. From the microscopic conservation equation (S1), we obtain that ρ and j must satisfy

$$\partial_t \rho = \sqrt{T} \left[j\left(x - \frac{1}{\sqrt{T}}, t\right) - j(x, t) \right]. \quad (\text{S3})$$

If instead of δ -functions we use smooth smearing functions (normalised to 1), j is continuous so that (S3) becomes the usual continuity equation,

$$\partial_t \rho + \partial_x j = 0, \quad (\text{S4})$$

for large T . From the definition of j (S2), we can express the integrated current through the origin as

$$Q_t \equiv Q_t^{(0)} = T \int_{-\frac{1}{\sqrt{2T}}}^{\frac{1}{2\sqrt{T}}} dx' \int_0^{t/T} dt' j(x', t'). \quad (\text{S5})$$

Replacing again j by a smooth function, this becomes for large T ,

$$Q_T = \sqrt{T} \int_0^1 j(0, t') dt', \quad (\text{S6})$$

which corresponds to Eq. (5) in the main text.

Similarly, we define the generating function of the correlations between the occupation numbers and the current Q_T as

$$\sum_{n=0}^{\infty} \frac{\lambda^n}{n!} \langle n_i(T) Q_T^n \rangle_c = \frac{\langle n_i(T) e^{\lambda Q_T} \rangle}{\langle e^{\lambda Q_T} \rangle} = \frac{\langle \rho(x, 1) e^{\lambda Q_T} \rangle}{\langle e^{\lambda Q_T} \rangle}, \quad x = \frac{i}{\sqrt{T}}, \quad (\text{S7})$$

which is Eq. (18) in the main text.

B. Continuous models

Let us consider now a model of particles on the real line, at positions $x_n(t)$ at time t , with $n \in \mathbb{Z}$ the label of the particle. We can construct density and current fields at the microscopic level,

$$\rho_{\text{micro}}(x, t) = \sum_n \delta(x - x_n(t)), \quad j_{\text{micro}}(x, t) = \sum_n \delta(x - x_n(t)) \frac{d}{dt} x_n(t), \quad (\text{S8})$$

which already verify the continuity equation at the microscopic scale,

$$\partial_t \rho_{\text{micro}} + \partial_x j_{\text{micro}} = 0. \quad (\text{S9})$$

The macroscopic fields are defined by rescaling space by \sqrt{T} and time by T ,

$$\rho(x, t) \equiv \rho_{\text{micro}}(x\sqrt{T}, tT), \quad j(x, t) \equiv \sqrt{T} j_{\text{micro}}(x\sqrt{T}, tT), \quad (\text{S10})$$

where the factor \sqrt{T} in the current is placed to ensure the continuity relation

$$\partial_t \rho + \partial_x j = 0. \quad (\text{S11})$$

In particular, it indicates implies that the microscopic integrated current though the origin up to T takes the form

$$Q_T \equiv \int_0^T j_{\text{micro}}(0, t') dt' = \frac{1}{\sqrt{T}} \int_0^T j\left(0, \frac{t'}{T}\right) dt' = \sqrt{T} \int_0^1 j(0, t') dt', \quad (\text{S12})$$

which is again Eq. (5) in the main text.

We also define the generating function of the correlations between the microscopic density and the integrated current Q_T as

$$\sum_{n=0}^{\infty} \frac{\lambda^n}{n!} \langle \rho_{\text{micro}}(y, T) Q_T^n \rangle_c = \frac{\langle \rho_{\text{micro}}(y, T) e^{\lambda Q_T} \rangle}{\langle e^{\lambda Q_T} \rangle} = \frac{\langle \rho(x, 1) e^{\lambda Q_T} \rangle}{\langle e^{\lambda Q_T} \rangle}, \quad x = \frac{y}{\sqrt{T}}, \quad (\text{S13})$$

which is again Eq. (18) in the main text.

II. MACROSCOPIC FLUCTUATION THEORY

In this Section, we briefly summarise the main tools needed to study both the current and tracer properties of different single-file systems.

A. The transport coefficients and the fluctuating hydrodynamics

Bertini et al. [S3] introduced a hydrodynamic description for out-of-equilibrium (both driven and non-driven) microscopic diffusive systems. This approach asserts that at large times and distances, the system is fully characterized by the coarse-grained density $\rho(x, t)$ and the local current $j(x, t)$ (which can be defined for lattice gases or continuous models, see Section I). Under local equilibrium, their evolution is governed by the macroscopic conservation equation and Fick's law:

$$\partial_t \rho = -\partial_x j \quad (\text{S14})$$

$$j = -D(\rho) \partial_x \rho + \nu \sigma(\rho) \quad (\text{S15})$$

where the transport coefficient have been noted $D(\rho)$ for the diffusivity and $\sigma(\rho)$ for the mobility (sometimes called conductivity) and ν is the driving force. The diffusivity, which depends in general of the local density, characterizes the relaxation time of the density. Similarly to the diffusivity, the mobility $\sigma(\rho)$ is a function of the local density, quantifying the response to an external field. We stress that, at the macroscopic level, all the microscopic details are encapsulated into these two transport coefficients $D(\rho)$ and $\sigma(\rho)$. Initially defined for a lattice gas [S4], they can be more intuitively understood [S5] for any system by considering a finite system of length L between two reservoirs with densities ρ_L and ρ_R [S2]. Let Q_t represent the total number of particles transferred from the left reservoir to the right one up to time t . The diffusion coefficient then measures the average current in the presence of a small density difference, and the mobility $\sigma(\rho)$ the response to the driving force ν ,

$$\lim_{t \rightarrow \infty} \frac{\langle Q_t \rangle}{t} = -\frac{D(\rho)}{L} (\rho_R - \rho_L) + \nu \sigma(\rho), \quad \text{for } \rho_R - \rho_L \ll \rho \equiv \frac{\rho_R + \rho_L}{2}. \quad (\text{S16})$$

Under local equilibrium, these two quantities are linked by the Einstein relation, a result derived from the fluctuation-dissipation theorem, which is verified at equilibrium [S3]

$$\frac{2D(\rho)}{\sigma(\rho)} = \frac{d^2 f(\rho)}{d\rho^2}, \quad (\text{S17})$$

where $f(\rho) = \int^\rho \mu(r) dr$ is the density free-energy at equilibrium and μ the chemical potential.

Importantly, the mobility also controls the fluctuations of the current at equilibrium,

$$\lim_{t \rightarrow \infty} \frac{\langle Q_t^2 \rangle}{t} = \frac{\sigma(\rho)}{L}, \quad \text{for } \rho = \rho_R = \rho_L \text{ and } \nu = 0. \quad (\text{S18})$$

The main idea of the macroscopic fluctuation theory is to treat ρ and j as stochastic fields, which still verify the continuity equation (S14), but completed by a stochastic version of the Fick law (S15),

$$j(x, t) = -D(\rho(x, t))\partial_x \rho(x, t) + \nu\sigma(\rho(x, t)) + \sqrt{\frac{\sigma(\rho(x, t))}{\sqrt{T}}}\zeta(x, t), \quad (\text{S19})$$

where ζ is a Gaussian white noise delta correlated in space and time, meaning $\langle \zeta(x, t)\zeta(x', t') \rangle = \delta(x - x')\delta(t - t')$. Note that the noise (S19) scales as $T^{-1/4}$ due to the rescaling of space and time performed to construct ρ and j [S3]. Combining equations (S14) and (S19), we obtain the stochastic equation satisfied by the density ρ (we recall that ρ describes the system at the macroscopic scale, see Section I),

$$\partial_t \rho = \partial_x \left[D(\rho)\partial_x \rho - \nu\sigma(\rho) + \sqrt{\frac{\sigma(\rho)}{\sqrt{T}}}\zeta \right]. \quad (\text{S20})$$

B. The current through the origin

As we have seen in Section I, the integrated current through the origin Q_t is expressed in terms of the macroscopic current j as

$$Q_t = \sqrt{T} \int_0^{t/T} dt' j(0, t'). \quad (\text{S21})$$

As we will see below, the MFT formalism is more conveniently implemented by expressing Q_t in terms of a different current

$$\tilde{Q}_t \equiv \int_0^\infty (\rho(x, t) - \rho(x, 0)) dx, \quad (\text{S22})$$

which has the advantage to involve only the density ρ at two different times ($t = 0$ and the macroscopic measurement time t). The two expressions (S21, S22) are related by the continuity equation (S14). This can be shown by taking the time derivative of (S22),

$$\frac{d\tilde{Q}_t}{dt} = \int_0^\infty \partial_t \rho(x, t) dx = - \int_0^\infty \partial_x j(x, t) dx. \quad (\text{S23})$$

Using that $\rho(x, t) \rightarrow \bar{\rho}$ when $x \rightarrow \infty$ and $j(x, t) \rightarrow \nu\sigma(\bar{\rho})$, we get:

$$\frac{d\tilde{Q}_t}{dt} = j(0, t) - j(+\infty, t) = j(0, t) - \nu\sigma(\bar{\rho}). \quad (\text{S24})$$

Integrating up to time t/T , and combining with (S21), we get:

$$Q_t = \sqrt{T} \int_0^{t/T} j(0, t') dt' = \sqrt{T} \int_0^{t/T} \nu\sigma(\bar{\rho}) dt' + \sqrt{T} \tilde{Q}_{t/T} = \nu\sigma(\bar{\rho}) \frac{t}{\sqrt{T}} + \sqrt{T} \tilde{Q}_{t/T}. \quad (\text{S25})$$

Setting $t = T$, we obtain

$$Q_T = \sqrt{T} \left(\nu\sigma(\bar{\rho}) + \tilde{Q}_1 \right) = \sqrt{T} \left(\nu\sigma(\bar{\rho}) + \int_0^\infty (\rho(x, 1) - \rho(x, 0)) dx \right). \quad (\text{S26})$$

C. MFT for the integrated current

We sketch here the main steps of the derivation of the MFT equations for the integrated current Q_T . From (S26), the integrated current can be written as a functional of the density as

$$Q_T = \sqrt{T}\hat{Q}[\rho], \quad \hat{Q}[\rho] \equiv \nu\sigma(\bar{\rho}) + \int_0^\infty (\rho(x,1) - \rho(x,0)) dx. \quad (\text{S27})$$

We can thus write the moment generating function of Q_T (S21) using a path integral formulation [S6],

$$\langle e^{\lambda Q_T} \rangle = \int \mathcal{D}\rho(x,t) \mathcal{D}H(x,t) \int \mathcal{D}\rho(x,0) e^{-\sqrt{T}(S[\rho,H] + F[\rho(x,0)] - \lambda\hat{Q}[\rho])}, \quad (\text{S28})$$

which is a reformulation of the stochastic equation (S20). We have denoted S the MFT action

$$S[\rho, H] = \int_{-\infty}^\infty dx \int_0^T dt \left[H\partial_t\rho + D(\rho)\partial_x\rho\partial_x H - \frac{\sigma(\rho)}{2}(\partial_x H)^2 - \nu\sigma(\rho)\partial_x H \right], \quad (\text{S29})$$

F the free-energy, which gives the distribution of the (annealed) initial condition picked from a steady state density $\bar{\rho}$,

$$F[\rho(x,0)] = \int_{-\infty}^\infty dx \int_{\bar{\rho}}^{\rho(x,0)} dr [\rho(x,0) - r] \frac{2D(r)}{\sigma(r)}, \quad (\text{S30})$$

and H a Lagrange multiplier that enforces the local conservation of particles (S14). Thanks to the factor \sqrt{T} in the exponential in (S28), the functional integrals can be evaluated by a saddle point method in the large time limit. Let us denote (q, p) the fields (ρ, H) which minimize $S + F - \lambda\hat{Q}$. They satisfy the MFT equations [S6]

$$\partial_t q = \partial_x [D(q)\partial_x q - \sigma(q)\partial_x p + \nu\sigma(q)], \quad (\text{S31})$$

$$\partial_t p = -D(q)\partial_x^2 p - \frac{1}{2}\sigma(q)(\partial_x p)^2 - \nu\sigma'(q)\partial_x p, \quad (\text{S32})$$

with the final and initial conditions

$$p(x,1) = \lambda\Theta(x), \quad p(x,0) = \lambda\Theta(x) + \int_{\bar{\rho}}^{q(x,0)} \frac{2D(r)}{\sigma(r)} dr. \quad (\text{S33})$$

In terms of the saddle point solution (q, p) , the cumulant generating function is directly obtained from (S28) as

$$\ln \langle e^{\lambda Q_T} \rangle \underset{T \rightarrow \infty}{\simeq} \sqrt{T}\hat{\psi}, \quad \hat{\psi} \equiv \lambda\hat{Q}[q] - F[q(x,0)] - S[q, p]. \quad (\text{S34})$$

In practice, this relation is rather difficult to use since it requires to compute a space-time integral for S (S29), which involves the solution at all times. It is much simpler to compute the derivative of the cumulant generating function with respect to λ , which reduces to

$$\frac{d\hat{\psi}}{d\lambda} = \hat{Q}[q], \quad (\text{S35})$$

since (q, p) is the minimum of $S + F - \lambda\hat{Q}$. Since \hat{Q} is given by (S27), it only involves the optimal field q at initial time $t = 0$ and final time $t = 1$. We will use this relation to determine the first cumulants of Q_T in the following.

D. Duality mapping

Recently, a general mapping between symmetric diffusive single-file systems has been discovered. It maps the position of a tracer in a single-file system onto the integrated current through the origin in a dual model [S7]. In this section, we extend this mapping to driven diffusive systems.

The general idea of this duality mapping is that the same model can be described in two equivalent ways, by looking either at the positions of the particles, or at the distance between two consecutive particles. At the macroscopic level,

this duality becomes a mapping between the stochastic equations (S20). Let us first consider a model described by the density ρ and current j which obey

$$\partial_t \rho + \partial_x j = 0, \quad j = -D(\rho) \partial_x \rho + \nu \sigma(\rho) + \frac{\sqrt{\sigma(\rho)}}{T^{1/4}} \zeta, \quad (\text{S36})$$

with ζ a Gaussian white noise, delta correlated in space and time. We introduce dual density and current fields $\tilde{\rho}$ and \tilde{j} , defined from the original ones as

$$\rho(x, t) = \frac{1}{\tilde{\rho}(k(x, t), t)}, \quad j(x, t) = -\frac{\tilde{j}(k(x, t), t)}{\tilde{\rho}(k(x, t), t)}, \quad \partial_x k(x, t) = \rho(x, t), \quad \partial_t k(x, t) = -j(x, t). \quad (\text{S37})$$

Inserting this transformation into the stochastic equations (S36), we obtain that $\tilde{\rho}$ and \tilde{j} obey the same equation,

$$\partial_t \tilde{\rho} + \partial_k \tilde{j} = 0, \quad \tilde{j} = -\tilde{D}(\tilde{\rho}) \partial_k \tilde{\rho} + \tilde{\nu} \tilde{\sigma}(\tilde{\rho}) + \frac{\sqrt{\tilde{\sigma}(\tilde{\rho})}}{T^{1/4}} \zeta, \quad (\text{S38})$$

with the new dual transport coefficients,

$$\tilde{D}(\rho) = \frac{1}{\rho^2} D\left(\frac{1}{\rho}\right), \quad \tilde{\sigma}(\rho) = \rho \sigma\left(\frac{1}{\rho}\right), \quad \tilde{\nu} = -\nu. \quad (\text{S39})$$

Note that this relies on the fact that ζ is a Gaussian white noise, so that $\zeta(x, t) \stackrel{(\text{law})}{=} -\zeta(x, t)$, and

$$\zeta(k(x, t), t) \stackrel{(\text{law})}{=} \sqrt{\frac{\partial x}{\partial k}} \zeta(x, t) = \frac{1}{\sqrt{\rho(x, t)}} \zeta(x, t). \quad (\text{S40})$$

Due to the transformation of the density (S37), the mean density $\bar{\rho}$ of the original model becomes $\tilde{\bar{\rho}} \equiv \frac{1}{\bar{\rho}}$ for the dual model.

Importantly, $k(x, t)$ can be interpreted as the label of the particle located at position x at time t (at the macroscopic scale, the labels become continuous due to the rescaling by \sqrt{T}). Therefore, inverting the transformation (S37) gives the position of the k^{th} particle at time t as

$$x(k, t) - x(k, 0) = \int_0^t \tilde{j}(k, t') dt'. \quad (\text{S41})$$

This relation indicates that the displacement of the k^{th} particle in the original system described by (S36) actually corresponds to the opposite of the integrated current through the position k in the dual model described by (S38), so that

$$\ln \left\langle e^{\lambda X_k(t)} \right\rangle \Big|_{D, \sigma} = \ln \left\langle e^{-\lambda Q_t(k)} \right\rangle \Big|_{\tilde{D}, \tilde{\sigma}}, \quad (\text{S42})$$

with X_k the displacement of the k^{th} particle, and $Q_t(k)$ the integrated current through k in the dual model. Expanding this relation in powers of λ , we can directly map the cumulants of the integrated current onto those of a tracer by performing the substitution (S39). This will be useful below.

III. SOLVING THE MFT EQUATIONS FOR THE FIRST CUMULANTS

We will solve the MFT equations (S31, S32) in powers of λ to obtain the first cumulants and associated correlation profiles. Because of the boundary conditions (S33), p is of order λ . We define the expansions

$$q = \bar{\rho} + \lambda q_1 + \lambda^2 q_2 + \mathcal{O}(\lambda^3), \quad p = \lambda p_1 + \lambda^2 p_2 + \mathcal{O}(\lambda^3). \quad (\text{S43})$$

When using these expansions in the bulk equations (S31, S32), we get a drift term $-\nu \sigma'(\bar{\rho}) \partial_x$ in both equations. We can absorb this term by defining the fields in a moving frame,

$$q_n(x, t) = Q_n(x - \nu \sigma'(\bar{\rho}) t, t), \quad p_n(x, t) = P_n(x - \nu \sigma'(\bar{\rho}) t, t). \quad (\text{S44})$$

A. First Order

At first order in λ , the MFT equations become

$$\partial_t P_1 = -D(\bar{\rho})\partial_x^2 P_1, \quad \partial_t Q_1 = D(\bar{\rho})\partial_x^2 Q_1 - \sigma(\bar{\rho})\partial_x^2 P_1, \quad (\text{S45})$$

with the boundary conditions (S33) which yield

$$P_1(x, 1) = \Theta(x + \nu\sigma'(\bar{\rho})), \quad Q_1(x, 0) = \frac{\sigma(\bar{\rho})}{2D(\bar{\rho})}(P_1(x, 0) - \Theta(x)). \quad (\text{S46})$$

The equation on P_1 gives straightforwardly

$$P_1(x, t) = \frac{1}{2} \operatorname{erfc} \left(-\frac{x + \nu\sigma'(\bar{\rho})}{2\sqrt{D(\bar{\rho})(1-t)}} \right). \quad (\text{S47})$$

For Q_1 it is convenient to define

$$Q_1(x, t) = \frac{\sigma(\bar{\rho})}{2D(\bar{\rho})}(P_1(x, t) + \tilde{Q}_1(x, t)), \quad (\text{S48})$$

so that

$$\partial_t \tilde{Q}_1 = D(\bar{\rho})\partial_x^2 \tilde{Q}_1, \quad \tilde{Q}_1(x, 0) = -\Theta(x). \quad (\text{S49})$$

We thus get

$$\tilde{Q}_1(x, t) = -\frac{1}{2} \operatorname{erfc} \left(-\frac{x}{2\sqrt{D(\bar{\rho})t}} \right). \quad (\text{S50})$$

In particular, we get from these expressions

$$q_1(x, 1) = \begin{cases} \frac{\sigma(\bar{\rho})}{4D(\bar{\rho})} \operatorname{erfc} \left(\frac{x - \nu\sigma'(\bar{\rho})}{2\sqrt{D(\bar{\rho})}} \right) & \text{for } x > 0, \\ -\frac{\sigma(\bar{\rho})}{4D(\bar{\rho})} \operatorname{erfc} \left(-\frac{x - \nu\sigma'(\bar{\rho})}{2\sqrt{D(\bar{\rho})}} \right) & \text{for } x < 0, \end{cases} \quad (\text{S51})$$

$$q_1(x, 0) = \begin{cases} -\frac{\sigma(\bar{\rho})}{4D(\bar{\rho})} \operatorname{erfc} \left(\frac{x + \nu\sigma'(\bar{\rho})}{2\sqrt{D(\bar{\rho})}} \right) & \text{for } x > 0, \\ \frac{\sigma(\bar{\rho})}{4D(\bar{\rho})} \operatorname{erfc} \left(-\frac{x + \nu\sigma'(\bar{\rho})}{2\sqrt{D(\bar{\rho})}} \right) & \text{for } x < 0. \end{cases} \quad (\text{S52})$$

B. Second order

At second order in λ , the MFT equations are:

$$\partial_t P_2 = -D(\bar{\rho})\partial_x^2 P_2 - \frac{1}{2}\sigma'(\bar{\rho})(\partial_x P_1)^2 - (\nu\sigma''(\bar{\rho})\partial_x P_1 + D'(\bar{\rho})\partial_x^2 P_1) Q_1,$$

$$\partial_t Q_2 = D(\bar{\rho})\partial_x^2 Q_2 - \sigma'(\bar{\rho})\partial_x P_1 \partial_x Q_1 - \sigma(\bar{\rho})\partial_x^2 P_2 - (\nu\sigma''(\bar{\rho})\partial_x Q_1 + \sigma'(\bar{\rho})\partial_x^2 P_1 - D'(\bar{\rho})\partial_x^2 Q_1) Q_1 + D'(\bar{\rho})(\partial_x Q_1)^2, \quad (\text{S53})$$

with the boundary conditions (S33) which yield

$$P_2(x, 1) = 0, \quad Q_2(x, 0) = \frac{\sigma(\bar{\rho})}{2D(\bar{\rho})} P_2(x, 0) + \left(\frac{\sigma'(\bar{\rho})}{2\sigma(\bar{\rho})} - \frac{D'(\bar{\rho})}{2D(\bar{\rho})} \right) Q_1(x, 0)^2. \quad (\text{S54})$$

Setting

$$\begin{aligned}
P_2(x, t) &= \frac{\sigma'(\bar{\rho})}{16D(\bar{\rho})} \operatorname{erfc}\left(\frac{x + \nu\sigma'(\bar{\rho})}{2\sqrt{D(\bar{\rho})(1-t)}}\right) \operatorname{erfc}\left(-\frac{x + \nu\sigma'(\bar{\rho})}{2\sqrt{D(\bar{\rho})(1-t)}}\right) \\
&+ \frac{\nu\sigma(\bar{\rho})\sigma''(\bar{\rho})}{8D(\bar{\rho})^2} \left(\sqrt{\frac{D(\bar{\rho})(1-t)}{\pi}} e^{-\frac{(x+\nu\sigma'(\bar{\rho}))^2}{4D(\bar{\rho})(1-t)}} + \frac{x + \nu\sigma'(\bar{\rho})}{2} \operatorname{erfc}\left(-\frac{x + \nu\sigma'(\bar{\rho})}{2\sqrt{D(\bar{\rho})(1-t)}}\right) \right) \operatorname{erfc}\left(\frac{x + \nu\sigma'(\bar{\rho})}{2\sqrt{D(\bar{\rho})(1-t)}}\right) + \tilde{P}_2(x, t),
\end{aligned} \tag{S55}$$

we find that \tilde{P}_2 obeys

$$\begin{aligned}
\partial_t \tilde{P}_2 + D(\bar{\rho}) \partial_x^2 \tilde{P}_2 &= \frac{\nu\sigma(\bar{\rho})\sigma''(\bar{\rho})}{8D(\bar{\rho})} \frac{e^{-\frac{(x+\nu\sigma'(\bar{\rho}))^2}{4D(\bar{\rho})(1-t)}}}{\sqrt{D(\bar{\rho})\pi(1-t)}} \operatorname{erfc}\left(-\frac{x}{2\sqrt{D(\bar{\rho})t}}\right) \\
&+ \frac{e^{-\frac{(x+\nu\sigma'(\bar{\rho}))^2}{4D(\bar{\rho})(1-t)}}}{16D(\bar{\rho})^{5/2}\sqrt{\pi}(1-t)^{3/2}} D'(\bar{\rho})\sigma(\bar{\rho})(x + \nu\sigma'(\bar{\rho})) \left(\operatorname{erfc}\left(\frac{x}{2\sqrt{D(\bar{\rho})t}}\right) - \operatorname{erfc}\left(\frac{x + \nu\sigma'(\bar{\rho})}{2\sqrt{D(\bar{\rho})(1-t)}}\right) \right), \quad \tilde{P}_2(x, 1) = 0.
\end{aligned} \tag{S56}$$

Then setting one more time:

$$\tilde{P}_2(2\sqrt{D(\bar{\rho})}x, t) = \frac{\sigma(\bar{\rho})D'(\bar{\rho})}{D(\bar{\rho})^{5/2}} \tilde{P}_{2d}(x, t) + \frac{\nu\sigma(\bar{\rho})\sigma''(\bar{\rho})}{D(\bar{\rho})^{3/2}} \tilde{P}_{2s}(x, t) + \frac{\sigma(\bar{\rho})D'(\bar{\rho})}{D(\bar{\rho})^2} \tilde{f}_{2p} \left(\frac{\frac{\nu\sigma'(\bar{\rho})}{2\sqrt{D(\bar{\rho})}} + x}{\sqrt{1-t}} \right), \tag{S57}$$

where we have defined

$$\tilde{f}_{2p}(x) = \frac{x e^{-x^2}}{8\sqrt{\pi}} (\operatorname{erfc}(x) - 2) - \frac{e^{-2x^2}}{8\pi}. \tag{S58}$$

Inserting that expression of \tilde{P}_2 in (S56) we find that \tilde{P}_{2d} and \tilde{P}_{2s} obey the following anti-diffusion equations:

$$\partial_t \tilde{P}_{2d} + \frac{1}{4} \partial_x^2 \tilde{P}_{2d} = \frac{\operatorname{erfc}\left(-\frac{x}{\sqrt{t}}\right) \left(2x\sqrt{D(\bar{\rho})} + \nu\sigma'(\bar{\rho})\right) \exp\left(-\frac{(2x\sqrt{D(\bar{\rho})} + \nu\sigma'(\bar{\rho}))^2}{4(1-t)D(\bar{\rho})}\right)}{16\sqrt{\pi}(1-t)^{3/2}}, \tag{S59}$$

$$\partial_t \tilde{P}_{2s} + \frac{1}{4} \partial_x^2 \tilde{P}_{2s} = \frac{\operatorname{erfc}\left(-\frac{x}{\sqrt{t}}\right) \exp\left(-\frac{(2x\sqrt{D(\bar{\rho})} + \nu\sigma'(\bar{\rho}))^2}{4(1-t)D(\bar{\rho})}\right)}{8\sqrt{\pi}(1-t)}. \tag{S60}$$

We can implement the initial condition (S54) for Q_2 by defining \tilde{Q}_2 as

$$\begin{aligned}
Q_2(x, t) &= \frac{\sigma(\bar{\rho})}{2D(\bar{\rho})} P_2(x, t) + \left(\frac{\sigma'(\bar{\rho})}{2\sigma(\bar{\rho})} - \frac{D'(\bar{\rho})}{2D(\bar{\rho})} \right) Q_1(x, t)^2 \\
&+ \frac{\sigma(\bar{\rho})\sigma'(\bar{\rho})}{32D(\bar{\rho})^2} \operatorname{erfc}\left(\frac{x}{2\sqrt{D(\bar{\rho})t}}\right) \operatorname{erfc}\left(\frac{-x}{2\sqrt{D(\bar{\rho})t}}\right) + \tilde{Q}_2\left(\frac{x}{2\sqrt{D(\bar{\rho})}}, t\right).
\end{aligned} \tag{S61}$$

with $\tilde{Q}_2(x, 0) = 0$.

As for \tilde{P}_2 we set:

$$\tilde{Q}_2(x, t) = \frac{\sigma(\bar{\rho})^2 D'(\bar{\rho}) \left(\tilde{Q}_{2d}(x, t) + \frac{1}{2} \tilde{f}_{2p}\left(\frac{x}{\sqrt{t}}\right) \right)}{D(\bar{\rho})^3} + \frac{\nu\sigma(\bar{\rho})^2 \sigma''(\bar{\rho}) \left(\tilde{Q}_{2s}(x, t) + \sqrt{t} \tilde{f}_{2s}\left(\frac{x}{\sqrt{t}}\right) \right)}{D(\bar{\rho})^{5/2}}, \tag{S62}$$

where we have introduced:

$$\tilde{f}_{2s}(x, t) = \frac{1}{16} \left(x(\operatorname{erfc}(x) - 2) - \frac{e^{-x^2}}{\sqrt{\pi}} \right) \operatorname{erfc}(x). \quad (\text{S63})$$

Injecting that expression in (S62), one deduces the following diffusion equations:

$$\partial_t \tilde{Q}_{2d} - \frac{1}{4} \partial_x^2 \tilde{Q}_{2d} = \frac{x e^{-\frac{x^2}{t}} \operatorname{erfc} \left(-\frac{2x\sqrt{D(\bar{\rho})} + \nu\sigma'(\bar{\rho})}{2\sqrt{(1-t)D(\bar{\rho})}} \right)}{16\sqrt{\pi}t^{3/2}}, \quad \tilde{Q}_{2d}(x, 0) = 0, \quad (\text{S64})$$

$$\partial_t \tilde{Q}_{2s} - \frac{1}{4} \partial_x^2 \tilde{Q}_{2s} = \frac{e^{-\frac{x^2}{t}} \operatorname{erfc} \left(-\frac{2x\sqrt{D(\bar{\rho})} + \nu\sigma'(\bar{\rho})}{2\sqrt{(1-t)D(\bar{\rho})}} \right)}{16\sqrt{\pi}\sqrt{t}}, \quad \tilde{Q}_{2s}(x, 0) = 0. \quad (\text{S65})$$

The solutions of the equations for $\tilde{P}_{2d}, \tilde{P}_{2s}, \tilde{Q}_{2d}$ and \tilde{Q}_{2s} (S59, S60, S64, S65) are related by

$$\tilde{P}_{2d}(x, t; \nu) = 2\sqrt{D(\bar{\rho})} \tilde{Q}_{2d} \left(\frac{\nu\sigma'(\bar{\rho})}{2\sqrt{D(\bar{\rho})}} + x, 1-t; -\nu \right) \quad (\text{S66})$$

$$\tilde{P}_{2s}(x, t; \nu) = -2\tilde{Q}_{2s} \left(x - \frac{\nu\sigma'(\bar{\rho})}{2\sqrt{D(\bar{\rho})}}, 1-t; -\nu \right) \quad (\text{S67})$$

where we have intraduced a third argument $\pm\nu$ to notice that in the expression of \tilde{Q}_{2d} and \tilde{Q}_{2s} one has to replace ν by $-\nu$ to deduce \tilde{P}_{2d} and \tilde{P}_{2s} . We now turn to the determination of these functions.

1. The term in σ''

Our goal in this section is to determine the function \tilde{Q}_{2s} . Inserting the change of variable $\nu \rightarrow \frac{2Y\sqrt{D(\bar{\rho})}}{\sigma'(\bar{\rho})}$ in (S65), one gets:

$$\partial_t \tilde{Q}_{2s} - \frac{1}{4} \partial_x^2 \tilde{Q}_{2s} = \frac{1}{16} \frac{e^{-\frac{x^2}{t}} \operatorname{erfc} \left(-\frac{x+Y}{\sqrt{1-t}} \right)}{\sqrt{\pi}\sqrt{t}} \quad (\text{S68})$$

Formally we can write the solution as

$$\tilde{Q}_{2s}(x, t|Y) = \int_0^t dt' \int_{-\infty}^{\infty} dy K(x-y|t-t') \frac{e^{-y^2/t'}}{16\sqrt{\pi}t'} \operatorname{erfc} \left(-\frac{y+Y}{\sqrt{1-t'}} \right), \quad (\text{S69})$$

with

$$K(x|t) = \frac{e^{-x^2/t}}{\sqrt{\pi}t} \quad (\text{S70})$$

the heat kernel. The integral over y can be performed analytically [S8]. Since we only need \tilde{P}_2 at $t = 0$ and \tilde{Q}_2 at $t = 1$, we only compute

$$\tilde{Q}_{2s}(x, 1|Y) = \frac{e^{-x^2}}{16\sqrt{\pi}} \int_0^1 dt \operatorname{erfc} \left(-\frac{Y+tx}{\sqrt{1-t^2}} \right) \equiv I(x, Y). \quad (\text{S71})$$

To determine this function, we notice that the integral can be simplified by computing

$$\partial_x I(x, Y) + 2xI(x, Y) = \int_0^1 dt \frac{e^{-\frac{x^2+2txY+Y^2}{1-t^2}}}{8\pi\sqrt{1-t^2}} dt. \quad (\text{S72})$$

Using the method of Refs. [S9, S10] which relies on a numerical evaluation of the integral, we find that

$$\begin{aligned} \partial_x I(x) + 2xI(x) &= \frac{1}{8\pi} \left[x \operatorname{erfc}(x \operatorname{sign}(x+Y)) - \operatorname{sign}(x+Y) \frac{e^{-x^2}}{\sqrt{\pi}} \right] \\ &\quad \times \left[Y \operatorname{erfc}(Y \operatorname{sign}(x+Y)) - \operatorname{sign}(x+Y) \frac{e^{-Y^2}}{\sqrt{\pi}} \right]. \end{aligned} \quad (\text{S73})$$

Integrating this differential equation, we find that

$$\tilde{Q}_{2s}(x, 1) = I(x, Y) = \Theta(x+Y) \frac{e^{-x^2}}{8\sqrt{\pi}} + \frac{1}{16} \operatorname{erfc}(y \operatorname{sign}(x+Y)) \left[Y \operatorname{erfc}(Y \operatorname{sign}(x+Y)) - \operatorname{sign}(x+Y) \frac{e^{-Y^2}}{\sqrt{\pi}} \right]. \quad (\text{S74})$$

Using the relation (S66) we obtain immediately the expression of \tilde{P}_{2s} at $t=0$, which will be useful below to compute $q(x, 0)$,

$$\tilde{P}_{2s}(x, 0) = -\frac{e^{-(x+Y)^2}}{4\sqrt{\pi}} \Theta(x) + \frac{1}{8} \operatorname{erfc}((x+Y) \operatorname{sign}(x)) \left[Y \operatorname{erfc}(-Y \operatorname{sign}(x)) + \operatorname{sign}(x) \frac{e^{-Y^2}}{\sqrt{\pi}} \right]. \quad (\text{S75})$$

2. The term in D'

We follow the same procedure to write \tilde{Q}_{2d} from (S64) as

$$\tilde{Q}_{2d}(x, 1) = \int_0^1 dt' \int_{-\infty}^{\infty} dy K(x-y|t-t') \frac{y e^{-\frac{y^2}{t'}} \operatorname{erfc}\left(-\frac{y+Y}{\sqrt{1-t'}}\right)}{16\sqrt{\pi}(t')^{3/2}}. \quad (\text{S76})$$

Performing the integral over y using [S8], we get,

$$\tilde{Q}_{2d}(x, 1) = x e^{-x^2} \int_0^1 \frac{dt}{16\sqrt{\pi}} \operatorname{erfc}\left(-\frac{tx+Y}{\sqrt{1-t^2}}\right) + e^{-x^2} \int_0^1 \frac{dt}{16\pi} \sqrt{\frac{1-t}{1+t}} e^{-\frac{(tx+Y)^2}{1-t^2}}. \quad (\text{S77})$$

We notice that this function can be expressed in terms of the integral $I(x, Y)$ (S71) computed previously as

$$\tilde{Q}_{2d}(x, 1) = \frac{1}{2} \partial_x (I(Y, x) - I(x, Y)). \quad (\text{S78})$$

Therefore, we directly obtain from (S74),

$$\tilde{Q}_{2d}(x, 1) = -\frac{e^{-x^2-Y^2}}{16\pi} + \frac{x e^{-x^2}}{8\sqrt{\pi}} \Theta(x+Y) + \frac{1}{32} \operatorname{erfc}(Y \operatorname{sign}(x+Y)) \left[\operatorname{erfc}(x \operatorname{sign}(x+Y)) + 2Y \operatorname{sign}(x+Y) \frac{e^{-x^2}}{\sqrt{\pi}} \right]. \quad (\text{S79})$$

Using (S66), we deduce

$$\begin{aligned} \frac{\tilde{P}_{2d}(x, 0)}{2\sqrt{D(\bar{\rho})}} &= -\frac{e^{-(x+Y)^2-Y^2}}{16\pi} + \frac{(x+Y) e^{-(x+Y)^2}}{8\sqrt{\pi}} \Theta(x) \\ &\quad + \frac{1}{32} \operatorname{erfc}(Y \operatorname{sign}(x)) \left[\operatorname{erfc}((x+Y) \operatorname{sign}(x)) + 2Y \operatorname{sign}(x) \frac{e^{-(x+Y)^2}}{\sqrt{\pi}} \right]. \end{aligned} \quad (\text{S80})$$

3. Summary

Using the relations (S61,S62,S74,S75) and (S44), we obtain the solution of the MFT equations at order 2 in λ , at initial and final time,

$$\begin{aligned}
q_2(x, 1) = & \frac{\sigma(\bar{\rho})}{32D(\bar{\rho})^3} \left(D(\bar{\rho})\sigma'(\bar{\rho}) \operatorname{erfc} \left(\frac{x - \nu\sigma'(\bar{\rho})}{2\sqrt{D(\bar{\rho})}} \right) \operatorname{erfc} \left(\frac{\nu\sigma'(\bar{\rho}) - x}{2\sqrt{D(\bar{\rho})}} \right) - (\sigma(\bar{\rho})d'(\bar{\rho}) - D(\bar{\rho})\sigma'(\bar{\rho})) \left(\operatorname{erfc} \left(\frac{\nu\sigma'(\bar{\rho}) - x}{2\sqrt{D(\bar{\rho})}} \right) - 2\Theta(x) \right) \right) \\
& + \frac{\sigma(\bar{\rho})d'(\bar{\rho})e^{-\frac{3\nu^2\sigma'(\bar{\rho})^2+2x^2}{4D(\bar{\rho})}}}{\pi\sqrt{D(\bar{\rho})}} \left(\sqrt{\pi}e^{\frac{2\nu^2\sigma'(\bar{\rho})^2+x^2+2\nu x\sigma'(\bar{\rho})}{4D(\bar{\rho})}} \left(\nu\sigma'(\bar{\rho}) \left(\operatorname{sign}(x) \operatorname{erfc} \left(\frac{\nu\operatorname{sign}(x)\sigma'(\bar{\rho})}{2\sqrt{D(\bar{\rho})}} \right) \right) \right. \right. \\
& \left. \left. - \operatorname{erfc} \left(\frac{x - \nu\sigma'(\bar{\rho})}{2\sqrt{D(\bar{\rho})}} \right) + 2 \right) + x \left(\operatorname{erfc} \left(\frac{x - \nu\sigma'(\bar{\rho})}{2\sqrt{D(\bar{\rho})}} \right) - 2 \right) + 2\Theta(x)(x - \nu\sigma'(\bar{\rho})) \right) \\
& + \sqrt{D(\bar{\rho})} \left(\pi e^{\frac{3\nu^2\sigma'(\bar{\rho})^2+2x^2}{4D(\bar{\rho})}} \operatorname{erfc} \left(\frac{\nu\operatorname{sign}(x)\sigma'(\bar{\rho})}{2\sqrt{D(\bar{\rho})}} \right) \operatorname{erfc} \left(\frac{\operatorname{sign}(x)(x - \nu\sigma'(\bar{\rho}))}{2\sqrt{D(\bar{\rho})}} \right) - 2 \left(e^{\frac{\nu\sigma'(\bar{\rho})(\nu\sigma'(\bar{\rho})+4x)}{4D(\bar{\rho})}} + e^{\frac{(\nu\sigma'(\bar{\rho})+x)^2}{4D(\bar{\rho})}} \right) \right) \\
& \frac{2\nu\sqrt{D(\bar{\rho})}\sigma(\bar{\rho})\sigma''(\bar{\rho})e^{-\frac{\nu^2\sigma'(\bar{\rho})^2+x^2}{4D(\bar{\rho})}} \left(\operatorname{sign}(x) e^{\frac{x^2}{4D(\bar{\rho})}} \operatorname{erfc} \left(\frac{\operatorname{sign}(x)(x - \nu\sigma'(\bar{\rho}))}{2\sqrt{D(\bar{\rho})}} \right) + e^{\frac{\nu x\sigma'(\bar{\rho})}{2D(\bar{\rho})}} \operatorname{erfc} \left(\frac{x - \nu\sigma'(\bar{\rho})}{2\sqrt{D(\bar{\rho})}} \right) - 2\Theta(x)e^{\frac{\nu x\sigma'(\bar{\rho})}{2D(\bar{\rho})}} \right)}{\sqrt{\pi}} \\
& + \nu\sigma(\bar{\rho})\sigma''(\bar{\rho}) \left(\nu\sigma'(\bar{\rho}) \left(\operatorname{erfc} \left(\frac{\nu\operatorname{sign}(x)\sigma'(\bar{\rho})}{2\sqrt{D(\bar{\rho})}} \right) \operatorname{erfc} \left(\frac{\operatorname{sign}(x)(x - \nu\sigma'(\bar{\rho}))}{2\sqrt{D(\bar{\rho})}} \right) - \operatorname{erfc} \left(\frac{x - \nu\sigma'(\bar{\rho})}{2\sqrt{D(\bar{\rho})}} \right)^2 + 2\operatorname{erfc} \left(\frac{x - \nu\sigma'(\bar{\rho})}{2\sqrt{D(\bar{\rho})}} \right) \right) \right. \\
& \left. x \left(\operatorname{erfc} \left(\frac{x - \nu\sigma'(\bar{\rho})}{2\sqrt{D(\bar{\rho})}} \right) - 2 \right) \operatorname{erfc} \left(\frac{x - \nu\sigma'(\bar{\rho})}{2\sqrt{D(\bar{\rho})}} \right) \right), \quad (\text{S81})
\end{aligned}$$

$$\begin{aligned}
q_2(x, 0) = & \frac{\sigma(\bar{\rho})}{32D(\bar{\rho})^3} \left(D(\bar{\rho})\sigma'(\bar{\rho}) \operatorname{erfc} \left(-\frac{\nu\sigma'(\bar{\rho}) + x}{2\sqrt{D(\bar{\rho})}} \right) \operatorname{erfc} \left(\frac{\nu\sigma'(\bar{\rho}) + x}{2\sqrt{D(\bar{\rho})}} \right) - \frac{2\sigma(\bar{\rho})D'(\bar{\rho})e^{-\frac{(\nu\sigma'(\bar{\rho})+x)^2}{2D(\bar{\rho})}}}{\pi} \right) \\
& + \frac{\sigma(\bar{\rho})D'(\bar{\rho})(\nu\sigma'(\bar{\rho}) + x)e^{-\frac{(\nu\sigma'(\bar{\rho})+x)^2}{4D(\bar{\rho})}} \left(\operatorname{erfc} \left(\frac{\nu\sigma'(\bar{\rho})+x}{2\sqrt{D(\bar{\rho})}} \right) - 2 \right) - (\sigma(\bar{\rho})D'(\bar{\rho}) - D(\bar{\rho})\sigma'(\bar{\rho})) \left(\operatorname{erfc} \left(-\frac{\nu\sigma'(\bar{\rho}) + x}{2\sqrt{D(\bar{\rho})}} \right) - 2\Theta(x) \right)^2}{\sqrt{\pi}\sqrt{D(\bar{\rho})}} \\
& + \frac{\sigma(\bar{\rho})D'(\bar{\rho})e^{-\frac{2\nu^2\sigma'(\bar{\rho})^2+x^2+2\nu x\sigma'(\bar{\rho})}{4D(\bar{\rho})}}}{\pi\sqrt{D(\bar{\rho})}} \left(\sqrt{\pi}e^{\frac{\nu^2\sigma'(\bar{\rho})^2}{4D(\bar{\rho})}} \left(2\Theta(x)(\nu\sigma'(\bar{\rho}) + x) - \nu\operatorname{sign}(x)\sigma'(\bar{\rho}) \operatorname{erfc} \left(-\frac{\nu\operatorname{sign}(x)\sigma'(\bar{\rho})}{2\sqrt{D(\bar{\rho})}} \right) \right) \right) \\
& + \sqrt{D(\bar{\rho})} \left(\pi e^{\frac{2\nu^2\sigma'(\bar{\rho})^2+x^2+2\nu x\sigma'(\bar{\rho})}{4D(\bar{\rho})}} \operatorname{erfc} \left(-\frac{\nu\operatorname{sign}(x)\sigma'(\bar{\rho})}{2\sqrt{D(\bar{\rho})}} \right) \operatorname{erfc} \left(\frac{\operatorname{sign}(x)(\nu\sigma'(\bar{\rho}) + x)}{2\sqrt{D(\bar{\rho})}} \right) - 2 \right) \\
& + \frac{2\nu\sqrt{D(\bar{\rho})}\sigma(\bar{\rho})\sigma''(\bar{\rho})e^{-\frac{(\nu\sigma'(\bar{\rho})+x)^2}{4D(\bar{\rho})}} \operatorname{erfc} \left(\frac{\nu\sigma'(\bar{\rho})+x}{2\sqrt{D(\bar{\rho})}} \right) + \nu\sigma(\bar{\rho})\sigma''(\bar{\rho})(\nu\sigma'(\bar{\rho}) + x) \operatorname{erfc} \left(-\frac{\nu\sigma'(\bar{\rho}) + x}{2\sqrt{D(\bar{\rho})}} \right) \operatorname{erfc} \left(\frac{\nu\sigma'(\bar{\rho}) + x}{2\sqrt{D(\bar{\rho})}} \right)}{\sqrt{\pi}} \\
& - \frac{2\nu\sqrt{D(\bar{\rho})}\sigma(\bar{\rho})\sigma''(\bar{\rho}) \left(\frac{1}{2} \operatorname{erfc} \left(\frac{\operatorname{sign}(x)(\nu\sigma'(\bar{\rho}) + x)}{2\sqrt{D(\bar{\rho})}} \right) \left(-\frac{\sqrt{\pi}\nu\sigma'(\bar{\rho}) \operatorname{erfc} \left(-\frac{\nu\operatorname{sign}(x)\sigma'(\bar{\rho})}{2\sqrt{D(\bar{\rho})}} \right)}{\sqrt{D(\bar{\rho})}} - 2\operatorname{sign}(x)e^{-\frac{\nu^2\sigma'(\bar{\rho})^2}{4D(\bar{\rho})}} \right) \right)}{\sqrt{\pi}} \\
& \left. + 2\Theta(x)e^{-\frac{(\nu\sigma'(\bar{\rho})+x)^2}{4D(\bar{\rho})}} \right). \quad (\text{S82})
\end{aligned}$$

IV. CUMULANTS AND PROFILES FOR THE INTEGRATED CURRENT

A. Cumulants

We can directly compute the cumulants of Q_T from the profiles at the first orders (S51,S52,S81,S82) using the relation (S35), which expanded in powers of λ reads

$$\frac{d\hat{\psi}}{d\lambda} = \sum_{n=0}^{\infty} \frac{\lambda^n}{n!} \hat{\kappa}_{n+1} = \nu\sigma(\bar{\rho}) + \sum_{n=1}^{\infty} \int_0^{\infty} [q_n(x,1) - q_n(x,0)] dx. \quad (\text{S83})$$

Note that the first cumulant is already determined by the first term of (S83), while the higher-order cumulants are encoded in the second. We thus get

$$\hat{\kappa}_1 = \nu\sigma(\bar{\rho}), \quad (\text{S84})$$

$$\hat{\kappa}_2 = \int_0^{\infty} (q_1(x,1) - q_1(x,0)) dx = \frac{\sigma(\bar{\rho})}{2\sqrt{D(\bar{\rho})}} \left(\frac{2}{\sqrt{\pi}} e^{-\frac{(\nu\sigma'(\bar{\rho}))^2}{4D(\bar{\rho})}} + \frac{\nu\sigma'(\bar{\rho})}{\sqrt{D(\bar{\rho})}} \operatorname{erf} \left(\frac{\nu\sigma'(\bar{\rho})}{2\sqrt{D(\bar{\rho})}} \right) \right), \quad (\text{S85})$$

$$\begin{aligned} \hat{\kappa}_3 = & \frac{\nu\sigma(\bar{\rho})}{16D(\bar{\rho})^3} \left(3\nu^2\sigma(\bar{\rho})\sigma'(\bar{\rho})^2\sigma''(\bar{\rho}) + 4D(\bar{\rho}) \left(\sigma'(\bar{\rho})^2 + \left(1 - \frac{3e^{-\frac{\nu^2\sigma'(\bar{\rho})^2}{2D(\bar{\rho})}}}{\pi} \right) \sigma(\bar{\rho})\sigma''(\bar{\rho}) \right) \right. \\ & \left. + 3\nu\sigma(\bar{\rho})\sigma'(\bar{\rho})\sigma''(\bar{\rho}) \operatorname{erf} \left(\frac{\nu\sigma'(\bar{\rho})}{2\sqrt{D(\bar{\rho})}} \right) \left(-\frac{4\sqrt{D(\bar{\rho})}e^{-\frac{\nu^2\sigma'(\bar{\rho})^2}{4D(\bar{\rho})}}}{\sqrt{\pi}} - \nu\sigma'(\bar{\rho}) \operatorname{erf} \left(\frac{\nu\sigma'(\bar{\rho})}{2\sqrt{D(\bar{\rho})}} \right) \right) \right) \\ & \left. + \frac{D'(\bar{\rho})\sigma(\bar{\rho})^2}{8D(\bar{\rho})^3} \left(\nu\sigma'(\bar{\rho}) - \frac{6\sqrt{D(\bar{\rho})} \operatorname{erf} \left(\frac{\nu\sigma'(\bar{\rho})}{2\sqrt{D(\bar{\rho})}} \right)}{\sqrt{\pi}} - 3\nu\sigma'(\bar{\rho}) \operatorname{erf} \left(\frac{\nu\sigma'(\bar{\rho})}{2\sqrt{D(\bar{\rho})}} \right) \right)^2 \right), \quad (\text{S86}) \end{aligned}$$

which correspond to Eqs. (10,11) in the main text.

B. Correlation profile

In order to get the correlation profiles at the microscopic large time T we remark that we can write their generating function as [S11],

$$\begin{aligned} \Phi(x) \equiv \frac{\langle \rho(x,1)e^{\lambda Q_T} \rangle}{\langle e^{\lambda Q_T} \rangle} &= \frac{\int \mathcal{D}\rho \mathcal{D}H \rho(x,1) e^{-\sqrt{T}(S[\rho,H]+F[\rho(x,0)]-\lambda\hat{Q}[\rho])}}{\int \mathcal{D}\rho \mathcal{D}H e^{-\sqrt{T}(S[\rho,H]+F[\rho(x,0)]-\lambda\hat{Q}[\rho])}} \\ &\simeq \frac{q(x,1) e^{-\sqrt{T}(S[q,p]+F[q(x,0)]-\lambda\hat{Q}[q])}}{e^{-\sqrt{T}(S[q,p]+F[q(x,0)]-\lambda\hat{Q}[q])}} \\ &= q(x,1) \end{aligned} \quad (\text{S87})$$

because (q,p) is the saddle point of $S[\rho,H] + F[\rho(x,0)] - \lambda\hat{Q}[\rho]$, using the expansion (S43) and the results of our computations for the profiles (S51,S81), we deduce

$$\langle \rho(x,1)Q_T \rangle_c \simeq q_1(x,1) = \operatorname{sign}(x) \frac{\sigma(\bar{\rho})}{4D(\bar{\rho})} \operatorname{erfc} \left(\operatorname{sign}(x) \frac{x - \nu\sigma'(\bar{\rho})}{2\sqrt{D(\bar{\rho})}} \right), \quad (\text{S88})$$

$$\langle \rho(x,1)Q_T^2 \rangle_c \simeq q_2(x,1) \quad (\text{S89})$$

where q_2 is given by (S81) and the index c refers to the connected moments (for example $\langle \rho(x,1)Q_T \rangle_c = \langle \rho(x,1)Q_T \rangle - \langle \rho(x,1) \rangle \langle Q_T \rangle$ is the covariance, and higher order connected moments generalize it).

V. CUMULANTS AND PROFILES FOR THE TRACER

A. Cumulants

Using the duality mapping presented in II D, we deduce the first three normalized cumulants of a tracer initially at the origin:

$$\frac{\langle X_T \rangle}{\sqrt{T}} \underset{T \rightarrow \infty}{\simeq} \nu \frac{\sigma(\bar{\rho})}{\bar{\rho}}, \quad (\text{S90})$$

$$\frac{\langle X_T^2 \rangle_c}{\sqrt{T}} \underset{T \rightarrow \infty}{\simeq} \frac{\sigma(\bar{\rho})}{2\bar{\rho}^3 D(\bar{\rho})} \left(\nu (\sigma(\bar{\rho}) - \bar{\rho}\sigma'(\bar{\rho})) \operatorname{erf} \left(\frac{\nu (\sigma(\bar{\rho}) - \bar{\rho}\sigma'(\bar{\rho}))}{2\sqrt{\bar{\rho}^2 D(\bar{\rho})}} \right) + \frac{2\sqrt{\bar{\rho}^2 D(\bar{\rho})} \exp \left(-\frac{\nu^2 (\sigma(\bar{\rho}) - \bar{\rho}\sigma'(\bar{\rho}))^2}{4\bar{\rho}^2 D(\bar{\rho})} \right)}{\sqrt{\pi}} \right) \quad (\text{S91})$$

$$\begin{aligned} \frac{\langle X_T^3 \rangle_c}{\sqrt{T}} \underset{T \rightarrow \infty}{\simeq} & \frac{\sigma(\bar{\rho})}{16\bar{\rho}^5 D(\bar{\rho})^3} \left((\bar{\rho}D'(\bar{\rho}) + 2D(\bar{\rho}) + \nu^2 (\bar{\rho}\sigma'(\bar{\rho}) - \sigma(\bar{\rho}))\sigma''(\bar{\rho})) \right. \\ & \times \frac{12\bar{\rho}\sqrt{D(\bar{\rho})}\sigma(\bar{\rho})}{\sqrt{\pi}} \operatorname{erf} \left(\frac{\nu (\sigma(\bar{\rho}) - \bar{\rho}\sigma'(\bar{\rho}))}{2\bar{\rho}\sqrt{D(\bar{\rho})}} \right) \exp \left(-\frac{\nu^2 (\sigma(\bar{\rho}) - \bar{\rho}\sigma'(\bar{\rho}))^2}{4\bar{\rho}^2 D(\bar{\rho})} \right) \\ & + \nu\sigma(\bar{\rho}) (\sigma(\bar{\rho}) - \bar{\rho}\sigma'(\bar{\rho})) \left(2(\bar{\rho}D'(\bar{\rho}) + 2D(\bar{\rho})) \left(3\operatorname{erf} \left(\frac{\nu (\sigma(\bar{\rho}) - \bar{\rho}\sigma'(\bar{\rho}))}{2\bar{\rho}\sqrt{D(\bar{\rho})}} \right)^2 - 1 \right) \right. \\ & \left. \left. - 3\nu^2 (\sigma(\bar{\rho}) - \bar{\rho}\sigma'(\bar{\rho}))\sigma''(\bar{\rho}) \left(\operatorname{erf} \left(\frac{\nu (\sigma(\bar{\rho}) - \bar{\rho}\sigma'(\bar{\rho}))}{2\bar{\rho}\sqrt{D(\bar{\rho})}} \right)^2 - 1 \right) \right) \right) \\ & \left. + 4\nu D(\bar{\rho}) \left(\bar{\rho}^2 \sigma(\bar{\rho})\sigma''(\bar{\rho}) \left(1 - \frac{3 \exp \left(-\frac{\nu^2 (\sigma(\bar{\rho}) - \bar{\rho}\sigma'(\bar{\rho}))^2}{2\bar{\rho}^2 D(\bar{\rho})} \right)}{\pi} \right) + (\sigma(\bar{\rho}) - \bar{\rho}\sigma'(\bar{\rho}))^2 \right) \right). \quad (\text{S92}) \end{aligned}$$

B. Correlation profiles

The duality mapping (S37) is written for the stochastic fields ρ and j . In the limit $T \rightarrow \infty$, the MFT action is dominated by a specific realisation of these fields, and in particular an optimal density given by $q(x, t)$. Therefore, we can deduce the correlation profiles in the reference frame of the tracer from those obtained for Q_T (S87) as

$$\Phi_T(x) \equiv \frac{\left\langle \rho \left(x + \frac{X_T}{\sqrt{T}}, 1 \right) e^{\lambda X_T} \right\rangle}{\left\langle e^{\lambda X_T} \right\rangle} = \frac{1}{\Phi(y(x))} \Big|_{D \rightarrow \bar{D}, \sigma \rightarrow \bar{\sigma}, \nu \rightarrow \bar{\nu}}, \quad y(x) = \int_0^x \Phi(x') dx', \quad (\text{S93})$$

where we have again used the scaling of Section I to express the correlations in terms of the macroscopic density $\rho(x, 1)$ which represents the microscopic system at the observation time T . For instance, we obtain at first order in λ ,

$$\left\langle X_T \rho \left(x + \frac{X_T}{\sqrt{T}}, 1 \right) \right\rangle_c \underset{T \rightarrow \infty}{\simeq} \frac{\sigma(\bar{\rho})}{4\bar{\rho}D(\bar{\rho})} \operatorname{erfc} \left(\frac{x - \nu \frac{\sigma(\bar{\rho}) - \bar{\rho}\sigma'(\bar{\rho})}{\bar{\rho}}}{2\sqrt{D(\bar{\rho})}} \right), \quad \text{for } x > 0. \quad (\text{S94})$$

As announced in the main text, the spatial dependence of this profile is identical to the one obtained in the case of the current (S88), but with $\sigma'(\bar{\rho})$ replaced by $\frac{\sigma(\bar{\rho}) - \bar{\rho}\sigma'(\bar{\rho})}{\bar{\rho}}$.

VI. LARGE ASYMMETRY: BALLISTIC LIMIT

The situation of driven diffusive system can be thought as representing the dynamics of driven ballistic systems at a large but finite time T . Indeed, for fixed F and large T , choosing $\nu = F\sqrt{T}$ should allow to describe the properties of the system at time T . An interesting question is to look at the limit of large asymmetry $\nu \rightarrow \infty$ at fixed F . Can we obtain the large times limit properties of the initial ballistic system by taking that limit? This convergence is not a priori obvious, as the hydrodynamic scalings in diffusive and ballistic systems are of different nature: for a system of size L , the time scales like L^2 for diffusive dynamics, and like L for the ballistic case. Then there is no reason that the limits $\nu \rightarrow \infty$ and $T \rightarrow \infty$ commute. Nevertheless setting $\nu = F\sqrt{T}$ and keeping only the dominant term in (S90), (S91) and (S92), we obtain for the tracer

$$\langle X_T \rangle \underset{T \rightarrow \infty}{\simeq} T F \frac{\sigma(\bar{\rho})}{\bar{\rho}}, \quad (\text{S95})$$

$$\langle X_T^2 \rangle_c \underset{T \rightarrow \infty}{\simeq} T |F| \frac{\sigma(\bar{\rho})}{2\bar{\rho}^3 D(\bar{\rho})} |\sigma(\bar{\rho}) - \bar{\rho}\sigma'(\bar{\rho})|, \quad (\text{S96})$$

$$\langle X_T^3 \rangle_c \underset{T \rightarrow \infty}{\simeq} T F \frac{\sigma(\bar{\rho})}{4\bar{\rho}^3 D(\bar{\rho})^2} \left[\sigma''(\bar{\rho})\sigma(\bar{\rho}) + (\sigma(\bar{\rho}) - \bar{\rho}\sigma'(\bar{\rho})) \left(\frac{3\sigma(\bar{\rho}) - \bar{\rho}\sigma'(\bar{\rho})}{\bar{\rho}^2} + \frac{\sigma(\bar{\rho})D'(\bar{\rho})}{\bar{\rho}D(\bar{\rho})} \right) \right]. \quad (\text{S97})$$

These results correspond to Eqs. (15-17) of the main text. Performing the same procedure for the integrated current given by equations (S84-S86), we obtain

$$\langle Q_T \rangle = \sqrt{T}\hat{\kappa}_1 \underset{T \rightarrow \infty}{\simeq} T |F| \sigma(\bar{\rho}) \quad (\text{S98})$$

$$\langle Q_T^2 \rangle_c = \sqrt{T}\hat{\kappa}_2 \underset{T \rightarrow \infty}{\simeq} T |F| \frac{|\sigma(\bar{\rho})\sigma'(\bar{\rho})|}{2D(\bar{\rho})} \quad (\text{S99})$$

$$\langle Q_T^3 \rangle_c = \sqrt{T}\hat{\kappa}_3 \underset{T \rightarrow \infty}{\simeq} T |F| \frac{\sigma(\bar{\rho})}{4D(\bar{\rho})^2} \left[\sigma(\bar{\rho}) \left(\sigma''(\bar{\rho}) - \frac{D'(\bar{\rho})\sigma'(\bar{\rho})}{D(\bar{\rho})} \right) + \sigma'(\bar{\rho})^2 \right]. \quad (\text{S100})$$

VII. EXTENSION: CORRELATIONS BETWEEN TWO TRACERS OR TWO CURRENTS

In this section, we demonstrate how our results on the correlation profiles can be used to derive, as a byproduct, the correlations between two integrated currents measured at different points. In turn, this allows to obtain the correlations between two tracers using the duality mapping (S41). We thus consider the integrated current through a point x up to time T , which reads in terms of the macroscopic fields,

$$Q_T(x) \equiv \sqrt{T} \int_0^t j(x, t') dt'. \quad (\text{S101})$$

Note that the macroscopic variable x corresponds to a microscopic position $x\sqrt{T}$. The joint cumulant generating function of two integrated currents thus takes the form,

$$\left\langle e^{\lambda Q_T(0) + \chi Q_T(\xi)} \right\rangle = \int \mathcal{D}\rho \mathcal{D}H e^{-\sqrt{T}(S[\rho, H] + F[\rho(x, 0)] - \lambda \hat{Q}_0[\rho] - \chi \hat{Q}_\xi[\rho])}, \quad (\text{S102})$$

where we have defined

$$\hat{Q}_x[\rho] = \nu \sigma(\bar{\rho}) + \int_x^\infty [\rho(x', 1) - \rho(x', 0)] dx', \quad (\text{S103})$$

by analogy with the $x = 0$ case (S27). In the long time limit $T \rightarrow \infty$, these integrals can again be evaluated by a saddle point method, which yields

$$\hat{\psi}_\xi(\lambda, \chi) = \lim_{T \rightarrow \infty} \frac{1}{\sqrt{T}} \ln \left\langle e^{\lambda Q_T(0) + \chi Q_T(\xi)} \right\rangle = \lambda \hat{Q}_0[q] + \chi \hat{Q}_\xi[q] - S[q, p] - F[q(x, 0)]. \quad (\text{S104})$$

where we have denoted (q, p) the saddle point of (ρ, H) , for given values of λ and χ . The key point is that, to obtain the covariance between the two currents at long time T , given by

$$\frac{\langle Q_T(0) Q_T(\xi) \rangle_c}{\sqrt{T}} \underset{T \rightarrow \infty}{\simeq} \left. \frac{\partial^2 \hat{\psi}_\xi}{\partial \lambda \partial \chi} \right|_{\lambda=0, \chi=0}, \quad (\text{S105})$$

we only need the solution for $\chi = 0$ computed in Section III. Indeed, since (q, p) minimise the action in (S102), we have that

$$\left. \frac{\delta}{\delta \rho} \left(S[\rho, H] + F[\rho(x, 0)] - \lambda \hat{Q}_0[\rho] - \chi \hat{Q}_\xi[\rho] \right) \right|_{(q,p)} = 0 \quad (\text{S106})$$

and

$$\left. \frac{\delta}{\delta H} \left(S[\rho, H] + F[\rho(x, 0)] - \lambda \hat{Q}_0[\rho] - \chi \hat{Q}_\xi[\rho] \right) \right|_{(q,p)} = 0. \quad (\text{S107})$$

Therefore, the derivative with respect to χ of (S104) is given by

$$\frac{d}{d\chi} \hat{\psi}_\xi = \hat{Q}_\xi[q]. \quad (\text{S108})$$

Using this into the expression of the covariance (S105), and using that $\hat{Q}_\xi[q]$ (S103) is linear in q , we obtain

$$\frac{\langle Q_T(0) Q_T(\xi) \rangle_c}{\sqrt{T}} \underset{T \rightarrow \infty}{\simeq} \left. \frac{\partial \hat{Q}_\xi[q]}{\partial \lambda} \right|_{\lambda=0, \chi=0} = \hat{Q}_\xi \left[\left. \frac{\partial q}{\partial \lambda} \right|_{\lambda=0, \chi=0} \right] = \hat{Q}_\xi \left[\left. \frac{\partial}{\partial \lambda} \left(q \Big|_{\chi=0} \right) \right|_{\lambda=0} \right]. \quad (\text{S109})$$

This last relation involves only the optimal profile for $\chi = 0$, which is the one computed previously in Section III. Inserting the expression of q at first order in λ (S51, S52), we directly obtain,

$$\begin{aligned} \frac{\langle Q_T(0) Q_T(\xi) \rangle_c}{\sqrt{T}} \simeq \frac{\sigma(\bar{\rho})}{4D(\bar{\rho})} \left(\frac{2\sqrt{D(\bar{\rho})} e^{-\frac{(\xi + \nu\sigma'(\bar{\rho}))^2}{4D(\bar{\rho})}}}{\sqrt{\pi}} - (\xi + \nu\sigma'(\bar{\rho})) \operatorname{erfc} \left(\frac{\xi + \nu\sigma'(\bar{\rho})}{2\sqrt{D(\bar{\rho})}} \right) \right) \\ + (\nu\sigma'(\bar{\rho}) - \xi) \operatorname{erfc} \left(\frac{\xi - \nu\sigma'(\bar{\rho})}{2\sqrt{D(\bar{\rho})}} \right) + \frac{2\sqrt{D(\bar{\rho})} e^{-\frac{(\xi - \nu\sigma'(\bar{\rho}))^2}{4D(\bar{\rho})}}}{\sqrt{\pi}} \end{aligned} \quad (\text{S110})$$

From the result (S110) and the duality mapping (S41), we obtain the correlation at large time between two tracers \tilde{X}_t and \tilde{Y}_t , initially placed at $\tilde{X}_0 = 0$ and $\tilde{Y}_0 = \xi\sqrt{T}$ by performing the substitution

$$D(\rho) \rightarrow \frac{1}{\rho^2} D \left(\frac{1}{\rho} \right), \quad \sigma(\rho) \rightarrow \rho \sigma \left(\frac{1}{\rho} \right), \quad \nu \rightarrow -\nu, \quad \bar{\rho} \rightarrow \frac{1}{\bar{\rho}}. \quad (\text{S111})$$

Explicitly, this gives,

$$\begin{aligned} \frac{\langle \tilde{X}_T \tilde{Y}_T \rangle_c}{\sqrt{T}} = \frac{\sigma(\bar{\rho})}{4\bar{\rho}^3 D(\bar{\rho})} \left(\xi \left(\operatorname{erf} \left(\frac{\xi - \nu\bar{\rho}\sigma'(\bar{\rho}) + \nu\sigma(\bar{\rho})}{2\sqrt{\bar{\rho}^2 D(\bar{\rho})}} \right) + \operatorname{erf} \left(\frac{\xi + \nu\bar{\rho}\sigma'(\bar{\rho}) - \nu\sigma(\bar{\rho})}{2\sqrt{\bar{\rho}^2 D(\bar{\rho})}} \right) - 2 \right) \right. \\ \left. + \nu(\sigma(\bar{\rho}) - \bar{\rho}\sigma'(\bar{\rho})) \left(\operatorname{erf} \left(\frac{\xi - \nu\bar{\rho}\sigma'(\bar{\rho}) + \nu\sigma(\bar{\rho})}{2\sqrt{\bar{\rho}^2 D(\bar{\rho})}} \right) - \operatorname{erf} \left(\frac{\xi + \nu\bar{\rho}\sigma'(\bar{\rho}) - \nu\sigma(\bar{\rho})}{2\sqrt{\bar{\rho}^2 D(\bar{\rho})}} \right) \right) \right) \\ \left. + \frac{2\sqrt{\bar{\rho}^2 D(\bar{\rho})} \left(1 + e^{\frac{\xi\nu(\sigma(\bar{\rho}) - \bar{\rho}\sigma'(\bar{\rho}))}{\bar{\rho}^2 D(\bar{\rho})}} \right) \exp \left(-\frac{(\xi - \nu\bar{\rho}\sigma'(\bar{\rho}) + \nu\sigma(\bar{\rho}))^2}{4\bar{\rho}^2 D(\bar{\rho})} \right)}{\sqrt{\pi}} \right). \end{aligned} \quad (\text{S112})$$

This corresponds to Eq. (20) in the main text.

VIII. THE WEAKLY ASYMMETRIC SIMPLE EXCLUSION PROCESS (WASEP)

In this section, we show how we can obtain exact macroscopic equations for the WASEP from the microscopic description of the asymmetric simple exclusion process (ASEP). These equations will be useful in Section IX B below to deduce a boundary relation for the cumulant generating function of the current and the density profiles. We also give the explicit form of the first cumulants of Q_T and X_T in the case of the WASEP, deduced from our general results.

A. Generalities on the ASEP

We consider an infinite one-dimensional lattice and introduce the occupation numbers $\eta_i = 0$ or 1 for site i . Particles can jump to the left with rate p_- and to the right with rate p_+ , only if the neighbouring site is empty. The master equation associated with the probability $P_t(\{\eta\})$, where $\{\eta\} \equiv \{\eta_i\}_{i \in \mathbb{Z}}$ is a given configuration of the system, reads

$$\partial_t P_t(\{\eta\}) = \sum_i [c_i(\{\eta\}_i^+) P_t(\{\eta\}_i^+) - c_i(\{\eta\}) P_t(\{\eta\})], \quad c_i(\{\eta\}) = p_+ \eta_i (1 - \eta_{i+1}) + p_- \eta_{i+1} (1 - \eta_i). \quad (\text{S113})$$

where $\{\eta\}_i^+$ is the configuration in which η_i and η_{i+1} have been exchanged. The stationary distribution is given by the product of independent Bernoulli random variables for each site (the same as for the SEP),

$$P_{\text{stat}}(\{\eta\}) = \prod_i P_{\text{Bern}}(\eta_i), \quad P_{\text{Bern}}(\eta_i = 1) = \bar{\rho}. \quad (\text{S114})$$

B. Microscopic equations for the ASEP

From the master equation (S113), we can write the time evolution of the generating function of the current through the origin Q_t , which counts the number of particles, that have jumped from site 0 to site 1, minus the number of particles that have jumped from site 1 to 0.

$$\begin{aligned} \partial_t \langle e^{\lambda Q_t} \rangle &= \partial_t \sum_{\{\eta\}} e^{\lambda Q_t[\{\eta\}]} P_t(\{\eta\}) \\ &= \sum_{\{\eta\}} P_t(\{\eta\}) c_0(\{\eta\}) e^{\lambda Q_t[\{\eta\}]} \left(e^{\lambda Q_t[\{\eta\}_0^+]} - e^{\lambda Q_t[\{\eta\}]} \right). \end{aligned} \quad (\text{S115})$$

Since the current is only changed when a particle jumps between sites 0 and 1, with the expression for c_i (S113), we get

$$\partial_t \langle e^{\lambda Q_t} \rangle = p_+ (e^\lambda - 1) \langle \eta_0 (1 - \eta_1) e^{\lambda Q_t} \rangle + p_- (e^{-\lambda} - 1) \langle \eta_1 (1 - \eta_0) e^{\lambda Q_t} \rangle. \quad (\text{S116})$$

And then,

$$\partial_t \ln \langle e^{\lambda Q_t} \rangle = p_+ (e^\lambda - 1) \frac{\langle \eta_0 (1 - \eta_1) e^{\lambda Q_t} \rangle}{\langle e^{\lambda Q_t} \rangle} + p_- (e^{-\lambda} - 1) \frac{\langle \eta_1 (1 - \eta_0) e^{\lambda Q_t} \rangle}{\langle e^{\lambda Q_t} \rangle}. \quad (\text{S117})$$

For $p_+ = p_-$, we recover the one for the SEP [S9]. Similarly, we can compute

$$\begin{aligned} \partial_t \langle \eta_1 e^{\lambda Q_t} \rangle &= p_- \langle \eta_2 (1 - \eta_1) e^{\lambda Q_t} \rangle - p_+ \langle \eta_1 (1 - \eta_2) e^{\lambda Q_t} \rangle \\ &\quad + p_+ e^\lambda \langle \eta_0 (1 - \eta_1) e^{\lambda Q_t} \rangle - p_- \langle \eta_1 (1 - \eta_0) e^{\lambda Q_t} \rangle. \end{aligned} \quad (\text{S118})$$

We can rewrite the second line of the above equation using (S117), since

$$\begin{aligned} p_+ e^\lambda \langle \eta_0 (1 - \eta_1) e^{\lambda Q_t} \rangle - p_- \langle \eta_1 (1 - \eta_0) e^{\lambda Q_t} \rangle \\ = -\frac{1}{e^{-\lambda} - 1} \left(p_+ (e^\lambda - 1) \frac{\langle \eta_0 (1 - \eta_1) e^{\lambda Q_t} \rangle}{\langle e^{\lambda Q_t} \rangle} + p_- (e^{-\lambda} - 1) \frac{\langle \eta_1 (1 - \eta_0) e^{\lambda Q_t} \rangle}{\langle e^{\lambda Q_t} \rangle} \right). \end{aligned} \quad (\text{S119})$$

We thus obtain

$$\partial_t \frac{\langle \eta_1 e^{\lambda Q_t} \rangle}{\langle e^{\lambda Q_t} \rangle} = p_- \frac{\langle \eta_2 (1 - \eta_1) e^{\lambda Q_t} \rangle}{\langle e^{\lambda Q_t} \rangle} - p_+ \frac{\langle \eta_1 (1 - \eta_2) e^{\lambda Q_t} \rangle}{\langle e^{\lambda Q_t} \rangle} - \partial_t \ln \langle e^{\lambda Q_t} \rangle \left(\frac{\langle \eta_1 e^{\lambda Q_t} \rangle}{\langle e^{\lambda Q_t} \rangle} + \frac{1}{e^{-\lambda} - 1} \right). \quad (\text{S120})$$

Using the same procedure, we get

$$\partial_t \frac{\langle \eta_0 e^{\lambda Q_t} \rangle}{\langle e^{\lambda Q_t} \rangle} = p_+ \frac{\langle \eta_{-1} (1 - \eta_0) e^{\lambda Q_t} \rangle}{\langle e^{\lambda Q_t} \rangle} - p_- \frac{\langle \eta_0 (1 - \eta_{-1}) e^{\lambda Q_t} \rangle}{\langle e^{\lambda Q_t} \rangle} - \partial_t \ln \langle e^{\lambda Q_t} \rangle \left(\frac{\langle \eta_0 e^{\lambda Q_t} \rangle}{\langle e^{\lambda Q_t} \rangle} + \frac{1}{e^\lambda - 1} \right). \quad (\text{S121})$$

We now use these equations to derive equations for the WASEP at the macroscopic scale.

C. Microscopic equations for the WASEP

We define the WASEP as follows. We choose an observation time $T > 0$, and set the hopping rates of the ASEP as

$$p_+ = 1 + \frac{\nu}{\sqrt{T}}, \quad p_- = 1 - \frac{\nu}{\sqrt{T}}, \quad (\text{S122})$$

with $\nu \in [-\sqrt{T}, \sqrt{T}]$. We then consider the integrated current Q_T up to time T or the displacement X_T of a tracer at that same time in the ASEP with (S122). Obtaining an equation for the time evolution of $\langle e^{\lambda Q_T} \rangle$ in the WASEP is therefore more tricky than in the ASEP, because the parameters of the model (S122) change with the observation time T . To handle this, we introduce the different notations for the cumulant generating functions of the ASEP and the WASEP

$$\psi_{\text{ASEP}}(\lambda, t, p_+) \equiv \ln \langle e^{\lambda Q_t} \rangle, \quad \psi_{\text{WASEP}}(\lambda, T, \nu) = \psi_{\text{ASEP}}\left(\lambda, T, 1 + \frac{\nu}{\sqrt{T}}\right), \quad (\text{S123})$$

where the averaging in the first equation is performed over the solution of the master equation (S113) for the ASEP with p_+ fixed. Note that we do not write the dependence on p_- since we use the convention $p_+ + p_- = 2$. Similarly, we define the correlation profiles in both cases as

$$w_{\text{ASEP}}(\lambda, t, r, p_+) \equiv \frac{\langle \eta_r(t) e^{\lambda Q_t} \rangle}{\langle e^{\lambda Q_t} \rangle}, \quad w_{\text{WASEP}}(\lambda, T, r, \nu) \equiv w_{\text{ASEP}}\left(\lambda, T, r, 1 + \frac{\nu}{\sqrt{T}}\right). \quad (\text{S124})$$

We can therefore deduce the equations satisfied in the WASEP from those obtained for the ASEP in Section VIII B. For instance, taking the time derivative of (S123), we get

$$\partial_T \psi_{\text{WASEP}} = \left(\partial_t \psi_{\text{ASEP}} - \frac{\nu}{2T^{3/2}} \partial_{p_+} \psi_{\text{ASEP}} \right) \Big|_{t=T, p_+=1+\nu/\sqrt{T}} = -\frac{\nu}{2T} \partial_\nu \psi_{\text{WASEP}} + \partial_t \psi_{\text{ASEP}} \Big|_{t=T, p_+=1+\nu/\sqrt{T}}, \quad (\text{S125})$$

where we have used that $\partial_\nu \psi_{\text{WASEP}} = \frac{1}{\sqrt{T}} \partial_{p_+} \psi_{\text{ASEP}}$. Similarly, we obtain that

$$\partial_r w_{\text{WASEP}} = -\frac{\nu}{2T} \partial_\nu w_{\text{WASEP}} + \partial_t w_{\text{ASEP}} \Big|_{t=T, p_+=1+\nu/\sqrt{T}}. \quad (\text{S126})$$

Combined with (S117) and (S120, S121), these equations describe the time evolution of the cumulants and the correlation profiles in the WASEP. We now use these results to derive equations describing the WASEP at the macroscopic scale.

D. Macroscopic equations for the WASEP

At the macroscopic scale, one can define the density and current fields for the WASEP by (S2) (with n_i replaced by η_i). Since this model can be described within the MFT formalism, we have that the cumulant generating function is given by (S34), i.e.,

$$\psi_{\text{WASEP}}(\lambda, T, \nu) \underset{T \rightarrow \infty}{\simeq} \sqrt{T} \hat{\psi}_{\text{WASEP}}(\lambda, \nu), \quad \text{with} \quad \frac{d}{d\lambda} \hat{\psi}_{\text{WASEP}} = \int_0^\infty [q(x, 1) - q(x, 0)] dx. \quad (\text{S127})$$

Importantly, the solution q of the MFT equation also describes the correlation profiles (S7), since from (S87),

$$w_{\text{WASEP}}(\lambda, T, r, \nu) \underset{T \rightarrow \infty}{\simeq} \frac{\langle \rho(x, 1) e^{\lambda Q_T} \rangle}{\langle e^{\lambda Q_T} \rangle} = q(x, 1) \equiv \Phi(x), \quad \text{with} \quad x = \frac{r - \frac{1}{2}}{\sqrt{T}}, \quad (\text{S128})$$

where we have introduced a shift $-\frac{1}{2}$ so that the site 0 corresponds to 0^- and the site 1 to 0^+ . Inserting the scaling forms (S127, S128) into the evolution equations (S125, S126) combined with (S117) and (S120, S121), we obtain, at leading order in T , boundary conditions satisfied by Φ ,

$$(e^\lambda - 1)\Phi(0^-)(1 - \Phi(0^+)) + (e^{-\lambda} - 1)\Phi(0^+)(1 - \Phi(0^-)) = 0, \quad (\text{S129})$$

$$\partial_x \Phi(0^\pm) - (\psi_{\text{WASEP}} + \nu \partial_\nu \psi_{\text{WASEP}}) \frac{1}{2} \left(\Phi(0^\pm) + \frac{1}{e^{\mp\lambda} - 1} \right) - 2\nu \Phi(0^\pm)(1 - \Phi(0^\pm)) = 0. \quad (\text{S130})$$

Equivalently, these equations can be rewritten as

$$e^\lambda = \frac{(1 - \Phi(0^-))\Phi(0^+)}{(1 - \Phi(0^+))\Phi(0^-)}, \quad \frac{\partial_x \Phi(0^+)}{\Phi(0^+)(1 - \Phi(0^+))} = \frac{\partial_x \Phi(0^-)}{\Phi(0^-)(1 - \Phi(0^-))}, \quad (\text{S131})$$

$$\psi_{\text{WASEP}} + \nu \partial_\nu \psi_{\text{WASEP}} = -2 \left(\frac{\partial_x \Phi(x)|_{0^+}}{2\Phi(0^+)(1 - \Phi(0^+))} - 2\nu \right) (\Phi(0^+) - \Phi(0^-)), \quad (\text{S132})$$

In this form, these boundary conditions will be a basis to obtain the boundary conditions for any driven diffusive system in Section IX.

E. Cumulants and correlations profiles

Inserting the transport coefficients of the WASEP $D(\rho) = 1$ and $\sigma(\rho) = 2\rho(1 - \rho)$ in the expressions obtained for the cumulants of Q_T for general driven diffusive systems (S84,S85,S86), we obtain for the WASEP,

$$\frac{\langle Q_T \rangle_c}{\sqrt{T}} \underset{T \rightarrow \infty}{\simeq} 2\bar{\rho}(1 - \bar{\rho}), \quad \frac{\langle Q_T^2 \rangle_c}{\sqrt{T}} \underset{T \rightarrow \infty}{\simeq} \bar{\rho}(1 - \bar{\rho}) \left(\frac{2}{\sqrt{\pi}} e^{-(\nu\sigma'(\bar{\rho}))^2/4} + \nu\sigma'(\bar{\rho}) \operatorname{erf} \left(\frac{\nu\sigma'(\bar{\rho})}{2} \right) \right), \quad (\text{S133})$$

$$\begin{aligned} \frac{\langle Q_T^3 \rangle_c}{\sqrt{T}} \underset{T \rightarrow \infty}{\simeq} &= 2\nu\bar{\rho}(1 - \bar{\rho}) - 12\nu\bar{\rho}^2(1 - \bar{\rho})^2 + 12\nu\bar{\rho}^2(1 - \bar{\rho})^2 \frac{e^{-\frac{(\nu\sigma'(\bar{\rho}))^2}{2}}}{\pi} - 12\nu^3\bar{\rho}^2(1 - \bar{\rho})^2(1 - 2\bar{\rho})^2 \\ &+ 24\nu^2\bar{\rho}^2(1 - \bar{\rho})^2(1 - 2\bar{\rho}) \frac{e^{-\frac{(\nu\sigma'(\bar{\rho}))^2}{4}}}{\sqrt{\pi}} \operatorname{erf} \left(\frac{\nu\sigma'(\bar{\rho})}{2} \right) + 12\nu^3\bar{\rho}^2(1 - \bar{\rho})^2(1 - 2\bar{\rho})^2 \operatorname{erf} \left(\frac{\nu\sigma'(\bar{\rho})}{2} \right)^2. \end{aligned} \quad (\text{S134})$$

Performing the same substitution into (S90,S91,S92), we obtain the cumulants of X_T for the WASEP,

$$\frac{\langle X_T \rangle_c}{\sqrt{T}} \underset{T \rightarrow \infty}{\simeq} 2\nu(1 - \bar{\rho}), \quad \frac{\langle X_T^2 \rangle_c}{\sqrt{T}} \underset{T \rightarrow \infty}{\simeq} \frac{2(1 - \bar{\rho})}{\bar{\rho}\sqrt{\pi}} \left(e^{-\nu^2\bar{\rho}^2} + \nu\bar{\rho}\sqrt{\pi} \operatorname{erf}(\nu\bar{\rho}) \right), \quad (\text{S135})$$

$$\begin{aligned} \frac{\langle X_T^3 \rangle_c}{\sqrt{T}} \underset{T \rightarrow \infty}{\simeq} &= \frac{2(1 - \bar{\rho})}{\bar{\rho}^2} \left(-\frac{3(1 - \bar{\rho})e^{-\nu^2\bar{\rho}^2} (4\nu^2\bar{\rho}^2 + 1) \operatorname{erf}(\nu\bar{\rho})}{\sqrt{\pi}} \right. \\ &\left. + \nu\bar{\rho}(1 - \bar{\rho}) \left((6\nu^2\bar{\rho}^2 + 3) \operatorname{erf}(\nu\bar{\rho})^2 - 6\nu^2\bar{\rho}^2 - 1 \right) + \frac{\nu\bar{\rho}e^{-2\nu^2\bar{\rho}^2} (\pi(3\bar{\rho} - 2)e^{2\nu^2\bar{\rho}^2} - 6\bar{\rho} + 6)}{\pi} \right). \end{aligned} \quad (\text{S136})$$

In particular, inserting that $\nu = \frac{p_+ - p_-}{2} \sqrt{T}$ from the definition of the WASEP, we recover the cumulants of the ASEP [S12],

$$\langle X_T \rangle \simeq \langle X_T^2 \rangle_c \simeq \langle X_T^3 \rangle_c \simeq (p_+ - p_-)(1 - \bar{\rho})T. \quad (\text{S137})$$

IX. BOUNDARY CONDITIONS FOR GENERAL DRIVEN DIFFUSIVE SYSTEMS

A. Derivation from MFT

In this section, we show that, in the case of Q_T , the correlation profile Φ satisfies the following boundary conditions,

$$\mu(\Phi)|_{0^+} - \mu(\Phi)|_{0^-} = \lambda, \quad \mu(\rho) = \int^\rho \frac{2D(r)}{\sigma(r)} dr, \quad (\text{S138})$$

$$\partial_x \mu(\Phi)|_{0^+} = \partial_x \mu(\Phi)|_{0^-}, \quad (\text{S139})$$

for any driven diffusive system. We adapt the procedure used in Ref. [S13] for non-driven systems, and introduce

$$q(x, t) = \hat{q}(x, 1 - t), \quad p(x, t) = -\hat{p}(x, 1 - t) + \mu(\hat{q}(x, 1 - t)) - 2\nu x. \quad (\text{S140})$$

Inserting these relations into the MFT equations (S31,S32), we find that \hat{q} and \hat{p} obey the exact same equations,

$$\partial_t \hat{q} = \partial_x [D(\hat{q})\partial_x \hat{q} - \sigma(\hat{q})\partial_x \hat{p} - \nu\sigma(\hat{q})], \quad (\text{S141})$$

$$\partial_t \hat{p} = -D(\hat{q})\partial_x^2 \hat{p} - \frac{1}{2}\sigma'(\hat{q})(\partial_x \hat{p})^2 - \nu\sigma'(\hat{q})\partial_x \hat{p}, \quad (\text{S142})$$

but with the new initial and final condition deduced from (S33),

$$\hat{p}(x, 0) = -\lambda\Theta(x) + \mu(\hat{q}(x, 0)) - 2\nu x, \quad \hat{p}(x, 1) = -\lambda\Theta(x) + \mu(\bar{\rho}) - 2\nu x. \quad (\text{S143})$$

The key point is that \hat{p} obeys an anti-diffusion equation, so that it is smooth at $t = 0$, hence,

$$\hat{p}(0^-, 0) = \hat{p}(0^+, 0), \quad \partial_x \hat{p}(0^-, 0) = \partial_x \hat{p}(0^+, 0). \quad (\text{S144})$$

Using the expression of $\hat{p}(x, 0)$ (S143) and using that $\hat{q}(x, 0) = q(x, 1) = \Phi(x)$, we obtain the boundary conditions (S138,S139) announced above. Note that in the case of the WASEP, $D(\rho) = 1$ and $\sigma(\rho) = 2\rho(1 - \rho)$, these equations reduce to (S131) obtained from microscopic considerations.

B. A conjecture for the cumulant generating function

In the case of the WASEP discussed in Section VIII D, the two boundary conditions (S138,S139) are completed by a third condition which relates $\hat{\psi}$ to $\Phi(0^\pm)$ and $\partial_x \Phi(0^\pm)$. In practice, this last relation greatly simplifies the computation of the cumulants from the correlation profile Φ since it bypasses the computation of the spatial integral in (S35). The WASEP relation (S132) can actually be written in terms of D and σ as

$$\hat{\psi} + \nu\partial_\nu \hat{\psi} = -2(\partial_x \mu(\Phi)|_{0^+} - 2\nu) \int_{\Phi(0^-)}^{\Phi(0^+)} D(r) dr. \quad (\text{S145})$$

We conjecture that this relation is valid for any driven diffusive system. This conjecture is verified for the WASEP by construction, and for any driven diffusive system up to order 3 in λ from our results on the cumulants (S84,S85,S86) and the correlation profiles (S51,S81). In the non-driven case $\nu = 0$, it reduces to the expression written in [S13].

Remarkably, the relation (S145) is invariant under the duality mapping described in Section II D, as we now proceed to show. The profile $\tilde{\Phi}$ in the dual system is related to the profile Φ in the original system by

$$\tilde{\Phi}(y(x)) = \frac{1}{\Phi(x)} \quad y(x, t) = \int_0^x \Phi(x') dx', \quad (\text{S146})$$

and the coefficients $D(\rho)$, $\sigma(\rho)$ and ν are mapped according to (S39). Applying that mapping to the term at the right hand side of the conjectured relation (S145), we obtain for the chemical potential

$$\partial_x \mu(q) = -\frac{\partial_y \tilde{q}}{\tilde{q}^3} \mu' \left(\frac{1}{\tilde{q}} \right) = \frac{\partial_y \tilde{q}}{\tilde{q}^3} \frac{2D(\frac{1}{\tilde{q}})}{\sigma(\frac{1}{\tilde{q}})} = -\partial_y \tilde{\mu}(\tilde{q}), \quad (\text{S147})$$

where $\tilde{\mu}$ is the chemical potential of the dual model. Setting the change of variable $u = \frac{1}{\tilde{r}}$, in the integral,

$$\int_{\Phi(0^-)}^{\Phi(0^+)} D(r) dr = - \int_{\frac{1}{\Phi(0^+)}}^{\frac{1}{\Phi(0^-)}} D \left(\frac{1}{u} \right) \frac{du}{u^2} = - \int_{\tilde{\Phi}(0^-)}^{\tilde{\Phi}(0^+)} \tilde{D}(u) du. \quad (\text{S148})$$

All together, we deduce that the term on the r.h.s. of (S145) is invariant under the duality mapping, which is consistent with the fact that $\hat{\psi}$ is also invariant. This further supports the validity of the relation (S145).

[S1] M. R. Evans and T. Hanney, *J. Phys. A* **38**, R195 (2005).

- [S2] B. Derrida, *J. Stat. Mech.* **2007**, P07023 (2007).
- [S3] L. Bertini, A. De Sole, D. Gabrielli, G. Jona-Lasinio, and C. Landim, *Rev. Mod. Phys.* **87**, 593 (2015).
- [S4] H. Spohn, *J. Phys. A* **16**, 4275 (1983).
- [S5] K. Mallick, *Physica A* **418**, 17 (2015), proceedings of the 13th International Summer School on Fundamental Problems in Statistical Physics.
- [S6] B. Derrida and A. Gerschenfeld, *J. Stat. Phys.* **137**, 978 (2009).
- [S7] P. Rizkallah, A. Grabsch, P. Illien, and O. Bénichou, *J. Stat. Mech: Theory Exp.* **2023**, 013202 (2023).
- [S8] D. B. Owen, *Commun. Stat. Simul. Comput.* **9**, 389 (1980).
- [S9] A. Grabsch, A. Poncet, P. Rizkallah, P. Illien, and O. Bénichou, *Sci. Adv.* **8**, eabm5043 (2022).
- [S10] A. Grabsch, P. Rizkallah, A. Poncet, P. Illien, and O. Bénichou, *Phys. Rev. E* **107**, 044131 (2023).
- [S11] A. Poncet, A. Grabsch, P. Illien, and O. Bénichou, *Phys. Rev. Lett.* **127**, 220601 (2021).
- [S12] A. De Masi and P. A. Ferrari, *J. Stat. Phys.* **38**, 603 (1985).
- [S13] A. Grabsch, P. Rizkallah, and O. Bénichou, *SciPost Phys.* **16**, 016 (2024).