THE CENTER OF THE WALLED BRAUER ALGEBRA $B_{r,1}(\delta)$

EIRINI CHAVLI, MAUD DE VISSCHER, ALISON PARKER, SARAH SALMON, AND ULRICA WILSON

To the memory of Emmelia Kokota

ABSTRACT. We show that the centre of the walled Brauer algebra $B_{r,1}(\delta)$ over the complex field \mathbb{C} , for any parameter $\delta \in \mathbb{C}$, is generated by the supersymmetric polynomials evaluated at the Jucys-Murphy elements. Moreover, we prove that its dimension is independent of the parameter δ .

1. Introduction

Let $V = \mathbb{C}^n$ be the natural representation of $\mathrm{GL}_n(\mathbb{C})$ and let r be a positive integer. There are two actions on the rth tensor product $V^{\otimes r}$: the first action is by $\mathrm{GL}_n(\mathbb{C})$, which is the (left) diagonal action. The second one is the (right) action of the symmetric group S_r which permutes the tensor factors. One can easily see that these two actions compute. However, there is a stronger relation between them, as asserted by Schur-Weyl duality [19], namely that the span of the image of S_r and $\mathrm{GL}_n(\mathbb{C})$ in $\mathrm{End}(V^{\otimes r})$ are centralizers of each other.

The Brauer algebra $B_r(\delta)$ was introduced by Brauer [1], in the context of classical invariant theory, to play the role of the symmetric group in a corresponding Schur-Weyl duality for the orthogonal groups (for δ positive integer) and for the symplectic groups (for δ negative integer).

The walled Brauer algebra $B_{r,s}(\delta)$ is a subalgebra of $B_{r+s}(\delta)$ and it was introduced independently by Turaev and by Koike [8, 17]. It appears in another generalisation of Schur-Weyl duality, when considering the action of $GL_n(\mathbb{C})$ on mixed tensor space $V^{\otimes r} \otimes (V^*)^{\otimes s}$.

Brauer algebras and walled Brauer algebras are particular examples of diagram algebras. Their representation theory has been studied extensively, see for example [5, 4, 3, 9, 13] and references therein. In particular, they are semisimple unless the parameter δ is a small integer compared to r (and s). In the semisimple case, there is an explicit construction of the simple modules. Even in the non-semisimple cases, their representation theory is well-understood: they are cellular algebras and we have an explicit combinatorial formula for their decomposition numbers. One question however remains open in the non-semisimple case, namely the description of the centre of these algebras. The aim of this paper is to make some progress in the case of the walled Brauer algebra.

Much of the work to study the structure of these diagram algebras has been done by analogy with the group algebra of the symmetric group. The study of their centres is no exception. So we start by recalling the classical results on the centre of $\mathbb{C}S_r$.

As for the centre of any group algebras of a finite group, the centre of $\mathbb{C}S_r$ has a basis given by the conjugacy class sums, indexed in this case by partitions of r. But there is another description, in terms of the so-called Jucys-Murphy elements, first introduced in [6, 11], defined by

$$L_k := \sum_{j=1}^{k-1} (j, k), \ 1 \le k \le r.$$

These elements play an essential role in the Okounkov-Vershik approach to the representation theory of the symmetric groups [12]. The center of $\mathbb{C}S_r$ can be described as the algebra of all the symmetric polynomials evaluated in the Jucys-Murphy elements (for more details, see [6, 10]).

We would like to thank the Women in Noncommutative Algebra and Representation Theory (WINART2) workshop, held at the University of Leeds in May 2019, where this project started.


Analogues of these Jucys-Murphy elements for the algebra $B_{r,s}(\delta)$ were introduced in [2, 14, 15] and conjectures were made about using some versions of symmetric polynomials in these to describe the centre of the walled Brauer algebra (see [2, Remark 2.6] and [15, Conjecture 7.3]). In [7] Jung and Kim, inspired by [15, Conjecture 7.3], introduced a renormalisation of these Jucys-Murphy elements, which we denote again as L_1, \ldots, L_{r+s} . They proved that supersymmetric polynomials in these elements are central in $B_{r,s}(\delta)$ for all values of the parameter δ and that moreover, when $B_{r,s}(\delta)$ is semisimple, this is in fact the whole centre. The key fact here is that when an algebra is semisimple, the dimension of its centre is given by the number of isomorphism classes of simple modules. Jung and Kim conjectured in [7, Conjecture 5.4] that the result still hold in all cases. The aim of this paper is to investigate this question. We give a strategy for the general case, following the work of Shalile [16] and proves this conjecture in the case when r is arbitrary and s = 1.

Structure of the paper. Section 2 contains the definition of the walled Brauer algebra in terms of its diagram basis and multiplication. One should note that $B_{r,s}(\delta)$ contains $\mathbb{C}(S_r \times S_s)$ as a subalgebra and that in fact, it is generated by this subalgebra together with one more element, which we denote by e. We also recall the definition of the Jucys-Murphy elements, supersymmetric polynomials and state Jung-Kim's result for the centre in the semisimple case. In Section 3, we obtain a lower bound for the dimension of the centre in the general case, using Jung-Kim's result and a base-change argument. In Section 4 we recall the construction of the generalised walled cycle types due to Shalile [16]. The sums of all diagrams with the same cycle type give a basis for the centraliser algebra of the product of symmetric group $S_r \times S_s$ in the walled Brauer algebra $B_{r,s}(\delta)$. Thus, in order to find the centre of the walled Brauer algebra, we only need to impose one additional condition, namely commutation with the extra generator e. This, together with the lower bound obtained in Section 4 gives a general strategy to find the centre. We then apply this strategy explicitly in Section 5 for the special case $B_{r,1}(\delta)$.

2. Preliminaries


- 2.1. Walled Brauer algebras. Let r and s be nonnegative integers. An (r,s)-walled Brauer diagram is a graph drawn in a rectangle with (r+s) vertices on its top and bottom edges, numbered $1, \ldots, r+s$ in order from left to right. Each vertex is connected by a strand to exactly one other vertex. In addition, there is a vertical wall separating the left r vertices from the right s vertices, such that the following conditions hold:
 - (1) A propagating line connects a vertex on the top row with one on the bottom row, and it cannot cross the wall.
 - (2) A northern arc (respectively, southern arc) connects vertices on the top row (respectively, on the bottom row), and it must cross the wall.

For example, the following graphs are (4, 2)-Brauer diagrams:




Let δ be a complex number. The walled Brauer algebra $B_{r,s}(\delta)$ is the \mathbb{C} -linear span of the (r,s)-walled Brauer diagrams with the multiplication defined as follows: The product of two (r,s)-walled Brauer diagrams d_1 and d_2 is determined by putting d_1 above d_2 and identifying the bottom vertices of d_1 with the top vertices of d_2 . Let n be the number of loops in the middle row so obtained. The product d_1d_2 is given by δ^n times the resulting diagram with loops omitted.

For example, for the diagrams x and y above we have:



The dimension of $B_{r,s}(\delta)$ equals (r+s)! (see, for example, [2, 2.2]).

We denote by s_i $(1 \le i \le r-1 \text{ or } r+1 \le i \le r+s-1)$ and e the following (r,s)-walled Brauer diagrams:



Note that $B_{0,n}(\delta) \simeq B_{n,0}(\delta) \simeq \mathbb{C}S_n$, the group algebra of the symmetric group S_n on n letters. It's easy to check that the algebra $B_{r,s}(\delta)$ is generated by the elements s_i $(1 \le i \le r-1)$ or $r+1 \le j \le r+s-1$) and e.

For any (r, s)-walled Brauer diagram d, we denote by d^* the flip diagram of d, obtained by horizontally flipping d. This define a \mathbb{C} -linear anti-automorphism $*: B_{r,s}(\delta) \to B_{r,s}(\delta)$.

The following theorem is Theorem 6.3 in [4] and it provides a criterion of the semisimplicity of the algebra $B_{r,s}(\delta)$.


Theorem 2.1. The walled Brauer algebra $B_{r,s}(\delta)$ is semisimple if and only if one of the following holds:

- (1) r = 0 or s = 0,
- (2) $\delta \notin \mathbb{Z}$,
- (3) $|\delta| > r + s 2$,
- (4) $\delta = 0$, and $(r, s) \in \{(1, 2), (1, 3), (2, 1), (3, 1)\}.$


Therefore, for a fixed pair (r, s) the algebra $B_{r,s}(\delta)$ is semisimple except for finitely many values $\delta \in \mathbb{C}$.

2.2. **Jucys-Murphy elements.** There are different definitions for Jucys-Murphy elements for the walled Brauer algebra $B_{r,s}(\delta)$. In this paper, we use Definition 2.1 of [7].

Consider the transposition (a, b) given by the diagram



Define also $e_{j,k}$, $(1 \le j \le r, r+1 \le k \le r+s)$ to be the diagram



For each $1 \le k \le r + s$ we define the Jucys-Murphy elements L_k of $B_{r,s}(\delta)$ as follows:

$$L_k := \begin{cases} 0 & \text{if } k = 1, \\ \sum\limits_{j=1}^{k-1} (j,k) & \text{if } 1 < k \le r, \\ -\sum\limits_{j=1}^{r} e_{j,k} + \sum\limits_{j=r+1}^{k-1} (j,k) + \delta & \text{if } r+1 \le k \le r+s. \end{cases}$$

- 2.3. Supersymmetric polynomials and central elements. Let m, n be nonnegative integers. An element p in the polynomial ring $\mathbb{C}[x_1, \ldots, x_m, y_1, \ldots, y_n]$ is supersymmetric if
 - (1) p is symmetric in x_1, \ldots, x_m and in y_1, \ldots, y_n separately.
 - (2) The substitution $x_m = t$, $y_1 = -t$ yields a polynomial in $x_1, \ldots, x_{m-1}, y_2, \ldots, y_m$ which is independent of t.

We denote by $S_{m,n}[x;y]$ the set of supersymmetric polynomials in $x_1,\ldots,x_m,y_1,\ldots,y_n$.

The following result is Corollary 2.9 in [7] and it provides us with some central elements of $B_{r,s}(\delta)$.

Proposition 2.2. For every supersymmetric polynomial p in $S_{r,s}[x;y]$, the element

$$p(L_1,\ldots,L_r,L_{r+1},\ldots,L_{r+s})$$

belongs to the centre of $B_{r,s}(\delta)$.

2.4. The semisimple case. We know that when an algebra is semisimple, the dimension of its centre is equal to the number of isomorphic classes of simple modules. For the walled Brauer algebra $B_{r,s}(\delta)$, these are indexed by the set $\Lambda_{r,s}$ of bipartitions defined as follows (see for example [4, Theorem 2.7]).

A partition is a decreasing sequence of non-negative integers $\lambda = (\lambda_1, \lambda_2, \dots)$. We write Λ for the set of all partitions. For $\lambda \in \Lambda$ we set $|\lambda| := \sum_{i>1} \lambda_i$. Now define

$$\Lambda_{r,s} := \coprod_{k=0}^{\min(r,s)} \{ (\lambda, \mu) \in \Lambda \times \Lambda \mid |\lambda| = r - k, \ |\mu| = s - k \}$$

Therefore, when $B_{r,s}(\delta)$ is semisimple, the dimension of its centre is the cardinality of $\Lambda_{r,s}$. Jung and Kim [7, Theorem 3.5] gave a basis of the centre of $B_{r,s}(\delta)$ in this case. More precisely, they proved the following theorem.

Theorem 2.3. If the walled Brauer algebra $B_{r,s}(\delta)$ is semisimple, the supersymmetric polynomials in L_1, \ldots, L_{r+s} generate the centre of $B_{r,s}(\delta)$. Moreover, there is a set of supersymmetric polynomials $\{p_{(\lambda,\mu)} \in S_{r,s}[x;y] \mid (\lambda,\mu) \in \Lambda_{r,s}\}$, such that the set $\{p_{(\lambda,\mu)}(L_1,\ldots,L_{r+s}) \mid (\lambda,\mu) \in \Lambda_{r,s}\}$ is a basis of the centre of $B_{r,s}(\delta)$.

One can find a precise description of these supersymmetric polynomials p_{λ} in [7, Section 3]. The goal of this paper to extend this result to the non-semisimple case. More precisely, we want to prove the following conjecture [7, Conjecture 5.4]:

Conjecture 2.4. For every $\delta \in \mathbb{C}$, the center of the walled Brauer algebra $B_{r,s}(\delta)$ is generated by the supersymmetric polynomials in the Jucys-Murphy elements L_1, \ldots, L_{r+s} .

3. A LOWER BOUND ON THE DIMENSION OF THE CENTRE OF $B_{r,s}(\delta)$

Let z be a formal parameter. We consider the walled Brauer algebra $B_{r,s}^{\mathbb{C}[z]}(z)$ over the polynomial ring $\mathbb{C}[z]$. We define the Jucys-Murphy elements \mathcal{L}_i , $1 \leq i \leq r+s$ as in Section 2.2, replacing δ by z. The proof of Proposition 2.2 goes through unchanged when working in this setting and we get the following proposition.

Proposition 3.1. All supersymmetric polynomials (with coefficients in $\mathbb{C}[z]$) in $\mathcal{L}_1, \ldots, \mathcal{L}_{r+s}$ are central in $B_{r,s}^{\mathbb{C}[z]}(z)$.

We now take a $\delta_0 \in \mathbb{C}$ ($\delta_0 \neq 0$), such that $B_{r,s}(\delta_0)$ is semisimple (see Theorem 2.1). Set $m = |\Lambda_{r,s}|$ and choose $p_i \in S_{r,s}[x;y]$ for $1 \leq i \leq m$ to be supersymmetric polynomials such that

$$\{p_i(L_1,\ldots,L_{r+s}) \mid 1 \le i \le m\}$$

form a \mathbb{C} -basis for the centre $Z(B_{r,s}(\delta_0))$ of $B_{r,s}(\delta_0)$.

Let $\mathbb{C}(z)$ be the field of fractions of $\mathbb{C}[z]$ and we consider the algebra

$$B_{r,s}^{\mathbb{C}(z)}(z) := B_{r,s}^{\mathbb{C}[z]}(z) \otimes_{\mathbb{C}[z]} \mathbb{C}(z).$$

Using the same arguments of Theorem 6.3 in [4], we have that $B_{r,s}^{\mathbb{C}(z)}(z)$ is a semisimple $\mathbb{C}(z)$ -algebra with m simple modules, hence

$$\dim_{\mathbb{C}(z)} Z(B_{r,s}^{\mathbb{C}(z)}(z)) = m.$$

Moreover,

$$Z(B_{r,s}^{\mathbb{C}(z)}(z)) = Z(B_{r,s}^{\mathbb{C}[z]}(z)) \otimes_{\mathbb{C}[z]} \mathbb{C}(z).$$

Therefore,

$$\operatorname{rank}_{\mathbb{C}[z]} Z(B^{\mathbb{C}[z]}_{r,s}(z)) = \dim_{\mathbb{C}(z)} Z(B^{\mathbb{C}(z)}_{r,s}(z)) = m.$$

Proposition 3.2. The set $\{p_i(\mathcal{L}_1,\ldots,\mathcal{L}_{r+s}) \mid 1 \leq i \leq m\}$ forms a $\mathbb{C}[z]$ -basis for $Z(B_{r,s}^{\mathbb{C}[z]}(z))$.

Proof. From Proposition 3.1 we have that $p_i(\mathcal{L}_1, \dots, \mathcal{L}_{r+s}) \in Z(B_{r,s}^{\mathbb{C}[z]}(z))$, for every $i = 1, \dots, m$. Moreover, $\operatorname{rank}_{\mathbb{C}[z]} Z(B_{r,s}^{\mathbb{C}[z]}(z)) = m$. Therefore, it is enough to show that this set is linearly independent.

Assume $\sum_{i=1}^{m} a_i(z)p_i(\mathcal{L}_1,\ldots,\mathcal{L}_{r+s}) = 0$, for some $a_i(z) \in \mathbb{C}[z]$, not all 0. Dividing by the highest power of $(z - \delta_0)$ which divides all the $a_i(z)$'s, we can assume that there exists some i_0 with $a_{i_0}(\delta_0) \neq 0$.

Now, specializing to $z = \delta_0$, and noting that under this specialization the elements \mathcal{L}_i 's become the L_i 's, we get

$$\sum_{i=1}^{m} a_i(\delta_0) p_i(L_1, \dots, L_{r+s}) = 0.$$

Now, as $\{p_i(L_1,\ldots,L_{r+s}) \mid 1 \leq i \leq m\}$ is a linearly independent set, we must have $a_i(\delta_0) = 0, \forall i = 1,\ldots,m$, which contradicts the fact that $a_{i_0}(\delta_0) \neq 0$.

We now take an arbitrary $\delta \in \mathbb{C}$ and we consider

$$B_{r,s}^{\mathbb{C}}(\delta) = B_{r,s}^{\mathbb{C}[z]}(z) \otimes_{\mathbb{C}[z]} \mathbb{C}[z]/\langle z - \delta \rangle.$$

For any $Q \in B_{r,s}^{\mathbb{C}[z]}(z)$, we write

$$\bar{Q} := Q \otimes_{\mathbb{C}[z]} \mathbb{C}[z]/\langle z - \delta \rangle \in B_{r,s}^{\mathbb{C}}(\delta).$$

To simplify notation we write

$$\mathcal{P}_i := p_i(\mathcal{L}_1, \dots, \mathcal{L}_{r+s}) \in B_{r,s}^{\mathbb{C}[z]}(z), \ i = 1, \dots, m.$$

Proposition 3.3. For any choice of $\delta \in \mathbb{C}$, the set $\{\bar{\mathcal{P}}_i \mid 1 \leq i \leq m\}$ is a linearly independent set in $Z(B_{r,s}^{\mathbb{C}}(z))$. In particular, $\dim_{\mathbb{C}} Z(B_{r,s}^{\mathbb{C}}(z)) \geq |\Lambda_{r,s}|$.

Proof. First note that as \mathcal{P}_i are central elements in $B_{r,s}^{\mathbb{C}[z]}(z)$ we have that $\bar{\mathcal{P}}_i$ are central in $B_{r,s}^{\mathbb{C}}(z)$. Now let

$$\sum_{i=1}^{m} a_i \bar{\mathcal{P}}_i = 0$$

for some $a_i \in \mathbb{C}$. Then, we have

$$\sum_{i=1}^{m} a_i \bar{\mathcal{P}}_i = \sum_{i=1}^{m} \overline{a_i \mathcal{P}_i} = 0$$

and so

(3.1)
$$\sum_{i=1}^{m} a_i \mathcal{P}_i = (z - \delta)R$$

for some $R \in Z(B_{r,s}^{\mathbb{C}[z]}(z))$. Thus, R can be written as $R = \sum_{i=1}^m b_i(z)\mathcal{P}_i$, for some $b_i(z) \in \mathbb{C}[z]$. Using now Equation (3.1) we get $\sum_{i=1}^m a_i\mathcal{P}_i = \sum_{i=1}^m (z-\delta)b_i(z)\mathcal{P}_i$. As $a_i \in \mathbb{C}$, $b_i(z) \in \mathbb{C}[z]$ and $\{\mathcal{P}_i \mid 1 \leq i \leq m\}$ are linearly independent, we must have $a_i = (z-\delta)b_i(z) = 0$, $\forall i = 1, \ldots, m$.

4. Shalile's cycle type and general strategy

- 4.1. Walled generalized cycle types. The following definition is Definition 7.3 in [16]. For a diagram $d \in B_{r,s}(\delta)$ we define the walled generalized cycle type c(d) of d to be a set of words (called parts) in the alphabet L, R, N and S, obtained as follows: We first connect each vertex in the top row of d with the vertex in the bottom row below it. The parts of c(d) correspond to the connected components of this new graph, as follows: We take a connected component of the new graph, we pick a vertex of it and we follow the path, until all the edges of the connected component have been read off once. Following the path, we record in order with the letters L, R, N and S the types of the edges of the diagram d which are traversed. More precisely, we record:
 - \bullet N if the type of the edge is a northern arc.
 - S if the type of the edge is a southern arc.
 - L if the type of the edge is a propagating line to the left of the wall.
 - R if the type of the edge is a propagating line to the right of the wall.


For example, for the diagrams x and y we saw in Section 2.1 we have $c(x) = \{LL, NSNS\}$ and $c(y) = \{L, L, NS, NS\}$.

Two parts are equivalent if one is obtained from the other by repeated cyclic permutation and/or reverse reading. Two walled generalized cycle types are equal if their parts are equivalent. For example, we also have $c(y) = \{L, L, SN, SN\}$ and $c(x) = \{LL, SNSN\}$.


Remark 4.1. From the definition of (equal) walled generalized cycle types we notice the following:

- (1) Each part of c(d) is of one of the following forms:
 - (a) L^a , $a \ge 1$.
 - (b) $R^b, b \ge 1$.
 - (c) $NR^{b_1}SL^{a_1}NR^{b_2}SL^{a_2}...$, for some $a_i, b_i \ge 0$.
- (2) The number of N's in c(d) equals the number of S's in c(d). We denote this number by t. Then, we also have that the number of L's in c(d) equals r-t, while the number of R's in c(d) equals s-t.

(3) In general, $c(d) \neq c(d^*)$. For example, we consider the following diagram d of the walled Brauer algebra $B_{3,3}(\delta)$:



We have $c(d) = \{NSNRSL\}$. On the other hand,



We have $c(d^*) = \{NSLNRS\}$. Note that $c(d) \neq c(d^*)$. A description of how to obtain $c(d^*)$ from c(d) in general is given in [16, Lemma 2.14].

The following theorem is Theorem 7.7 in [16].

Theorem 4.2. Two diagrams of $B_{r,s}(\delta)$ are $S_r \times S_s$ -conjugate if and only if they have equal walled generalized cycle types.

4.2. A general strategy. Let \mathcal{B} be the diagram basis of $B_{r,s}(\delta)$. We denote by $C_{r,s}$ the set of all generalised walled cycle types for $B_{r,s}(\delta)$. For each walled generalized cycle type $\mu \in C_{r,s}$ we set $\mathcal{B}_{\mu} := \{d \in \mathcal{B} \mid c(d) = \mu\}$. It follows from Theorem 4.2 that the centralizer of $Z_{B_{r,s}(\delta)}(\mathbb{C}(\mathfrak{S}_r \times \mathfrak{S}_s))$ has a basis consisting of all elements of the form $\sum_{d \in \mathcal{B}_{\mu}} d$ where μ runs over all walled generalised

cycle types. Recall from Section 2.1 that the walled Brauer algebra $B_{r,s}(\delta)$ is generated by the generators of $S_r \times S_s$ and by the generator e. A central element of $B_{r,s}(\delta)$ is therefore an element of $Z_{B_{r,s}(\delta)}(\mathbb{C}(S_r \times S_s))$ which commutes with the generator e. Let us consider an arbitrary element of $Z_{B_{r,s}(\delta)}(\mathbb{C}(S_r \times S_s))$, which is of the form $\sum_{\mu} a_{\mu} \sum_{d \in \mathcal{B}_{\mu}} d$, for some $a_{\mu} \in \mathbb{C}$. In order for this element

to be in the centre of $B_{r,s}(\delta)$, it needs to satisfy

(4.1)
$$\sum_{\mu} a_{\mu} \sum_{d \in \mathcal{B}_{\mu}} (de - ed) = \sum_{x \in \mathcal{B}} b_x x = 0.$$

So we need to consider the system

$$\{b_x = 0 \mid x \in \mathcal{B}\}\$$

as a system of equations in the variables a_{μ} , $\mu \in C_{r,s}$. Showing that the rank of this system is $|C_{r,s}| - |\Lambda_{r,s}|$ would prove Conjecture 2.4. In fact, using Proposition 3.3, it is enough to show that the rank is at least $|C_{r,s}| - |\Lambda_{r,s}|$. Thus our strategy is to find $|C_{r,s}| - |\Lambda_{r,s}|$ linearly independent equations among $\{b_x = 0 \mid x \in \mathcal{B}\}$. To identify which equations we should be considering, let us start by making the following observation.

Definition 4.3. Let $\mu \in C_{r,s}$. We call a part of μ of type (a), (b), or (c) of the form NS in Remark 4.1(1) a trivial part.

Proposition 4.4. There is a bijection between the set $\Lambda_{r,s}$ and the set of diagrams in $B_{r,s}(\delta)$ having only trivial parts.

Proof. Let $k \in \mathbb{N}_0$ and let $\lambda = (\lambda_1, \lambda_2, \dots)$, $\mu = (\mu_1, \mu_2, \dots)$ partitions of r-k and s-k, respectively. The following map is a bijection:

$$(\lambda,\mu) \mapsto \left\{ d \in B_{r,s}(\delta) \mid c(d) = \{L^{\lambda_1}, L^{\lambda_2}, \dots, R^{\mu_1}, R^{\mu_2}, \dots, \underbrace{NS, NS, \dots, NS}_{k \text{ terms}} \} \right\}.$$

Thus in order to prove Conjecture 2.4, it would be enough to pick one diagram $x \in \mathcal{B}_{\mu}$ for each cycle type $\mu \in C_{r,s}$ containing at least one non-trivial part and show that the set of equations $b_x = 0$ for all such x's are linearly independent. We will do this in the case s = 1 in the next section.

5. The center of
$$B_{r,1}(\delta)$$

Let \mathcal{B} be the diagram basis of $B_{r,1}(\delta)$, consisting of (r+1)! diagrams. We recall that for each walled generalized cycle type μ we set $\mathcal{B}_{\mu} = \{d \in \mathcal{B} \mid c(d) = \mu\}$.

The following lemma is easy to see but note that it is true only for the case of $B_{r,1}(\delta)$ (see Remark 4.1 (3)).

Lemma 5.1. The flip map * gives a bijection $\mathcal{B}_{\mu} \to \mathcal{B}_{\mu}$, $d \mapsto d^*$ for each $\mu \in C_{r,s}$.

We now consider equation (4.1) and system (4.2). Our first attempt to simplify this system is to find diagrams $x \in B_{r,1}(\delta)$ for which we have $b_x = 0$. The first result in this direction is the following proposition:

Proposition 5.2. For any $x \in \mathcal{B}$ we have $b_{x^*} = -b_x$. In particular, if $x = x^*$ then $b_x = 0$.

Proof. We apply the anti-automorphism * to both sides of (4.1) and we have:

$$\sum_{\mu} a_{\mu} \sum_{d \in C_{\mu}} (de - ed)^{*} = \sum_{x \in \mathcal{B}} b_{x} x^{*} \Longrightarrow$$

$$\sum_{\mu} a_{\mu} \sum_{d \in C_{\mu}} (e^{*}d^{*} - d^{*}e^{*}) = \sum_{x \in \mathcal{B}} b_{x} x^{*} \stackrel{e^{*} = e}{\Longrightarrow}$$

$$\sum_{\mu} a_{\mu} \sum_{d \in C_{\mu}} (ed^{*} - d^{*}e) = \sum_{x \in \mathcal{B}} b_{x} x^{*} \stackrel{5.1}{\Longrightarrow}$$

$$\sum_{\mu} a_{\mu} \sum_{d \in C_{\mu}} (ed - de) = \sum_{x \in \mathcal{B}} b_{x} x^{*} \Longrightarrow$$

$$-\sum_{\mu} a_{\mu} \sum_{d \in C_{\mu}} (de - ed) = \sum_{x \in \mathcal{B}} b_{x} x^{*} \Longrightarrow$$

$$-\sum_{x \in \mathcal{B}} b_{x} x = \sum_{x \in \mathcal{B}} b_{x} x^{*} \Longrightarrow$$

$$-\sum_{x \in \mathcal{B}} b_{x} x = \sum_{x \in \mathcal{B}} b_{x} x^{*} \Longrightarrow$$

$$-\sum_{x \in \mathcal{B}} b_{x} x = \sum_{x \in \mathcal{B}} b_{x} x^{*}.$$

The next proposition is another case of a basis diagram x, such that $b_x = 0$.

Proposition 5.3. Let x = Q $\in B_{r,1}(\delta)$ for some $Q \in S_{r-1}$ then $b_x = 0$.

Proof. Let $d \in \mathcal{B}$ such that $de = \delta^m x$, $m \in \{0,1\}$. We have the following cases:

- (1) $d = \mathcal{Q} \in S_{r-1} \subseteq B_{r,1}(\delta)$, with \mathcal{Q} as defined in the diagram of x. We have de = x.
- (2) d = x. We have $de = \delta x$.
- (3) d = x(i, r), where (i, r) corresponds to the permutation diagram swapping i and r. We have de = x.


We notice that for every diagram d as described above, there is a diagram $d' \in \mathcal{B}$ such that $ed' = \delta^m x$, $m \in \{0,1\}$. More precisely, for the cases (1) and (2), the diagram d' = d. For case (3), we have d' = (i, r)x. We notice that the diagrams d' we have here are all the diagrams $d' \in \mathcal{B}$ such that $de = \delta^m x$, $m \in \{0,1\}$. In order to prove that $b_x = 0$ it is enough to prove that in each case, d and d' have the same walled generalized cycle type. Then the terms in the sum in equation (4.1) cancel out and, hence, $b_x = 0$. For cases (1) and (2) this is obvious. It remains to prove that the diagrams x(i, r) and (i, r)x have the same walled generalized cycle type.

There are two types of parts in the walled generalized cycle type c(x) of the diagram x: The part $p_1 = NS$ and the parts p_j , j = 2, 3, ..., k, which are of one of the forms: L, LL, LLL, Let p_{j_0} be the part, which belongs to the connected component with vertex i on top.

In the diagram of d there is a northern arc, which connects the vertices r and r+1 on top row and a southern arc, which connects the vertices i and r on bottom row. The propagating lines are the same as the ones appearing in the diagram of x, with one difference: The propagating line which connects the vertex i on bottom row with the vertex i' on top row in the diagram of x, it connects now the vertex r on bottom row with the vertex i' on top row. Therefore, $c(d) = \{NSp_{j_0}, p_2, p_3, p_{j_0-1}, p_{j_0+1}, \ldots, p_k\}.$


Similarly, in the diagram of d' there are two arcs, which are the flipped arcs of the diagram of d and the propagating lines remaining from the diagram of x with again one difference: the propagating line which connects the vertex i on top row with the vertex i' on bottom row in the diagram of x, it connects now the vertex r on top row with the vertex i' on the bottom row. Therefore, we have $c(d') = \{NSp_{j_0}, p_2, p_3, p_{j_0-1}, p_{j_0+1}, \ldots, p_k\} = c(d)$.

We consider now the system (4.2). According to Proposition 5.3, the only possible diagrams x with $b_x \neq 0$ are of the form



Note that the second case can be obtained from the first by applying the flip map *. Therefore, by Proposition 5.2, it is enough to consider only the first case.

Proposition 5.4. Let $x, y \in B_{r,1}(\delta)$ be of the form



with $y = \sigma x \sigma^{-1}$, for some $\sigma \in S_r$. Then, $b_x = b_y$.

Proof. We first notice that, since both diagrams x and y have the northern arc that connects the vertices r and r+1, we must have $\sigma \in S_{r-1}$. Therefore, $\sigma e = e\sigma$ and $\sigma^{-1}e = e\sigma^{-1}$.

We now conjugate Equation (5.1) by σ and we get:

$$\sum_{\mu} a_{\mu} \sum_{d \in C_{\mu}} \sigma^{-1} (de - ed) \sigma = \sum_{x \in \mathcal{B}} b_{x} \sigma^{-1} x \sigma \implies$$


$$\sum_{\mu} a_{\mu} \sum_{d \in C_{\mu}} (\sigma^{-1} de \sigma - \sigma^{-1} ed \sigma) = \sum_{x \in \mathcal{B}} b_{x} \sigma^{-1} x \sigma \implies$$

$$\sum_{\mu} a_{\mu} \sum_{d \in C_{\mu}} (\sigma^{-1} d\sigma e - e\sigma^{-1} d\sigma) = \sum_{x \in \mathcal{B}} b_{x} \sigma^{-1} x \sigma \implies$$

$$\sum_{\mu} a_{\mu} \sum_{d \in C_{\mu}} (de - ed) = \sum_{x \in \mathcal{B}} b_{x} \sigma^{-1} x \sigma \implies$$

$$\sum_{\mu} b_{x} x = \sum_{x \in \mathcal{B}} b_{\sigma x \sigma^{-1}} x.$$

Definition 5.5. Let $x \in B_{r,1}(\delta)$ be of the form



where $i_x \neq r$. The diagram x defines the following bijection, given by the propagating lines in the diagram x:

$$f_x: \{1, \dots, r-1\} \to \{1, \dots, \widehat{i_x}, \dots, r\}.$$

We associate two diagrams in $B_{r,1}(\delta)$ to x as follows:


(i) Define $\sigma_x \in S_r \times S_1 \subseteq B_{r,1}(\delta)$ by

$$\sigma_x(k) = \begin{cases} f_x(k), & \text{if } k \in \{1, \dots, r-1\} \\ i_x, & \text{if } k = r \\ r+1, & \text{if } k = r+1 \end{cases}$$


(ii) Define $z_x \in B_{r,1}(\delta)$ as follows: The vertices i_x and r+1 on top row (respectively, on bottom row) are connected by a northern arc (respectively, a southern arc). The propagating lines are given by the permutation $\tau_x : \{1, \ldots, \widehat{i_x}, \ldots, r\} \to \{1, \ldots, \widehat{i_x}, \ldots, r\}$, defined by

$$\tau_x(k) = \begin{cases} f_x(k), & \text{if } k \neq r \\ f_x(i_x), & \text{if } k = r \end{cases}$$

For example, let $x \in B_{6,1}(\delta)$ be the diagram



Then, we have: $f_x(1) = 4$, $f_x(2) = 1$, $f_x(3) = 2$, $f_x(4) = 5$, $f_x(5) = 6$. Therefore:



Proposition 5.6. Let x, σ_x and z_x be as in 5.5, then we have:

- (i) $e\sigma_x = x$. Moreover, $c(\sigma_x)$ is obtained from c(x) by replacing NS by L and σ_x is the unique diagram $y \in B_{r,1}(\delta)$ with c(y) without NS satisfying ey = x.
- (ii) $ez_x = x$. Moreover, $c(z_x)$ is obtained from c(x) by removing NS and adding it back as a trivial part and z_x is the unique diagram $y \in B_{r,1}(\delta)$ with c(y) having NS only as trivial part satisfying ey = x.

Proof. The result follows by definition of σ_x , z_x , concatenation of diagrams and the definition of walled generalized cycle type.

Corollary 5.7. The set of equations

viewed as equations in variables a_{μ} as defined in equation (4.1) are linearly independent.

Proof. We represent the aforementioned equations by a matrix, whose rows are the b_x 's and columns the coefficients a_u .

By Proposition 5.6, each b_x has a unique factor a_μ appearing with coefficient 1, such that μ has no NS, namely $a_{c(\sigma_x)}$, and a unique factor $a_{\mu'}$ (appearing also with coefficient 1), such that μ' has NS as a trivial part, namely $a_{c(z_x)}$.

Now, note that $c(\sigma_x)$ and $c(z_x)$ determine c(x), therefore the matrix cannot have the following form:

Therefore, the matrix is of the following form and, hence, the equations linearly independent.

Theorem 5.8. For any $\delta \in \mathbb{C}$, the centre of $B_{r,1}(\delta)$ is given by the algebra of supersymmetric polynomials evaluated at the Jucys-Murphy elements. Moreover, its dimension is given by $\dim Z(B_{r,1}(\delta)) = |\Lambda_{r,1}|$.

References

- [1] R.. Brauer, On algebras which are connected with the semisimple continuous groups, Ann. of Math.38 (1937), 857–872.
- [2] J. Brundan, C. Stroppel, Gradings on walled Brauer algebras and Khovanov's arc algebra, Adv. Math. 231(2) (2012), 709—773.
- [3] A. Cox, M. De Visscher, Diagrammatic Kazhdan-Lusztig theory for the (walled) Brauer algebra, J. Algebra **340**(1) (2011), pp. 151–181
- [4] A. Cox, M. De Visscher, S. Doty, P. Martin, On the blocks of the walled Brauer algebra, J. Algebra 320(1) (2008), 169-212.
- [5] A. Cox, M. De Visscher, P. Martin, The blocks of the Brauer algebra in characteristic zero, Representation Theory 13 (2009), pp. 272–308
- [6] A. A. Jucys, Symmetric polynomials and the centre of the symmetric group ring, Rep. Mat. Phys. 5 (1974), 107–112.
- [7] J. H. Jung, M. Kim, M., Supersymmetric polynomials and the center of the walled Brauer algebra, Algebr. and Representat. Theory 23 (2020), 1945–1975.
- [8] K. Koike, On the decomposition of tensor products of the representations of classical groups: by means of universal characters, Adv. Math., 74 (1989), 57–86.
- [9] P. P. Martin, The decomposition matrices of the Brauer algebra over the complex field, Transactions of the American Mathematical Society 367, No. 3 (2015), pp. 1797-1825
- [10] G. E. Murphy, A new construction of Young's seminormal representation of the symmetric group,, J. Algebra 69 (1981), 287–291.
- [11] G. E. Murphy, The idempotents of the symmetric group and Nakayama's conjecture, J. Algebra 81 (1983), 258–265.
- [12] A. Okounkov, A. Vershik, A new approach to the representation theory of symmetric groups, Selecta. Math. New Ser. 2 (1996) 581-605.
- [13] H. Rui, A criterion on the semisimple Brauer algebras, J. Comb. Theory Ser. A 111 (2005), 78–88.
- [14] H. Rui, Y. Su, Affine walled Brauer algebras and super Schur-Weyl duality, Adv. Math. 285 (2015), 28–71.
- [15] A. Sartori, C. Stroppel, Walled Brauer algebras as idempotent truncations of level 2 cyclotomic quotients, J. Algebra 440 (2015), 602–638.
- [16] A. Shalile, On the center of the Brauer algebra, Algebr. and Representat. Theory 16 (2013) 65–100.
- [17] V. G. Turaev, Operator invariants of tangles, and R-matrices, Izv. Akad. Nauk SSSR Ser. Mat., 53(5) (1989), 1073—1107.
- [18] H. Wenzl, On the structure of Brauer's centralizer algebras, Ann. of Math. 128(1), (1988) 173–193.
- [19] H. Weyl, The classical groups, their invariants and representations, Princeton University Press, 1946.

Institut für Diskrete Strukturen und Symbolisches Rechnen, Universität Stuttgart, Pfaffenwaldring 57, 70569 Stuttgart, Germany.

Email address: eirini.chavli@mathematik.uni-stuttgart.de

DEPARTMENT OF MATHEMATICS, CITY UNIVERSITY OF LONDON, NORTHAMPTON SQUARE, LONDON EC1V 0HB, UNITED KINGDOM.

Email address: maud.devisscher.1@city.ac.uk

SCHOOL OF MATHEMATICS, UNIVERSITY OF LEEDS, LS2 9JT, UNITED KINGDOM.

Email address: a.e.parker@leeds.ac.uk

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF COLORADO BOULDER, CAMPUS BOX 395, 80309-0395, UNITED STATES.

Email address: Sarah.Salmon@colorado.edu

Science, Technology, Engineering and Mathematics Division Faculty, Morehouse College,830 Westview Drive, S.W. Atlanta, GA 30314, United States.

Email address: ulrica.wilson@morehouse.edu