PHASE MIXING FOR THE HARTREE EQUATION AND LANDAU DAMPING IN THE SEMICLASSICAL LIMIT

MARNIE SMITH

ABSTRACT. The asymptotic behaviour of the Hartree equation is studied for short-range interaction potentials near translation-invariant steady states satisfying the Penrose stability condition. Phase-mixing estimates in finite regularity are derived, demonstrating density decay and scattering of solutions in weighted quantum Sobolev spaces. These results provide a quantum analogue of Landau damping in classical plasma physics and remain uniform in the semiclassical limit, bridging the quantum and classical regimes.

CONTENTS

1.	Introduction	T
2.	Linearised phase mixing	14
3.	Energy estimates on the density	22
4.	Energy estimates on the density matrix	28
Re	References	

1. INTRODUCTION

1.1. The Hartree equation. The *Hartree equation*, a mean-field model describing the dynamics of infinitely many interacting fermions, is expressed as:

$$\begin{cases} i\hbar\partial_t\gamma = \left[-\frac{\hbar^2}{2}\Delta + w * \rho_\gamma, \gamma\right],\\ \gamma(0) = \gamma_{\rm in}, \end{cases}$$
(H)

1

where $\gamma(t)$ is a positive, self-adjoint operator on $L^2(\mathbb{R}^d; \mathbb{C})$. The function $\rho_{\gamma}(t, x) = \gamma(t, x, x)$ represents the density, where $\gamma(t, x, y)$ denotes the integral kernel of $\gamma(t)$. The nonlinear convolution term $w * \rho_{\gamma}$ describes the mean-field interaction between the particles, where $w : \mathbb{R}^d \to \mathbb{R}$ is a specified two-body interaction potential function. Only short-range potentials $w \in L^1(\mathbb{R}^d)$ are considered, excluding the Coulomb potential. For the nonlinear statement, an additional condition is required:

$$|\widehat{w}(k)| \lesssim \langle k \rangle^{-M-1/2}$$

where $M := \lfloor \frac{d+1}{2} \rfloor$. The parameter \hbar represents the Planck constant, and the semiclassical limit involves taking $\hbar \to 0$.

For any function $g \in L^{\infty}(\mathbb{R}^d; \mathbb{R}_+)$, define the translation-invariant operator

$$\gamma_g := \hbar^d g(-i\hbar\nabla)$$

Introduction

Department of Pure Mathematics and Mathematical Statistics, University of Cambridge $\it E-mail~address:~ms2724@cam.ac.uk.$

Date: 20/12/2024.

on $L^2(\mathbb{R}^d;\mathbb{C})$. The integral kernel of γ_g is

$$\gamma_g(x,y) = \frac{1}{(2\pi)^d} \widehat{g}\left(\frac{y-x}{\hbar}\right),$$

and its constant density is given by

$$\rho_{\gamma_g}=\frac{1}{(2\pi)^d}\widehat{g}(0)=\frac{1}{(2\pi)^d}\int_k g(k)\,dk.$$

These operators represent a class of steady-state solutions to the Hartree equation (H), as they satisfy commutation relations $[-\Delta, \gamma_g] = 0$ and $[w * \rho_{\gamma_g}, \gamma_g] = 0$, where $w * \rho_{\gamma_g}$ is constant.

The potential and the steady state are assumed to satisfy a linear stability criterion, known in kinetic theory as the *Penrose condition*.

Definition 1.1 (Penrose condition). A potential $w \colon \mathbb{R}^d \to \mathbb{R}$ and steady state $\gamma_g = \hbar^d g(-i\hbar\nabla)$ satisfy the Penrose condition if there exists a constant $\kappa > 0$ such that the linear operator

$$\widetilde{\mathscr{L}}(\lambda,k) = \frac{1}{(2\pi)^d} \frac{2}{\hbar} \widehat{w}(k) \sin\left(\frac{1}{2}\hbar t |k|^2\right) \widehat{g}(kt) \,.$$
$$\inf_{\substack{\mathrm{Re}\,\lambda \ge 0\\k \in \mathbb{R}^d}} \left| 1 + \widetilde{\mathscr{L}}(\lambda,k) \right| \ge \kappa.$$
(P)

satisfies

Proposition 2.6 provides sufficient conditions on
$$w$$
 and g that guarantee the Penrose condition (P). When
these conditions do not include a smallness assumption, only defocusing potentials $\hat{w}(k) \geq 0$ can be treated.

The trace $\operatorname{Tr} \gamma$ determines the total number of particles. For the translation-invariant steady states γ_g , this number is infinite due to the constant positive density, resulting in $\operatorname{Tr} \gamma_g = \int \rho_{\gamma_g} dx = +\infty$. Define the perturbation $Q := \gamma - \gamma_g$, where γ is a solution of (H). The perturbation Q satisfies the *nonlinear Hartree equation*:

$$\begin{cases} i\hbar\partial_t Q = \left[-\frac{\hbar^2}{2}\Delta + w * \rho_Q, Q + \gamma_g \right],\\ Q(0) = Q_{\rm in}. \end{cases}$$
(NH)

The main result establishes phase-mixing estimates for Q. Theorem 1.5 demonstrates mode-by-mode decay of the density ρ_Q and the scattering behaviour of Q, with these estimates remaining uniform in the semiclassical limit $\hbar \to 0$.

1.2. Existing results. The well-posedness of the Hartree–Fock model, which includes an exchange term, was established by Bove, Da Prato, and Fano [5, 6], and independently by Chadam [7], under the assumption that the initial data is a trace-class operator. Lewin and Sabin [15] extended the theory to the Hartree equation with initial conditions as perturbations of a translation-invariant state. Unlike earlier works, their model accommodates infinitely many particles and relies on the conservation of relative free energy to ensure well-defined evolution. Subsequently, in [16], they studied the asymptotic stability of stationary states in two dimensions, showing weak convergence of the system to the stationary state over time. These findings were later generalised to higher dimensions [9] and extended to the nonlinear Schrödinger system [8, 11]. Nguyen and You extended the analysis of the Hartree equation to the Coulomb potential, a case that differs significantly because the Penrose condition is never satisfied for any steady state. Their work first addressed the linearised behaviour [19] and later established decay estimates and a modified scattering result for the nonlinear equation near vacuum [18].

The classical analogue is *Landau damping*, a well-studied phenomenon in plasma physics where small perturbations in plasma density decay over time due to collective particle motion. This behaviour is described by the Vlasov equation

$$\partial_t f + v \cdot \nabla_x f + \nabla_x (w * \rho_f) \cdot \nabla_v f = 0, \tag{1}$$

where f = f(t, x, v) is a distribution function, w is an interaction potential and $\rho_f = \int_v f dv$ is the spatial density. In his foundational work, Landau demonstrated that solutions of the linearised Vlasov equation around homogeneous steady states, under specific stability conditions, exhibit this time decay [13]. The first proof of Landau damping for the nonlinear Vlasov equation was provided by Mouhot and Villani in analytic regularity [17], and has remained an active area of research since, e.g. [3, 1, 10, 2, 12].

The Planck constant $\hbar \approx 6.626 \times 10^{-34}$ J s serves as a bridge between the macroscopic world one experiences daily and the microscopic realm of quantum mechanics. Its small value allows classical mechanics to effectively describe the behaviour of large-scale systems, with quantum effects viewed as perturbations of classical behaviour. Establishing results for the Hartree equation without assuming the Planck constant is a fixed positive value is important, as this allows for the recovery of classical behaviour in the semiclassical limit. The relationship between the Hartree and Vlasov equations is explicitly established in this limit: as the Planck constant approaches zero, the solution of the Hartree equation converge to those of the Vlasov equation.



Lewin and Sabin, working within the framework of the nonlinear Hartree equation near a translation-invariant stationary state with a short-range interaction potential, proved that solutions of the Hartree equation converge to solutions of the Vlasov equation in the semiclassical limit [14].

In a recent preprint, You [20] has presented related results to this present paper, also establishing phasemixing estimates for the nonlinear Hartree equation with short-range potentials. The works were completed independently and have fundamental differences. You demonstrates the decay of density and scattering in the Hilbert–Schmidt norm, under weaker assumptions than those required in this paper. The analysis exploits the finite difference structure of the Hartree equation, which uses a commutator instead of a derivative. To illustrate this point, upon taking suitable transforms as described below, the Hartree equation can be expressed in terms of a quantity f^{\hbar} , with the nonlinear term taking the approximate form:

$$\frac{f^{\hbar}(v+\frac{\hbar}{2}\nabla_x)-f^{\hbar}(v-\frac{\hbar}{2}\nabla_x)}{\hbar}(w*\rho_{f^{\hbar}}).$$
(2)

Heuristically, if $f^{\hbar} \to f$ as $\hbar \to 0$, then the quantity (2) converges to

 $\nabla_v f \cdot \nabla_x (w * \rho_f),$

which is the nonlinear term in the Vlasov equation (1). In [20], \hbar is fixed, and hence both terms in (2) can be treated separately, allowing the Hartree equation to be viewed as "one derivative better" than the Vlasov equation, resulting in less loss of regularity. However, the trade-off is that the estimates are no longer valid in the semiclassical limit. The present work extends methods developed by Bedrossian, Masmoudi and Mouhot for the Vlasov equation [1], enabling uniform estimates in the limit $\hbar \to 0$. Additionally, scattering is established in a stronger norm, specifically within a weighted quantum Sobolev space \mathcal{H}_M^{σ} .



FIGURE 1. Semiclassical limit and scattering map.

4 PHASE MIXING FOR THE HARTREE EQUATION AND LANDAU DAMPING IN THE SEMICLASSICAL LIMIT

A corollary of the uniform scattering result for the Hartree equation established here is the construction of a map to the scattering result for the Vlasov equation via the semiclassical limit, as illustrated in Figure 1. This work demonstrates that a solution $\gamma(t)$ to the Hartree equation with appropriate initial data γ_{in} scatters along the free flow to an operator γ_{∞} as $t \to \infty$, uniformly in \hbar . The analogous result for the Vlasov equation, where a solution f(t) with suitable initial data f_{in} scatters along the free flow to an operator f_{∞} as $t \to \infty$, was established in [1]. The semiclassical limit in this context, as shown in [14], asserts that if $\gamma_{in} \to f_{in}$ as $\hbar \to 0$, then $\gamma(t) \to f(t)$ as $\hbar \to 0$ for all $t \ge 0$. The novel conclusion, that $\gamma_{\infty} \to f_{\infty}$ as $\hbar \to 0$, thus completing Figure 1, follows from the Moore–Osgood theorem by leveraging the uniform convergence in the Hartree scattering statement.

1.3. Quantum operators and norms. The quantum analogue of spatial and velocity derivatives and weights are defined as follows.

Definition 1.2 (Quantum operators). The quantum analogue of spatial and velocity derivatives are

$$\boldsymbol{\nabla}_x P := [\nabla, P], \quad (\boldsymbol{\nabla}_x P)(x, y) = (\nabla_x + \nabla_y) P(x, y)$$

$$\boldsymbol{\nabla}_{\xi} P := [\frac{x}{i\hbar}, P], \quad (\boldsymbol{\nabla}_{\xi} P)(x, y) = (\frac{x-y}{i\hbar}) P(x, y).$$

The quantum analogue of spatial and velocity weights are

$$\begin{split} \boldsymbol{x}P &:= \frac{1}{2}(\boldsymbol{x}P + P\boldsymbol{x}), \qquad (\boldsymbol{x}P)(\boldsymbol{x},\boldsymbol{y}) = (\frac{\boldsymbol{x}+\boldsymbol{y}}{2})P(\boldsymbol{x},\boldsymbol{y}), \\ \boldsymbol{\xi}P &:= -\frac{i\hbar}{2}(\nabla P + P\nabla), \quad (\boldsymbol{\xi}P)(\boldsymbol{x},\boldsymbol{y}) = -\frac{i\hbar}{2}(\nabla_{\boldsymbol{x}} - \nabla_{\boldsymbol{y}})P(\boldsymbol{x},\boldsymbol{y}) \end{split}$$

The weighted Sobolev norm $\|\cdot\|_{H^{\sigma}_{M}}$ is now defined.

Definition 1.3 (Weighted Sobolev norm). Let $\sigma \ge 0$ and let $M \ge 0$ be an integer. For a function g = g(v), the weighted Sobolev norm H_M^{σ} is defined as

$$\|g\|_{H^{\sigma}_{M}} := \sum_{|\alpha| \leq M} \|v^{\alpha}g\|_{H^{\sigma}} = \sum_{|\alpha| \leq M} \|\langle \nabla \rangle^{\sigma} (v^{\alpha}g)\|_{L^{2}}.$$

For a function f = f(z, v), the weighted Sobolev norm H_M^{σ} is defined as,

$$\|f\|_{H^{\sigma}_{M}} := \sum_{|\alpha| \le M} \|v^{\alpha}f\|_{H^{\sigma}} = \sum_{|\alpha| \le M} \|\langle \nabla_{x}, \nabla_{v} \rangle^{\sigma} (v^{\alpha}f)\|_{L^{2}}$$

where the weight M applies only to the v-variable.

Operator norms, quantum analogous to Lebesgue and Sobolev norms, are introduced.

Definition 1.4 (Quantum norms). For an operator Q on $L^2(\mathbb{R}^d; \mathbb{C})$ with integral kernel Q = Q(x, y), the following norms are defined.

(a) Quantum Lebesgue norm \mathcal{L}^2 :

$$||Q||_{\mathcal{L}^2} := (2\pi\hbar)^{-d/2} ||Q(x,y)||_{L^2_{x,y}}$$

(b) Quantum Sobolev norm \mathcal{H}^{σ} :

$$\|Q\|_{\mathcal{H}^{\sigma}} := \|\langle \boldsymbol{\nabla}_x, \boldsymbol{\nabla}_{\xi} \rangle^{\sigma} Q\|_{\mathcal{L}^2}.$$

(c) Weighted quantum Sobolev norm \mathcal{H}_M^{σ} :

$$\|Q\|_{\mathcal{H}^{\sigma}_{M}} := \sum_{|lpha|,|eta| \leq M} \left\| oldsymbol{x}^{eta} oldsymbol{\xi}^{lpha} Q
ight\|_{\mathcal{H}^{\sigma}}$$

Remark. The norm $\|Q(x,y)\|_{L^2_{x,y}}$ is equivalent to both the Hilbert–Schmidt norm and the \mathfrak{S}^2 -Schatten norm, implying that the quantum Lebesgue norm is an \hbar -scaled adjustment of these norms. Additionally, unlike the classical norm H^{σ}_{M} , which applies weights only to the velocity variable, the quantum Sobolev norm \mathcal{H}^{σ}_{M} incorporates weights in both the position variable \boldsymbol{x} and the momentum variable $\boldsymbol{\xi}$.

1.4. **Main result.** The main result establishes decay and scattering properties for solutions of the nonlinear Hartree equation. The theorem is stated as follows:

Theorem 1.5 (Nonlinear phase mixing). Fix dimension $d \ge 3$ and $M := \lceil (d+1)/2 \rceil$. Assume $0 < \hbar \le 1$ and constants $N_0 > \sigma_1 > \sigma_0$ such that

$$\sigma_1 \ge d+7, \quad N_0 - \sigma_1 > 3d/2 + 5/2 \quad and \quad \sigma_1 - \sigma_0 > d+1.$$

Let $g \in H^{N_0+7/2+\delta_0}_{2+\lceil d/2\rceil}(\mathbb{R}^d;\mathbb{R}_+)$ for some $0 < \delta_0 \ll 1$, and let $w \in L^1(\mathbb{R}^d)$ satisfy

$$|\widehat{w}(k)| \leq \langle k \rangle^{-M-1/2}$$

for all $k \in \mathbb{R}^d$. Assume g and w satisfy the Penrose condition (P) with constant κ . Then there exist constants $C = C(N_0, w, g, \kappa, d, \delta_0) > 0$ and $\varepsilon_0 > 0$ such that, for any $0 < \varepsilon \leq \varepsilon_0$, if Q_{in} satisfies

$$\|Q_{\mathrm{in}}\|_{\mathcal{H}^{N_0}} \leq \varepsilon$$

the solution Q(t) to the nonlinear Hartree equation (NH) satisfies the following.

• Density decay: for all $k \in \mathbb{R}^d$ and $t \ge 0$,

$$\langle k, kt \rangle^{\sigma_1} |\widehat{\rho_Q}(t,k)| \le C\varepsilon.$$
 (T1)

• Scattering: there exists an operator $Q_{\infty} \in \mathcal{H}_{M}^{\sigma_{0}}(\mathbb{R}^{d};\mathbb{C})$ such that, for all $t \geq 0$,

$$\left\| e^{-i\frac{\hbar}{2}t\Delta}Q(t)e^{i\frac{\hbar}{2}t\Delta} - Q_{\infty} \right\|_{\mathcal{H}_{M}^{\sigma_{0}}} \le C\varepsilon\langle t\rangle^{-d/2}.$$
 (T2)

Remark. The constant C is independent of $0 < \hbar \leq 1$, ensuring that the damping and scattering results (T1) and (T2) remain valid in the semiclassical limit $\hbar \to 0$.

Theorem 1.5 implies physical decay of the density in L^p , $p \in [2, \infty]$.

Corollary 1.6 (Physical decay). Let $p \in [2, \infty]$ assume that the assumptions in Theorem 1.5 are satisfied. Then, for any $n < \sigma_1 - d(1 - 1/p)$,

$$\langle t \rangle^{d(1-\frac{1}{p})} \| \langle \nabla_x, t \nabla_x \rangle^n \rho(t, \cdot) \|_{L^p(\mathbb{R}^d)} \lesssim \varepsilon.$$

Proof. By the Hausdorff–Young inequality,

$$\left\| \langle \nabla_x, t \nabla_x \rangle^n \rho_Q(t, \cdot) \right\|_{L^p} \lesssim_p \left(\int_k \langle k, kt \rangle^{p'n} |\widehat{\rho_Q}(t, k)|^{p'} dk \right)^{\frac{1}{p'}} \lesssim \left(\int_k \langle k, kt \rangle^{p'(n-\sigma_1)} dk \right)^{\frac{1}{p'}} \lesssim \langle t \rangle^{-d(1-\frac{1}{p})}. \qquad \Box$$

1.5. Notations and tools.

1.5.1. Fourier transform. The following conventions are adopted for the Fourier transform of functions g = g(v) and f = f(z, v):

$$\widehat{g}(p) := \int_{v} e^{-ip \cdot v} g(v) \, dv, \qquad \widehat{f}(k, p) := \int_{z} \int_{v} e^{-ik \cdot z} e^{-ip \cdot v} f(z, v) \, dv dz.$$

On the Fourier side, the operators defined in Definition 1.2 obey the following relations:

$$\begin{split} \widehat{\nabla_{x}P}(k,p) &= i(k+p)\widehat{P}(k,p), \\ \widehat{\nabla_{\xi}P}(k,p) &= \frac{1}{\hbar}(\nabla_{k}-\nabla_{p})\widehat{P}(k,p), \\ \widehat{\nabla_{\xi}P}(k,p) &= \frac{1}{\hbar}(\nabla_{k}-\nabla_{p})\widehat{P}(k,p), \\ \widehat{\nabla_{\xi}P}(k,p) &= \frac{i}{2}(\nabla_{k}+\nabla_{p})\widehat{P}(k,p), \\ \widehat{\xi}\widehat{P}(k,p) &= \frac{i}{2}(\nabla_{k}+\nabla_{p})\widehat{P}(k,p), \\ \widehat{\xi}\widehat{P}(k,p) &= \frac{\hbar}{2}(k-p)\widehat{P}(k,p), \\ \widehat{\xi}\widehat{P}(k,p) &= \frac{\hbar}{2}(k-p)\widehat{P}(k,p)$$

1.5.2. Density of an operator. The density of an operator Q is defined as the integral kernel along the diagonal, $\rho_Q(x) := Q(x, x)$. Its Fourier transform can be computed from the Fourier transform of the integral kernel \hat{Q} , as shown in the following lemma.

Lemma 1.7. Let Q be an operator on $L^2(\mathbb{R}^d; \mathbb{C})$ identified with its integral kernel Q = Q(x, y). Then its density $\rho_Q(x) := Q(x, x)$ has Fourier representation

$$\widehat{\rho_Q}(t,k) = \frac{1}{(2\pi)^d} \int_p \widehat{Q}(k-p,p) \, dp = \frac{1}{(2\pi)^d} \int_p \widehat{Q}\left(\frac{k}{2} + p, \frac{k}{2} - p\right) \, dp.$$

Proof. Writing Q(x, x) as the inverse Fourier transform of \widehat{Q} evaluated at (x, x),

$$\begin{split} \widehat{\rho_Q}(t,k) &= \int_x e^{-ik \cdot x} \left[\frac{1}{(2\pi)^{2d}} \int_{k'} \int_p e^{ik' \cdot x} e^{ip \cdot x} \widehat{Q}(t,k',p) \, dp dk' \right] \, dx \\ &= \frac{1}{(2\pi)^d} \int_{k'} \int_p \widehat{Q}(t,k',p) \left(\frac{1}{(2\pi)^d} \int_x e^{i(k'+p-k) \cdot x} \, dx \right) \, dp dk' \\ &= \frac{1}{(2\pi)^d} \int_{k'} \int_p \widehat{Q}(t,k',p) \delta(k'+p-k) \, dp dk' \\ &= \frac{1}{(2\pi)^d} \int_p \widehat{Q}(t,k-p,p) \, dp. \end{split}$$

The second representation in the lemma follows from the change of variable $p \mapsto k/2 - p$.

1.5.3. *Wigner transform.* The Wigner transform facilitates the representation of the density matrix in the phase space.

Definition 1.8 (Wigner transform). For an operator P on $L^2(\mathbb{R}^d; \mathbb{C})$ with integral kernel P(x, y), define the Wigner transform of P, $W[P]: \mathbb{R}^d \times \mathbb{R}^d \to \mathbb{C}$, by

$$W[P](x,\xi) = \frac{1}{(2\pi)^d} \frac{1}{\hbar^d} \int_{\mathbb{R}^d} e^{-i\frac{\xi}{\hbar} \cdot y} P\left(x + \frac{y}{2}, x - \frac{y}{2}\right) \, dy.$$

The Fourier transform of the Wigner transform of P satisfies

$$\widehat{W[P]}(k,\eta) = \int_{x} e^{-ik \cdot x} P\left(x - \frac{\hbar\eta}{2}, x + \frac{\hbar\eta}{2}\right) \, dx = \frac{1}{(2\pi)^d} \int_{p} e^{-i\hbar\eta \cdot p} \widehat{W[P]}\left(\frac{k}{2} + p, \frac{k}{2} - p\right) \, dp$$

and can be controlled by the \mathcal{H}^0_M norm of P via the Sobolev embedding.

Lemma 1.9. Suppose $P \in \mathcal{H}^0_M(\mathbb{R}^d; \mathbb{C})$. Then the Wigner transform of P satisfies, for all $k, \ell \in \mathbb{R}^d$,

$$\widetilde{W}[\widetilde{P}](k,\ell) \bigg| \lesssim_d \|P\|_{\mathcal{H}^0_M}$$

Proof. By Definition 1.2, Cauchy–Schwarz and the Sobolev embedding, for $0 < \tilde{\delta} \leq 1/2$,

$$\begin{split} \widehat{W[P]}(k,\eta) \bigg| &= \left| \int_{x} e^{-ik \cdot x} P\left(x - \frac{\hbar}{2}\eta, x + \frac{\hbar}{2}\eta\right) dx \right| \\ &\lesssim \left(\int_{x} \left| \langle x \rangle^{\frac{d}{2} + \widetilde{\delta}} P\left(x - \frac{\hbar}{2}\eta, x + \frac{\hbar}{2}\eta\right) \right|^{2} dx \right)^{\frac{1}{2}} \\ &\lesssim \left(\int_{x,y} \left| \langle \nabla_{y} \rangle^{\frac{d}{2} + \widetilde{\delta}} \left[\langle x \rangle^{\frac{d}{2} + \widetilde{\delta}} P\left(x - \frac{\hbar}{2}y, x + \frac{\hbar}{2}y\right) \right] \right|^{2} dx dy \right)^{\frac{1}{2}} \\ &= \left(\int_{x,y} \left| \langle \frac{\hbar}{2} (\nabla_{x} - \nabla_{y}) \rangle^{\frac{d}{2} + \widetilde{\delta}} \langle x \rangle^{\frac{d}{2} + \widetilde{\delta}} P\left(x - \frac{\hbar}{2}y, x + \frac{\hbar}{2}y\right) \right|^{2} dx dy \right)^{\frac{1}{2}} \\ &= \hbar^{-d/2} \left\| \langle \boldsymbol{\xi} \rangle^{\frac{d}{2} + \widetilde{\delta}} \langle x \rangle^{\frac{d}{2} + \widetilde{\delta}} P(x, y) \right\|_{L^{2}_{x,y}} \\ &\lesssim \hbar^{-d/2} \sum_{|\alpha|, |\beta| \leq M} \left\| \boldsymbol{\xi}^{\alpha} \boldsymbol{x}^{\beta} P(x, y) \right\|_{L^{2}_{x,y}}. \end{split}$$

The lemma then follows from the definition of the quantum Sobolev norm.

The operators defined in Definition 1.2 interact nicely with the Wigner transform:

Lemma 1.10. Suppose $P \in \mathcal{L}^2(\mathbb{R}^d; \mathbb{C})$. Then the Wigner transform of P satisfies the following:

$$\begin{aligned} \nabla_x W[P](x,\xi) &= W[\boldsymbol{\nabla}_x P](x,\xi), & ik \overline{W[P]}(k,\eta) = \overline{W[\boldsymbol{\nabla}_x P]}(k,\eta), \\ \nabla_\xi W[P](x,\xi) &= W[\boldsymbol{\nabla}_\xi] P(x,\xi), & i\eta \overline{W[P]}(k,\eta) = \overline{W[\boldsymbol{\nabla}_\xi P]}(k,\eta), \\ x W[P](x,\xi) &= W[\boldsymbol{x} P](x,\xi), & i\nabla_k \overline{W[P]}(k,\eta) = \overline{W[\boldsymbol{x} P]}(k,\eta), \\ \xi W[P](x,\xi) &= W[\boldsymbol{\xi} P](x,\xi), & i\nabla_\eta \widehat{W[P]}(k,\eta) = \overline{W[\boldsymbol{\xi} P]}(k,\eta). \end{aligned}$$

Proof. These statements follow directly from the definitions of the Wigner transform (Definition 1.8) and the quantum operators (Definition 1.2). \Box

The L^2 norm of W[P] is related to the \mathcal{L}^2 norm of P via the following relation:

Lemma 1.11. Suppose $P \in \mathcal{L}^2(\mathbb{R}^d; \mathbb{C})$. Then the Wigner transform of P satisfies

$$\|W[P](x,\xi)\|_{L^2_{x,\xi}} = (2\pi\hbar)^{-d/2} \|P(x,y)\|_{L^2_{x,y}} = (2\pi)^{-d/2} \|P\|_{\mathcal{L}^2}.$$

Proof. The result follows from the observation that

$$W[P](x,\xi) = (2\pi\hbar)^{-d} \mathcal{F}\left[P\left(x + \frac{1}{2}, x - \frac{1}{2}\right)\right] (\xi/\hbar)$$

combined with Plancherel's theorem.

1.6. Free phase mixing. Consider the free Hartree equation

$$\begin{cases} i\hbar\partial_t Q_{\rm FH} = \left[-\frac{\hbar^2}{2} \Delta, Q_{\rm FH} \right] \\ Q_{\rm FH}(0) = Q_{\rm in}, \end{cases}$$
(FH)

where $Q_{\rm FH}(t)$ is an operator on $L^2(\mathbb{R}^d;\mathbb{C})$. The free Hartree equation (FH) has the explicit solution

$$Q_{\rm FH}(t) = e^{i\frac{\hbar}{2}t\Delta}Q_{\rm in}e^{-i\frac{\hbar}{2}t\Delta},\tag{3}$$

with integral kernel

$$\begin{aligned} Q_{\rm FH}(t,x,y) &= e^{i\frac{\hbar}{2}t(\Delta_x - \Delta_y)}Q_{\rm in}(x,y),\\ \widehat{Q_{\rm FH}}(t,k,p) &= e^{-i\frac{\hbar}{2}t(|k|^2 - |p|^2)}\widehat{Q_{\rm in}}(k,p) \end{aligned}$$

The phase-mixing estimates in Theorem 1.5 show that, under the given assumptions, the solution to the nonlinear Hartree equation approaches the behaviour of a solution to the free Hartree equation as $t \to \infty$.

The density of the solution $Q_{\rm FH}$ is defined as $\rho_{Q_{\rm FH}}(t,x) := Q_{\rm FH}(t,x,x)$. Using Lemma 1.7, its Fourier transform is given by

$$\widehat{\rho_{Q_{\rm FH}}}(t,k) = \frac{1}{(2\pi)^d} \int_p e^{-i\hbar p \cdot kt} \widehat{Q_{\rm in}} \left(\frac{k}{2} + p, \frac{k}{2} - p\right) \, dp = \widehat{W[Q_{\rm in}]}(k,kt).$$

The following proposition establishes phase-mixing estimates for the density of solutions to the free Hartree equation, providing a bound in terms of the initial data.

Proposition 1.12 (Phase mixing for the free Hartree equation). Fix $N_0 \ge 0$. Let Q_{in} be an operator on $L^2(\mathbb{R}^d;\mathbb{C})$ identified with its integral kernel $Q_{\text{in}} = Q_{\text{in}}(x,y)$. The density of the solution Q_{FH} to the free Hartree equation (FH) satisfies

$$\langle k, kt \rangle^{N_0} \left| \widehat{\rho_{Q_{\mathrm{FH}}}}(t,k) \right| \le \|Q_{\mathrm{in}}\|_{\mathcal{H}^{N_0}}$$

7

8 PHASE MIXING FOR THE HARTREE EQUATION AND LANDAU DAMPING IN THE SEMICLASSICAL LIMIT

Proof. First observe that, using Lemma 1.10, for any $n \ge 0$ and $m \ge 0$,

$$|k|^{n}\widehat{\rho_{Q_{\mathrm{FH}}}}(t,k) = \overline{W\left[|\boldsymbol{\nabla}_{x}|^{n}Q_{\mathrm{in}}\right]}(k,kt) \quad \text{and} \quad |kt|^{m}\widehat{\rho_{Q_{\mathrm{FH}}}}(t,k) = \overline{W\left[|\boldsymbol{\nabla}_{\xi}|^{n}Q_{\mathrm{in}}\right]}(k,kt).$$

Combining these estimates gives

$$\langle k, kt \rangle^{N_0} \widehat{\rho_{Q_{\rm FH}}}(t, k) = \overline{W\left[\langle \boldsymbol{\nabla}_x, \boldsymbol{\nabla}_\xi \rangle^{N_0} Q_{\rm in} \right]}(k, kt)$$

The result then follows from Lemma 1.9.

The same estimates on the density of the nonlinear Hartree equation are obtained in Theorem 1.5, with a greater loss of regularity on the initial data. Moreover, the scattering statement (T2) indicates that the solution Q(t) of (NH) behaves like a solution of the free Hartree equation as $t \to \infty$.

1.7. **Outline of proof.** The proof strategy for Theorem 1.5 is outlined, including the Fourier and Wigner transforms of (NH) and the techniques employed. Bootstrap hypotheses are introduced and shown to imply the phase-mixing estimates and scattering properties.

1.7.1. Fourier transform of (NH). Let Q = Q(t, x, y) be the time-dependent integral kernel of the perturbation Q. The nonlinear equation (H) can be written as

$$i\hbar\partial_t Q(t,x,y) = -\frac{\hbar^2}{2} \left(\Delta_x - \Delta_y\right) Q(t,x,y) + \left(V(t,x) - V(t,y)\right) \left(\gamma_g(x,y) + Q(t,x,y)\right),$$

where $V(s, x) = w * \rho_Q(s, x)$. Applying Duhamel's formula provides an implicit solution to the linearised Hartree equation:

$$Q(t, x, y) = e^{i\frac{\hbar}{2}(\Delta_x - \Delta_y)t} Q_{in}(x, y) - \frac{i}{\hbar} \int_0^t e^{i\frac{\hbar}{2}(\Delta_x - \Delta_y)(t-s)} \left[V(s, x) - V(s, y) \right] \left(\gamma_g(x, y) + Q(s, x, y) \right) \, ds.$$
(4)

The Fourier transform representation of equation (4) is

$$\begin{split} \widehat{Q}(t,k,p) &= e^{-i\frac{\hbar}{2}(|k|^2 - |p|^2)t} \widehat{Q_{\text{in}}}(k,p) \\ &- \frac{i}{\hbar} \frac{1}{(2\pi)^d} \int_0^t e^{-i\frac{\hbar}{2}(|k|^2 - |p|^2)(t-s)} \left[\widehat{V}(s,k) *_k - \widehat{V}(s,p) *_p \right] \left(\widehat{\gamma_g}(k,p) + \widehat{Q}(s,k,p) \right) \, ds. \end{split}$$

Observing that $\widehat{\gamma}_{g}(k,p) = (2\pi)^{d} \hbar^{d} g(\hbar k) \delta(p = -k)$ and $\widehat{V}(t,k) = \widehat{w}(k) \widehat{\rho}_{Q}(t,k)$, the expression becomes

$$\begin{split} &Q(t,k,p) \\ &= e^{-i\frac{\hbar}{2}(|k|^2 - |p|^2)t}\widehat{Q_{\mathrm{in}}}(k,p) - i\hbar^{d-1}\widehat{w}(k+p)\left[g(-\hbar p) - g(\hbar k)\right]\int_0^t e^{-i\frac{\hbar}{2}(|k|^2 - |p|^2)(t-s)}\widehat{\rho_Q}(s,k+p)\,ds \\ &- \frac{i}{\hbar}\frac{1}{(2\pi)^d}\int_0^t \int_\ell e^{-i\frac{\hbar}{2}(|k|^2 - |p|^2)(t-s)}\widehat{w}(\ell)\widehat{\rho_Q}(s,\ell)\left(\widehat{Q}(s,k-\ell,p) - \widehat{Q}(s,k,p-\ell)\right)\,d\ell ds. \end{split}$$

1.7.2. Coordinate shift. Define the operator P to be the solution Q along the free flow:

$$Q(t,x,y) = e^{i\frac{\hbar}{2}(\Delta_x - \Delta_y)t} P(t,x,y), \qquad \widehat{Q}(t,k,p) = e^{-i\frac{\hbar}{2}(|k|^2 - |p|^2)t} \widehat{P}(t,k,p)$$

Then, P satisfies the equation

$$\begin{cases} \widehat{P}(t,k,p) = \widehat{Q_{\text{in}}}(k,p) + \int_{0}^{t} \widehat{P_{\text{L}}}(s,k,p) \, ds + \int_{0}^{t} \widehat{P_{\text{NL}}}(s,k,p) \, ds, \\ \widehat{P_{\text{L}}}(s,k,p) = -i\hbar^{d-1}\widehat{w}(k+p) \left[g(-\hbar p) - g(\hbar k)\right] e^{i\frac{\hbar}{2}(|k|^{2} - |p|^{2})s} \widehat{\rho_{Q}}(s,k+p), \\ \widehat{P_{\text{NL}}}(s,k,p) = -\frac{i}{\hbar} \frac{1}{(2\pi)^{d}} \int_{\ell} \widehat{w}(\ell) \widehat{\rho_{Q}}(s,\ell) \left(e^{-i\frac{\hbar}{2}(|k-\ell|^{2} - |k|^{2})s} \widehat{P}(s,k-\ell,p) - e^{-i\frac{\hbar}{2}(|p|^{2} - |p-\ell|^{2})s} \widehat{P}(s,k,p-\ell)\right) \, d\ell \\ (\text{NLa}) \end{cases}$$

An implicit equation for the density in terms of P,

$$\widehat{\rho_Q}(t,k) = \frac{1}{(2\pi)^d} \int_p e^{-i\hbar p \cdot kt} \widehat{P}(t,\frac{k}{2}+p,\frac{k}{2}-p) \, dp,$$

is

$$\begin{split} \widehat{\rho_Q}(t,k) \\ &= \widehat{\rho_{Q_{\rm FH}}}(t,k) - \frac{i\hbar^{d-1}}{(2\pi)^d} \widehat{w}(k) \int_p \left[g\left(\hbar(p-\frac{k}{2})\right) - g\left(\hbar(p+\frac{k}{2})\right) \right] \int_0^t e^{-i\hbar p \cdot k(t-s)} \widehat{\rho_Q}(s,k) \, ds dp \\ &- \frac{i\hbar^{-1}}{(2\pi)^{2d}} \int_0^t \int_\ell \widehat{w}(\ell) \widehat{\rho_Q}(s,\ell) \left(e^{-i\frac{\hbar}{2}\ell \cdot k(t-s)} - e^{i\frac{\hbar}{2}\ell \cdot k(t-s)} \right) \int_p e^{-i\hbar p \cdot (kt-\ell s)} \widehat{P}\left(s, \frac{k-\ell}{2} + p, \frac{k-\ell}{2} - p\right) \, dp d\ell ds. \end{split}$$
(NLb)

1.7.3. Wigner-Fourier transform of (NH). By introducing a phase-space representation of quantum density matrices through the Wigner transform, a connection to classical equations is established. In this context, the Hartree equation exhibits several similarities with the Vlasov equation. The Wigner transform of P, defined in Definition 1.8, satisfies

$$\begin{cases} \partial_t \widehat{W[P]}(t,k,\eta) = \widehat{W[P_L]}(t,k,\eta) + \widehat{W[P_{NL}]}(t,k,\eta), \\ \widehat{W[P_L]}(t,k,\eta) = -\frac{i\hbar^{-1}}{(2\pi)^d} \widehat{w}(k)\widehat{\rho}(t,k) \left(e^{-i\frac{\hbar}{2}k\cdot(\eta-kt)} - e^{i\frac{\hbar}{2}k\cdot(\eta-kt)}\right)\widehat{g}(\eta-kt), \\ \widehat{W[P_{NL}]}(t,k,\eta) = -\frac{i\hbar^{-1}}{(2\pi)^d} \int_{\ell} \widehat{w}(\ell)\widehat{\rho_Q}(t,\ell) \left(e^{-i\frac{\hbar}{2}\ell\cdot(\eta-kt)} - e^{i\frac{\hbar}{2}\ell\cdot(\eta-kt)}\right)\widehat{W[P]}(t,k-\ell,\eta-\ell t) d\ell, \end{cases}$$

$$(5)$$

where $\widehat{W[P_{\rm L}]}$ (resp. $\widehat{W[P_{\rm NL}]}$) is the Fourier transform of the Wigner transform of $P_{\rm L}$ (resp. $P_{\rm NL}$).

The following lemma provides a bound on the D^{α}_{η} -derivatives of $\widehat{W[P]}(k,\eta)$, establishing control over these derivatives in terms of $\boldsymbol{\xi}^{\alpha}$ -weighted norms of P. This result, derived using Lemma 1.10, is fundamental for verifying the bootstrap assumptions (B1) and (B3). If $\alpha \geq 2$, then additional terms of order $O(\hbar)$ or higher appear.

Lemma 1.13. Let α be a multi-index. The Wigner transform of $P_{\rm L}$ satisfies

$$\begin{split} \left| D^{\alpha}_{\eta} \widehat{W[P_{\mathrm{L}}]}(t,k,\eta) \right| &\lesssim |k| |\widehat{w}(k)| |\widehat{\rho_{Q}}(t,k)| |\eta - kt| \left| D^{\alpha}_{\eta} \widehat{g}(\eta - kt) \right| \\ &+ |k| \langle \hbar k \rangle^{|\alpha| - 1} |\widehat{w}(k)| |\widehat{\rho_{Q}}(t,k)| \sum_{|\beta| \leq |\alpha| - 1} \left| D^{\beta}_{\eta} \widehat{g}(\eta - kt) \right|. \end{split}$$

The Wigner transform of $P_{\rm NL}$ satisfies

$$\begin{split} \left| \widehat{D_{\eta}^{\alpha} W[P_{\mathrm{NL}}]}(t,k,\eta) \right| \lesssim \int_{\ell} |\ell| |\widehat{w}(\ell)| |\widehat{\rho_Q}(t,\ell)| |\eta - kt| \left| D_{\eta}^{\alpha} \widehat{W[P]}(t,k-\ell,\eta-\ell t) \right| d\ell \\ + \sum_{|\beta| \le |\alpha| - 1} \int_{\ell} |\ell| \langle \hbar \ell \rangle^{|\alpha| - 1} |\widehat{w}(\ell)| |\widehat{\rho_Q}(t,\ell)| \left| D_{\eta}^{\beta} \widehat{W[P]}(t,k-\ell,\eta-\ell t) \right| d\ell. \end{split}$$

Remark. Observe that $D^{\alpha}_{\eta}W[P_{\rm L}]$ and $D^{\alpha}_{\eta}W[P_{\rm NL}]$ are $O(\hbar^0)$ as $\hbar \to 0$.

Proof. Using the Leibniz rule,

$$D_{\eta}^{\alpha} \left[\left(e^{-i\frac{\hbar}{2}k \cdot (\eta - kt)} - e^{i\frac{\hbar}{2}k \cdot (\eta - kt)} \right) \widehat{g}(\eta - kt) \right] = \sum_{\beta \le \alpha} \binom{\alpha}{\beta} D_{\eta}^{\alpha - \beta} \left[\left(e^{-i\frac{\hbar}{2}k \cdot (\eta - kt)} - e^{i\frac{\hbar}{2}k \cdot (\eta - kt)} \right) \right] D_{\eta}^{\beta} \widehat{g}(\eta - kt).$$

For the term in the summation with $\beta = \alpha$, applying the Mean Value Theorem leads to the bound

$$\left|e^{-i\frac{\hbar}{2}k\cdot(\eta-kt)}-e^{i\frac{\hbar}{2}k\cdot(\eta-kt)}\right|\leq \hbar|k||\eta-kt|$$

For other terms $\beta < \alpha$ in the summation the bound

$$\left| e^{-i\frac{\hbar}{2}k \cdot (\eta - kt)} \pm e^{i\frac{\hbar}{2}k \cdot (\eta - kt)} \right| \le 2$$

is applied. This leads to the control.

$$\left| D_{\eta}^{\alpha} \left[\left(e^{-i\frac{\hbar}{2}k \cdot (\eta - kt)} - e^{i\frac{\hbar}{2}k \cdot (\eta - kt)} \right) \widehat{g}(\eta - kt) \right] \right| \lesssim \hbar |k| |\eta - kt| \left| D_{\eta}^{\alpha} \widehat{g}(\eta - kt) \right| + \sum_{\beta < \alpha} (\hbar |k|)^{|\alpha - \beta|} \left| D^{\beta} \widehat{g}(\eta - kt) \right|.$$

The result follows from the observation that $(\hbar|k|)^{|\alpha-\beta|} \leq \hbar|k|\langle\hbar k\rangle^{|\alpha|-1}$ for all $\beta < \alpha$. The inequality for $W[P_{\rm NL}]$ follows similarly.

1.7.4. Bootstrap assumptions. Fix positive constants $\sigma_4 > \sigma_3 > \sigma_2 > \sigma_1$ such that

$$\sigma_1 \ge d+7, \quad \sigma_2 - \sigma_1 > 3/2, \quad \sigma_3 - \sigma_2 > d/2, \quad \sigma_4 - \sigma_3 > (d+1)/2, \quad \sigma_4 < N_0 - (d+1)/2 \tag{6}$$

and constants $K_i \ge 1$ determined by the proof. Set $0 < \delta \ll 1/2$ such that $\sigma_4 - \sigma_3 > (d+1)/2 + \delta$. Let $I = [0, T^*]$ be the largest connected interval containing zero such that the following bootstrap controls hold:

$$\hbar^{-d} \sum_{|\alpha| \le M} \|\boldsymbol{\xi}^{\alpha} \langle \boldsymbol{\nabla}_x, \boldsymbol{\nabla}_{\boldsymbol{\xi}} \rangle^{\sigma_4} \langle t \boldsymbol{\nabla}_x, \boldsymbol{\nabla}_{\boldsymbol{\xi}} \rangle P(t) \|_{L^2_{x,y}}^2 \le 4K_1 \langle t \rangle^{2d-1} \varepsilon^2, \tag{B1}$$

$$\left\| |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_4} \widehat{\rho_Q}(t,k) \right\|_{L^2_{t,k}}^2 \le 4K_2 \varepsilon^2, \tag{B2}$$

$$\hbar^{-d} \sum_{|\alpha| \le M} \left\| \boldsymbol{\xi}^{\alpha} \langle \boldsymbol{\nabla}_{x}, \boldsymbol{\nabla}_{\xi} \rangle^{\sigma_{3}} | \boldsymbol{\nabla}_{x} |^{\delta} P(t) \right\|_{L^{2}_{x,y}}^{2} \le 4K_{3}\varepsilon^{2}, \tag{B3}$$

$$\left\| |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_2} \widehat{\rho_Q}(t, k) \right\|_{L^{\infty}_k L^2_t}^2 \le 4K_4 \varepsilon^2, \tag{B4}$$

$$\langle k,\eta\rangle^{2\sigma_1} \left| \int_p e^{-i\hbar\eta \cdot p} \widehat{P}\left(\frac{k}{2} + p, \frac{k}{2} - p\right) dp \right|^2 \le 4K_5\varepsilon^2.$$
 (B5)

Observe that (B1) and (B3) hold initially under the assumption $\|Q_{\text{in}}\|_{\mathcal{H}_M^{N_0}} \leq \varepsilon$. Moreover, at t = 0, (B5) can be rewritten in terms of the Wigner transform as

$$\langle k,\eta\rangle^{2\sigma_1} \left|\widehat{W[Q_{\rm in}]}(k,\eta)\right|^2 = \left|\overline{W[\langle \boldsymbol{\nabla}_x, \boldsymbol{\nabla}_\xi\rangle^{\sigma_1} Q_{\rm in}]}(k,\eta)\right|^2 \lesssim \|Q_{\rm in}\|_{\mathcal{H}_M^{\sigma_1}}^2 \le \varepsilon^2,$$

where Lemma 1.9 has been used. Bootstrap assumptions (B2) and (B4) are trivially satisfied if $I = \{0\}$. It is now shown that the norms (B2) and (B4) are natural, in the sense that, for the free Hartree density, they are controlled by the assumption on the initial data Q_{in} in Theorem 1.5.

Lemma 1.14. Fix $\sigma > 0$ and let $\rho_{Q_{\text{FH}}}$ be the density of the solution to the free Hartree equation (FH), defined in (3). Then, for $s_1 > (d+1)/2$ and $s_2 > 1/2$,

$$\begin{aligned} \left\| |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma} \widehat{\rho_{Q_{\mathrm{FH}}}}(t, k) \right\|_{L^{2}_{k}L^{2}_{t}} \lesssim_{s_{1}} \left\| Q_{\mathrm{in}} \right\|_{\mathcal{H}^{\sigma+s_{1}}_{M}} \\ \left\| |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma} \widehat{\rho_{Q_{\mathrm{FH}}}}(t, k) \right\|_{L^{\infty}_{k}L^{2}_{t}} \lesssim_{s_{2}} \left\| Q_{\mathrm{in}} \right\|_{\mathcal{H}^{\sigma+s_{2}}_{M}}. \end{aligned}$$

Proof. First recall that $\widehat{\rho_{Q_{\text{FH}}}}(t,k) = \widehat{W[Q_{\text{in}}]}(k,kt).$

$$\begin{split} \left\| |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma} \widehat{\rho_Q}(t,k) \right\|_{L^2_{t,k}} &= \left(\int_{t=0}^{T^*} \int_k |k| \langle k, kt \rangle^{2\sigma} \Big| \widehat{W[Q_{\text{in}}]}(k,kt) \Big|^2 \, dk dt \right)^{\frac{1}{2}} \\ &\leq \sup_{k,\eta} \left| \langle k, \eta \rangle^{\sigma+s_1} \widehat{W[Q_{\text{in}}]}(k,\eta) \Big| \left(\int_k \int_{t=0}^{\infty} \frac{|k|}{\langle k, kt \rangle^{2s_1}} \, dt dk \right)^{\frac{1}{2}}. \end{split}$$

Then, by making the change of variables $t \mapsto t/|k|$, and denoting $\overline{\varepsilon} = s_1 - \frac{d+1}{2}$,

$$\int_{k} \int_{t=0}^{\infty} \frac{|k|}{\langle k, kt \rangle^{s_{1}}} \, dt dk = \int_{k} \int_{t=0}^{\infty} \frac{1}{\langle k, t \rangle^{d+1+2\overline{\varepsilon}}} \, dt dk \lesssim \int_{k} \frac{1}{\langle k \rangle^{d+\overline{\varepsilon}}} \, dk \int_{t} \frac{1}{\langle t \rangle^{1+\overline{\varepsilon}}} \, dt \lesssim 1.$$

Furthermore, using the Sobolev embedding, for any $0 < \tilde{\delta} \leq 1/2$,

$$\begin{split} \sup_{k,\eta} \left| \langle k,\eta \rangle^{\sigma+s_1} \widehat{W[Q_{\mathrm{in}}]}(k,\eta) \right| \lesssim \left\| \langle \nabla_\eta \rangle^{\frac{d}{2} + \widetilde{\delta}} \langle \nabla_k \rangle^{\frac{d}{2} + \widetilde{\delta}} \left(\langle k,\eta \rangle^{\sigma+s_1} \widehat{W[Q_{\mathrm{in}}]}(k,\eta) \right) \right\|_{L^2_{k,\eta}} \\ &\sim \left\| W \left[\langle \boldsymbol{\xi} \rangle^{\frac{d}{2} + \widetilde{\delta}} \langle \boldsymbol{x} \rangle^{\frac{d}{2} + \widetilde{\delta}} \langle \boldsymbol{\nabla}_x, \boldsymbol{\nabla}_{\boldsymbol{\xi}} \rangle^{\sigma+s_1} Q_{\mathrm{in}} \right] \right\|_{L^2_{x,\boldsymbol{\xi}}} \\ &\lesssim \hbar^{-d/2} \sum_{|\alpha|, |\beta| \leq M} \left\| \boldsymbol{\xi}^{\alpha} \boldsymbol{x}^{\beta} \langle \boldsymbol{\nabla}_x, \boldsymbol{\nabla}_{\boldsymbol{\xi}} \rangle^{\sigma+s_1} Q_{\mathrm{in}} \right\|_{L^2_{x,y}}. \end{split}$$
stimate is proved similarly.

The second estimate is proved similarly.

The remainder of this paper focuses on improving the bootstrap assumptions (B1)-(B5).

Proposition 1.15 (Bootstrap). Let the assumptions on w, g and Q_{in} be as in Theorem 1.5. There exist $\varepsilon_0 = \varepsilon_0(N_0, w, g, \kappa, d, \delta_0, \sigma_i)$ and $K_i = K_i(N_0, w, g, \kappa, d, \delta_0, \sigma_i)$ such that if the bootstrap controls (B1)-(B5) hold on some time interval $[0, T^*]$ and $\varepsilon < \varepsilon_0$, then

$$\hbar^{-d} \sum_{|\alpha| \le M} \| \boldsymbol{\xi}^{\alpha} \langle \boldsymbol{\nabla}_x, \boldsymbol{\nabla}_{\boldsymbol{\xi}} \rangle^{\sigma_4} \langle t \boldsymbol{\nabla}_x, \boldsymbol{\nabla}_{\boldsymbol{\xi}} \rangle P(t) \|_{L^2_{x,y}}^2 \le 2K_1 \langle t \rangle^{2d-1} \varepsilon^2, \tag{I1}$$

$$\left\| |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_4} \widehat{\rho_Q}(t,k) \right\|_{L^2_{t,k}}^2 \le 2K_2 \varepsilon^2, \tag{I2}$$

$$\hbar^{-d} \sum_{|\alpha| \le M} \left\| \boldsymbol{\xi}^{\alpha} \langle \boldsymbol{\nabla}_{x}, \boldsymbol{\nabla}_{\xi} \rangle^{\sigma_{3}} |\boldsymbol{\nabla}_{x}|^{\delta} P(t) \right\|_{L^{2}_{x,y}}^{2} \le 2K_{3}\varepsilon^{2}, \tag{I3}$$

$$\left\| |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_2} \widehat{\rho_Q}(t,k) \right\|_{L_k^\infty L_t^2}^2 \le 2K_4 \varepsilon^2, \tag{I4}$$

$$\langle k,\eta\rangle^{2\sigma_1} \left| \int_p e^{-i\hbar\eta \cdot p} \widehat{P}\left(\frac{k}{2} + p, \frac{k}{2} - p\right) dp \right|^2 \le 2K_5\varepsilon^2$$
 (I5)

for all $t \in [0, T^*]$. It follows that $T^* = \infty$.

The improvement of each bootstrap assumption in Proposition 1.15 is established in Sections 3.2, 3.3, 4.1, 4.2, and 4.3, with the conclusion presented in Section 1.7.6. Figure 2 illustrates which bootstrap assumptions are required for each improvement. For example, (B1) points to (B2) because bootstrap assumption (B1) is necessary to establish bootstrap improvement (I2). In Lemmas 1.16 and 1.17, it is shown that the decay of the density (T1) in Theorem 1.5 relies solely on bootstrap improvement (I5), and the scattering statement (T2) in Theorem 1.5 is established using only bootstrap assumptions (I3) and (I5).



FIGURE 2. Bootstrap implications.

The most challenging bootstrap improvement to establish is (I2), as it permits no regularity loss or time growth. Achieving this requires precise control over the nonlinear interaction in (NLb), which includes a bilinear term in the perturbation. The density mode $\widehat{\rho_Q}(t,k)$ can be influenced by earlier modes $\widehat{\rho_Q}(s,\ell)$, $0 \leq s \leq t$, and in some cases, this interaction may cascade, producing *resonances* at arbitrarily large times. Such growth is driven by an unmixing mechanism caused by reverse-in-time phase mixing, as discussed in [4]. For the density of the Hartree equation in \mathbb{R}^d with $d \geq 3$ and potential $|\widehat{w}(\ell)| \leq \langle \ell \rangle^{-2}$, resonances are controlled by a dispersive mechanism. This mechanism relies on the set of modes capable of driving the mode $\widehat{\rho_Q}(t,k)$ shrinking over time; these are modes $\widehat{\rho_Q}(s,\ell)$ satisfying $|k \cdot \ell| \leq \langle s \rangle^{-\zeta}$ for some $\zeta > 0$, meaning they become increasingly "almost collinear" with k over time. The kernel capturing the maximal impact of these resonances matches that used in the Vlasov–Poisson equation, as detailed in [1], and is controlled similarly in Section 3.2.2.

1.7.5. *Deduction of Main Theorem*. The bootstrap assumption (B5) implies the mode-by-mode decay of the density (T1).

Lemma 1.16. Proposition 1.15 implies the damping estimate (T1) in Theorem 1.5.

Proof. Recall the definition of the density is $\widehat{\rho_Q}(t,k) := \frac{1}{(2\pi)^d} \int_p \widehat{Q}(\frac{k}{2}+p,\frac{k}{2}-p) dp$. In terms of P, the integral kernel of Q is $\widehat{Q}(t,\frac{k}{2}+p,\frac{k}{2}-p) = e^{-i\hbar p \cdot kt} \widehat{P}(t,\frac{k}{2}+p,\frac{k}{2}-p)$. The bootstrap control (B5) with $\eta = kt$ yields precisely

$$\langle k, kt \rangle^{\sigma_1} |\widehat{\rho_Q}(t, k)| = \langle k, kt \rangle^{\sigma_1} \left| \frac{1}{(2\pi)^d} \int_p e^{-i\hbar p \cdot kt} \widehat{P}(t, \frac{k}{2} + p, \frac{k}{2} - p) \, dp \right| \lesssim \sqrt{K_5} \varepsilon.$$

Furthermore, the bootstrap controls (B3) and (B5) are sufficient to deduce the scattering statement (T2).

Lemma 1.17. Proposition 1.15 implies the scattering statement (T2) in Theorem 1.5.

Proof. To prove decay of the linear term, the analysis is carried out on the Fourier side of the Wigner transform. Let $|\alpha| \leq M$. Then, by Lemma 1.11,

$$\|\langle \boldsymbol{\nabla}_{x}, \boldsymbol{\nabla}_{\xi} \rangle^{\sigma_{0}}(\boldsymbol{\xi}^{\alpha} P_{\mathrm{L}}(t))\|_{L^{2}_{x,y}}^{2} = (2\pi)^{d} \hbar^{d} \|\langle \nabla_{x}, \nabla_{\xi} \rangle^{\sigma_{0}}(\xi^{\alpha} W[P_{\mathrm{L}}](t))\|_{L^{2}_{x,\xi}}^{2} = (2\pi)^{2d} \hbar^{d} \|\langle k, \eta \rangle^{\sigma_{0}} D_{\eta}^{\alpha}(\widehat{W[P_{\mathrm{L}}]}(t))\|_{L^{2}_{k,\eta}}^{2}.$$

Using Lemma 1.13 and $\langle k, \eta \rangle^{\sigma_0} \lesssim \langle k, kt \rangle^{\sigma_0} \langle \eta - kt \rangle^{\sigma_0}$,

$$\begin{split} \left\| \langle k, \eta \rangle^{\sigma_0} D_{\eta}^{\alpha}(\widehat{W[P_{\mathrm{L}}]}(t)) \right\|_{L^2_{k,\eta}}^2 \lesssim \int_{k,\eta} |\widehat{w}(k)|^2 |k|^2 \langle k, kt \rangle^{2\sigma_0} |\widehat{\rho_Q}(t,k)|^2 \langle \eta - kt \rangle^{2\sigma_0} |\eta - kt|^2 \left| D_{\eta}^{\alpha} \widehat{g}(\eta - kt) \right|^2 dk d\eta \\ &+ \langle \hbar \rangle^2 \sum_{|\beta| \le |\alpha| - 1} \int_{k,\eta} \langle k \rangle^2 |\widehat{w}(k)| |k|^2 \langle k, kt \rangle^{2\sigma_0} |\widehat{\rho_Q}(t,k)|^2 \langle \eta - kt \rangle^{2\sigma_0} \left| D_{\eta}^{\beta} \widehat{g}(\eta - kt) \right| dk d\eta \\ &\lesssim \|g\|_{H^{\sigma_1}_M}^2 \int_k |k|^2 \langle k, kt \rangle^{2\sigma_0} |\widehat{\rho_Q}(t,k)|^2 dk \\ &\lesssim K_5 \varepsilon^2 \|g\|_{H^{\sigma_0+1}_M}^2 \int_k \frac{|k|^2}{\langle k, kt \rangle^{2\sigma_1 - 2\sigma_0}} dk \\ &\lesssim \frac{K_5 \varepsilon^2}{\langle t \rangle^{d+2}}, \end{split}$$

by the bootstrap assumption (B5) if $2\sigma_1 - 2\sigma_0 > d + 2$. Therefore,

$$\|\langle \boldsymbol{\nabla}_x, \boldsymbol{\nabla}_{\boldsymbol{\xi}} \rangle^{\sigma_0}(\boldsymbol{\xi}^{\alpha} P_{\mathrm{L}}(t))\|_{L^2_{x,y}} \lesssim \hbar^{d/2} \frac{\sqrt{K_5}\varepsilon}{\langle t \rangle^{d/2+1}}.$$

For the nonlinear term, again using Lemma 1.13 and $\langle k, \eta \rangle^{\sigma_0} \lesssim \langle \ell, \ell t \rangle^{\sigma_0} \langle k - \ell, \eta - \ell t \rangle^{\sigma_0}$ and $|\eta - kt| \lesssim \langle t \rangle \langle k - \ell, \eta - \ell t \rangle$,

$$\begin{split} \left\| \langle k, \eta \rangle^{\sigma_{0}} D_{\eta}^{\alpha}(\widehat{W[P_{\mathrm{NL}}]}(t)) \right\|_{L^{2}_{k,\eta}}^{2} \\ &\lesssim \int_{k,\eta} \left(\langle k, \eta \rangle^{\sigma_{0}} \int_{\ell} |\ell| |\widehat{w}(\ell)| |\widehat{\rho_{Q}}(t,\ell)| |\eta - kt| \left| D_{\eta}^{\alpha} \widehat{W[P]}(t,k-\ell,\eta-\ell t) \right| d\ell \right)^{2} dk d\eta \\ &+ \sum_{|\beta| \leq |\alpha| - 1} \int_{k,\eta} \left(\langle k, \eta \rangle^{\sigma_{0}} \int_{\ell} \langle \hbar \ell \rangle^{|\alpha| - 1} |\ell| |\widehat{w}(\ell)| |\widehat{\rho_{Q}}(t,\ell)| \left| D_{\eta}^{\beta} \widehat{W[P]}(t,k-\ell,\eta-\ell t) \right| d\ell \right)^{2} dk d\eta \\ &\lesssim \langle t \rangle^{2} \sum_{|\beta| \leq M} \int_{k,\eta} \left(\int_{\ell} |\ell| \langle \ell, \ell t \rangle^{\sigma_{0}} |\widehat{\rho_{Q}}(t,\ell)| \langle k-\ell,\eta-\ell t \rangle^{\sigma_{0}+1} \left| D_{\eta}^{\beta} \widehat{W[P]}(t,k-\ell,\eta-\ell t) \right| d\ell \right)^{2} dk d\eta \\ &\lesssim K_{5} \varepsilon^{2} \langle t \rangle^{2} \sum_{|\beta| \leq M} \int_{k,\eta} \left(\int_{\ell} \frac{|\ell|}{\langle \ell, \ell t \rangle^{\sigma_{1} - \sigma_{0}}} \left| \overline{W[\langle \nabla_{x}, \nabla_{\xi} \rangle^{\sigma_{0} + 1}(\xi^{\beta} P)]}(t,k-\ell,\eta-\ell t) \right| d\ell \right)^{2} dk d\eta \end{split}$$

if $0 < \hbar \leq 1$ and $|\widehat{w}(\ell)| \leq \langle \ell \rangle^{-M-1/2}$, by the bootstrap estimate (B5). Then, using Cauchy–Schwarz in ℓ , for $0 < \delta \ll 1$,

$$\begin{split} \left\| \langle k, \eta \rangle^{\sigma_0} D_{\eta}^{\alpha}(\widehat{W[P_{\mathrm{NL}}]}(t)) \right\|_{L^2_{k,\eta}}^2 \\ \lesssim K_5 \varepsilon^2 \langle t \rangle^2 \sum_{|\beta| \le M} \int_{k,\eta} \left[\int_{\ell} \frac{1}{|k-\ell|^{2\delta}} \frac{|\ell|}{\langle \ell, \ell t \rangle^{\sigma_1 - \sigma_0}} d\ell \right] \\ \times \int_{\ell} \frac{|\ell|}{\langle \ell, \ell t \rangle^{\sigma_1 - \sigma_0}} |k-\ell|^{2\delta} \Big| \overline{W[\langle \nabla_x, \nabla_{\xi} \rangle^{\sigma_0 + 1}(\xi^{\beta} P)]}(t, k-\ell, \eta - \ell t) \Big|^2 d\ell dk d\eta \\ \lesssim K_5 \varepsilon^2 \langle t \rangle^2 \sup_{k'} \left[\int_{\ell} \frac{1}{|k'-\ell|^{2\delta}} \frac{|\ell|}{\langle \ell, \ell t \rangle^{\sigma_1 - \sigma_0}} d\ell \right] \left(\int_{\ell} \frac{|\ell|}{\langle \ell, \ell t \rangle^{\sigma_1 - \sigma_0}} d\ell \right) \sum_{|\beta| \le M} \left\| W[|\nabla_x|^{\delta} \langle \nabla_x, \nabla_{\xi} \rangle^{\sigma_0 + 1}(\xi^{\beta} P)] \right\|_{L^2_{x,\xi}}^2 \\ = K_5 \varepsilon^2 \langle t \rangle^2 \left[\int_{\ell} \frac{|\ell|^{1 - 2\delta}}{\langle \ell, \ell t \rangle^{\sigma_1 - \sigma_0}} d\ell \right] \left(\int_{\ell} \frac{|\ell|}{\langle \ell, \ell t \rangle^{\sigma_1 - \sigma_0}} d\ell \right) \frac{1}{\hbar^d} \sum_{|\beta| \le M} \left\| |\nabla_x|^{\delta} \langle \nabla_x, \nabla_{\xi} \rangle^{\sigma_0 + 1}(\xi^{\beta} P) \right\|_{L^2_{x,y}}^2 \\ \lesssim K_5 K_3 \varepsilon^4 \langle t \rangle^2 \frac{1}{\langle t \rangle^{d+1 - 2\delta}} \frac{1}{\langle t \rangle^{d+1}}, \end{split}$$

if $\sigma_1 - \sigma_0 > d + 1$ and $\sigma_0 + 1 \le \sigma_3$. Thus for the nonlinear term, using Lemma 1.11,

$$\left\| \langle \boldsymbol{\nabla}_x, \boldsymbol{\nabla}_{\boldsymbol{\xi}} \rangle^{\sigma_0}(\boldsymbol{\xi}^{\alpha} P_{\mathrm{NL}}(t)) \right\|_{L^2_{x,y}} = (2\pi)^d \hbar^{d/2} \left\| \langle k, \eta \rangle^{\sigma_0} D_{\eta}^{\alpha}(\widehat{W[P_{\mathrm{NL}}]}(t)) \right\|_{L^2_{k,\eta}} \lesssim \hbar^{d/2} \frac{\sqrt{K_5 K_3 \varepsilon^2}}{\langle t \rangle^{d-2\delta}}.$$

Therefore, both of the time integrals in (NLa) are absolutely convergent in $H_M^{\sigma_0}$. Hence, define

$$\widehat{Q_{\infty}}(k,p) = \widehat{Q_{\text{in}}}(k,p) - \int_0^\infty \widehat{P_{\text{L}}}(s,k,p) \, ds - \int_0^\infty \widehat{P_{\text{NL}}}(s,k,p) \, ds.$$

Inequality (T2) then follows from

$$\begin{split} \|P(t) - Q_{\infty}\|_{\mathcal{H}_{M}^{\sigma_{0}}} &\lesssim \int_{t}^{\infty} \|\langle \boldsymbol{\nabla}_{x}, \boldsymbol{\nabla}_{\xi} \rangle^{\sigma_{0}} (\boldsymbol{\xi}^{\alpha} P_{\mathrm{L}}(s))\|_{\mathcal{L}^{2}_{x,y}} \, ds + \int_{t}^{\infty} \|\langle \boldsymbol{\nabla}_{x}, \boldsymbol{\nabla}_{\xi} \rangle^{\sigma_{0}} (\boldsymbol{\xi}^{\alpha} P_{\mathrm{NL}}(s))\|_{\mathcal{L}^{2}_{x,y}} \, ds \\ &\lesssim \sqrt{K_{5}} \varepsilon \int_{t}^{\infty} \frac{1}{\langle s \rangle^{d/2+1}} \, ds + \sqrt{K_{5}K_{3}} \varepsilon^{2} \int_{t}^{\infty} \frac{1}{\langle s \rangle^{d-2\delta}} \, ds \\ &\lesssim \frac{\varepsilon}{\langle t \rangle^{d/2}}, \end{split}$$

for $d \geq 3$.

1.7.6. Conclusion of bootstrap. The bootstrap improvements in Proposition 1.15, and the deduction that $T^* = \infty$, are concluded as follows.

Proof of Proposition 1.15. Fix dimension $d \geq 3$, constants σ_0 , σ_1 , N_0 , potential w, steady state g and initial perturbation Q_{in} as in Theorem 1.5 and σ_2 , σ_3 , σ_4 to satisfy (6). Section 3 demonstrates that the constants K_2 and K_4 appearing in the bootstrap assumptions (B2) and (B4) can be chosen based solely on the setup. In Section 4, it is shown that the constants K_1 , K_3 , and K_5 , which appear in bootstrap assumptions (B1), (B3), and (B5), can also be selected based only on the setup and on K_2 and K_4 . If ε_0 satisfies

$$0 < \varepsilon_0 \le \min\left\{\frac{1}{\sqrt{K_3K_5 + K_2K_5}}, \frac{1}{\sqrt{K_4K_5} + \sqrt{K_3K_5}} + \frac{1}{\sqrt{K_2K_3} + \sqrt{K_1K_5} + \sqrt{K_3K_5}}\right\},\$$

then, by Sections 3.2, 3.3, 4.1, 4.2 and 4.3, the bootstrap improvements (I1)-(I5) hold.

The proof is concluded by showing that $T^* = \infty$ using a standard bootstrap argument. Fix ε sufficiently small to ensure that the improvement stated in Proposition 1.15 holds. Define $\mathcal{A} \subseteq [0, \infty)$ as the set consisting of those values T^* for which the bootstrap assumptions (B1)-(B5) hold. First, note that \mathcal{A} is non-empty, as $0 \in \mathcal{A}$. The set \mathcal{A} is also connected and closed by the definition of \mathcal{A} . Finally, \mathcal{A} is open: if $T \in \mathcal{A}$, the improvement of the bootstrap assumptions (as per Proposition 1.15) indicates that there exists $\tilde{\varepsilon} > 0$ such that $T + \tilde{\varepsilon} \in \mathcal{A}$. The only set that satisfies these properties is $\mathcal{A} = [0, \infty)$, which implies that $T^* = \infty$. \Box

2. LINEARISED PHASE MIXING

In this section, the *linearised Hartree equation* is considered, which is found by removing the quadraticin-Q term $[w * \rho_Q, Q]$ from (NH) which is expected to be small:

$$\begin{cases} i\hbar\partial_t Q_{\rm L} = \left[-\frac{\hbar^2}{2} \Delta, Q_{\rm L} \right] + \left[w * \rho_{Q_{\rm L}}, \gamma_g \right], \\ Q_{\rm L}(0) = Q_{\rm in}. \end{cases}$$
(LH)

2.1. Fourier-Laplace transform. Let $Q_{\rm L} = Q_{\rm L}(t, x, y)$ denote the time-dependent integral kernel of the perturbation $Q_{\rm L}$. The linearised equation (LH) can be expressed as

$$i\hbar\partial_t Q_{\mathrm{L}}(t,x,y) = -\frac{\hbar^2}{2} \left(\Delta_x - \Delta_y\right) Q_{\mathrm{L}}(t,x,y) + \frac{1}{(2\pi)^d} \left(w * \rho_{Q_{\mathrm{L}}}(t,x) - w * \rho_{Q_{\mathrm{L}}}(t,y)\right) \widehat{g}\left(\frac{y-x}{\hbar}\right).$$

Applying Duhamel's formula yields an implicit formula for the solution of the linearised Hartree equation:

$$Q_{\mathrm{L}}(t,x,y) = e^{i\frac{\omega}{2}(\Delta_x - \Delta_y)t}Q_{\mathrm{in}}(x,y) - \frac{i}{(2\pi)^d}\frac{1}{\hbar}\int_0^t e^{i\frac{\hbar}{2}(\Delta_x - \Delta_y)(t-s)} \left[w * \rho_{Q_{\mathrm{L}}}(s,x) - w * \rho_{Q_{\mathrm{L}}}(s,y)\right]\widehat{g}\left(\frac{y-x}{\hbar}\right) ds.$$

$$\tag{7}$$

The Fourier transform of (7) is

· ħ / •

$$\widehat{Q_{\rm L}}(t,k,p) = e^{-i\frac{\hbar}{2}(|k|^2 - |p|^2)t} \widehat{Q_{\rm in}}(k,p) - i\hbar^{d-1}\widehat{w}(k+p) \left[g(-\hbar p) - g(\hbar k)\right] \int_0^t e^{-i\frac{\hbar}{2}(|k|^2 - |p|^2)(t-s)} \widehat{\rho_{Q_{\rm L}}}(s,k+p) \, ds.$$
(8)

and so using the formula for the Fourier transform of the density $\widehat{\rho_{Q_{\rm L}}}(k) = \frac{1}{(2\pi)^d} \int_p \widehat{Q_{\rm L}}(\frac{k}{2} + p, \frac{k}{2} - p) \, dp$, it follows that

$$\widehat{\rho_{Q_{\mathrm{L}}}}(t,k) = \frac{1}{(2\pi)^d} \int_{\mathbb{R}_p^d} e^{-i\hbar kt \cdot p} \widehat{Q_{\mathrm{in}}} \left(\frac{k}{2} + p, \frac{k}{2} - p\right) dp - \frac{i\hbar^{d-1}}{(2\pi)^d} \widehat{w}(k) \int_{\mathbb{R}_p^d} \int_0^t e^{-i\hbar(t-s)k \cdot p} \left[g\left(\hbar(p-\frac{k}{2})\right) - g\left(\hbar(p+\frac{k}{2})\right)\right] \widehat{\rho_{Q_{\mathrm{L}}}}(s,k) dp ds.$$

This is a Volterra equation:

$$\widehat{\rho_{Q_{\rm L}}}(t,k) = \widehat{\rho_{Q_{\rm FH}}}(t,k) - \widehat{\mathscr{L}} *_t \widehat{\rho_{Q_{\rm L}}}(t,k), \tag{9}$$

where $\rho_{Q_{\rm FH}}(t,k)$ is the density of the free Hartree equation (FH) and

$$\begin{aligned} \widehat{\mathscr{L}}(t,k) &:= \frac{1}{(2\pi)^d} i\hbar^{d-1} \widehat{w}(k) \int_{\mathbb{R}_p^d} e^{-i\hbar kt \cdot p} \left[g\left(\hbar(p-\frac{k}{2})\right) - g\left(\hbar(p+\frac{k}{2})\right) \right] dp \\ &= \frac{1}{(2\pi)^d} i\hbar^{d-1} \widehat{w}(k) \mathcal{F}_p \left[g\left(\hbar(p-\frac{k}{2})\right) - g\left(\hbar(p+\frac{k}{2})\right) \right] (\hbar kt) \\ &= \frac{1}{(2\pi)^d} \frac{i}{\hbar} \widehat{w}(k) \left(e^{-i\frac{\hbar}{2}|k|^2 t} - e^{i\frac{\hbar}{2}|k|^2 t} \right) \widehat{g} (kt) \\ &= \frac{1}{(2\pi)^d} \frac{2}{\hbar} \widehat{w}(k) \sin\left(\frac{1}{2}\hbar t|k|^2\right) \widehat{g} (kt) . \end{aligned}$$
(10)

Note also that $w, g \in L^1(\mathbb{R}^d)$ implies that the zero mode $\widehat{\mathscr{L}}(t, k = 0) = 0$ for all $t \ge 0$.

In order to find an explicit equation for $\widehat{\rho_{Q_L}}$, the Laplace transform in time of the Volterra equation is taken. Denoting $\widetilde{f}(\lambda, k) := \int_0^\infty e^{-\lambda t} \widehat{f}(t, k) dt$, the Laplace transform of (9) is

$$\left(1 + \widetilde{\mathscr{L}}(\lambda, k)\right)\widetilde{\rho_{Q_{\rm L}}}(\lambda, k) = \widetilde{\rho_{Q_{\rm FH}}}(\lambda, k),\tag{11}$$

where

$$\widetilde{\mathscr{L}}(\lambda,k) = \frac{1}{(2\pi)^d} \frac{2}{\hbar} \widehat{w}(k) \int_0^\infty e^{-\lambda t} \sin(\hbar t |k|^2) \widehat{g}(kt) dt$$

$$= \frac{1}{(2\pi)^d} \frac{2}{\hbar |k|} \widehat{w}(k) \int_0^\infty e^{-\lambda \frac{s}{|k|}} \sin(\hbar |k|s) \widehat{g}(\frac{k}{|k|}s) ds$$
(12)

if $k \neq 0$. For $\operatorname{Re} \lambda > 0$, applying $\int_0^\infty e^{-\lambda t} e^{zt} dt = \frac{1}{\lambda - z}$ if $\operatorname{Re} \lambda > \operatorname{Re} z$ to (10) also yields the representation

$$\widetilde{\mathscr{L}}(\lambda,k) = \frac{\widehat{w}(k)}{(2\pi)^d} \frac{1}{\hbar|k|} \int_{p_1 \in \mathbb{R}} g_k(p_1) \left[\frac{1}{p_1 + \frac{i\lambda}{|k|} + \frac{\hbar|k|}{2}} - \frac{1}{p_1 + \frac{i\lambda}{|k|} - \frac{\hbar|k|}{2}} \right] dp_1,$$

where $g_k \colon \mathbb{R} \to \mathbb{R}$ denotes the marginal of g along the vector $k \in \mathbb{R}^d$, defined by

$$g_k(p_1) = \int_{\{k\}^{\perp}} g\left(\frac{k}{|k|}p_1 + |\widetilde{p}|^2\right) d\widetilde{p}.$$

On the other hand, if $\operatorname{Re} \lambda = 0$, $\lambda = i\tau$, then the Plemelj formula implies

$$\widetilde{\mathscr{L}}(i\tau,k) = \frac{\widehat{w}(k)}{(2\pi)^d} \frac{1}{\hbar|k|} \left(i\pi \left[g_k \left(\frac{\tau}{|k|} + \frac{\hbar|k|}{2} \right) - g_k \left(\frac{\tau}{|k|} - \frac{\hbar|k|}{2} \right) \right] + \text{p.v.} \int_{p_1 \in \mathbb{R}} \left[\frac{g_k(p_1)}{p_1 - \frac{\tau}{|k|} + \frac{\hbar|k|}{2}} - \frac{g_k(p_1)}{p_1 - \frac{\tau}{|k|} - \frac{\hbar|k|}{2}} \right] dp_1 \right).$$

$$(13)$$

2.2. **Penrose condition.** In this section, sufficient conditions on w and g are established to ensure that the Penrose condition (P) is satisfied. For all conditions, the potential is required to be $w \in L^1(\mathbb{R}^d)$, which excludes the Coulomb potential. In fact, if w is the Coulomb potential, then the Penrose condition does not hold for any steady state g (see [19]).

Firstly, a result is presented that establishes the flexibility to order moments and derivatives in the H_M^{σ} in whichever manner is most convenient.

Lemma 2.1 (Ordering of moments). Let $\sigma \geq 0$, let M be a non-negative integer, and let $g \in H^{\sigma}_{M}(\mathbb{R}^{d})$. Then

$$\|g\|_{H^{\sigma}_{M}} := \sum_{|\alpha| \le M} \|\langle \nabla_{k} \rangle^{\sigma} (k^{\alpha}g)\|_{L^{2}} \sim \sum_{|\alpha| \le M} \|k^{\alpha} \langle \nabla_{k} \rangle^{\sigma}g\|_{L^{2}}$$

The Sobolev L^2 trace lemma is also recalled.

Lemma 2.2 (L^2 trace [1]). Let $g \in H^s(\mathbb{R}^d)$ with $s > \frac{d-1}{2}$ and $C \subset \mathbb{R}^d$ be an arbitrary straight line. Then, there holds

$$||g||_{L^2(C)} \lesssim_{s,d} ||g||_{H^s(\mathbb{R}^d)}$$

Using these results, the following can be deduced:

Lemma 2.3. If $n \ge 0$ and α is a multi-index, then

$$\int_0^\infty \langle s \rangle^n \left| \partial_k^\alpha \widehat{g}\left(\frac{k}{|k|}s\right) \right| ds \lesssim_{d,\delta_0,n,|\alpha|} \|g\|_{H^{n+\frac{1}{2}+\delta_0}_{|\alpha|+\lceil d/2\rceil}}$$

Proof. Applying Cauchy–Schwarz followed by Lemma 2.2,

$$\begin{split} \int_0^\infty \langle s \rangle^n \Big| \partial_k^\alpha \widehat{g} \left(\frac{k}{|k|} s \right) \Big| \, ds \lesssim_{\delta_0} \left(\int_0^\infty \langle s \rangle^{2n+1+2\delta_0} \Big| \partial_k^\alpha \widehat{g} \left(\frac{k}{|k|} s \right) \Big|^2 \, ds \right)^{\frac{1}{2}} \\ \lesssim_{\delta_0} \left(\int_{\mathbb{R}_k^d} \Big| \langle \nabla_k \rangle^{\frac{d-1}{2}+\delta_0} \left(\langle k \rangle^{n+\frac{1}{2}+\delta_0} \partial_k^\alpha \widehat{g}(k) \right) \Big|^2 \, dk \right)^{\frac{1}{2}}. \end{split}$$

Then, by Plancherel and Lemma 2.1, the following holds:

$$\int_{0}^{\infty} \langle s \rangle^{n} \left| \partial_{k}^{\alpha} \widehat{g} \left(\frac{k}{|k|} s \right) \right| ds \lesssim_{d} \left\| \langle x \rangle^{\frac{d-1}{2} + \delta_{0}} \langle \nabla_{x} \rangle^{n + \frac{1}{2} + \delta_{0}} \left(x^{\alpha} g(x) \right) \right\|_{L^{2}}$$

$$\lesssim \sum_{|\beta| \leq \lceil d/2 \rceil} \left\| x^{\beta} \langle \nabla_{x} \rangle^{n + \frac{1}{2} + \delta_{0}} \left(x^{\alpha} g(x) \right) \right\|_{L^{2}}$$

$$\lesssim_{n} \sum_{|\beta| \leq \lceil d/2 \rceil} \left\| \langle \nabla_{x} \rangle^{n + \frac{1}{2} + \delta_{0}} \left(x^{\beta + \alpha} g(x) \right) \right\|_{L^{2}}$$

$$\lesssim_{\alpha} \left\| g \right\|_{H^{n + \frac{1}{2} + \delta_{0}}_{|\alpha| + \lceil d/2 \rceil}}.$$

Using Lemma 2.3, boundedness and decay properties of $\widetilde{\mathscr{L}}$ are established:

Lemma 2.4. Suppose $w \in L^1(\mathbb{R}^d)$ and $g \in H^{\frac{3}{2}+\delta_0}_{\lceil d/2 \rceil}(\mathbb{R}^d)$.

- (a) For fixed $\operatorname{Re} \lambda \geq 0$, $\widetilde{\mathscr{L}}(\lambda, k) \to 0$ as $|k| \to \infty$ at a rate independent of λ . (b) There exists $C_1 = C_1(d, \delta_0)$ such that, for all $\operatorname{Re} \lambda \geq 0$ and $k \in \mathbb{R}^d$,

$$\left|\widetilde{\mathscr{L}}(\lambda,k)\right| \leq C_1 \|w\|_{L^1} \|g\|_{H^{\frac{3}{2}+\delta_0}_{\lceil d/2\rceil}}$$

(c) If additionally $g \in H^{\frac{3}{2}+\delta_0}_{1+\lceil d/2 \rceil}(\mathbb{R}^d)$, then, for all non-zero $\operatorname{Re} \lambda \geq 0$ and $k \in \mathbb{R}^d$,

$$\widetilde{\mathscr{L}}(\lambda,k)\Big|\lesssim \frac{|k|}{|\lambda|}\|w\|_{L^1}\|g\|_{H^{\frac{3}{2}+\delta_0}_{1+\lceil d/2\rceil}}$$

Proof. Recall that $\widetilde{\mathscr{L}}(\lambda, k = 0) = 0$. From the formula (12) for $\widetilde{\mathscr{L}}(\lambda, k)$ and Lemma 2.3, if $k \neq 0$ and $\operatorname{Re}\lambda \geq 0$,

$$\begin{split} \left| \widetilde{\mathscr{L}}(\lambda,k) \right| \lesssim_{d} |\widehat{w}(k)| \frac{1}{\hbar |k|} \int_{0}^{\infty} \left| \sin(\frac{\hbar}{2} |k|s) \widehat{g}(\frac{k}{|k|}s) \right| ds \\ \lesssim |\widehat{w}(k)| \int_{0}^{\infty} s \left| \widehat{g}(\frac{k}{|k|}s) \right| ds \\ \lesssim_{d,\delta_{0}} |\widehat{w}(k)| \|g\|_{H^{\frac{3}{2}+\delta_{0}}_{\lceil d/2 \rceil}}. \end{split}$$

If $w \in L^1(\mathbb{R}^d)$, then $\widehat{w}(k) \to 0$ as $|k| \to \infty$, which proves (a). Moreover $|\widehat{w}(k)| \le ||w||_{L^1}$ for all $k \in \mathbb{R}^d$ which proves (b).

For (c), if $q \in L^1(\mathbb{R}^d)$, (12) can be integrated by parts with no boundary data to obtain

$$\begin{aligned} \widetilde{\mathscr{L}}(\lambda,k) &= \frac{2}{(2\pi)^d} \widehat{w}(k) \frac{1}{\hbar|k|} \left(-\frac{|k|}{\lambda} \right) \int_0^\infty \partial_s \left[e^{-\lambda \frac{s}{|k|}} \right] \sin(\hbar|k|s) \widehat{g}(\frac{k}{|k|}s) \, ds \\ &= \frac{2}{(2\pi)^d} \widehat{w}(k) \frac{1}{\hbar\lambda} \int_0^\infty e^{-\lambda \frac{s}{|k|}} \left[\frac{\hbar}{2} |k| \cos(\frac{\hbar}{2}|k|s) \widehat{g}(\frac{k}{|k|}s) + \sin(\frac{\hbar}{2}|k|s) \frac{k}{|k|} \cdot \nabla_k \widehat{g}(\frac{k}{|k|}s) \right] \, ds \end{aligned}$$

and, by Lemma 2.3,

$$\begin{split} \left| \widetilde{\mathscr{L}}(\lambda,k) \right| \lesssim \frac{|k|}{|\lambda|} \|w\|_{L^1} \int_0^\infty \left[\left| \widehat{g}(\frac{k}{|k|}s) \right| + s \left| \nabla_k \widehat{g}(\frac{k}{|k|}s) \right| \right] \, ds \\ \lesssim_{d,\delta_0} \frac{|k|}{|\lambda|} \|w\|_{L^1} \|g\|_{H^{\frac{3}{2}+\delta_0}_{1+\lceil d/2 \rceil}}. \end{split}$$

Next, using the argument principle, a condition on the behaviour of $\widetilde{\mathscr{L}}$ along the imaginary axis is shown to suffice for the Penrose condition.

Lemma 2.5. If $w \in L^1(\mathbb{R}^d)$, $g \in H^{\frac{3}{2}+\delta_0}_{1+\lceil d/2 \rceil}(\mathbb{R}^d)$ and the following implication holds:

$$\forall \tau \in \mathbb{R}, k \in \mathbb{R}^d, \quad \operatorname{Im} \widehat{\mathscr{L}}(i\tau, k) = 0 \implies \operatorname{Re} \widehat{\mathscr{L}}(i\tau, k) > -1, \tag{14}$$

then the Penrose condition (P) is satisfied.

Proof. The assumptions $w \in L^1(\mathbb{R}^d)$, $g \in H^{3/2+\delta_0}_{1+\lceil d/2 \rceil}(\mathbb{R}^d)$ imply that $\lambda \mapsto \widetilde{\mathscr{L}}(\lambda, k)$ is analytic in $\{\operatorname{Re} \lambda > 0\}$. Using this analyticity, the argument principle can be applied: the number of zeros of $(1 + \widetilde{\mathscr{L}}(\lambda, k))$ in $\{\operatorname{Re} \lambda > 0\}$ is equal to the winding number of curve $\Gamma = \{\widetilde{\mathscr{L}}(i\omega, k) : \omega \in \mathbb{R}\}$ around -1. But the implication (14) prevents Γ from crossing the real line below -1. So the winding number is 0 (that is, $\mathscr{L}(i\mathbb{R},k)$ does not encircle -1) and therefore the number of zeros of $(\widetilde{\mathscr{L}}(\lambda, k) + 1)$ must be zero in $\{\operatorname{Re} \lambda > 0\}$. Finally, using Lemma 2.4, the reduction to a compact set implies the existence of a $\kappa > 0$ such that (P) holds.

Three sufficient conditions on the potential w and steady state q are now presented, each of which implies the Penrose condition (P). These are: a smallness condition, repulsive interaction (i.e. defocusing) and decreasing marginal, and a generalised Penrose condition.

Proposition 2.6. Suppose any of the three assumptions are satisfied:

(i) smallness: $w \in L^1(\mathbb{R}^d)$ and $g \in H^{\frac{3}{2}+\delta_0}_{\lceil d/2 \rceil}(\mathbb{R}^d)$ (for some $0 < \delta_0 \ll \frac{1}{2}$) are sufficiently small such that $C_1 \|w\|_{L^1} \|g\|_{H^{\frac{3}{2}+\delta_0}_{\lceil d/2 \rceil}} < 1,$

where C_1 is the constant in Lemma 2.4, (ii) repulsive interaction and decreasing marginal: $w \in L^1(\mathbb{R}^d)$ with $\widehat{w}(k) \geq 0$ for all $k \in \mathbb{R}^d$, and $g \in H^{\frac{3}{2}+\delta_0}_{1+\lceil d/2\rceil}(\mathbb{R}^d)$ is radial and satisfies

$$\begin{cases} g'_k(u) < 0 & \text{if } u > 0, \\ g'_k(u) > 0 & \text{if } u < 0, \end{cases}$$
(15)

for all $k \in \mathbb{R}^d$, where $g_k(u) := \int_{\{k\}^\perp} g\left(\frac{k}{|k|}u + |w|^2\right) dw$,

(iii) generalised Penrose condition: $w \in L^1(\mathbb{R}^d)$ with $\widehat{w}(k) \ge 0$ for all $k \in \mathbb{R}^d$, and $g \in H^{\frac{3}{2}+\delta_0}_{1+\lceil d/2 \rceil}(\mathbb{R}^d)$, and, for all $\tau \in \mathbb{R}$ and $k \ne 0$, the following holds,

$$g_k\left(\frac{\tau}{|k|} + \frac{\hbar|k|}{2}\right) = g_k\left(\frac{\tau}{|k|} - \frac{\hbar|k|}{2}\right)$$
$$\implies \frac{\widehat{w}(k)}{(2\pi)^d} \frac{1}{\hbar|k|} \text{p.v.} \int_{p_1 \in \mathbb{R}} g_k(p_1) \left[\frac{1}{p_1 - \frac{\tau}{|k|} + \frac{\hbar|k|}{2}} - \frac{1}{p_1 - \frac{\tau}{|k|} - \frac{\hbar|k|}{2}}\right] dp_1 > -1.$$

Then, the implication (14) holds and hence the Penrose condition (P) is satisfied.

Proof. For (i), Lemma 2.4 (b) is used to see that, for all $\operatorname{Re} \lambda \geq 0$ and $k \in \mathbb{R}^d$, $\left| \widetilde{\mathscr{L}}(\lambda, k) \right| \leq C_1 \|w\|_{L^1} \|g\|_{H^{\frac{3}{2}+\delta_0}}$. which implies (14).

Recall that if $g \in L^1(\mathbb{R}^d)$ then $\widetilde{\mathscr{L}}(i\tau, k=0) = 0$ for all $\tau \in \mathbb{R}$. For $k \neq 0$, using the Plemelj formula (13), one can see that the generalised Penrose condition (iii) is exactly the implication (14). Finally, for (ii), observe that if g is radial and satisfies (15), then, for $k \neq 0$, $g_k(\frac{\tau}{|k|} + \frac{|k|}{2}) = g_k(\frac{\tau}{|k|} - \frac{|k|}{2})$ if and only if $\tau = 0$. But, when $\tau = 0$, the Plemelj formula (13) implies

$$\operatorname{Re}\widetilde{\mathscr{L}}(0,k) = \frac{\widehat{w}(k)}{(2\pi)^d} \frac{2}{\hbar|k|} \lim_{\varepsilon \downarrow 0} \int_{\varepsilon}^{\infty} \frac{g_k(p_1 - \frac{\hbar|k|}{2}) - g_k(p_1 + \frac{\hbar|k|}{2})}{p_1} \, dp_1$$

which is always non-negative by the assumption that $g'_k(u) < 0$ for all u > 0, leading to (14).

2.3. Inverse Laplace transform of Green function. Throughout the remainder of the paper, w and g are assumed to satisfy the Penrose condition (P). Now, when $\operatorname{Re} \lambda \geq 0$, division by $1 + \widetilde{\mathscr{L}}(\lambda, k)$ in (11) becomes possible, yielding

$$\widetilde{\rho_{Q_{\rm L}}}(\lambda,k) = \frac{1}{1 + \widetilde{\mathscr{L}}(\lambda,k)} \widetilde{\rho_{Q_{FT}}}(\lambda,k).$$
(16)

The analysis of the inverse linearised operator, or the *Green function* of the linearised equation, now conducted, follows the approach outlined by Nguyen and You [19]. Specifically, this includes Lemma 2.7, Lemma 2.8, and Proposition 2.9.

Define the *Lindhard* function m_g by

$$m_g(\lambda, k) = \frac{1}{(2\pi)^d} \frac{2}{\hbar} \int_0^\infty e^{-\lambda t} \sin(\frac{\hbar}{2} t |k|^2) \widehat{g}(kt) dt$$

$$= \frac{1}{(2\pi)^d} \frac{2}{\hbar |k|} \int_0^\infty e^{-\lambda \frac{s}{|k|}} \sin(\frac{\hbar}{2} |k|s) \widehat{g}(\frac{k}{|k|}s) ds,$$
(17)

if $k \neq 0$, such that $\widetilde{\mathscr{L}}(\lambda, k) = \widehat{w}(k)m_g(\lambda, k)$. The Lindhard function plays an important role in physics literature; see [16].

Lemma 2.7 (Decay of $m_g(\lambda, k)$). For any $n \ge 0$, if $g \in H^{n+3/2+\delta_0}_{2+\lceil d/2 \rceil}(\mathbb{R}^d)$, then

 $\begin{array}{ll} (i) & |\partial_{\lambda}^{n}m_{g}(i\tau,k)| \lesssim_{n,d,\delta_{0}} |k|^{-n} \|g\|_{H^{n+3/2+\delta_{0}}_{\lceil d/2 \rceil}}, \ and \\ (ii) & |\partial_{\lambda}^{n}m_{g}(i\tau,k)| \lesssim_{n,d,\delta_{0}} \frac{|k|^{2-n}}{|k|^{2+\tau^{2}}} \|g\|_{H^{n+3/2+\delta_{0}}_{2+\lceil d/2 \rceil}}. \end{array}$

Proof. Differentiating (17) n times in λ ,

$$(-1)^n \partial_{\lambda}^n m_g(\lambda, k) = \frac{1}{(2\pi)^d} \frac{1}{|k|^n} \frac{2}{\hbar |k|} \int_0^\infty e^{-\lambda \frac{s}{|k|}} s^n \sin(\frac{\hbar}{2} |k|s) \widehat{g}(\frac{k}{|k|}s) \, ds,$$

So, for any $\operatorname{Re} \lambda \geq 0$,

$$\left|\partial_{\lambda}^{n}m_{g}(\lambda,k)\right| \lesssim_{n} \frac{1}{\left|k\right|^{n}} \int_{0}^{\infty} s^{n+1} \left|\widehat{g}(\frac{k}{\left|k\right|}s)\right| ds \lesssim_{n,d,\delta_{0}} \frac{1}{\left|k\right|^{n}} \|g\|_{H^{n+3/2+\delta_{0}}_{\left\lceil d/2 \right\rceil}}$$

by Lemma 2.3, which proves (i).

Next, for any $\tau \in \mathbb{R}$ and $n \geq 2$, integrating by parts twice,

$$\begin{split} \tau^{2}(-1)^{n}\partial_{\lambda}^{n}m_{g}(i\tau,k) \\ &= \frac{1}{(2\pi)^{d}}\frac{1}{|k|^{n}}\frac{2}{\hbar|k|}\int_{0}^{\infty}\tau^{2}e^{-i\tau\frac{s}{|k|}}s^{n}\sin(\frac{\hbar}{2}|k|s)\widehat{g}(\frac{k}{|k|}s)\,ds \\ &= \frac{1}{(2\pi)^{d}}\frac{|k|^{2}}{|k|^{n}}\frac{2}{\hbar|k|}\int_{0}^{\infty}\partial_{s}^{2}\left(e^{-i\tau\frac{s}{|k|}}\right)s^{n}\sin(\frac{\hbar}{2}|k|s)\widehat{g}(\frac{k}{|k|}s)\,ds \\ &= \frac{1}{(2\pi)^{d}}\frac{|k|^{2}}{|k|^{n}}\frac{2}{\hbar|k|}\int_{0}^{\infty}e^{-i\tau\frac{s}{|k|}}\left[-\frac{\hbar^{2}}{4}|k|^{2}s^{n}\sin(\frac{\hbar}{2}|k|s)\widehat{g}(\frac{k}{|k|}s)+n(n-1)s^{n-2}\sin(\frac{\hbar}{2}|k|s)\widehat{g}(\frac{k}{|k|}s)\right. \\ &\quad +2ns^{n-1}\sin(\frac{\hbar}{2}|k|s)\frac{k}{|k|}\cdot\nabla_{k}\widehat{g}(\frac{k}{|k|}s)+\frac{\hbar}{2}|k|ns^{n-1}\cos(\frac{\hbar}{2}|k|s)\widehat{g}(\frac{k}{|k|}s) \\ &\quad +\frac{\hbar}{2}|k|s^{n}\cos(\frac{\hbar}{2}|k|s)\frac{k}{|k|}\cdot\nabla_{k}\widehat{g}(\frac{k}{|k|}s)+s^{n}\sin(\frac{\hbar}{2}|k|s)(\frac{k}{|k|}\cdot\nabla_{k})^{2}\widehat{g}(\frac{k}{|k|}s)\right]ds. \end{split}$$

The first term is exactly $-(-1)^n \frac{\hbar^2}{2} |k|^4 \partial_{\lambda}^n m_g(i\tau, k)$. Moreover, all the terms in the integrand are $O(\hbar |k|)$, and hence

$$\begin{aligned} (\hbar^2|k|^4 + \tau^2) |\partial_{\lambda}^n m_g(i\tau, k)| &\lesssim_n |k|^{2-n} \int_0^\infty \langle s \rangle^{n+1} \left(\left| \widehat{g}(\frac{k}{|k|}s) \right| + \left| \nabla_k \widehat{g}(\frac{k}{|k|}s) \right| + \left| \nabla_k^2 \widehat{g}(\frac{k}{|k|}s) \right| \right) \, ds \\ &\lesssim_{n,d,\delta_0} |k|^{2-n} \|g\|_{H^{n+3/2+\delta_0}_{2+\lceil d/2\rceil}} \end{aligned}$$

by Lemma 2.3. The cases n = 0 and n = 1 give rise to fewer terms upon the integration by parts but are otherwise identical.

To conclude (ii), observe that $\tau^2 |\partial_{\lambda}^n m_g(i\tau,k)| \lesssim |k|^{2-n}$ together with the estimate $|\partial_{\lambda}^n m_g(i\tau,k)| \lesssim |k|^{-n}$ from (17) imply $(|k|^2 + \tau^2) |\partial_{\lambda}^n m_g(i\tau,k)| \lesssim |k|^{2-n}$.

Upon taking the inverse Laplace transform of (16), the following is obtained:

$$\widehat{\rho_{Q_{\rm L}}}(t,k) = \widehat{\rho_{Q_{\rm FH}}}(t,k) - \int_0^t \widehat{\mathscr{G}^r}(t-s,k)\widehat{\rho_{Q_{\rm FH}}}(s,k)\,ds,\tag{18}$$

where

$$\widetilde{\mathscr{G}}(\lambda,k) := \frac{1}{1 + \widetilde{\mathscr{L}}(\lambda,k)}.$$
(19)

Define

$$\widetilde{\mathscr{G}^{r}}(\lambda,k) := \frac{\widetilde{\mathscr{L}}(\lambda,k)}{1+\widetilde{\mathscr{L}}(\lambda,k)} = \frac{\widehat{w}(k)m_{g}(\lambda,k)}{1+\widehat{w}(k)m_{g}(\lambda,k)}$$

so that $\widetilde{\mathscr{G}}(\lambda, k) = 1 - \widetilde{\mathscr{G}^r}(\lambda, k).$

Lemma 2.8 (Decay of $\widetilde{\mathscr{G}^r}(\lambda, k)$). Let $n \ge 0$ and suppose $w \in L^1(\mathbb{R}^d)$ and $g \in H^{n+3/2+\delta_0}_{2+\lceil d/2 \rceil}(\mathbb{R}^d)$ satisfy the Penrose condition (P) with constant κ . Then, there exists a constant $C = C(n, w, g, \kappa, d, \delta_0)$ such that

$$\left|\partial_{\lambda}^{n}\widetilde{\mathscr{G}^{r}}(i\tau,k)\right| \leq C \frac{\left|k\right|^{2-n}}{\left|k\right|^{2}+\tau^{2}}.$$

Proof. Using the Faà di Bruno formula, for any $n \ge 1$,

$$\frac{d^n}{dx^n} \left[\frac{f(x)}{1+f(x)} \right] = \sum_{\pi \in \Pi_n} \frac{(-1)^{|\pi|} |\pi|!}{(1+f(x))^{|\pi|+1}} \prod_{B \in \pi} f^{(|B|)}(x),$$

where π runs through the set Π_n of all the partitions of the set $\{1, ..., n\}$. So,

$$\begin{split} \partial_{\lambda}^{n}\widetilde{\mathscr{G}^{r}}(i\tau,k) &= \frac{d^{n}}{d\lambda^{n}} \left[\frac{\widehat{w}(k)m_{g}(\lambda,k)}{1+\widehat{w}(k)m_{g}(\lambda,k)} \right] \bigg|_{\lambda=i\tau} \\ &= \sum_{\pi \in \Pi_{n}} \frac{(-1)^{|\pi|} |\pi|!}{(1+\widehat{w}(k)m_{g}(i\tau,k))^{|\pi|+1}} \prod_{B \in \pi} \widehat{w}(k) \partial_{\lambda}^{|B|} m_{g}(i\tau,k). \end{split}$$

Moreover, writing $\pi = \{B_0\} \cup (\pi \setminus \{B_0\})$ and using Lemma 2.7 (ii) to bound the $\partial_{\lambda}^{|B_0|} m_g(i\tau, k)$ and Lemma (ii) to bound the (possibly empty) $\partial_{\lambda}^{|B|} m_g(i\tau, k)$ terms in $\pi \setminus \{B_0\}$,

$$\begin{split} \left| \partial_{\lambda}^{n} \widetilde{\mathscr{G}^{r}}(i\tau,k) \right| \lesssim_{n,d,\delta_{0}} \lesssim \sum_{\pi \in \Pi_{n}} \frac{1}{\kappa^{|\pi|+1}} \frac{|k|^{2-|B_{0}|}}{|k|^{2}+\tau^{2}} |\widehat{w}(k)| \|g\|_{H^{|B_{0}|+3/2+\delta_{0}}_{2+\lceil d/2\rceil}} \prod_{B \in \pi \setminus \{B_{0}\}} |\widehat{w}(k)| |k|^{-|B|} \|g\|_{H^{|B|+3/2+\delta_{0}}_{\lceil d/2\rceil}} \\ & \leq \frac{|k|^{2-n}}{|k|^{2}+\tau^{2}} \sum_{\pi \in \Pi_{n}} \frac{|\widehat{w}(k)|^{|\pi|}}{\kappa^{|\pi|+1}} \prod_{B \in \pi} \|g\|_{H^{|B|+3/2+\delta_{0}}_{2+\lceil d/2\rceil}}. \end{split}$$

The remainder of the proof of the lemma focuses on obtaining an improved constant C. Using Sobolev-space interpolation (with $|B|+3/2+\delta_0 = (1-\theta)\cdot 0 + \theta \cdot (n+3/2+\delta_0)$ giving $\theta = \frac{|B|+3/2+\delta_0}{n+3/2+\delta_0}$ and $1-\theta = \frac{n-|B|}{n+3/2+\delta_0}$),

$$\|g\|_{H^{|B|+3/2+\delta_0}} \le \|g\|_{L^2}^{\frac{n-|B|}{n+3/2+\delta_0}} \|g\|_{H^{n+3/2+\delta_0}}^{\frac{|B|+3/2+\delta_0}{n+3/2+\delta_0}}.$$

Therefore,

$$\begin{split} \left| \partial_{\lambda}^{n} \widetilde{\mathscr{G}^{r}}(i\tau,k) \right| \lesssim_{n,d,\delta_{0}} \frac{|k|^{2-n}}{|k|^{2} + \tau^{2}} \sum_{\pi \in \Pi_{n}} \frac{|\widehat{w}(k)|^{|\pi|}}{\kappa^{|\pi|+1}} \prod_{B \in \pi} \|g\|_{L_{2+\lceil d/2]}^{\frac{n-|B|}{n+3/2+\delta_{0}}} \|g\|_{H_{n+3/2+\delta_{0}}^{\frac{n+3/2+\delta_{0}}{n+3/2+\delta_{0}}} \\ &= \frac{|k|^{2-n}}{|k|^{2} + \tau^{2}} \sum_{\pi \in \Pi_{n}} \frac{|\widehat{w}(k)|^{|\pi|}}{\kappa^{|\pi|+1}} \|g\|_{L_{2+\lceil d/2]}^{\frac{(|\pi|-1)n}{n+3/2+\delta_{0}}} \|g\|_{H_{2+\lceil d/2]}^{\frac{n+3/2+\delta_{0}}{n+3/2+\delta_{0}}} \\ &= \frac{|k|^{2-n}}{|k|^{2} + \tau^{2}} \frac{|\widehat{w}(k)|}{\kappa^{2}} \|g\|_{H_{2+\lceil d/2]}^{n+3/2+\delta_{0}}} \sum_{\pi \in \Pi_{n}} \left(\frac{|\widehat{w}(k)|}{\kappa} \|g\|_{L_{2+\lceil d/2]}^{\frac{n+3/2+\delta_{0}}{n+3/2+\delta_{0}}}} \|g\|_{H_{2+\lceil d/2]}^{\frac{n+3/2+\delta_{0}}{n+3/2+\delta_{0}}} \right)^{|\pi|-1} \\ &\lesssim \frac{|k|^{2-n}}{|k|^{2} + \tau^{2}} \frac{|\widehat{w}(k)|}{\kappa^{2}} \|g\|_{H_{2+\lceil d/2]}^{n+3/2+\delta_{0}}} \left(1 + \frac{|\widehat{w}(k)|}{\kappa} \|g\|_{L_{2+\lceil d/2]}^{\frac{n+3/2+\delta_{0}}{n+3/2+\delta_{0}}}} \|g\|_{H_{2+\lceil d/2]}^{\frac{n+3/2+\delta_{0}}{n+3/2+\delta_{0}}} \right)^{n-1} \\ &\lesssim \frac{|k|^{2-n}}{|k|^{2} + \tau^{2}} \frac{1}{\kappa} \left\langle \frac{\widehat{w}(k)}{\kappa} \right\rangle^{n} \left\langle \|g\|_{L_{2+\lceil d/2]}^{2}} \right\rangle^{n-1} \left\langle \|g\|_{H_{2+\lceil d/2]}^{n+3/2+\delta_{0}}} \right)^{1+\frac{3}{2}+\delta_{0}}}. \end{split}$$

The decay of the Green function on the Fourier side is now demonstrated by taking the inverse Laplace transform.

Proposition 2.9 (Decay of $\widehat{\mathscr{G}^r}(t,k)$). Let $n \ge 0$ and assume $w \in L^1(\mathbb{R}^d)$ and $g \in H^{n+3/2+\delta_0}_{2+\lceil d/2 \rceil}(\mathbb{R}^d)$ satisfy the Penrose condition (P) with constant κ . Then, there exists a constant $C = C(n, w, g, \kappa, d, \delta_0)$ such that

$$\left|\widehat{\mathscr{G}^r}(t,k)\right| \le C \frac{|k|}{\langle kt \rangle^n}.$$

Proof. By the Penrose condition (P), $\lambda \mapsto \widetilde{\mathscr{G}^r}(\lambda, k)$ is analytic in {Re $\lambda > 0$ }. Therefore, the contour in the inverse Laplace transform can be deformed to the imaginary axis:

$$\widehat{\mathscr{G}^{r}}(t,k) = \frac{1}{2\pi i} \int_{\operatorname{Re}\lambda=0} e^{\lambda t} \widetilde{\mathscr{G}^{r}}(\lambda,k) \, d\lambda = \frac{1}{2\pi} \int_{\mathbb{R}} e^{i\tau t} \widetilde{\mathscr{G}^{r}}(i\tau,k) \, d\tau$$

Integrating by parts n times and using Lemma 2.8,

$$\begin{split} \left|\widehat{\mathscr{G}^{r}}(t,k)\right| &= \frac{1}{2\pi t^{n}} \left| \int_{\mathbb{R}} \partial_{\tau}^{n} \left(e^{-i\tau t} \right) \widetilde{\mathscr{G}}(i\tau,k) \, d\tau \right| = \frac{1}{2\pi t^{n}} \left| \int_{\mathbb{R}} e^{-i\tau t} \partial_{\tau}^{n} \widetilde{\mathscr{G}}(i\tau,k) \, d\tau \right| \\ &\lesssim \frac{1}{(|k|t)^{n}} \int_{\mathbb{R}} \frac{|k|^{2}}{|k|^{2} + \tau^{2}} \, d\tau \\ &= \pi \frac{|k|}{(|k|t)^{n}}, \end{split}$$

where it was used that $\int_{\mathbb{R}} \frac{a}{a^2 + x^2} dx = \pi$ if a > 0. Note that there are no boundary terms by Proposition 2.8.

2.4. Linearised phase mixing. This section proves density decay and scattering for the linearised Hartree equation.

Proposition 2.10 (Linearised phase mixing). Let $N_0 > N_1 \ge 0$ be constants such that

$$0 \le N_1 < N_0 - d/2 - 1.$$

Assume $w \in L^1(\mathbb{R}^d)$ and $g \in H^{N_0+7/2+\delta_0}_{2+\lceil d/2 \rceil}(\mathbb{R}^d)$, for some $0 < \delta_0 \ll 1$, satisfy the Penrose condition (P) with constant κ . Assume the initial perturbation satisfies

$$\|Q_{\rm in}\|_{\mathcal{H}^{N_0}_M} < \infty.$$

Then, there exists a constant $C = C(N_0, w, g, \kappa, d, \delta_0)$ such that the solution Q_L to the linearised Hartree equation (LH) with initial data $Q_{\rm L}(0) = Q_{\rm in}$, satisfies the following.

• Density decay: for all $k \in \mathbb{R}^d$ and $t \ge 0$,

$$\langle k, kt \rangle^{N_0} | \widehat{\rho_{Q_L}}(t,k) | \le C \| Q_{\mathrm{in}} \|_{\mathcal{H}^{N_0}_M}$$

• Scattering: there exists an operator $Q_{\infty} \in \mathcal{H}^{N_1}(\mathbb{R}^d;\mathbb{C})$ such that, for $t \geq 0$,

$$\left\| \langle t \boldsymbol{\nabla}_x, \boldsymbol{\nabla}_\xi \rangle^{N_1} \left[e^{-i\frac{\hbar}{2}t\Delta} Q_{\mathrm{L}}(t) e^{i\frac{\hbar}{2}t\Delta} - Q_\infty \right] \right\|_{\mathcal{L}^2} \lesssim \frac{1}{\langle t \rangle^{\frac{d}{2}}}.$$
 (20)

Remark. Observe that each spatial derivative of $e^{-i\frac{\hbar}{2}t\Delta}Q_{\rm L}(t)e^{i\frac{\hbar}{2}t\Delta}$ improves the convergence to Q_{∞} by a power of t. This is not attained for the nonlinear equation in Theorem 1.5.

Proof. Applying Proposition 1.12 for the decay of $\widehat{\rho_{Q_{\text{FH}}}}$, and Proposition 2.9 for the decay of $\widehat{\mathscr{G}^r}$, to the equation (18), yields

$$\begin{split} |\widehat{\rho_{Q_{\mathrm{L}}}}(t,k)| &= \left|\widehat{\rho_{Q_{\mathrm{FH}}}}(t,k)\right| + \int_{0}^{t} \left|\widehat{\mathscr{G}^{r}}(s,k)\right| \left|\widehat{\rho_{Q_{\mathrm{FH}}}}(t-s,k)\right| ds \\ &\lesssim \frac{\|Q_{\mathrm{in}}\|_{\mathcal{H}_{M}^{N_{0}}}}{\langle k,kt\rangle^{N_{0}}} + \int_{0}^{t} \frac{|k|}{\langle ks\rangle^{N_{0}+2}} \frac{\|Q_{\mathrm{in}}\|_{\mathcal{H}_{M}^{N_{0}}}}{\langle k,k(t-s)\rangle^{N_{0}}} ds \\ &\lesssim \frac{\|Q_{\mathrm{in}}\|_{\mathcal{H}_{M}^{N_{0}}}}{\langle k,kt\rangle^{N_{0}}} + \frac{\|Q_{\mathrm{in}}\|_{\mathcal{H}_{M}^{N_{0}}}}{\langle k,kt\rangle^{N_{0}}} \int_{0}^{t} \frac{|k|}{\langle ks\rangle^{2}} ds \\ &\lesssim \frac{1}{\langle k,kt\rangle^{N_{0}}} \|Q_{\mathrm{in}}\|_{\mathcal{H}_{M}^{N_{0}}}, \end{split}$$

where it was used that $\langle k, kt \rangle \lesssim \langle k, k(t-s) \rangle \langle ks \rangle$ and $\int_0^t \frac{|k|}{\langle ks \rangle^2} ds = \int_0^{|k|t} \frac{1}{\langle r \rangle^2} dr \lesssim 1$.

For the scattering statement, the change of variables $(k, p) \mapsto (\frac{k}{2} + p, \frac{k}{2} - p)$ is performed in (8): $e^{i\hbar k \cdot pt}\widehat{Q_{\mathrm{L}}}(t,\frac{k}{2}+p,\frac{k}{2}-p) - \widehat{Q_{\mathrm{in}}}(\frac{k}{2}+p,\frac{k}{2}-p) = -i\hbar^{d-1}\widehat{w}(k)\left[g\left(\hbar(p-\frac{k}{2})\right) - g\left(\hbar(p+\frac{k}{2})\right)\right]\int_{0}^{t} e^{i\hbar k \cdot ps}\widehat{\rho_{Q_{\mathrm{L}}}}(s,k)\,ds.$

For any non-negative constant σ and multi-index α such that $\sigma + |\alpha| \leq N_1$, applying the inequality

$$\int_{x} \left| f(x+\ell) - f(x) \right|^2 dx \lesssim \left| \ell \right|^2 \int_{x} \left| \nabla f(x) \right|^2 dx,$$

yields the estimate:

$$\begin{split} \left\| |k|^{\sigma} \left(\frac{1}{\hbar} \nabla_{p} \right)^{\alpha} \left[e^{i\hbar k \cdot pt} \widehat{Q_{L}}(t, \frac{k}{2} + p, \frac{k}{2} - p) - \widehat{Q_{in}}(\frac{k}{2} + p, \frac{k}{2} - p) \right] \right\|_{L^{2}_{k,p}} \\ &\lesssim \hbar^{d-1} \sum_{\beta \leq \alpha} \left(\int_{k} |k|^{2\sigma} |\widehat{w}(k)|^{2} |\widehat{\rho_{Q_{L}}}(s,k)|^{2} \int_{p} |ks|^{2|\alpha-\beta|} |\nabla^{\beta}g \left(\hbar(p - \frac{k}{2}) \right) - \nabla^{\beta}g \left(\hbar(p + \frac{k}{2}) \right) \Big|^{2} dp dk \right)^{\frac{1}{2}} \\ &\lesssim \hbar^{\frac{d}{2}} \|w\|_{L^{1}} \sum_{\beta \leq \alpha} \|\nabla^{\beta}g\|_{\dot{H}^{1}} \left(\int_{k} \frac{|k|^{2\sigma} |k|^{2} |ks|^{2|\alpha-\beta|}}{\langle k, ks \rangle^{2N_{0}}} dk \right)^{\frac{1}{2}} \\ &= \frac{\hbar^{\frac{d}{2}}}{\langle s \rangle^{\frac{d}{2} + \sigma + 1}} \|w\|_{L^{1}} \sum_{\beta \leq \alpha} \|\nabla^{\beta}g\|_{\dot{H}^{1}} \left(\int_{k} \frac{|k|^{2\sigma} |k|^{2} |k|^{2|\alpha-\beta|}}{\langle \frac{k}{s}, k \rangle^{2N_{0}}} dk \right)^{\frac{1}{2}} \\ &\lesssim \frac{\hbar^{\frac{d}{2}}}{\langle s \rangle^{\frac{d}{2} + \sigma + 1}} \|w\|_{L^{1}} \|g\|_{H^{1+|\alpha|}} \\ N_{0} - 2\sigma - 2 - 2|\alpha| > d. \end{split}$$

Define the operator Q_{∞} by

if 2

$$\widehat{Q}_{\infty}(k,p) := \widehat{Q}_{\text{in}}(k,p) + i\hbar^{d-1}\widehat{w}(k+p) \left[g(-\hbar p) - g(\hbar k)\right] \int_{0}^{\infty} e^{i\frac{\hbar}{2}(|k|^{2} - |p|^{2})s} \widehat{\rho}_{Q_{\text{L}}}(s,k+p) \, ds$$

This leads to the convergence

$$\frac{1}{\hbar^{\frac{d}{2}}} \left\| |k|^{\sigma} \left(\frac{1}{\hbar} \nabla_p \right)^{\alpha} \left[e^{i\hbar k \cdot pt} \widehat{Q} \left(t, \frac{k}{2} + p, \frac{k}{2} - p \right) - \widehat{Q}_{\infty} \left(\frac{k}{2} + p, \frac{k}{2} - p \right) \right] \right\|_{L^2_{k,p}} \lesssim \int_t^{\infty} \frac{1}{\langle s \rangle^{\frac{d}{2} + \sigma + 1}} \, ds \lesssim \frac{1}{\langle t \rangle^{\frac{d}{2} + \sigma}}.$$

Since the convergence holds for all $\sigma + |\alpha| \leq N_1$, the scattering statement (20) follows.

3. Energy estimates on the density

This section presents a proof of the bootstrap improvements (I2) and (I4). The argument begins by applying results from Section 2 to the nonlinear equation.

3.1. Linear damping in L_t^2 . The linear theory is applied to the nonlinear equation to invert the linear operator. Consider the equation

$$\phi(t,k) = H(t,k) - \int_0^t \widehat{\mathscr{L}}(t-s,k)\phi(s,k)\,ds, \quad \widehat{\mathscr{L}}(t,k) = \frac{1}{(2\pi)^d} \frac{2}{\hbar}\widehat{w}(k)\sin(\frac{\hbar}{2}t|k|^2)\widehat{g}\left(kt\right). \tag{21}$$

Based on the linear analysis in Section 2, weighted control of the solution to the system (21) in L_t^2 is deduced, analogous to Proposition 2.2 in [1].

Proposition 3.1. Fix $T^* > 0$ and assume that H(t, k) = 0 for $t > T^*$. Let $m \in \mathbb{R}$ and $\sigma \ge 0$ an integer. Suppose $w \in L^1(\mathbb{R}^d)$ and $g \in H^{\sigma+3/2+\delta_0}_{2+\lceil d/2 \rceil}(\mathbb{R}^d)$ satisfy the Penrose condition (P) with constant κ . Assume

$$\||k|^m \langle k, kt \rangle^\sigma H(t,k)\|_{L^2_t(I)} < \infty$$

where $I = [0, T^*)$. Then, there exists a constant $C_{\text{LD}} = C_{\text{LD}}(\sigma_i, d, \delta_0, w, g, \kappa)$, such that the solution $\phi(t, k)$ to the system (21) satisfies the pointwise-in-k estimate

$$|||k|^{m} \langle k, kt \rangle^{\sigma} \phi(t,k)||_{L^{2}_{t}(I)} \leq C_{\mathrm{LD}} |||k|^{m} \langle k, kt \rangle^{\sigma} H(t,k)||_{L^{2}_{t}(I)}$$

Proof. Taking the Laplace transform of (21) yields

$$\widetilde{\phi}(\lambda,k) = \widetilde{\mathscr{G}}(\lambda,k)\widetilde{H}(\lambda,k), \tag{22}$$

where $\widetilde{\mathscr{G}}$ is defined in (19). For any $0 \leq n \leq \sigma$,

$$|k|^{n}\partial_{\lambda}^{n}\widetilde{\phi}(\lambda,k) = \sum_{j\leq n} \binom{n}{j} |k|^{j}\partial_{\lambda}^{j}\widetilde{\mathscr{G}}(\lambda,k)|k|^{n-j}\partial_{\lambda}^{n-j}\widetilde{\phi}(\lambda,k)$$
$$= \widetilde{\mathscr{G}}(\lambda,k)|k|^{n}\partial_{\lambda}^{n}H(\lambda,k) - \sum_{1\leq j\leq n} \binom{n}{j} |k|^{j}\partial_{\lambda}^{j}\widetilde{\mathscr{G}}^{r}(\lambda,k)|k|^{n-j}\partial_{\lambda}^{n-j}\widetilde{H}(\lambda,k),$$
(23)

recalling that $\widetilde{\mathscr{G}}(\lambda, k) = 1 - \widetilde{\mathscr{G}^r}(\lambda, k)$. For the first term on the right-hand side, note that $\left|\widetilde{\mathscr{G}}(i\tau, k)\right| \leq \kappa^{-1}$, which follows from w and g satisfying the Penrose condition (P). Along the imaginary axis $\lambda = i\tau$, Lemma 2.8 can be applied to the other terms in (23) to bound $|k|^j \left|\partial_{\lambda}^j \widetilde{\mathscr{G}^r}(i\tau, k)\right| \leq 1$. Therefore,

$$\left|k\right|^{n}\left|\partial_{\lambda}^{n}\widetilde{\phi}(i\tau,k)\right| \lesssim \sum_{j \leq n} \left|k\right|^{j}\left|\partial_{\lambda}^{j}\widetilde{H}(i\tau,k)\right|.$$

Taking the L^2 norm in τ and applying Plancherel's theorem, yields

$$\||kt|^n \phi(t,k)\|_{L^2_t(I)} \lesssim \|\langle kt \rangle^n H(t,k)\|_{L^2_t(I)}$$

Moreover,

$$\langle k \rangle^{\sigma} \Big| \widetilde{\phi}(i\tau,k) \Big| = \langle k \rangle^{\sigma} \Big| \widetilde{\mathscr{G}}(i\tau,k) \widetilde{H}(i\tau,k) \Big| \lesssim \kappa^{-1} \langle k \rangle^{\sigma} \Big| \widetilde{H}(i\tau,k) \Big|.$$

Using $\langle k, kt \rangle^{\sigma} \sim \langle k \rangle^{\sigma} + |kt|^{\sigma}$, it follows that

 $\|\langle k, kt \rangle^{\sigma} \phi(t,k)\|_{L^2_t(I)} \lesssim \|\langle k, kt \rangle^{\sigma} H(t,k)\|_{L^2_t(I)}.$

Multiplying both sides of (22) by $|k|^m$ yields the full result.

3.2. Bootstrap improvement (I2). The nonlinear equation (NLb) takes the form

$$\widehat{\rho_Q}(t,k) = H(t,k) - \int_0^t \mathscr{L}(t-s,k)\widehat{\rho_Q}(s,k)\,ds, \quad \mathscr{L}(t,k) = \frac{1}{(2\pi)^d} \frac{2}{\hbar} \widehat{w}(k) \sin(\frac{\hbar}{2}t|k|^2) \widehat{g}\left(kt\right) + \frac{1}{2} \widehat{g}\left(kt\right) + \frac{1}$$

with

$$H(t,k) = \widehat{\rho_{Q_{\mathrm{FH}}}}(t,k) - \frac{i\hbar^{-1}}{(2\pi)^{2d}} \int_0^t \int_\ell \widehat{w}(\ell) \widehat{\rho_Q}(s,\ell) \left(e^{-i\frac{\hbar}{2}\ell \cdot k(t-s)} - e^{i\frac{\hbar}{2}\ell \cdot k(t-s)} \right) \widehat{W[P]}(s,k-\ell,kt-\ell s) \, d\ell ds$$

Under the assumptions of Theorem 1.5, $w \in L^1(\mathbb{R}^d)$ and $g \in H^{\sigma_4+3/2+\delta_0}_{2+\lceil d/2\rceil}$ satisfy the Penrose condition (P). Therefore, the linear damping result (Proposition 3.1) can be applied to the nonlinear equation (NLb) to obtain:

$$\left\| |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_4} \widehat{\rho_Q}(t,k) \right\|_{L^2_{t,k}}^2 \le C_{\mathrm{LD}} \left(\left\| |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_4} \widehat{\rho_{Q_{\mathrm{FH}}}}(t,k) \right\|_{L^2_{t,k}}^2 + \mathcal{NL}^{(\mathrm{B1})} \right),$$

where

 $\mathcal{NL}^{(B1)}$

$$=\int_{t=0}^{T^*}\int_k \left[|k|^{\frac{1}{2}}\langle k,kt\rangle^{\sigma_4}\int_0^t\int_\ell \widehat{w}(\ell)\widehat{\rho_Q}(s,\ell)\hbar^{-1}\left(e^{-i\frac{\hbar}{2}\ell\cdot k(t-s)}-e^{i\frac{\hbar}{2}\ell\cdot k(t-s)}\right)\widehat{W[P]}\left(s,k-\ell,kt-\ell s\right)\,dpd\ell ds\right]^2dkdt.$$

The initial data term is controlled by Lemma 1.14 and the assumption $\|Q_{\rm in}\|_{\mathcal{H}_{M}^{N_0}} \leq \varepsilon$,

$$\left\| |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_4} \widehat{\rho_{Q_{\mathrm{FH}}}}(t,k) \right\|_{L^2_{t,k}}^2 \lesssim \sum_{|\alpha|,|\beta| \le M} \left\| \boldsymbol{\xi}^{\alpha} \boldsymbol{x}^{\beta} Q_{\mathrm{in}} \right\|_{\mathcal{H}^{N_0}}^2 \lesssim \varepsilon^2,$$

if $\sigma_4 < N_0 - (d+1)/2$.

The nonlinear term, bilinear in the perturbation, is now controlled. The triangle inequality $\langle k, kt \rangle^{\sigma_4} \lesssim \langle k - \ell, kt - \ell s \rangle^{\sigma_4} + \langle \ell, \ell s \rangle^{\sigma_4}$ is used to divide into transport and reaction parts in the terminology of [1], $\mathcal{NL}^{(B1)} \lesssim \mathcal{T}^{(B1)} + \mathcal{R}^{(B1)}$,

$$\mathcal{T}^{(\mathrm{B1})} \lesssim \int_{t=0}^{T^*} \int_k \left[\int_0^t \int_\ell |k|^{\frac{1}{2}} \langle k-\ell, kt-\ell s \rangle^{\sigma_4} |\widehat{w}(\ell)| |\widehat{\rho_Q}(s,\ell)| |\ell| |k(t-s)| \left| \widehat{W[P]}(s,k-\ell,kt-\ell s) \right| d\ell ds \right]^2 dk dt$$

$$\mathcal{R}^{(\mathrm{B1})} \lesssim \int_{t=0}^{T^*} \int_k \left[\int_0^t \int_\ell |k|^{\frac{1}{2}} \langle \ell,\ell s \rangle^{\sigma_4} |\widehat{w}(\ell)| |\widehat{\rho_Q}(s,\ell)| |\ell| |k(t-s)| \left| \widehat{W[P]}(s,k-\ell,kt-\ell s) \right| d\ell ds \right]^2 dk dt.$$

The regularity of the solution moves along the flow, allowing control of the transport term through the bootstrap assumption (B1). The reaction term, however, introduces the *time response kernel* and requires careful handling of resonances.

3.2.1. Control of transport term. To control the transport term, bootstrap assumptions (B1) and (B5) are used. First, by controlling $|k(t-s)| = |kt - \ell s - s(k-\ell)| \leq \langle s(k-\ell), kt - \ell s \rangle$, the weights are absorbed onto P using Definition 1.2,

$$\langle s(k-\ell), kt-\ell s \rangle \langle k-\ell, kt-\ell s \rangle^{\sigma_4} \left| \widehat{W[P]}(s,k-\ell,kt-\ell s) \right|$$
$$= \left| \overline{W[\langle s \nabla_x, \nabla_\xi \rangle \langle \nabla_x, \nabla_\xi \rangle^{\sigma_4} P]}(s,k-\ell,kt-\ell s) \right|.$$

By Cauchy–Schwarz in s and ℓ and $|\widehat{w}(\ell)| \leq 1$,

$$\mathcal{T}^{(\mathrm{B1})} \lesssim \int_{t=0}^{T^{*}} \int_{k} \left[\int_{s=0}^{t} \int_{\ell} \langle s \rangle^{d-\frac{1}{2}} |\ell| |\widehat{\rho_{Q}}(s,\ell)| \, d\ell ds \right] \\ \times \left[\int_{s=0}^{t} \int_{\ell} \langle s \rangle^{-(d-\frac{1}{2})} |\widehat{\rho_{Q}}(s,\ell)| |\ell| |k| \left| \overline{W\left[\langle s \nabla_{x}, \nabla_{\xi} \rangle \langle \nabla_{x}, \nabla_{\xi} \rangle^{\sigma_{4}} P \right]} \left(s, k-\ell, kt-\ell s \right) \right|^{2} d\ell ds \right] dk dt.$$

Then (B5) (via Lemma 1.16) implies

$$\int_{s=0}^{t} \langle s \rangle^{d-\frac{1}{2}} \int_{\ell} |\ell| |\widehat{\rho_Q}(s,\ell)| \, d\ell ds \lesssim \sqrt{K_5} \varepsilon \int_{s=0}^{\infty} \langle s \rangle^{d-\frac{1}{2}} \int_{\ell} \frac{|\ell|}{\langle \ell, \ell s \rangle^{\sigma_1}} \, d\ell ds \lesssim \sqrt{K_5} \varepsilon \int_{0}^{\infty} \frac{\langle s \rangle^{d-\frac{1}{2}}}{\langle s \rangle^{d+1}} \, ds \lesssim \sqrt{K_5} \varepsilon$$

if $\sigma_1 > d + 1$, which allows control over the first *s*- ℓ -integral. For the second, the limits of integration are changed, and Fubini's theorem is applied,

$$\mathcal{T}^{(\mathrm{B1})} \lesssim \sqrt{K_5} \varepsilon \int_{t=0}^{T^*} \int_k \int_{s=0}^t \int_{\ell} \langle s \rangle^{-(d-\frac{1}{2})} |\ell| |\widehat{\rho_Q}(s,\ell)| |k| \left| \overline{W[\langle s \nabla_x, \nabla_\xi \rangle \langle \nabla_x, \nabla_\xi \rangle^{\sigma_4} P]} (s,k-\ell,kt-\ell s) \right|^2 d\ell ds dk dt$$

$$\lesssim \sqrt{K_5} \varepsilon \int_{s=0}^{T^*} \int_{\ell} \langle s \rangle^{d-\frac{1}{2}} |\ell| |\widehat{\rho_Q}(s,\ell)| d\ell ds$$

$$\times \sup_{s \in [0,T^*], \ell \in \mathbb{R}^d} \langle s \rangle^{-2d+1} \int_k \int_{t=-\infty}^{\infty} |k| \left| \overline{W[\langle s \nabla_x, \nabla_\xi \rangle \langle \nabla_x, \nabla_\xi \rangle^{\sigma_4} P]} (s,k-\ell,kt-\ell s) \right|^2 dt dk dt$$

The s- ℓ -integral is controlled again using (B5). Finally, the change of variable $t \mapsto t/|k|$ is made, and the L^2 trace lemma (Lemma 2.2) is applied to the η -variable:

$$\mathcal{T}^{(\mathrm{B1})} \lesssim K_{5}\varepsilon^{2} \sup_{s \in [0,T^{*}], \ell \in \mathbb{R}^{d}} \langle s \rangle^{-2d+1} \int_{k} \int_{t=-\infty}^{\infty} \left| \overline{W\left[\langle s \nabla_{x}, \nabla_{\xi} \rangle \langle \nabla_{x}, \nabla_{\xi} \rangle^{\sigma_{4}} P \right]} \left(s, k - \ell, \frac{k}{|k|} t - \ell s \right) \right|^{2} dt dk$$

$$\lesssim K_{5}\varepsilon^{2} \sup_{s \in [0,T^{*}]} \langle s \rangle^{-2d+1} \int_{k} \sup_{\omega \in \mathbb{S}^{d-1}, x \in \mathbb{R}^{d}} \int_{t=-\infty}^{\infty} \left| \overline{W\left[\langle s \nabla_{x}, \nabla_{\xi} \rangle \langle \nabla_{x}, \nabla_{\xi} \rangle^{\sigma_{4}} P \right]} \left(s, k, \omega t - x \right) \right|^{2} dt dk$$

$$\lesssim K_{5}\varepsilon^{2} \sup_{s \in [0,T^{*}]} \langle s \rangle^{-2d+1} \int_{k} \left\| \langle \nabla_{\eta} \rangle^{q} \overline{W\left[\langle s \nabla_{x}, \nabla_{\xi} \rangle \langle \nabla_{x}, \nabla_{\xi} \rangle^{\sigma_{4}} P \right]} \left(s, k, \eta \right) \right\|_{L^{2}_{\eta}}^{2} dk$$

$$\lesssim K_{5}\varepsilon^{2} \sup_{s \in [0,T^{*}]} \langle s \rangle^{-2d+1} \sum_{|\alpha| \leq M} \left\| v^{\alpha} W\left[\langle s \nabla_{x}, \nabla_{\xi} \rangle \langle \nabla_{x}, \nabla_{\xi} \rangle^{\sigma_{4}} P \right] \left(s, z, v \right) \right\|_{L^{2}_{z,v}}^{2},$$

where $(d-1)/2 < q \leq M$. Using Definition 1.2, it is observed that this quantity is exactly bootstrap assumption (B1), and hence altogether it has been shown that the transport term is controlled by

$$\mathcal{T}^{(\mathrm{B1})} \lesssim K_1 K_5 \varepsilon^4.$$

3.2.2. Control of resonances. For the reaction term $\mathcal{R}^{(B1)}$, the bounds $|\widehat{w}(\ell)| \leq \langle \ell \rangle^{-2}$ and $|k(t-s)| \leq \langle s \rangle \langle k - \ell, kt - \ell s \rangle$ are utilised, and bootstrap assumption (B5) is used to control

$$\mathcal{R}^{(\mathrm{B1})} \lesssim K_5 \varepsilon^2 \int_{t=0}^{T^*} \int_k \left[\int_0^t \int_\ell \frac{\langle s \rangle |k|^{\frac{1}{2}} |\ell|^{\frac{1}{2}}}{\langle \ell \rangle^2 \langle k-\ell, kt-\ell s \rangle^{\sigma_1-1}} |\ell|^{\frac{1}{2}} \langle \ell, \ell s \rangle^{\sigma_4} |\widehat{\rho_Q}(s,\ell)| \, d\ell ds \right]^2 dk dt.$$

Hence,

$$\mathcal{R}^{(\mathrm{B1})} \lesssim K_5 \varepsilon^2 \left\| \overline{R} \left[|k|^{\frac{1}{2}} \langle k, ks \rangle^{\sigma_4} \widehat{\rho_Q}(s, k) \right] \right\|_{L^2_{t,k}}^2$$
(24)

where the integral operator $\overline{R} \colon L^2_t L^2_k \to L^2_t L^2_k$ is defined by

$$\overline{R}[f](t,k) = \int_0^t \int_\ell \frac{\langle s \rangle |k|^{\frac{1}{2}} |\ell|^{\frac{1}{2}}}{\langle \ell \rangle^2 \langle k-\ell, kt-\ell s \rangle^{\sigma_1-1}} f(s,\ell) \, d\ell ds.$$
(25)

Through this kernel, density modes at earlier times $0 \le s \le t$ can influence the k-th mode of the density at time t. Preventing a cascade of resonances is crucial: the ℓ -th mode of the density at time s may sequentially drive the k-th mode at time t, and so on, potentially causing growth in the nonlinear interaction at arbitrarily large times. Schur's test implies that

$$\begin{aligned} \left\|\overline{R}\right\|_{L^{2}_{t}L^{2}_{k} \to L^{2}_{t}L^{2}_{k}}^{2} &\leq \left(\sup_{t \in [0, T^{*}], k \in \mathbb{R}^{d}} \int_{0}^{t} \int_{\ell} \frac{\langle s \rangle |k|^{\frac{1}{2}} |\ell|^{\frac{1}{2}}}{\langle \ell \rangle^{2} \langle k - \ell, kt - \ell s \rangle^{\sigma_{1} - 1}} \, d\ell ds \right) \\ &\times \left(\sup_{s \in [0, T^{*}], \ell \in \mathbb{R}^{d}} \int_{s}^{T^{*}} \int_{k} \frac{\langle s \rangle |k|^{\frac{1}{2}} |\ell|^{\frac{1}{2}}}{\langle \ell \rangle^{2} \langle k - \ell, kt - \ell s \rangle^{\sigma_{1} - 1}} \, dk dt \right). \end{aligned}$$

$$(26)$$

The following lemma provides control over these integrals, analogous to [1, Lemmas 3.1 and 3.2], with the proof extended to $d \ge 3$. The main challenge in managing the integrals in (26) arises from the $\langle s \rangle$ -growth in the numerator, which becomes difficult to handle due to potential cancellations in $\langle k - \ell, kt - \ell s \rangle$ when k and ℓ are nearly collinear. To address this, a *resonant region* in \mathbb{R}^d is identified, where k and ℓ are nearly

collinear, and which diminishes over time. The integral over this resonant region converges as it shrinks at a sufficient rate to ensure convergence in the time variable. Outside the resonant region, where ℓ and k are sufficiently non-collinear, additional decay is extracted from the denominator through powers of $|kt - \ell s|$.

Lemma 3.2 (Control of resonances). Let $T^* \ge 0$ and $\sigma_1 \ge d+7$. The following integral inequalities hold:

$$\sup_{t \in [0,T^*], k \in \mathbb{R}^d} \int_0^t \int_{\ell \in \mathbb{R}^d} \frac{\langle s \rangle |k|^{\frac{1}{2}} |\ell|^{\frac{1}{2}}}{\langle \ell \rangle^2 \langle k - \ell, kt - \ell s \rangle^{\sigma_1 - 1}} d\ell ds \lesssim 1$$
(27)

and

$$\sup_{s\in[0,T^*],\ell\in\mathbb{R}^d}\int_s^{T^*}\int_{k\in\mathbb{R}^d}\frac{\langle s\rangle|k|^{\frac{1}{2}}|\ell|^{\frac{1}{2}}}{\langle\ell\rangle^2\langle k-\ell,kt-\ell s\rangle^{\sigma_1-1}}dkdt\lesssim 1.$$
(28)

As a consequence, Schur's test establishes that the operator \overline{R} , defined in (25), satisfies $\|\overline{R}\|_{L_t^2 L_k^2 \to L_t^2 L_k^2} \lesssim 1$.

Proof. To establish inequality (27), fix $t \in [0, T^*]$ and $k \in \mathbb{R}^d$. Let $\beta := \sigma_1 - 1$ for simplicity. The integral over early times is controlled as follows:

$$\int_0^1 \int_{\ell} \frac{\langle s \rangle |k|^{\frac{1}{2}} |\ell|^{\frac{1}{2}}}{\langle \ell \rangle^2 \langle k-\ell, kt-\ell s \rangle^{\beta}} d\ell ds \lesssim \int_{\ell} \frac{|k|^{\frac{1}{2}} |\ell|^{\frac{1}{2}}}{\langle \ell \rangle^2 \langle k-\ell \rangle^{\beta}} d\ell \lesssim \int_{\ell} \frac{|\ell|^{\frac{1}{2}}}{\langle \ell \rangle^{\frac{3}{2}} \langle k-\ell \rangle^{\beta-\frac{1}{2}}} d\ell \lesssim \int_{\ell} \frac{|\ell|^{\frac{1}{2}}}{\langle \ell \rangle^{\beta+1}} d\ell \lesssim 1,$$

where $\langle k \rangle \lesssim \langle \ell \rangle \langle k - \ell \rangle$ was used.

Assume $T^* > 1$. Decompose $\ell \in \mathbb{R}^d$ into parallel and orthogonal components relative to k as $\ell_{\parallel} := (k \cdot \ell)k/|k|^2$ and $\ell_{\perp} := \ell - \ell_{\parallel}$. Define $I_R \subset \mathbb{R}^d$ as a cylinder around k with radius shrinking over time, and set $I_{NR} = I_R^c$, where

$$I_R = \left\{ \ell : |\ell_{\perp}| < \frac{|k|^b}{\langle s \rangle^{\zeta}} \right\} \quad \text{and} \quad I_{NR} = \left\{ \ell : |\ell_{\perp}| \ge \frac{|k|^b}{\langle s \rangle^{\zeta}} \right\},$$

with $\zeta := 2/(d-1/2) + \text{ and } b := (1-\zeta)/2$. Here, $x = a \pm \text{ denotes } x = a \pm \overline{\varepsilon}$ for arbitrary $0 < \overline{\varepsilon} \ll 1$.

The integral over the cylinder I_R satisfies, using $|\ell|^{\frac{1}{2}} \lesssim |\ell_{\parallel}|^{\frac{1}{2}} + |\ell_{\perp}|^{\frac{1}{2}}$,

$$\mathcal{I}_{R} := \int_{1}^{t} \int_{I_{R}} \frac{\langle s \rangle |k|^{\frac{1}{2}} |\ell|^{\frac{1}{2}}}{\langle \ell \rangle^{2} \langle k - \ell, kt - \ell s \rangle^{\beta}} \, d\ell ds \lesssim \int_{1}^{t} \int_{I_{R}} \frac{\langle s \rangle |k|^{\frac{1}{2}}}{\langle \ell_{\parallel} \rangle^{2} \langle k - \ell_{\parallel}, kt - \ell_{\parallel} s \rangle^{\beta}} \left(|\ell_{\parallel}|^{\frac{1}{2}} + |\ell_{\perp}|^{\frac{1}{2}} \right) \, d\ell ds.$$

Applying the bound for $|\ell_{\perp}|$ on I_R and the volume of a cylinder on \mathbb{R}^d gives

$$\mathcal{I}_R \lesssim \int_1^t \int_{-\infty}^\infty \frac{\langle s \rangle |k|^{\frac{1}{2}}}{\langle \ell_{\parallel} \rangle^2 \langle k - \ell_{\parallel}, kt - \ell_{\parallel} s \rangle^{\beta}} \frac{|k|^{(d-1)b}}{\langle s \rangle^{(d-1)\zeta}} \left(\left| \ell_{\parallel} \right|^{\frac{1}{2}} + \frac{|k|^{\frac{b}{2}}}{\langle s \rangle^{\frac{\zeta}{2}}} \right) d\ell_{\parallel} ds$$

leading to two contributions \mathcal{I}_1 and \mathcal{I}_2 . For \mathcal{I}_1 , the change of variable $s \mapsto s/|\ell_{\parallel}|$ and $(d-1)\zeta \geq 1$ yield convergence under the condition $\beta - 2(d-1)b > 1/2$. The second term \mathcal{I}_2 converges due to the choice of ζ , ensuring that $1/\langle s \rangle^{(d-1/2)\zeta-1}$ is integrable, if $\beta > 2(d-1/2)b$.

On I_{NR} , for $t \ge 1$, it holds that $|kt - \ell s| \gtrsim s |\ell_{\perp}| \gtrsim |k|^b \langle s \rangle^{1-\zeta}$, so

$$\mathcal{I}_{NR} := \int_{1}^{t} \int_{I_{NR}} \frac{\langle s \rangle |k|^{\frac{1}{2}} |\ell|^{\frac{1}{2}}}{\langle \ell \rangle^{2} \langle k-\ell, kt-\ell s \rangle^{\beta}} \, d\ell ds \lesssim \int_{1}^{t} \int_{I_{NR}} \frac{\langle s \rangle |k|^{\frac{1}{2}} |\ell|^{\frac{1}{2}}}{\langle \ell \rangle^{2} \langle k-\ell, kt-\ell s \rangle^{\beta-\frac{1}{2b}} |k|^{\frac{1}{2}} \langle s \rangle^{\frac{1}{2b}(1-\zeta)}} \, d\ell ds.$$

Using $(1 - \zeta)/(2b) - 1 = 0$ and $\beta - 1/(2b) > d$, and applying the change of variable $s \mapsto s/|\ell|$, the integral \mathcal{I}_{NR} converges if $\beta - 1/(2b) + 3/2 > d$. Hence, inequality (27) holds if β is sufficiently large.

To establish inequality (28), it is observed that:

$$\sup_{s \in [0,T^*], \ell \in \mathbb{R}^d} \int_s^{T^*} \int_{k \in \mathbb{R}^d} \frac{\langle s \rangle |k|^{\frac{1}{2}} |\ell|^{\frac{1}{2}}}{\langle \ell \rangle^2 \langle k - \ell, kt - \ell s \rangle^\beta} dk dt \le \sup_{s \in [0,T^*], \ell \in \mathbb{R}^d} \int_0^{T^*} \int_{k \in \mathbb{R}^d} \frac{\langle t \rangle |k|^{\frac{1}{2}} |\ell|^{\frac{1}{2}}}{\langle \ell \rangle^2 \langle k - \ell, kt - \ell s \rangle^\beta} dk dt \le \frac{1}{2} \left(\int_s^{T^*} \int_{k \in \mathbb{R}^d} \frac{\langle t \rangle |k|^{\frac{1}{2}} |\ell|^{\frac{1}{2}}}{\langle \ell \rangle^2 \langle k - \ell, kt - \ell s \rangle^\beta} dk dt \le \frac{1}{2} \left(\int_s^{T^*} \int_{k \in \mathbb{R}^d} \frac{\langle t \rangle |k|^{\frac{1}{2}} |\ell|^{\frac{1}{2}}}{\langle \ell \rangle^2 \langle k - \ell, kt - \ell s \rangle^\beta} dk dt \le \frac{1}{2} \left(\int_s^{T^*} \int$$

For fixed $s \in [0, T^*]$ and $\ell \in \mathbb{R}^d$, the integral over early times is estimated as

$$\int_0^1 \int_k \frac{\langle t \rangle |k|^{\frac{1}{2}} |\ell|^{\frac{1}{2}}}{\langle \ell \rangle^2 \langle k-\ell, kt-\ell s \rangle^\beta} dk dt \lesssim \int_k \frac{|k|^{\frac{1}{2}} |\ell|^{\frac{1}{2}}}{\langle \ell \rangle^2 \langle k-\ell \rangle^\beta} dk \lesssim \int_k \frac{|k|^{\frac{1}{2}}}{\langle k \rangle^\beta} dk \lesssim 1.$$

Assume $T^* > 1$. Decompose $k \in \mathbb{R}^d$ into parallel and orthogonal components relative to ℓ , defining $k_{\parallel} := (\ell \cdot k)\ell/|\ell|^2$ and $k_{\perp} := k - k_{\parallel}$, and partition

$$J_R = \left\{ k : |k_{\perp}| < \frac{|\ell|^c}{\langle s \rangle^{\zeta}} \right\} \quad \text{and} \quad J_{NR} = \left\{ k : |k_{\perp}| \ge \frac{|\ell|^c}{\langle s \rangle^{\zeta}} \right\},$$

where $\zeta := 2/(d - 1/2) + \text{ and } c = \min \{3/(2(d - 1/2)), (1 - \zeta)/2\}.$

The integral over J_R satisfies

$$\begin{aligned} \mathcal{J}_R &:= \int_1^{T^*} \int_{J_R} \frac{\langle t \rangle |\ell|^{\frac{1}{2}}}{\langle \ell \rangle^2 \langle k_{\parallel} - \ell, k_{\parallel} t - \ell s \rangle^{\beta}} \left(\left| k_{\parallel} \right|^{\frac{1}{2}} + \left| k_{\perp} \right|^{\frac{1}{2}} \right) \, dk dt \\ &\lesssim \int_1^{T^*} \int_{\mathbb{R}} \frac{|\ell|^{\frac{1}{2}}}{\langle \ell \rangle^2 \langle k_{\parallel} - \ell, k_{\parallel} t - \ell s \rangle^{\beta}} \frac{|\ell|^{(d-1)b}}{\langle t \rangle^{(d-1)\zeta - 1}} \left(\left| k_{\parallel} \right|^{\frac{1}{2}} + \frac{|\ell|^{\frac{b}{2}}}{\langle t \rangle^{\frac{\zeta}{2}}} \right) \, dk_{\parallel} dt, \end{aligned}$$

leading to two contributions, \mathcal{J}_1 and \mathcal{J}_2 . Following the change of variable $t \mapsto t/|k_{\parallel}|$ in \mathcal{J}_1 , the integrals converge under the conditions $c \leq 3/(2(d-1/2))$ and $(d-1/2)\zeta - 1 > 1$.

On J_{NR} for $t \ge 1$ it holds that $|kt - \ell s| \gtrsim t |k_{\perp}| \gtrsim |\ell|^c \langle t \rangle^{1-\zeta}$, yielding

$$\begin{aligned} \mathcal{J}_{NR} &:= \int_{1}^{T^{*}} \int_{J_{NR}} \frac{\langle t \rangle |\ell|^{\frac{1}{2}} |k|^{\frac{1}{2}}}{\langle \ell \rangle^{2} \langle k - \ell, kt - \ell s \rangle^{\beta}} dk dt \\ &\lesssim \int_{1}^{T^{*}} \int_{I_{NR}} \frac{|\ell|^{\frac{1}{2}} |k|^{\frac{1}{2}}}{\langle \ell \rangle^{2} \langle k - \ell, kt - \ell s \rangle^{\beta - x} |\ell|^{cx} \langle t \rangle^{x(1 - \zeta) - 1}} \, dk dt. \end{aligned}$$

Choosing $x = 1/(1-\zeta)$ yields $x(1-\zeta) - 1 = 0$ and $cx \le \frac{1}{2}$. Applying the change of variable $t \mapsto t/|k|$, convergence holds if $\beta - x - \frac{1}{2} > d$, requiring $\beta > d + (d - 1/2)(d - 5/2) + 1/2$. All conditions on β are satisfied if $\beta \ge d + 6$ for $d \ge 3$.

Applying Lemma 3.2 to (24), it follows that the reaction term is controlled by

$$\mathcal{R}^{(\mathrm{B1})} \lesssim K_5 \varepsilon^2 \left\| |k|^{\frac{1}{2}} \langle k, ks \rangle^{\sigma_4} \widehat{\rho_Q}(s, k) \right\|_{L^2_{t,k}}^2$$

3.2.3. Conclusion of bootstrap improvement (I2). Altogether, under the bootstrap assumptions (B1) and (B5), it has been shown that there exists a positive constant $K_{\rm LD} = K_{\rm LD}(C_{\rm LD}, \sigma_i, d, w, Q_{\rm in})$, independent of $0 < \hbar \leq 1$ and K_i , such that

$$\left\| |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_4} \widehat{\rho_Q}(t,k) \right\|_{L^2_{t,k}}^2 \le K_{\mathrm{LD}} \left(\varepsilon^2 + K_1 K_5 \varepsilon^4 + K_5 \varepsilon^2 \left\| |k|^{\frac{1}{2}} \langle k, ks \rangle^{\sigma_4} \widehat{\rho_Q}(t,k) \right\|_{L^2_{t,k}}^2 \right).$$

By choosing ε sufficiently small such that $K_{\rm LD}K_5\varepsilon^2 \leq 1/2$, the last term on the right-hand side can be absorbed onto the left-hand side,

$$\left\| |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_4} \widehat{\rho_Q}(t,k) \right\|_{L^2_{t,k}}^2 \leq 2K_{\mathrm{LD}} \left(\varepsilon^2 + K_1 K_5 \varepsilon^4 \right).$$

Moreover, by choosing ε sufficiently small such that $K_1 K_5 \varepsilon^2 \leq 1$ and $K_2 := 2K_{\text{LD}}$, it can be concluded that

$$\left\| |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_4} \widehat{\rho_Q}(t,k) \right\|_{L^2_{t,k}}^2 \le 2K_2 \varepsilon^2.$$

Note that K_2 does not depend on the other K_i , $i \neq 2$. Therefore, the improvement (I2) in Proposition 1.15 is established.

3.3. Bootstrap improvement (I4). As in Section 3.2, the linear damping result (Proposition 3.1) is applied to the nonlinear equation, which gives

$$\left\| |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_2} \widehat{\rho_Q}(t,k) \right\|_{L^2_t}^2 \lesssim \left\| |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_2} \widehat{\rho_{Q_{\mathrm{FH}}}}(t,k) \right\|_{L^2_t}^2 + \mathcal{NL}^{(\mathrm{B4})}(k),$$

where

$$\mathcal{NL}^{(\mathrm{B4})}(k) = \int_{t=0}^{T^*} \left[|k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_2} \int_0^t \int_\ell \widehat{w}(\ell) \widehat{\rho_Q}(s,\ell) \hbar^{-1} \left(e^{-i\frac{\hbar}{2}\ell \cdot k(t-s)} - e^{i\frac{\hbar}{2}\ell \cdot k(t-s)} \right) \widehat{W[P]}(s,k-\ell,kt-\ell s) \, dp d\ell ds \right]^2 dt.$$

The initial data term is controlled by Lemma 1.14 and the assumption on the initial data,

$$\left\||k|^{\frac{1}{2}}\langle k,kt\rangle^{\sigma_{2}}\widehat{\rho_{Q_{\mathrm{FH}}}}(t,k)\right\|_{L^{\infty}_{k}L^{2}_{t}}^{2}\lesssim \sum_{|\alpha|,|\beta|\leq M}\left\|\boldsymbol{\xi}^{\alpha}\boldsymbol{x}^{\beta}Q_{\mathrm{in}}\right\|_{\mathcal{H}^{N_{0}}}^{2}\lesssim \varepsilon^{2}$$

if $\sigma_2 < N_0 - 1/2$.

As before, to control the nonlinear part, the division into transport and reaction parts is performed, $\mathcal{NL}^{(B4)} \leq \mathcal{T}^{(B4)} + \mathcal{R}^{(B4)}$, where

$$\mathcal{T}^{(\mathrm{B4})} \lesssim \int_{t=0}^{T^*} \left[\int_0^t \int_{\ell} |k|^{\frac{1}{2}} \langle k-\ell, kt-\ell s \rangle^{\sigma_2} |\widehat{w}(\ell)| |\widehat{\rho_Q}(s,\ell)| |\ell| |k(t-s)| \left| \widehat{W[P]}(s,k-\ell,kt-\ell s) \right| d\ell ds \right]^2 dt \\ \mathcal{R}^{(\mathrm{B4})} \lesssim \int_{t=0}^{T^*} \left[\int_0^t \int_{\ell} |k|^{\frac{1}{2}} \langle \ell,\ell s \rangle^{\sigma_2} |\widehat{w}(\ell)| |\widehat{\rho_Q}(s,\ell)| |\ell| |k(t-s)| \left| \widehat{W[P]}(s,k-\ell,kt-\ell s) \right| d\ell ds \right]^2 dt.$$

To control the transport term, bootstrap assumptions (B3) and (B5) are used. First, the assumption $|\widehat{w}(\ell)| \leq 1$ is applied, and $|k(t-s)| \leq |k|t$ is controlled. Then

$$\begin{aligned} \mathcal{T}^{(\mathrm{B4})} &\lesssim K_5 \varepsilon^2 |k|^3 \int_{t=0}^{T^*} t^2 \left[\int_{\ell}^t \int_{\ell} \frac{|\ell|}{\langle \ell, \ell s \rangle^{\sigma_1}} \langle k - \ell, kt - \ell s \rangle^{\sigma_2} \left| \widehat{W[P]} \left(s, k - \ell, kt - \ell s \right) \right| d\ell ds \right]^2 dt \\ &\lesssim K_5 \varepsilon^2 |k|^3 \int_{t=0}^{T^*} t^2 \left[\int_{s=0}^t \left(\int_{\ell} \frac{|\ell|^2}{\langle \ell, \ell s \rangle^{2\sigma_1}} \frac{1}{|k - \ell|^{2\delta} \langle k - \ell, kt - \ell s \rangle^{2(\sigma_3 - \sigma_2)}} d\ell \right)^{\frac{1}{2}} \\ & \times \left(\int_{\ell} |k - \ell|^{2\delta} \langle k - \ell, kt - \ell s \rangle^{2\sigma_3} \left| \widehat{W[P]} \left(s, k - \ell, kt - \ell s \right) \right|^2 d\ell \right)^{\frac{1}{2}} ds \right]^2 dt, \end{aligned}$$

where Cauchy–Schwarz was performed in ℓ in the second inequality. To control the integral containing $\widehat{W}[\widehat{P}]$, the Sobolev embedding is used together with Lemma 1.11 and bootstrap assumption (B3),

$$\begin{split} \int_{\ell} \sup_{\eta} \left| \overline{W[|\boldsymbol{\nabla}_{x}|^{\delta} \langle \boldsymbol{\nabla}_{x}, \boldsymbol{\nabla}_{\xi} \rangle^{\sigma_{3}} P]}(\ell, \eta) \right|^{2} d\ell &\lesssim \int_{\ell} \int_{\eta} \left| \langle \nabla_{\eta} \rangle^{s} \overline{W[|\boldsymbol{\nabla}_{x}|^{\delta} \langle \boldsymbol{\nabla}_{x}, \boldsymbol{\nabla}_{\xi} \rangle^{\sigma_{3}} P]}(\ell, \eta) \right|^{2} d\eta d\ell \\ &\lesssim \sum_{|\alpha| \leq M} \left\| W \left[\boldsymbol{\xi}^{\alpha} |\boldsymbol{\nabla}_{x}|^{\delta} \langle \boldsymbol{\nabla}_{x}, \boldsymbol{\nabla}_{\xi} \rangle^{\sigma_{3}} P \right] \right\|_{L^{2}_{z,v}}^{2} \\ &\lesssim K_{3} \varepsilon^{2} \end{split}$$

if $d/2 < s \leq \lceil (d+1)/2 \rceil$. For the other integral, by the property of the Japanese bracket, $\langle \ell, \ell s \rangle \langle k-\ell, kt-\ell s \rangle \gtrsim \langle k, kt \rangle$. Hence, if $j \leq \min \{\sigma_1, \sigma_3 - \sigma_2\}$, then

$$\frac{1}{\langle \ell, \ell s \rangle^{2\sigma_1} \langle k - \ell, kt - \ell s \rangle^{2(\sigma_3 - \sigma_2)}} \lesssim \frac{1}{\langle \ell, \ell s \rangle^{2\sigma_1 - 2j} \langle k, kt \rangle^{2j}}.$$

Hence

$$\mathcal{T}^{(\mathrm{B4})} \lesssim K_3 K_5 \varepsilon^4 |k|^3 \int_{t=0}^{T^*} \frac{t^2}{\langle k, kt \rangle^{2j}} dt \left[\int_{s=0}^t \left(\int_{\ell} \frac{|\ell|^2}{\langle \ell, \ell s \rangle^{2\sigma_1 - 2j}} \frac{1}{|k-\ell|^{2\delta}} d\ell \right)^{\frac{1}{2}} ds \right]^2$$
$$\lesssim K_3 K_5 \varepsilon^4 \int_{t=0}^{|k|T^*} \frac{t^2}{\langle k, t \rangle^{2j}} dt \left[\int_{s=0}^t \frac{1}{\langle s \rangle^{\frac{d}{2} + 1 - \delta}} ds \right]^2$$
$$\lesssim K_3 K_5 \varepsilon^4$$

if $\sigma_1 - j > d/2 + 1 - \delta$ and j > 3/2. This leads to the assumption $\sigma_3 - \sigma_2 > d/2$ and $\sigma_1 > d/2 + 5/2 - \delta$.

To control the reaction term, bootstrap assumptions (B2) and (B5) are used. Using the assumption $|\hat{w}(\ell)| \leq 1$ and starting with Cauchy–Schwarz in ℓ and s,

$$\mathcal{R}^{(\mathrm{B4})} \lesssim \int_{t=0}^{T^*} |k|^3 t^2 \left(\int_{s=0}^t \int_{\ell} |\ell| \langle \ell, \ell s \rangle^{2\sigma_4} |\widehat{\rho_Q}(s,\ell)|^2 \, d\ell ds \right) \\ \times \left(\int_{s=0}^t \int_{\ell} |\ell| \langle \ell, \ell s \rangle^{-2(\sigma_4 - \sigma_2)} \Big| \widehat{W[P]}(s,k-\ell,kt-\ell s) \Big|^2 \, d\ell ds \right) dt \\ \lesssim K_2 K_5 \varepsilon^4 \int_{t=0}^{T^*} |k|^3 t^2 \int_{s=0}^t \int_{\ell} \frac{|\ell|}{\langle \ell, \ell s \rangle^{2(\sigma_4 - \sigma_2)} \langle k - \ell, kt - \ell s \rangle^{2\sigma_1}} \, d\ell ds dt.$$

Letting $j \leq \min \{\sigma_1, \sigma_4 - \sigma_2\}$, it follows that

$$\frac{1}{\langle \ell, \ell s \rangle^{2(\sigma_4 - \sigma_2)} \langle k - \ell, kt - \ell s \rangle^{2\sigma_1}} \lesssim \frac{1}{\langle \ell, \ell s \rangle^{2(\sigma_4 - \sigma_2 - j)} \langle k, kt \rangle^{2j}}$$

Hence

$$\mathcal{R}^{(\mathrm{B4})} \lesssim K_2 K_5 \varepsilon^4 \int_{t=0}^{T^*} \frac{|k|^3 t^2}{\langle k, kt \rangle^{2j}} dt \int_{s=0}^t \int_{\ell} \frac{|\ell|}{\langle \ell, \ell s \rangle^{2(\sigma_4 - \sigma_2 - j)}} d\ell ds$$
$$\lesssim K_2 K_5 \varepsilon^4$$

where the ℓ -s-integral converges if $j < \sigma_4 - \sigma_2 - (d+1)/2$ and the t-integral converges if j > 3/2. For such a j to exist, the assumption $\sigma_4 - \sigma_2 > d/2 + 2$ is needed.

3.3.1. Conclusion of bootstrap improvement (I4). Under the bootstrap assumptions (B2), (B3), and (B5), a positive constant $\widetilde{K} = \widetilde{K}(d, g, w, \sigma_i, Q_{in})$ exists, such that

$$\left\| |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_2} \widehat{\rho_Q}(t,k) \right\|_{L^2_t}^2 \leq \widetilde{K} \varepsilon^2 \left(1 + (K_3 K_5 + K_2 K_5) \varepsilon^4 \right).$$

For $\varepsilon^2 \leq (K_3K_5 + K_2K_5)^{-1}$ and $K_4 := \widetilde{K}$ the improvement (I4) in Proposition 1.15 is achieved:

$$\left\| |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_2} \widehat{\rho_Q}(t,k) \right\|_{L^2_t}^2 \le 2K_4 \varepsilon^2$$

The constant K_4 is independent of the other K_i , $i \neq 4$.

4. Energy estimates on the density matrix

This section presents the proof of the bootstrap improvements (I5), (I1), and (I3) on P, the density matrix of the solution of the nonlinear Hartree equation along the free flow.

A useful lemma for the energy estimates from [1] is now presented.

Lemma 4.1. (a) Let $f_1, f_2 \in L^2(\mathbb{R}^d_k \times \mathbb{R}^d_\eta)$ and $r \in L^1(\mathbb{R}^d)$. Then

$$\left| \int_{\eta} \int_{k} \int_{\ell} f_{1}(k,\eta) r(\ell) f_{2}(k-\ell,\eta-\ell t) \, d\ell d\ell d\eta \right| \lesssim \|f_{1}\|_{L^{2}_{k,\eta}} \|f_{2}\|_{L^{2}_{k,\eta}} \|r\|_{L^{1}_{k}}$$

(b) Let
$$f_1 \in L^2(\mathbb{R}^d_k \times \mathbb{R}^d_\eta)$$
 and $f_2 \in L^1(\mathbb{R}^d_k; L^2(\mathbb{R}^d_\eta))$ and $r \in L^2(\mathbb{R}^d)$. Then
$$\left| \int_{\eta} \int_k \int_{\ell} f_1(k, \eta) r(\ell) f_2(k - \ell, \eta - \ell t) \, d\ell dk d\eta \right| \lesssim \|f_1\|_{L^2_{k,\eta}} \|f_2\|_{L^1_k(L^2_\eta)} \|r\|_{L^2_k}.$$

Proof. The proof of (a) relies on the Cauchy–Schwarz inequality, while the proof of (b) incorporates Young's convolution inequality. \Box

4.1. Bootstrap improvement (I5). This section demonstrates the improvement of the bootstrap assumption (B5). By the definition of the Wigner transform, it is equivalent to prove that

$$\langle k,\eta\rangle^{\sigma_1} \left|\widehat{W[P]}(k,\eta)\right| \leq \sqrt{2}K_5\varepsilon,$$

for ε sufficiently small.

Recalling the definitions of $W[P_{\rm L}]$ and $W[P_{\rm NL}]$ in (5), the triangle inequality is applied to bound

$$\begin{split} \langle k,\eta\rangle^{\sigma_1} \left| \widehat{W[P]}(k,\eta) \right| \\ &\leq \langle k,\eta\rangle^{\sigma_1} \left| \widehat{W[Q_{\mathrm{in}}]}(k,\eta) \right| + \langle k,\eta\rangle^{\sigma_1} \int_0^t \left| \widehat{W[P_{\mathrm{L}}]}(s,k,\eta) \right| ds + \langle k,\eta\rangle^{\sigma_1} \int_0^t \left| \widehat{W[P_{\mathrm{NL}}]}(s,k,\eta) \right| ds \\ &= \mathcal{I}^{(\mathrm{B5})} + \mathcal{L}^{(\mathrm{B5})} + \mathcal{N}\mathcal{L}^{(\mathrm{B5})}. \end{split}$$

The initial data term satisfies $\mathcal{I} \lesssim \varepsilon$, using

$$\langle k,\eta\rangle^{\sigma_1} \left| \widehat{W[Q_{\mathrm{in}}]}(k,\eta) \right| = \left| \widehat{W[\langle \boldsymbol{\nabla}_x, \boldsymbol{\nabla}_{\boldsymbol{\xi}} \rangle^{\sigma_1} Q_{\mathrm{in}}]}(k,\eta) \right|,$$

and Lemma 1.9 together with the assumption on $Q_{\rm in}$ in Theorem 1.5.

Turning next to the linear term, the triangle inequality $\langle k, \eta \rangle^{\sigma_1} \lesssim \langle k, ks \rangle^{\sigma_1} \langle \eta - ks \rangle^{\sigma_1}$ and bootstrap assumption (B4) yield

$$\begin{split} \langle k,\eta\rangle^{\sigma_1} \left| \int_0^t \widehat{W[P_{\mathrm{L}}]}(s,k,\eta) \, ds \right| &= \langle k,\eta\rangle^{\sigma_1} \left| \int_0^t \frac{-i\hbar^{-1}}{(2\pi)^d} \widehat{w}(k) \widehat{\rho}(s,k) \left(e^{-i\frac{\hbar}{2}k \cdot (\eta-ks)} - e^{i\frac{\hbar}{2}k \cdot (\eta-ks)} \right) \widehat{g}(\eta-ks) \, ds \right| \\ &\lesssim \int_0^t \langle k,ks\rangle^{\sigma_1} |\widehat{w}(k)| |\widehat{\rho_Q}(s,k)| |k| \langle \eta-ks\rangle^{\sigma_1} |\eta-ks| |\widehat{g}(\eta-ks)| \, ds \\ &\lesssim \left\| |k|^{\frac{1}{2}} \langle k,kt\rangle^{\sigma_2} \widehat{\rho_Q}(t,k) \right\|_{L^2_t} \left(\int_0^t |k| \langle \eta-ks\rangle^{2\sigma_1+2} |\widehat{g}(\eta-ks)|^2 \, ds \right)^{\frac{1}{2}} \\ &\lesssim \sqrt{K_4} \varepsilon \left(\int_0^\infty \langle \eta-\frac{k}{|k|}s\rangle^{2\sigma_1+2} \left| \widehat{g}(\eta-\frac{k}{|k|}s) \right|^2 \, ds \right)^{\frac{1}{2}} \\ &\lesssim \sqrt{K_4} \varepsilon \|g\|_{H^{\sigma_1+1}_M}, \end{split}$$

where the last inequality utilises the L^2 trace lemma (Lemma 2.2).

For the nonlinear term, the triangle inequality $\langle k, \eta \rangle^{\sigma_1} \lesssim \langle \ell, \ell s \rangle^{\sigma_1} + \langle k - \ell, \eta - \ell s \rangle^{\sigma_1}$ facilitates division into reaction and transport parts, yielding $\mathcal{NL}^{(B5)} \lesssim \mathcal{R}^{(B5)} + T^{(B5)}$, where

$$\mathcal{R}^{(\mathrm{B5})} \lesssim \int_{0}^{t} \int_{\ell} \langle \ell, \ell s \rangle^{\sigma_{1}} |\widehat{w}(\ell)| |\widehat{\rho_{Q}}(s,\ell)| |\ell| |\eta - ks| \left| \widehat{W[P]}(s,k-\ell,\eta-\ell s) \right| d\ell \, ds$$
$$T^{(\mathrm{B5})} \lesssim \int_{0}^{t} \int_{\ell} |\widehat{w}(\ell)| |\widehat{\rho_{Q}}(s,\ell)| |\ell| |\eta - ks| \langle k-\ell,\eta-\ell s \rangle^{\sigma_{1}} \left| \widehat{W[P]}(s,k-\ell,\eta-\ell s) \right| d\ell \, ds.$$

For the reaction term, control of $|\eta - ks| \leq \langle s \rangle \langle k - \ell, \eta - \ell s \rangle$ and $|\widehat{w}(\ell)| \leq 1$ allows application of Cauchy–Schwarz in t, followed by bootstrap assumptions (B4) and (B5):

$$\begin{aligned} \mathcal{R}^{(\mathrm{B5})} &\lesssim \left\| |\ell|^{\frac{1}{2}} \langle \ell, \ell t \rangle^{\sigma_2} \widehat{\rho_Q}(t,\ell) \right\|_{L^2_t} \\ & \qquad \times \int_{\ell} \left(\int_0^t \langle s \rangle^2 \langle \ell, \ell s \rangle^{-2(\sigma_2 - \sigma_1)} |\ell| \langle k - \ell, \eta - \ell s \rangle^2 \Big| \widehat{W[P]}(s,k-\ell,\eta-\ell s) \Big|^2 \, d\ell \, ds \right)^{\frac{1}{2}} d\ell, \\ &\lesssim \sqrt{K_4 K_5} \varepsilon^2 \int_{\ell} \left(\int_0^t \langle s \rangle^2 \frac{|\ell|}{\langle \ell, \ell s \rangle^{2(\sigma_2 - \sigma_1)} \langle k - \ell, \eta - \ell s \rangle^{2\sigma_1 - 2}} \, ds \right)^{\frac{1}{2}} d\ell. \end{aligned}$$

Let $2(\sigma_2 - \sigma_1) = j_1$ and $2\sigma_1 - 2 = j_2$. Thus, $\langle \ell, \ell s \rangle^{2(\sigma_2 - \sigma_1)} \ge \langle \ell s \rangle^{2j_1}$ and $\langle k - \ell, \eta - \ell s \rangle^{2\sigma_1 - 2} \ge \langle k - \ell \rangle^{2j_2}$, yielding:

$$\begin{split} \mathcal{R}^{(\mathrm{B5})} &\lesssim \sqrt{K_4 K_5} \varepsilon^2 \int_{\ell} \frac{1}{\langle k - \ell \rangle^{j_2}} \left(\int_0^t |\ell| \frac{\langle s \rangle^2}{\langle \ell s \rangle^{2j_1}} \, ds \right)^{\frac{1}{2}} d\ell \\ &\lesssim \sqrt{K_4 K_5} \varepsilon^2 \int_{\ell} \frac{1}{\langle k - \ell \rangle^{j_2}} \left(\int_0^t \frac{\langle \frac{s}{|\ell|} \rangle^2}{\langle s \rangle^{j_1}} \, ds \right)^{\frac{1}{2}} d\ell \\ &\lesssim \sqrt{K_4 K_5} \varepsilon^2 \int_{\ell} \frac{\max\left\{ 1, |\ell|^{-1} \right\}}{\langle \ell \rangle^{j_2}} d\ell, \end{split}$$

where the s-integral converges if $j_1 > 3$ (requiring $\sigma_2 - \sigma_1 > 3/2$) and the ℓ -integral converges if $j_2 > d$ (requiring $\sigma_1 > d/2 + 1$).

For the transport term, control of $|\eta - ks| \leq \langle s \rangle \langle k - \ell, \eta - \ell s \rangle$ and $|\widehat{w}(\ell)| \leq 1$, followed by Cauchy–Schwarz in ℓ and application of bootstrap assumption (B5), leads to

By symmetry, the following relationship holds:

$$\sup_{k} \int_{\ell} \frac{|\ell|^2}{\langle \ell, \ell s \rangle^{2\sigma_1} |k-\ell|^{2\delta}} \, d\ell = \int_{\ell} \frac{|\ell|^{2-2\delta}}{\langle \ell, \ell s \rangle^{2\sigma_1}} \, d\ell \lesssim \frac{1}{\langle s \rangle^{d+2-2\delta}} \int_{\ell} \frac{|\ell|^{2-2\delta}}{\langle \ell \rangle^{2\sigma_1}} \, d\ell \lesssim \frac{1}{\langle s \rangle^{d+2-2\delta}},$$

where the change of variable $\ell \mapsto \ell/s$ has been applied. The integral converges if $\sigma_1 > d/2 + 1 - \delta$. For the integral involving $\widehat{W[P]}$, the Sobolev embedding is employed to apply bootstrap assumption (B3),

$$\begin{split} \left\| |k-\ell|^{\delta} \langle k-\ell, \eta-\ell s \rangle^{\sigma_{1}+1} \widehat{W[P]}(s,k-\ell,\eta-\ell s) \right\|_{L^{\infty}_{\eta}L^{2}_{\ell}} &= \left\| |k|^{\delta} \langle k,\eta \rangle^{\sigma_{3}} \widehat{W[P]}(s,k,\eta) \right\|_{L^{\infty}_{\eta}L^{2}_{k}} \\ &\lesssim \sum_{|\alpha| \leq M} \left\| D^{\alpha}_{\eta} \left(|k|^{\delta} \langle k,\eta \rangle^{\sigma_{3}} \widehat{W[P]}(s,k,\eta) \right) \right\|_{L^{2}_{k,\eta}} \\ &\lesssim \sqrt{K_{3}} \varepsilon. \end{split}$$

Therefore,

$$T^{(\mathrm{B5})} \lesssim \sqrt{K_3 K_5} \varepsilon^2 \int_0^t \frac{1}{\langle s \rangle^{\frac{d}{2} - \delta}} \, ds \lesssim \sqrt{K_3 K_5} \varepsilon^2.$$

4.1.1. Conclusion of bootstrap improvement (I5). Under the bootstrap assumptions (B3), (B4) and (B5), it has been shown that there exists a constant $\widetilde{K} = \widetilde{K}(d, g, w, \sigma_i, Q_{\rm in})$ (and independent of K_i and $0 < \hbar \leq 1$), such that

$$\langle k,\eta\rangle^{\sigma_1} \left| \int_p e^{-i\hbar\eta \cdot p} \widehat{P}\left(\frac{k}{2} + p, \frac{k}{2} - p\right) dp \right| \leq \widetilde{K} \left(\varepsilon + \sqrt{K_4}\varepsilon + \left(\sqrt{K_4}K_5 + \sqrt{K_3}K_5\right)\varepsilon^2 \right).$$

Choose ε sufficiently small such that $\left(\sqrt{K_4K_5} + \sqrt{K_3K_5}\right)\varepsilon \leq 1$. Then

$$\langle k,\eta\rangle^{\sigma_1} \left| \int_p e^{-i\hbar\eta \cdot p} \widehat{P}\left(\frac{k}{2} + p, \frac{k}{2} - p\right) dp \right| \leq \widetilde{K}\left(2 + \sqrt{K_4}\right) \varepsilon$$

Thus, by fixing

$$\sqrt{K_5} := \frac{\widetilde{K}(2 + \sqrt{K_4})}{\sqrt{2}},$$

the improvement (I5) of (B5) is obtained:

$$\langle k,\eta\rangle^{\sigma_1} \left| \int_p e^{-i\hbar\eta \cdot p} \widehat{P}\left(\frac{k}{2}+p,\frac{k}{2}-p\right) dp \right| \leq \sqrt{2}\sqrt{K_5}\varepsilon.$$

4.2. Bootstrap improvement (I1). In this section, the improvement of (B1) is shown. The work is conducted on the Wigner transform, using that

$$\hbar^{-d} \sum_{|\alpha| \le M} \|\boldsymbol{\xi}^{\alpha} \langle \boldsymbol{\nabla}_{x}, \boldsymbol{\nabla}_{\xi} \rangle^{\sigma_{4}} \langle t \boldsymbol{\nabla}_{x}, \boldsymbol{\nabla}_{\xi} \rangle P(t) \|_{L^{2}_{x,y}}^{2} = (2\pi)^{d} \| \langle t \nabla_{z}, \nabla_{v} \rangle W[P] \|_{H^{\sigma_{4}}_{M}}^{2}$$

by Lemma 1.11.

4.2.1. Estimate on the velocity derivative. Let $|\alpha| \leq M$ be a multi-index. An energy estimate yields

$$\begin{split} \frac{1}{2} \frac{d}{dt} \| \langle \nabla_v \rangle (v^{\alpha} W[P]) \|_{H^{\sigma_4}}^2 = &\operatorname{Re} \int_{k,\eta} \langle \eta \rangle \langle k,\eta \rangle^{\sigma_4} \overline{D_{\eta}^{\alpha} \widehat{W[P]}(t,k,\eta)} \langle \eta \rangle \langle k,\eta \rangle^{\sigma_4} D_{\eta}^{\alpha} \widehat{W[P_{\mathrm{L}}]}(t,k,\eta) \, d\eta dk \\ &+ \operatorname{Re} \int_{k,\eta} \langle \eta \rangle \langle k,\eta \rangle^{\sigma_4} \overline{D_{\eta}^{\alpha} \widehat{W[P]}(t,k,\eta)} \langle \eta \rangle \langle k,\eta \rangle^{\sigma_4} D_{\eta}^{\alpha} \widehat{W[P_{\mathrm{NL}}]}(t,k,\eta) \, d\eta dk \\ &= \mathcal{L}^{(\mathrm{B1})} + \mathcal{NL}^{(\mathrm{B1})}, \end{split}$$

where $\widehat{W[P_{\rm L}]}(t,k,\eta)$ and $\widehat{W[P_{\rm NL}]}(t,k,\eta)$ are defined in (5).

For $\mathcal{L}^{(B1)}$, Lemma 1.13 is used to separate the highest-order and remainder terms arising from the v^{α} -weights,

$$\begin{split} \left| \mathcal{L}^{(\mathrm{B1})} \right| \lesssim \int_{k,\eta} \langle \eta \rangle \langle k,\eta \rangle^{\sigma_4} \left| D_{\eta}^{\alpha} \widehat{W[P]}(t,k,\eta) \right| \langle \eta \rangle \langle k,\eta \rangle^{\sigma_4} |k| |\widehat{w}(k)| |\widehat{\rho_Q}(t,k)| |\eta - kt| \left| D_{\eta}^{\alpha} \widehat{g}(\eta - kt) \right| d\eta dk \\ &+ \sum_{|\beta| \le 1} \int_{k,\eta} \langle \eta \rangle \langle k,\eta \rangle^{\sigma_4} \left| D_{\eta}^{\alpha} \widehat{W[P]}(t,k,\eta) \right| \langle \eta \rangle \langle k,\eta \rangle^{\sigma_4} \langle \hbar k \rangle^{|\alpha| - 1} |k| |\widehat{w}(k)| |\widehat{\rho_Q}(t,k)| \left| D_{\eta}^{\beta} \widehat{g}(\eta - kt) \right| d\eta dk \\ &= \mathcal{L}_0^{(\mathrm{B1})} + \mathcal{L}_M^{(\mathrm{B1})}. \end{split}$$

For $\mathcal{L}_0^{(B1)}$, the triangle inequalities $\langle \eta \rangle \lesssim \langle kt \rangle \langle \eta - kt \rangle$ and $\langle k, \eta \rangle^{\sigma_4} \lesssim \langle k, kt \rangle^{\sigma_4} \langle \eta - kt \rangle^{\sigma_4}$ are used, along with $\langle kt \rangle |k| |\widehat{w}(k)| \lesssim \langle t \rangle |k|^{\frac{1}{2}}$, are applied to estimate

$$\begin{split} \left| \mathcal{L}_{0}^{(\mathrm{B1})} \right| &\lesssim \langle t \rangle \int_{k,\eta} \langle \eta \rangle \langle k,\eta \rangle^{\sigma_{4}} \left| D_{\eta}^{\alpha} \widehat{W[P]}(t,k,\eta) \right| |k|^{\frac{1}{2}} \langle k,kt \rangle^{\sigma_{4}} |\widehat{\rho_{Q}}(t,k)| \langle \eta-kt \rangle^{\sigma_{4}+2} \left| D_{\eta}^{\alpha} \widehat{g}(\eta-kt) \right| d\eta dk \\ &\lesssim \langle t \rangle \| \langle \nabla_{v} \rangle W[P] \|_{H_{M}^{\sigma_{4}}} \left\| |k|^{\frac{1}{2}} \langle k,kt \rangle^{\sigma_{4}} \widehat{\rho_{Q}}(t,k) \right\|_{L_{k}^{2}} \|g\|_{H_{M}^{\sigma_{4}+2}}. \end{split}$$

Similarly, for $\mathcal{L}_{M}^{(\mathrm{B1})}$,

$$\langle kt\rangle \langle \hbar k\rangle^{|\alpha|-1} |k| |\widehat{w}(k)| \lesssim \langle t\rangle |k| \langle k\rangle^M |\widehat{w}(k)| \lesssim \langle t\rangle |k|^{\frac{1}{2}}$$

if $|\widehat{w}(k)| \leq \langle k \rangle^{-M-1/2}$ and $0 < \hbar \leq 1$. Then Cauchy–Schwarz implies

$$\mathcal{L}_{M}^{(\mathrm{B1})} \lesssim \langle t \rangle \| \langle \nabla_{v} \rangle W[P] \|_{H_{M}^{\sigma_{4}}} \left\| |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_{4}} \widehat{\rho_{Q}}(t,k) \right\|_{L_{k}^{2}} \|g\|_{H_{M}^{\sigma_{4}+1}}.$$

For $\mathcal{NL}^{(B1)}$, Lemma 1.13 is first used to extract the highest-order term from the rest, $\mathcal{NL}^{(B1)} = \mathcal{NL}_0^{(B1)} + \mathcal{NL}_M^{(B1)}$ where

$$\begin{split} \mathcal{NL}_{0}^{(\mathrm{B1})} &= \mathrm{Im} \int_{k,\eta} \langle \eta \rangle \langle k,\eta \rangle^{\sigma_{4}} \overline{D_{\eta}^{\alpha} \widehat{W[P]}(t,k,\eta)} \langle \eta \rangle \langle k,\eta \rangle^{\sigma_{4}} \\ & \times \left[\frac{\hbar^{-1}}{(2\pi)^{d}} \int_{l} \widehat{w}(l) \widehat{\rho_{Q}}(t,\ell) \left(e^{-i\frac{\hbar}{2}\ell \cdot (\eta-kt)} - e^{i\frac{\hbar}{2}\ell \cdot (\eta-kt)} \right) D_{\eta}^{\alpha} \widehat{W[P]}(t,k-\ell,\eta-\ell t) \, d\ell \right] dk d\eta \end{split}$$

and a remainder term which is controlled by

$$\begin{split} \left| \mathcal{NL}_{M}^{(\mathrm{B1})} \right| &\lesssim \sum_{|\beta| \leq |\alpha| - 1} \int_{k,\eta,\ell} \langle \eta \rangle \langle k,\eta \rangle^{\sigma_{4}} \left| D_{\eta}^{\alpha} \widehat{W[P]}(t,k,\eta) \right| \langle \eta \rangle \langle k,\eta \rangle^{\sigma_{4}} \\ &\times \left| \widehat{w}(\ell) \right| \left| \ell \right| \langle \ell \rangle^{|\alpha| - 1} \left| \widehat{\rho_{Q}}(t,\ell) \right| \left| D_{\eta}^{\beta} \widehat{W[P]}(t,k-\ell,\eta-\ell t) \right| d\ell dk d\eta \end{split}$$

 $\text{if } 0 < \hbar \leq 1.$

To bound the term $\mathcal{NL}_0^{(B1)}$, an integral which evaluates to zero is "subtracted":

$$\mathcal{NL}_{0}^{(\mathrm{B1})} = \mathrm{Im} \int_{k,\eta} \langle \eta \rangle \langle k,\eta \rangle^{\sigma_{4}} \overline{D_{\eta}^{\alpha} \widehat{W[P]}(t,k,\eta)} \left(\langle \eta \rangle \langle k,\eta \rangle^{\sigma_{4}} - \langle \eta - \ell t \rangle \langle k - \ell,\eta - \ell t \rangle^{\sigma_{4}} \right) \\ \times \left[\frac{\hbar^{-1}}{(2\pi)^{d}} \int_{\ell} \widehat{w}(\ell) \widehat{\rho_{Q}}(t,\ell) \left(e^{-i\frac{\hbar}{2}\ell \cdot (\eta - kt)} - e^{i\frac{\hbar}{2}\ell \cdot (\eta - kt)} \right) D_{\eta}^{\alpha} \widehat{W[P]}(t,k-\ell,\eta-\ell t) \, d\ell \right] dk d\eta.$$
⁽²⁹⁾

The integral that has been subtracted is now shown to be zero. On the physical side, by computing the inverse Fourier transforms of $\widehat{W[P_{\rm L}]}$ and $\widehat{W[P_{\rm NL}]}$, defined in (5), the Wigner transform of P satisfies the equation:

$$\begin{cases} \partial_{t} W[P](t,z,v) = W[P_{\rm L}](t,z,v) + W[P_{\rm NL}](t,z,v) \\ W[P_{\rm L}](t,z,v) = -\frac{i\hbar^{-1}}{(2\pi)^{d}} \int_{k} e^{ik \cdot (z+vt)} \widehat{w}(k) \widehat{\rho_{Q}}(t,k) \left(g(v-\frac{\hbar}{2}k) - g(v+\frac{\hbar}{2}k)\right) dk, \\ W[P_{\rm NL}](t,z,v) = -\frac{i\hbar^{-1}}{(2\pi)^{d}} \int_{\ell} e^{i\ell \cdot (z+vt)} \widehat{w}(\ell) \widehat{\rho_{Q}}(t,\ell) \left(W[P](t,z+\frac{\hbar}{2}\ell t,v-\frac{\hbar}{2}\ell) - W[P](t,z-\frac{\hbar}{2}\ell t,v+\frac{\hbar}{2}\ell)\right) d\ell. \end{cases}$$
(30)

Then, by applying Plancherel's theorem and utilising (5) and (30), the integral subtracted from (29) can be expressed on the physical side as:

$$\begin{split} \int_{k,\eta} \langle \eta \rangle \langle k,\eta \rangle^{\sigma_4} \overline{D_{\eta}^{\alpha} W[P]}(t,k,\eta) \langle \eta - \ell t \rangle \langle k - \ell,\eta - \ell t \rangle^{\sigma_4} \\ & \times \left[\frac{\hbar^{-1}}{(2\pi)^d} \int_{\ell} \widehat{w}(\ell) \widehat{\rho_Q}(t,\ell) \left(e^{-i\frac{\hbar}{2}\ell \cdot (\eta - kt)} - e^{i\frac{\hbar}{2}\ell \cdot (\eta - kt)} \right) D_{\eta}^{\alpha} \widehat{W[P]}(t,k-\ell,\eta - \ell t) \, d\ell \right] dk d\eta \\ &= \int_{z,v} \overline{\langle \nabla_v \rangle \langle \nabla \rangle^{\sigma_4} \left((-iv)^{\alpha} W[P] \right) (t,z,v)} \hbar^{-1} \int_{\ell} e^{i\ell \cdot (z + vt)} \widehat{w}(\ell) \widehat{\rho_Q}(t,\ell) \\ & \times \left[\left(\langle \nabla_v \rangle \langle \nabla \rangle^{\sigma_4} ((-iv)^{\alpha} W[P] \right) \right) (t,z + \frac{\hbar}{2}\ell t, v - \frac{\hbar}{2}\ell \right) - \left(\langle \nabla_v \rangle \langle \nabla \rangle^{\sigma_4} ((-iv)^{\alpha} W[P]) \right) (t,z - \frac{\hbar}{2}\ell t, v + \frac{\hbar}{2}\ell) \right] d\ell dz dv. \end{split}$$

where $\langle \nabla \rangle = \langle \nabla_z, \nabla_v \rangle$ in an abuse of notation. To see that this integral has zero imaginary part, observe that it is of the form

$$I := \hbar^{-1} \int_{z,v,\ell} \overline{f(z,v)} e^{i\ell \cdot (z+tv)} \widehat{\rho_Q}(t,\ell) \widehat{w}(\ell) \left[f(z - \frac{\hbar}{2}\ell t, v + \frac{\hbar}{2}\ell) - f(z + \frac{\hbar}{2}\ell t, v - \frac{\hbar}{2}\ell) \right] d\ell dz dv$$

where $f(z,v) = \langle \nabla_v \rangle \langle \nabla_z, \nabla_v \rangle^{\sigma_4} ((-iv)^{\alpha} W[P])(t,z,v)$. By making the change of variables $z \mapsto z + \frac{\hbar}{2} \ell t$, $v \mapsto v - \frac{\hbar}{2} \ell$ in the first term, and $z \mapsto z - \frac{\hbar}{2} \ell t$, $v \mapsto v + \frac{\hbar}{2} \ell$ in the second,

$$\begin{split} I &= \hbar^{-1} \int_{z,v,\ell} \overline{f(z,v)} e^{i\ell \cdot (z+tv)} \widehat{\rho_Q}(t,\ell) \widehat{w}(\ell) \left[f(z - \frac{\hbar}{2}\ell t, v + \frac{\hbar}{2}\ell) - f(z + \frac{\hbar}{2}\ell t, v - \frac{\hbar}{2}\ell) \right] d\ell dz dv \\ &= \hbar^{-1} \int_{z,v,\ell} \overline{\left(f(z + \frac{\hbar}{2}t\ell, v - \frac{\hbar}{2}\ell) - f(z - \frac{\hbar}{2}\ell t, v + \frac{\hbar}{2}\ell) \right)} e^{i\ell \cdot (z+tv)} \widehat{\rho_Q}(t,\ell) \widehat{w}(\ell) f(z,v) d\ell dv dz \\ &= \hbar^{-1} \int_{z,v,\ell} \overline{\left(f(z - \frac{\hbar}{2}t\ell, v + \frac{\hbar}{2}\ell) - f(z + \frac{\hbar}{2}\ell t, v - \frac{\hbar}{2}\ell) \right)} e^{-i\ell \cdot (z+tv)} \widehat{\rho_Q}(t,-\ell) \widehat{w}(-\ell) f(z,v) d\ell dv dz \\ &= \hbar^{-1} \overline{\int_{z,v,\ell} \overline{f(z,v)} e^{i\ell \cdot (z+tv)} \widehat{\rho_Q}(t,\ell) \widehat{w}(\ell) \left[f(z - \frac{\hbar}{2}\ell t, v + \frac{\hbar}{2}\ell) - f(z + \frac{\hbar}{2}\ell t, v - \frac{\hbar}{2}\ell) \right] d\ell dz dv} \\ &= \overline{h}. \end{split}$$

where in penultimate step it was used that w and ρ_Q are real (Q is self-adjoint), which implies that $\overline{\hat{w}(-\ell)} = \widehat{w}(\ell)$ and $\overline{\widehat{\rho_Q}(t,-\ell)} = \widehat{\rho_Q}(t,\ell)$. Hence Im I = 0.

Therefore, $\mathcal{NL}_{0}^{(B1)}$ can be controlled using (29),

$$\begin{split} \left| \mathcal{NL}_{0}^{(\mathrm{B1})} \right| \lesssim \int_{k,\eta,\ell} \langle \eta \rangle \langle k,\eta \rangle^{\sigma_{4}} \left| D_{\eta}^{\alpha} \widehat{W[P]}(t,k,\eta) \right| |\langle \eta \rangle \langle k,\eta \rangle^{\sigma_{4}} - \langle \eta - \ell t \rangle \langle k - \ell,\eta - \ell t \rangle^{\sigma_{4}} | \\ \times |\widehat{w}(\ell)| |\ell| |\widehat{\rho_{Q}}(t,\ell)| |\eta - kt| \left| D_{\eta}^{\alpha} \widehat{W[P]}(t,k-\ell,\eta - \ell t) \right| d\ell dk d\eta. \end{split}$$

By the Mean Value Theorem,

$$\begin{split} |\langle \eta \rangle \langle k, \eta \rangle^{\sigma_4} - \langle \eta - \ell t \rangle \langle k - \ell, \eta - \ell t \rangle^{\sigma_4} | \\ &\leq |\langle \eta \rangle - \langle \eta - \ell t \rangle |\langle k, \eta \rangle^{\sigma_4} + \langle \eta - \ell t \rangle |\langle k, \eta \rangle^{\sigma_4} - \langle k - \ell, \eta - \ell t \rangle^{\sigma_4} | \\ &\lesssim \langle \ell t \rangle \langle k, \eta \rangle^{\sigma_4} + \langle \eta - \ell t \rangle |\ell, \ell t| \sup_{|\ell'| \leq |\ell|} \langle k - \ell', \eta - \ell' t \rangle^{\sigma_4 - 1} \\ &\lesssim \langle \ell t \rangle \left(\langle \ell, \ell t \rangle^{\sigma_4} + \langle k - \ell, \eta - \ell t \rangle^{\sigma_4} \right) + \langle \eta - \ell t \rangle |\ell, \ell t| \sup_{|\ell'| \leq |\ell|} \left(\langle k - \ell, \eta - \ell t \rangle^{\sigma_4 - 1} + \langle \ell - \ell', (\ell - \ell') t \rangle^{\sigma_4 - 1} \right) \\ &= \langle \ell t \rangle \langle \ell, \ell t \rangle^{\sigma_4} + \langle \eta - \ell t \rangle |\ell, \ell t| \langle \ell, \ell t \rangle^{\sigma_4 - 1} + \langle \ell t \rangle \langle k - \ell, \eta - \ell t \rangle^{\sigma_4} + \langle \eta - \ell t \rangle |\ell, \ell t| \langle k - \ell, \eta - \ell t \rangle^{\sigma_4 - 1}. \end{split}$$

This leads to four terms $\left|\mathcal{NL}_{0}^{(B1)}\right| \lesssim \mathcal{R}_{0,1}^{(B1)} + \mathcal{R}_{0,2}^{(B1)} + \mathcal{T}_{0,1}^{(B1)} + \mathcal{T}_{0,2}^{(B1)}$.

For $\mathcal{R}_{0,1}^{(B1)}$, $|\eta - kt| \lesssim \langle t \rangle \langle k - \ell, \eta - \ell t \rangle$ is utilised, along with the regularity of the potential \hat{w} which leads to $\langle \ell t \rangle |\ell| |\hat{w}(\ell)| \lesssim \langle t \rangle |\ell|^{\frac{1}{2}}$. Additionally, Lemma 4.1 (b) is applied:

$$\begin{split} \mathcal{R}_{0,1}^{(\mathrm{B1})} \lesssim & \int_{k,\eta,\ell} \langle \eta \rangle \langle k,\eta \rangle^{\sigma_4} \left| D_{\eta}^{\alpha} \widehat{W[P]}(t,k,\eta) \right| \langle \ell t \rangle \langle \ell,\ell t \rangle^{\sigma_4} | \widehat{w}(\ell) | |\ell| | \widehat{\rho_Q}(t,\ell) | \\ & \times |\eta - kt| \left| D_{\eta}^{\alpha} \widehat{W[P]}(t,k-\ell,\eta-\ell t) \right| d\ell dk d\eta \\ \lesssim \langle t \rangle^2 \int_{k,\eta,\ell} \langle \eta \rangle \langle k,\eta \rangle^{\sigma_4} \left| D_{\eta}^{\alpha} \widehat{W[P]}(t,k,\eta) \right| \langle \ell,\ell t \rangle^{\sigma_4} |\ell|^{\frac{1}{2}} | \widehat{\rho_Q}(t,\ell) | \\ & \times \langle k-\ell,\eta-\ell t \rangle \left| D_{\eta}^{\alpha} \widehat{W[P]}(t,k-\ell,\eta-\ell t) \right| d\ell dk d\eta \\ \lesssim \langle t \rangle^2 \| \langle \nabla_v \rangle W[P] \|_{H_{M}^{\sigma_4}} \left\| |k|^{\frac{1}{2}} \langle k,kt \rangle^{\sigma_4} \widehat{\rho_Q}(t,k) \right\|_{L^2_k} \int_k \left\| \langle k,\eta \rangle D_{\eta}^{\alpha} \widehat{W[P]}(t,k,\eta) \right\|_{L^2_\eta} dk \\ \lesssim \langle t \rangle^2 \| \langle \nabla_v \rangle W[P] \|_{H_{M}^{\sigma_4}} \left\| |k|^{\frac{1}{2}} \langle k,kt \rangle^{\sigma_4} \widehat{\rho_Q}(t,k) \right\|_{L^2_k} \left\| |k|^{\delta} \langle k \rangle^d \langle k,\eta \rangle D_{\eta}^{\alpha} \widehat{W[P]}(t,k,\eta) \right\|_{L^2_{k,\eta}} \\ \lesssim \sqrt{K_3} \varepsilon \langle t \rangle^2 \| \langle \nabla_v \rangle W[P] \|_{H_{M}^{\sigma_4}} \left\| |k|^{\frac{1}{2}} \langle k,kt \rangle^{\sigma_4} \widehat{\rho_Q}(t,k) \right\|_{L^2_k} \end{split}$$

if $\sigma_3 \ge d+1$ (using bootstrap estimate (B3)). The second reaction term $\mathcal{R}_{0,2}^{(B1)}$ is controlled similarly by

$$\mathcal{R}_{0,2}^{(\mathrm{B1})} \lesssim \sqrt{K_3} \varepsilon \langle t \rangle \| \langle \nabla_v \rangle W[P] \|_{H_M^{\sigma_4}} \left\| |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_4} \widehat{\rho_Q}(t,k) \right\|_{L_k^2}.$$

For the transport terms, the estimate $|\eta - kt| \leq \langle t(k - \ell), \eta - \ell t \rangle$ is used, along with Lemma 4.1 (a) and the bootstrap estimate (B5):

$$\begin{split} \mathcal{T}_{0,1}^{(\mathrm{B1})} \lesssim & \int_{k,\eta,\ell} \langle \eta \rangle \langle k,\eta \rangle^{\sigma_4} \left| D_{\eta}^{\alpha} \widehat{W[P]}(t,k,\eta) \right| \langle \ell t \rangle \langle k-\ell,\eta-\ell t \rangle^{\sigma_4} \\ & \times |\widehat{w}(\ell)| |\ell| |\widehat{\rho_Q}(t,\ell)| |\eta-kt| \left| D_{\eta}^{\alpha} \widehat{W[P]}(t,k-\ell,\eta-\ell t) \right| d\ell dk d\eta \\ \lesssim \| \langle \nabla_v \rangle W[P] \|_{H_M^{\sigma_4}} \| \langle t \nabla_z, \nabla_v \rangle W[P] \|_{H_M^{\sigma_4}} \int_l \langle l t \rangle |\ell| |\widehat{\rho_Q}(t,\ell)| |\widehat{w}(\ell)| d\ell \\ \lesssim \frac{\sqrt{K_5} \varepsilon}{\langle t \rangle^{d+1}} \| \langle \nabla_v \rangle W[P] \|_{H_M^{\sigma_4}} \| \langle t \nabla_z, \nabla_v \rangle W[P] \|_{H_M^{\sigma_4}} \end{split}$$

if $\sigma_1 > d+2$. Proceeding identically, the second transport term $\mathcal{T}_{0,2}^{(B_1)}$ is controlled by

$$\mathcal{T}_{0,1}^{(B1)} \lesssim \frac{\sqrt{K_5}\varepsilon}{\langle t \rangle^{d+1}} \| \langle \nabla_v \rangle W[P] \|_{H_M^{\sigma_4}} \| \langle t \nabla_z, \nabla_v \rangle W[P] \|_{H_M^{\sigma_4}}.$$

The remainder $\mathcal{NL}_M^{(B1)}$ defined above remains to be dealt with. The weight $\langle \eta \rangle \langle k, \eta \rangle^{\sigma_4}$ is separated into transport and reaction components:

$$\langle \eta \rangle \langle k, \eta \rangle^{\sigma_4} \lesssim \left(\langle \ell t \rangle + \langle \eta - \ell t \rangle \right) \langle \ell, \ell t \rangle^{\sigma_4} + \left(\langle \ell t \rangle + \langle \eta - \ell t \rangle \right) \langle k - \ell, \eta - \ell t \rangle^{\sigma_4},$$

resulting in four terms: $\left|\mathcal{NL}_{M}^{(\text{B1})}\right| \lesssim \mathcal{R}_{M,1}^{(\text{B1})} + \mathcal{R}_{M,2}^{(\text{B1})} + \mathcal{T}_{M,1}^{(\text{B1})} + \mathcal{T}_{M,2}^{(\text{B1})}$. For the reaction terms, if $|\widehat{w}(\ell)| \lesssim \langle \ell \rangle^{-M-1/2}$, then proceeding as in the estimate $\mathcal{R}_{0,1}^{(\text{B1})}$, yields

$$\mathcal{R}_{M,1}^{(\mathrm{B1})} + \mathcal{R}_{M,2}^{(\mathrm{B1})} \lesssim \sqrt{K_3} \varepsilon \langle t \rangle \| \langle \nabla_v \rangle W[P] \|_{H_M^{\sigma_4}} \left\| |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_4} \widehat{\rho_Q}(t,k) \right\|_{L_k^2}$$

Finally, the transport terms are controlled similarly to $\mathcal{T}_{0,1}^{(B_1)}$ and $\mathcal{T}_{0,2}^{(B_1)}$:

$$\mathcal{T}_{M,1}^{(\mathrm{B1})} + \mathcal{T}_{M,2}^{(\mathrm{B1})} \lesssim \frac{\sqrt{K_5\varepsilon}}{\langle t \rangle^{d+1}} \| \langle \nabla_v \rangle W[P](t) \|_{H_M^{\sigma_4}}^2.$$

4.2.2. Conclusion of the velocity derivative. The analysis has demonstrated that

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \| \langle \nabla_v \rangle W[P](t) \|_{H^{\sigma_4}_M}^2 &\lesssim \langle t \rangle \| \langle \nabla_v \rangle W[P](t) \|_{H^{\sigma_4}} \left\| |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_4} \widehat{\rho_Q}(t,k) \right\|_{L^2_k} \\ &+ \sqrt{K_3} \varepsilon \langle t \rangle^2 \| \langle \nabla_v \rangle W[P](t) \|_{H^{\sigma_4}_M} \left\| |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_4} \widehat{\rho_Q}(t,k) \right\|_{L^2_k} \\ &+ \frac{\sqrt{K_5} \varepsilon}{\langle t \rangle^{d+1}} \| \langle \nabla_v \rangle W[P](t) \|_{H^{\sigma_4}_M} \| \langle t \nabla_z, \nabla_v \rangle W[P](t) \|_{H^{\sigma_4}_M}. \end{aligned}$$

This implies that

$$\frac{d}{dt} \| \langle \nabla_v \rangle W[P](t) \|_{H^{\sigma_4}_M} \lesssim \left(\langle t \rangle + \sqrt{K_3} \varepsilon \langle t \rangle^2 \right) \left\| |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_4} \widehat{\rho_Q}(t,k) \right\|_{L^2_k} + \frac{\sqrt{K_5} \varepsilon}{\langle t \rangle^{d+1}} \| \langle t \nabla_z, \nabla_v \rangle W[P](t) \|_{H^{\sigma_4}_M}$$

Integration, combined with the bootstrap assumptions (B1) and (B2), yields

$$\begin{split} \|\langle \nabla_v \rangle W[P](t)\|_{H_M^{\sigma_4}} \\ \lesssim \int_0^t \left(\langle s \rangle + \sqrt{K_3} \varepsilon \langle s \rangle^2 \right) \left\| |k|^{\frac{1}{2}} \langle k, ks \rangle^{\sigma_4} \widehat{\rho_Q}(s, k) \right\|_{L_k^2} ds + \int_0^t \frac{\sqrt{K_5} \varepsilon}{\langle s \rangle^{d+1}} \|\langle s \nabla_z, \nabla_v \rangle W[P](s)\|_{H_M^{\sigma_4}} ds \\ \lesssim \left(\int_0^t \left(\langle s \rangle + \sqrt{K_3} \varepsilon \langle s \rangle^2 \right)^2 ds \right)^{\frac{1}{2}} \left\| |k|^{\frac{1}{2}} \langle k, ks \rangle^{\sigma_4} \widehat{\rho_Q}(s, k) \right\|_{L_{t,k}^2} + \sqrt{K_1 K_5} \varepsilon^2 \int_0^t \frac{\langle s \rangle^{d-\frac{1}{2}}}{\langle s \rangle^{d+1}} ds \\ \lesssim \sqrt{K_2} \varepsilon \left(1 + \sqrt{K_3} \varepsilon \langle t \rangle \right) \langle t \rangle^{\frac{3}{2}} + \sqrt{K_1 K_5} \varepsilon^2. \end{split}$$

Thus, there exists $\widetilde{K}_V = \widetilde{K}_V(d, g, w, \sigma_i, Q_{\rm in})$, such that

$$\hbar^{-d} \sum_{|\alpha| \le M} \|\boldsymbol{\xi}^{\alpha} \langle \boldsymbol{\nabla}_{x}, \boldsymbol{\nabla}_{\xi} \rangle^{\sigma_{4}} \langle \boldsymbol{\nabla}_{\xi} \rangle P(t)\|_{L^{2}_{x,y}}^{2} \le \widetilde{K}_{V} \varepsilon^{2} \left[\sqrt{K_{2}} + \left(\sqrt{K_{2}K_{3}} + \sqrt{K_{1}K_{5}} \varepsilon \right) \right]^{2} \langle t \rangle^{5}.$$
(31)

4.2.3. Estimate on the spatial derivative. Let $|\alpha| \leq M$ be a multi-index. An energy estimate yields

$$\begin{split} \frac{1}{2} \frac{d}{dt} \| \langle \nabla_z \rangle v^{\alpha} W[P] \|_{H^{\sigma_4}}^2 = &\operatorname{Re} \int_{k,\eta} \langle k \rangle \langle k,\eta \rangle^{\sigma_4} \overline{D_{\eta}^{\alpha} \widehat{W[P]}(t,k,\eta)} \langle k \rangle \langle k,\eta \rangle^{\sigma_4} D_{\eta}^{\alpha} \widehat{W[P_{\mathrm{L}}]}(t,k,\eta) \, d\eta dk \\ &+ \operatorname{Re} \int_{k,\eta} \langle k \rangle \langle k,\eta \rangle^{\sigma_4} \overline{D_{\eta}^{\alpha} \widehat{W[P]}(t,k,\eta)} \langle k \rangle \langle k,\eta \rangle^{\sigma_4} D_{\eta}^{\alpha} \widehat{W[P_{\mathrm{NL}}]}(t,k,\eta) \, d\eta dk \\ &= L^{(B1),Z} + NL^{(B1),Z}. \end{split}$$

For $L^{(B1),Z}$, Lemma 1.13 is used to control $\widehat{W[P_L]}$:

$$\begin{split} \mathcal{L}^{\,(\mathrm{B1})} &\lesssim \int_{k,\eta} \langle k \rangle \langle k,\eta \rangle^{\sigma_4} \left| D_{\eta}^{\alpha} \widehat{W[P]}(t,k,\eta) \right| \langle k \rangle \langle k,\eta \rangle^{\sigma_4} |k| |\widehat{w}(k)| |\widehat{\rho_Q}(t,k)| |\eta - kt| \left| D_{\eta}^{\alpha} \widehat{g}(\eta - kt) \right| d\eta dk \\ &+ \sum_{|\beta| \leq |\alpha| - 1} \int_{k,\eta} \langle k \rangle \langle k,\eta \rangle^{\sigma_4} \left| D_{\eta}^{\alpha} \widehat{W[P]}(t,k,\eta) \right| \langle k \rangle \langle k,\eta \rangle^{\sigma_4} \langle k \rangle^{|\alpha| - 1} |k| |\widehat{w}(k)| |\widehat{\rho_Q}(t,k)| \left| D_{\eta}^{\beta} \widehat{g}(\eta - kt) \right| d\eta dk \\ &= \mathcal{L}_0^{\,(B1),Z} + \mathcal{L}_M^{\,(B1),Z}. \end{split}$$

For $\mathcal{L}_0^{(B1),Z}$, the inequality $\langle k,\eta\rangle^{\sigma_4} \lesssim \langle k,kt\rangle^{\sigma_4} \langle \eta-kt\rangle^{\sigma_4}$ is applied, along with $\langle k\rangle |k| |\widehat{w}(k)| \lesssim |k|^{\frac{1}{2}}$,

$$\begin{split} \mathcal{L}_{0}^{(B1),Z} &\lesssim \int_{k,\eta} \langle k \rangle \langle k,\eta \rangle^{\sigma_{4}} \Big| D_{\eta}^{\alpha} \widehat{W[P]}(t,k,\eta) \Big| |k|^{\frac{1}{2}} \langle k,kt \rangle^{\sigma_{4}} |\widehat{\rho_{Q}}(t,k)| \langle \eta-kt \rangle^{\sigma_{4}+1} \big| D_{\eta}^{\alpha} \widehat{g}(\eta-kt) \big| \, d\eta dk \\ &\lesssim \Big\| \langle \nabla_{z} \rangle \widehat{W[P]} \Big\|_{H^{\sigma_{4}}} \Big\| |k|^{\frac{1}{2}} \langle k,kt \rangle^{\sigma_{4}} \widehat{\rho_{Q}}(t,k) \Big\|_{L^{2}_{k}} \|g\|_{H^{\sigma_{4}+1}_{M}} \end{split}$$

upon performing Cauchy–Schwarz in η and then k. Similarly, for $\mathcal{L}_{M}^{(B1)}$, if $|\hat{w}(k)| \leq \langle k \rangle^{-M-1/2}$ and $0 < \hbar \leq 1$, then

$$\mathcal{L}_{M}^{(B1),Z} \lesssim \left\| \langle \nabla_{v} \rangle W[P] \right\|_{H^{\sigma_{4}}} \left\| |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_{4}} \widehat{\rho_{Q}}(t,k) \right\|_{L^{2}_{k}} \|g\|_{H^{\sigma_{4}}_{M}}.$$

For the nonlinear term, an identical approach to Section 4.2.1 is used to decompose $NL^{(B1),Z} = \mathcal{NL}_0^{(B1),Z} + \mathcal{NL}_M^{(B1),Z}$ where

$$\begin{split} \left| \mathcal{NL}_{0}^{(B1),Z} \right| \lesssim \int_{k,\eta,\ell} \langle k \rangle \langle k,\eta \rangle^{\sigma_{4}} \left| D_{\eta}^{\alpha} \widehat{W[P]}(t,k,\eta) \right| |\langle k \rangle \langle k,\eta \rangle^{\sigma_{4}} - \langle k-\ell \rangle \langle k-\ell,\eta-\ell t \rangle^{\sigma_{4}} | \\ \times |\widehat{w}(\ell)| |\ell| |\widehat{\rho_{Q}}(t,\ell)| |\eta-kt| \left| D_{\eta}^{\alpha} \widehat{W[P]}(t,k-\ell,\eta-\ell t) \right| d\ell dk d\eta \end{split}$$

and

$$\begin{split} \left| \mathcal{NL}_{M}^{(B1),Z} \right| \lesssim \sum_{|\beta| \le |\alpha| - 1} \int_{k,\eta,\ell} \langle k \rangle \langle k,\eta \rangle^{\sigma_4} \left| D_{\eta}^{\alpha} \widehat{W[P]}(t,k,\eta) \right| \langle k \rangle \langle k,\eta \rangle^{\sigma_4} \\ & \times |\widehat{w}(\ell)| |\ell| \langle \ell \rangle^{|\alpha| - 1} |\widehat{\rho_Q}(t,\ell)| \left| D_{\eta}^{\beta} \widehat{W[P]}(t,k-\ell,\eta-\ell t) \right| d\ell dk d\eta. \end{split}$$

For $\mathcal{NL}_0^{(B1),Z}$, the Mean Value Theorem implies that

$$\begin{split} |\langle k \rangle \langle k, \eta \rangle^{\sigma_4} - \langle k - \ell \rangle \langle k - \ell, \eta - \ell t \rangle^{\sigma_4} | \\ \lesssim \langle \ell \rangle \langle \ell, \ell t \rangle^{\sigma_4} + \langle k - \ell \rangle |\ell, \ell t| \langle \ell, \ell t \rangle^{\sigma_4 - 1} + \langle \ell \rangle \langle k - \ell, \eta - \ell t \rangle^{\sigma_4} + \langle k - \ell \rangle |\ell, \ell t| \langle k - \ell, \eta - \ell t \rangle^{\sigma_4 - 1}, \end{split}$$

leading to four terms $\left|\mathcal{NL}_{0}^{(B1),Z}\right| \lesssim \mathcal{R}_{0,1}^{(B1),Z} + \mathcal{R}_{0,2}^{(B1),Z} + \mathcal{T}_{0,1}^{(B1),Z} + \mathcal{T}_{0,2}^{(B1),Z}$. These terms are similar to those in Section 4.2.1, and are controlled by the same estimates, except that $\mathcal{R}_{0,1}^{(B1),Z}$ leads to one less power of $\langle t \rangle$. In summary,

$$\left|\mathcal{NL}_{0}^{(B1),Z}\right| \lesssim \sqrt{K_{3}}\varepsilon\langle t\rangle \|\langle \nabla_{z}\rangle W[P]\|_{H_{M}^{\sigma_{4}}} \left\||k|^{\frac{1}{2}}\langle k,kt\rangle^{\sigma_{4}}\widehat{\rho_{Q}}(t,k)\right\|_{L_{k}^{2}} + \frac{\sqrt{K_{5}}\varepsilon}{\langle t\rangle^{d+1}} \|\langle \nabla_{z}\rangle W[P]\|_{H_{M}^{\sigma_{4}}} \|\langle t\nabla_{z},\nabla_{v}\rangle W[P]\|_{H_{M}^{\sigma_{4}}} \|\langle t\nabla_{v}\rangle W[P]\|_{H_{M}^{\sigma_{4$$

provided that $\sigma_3 > d + 2$. The same proof can be applied to $\mathcal{NL}_M^{(B1),Z}$ under the assumption $|\widehat{w}(\ell)| \leq \langle \ell \rangle^{-M-1/2}$.

4.2.4. Conclusion of the spatial derivative. Proceeding as in Section 4.2.2, the analysis demonstrates that

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \| \langle \nabla_z \rangle W[P](t) \|_{H^{\sigma_4}}^2 &\lesssim \| \langle \nabla_z \rangle W[P](t) \|_{H^{\sigma_4}} \left\| |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_4} \widehat{\rho_Q}(t,k) \right\|_{L^2_k} \\ &+ \sqrt{K_3} \varepsilon \langle t \rangle \| \langle \nabla_z \rangle W[P](t) \|_{H^{\sigma_4}_M} \left\| |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_4} \widehat{\rho_Q}(t,k) \right\|_{L^2_k} \\ &+ \frac{\sqrt{K_5} \varepsilon}{\langle t \rangle^{d+1}} \| \langle \nabla_z \rangle W[P](t) \|_{H^{\sigma_4}_M} \| \langle t \nabla_z, \nabla_v \rangle W[P](t) \|_{H^{\sigma_4}_M}. \end{aligned}$$

which implies that

$$\frac{d}{dt} \| \langle \nabla_z \rangle W[P](t) \|_{H^{\sigma_4}_M} \lesssim \left(1 + \sqrt{K_3} \varepsilon \langle t \rangle \right) \left\| |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_4} \widehat{\rho_Q}(t,k) \right\|_{L^2_k} + \frac{\sqrt{K_5} \varepsilon}{\langle t \rangle^{d+1}} \| \langle t \nabla_z, \nabla_v \rangle W[P](t) \|_{H^{\sigma_4}_M}.$$

Integrating and using the bootstrap assumptions (B1) and (B2),

$$\begin{split} \|\langle \nabla_z \rangle W[P](t)\|_{H^{\sigma_4}_M} \\ \lesssim \int_0^t \left(1 + \sqrt{K_3} \varepsilon \langle s \rangle \right) \left\| |k|^{\frac{1}{2}} \langle k, ks \rangle^{\sigma_4} \widehat{\rho_Q}(s, k) \right\|_{L^2_k} ds + \int_0^t \frac{\sqrt{K_5} \varepsilon}{\langle s \rangle^{d+1}} \|\langle s \nabla_z, \nabla_v \rangle W[P](s)\|_{H^{\sigma_4}_M} ds \\ \lesssim \left(\int_0^t \left(1 + \sqrt{K_3} \varepsilon \langle s \rangle \right)^2 ds \right)^{\frac{1}{2}} \left\| |k|^{\frac{1}{2}} \langle k, ks \rangle^{\sigma_4} \widehat{\rho_Q}(s, k) \right\|_{L^2_{t,k}} + \sqrt{K_1 K_5} \varepsilon^2 \int_0^t \frac{\langle s \rangle^{d-\frac{1}{2}}}{\langle s \rangle^{d+1}} ds \\ \lesssim \sqrt{K_2} \varepsilon \left(1 + \sqrt{K_3} \varepsilon \langle t \rangle \right) \langle t \rangle + \sqrt{K_1 K_5} \varepsilon^2. \end{split}$$

Hence there exists a positive constant $\widetilde{K}_Z = \widetilde{K}_Z(d, g, w, \sigma_i, Q_{in})$, such that

$$\hbar^{-d} \sum_{|\alpha| \le M} \left\| \boldsymbol{\xi}^{\alpha} \langle \boldsymbol{\nabla}_x, \boldsymbol{\nabla}_{\boldsymbol{\xi}} \rangle^{\sigma_4} \langle \boldsymbol{\nabla}_x \rangle P(t) \right\|_{L^2_{x,y}}^2 \le \widetilde{K}_Z \varepsilon^2 \left[\sqrt{K_2} + \varepsilon \left(\sqrt{K_2 K_3} + \sqrt{K_1 K_5} \right) \right]^2 \langle t \rangle^4.$$
(32)

4.2.5. Conclusion of bootstrap improvement (I1). By selecting $K_1 := 2 \max \left\{ \widetilde{K}_V, \widetilde{K}_Z \right\} \left(\sqrt{K_2} + 1 \right)^2$ and imposing $\varepsilon \leq (\sqrt{K_2 K_3} + \sqrt{K_1 K_5})^{-1}$, the following bounds are obtained from (31) and (32):

$$\hbar^{-d} \sum_{|\alpha| \le M} \left\| \boldsymbol{\xi}^{\alpha} \langle \boldsymbol{\nabla}_{x}, \boldsymbol{\nabla}_{\xi} \rangle^{\sigma_{4}} \langle \boldsymbol{\nabla}_{\xi} \rangle P(t) \right\|_{L^{2}_{x,y}}^{2} \le \frac{1}{2} K_{1} \varepsilon^{2} \langle t \rangle^{5}$$

and

$$\hbar^{-d} \sum_{|\alpha| \le M} \left\| \boldsymbol{\xi}^{\alpha} \langle \boldsymbol{\nabla}_x, \boldsymbol{\nabla}_{\boldsymbol{\xi}} \rangle^{\sigma_4} \langle \boldsymbol{\nabla}_x \rangle P(t) \right\|_{L^2_{x,y}}^2 \le \frac{1}{2} K_1 \varepsilon^2 \langle t \rangle^4.$$

Therefore,

$$\hbar^{-d} \sum_{|\alpha| \le M} \left\| \boldsymbol{\xi}^{\alpha} \langle \boldsymbol{\nabla}_{x}, \boldsymbol{\nabla}_{\xi} \rangle^{\sigma_{4}} \langle t \boldsymbol{\nabla}_{x}, \boldsymbol{\nabla}_{\xi} \rangle P(t) \right\|_{L^{2}_{x,y}}^{2} \le 2K_{1} \langle t \rangle^{5} \varepsilon^{2} \le 2K_{1} \langle t \rangle^{2d-1} \varepsilon^{2}$$

for $d \geq 3$, which concludes the proof of (I1).

4.3. Bootstrap improvement (I3). This section establishes bootstrap improvement (I3). The analysis is conducted on the Wigner transform, leveraging the equivalence given by

$$\hbar^{-d} \sum_{|\alpha| \le M} \left\| \boldsymbol{\xi}^{\alpha} \langle \boldsymbol{\nabla}_{x}, \boldsymbol{\nabla}_{\xi} \rangle^{\sigma_{4}} | \boldsymbol{\nabla}_{x} |^{\delta} P(t) \right\|_{L^{2}_{x,y}}^{2} \sim \left\| \langle t \nabla_{z}, \nabla_{v} \rangle W[P] \right\|_{H^{\sigma_{4}}_{M}}$$

Let $|\alpha| \leq M$ be a multi-index. An energy estimate yields

$$\begin{split} \frac{1}{2} \frac{d}{dt} \Big\| |\nabla_z|^{\delta} \langle \nabla_z, \nabla_v \rangle^{\sigma_3} (v^{\alpha} W[P]) \Big\|_{L^2}^2 &= \operatorname{Re} \int_{k,\eta} |k|^{\delta} \langle k, \eta \rangle^{\sigma_3} \overline{D_{\eta}^{\alpha} \widehat{W[P]}(t,k,\eta)} |k|^{\delta} \langle k, \eta \rangle^{\sigma_3} D_{\eta}^{\alpha} \widehat{W[P_{\mathrm{L}}]}(t,k,\eta) \, d\eta dk \\ &+ \operatorname{Re} \int_{k,\eta} |k|^{\delta} \langle k, \eta \rangle^{\sigma_3} \overline{D_{\eta}^{\alpha} \widehat{W[P]}(t,k,\eta)} |k|^{\delta} \langle k, \eta \rangle^{\sigma_3} D_{\eta}^{\alpha} \widehat{W[P_{\mathrm{NL}}]}(t,k,\eta) \, d\eta dk \\ &= \mathcal{L}^{(\mathrm{B3})} + \mathcal{N} \mathcal{L}^{(\mathrm{B3})}, \end{split}$$

where $\widehat{W[P_{L}]}(t,k,\eta)$ and $\widehat{W[P_{NL}]}(t,k,\eta)$ are defined in (5).

For $\mathcal{L}^{(B3)}$, Lemma 1.13 is utilised to separate the highest-order and remainder terms arising from the v^{α} -weights,

$$\begin{split} \left| \mathcal{L}^{(\mathrm{B3})} \right| \lesssim & \int_{k,\eta} |k|^{\delta} \langle k,\eta \rangle^{\sigma_{3}} \left| D_{\eta}^{\alpha} \widehat{W[P]}(t,k,\eta) \right| \langle k,\eta \rangle^{\sigma_{3}} |k|^{1+\delta} |\widehat{w}(k)| |\widehat{\rho_{Q}}(t,k)| |\eta - kt| \left| D_{\eta}^{\alpha} \widehat{g}(\eta - kt) \right| d\eta dk \\ &+ \sum_{|\beta| \le |\alpha| - 1} \int_{k,\eta} |k|^{\delta} \langle k,\eta \rangle^{\sigma_{3}} \left| D_{\eta}^{\alpha} \widehat{W[P]}(t,k,\eta) \right| \langle k,\eta \rangle^{\sigma_{3}} |k|^{1+\delta} \langle \hbar k \rangle^{|\alpha| - 1} |\widehat{w}(k)| |\widehat{\rho_{Q}}(t,k)| \left| D_{\eta}^{\beta} \widehat{g}(\eta - kt) \right| d\eta dk \\ &= \mathcal{L}_{0}^{(\mathrm{B3})} + \mathcal{L}_{M}^{(\mathrm{B3})}. \end{split}$$

To control $\mathcal{L}_0^{(B3)}$, the inequality $\langle k, \eta \rangle^{\sigma_3} \lesssim \langle k, kt \rangle^{\sigma_3} \langle \eta - kt \rangle^{\sigma_3}$ is used along with

$$\langle k, kt \rangle^{\sigma_3} |k|^{1+\delta} = |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_4} \frac{|k|^{\frac{1}{2}+\delta}}{\langle k, kt \rangle^{\sigma_4-\sigma_3}} \le |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_4} \frac{|k|^{\frac{1}{2}+\delta}}{|k, kt|^{\frac{1}{2}+\delta}} \le \frac{|k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_4}}{t^{\frac{1}{2}+\delta}}$$

if $\sigma_4 - \sigma_3 \geq \frac{1}{2} + \delta$. Hence,

$$\begin{split} \left| \mathcal{L}_{0}^{(\mathrm{B3})} \right| &\lesssim \frac{1}{\langle t \rangle^{\frac{1}{2} + \delta}} \int_{k,\eta} |k|^{\delta} \langle k, \eta \rangle^{\sigma_{3}} \left| D_{\eta}^{\alpha} \widehat{W[P]}(t,k,\eta) \right| |k|^{\frac{1}{2}} \langle k,kt \rangle^{\sigma_{4}} |\widehat{\rho_{Q}}(t,k)| \langle \eta - kt \rangle^{\sigma_{3}+1} \left| D_{\eta}^{\alpha} \widehat{g}(\eta - kt) \right| d\eta dk \\ &\lesssim \frac{1}{\langle t \rangle^{\frac{1}{2} + \delta}} \left\| |\nabla_{z}|^{\delta} \langle \nabla_{z}, \nabla_{v} \rangle^{\sigma_{3}} (v^{\alpha} W[P]) \right\|_{L^{2}} \left\| |k|^{\frac{1}{2}} \langle k,kt \rangle^{\sigma_{4}} \widehat{\rho_{Q}}(t,k) \right\|_{L^{2}_{k}} \|g\|_{H^{\sigma_{3}+1}_{M}}, \end{split}$$

upon performing Cauchy–Schwarz in η and then k. Similarly, for $0 < \hbar \leq 1$, and under the assumption $|\widehat{w}(k)| \lesssim \langle k \rangle^{-M-1/2}$,

$$\left|\mathcal{L}_{M}^{(\mathrm{B3})}\right| \lesssim \frac{1}{\langle t \rangle^{\frac{1}{2} + \delta}} \left\| |\nabla_{z}|^{\delta} \langle \nabla_{z}, \nabla_{v} \rangle^{\sigma_{3}} (v^{\alpha} W[P]) \right\|_{L^{2}} \left\| |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_{4}} \widehat{\rho_{Q}}(t, k) \right\|_{L^{2}_{k}} \|g\|_{H^{\sigma_{3}}_{M}}.$$

For the nonlinear term, using an identical approach to Section 4.2.1, the decomposition $\mathcal{NL}^{(B3)} = \mathcal{NL}_0^{(B3)} + \mathcal{NL}_M^{(B1)}$ is performed, where

$$\begin{split} \left| \mathcal{NL}_{0}^{(\mathrm{B3})} \right| \lesssim \int_{k,\eta,\ell} |k|^{\delta} \langle k,\eta \rangle^{\sigma_{3}} \left| D_{\eta}^{\alpha} \widehat{W[P]}(t,k,\eta) \right| \Big| |k|^{\delta} \langle k,\eta \rangle^{\sigma_{3}} - |k-\ell|^{\delta} \langle k-\ell,\eta-\ell t \rangle^{\sigma_{3}} \Big| \\ \times |\widehat{w}(\ell)| |\ell| |\widehat{\rho_{Q}}(t,\ell)| |\eta-kt| \Big| D_{\eta}^{\alpha} \widehat{W[P]}(t,k-\ell,\eta-\ell t) \Big| \, d\ell dk d\eta \end{split}$$

and

$$\begin{split} \left| \mathcal{NL}_{M}^{(\mathrm{B3})} \right| \lesssim \sum_{|\beta| \le |\alpha| - 1} \int_{k,\eta,\ell} |k|^{\delta} \langle k,\eta \rangle^{\sigma_{3}} \Big| D_{\eta}^{\alpha} \widehat{W[P]}(t,k,\eta) \Big| \\ \times |k|^{\delta} \langle k,\eta \rangle^{\sigma_{3}} |\widehat{w}(\ell)| |\ell| \langle \hbar \ell \rangle^{|\alpha| - 1} |\widehat{\rho_{Q}}(t,\ell)| \Big| D_{\eta}^{\beta} \widehat{W[P]}(t,k-\ell,\eta-\ell t) \Big| \, d\ell dk d\eta \end{split}$$

Now, using the Mean Value Theorem,

$$\begin{split} \left| |k|^{\delta} \langle k, \eta \rangle^{\sigma_{3}} - |k - \ell|^{\delta} \langle k - \ell, \eta - \ell t \rangle^{\sigma_{3}} \right| \\ & \leq \left| |k|^{\delta} - |k - \ell|^{\delta} \Big| \langle k, \eta \rangle^{\sigma_{3}} + |k - \ell|^{\delta} |\langle k, \eta \rangle^{\sigma_{3}} - \langle k - \ell, \eta - \ell t \rangle^{\sigma_{3}} | \\ & \lesssim |\ell|^{\delta} \langle k, \eta \rangle^{\sigma_{3}} + |k - \ell|^{\delta} |\ell, \ell t| \sup_{|\ell'| \leq |\ell|} \langle k - \ell', \eta - \ell' t \rangle^{\sigma_{3} - 1} \\ & \lesssim \left(|\ell|^{\delta} + |k - \ell|^{\delta} \right) \langle \ell, \ell t \rangle^{\sigma_{3}} + |\ell|^{\delta} \langle k - \ell, \eta - \ell t \rangle^{\sigma_{3}} + |k - \ell|^{\delta} |\ell, \ell t| \langle k - \ell, \eta - \ell t \rangle^{\sigma_{3} - 1} \end{split}$$

This leads to three terms $|\mathcal{NL}_0^{(B3)}| \lesssim \mathcal{R}_0^{(B3)} + \mathcal{T}_{0,1}^{(B3)} + \mathcal{T}_{0,2}^{(B3)}$, with designations of "reaction" and "transport" assigned to the leading order weight. By employing the inequality $|\eta - kt| \leq |\eta - \ell t| + t|k - \ell|$, the reaction term is decomposed into four components:

$$\begin{split} \mathcal{R}_{0}^{(\mathrm{B3})} \lesssim & \int_{k,\eta,\ell} |k|^{\delta} \langle k,\eta \rangle^{\sigma_{3}} \left| D_{\eta}^{\alpha} \widehat{W[P]}(t,k,\eta) \right| |\ell|^{\delta} \langle \ell,\ell t \rangle^{\sigma_{3}} \\ & \times |\widehat{w}(\ell)| |\ell| |\widehat{\rho_{Q}}(t,\ell)| \left(|\eta-\ell t|+t|k-\ell| \right) \left| D_{\eta}^{\alpha} \widehat{W[P]}(t,k-\ell,\eta-\ell t) \right| d\ell dk d\eta \\ & + \int_{k,\eta,\ell} |k|^{\delta} \langle k,\eta \rangle^{\sigma_{3}} \left| D_{\eta}^{\alpha} \widehat{W[P]}(t,k,\eta) \right| |k-\ell|^{\delta} \langle \ell,\ell t \rangle^{\sigma_{3}} \\ & \times |\widehat{w}(\ell)| |\ell| |\widehat{\rho_{Q}}(t,\ell)| \left(|\eta-\ell t|+t|k-\ell| \right) \left| D_{\eta}^{\alpha} \widehat{W[P]}(t,k-\ell,\eta-\ell t) \right| d\ell dk d\eta \\ & = \mathcal{R}_{1;V}^{(\mathrm{B3})} + \mathcal{R}_{1;Z}^{(\mathrm{B3})} + \mathcal{R}_{2;V}^{(\mathrm{B3})}, \end{split}$$

where the sub-divisions $\mathcal{R}_{i;V}$ and $\mathcal{R}_{i;Z}$ correspond to the terms involving $|\eta - \ell t|$ and $t|k - \ell|$, respectively. The term $\mathcal{R}_{1;V}^{(B3)}$ is the most challenging to control due to the absence of a factor of $|k - \ell|$ in its integrated, making it more difficult to apply (B3). Applying the bootstrap assumption (B2) to control the ρ terms leads to additional decay with respect to t:

$$|\ell|^{1+\delta} \langle \ell, \ell t \rangle^{\sigma_3} = |\ell|^{\frac{1}{2}} \langle \ell, \ell t \rangle^{\sigma_4} \frac{|\ell|^{\frac{1}{2}+\delta}}{\langle \ell, \ell t \rangle^{\sigma_4-\sigma_3}} \le |\ell|^{\frac{1}{2}} \langle \ell, \ell t \rangle^{\sigma_4} \frac{|\ell|^{\frac{1}{2}+\delta}}{|\ell, \ell t|^{\frac{1}{2}+\delta}} = \frac{|\ell|^{\frac{1}{2}} \langle \ell, \ell t \rangle^{\sigma_4}}{\langle t \rangle^{\frac{1}{2}+\delta}}$$

given that $\sigma_4 - \sigma_3 \geq \frac{1}{2} + \delta$. By applying Lemma 4.1 (b) with the the L^1 norm on the function $(k,\eta) \mapsto \eta D_{\eta}^{\alpha} \widehat{W}[P](t,k,\eta)$, and using that $|k|^{-2\delta} \langle k \rangle^{-2\sigma_3}$ is integrable in $d \geq 3$, it follows that

$$\begin{aligned} \mathcal{R}_{1;V}^{(\mathrm{B3})} &\lesssim \frac{1}{\langle t \rangle^{\frac{1}{2} + \delta}} \left\| |\nabla_z|^{\delta} \langle \nabla_z, \nabla_v \rangle^{\sigma_3} (v^{\alpha} W[P]) \right\|_{L^2} \left\| |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_4} \widehat{\rho_Q}(t,k) \right\|_{L^2_k} \int_k \left\| \langle \eta \rangle D_{\eta}^{\alpha} \widehat{W}[P](t,k,\eta) \right\|_{L^2_{\eta}} dk \\ &\lesssim \frac{1}{\langle t \rangle^{\frac{1}{2} + \delta}} \left\| |\nabla_z|^{\delta} \langle \nabla_z, \nabla_v \rangle^{\sigma_3} (v^{\alpha} W[P]) \right\|_{L^2}^2 \left\| |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_4} \widehat{\rho_Q}(t,k) \right\|_{L^2_k}. \end{aligned}$$

Each of the remaining terms $\mathcal{R}_{1;Z}^{(B3)}$, $\mathcal{R}_{2;V}^{(B3)}$ and $\mathcal{R}_{2;Z}^{(B3)}$ contains a factor of $|k - \ell|$ in its integrand, allowing the use of Lemma 4.1 (a) with the L^1 term applied to ρ with its weights. For $\mathcal{R}_{1;Z}^{(B3)}$, the inequality

$$\begin{split} |k-\ell| &\leq |k-\ell|^{\delta} \langle k-\ell, \eta-\ell t \rangle^{\sigma_{3}} \text{ is employed,} \\ \mathcal{R}_{1;Z}^{(\mathrm{B3})} \lesssim t \Big\| |\nabla_{z}|^{\delta} \langle \nabla_{z}, \nabla_{v} \rangle^{\sigma_{3}} (v^{\alpha} W[P]) \Big\|_{L^{2}}^{2} \int_{\ell} |\ell|^{1+\delta} |\widehat{w}(\ell)| \langle \ell, \ell t \rangle^{\sigma_{3}} |\widehat{\rho_{Q}}(t,\ell)| \, d\ell \\ &\lesssim t \Big\| |\nabla_{z}|^{\delta} \langle \nabla_{z}, \nabla_{v} \rangle^{\sigma_{3}} (v^{\alpha} W[P]) \Big\|_{L^{2}}^{2} \Big\| |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_{4}} \widehat{\rho_{Q}}(t,k) \Big\|_{L^{2}_{k}} \left(\int_{\ell} \frac{|\ell|^{1+2\delta}}{\langle \ell, \ell t \rangle^{2\sigma_{4}-2\sigma_{3}}} \, d\ell \right)^{\frac{1}{2}} \\ &\lesssim \frac{1}{\langle t \rangle^{\frac{d}{2}-\frac{1}{2}+\delta}} \Big\| |\nabla_{z}|^{\delta} \langle \nabla_{z}, \nabla_{v} \rangle^{\sigma_{3}} (v^{\alpha} W[P]) \Big\|_{L^{2}}^{2} \Big\| |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_{4}} \widehat{\rho_{Q}}(t,k) \Big\|_{L^{2}_{k}} \left(\int_{\ell} \frac{|\ell|^{1+2\delta}}{\langle \ell, \ell t \rangle^{2\sigma_{4}-2\sigma_{3}}} \, d\ell \right)^{\frac{1}{2}} \end{split}$$

if $\sigma_4 - \sigma_3 > d/2 + 1/2 + \delta$. The sub-divisions $\mathcal{R}_{2;V}^{(B3)}$ and $\mathcal{R}_{2;Z}^{(B3)}$ can be controlled similarly to obtain:

$$\mathcal{R}_{2;V}^{(\mathrm{B3})} + \mathcal{R}_{2;Z}^{(\mathrm{B3})} \lesssim \frac{1}{\langle t \rangle} \left\| |\nabla_z|^{\delta} \langle \nabla_z, \nabla_v \rangle^{\sigma_3} (v^{\alpha} W[P]) \right\|_{L^2}^2 \left\| |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_4} \widehat{\rho_Q}(t,k) \right\|_{L^2_k}$$

Next, the transport terms $\mathcal{T}_{0,1}^{(B3)}$ and $\mathcal{T}_{0;2}^{(B3)}$ are addressed. For $\mathcal{T}_{0,1}^{(B3)}$, the additional factor of $|\eta - kt|$ necessitates the use of the bootstrap estimate (B1), which behaves like $\langle t \rangle^{d-\frac{1}{2}}$. Thus, decay must be extracted from $|\widehat{\rho_Q}(t,\ell)| \leq \sqrt{K_5} \varepsilon \langle \ell, \ell t \rangle^{-\sigma_1}$ to control this term. Bounding $|\eta - kt| \leq \langle t(k-\ell), \eta - \ell t \rangle$ leads to the following:

$$\begin{split} \mathcal{T}_{0,1}^{(\mathrm{B3})} \lesssim & \int_{k,\eta,\ell} |k|^{\delta} \langle k,\eta \rangle^{\sigma_{3}} \left| D_{\eta}^{\alpha} \widehat{W[P]}(t,k,\eta) \right| |\ell|^{1+\delta} |\widehat{\rho_{Q}}(t,\ell)| \\ & \times \langle t(k-\ell),\eta-\ell t \rangle \langle k-\ell,\eta-\ell t \rangle^{\sigma_{3}} \left| D_{\eta}^{\alpha} \widehat{W[P]}(t,k-\ell,\eta-\ell t) \right| dl dk d\eta \\ \lesssim \left\| |\nabla_{z}|^{\delta} \langle \nabla_{z},\nabla_{v} \rangle^{\sigma_{3}} (v^{\alpha} W[P]) \right\|_{L^{2}} \| \langle t \nabla_{z},\nabla_{v} \rangle W[P] \|_{H_{M}^{\sigma_{4}}} \int_{\ell} |\ell|^{1+\delta} |\widehat{\rho_{Q}}(t,\ell)| \, d\ell \\ \lesssim \frac{\sqrt{K_{5}}\varepsilon}{\langle t \rangle^{d+1+\delta}} \left\| |\nabla_{z}|^{\delta} \langle \nabla_{z},\nabla_{v} \rangle^{\sigma_{3}} (v^{\alpha} W[P]) \right\|_{L^{2}} \| \langle t \nabla_{z},\nabla_{v} \rangle W[P] \|_{H_{M}^{\sigma_{4}}}. \end{split}$$

For $\mathcal{T}_{0,2}^{(B3)}$, the additional flexibility allows for the expression $|\eta - kt| \lesssim \langle t \rangle \langle k - \ell, \eta - \ell t \rangle$ to be utilised, leading to control via the (B3) norm:

$$\begin{split} \mathcal{T}_{0,2}^{(\mathrm{B3})} \lesssim \langle t \rangle \int_{k,\eta,\ell} |k|^{\delta} \langle k,\eta \rangle^{\sigma_3} \Big| D_{\eta}^{\alpha} \widehat{W[P]}(t,k,\eta) \Big| |\ell| |\ell,\ell t| |\widehat{\rho_Q}(t,\ell)| \\ \times \langle k-\ell,\eta-\ell t \rangle^{\sigma_3} |k-\ell|^{\delta} \Big| D_{\eta}^{\alpha} \widehat{W[P]}(t,k-\ell,\eta-\ell t) \Big| \, d\ell dk d\eta \\ \lesssim \langle t \rangle \Big\| |\nabla_z|^{\delta} \langle \nabla_z,\nabla_v \rangle^{\sigma_3} (v^{\alpha} W[P]) \Big\|_{L^2}^2 \int_{\ell} |\ell| |\ell,\ell t| |\widehat{\rho_Q}(t,\ell)| \, d\ell \\ \lesssim \frac{\sqrt{K_5} \varepsilon}{\langle t \rangle^d} \Big\| |\nabla_z|^{\delta} \langle \nabla_z,\nabla_v \rangle^{\sigma_3} (v^{\alpha} W[P]) \Big\|_{L^2}^2. \end{split}$$

This holds under the condition $\sigma_1 > d+2$.

For the remainder term $\mathcal{NL}_{M}^{(B3)}$, the triangle inequality yields $|k|^{\delta} \leq |\ell|^{\delta} + |k - \ell|^{\delta}$ and $\langle k, \eta \rangle^{\sigma_{3}} \lesssim \langle \ell, \ell t \rangle^{\sigma_{3}} + \langle k - \ell, \eta - \ell t \rangle^{\sigma_{3}}$. This allows for the following decomposition:

$$|k|^{\delta} \langle k, \eta \rangle^{\sigma_3} \lesssim \left(|\ell|^{\delta} + |k - \ell|^{\delta} \right) + |\ell|^{\delta} \langle k - \ell, \eta - \ell t \rangle^{\sigma_3} + |k - \ell|^{\delta} \langle k - \ell, \eta - \ell t \rangle^{\sigma_3}.$$

Consequently, three terms are obtained such that $\left|\mathcal{NL}_{M}^{(B3)}\right| \lesssim \mathcal{R}_{M}^{(B3)} + \mathcal{T}_{M,1}^{(B3)} + \mathcal{T}_{M,2}^{(B3)}$. The reaction term is treated in the same manner as the highest-order term $\mathcal{R}_{0}^{(B3)}$ above:

$$\mathcal{R}_{M}^{(\mathrm{B3})} \lesssim \frac{1}{\langle t \rangle^{\frac{1}{2} + \delta}} \left\| |\nabla_{z}|^{\delta} \langle \nabla_{z}, \nabla_{v} \rangle^{\sigma_{3}} (v^{\alpha} W[P]) \right\|_{L^{2}} \left\| |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_{4}} \widehat{\rho_{Q}}(t, k) \right\|_{L^{2}_{k}} \left\| |\nabla_{z}|^{\delta} W[P] \right\|_{H^{\sigma_{3}}_{M}}$$

The transport term $\mathcal{T}_{M,1}^{(\mathrm{B3})}$ is treated similarly to $\mathcal{T}_{0,1}^{(\mathrm{B3})}$, leading to

$$\mathcal{T}_{M,1}^{(\mathrm{B3})} \lesssim \frac{\sqrt{K_5\varepsilon}}{\langle t \rangle^{d+1+\delta}} \left\| |\nabla_z|^{\delta} \langle \nabla_z, \nabla_v \rangle^{\sigma_3} (v^{\alpha} W[P]) \right\|_{L^2} \| \langle t \nabla_z, \nabla_v \rangle W[P] \|_{H^{\sigma_4}_M}$$

Finally, $\mathcal{T}_{M,2}^{(B3)}$ is handled as in $\mathcal{T}_{0,2}^{(B3)}$,

$$\mathcal{T}_{M,2}^{(\mathrm{B3})} \lesssim \frac{\sqrt{K_5}\varepsilon}{\langle t \rangle^{d+1}} \left\| \left| \nabla_z \right|^{\delta} \langle \nabla_z, \nabla_v \rangle^{\sigma_3} (v^{\alpha} W[P]) \right\|_{L^2} \left\| \left| \nabla_z \right|^{\delta} W[P] \right\|_{H_M^{\sigma_3}}$$

4.3.1. Conclusion of bootstrap improvement (I3). Combining all estimates shows that for each $|\alpha| \leq M$ and $d \geq 3$,

$$\begin{split} & \frac{d}{dt} \left\| |\nabla_{z}|^{\delta} \langle \nabla_{z}, \nabla_{v} \rangle^{\sigma_{3}} (v^{\alpha} W[P](t)) \right\|_{L^{2}} \\ & \lesssim \frac{1}{\langle t \rangle^{\frac{1}{2} + \delta}} \left\| |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_{4}} \widehat{\rho_{Q}}(t, k) \right\|_{L^{2}_{k}} + \frac{1}{\langle t \rangle^{\frac{1}{2} + \delta}} \left\| |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_{4}} \widehat{\rho_{Q}}(t, k) \right\|_{L^{2}_{k}} \left\| |\nabla_{z}|^{\delta} W[P](t) \right\|_{H^{\sigma_{3}}_{M}} \\ & \quad + \frac{\sqrt{K_{5}}\varepsilon}{\langle t \rangle^{d+1+\delta}} \| \langle t \nabla_{z}, \nabla_{v} \rangle W[P](t) \|_{H^{\sigma_{4}}_{M}} + \frac{\sqrt{K_{5}}\varepsilon}{\langle t \rangle^{d}} \left\| |\nabla_{z}|^{\delta} W[P](t) \right\|_{H^{\sigma_{3}}_{M}}. \end{split}$$

Integrating and utilising the bootstrap assumptions (B1), (B2), and (B3) results in:

$$\begin{split} |\nabla_{z}|^{\delta} \langle \nabla_{z}, \nabla_{v} \rangle^{\sigma_{3}} (v^{\alpha} W[P](t)) \Big\|_{L^{2}} \\ &\lesssim (1 + \sqrt{K_{3}}\varepsilon) \int_{0}^{t} \frac{1}{\langle s \rangle^{\frac{1}{2} + \delta}} \Big\| |k|^{\frac{1}{2}} \langle k, ks \rangle^{\sigma_{4}} \widehat{\rho_{Q}}(s, k) \Big\|_{L^{2}_{k}} ds \\ &+ \sqrt{K_{5}}\varepsilon \int_{0}^{t} \frac{1}{\langle s \rangle^{\frac{3}{2} + \delta}} \sup_{0 \le s \le T^{*}} \left(\frac{\| \langle s \nabla_{z}, \nabla_{v} \rangle W[P](s) \|_{H^{\sigma_{4}}_{M}}}{\langle s \rangle^{d - \frac{1}{2}}} \right) ds + \sqrt{K_{3}} K_{5} \varepsilon^{2} \int_{0}^{t} \frac{1}{\langle s \rangle^{d}} ds \\ &\lesssim (1 + \sqrt{K_{3}}\varepsilon) \left(\int_{0}^{t} \frac{1}{\langle s \rangle^{1 + 2\delta}} ds \right)^{\frac{1}{2}} \Big\| |k|^{\frac{1}{2}} \langle k, kt \rangle^{\sigma_{4}} \widehat{\rho_{Q}}(t, k) \Big\|_{L^{2}_{t,k}} + \sqrt{K_{1}} K_{5} \varepsilon^{2} + \sqrt{K_{3}} K_{5} \varepsilon^{2} \\ &\lesssim \varepsilon \left[\sqrt{K_{2}} + \left(\sqrt{K_{2}} K_{3} + \sqrt{K_{1}} K_{5} + \sqrt{K_{3}} K_{5} \right) \varepsilon \right]. \end{split}$$

Hence, there exists a positive constant $\widetilde{K} = \widetilde{K}(d, g, w, \sigma_i, Q_{in})$ (independent of $0 < \hbar \leq 1$), such that

$$\frac{1}{\hbar^d} \sum_{|\alpha| \le M} \left\| \langle \boldsymbol{\nabla}_x, \boldsymbol{\nabla}_{\boldsymbol{\xi}} \rangle^{\sigma_3} | \boldsymbol{\nabla}_x |^{\delta} \boldsymbol{\xi}^{\alpha} P(t) \right\|_{L^2_{x,y}}^2 \le \widetilde{K} \varepsilon^2 \left[\sqrt{K_2} + \left(\sqrt{K_2 K_3} + \sqrt{K_1 K_5} + \sqrt{K_3 K_5} \right) \varepsilon \right]^2 \right]^2$$

By selecting $K_3 := \widetilde{K} \left(\sqrt{K_2} + 1\right)^2 / 2$ and $\varepsilon \leq \left(\sqrt{K_2 K_3} + \sqrt{K_1 K_5} + \sqrt{K_3 K_5}\right)^{-1}$, the improvement (I3) is obtained.

Acknowledgement. Thanks are due to Clément Mouhot for suggesting this topic, as well as for numerous helpful discussions and ongoing encouragement throughout this work.

References

- J. Bedrossian, N. Masmoudi, and C. Mouhot. "Landau damping in finite regularity for unconfined systems with screened interactions". In: Comm. Pure Appl. Math. 71.3 (2018). DOI: 10.1002/cpa.21730.
- [2] J. Bedrossian, N. Masmoudi, and C. Mouhot. "Linearized Wave-Damping Structure of Vlasov-Poisson in ℝ³". In: SIAM Journal on Mathematical Analysis 54.4 (2022). DOI: 10.1137/20M1386141.
- [3] J. Bedrossian, N. Masmoudi, and Clément Mouhot. "Landau damping: paraproducts and Gevrey regularity". In: Ann. PDE 2.1 (2016). DOI: 10.1007/s40818-016-0008-2.
- [4] Jacob Bedrossian. "Nonlinear echoes and Landau damping with insufficient regularity". In: *Tunisia Journal of Mathematics* 3.1 (2021). DOI: 10.2140/tunis.2021.3.121.

- [5] A. Bove, G. Da Prato, and G. Fano. "An existence proof for the Hartree-Fock time-dependent problem with bounded two-body interaction". In: *Comm. Math. Phys.* 37 (1974). ISSN: 0010-3616.
- [6] A. Bove, G. Da Prato, and G. Fano. "On the Hartree-Fock time-dependent problem". In: Comm. Math. Phys. 49.1 (1976). ISSN: 0010-3616.
- [7] J. M. Chadam. "The time-dependent Hartree-Fock equations with Coulomb two-body interaction". In: Comm. Math. Phys. 46.2 (1976). ISSN: 0010-3616.
- [8] T. Chen, Y. Hong, and N. Pavlović. "Global well-posedness of the NLS system for infinitely many fermions". In: Arch. Ration. Mech. Anal. 224 (2017). DOI: 10.1007/s00205-016-1068-x.
- [9] T. Chen, Y. Hong, and N. Pavlović. "On the scattering problem for infinitely many fermions in dimensions d ≥ 3 at positive temperature". In: Ann. Inst. H. Poincaré C Anal. Non Linéaire 35.2 (2018). DOI: 10.1016/j.anihpc.2017.05.002.
- [10] E. Grenier, T. T. Nguyen, and I. Rodnianski. "Landau damping for analytic and Gevrey data". In: Mathematical Research Letters 28.6 (2021). DOI: 10.4310/mrl.2021.v28.n6.a3.
- [11] S. Hadama. "Asymptotic stability of a wide class of stationary solutions for the Hartree and Schrödinger equations for infinitely many particles". In: *arXiv e-prints* (2024). arXiv: 2308.15929 [math.AP].
- [12] I. Ionescu et al. "Nonlinear Landau damping for the Vlasov-Poisson system in ℝ³: the Poisson equilibrium". In: Ann. PDE 10.2 (2024). DOI: 10.1007/s40818-023-00161-w.
- [13] L. Landau. "On the vibrations of the electronic plasma". In: Acad. Sci. USSR. J. Phys. 10 (1946).
- [14] M. Lewin and J. Sabin. "The Hartree and Vlasov equations at positive density". In: Comm. Partial Differential Equations 45.12 (2020). DOI: 10.1080/03605302.2020.1803355.
- M. Lewin and J. Sabin. "The Hartree equation for infinitely many particles I. Well-posedness theory". In: Comm. Math. Phys. 334.1 (2015). DOI: 10.1007/s00220-014-2098-6.
- [16] M. Lewin and J. Sabin. "The Hartree equation for infinitely many particles, II: Dispersion and scattering in 2D". In: Anal. PDE 7.6 (2014). DOI: 10.2140/apde.2014.7.1339.
- [17] C. Mouhot and C. Villani. "On Landau damping". In: Acta Math 207.1 (2011), pp. 29–201. DOI: 10.1007/s11511-011-0068-9.
- [18] T. T. Nguyen and C. You. "Modified scattering for long-range Hartree equations of infinite rank near vacuum". In: arXiv e-prints (2024). arXiv: 2408.15860 [math.AP].
- [19] T. T. Nguyen and C. You. "Plasmons for the Hartree equations with Coulomb interaction". In: arXiv e-prints (2023). arXiv: 2306.03800 [math.AP].
- [20] C. You. "Phase mixing estimates for the nonlinear Hartree equation of infinite rank". In: arXiv e-prints (2024). arXiv: 2408.15972 [math.AP].