A UNIVERSAL CHARACTERIZATION OF UNCURVING

GUILLAUME LAPLANTE-ANFOSSI, ADRIAN PETR, AND VIVEK SHENDE

ABSTRACT. We introduce a category of non-symmetric operads with distinguished curvature term, and determine the right adjoint to the functor forgetting the distinction. As a result we give a universal characterization of the procedure of "twisting by a Maurer-Cartan element", and also of the curved A-infinity operad itself.

1. INTRODUCTION

The operad A_{∞} , which encodes the notion of associative algebras up to homotopy, can be presented as the (cellular chains on the) operad of stable holomorphic disks with $n \geq 3$ boundary marked points. As such, the operad A_{∞} is implicated in geometric questions involving such moduli; in particular, and famously, in Lagrangian Floer theory [5]. However, due to the existence of stable maps from disks with unstable domains (zero, one, or two boundary marked points), the Fukaya category does not generally begin life with an A_{∞} structure, but rather with a somewhat more complicated *curved* A_{∞} structure [6].

Fortunately for those who wish to return to the world of usual categories, there is a standard procedure, namely twisting by a Maurer-Cartan element, which turns curved A_{∞} structures into ordinary A_{∞} structures. To describe this construction, let us introduce some notation¹. We write A_{∞}^+ for the variant of the operad A_{∞} containing the degree (-1) arity 1 operation μ_1 (so there's an equivalence between representations of A_{∞}^+ in graded modules, and representations of A_{∞} in dg modules). We write \mathcal{T} the operad freely generated by a degree (-1) arity 0 element μ_0 and a degree 0 arity 0 element α , and equip it with differential $d_{\mathcal{T}}\alpha = \mu_0$. The coproduct of operads $A_{\infty}^+ \vee \mathcal{T}$ admits a natural map $A_{\infty}^+ \to A_{\infty}^+ \vee \mathcal{T}$ taking μ_n to μ_n , so any $A_{\infty}^+ \vee \mathcal{T}$ -algebra carries the structure of an A_{∞}^+ -algebra.

Now consider the curved A_{∞} operad cA_{∞} , which contains also the degree (-1) arity 0 curvature term μ_0 . The twisting procedure is encoded by the following morphism

(1.1)
$$\begin{array}{cccc} \operatorname{Tw}: cA_{\infty} & \longrightarrow & A_{\infty}^{+} \lor \mathcal{T} \\ \mu_{0} & \mapsto & \sum_{k} \mu_{k}(\alpha, \cdots, \alpha) \\ \mu_{n} & \mapsto & \sum_{i_{1}, \dots, i_{n+1}} \mu_{n+\sum i_{k}}(\alpha^{i_{1}}, -, \alpha^{i_{2}}, -, \cdots, -, \alpha^{i_{n+1}}) \end{array},$$

along with the fact (which can be seen readily from the formula for Tw) that an $A^+_{\infty} \vee \mathcal{T}$ algebra is a cA_{∞} -algebra plus the datum of a Maurer-Cartan element (to which α will map).

The work of the authors is supported by Novo Nordisk Foundation grant NNF20OC0066298, Villum Fonden Villum Investigator grant 37814, and Danish National Research Foundation grant DNRF157. GLA was supported by the Andrew Sisson Fund and the Australian Research Council Future Fellowship FT210100256.

¹We work with non-symmetric operads in complete filtered differential graded modules over \mathbb{Z} and use homological and shifted grading conventions, see Section 2.

That is, given a cA_{∞} -algebra with Maurer-Cartan element, one has also the structure of an $A_{\infty}^+ \vee \mathcal{T}$ -algebra, hence by restriction the structure of an A_{∞} -algebra.

The twisting procedure is usually motivated by extension or parallel with the corresponding procedure for (uncurved) Lie algebras, which in turn is well motivated by deformation theoretic considerations and deeply rooted in the literature, going back, apparently, to Quillen [8]. Let us mention also Kontsevich's interpretation of A_{∞} structures and twisting in terms of non-commutative vector fields [7].

The purpose of this note is to record a different sort of motivation: a universal property of uncurving, and, relatedly, a universal property of the operad cA_{∞} itself.

If \mathcal{P} and \mathcal{Q} are operads, then the operad coproduct $\mathcal{P} \vee \mathcal{Q}$ has two increasing filtrations: a \mathcal{P} -filtration where the filtration level of a given element x is the minimal number of operations from \mathcal{P} needed to express x, and similarly a \mathcal{Q} -filtration. We write [q] for an operad generated by the single nonvanishing operation q.

Definition 1.1. We denote by Curv the category defined as follows:

- (1) An object of Curv is a tuple $(\mathcal{Q}, \mathcal{Q}_0, \mu_0, d_\mathcal{Q})$ where $(\mathcal{Q}, d_\mathcal{Q})$ is a dg operad, $\mathcal{Q}_0 \subset \mathcal{Q}$ is a graded (not dg) suboperad, and $\mu_0 \in \mathcal{Q}$ is a nonzero element of degree (-1) and arity 0, such that the natural map of graded operads $\mathcal{Q}_0 \vee [\mu_0] \to \mathcal{Q}$ is an isomorphism. Writing $\mathcal{Q}_0 \subset \mathcal{Q}_1 \subset \cdots$ for the resulting μ_0 -filtration on \mathcal{Q} and $\operatorname{Gr}(d_\mathcal{Q}) : \operatorname{Gr}_k \mathcal{Q} \to \operatorname{Gr}_{k+1} \mathcal{Q}$ for the resulting map of submodules (not operads), we further require that
 - (a) $d_{\mathcal{Q}}\mathcal{Q}_k \subset \mathcal{Q}_{k+1}$, and
 - (b) $\operatorname{Gr}(d_{\mathcal{Q}})(\mu_0) = 0$, i.e. $d_{\mathcal{Q}}(\mu_0) \in \mathcal{Q}_1$.
- (2) A morphism in Curv is a morphism $\Phi: \mathcal{Q} \to \mathcal{Q}'$ of dg operads such that
 - (a) $\Phi(\mathcal{Q}_k) \subset \mathcal{Q}'_k$, and
 - (b) $Gr(\Phi)(\mu_0) = \mu'_0$, i.e. $(\Phi(\mu_0) \mu'_0) \in \mathcal{Q}'_0$.

Example 1.2. Writing $cA_{\infty}^{>0} \subset cA_{\infty}$ for the (not dg) sub-operad spanned by operations of arity > 0, we have $(cA_{\infty}, cA_{\infty}^{>0}, \mu_0, d_{cA_{\infty}}) \in Curv$.

Example 1.3. The tuple $(\mathcal{T}, [\alpha], \mu_0, d_{\mathcal{T}})$ is an object of Curv. More generally, if $(\mathcal{P}, d_{\mathcal{P}})$ is a dg operad, then we have $(\mathcal{P} \lor \mathcal{T}, \mathcal{P} \lor [\alpha], \mu_0, d_{\mathcal{P}} \lor d_{\mathcal{T}}) \in \text{Curv}.$

Example 1.4. The map Tw of (1.1) is a morphism in Curv,

$$\Gamma w : (cA_{\infty}, cA_{\infty}^{>0}, \mu_0, d_{cA_{\infty}}) \to (A_{\infty}^+ \lor \mathcal{T}, A_{\infty}^+ \lor [\alpha], \mu_0, d_{A_{\infty}^+} \lor d_{\mathcal{T}})$$

We write Op for the category of dg operads. There is a forgetful functor \mathcal{L} : Curv \rightarrow Op carrying $(\mathcal{Q}, \mathcal{Q}_0, \mu_0, d_{\mathcal{Q}}) \mapsto (\mathcal{Q}, d_{\mathcal{Q}})$. We will show:

Theorem 1.5. The forgetful functor \mathcal{L} has a right adjoint $\mathcal{R} : \mathrm{Op} \to \mathrm{Curv}$, and the co-unit $\mathcal{L} \circ \mathcal{R} \to id_{\mathrm{Op}}$ admits a section.

The existence of said structures imply the existence of a twisting procedure:

Construction 1.6. Given $Q = (Q, Q_0, \mu_0, d_Q) \in \text{Curv}$, we write Q/μ_0 for the quotient by the dg-operadic ideal generated by μ_0 . We write $\text{Tw}_Q : \mathcal{L}Q \to \mathcal{LR}(Q/\mu_0)$ for the image of the quotient map $\pi : \mathcal{L}Q \to Q/\mu_0$ under

 $\hom_{\mathrm{Op}}(\mathcal{LQ}, \mathcal{Q}/\mu_0) = \hom_{\mathrm{Curv}}(\mathcal{Q}, \mathcal{R}(\mathcal{Q}/\mu_0)) \to \hom_{\mathrm{Op}}(\mathcal{LQ}, \mathcal{LR}(\mathcal{Q}/\mu_0)).$

Then a \mathcal{Q} -algebra structure which factors through $\operatorname{Tw}_{\mathcal{Q}}$ defines a \mathcal{Q}/μ_0 -algebra through the section of the co-unit of the adjunction $\mathcal{Q}/\mu_0 \to \mathcal{LR}(\mathcal{Q}/\mu_0)$.

An equivalent formulation: in the comma category associated to \mathcal{L} : Curv \rightarrow Op, it follows from the adjunction that the map $\mathcal{LR}(\mathcal{Q}/\mu_0) \rightarrow \mathcal{Q}/\mu_0$ is terminal among objects over \mathcal{Q}/μ_0 . As $\mathcal{LQ} \rightarrow \mathcal{Q}/\mu_0$ is another such object, there is a canonical map $\mathcal{LQ} \rightarrow \mathcal{LR}(\mathcal{Q}/\mu_0)$.

Remark 1.7. Forgetful functors more commonly have *left* adjoints, which produce objects that are 'free', i.e. initial for some mapping property. Correspondingly, a right adjoint of a (forgetful) functor produces objects which are terminal for some mapping property.

Remark 1.8. The notation Tw usually denotes an endofunctor on the category of multiplicative operads, see for instance [3, 4]. The analogue in our curved context is the functor MC from Definition-Proposition 5.1.

In fact, we give \mathcal{R} explicitly simultaneously with establishing its existence. (We stated existence separately in order to emphasize that the abstract form of the twisting procedure given in Construction 1.6 depends only on the existence of \mathcal{R} and section of the counit.)

Theorem 1.9. The functor \mathcal{R} is $(\mathcal{P}, d_{\mathcal{P}}) \mapsto (\mathcal{P} \vee \mathcal{T}, \mathcal{P} \vee [\alpha], \mu_0, d_{\mathcal{P}} \vee d_{\mathcal{T}})$. The co-unit of the adjunction is the quotient $(\mathcal{P}, d_{\mathcal{P}}) \vee (\mathcal{T}, d_{\mathcal{T}}) \rightarrow (\mathcal{P}, d_{\mathcal{P}})$ sending $\mu_0, \alpha \mapsto 0$, which thus admits a section given by the inclusion into a coproduct $(\mathcal{P}, d_{\mathcal{P}}) \rightarrow (\mathcal{P}, d_{\mathcal{P}}) \vee (\mathcal{T}, d_{\mathcal{T}})$.

Corollary 1.10. The map $Tw_{cA_{\infty}}$ of Construction 1.6 is given by the formula (1.1).

Proof. By the adjunction, $\operatorname{Tw}_{cA_{\infty}} : cA_{\infty} \to A_{\infty}^+ \lor \mathcal{T}$ is uniquely characterized by the fact that it is a map in Curv whose composition with the co-unit, $cA_{\infty} \to A_{\infty}^+ \lor \mathcal{T} \to A_{\infty}^+$, recovers the natural quotient $cA_{\infty} \to A_{\infty}^+$. As the same is true for the map Tw of (1.1), we have $\operatorname{Tw}_{cA_{\infty}} = \operatorname{Tw}$.

Remark 1.11. The proof of Theorem 1.9 in fact gives an inductive procedure computing the adjunction, which could have been used to determine $Tw_{cA_{\infty}}$.

Perhaps even more remarkably, the operad cA_{∞} itself has a universal characterization in terms of the following:

Definition 1.12. We write Curv₀ for the not-full subcategory of Curv where we restrict to the objects for which $d\mu_0 \equiv 0 \pmod{\mu_0}$, and to the morphisms for which $\Phi(\mu_0) - \mu'_0 = 0$.

Theorem 1.13. The operad cA_{∞} is the initial object of Curv₀.

Remark 1.14. Because $\operatorname{Tw}_{cA_{\infty}}$ is in $\operatorname{hom}_{\operatorname{Curv}}(cA_{\infty}, A_{\infty}^+ \vee \mathcal{T})$ but not in $\operatorname{hom}_{\operatorname{Curv}_0}(cA_{\infty}, A_{\infty}^+ \vee \mathcal{T})$, the operad cA_{∞} cannot also be initial in Curv. Indeed, the canonical map $cA_{\infty} \to A_{\infty}^+ \vee \mathcal{T}$ produced by Theorem 1.13 sends $\mu_0 \mapsto \mu_0$ and $\mu_i \mapsto 0$ for i > 0.

Finally, in Theorem 5.2, we match our constructions with the more conventional account of twisting in terms of Maurer-Cartan operads, as found, e.g. in the textbook [4].

Acknowledgements. We would like to thank Alexander Berglund, Coline Emprin, Hugo Pourcelot, Alex Takeda and Bruno Vallette for useful discussions.

2. Preliminaries

2.1. Conventions. Throughout the text, we will work with non-symmetric operads in filtered complete differential graded modules over \mathbb{Z} , and will use homological conventions. For more details, we refer to [4, Chap. 2]. Unless otherwise stated, when we write "operad" or "morphisms of operads" we mean objects and morphisms in this category. 2.2. **Basic definitions.** Let A be a filtered complete graded \mathbb{Z} -module. We consider it as a dg \mathbb{Z} -module with trivial differential (A, 0). The *endomorphism operad* of (A, 0) is the operad given in arity n by the dg \mathbb{Z} -module

$$\operatorname{End}_{(A,0)}(n) := (\operatorname{hom}(A^{\otimes n}, A), \partial),$$

where the differential is given by $\partial(f) := d_A f - (-1)^{|f|} f d_{A^{\otimes n}}$. Operadic composition is given by composition of functions.

Definition 2.1. The shifted A-infinity operad, denoted A^+_{∞} , is the free operad on generators μ_n , $n \ge 1$, of arity n and degree (-1), endowed with the differential

(2.1)
$$-d_{\mathcal{A}^+_{\infty}}(\mu_n) := \sum_{\substack{p+q+r=n\\p+r \ge 0, q \ge 1}} \mu_{p+1+r} \circ_{p+1} \mu_q \; .$$

Algebras over A^+_{∞} , that is morphisms of operads $A^+_{\infty} \to \operatorname{End}_{(A,0)}$ are shifted A_{∞} -algebras.

Remark 2.2. Contrary to the usual algebraic setting, we consider A_{∞} -operads where the differential is part of the operations. We use the notation A_{∞}^+ to emphasize this fact. Note that any A_{∞}^+ -algebra $f : A_{\infty}^+ \to \operatorname{End}_{(A,0)}$ gives rise to an ordinary A_{∞} -algebra: denoting by $d := f(\mu_1)$ the image of μ_1 and twisting by the operadic Maurer-Cartan element μ_1 gives a morphism of operads $A_{\infty} \to (A_{\infty}^+)^{\mu_1} \to (\operatorname{End}_{(A,0)})^d = \operatorname{End}_{(A,d)}$, see [4, Ex. 5.6].

Definition 2.3. The shifted *curved A-infinity operad*, denoted cA_{∞} , is the free operad on generators μ_n , $n \ge 0$, of arity n and degree -1, endowed with the differential

(2.2)
$$-d_{cA_{\infty}}(\mu_n) := \sum_{\substack{p+q+r=n\\p,q,r\geq 0}} \mu_{p+1+r} \circ_{p+1} \mu_q .$$

The generator μ_0 is in positive filtration.

Algebras over cA_{∞} , that is morphisms of operads $cA_{\infty} \to End_{(A,0)}$ are shifted curved A_{∞} -algebras.

Remark 2.4. The A_{∞} -operads that we consider here are shifted, an adjective we will omit from now on. One can recover the usual degree and signs conventions for (curved) A_{∞} algebras by de-suspension, see [4, Sec. 4.1] for details.

Remark 2.5. Having the differential as part of the operations of A_{∞}^+ and cA_{∞} makes them quasi-isomorphic to the trivial operad [2, Thm. 5.7], see also [1, Rem. 2.53]. This is why one works with filtered objects, and instead consider quasi-isomorphisms at the level of the associated graded [1, Sec. 2.2].

Finally, let \mathcal{P} be an operad. We will make use of the *operadic bracket*, defined on $\mu \in \mathcal{P}(m)$ and $\nu \in \mathcal{P}(n)$ by

$$[\mu,\nu] := \sum_{i=1}^m \mu \circ_i \nu - (-1)^{|\nu||\mu|} \sum_{i=1}^n \nu \circ_i \mu.$$

3. Proof of Theorems 1.5 and 1.9

Any operad in Curv comes equipped with a distinguished element μ_0 of arity zero and degree (-1), which we assume is in positive filtration. Recall that $\mathcal{T} \in \text{Curv}$ is the quasi-free operad generated by μ_0 and an element α of arity 0 and degree 0, with $d_{\mathcal{T}}\alpha = \mu_0$. Here, α is also assumed to be in positive filtration. Working with operads in complete filtered modules will allow us to consider infinite sums involving α and μ_0 .

The task is to show that the forgetful functor

$$\mathcal{L} : \operatorname{Curv} o \operatorname{Op} \ (\mathcal{Q}, \mathcal{Q}_0, \mu_0, d_{\mathcal{Q}}) o (\mathcal{Q}, d_{\mathcal{Q}})$$

has right adjoint

$$\begin{array}{rcl} \mathcal{R} & : & \mathrm{Op} & \to & \mathrm{Curv} \\ & & (\mathcal{P}, d_{\mathcal{P}}) & \mapsto & (\mathcal{P} \lor \mathcal{T}, \mathcal{P} \lor [\alpha], \mu_0, d_{\mathcal{P}} \lor d_{\mathcal{T}}) \ , \end{array}$$

and to show that the co-unit is the map $\pi : \mathcal{P} \vee \mathcal{T} \to \mathcal{P}$ which sends α and μ_0 to 0.

Let us first check, as a special case, that \mathcal{T} is a terminal object of Curv. For any object \mathcal{Q} of Curv, we must show there is a unique morphism $\Phi : \mathcal{Q} \to \mathcal{T}$. Note the module underlying \mathcal{T} has only two generators, α and μ_0 , both of arity zero, and of degrees 0 and (-1), respectively. By definition of Curv and the properties of \mathcal{T} , said morphism must have:

- (1) $\Phi(\mu_0) = \mu_0$,
- (2) if $\nu \in \mathcal{Q}_0$ is of non-zero arity or non-zero degree, then $\Phi(\nu) = 0$,
- (3) if $\nu \in \mathcal{Q}_0$ is of arity 0 and degree 0, there are $c \in \mathbb{Z}$ and $\nu' \in \mathcal{Q}_0(0)$ such that $d_{\mathcal{Q}}(\nu) = \nu' + c\mu_0$, and $\Phi(\nu) = c\alpha$.

On the other hand, one can always define a morphism $\Phi: \mathcal{Q} \to \mathcal{T}$ by the above formulas.

Before returning to the general case, we introduce some notations. Given $\mathcal{P} \in \text{Op}$ and $k, \ell \in \mathbb{Z}_{\geq 0}$, we denote by $\mathcal{R}^k(\mathcal{P})$, respectively $\mathcal{R}^k_\ell(\mathcal{P})$, the sub-N-module (i.e. family of graded modules indexed by N) of $\mathcal{P} \vee \mathcal{T}$ whose elements are sums of terms containing exactly k of α , respectively k of α and ℓ of μ_0 .

Lemma 3.1. Let $k \in \mathbb{Z}_{>0}$ and $\lambda \in \mathcal{R}_0^{k+1}(\mathcal{P})$. If $d_{\mathcal{T}}(\lambda) = 0$, then $\lambda = 0$.

Proof. An element $\lambda \in \mathcal{R}_0^{k+1}(\mathcal{P})$ is a sum of trees with exactly (k+1) leaves decorated by α . Let t denote one of those trees. Its image $d_{\mathcal{T}}(t)$ is the sum of all the trees t' that can be obtained from t by replacing a leaf's decoration α by μ_0 (there are exactly (k + 1) such trees t' in $d_{\mathcal{T}}(t)$). Since $\mathcal{R}(\mathcal{P}) = \mathcal{P} \vee \mathcal{T}$, no two of these trees can cancel each other. For the same reason, since the trees appearing in the images $d_{\mathcal{T}}(s), d_{\mathcal{T}}(t)$ of two distinct trees s, t in λ are all distinct, no two of them can cancel each other. Therefore, if $d_{\mathcal{T}}(\lambda) = 0$, we must have that $\lambda = 0$.

Lemma 3.2. Let $k \in \mathbb{Z}_{\geq 0}$ and $\lambda \in \mathcal{R}_1^k(\mathcal{P})$. If $d_{\mathcal{T}}(\lambda) = 0$, then there exists a unique $\kappa \in \mathcal{R}_0^{k+1}(\mathcal{P})$ such that $d_{\mathcal{T}}(\kappa) = \lambda$.

Proof. Let t_0 be a tree in λ with one leaf ℓ_0 decorated by μ_0 and k leaves ℓ_1, \ldots, ℓ_k decorated by α . The image $d_{\mathcal{T}}(t_0)$ has one term which is the tree t'_0 with ℓ_0 and ℓ_1 decorated by μ_0 , and the other leaves decorated by α . Since $d_{\mathcal{T}}(\lambda) = 0$ and $\mathcal{R}(\mathcal{P}) = \mathcal{P} \vee \mathcal{T}$, the sum λ must also contain the only other tree t_1 whose image $d_{\mathcal{T}}(t_1)$ contains t'_0 , that is, the tree t_1 with ℓ_1 decorated by μ_0 and the other leaves decorated by α . Now, the image $d_{\mathcal{T}}(t_1)$ has one term which is the tree t'_1 with leaves ℓ_1 and ℓ_2 decorated by μ_0 , and the other leaves decorated by α . Since $d_{\mathcal{T}}(\lambda) = 0$, the sum λ must also contain the tree t_2 , with the leaf ℓ_2 decorated by μ_0 and the other leaves decorated by α . Continuing in this fashion, we obtain that λ must contain $d_{\mathcal{T}}(t^{\alpha})$, where t^{α} is the tree t with all the leaves ℓ_0, \ldots, ℓ_k decorated by α . Repeating the process for every distinct tree t in λ , and using again that $\mathcal{R}(\mathcal{P}) = \mathcal{P} \vee \mathcal{T}$, we obtain the desired element $\kappa := \sum_{t \in \lambda} t^{\alpha}$. The uniqueness of κ follows from Lemma 3.1.

Let $\mathcal{Q} \in \text{Curv}$, $\mathcal{P} \in \text{Op}$, and $f : \mathcal{L}(\mathcal{Q}) \to \mathcal{P}$. We want to prove that there exists a unique morphism $\Phi \in \text{hom}_{\text{Curv}}(\mathcal{Q}, \mathcal{R}(\mathcal{P}))$ such that $\pi \circ \Phi = f$. Given a morphism of graded operads (non-dg) $\Phi : \mathcal{Q} \to \mathcal{R}(\mathcal{P})$ and $k \in \mathbb{Z}_{\geq 0}$, we denote by Φ^k the induced map from \mathcal{Q} to $\mathcal{R}^k(\mathcal{P})$.

We start with uniqueness. Suppose given two maps with the desired property, i.e. some $\Phi_{-}, \Phi_{+} \in \hom_{Curv}(\mathcal{Q}, \mathcal{R}(\mathcal{P}))$ such that $\pi \circ \Phi_{\pm} = f$. We prove recursively that $\Phi_{+}^{k} = \Phi_{-}^{k}$ for every $k \in \mathbb{Z}_{\geq 0}$. Observe that, since the natural map of graded operads $\mathcal{Q}_{0} \lor [\mu_{0}] \to \mathcal{Q}$ is an isomorphism, and since Φ_{\pm} are morphisms of operads, it is enough to check the latter equality on elements of $\mathcal{Q}_{0} \cup {\{\mu_{0}\}}$. For k = 0, since Φ_{\pm} are morphisms in Curv, and $\pi \circ \Phi_{\pm}^{0} = \pi \circ \Phi_{\pm} = f$, we have

$$\Phi_{\pm}^{0}(\nu) = \begin{cases} f(\mu_{0}) + \mu_{0} & \text{if } \nu = \mu_{0}, \\ f(\nu) & \text{if } \nu \in \mathcal{Q}_{0}. \end{cases}$$

Assume now that, given $k \in \mathbb{Z}_{\geq 0}$, we proved $\Phi^j_+ = \Phi^j_-$ for every $j \in \{0, \ldots, k\}$. Using that Φ_{\pm} commute with the differentials, we get

$$d_{\mathcal{T}} \circ \Phi^{k+1}_{+} = d_{\mathcal{T}} \circ \Phi^{k+1}_{-} = \Phi^{k}_{\pm} \circ d_{\mathcal{Q}} - d_{\mathcal{P}} \circ \Phi^{k}_{\pm}$$

Let $\nu \in \mathcal{Q}_0 \cup \{\mu_0\}$. Since $\Phi_{\pm}(\mathcal{Q}_0) \subset \mathcal{P} \vee [\alpha]$ and $(\Phi_{\pm}(\mu_0) - \mu_0) \in \mathcal{P} \vee [\alpha]$, we get that $(\Phi_{\pm}^{k+1}(\nu) - \Phi_{\pm}^{k+1}(\nu)) \in \mathcal{R}_0^{k+1}(\mathcal{P})$ and is in the kernel of $d_{\mathcal{T}}$. According to Lemma 3.1, we get $\Phi_{\pm}^{k+1}(\nu) = \Phi_{\pm}^{k+1}(\nu)$. This concludes the proof of uniqueness.

We now prove the existence. More precisely, we recursively prove that, for every $k \in \mathbb{Z}_{\geq 0}$, there exists a family of N-modules morphisms $(\Phi^j : \mathcal{Q} \to \mathcal{R}^j(\mathcal{P}))_{0 \leq j \leq k}$ such that:

(1) We have

$$\Phi^{0}(\nu) = \begin{cases} f(\mu_{0}) + \mu_{0} & \text{if } \nu = \mu_{0}, \\ f(\nu) & \text{if } \nu \in \mathcal{Q}_{0}. \end{cases}$$

- (2) For every $j \in \{1, \ldots, k\}$ and $\nu \in \mathcal{Q}_0 \cup \{\mu_0\}$, we have $\Phi^j(\nu) \in \mathcal{R}_0^j(\mathcal{P})$.
- (3) For every $j \in \{0, \ldots, k\}$ and $\nu_1, \nu_2 \in \mathcal{Q}$, we have

$$\Phi^{j}(\nu_{1} \circ_{i} \nu_{2}) = \sum_{\substack{p,q \ge 0\\ p+q=j}} \Phi^{p}(\nu_{1}) \circ_{i} \Phi^{q}(\nu_{2}).$$

(4) For every $j \in \{0, ..., k - 1\}$, we have

$$d_{\mathcal{T}} \circ \Phi^{j+1} = \Phi^j \circ d_{\mathcal{Q}} - d_{\mathcal{P}} \circ \Phi^j.$$

It is then straightforward to show that we get a morphism $\Phi := \sum_{j\geq 0} \Phi^j : \mathcal{Q} \to \mathcal{R}(\mathcal{P})$ such that $\pi \circ \Phi = f$.

For k = 0, we define Φ^0 so that it satisfies (1) and (3) (these conditions define Φ^0 uniquely). Assume now that, given $k \in \mathbb{Z}_{\geq 0}$, we defined a family $(\Phi^j : \mathcal{Q} \to \mathcal{R}^j(\mathcal{P}))_{0 \leq j \leq k}$ satisfying the properties above. Given $\nu \in \mathcal{Q}_0 \cup {\mu_0}$, we consider

$$\lambda := (\Phi^k \circ d_{\mathcal{Q}} - d_{\mathcal{P}} \circ \Phi^k)(\nu).$$

First observe that $d_{\mathcal{T}}(\lambda) = 0$ if k = 0, and if $k \ge 1$ we also have

$$d_{\mathcal{T}}(\lambda) = (d_{\mathcal{T}} \circ \Phi^k \circ d_{\mathcal{Q}} - d_{\mathcal{T}} \circ d_{\mathcal{P}} \circ \Phi^k)(\nu)$$

= $(d_{\mathcal{T}} \circ \Phi^k \circ d_{\mathcal{Q}} + d_{\mathcal{P}} \circ d_{\mathcal{T}} \circ \Phi^k)(\nu)$
= $((\Phi^{k-1} \circ d_{\mathcal{Q}} - d_{\mathcal{P}} \circ \Phi^{k-1}) \circ d_{\mathcal{Q}} + d_{\mathcal{P}} \circ (\Phi^{k-1} \circ d_{\mathcal{Q}} - d_{\mathcal{P}} \circ \Phi^{k-1}))(\nu)$
= 0

We now proceed to prove that $\lambda \in \mathcal{R}_1^k(\mathcal{P})$. One key observation is that, since $\mathcal{Q} \in \text{Curv}$ and $\nu \in \mathcal{Q}_0 \cup \{\mu_0\}$, we have $\nu, d_{\mathcal{Q}}\nu \in \mathcal{Q}_1$. We now treat two cases separately:

(1) If k = 0, first observe that

$$\Phi^{0}(\nu) = f(\nu) + \xi$$
 and $\Phi^{0}(d_{\mathcal{Q}}\nu) = f(d_{\mathcal{Q}}\nu) + \xi'$,

where $\xi, \xi' \in \mathcal{R}^0_1(\mathcal{P})$. Then we have

$$\lambda = \Phi^0(d_{\mathcal{Q}}\nu) - d_{\mathcal{P}}(\Phi^0(\nu))$$

= $f(d_{\mathcal{Q}}\nu) + \xi' - d_{\mathcal{P}}(f(\nu)) - d_{\mathcal{P}}(\xi)$
= $d_{\mathcal{P}}f(\nu) + \xi' - d_{\mathcal{P}}f(\nu) - d_{\mathcal{P}}(\xi)$
= $\xi' - d_{\mathcal{P}}(\xi).$

Therefore we have $\lambda \in \mathcal{R}^0_1(\mathcal{P})$.

(2) If $k \geq 1$, first observe that, since $\Phi^k(\mathcal{Q}_1) \subset \mathcal{R}_0^k(\mathcal{P}) \oplus \mathcal{R}_1^k(\mathcal{P})$ (because of assumption (2)) and $\nu, d_{\mathcal{Q}}\nu \in \mathcal{Q}_1$, we have $\lambda \in \mathcal{R}_0^k(\mathcal{P}) \oplus \mathcal{R}_1^k(\mathcal{P})$. We write $\lambda = \lambda_0 + \lambda_1$ accordingly. Since $d_{\mathcal{T}}(\lambda) = 0$ and $d_{\mathcal{T}}(\mathcal{R}_{\ell}^k(\mathcal{P})) \subset \mathcal{R}_{\ell+1}^{k-1}(\mathcal{P})$, we get that $d_{\mathcal{T}}(\lambda_0) = d_{\mathcal{T}}(\lambda_1) = 0$. It follows from Lemma 3.1 that $\lambda_0 = 0$, and therefore we have $\lambda = \lambda_1 \in \mathcal{R}_1^k(\mathcal{P})$.

Now, according to Lemma 3.2, there exists a unique $\kappa \in \mathcal{R}_0^{k+1}(\mathcal{P})$ such that $\lambda = d_{\mathcal{T}}(\kappa)$. We define $\Phi^{k+1}(\nu) := \kappa$.

We have now defined Φ^{k+1} on $\mathcal{Q}_0 \cup \{\mu_0\}$, and it satisfies (2). Using the assumptions satisfied by $(\Phi^j)_{0 \leq j \leq k}$, it is straightforward to show that, for every $\nu_1, \nu_2 \in \mathcal{Q}_0$, the sum

$$\sum_{\substack{p,q\geq 0\\p+q=k+1}} \Phi^p(\nu_1) \circ_i \Phi^q(\nu_2)$$

is in $\mathcal{R}_0^{k+1}(\mathcal{P})$ and is mapped by $d_{\mathcal{T}}$ to $(\Phi^k \circ d_{\mathcal{Q}} - d_{\mathcal{P}} \circ \Phi^k)(\nu_1 \circ_i \nu_2)$. By uniqueness, this implies that, for every $\nu_1, \nu_2 \in \mathcal{Q}_0$, we have

$$\Phi^{k+1}(\nu_1 \circ_i \nu_2) = \sum_{\substack{p,q \ge 0\\ p+q=k+1}} \Phi^p(\nu_1) \circ_i \Phi^q(\nu_2).$$

We can now extend Φ^{k+1} to \mathcal{Q} so that it satisfies (3). Finally, Φ^{k+1} satisfies (4) by definition. This completes the proof. \Box

4. PROOF OF THEOREM 1.13

We must show that for any $\mathcal{Q} \in \text{Curv}$ such that $d_{\mathcal{Q}}(\mu_0)$ is in the non-dg operadic ideal generated by μ_0 , there is a unique morphism $\Phi_{\mathcal{Q}} \in \text{hom}_{\text{Curv}}(cA_{\infty}, \mathcal{Q})$ such that $\Phi_{\mathcal{Q}}(\mu_0) = \mu_0$.

First observe that, since $d_{\mathcal{Q}}(\mu_0)$ belongs to the non-dg ideal (μ_0) , the natural morphism of non-dg operads $\mathcal{Q}_0 \to \mathcal{Q}/(\mu_0)$ is an isomorphism. This gives a dg-operad structure on \mathcal{Q}_0 .

We start by proving uniqueness. Let Φ_+ , Φ_- be two elements of $\hom_{Curv}(cA_{\infty}, \mathcal{Q})$ such that $\Phi_{\pm}(\mu_0) = \mu_0$. We recursively prove that, for every $k \in \mathbb{Z}_{\geq 0}$, we have $\Phi_+(\mu_j) = \Phi_-(\mu_j)$

for every $j \in \{0, ..., k\}$. For k = 0, this follows by assumption. Assume that, given $k \in \mathbb{Z}_{\geq 0}$, we proved that $\Phi_+(\mu_j) = \Phi_-(\mu_j)$ for every $j \in \{0, ..., k\}$. Recall that the differential of cA_{∞} is given by

$$-d_{cA_{\infty}}(\mu_{k}) = \sum_{\substack{p,r \ge 0, q \ge 1\\ p+q+r=k}} \mu_{p+1+r} \circ_{p+1} \mu_{q} + [\mu_{k+1}, \mu_{0}].$$

Since Φ_{\pm} are morphisms of dg-operads, we get

$$[\Phi_{+}(\mu_{k+1}),\mu_{0}] = [\Phi_{-}(\mu_{k+1}),\mu_{0}] = -d_{\mathcal{Q}}(\Phi_{\pm}(\mu_{k})) - \sum_{\substack{p,r \ge 0, q \ge 1\\p+q+r=k}} \Phi_{\pm}(\mu_{p+1+r}) \circ_{p+1} \Phi_{\pm}(\mu_{q}).$$

It follows that $\Phi_+(\mu_{k+1}) = \Phi_-(\mu_{k+1})$, which proves uniqueness.

Before proving existence, we observe that, given $\nu \in \mathcal{Q}_0(k)$, there exists a family of operations $(\nu^i)_{1 \leq i \leq k+1}$ in $\mathcal{Q}_0(k+1)$ such that

(4.1)
$$-d_{\mathcal{Q}}(\nu) = -d_{\mathcal{Q}_0}(\nu) + \sum_{i=1}^{k+1} \nu^i \circ_i \mu_0.$$

Indeed one can write $-d_{\mathcal{Q}}(\nu) = \lambda_0 + \lambda_1$ with $\lambda_0 \in \mathcal{Q}_0$ and $\lambda_1 \in \mathcal{Q}_1 \cap (\mu_0)$. We already know that λ_1 is of the desired form. Now we use the fact that $\pi : \mathcal{Q} \to \mathcal{Q}/(\mu_0)$ is a morphism of dg-operads to write

$$\pi(\lambda_0) = \pi(\lambda_0 + \lambda_1) = \pi(-d_{\mathcal{Q}}(\nu)) = -d_{\mathcal{Q}/(\mu_0)}(\pi(\nu))$$

We now recursively prove that, for every $k \in \mathbb{Z}_{\geq 1}$, there exists a family of operations $(\nu_j)_{0 \leq j \leq k}$ such that

- (1) $\nu_0 = \mu_0$,
- (2) for every $j \in \{1, \ldots, k\}$, the operation ν_j has degree (-1) and is in $\mathcal{Q}_0(j)$,
- (3) for every $j \in \{0, \ldots, k-1\}$, we have

$$-d_{\mathcal{Q}}(\nu_j) = \sum_{\substack{p,q,r \ge 0\\ p+q+r=j}} \nu_{p+1+r} \circ_{p+1} \nu_q.$$

It will follow that the assignment $\mu_k \mapsto \nu_k$ defines an element $\Phi \in \hom_{Curv}(cA_{\infty}, \mathcal{Q})$ such that $\Phi(\mu_0) = \mu_0$.

For k = 1, we set $\nu_0 := \mu_0$. Since $\mathcal{Q} \in \text{Curv}$ and $d_{\mathcal{Q}}(\mu_0) \in (\mu_0)$, there exists $\nu_1 \in \mathcal{Q}_0(1)$ of degree (-1) such that $-d_{\mathcal{Q}}(\mu_0) = \nu_1(\mu_0)$.

Assume now that, given $k \in \mathbb{Z}_{\geq 1}$, we have defined a family $(\nu_j)_{0 \leq j \leq k}$ satisfying the properties above. According to Eq. (4.1), there is a family $(\nu_{k+1}^i)_{1 \leq i \leq k+1}$ in $\mathcal{Q}_0(k+1)$ such that

(4.2)
$$-d_{\mathcal{Q}}(\nu_k) = -d_{\mathcal{Q}_0}(\nu_k) + \sum_{i=1}^{k+1} \nu_{k+1}^i \circ_i \mu_0.$$

Moreover, for any $0 \le j \le k - 1$ we have

(4.3)
$$-d_{\mathcal{Q}_0}(\nu_j) = \sum_{\substack{p,r \ge 0, q \ge 1\\p+q+r=j}} \nu_{p+1+r} \circ_{p+1} \nu_q.$$

Unravelling the consequences of the relation $d_{\mathcal{Q}}^2 = 0$ we have

$$\begin{aligned} 0 &= d_{\mathcal{Q}}^{2}(\nu_{k-1}) \\ &= d_{\mathcal{Q}}\left(-\sum_{\substack{p,q,r \geq 0\\p+q+r=k-1}} \nu_{p+1+r} \circ_{p+1} \nu_{q}\right) \\ &= d_{\mathcal{Q}}(-[\nu_{k},\mu_{0}]) - d_{\mathcal{Q}}\left(\sum_{\substack{p,r \geq 0,q \geq 1\\p+q+r=k-1}} \nu_{p+1+r} \circ_{p+1} \nu_{q}\right) \\ &= \left[-d_{\mathcal{Q}}(\nu_{k}),\mu_{0}\right] + \left[\nu_{k},d_{\mathcal{Q}}(\mu_{0})\right] - \sum_{\substack{p,r \geq 0,q \geq 1\\p+q+r=k-1}} (d_{\mathcal{Q}}(\nu_{p+1+r}) \circ_{p+1} \nu_{q} - \nu_{p+1+r} \circ_{p+1} d_{\mathcal{Q}}(\nu_{q})) \\ &= \left[-d_{\mathcal{Q}}(\nu_{k}),\mu_{0}\right] + \left[\nu_{k},d_{\mathcal{Q}}(\mu_{0})\right] \\ &- \sum_{\substack{p,r \geq 0,q \geq 1\\p+q+r=k-1}} ((d_{\mathcal{Q}_{0}}(\nu_{p+1+r}) - [\nu_{p+2+r},\mu_{0}]) \circ_{p+1} \nu_{q} - \nu_{p+1+r} \circ_{p+1} (d_{\mathcal{Q}_{0}}(\nu_{q}) - [\nu_{q+1},\mu_{0}])) \\ &= \left[-d_{\mathcal{Q}}(\nu_{k}),\mu_{0}\right] + \left[\nu_{k},d_{\mathcal{Q}}(\mu_{0})\right] + \sum_{\substack{p,r \geq 0,q \geq 1\\p+q+r=k-1}} ([\nu_{p+2+r},\mu_{0}] \circ_{p+1} \nu_{q} - \nu_{p+1+r} \circ_{p+1} [\nu_{q+1},\mu_{0}])) \end{aligned}$$

since $d_{\mathcal{Q}_0}(\nu_{k-1})$ is given by the A^+_{∞} -relations Eq. (4.3) and $d^2_{\mathcal{Q}_0}(\nu_{k-1}) = 0$. Further, using the decomposition Eq. (4.2), we have that

$$0 = \left[-d_{\mathcal{Q}_{0}}(\nu_{k}), \mu_{0}\right] + \left[\sum_{i=1}^{k+1} \nu_{k+1}^{i} \circ_{i} \mu_{0}, \mu_{0}\right] - \left[\nu_{k}, \left[\nu_{1}, \mu_{0}\right]\right] \\ + \sum_{\substack{p,r \geq 0, q \geq 1 \\ p+q+r=k-1}} \left(\left[\nu_{p+2+r}, \mu_{0}\right] \circ_{p+1} \nu_{q} - \nu_{p+1+r} \circ_{p+1} \left[\nu_{q+1}, \mu_{0}\right]\right)$$

Now we have two types of terms in this expression: terms with exactly one μ_0 , and terms with exactly two μ_0 . Both of these groups of terms must vanish separately. In the second case, we have

$$\left[\sum_{i=1}^{k+1} \nu_{k+1}^i \circ_i \mu_0, \mu_0\right] = 0$$

which implies that we must have $\nu_{k+1}^i = \nu_{k+1}^j$ for all $1 \leq i, j \leq k+1$. Therefore, we define $\nu_{k+1} := \nu_{k+1}^i = \nu_{k+1}^j$. Looking back at Eq. (4.2) this implies that

(4.4)
$$-d_{\mathcal{Q}}(\nu_k) = -d_{\mathcal{Q}_0}(\nu_k) + [\nu_{k+1}, \mu_0].$$

As for the first type of terms, using the Jacobi identity we get

$$\begin{array}{ll} 0 & = & \left[-d_{\mathcal{Q}_{0}}(\nu_{k}) - [\nu_{k}, \nu_{1}], \mu_{0} \right] - \left[[\nu_{k}, \mu_{0}], \nu_{1} \right] \\ & + \sum_{\substack{p,r \geq 0, q \geq 1 \\ p+q+r=k-1}} \left(\left[\nu_{p+2+r}, \mu_{0} \right] \circ_{p+1} \nu_{q} - \nu_{p+1+r} \circ_{p+1} \left[\nu_{q+1}, \mu_{0} \right] \right) \right) \\ & = & \left[-d_{\mathcal{Q}_{0}}(\nu_{k}) - [\nu_{k}, \nu_{1}], \mu_{0} \right] \\ & + \sum_{\substack{p,r \geq 0, q \geq 2 \\ p+q+r=k-1}} \left[\nu_{p+2+r}, \mu_{0} \right] \circ_{p+1} \nu_{q} - \sum_{\substack{p,r \geq 0, 1 \leq q \leq k-2 \\ p+q+r=k-1}} \nu_{p+1+r} \circ_{p+1} \left[\nu_{q+1}, \mu_{0} \right] . \end{array}$$

We claim that we have

(4.5)
$$\sum_{\substack{p,r \ge 0,q \ge 2\\p+q+r=k-1}} [\nu_{p+2+r},\mu_0] \circ_{p+1} \nu_q - \sum_{\substack{p,r \ge 0,1 \le q \le k-2\\p+q+r=k-1}} \nu_{p+1+r} \circ_{p+1} \mu_q]$$
$$= \left[-\sum_{\substack{p,r \ge 0,q \ge 2\\p+q+r=k}} \nu_{p+1+r} \circ_{p+1} \nu_q,\mu_0 \right],$$

which will end the proof since we will then have

$$0 = \left[-d_{\mathcal{Q}_0}(\nu_k) - \sum_{\substack{p,r \ge 0, q \ge 1 \\ p+q+r=k}} \nu_{p+1+r} \circ_{p+1} \nu_q, \mu_0 \right],$$

which implies that

(4.6)
$$-d_{\mathcal{Q}_0}(\nu_k) = \sum_{\substack{p,r \ge 0, q \ge 1\\ p+q+r=k}} \nu_{p+1+r} \circ_{p+1} \nu_q ,$$

and combining this with Eq. (4.4), we get that ν_k satisfies the cA_{∞} -relations in Q. Thus, it remains to prove Eq. (4.5). We have

$$\begin{bmatrix} -\sum_{\substack{p,r \ge 0,q \ge 2\\p+q+r=k}} \nu_{p+1+r} \circ_{p+1} \nu_q, \mu_0 \end{bmatrix} + \sum_{\substack{p,r \ge 0, \ 1 \le q \le k-2\\p+q+r=k-1}} \nu_{p+1+r} \circ_{p+1} [\nu_{q+1}, \mu_0] \\ = \begin{bmatrix} -\sum_{\substack{p,r \ge 0,q \ge 2\\p+q+r=k}} \nu_{p+1+r} \circ_{p+1} \nu_q, \mu_0 \end{bmatrix} + \sum_{\substack{p,r \ge 0,q \ge 2\\p+q+r=k}} \nu_{p+1+r} \circ_{p+1} [\nu_q, \mu_0] \\ = \sum_{\substack{p,r \ge 0,q \ge 2\\p+q+r=k}} \left(-\sum_{i=1}^p (\nu_{p+1+r} \circ_{p+1} \nu_q) \circ_i \mu_0 - \sum_{i=p+q+1}^k (\nu_{p+1+r} \circ_{p+1} \nu_q) \circ_i \mu_0 \right),$$

where the last equality is obtained from the previous one by keeping the terms in the first sum not cancelled by those in the second. We further have

$$\begin{split} &\sum_{\substack{p,r \geq 0,q \geq 2\\p+q+r=k}} \left(-\sum_{i=1}^{p} (\nu_{p+1+r} \circ_{p+1} \nu_q) \circ_i \mu_0 - \sum_{i=p+q+1}^{k} (\nu_{p+1+r} \circ_{p+1} \nu_q) \circ_i \mu_0 \right) \\ &= \sum_{\substack{p,r \geq 0,q \geq 2\\p+q+r=k}} \left(\sum_{i=1}^{p} (\nu_{p+1+r} \circ_i \mu_0) \circ_p \nu_q + \sum_{i=p+q+1}^{k} (\nu_{p+1+r} \circ_{i-q+1} \mu_0) \circ_{p+1} \nu_q \right) \\ &= \sum_{\substack{p,r \geq 0,q \geq 2\\p+q+r=k}} \left(\sum_{i=1}^{p} (\nu_{p+1+r} \circ_i \mu_0) \circ_p \nu_q + \sum_{i=p+2}^{p+1+r} (\nu_{p+1+r} \circ_i \mu_0) \circ_{p+1} \nu_q \right) \\ &= \sum_{\substack{p\geq 1,r \geq 0,q \geq 2\\p+q+r=k}} \left(\sum_{i=1}^{p} (\nu_{p+1+r} \circ_i \mu_0) \circ_p \nu_q \right) + \sum_{\substack{p\geq 0,r \geq 1,q \geq 2\\p+q+r=k}} \left(\sum_{i=p+2}^{p+1+r} (\nu_{p+1+r} \circ_i \mu_0) \circ_{p+1} \nu_q \right) \\ &= \sum_{\substack{p,r \geq 0,q \geq 2\\p+q+r=k-1}} \left(\sum_{i=1}^{p+1} (\nu_{p+2+r} \circ_i \mu_0) \circ_{p+1} \nu_q + \sum_{i=p+2}^{p+2+r} (\nu_{p+2+r} \circ_i \mu_0) \circ_{p+1} \nu_q \right) \\ &= \sum_{\substack{p,r \geq 0,q \geq 2\\p+q+r=k-1}} \left[\nu_{p+2+r}, \mu_0 \right] \circ_{p+1} \nu_q \,, \end{split}$$

which yields the desired equality. \Box

Given an operad \mathcal{Q} in Curv, we constructed a unique morphism $\operatorname{Tw}_{\mathcal{Q}} : \mathcal{Q} \to \mathcal{Q}/\mu_0 \lor \mathcal{T}$ in the comma category $(\mathcal{L} \downarrow \mathcal{Q}/\mu_0)$. In the case where $d_{\mathcal{Q}}(\mu_0)$ is in the operadic ideal generated by μ_0 , Theorem 1.13 gives us a unique morphism $\Phi_{\mathcal{Q}} : cA_{\infty} \to \mathcal{Q}$ such that $\Phi_{\mathcal{Q}}(\mu_0) = \mu_0$. The composition $\operatorname{Tw}_{\mathcal{Q}} \circ \Phi_{\mathcal{Q}}$ has an explicit description.

Corollary 4.1. Let $Q \in \text{Curv}$ such that $d_Q(\mu_0)$ belongs to the non-dg ideal generated by μ_0 . Then the composite

$$cA_{\infty} \xrightarrow{\Phi_{\mathcal{Q}}} \mathcal{Q} \xrightarrow{Tw_{\mathcal{Q}}} \mathcal{Q}/\mu_0 \vee \mathcal{T}$$

is given by the assignment

$$\mu_n \mapsto \mu_n^{\alpha} := \sum_{r_0, \dots, r_n \ge 0} \mu_{n+r_0+\dots+r_n} (\alpha^{r_0}, -, \alpha^{r_1}, -, \dots, \alpha^{r_n}) ,$$

for all $n \geq 0$ (here we abuse notations by writing μ_n for both the element in cA_{∞} and its image in \mathcal{Q} under $\Phi_{\mathcal{Q}}$). In particular, we have $Tw_{\mathcal{Q}}(\mu_0) = \mu_0^{\alpha}$.

Proof. Consider the quotient maps $\pi_{\mathcal{Q}} : \mathcal{Q} \to \mathcal{Q}/\mu_0$ and $\pi_{\mathcal{Q}/\mu_0} : \mathcal{Q}/\mu_0 \lor \mathcal{T} \to \mathcal{Q}/\mu_0$, and recall that $\operatorname{Tw}_{\mathcal{Q}}$ is (by definition) the unique morphism in the comma category $(\mathcal{L} \downarrow \mathcal{Q}/\mu_0)$ from $(\mathcal{Q}, \pi_{\mathcal{Q}})$ to $(\mathcal{Q}/\mu_0 \lor \mathcal{T}, \pi_{\mathcal{Q}/\mu_0})$. It follows that $\operatorname{Tw}_{\mathcal{Q}} \circ \Phi_{\mathcal{Q}}$ is the unique morphism in $(\mathcal{L} \downarrow \mathcal{Q}/\mu_0)$ from $(\operatorname{cA}_{\infty}, \pi_{\mathcal{Q}} \circ \Phi_{\mathcal{Q}})$ to $(\mathcal{Q}/\mu_0 \lor \mathcal{T}, \pi_{\mathcal{Q}/\mu_0})$. Therefore the result will follow if we show that the assignment

$$\Psi: \mu_n \mapsto \mu_n^{\alpha} := \sum_{r_0, \dots, r_n \ge 0} \mu_{n+r_0+\dots+r_n}(\alpha^{r_0}, -, \alpha^{r_1}, -, \dots, \alpha^{r_n})$$

defines a morphism in $(\mathcal{L} \downarrow \mathcal{Q}/\mu_0)$ from $(cA_{\infty}, \pi_{\mathcal{Q}} \circ \Phi_{\mathcal{Q}})$ to $(\mathcal{Q}/\mu_0 \lor \mathcal{T}, \pi_{\mathcal{Q}/\mu_0})$.

Since $\Phi_{\mathcal{Q}}(\mu_0) = \mu_0$, we can write $\pi_{\mathcal{Q}} \circ \Phi_{\mathcal{Q}} = \Phi_{\mathcal{Q}/\mu_0} \circ \pi_{cA_{\infty}}$ with $\Phi_{\mathcal{Q}/\mu_0} : A^+_{\infty} \to \mathcal{Q}/\mu_0$. Now the key observation is that, according to Corollary 1.10, we have

$$\Psi = (\Phi_{\mathcal{Q}/\mu_0} \lor \mathrm{id}_{\mathcal{T}}) \circ \mathrm{Tw}_{\mathrm{cA}_{\infty}}$$

Since $\operatorname{Tw}_{cA_{\infty}}$ is the unique morphism in $(\mathcal{L} \downarrow A_{\infty}^+)$ from $(cA_{\infty}, \pi_{cA_{\infty}})$ to $(A_{\infty}^+ \lor \mathcal{T}, \pi_{A_{\infty}^+})$, it follows that Ψ is a morphism in $(\mathcal{L} \downarrow \mathcal{Q}/\mu_0)$ from $(cA_{\infty}, \Phi_{\mathcal{Q}/\mu_0} \circ \pi_{cA_{\infty}}) = (cA_{\infty}, \pi_{\mathcal{Q}} \circ \Phi_{\mathcal{Q}})$ to $(\mathcal{Q}/\mu_0 \lor \mathcal{T}, \pi_{\mathcal{Q}/\mu_0})$. This concludes the proof.

5. MAURER-CARTAN ELEMENTS

We can now define the *Maurer-Cartan operad* associated with any operad Q in Curv such that $d_Q(\mu_0)$ belongs to the non-dg ideal generated by μ_0 . Recall that $[\alpha]$ denotes the operad with a single nonvanishing operation α of arity 0 and degree 0 (in positive filtration).

Definition-Proposition 5.1. Let $\mathcal{Q} \in \text{Curv}$ such that $d_{\mathcal{Q}}(\mu_0)$ belongs to the non-dg ideal generated by μ_0 . Its *Maurer-Cartan operad* MC(\mathcal{Q}) is the operad with underlying non-dg operad $\mathcal{Q} \vee [\alpha]$ and differential

$$(d_{\mathrm{MC}(\mathcal{Q})})|_{\mathcal{Q}} := d_{\mathcal{Q}}, \quad d_{\mathrm{MC}(\mathcal{Q})}(\alpha) := -\mu_0^{\alpha}.$$

Proof. We have to check that $d^2_{MC(\mathcal{Q})}(\alpha) = 0$. We compute

$$\begin{aligned} d_{\mathrm{MC}(\mathcal{Q})}^{2}(\alpha) &= d_{\mathrm{MC}(\mathcal{Q})}\left(-\sum_{n\geq 0}\mu_{n}(\alpha^{n})\right) \\ &= \sum_{n\geq 0}\sum_{\substack{p,q,r\geq 0\\p+q+r=n}}(\mu_{p+1+r}\circ_{p+1}\mu_{q})(\alpha^{n}) - \sum_{n\geq 0}\sum_{p=0}^{n-1}\mu_{n}\left(\alpha^{p},\sum_{q\geq 0}\mu_{q}(\alpha^{q}),\alpha^{n-p-1}\right) \\ &= \sum_{k\geq 0}\left(\sum_{\substack{p,q,r\geq 0\\p+q+r=k}}(\mu_{p+1+r}\circ_{p+1}\mu_{q})(\alpha^{k}) - \sum_{\substack{p,q\geq 0,n\geq p+1\\q+n-1=k}}(\mu_{n}\circ_{p+1}\mu_{q})(\alpha^{k})\right) \\ &= 0\end{aligned}$$

The operad $MC(\mathcal{Q})$ is naturally an object of Curv, and moreover of the comma category $(\mathcal{L} \downarrow \mathcal{Q}/\mu_0)$. Its algebras are \mathcal{Q} -algebras A together with an element $\alpha \in A$ of degree 0 such that

 \square

$$\mu_0^{\alpha} = \mu_0 + \mu_1(\alpha) + \mu_2(\alpha, \alpha) + \dots = 0.$$


We call such an $\alpha \in A$ a *Maurer-Cartan element* of the *Q*-algebra *A*.

Theorem 5.2. Let $\mathcal{Q} \in \text{Curv}$ such that $d_{\mathcal{Q}}(\mu_0)$ belongs to the non-dg ideal generated by μ_0 . The unique morphism in the comma category $(\mathcal{L} \downarrow \mathcal{Q}/\mu_0)$ from MC(\mathcal{Q}) to $\mathcal{Q}/\mu_0 \lor \mathcal{T}$ is

$$\operatorname{Tw}_{\mathcal{Q}} \lor (-\operatorname{id}_{\alpha}) : \operatorname{MC}(\mathcal{Q}) \to \mathcal{Q}/\mu_0 \lor \mathcal{T}$$

and is an isomorphism. Therefore, MC(Q) is terminal in $(\mathcal{L} \downarrow Q/\mu_0)$ and the datum of a $Q/\mu_0 \lor \mathcal{T}$ -algebra is equivalent to the datum of a Q-algebra with Maurer-Cartan element.

Proof. Denote by $\psi_{\mathcal{Q}}$ the unique morphism in the comma category $(\mathcal{L} \downarrow \mathcal{Q}/\mu_0)$ from MC(\mathcal{Q}) to $\mathcal{Q}/\mu_0 \lor \mathcal{T}$. First we check that $\psi_{\mathcal{Q}} = \text{Tw}_{\mathcal{Q}} \lor (-\text{id}_{\alpha})$. Consider the commutative diagram:



As $\operatorname{Tw}_{\mathcal{Q}}$ is the unique morphism from \mathcal{Q} to $\mathcal{Q}/\mu_0 \vee \mathcal{T}$ in the comma category $(\mathcal{L} \downarrow \mathcal{Q}/\mu_0)$, we must have that $\psi_{\mathcal{Q}}$ is equal to $\operatorname{Tw}_{\mathcal{Q}}$ on \mathcal{Q} . We claim that $\psi_{\mathcal{Q}}(\alpha) = -\alpha$. The only non-trivial thing to check is that $\operatorname{Tw}_{\mathcal{Q}} \vee (-\operatorname{id}_{\alpha})$ commutes with the differentials. This is immediate for elements of \mathcal{Q} ; let us check it for α as well, i.e. that $d_{\mathcal{Q}/\mu_0 \vee \mathcal{T}}(-\alpha) = (\operatorname{Tw}_{\mathcal{Q}} \vee - \operatorname{id}_{\alpha})(d_{\operatorname{MC}(\mathcal{Q})}\alpha)$. Using the explicit description of $\operatorname{Tw}_{\mathcal{Q}}$ in Corollary 4.1, we need to check that

$$-\mu_0 = (\mathrm{Tw}_{\mathcal{Q}} \vee -\mathrm{id}_\alpha) \left(-\sum_{n\geq 0} \mu_n(\alpha^n)\right) = -\sum_{n\geq 0} (-1)^n \mu_n^\alpha(\alpha^n).$$

This is a straightforward computation: we have

$$\sum_{n\geq 0} (-1)^n \mu_n^{\alpha}(\alpha^n) = \sum_{n\geq 0} (-1)^n \sum_{r_0,\dots,r_n\geq 0} \mu_{n+r_0+\dots+r_n}(\alpha^{r_0},-,\alpha^{r_2},\dots,-,\alpha^{r_n})(\alpha^n)$$
$$= \sum_{k\geq 0} \sum_{n=0}^k (-1)^n \binom{k}{n} \mu_k(\alpha^k)$$
$$= \sum_{k\geq 0} (1-1)^k \mu_k(\alpha^k)$$
$$= \mu_0.$$

We now prove that $\operatorname{Tw}_{\mathcal{Q}} \lor (-\operatorname{id}_{\alpha}) : \operatorname{MC}(\mathcal{Q}) \to \mathcal{Q}/\mu_0 \lor \mathcal{T}$ is an isomorphism. In order to do that, we consider on both sides the (decreasing) filtration by the number of α and show that the morphism induced on the associated graded is an isomorphism. Writing $\operatorname{Tw}_{\mathcal{Q}} = \sum_{k\geq 0} \operatorname{Tw}_{\mathcal{Q}}^k$ according to the grading by the number of α , the induced morphism between associated graded is $\operatorname{Tw}_{\mathcal{Q}}^0 \lor (-\operatorname{id}_{\alpha})$. It remains to observe that $\operatorname{Tw}_{\mathcal{Q}}^0 = \operatorname{id}_{\mathcal{Q}}$ since we have $(\operatorname{Tw}_{\mathcal{Q}}(\mu_0) - \mu_0) \in \mathcal{Q}/\mu_0$ and $\pi_{\mathcal{Q}/\mu_0} \circ \operatorname{Tw}_{\mathcal{Q}}^0 = \pi_{\mathcal{Q}/\mu_0} \circ \operatorname{Tw}_{\mathcal{Q}} = \pi_{\mathcal{Q}}$.

References

- [1] Damien Calaque, Ricardo Campos, and Joost Nuiten. Lie algebroids are curved Lie algebras. arXiv:2103.10728.
- [2] Joseph Chuang and Andrey Lazarev. Combinatorics and Formal Geometry of the Maurer-Cartan Equation. Letters in Mathematical Physics, 103(1):79–112, 2013.
- [3] Vasily Dolgushev and Thomas Willwacher. Operadic twisting With an application to Deligne's conjecture. Journal of Pure and Applied Algebra, 219(5):1349–1428, 2015.
- [4] Vladimir Dotsenko, Sergey Shadrin, and Bruno Vallette. Maurer-Cartan Methods in Deformation Theory: The Twisting Procedure. London Mathematical Society Lecture Note Series. Cambridge University Press, Cambridge, 2024.
- [5] Kenji Fukaya. Morse homotopy, A[∞]-category, and Floer homologies. In Proc. of the GARC Workshop on Geometry and Topology'93, Seoul, 1993, pages 1–102. Seoul Nat. Univ., 1993.

- [6] Kenji Fukaya, Yong-Geun Oh, Hiroshi Ohta, and Kaoru Ono. Lagrangian intersection Floer theory: anomaly and obstruction, Part I. American Mathematical Soc., 2010.
- [7] Maxim Kontsevich and Yan Soibelman. Notes on A_∞-algebras, A_∞-categories and non-commutative geometry. In *Homological Mirror Symmetry: New Developments and Perspectives*, pages 1–67. Springer, 2008.
- [8] Daniel Quillen. Rational homotopy theory. Annals of Mathematics, 90(2):205–295, 1969.