

# Expansion into vacuum of stochastic gases with long-range interactions

P. L. Krapivsky<sup>1,2</sup> and Kirone Mallick<sup>3</sup>

<sup>1</sup>*Department of Physics, Boston University, Boston, Massachusetts 02215, USA*

<sup>2</sup>*Santa Fe Institute, Santa Fe, New Mexico 87501, USA*

<sup>3</sup>*Institut de Physique Théorique, CEA/Saclay, F-91191 Gif-sur-Yvette Cedex, France*

We study the evolution of a system of many point particles initially concentrated in a small region in  $d$  dimensions. Particles undergo overdamped motion caused by pairwise interactions through the long-ranged repulsive  $r^{-s}$  potential; each particle is also subject to Brownian noise. When  $s < d$ , the expansion is governed by non-local hydrodynamic equations. In the one-dimensional case, we deduce self-similar solutions for all  $s \in (-2, 1)$ . The expansion of Coulomb gases remains well-defined in the infinite-particle limit: The density is spatially uniform and inversely proportional to time independent of the spatial dimension.

## I. INTRODUCTION

Many-particle systems with long-range interactions [1, 2] display unusual collective behaviors and fluctuations. Probing these behaviors requires an extension of the techniques (such as hydrodynamic limits and large deviations) initially developed studies of systems with short-range forces.

In the present work, we consider point particles in  $\mathbb{R}^d$  undergoing independent Brownian motions and an overdamped motion caused by the repulsive Riesz potential

$$V(\mathbf{r}) = \frac{g}{sr^s} \text{ with } g > 0. \quad (1)$$

When the decay exponent  $s$  is smaller than the dimension of space,  $s < d$ , the interaction is long-ranged, viz., it dominates diffusion for a sufficiently long time, even forever, if  $s < 0$ . In the present work, we consider long-ranged Riesz potentials.

In the  $s \rightarrow 0$  limit, the Riesz potential becomes logarithmic

$$V(\mathbf{r}) = -g \ln r \quad (2)$$

In one dimension, a gas of Brownian particles interacting via logarithmic potential was introduced by Dyson [3] in the context of random matrices [3, 4] where eigenvalues played the role of particles. In two dimensions, a gas of Brownian particles interacting via logarithmic potential was introduced by Ginibre [5] also in the context of random matrices, with complex eigenvalues playing the role of particles [4–6]. In the Dyson and Ginibre gases, the particles undergo an overdamped motion caused by the logarithmic potential and are subject to Brownian noises. The Dyson and Ginibre gases suggest exploring stochastic Riesz gases with arbitrary long-ranged Riesz potential and in arbitrary dimensions. Deterministic Riesz gases with particles moving deterministically and interacting through Riesz potential is a more explored subject [7, 8], but in this paper, we consider only *stochastic* Riesz gases.

Apart from logarithmic gases corresponding to  $s = 0$ , several other values of the exponent  $s$  correspond to interesting classical models. The  $r^{-2}$  potential is the Calogero gas mostly studied in one dimension [9, 10], albeit it

makes sense in arbitrary dimension  $d$ ; the Calogero gas is long-ranged when  $d > 2$ . When  $s = d - 2$ , the Riesz potential reduces to the Coulomb potential

$$V(\mathbf{r}) = \frac{g}{(d-2)r^{d-2}} \quad (3)$$

In two dimensions, the Coulomb potential is logarithmic.

The equation of motion of the position  $\mathbf{x}_i(t) \in \mathbb{R}^d$  of the particle with label  $i$  is

$$\frac{d\mathbf{x}_i}{dt} = g \sum_{j \neq i} \frac{\mathbf{x}_i - \mathbf{x}_j}{|\mathbf{x}_i - \mathbf{x}_j|^{2+s}} + \boldsymbol{\eta}_i(t) \quad (4)$$

In addition to the force caused by pairwise interactions, each particle is subjected to the white noise  $\boldsymbol{\eta}_i(t)$ . The noises are independent and cause each particle to diffuse with diffusion coefficient  $D$  if a particle is alone.

We shall always assume that the number of particles is large,  $N \gg 1$ . In this situation, the collective behavior of the system is well described by a continuous (hydrodynamic) framework. The chief difference with standard hydrodynamic equations is that the governing equations are non-local. The influence of noise is limited or negligible throughout the evolution, as will become clear from our analysis.

For a logarithmic potential, the interaction strength  $g$  and the diffusion coefficient  $D$  have the same dimension [length]<sup>2</sup>/[time]. Their relative strength is quantified by the (dimensionless) ratio

$$\beta = \frac{g}{D} \quad (5)$$

playing the role of an inverse temperature [7]. For the one-dimensional Dyson gas [3] introduced in the context of random matrices, the exceptional values  $\beta = 1, 2, 4$  correspond to canonical ensembles (symmetric, hermitian, symplectic) of random matrices [3, 4, 11].

Our goal is to analyze the expansion into the vacuum of a large number of particles initially concentrated in a small spatial region. We rely on a continuous framework, namely on non-local hydrodynamic equations accounting for long-ranged interactions and ignoring diffusion. In dimension, the expansion of a long-ranged Riesz gas with

$s \in (-2, 1)$  can be solved exactly. In higher dimensions,  $d \geq 2$ , the Coulomb gases are tractable (thanks to the Gauss theorem). The case of non-Coulombic Riesz gases,  $s \neq d - 2$ , in  $d \geq 2$  dimensions remains open.

Our results can be summarized as follows. Qualitatively, the density profiles have compact support for Riesz gases with long-range potential ( $s < d$ ). For the one-dimensional Riesz gas with  $s \in (-2, 1)$ , the density

$$\rho(x, t) = \frac{B_s}{gt} (L^2 - x^2)^{\frac{s+1}{2}}, \quad B_s = \frac{\cos(\pi s/2)}{\pi(s+1)(s+2)} \quad (6)$$

is non-vanishing on the interval  $[-L(t), L(t)]$  with

$$L(t) = \left[ \frac{\sqrt{\pi} (2+s)^2 \Gamma(1 + \frac{s}{2})}{\cos(\frac{\pi s}{2}) \Gamma(\frac{1+s}{2})} Ngt \right]^{\frac{1}{s+2}} \quad (7)$$

For the Dyson gas ( $s = 0$ ), the density profile (6) is the celebrated Wigner semi-circle (expanding in the present case without confining potential).

For Coulomb gases in arbitrary dimension,  $s = d - 2$ , the radius of the expanding ball is

$$R = (dNgt)^{1/d} \quad (8)$$

The density profiles are uniform inside the expanding ball of radius  $R(t)$  and vanish outside the ball. The density of stochastic Coulomb gases inside the expanding ball is remarkably universal, viz., independent on  $N$  and inversely proportional to time in arbitrary spatial dimension:

$$\rho(r, t) = (\Omega_d gt)^{-1} \quad (9)$$

where  $\Omega_d$  is the volume of the unit sphere  $\mathbb{S}^{d-1}$ .

Intriguingly, the expansion of stochastic Coulomb gases remains well-defined when  $N = \infty$ . For non-Coulombic stochastic Riesz gases with infinitely many particles, the problem is ill-defined: the density vanishes when  $s < d - 2$  and becomes infinite when  $s \in (d - 2, d)$ .

We shall also demonstrate how to account for harmonic confining potential  $\epsilon r^2/2$ . In the Coulomb case

$$R(t) = \left[ \frac{gN}{\epsilon} (1 - e^{-det}) \right]^{1/d} \quad (10)$$

generalizes (8) and

$$\rho(r, t) = \frac{\epsilon d}{\Omega_d g} \frac{1}{1 - e^{-det}} \quad (11)$$

generalizes (9). The problem thus remains well-defined when  $N = \infty$  and the density is again uniform in the entire space  $\mathbb{R}^d$ . In the earlier time regime,  $t \ll \epsilon^{-1}$ , the density decays according to (9), then it saturates to the finite value  $\rho \rightarrow \rho_\infty = \frac{\epsilon d}{\Omega_d g}$ , when  $t \gg \epsilon^{-1}$ .

The outline of this work is as follows. In Sec. II, we solve the expansion problem for one-dimensional long-ranged Riesz gases with  $s \in (-2, 1)$ . In Sec. III, we analyze the expansion problem for the Coulomb gases

in arbitrary dimensions. We provide a detailed derivation in two spatial dimensions, i.e., for the Ginibre gas (Sec. III A), and in three spatial dimensions, i.e., for the classical Coulomb gas (Sec. III B). The general case is then easy to appreciate (Sec. III C). In the last section, we discuss open problems and mention amusing analogies between our results and formulas arising in Newtonian cosmology.

## II. ONE DIMENSION

At the coarse grained level, the behavior of the large number of particles evolving according to Eqs. (4) is described by continuum field theory for a single scalar field, the density  $\rho(x, t)$ , satisfying the continuity equation which in one dimension reads

$$\partial_t \rho + \partial_x J = 0 \quad (12)$$

The local current  $J(x, t)$  depends on the density and the inter-particle potential. The conservation equation (12) ensures that the total number of particles remains constant throughout the evolution:

$$\int_{-\infty}^{\infty} dx \rho(x, t) = N \quad (13)$$

at all  $t \geq 0$ .

To close the continuity equation (12), we must express  $J$  through the density and the interaction potential. The current contains the standard diffusion term,  $-D\partial_x \rho$ , resulting from the entropic contribution at microscopic level. A general long-range interaction potential  $V$  gives rise to a non-local deterministic contribution

$$J_V(x, t) = \rho(x, t) \int_{-\infty}^{\infty} dy \nabla V(x - y) \rho(y, t) \quad (14)$$

to the current. There is also a stochastic component to the current that can be written as  $\sqrt{2\rho}\eta$ , where the noise  $\eta(x, t)$  is a Gaussian noise, white in space and time. The amplitude  $\sqrt{2\rho}$  of this noise originates from the Brownian character of the particles, see [12]. Thus, the total current  $J(x, t)$  can be written as

$$J = -D\partial_x \rho + J_V + \sqrt{2\rho}\eta \quad (15)$$

A formal derivation of (15) by taking the continuous limit of the discrete set of equations can be obtained by following the strategy developed by Kawasaki and Dean [13, 14] (see [15, 16] for recent expositions).

We consider one-dimensional stochastic Riesz gases with  $s \in (-2, 1)$ . If  $s > 1$ , the interactions are effectively short-ranged (see [17, 18]), the governing hydrodynamic equations are local, and the emerging behaviors are more standard. The usual justification of the lower bound is that the original appearance of the Dyson gas in the context of random matrices involves the harmonic

confining  $r^2$  potential, and one does not want the pairwise interactions to overwhelm the confining potential. Even without confining potential, the systems with repulsive  $r^{-s}$  interactions are very peculiar when  $s < -2$ . Already for two particles, the inter-particle separation  $\ell(t)$  satisfies  $\frac{d\ell}{dt} = g\ell^{-s-1}$ , so  $\ell$  diverges in a finite time if  $s < -2$ . (This happens in arbitrary spatial dimension; taking diffusion into account does not prevent the finite-time blow-up.) These arguments explain why we study Riesz gases with exponents  $s \in (-2, 1)$ , and generally  $s \in (-2, d)$  in  $d$  dimensions.

In the present work, we focus on the typical gas expansion. We will see that the influence of diffusion on the expansion of Riesz gases with  $N \gg 1$  particles depends on whether  $s < 0$  or  $s > 0$ . When  $-2 < s < 0$ , the expansion is super-diffusive, namely  $L(t) \sim (Ngt)^{\frac{1}{s+2}}$ , cf. Eq. (7), and diffusion is effectively irrelevant. When  $0 < s < 1$ , the expansion is sub-diffusive. The diffusion length  $\sqrt{Dt}$  catches  $L(t)$  at a crossover time  $t_*$  estimated from  $\sqrt{Dt_*} \sim (Ngt_*)^{\frac{1}{s+2}}$ . Thus, the crossover time and size  $L_* = L(t_*)$  of the occupied region are

$$t_* \sim D^{-1} \left( \frac{Ng}{D} \right)^{2/s}, \quad L_* \sim \left( \frac{Ng}{D} \right)^{1/s} \quad (16)$$

This crossover time diverges when  $N \rightarrow \infty$ , as expected. For  $t \ll t_*$ , the profile evolves as (6) whereas for  $t \gg t_*$ , the ordinary diffusion prevails.

The hydrodynamic description predicts that the occupied region is compact,  $[-L(t), L(t)]$  if we ignore diffusion. For Riesz gases with  $s \in (0, 1)$ , diffusion cannot be ignored when  $t \gtrsim t_*$ . For Riesz gases with  $s \in (-2, 0)$ , the continuum description is inapplicable outside the  $[-L(t), L(t)]$  region. The analogy with random matrices suggests the position of the right-most [resp. left-most] particle is close to  $L(t)$  [resp.  $-L(t)$ ]. Studying the distributions of the positions of the right-most and left-most particles requires different techniques, so we shall merely present a few estimates but mostly rely on the hydrodynamic description accounting only the non-local deterministic contribution  $J_V$  to the current (15).

The Dyson gas separates the  $s \in (-2, 0)$  range where diffusion irrelevant and  $s \in (0, 1)$  range where diffusion is relevant when  $t \gtrsim t_*$ . Let us keep the diffusive contribution in the marginal case of the Dyson gas. The current in the Dyson gas caused by interactions is deduced from (14) by specializing to logarithmic interactions. One gets (see also [15, 16])

$$J_V(x, t) = g\rho(x, t) \mathcal{f}_{-\infty}^{\infty} dy \frac{\rho(y, t)}{x-y} \quad (17)$$

where  $\mathcal{f}$  denotes the Cauchy principal value

$$\mathcal{f}_{-\infty}^{\infty} dy \frac{\rho(y, t)}{x-y} = \lim_{\delta \rightarrow 0} \int_{|x-y| > \delta} dy \frac{\rho(y, t)}{x-y}.$$

Recalling the definition of the Hilbert transform

$$\mathcal{H}[\rho](x, t) = \frac{1}{\pi} \mathcal{f}_{-\infty}^{\infty} dy \frac{\rho(y, t)}{x-y} \quad (18)$$

and adding the diffusion term we arrive at

$$J = -D\partial_x \rho + \pi g \rho \mathcal{H}[\rho] \quad (19)$$

Combining this current with (12) we obtain

$$\partial_t \rho + \partial_x (\pi g \rho \mathcal{H}[\rho]) - D\partial_{xx} \rho = 0 \quad (20)$$

The expansion of the Dyson gas into the vacuum begins with particles concentrated in a tiny spatial region. We thus postulate that all particles are initially at the origin:

$$\rho(x, 0) = N\delta(x) \quad (21)$$

The initial-value problem (20)–(21) is invariant under the one-parameter group of transformations

$$\rho \rightarrow a^{-1} \rho, \quad x \rightarrow ax, \quad t \rightarrow a^2 t. \quad (22)$$

This implies that  $\sqrt{t}\rho(x, t)$  depends on the single variable  $x/\sqrt{t}$ . A more carefully chosen self-similar form

$$\rho(x, t) = \frac{N}{\sqrt{Ngt}} F(X) \quad \text{with} \quad X = \frac{x}{\sqrt{Ngt}} \quad (23)$$

has an advantage that both  $N$  and  $g$  disappear from the leading terms in the governing equation for the scaled density  $F(X)$  and from the conservation law. Indeed, the conservation law (13) turns into the conservation law

$$\int_{-\infty}^{\infty} dX F(X) = 1 \quad (24)$$

Substituting the scaling ansatz (23) into (20) we find that the scaled density satisfies

$$\frac{d(XF)}{dX} + \frac{2}{\beta N} \frac{d^2 F}{dX^2} = 2 \frac{d}{dX} \left[ F(X) \mathcal{f}_{-\infty}^{\infty} dY \frac{F(Y)}{X-Y} \right]$$

that we integrate to find

$$XF + \frac{2}{\beta N} \frac{dF}{dX} = 2F(X) \mathcal{f}_{-\infty}^{\infty} dY \frac{F(Y)}{X-Y} \quad (25)$$

The non-linear integro-differential equation (25) is not solvable analytically. However, we are interested in the behavior of a large number particles,  $N \gg 1$ . Hence, we neglect the  $1/N$  term in Eq. (25) and arrive at

$$F(X) \left[ X - 2 \mathcal{f}_{-\infty}^{\infty} dY \frac{F(Y)}{X-Y} \right] = 0 \quad (26)$$

We seek a symmetric solution,  $F(X) = F(-X)$ , vanishing in the  $|X| \rightarrow \infty$  limits. Such solution of Eq. (26) has a compact support:  $F = 0$  for  $|X| > R$ ; when  $|X| < R$ , the scaled density satisfies

$$X = 2 \mathcal{f}_{-R}^R dY \frac{F(Y)}{X-Y}. \quad (27)$$

The solution of the linear integral equation (27) is well-known,  $F = (2\pi)^{-1} \sqrt{R^2 - X^2}$ . [See [19–21] for the descriptions of the methods of solving Eq. (27) and similar

linear singular integral equations.] Using the conservation law (24), we fix  $R = 2$ . Summarizing

$$F = \begin{cases} \frac{1}{2\pi} \sqrt{4 - X^2} & |X| < 2 \\ 0 & |X| > 2 \end{cases} \quad (28)$$

In the original variables, the density profile is the Wigner semi-circle

$$\rho(x, t) = \frac{1}{2\pi gt} \sqrt{4Ngt - x^2} \quad (29)$$

as for the equilibrium Dyson gas in a harmonic confining potential. The difference is that the span now diffusively expands with time:  $|x| \leq 2\sqrt{Ngt}$ .

One can employ asymptotic techniques [22] to obtain more precise solution of Eq. (25) inside the occupied  $|X| < 2$  region. One seeks the solution as an expansion in a small parameter  $(\beta N)^{-1} \ll 1$ :

$$F(X) = \frac{1}{2\pi} \sqrt{4 - X^2} + \frac{1}{\beta N} F_1(X) + \dots \quad (30)$$

Sustituting (30) into Eq. (25) we find that the leading correction satisfies

$$\int_{-2}^2 dY \frac{F_1(Y)}{X - Y} = -\frac{X}{4 - X^2} \quad (31)$$

We solve the linear singular integral equation (31) using the methods described in [19–21] and obtain

$$F_1(X) = -\frac{1}{\pi^2 \sqrt{4 - X^2}} \left[ 4 + X \ln \frac{2 - X}{2 + X} \right] \quad (32)$$

The leading and the sub-leading terms in the expansion (30) become comparable when  $2 - |X| \sim N^{-1}$ . Therefore, Eq. (30) gives an outer expansion, which should be matched with inner expansions valid in the internal  $2 - |X| \sim N^{-1}$  layers. We do not perform such an analysis since, inside the internal layers, the deterministic framework is no longer justified. (Relying on the deterministic continuum framework for  $|X| > 2$  is even more erroneous.) Indeed, using (29), one estimates the positions  $x_{\pm}(t)$  of the right-most and left-most particles:

$$\frac{x_{\pm}(t)}{L(t)} = \pm 1 + N^{-\frac{2}{3}} \xi_{\pm}(\beta) \quad (33)$$

with  $L(t) = 2\sqrt{Ngt}$  and random variables  $\xi_{\pm}(\beta)$  that are expected to be asymptotically stationary with Tracy-Widom type distributions dependent on  $\beta = g/D$ . (The Tracy-Widom distributions appearing in the context of random matrices are explicitly known for  $\beta = 1, 2, 4$  characterizing ensembles of random matrices [4, 11, 23]. In our case,  $\beta > 0$  is arbitrary.)

We now consider general one-dimensional Riesz gases with long-ranged interactions,  $-2 < s < 1$ . We ignore diffusion and comment below when this approximation is

justifiable. The contribution (14) to the current due to interactions becomes (see also [18])

$$J(x, t) = g\rho(x, t) \int_{-\infty}^{\infty} dy \frac{x - y}{|x - y|^{s+2}} \rho(y, t) \quad (34)$$

Combining the continuity equation (12) with (34) leads to the following integro-differential equation

$$\partial_t \rho + \partial_x \left[ g\rho \int_{-\infty}^{\infty} dy \frac{x - y}{|x - y|^{s+2}} \rho(y, t) \right] = 0 \quad (35)$$

for the density profile. The governing equation (35) and the conservation law (13) are invariant under the one-parameter group of transformations

$$\rho \rightarrow a^{-\frac{1}{s+2}} \rho, \quad x \rightarrow a^{\frac{1}{s+2}} x, \quad t \rightarrow at. \quad (36)$$

Using the self-similar form

$$\rho(x, t) = \frac{N}{(Ngt)^{\frac{1}{s+2}}} F(X) \quad \text{with} \quad X = \frac{x}{(Ngt)^{\frac{1}{s+2}}} \quad (37)$$

we find that the scaled density satisfies

$$\frac{F + X \frac{dF}{dX}}{s + 2} = \frac{d}{dX} \left[ F \int_{-\infty}^{\infty} dY \frac{X - Y}{|X - Y|^{s+2}} F(Y) \right] \quad (38)$$

Integrating Eq. (38) we arrive at the integral equation

$$\frac{X}{s + 2} = \int_{-R}^R dY \frac{X - Y}{|X - Y|^{s+2}} F(Y) \quad (39)$$

inside the  $|X| < R$  interval where  $F(X) > 0$ ; the density vanishes outside that interval. Integral equations such as (39) arise in many problems, see, e.g., [17, 21]. The solution has again a compact support and appears in textbooks [19, 20]. For  $|X| < R$ , we have

$$F = B_s (R^2 - X^2)^{\frac{s+1}{2}}, \quad B_s = \frac{\cos(\pi s/2)}{\pi(s+1)(s+2)} \quad (40)$$

The normalization condition is again given by (24) which we combine with (40) to yield

$$R = \left[ \frac{\sqrt{\pi} (2 + s)^2 \Gamma(1 + \frac{s}{2})}{\cos(\frac{\pi s}{2}) \Gamma(\frac{1+s}{2})} \right]^{\frac{1}{s+2}} \quad (41)$$

The density is non-vanishing when  $x \in [-L(t), L(t)]$  with  $L(t) = R(Ngt)^{\frac{1}{s+2}}$  as was announced in Eq. (7). The density profile (40) reduces to the announced result (6) in the original variables. In Fig. 1, we show the density profiles for  $s = -1, 0, 7/8$ .

We relied on Eq. (35) ignoring diffusion. If  $0 < s < 1$ , the diffusion is irrelevant when  $t \ll t_*$  with crossover time given by (16). In the  $-2 < s < 0$  range, the expansion is super-diffusive and diffusion is essentially irrelevant. Using (6)–(7), one estimates the positions of the right-most




FIG. 1. The spatial dependence of the density re-scaled to be equal to unity at the origin:  $F/(B_s R^{s+1})$ , with  $F$  given by Eq. (40), versus  $X/R$ . The density vanishes when  $|X| > R$ . The shown curves (top to bottom) correspond to the Riesz exponent  $s = -3/2, -1, 0, 7/8$ . The density is flat for the one-dimensional Coulomb gas ( $s = -1$ ). The semi-circle represents the density of the Dyson gas ( $s = 0$ ).

and left-most particles and arrives to the result similar to (33), with  $N^{-2/3}$  replaced by  $N^{-\frac{2}{s+3}}$ .

In the presence of a harmonic confining potential, the density becomes stationary in the long time limit. The stationary density satisfies

$$g \int_{-\infty}^{\infty} dy \frac{x-y}{|x-y|^{s+2}} \rho(y) = \epsilon x \quad (42)$$

which is solved similarly to (39) to give

$$\rho = \frac{\epsilon}{g} \frac{\cos(\pi s/2)}{\pi(s+1)} (r^2 - x^2)^{\frac{s+1}{2}} \quad (43)$$

from which we deduce that the span is given by

$$r = \left[ \frac{Ng}{\epsilon} \frac{\sqrt{\pi} (2+s) \Gamma(1 + \frac{s}{2})}{\cos(\frac{\pi s}{2}) \Gamma(\frac{1+s}{2})} \right]^{\frac{1}{s+2}} \quad (44)$$

To determine the evolution in the presence of the harmonic confining potential one ought to solve

$$\partial_t \rho + \partial_x \left[ g \rho \int_{-\infty}^{\infty} dy \frac{x-y}{|x-y|^{s+2}} \rho(y, t) - \epsilon \rho x \right] = 0 \quad (45)$$

The similarity of the solution (37) and (40), and the stationary solution (6) in the harmonic confining potential, suggests to seek the solution of (45) in the same functional form:

$$\rho(x, t) = \frac{NC_s}{R^{s+2}} (R^2 - x^2)^{\frac{s+1}{2}}, \quad C_s = \frac{\Gamma(2 + \frac{s}{2})}{\sqrt{\pi} \Gamma(\frac{3+s}{2})} \quad (46)$$

with amplitude fixed by normalization. Substituting (46) into (45), we find that (46) provides a consistent solution if the span satisfies

$$\frac{dR}{dt} = \frac{gNC_s}{R^{s+1}} - \epsilon R \quad (47)$$

Solving (47) subject to  $R(0) = 0$  gives

$$R = \left\{ \frac{gNC_s}{\epsilon} \left[ 1 - e^{-\epsilon(s+2)t} \right] \right\}^{\frac{1}{s+2}} \quad (48)$$

Thus, for the Riesz gas in the confining harmonic potential, the evolving density profile describing the expansion into the vacuum has the form (46) throughout the evolution, with span given by (48). The evolving density profile does not ‘feel’ the confining potential when  $t \ll t_c \sim \epsilon^{-1}$ ; the density profile approaches to the stationary density (43)–(44) when  $t \gg t_c$ .

### III. THE COULOMB GAS IN HIGHER DIMENSIONS

#### A. The case $d = 2$ (Ginibre gas)

The expansion is isotropic. Hence  $\rho(\mathbf{r}, t) = \rho(r, t)$  and the continuity equation reduces to

$$\partial_t \rho + r^{-1} \partial_r (rJ) = 0 \quad (49)$$

A simple expression for the current

$$J(r, t) = g \rho(r, t) r^{-1} \int_0^r dr' 2\pi r' \rho(r', t) \quad (50)$$

reflects the rotational symmetry and the Coulomb nature of the logarithmic potential in two dimensions: the force at  $r$  is the same as if the entire charge of the disk of radius  $r$  was at the origin (Gauss theorem). Introducing an auxiliary variable

$$F(r, t) = 2\pi r \rho(r, t) \quad (51)$$

we re-write (49)–(50) as

$$\partial_t F(r, t) + g \partial_r \left[ r^{-1} F(r, t) \int_0^r dr' F(r', t) \right] = 0 \quad (52)$$

The conservation law  $\int d\mathbf{r} \rho(\mathbf{r}, t) = N$  simplifies to

$$\int_0^\infty dr F(r, t) = N \quad (53)$$

Equations (52)–(53) are invariant under the one-parameter group of transformations

$$F \rightarrow a^{-1} F, \quad r \rightarrow ar, \quad t \rightarrow a^2 t \quad (54)$$

implying that the solution has a self-similar form

$$F(r, t) = \sqrt{\frac{N}{gt}} \Phi(\xi) \quad \text{with} \quad \xi = \frac{r}{\sqrt{Ngt}} \quad (55)$$

chosen to ensure that both  $N$  and  $g$  disappear from the governing equation and the conservation law for  $\Phi(\xi)$ . Substituting the scaling ansatz (55) into (52) yields

$$\Phi + \xi \frac{d\Phi}{d\xi} = 2 \frac{d}{d\xi} \left[ \frac{\Phi \Psi}{\xi} \right] \quad (56)$$



where

$$\Psi(\xi) = \int_0^\xi d\xi' \Phi(\xi') \quad (57)$$

Integrating (56) gives

$$\Phi\left(\xi - \frac{2\Psi}{\xi}\right) = 0 \quad (58)$$

The solution reads

$$\Phi = \begin{cases} \xi & \xi \leq \xi_0 \\ 0 & \xi > \xi_0 \end{cases} \quad (59)$$

Substituting this solution into the conservation law

$$\int_0^\infty d\xi \Phi(\xi) = 1 \quad (60)$$

we fix  $\xi_0 = \sqrt{2}$ . Thus, at any time the density is uniform inside the growing disk

$$\rho(r, t) = \begin{cases} (2\pi gt)^{-1} & r \leq \sqrt{2Ngt} \\ 0 & r > \sqrt{2Ngt} \end{cases} \quad (61)$$

The radius of the disk grows diffusively as for the one-dimensional Dyson gas; the extra  $N$  factor ensures again that diffusion plays a minor role. The diffusive growth of the ball indicates that the scaling ansatz (55) remains applicable even when we take diffusion into account. The auxiliary function (57) satisfies a neat ordinary differential equation

$$\frac{1}{2\beta N} \eta \Psi'' = \Psi'(\Psi - \eta) \quad (62)$$

where  $\eta = \xi^2/2$  and prime denotes a derivative with respect to  $\eta$ . This equation is simpler than the integro-differential equation (25) describing the Dyson gas. Still, Eq. (62) is non-linear and analytically intractable.

Generalizing (52) to account for the confining harmonic potential yields

$$\partial_t F + \partial_r \left[ \frac{g}{r} F \int_0^r dr' F(r', t) - \epsilon r F \right] = 0 \quad (63)$$

We anticipate that the density profile is uniform inside an expanding disk and vanishes outside the disk  $r > R(t)$ . Substituting  $F(r, t) = r f(t)$  into (63), one finds a closed equation for  $f(t)$ . Thus, we conclude that there is a uniform density profile:

$$\rho(r, t) = \frac{\epsilon}{\pi g} \frac{1}{1 - e^{-2\epsilon t}} \quad (64)$$

inside the disk of radius  $R(t)$ . This radius is fixed by the mass conservation law and is given by

$$R(t) = \sqrt{\frac{gN}{\epsilon} (1 - e^{-2\epsilon t})} \quad (65)$$

Again, we observe an exponential convergence to the stationary case, with a crossover time of order  $1/\epsilon$ .

## B. The three dimensional case

In the spherically symmetric case, the continuity equation reads

$$\partial_t \rho + r^{-2} \partial_r (r^2 J) = 0 \quad (66)$$

For the Coulomb gas, we have  $V(\mathbf{r}) = g/r$  and the current is given by

$$J(r, t) = g\rho(r, t)r^{-2} \int_0^r dr' 4\pi r'^2 \rho(r', t) \quad (67)$$

This expression reflects rotational symmetry and we have used again the Coulomb nature of the potential: the force at  $r$  is the same as if the entire mass of the ball of radius  $r$  was at the origin. Using

$$F(r, t) = 4\pi r^2 \rho(r, t) \quad (68)$$

we re-write (66)–(67) as

$$\partial_t F(r, t) + g\partial_r \left[ r^{-2} F(r, t) \int_0^r dr' F(r', t) \right] = 0 \quad (69)$$

The conservation law has the same form (60) as in two dimensions. Equations (69) and (60) are invariant under the one-parameter group of transformations

$$F \rightarrow a^{-1}F, \quad r \rightarrow ar, \quad t \rightarrow a^3t \quad (70)$$

implying that the solution has a self-similar form

$$F(r, t) = \frac{N}{\sqrt[3]{Ngt}} \Phi(\xi) \quad \text{with} \quad \xi = \frac{r}{\sqrt[3]{Ngt}} \quad (71)$$

Substituting the scaling ansatz (71) into (69) yields

$$\Phi + \xi \frac{d\Phi}{d\xi} = 3 \frac{d}{d\xi} \left[ \frac{\Phi\Psi}{\xi^2} \right] \quad (72)$$

which is integrated to give

$$\Phi\left(\xi - \frac{3\Psi}{\xi^2}\right) = 0 \quad (73)$$

The solution reads

$$\Phi = \begin{cases} \xi^2 & \xi \leq \xi_0 \\ 0 & \xi > \xi_0 \end{cases} \quad (74)$$

Substituting this solution into the conservation law (60) we fix  $\xi_0 = \sqrt[3]{3}$ . Thus at any time the density is uniform inside the growing disk

$$\rho(r, t) = \begin{cases} (4\pi gt)^{-1} & r \leq \sqrt[3]{3Ngt} \\ 0 & r > \sqrt[3]{3Ngt} \end{cases} \quad (75)$$

The radius of the ball grows sub-diffusively, so the above results are valid up to a crossover time  $t_* \sim (Ng)^2/D^3$ .

To account for the confining harmonic potential we must solve

$$\partial_t F + \partial_r \left[ \frac{g}{r^2} F \int_0^r dr' F(r', t) - \epsilon r F \right] = 0 \quad (76)$$

We anticipate that the density profile is uniform inside an expanding ball and vanishes outside the ball,  $r > R(t)$ . Seeking such a uniform solution gives

$$\rho(r, t) = \frac{3\epsilon}{4\pi g} \frac{1}{1 - e^{-3\epsilon t}} \quad (77)$$

inside the ball and the conservation law fixes the radius of the ball

$$R(t) = \sqrt[3]{\frac{gN}{\epsilon} (1 - e^{-3\epsilon t})} \quad (78)$$

### C. Arbitrary spatial dimension

The same analysis as in Secs. III A–III B can be carried in arbitrary dimensions. We obtain

$$\rho(r, t) = \begin{cases} (\Omega_d g t)^{-1} & r \leq (dNgt)^{1/d} \\ 0 & r > (dNgt)^{1/d} \end{cases} \quad (79)$$

where  $\Omega_d = 2\pi^{d/2}/\Gamma(d/2)$  is the volume of the unit sphere  $\mathbb{S}^{d-1}$ . The density remains uniform in the presence of the confining harmonic potential. One finds

$$\rho(r, t) = \frac{\epsilon d}{\Omega_d g} \frac{1}{1 - e^{-d\epsilon t}} \quad (80)$$

inside an expanding  $d$ -dimensional ball,  $r < R(t)$ . The radius of the ball is given by the announced formula (10).

Consider now the expansion into the vacuum of an infinite free gas with  $N = \infty$  and without an external potential. Stunningly, the Coulomb gas is the only Riesz gas with a well-defined behavior in this limit. This claim is easy to prove for the one-dimensional Riesz gas solved in Sec. II. For instance, the density at the origin in the system with a large but finite number of particles

$$\rho(0, t) = (NC_s)^{\frac{s+1}{s+2}} [g(s+2)t]^{-\frac{1}{s+2}} \quad (81)$$

diverges in the  $N \rightarrow \infty$  limit if  $s > -1$  and vanishes if  $s < -1$ . A well-defined finite answer emerges only when  $s = -1$ , i.e., for the one-dimensional Coulomb potential. More generally, for the Coulomb gas in arbitrary dimension, the density is well-defined, uniform in the entire infinite space  $\mathbb{R}^d$ , and decays inversely proportional to time when the number of particles is infinite,  $\rho(r, t) = (\Omega_d g t)^{-1}$ .

## IV. CONCLUDING REMARKS

We have analyzed the expansion of the stochastic Riesz gases with long-ranged interactions in one dimension and

of the Coulomb gases in an arbitrary dimension. Stochastic Coulomb gases remain well-defined in the infinite-particle limit,  $N = \infty$ . The emerging spatially uniform solution (9) bears an amusing resemblance with the uniform decaying density in the expanding Universe. The gravitational (Newton) potential is Coulombic yet attractive, but the motion is governed by Newton's laws (not overdamped). In the Universe filled with ‘‘dust’’ (i.e., neglecting the pressure), the density decays as  $t^{-2}$  both in Newtonian cosmology and Einsteinian cosmology in the flat Universe (see, e.g., [24]). The  $t^{-2}$  decay is independent on the spatial dimension as the  $t^{-1}$  decay (9) in an overdamped ‘Universe’. Instead of the Hubble law for the velocity,  $\mathbf{v} = H(t)\mathbf{r}$  with  $H = \frac{2}{d}t^{-1}$ , we have  $\mathbf{J}(\mathbf{r}, t) \propto \mathbf{r}/t^2$ . The underlying reason for these striking similarities is the peculiar nature of the Coulomb potential. Besides, adding the confining harmonic potential is akin to adding the cosmological constant. The signs are reversed: the interactions are repulsive (in contrast with the attractive gravitational interactions), and the ‘cosmological constant’ ( $-\epsilon$ ) is negative rather than positive. This analogy can be carried further [25, 26].

We have not been able to analyze the expansion of the Riesz gas in  $d > 1$  dimensions, and we now explain the technical challenge. The governing equation is given by

$$\partial_t \rho + \nabla \cdot \mathbf{J} = 0 \quad (82)$$

with

$$\mathbf{J}(\mathbf{r}, t) = g\rho(\mathbf{r}, t) \int_{B_R} d\mathbf{y} \frac{\mathbf{r} - \mathbf{y}}{|\mathbf{r} - \mathbf{y}|^{s+2}} \rho(\mathbf{y}, t) \quad (83)$$

We expect that the density  $\rho(\mathbf{r}, t)$  vanishes outside the ball  $B_R$  of radius  $R$ , so we integrate over this ball, i.e.,  $|\mathbf{y}| < R$ , in (83). The rotational symmetry implies  $\rho(\mathbf{r}, t) = \rho(r, t)$  and  $\mathbf{J}(\mathbf{r}, t) = \frac{\mathbf{r}}{r} J(r, t)$ , so Eq. (82) simplifies to

$$\partial_t \rho + r^{-(d-1)} \partial_r (r^{d-1} J) = 0 \quad (84)$$

The functional  $J[\rho]$  involves a  $d$ -fold integral of  $\rho$ , which can not be simplified further because for non-Coulombic gases where we cannot rely on the Gauss theorem.

The self-similar form (37) generalizes to

$$\rho(r, t) = \frac{N}{(Ngt)^{\frac{d}{s+2}}} F(\xi) \quad \text{with} \quad \xi = \frac{r}{(Ngt)^{\frac{1}{s+2}}} \quad (85)$$

The Riesz exponent varies in the  $s \in (-2, d)$  range.

The chief technical challenge is to derive an integro-differential equation for  $F(\xi)$ . It is perhaps possible to advance when  $d = 3$ . The idea is to rely on the neat identity [21]

$$\int_{B_R} d\mathbf{y} \frac{\rho(y)}{|\mathbf{r} - \mathbf{y}|^s} = \frac{2\pi}{s-2} \int_{-R}^R dy y \rho(y) \frac{|r-y|^{2-s}}{r} \quad (86)$$

valid for an arbitrary spherically-symmetric function  $\rho(y)$  but only in three dimensions. Differentiating (86), we

obtain another identity with the left-hand side equal to the integral in Eq. (83). In contrast, the right-hand side involves the integration over the interval  $(-R, R)$  rather than the three-dimensional ball. This strategy may lead to the solution of the problem in three dimensions.

Stepping away from the unconstrained hydrodynamic behaviors, we mention fluctuations. For one-dimensional Riesz gases in extremely shallow confining potential, the fluctuations of the displacement of the tagged particle in one-dimensional Riesz gases have been studied in [18] both in the long-ranged regime  $0 < s < 1$  and effectively short-range regime  $s > 1$ . The fluctuations of the local current in the Dyson gas ( $s = 0$ ) have also been investigated [16]. One anticipates that the tagged particle behaves diffusively when  $s < -1$ ; the self-diffusion coefficient is unknown.

An interesting challenge concerns large deviations parallel to those investigated in equilibrium [27, 28]. For in-

stance, the Dyson gas expands diffusively,  $|x| \leq 2\sqrt{Ngt}$ , and the density profile is the Wigner semi-circle (29). It would be interesting to compute the probability of atypically slow expansion,  $|x| \leq A\sqrt{Ngt}$  with some fixed  $A < 2$  during the time interval  $t \in (0, T)$ , and the density profile during such constrained expansion. Atypically fast expansions,  $|x| \geq A\sqrt{Ngt}$  with some fixed  $A > 2$  during the time interval  $t \in (0, T)$ , are equally interesting. Alternatively, one can seek the probability that the occupied region is atypically large or small,  $|x(T)| \leq A\sqrt{NgT}$  with  $A \neq 2$ , only at the final moment.

The work of KM is supported by the project RETENU ANR-20-CE40-0005-01 of the French National Research Agency (ANR). KM is thankful to the participants of the Indo-French workshop on classical and quantum dynamics organized by ICTS Bengaluru for discussions. PLK is grateful to IPhT for excellent working conditions.

- 
- [1] A. Campa, T. Dauxois, and S. Ruffo, Statistical mechanics and dynamics of solvable models with long-range interactions, *Physics Reports* **480**, 57 (2009).
  - [2] A. Campa, T. Dauxois, D. Fanelli, and S. Ruffo, *Physics of Long-Range Interacting Systems* (Oxford University Press, Oxford, UK, 2014).
  - [3] F. J. Dyson, A Brownian motion model for the eigenvalues of a random matrix, *J. Math. Phys.* **3**, 1191 (1962).
  - [4] P. J. Forrester, *Log-gases and random matrices* (Princeton University Press, Princeton, NJ, 2010).
  - [5] J. Ginibre, Statistical ensembles of complex, quaternion, and real matrices, *J. Math. Phys.* **6**, 440 (1965).
  - [6] S.-S. Byun and P. J. Forrester, *Progress on the study of the Ginibre ensembles* (Springer, Singapore, 2024).
  - [7] M. Lewin, Coulomb and Riesz gases: The known and the unknown, *J. Math. Phys.* **63**, 061101 (2022).
  - [8] S. Serfaty, Lectures on Coulomb and Riesz gases, [arXiv:2407.21194](https://arxiv.org/abs/2407.21194) (2024).
  - [9] F. Calogero, Ground state of the one-dimensional N-body system, *J. Math. Phys.* **10**, 2197 (1969).
  - [10] A. P. Polychronakos, The physics and mathematics of Calogero particles, *J. Phys. A* **39**, 12793 (2006).
  - [11] G. Livan, M. Novaes, and P. Vivo, *Introduction to Random Matrices* (Springer, Cham, Switzerland, 2018).
  - [12] H. Spohn, *Large Scale Dynamics of Interacting Particles* (Springer, Berlin, 1991).
  - [13] K. Kawasaki, Stochastic model of slow dynamics in supercooled liquids and dense colloidal suspensions, *Physica A* **208**, 35 (1994).
  - [14] D. S. Dean, Langevin equation for the density of a system of interacting Langevin processes, *J. Phys. A* **29**, L613 (1996).
  - [15] L. Touzo, P. Le Doussal, and G. Schehr, Interacting, running and tumbling: The active Dyson Brownian motion, *EPL* **142**, 61004 (2023).
  - [16] R. Dandekar, P. L. Krapivsky, and K. Mallick, Current fluctuations in the Dyson gas, [arXiv:2409.06881](https://arxiv.org/abs/2409.06881) (2024).
  - [17] S. Agarwal, A. Dhar, M. Kulkarni, A. Kundu, S. N. Majumdar, D. Mukamel, and G. Schehr, Harmonically confined particles with long-range repulsive interactions, *Phys. Rev. Lett.* **123**, 100603 (2019).
  - [18] R. Dandekar, P. L. Krapivsky, and K. Mallick, Dynamical fluctuations in the Riesz gas, *Phys. Rev. E* **107**, 044129 (2023).
  - [19] R. Estrada and R. P. Kanwal, *Singular Integral Equations* (Birkhäuser, Boston, MA, 2000).
  - [20] A. D. Polyanin and A. V. Manzhirov, *Handbook of Integral Equations* (Chapman and Hall/CRC, Boca Raton, FL, 2008).
  - [21] J. A. Carrillo and Y. Huang, Explicit equilibrium solutions for the aggregation equation with power-law potentials, *Kinet. Relat. Models* **10**, 171 (2017).
  - [22] C. M. Bender and S. A. Orszag, *Advanced Mathematical Methods for Scientists and Engineers* (McGraw-Hill, New York, 1978).
  - [23] S. N. Majumdar and G. Schehr, Top eigenvalue of a random matrix: large deviations and third order phase transition, *J. Stat. Mech.* **2014**, P01012 (2014).
  - [24] V. Mukhanov, *Physical Foundations of Cosmology* (Cambridge University Press, Cambridge, UK, 2010).
  - [25] E. B. Kolomeisky, Natural analog to cosmology in basic condensed matter physics, *Phys. Rev. B* **100**, 140301 (2019).
  - [26] E. B. Kolomeisky, Analog de Sitter space in a controlled Coulomb explosion, *Phys. Rev. A* **103**, 043101 (2021).
  - [27] D. S. Dean and S. N. Majumdar, Large deviations of extreme eigenvalues of random matrices, *Phys. Rev. Lett.* **97**, 160201 (2006).
  - [28] D. S. Dean and S. N. Majumdar, Extreme value statistics of eigenvalues of Gaussian random matrices, *Phys. Rev. E* **77**, 041108 (2008).