Predictions of masses for light hybrid baryons

Qi-Nan Wang^{1,2,3}, Ding-Kun Lian¹, Wei Chen^{1,4},* Hui-Min Yang⁵, Hua-Xing Chen⁶,[†] J. Ho⁷,[‡] and T. G. Steele⁸

¹School of Physics, Sun Yat-Sen University, Guangzhou 510275, China

²Editorial Department of Journal, Bohai University, Jinzhou 121013, China

³College of physical science and technology, Bohai University, Jinzhou 121013, China

⁴Southern Center for Nuclear-Science Theory (SCNT), Institute of Modern Physics,

Chinese Academy of Sciences, Huizhou 516000, Guangdong Province, China

⁵School of Physics and Center of High Energy Physics, Peking University, Beijing 100871, China

⁶School of Physics, Southeast University, Nanjing 210094, China

⁷Department of Physics, Dordt University, Sioux Center, Iowa, 51250, USA

⁸Department of Physics and Engineering Physics, University of Saskatchewan, Saskatoon, SK, S7N 5E2, Canada

Within the method of parity-projected QCD sum rules, we study the mass spectra of light hybrid baryons with $I(J^P) = 1/2(1/2^{\pm}), 3/2(1/2^{\pm}), 1/2(3/2^{\pm}), 3/2(3/2^{\pm})$ by constructing the local *qqqg* interpolating currents. We calculate the correlation functions up to dimension eight condensates at the leading order of α_s . The stable QCD Lapalce sum rules can be established for the positive-parity $N_{1/2^+}, \Delta_{3/2^+}, \Delta_{1/2^+}$ and negative-parity $N_{1/2^-}, N_{3/2^-}, \Delta_{1/2^-}$ channels to extract their mass spectra. The lowest-lying hybrid baryons are predicted to be the negative-parity $N_{1/2^-}$ state around 2.28 GeV and $\Delta_{1/2^-}$ state around 2.64 GeV. These hybrid baryons mainly decay into conventional baryon plus meson final states. We propose to search for the light hybrid baryons through the χ_{cJ}/Υ decays via the three-gluon emission mechanism in BESIII and BelleII experiments. Our studies of the light hybrid baryons will be useful for understanding the excited baryon spectrum and the behavior of gluonic degrees of freedom in QCD.

PACS numbers: 12.39.Mk, 12.38.Lg, 14.40.Ev, 14.40.Rt Keywords: Hybrid baryon, QCD sum rules

Introduction.—As a distinctive prediction of QCD, the existence of hybrid states is a longstanding issue in both experimental and theoretical aspects in the past several decades. Different from the conventional $q\bar{q}$ meson and qqq baryon [1, 2], hybrid states contain the gluonic degrees of freedom in the configurations, such as $\bar{q}gq$ hybrid mesons and qqqg hybrid baryons. It is of crucial importance to identify the existence of hybrid states for understanding the nonperturbative properties of QCD [3–10].

In the past several decades, there are numerous investigations on hybrid mesons in various theoretical methods, such as the MIT bag model [11, 12], Coulomb gauge QCD [13], LQCD [14–22], the flux tube model [23–25], Bethe-Salpeter equation [26, 27], QCD sum rules [28–42] and so on. The most intriguing candidates for light hybrid mesons are $\pi_1(1400)$ [43], $\pi_1(1600)$ [44], $\pi_1(2015)$ [45] and $\eta_1(1855)$ [46, 47] with exotic $J^{PC} = 1^{-+}$. Comparing to hybrid mesons, a hybrid baryon is composed of a color octet triquark field and a color octet gluonic excitation. To date, the studies of hybrid baryons are much less appealing since they cannot carry exotic quantum numbers. In general, QCD allows arbitrarily strong mixing effect between qqqg hybrid baryons and excited qqq states with the same quantum numbers.

In Refs. [48–50], the Roper resonance N(1440) was studied as a hybrid baryon with $J^{P} = 1/2^{+}$. Such a possibility is

excluded according to the meson-cloud theory from the constituent quark model [51]. In Ref. [48], the masses of light hybrid baryon states with spin-parity $J^P = 1/2^+, 3/2^+, 5/2^+$ were calculated in the MIT bag model. They predicted the masses of these hybrids in the range of 1.6 - 2.5 GeV, among which the lightest one is the $1/2^+$ state. In the flux-tube model, the lightest hybrid baryons are the mass-degenerate states with $J^{P} = 1/2^{+}$ and $3/2^{+}$ near 1.87 GeV [52, 53]. Using the relativistic quark model, the authors of Ref. [54] studied the mass spectra of the nucleon and delta hybrids and suggested that N(1710) and $\Delta(1600)$ could be hybrid baryons. In Refs. [55, 56], the mass spectra of positiveparity hybrid baryons were calculated in LQCD, in which $N_{1/2^+}$ and $N_{3/2^+}$ were predicted to be the lowest-lying hybrid states around 2.3 - 2.6 GeV. The QCD sum rule method has also been applied to study the hybrid baryons [50, 57-59]. In Ref. [57], a light hybrid baryon $N_{1/2^+}$ was studied and found to be approximately 2.1 GeV. Using the same interpolating current, the calculation was improved to give different results of the gluon, trigluon and mixed condensates, while the hybrid mass was found to be around the observed mass of the Roper resonance [50]. In QCD sum rules, the $\Lambda(1600)$ and $\Lambda(1405)$ were suggested as strange hybrid baryons with $I(J^{P}) = 0(1/2^{+})$ [58] and $I(J^{P}) = 0(1/2^{-})$ [59], respectively. Recently, the triply heavy cccg and bbbg hybrids were studied in QCD sum rules [60] and a semirelativistic potential model [61].

In this work, we systematically study the mass spectra of nucleon and delta hybrid baryons with both positive and negative parities by using the parity-projected QCD sum rules. We construct the $[qqq]_{8c}[G]_{8c}$ local hybrid baryon interpolating currents without the covariant derivative operators and introduce the parity projected sum rules to separate the hybrid

^{*}Electronic address: chenwei29@mail.sysu.edu.cn

[†]Electronic address: hxchen@seu.edu.cn

[‡]Electronic address: jason.ho@dordt.edu

[§]Electronic address: tom.steele@usask.ca

[¶]Qi-Nan Wang and Ding-Kun Lian equally contribute to this work.

2

baryons with different parities. We find stable mass sum rules and establish extensive mass spectra for the positive-parity $N_{1/2^+}, \Delta_{3/2^+}, \Delta_{1/2^+}$ and negative-parity $N_{1/2^-}, N_{3/2^-}, \Delta_{1/2^-}$ hybrid baryons. Our results show that the lowest-lying hybrid baryon is predicted to appear in the negative-parity $N_{1/2^-}$ channel around 2.28 GeV, while the lightest $\Delta_{1/2^-}$ state is about 300 MeV higher. We propose to search for these light hybrid baryons through the χ_{cJ}/Υ decays via the three-gluon emission mechanism in BESIII and BelleII experiments.

Hybrid baryon currents and parity-projected QCD sum rules.— A hybrid baryon current is composed of a color-octet triquark operator $[qqq]_{8c}$ and a color-octet excited gluon field. To investigate the low-lying states, we use the S-wave triquark fields introduced as "Ioffe currents" [62]. The gluon field is combined to compose the color-singlet hybrid baryon interpolating currents with $J^P = 1/2^+$ and $3/2^+$

$$\begin{split} J_{1} &= \varepsilon^{abc} g_{s} [(u^{aT} C \gamma^{\mu} u^{b}) \gamma^{\nu} \gamma^{5} (G_{\mu\nu} d)^{c} - (u^{aT} C \gamma^{\mu} d^{b}) \gamma^{\nu} \gamma^{5} (G_{\mu\nu} u)^{c}] \\ J_{2} &= \varepsilon^{abc} g_{s} [(u^{aT} C \gamma^{\mu} u^{b}) \gamma^{\nu} \gamma^{5} (G_{\mu\nu} d)^{c} + 2(u^{aT} C \gamma^{\mu} d^{b}) \gamma^{\nu} \gamma^{5} (G_{\mu\nu} u)^{c}] , \\ J_{3} &= \varepsilon^{abc} g_{s} [(u^{aT} C \sigma^{\mu\nu} u^{b}) \gamma^{5} (G_{\mu\nu} d)^{c} - (u^{aT} C \sigma^{\mu\nu} d^{b}) \gamma^{5} (G_{\mu\nu} u)^{c}] , \\ J_{4} &= \varepsilon^{abc} g_{s} [(u^{aT} C \sigma^{\mu\nu} u^{b}) \gamma^{5} (G_{\mu\nu} d)^{c} + 2(u^{aT} C \sigma^{\mu\nu} d^{b}) \gamma^{5} (G_{\mu\nu} u)^{c}] , \\ J_{5} &= \varepsilon^{abc} g_{s} [(u^{aT} C \gamma^{\mu} u^{b}) \gamma^{\nu} (\tilde{G}_{\mu\nu} d)^{c} - (u^{aT} C \gamma^{\mu} d^{b}) \gamma^{\nu} (\tilde{G}_{\mu\nu} u)^{c}] , \\ J_{6} &= \varepsilon^{abc} g_{s} [(u^{aT} C \gamma^{\mu} u^{b}) \gamma^{\nu} (\tilde{G}_{\mu\nu} d)^{c} + 2(u^{aT} C \gamma^{\mu} d^{b}) \gamma^{\nu} (\tilde{G}_{\mu\nu} u)^{c}] , \\ J_{7} &= \varepsilon^{abc} g_{s} [(u^{aT} C \sigma^{\mu\nu} u^{b}) (\tilde{G}_{\mu\nu} d)^{c} - (u^{aT} C \sigma^{\mu\nu} d^{b}) (\tilde{G}_{\mu\nu} u)^{c}] , \\ J_{8} &= \varepsilon^{abc} g_{s} [(u^{aT} C \sigma^{\mu\nu} u^{b}) (\tilde{G}_{\mu\nu} d)^{c} + 2(u^{aT} C \sigma^{\mu\nu} d^{b}) (\tilde{G}_{\mu\nu} u)^{c}] , \\ J_{1\mu} &= \varepsilon^{abc} g_{s} [(u^{aT} C \gamma^{\nu} u^{b}) (G_{\mu\nu} d)^{c} - (u^{aT} C \gamma^{\nu} d^{b}) (G_{\mu\nu} u)^{c}] , \\ J_{2\mu} &= \varepsilon^{abc} g_{s} [(u^{aT} C \gamma^{\nu} u^{b}) (G_{\mu\nu} d)^{c} - (u^{aT} C \gamma^{\nu} d^{b}) (G_{\mu\nu} u)^{c}] , \\ J_{3\mu} &= \varepsilon^{abc} g_{s} [(u^{aT} C \gamma^{\nu} u^{b}) (\gamma^{5} (\tilde{G}_{\mu\nu} d)^{c} - (u^{aT} C \gamma^{\nu} d^{b}) (G_{\mu\nu} u)^{c}] , \\ J_{4\mu} &= \varepsilon^{abc} g_{s} [(u^{aT} C \gamma^{\nu} u^{b}) \gamma^{5} (\tilde{G}_{\mu\nu} d)^{c} - (u^{aT} C \gamma^{\nu} d^{b}) \gamma^{5} (\tilde{G}_{\mu\nu} u)^{c}] , \\ (1) \end{split}$$

where u/d is the up/down quark field, $G_{\mu\nu}$ is the gluon field strength, and $\tilde{G}_{\mu\nu} = \frac{1}{2} \varepsilon_{\mu\nu\alpha\beta} G^{\alpha\beta}$ the dual gluon field strength. The currents $J_{1,3,5,7,1\mu,3\mu}$ can couple to the nucleon hybrids $N_{1/2^{\pm}}, N_{3/2^{\pm}}$ with I = 1/2, while $J_{2,4,6,8,2\mu,4\mu}$ couple to the delta hybrids $\Delta_{1/2^{\pm}}, \Delta_{3/2^{\pm}}$ with I = 3/2. The couplings between currents and the corresponding positive-parity states can be written as

$$\langle 0|J_i|H_+(p)\rangle = f_+u(p),$$

$$\langle 0|J_{i\mu}|H_+(p)\rangle = f_+u_\mu(p),$$

$$(2)$$

in which u(p) and $u_{\mu}(p)$ are the Dirac spinor and Rarita-Schwinger vector, respectively, and f_+ is the coupling constant. These currents can also couple to the negative-parity states via different coupling relations

$$\langle 0|J_i|H_-(p)\rangle = f_-\gamma_5 u(p), \langle 0|J_{i\mu}|H_-(p)\rangle = f_-\gamma_5 u_\mu(p).$$
 (3)

The two-point correlation functions can be written as

$$\Pi(p^{2}) = i \int d^{4}x e^{ip \cdot x} \langle 0|T \left[J(x)\bar{J}(0)\right]|0\rangle = \Pi_{1/2}(p^{2}),$$

$$\Pi_{\mu\nu}(p^{2}) = i \int d^{4}x e^{ip \cdot x} \langle 0|T \left[J_{\mu}(x)\bar{J}_{\nu}(0)\right]|0\rangle$$

$$= -g_{\mu\nu}\Pi_{3/2}(p^{2}) + \cdots,$$
(4)

where $\Pi_{1/2}(p^2)$ and $\Pi_{3/2}(p^2)$ correspond to the hadron states with spin-1/2 and 3/2 respectively. They contain information of both positive- and negative-parity states

$$\Pi(p^{2}) = p \Pi_{p} + \Pi_{I}$$

$$= \left(\frac{p}{\sqrt{p^{2}}} + 1\right) \Pi_{+}(p^{2}) + \left(\frac{p}{\sqrt{p^{2}}} - 1\right) \Pi_{-}(p^{2}) \quad (5)$$

in which $p = \gamma_{\mu} p^{\mu}$. Π_+ and Π_- are the correlation function for positive- and negative-parity states respectively, which satisfy the following equation,

$$\Pi_{\pm}(p^2) = \frac{f_{\pm}'^2}{M_{\pm}^2 - p^2} + \cdots, \qquad (6)$$

where " \cdots " denote the excited states and continuum contributions, $f_{\pm}^{\prime 2} = M_{\pm}f_{\pm}^2$. At the hadronic level, we use the dispersion relation to describe the invariant function as

$$\Pi(p^2) = \frac{(p^2)^N}{\pi} \int_0^\infty \frac{\operatorname{Im}\Pi(s)}{s^N \left(s - p^2 - i\epsilon\right)} ds + \sum_{n=0}^{N-1} b_n (p^2)^n \,. \tag{7}$$

The spectral function can be defined in the "narrow resonance" approximation

$$\rho(s) \equiv \frac{\mathrm{Im}\Pi(s)}{\pi} = p \rho^{p}(s) + \rho^{I}(s)$$
(8)
= $f_{+}^{\prime 2} (\frac{p}{\sqrt{s}} + 1) \delta(s - M_{+}^{2}) + f_{-}^{\prime 2} (\frac{p}{\sqrt{s}} - 1) \delta(s - M_{-}^{2}) + \cdots$

where

$$\rho^{p}(s) = \frac{f_{+}^{\prime 2}}{\sqrt{s}} \delta\left(s - M_{+}^{2}\right) + \frac{f_{-}^{\prime 2}}{\sqrt{s}} \delta\left(s - M_{-}^{2}\right) + \cdots,$$

$$\rho^{I}(s) = f_{+}^{\prime 2} \delta\left(s - M_{+}^{2}\right) - f_{-}^{\prime 2} \delta\left(s - M_{-}^{2}\right) + \cdots.$$
(9)

The negative- and positive-parity states can be extracted as

$$\rho_{\pm}(s) = \sqrt{s}\rho^{p}(s) \pm \rho^{I}(s). \tag{10}$$

Then the QCD sum rules can be established as

$$\Pi_{\pm}(M_B^2, s_0) = \int_{(3m_q)^2}^{s_0} \rho_{\pm}(s) e^{-s/M_B^2} ds = 2f_{\pm}'^2 e^{-M_{\pm}^2/M_B^2}, \quad (11)$$

where $\rho_{\pm}(s) = \text{Im}\Pi_{\pm}(s)/\pi$, m_q is the light quark mass (q = u/d), M_B and s_0 are the Borel mass and continuum threshold respectively. It is clear that $\rho_{\pm}(s)$ contains only contributions from positive/negative-parity baryon states. The lowest-lying hadron mass can be obtained as

$$M_{\pm}\left(M_B^2, s_0\right) = \sqrt{\frac{\frac{\partial}{\partial \left(-1/M_B^2\right)} \Pi_{\pm}\left(M_B^2, s_0\right)}{\Pi_{\pm}\left(M_B^2, s_0\right)}}.$$
 (12)

At the QCD side, the correlation functions and the spectral functions can be evaluated via the operator product expansion (OPE). In this work, we calculate the correlation functions at the leading order of α_s up to dimension-eight condensates,



FIG. 1: Feynman diagrams involved in the calculations.

including the Feynman diagrams in Fig. 1. As an example, we show the spectral function for the current J_3 as

$$\rho_{3}^{p}(s) = -\frac{19\left\langle\alpha_{s}GG\right\rangle^{2}}{3072\pi^{2}} + \frac{5\left\langle g_{s}^{3}G^{3}\right\rangle M^{2}}{2048\pi^{4}}$$
(13)
$$-\frac{151\alpha_{s}\left\langle g_{s}\bar{q}\sigma Gq\right\rangle \left\langle \bar{q}q\right\rangle}{576\pi} + \frac{\alpha_{s}\left\langle \bar{q}q\right\rangle^{2}s}{12\pi} + \frac{3\alpha_{s}s^{4}}{20480\pi^{5}},$$

$$\rho_{3}^{I}(s) = -\frac{\left\langle\alpha_{s}GG\right\rangle \left\langle \bar{q}q\right\rangle s}{32\pi} - \frac{13\alpha_{s}\left\langle g_{s}\bar{q}\sigma Gq\right\rangle s^{2}}{256\pi^{3}} + \frac{5\alpha_{s}\left\langle \bar{q}q\right\rangle s^{3}}{192\pi^{3}}.$$
(14)

At the leading order of α_s , the dimension-4 and dimension-6 gluon condensates give no contribution for the spin-1/2 currents $J_1 - J_8$. We calculate the $\langle \bar{q}q \rangle \langle \bar{q}q \rangle$ and $\langle \bar{q}q \rangle \langle \alpha_s GG \rangle$ condensates by applying the factorization assumption, which is usually adopted in QCD sum rules to estimate the values of the high dimension condensates [64, 65]. Generally speaking, there are various Feynman diagrams contributing to the dimension-8 gluon condensates and the calculations are quite complicated. In this work, we consider part of these diagrams as shown in Fig. 1. The omission of any additional diagram will be suppressed because the Borel window criterion of Eq. (17) will be chosen to be be less than a 10% effect in our following analyses. One can consult Refs. [66, 67] for the methods of such calculations. There are seven independent gluon condensates for dimension-8 operators[68]

$$O_{1} = \operatorname{Tr} \left(G_{\mu\nu} G_{\mu\nu} G_{\alpha\beta} G_{\alpha\beta} \right),$$

$$O_{2} = \operatorname{Tr} \left(G_{\mu\nu} G_{\alpha\beta} G_{\mu\nu} G_{\alpha\beta} \right),$$

$$O_{3} = \operatorname{Tr} \left(G_{\mu\nu} G_{\nu\alpha} G_{\alpha\beta} G_{\beta\mu} \right),$$

$$O_{4} = \operatorname{Tr} \left(G_{\mu\nu} G_{\alpha\beta} G_{\nu\alpha} G_{\beta\mu} \right),$$
(15)

$$O_5 = f_{abc} G^a_{\mu\nu} J^b_{\mu} J^c_{\nu},$$

$$O_7 = J^a_{\mu} D^2 J^a_{\mu},$$

$$O_8 = f^{abc} G^a_{\mu\nu} G^b_{\nu\lambda} D^2 G^c_{\lambda\mu},$$

where $J^a_{\mu} = g_s \sum_{u,d,s} \bar{q} \gamma_{\mu} (\lambda^a/2) q$. We also apply the factorization assumption for these operators to simplify the calculations, as discussed in Refs. [68, 69]

$$g_{s}^{4}O_{1} = \frac{16\pi^{2}}{12} \langle \alpha_{s}GG \rangle^{2},$$

$$g_{s}^{4}O_{2} = -\frac{5 \times 16\pi^{2}}{48} \langle \alpha_{s}GG \rangle^{2} + 2\phi,$$

$$g_{s}^{4}O_{3} = g_{s}^{4}O_{4} = \frac{16\pi^{2}}{192} \langle \alpha_{s}GG \rangle^{2} + \frac{1}{2}\phi,$$

$$g_{s}^{3}O_{5} = -\frac{3}{2}g_{s}^{4} \langle \bar{q}q \rangle^{2}M^{2},$$

$$g_{s}^{2}O_{7} = -\frac{4}{3}g_{s}^{4} \langle \bar{q}q \rangle^{2}M^{2},$$

$$g_{s}^{3}O_{8} = \langle g_{s}^{3}G^{3} \rangle M^{2},$$
(16)

where ϕ is chosen as $\frac{7 \times 16\pi^2}{96} \langle \alpha_s GG \rangle^2$ [70] and $M^2 = 0.3 \text{ GeV}^2$ [71].

In Eq. (12), one should find reasonable working regions of s_0 and M_B^2 to determine the hadron mass. We require a good OPE convergence to obtain the lower bound of M_B^2

$$R_{D>6} = \left| \frac{\Pi_{\pm}^{D>6} \left(M_B^2, \infty \right)}{\Pi_{\pm}^{tot} \left(M_B^2, \infty \right)} \right|, \qquad (17)$$

where $\Pi_{\pm}^{D>6}(M_B^2, \infty)$ represents the contributions from D > 6 condensates. The upper bound of M_B^2 and the optimal value of s_0 are found by ensuring a sufficiently large pole contribution

$$PC = \frac{\Pi_{\pm} \left(M_B^2, s_0 \right)}{\Pi_{\pm} \left(M_B^2, \infty \right)}.$$
 (18)

Numerical analyses.—To perform numerical analyses, we use the following values for various QCD parameters at the renormalization scale $\mu = 1$ GeV and $\Lambda_{QCD} = 300$ MeV [69, 72–74]:

$$m_{u} = m_{d} = m_{q} = 0,$$

$$\alpha_{s}(\mu) = \frac{4\pi}{29/3 \ln \left(\mu^{2}/\Lambda_{\text{QCD}}^{2}\right)},$$

$$\left\langle \bar{q}q \right\rangle = -(0.24 \pm 0.01)^{3} \text{ GeV}^{3},$$

$$\left\langle g_{s}\bar{q}\sigma Gq \right\rangle = (0.8 \pm 0.2) \times \left\langle \bar{q}q \right\rangle \text{ GeV}^{2},$$

$$\left\langle \alpha_{s}GG \right\rangle = (6.35 \pm 0.35) \times 10^{-2} \text{ GeV}^{4},$$

$$\left\langle g_{s}^{3}G^{3} \right\rangle = (8.2 \pm 1.0) \times \left\langle \alpha_{s}GG \right\rangle \text{ GeV}^{2}.$$
(19)

As an example, we show our numerical analysis for the current J_3 , which can couple to both negative- and positive-parity



FIG. 2: Variations of hadron mass with (a) continuum threshold s_0 and (b) Borel parameter M_B^2 for the hybrid baryon with $I(J^P) = 1/2(1/2)^-$ from J_3 .

hybrid baryons with I(J) = 1/2(1/2). We firstly investigate the hybrid baryon with $J^P = 1/2^-$. We require $R_{D>6} \le 10\%$ to guarantee the good OPE convergence, leading to the lower bound of the Borel mass $M_B^2 \ge 1.59 \text{ GeV}^2$. By using this value and requiring that the pole contribution be larger than 20%, the threshold value s_0 can be determined around 6.5 GeV². We set 5% uncertainty of s_0 in our analyses. In Fig. 2(a), we show the variations of the hybrid baryon mass m_X with s_0 for various Borel mass M_B^2 , from which the optimal working region is $6.18 \le s_0 \le 6.83 \text{ GeV}^2$. Using the central value of s_0 , the upper bound of M_B^2 can be obtained by requiring the pole contribution be larger than 20%. Finally, the Borel parameter can be determined to be $1.59 \le M_B^2 \le 2.05 \text{ GeV}^2$.

As shown in Fig. 2(b), the Borel curves are very stable in the working parameter regions so that the mass predictions are reliable. The mass of the hybrid baryon with $I(J^P) = 1/2(1/2)^-$ is predicted as

$$M_{-} = 2.28^{+0.07}_{-0.06} \,\text{GeV}\,,\tag{20}$$

where the errors come from s_0 and various condensates in Eq. (19).

For all interpolating currents in Eq. (1), the numerical analyses show that the OPE convergences and pole contributions may vary significantly for different channels. For instance, the pole contributions for the positive-parity channels from currents J_6 and $J_{2\mu}$ are significant so that we set $PC \ge 50\%$ in the analyses. Besides, the OPE convergence behavior for the positive-parity channel from $J_{2\mu}$ is much better than all other ones. These channels can actually provide more reliable mass predictions.

In Table I, we collect the numerical results for the hybrid baryons by using all interpolating currents in Eq. (1). It is shown that there are reliable mass predictions for the positive-parity $N_{1/2^+}, \Delta_{3/2^+}, \Delta_{1/2^+}$ and negative-parity $N_{1/2^-}, N_{3/2^-}, \Delta_{1/2^-}$ hybrid baryons. For some currents, the QCD sum rules are unstable so that the mass predictions are unreliable for these channels, e.g., $N_{3/2^+}$ and $\Delta_{3/2^-}$, which are denoted as "-" in Table I. As depicted in Fig. 3, the lightest hybrid baryon is predicted to appear in the negative-parity $N_{1/2^-}$ channel, while the lightest $\Delta_{1/2^-}$ state is about 300 MeV higher. Such a mass splitting is almost the same as the mass difference $m_{\Delta} - m_N$ for conventional baryons. For the other channels, the hybrid baryons are predicted to be above 3 GeV.

 TABLE I: Mass spectra for all hybrid baryons. The notation

 "-" means no reliable mass prediction in this channel.

Current	N/Δ	$s_0[\text{GeV}^2]$	M_B^2 [GeV ²]	Mass[GeV]	PC[%]
J_1	$N_{1/2^{-}}$	11.0(±5%)	2.75-2.87	$3.01^{+0.06}_{-0.06}$	21.2
	$N_{1/2^{+}}$	-	-	-	-
J_2	$\Delta_{1/2^-}$	16.5(±5%)	2.53-4.25	$3.64^{+0.08}_{-0.08}$	34.7
	$\Delta_{1/2^+}$	-	-	-	-
J_3	$N_{1/2^{-}}$	6.5(±5%)	1.59-2.05	$2.28^{+0.07}_{-0.06}$	26.8
	$N_{1/2^{+}}$	13.0(±5%)	2.00-2.87	$3.37^{+0.28}_{-0.15}$	29.3
J_4	$\Delta_{1/2^-}$	8.0(±5%)	2.43-2.57	$2.64^{+0.39}_{-0.15}$	10.9
	$\Delta_{1/2^+}$	-	-	-	-
J_5	$N_{1/2^{-}}$	10.5(±5%)	2.78-3.06	$2.91^{+0.05}_{-0.05}$	22.6
	$N_{1/2^{+}}$	-	-	-	-
J_6	$\Delta_{1/2^{-}}$	-	-	-	-
	$\Delta_{1/2^+}$	20.0(±5%)	3.09-3.32	$4.03^{+0.13}_{-0.11}$	52.5
J_7	N _{1/2} -	14.5(±5%)	2.21-3.44	$3.48^{+0.14}_{-0.11}$	32.3
	$N_{1/2^{+}}$	-	-	-	-
J_8	$\Delta_{1/2^{-}}$	12.5(±5%)	2.02-3.56	$3.15^{+0.06}_{-0.06}$	36.1
	$\Delta_{1/2^+}$	-	-	-	-
$J_{1\mu}$	N _{3/2} -	17.5(±5%)	3.28-4.58	$3.76^{+0.10}_{-0.09}$	29.6
	$N_{3/2^{+}}$	-	-	-	-
$J_{2\mu}$	$\Delta_{3/2^{-}}$	-	-	-	-
	$\Delta_{3/2^+}$	15.0(±5%)	2.39-2.76	$3.43^{+0.06}_{-0.06}$	55.6
$J_{3\mu}$	N _{3/2} -	12.0(±5%)	2.88-3.44	$3.12^{+0.08}_{-0.08}$	24.9
	$N_{3/2^{+}}$	-	-	-	-
$J_{4\mu}$	$\Delta_{3/2^{-}}$	-	-	-	-
	$\Delta_{3/2^+}$	-	-	-	-
5.0				-	
-				• N	egative-parity
4.5				Po	ositive-parity
4.0					I
- ^{4.0}		Ŧ	_		I
			T		



FIG. 3: Mass spectra for both negative-parity and positive-parity hybrid baryons.

Considering the renormalization scale at $\mu = 2$ GeV, the predicted hybrid masses will be slightly increased without changing the overall spectrum structure. One may wonder that the traditional baryon states with the same quantum numbers also contribute to the correlation functions. Following the analysis of Ref. [34], we note that mixing with (lighter) conventional baryons that are weakly coupled to the hybrid currents can only raise the hybrid mass predictions in Table I.

Production and decay mechanisms.-The decay of char-



FIG. 4: Possible production mechanisms of light hybrid baryons from Υ/χ_{cJ} decays.



FIG. 5: Possible decay mechanisms of light hybrid baryon to gluon-rich channels.

monium and bottomonium mesons can provide an ideal platform for producing gluon-rich hadron states. For instance, the well known 1⁻⁺ hybrid candidate $\eta_1(1855)$ was observed in the radiative decays of J/ψ [46, 75]. Similarly, the charmonium/bottomonium decay processes may produce light hybrid baryons, if kinematically allowed. According to our results, all predicted hybrid baryons can be generated through the bottomonium decays via the three-gluon emission processes, such as $\Upsilon \rightarrow \bar{N}N_{hyb}/\bar{\Delta}\Delta_{hyb} + c.c.$. Besides, the lightest $N_{1/2^-}$ hybrid state can also be generated through the charmonium decays $\chi_{cJ} \rightarrow \bar{N}N_{1/2^-} + c.c.$. We propose such hybrid baryon production mechanisms in Fig. 4.

Carrying the same quantum numbers, the hybrid baryons could always mix with traditional baryon states, so that their decay channels to the conventional baryon plus meson are similar to the latter ones, as shown in Table II. Moreover, a hybrid baryon may decay into final states containing gluonrich hadrons such as hybrid mesons or glueballs, as depicted in Fig. 5. Such characteristic decays may be useful for distinguishing hybrid baryons from conventional states and pentaquark states.

Summary.—We calculate the masses of the light hybrid baryons with quantum numbers $I(J^P) = 1/2(1/2^{\pm}), 3/2(1/2^{\pm}), 1/2(3/2^{\pm})$ and $3/2(3/2^{\pm})$ in parityprojected QCD sum rule method by constructing local interpolating currents. The correlation functions and spectral densities are evaluated up to dimension eight condensates. We establish stable mass sum rules for the positive-parity $N_{1/2^+}, \Delta_{3/2^+}, \Delta_{1/2^+}$ and negative-parity $N_{1/2^-}, N_{3/2^-}, \Delta_{1/2^-}$ hybrid baryons, so that the mass predictions in these channels are reliable.

TABLE II: Some possible decay modes into conventional baryon-meson final states for light hybrid baryons.

Current	Hybrid	Decay Modes
J_1	$N_{1/2^{-}}$	$\frac{N\pi/\eta^{(\prime)}/\rho/\omega, N(1440)\pi/\eta'/\rho/\omega,}{\Delta\rho, \Delta(1600)\rho, N(1535)f_0(980)/a_0(980)}$
J_2	$\Delta_{1/2^{-}}$	$N\pi/ ho, N(1440)\pi/ ho, \\ \Delta(1232) ho, N(1535)a_0(980)$
J_3	N _{1/2} -	$N\pi/\eta^{(\prime)}/\rho/\omega, N(1440)\pi/\eta^{(\prime)}, \Delta(1232)\rho$
	$N_{1/2^{+}}$	$N f_0(980)/a_0(980), \Delta(1232)\pi/\rho, N(1440)f_0(980)/a_0(980)$
J_4	$\Delta_{1/2^{-}}$	$N\pi/\rho, N(1440)\pi/\rho, \Delta(1232)\rho, N(1535)a_0(980)$
J_5	N _{1/2} -	$\frac{N\pi/\eta^{(\prime)}/\rho/\omega, N(1440)\pi/\eta^{(\prime)}/\rho/\omega,}{\Delta(1232)\rho, \Delta(1600)\rho, N(1535)f_0(980)/a_0(980)}$
J_6	$\Delta_{1/2^+}$	$Na_0(980), N(1440)a_0(980), \Delta(1232)a_0(980)f_0(980)$
J_7	N _{1/2} -	$\frac{N\pi/\eta^{(\prime)}/\rho/\omega, N(1440)\pi/\eta'/\rho/\omega,}{\Delta(1232)\rho, \Delta(1600)\rho, N(1535)f_0(980)/a_0(980)}$
J_8	$\Delta_{1/2^-}$	$N\pi/\rho, N(1440)\pi/\rho, \Delta(1232)\rho, N(1535)a_0(980)$
${J}_{1\mu}$	N _{3/2} -	$N\rho/\omega, N(1400)\rho/\omega, \Delta(1232)\rho$
$J_{2\mu}$	$\Delta_{3/2^{+}}$	$Nb_1/\rho, N(1440)b_1/\rho, \Delta(1232)b_1/h_1/\rho$
$J_{3\mu}$	N _{3/2} -	$N\rho/\omega, N(1400)\rho/\omega, \Delta(1232)\rho$

The lightest hybrid baryons are predicted to be the negativeparity states $N_{1/2^-}$ and $\Delta_{1/2^-}$ around 2.28 GeV and 2.64 GeV, respectively. This is in contrast to the prediction from LQCD, which suggests that the lightest hybrid baryons are the positive-parity states [55]. However, recent investigations using a semirelativistic potential model suggest that heavy hybrid baryons with negative-parity $J^P = 1/2^-$, $3/2^-$ are ground states, featuring a $J_C^{Pc} = 1/2^+$ triquark core and a $J_g^{Pg} = 1^$ gluon excitation [61], which supports our calculations of the mass spectra to some extent. All other hybrids in our results are predicted to be above 3 GeV. These hybrid baryons may primarily decay into the conventional baryon-meson final states. We suggest searching for such hybrids through the gluon-rich processes in the BESIII and BelleII experiments.

ACKNOWLEDGMENTS: This work is supported by the National Natural Science Foundation of China under Grant No. 12305147, No. 12175318 and No. 12075019, the Natural Science Foundation of Guangdong Province of China under Grants No. 2022A1515011922. TGS is grateful for research funding from the Natural Sciences & Engineering Research Council of Canada (NSERC). JH was supported by the Mitacs Globalink Research Award and is grateful for the support and hospitality of SYSU. opments in the Quark Theory of Hadrons, volume 1 (1964)

- [3] C. Meyer and E. Swanson, Prog. Part. Nucl. Phys. 82, 21 (2015)
- [4] H.-X. Chen, W. Chen, X. Liu, and S.-L. Zhu, Phys. Rep. 639, 1 (2016)
- [5] A. Esposito, A. Pilloni, and A. D. Polosa, Phys. Rept. **668**, 1 (2017)
- [6] F.-K. Guo, C. Hanhart, U.-G. Meißner, Q. Wang, Q. Zhao, and B.-S. Zou, Rev. Mod. Phys. 90, 015004 (2018)
- [7] Y.-R. Liu, H.-X. Chen, W. Chen, X. Liu, and S.-L. Zhu, Prog. Part. Nucl. Phys. 107, 237 (2019)
- [8] N. Brambilla et al., Phys. Rep. 873, 1 (2020)
- [9] H.-X. Chen, W. Chen, X. Liu, Y.-R. Liu, and S.-L. Zhu, Rept. Prog. Phys. 86, 026201 (2023)
- [10] L. Meng, B. Wang, G.-J. Wang, and S.-L. Zhu, Phys. Rept. 1019, 1 (2023)
- [11] T. Barnes, F. E. Close, and F. de Viron, Nucl. Phys. B 224, 241 (1983)
- [12] M. S. Chanowitz and S. R. Sharpe, Nucl. Phys. B 222, 211 (1983)
- [13] P. Guo, A. P. Szczepaniak, G. Galata, A. Vassallo, and E. Santopinto, Phys. Rev. D 78, 056003 (2008)
- [14] P. Lacock, C. Michael, P. Boyle, and P. Rowland, Phys. Lett. B 401, 308 (1997)
- [15] P. Lacock, C. Michael, P. Boyle, and P. Rowland, Phys. Rev. D 54, 6997 (1996)
- [16] J. J. Dudek, R. G. Edwards, M. J. Peardon, D. G. Richards, and C. E. Thomas, Phys. Rev. Lett. **103**, 262001 (2009)
- [17] J. J. Dudek, R. G. Edwards, M. J. Peardon, D. G. Richards, and C. E. Thomas, Phys. Rev. D 82, 034508 (2010)
- [18] J. J. Dudek, R. G. Edwards, B. Joo, M. J. Peardon, D. G. Richards, and C. E. Thomas, Phys. Rev. D 83, 111502 (2011)
- [19] J. J. Dudek, Phys. Rev. D 84, 074023 (2011)
- [20] L. Liu et al., JHEP 07, 126 (2012)
- [21] J. J. Dudek, R. G. Edwards, P. Guo, and C. E. Thomas, Phys. Rev. D 88, 094505 (2013)
- [22] Y. Ma, Y. Chen, M. Gong, and Z. Liu, Chin. Phys. C 45, 013112 (2021)
- [23] N. Isgur and J. E. Paton, Phys. Rev. D **31**, 2910 (1985)
- [24] N. Isgur, R. Kokoski, and J. Paton, Phys. Rev. Lett. 54, 869 (1985)
- [25] T. J. Burns and F. E. Close, Phys. Rev. D 74, 034003 (2006)
- [26] C. J. Burden and M. A. Pichowsky, Few Body Syst. 32, 119 (2002)
- [27] C. J. Burden, L. Qian, C. D. Roberts, P. C. Tandy, and M. J. Thomson, Phys. Rev. C 55, 2649 (1997)
- [28] I. I. Balitsky, D. Diakonov, and A. V. Yung, Phys. Lett. B 112, 71 (1982)
- [29] J. Govaerts, F. de Viron, D. Gusbin, and J. Weyers, Phys. Lett. B 128, 262 (1983)
- [30] J. I. Latorre, S. Narison, P. Pascual, and R. Tarrach, Phys. Lett. B 147, 169 (1984)
- [31] J. Govaerts, F. de Viron, D. Gusbin, and J. Weyers, Nucl. Phys. B 248, 1 (1984)
- [32] J. Govaerts, L. J. Reinders, H. R. Rubinstein, and J. Weyers, Nucl. Phys. B 258, 215 (1985)
- [33] I. I. Balitsky, D. Diakonov, and A. V. Yung, Z. Phys. C 33, 265 (1986)
- [34] W. Chen, R. T. Kleiv, T. G. Steele, B. Bulthuis, D. Harnett, J. Ho, T. Richards and S. L. Zhu, JHEP 09 (2013), 019

- [35] J. Ho, R. Berg, W. Chen, D. Harnett, and T. G. Steele, Phys. Rev. D 98, 096020 (2018)
- [36] J. Ho, R. Berg, T. G. Steele, W. Chen, and D. Harnett, Phys. Rev. D 100, 034012 (2019)
- [37] A. Palameta, D. Harnett, and T. G. Steele, Phys. Rev. D 98, 074014 (2018)
- [38] B. Barsbay, K. Azizi, and H. Sundu, Eur. Phys. J. C 82, 1086 (2022)
- [39] H.-X. Chen, W. Chen, and S.-L. Zhu, Phys. Rev. D 105, L051501 (2022)
- [40] H.-X. Chen, N. Su, and S.-L. Zhu, Chin. Phys. Lett. 39, 051201 (2022)
- [41] Q.-N. Wang, D.-K. Lian, and W. Chen, Phys. Rev. D 108, 114010 (2023)
- [42] D.-K. Lian, Q.-N. Wang, X.-L. Chen, P.-F. Yang, W. Chen, and H.-X. Chen, JHEP 06, 173 (2024)
- [43] D. Alde et al., Phys. Lett. B 205, 397 (1988)
- [44] E. I. Ivanov et al., Phys. Rev. Lett. 86, 3977 (2001)
- [45] J. Kuhn et al., Phys. Lett. B 595, 109 (2004)
- [46] M. Ablikim et al., Phys. Rev. Lett. 129, 192002 (2022)
- [47] M. Ablikim et al., Phys. Rev. D 106, 072012 (2022)
- [48] T. Barnes and F. E. Close, Phys. Lett. B 123, 89 (1983)
- [49] E. Golowich, E. Haqq, and G. Karl, Phys. Rev. D 28, 160 (1983)
- [50] L. S. Kisslinger and Z. P. Li, Phys. Rev. D **51**, R5986 (1995)
- [51] V. D. Burkert and C. D. Roberts, Rev. Mod. Phys. 91, 011003 (2019)
- [52] S. Capstick and P. R. Page, Phys. Rev. D 60, 111501 (1999)
- [53] S. Capstick and P. R. Page, Phys. Rev. C 66, 065204 (2002)
- [54] S. M. Gerasyuta and V. I. Kochkin, Phys. Rev. D 66, 116001 (2002)
- [55] J. J. Dudek and R. G. Edwards, Phys. Rev. D 85, 054016 (2012)
- [56] T. Khan, D. Richards, and F. Winter, Phys. Rev. D 104, 034503 (2021)
- [57] A. P. Martynenko, Sov. J. Nucl. Phys. 54 (1991), 488-490
- [58] L. S. Kisslinger, Phys. Rev. D 69, 054015 (2004)
- [59] K. Azizi, B. Barsbay, and H. Sundu, Eur. Phys. J. Plus 133, 121 (2018)
- [60] Y.-C. Zhao, C.-M. Tang, and L. Tang, Eur. Phys. J. C 83, 654 (2023)
- [61] L. Cimino, C. T. Willemyns, and C. Semay, Phys. Rev. D 110, 034032 (2024)
- [62] B. Ioffe, Nucl. Phys. B 188, 317 (1981)
- [63] D. Jido, N. Kodama, and M. Oka, Phys. Rev. D 54, 4532 (1996)
- [64] M. Shifman, A. Vainshtein, and V. Zakharov, Nucl. Phys. B 147, 385 (1979)
- [65] L. Reinders, H. Rubinstein, and S. Yazaki, Phys. Rep. 127, 1 (1985)
- [66] A. V. Pimikov, Phys. Rev. D 106, 056011 (2022)
- [67] A. G. Grozin, Int. J. Mod. Phys. A 10, 3497 (1995)
- [68] Z.-R. Huang, H.-Y. Jin, and Z.-F. Zhang, JHEP 04, 004 (2015)
- [69] S. Narison, Phys. Lett. B 706, 412 (2012)
- [70] E. Bagan, J. I. Latorre, P. Pascual, and R. Tarrach, Nucl. Phys. B 254, 555 (1985)
- [71] S. N. Nikolaev, A. V. Radyushkin, Phys. Lett. B 124, 243 (1983)
- [72] R. L. Workman et al., PTEP **2022**, 083C01 (2022)
- [73] S. Narison, Int. J. Mod. Phys. A 33, 1850045 (2018)
- [74] M. Jamin, Phys. Lett. B **538**, 71 (2002)
- [75] F. Chen et al., Phys. Rev. D 107, 054511 (2023)