

# Quadratic maps between non-abelian groups

Asgar Jamneshan and Andreas Thom

ABSTRACT. The aim of this article is to study quadratic maps, in the sense of Leibman, between non-abelian groups. As a key result, we provide a complete characterization of quadratic maps through a universal construction that can be computed explicitly. Applications include the classification of quadratic maps on finite abelian groups, proving the non-existence of polynomial maps of higher degree on perfect groups, establishing stability results for polynomial maps, and presenting a general inverse theorem for the Gowers norm with non-abelian coefficients on perfect groups with bounded commutator width.

## CONTENTS

1. Introduction	1
2. Basic results	4
3. The computation of $\text{Pol}_2(G)$ and its abelianization	7
4. Some computations and consequences	10
5. Approximate polynomial maps	13
6. The Gowers norms and an inverse theorem	16
Acknowledgments	20
References	21

## 1. Introduction

The notion of a polynomial map, for instance between vector spaces, can be defined independently of the multiplication of coordinates. Specifically, polynomial maps of degree 0 are constant, while polynomial maps  $f: \mathbb{R}^n \rightarrow \mathbb{R}^m$  of degree  $d + 1$  are precisely those for which the finite differences  $(\Delta_y f)(x) := f(y + x) - f(x)$  are polynomials of degree  $d$  for all  $y \in \mathbb{R}^n$ .

Various generalizations of this idea to the non-abelian setting have been proposed. In this article, we follow the framework established by Leibman

[11, 12]. Let  $G$  and  $H$  be groups. For a map  $\varphi: G \rightarrow H$  and  $k \in G$ , we define the non-abelian analogue of finite differences as follows:

$$(\Delta_k \varphi)(g) := \varphi(kg)\varphi(g)^{-1}.$$

DEFINITION 1.1. *A map  $\varphi: G \rightarrow H$  is said to be polynomial of degree  $-1$  if  $\varphi(g) = 1$  for all  $g \in G$ . For  $d \geq -1$ , a map  $\varphi: G \rightarrow H$  is said to be polynomial of degree  $d+1$  if, for all  $k \in G$ , the map  $\Delta_k(\varphi)$  is polynomial of degree  $d$ .*

It is clear from the definition that the composition of a polynomial of degree  $d$  with a homomorphism in either order is again a polynomial map of degree  $d$ . It also follows easily that polynomial maps of degree 0 are constant functions, and that polynomial maps of degree 1 are the product of a homomorphism and a constant function (see Lemma 2.1). The primary focus of this paper is the study of the first non-trivial case: quadratic maps, i.e., polynomials of degree at most two. It also follows directly from the definition that if  $\varphi: G \rightarrow H$  is a quadratic map, then the map  $\varphi': G \rightarrow H$  defined by  $\varphi'(g) = \varphi(g)h$ , for some fixed  $h \in H$ , is also quadratic. Consequently, without loss of generality, we may assume that  $\varphi(1) = 1$ . We call a map  $\varphi: G \rightarrow H$  *unital* if it satisfies  $\varphi(1) = 1$ , and denote by  $\text{Quad}(G, H)$  the set of unital quadratic maps from  $G$  to  $H$ . For a fixed group  $G$ , we can then interpret  $\text{Quad}(G, ?)$  as a functor from the category of groups to the category of sets.

Our main result provides a characterization of quadratic maps in terms of group homomorphisms, as follows:

THEOREM 1.2. *Let  $G$  be a group. There exists a group  $\text{Pol}_2(G)$  and a natural bijection*

$$\text{hom}(\text{Pol}_2(G), ?) \rightarrow \text{Quad}(G, ?),$$

*i.e., the functor  $\text{Quad}(G, ?)$  is representable. More specifically, there exists a universal unital quadratic map  $\varphi: G \rightarrow \text{Pol}_2(G)$  such that every unital quadratic map  $\varphi': G \rightarrow H$  can be expressed as  $\varphi' = \psi \circ \varphi$  for a unique homomorphism  $\psi: \text{Pol}_2(G) \rightarrow H$ .*

We can describe the group  $\text{Pol}_2(G)$  in concrete terms as follows. Here,  $\omega(G)$  denotes the augmentation ideal of the group ring  $\mathbb{Z}[G]$ , and  $G^{ab}$  denotes the abelianization of  $G$ .

THEOREM 1.3. *The group  $\text{Pol}_2(G)$  is isomorphic to the twisted crossed product group*

$$(\omega(G) \otimes_{\mathbb{Z}} G^{ab}) \rtimes_{\psi} G,$$

where  $G$  acts on  $\omega(G)$  by left multiplication and trivially on  $G^{ab}$ . The 2-cocycle  $\psi$  is given by  $c \otimes \text{ab}$ , the exterior product of the tautological 1-cocycle  $c: G \rightarrow \omega(G)$  and the trivial 1-cocycle  $\text{ab}: G \rightarrow G^{ab}$ .

One may naturally inquire about the classification of polynomial maps of higher degree. While the definition of  $\text{Pol}_k(G)$  is straightforward, obtaining an explicit description such as in Theorem 1.3 seems to be more challenging.

Theorem 1.3 has several noteworthy consequences. In particular, it provides a concrete description of the abelianization of  $\text{Pol}_2(G)$ , as detailed below. This result is proved in Section 3. Here,  $\text{Sym}^2(G)$  denotes the symmetric product of an abelian group  $G$ .

**COROLLARY 1.4.** *Let  $G$  be an abelian group. The abelianization of  $\text{Pol}_2(G)$  fits into an extension*

$$1 \rightarrow \text{Sym}^2(G) \rightarrow \text{Pol}_2^{ab}(G) \rightarrow G \rightarrow 1,$$

so that  $\text{Pol}_2^{ab}(G)$  is isomorphic to the twisted crossed product group

$$\text{Sym}^2(G) \rtimes_{\sigma} G$$

for the 2-cocycle  $\sigma: G \rightarrow \text{Sym}^2(G)$  given by  $\sigma(g, h) = gh$ .

This corollary contributes to the classification of polynomial maps from a finite abelian group  $G$  to the torus  $\mathbb{T}$ . Such a classification is a key aspect of the inverse theory for the Gowers norms on abelian groups in higher-order Fourier analysis. For example, if  $G = k^n$  is a finite-dimensional vector space over a finite field  $k$ , the polynomial maps on  $G$  with values in  $\mathbb{T}$  act as obstructions in the inverse theorem for the Gowers norms on such groups [15, 16]. Their complete classification is provided in [16, Lemma 1.7]. We elaborate on this connection in Section 4.

Recall that a group  $G$  is said to be perfect if  $G$  is equal to its commutator subgroup  $[G, G]$ , equivalently,  $G^{ab}$  is trivial.

**COROLLARY 1.5.** *If  $G$  is a perfect group, then the natural homomorphism  $\pi: \text{Pol}_2(G) \rightarrow G$  is an isomorphism. Consequently, every unital polynomial map defined on a perfect group is automatically a homomorphism.*

A group  $G$  is said to have commutator width  $k$  if every element  $g \in G$  can be expressed as a product of at most  $k$  commutators. The remainder of the paper focuses on perfect groups  $G$  of bounded commutator width and addresses approximate polynomial maps with values in metric groups, see Theorem 5.5, and an inverse theorem for the Gowers norm for functions with unitary values, see Theorem 6.1. The latter section builds on the work of Gowers–Hatami [6]. We demonstrate that, in this specific setting, there is a complete collapse of the hierarchy in higher-order Fourier analysis. More

specifically any map  $\varphi: G \rightarrow U(n)$  that has  $k$ -th Gowers norm close enough to 1 is close to a homomorphism in a suitable sense; where the estimates only depend on  $k$  and the commutator width and are independent of  $G$  and  $n$ . This includes, for example, uniform estimates for the class of non-abelian simple groups, whose commutator width is well-known to be bounded, as shown already in the work of Wilson [17].

## 2. Basic results

Before we get to quadratic maps, we recall a basic result on polynomial maps of degree one.

LEMMA 2.1. *A map  $\varphi: G \rightarrow H$  is polynomial of degree one if and only if there exists  $c \in H$  and a homomorphism  $\psi: G \rightarrow H$ , such that*

$$\varphi(g) = \psi(g)c, \quad \forall g \in G.$$

PROOF. We set  $\psi(g) := \varphi(g)\varphi(1)^{-1}$ . Note that  $(\Delta_g\varphi)(h) = \varphi(gh)\varphi(h)^{-1}$  is of degree zero and therefore a constant equal to  $\varphi(g)\varphi(1)^{-1}$ . Thus, we obtain

$$\psi(gh) = \varphi(gh)\varphi(1)^{-1} = \varphi(g)\varphi(1)^{-1}\varphi(h)\varphi(1)^{-1} = \psi(g)\psi(h)$$

and conclude that  $\psi$  is a homomorphism, and set  $c := \varphi(1)$ .  $\square$

We will freely use the terminology of group actions and cocycles, see [2] for details and background. Note that there is a natural right action of  $G$  on the set of functions  $f$  from  $G$  to  $H$ , given by  $f^k(g) = f(kg)$  for all  $k, g \in G$ . Indeed, we compute

$$f^{k_1k_2}(g) = f(k_1k_2g) = f^{k_1}(k_2g) = (f^{k_1})^{k_2}(g).$$

In terms of this action, the finite differences satisfy the following cocycle relation.

LEMMA 2.2. *Let  $\varphi: G \rightarrow H$  be a map and  $k_1, k_2 \in G$ , then the following 1-cocycle identity holds:*

$$(1) \quad \Delta_{k_1k_2}\varphi = (\Delta_{k_1}\varphi)^{k_2} \cdot \Delta_{k_2}\varphi$$

PROOF. We compute

$$\begin{aligned} (\Delta_{k_1k_2}\varphi)(g) &= \varphi(k_1k_2g)\varphi(g)^{-1} \\ &= \varphi(k_1k_2g)\varphi(k_2g)^{-1}\varphi(k_2g)\varphi(g)^{-1} \\ &= (\Delta_{k_1}\varphi)(k_2g) \cdot (\Delta_{k_2}\varphi)(g) \end{aligned}$$

$\square$

Let  $\varphi: G \rightarrow H$  be a unital quadratic map. Using Lemma 2.1, for each  $k \in G$  there exists a homomorphism  $\psi_k: G \rightarrow H$  and a constant  $c_k \in H$ , such that

$$(\Delta_k \varphi)(g) = \varphi(kg)\varphi(g)^{-1} = \psi_k(g)c_k, \quad \forall g, k \in G.$$

For  $g = 1$ , we obtain  $c_k = \varphi(k)$ .

Setting

$$\beta_k(g) := \varphi(k)^{-1}\psi_k(g)\varphi(k) = \varphi(k)^{-1}(\Delta_k \varphi)(g)$$

we obtain from Equation(1) the following identity:

$$(2) \quad \beta_{k_1 k_2}(g) = \varphi(k_2)^{-1}\beta_{k_1}(g)\varphi(k_2) \cdot \beta_{k_2}(g).$$

Indeed, we compute as follows:

$$\begin{aligned} \beta_{k_1 k_2}(g) &= \varphi(k_1 k_2)^{-1}(\Delta_{k_1 k_2} \varphi)(g) \\ &= \varphi(k_1 k_2)^{-1}(\Delta_{k_1} \varphi)^{k_2}(g)(\Delta_{k_2} \varphi)(g) \\ &= \varphi(k_1 k_2)^{-1}(\Delta_{k_1} \varphi)(k_2 g)(\Delta_{k_2} \varphi)(g) \\ &= \varphi(k_1 k_2)^{-1}\varphi(k_1)\beta_{k_1}(k_2 g)\varphi(k_2)\beta_{k_2}(g) \\ &= \varphi(k_1 k_2)^{-1}\varphi(k_1)\beta_{k_1}(k_2)\beta_{k_1}(g)\varphi(k_2)\beta_{k_2}(g) \\ &= \varphi(k_2)^{-1}\beta_{k_1}(g)\varphi(k_2) \cdot \beta_{k_2}(g). \end{aligned}$$

Note that  $\beta_k: G \rightarrow H$  is also a homomorphism for  $k \in G$  and  $\beta_1$  is trivial. Moreover, we conclude from Equation (2) that the pointwise product of the homomorphism  $g \mapsto \varphi(k_2)^{-1}\beta_{k_1}(g)\varphi(k_2)$  and the homomorphism  $g \mapsto \beta_{k_2}(g)$  is again a homomorphism. The following folklore lemma shows that this is a rather special situation.

LEMMA 2.3. *Let  $\alpha, \beta: G \rightarrow H$  be homomorphisms.*

- (i) *If the pointwise product  $\alpha\beta$  is again a homomorphism, then the ranges of  $\alpha$  and  $\beta$  commute.*
- (ii) *If the pointwise inverse  $\alpha^{-1}$  is again a homomorphism, then the range of  $\alpha$  is abelian.*

PROOF. We first prove Claim (i) and compute:

$$\begin{aligned} \alpha(g)\alpha(h)\beta(g)\beta(h) &= \alpha(gh)\beta(gh) = (\alpha\beta)(gh) \\ &= (\alpha\beta)(g)(\alpha\beta)(h) = \alpha(g)\beta(g)\alpha(h)\beta(h). \end{aligned}$$

Hence,  $\alpha(h)\beta(g) = \beta(g)\alpha(h)$  for all  $g, h \in G$ . Claim (ii) follows since the pointwise product of  $\alpha$  and  $\alpha^{-1}$  is the trivial homomorphism and the ranges of  $\alpha$  and  $\alpha^{-1}$  agree.  $\square$

PROPOSITION 2.4. *The range of  $\beta_k$  is abelian for all  $k \in G$  and the ranges of the homomorphisms  $\beta_{k_1}$  and  $\beta_{k_2}$  commute for all  $k_1, k_2 \in G$ .*

PROOF. Indeed, Equation (2) shows for  $k_1 = k, k_2 = k^{-1}$  that the map  $g \mapsto \beta_k(g)^{-1}$  is also a homomorphism. By Lemma 2.3(ii), this implies that the range of  $\beta_k$  is abelian for all  $k \in G$ . Now, Lemma 2.3(i) implies that the ranges of  $g \mapsto \varphi(k_2)^{-1}\beta_{k_1}(g)\varphi(k_2)$  and  $\beta_{k_2}$  commute. Hence, we conclude also that their product  $\beta_{k_1k_2}$  commutes with  $\beta_{k_2}$ . Since  $k_1, k_2$  were arbitrary, this proves the claim.  $\square$

We can already prove our first result.

PROOF OF THEOREM 1.2: Let  $G$  be a group. We define a group  $\text{Pol}_2(G)$  on formal generators  $\varphi(g)$  for  $g \in G$ , subject to relations

$$(3) \quad \varphi(g_1g_2g_3) = \varphi(g_1g_2)\varphi(g_2)^{-1}\varphi(g_1)^{-1}\varphi(g_1g_3)\varphi(g_3)^{-1}\varphi(g_2g_3)$$

for all  $g_1, g_2, g_3 \in G$ . We claim that the tautological map  $\varphi: G \rightarrow \text{Pol}_2(G)$  is the universal unital quadratic map. Applying the relation to  $(g_1, g_2, g_3) = (1, 1, 1)$ , we see that  $\varphi(1) = 1$ , so that  $\varphi$  is unital.

The requirement for  $\varphi$  to be quadratic is precisely that  $\Delta_{g_1}\varphi$  is linear for every  $g_1$ . Recall that

$$(\Delta_{g_1}\varphi)(g) := \varphi(g_1g)\varphi(g)^{-1},$$

so that  $(\Delta_{g_1}\varphi)(1) = \varphi(g_1)$ . We set

$$\psi_{g_1}(g) := \varphi(g_1g)\varphi(g)^{-1}\varphi(g_1)^{-1}.$$

It remains to show that  $\psi_{g_1}$  is a homomorphism for every  $g_1$ . However,

$$\begin{aligned} \psi_{g_1}(g_2)\psi_{g_1}(g_3) &= \varphi(g_1g_2)\varphi(g_2)^{-1}\varphi(g_1)^{-1}\varphi(g_1g_3)\varphi(g_3)^{-1}\varphi(g_1)^{-1} \\ &\stackrel{(3)}{=} \varphi(g_1g_2g_3)\varphi(g_2g_3)^{-1}\varphi(g_1)^{-1} \\ &= \psi_{g_1}(g_2g_3). \end{aligned}$$

This proves the claim and shows that  $\varphi$  is indeed quadratic. At the same time, we see that all relations are necessary so that it follows that  $\varphi$  is indeed the universal quadratic map.  $\square$

REMARK 2.5. It is clear from the construction that a similar approach would define  $\text{Pol}_k(G)$ , the relations to put are just  $((\Delta_{g_1} \circ \dots \circ \Delta_{g_{k+1}})\varphi)(g_{k+2})$  for all  $g_1, \dots, g_{k+2} \in G$  and  $\varphi(1) = 1$ .

PROPOSITION 2.6. *The assignment  $G \mapsto \text{Pol}_2(G)$  is a functor on the category of groups.*

The following corollary will allow us to compute  $\text{Pol}_2(G)$  in various special cases.

**COROLLARY 2.7.** *Let  $G$  be a group with symmetric generating set  $S \subseteq G$  and let  $N \leq G$  be a normal subgroup generated by a symmetric set  $\Sigma \subseteq N$ . Then  $\text{Pol}_2(G/N)$  is canonically isomorphic to the quotient of  $\text{Pol}_2(G)$  by the normal subgroup generated by the elements  $\varphi(\sigma)$  and  $\varphi(\sigma s)\varphi(s)^{-1}$  for  $\sigma \in \Sigma, s \in S$ .*

**PROOF.** We set  $K_N := \langle \langle \varphi(\sigma), \varphi(\sigma s)\varphi(s)^{-1} : \sigma \in \Sigma, s \in S \rangle \rangle$  and aim to show that  $\text{Pol}_2(G)/K_N$  has the required universal property. Note that  $K_N$  lies in the kernel of the canonical map  $\text{Pol}_2(G) \rightarrow \text{Pol}_2(G/N)$ . It remains to provide an inverse to the induced homomorphism  $\text{Pol}_2(G)/K_N \rightarrow \text{Pol}_2(G/N)$ . In order to do so we show that the canonical quadratic map  $\bar{\varphi}: G \rightarrow \text{Pol}_2(G)/K_N$  is well-defined on  $G/N$ .

Now, it follows that  $\beta_\sigma(s) = \varphi(\sigma)^{-1}\varphi(\sigma s)\varphi(s)^{-1} \in K_N$  for  $\sigma \in \Sigma, s \in S$  and hence  $\beta_\sigma(g) \in K_N$  for all  $g \in G$ . We conclude from Equation (2) and the fact that  $K_N$  is normal, that  $\beta_h(g) \in K_N$  for all  $h \in N$  and  $g \in G$ . Since  $\beta_h(h') = \varphi(h)^{-1}\varphi(hh')\varphi(h')^{-1}$  it follows by induction on the word length that  $\varphi(h) \in K_N$  for all  $h \in N$ . Finally, this implies that  $\varphi(hg)\varphi(g)^{-1} \in K_N$  for all  $h \in N, g \in G$ . Thus,  $\bar{\varphi}$  is indeed well-defined on  $G/N$ .  $\square$

### 3. The computation of $\text{Pol}_2(G)$ and its abelianization

This section is concerned with the proof of our main result about the structure of  $\text{Pol}_2(G)$ . This is the basis for later sections, in which we exploit this structure result in order to derive explicit computations of  $\text{Pol}_2(G)$  in special cases and other consequences.

**PROOF OF THEOREM 1.3.** We have an extension

$$1 \rightarrow N(G) \rightarrow \text{Pol}_2(G) \xrightarrow{\pi} G \rightarrow 1$$

where  $\pi$  is the homomorphism that classifies the unital quadratic map  $\text{id}: G \rightarrow G$ , and the group  $N(G)$  is defined to be  $\ker(\pi)$ . Since  $\pi(\varphi(g)) = g$ , we have that  $\varphi: G \rightarrow \text{Pol}_2(G)$  is a canonical set-theoretic section of  $\pi$ . Recall that  $\beta_k(g) = \varphi(k)^{-1}\varphi(kg)\varphi(g)^{-1}$  for all  $k, g \in G$ .

**PROPOSITION 3.1.** *The group  $N(G)$  is abelian and generated as a group by the set  $\{\beta_k(g): g, k \in G \setminus \{1\}\}$ .*

**PROOF.** We obtain from Equation (2) that

$$\varphi(k_2)^{-1}\beta_{k_1}(g)\varphi(k_2) = \beta_{k_1k_2}(g)\beta_{k_2}(g)^{-1}.$$

Since  $\text{Pol}_2(G)$  is generated by  $\varphi(G)$ , we conclude that the group  $\langle \beta_k(g): g, k \in G \setminus \{1\} \rangle$  is normal in  $G$ . Now, since the relations of the form  $\beta_k(g)$  define the quotient  $G = \text{Pol}_2(G)/N(G)$ , we obtain  $N(G) = \langle \beta_k(g): g, k \in G \setminus \{1\} \rangle$  as claimed. By Proposition 2.4,  $N(G)$  is abelian.  $\square$

Let's try to analyze the extension

$$1 \rightarrow N(G) \rightarrow \text{Pol}_2(G) \rightarrow G \rightarrow 1$$

more closely. The study of extensions with abelian kernel is a classical topic, see, e.g., [2, p.90] for background.

First observe, that there is a homomorphism  $\mu: G \rightarrow \text{Aut}(N(G))$ , given by the formula  $\mu_g(h) = \varphi(g)h\varphi(g)^{-1}$  for  $h \in N(G), g \in G$ . Indeed, this is a homomorphism since  $\varphi(g)^{-1}\varphi(gh)\varphi(h)^{-1} = \beta_g(h) \in N(G)$  and  $N(G)$  is abelian.

Now, if we write the cocycle relation in additive notation, we obtain:

$$(4) \quad \beta_{k_1 k_2}(g) = \mu_{k_2}^{-1}(\beta_{k_1}(g)) + \beta_{k_2}(g).$$

or equivalently

$$\mu_{k_2}^{-1}(\beta_{k_1}(g)) = \beta_{k_1 k_2}(g) - \beta_{k_2}(g).$$

There is also a 2-cocycle  $\psi: G \times G \rightarrow N(G)$  with respect to the action just defined. The formula for  $\psi$  is given by

$$\psi(g, h) = -\mu_g(\beta_g(h)) = \varphi(g)\varphi(h)\varphi(gh)^{-1}.$$

Recall, that in formulas being a 2-cocycle means:

$$\mu_{g_1}(\psi(g_2, g_3)) - \psi(g_1 g_2, g_3) + \psi(g_1, g_2 g_3) - \psi(g_1, g_2) = 0$$

for all  $g_1, g_2, g_3 \in G$ , which is easily checked. Indeed, for sake of completeness, we compute:

$$\begin{aligned} & \mu_{g_1}(\psi(g_2, g_3)) - \psi(g_1 g_2, g_3) + \psi(g_1, g_2 g_3) - \psi(g_1, g_2) \\ &= -\mu_{g_1}(\mu_{g_2}(\beta_{g_2}(g_3))) + \mu_{g_1 g_2}(\beta_{g_1 g_2}(g_3)) - \mu_{g_1}(\beta_{g_1}(g_2 g_3)) + \mu_{g_1}(\beta_{g_1}(g_2)) \\ &= \mu_{g_1}(-\mu_{g_2}(\beta_{g_2}(g_3)) + \mu_{g_2}(\beta_{g_1 g_2}(g_3)) - \beta_{g_1}(g_2 g_3) + \beta_{g_1}(g_2)) \\ &= \mu_{g_1}(-\mu_{g_2}(\beta_{g_2}(g_3)) + \mu_{g_2}(\beta_{g_1 g_2}(g_3)) - \beta_{g_1}(g_3)) \\ &= \mu_{g_1 g_2}(-\beta_{g_2}(g_3) + \beta_{g_1 g_2}(g_3) - \mu_{g_2}^{-1}(\beta_{g_1}(g_3))) \stackrel{(4)}{=} 0. \end{aligned}$$

The extension would be described completely as

$$\text{Pol}_2(G) = N(G) \rtimes_{\mu, \psi} G,$$

after determining the  $\mathbb{Z}[G]$ -module  $N(G)$  and understanding the cohomology class  $[\psi] \in H^2(G, N(G))$  more explicitly. Recall that the multiplication in the twisted crossed product is given by

$$(\xi_1, g_1)(\xi_2, g_2) = (\xi_1 + \mu_{g_1}(\xi_2) + \psi(g_1, g_2), g_1 g_2).$$

We denote by  $\omega(G)$  the augmentation ideal of  $\mathbb{Z}[G]$ , i.e.,  $\omega(G) := \{a \in \mathbb{Z}[G], \varepsilon(a) = 0\}$  with  $\varepsilon(\sum_g a_g g) := \sum_g a_g$ . There is a natural free basis for  $\omega(G)$  consisting of the set  $\{g - 1 \mid g \in G, g \neq 1\}$ . We set  $c(g) := g - 1$  and



note that  $gc(h) = gh - g = c(gh) - c(g)$ . This is the tautological 1-cocycle  $c: G \rightarrow \omega(G)$ .

The key to understand  $N(G)$  is to note that there exists a  $G$ -equivariant surjection  $\alpha: \omega(G) \otimes_{\mathbb{Z}} G^{ab} \rightarrow N(G)$  given by the formula  $\alpha(c(h) \otimes \bar{g}) = \beta_{h^{-1}}(g)$ , where  $\bar{g}$  denotes the image of  $g \in G$  in  $G^{ab}$ . The  $G$ -equivariance is checked by the following computation:

$$\begin{aligned} \alpha(g_1(c(h) \otimes \bar{g})) &= \alpha(c(g_1h) \otimes \bar{g}) - \alpha(c(g_1) \otimes \bar{g}) \\ &= \beta_{h^{-1}g_1^{-1}}(g) - \beta_{g_1^{-1}}(g) \\ &= \mu_{g_1}(\beta_{h^{-1}}(g)) \\ &= \mu_{g_1}(\alpha(c(h) \otimes \bar{g})). \end{aligned}$$

We claim that  $\alpha$  is an isomorphism. In order to show this, we would like to identify a 2-cocycle  $\psi': G \times G \rightarrow \omega(G) \otimes_{\mathbb{Z}} G^{ab}$  compatible with  $\alpha$  and  $\psi$ , so that the natural map

$$(\omega(G) \otimes_{\mathbb{Z}} G^{ab}) \rtimes_{\psi'} G \rightarrow \text{Pol}_2(G)$$

is an isomorphism. This then also gives the desired computation of  $\text{Pol}_2(G)$ .

First of all, we need a good formula for the 2-cocycle  $\psi': G \times G \rightarrow \omega(G) \otimes_{\mathbb{Z}} G^{ab}$ . Since we have  $\psi(g, h) = \varphi(g)\varphi(h)\varphi(gh)^{-1} = \mu_g(\beta_g(h))^{-1}$  a natural guess for a 2-cocycle  $\psi': G \times G \rightarrow \omega(G) \otimes_{\mathbb{Z}} G^{ab}$  would be

$$\psi'(g, h) = -g(c(g^{-1}) \otimes \bar{h}) = c(g) \otimes \bar{h}.$$

This is indeed a 2-cocycle, namely the exterior product of the tautological 1-cocycle  $c: G \rightarrow \omega(G)$  and the trivial 1-cocycle  $\text{ab}: G \rightarrow G^{ab}$ , where  $G$  acts trivially on the abelian group  $G^{ab}$ . Moreover, if we compose the 2-cocycle  $\psi': G \times G \rightarrow \omega(G) \otimes_{\mathbb{Z}} G^{ab}$  with the surjective  $G$ -module homomorphism  $\alpha: \omega(G) \otimes_{\mathbb{Z}} G^{ab} \rightarrow N(G)$ , then we obtain  $\psi$  by construction.

Thus, we may consider the twisted crossed product group  $(\omega(G) \otimes_{\mathbb{Z}} G^{ab}) \rtimes_{\psi'} G$  with its natural multiplication

$$(\xi_1, g_1)(\xi_2, g_2) := (\xi_1 + g_1(\xi_2) + \psi'(g_1, g_2), g_1g_2).$$

We obtain a natural surjection

$$\alpha': (\omega(G) \otimes_{\mathbb{Z}} G^{ab}) \rtimes_{\psi'} G \rightarrow \text{Pol}_2(G)$$

given by  $\alpha'((c(h) \otimes \bar{g}_1), g_2) = \beta_{h^{-1}}(g_1)\varphi(g_2)$ .

In order to find the inverse of  $\alpha'$ , we use the universal property of  $\text{Pol}_2(G)$  and just construct a quadratic map

$$\varphi': G \rightarrow (\omega(G) \otimes_{\mathbb{Z}} G^{ab}) \rtimes_{\psi'} G.$$

Our philosophy suggests that it must be  $\varphi'(g) := (0, g)$  in the notation above. Indeed, if  $\varphi'$  is quadratic, it is easy to see that its classifying map is

the inverse of  $\alpha$ . The inverse of  $\varphi'(g)$  is naturally computed as  $\varphi'(g)^{-1} = (-\psi'(g^{-1}, g), g^{-1})$ .

Now, for any  $g \in G$ , we obtain

$$\begin{aligned}
\beta'_g(h) &= \varphi'(g)^{-1} \varphi'(gh) \varphi'(h)^{-1} \\
&= (-\psi'(g^{-1}, g), g^{-1})(0, gh)(-\psi'(h^{-1}, h), h^{-1}) \\
&= (-\psi'(g^{-1}, g) + \psi'(g^{-1}, gh), h)(-\psi'(h^{-1}, h), h^{-1}) \\
&= (-\psi'(g^{-1}, g) + \psi'(g^{-1}, gh) - h\psi'(h^{-1}, h) + \psi(h, h^{-1}), 1) \\
&= -c(g^{-1}) \otimes \bar{g} + c(g^{-1}) \otimes (\bar{g} + \bar{h}) - hc(h^{-1}) \otimes \bar{h} - c(h) \otimes h \\
&= c(g^{-1}) \otimes \bar{h}.
\end{aligned}$$

This is clearly a homomorphism in  $h$ . Thus,  $\varphi'$  is quadratic and the proof of Theorem 1.3 is complete.  $\square$

Having Theorem 1.3 at hand it is not difficult to describe its abelianization in concrete terms.

**PROOF OF COROLLARY 1.4.** Let  $G$  be an abelian group. We need to compute the abelianization of the group  $(\omega(G) \otimes_{\mathbb{Z}} G) \rtimes_{\psi} G$ , where  $\psi: G \times G \rightarrow \omega(G) \otimes_{\mathbb{Z}} G^{ab}$  is given by  $\psi(g, h) = c(g) \otimes \bar{h}$ . We have

$$(\xi_1, g_1)(\xi_2, g_2) := (\xi_1 + g_1(\xi_2) + \psi'(g_1, g_2), g_1 g_2)$$

for all  $\xi_1, \xi_2 \in \omega(G) \otimes_{\mathbb{Z}} G$  and  $g_1, g_2 \in G$ . Now, clearly  $[(0, g), (\xi, 1)] = (g\xi - \xi, 1)$ , so that  $\omega(G)^2 \otimes_{\mathbb{Z}} G$  lies in the commutator subgroup. Here  $\omega(G)^2$  denotes the square of the ideal  $\omega(G) \subseteq \mathbb{Z}[G]$ . However,  $\omega(G)/\omega(G)^2 = G$ , so that the map to the abelianization factorizes via  $(G \otimes_{\mathbb{Z}} G) \rtimes_{\psi''} G$  with the two-cocycle  $\psi''(g, h) = g \otimes h$ . This group is not yet abelian, but

$$\begin{aligned}
[(0, g), (0, h)] &= (0, g)(0, h)(\psi''(g, g), -g)(\psi''(h, h), -h) \\
&= (\psi''(g, h), g + h)(\psi''(g, g) + \psi(h, h) + \psi''(g, h), -g - h) \\
&= (\psi''(g, h) - \psi''(h, g), 0).
\end{aligned}$$

Hence, the abelianization factorizes via  $\text{Sym}^2(G) \rtimes_{\sigma} G$  with  $\sigma: G \times G \rightarrow \text{Sym}^2(G)$  given by  $\sigma(g, h) = g \cdot h$ . This group is easily checked to be abelian and hence isomorphic to the abelianization of  $\text{Pol}_2(G)$ . This finishes the proof.  $\square$

#### 4. Some computations and consequences

This section contains various concrete computations of  $\text{Pol}_2(G)$  and its abelianization. The following corollary is already interesting for  $d = 1$ .

**COROLLARY 4.1.** *Let  $\mathbb{F}_d$  be the free group on  $d$  generators. The group  $\text{Pol}_2(\mathbb{F}_d)$  is isomorphic to  $(\mathbb{Z}[\mathbb{F}_d]^{\oplus d^2}) \rtimes \mathbb{F}_d$ . Moreover, the abelianization is isomorphic to  $\mathbb{Z}^{d^2+d}$ .*

**PROOF.** It is well-known that  $\omega(\mathbb{F}_d) = \mathbb{Z}[\mathbb{F}_d]^{\oplus d}$  and hence  $\omega(\mathbb{F}_d) \otimes_{\mathbb{Z}} \mathbb{Z}^d = \mathbb{Z}[\mathbb{F}_d]^{\oplus d^2}$  as  $\mathbb{F}_d$ -modules. Moreover, the cohomological dimension of  $\mathbb{F}_d$  is less than two, i.e., all cohomology in dimension 2 and above vanishes. Both of these facts are consequences of the existence of a free resolution

$$0 \rightarrow \mathbb{Z}[\mathbb{F}_d]^{\oplus d} \rightarrow \mathbb{Z}[\mathbb{F}_d] \xrightarrow{\varepsilon} \mathbb{Z} \rightarrow 0,$$

see, e.g., [2, p.16]. Consequently, the class of the cocycle

$$[\psi] \in \text{H}^2(\mathbb{F}_d, \omega(\mathbb{F}_d) \otimes_{\mathbb{Z}} \mathbb{F}_d^{ab})$$

is trivial and the extension from Theorem 1.3 is split. This implies the claim.  $\square$

**REMARK 4.2.** By Corollary 4.1,  $\text{Pol}_2(\mathbb{Z})$  is isomorphic to the lamplighter group  $\mathbb{Z} \wr \mathbb{Z}$ . It is well-known that this group is not nilpotent but residually nilpotent. As a consequence, for every  $n \in \mathbb{N}$ , there is a quadratic map  $\varphi_n: \mathbb{Z} \rightarrow G_n$  with  $G_n$  nilpotent of class  $n$  and such that the image of  $\varphi_n$  generates  $G_n$ .

**EXAMPLE 4.3.** The abelianization of the group  $\text{Pol}_2(\mathbb{F}_d)$  is  $\mathbb{Z}^{d^2+d}$  by Corollary 4.1, whereas the abelianization of  $\text{Pol}_2(\mathbb{Z}^d)$  is  $\mathbb{Z}^{(d^2+3d)/2}$  by Corollary 1.4. The numbers in the exponent differ for  $d \geq 2$ . This implies that there are exotic unital quadratic  $\mathbb{Z}$ -valued maps on  $\mathbb{F}_2$  that do not factor through the abelianization  $\mathbb{F}_2 \rightarrow \mathbb{Z}^2$ . One such map arises from the canonical homomorphism  $\mathbb{F}_2$  to  $\text{H}_3(\mathbb{Z}) = \mathbb{F}_2/[\mathbb{F}_2, [\mathbb{F}_2, \mathbb{F}_2]]$  composed with the map  $\text{H}_3(\mathbb{Z}) \rightarrow \mathbb{Z}$  that sends the  $3 \times 3$ -matrix to the entry in the upper right corner. More concretely, this map is given by the formula

$$\varphi(a^{n_1} b^{m_1} \dots a^{n_k} b^{m_k}) := \sum_{i < j} m_i n_j,$$

i.e., it counts the number of  $b$ 's on the left of  $a$ 's. To put this into perspective, Leibman [12] showed that every unital quadratic map on any group with values in  $\mathbb{Z}$  factors through a nilpotent quotient.

**PROOF OF COROLLARY 1.5:** By Theorem 1.3, we have

$$\ker(\pi: \text{Pol}_2(G) \rightarrow G) = \omega(G) \otimes_{\mathbb{Z}} G^{ab}.$$

Hence,  $\pi$  must be an isomorphism if  $G$  is perfect. Thus, every unital quadratic map is a homomorphism. The general claim follows then by the inductive nature of the definition of the degree of a polynomial.  $\square$

Let's record another corollary of Theorem 1.3.

COROLLARY 4.4. *If  $G$  is finite, then  $\text{Pol}_2(G)$  is finite.*

PROOF. Indeed,  $G^{ab}$  is finite, so that  $\omega(G) \otimes_{\mathbb{Z}} G^{ab}$  is finite as well. It follows that  $\text{Pol}_2(G)$  is a finite group.  $\square$

REMARK 4.5. While it is relatively easy to compute the abelianization of  $\text{Pol}_2(\mathbb{Z}/n\mathbb{Z})$  from Corollary 1.4, it is more challenging to pin down  $\text{Pol}_2(\mathbb{Z}/n\mathbb{Z})$ . Clearly, Theorem 1.3 implies that  $\text{Pol}_2(\mathbb{Z}/n\mathbb{Z})$  is a metabelian group of order  $n^n$ . We have  $\text{Pol}_2(\mathbb{Z}/2\mathbb{Z}) = \mathbb{Z}/4\mathbb{Z}$ . One can compute  $\text{Pol}_2(\mathbb{Z}/3\mathbb{Z})$  to be the Heisenberg group over the ring  $\mathbb{Z}/3\mathbb{Z}$ .

We now turn to some non-trivial applications for  $G$  abelian and for abelian targets. Of particular interest is the case  $H = S^1$ . Note that if  $G$  is finite then the group  $\text{Pol}_2^{ab}(G)$  is isomorphic to  $\text{Quad}(G, S^1) = \text{hom}(\text{Pol}_2(G), S^1)$  by Pontryagin duality and Corollary 4.4.

PROPOSITION 4.6. *If  $G$  is an abelian group of bounded exponent and  $G = \bigoplus_p G_p$  is its primary decomposition, then*

$$\text{Pol}_2^{ab}(G) = \bigoplus_p \text{Pol}_2^{ab}(G_p).$$

PROOF. This is a basic consequence of the the isomorphism  $\text{Sym}^2(G) = \bigoplus_p \text{Sym}^2(G_p)$  in this case.  $\square$

PROPOSITION 4.7. *If  $G$  is a 2-divisible finite abelian group, then the extension in Corollary 1.4 is split.*

PROOF. Note that if  $G$  is finite abelian and divisible by 2, then  $\nu: G \rightarrow \text{Sym}^2(G)$  given by  $\nu(g) = g^2/2$  satisfies  $\sigma(g, h) = \nu(g + h) - \nu(g) - \nu(h)$ . In particular, the 2-cocycle  $\sigma$  is a coboundary and the extension in Corollary 1.4 is split.  $\square$

COROLLARY 4.8. *If  $G$  is a finite abelian and 2-divisible group, then  $\text{Pol}_2^{ab}(G) = \text{Sym}^2(G) \oplus G$ . This applies to all finite abelian groups without 2-primary part.*

REMARK 4.9. The case of finite 2-groups is more subtle. As mentioned before, it is well-known that  $\text{Pol}_2^{ab}(\mathbb{Z}/2\mathbb{Z}) = \mathbb{Z}/4\mathbb{Z}$ . Moreover, we obtain for example that

$$\text{Pol}_2^{ab}(\mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/4\mathbb{Z}) = (\mathbb{Z}/2\mathbb{Z})^{\oplus 2} \oplus \mathbb{Z}/4\mathbb{Z} \oplus \mathbb{Z}/8\mathbb{Z}.$$

Here,  $\mathbb{Z}/4\mathbb{Z}$  corresponds to the unital quadratic maps factoring through  $\mathbb{Z}/2\mathbb{Z}$ ,  $\mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/8\mathbb{Z}$  corresponds to the unital quadratic maps factoring through  $\mathbb{Z}/4\mathbb{Z}$ , and another copy of  $\mathbb{Z}/2\mathbb{Z}$  corresponds to the unital quadratic map  $\mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/4\mathbb{Z} \ni (x, y) \mapsto xy \in \mathbb{Z}/2\mathbb{Z}$ .

The precise computation of  $\text{Pol}_2^{ab}(G)$  as well as  $\text{Pol}_k^{ab}(G)$  for  $k \geq 3$  is subject of further stu

Finally, for illustration of the result, we consider a classical infinitary situation: Let  $d \geq 1$  be an integer. For  $n = (n_1, \dots, n_d) \in \mathbb{Z}^d$  and  $j = (j_1, \dots, j_d) \in \mathbb{Z}_{\geq 0}^d$ , define the multi-binomial coefficient

$$B_j(n) := \binom{n_1}{j_1} \cdots \binom{n_d}{j_d}.$$

Let  $J$  denote the collection of  $j = (j_1, \dots, j_d) \in \mathbb{Z}_{\geq 0}^d$  such that  $1 \leq \sum_{i=1}^d j_i \leq 2$ . We use the notation  $\exp(x) := e^{2\pi i x}$  to represent the fundamental character from  $\mathbb{R}/\mathbb{Z}$  to  $S^1$ .

By the Taylor expansion formula for polynomial maps between filtered groups (see, e.g., [4, Lemma B.9], and its Appendix B for background on such polynomial maps), every quadratic  $\varphi \in \text{Quad}(\mathbb{Z}^d, S^1)$  is of the form

$$\varphi(n) = \prod_{j \in J} \exp(c_j)^{B_j(n)}$$

for some  $c_j \in \mathbb{R}/\mathbb{Z}$ ,  $j \in J$ .

The cardinality of  $J$ , which is the number of basis polynomials  $n \mapsto \exp(c_j)^{B_j(n)}$ , is  $(d^2 + 3d)/2$ . The abelianization of  $\text{Pol}_2(\mathbb{Z}^d)$  is  $\mathbb{Z}^{(d^2+3d)/2}$ , which is the Pontryagin dual of  $(\mathbb{R}/\mathbb{Z})^{(d^2+3d)/2}$ .

## 5. Approximate polynomial maps

This is the first of two sections where we study the case of perfect groups more closely. We start with an investigation of approximate polynomial maps with values in groups that carry a bi-invariant metric. Let's recall the following definition:

**DEFINITION 5.1.** *We say that a perfect group  $G$  is of commutator width  $k$ , if each element of  $G$  is a product of at most  $k$  commutators.*

**REMARK 5.2.** Note that a finite group is automatically of finite commutator width. For interesting classes of finite perfect groups there exist uniform bounds on the commutator width. For example, as mentioned already in the introduction, Wilson [17] showed that the commutator width is uniformly bounded for *all* finite simple groups. In fact, the commutator width is now known to be exactly 1, as established by the more recent solution to the Ore problem [13]. See also [14] for even broader classes of finite perfect groups of bounded commutator width. It was shown by Holt–Plesken [8, Lemma 2.1.10], that the commutator width of a finite perfect group can become arbitrarily large. This also implies that an infinite perfect group does not need to have finite commutator width.

We will now derive a useful consequence of Corollary 1.5:

COROLLARY 5.3. *Let  $k \in \mathbb{N}$ . Let  $G$  be a perfect group of commutator width  $k$ . Consider the free group on the set  $\varphi(G)$ . There exists a constant  $c := c(k) \geq 0$ , such that for every  $g, h \in G$ , the element  $\varphi(g_1 g_2) \varphi(g_2)^{-1} \varphi(g_1)^{-1}$  is a product of  $c$  conjugates of elements of the form*

$$\varphi(g_1 g_2 g_3)^{-1} \varphi(g_1 g_2) \varphi(g_2)^{-1} \varphi(g_1)^{-1} \varphi(g_1 g_3) \varphi(g_3)^{-1} \varphi(g_2 g_3)$$

or their inverses for  $g_1, g_2, g_3 \in G$ .

PROOF. We consider subsets  $\Sigma_1 := \{\varphi(g_1 g_2) \varphi(g_2)^{-1} \varphi(g_1)^{-1} \mid g_1, g_2 \in G\}$  and  $\Sigma_2 := \{\varphi(g_1 g_2 g_3)^{-1} \varphi(g_1 g_2) \varphi(g_2)^{-1} \varphi(g_1)^{-1} \varphi(g_1 g_3) \varphi(g_3)^{-1} \varphi(g_2 g_3) \mid g_1, g_2, g_3 \in G\}$  of the free group generated by  $\varphi(G)$ . Since  $G$  is assumed to be perfect, by Corollary 1.5, both sets of relations define the same normal subgroup, i.e.,  $\langle\langle \Sigma_1 \rangle\rangle = \langle\langle \Sigma_2 \rangle\rangle$ . Thus, each word of the form  $\varphi(g_1 g_2) \varphi(g_2)^{-1} \varphi(g_1)^{-1}$  for  $g_1, g_2 \in G$  must be a product of conjugates of elements of the form  $\varphi(g_1 g_2 g_3)^{-1} \varphi(g_1 g_2) \varphi(g_2)^{-1} \varphi(g_1)^{-1} \varphi(g_1 g_3) \varphi(g_3)^{-1} \varphi(g_2 g_3)$  or their inverses. The uniform estimate  $c(k)$  on the number of factors follows from an easy ultra-product argument noting that the commutator width is preserved by the operation of taking an ultra-product of groups.  $\square$

Our focus is now on almost structure preserving maps to metric groups. Let  $(H, \partial)$  be a metric group. It is of some interest in geometry as well as algebra to consider maps which are almost homomorphisms  $\varphi: G \rightarrow H$  in the sense that

$$\partial(\varphi(gh), \varphi(g)\varphi(h)) < \varepsilon$$

uniformly for all  $g, h \in G$ . This has been studied in various situations, see for example [1, 3, 6, 7, 10]) and the references therein. Note that

$$(\Delta_h(\Delta_g \varphi))(1) = (\Delta_g \varphi)(h) (\Delta_g \varphi)(1)^{-1} = \varphi(gh) \varphi(h)^{-1} \varphi(g)^{-1}$$

so that being an almost homomorphism as above can be easily expressed in terms of a condition on iterated finite differences of  $\varphi$ . We intend to generalize this idea and study arbitrary iterated differences of  $\varphi$ . For  $g_1, \dots, g_{d+1}$ , we set  $\Delta_{g_1, \dots, g_{d+1}} := \Delta_{g_1} \circ \dots \circ \Delta_{g_{d+1}}$ .

DEFINITION 5.4. *Let  $(H, \partial)$  be a group with a bi-invariant metric. We say that a unital map  $\varphi: G \rightarrow H$  is a uniform  $\varepsilon$ -polynomial of degree  $d$  if*

$$\partial((\Delta_{g_1, \dots, g_{d+1}} \varphi)(1), 1) \leq \varepsilon$$

for all  $g_1, \dots, g_{d+1} \in G$ .

THEOREM 5.5. *Let  $k \in \mathbb{N}$ . There exists a constant  $c := c(k) > 0$  such that the following holds: Let  $d \in \mathbb{N}$  and  $(H, \partial)$  be a group with a bi-invariant metric and  $G$  a group of commutator length at most  $k$ . Every unital uniform  $\varepsilon$ -polynomial  $\varphi: G \rightarrow H$  of degree  $d$  is a uniform  $c(k)^{d-1} \varepsilon$ -homomorphism.*

PROOF. The proof proceeds by induction on  $d$ , where the case  $d = 1$  is trivial. The case  $d = 2$  requires special attention. Let's compute:

$$\begin{aligned}
 (\Delta_{g_1, g_2, g_3} \varphi)(g_0) &= (\Delta_{g_1}(\Delta_{g_2, g_3} \varphi))(g_0) \\
 &= (\Delta_{g_2, g_3} \varphi)(g_1 g_0) ((\Delta_{g_2, g_3} \varphi)(g_0))^{-1} \\
 &= \varphi(g_3 g_2 g_1 g_0) \varphi(g_2 g_1 g_0)^{-1} \varphi(g_1 g_0) \varphi(g_3 g_1 g_0)^{-1} \\
 &\quad \cdot (\varphi(g_3 g_2 g_0) \varphi(g_2 g_0)^{-1} \varphi(g_0) \varphi(g_3 g_0)^{-1})^{-1} \\
 &= \varphi(g_3 g_2 g_1 g_0) \varphi(g_2 g_1 g_0)^{-1} \varphi(g_1 g_0) \varphi(g_3 g_1 g_0)^{-1} \\
 &\quad \cdot \varphi(g_3 g_0) \varphi(g_0)^{-1} \varphi(g_2 g_0) \varphi(g_3 g_2 g_0)^{-1}
 \end{aligned}$$

Thus, for  $\varphi$  unital, we obtain:

$$(\Delta_{g_1, g_2, g_3} \varphi)(1) = \varphi(g_3 g_2 g_1) \varphi(g_2 g_1)^{-1} \varphi(g_1) \varphi(g_3 g_1)^{-1} \varphi(g_3) \varphi(g_2) \varphi(g_3 g_2)^{-1}.$$

By our assumption, we obtain

$$\partial(\varphi(g_3 g_2 g_1) \varphi(g_2 g_1)^{-1} \varphi(g_1) \varphi(g_3 g_1)^{-1} \varphi(g_3) \varphi(g_2) \varphi(g_3 g_2)^{-1}, 1) \leq \varepsilon$$

for all  $g_1, g_2, g_3 \in G$ . Note that the same estimate holds for conjugates and inverses by bi-invariance of the metric. Thus, Corollary 5.3 implies that  $\partial(\varphi(g_1 g_2) \varphi(g_2)^{-1} \varphi(g_1)^{-1}, 1) \leq c(k)\varepsilon$  and equivalently

$$\partial((\Delta_{g_1, g_2} \varphi)(1), 1) \leq c(k)\varepsilon.$$

As required, we conclude that  $\varphi$  is a uniform  $c(k)\varepsilon$ -polynomial of degree 1. The general case follows by induction. Indeed, note that for  $d \geq 2$ ,  $\varphi$  is a uniform  $\varepsilon$ -polynomial of degree  $d + 1$  if and only if  $\Delta_g \varphi$  is a uniform  $\varepsilon$ -polynomial of degree  $d$  for all  $g \in G$ .  $\square$

Note that the previous result is particularly useful when a stability result is present for uniform  $\varepsilon$ -homomorphisms, which is the case in various situations, see the references above. In those cases, every  $\varepsilon$ -polynomial for  $\varepsilon > 0$  small enough is uniformly close to a homomorphism.

A typical example where this happens is the group  $U(n)$  with the 2-norm and  $G$  amenable, see [3]. Here, for  $a \in M_n \mathbb{C}$  we denote by  $\|a\|_2 = ((1/n) \sum_{ij} |a_{ij}|^2)^{1/2}$  the 2-norm or normalized Hilbert-Schmidt norm. Thus, we obtain:

**THEOREM 5.6.** *Let  $d, k \geq 0$ . For all  $\varepsilon > 0$ , there exists  $\delta := \delta(d, k) > 0$ , such that the following holds: Let  $G$  be an amenable group of commutator width  $k$  and let  $\varphi: G \rightarrow U(n)$  be a unital uniform  $\delta$ -polynomial of degree  $d$ . Then there exists  $n' \in [n, (1 + \varepsilon)n] \cap \mathbb{N}$  and a homomorphism  $\beta: G \rightarrow U(n')$ , such that  $\sup_g \|(\varphi(g) \oplus 1_{n'-n}) - \beta(g)\|_2 < \varepsilon$ .*

**REMARK 5.7.** The previous corollary and theorem applies uniformly to all finite simple groups and fixed  $d \in \mathbb{N}$ . An example of an infinite perfect

amenable group with bounded commutator width is the group  $\text{Alt}_{\text{fin}}(\mathbb{N})$  consisting of all alternating permutations of  $\mathbb{N}$  with finite support.

REMARK 5.8. We did not succeed to derive an explicit bound for  $c(k)$ , even in the case  $k = 1$ .

## 6. The Gowers norms and an inverse theorem

Fourier analysis studies how general functions can be decomposed into characters, which, in the formalism of this paper, correspond to unital linear maps for abelian groups with values in the torus and for non-abelian groups with values in unitary groups. In the context of Szemerédi's theorem in additive combinatorics, a more general version of Fourier analysis has proven necessary, one that studies more subtle forms of correlation with polynomial patterns [5]. This broader framework is now known as higher-order Fourier analysis.

Higher-order Fourier analysis naturally introduces a hierarchy of norms, known as the Gowers norms, on spaces of functions defined on groups. This hierarchy is related to the degree of polynomials. Typically, the studied groups are abelian (or at least nilpotent), and the functions are scalar-valued; see [4, 9, 15, 16] and the references therein. Gowers and Hatami [6] extended the definition of the second Gowers norm to functions on arbitrary finite groups with values in matrices. Their work established a connection between the study of unitary-valued functions with large Gowers norms and stability questions for almost representations; see also [3].

In this section we define the Gowers norms of arbitrary degree for functions on a finite group with values in  $M_n\mathbb{C}$ . We follow the inspiration taken from the work of Gowers–Hatami [6]. We set for a matrix valued function  $\varphi: G \rightarrow M_n\mathbb{C}$

$$(\Delta_g\varphi)(h) := \varphi(gh)\varphi(h)^*.$$

Note that if the function takes values in the group of unitaries, then this is just the old finite difference. Recall our definition  $\Delta_{g_1, \dots, g_k} := \Delta_{g_1} \circ \dots \circ \Delta_{g_k}$  and set

$$\|\varphi\|_{U^k}^{2k} := \mathbb{E}_{g_0, g_1, \dots, g_k} \text{tr}((\Delta_{g_1, \dots, g_k}\varphi)(g_0)),$$

where  $\text{tr}: M_n\mathbb{C} \rightarrow \mathbb{C}$  denotes the *normalized* trace and  $\mathbb{E}$  denotes the expectation with respect to the normalized counting measure on  $G^{k+1}$ .

For  $k = 2$ , we compute

$$\begin{aligned} (\Delta_{g_1, g_2}\varphi)(g_0) &= (\Delta_{g_2}\varphi)(g_1g_0)(\Delta_{g_1}\varphi)(g_0)^* \\ &= \varphi(g_2g_1g_0)\varphi(g_1g_0)^*\varphi(g_0)\varphi(g_2g_0)^* \end{aligned}$$



so that

$$\mathbb{E}_{g_0, g_1, g_2} \operatorname{tr}((\Delta_{g_1, g_2} \varphi)(g_0)) = \mathbb{E}_{xy^{-1}zw^{-1}=1} \operatorname{tr}(\varphi(x)\varphi(y)^* \varphi(z)\varphi(w)^*).$$

We see from this equality that our definition agrees with the  $U^2$ -Gowers norm as defined in [6], modulo the normalization of the trace. The assignment  $\varphi \mapsto \|\varphi\|_{U^k}$  is called  $k$ -th Gowers norm.

Clearly, any polynomial  $\varphi: G \rightarrow \mathbb{U}(n)$  of degree  $k-1$  has a  $k$ -th Gowers norm equal to 1 and it is interesting to study the question whether functions with unitary values and  $k$ -th Gowers norm near 1 must be perturbations of polynomial maps in a suitable sense. Such a result is called an inverse theorem in the 99% regime. Our main result is the following strong form of an inverse theorem in the 99% regime that comes along with a collapse of the higher-order Fourier hierarchy. Such results can only be expected under strong additional hypothesis and we prove it assuming that the finite group  $G$  is perfect.

For a function  $\varphi: G \rightarrow M_n \mathbb{C}$ , we set  $\|\varphi\|_\infty := \sup_{g \in G} \|\varphi(g)\|$ , where  $\|a\|$  for  $a \in M_n \mathbb{C}$  denotes the usual operator norm of a matrix.

**THEOREM 6.1.** *Let  $k_0, k \in \mathbb{N}$ . For all  $\delta > 0$  there exists  $\varepsilon = \varepsilon(\delta, k, k_0) > 0$ , such that the following holds: Let  $G$  be a finite perfect group of commutator width  $k_0$  and let  $\varphi: G \rightarrow M_n \mathbb{C}$  be map. If  $\|\varphi\|_\infty \leq 1$  and  $\|\varphi\|_{U^k}^{2k} \geq 1 - \varepsilon$ , then there exists a homomorphism  $\beta: G \rightarrow \mathbb{U}(n')$  with  $n' \in [n, (1 + \delta)n] \cap \mathbb{N}$  such that*

$$\mathbb{E}_g \|(\varphi(1)^* \varphi(g) \oplus 1_{n'-n}) - \beta(g)\|_2^2 < \delta.$$

First of all, let us note that we may assume without loss of generality that the function  $\varphi$  takes unitary values and  $\varphi(1) = 1_n$ . Indeed, this is a consequence of the following two lemmas.

**LEMMA 6.2.** *Let  $k \geq 1$  be a natural number. For all  $\varepsilon > 0$  there exists  $\delta > 0$  such that if  $v_1, \dots, v_k \in M_n \mathbb{C}$  are contractions with*

$$\operatorname{tr}(v_1 \cdots v_k) \geq 1 - \delta,$$

*then there exists unitaries  $u_1, \dots, u_k \in \mathbb{U}(n)$  such that  $\|v_i - u_i\|_2 < \varepsilon$  for all  $1 \leq i \leq k$  and  $u_1 \cdots u_k = 1_n$ .*

**PROOF.** Suppose by contradiction that for some  $\varepsilon > 0$ , there exists sequences of contractions  $(v_{i,m})_m$  for  $1 \leq i \leq k$  with  $v_{i,n} \in M_{n_m} \mathbb{C}$  such that

$$\operatorname{tr}(v_{1,m} \cdots v_{k,m}) \geq 1 - \frac{1}{m}$$

and at the same time, there do not exist unitaries  $u_1, \dots, u_k$  which are  $\varepsilon$ -close to  $v_1, \dots, v_k$  in the 2-norm.

Consider the tracial ultraproduct of von Neumann algebras.

$$(M, \tau) := \prod_{m \rightarrow \mathcal{U}} (M_{n_m} \mathbb{C}, \text{tr}).$$

It is well-known that  $(M, \tau)$  is a tracial von Neumann algebra, in fact it is a factor of type  $\text{II}_1$ . Consider the contractions  $v_i \in M$  represented by the sequences  $(v_{i,m})_m$ . From our assumption, we obtain

$$\tau(v_1 \cdots v_k) = 1.$$

Now, let  $x \in M$  be an arbitrary contraction with  $\tau(x) = 1$ . Since,  $\|1 - x\|_2^2 = 1 + \tau(x^*x) - \tau(x) - \tau(x^*) \leq 0$  we obtain  $x = 1$ . In particular,  $v_1$  is a contraction whose inverse is also a contraction, being equal to  $v_2 \cdots v_k$ . This implies that  $v_1$  is isometric and hence unitary. By induction, we conclude that  $v_1, \dots, v_k$  are all unitaries. We conclude that  $v_i$  can also be represented by a sequence of unitaries  $(u_{i,m})_m$  with

$$\lim_{m \rightarrow \mathcal{U}} \|u_{i,m} - v_{i,m}\|_2 = 0 \quad \text{for } 1 \leq i \leq k-1.$$

We put  $u_{i,k} := (u_{i,1} \cdots u_{i,k-1})^{-1}$ . This is a contradiction and thus, proves the claim.  $\square$

LEMMA 6.3. *Let  $k \in \mathbb{N}$  and  $\varepsilon > 0$ . There exists  $\delta := \delta(k, \varepsilon) > 0$ , such that the following is true: Let  $G$  be a finite group and let  $\varphi: G \rightarrow M_n \mathbb{C}$  with*

$$\|\varphi\|_\infty \leq 1 \quad \text{and} \quad \|\varphi\|_{U^k}^{2^k} > 1 - \delta.$$

*There exists a function  $\varphi': G \rightarrow U(n)$  such that*

$$\mathbb{E}_g \|\varphi(g) - \varphi'(g)\|_2^2 < \varepsilon \quad \text{and} \quad \|\varphi'\|_{U^k}^{2^k} > 1 - \varepsilon.$$

PROOF. We have

$$\|\varphi\|_{U^k}^{2^k} = \mathbb{E}_{g_0, g_2, g_3, \dots, g_k} \text{tr}((\Delta_{g_1, \dots, g_k} \varphi)(g_0)) > 1 - \delta$$

while  $(\Delta_{g_1, \dots, g_k} \varphi)(g_0)$  is a product of  $2^k$  contractions from the set  $\varphi(G)$ . The lower bound implies that the set  $\Sigma \subset G$  of those  $g \in G$  for which  $\varphi(g)$  appears in such a product whose trace is at least  $1 - \delta^{1/2}$  is of density at least  $1 - \delta^{1/2}$ . By Lemma 6.2, provided  $\delta > 0$  is small enough, any such  $\varphi(g)$  can be replaced by a unitary  $\varphi'(g)$  such that  $\|\varphi(g) - \varphi'(g)\|_2 < \delta'$ . For all  $g \notin \Sigma$ , we define  $\varphi'(g) = 1_n$ .

Now, note that the set of  $(k+1)$ -tuples  $g_0, \dots, g_k$  such that all contractions arising in factors of  $(\Delta_{g_1, \dots, g_k} \varphi)(g_0)$  lie in  $\Sigma$  is of density at least  $1 - 2^k \delta^{1/2}$ . For those  $(k+1)$ -tuples, we obtain

$$\|(\Delta_{g_1, \dots, g_k} \varphi)(g_0) - (\Delta_{g_1, \dots, g_k} \varphi')(g_0)\|_2 \leq 2^k \delta'$$

and hence

$$|\text{tr}((\Delta_{g_1, \dots, g_k} \varphi)(g_0) - (\Delta_{g_1, \dots, g_k} \varphi')(g_0))| \leq 2^{k+1} \delta'.$$

Finally, we conclude

$$\left| \|\varphi\|_{U^k}^{2^k} - \|\varphi'\|_{U^k}^{2^k} \right| < 2^{k+1}\delta' + 2^{k+1}\delta^{1/2}.$$

Now,  $\delta' > 0$  is as small as we wish, when  $\delta$  is small. And  $\delta > 0$  can be taken small enough so that also the second summand does not contribute. This proves the claim.  $\square$

Before we start with the actual proof of Theorem 6.1, let us recall that the case  $k = 2$  is a consequence of [6, Theorem 5.6]. Indeed, assuming that  $\varphi$  takes values in  $U(n)$ , we compute

$$\|(\varphi(g) \oplus 1_{n'-n}) - \beta(g)\|_2^2 = 2 - 2\Re\langle \varphi(g) \oplus 1_{n'-n}, \beta(g) \rangle,$$

so that large correlation of unitary-valued maps is directly tied to a small 2-norm of the difference. In view of Corollary 1.5, all unital polynomial maps are automatically homomorphisms, so that we can view Theorem 6.1 as a generalization of [6, Theorem 5.6] from the case  $k = 2$  to higher degrees.

**PROOF OF THEOREM 6.1.** We may assume that  $\varphi$  takes values in unitaries and upon replacing  $\varphi$  by  $g \mapsto \varphi(1)^*\varphi(g)$ , we may assume that  $\varphi$  is unital.

We prove the claim by induction on  $k$ , where the case  $k = 2$  was proved in [6] as noted above. So let  $k \geq 3$ ,  $\delta > 0$  and  $\varphi: G \rightarrow U(n)$  be arbitrary and assume that  $\|\varphi\|_{U^k}^{2^k} \geq 1 - \varepsilon$  for some  $\varepsilon > 0$  to be determined later. Now, the inequality just says that

$$\mathbb{E}_{g_0, g_2, g_3, \dots, g_k} \operatorname{tr}((\Delta_{g_1, \dots, g_k} \varphi)(g_0)) = \|\varphi\|_{U^k}^{2^k} \geq 1 - \varepsilon.$$

Note that  $\|\varphi\|_\infty \leq 1$  and we conclude by Markov's inequality, that the set  $A \subseteq G$  of those  $g \in G$ , such that

$$\mathbb{E}_{g_0, g_2, g_3, \dots, g_{k-1}} \operatorname{tr}((\Delta_{g_1, \dots, g_{k-1}, g} \varphi)(g_0)) = \|\Delta_g \varphi\|_{U^{k-1}}^{2^{k-1}} \geq 1 - \varepsilon^{1/2}$$

has cardinality at least  $(1 - \varepsilon^{1/2})|G|$ .

Let  $\eta > 0$  to be determined later. For  $\varepsilon > 0$  such that  $\varepsilon^{1/2} \leq \varepsilon(\eta, k - 1, k_0)$ , by induction, we find  $n' \in [n, (1 + \eta)n] \cap \mathbb{N}$ , homomorphisms  $\beta_g: G \rightarrow U(n')$  such that

$$\mathbb{E}_h \|(\varphi(g)^*(\Delta_g \varphi)(h) \oplus 1_{n'-n}) - \beta_g(h)\|_2^2 < \eta$$

for all  $g \in A$ .

For  $g \in A$ , again by Markov's inequality, there exists a subset  $B(g) \subseteq G$  of density at least  $1 - \eta^{1/2}$  such that  $\|(\varphi(g)^*(\Delta_g \varphi)(h) \oplus 1_{n'-n}) - \beta_g(h)\|_2 < \eta^{1/4}$  for all  $h \in B(g)$ .

Let us consider  $g \in A \cap A^{-1}$ . For ease of notation, we will write  $u \stackrel{\nu}{=} v$  if  $\|u - v\|_2 < \nu$ . Now, by the proof of Equation (2), we obtain

$$1_{n'} \stackrel{2\eta^{1/4}}{=} \varphi(g)^{-1} \beta_{g^{-1}}(h) \varphi(g) \cdot \beta_g(h)$$

for all  $h \in g^{-1}B(g^{-1}) \cap B(g)$ . Thus, if  $h_1, h_2, h_2h_1 \in g^{-1}B(g^{-1}) \cap B(g)$ , we obtain

$$\begin{aligned} \beta_g(h_1h_2) &= \beta_g(h_1)\beta_g(h_2) \\ &\stackrel{4\eta^{1/4}}{=} (\varphi(g)^{-1}\beta_{g^{-1}}(h_1)\varphi(g))^{-1}(\varphi(g)^{-1}\beta_{g^{-1}}(h_2)\varphi(g))^{-1} \\ &= (\varphi(g)^{-1}\beta_{g^{-1}}(h_2)\beta_{g^{-1}}(h_1)\varphi(g))^{-1} \\ &= (\varphi(g)^{-1}\beta_{g^{-1}}(h_2h_1)\varphi(g))^{-1} \\ &\stackrel{2\eta^{1/4}}{=} \beta_g(h_2h_1). \end{aligned}$$

We conclude from this computation that the set  $C(g)$  of pairs  $(h_1, h_2) \in G \times G$  with

$$\beta_g([h_1, h_2]) \stackrel{6\eta^{1/4}}{=} 1_{n'}$$

has density at least  $1 - 6\eta^{1/2}$ . If  $1 - 6\eta^{1/2} > 1/2$ , then  $C(g)^2 = G \times G$ . Indeed, we have  $C(g)^{-1} = C(g)$  and under this condition on the intersection  $(h_1, h_2)C(g)^{-1} \cap C(g)$  must be non-empty for each  $(h_1, h_2) \in G \times G$ . Note also that commutators satisfy the following universal identities:

$$[ab, cd] = [ab, c] \cdot [ab, d]^c = [a, c]^b [b, c] [a, d]^{bc} [b, d]^c$$

for all  $a, b, c, d \in G$ . Thus, we obtain from  $C(g)^2 = G \times G$  that  $\beta_g([h_1, h_2]) \stackrel{24\eta^{1/4}}{=} 1_{n'}$  for all  $h_1, h_2 \in G$ .

Since  $G$  has commutator width bounded by  $k_0$ , this implies  $\|\beta_g(h) - 1_{n'}\| < 24k_0\eta^{1/4}$  for all  $h \in G$ . Thus,  $\Delta_g\varphi$  is almost constant for  $g \in A \cap A^{-1}$ . Since  $A \cap A^{-1}$  has density at least  $1 - 2\varepsilon^{1/2}$  and  $\|\varphi\|_\infty \leq 1$ , this implies that  $\|\varphi\|_{U^2}$  is close to 1. Indeed, we obtain

$$\|\varphi\|_{U^2}^4 = \mathbb{E}_{g_0, g_1, g_2} \text{tr}(\Delta_{g_1, g_2}\varphi)(g_0) \geq 1 - 48k_0\eta^{1/4} - 4\varepsilon^{1/2}.$$

Thus, taking  $\eta$  and  $\varepsilon$  sufficiently small, so that  $48k_0\eta^{1/4} + 4\varepsilon^{1/2} \leq \varepsilon(\delta, 2, k_0)$  we are done by the results of Gowers–Hatami for the case  $k = 2$ .  $\square$

REMARK 6.4. The result of Gowers–Hatami [6, Theorem 5.6] for  $k = 2$  is more general since it is also an inverse theorem in the 1% regime, i.e., a positive correlation with a linear map is derived only from  $\|\varphi\|_{U^2} \geq \varepsilon$ . We were not able to generalize this to hold for arbitrary  $k \geq 3$ .

### Acknowledgments

Andreas Thom acknowledges funding by the Deutsche Forschungsgemeinschaft (SPP 2026 “Geometry at infinity”).

## References

- [1] Oren Becker and Michael Chapman, *Stability of approximate group actions: uniform and probabilistic*, J. Eur. Math. Soc. (JEMS) **25** (2023), no. 9, 3599–3632. [↑5](#)
- [2] Kenneth S. Brown, *Cohomology of groups*, Graduate Texts in Mathematics, vol. 87, Springer-Verlag, New York, 1994. Corrected reprint of the 1982 original. [↑2](#), [3](#), [4](#)
- [3] Marcus De Chiffre, Narutaka Ozawa, and Andreas Thom, *Operator algebraic approach to inverse and stability theorems for amenable groups*, Mathematika **65** (2019), no. 1, 98–118. [↑5](#), [5](#), [6](#)
- [4] Ben Green, Terence Tao, and Tamar Ziegler, *An inverse theorem for the Gowers  $U^{k+1}[N]$ -norm*, Ann. Math. **176** (2012), 1231–1372. [↑4](#), [6](#)
- [5] W. T. Gowers, *A new proof of Szemerédi’s theorem*, Geom. Funct. Anal. **11** (2001), no. 3, 465–588 (2001); erratum 11, no. 4, 869. [↑6](#)
- [6] Timothy Gowers and Omid Hatami, *Inverse and stability theorems for approximate representations of finite groups*, Mat. Sb. **208** (2017), no. 12, 70–106 (Russian, with Russian summary); English transl., Sb. Math. **208** (2017), no. 12, 1784–1817. [↑1](#), [5](#), [6](#), [6](#), [6.4](#)
- [7] Karsten Grove, Hermann Karcher, and Ernst A. Ruh, *Group actions and curvature*, Bull. Amer. Math. Soc. **81** (1975), 89–92. [↑5](#)
- [8] Derek Holt and Wilhelm Plesken, *Perfect groups*, Oxford Math. Monogr. (Oxford University Press, New York, 1989). [↑5.2](#)
- [9] Asgar Jamneshan and Terence Tao, *The inverse theorem for the  $U^3$  Gowers uniformity norm on arbitrary finite abelian groups: Fourier-analytic and ergodic approaches*, Discrete Anal. (2023), Paper No. 11, 48. [↑6](#)
- [10] David Kazhdan, *On  $\varepsilon$ -representations*, Israel J. Math. **43** (1982), no. 4, 315–323. [↑5](#)
- [11] Alexander Leibman, *Polynomial sequences in groups*, J. Algebra **201** (1998), no. 1, 189–206. [↑1](#)
- [12] ———, *Polynomial mappings of groups*, Israel J. Math. **129** (2002), 29–60. [↑1](#), [4.3](#)
- [13] Martin W. Liebeck, E. A. O’Brien, Aner Shalev, and Pham Huu Tiep, *The Ore conjecture*, J. Eur. Math. Soc. (JEMS) **12** (2010), no. 4, 939–1008. [↑5.2](#)
- [14] Nikolay Nikolov, *On the commutator width of perfect groups*, Bull. London Math. Soc. **36** (2004), 30–36. [↑5.2](#)
- [15] Terence Tao and Tamar Ziegler, *The inverse conjecture for the Gowers norm over finite fields via the correspondence principle*, Anal. PDE **3** (2010), no. 1, 1–20. [↑1](#), [6](#)
- [16] ———, *The inverse conjecture for the Gowers norm over finite fields in low characteristic*, Ann. Comb. **16** (2012), no. 1, 121–188. [↑1](#), [6](#)
- [17] John Wilson, *First-order group theory*, Infinite groups 1994 (Ravello), de Gruyter, Berlin, 1996, pp. 301–314. [↑1](#), [5.2](#)

ASGAR JAMNESHAN, TU DRESDEN, 01062 DRESDEN  
 Email address: `asgar.jamneshan@tu-dresden.de`

ANDREAS THOM, TU DRESDEN, 01062 DRESDEN  
 Email address: `andreas.thom@tu-dresden.de`