

The Deep Underground Neutrino Experiment (DUNE) program

Inés Gil-Botella* on behalf of the DUNE collaboration

CIEMAT, Centro de Investigaciones Energéticas Medioambientales y Tecnológicas, Av. Complutense, 40 - E-28040 Madrid, Spain

E-mail: ines.gil@ciemat.es

The Deep Underground Neutrino Experiment (DUNE) is a next-generation long-baseline neutrino oscillation experiment aimed at determining the neutrino mass hierarchy and the CP-violating phase. The DUNE physics program also includes the detection of astrophysical neutrinos and the search for signatures beyond the Standard Model, such as nucleon decays. DUNE consists of a near detector complex located at Fermilab and four 17 kton Liquid Argon Time Projection Chamber (LArTPC) far detector modules to be built 1.5 km underground at SURF, approximately 1300 km away. The detectors are exposed to a wideband neutrino beam generated by a 1.2 MW proton beam with a planned upgrade to > 2 MW. Two 770 ton LArTPCs (ProtoDUNEs) have been operated at CERN for over 2 years as a testbed for DUNE far detectors and have been optimized to take new cosmic and test-beam data in 2024-2025. The DUNE and ProtoDUNE experiments and physics goals, as well as recent progress and results, are presented.

42nd International Conference on High Energy Physics (ICHEP2024) 18-24 July 2024 Prague, Czech Republic

*Speaker

1. The DUNE physics program

The Deep Underground Neutrino Experiment (DUNE) [1] is a world-leading liquid argon TPC (LArTPC) neutrino oscillation experiment with the potential to deliver groundbreaking results as the unambiguous determination of the mass ordering, and the discovery of leptonic CP violation. The DUNE physics program also includes the detection of astrophysical neutrinos and the search for physics Beyond the Standard Model (BSM).

DUNE will detect neutrinos from the most powerful neutrino beam in the world (> 2 MW) produced at Fermilab, and sent to the Sanford Underground Research Facility in Lead, South Dakota (SURF) along 1300 km of distance. A suit of four Far Detector (FD) modules (70 kton LAr TPCs) will be installed 1.5 km underground at SURF to observe the neutrino interactions. A Near Detector (ND) complex at 560 m from the source will measure the neutrino flux and provide an unprecedented control of the systematic uncertainties to enable precision studies of neutrino oscillations.

The measurement of the neutrino and antineutrino energy spectra at the FD modules will enable the determination of the CP phase, neutrino mass ordering and θ_{23} mixing angle [2, 3]. Figures 1 and 2 show the DUNE neutrino mass ordering and CP violation sensitivity as a function of the years for the currently assumed staging scenario. For the best case ($\delta_{CP} = -\pi/2$), DUNE will reach a 5σ mass ordering sensitivity in 1 year of data and 3σ in 3.5 years. For the worst case, DUNE will need only 3 years to reach 5σ sensitivity for the mass ordering and in long-term, DUNE can establish CP violation over 75% of the δ_{CP} values at more than 3σ .




Figure 1: The significance for DUNE to establish the mass ordering for the best case (left panel) and worse case (right) as a function of running time.

The ultimate goal of DUNE is to provide high precision measurements of the oscillation parameters in a single experiment. DUNE will reach between 6-16 degrees resolution to δ_{CP} (with external input for only solar parameters) depending on the true δ_{CP} value and world-leading precision in the θ_{13} measurement by a long-baseline experiment, to be compared with the θ_{13} measurement by future reactor neutrino experiments (Figure 3).

Besides the long-baseline neutrino oscillation program, DUNE has a broad non-beam physics program. The DUNE underground location allows for astrophysical neutrino measurements, in particular neutrinos from core-collapse supernovae and the sun. DUNE has unique sensitivity to MeV electron neutrinos, thanks to the ν_e CC absorption channel on 40 Ar. DUNE expects to detect




Figure 2: The significance for DUNE to establish CP violation for the best case (left panel) and 75% (right) of δ_{CP} values as a function of running time.




Figure 3: (Left panel) The DUNE resolution to δ_{CP} for as a function of the true δ_{CP} value for different integrated exposures. (Right panel) The DUNE resolution to $\sin^2 2\theta_{13}$ as a function of exposure in kt-MW-yrs.

several thousand events from a galactic supernova burst. As shown in Figure 4-left, DUNE can detect the supernova neutronization burst, which is an expected early emission of ν_e , and measure the neutrino mass ordering by looking the suppression or not of this burst [4]. DUNE is also sensitive to all neutrino flavors through the elastic-scattering (ES) channel, which provides pointing capabilities, being able to achieve ~5 degrees pointing resolution [5].

DUNE has the potential to detect a huge amount of electron neutrinos from the sun (\sim 10k/year CC; 2k ES/year) together with radioactive background events expanding up to \sim 14 MeV (mainly neutron-capture events from the cavern surrounding the detector at SURF). The 8B solar neutrino flux is visible above 9-10 MeV and the hep flux can be measured for the first time. The measurement of the 8B neutrinos will improve current solar measurements of Δm_{21}^2 because of the day-night asymmetry induced by Earth matter effects.

DUNE provides enormous opportunities to probe phenomena beyond the Standard Model (BSM) [6]. By measuring neutrino and antineutrino spectra, DUNE can look for neutrino oscillation




Figure 4: Expected event rates as a function of time for 40 kton of argon during early stages of a supernova burst at 10 kpc (left). DUNE sensitivity to inelastic boosted dark matter search in the *p*-scattering channel [6] (right).

scenarios beyond the standard three-flavor picture, including sterile neutrinos, and PMNS non-unitarity. The high statistics in neutrino and antineutrino measurements allow for CPT violation searches. DUNE has unique sensitivity to non-standard interactions due to long-baseline matter effects. Other BSM searches in ND and FD are possible such as the search for inelastic dark matter scattering at the FD (Figure 4-right), low-mass dark matter at the ND, nucleon decay, neutron-antineutron oscillations, heavy-neutral leptons, neutrino tridents, among others.

2. The DUNE detectors

To accomplish this physics program, DUNE is currently building two LArTPC FD modules. One module (FD-HD) has an horizontal drift (3.6 m drift) and uses wire readout planes with 4 drift regions (Figure 5-left) [7]. In the other FD (FD-VD) [8], the ionizing electrons drift vertically in two 6 m drift regions (Figure 5-center). This second technology is simpler to install and is the baseline for the third and forth FD modules. New caverns to allocate these huge detectors have been excavated at SURF (Figure 5-right).




Figure 5: The horizontal drift FD-HD module (left), vertical drift FD-VD module (center), and one of the SURF caverns that will host the DUNE far detectors (right).

The DUNE ND complex [9] will enable the prediction of the FD reconstructed spectra. The ND will use a movable detector system composed of a LAr TPC (ND-LAr) and a muon spectrometer (TMS). They will collect high statistics of the neutrino beam. The off-axis data of this system will

allow to constrain the energy dependence of the neutrino cross sections. The use of the same target and technology will inform predictions of the neutrino reconstructed energy in the FD. The final component of the ND complex will be an on-axis magnetized detector (SAND) to monitor the beam stability and perform measurements of neutrino interactions.

DUNE has developed a two-phase strategy toward the implementation of the leading-edge, large-scale science projects. The construction of the initial experiment configuration (DUNE Phase I) is well underway. It includes the two FD modules and the ND components already described, with a 1.2 MW proton beam. DUNE Phase-II [10] consists of a third and fourth FD module, an upgraded ND complex, and an enhanced 2.1 MW beam.

3. The DUNE prototypes

The DUNE collaboration is developing a very active prototyping program for both ND and FDs at Fermilab and CERN in order to test at large scale the main components and demonstrate the detector performance. In particular, one key aspect of the ND-LAr operation is the ability to cope with a large number of neutrino interactions in each spill. The expected high rate at the near site motivates the use of a pixelated readout and optical modularity in the ND. In order to test this new technology, four LAr TPC modules (Figure 6-left) have been built and operated in LAr at the University of Bern and then moved to Fermilab and installed in the former location of the MINOS ND. The system also includes upstream and downstream MINERvA scintillator and calorimeter planes. The ND-LAr 2x2 demonstrator has detected the first neutrino interactions and will continue collecting data with the goal of proving the 3D readout in a neutrino beam with similar event rate to DUNE.




Figure 6: One of the four modules of the ND-LAr 2x2 demonstrator (left panel). Inner view of ProtoDUNE-HD (center) and ProtoDUNE-VD (right) at the CERN Neutrino Platform.

In order to demonstrate the technical feasibility of the FDs and show the expected performance, two full-scale LAr prototypes (770 ton each) were built and operated at CERN between 2018 and 2020 [11]. After a successful operation with cosmic muons and charged particles, ProtoDUNEs are now entering a new phase. Two new prototypes were installed at the CERN Neutrino Platform, incorporating the final technical solutions for FD-HD and FD-VD (Figure 6-center, right). ProtoDUNE-HD was filled with LAr and started taking cosmic rays and test-beam data at CERN in May 2024. This new run will increase beam data statistics for cross section measurements, particle identification, calibration, and reconstruction developments. After the operation of ProtoDUNE-HD, LAr will be transferred to ProtoDUNE-VD in fall 2024 for running in early 2025.

The VD technology is the baseline design for the DUNE FD Phase II modules. Some improvements to the light collection for FD3 are already being explored. But the DUNE phased construction allows for more ambitious detector improvements that allow to expand the DUNE physics scope, in particular, the low-energy physics program. The collaboration is pursuing an active R&D program for the so-called Module of Opportunity FD4 [10], including pixel readout in the FD, integrated charge-light readout, as the SoLAr proposal [12], low background modules or non-LAr technologies.

References

- [1] B. Abi et al. [DUNE], Deep Underground Neutrino Experiment (DUNE), Far Detector Technical Design Report, Volume I Introduction to DUNE, JINST 15 (2020) no.08, T08008 [arXiv:2002.02967].
- [2] B. Abi et al. [DUNE], Long-baseline neutrino oscillation physics potential of the DUNE experiment Eur. Phys. J. C 80 (2020) no.10, 978 [arXiv:2006.16043].
- [3] A. Abed Abud et al. [DUNE], Low exposure long-baseline neutrino oscillation sensitivity of the DUNE experiment, Phys. Rev. D 105 (2022) no.7, 072006 [arXiv:2109.01304].
- [4] B. Abi et al. [DUNE], Supernova neutrino burst detection with the Deep Underground Neutrino Experiment, Eur. Phys. J. C 81 (2021) no.5, 423 [arXiv:2008.06647].
- [5] A. Abed Abud *et al.* [DUNE], *Supernova Pointing Capabilities of DUNE*, [arXiv:2407.10339].
- [6] B. Abi et al. [DUNE], Prospects for beyond the Standard Model physics searches at the Deep Underground Neutrino Experiment, Eur. Phys. J. C 81 (2021) no.4, 322 [arXiv:2008.12769].
- [7] B. Abi et al. [DUNE], Deep Underground Neutrino Experiment (DUNE), Far Detector Technical Design Report, Volume IV: Far Detector Single-phase Technology, JINST 15 (2020) no.08, T08010 [arXiv:2002.03010].
- [8] A. Abed Abud et al. [DUNE], The DUNE Far Detector Vertical Drift Technology. Technical Design Report, JINST 19 (2024) no.08, T08004 [arXiv:2312.03130].
- [9] V. Hewes et al. [DUNE], Deep Underground Neutrino Experiment (DUNE) Near Detector Conceptual Design Report, Instruments 5 (2021) no.4, 31 [arXiv:2103.13910].
- [10] A. Abed Abud et al. [DUNE], DUNE Phase II: Scientific Opportunities, Detector Concepts, Technological Solutions, [arXiv:2408.12725].
- [11] A. Abed Abud et al. [DUNE], Design, construction and operation of the ProtoDUNE-SP Liquid Argon TPC, JINST 17 (2022) no.01, P01005 [arXiv:2108.01902].
- [12] N. Anfimov et al. [SoLAr], First Demonstration of a Combined Light and Charge Pixel Readout on the Anode Plane of a LArTPC, [arXiv:2406.14121].