ℓ^1 -BASES, ALGEBRAIC STRUCTURE AND STRONG ARENS **IRREGULARITY OF BANACH ALGEBRAS IN HARMONIC** ANALYSIS¹

M. FILALI AND J. GALINDO

ABSTRACT. A long standing problem in abstract harmonic analysis concerns the strong Arens irregularity (sAir, for short) of the Fourier algebra A(G) of a locally compact group G. The groups for which A(G) is known to be sAir are all amenable. So far this class includes the abelian groups, the discrete amenable groups, the second countable amenable groups G such that [G, G]is not open in G, the groups of the form $\prod_{i=0}^{\infty} G_i$ where each $G_i, i \ge 1$, is a non-trivial metrizable compact group and G_0 is an amenable second countable locally compact group, the groups of the form $G_0 \times G$, where G is a compact group whose local weight w(G) has uncountable cofinality and G_0 is any locally compact amenable group with $w(G_0) \leq w(G)$, and the compact group SU(2).

We were primarily concerned with the groups for which A(G) is sAir. We introduce a new class of ℓ^1 -bases in Banach algebras. These new ℓ^1 -bases enable us, among other results, to unify most of the results related to Arens products proved in the past seventy years since Arens defined his products. This includes the strong Arens irregularity of algebras in harmonic analysis, and in particular almost all the cases mentioned above for the Fourier algebras. In addition, we also show that A(G) is sAir for compact connected groups with an infinite dual rank. The ℓ^1 -bases for the Fourier algebra are constructed with coefficients of certain irreducible representations of the group. With this new approach using ℓ^1 -bases, the rich algebraic structure of the algebras and semigroups under study such as the second dual of Banach algebras with an Arens product or certain semigroup compactifications (the Stone-Cech compactification of an infinite discrete group, for instance) is also unveiled

CONTENTS

1.	Introduction	2
2.	Some basic definitions and notation	3
3.	Factorizations of \mathcal{A}^* and $\mathcal{LUC}(\mathcal{A}^*)$ and topological centres	7
4.	$F\ell^1(\eta)$ -bases in \mathcal{A} and factorizations of \mathcal{A}^* , $\mathcal{LUC}(\mathcal{A})$ and $M_r(\mathcal{A})^*$	11
5.	The algebraic structure of \mathcal{A}^{**} , $\mathcal{LUC}(\mathcal{A})^*$ and $M_r(\mathcal{A})^{**}$	21
6.	Application I: Group algebra, semigroup algebra and measure algebra	26
7.	Application II: Weighted group, semigroup and measure algebras	34
8.	Application III: Fourier algebra	38
9.	Factorizations of $VN(G \times H)$ and applications	52
Re	References	
م		
Date: Decemper 20, 2024.		

2010 Mathematics Subject Classification. Primary 46H10, 43A30, 22D15, 22D10, 43A75; Secondary 22D10, 43A20, 43A10.

Key words and phrases. Arens products, Arens-regular algebra, Banach algebra, F ℓ^1 -base, Factorization, strongly Arens irregular, group algebra, measure algebra, weighted group algebra, weighted measure algebra, Fourier algebra, compact non-metrizable groups, infinite product of compact groups, compact connected groups.

 $^{^{1}}$ The results were presented in 2019 IMAC AHA Days (Castellon, Spain) and in the conference on Banach Algebras and Applications, 2017 (Oulu, Finland).

1. INTRODUCTION

A Banach algebra \mathcal{A} is Arens regular when the two Arens products defined on its second Banach dual \mathcal{A}^{**} coincide. This is equivalent to saying that one of the products (and so both products are) is separately weak*-continuous making \mathcal{A}^{**} a semitopological algebra when equipped with the weak*-topology. It is worthwhile to note, that due to Grothendieck's criterion, this is in turn equivalent to that every element in the Banach dual \mathcal{A}^* is weakly almost periodic, i.e., $\mathcal{A}^* = \mathcal{WAP}(\mathcal{A})$. This fact is not going to be used in the present article, but was essential in our study of Arens irregularity of Banach algebras in harmonic analysis in the articles [40] and [41]. As seen already in these past two articles and as we shall further see presently, unlike C^{*}-algebras, the usual algebras in harmonic analysis such as the group algebra, and the Fourier algebra in most cases, have turned out to be all non-Arens regular (even in an extreme way). At the very beginning, about seventy years ago, Richard Arens proved in [1] that the convolution algebra ℓ^1 is Arens irregular. A long exciting story followed, and so much has already been written on the subject ever since. For more details, see the surveys [23] and [51], the book [15], the memoirs [16] and [17], and our most recent survey [39] where the results of the actual article were announced.

There are two different types of utmost non-Arens regularities, arising naturally from the way the algebras are decided to be non-Arens regular.

Since \mathcal{A} is Arens regular if and only if $\mathcal{A}^* = \mathcal{WAP}(\mathcal{A})$, the degree of non-Arens regularity of an algebra \mathcal{A} may be measured by the size of the quotient space $\frac{\mathcal{A}^*}{\mathcal{WAP}(\mathcal{A})}$. In the extreme situation (i.e., when $\frac{\mathcal{A}^*}{\mathcal{WAP}(\mathcal{A})}$ is as large as \mathcal{A}^*) Granirer called such an algebra \mathcal{A} extremely non-Arens regular (enAr for short) and proved that this was the case for the Fourier algebra of a non-discrete second countable locally compact group G (see [58, Corollaries 6 and 7]). Granirer's result was extended by Hu in [68] to locally compact groups G having their local weight w(G)greater or equal to their compact covering $\kappa(G)$.

On the dual hand, the group algebra is isometrically enAr, and so enAr, for any infinite locally compact group G. This was proved in [52] when G is discrete, in [8] when $\kappa(G) \geq w(G)$, and finally in [37] for the general case. The semigroup algebra $\ell^1(S)$ is isometrically enAr for any infinite weakly cancellative discrete semigroup (see [52]). Here, a Banach algebra is isometrically enAr when $\frac{\mathcal{A}^*}{W\mathcal{AP}(\mathcal{A})}$ contains an isometric copy of \mathcal{A}^* .

This topic was our main focus in [40] and [41], where ℓ^1 -bases approximating lower and upper triangles of products of elements in \mathcal{A} were the reason behind the non-Arens regularity of \mathcal{A} , and in many cases the extreme non-Arens regularity of \mathcal{A} . This unified approach dealing with the extreme non-Arens regularity helped us to improve and extend the above-cited results to a large class of Banach algebras. The combinatorics of the locally compact group G and its dual object \hat{G} were the tools to construct these ℓ^1 -bases for many algebras which are associated to G in harmonic analysis.

Non-Arens regularity may also be measured by the degree of defect of the separate weak*-continuity of Arens products. In the extreme situation (i.e., when separate weak*-continuity of the product in \mathcal{A}^{**} happens only when one of the elements in the product is in \mathcal{A}), Dales and Lau called the algebra \mathcal{A} strongly Arens irregular (see [16]). This will be our main focus in the present paper. A more refined ℓ^1 -bases in \mathcal{A} are necessary to show the strong Arens irregularity of Banach algebras associated to a locally compact group G. Again the reason making these refined ℓ^1 -bases available are the combinatorics in G and \widehat{G} . The paper is organized as follows. After a preliminary section recalling the basic definitions and results needed in the paper, we present our general theory in Sections 3 and 4. The new definitions of some special ℓ^1 -bases, we call $F\ell^1(\eta)$ -base, are given for any Banach algebra \mathcal{A} (η is an infinite cardinal which is going to be the Mazur property level of the algebra \mathcal{A} when we deal with the topological centres). The theorems in these two sections show how a presence of these $F\ell^1(\eta)$ -bases in \mathcal{A} yield strong Arens irregularity of the algebra \mathcal{A} and induces a rich algebraic structure in \mathcal{A}^{**} . Similar results are also proved in $\mathcal{LUC}(\mathcal{A})^*$ when \mathcal{A} has a bounded approximate identity. Under certain conditions the theorems are proved also for the right multiplier algebra $M_r(\mathcal{A})$ of \mathcal{A} .

We illustrate our general theory with three sections, starting with the group algebra and the measure algebra of a non-compact locally compact group and the semigroup algebra of an infinite weakly cancellative, right cancellative, discrete semigroup. Then come the weighted analogues of these algebras. These are studied when the weight is diagonally bounded. Using some special subsets of G constructed inductively, the required $F\ell^1(\eta)$ -bases are produced in each case, where η is the compact covering of the locally compact group, or the cardinality of the discrete semigroup.

The penultimate section deals with the Fourier algebra of a compact group. The cases we study first are compact groups with local weight having uncountable cofinality, then infinite product of compact groups, and finally compact connected groups with a certain infinite dimension (this will be defined in time). Using the irreducible representations of G, we shall produce in each case the required $F\ell^1(\eta)$ -bases, where η is the local weight of G.

With the availability of the required $F\ell^1(\eta)$ -base in each of the cases studied in Sections 6, 7 and 8, the topological centers as well as the algebraic structure of all the algebras in question will be an immediate consequence of Sections 3, 4 and 5.

In the last section, we shall extend our results to the case when the group is the product $G \times H$, where G is any of the previous compact groups and H is an amenable locally compact group with local weight smaller or equal to that of G.

The algebraic structures and the topological centres of the Banach algebras $A(G \times H)^{**}$, $\mathfrak{UC}_2(G \times H)^*$ or $C^*(G \times H)^{\perp}$ will be our final conclusion.

Some of these results have been proved in the past by various authors as we shall indicate as we go along. In addition to the unification of all these results, there will be many other new results.

2. Some basic definitions and notation

Let \mathcal{A} be a Banach algebra. The elements in \mathcal{A} will be denoted by miniscule Roman letters. The elements in the Banach dual \mathcal{A}^* of \mathcal{A} will be denoted by majuscule Roman letters. Majuscule Greek letters will be used to denote the elements in the second dual of the algebra. With the exceptions of the product in \mathcal{A} and its first Arens extensions to \mathcal{A}^{**} and $\mathcal{LUC}(\mathcal{A})^*$, all the module actions will be denoted by ".". With the elements denoted as such, there will be no confusion as to which module action is considered.

2.1. Arens products. Recall that the Arens module actions are defined via the adjoint of continuous bilinear mappings (see [1] and [2]). That is, if

 $\mathcal{E}_1 \times \mathcal{E}_2 \to \mathcal{E}_3$ is a continuous bilinear mapping

on some normed spaces \mathcal{E}_i (i = 1, 2, 3), then its (312)-adjoint given by

 $\mathcal{E}_3^* \times \mathcal{E}_1 \to \mathcal{E}_2^*$ is again a continuous bilinear mapping,

and so in turn, its (312)-adjoint may be be taken. The process can be repeated indefinitely.

Here are the module actions we shall be using throughout the paper. We start with a Banach algebra \mathcal{A} and regard it as a left \mathcal{A} -module with its operation as the left module action of \mathcal{A} on itself; that is

$$\mathcal{A} \times \mathcal{A} \to \mathcal{A}$$
, where $a \cdot b = ab$.

Then by taking the (312)-adjoint, we shall regard \mathcal{A}^* as a right \mathcal{A} -module, where the natural right module action of \mathcal{A} on \mathcal{A}^* is given by

(1)
$$\mathcal{A}^* \times \mathcal{A} \to \mathcal{A}^*$$
, where $\langle S \cdot a, b \rangle = \langle S, ab \rangle$ for $S \in \mathcal{A}^*$, $a, b \in \mathcal{A}$.

This is followed (taking again the (312)-adjoint) by regarding \mathcal{A}^* as a left \mathcal{A}^{**-} module, where the natural left module action of \mathcal{A}^{**} on \mathcal{A}^{*} is given by

(2) $\mathcal{A}^{**} \times \mathcal{A}^* \to \mathcal{A}^*$, where $\langle \Psi \cdot S, a \rangle = \langle \Psi, S \cdot a \rangle$ for $\Psi \in \mathcal{A}^{**}$, $S \in \mathcal{A}^*, a \in \mathcal{A}$.

This induces the first (left) Arens product in \mathcal{A}^{**} making \mathcal{A}^{**} a Banach algebra. For $\Phi, \Psi \in \mathcal{A}^{**}$, the product is given by

(3)
$$\mathcal{A}^{**} \times \mathcal{A}^{**} \to \mathcal{A}^{**}$$
, where $\langle \Phi \cdot \Psi, S \rangle = \langle \Phi, \Psi \cdot S \rangle$ for $S \in \mathcal{A}^*$.

The first Arens product in \mathcal{A}^{**} is the unique extension of the product in \mathcal{A} such that

(i) $\Phi \mapsto \Phi \Psi : \mathcal{A}^{**} \to \mathcal{A}^{**}$ is weak*-weak*-continuous for each $\Psi \in \mathcal{A}^{**}$,

(ii) $\Phi \mapsto a\Phi : \mathcal{A}^{**} \to \mathcal{A}^{**}$ is weak*-weak*-continuous for each $a \in \mathcal{A}$.

The first topological centre of \mathcal{A}^{**} may therefore be defined as

 $T_1Z(\mathcal{A}^{**}) = \{ \Psi \in \mathcal{A}^{**} : \Phi \mapsto \Psi \Phi : \mathcal{A}^{**} \to \mathcal{A}^{**} \text{ is weak}^* \text{-continuous} \}.$ Note that Item (ii) above means that $\mathcal{A} \subseteq T_1 Z(\mathcal{A}^{**})$.

If we start with A as a right A-module with the right module action given by the transpose of the operation in \mathcal{A} , that is $a \odot b = ba$ for $a, b \in \mathcal{A}$, then taking the transpose of the left action of \mathcal{A}^{**} on itself reached at the third step, will give the second (right) Arens product. We make an exception here and denote the second Arens product of two elements Φ and Ψ in \mathcal{A}^{**} by $\Phi \diamond \Psi$ and the actions leading to this product by " \odot " so that, for every $S \in \mathcal{A}^*$ and $a, b \in \mathcal{A}$, we have

(4)

$$\langle \Phi \diamond \Psi, S \rangle = \langle \Psi \odot \Phi, S \rangle = \langle \Psi, \Phi \odot S \rangle, \text{ where}$$

 $\langle \Phi \odot S, a \rangle = \langle \Phi, S \odot a \rangle \text{ and}$
 $\langle S \odot a, b \rangle = \langle S, a \odot b \rangle = \langle S, ba \rangle$

/ **T**

We note that the second Arens product in \mathcal{A}^{**} is the unique extension of the product in \mathcal{A} such that

- (i) $\Psi \mapsto \Phi \diamond \Psi : \mathcal{A}^{**} \to \mathcal{A}^{**}$ is weak*-weak*-continuous for each $\Phi \in \mathcal{A}^{**}$,
- (ii) $\Phi \mapsto \Phi \diamond a : \mathcal{A}^{**} \to \mathcal{A}^{**}$ is weak*-weak*-continuous for each $a \in \mathcal{A}$.

The second topological centre of \mathcal{A}^{**} may therefore be defined as

$$T_2Z(\mathcal{A}^{**}) = \{ \Psi \in \mathcal{A}^{**} : \Phi \mapsto \Phi \diamond \Psi : \mathcal{A}^{**} \to \mathcal{A}^{**} \text{ is weak*-weak*-continuous} \}.$$

Again Item (ii) means that $\mathcal{A} \subseteq T_2Z(\mathcal{A}^{**}).$

When these two ways yield the same product, i.e., if these two products coincide, we say that the algebra \mathcal{A} is Arens regular. It is clear that the algebra is Arens regular if and only if one of the Arens products (and so both are) is separately weak*-weak*-continuous, i.e.,

$$T_1Z(\mathcal{A}^{**}) = T_2Z(\mathcal{A}^{**}) = \mathcal{A}^{**}.$$

Such a situation happens for instance when \mathcal{A} is a C^* -algebra. This was first deduced from the fact that the second dual of a C^* -algebra can be seen as an algebra of operators on a Hilbert space, a result proved by Sherman in [108] and Takeda in [110]. Later on, in their seminal paper [12], Civin and Yood reproduced Takeda's proof and brought up explicitly the Arens regularity of C^* -algebras. A different proof of this can also be found in [7, Theorem 38.19].

When the Banach algebra is not Arens regular, i.e., when

$$\mathcal{A} \subseteq T_i Z(\mathcal{A}^{**}) \subsetneq \mathcal{A}^{**}$$
 for $i = 1$ or 2,

or equivalently when $\mathcal{A}^* \neq \mathcal{WAP}(\mathcal{A})$, we say that the algebra is Arens irregular or non-Arens regular.

In the extreme situation of $T_1Z(\mathcal{A}^{**}) = \mathcal{A}$ (respectively, $T_2Z(\mathcal{A}^{**}) = \mathcal{A}$), Dales and Lau called the algebra \mathcal{A} left (respectively, right) strongly Arens irregular. In the very extreme situation of $T_1Z(\mathcal{A}^{**}) = T_2Z(\mathcal{A}^{**}) = \mathcal{A}$, the algebra \mathcal{A} is strongly Arens irregular (sAir), see [16]. Note that when \mathcal{A} is commutative (as it is the case with the Fourier algebra $\mathcal{A}(G)$ studied in the last two sections), $T_1Z(\mathcal{A}^{**})$ and $T_2Z(\mathcal{A}^{**})$ coincide with the algebraic centre of \mathcal{A}^{**} , that is,

$$T_1 Z(\mathcal{A}^{**}) = T_2 Z(\mathcal{A}^{**}) = \{ \Psi \in \mathcal{A}^{**} : \Phi \Psi = \Psi \Phi \text{ for every } \Phi \in \mathcal{A}^{**} \}$$
$$= \{ \Psi \in \mathcal{A}^{**} : \Phi \diamond \Psi = \Psi \diamond \Phi \text{ for every } \Phi \in \mathcal{A}^{**} \}.$$

In this case, we will be talking just about the strongly Arens irregularity.

It should be also recalled that $T_1Z(\mathcal{A}^{**})$ and $T_2Z(\mathcal{A}^{**})$ may be different, and it may even happen that $T_1Z(\mathcal{A}^{**}) = \mathcal{A} \neq T_2Z(\mathcal{A}^{**})$, i.e., \mathcal{A} may be only one-sided strongly Arens irregular (see [16], [26], [55] and [104]).

2.2. The space $\mathcal{LUC}(\mathcal{A})$. We shall also be concerned with the dual of the following closed subspace of \mathcal{A}^* . This is the closed linear span in \mathcal{A}^* of $\mathcal{A}^* \cdot \mathcal{A}$, which we denote as usual by $\mathcal{LUC}(\mathcal{A})$. By Cohen's factorization theorem, this is $\mathcal{A}^* \cdot \mathcal{A}$ when \mathcal{A} has a bounded right approximate identity.

Now since $(S \cdot a) \cdot b = S \cdot (ab)$ for every $S \in \mathcal{A}^*$ and $a, b \in \mathcal{A}$, we see that $\mathcal{LUC}(\mathcal{A})$ is an \mathcal{A}^* -submodule, where the module action is given as in (1) (but with $\mathcal{LUC}(\mathcal{A})$ instead of \mathcal{A}^*) by

$$\mathcal{LUC}(\mathcal{A}) \times \mathcal{A} \to \mathcal{LUC}(\mathcal{A}) : (S \cdot a, b) \mapsto S \cdot (ab).$$

After taking the (312)-adjoint, we obtain the action $\mathcal{LUC}(\mathcal{A})^* \times \mathcal{LUC}(\mathcal{A}) \to \mathcal{A}^*$. But as known and easy to check (noting that $\Phi \cdot (S \cdot a) = (\Phi \cdot S) \cdot a$ for every $\Phi \in \mathcal{LUC}(\mathcal{A})^*$, $S \in \mathcal{A}^*$ and $a \in \mathcal{A}$), the range of this mapping is in $\mathcal{LUC}(\mathcal{A})$. In other words, we have in fact $\mathcal{LUC}(\mathcal{A})$ as a right $\mathcal{LUC}(\mathcal{A})^*$ -module

$$\mathcal{LUC}(\mathcal{A})^* \times \mathcal{LUC}(\mathcal{A}) \to \mathcal{LUC}(\mathcal{A})$$

(in the literature, a subspace of \mathcal{A}^* with such a property is said to be left introverted), and so as in (3), this induces by taking the (312)-adjoint, the first Arens product in $\mathcal{LUC}(\mathcal{A})^*$ making $\mathcal{LUC}(\mathcal{A})^*$ a Banach algebra. As in (3), the product of $\Phi, \Psi \in \mathcal{LUC}(\mathcal{A})^*$ is given by

$$\langle \Phi \Psi, S \rangle = \langle \Phi, \Psi \cdot S \rangle$$
 for all $S \in \mathcal{LUC}(\mathcal{A})$.

This product is the unique extension of the product in \mathcal{A} to $\mathcal{LUC}(\mathcal{A})^*$ such that

- (i) $\Phi \mapsto \Phi \Psi : \mathcal{LUC}(\mathcal{A})^* \to \mathcal{LUC}(\mathcal{A})^*$ is weak*-weak*-continuous for each $\Psi \in \mathcal{LUC}(\mathcal{A})^*$,
- (ii) $\Phi \mapsto a\Phi : \mathcal{LUC}(\mathcal{A})^* \to \mathcal{LUC}(\mathcal{A})^*$ is weak*-weak*-continuous for each $a \in \mathcal{A}$.

The (first) topological centre of $\mathcal{LUC}(\mathcal{A})^*$ may therefore be defined as

$$T_1Z(\mathcal{LUC}(\mathcal{A})^*) = \{ \Psi \in \mathcal{LUC}(\mathcal{A})^* : \Phi \mapsto \Psi \Phi : \mathcal{LUC}(\mathcal{A})^* \to \mathcal{LUC}(\mathcal{A})^*$$
 is weak*-weak*-continuous}.

It may be also worthwhile to note that when \mathcal{A} has a brai (the definition is given is Section 3), $\mathcal{LUC}(\mathcal{A})^*$ and the quotient $\frac{\mathcal{A}^{**}}{\mathcal{LUC}(\mathcal{A})^{\perp}}$ are isomorphic as Banach algebras, where

$$\mathcal{LUC}(\mathcal{A})^{\perp} = \{ \Psi \in \mathcal{A}^{**} : \langle \Psi, S \rangle = 0 \quad \text{for every} \quad S \in \mathcal{LUC}(\mathcal{A}) \}$$

is a closed ideal in \mathcal{A}^{**} . For more details, see [5].

Note finally that by the definition of $\mathcal{LUC}(\mathcal{A})$, the module action (1) may be put as

$$\mathcal{A}^* \times \mathcal{A} \to \mathcal{LUC}(\mathcal{A}),$$

and so the (312)-adjoint of this bilinear mapping makes \mathcal{A}^* as a left $\mathcal{LUC}(\mathcal{A})^*$ -module, where the left action of $\mathcal{LUC}(\mathcal{A})^*$ on \mathcal{A}^* is given by

(5)
$$\mathcal{LUC}(\mathcal{A})^* \times \mathcal{A}^* \to \mathcal{A}^*, \text{ where} \\ \langle \Psi \cdot S, a \rangle = \langle \Psi, S \cdot a \rangle, \text{ for } \Psi \in \mathcal{LUC}(\mathcal{A})^*, S \in \mathcal{A}^*, a \in \mathcal{A}.$$

Of course this action extends in turn to

(6)
$$\mathcal{A}^{**} \times \mathcal{LUC}(\mathcal{A})^* \to \mathcal{A}^{**}, \text{ where} \\ \langle \Phi.\Psi, S \rangle = \langle \Phi, \Psi.S \rangle \text{ for } \Phi \in \mathcal{A}^{**}, \Psi \in \mathcal{LUC}(\mathcal{A})^*, S \in \mathcal{A}^*.$$

The actions (5) and (6) do not conflict with the actions (2) and (3) giving the first Arens product for \mathcal{A}^{**} . In fact, if $\pi : \mathcal{A}^{**} \to \mathcal{LUC}(\mathcal{A})^*$ is the adjoint of the inclusion map $\mathcal{LUC}(\mathcal{A}) \hookrightarrow \mathcal{A}^*$, then for $S \in \mathcal{A}^*$, $\Phi \in \mathcal{A}^{**}$ and $\Psi \in \mathcal{LUC}(\mathcal{A})^*$, we have

(7)
$$\Psi.S = \Psi.S$$
 and $\Phi.\Psi = \Phi\Psi$,

where Ψ is picked in \mathcal{A}^{**} such that $\pi(\Psi) = \Psi$ (any Hahn-Banach extension of Ψ to \mathcal{A}^* will do). So we keep, as agreed, the dot to denote all the actions in the paper. No tilde will be used for the elements in \mathcal{A} , so miniscule Roman letter will be used whether \mathcal{A} is regarded as a subalgebra of \mathcal{A}^{**} or $\mathcal{LUC}(\mathcal{A})^*$.

Note that if (Ψ_{α}) is a bounded net in $\mathcal{LUC}(\mathcal{A})^*$ with weak*-limit Ψ in $\mathcal{LUC}(\mathcal{A})^*$ and $(\widetilde{\Psi_{\alpha}})$ is the corresponding bounded net of extensions in \mathcal{A}^{**} with weak*-limit Φ in \mathcal{A}^{**} , then by (7)

 $\widetilde{\Psi_{\alpha}} \cdot S = \Psi_{\alpha} \cdot S$ for every α and $S \in \mathcal{A}^*$,

and so

 $\Phi \cdot S = \Psi \cdot S$ for every $S \in \mathcal{A}^*$.

The analogue of the second Arens product defined on \mathcal{A}^{**} in (4) and so the (second) topological centre of $\mathcal{LUC}(\mathcal{A})^*$ cannot be defined for $\mathcal{LUC}(\mathcal{A})^*$ unless $S \odot a$ and $\Phi \odot S$ are in $\mathcal{LUC}(\mathcal{A})$ for every $a \in \mathcal{A}, S \in \mathcal{A}$ and $\Phi \in \mathcal{LUC}(\mathcal{A})^*$, which happens for example when \mathcal{A} is commutative. Here, we must use the space $\mathcal{RUC}(\mathcal{A})$ instead. This is the closed linear span in $\mathcal{A}^* \odot \mathcal{A}$, which again by Cohen's factorization theorem, is equal to $\mathcal{A}^* \odot \mathcal{A}$ when \mathcal{A} has a bounded left approximate identity.

Since $\mathcal{RUC}(\mathcal{A}) \odot \mathcal{A} \subseteq \mathcal{RUC}(\mathcal{A})$ and $\mathcal{RUC}(\mathcal{A})^* \odot \mathcal{RUC}(\mathcal{A}) \subseteq \mathcal{RUC}(\mathcal{A})$, the same procedure as before yields the second Arens product in $\mathcal{RUC}(\mathcal{A})^*$ making $\mathcal{RUC}(\mathcal{A})^*$ a Banach algebra: As in (4), the product of $\Phi, \Psi \in \mathcal{RUC}(\mathcal{A})^*$ is given by

$$\langle \Phi \diamond \Psi, S \rangle = \langle \Psi \odot \Phi, S \rangle = \langle \Psi, \Phi \odot S \rangle$$
 for all $S \in \mathcal{RUC}(\mathcal{A})$.

This product is the unique extension of the product in \mathcal{A} to $\mathcal{RUC}(\mathcal{A})^*$ such that

- (i) $\Psi \mapsto \Phi \diamond \Psi : \mathcal{RUC}(\mathcal{A})^* \to \mathcal{RUC}(\mathcal{A})^*$ is weak*-weak*-continuous for each $\Phi \in \mathcal{LUC}(\mathcal{A})^*$,
- (ii) $\Phi \mapsto \Phi \diamond a : \mathcal{RUC}(\mathcal{A})^* \to \mathcal{RUC}(\mathcal{A})^*$ is weak*-weak*-continuous for each $a \in \mathcal{A}$.

The (second) topololgical centre of $\mathcal{RUC}(\mathcal{A})^*$ may therefore be defined as $T_2Z(\mathcal{RUC}(\mathcal{A})^*) = \{\Psi \in \mathcal{RUC}(\mathcal{A})^* : \Phi \mapsto \Psi \diamond \Phi : \mathcal{RUC}(\mathcal{A})^* \to \mathcal{RUC}(\mathcal{A})^*$

 $I_{2Z}(\mathcal{KUC}(\mathcal{A})) = \{\Psi \in \mathcal{KUC}(\mathcal{A}) : \Psi \mapsto \Psi \diamond \Psi : \mathcal{KUC}(\mathcal{A}) \to \mathcal{KUC}(\mathcal{A})\}$

is weak*-weak*-continuous}.

Note again that $\mathcal{A} \subseteq T_2 Z(\mathcal{RUC}(\mathcal{A})^*)$.

Note also if $\Phi \odot S$ and $\Phi \odot \Psi$ are the actions of $\mathcal{RUC}(\mathcal{A})^*$ on \mathcal{A}^* and \mathcal{A}^{**} as defined with $\mathcal{LUC}(\mathcal{A})$ in (5) and (6), respectively, then

$$\Phi \odot S = \widetilde{\Phi} \odot S \quad \text{and} \quad \Psi \odot \Phi = \widetilde{\Phi} \diamond \Psi,$$

where $\widetilde{\Phi}$ is picked by Hahn-Banach Theorem in \mathcal{A}^{**} such that $\pi(\widetilde{\Phi}) = \Phi$. Here $\pi : \mathcal{A}^{**} \to \mathcal{RUC}(\mathcal{A})^*$ is the adjoint of the inclusion map $\mathcal{RUC}(\mathcal{A}) \hookrightarrow \mathcal{A}^*$.

3. Factorizations of \mathcal{A}^* and $\mathcal{LUC}(\mathcal{A}^*)$ and topological centres

Let \mathcal{A} be a Banach algebra and η be a cardinal number. Then the Banach dual \mathcal{A}^* of \mathcal{A} (respectively, $\mathcal{LUC}(\mathcal{A})$) is said to have the η -factorization property if for every bounded family of functionals $(S_{\lambda})_{\lambda < \eta}$ contained in \mathcal{A}^* , (respectively, $\mathcal{LUC}(\mathcal{A})$), there exist a bounded family $(\Psi_{\lambda})_{\lambda < \eta}$ in $\mathcal{LUC}(\mathcal{A})^*$ (usually in the unit ball of $\mathcal{LUC}(\mathcal{A})^*$) and a single functional $T \in \mathcal{A}^*$ (respectively, in $\mathcal{LUC}(\mathcal{A})$) such that the factorization formula $S_{\lambda} = \Psi_{\lambda} \cdot T$ holds for all $\lambda < \eta$.

This factorization was obtained and used, together with the so-called Mazur property, by Neufang in [100] to show that the group algebra of a non-compact locally compact group is strongly Arens irregular. In some cases, the same result was also obtained by the same author for the measure algebra in [101].

The factorization property (and some of its variants) was also used to study ideals and right cancellation in Banach algebras and semigroup compactifications with an Arens product such as $L^1(G)^{**}$, $\mathcal{LUC}(G)^*$ and the \mathcal{LUC} -compactification G^{LUC} of non-compact locally compact groups, $\ell^{\infty}(S)^*$ and the Stone-Čech compactification of infinite discrete semigroups S having some cancellation properties, $A(G)^{**}$ and $\mathcal{UC}_2(G)^*$ for a class of non-metrizable locally compact groups. See for example, [36], [45], [46], [47] and [48].

We define next a weaker factorization property, but we keep the same terminology. This weak η -factorization property was introduced and used in [44]. It was used, together with the Mazur property, to show that the Fourier algebra of a certain class of non-metrizable groups is strongly Arens irregular.

Definition 3.1. Let \mathcal{A} be a Banach algebra, η be a cardinal number and $X_1, ..., X_m$ be subsets of $\mathcal{LUC}(\mathcal{A})^*$ each with cardinality at least η . Then the Banach dual \mathcal{A}^* of \mathcal{A} is said to have the η -factorization property through $\prod_{k=1}^m X_k$ if for every bounded family of functionals $\{S_{\lambda} : \lambda < \eta\}$ contained in \mathcal{A}^* , there exist a bounded family $\{(\Psi_{\lambda k})_{k=1}^m : \lambda < \eta\}$ in $\prod_{k=1}^m X_k$ and a finite family of functionals $\{T_k : k = 1, ..., m\}$ in \mathcal{A}^* such that the factorization formula

(8)
$$S_{\lambda} = \sum_{k=1}^{m} \Psi_{\lambda k} \cdot T_{k}$$

holds for all $\lambda < \eta$.

If the elements $\{\Psi_{\lambda} = (\Psi_{\lambda k})_{k=1}^{m} : \lambda < \eta\}$ can be chosen in $\prod_{k=1}^{m} X_{k}$ independently of the elements S_{λ} , then we say that \mathcal{A}^{*} has the uniform η -factorization property. (This is going to be so in all the cases treated in the paper.)

Remarks 3.2.

(i) Putting $T = (T_k)_{k=1}^m \in \mathcal{A}^{*m}$, we may write the factorization formula (8) as

$$S_{\lambda} = \Psi_{\lambda} \cdot T$$
 for all $\lambda < \eta$.

(ii) The η-factorization property just defined is one-sided and should have been labeled "left" since we are using the left module action Ψ · T of ℒUC(A)* on A* (see 5). The mirror η-factorization property may also be defined using the action Ψ ⊙ T of 𝔅UC(A)* on A*. Noting that the left action Ψ ⊙ T of 𝔅UC(A)* on A* is the same as the right action T · Ψ of 𝔅UC(A)* on A* induced by the left action a · T of 𝔅 on A*, where ⟨a · T, b⟩ = ⟨T, ba⟩ for b ∈ 𝔅, we may say that 𝔅 has the right η-factorization property through Π^m_{k=1} X_k when the identity (8) in Definition 3.1 is changed to

$$S_{\lambda} = \sum_{k=1}^{m} \Psi_{\lambda k} \odot T_{k} = \sum_{k=1}^{m} T_{k} \cdot \Psi_{\lambda k} = T \cdot \Psi_{\lambda}$$

for all $\lambda < \eta$.

In order to apply this factorization property in Theorems 3.5 and 3.8 dealing with the topological centres of \mathcal{A}^{**} and $\mathcal{LUC}(\mathcal{A})^*$, we recall the following definition from [101].

Definition 3.3. Let \mathcal{E} be a Banach space and η be an infinite cardinal number.

- (i) A functional Ψ ∈ E^{**} is called η-weak^{*}-continuous if for every infinite net (S_λ)_λ in the unit ball of E^{*} of cardinality at most η with S_λ → 0 in the weak^{*}-topology, we have ⟨Ψ, S_λ⟩ → 0.
- (ii) We say that ε has the η-Mazur property if every η-weak*-continuous functional Ψ ∈ ε^{**} is an element of ε.

Notation 3.4. Throughout the paper, with the set $\prod_{k=1}^{m} X_k$ in $\mathcal{LUC}(\mathcal{A})^*$ through which the factorization of \mathcal{A}^* is made, we let $X = \bigcup_{k=1}^{m} X_k$ and \widetilde{X} be the set made of some fixed extensions of the elements in X to \mathcal{A}^* such that $\|\widetilde{\Psi}\| = \|\Psi\|$ as already mentioned in (7).

Theorems 3.5 and 3.8 below were proved by Neufang (see [101, Theorem 2.3 and Theorem 4.2]) with with m = 1, $\mathcal{S} = \mathcal{A}^{**}$ and X in the unit ball of $\mathcal{LUC}(\mathcal{A})^*$. The proof of the first theorem is a modification of the proof given in [101, Theorem 2.3], while we rely on [91] for the proof of the second theorem. Both proofs are included so that the role of the set $\prod_{k=1}^{m} X_k$ through which the factorization of \mathcal{A}^* is made for the determination of the topological centres becomes clear. This will be needed in our applications in later sections. For the proof of the second theorem as well as many other theorems in the paper, we shall need the right multiplier algebra $M_r(\mathcal{A})$ of \mathcal{A} , this is the algebra of all bounded operators R on \mathcal{A} satisfying

$$R(ab) = aR(b)$$
 for each a and b in \mathcal{A} .

The operation in $M_r(\mathcal{A})$ is the opposite of the operation \circ of composition of operators, i.e., $R_1R_2 = R_2 \circ R_1$.

We shall also prove in Theorems 3.6 and 3.10 the analogue of these results for the second toplogical centres. For this we need the set X to be in the unit ball of $\mathcal{RUC}(\mathcal{A})^*$ and we use the left multiplier algebra $M_l(\mathcal{A})$ of \mathcal{A} . A bounded operator on \mathcal{A} is a *left multiplier* of \mathcal{A} if L(ab) = L(a)b for $a, b \in \mathcal{A}$. The composition of operators is the operation in $M_l(\mathcal{A})$.

Recall also that in a Banach algebra \mathcal{A} , a bounded right approximate identity (brai for short) is a bounded net (e_{α}) with

$$\lim_{\alpha} \|ae_{\alpha} - a\| = 0 \quad \text{for each} \quad a \in \mathcal{A}.$$

Any weak^{*}-limit of a brai is a right identity in \mathcal{A}^{**} when \mathcal{A}^{**} has the first Arens product. Bounded left approximate identities (blai) are defined analogously. The net (e_{α}) is a bounded approximate identity (bai) if it is both brai and blai.

When \mathcal{A} has a bai bounded by 1, \mathcal{A} is a closed right ideal in $M_r(\mathcal{A})$ and a closed left ideal in $M_l(\mathcal{A})$. It will also be seen by Lemmas 3.7 and 3.9 that $M_r(\mathcal{A})$ and $M_l(\mathcal{A})$ are closed subalgebras of $\mathcal{LUC}(\mathcal{A})^*$ and $\mathcal{RUC}(\mathcal{A})^*$, respectively.

For more details on $M_r(\mathcal{A})$ and $M_l(\mathcal{A})$, see [15], particularly Theorems 2.5.12 and 2.9.49.

Theorem 3.5. Let \mathcal{A} be a Banach algebra, η be an infinite cardinal and $X_1, ..., X_m$ be subsets of $\mathcal{LUC}(\mathcal{A})^*$ each of cardinality at least η . Suppose that \mathcal{A} has the η -Mazur property and its dual \mathcal{A}^* has the η -factorization property through $\prod_{k=1}^m X_k$. Then

- (i) for any weak^{*}-closed subsemigroup S of A^{**} with $\widetilde{X} \subseteq S$, we have $T_1Z(S) = S \cap A$.
- (ii) In particular, $T_1Z(\mathcal{A}^{**}) = \mathcal{A}$, that is, \mathcal{A} is left strongly Arens irregular.

Proof. Let $\Psi \in T_1Z(\mathbb{S})$ and $(S_{\lambda})_{\lambda}$ be a net in the unit ball of \mathcal{A}^* of cardinality at most η with $S_{\lambda} \to 0$ in the weak*-topology. Factorize $(S_{\lambda})_{\lambda}$ through $\prod_{k=1}^{m} X_k$ as

$$S_{\lambda} = \sum_{k=1}^{m} \Psi_{\lambda k} \cdot T_k$$

for each $\lambda < \eta$, where $\{(\Psi_{\lambda k})_{k=1}^m : \lambda < \eta\}$ is a bounded net in $\prod_{k=1}^m X_k$ and $\{T_k : k = 1, ..., m\} \subset \mathcal{A}^*$. Let $\widetilde{\Psi_{\lambda k}}$ be the corresponding extensions in \widetilde{X} , and take m subnets of $(\widetilde{\Psi_{\lambda k}})_{\lambda < \eta}$ (indexed by the same set) with weak*-limit $\widetilde{\Psi_k}$ in S for each k = 1, ..., m. Denote for simplicity the m subnets also by $(\widetilde{\Psi_{\lambda k}})_{\lambda < \eta}$, k = 1, ..., m, and keep (S_{λ}) to denote the corresponding subnet of the original net in the unit ball of \mathcal{A}^* . It is then easy to check that $\sum_{k=1}^m \widetilde{\Psi_k} \cdot T_k$ is the weak*-limit of the subnet (S_{λ}) , and so it is zero. Since $\Psi \in T_1Z(S)$, using (7), this gives

(9)
$$\lim_{\lambda} \langle \Psi, S_{\lambda} \rangle = \lim_{\lambda} \left\langle \Psi, \sum_{k=1}^{m} \Psi_{\lambda k} . T_{k} \right\rangle = \lim_{\lambda} \sum_{k=1}^{m} \left\langle \Psi . \Psi_{\lambda k} , T_{k} \right\rangle$$
$$= \sum_{k=1}^{m} \lim_{\lambda} \left\langle \Psi \widetilde{\Psi_{\lambda k}} , T_{k} \right\rangle = \sum_{k=1}^{m} \left\langle \Psi \widetilde{\Psi_{k}} , T_{k} \right\rangle$$
$$= \sum_{k=1}^{m} \left\langle \Psi, \widetilde{\Psi_{k}} \cdot T_{k} \right\rangle = \left\langle \Psi, \sum_{k=1}^{m} \widetilde{\Psi_{k}} \cdot T_{k} \right\rangle = 0.$$

The η -Mazur property implies then that $\Psi \in \mathcal{A}$, as wanted.

Theorem 3.6. Let \mathcal{A} be a Banach algebra, η be an infinite cardinal and $X_1, ..., X_m$ be subsets of $\mathcal{RUC}(\mathcal{A})^*$ each of cardinality at least η . Suppose that \mathcal{A} has the η -Mazur property and its dual \mathcal{A}^* has the right η -factorization property through $\prod_{k=1}^m X_k$. Then

- (i) for any weak*-closed subsemigroup S of A** with X̃ ⊆ S, we have T₂Z(S) = S ∩ A.
- (ii) In particular, $T_2Z(\mathcal{A}^{**}) = \mathcal{A}$, that is, \mathcal{A} is right strongly Arens irregular.

Proof. Proceed as in the proof of Theorem 3.5.

For the proof of the second theorem we need the following lemma due essentially to Lau and Ülger in [91]. Let

$$I(\mathcal{A}) = \{ \Psi \in \mathcal{LUC}(\mathcal{A})^* : \mathcal{A}\Psi \subseteq \mathcal{A} \}.$$

Lemma 3.7. If \mathcal{A} is a Banach algebra with a bai (e_{α}) , then $I(\mathcal{A}) \subseteq T_1Z(\mathcal{LUC}(\mathcal{A})^*)$. When the bai is bounded by 1, $M_r(\mathcal{A}) \cong I(\mathcal{A})$.

Proof. It is clear that $I(\mathcal{A})$ is a norm-closed subalgebra of $\mathcal{LUC}(\mathcal{A})^*$. To see that $I(\mathcal{A}) \subseteq T_1 Z(\mathcal{LUC}(\mathcal{A})^*)$, let $\Psi \in I(\mathcal{A})$ and (Φ_α) be a net in $\mathcal{LUC}(\mathcal{A})^*$ with weak^{*}-limit Φ . Then for every $S \in \mathcal{A}^*$ and $a \in \mathcal{A}$, we have

$$\begin{split} \lim_{\alpha} \langle \Psi \Phi_{\alpha}, S.a \rangle &= \lim_{\alpha} \langle (a\Psi) \Phi_{\alpha}, S \rangle = \lim_{\alpha} \langle (a\Psi), \Phi_{\alpha} \cdot S \rangle = \lim_{\alpha} \langle \Phi_{\alpha}, S \cdot (a\Psi) \rangle \\ &= \langle \Phi, S \cdot (a\Psi) \rangle = \langle (a\Psi) \Phi, S \rangle = \langle \Psi \Phi, S.a \rangle, \end{split}$$

where the fourth equality is due to the fact that $a\Psi \in \mathcal{A}$. Therefore, Ψ is in the first topological centre of $\mathcal{LUC}(\mathcal{A})^*$, as wanted.

Next we proceed to identify $I(\mathcal{A})$ with $M_r(\mathcal{A})$. Note first that (e_α) is a weak^{*}convergent net in $\mathcal{LUC}(\mathcal{A})^*$ and has the identity in $\mathcal{LUC}(\mathcal{A})^*$ as its limit. To see this, take any weak^{*}-cluster point E of (e_α) in $\mathcal{LUC}(\mathcal{A})^*$. Since (e_α) is a brai, E is a right identity in $\mathcal{LUC}(\mathcal{A})^*$, and so E belongs in particular to $I(\mathcal{A})$. Since (e_α) is blai, we also have Ea = a for every $a \in \mathcal{A}$. Therefore $E\Psi = \Psi$ for every $\Psi \in \mathcal{LUC}(\mathcal{A})^*$ since $E \in I(\mathcal{A}) \subseteq T_1Z(\mathcal{LUC}(\mathcal{A})^*)$. Being the identity, E must be the unique weak^{*}-limit of (e_α) .

Now if $T \in M_r(\mathcal{A})$, then (Te_α) is a net in \mathcal{A} which is weak*-convergent in $\mathcal{LUC}(\mathcal{A})^*$. To see this, take two subnets (Te_β) and (Te_γ) with weak*-limits Ψ and Ψ' in $\mathcal{LUC}(\mathcal{A})^*$, respectively (this is possible since (Te_α) is bounded in \mathcal{A}).

Then again since (e_{β}) is a brai, we have

$$Te_{\gamma} = \lim_{\rho} T(e_{\gamma}e_{\beta}) \text{ for each } \gamma,$$

and so for every $S.a \in \mathcal{LUC}(\mathcal{A})$, we have

$$\langle \Psi, S.a \rangle = \langle E\Psi, S.a \rangle = \lim_{\gamma} \lim_{\beta} \langle e_{\gamma} T e_{\beta}, S.a \rangle = \lim_{\gamma} \lim_{\beta} \langle T(e_{\gamma} e_{\beta}), S.a \rangle$$
$$= \lim_{\gamma} \langle T e_{\gamma}, S.a \rangle = \langle \Psi', S.a \rangle,$$

as wanted. Denote the weak*-limit of (Te_{α}) by Ψ (note that Ψ depends on T). Consider now for each $\Phi \in I(\mathcal{A})$, the right multiplier $R_{\Phi} \in M_r(\mathcal{A})$ given by

 $R_{\Phi}(a) = a\Phi$ for every $a \in \mathcal{A}$

and consider the linear map

$$\mathfrak{I}: I(\mathcal{A}) \to M_r(\mathcal{A}), \quad \mathfrak{I}(\Phi) = R_{\Phi}.$$

Since, for every $T \in M_r(\mathcal{A})$ and $a \in \mathcal{A}$, we have

$$Ta = \lim_{\alpha} T(ae_{\alpha}) = \lim_{\alpha} aTe_{\alpha} = a\Psi = R_{\Psi}(a),$$

we see that \mathcal{I} is surjective (remember Ψ depends on T). Moreover

$$\|\mathfrak{I}(\Phi)\| \le \|\Phi\| = \|E\Phi\| = \lim_{\alpha} \|R_{\Phi}e_{\alpha}\| \le \|R_{\Phi}\| = \|\mathfrak{I}(\Phi)\|,$$

where the second inequality is due to the condition that (e_{α}) is bounded by 1. So when the bai is bounded by 1, $I(\mathcal{A})$ and $M_r(\mathcal{A})$ are isometrically isomorphic and $I(\mathcal{A}) \subseteq T_1 Z(\mathcal{LUC}(\mathcal{A})^*)$.

Theorem 3.8. Let \mathcal{A} be a Banach algebra with a bai, η be an infinite cardinal and $X_1, ..., X_m$ be subset of $\mathcal{LUC}(\mathcal{A})^*$ each with cardinality at least η . If \mathcal{A} has the η -Mazur property and its dual \mathcal{A}^* has the η -factorization property through $\prod_{k=1}^m X_k$, then

- (i) for any weak*-closed subsemigroup S of $\mathcal{LUC}(\mathcal{A})^*$ with $X \subseteq S$, we have $T_1Z(S) = S \cap I(\mathcal{A})$.
- (ii) If in addition the bai is bounded by 1, then T₁Z(S) is isometrically isomorphic to S ∩ M_r(A).

(iii) In particular, if the bai is bounded by 1, then $T_1Z(\mathcal{LUC}(\mathcal{A})^*)$ is isometrically isomorphic to the right multiplier algebra $M_r(\mathcal{A})$.

Proof. Let $\Psi \in T_1Z(S)$ and a be any element in \mathcal{A} . Let $(S_{\lambda})_{\lambda < \eta}$ be any net in \mathcal{A}^* with weak^{*}-limit 0 and let its elements be factorized as in the previous theorem as

$$S_{\lambda} = \sum_{k=1}^{m} \Psi_{\lambda k} \cdot T_k \quad \text{for each} \quad \lambda < \eta,$$

where $\{(\Psi_{\lambda k})_{k=1}^m : \lambda < \eta\}$ is a bounded net in $\prod_{k=1}^m X_k$ and $\{T_k : k = 1, ..., m\} \subset \mathcal{A}^*$. Take *m* subnets of $(\Psi_{\lambda k})_{\lambda} < \eta$ with weak*-limit Ψ_k in S for each k = 1, ..., m. Denote the subnets also by $(\Psi_{\lambda k})_{\lambda < \eta}$. It is then easy to check that $\sum_{k=1}^m \Psi_k \cdot T_k$ is the weak*-limit of the net $(S_{\lambda})_{\lambda}$, and so it is zero. Then, as in (9), since $\Psi \in T_1Z(S)$, we have

$$\begin{split} \lim_{\lambda} \langle a\Psi, S_{\lambda} \rangle &= \lim_{\lambda} \left\langle a\Psi, \sum_{k=1}^{m} \Psi_{\lambda k} \cdot T_{k} \right\rangle = \lim_{\lambda} \sum_{k=1}^{m} \langle (a\Psi)\Psi_{\lambda k}, T_{k} \rangle \\ &= \lim_{\lambda} \sum_{k=1}^{m} \langle \Psi\Psi_{\lambda k}, T_{k} \cdot a \rangle = \sum_{k=1}^{m} \lim_{\lambda} \langle \Psi\Psi_{\lambda k}, T_{k} \cdot a \rangle \\ &= \sum_{k=1}^{m} \langle \Psi\Psi_{k}, T_{k} \cdot a \rangle = \sum_{k=1}^{m} \langle a(\Psi\Psi_{k}), T_{k} \rangle \\ &= \langle a\Psi, \sum_{k=1}^{m} \Psi_{k} \cdot T_{k} \rangle = 0. \end{split}$$

The η -Mazur property implies then that $a\Psi \in \mathcal{A}$, and so $\Psi \in S \cap I(\mathcal{A})$. Lemma 3.7 finishes the proof of the theorem.

Let

$$J(\mathcal{A}) = \{ \Psi \in \mathfrak{RUC}(\mathcal{A})^* : \mathcal{A} \diamond \Psi \subseteq \mathcal{A} \}.$$

As in Lemma 3.7

Lemma 3.9. If \mathcal{A} is a Banach algebra with a bai (e_{α}) , then $J(\mathcal{A}) \subseteq T_2Z(\mathcal{RUC}(\mathcal{A})^*)$. When the bai is bounded by 1, $M_l(\mathcal{A}) \cong J(\mathcal{A})$.

Theorem 3.10. Let \mathcal{A} be a Banach algebra with a bai, η be an infinite cardinal and $X_1, ..., X_m$ be subset of $\mathcal{RUC}(\mathcal{A})^*$ each with cardinality at least η . If \mathcal{A} has the η -Mazur property and its dual \mathcal{A}^* has the right η -factorization property through $\prod_{k=1}^m X_k$, then

- (i) for any weak*-closed subsemigroup S of $\mathcal{RUC}(\mathcal{A})^*$ with $X \subseteq S$, we have $T_2Z(S) = S \cap J(\mathcal{A})$.
- (ii) If in addition the bai is bounded by 1, then T₂Z(S) is isometrically isomorphic to S ∩ M_l(A).
- (iii) In particular, if the bai bounded is by 1, then T₂Z(RUC(A)*) is isometrically isomorphic to the algebra M_l(A).

Proof. Proceed as in the proof of Lemma 3.7 and Theorem 3.8.

4. $F\ell^1(\eta)$ -bases in \mathcal{A} and factorizations of \mathcal{A}^* , $\mathcal{LUC}(\mathcal{A})$ and $M_r(\mathcal{A})^*$

In a given Banach algebra \mathcal{A} with a bai, we introduce next certain sets of cardinality η which will provide \mathcal{A}^* , as well as $M_r(\mathcal{A})^*$ in certain cases, with the η -factorization property. These sets are based on the standard ℓ^1 -bases, and are formed using some especially constructed sets in the multiplier algebra $M_r(\mathcal{A})$ together with an increasing family of sets in \mathcal{A} . We shall refer to them as $F\ell^1(\eta)$ -bases, where the letter "F" stands for factorization. The elements $\Psi_{\lambda k}$ needed for the factorization of operators in \mathcal{A}^* in Definition 3.1 are taken from the weak*-closure in $\mathcal{LUC}(\mathcal{A})^*$ of the sets constructed in the multiplier algebra $M_r(\mathcal{A})$.

In the three definitions given below, η is an infinite cardinal. We start in our first definition by adapting the standard definition of an ℓ^1 -base which was originally given by Rosenthal for sequences in [107].

Definition 4.1. $\ell^1(\eta)$ -BASE: Let η be an infinite cardinal. In a normed space \mathcal{E} , a bounded set $\mathcal{L} := \{a_\alpha : \alpha < \eta\}$ is an $\ell^1(\eta)$ -base with a constant K > 0 when the inequality

$$\sum_{n=1}^{p} |z_n| \le K \left\| \sum_{n=1}^{p} z_n a_{\alpha_n} \right\|$$

holds for all $p \in \mathbb{N}$ and for every possible choice of scalars $z_1, ..., z_p$ and (distinct) elements $a_{\alpha_1}, ..., a_{\alpha_p}$ in \mathcal{L} .

Remarks 4.2. If $\mathcal{L} := \{a_{\alpha} : \alpha < \eta\}$ is an $\ell^{1}(\eta)$ -base with a constant K > 0 in a normed space \mathcal{E} , then we note the following:

(i) For every function $c = (c_{\alpha})_{\alpha < \eta}$ in $\ell^{\infty}(\eta)$, there corresponds $T \in \mathcal{E}^*$ such that

$$\langle T, a_{\alpha} \rangle = c_{\alpha}$$
 for every $\alpha < \eta$.

This is obtained simply by letting

$$\left\langle T, \sum_{n=1}^{p} z_n a_{\alpha_n} \right\rangle = \sum_{n=1}^{p} z_n c_{\alpha_n} \quad \text{for every} \quad \sum_{n=1}^{p} z_n a_{\alpha_n} \in \langle \mathcal{L} \rangle,$$

and noting that

$$\left| \langle T, \sum_{n=1}^p z_n a_{\alpha_n} \rangle \right| = \left| \sum_{n=1}^p z_n c_{\alpha_n} \right| \le \|c\|_{\infty} \sum_{n=1}^p |z_n| \le \|c\|_{\infty} K \left\| \sum_{n=1}^p z_n a_{\alpha_n} \right\|.$$

Therefore we have a continuous linear functional $\langle \mathcal{L} \rangle \to \mathbb{C}$, which may be extended by Hahn-Banach Theorem to an element in \mathcal{E}^* .

- (ii) The closed linear span of an $\ell^1(\eta)$ -base is isomorphic to $\ell^1(\eta)$.
- (iii) The elements a_{α} in \mathcal{L} satisfy $||a_{\alpha}|| \geq \frac{1}{K}$ for every $\alpha < \eta$.

Next we relax a bit the definition of $\ell^1(\eta)$ -bases. This will enable us to include more algebras that are strongly Arens irregular in our study. Observe that if $\mathcal{L}_{\alpha} = \{a_{\alpha}\}$ and for some constant K > 0, the inequality $||a_{\alpha}|| \geq \frac{1}{K}$ holds for every $\alpha < \eta$, then next definition is precisely Definition 4.1. For,

$$\sum_{n=1}^{p} |z_n| = \sum_{n=1}^{p} \frac{1}{\|a_{\alpha_n}\|} \|z_n a_{\alpha_n}\| \le K \sum_{n=1}^{p} \|z_n a_{\alpha_n}\| \le K \left\| \sum_{n=1}^{p} z_n a_{\alpha_n} \right\|$$

The condition that $||a_{\alpha}|| \geq \frac{1}{K}$ for some constant K > 0 and other similar conditions (see (F2) and (F3) in the definitions below) will be crucial in the study. The conditions are automatically satisfied when we are dealing with ℓ^1 -bases by Remark 4.2 (iii). In general, however, as the reader will see, these conditions are the thorns in the investigation.

Definition 4.3. $\ell^1(\eta)$ -SELECTIVE BASE: In a normed space \mathcal{E} , we say that a bounded set \mathcal{L} is an $\ell^1(\eta)$ -selective base with constant $K_1 > 0$ when it is of the form $\bigcup_{\alpha < \eta} \mathcal{L}_{\alpha}$, where $\{\mathcal{L}_{\alpha} : \alpha < \eta\}$ is a family of nonempty subsets of \mathcal{E} such that

(i) the spans $\langle \mathcal{L}_{\alpha} \rangle$, $\alpha < \eta$, have zero pairwise intersections,

(ii) the inequality
(10)
$$\sum_{n=1}^{p} ||a_{n}|| \leq K_{1} \left\| \sum_{n=1}^{p} a_{n} \right\|$$
holds for every $\sum_{n=1}^{p} a_{n} \in \langle \mathcal{L} \rangle$, where $a_{n} \in \langle \mathcal{L}_{\alpha_{n}} \rangle$ for each $n = 1$,

Remarks 4.4.

(i) Clearly every $\ell^1(\eta)$ -base is an $\ell^1(\eta)$ -selective base since we have

$$\sum_{n=1}^{p} \|a_n\| = \sum_{n=1}^{p} \|z_n a_{\alpha_n}\| \le \sup_{\alpha < \eta} \|a_\alpha\| \sum_{n=1}^{p} |z_n| \le K \sup_{\alpha < \eta} \|a_\alpha\| \left\| \sum_{n=1}^{p} z_n a_{\alpha_n} \right\| = K \sup_{\alpha < \eta} \|a_\alpha\| \left\| \sum_{n=1}^{p} a_n \right\|$$

for every $a_n = z_n a_{\alpha_n} \in \langle a_{\alpha_n} \rangle$, n = 1, ..., p. (ii) Let $\mathcal{L} = \bigcup_{i=1}^{n} \mathcal{L}_{\alpha}$ be an $\ell^1(\eta)$ -selective base in a normed space \mathcal{E} . Pick for each $\alpha < \eta$, $a_{\alpha} \in \mathcal{L}_{\alpha}$ and consider the set $\{a_{\alpha} : \alpha < \eta\}$. Then, as in Remark 4.2(i), every function $c = (c_{\alpha})_{\alpha < \eta}$ in $\ell^{\infty}(\eta)$ defines a continuous linear functional $T \in \mathcal{E}^*$ such that

$$\langle T, a_{\alpha} \rangle = c_{\alpha}$$
 for every $\alpha < \eta$.

Lemma 4.5. Let \mathcal{E} be a Banach space with an $\ell^1(\eta)$ -selective base $\mathcal{L} = \bigcup_{\alpha < \eta} \mathcal{L}_{\alpha}$ with constant $K_1 > 0$. Let, for each $\alpha < \eta$, $\mathcal{M}_{\alpha} = \overline{\langle \mathcal{L}_{\alpha} \rangle}$. Then, for each $(S_{\alpha})_{\alpha < \eta} \in$ $\oplus_{\infty}\mathcal{M}^*_{\alpha}$, there exists $T \in \mathcal{E}^*$ such that $T_{|\mathcal{M}_{\alpha}} = S_{\alpha}$.

Proof. Let $(S_{\alpha})_{\alpha < \eta} \in \bigoplus_{\infty} \mathcal{M}_{\alpha}^*$. Since the spans $\langle \mathcal{L}_{\alpha} \rangle$ have zero pairwise intersections, each $x \in \langle \mathcal{L} \rangle$ is of the form $\sum_{n=1}^{p} a_n$ for some $p \in \mathbb{N}$ and $a_n \in \langle \mathcal{L}_{\alpha_n} \rangle$, n = 1, ..., p, and so we may naturally define T on $\langle \mathcal{L} \rangle$ by

$$\langle T, \sum_{n=1}^{p} a_n \rangle = \sum_{n=1}^{p} \langle S_{\alpha_n}, a_n \rangle.$$

The map T is well defined and linear on $\langle \mathcal{L} \rangle$. To check that it is bounded, let $x = \sum_{n=1}^{p} a_n \in \langle \mathcal{L} \rangle$, where $a_n \in \mathcal{L}_{\alpha_n}$ for each n = 1, ..., p. Then

$$|\langle T, x \rangle| = \left| \langle T, \sum_{n=1}^{p} a_n \rangle \right| = \left| \sum_{n=1}^{p} \langle S_{\alpha_n}, a_n \rangle \right|$$

$$\leq \sup_{\alpha} \|S_{\alpha}\| \sum_{n=1}^{p} \|a_n\|$$

$$\leq K_1 \sup_{\alpha} \|S_{\alpha}\| \|x\|,$$

where the last inequality is due to the fact that \mathcal{L} is an $\ell^1(\eta)$ -selective base with constant K_1 . This shows that T is bounded on $\langle \mathcal{L} \rangle$, and so it has a bounded extension to $\overline{\mathcal{L}}$. Extend it then to a bounded functional, denoted again by T, to the whole space by Hahn-Banach theorem. By definition, it is clear that $T_{|\mathcal{M}_{\alpha}} = S_{\alpha}$.

Now we come to the main definitions needed in the paper. Recall that \mathcal{A} is a right ideal in $M_r(\mathcal{A})$ when \mathcal{A} is faithful, so in the definitions below, the sets $A_{\alpha}a_{\alpha}^{\lambda}$ and $A_{\alpha}a_{\alpha}$ are contained in \mathcal{A} for each $\alpha < \eta$ and k = 1, ..., m.

 $\dots, p.$

Definition 4.6. In a Banach algebra \mathcal{A} with a bai, a zooming set is a set of the form $\mathcal{L} = \bigcup A_{\alpha} a_{\alpha}$, where

- (i) {A_α}_{α<η} is an increasing family of subsets of A such that A_η = U_{α<η} A_α is bounded (we may assume the bound is 1) with its closed linear span equal to A,
- (ii) $\{a_{\alpha} : \alpha < \eta\}$ is a bounded set in the multiplier algebra $M_r(\mathcal{A})$,
- (iii) the spans $\langle A_{\alpha}a_{\alpha}\rangle$, $\alpha < \eta$, have zero pairwise intersections.

Definition 4.7. $F\ell^1(\eta)$ -BASE OF TYPE 1. In a Banach algebra \mathcal{A} with a bai, a zooming $\ell^1(\eta)$ -base is called an $F\ell^1(\eta)$ -base of type 1.

Note that when \mathcal{L} is an $\mathbb{F}\ell^1(\eta)$ -base of type 1, then the property $1 \leq K ||aa_{\alpha}||$ for every $\alpha < \eta$ satisfied by the elements $a \in A_{\alpha}$, $\alpha < \eta$ (see Remark 4.2 (iii)), is equivalent to

(F2)
$$||a|| \le K ||aa_{\alpha}||$$
 for every $a \in \langle A_{\alpha} \rangle$ and $\alpha < \eta$.

For, since we are assuming that A_{η} is bounded by 1, we find that for every scalars $z_1, ..., z_p$ and (distinct) elements $a_1, ..., a_p$ in A_{α} ,

$$\left\|\sum_{1}^{p} z_{n} a_{n}\right\| \leq \sum_{n=1}^{1} |z_{n}| \leq K \left\|\sum_{n=1}^{p} z_{n} a_{n} a_{\alpha}\right\|.$$

In other words,

 $||a|| \le K ||aa_{\alpha}|| \quad \text{for every} \quad a \in \langle A_{\alpha} \rangle.$

In our applications to the algebraic structure and the topological centres of the Banach algebras with an Arens product, we will see that $F\ell^1(\eta)$ -base of type 1 are available in the group algebra of infinite discrete groups and the semigroup algebra of some infinite discrete semigroups as well as their weighted versions when the weight is diagonally bounded. But in general, these $F\ell^1(\eta)$ -bases are not available. The next definition relaxes the condition on being an $\ell^1(\eta)$ -base to an $\ell^1(\eta)$ -selective base having Property (F2). This second type of $F\ell^1(\eta)$ -base will be found in more general situations.

Definition 4.8. $F\ell^1(\eta)$ -BASE OF TYPE 2. In a Banach algebra \mathcal{A} with a bai, an $F\ell^1(\eta)$ -base of type 2 is a zooming $\ell^1(\eta)$ -selective base (with constant $K_1 > 0$) such that for some constant $K_2 > 0$,

(F2) $||a|| \le K_2 ||aa_{\alpha}||$ for every $a \in \langle A_{\alpha} \rangle$ and $\alpha < \eta$.

In certain cases (see Subsection 8.2), although zooming $\ell^1(\eta)$ -selective bases are available, we are not able to check the validity of Inequality (F2) above. This has forced us to relax this inequality to Condition (F3) below, which is easier to have. Condition (iii) in Definition 4.6 for the zooming property means in the definition below that the spans $\langle A_{\alpha} \rangle a_{\alpha}^k$ have zero pairwise intersections for each k = 1, ..., m.

Definition 4.9. $F\ell^1(\eta)$ -BASE OF TYPE 3. In a Banach algebra \mathcal{A} with a bai, an $F\ell^1(\eta)$ -base of type 3 is a zooming $\ell^1(\eta)$ -selective base (with constant $K_1 > 0$) of the form

$$\mathcal{L} = \bigcup_{\alpha < \eta} \bigcup_{k=1}^{m} A_{\alpha} a_{\alpha}^{k} = \bigcup_{\alpha < \eta} A_{\alpha} \{ a_{\alpha}^{k} : k = 1, ...m \},$$

where $\{\mathbf{a}_{\alpha} = (a_{\alpha}^{k})_{k=1}^{m} : \alpha < \eta\}$ is a bounded set in $M_{r}(\mathcal{A})^{m}$ such that for some $K_{2} > 0$,

(F3)
$$||a|| \le K_2 ||a\mathbf{a}_{\alpha}||$$
 for every $a \in \langle A_{\alpha} \rangle$ and $\alpha < \eta$.

Here $a\mathbf{a}_{\alpha} = (aa^k_{\alpha})^m_{k=1} \in \mathcal{A}^m$ and $||a\mathbf{a}_{\alpha}||$ is any of the p-norms of the vector $a\mathbf{a}_{\alpha}$ in $M_r(\mathcal{A})^m$, $p = 1, 2, ..., \infty$.

Remarks 4.10.

(i) The importance of Conditions (F2) and (F3) is not a mere surmise. Already with m = 1, the condition yields the inequality

$$\frac{1}{K_2} \|\Phi\| \le \|\Phi\Psi\| \le \|\Phi\| \quad \text{for every} \quad \Phi \in \mathcal{A}^{**}$$

whenever Ψ is weak*-cluster point of (a_{α}) . When $K_2 = 1$, these elements satisfy $\|\Phi\Psi\| = \|\Phi\|$ for every $\Phi \in \mathcal{A}^{**}$, they are called "right isometries" in the literature. They were essential in each of the proofs on the first topological centres of $\mathcal{LUC}(G)^*$ and $L^1(G)^{**}$ given in the past by various authors, see for instance [62], [80], [83], [50] and [9]. They came again in [49] studying the first topological centres of weighted algebras, with the constant K_2 as the diagonal bound of the weight (and so not necessarily equal to 1). Further details can be found in [50].

The condition (F3) (and so (F2) when m = 1) is essential and needed right at the beginning at Inequality (13) in the proof of the factorization Theorem 4.14.

(ii) The right $F\ell^1(\eta)$ -bases are defined similarly as in the three definitions above but with $a^k_{\alpha}A_{\alpha}$ and $a^k_{\alpha} \in M_l(\mathcal{A}), \ \alpha < \eta$ and k = 1, ..., m.

Notation 4.11.

(i) We use the term Fℓ¹(η)-base to designate any of the three types of Fℓ¹(η)-bases. For an Fℓ¹(η)-base in A, let η have the discrete topology and βη be the Stone-Čech compactification of η and let C(η) be the set of all cofinal ultrafilters on η. Recall that an ultrafilter p on η is cofinal if every set I in p is cofinal in η, that is, sup I = η (cf. [111]). By a result of van Douwen [111]

$$|\mathfrak{C}(\eta)| = 2^{2^{\eta}}.$$

Since for each k = 1, ..., m, the set $\{a_{\alpha}^k : \alpha < \eta\}$ is bounded in $M_r(\mathcal{A})$, the bounded map

(11)

$$\eta \to \mathcal{LUC}(\mathcal{A})^* \colon \quad \alpha \mapsto a_\alpha^k$$

has a continuous extension to $\beta\eta$. So for each k = 1, ..., m and each ultrafilter p in $\beta\eta$, there corresponds an element Ψ_{pk} in $\mathcal{LUC}(\mathcal{A})^*$. (If the elements a^k_{α} are bounded by 1, then the elements Ψ_{pk} are in the closed unit ball in $\mathcal{LUC}(\mathcal{A})^*$.) For each k = 1, ...m, we may consider the sets in the unit ball of $\mathcal{LUC}(\mathcal{A})^*$ given by

$$X_k = \{\Psi_{pk} : p \in \mathcal{C}(\eta)\}$$
 and let $X = \bigcup_{k=1}^m X_k$.

- (ii) When we deal with \mathcal{A}^{**} , as already agreed \widetilde{X} is the set made of fixed extensions of the elements in X to \mathcal{A}^* .
- (iii) When our concern is with $M_r(\mathcal{A})$, instead of the map (11), we consider the bounded map

$$\eta \to M_r(\mathcal{A})^{**}: \quad \alpha \mapsto a_\alpha^k,$$

and its extension to $\beta\eta$. So our elements Ψ_{pk} corresponding to the cofinal ultrafilters, as well as the set X, shall be regarded in the unit ball of $M_r(\mathcal{A})^{**}$.

Remarks 4.12.

(i) As already noted in Remark 4.4(i), every l¹(η)-base is an l¹(η)-selective base. Moreover, as noted after Definition 4.7, Condition (F2) in Definition 4.8 is automatically satisfied for an Fl¹(η)-base of type 1. Accordingly, an Fl¹(η)-base of type 1 is also of type 2.

The converse is however not true in general. The $F\ell^1(\eta)$ -bases found in Theorem 6.1 below are of type 2 but they are not usually of type 1 since they are not $\ell^1(\eta)$ -bases in general. To see this, let in Theorem 6.1 our group be \mathbb{R} and keep the same notation. With no loss of generality, we may just consider $V_0 =] -1, 1[, x_0 = 0$ the identity in \mathbb{R} , and consider the sequence $(\varphi_n)_{n < w}$ in A_0 be given by

$$\varphi_n(x) = \begin{cases} \frac{1}{2} - \frac{1}{n}, & \text{if } 0 \le x \le \frac{1}{2} - \frac{1}{n} \\ x, & \text{if } \frac{1}{2} - \frac{1}{n} \le x \le \frac{1}{2} + \frac{1}{n} \\ \frac{1}{2} + \frac{1}{n}, & \text{if } \frac{1}{2} + \frac{1}{n} \le x \le 1 \\ 0, & \text{otherwise.} \end{cases}$$

Then for every odd natural number p,

$$\sum_{n=1}^{p} (-1)^{n+1} \varphi_n(x) = \begin{cases} 1 - \sum_{n=1}^{p} (-1)^{n+1} \frac{1}{n}, & \text{if } 0 \le x \le \frac{1}{2} - \frac{1}{n} \\ x, & \text{if } \frac{1}{2} - \frac{1}{n} \le x \le \frac{1}{2} + \frac{1}{n} \\ 1 + \sum_{n=1}^{p} (-1)^{n+1} \frac{1}{n}, & \text{if } \frac{1}{2} + \frac{1}{n} \le x \le 1 \\ 0, & \text{otherwise,} \end{cases}$$

and so $\left|\left|\sum_{n=1}^{p}(-1)^{n+1}\varphi_{n}\right|\right|_{1}$ is close to 1 when p is sufficiently large. Therefore, there no K>0 for which

$$p \le K \left\| \sum_{n=1}^p (-1)^{n+1} \varphi_n \right\|_1$$

for every odd natural number p.

(ii) If together with the bounded set $\{\mathbf{a}_{\alpha} = (a_{\alpha}^{k})_{k=1}^{m} : \alpha < \eta\}$ in $M_{r}(\mathcal{A})^{m}$, there is another bounded set $\{\mathbf{b}_{\alpha} = (b_{\alpha}^{k})_{k=1}^{m} : \alpha < \eta\}$ in $M_{r}(\mathcal{A})^{m}$ such that

(f3)
$$\mathbf{a}_{\alpha}\mathbf{b}_{\alpha} = \sum_{k=1}^{m} a_{\alpha}^{k} b_{\alpha}^{k} = 1$$
 (the identity in $M_{r}(\mathcal{A})$) for each $\alpha < \eta$,

then Condition (F3) holds easily since

$$||a|| = ||a\mathbf{a}_{\alpha}\mathbf{b}_{\alpha}|| \le ||a\mathbf{a}_{\alpha}||_{p} ||\mathbf{b}_{\alpha}||_{q} \le K ||a\mathbf{a}_{\alpha}||_{p} \quad \text{for every} \quad a \in \langle A_{\alpha} \rangle \text{ and } \alpha < \eta,$$

where $\frac{1}{p} + \frac{1}{q} = 1$ when p > 1, $q = \infty$ when p = 1, and $K = \sup_{\alpha} \|\mathbf{b}_{\alpha}\|_{q}$.

Therefore a zooming $\ell^1(\eta)$ -selective base with Condition (f3) is an $F\ell^1(\eta)$ base with constant $K_2 = \sup_{\alpha} ||\mathbf{b}_{\alpha}||$, it is of type 2 if m = 1 and of type 3 if m is any positive integer.

Conditions (F3) and (f3) are however not equivalent in general (even for m = 1) as Condition (F3) holds in each of the cases studied in Sections 6, 7 and 8, but Condition (f3) is clearly not true in many situations starting with Theorem 6.2 when G is not a group.

(iii) In most examples in our application sections 6, 7 and 8, we will be needing just the lighter Definitions 4.7 and 4.8. The rather technical conditions in Definition 4.9 were forced into our approach because of Theorem 8.5 and the results related to this theorem, where we cannot find $F\ell^1(\eta)$ -bases of type 1 or 2.

Remark 4.13. Let $\mathcal{L} = \bigcup_{\alpha < \eta} A_{\alpha} \{ a_{\alpha}^{k} : k = 1, ..., m \}$ be an $\mathbb{F}\ell^{1}(\eta)$ -base in \mathcal{A} , and suppose that \mathcal{L} is of type 1 or 2 if m = 1, and it is of type 3, otherwise. Fix $a \in A_{0}$. Then $aa_{\alpha}^{k} \in A_{\alpha}a_{\alpha}^{k}$ for each $\alpha < \eta$ and each k = 1, ..., m. Since $\bigcup_{\alpha < \eta} A_{\alpha}a_{\alpha}^{k}$ is an $\ell^{1}(\eta)$ -selective base, it follows from Remark 4.4(ii), that for each k = 1, ..., m, every function $c = (c_{\alpha})_{\alpha < \eta}$ in $\ell^{\infty}(\eta)$ defines a continuous linear functional $T_{k} \in \mathcal{A}^{*}$ such that

$$\langle T_k, aa^k_\alpha \rangle = c_\alpha \quad \text{for every} \quad \alpha < \eta.$$

In other words, for each k = 1, ..., m, the function $c \in \ell^{\infty}(\eta)$ defines an operator $T_k \cdot a$ in $\mathcal{LUC}(\mathcal{A})$ such that

$$\langle T_k \cdot a, a_{\alpha}^k \rangle = \langle T_k, aa_{\alpha}^k \rangle = c_{\alpha} \text{ for every } \alpha < \eta.$$

Similarly, if \mathcal{L} is a right $\mathbb{F}\ell^1(\eta)$ -base in \mathcal{A} , the function $c \in \ell^{\infty}(\eta)$ defines for each k an operator $T_k \odot a = a \cdot T_k$ in $\mathcal{RUC}(A)$ such that

$$\langle a \cdot T_k, a_{\alpha}^k \rangle = \langle T_k, a_{\alpha}^k a \rangle = c_{\alpha}$$
 for every $\alpha < \eta$.

Theorem 4.14. Let \mathcal{A} be a Banach algebra with a bai and an $F\ell^1(\eta)$ -base for some infinite cardinal η and some $m \in \mathbb{N}$, and let $X_1, ..., X_m$ be the subsets in $\mathcal{LUC}(\mathcal{A})^*$ as defined in Notation 4.11. Then \mathcal{A}^* has the uniform η -factorization property through $\prod_{k=1}^m X_k$.

Proof. We prove first that \mathcal{A}^* has the 1-factorization property, i.e., for every $S \in \mathcal{A}^*$, there exists a family of functionals $\{T_k : k = 1, ..., m\}$ in \mathcal{A}^* such that the factorization formula

(12)
$$\Psi \cdot T = \sum_{k=1}^{m} \Psi_k \cdot T_k = S$$

holds for every element $(\Psi_k)_{k=1}^m$ in $\prod_{k=1}^m X_k$, where m = 1 if the $F\ell^1(\eta)$ -base is of type 1 or 2, and m is a finite integer if the $F\ell^1(\eta)$ -base is of type 3.

Let our $F\ell^1(\eta)$ -base \mathcal{L} have the form $\bigcup_{k=1}^m \bigcup_{\alpha < \eta} A_\alpha a_\alpha^k$, and consider in the Banach algebra \mathcal{A}^m (the operations are defined coordinatewise), the set $\mathbf{L} = \bigcup_{\alpha < \eta} A_\alpha \mathbf{a}_\alpha$, where $A_\alpha \mathbf{a}_\alpha = \{a\mathbf{a}_\alpha = (aa_\alpha^1, ..., aa_\alpha^m) : a \in A_\alpha\}$. Since \mathcal{L} is an $\ell^1(\eta)$ -selective base in \mathcal{A} , it is easy to check that \mathbf{L} is an $\ell^1(\eta)$ -selective base in \mathcal{A}^m .

As in Lemma 4.5, we consider in \mathcal{A}^m the subspaces

$$\mathcal{M}_{\alpha} = \overline{\langle A_{\alpha} \mathbf{a}_{\alpha} \rangle} = \overline{\langle A_{\alpha} \rangle} \mathbf{a}_{\alpha} \text{ for each } \alpha < \eta.$$

Let S be any element in \mathcal{A}^* . Then, for each $\alpha < \eta$, define S^{α} on $A_{\alpha} \mathbf{a}_{\alpha}$ by setting

$$\langle S^{\alpha}, a\mathbf{a}_{\alpha} \rangle = \langle S, a \rangle$$
 for each $a \in A_{\alpha}$.

(We are changing the notation S_{α} used in Lemma 4.5 to S^{α} to avoid confusion with the operators in the family $\{S_{\lambda}\}_{\lambda < \eta}$ coming up later on in the proof.) Then, we have

(13)
$$|\langle S^{\alpha}, a\mathbf{a}_{\alpha}\rangle| = |\langle S, a\rangle| \le ||S|| ||a|| \le K_2 ||S|| ||a\mathbf{a}_{\alpha}||$$
 for every $a \in \langle A_{\alpha}\rangle$,

where $K_2 > 0$ is the constant required by Inequality (F2) if m = 1 and by (F3) if $m \in \mathbb{N}$ is arbitrary.

Thus, each S^{α} is linear and bounded on $\langle A_{\alpha} \mathbf{a}_{\alpha} \rangle$, and so it may be extended to an element in \mathcal{M}^*_{α} which we denote again by S^{α} . We also deduce from the inequality above that $(S^{\alpha})_{\alpha < \eta} \in \bigoplus_{\infty} \mathcal{M}^*_{\alpha}$. Since **L** is an $\ell^1(\eta)$ -selective base in \mathcal{A}^m , it follows

from Lemma 4.5, there is $T = (T_k)_{k=1}^m \in \mathcal{A}^{m^*} = \mathcal{A}^{*m}$ such that $T_{|\mathcal{M}_{\alpha}|} = S^{\alpha}$ for every $\alpha < \eta$, that is,

(14) $\langle T, a\mathbf{a}_{\alpha} \rangle = \langle S^{\alpha}, a\mathbf{a}_{\alpha} \rangle = \langle S, a \rangle$, for every $a \in A_{\alpha}$ and $\alpha < \eta$, where

(15)
$$||T|| \le K_1 \sup ||S^{\alpha}|| \le K_1 K_2 ||S||.$$

Let now $\Psi = (\Psi_k)_{k=1}^m$ be in $\prod_{k=1}^m X_k$, We claim that

$$\langle \sum_{k=1}^{m} \Psi_k \cdot T_k, a \rangle = \langle \Psi \cdot T, a \rangle = \langle S, a \rangle \quad \text{for every} \quad a \in \mathcal{A}.$$

Pick a net from $\{\mathbf{a}_{\alpha} : \alpha < \eta\}$ that weak*-converges to Ψ in $(\mathcal{LUC}(\mathcal{A})^*)^m$. This net may be written as $(\mathbf{a}_{h(\gamma)})_{\gamma \in \Lambda}$, where Λ is a directed set and $h : \Lambda \to \eta$ is a monotone cofinal function. Let a be arbitrary in A_{η} . Then $a \in A_{\beta}$ for some $\beta < \eta$, and so $a \in A_{\alpha}$ for every $\alpha \geq \beta$. Accordingly,

(16)
$$\langle \Psi \cdot T, a \rangle = \langle \Psi, T \cdot a \rangle = \lim_{\gamma} \langle \mathbf{a}_{h(\gamma)}, T \cdot a \rangle = \lim_{\gamma} \langle T, a \mathbf{a}_{h(\gamma)} \rangle.$$

Note now that if $\gamma \in \Lambda$ is such that $h(\gamma) \geq \beta$, then $a\mathbf{a}_{h(\gamma)} \in A_{h(\gamma)}\mathbf{a}_{h(\gamma)}$, and so (14) implies that

$$\langle T, a\mathbf{a}_{h(\gamma)} \rangle = \langle S^{h(\gamma)}, a\mathbf{a}_{h(\gamma)} \rangle = \langle S, a \rangle.$$

Thus, continuing the argument in (16), we get

 $\langle \Psi \cdot T, a \rangle = \lim_{\gamma} \langle T, a \mathbf{a}_{h(\gamma)} \rangle = \langle S, a \rangle$ for every $a \in A_{\eta}$.

Since $\sum_{k=1}^{m} \Psi_k \cdot T_k$ and S agree on A_η and the linear span of A_η is dense in \mathcal{A} , we see that $\sum_{k=1}^{m} \Psi_k \cdot T_k$ and S are equal.

We prove now that \mathcal{A}^* has the uniform η -factorization property. So let $\mathcal{S} = \{S_{\lambda}\}_{\lambda < \eta}$ be a bounded family of functionals in \mathcal{A}^* . Then partition η into η many subsets I_{λ} such that each I_{λ} is cofinal in η (see [111, Lemma, page 61]). We see our $\mathbb{F}\ell^1(\eta)$ -base as

$$\mathcal{L} = \bigcup_{\lambda < \eta} \bigcup_{\alpha \in I_{\lambda}} A_{\alpha} \{ a_{\alpha}^{k} : k = 1, ..., m \}$$

and its corresponding $\ell^1(\eta)$ -selective base in \mathcal{A}^m as

$$\mathbf{L} = \bigcup_{\lambda < \eta} \bigcup_{\alpha \in I_{\lambda}} A_{\alpha} \mathbf{a}_{\alpha}.$$

Note, that for each $\lambda < \eta$, the sets $A_{\alpha} \mathbf{a}_{\alpha}$, $\alpha \in I_{\lambda}$, have zero pairwise intersections, and the family $\{A_{\alpha}\}_{\alpha \in I_{\lambda}}$ is increasing with $A_{\eta} = \bigcup_{\alpha \in I_{\lambda}} A_{\alpha}$. So we may proceed as in the previous case with the 1-factorization, but this time by taking for every $\lambda < \eta$, the element $\Psi_{\lambda} = (\Psi_{\lambda k})_{k=1}^{m}$ in $\prod_{k=1}^{m} X_{k}$ as any weak*-cluster point of the set $\{\mathbf{a}_{\alpha} : \alpha \in I_{\lambda}\}$, i.e., based on our Notation 4.11, $\Psi_{\lambda} = \Psi_{p}$ for some cofinal ultrafilter p having I_{λ} as a member. Then let $\{\mathbf{a}_{h_{\lambda}(\gamma)}\}_{\gamma \in \Lambda}$ be the net converging to Ψ_{λ} (here $h_{\lambda} : \Lambda \to I_{\lambda}$ is monotone and cofinal.)

Then, define for each $\alpha \in I_{\lambda}$ and $\lambda < \eta$, S^{α} on $A_{\alpha}\mathbf{a}_{\alpha}$ by

(17)
$$\langle S^{\alpha}, a\mathbf{a}_{\alpha} \rangle = \langle S_{\lambda}, a \rangle$$
 for every $a \in A_{\alpha}$

Again applying Lemma 4.5, we obtain as in (14), a functional $T = (T_k)_{k=1}^m \in \mathcal{A}^{*m}$ such that

(18)
$$\langle T, a\mathbf{a}_{\alpha} \rangle = \langle S^{\alpha}, a\mathbf{a}_{\alpha} \rangle = \langle S_{\lambda}, a \rangle$$
 for every $a \in A_{\alpha}, \ \alpha \in I_{\lambda}$ and $\lambda < \eta$.

We claim that

$$\Psi_{\lambda} \cdot T = S_{\lambda} \quad \text{for every} \quad \lambda < \eta$$

So fix $\lambda < \eta$, let $a \in A_{\eta}$ be arbitrary and pick $\beta < \eta$ with $a \in A_{\beta}$. Then there exists $\gamma_0 \in \Lambda$ such that $h_{\lambda}(\gamma) \ge \beta$ for every $\gamma \ge \gamma_0$ (possible since I_{λ} is cofinal in η), and so $a \in A_{h_{\lambda}(\gamma)}$ for every $\gamma \ge \gamma_0$. Since $h_{\lambda}(\gamma) \in I_{\lambda}$, we deduce from (18), that

$$\langle \Psi_{\lambda} \cdot T, a \rangle = \lim_{\gamma} \langle \mathbf{a}_{h_{\lambda}(\gamma)} \cdot T, a \rangle = \lim_{\gamma} \langle T, a \mathbf{a}_{h_{\lambda}(\gamma)} \rangle = \langle S_{\lambda}, a \rangle \quad \text{for every} \quad a \in \mathcal{A}_{\eta}.$$

As previously with the 1-factorization, the wanted factorization

$$\langle \sum_{k=1}^{m} \Psi_{\lambda k} \cdot T_k, a \rangle = \langle \Psi_{\lambda} \cdot T, \mathbf{a} \rangle = \langle S_{\lambda}, a \rangle \quad \text{for every} \quad a \in \mathcal{A}$$

follows for every $\lambda < \eta$.

Remarks 4.15.

(i) Let for some $\lambda < \eta$, I_{λ} be one of the cofinal subsets from the partition of η . Then instead of the element S^{α} in (17) defined on A_{α} for $\alpha < \eta$, we may define S^{λ} on $\mathcal{L} = \bigcup_{\alpha < \eta} A_{\alpha} \mathbf{a}_{\alpha}$ by

(19)
$$\langle S^{\lambda}, a\mathbf{a}_{\alpha} \rangle = \begin{cases} \langle S_{\lambda}, a \rangle, & \text{if } \alpha \in I_{\lambda} \\ 0, & \text{if } \alpha \notin I_{\lambda}, \end{cases}$$

i.e., starting in (17) with $S_{\mu} = 0$ for every $\mu \neq \lambda$.

If this is the case, then the operator $T = (T_k)_{k=1}^m \in \mathcal{A}^{*m}$ comes out defined on $\bigcup_{\alpha < n} A_\alpha \mathbf{a}_\alpha$ as

$$\langle T, a\mathbf{a}_{\alpha} \rangle = \begin{cases} \langle S^{\lambda}, a\mathbf{a}_{\alpha} \rangle = \langle S_{\lambda}, a \rangle & \text{for every} \quad \alpha \in I_{\lambda} \\ 0, & \text{if} \quad \alpha \notin I_{\lambda}, \end{cases}$$

If instead of T, we denote this operator by T_{λ} , then the 1-factorization identity (12) becomes

$$\Psi_{\mu} \cdot T_{\lambda} = \begin{cases} S_{\lambda}, & \text{if } \mu = \lambda \\ 0, & \text{otherwise.} \end{cases}$$

When \mathcal{L} is of type 3, one sees from the way the operator $T_{\lambda} = (T_{\lambda k})_{k=1}^{m} \in \mathcal{A}^{*m}$ is defined that

$$\Psi_{\lambda} \cdot T_{\lambda} = \sum_{k=1}^{m} \Psi_{\lambda k} \cdot T_{\lambda k} = S_{\lambda}, \text{ but}$$
$$\Psi_{\mu k} \cdot T_{\lambda k} = 0 \text{ for each } k = 1, ..., m, \text{ if } \mu \neq \lambda.$$

This means that for every $p \in \mathcal{C}(\eta)$, we have

$$\Psi_{p} \cdot T_{\lambda} = \sum_{k=1}^{m} \Psi_{pk} \cdot T_{\lambda k} = S_{\lambda} \quad \text{if} \quad I_{\lambda} \in p, \quad \text{but}$$
$$\Psi_{pk} \cdot T_{\lambda k} = 0 \quad \text{for each } k = 1, ..., m, \text{ if } I_{\lambda} \notin p.$$

The same modification may also be applied for the factorization identities in $\mathcal{LUC}(\mathcal{A})$ and $M_r(\mathcal{A})^*$ obtained below in Theorems 4.16 and 4.17. We shall make use of these remarks in the theorems in the forthcoming sections studying the algebraic structure of the algebras.

(ii) The elements S^{λ} defined in (19) for the 1-factorization of each S_{λ} can be used to provide a slightly different proof for the η -factorization in \mathcal{A}^* . For, let

(20)
$$S = \sum_{\lambda \le n} S^{\lambda}$$

Then this is well defined on \mathcal{L} . Furthermore, if $x \in \langle \mathcal{L} \rangle$, then

$$x = \sum_{j=1}^{p} \sum_{n \in I_j} a_n a_{\alpha_n},$$

where $I_j = \{n : \alpha_n \in I_{\lambda_j}\}$ and $a_n \in \langle A_{\alpha_n} \rangle$ for j = 1, ..., p. Accordingly,

$$\begin{aligned} |\langle S, x \rangle| &= |\langle \sum_{\lambda < \eta} S^{\lambda}, \sum_{j=1}^{P} \sum_{n \in I_{j}} a_{n} a_{\alpha_{n}} \rangle| = |\langle \sum_{j=1}^{P} \sum_{n \in I_{j}} S_{\lambda_{j}}, a_{n} \rangle| \\ &\leq \sup \|S_{\lambda}\| \sum_{j=1}^{p} \sum_{n \in I_{j}} \|a_{n}\| \leq K_{2} \sup \|S_{\lambda}\| \sum_{j=1}^{p} \sum_{n \in I_{j}} \|a_{n} a_{\alpha_{n}}\| \\ &\leq K_{1} K_{2} \sup \|S_{\lambda}\| \|x\|, \end{aligned}$$

where $K_1 > 0$ is the constant as in Lemma 4.5 and $K_2 > 0$ is the constant required by Inequality (F2) if m = 1 and by (F3) if $m \in \mathbb{N}$ is arbitrary. Since S is a bounded operator on $\langle \mathcal{L} \rangle$, we may extend it to $T \in \mathcal{A}^*$.

Then, using (19), we find for every $\lambda < \eta$ and $\alpha \in I_{\lambda}$,

$$\langle T, a\mathbf{a}_{\alpha} \rangle = \langle S_{\lambda}, a \rangle.$$

Therefore,

$$\langle \Psi_{\lambda} \cdot T, a \rangle = \lim_{\gamma} \langle \mathbf{a}_{h_{\lambda}(\gamma)} \cdot T, a \rangle = \lim_{\gamma} \langle T, a \mathbf{a}_{h_{\lambda}(\gamma)} \rangle = \langle S_{\lambda}, a \rangle,$$

as wanted.

(iii) When a Banach algebra \mathcal{A} is equipped with an $\mathbb{F}\ell^1(\eta)$ -base, \mathcal{A}^* has the η -factorization property as it has just been proved, but we have been unable to deduce that $\mathcal{LUC}(\mathcal{A})$ satisfies this property as well, that is for instance when m = 1, whether any bounded family of functionals $\{S_{\lambda} : \lambda < \eta\}$ contained in $\mathcal{LUC}(\mathcal{A})$ can be fatorized as $\Psi_{\lambda} \cdot T$ for some bounded family $\Psi_{\lambda} \in \mathcal{LUC}(\mathcal{A})^*$ and $T \in \mathcal{LUC}(\mathcal{A})$ for every $\lambda < \eta$.

We can nevertheless show below that the 1-factorization is true in $\mathcal{LUC}(\mathcal{A})$ whenever an $F\ell^1(\eta)$ -base is available in the algebra \mathcal{A} , which is enough as we shall see for our study of the algebra and the topological centre of $\mathcal{LUC}(\mathcal{A})^*$. To be able to deduce the η -factorization for $\mathcal{LUC}(\mathcal{A})$ one needs to make sure that the extension T of the operator S defined in (20) is in $\mathcal{LUC}(\mathcal{A})$.

Theorem 4.16. If a Banach algebra \mathcal{A} with a bai has an $F\ell^1(\eta)$ -base for some infinite cardinal η , then $\mathcal{LUC}(\mathcal{A})$ has the 1-factorization property through $\prod_{k=1}^m X_k$.

Proof. Suppose that the $F\ell^1(\eta)$ -base is of type 2 (type 1 is then inculded). Let $P \in \mathcal{LUC}(\mathcal{A})$ be arbitrary and choose $S \in \mathcal{A}^*$ and $a \in \mathcal{A}$ such that $P = S \cdot a$. Then, factorize S as $S = \Psi \cdot T$, where $\Psi \in X$ and $T \in \mathcal{A}^*$ are as obtained in Theorem 4.14. Since $T \cdot a \in \mathcal{LUC}(\mathcal{A})$ by definition, we only need to check that $P = \Psi \cdot (T \cdot a)$. So let $b \in \mathcal{A}$ be arbitrary. Then

$$\langle \Psi \cdot (T \cdot a), b \rangle = \langle \Psi, (T.a), b \rangle = \langle \Psi, T.(ab) \rangle = \langle \Psi \cdot T, ab \rangle = \langle S, ab \rangle$$

= $\langle S.a, b \rangle = \langle P, b \rangle$,

as wanted.

The proof is similar when the $F\ell^1(\eta)$ -base is of type 3.

Recall that a dual Banach algebra is a Banach algebra \mathcal{B} such that $\mathcal{B} = \mathcal{C}^*$ for some Banach space \mathcal{C} (not necessarily unique), which is a submodule of the dual \mathcal{B} -bimodule \mathcal{B}^* (see [54, page 194]).

The condition imposed below on the net in \mathcal{C} happens for instance when the net is a left bounded approximate identity and \mathcal{C} is Arens regular.

Our set X defined in Notation 4.11 is now seen as a bounded subset in $M_r(\mathcal{A})^{**}$.

Theorem 4.17. Let \mathcal{A} be a Banach algebra with a bai and an $F\ell^1(\eta)$ -base for some infinite cardinal η . Suppose that $M_r(\mathcal{A})$ is a dual Banach algebra with predual algebra \mathcal{C} having a net in $\mathcal{A} \cap \mathcal{C}$ weak^{*}-converging to a left identity in $M_r(\mathcal{A})^*$. Then $M_r(\mathcal{A})^*$ has the η -factorization property.

Proof. Let $\mathcal{L} := \bigcup_{\alpha < \eta} A_{\alpha} \{ a_{\alpha}^{k} : n = 1, ..., m \}$ be an $F\ell^{1}(\eta)$ -base in \mathcal{A} . The case m = 1 will correspond to both types of the $F\ell^{1}(\eta)$ -bases 1 and 2. We prove that $M(\mathfrak{A})^{*}$ has the 1 factorization property. So let S be any element in $M(\mathfrak{A})^{*}$ and

 $M_r(\mathcal{A})^*$ has the 1-factorization property. So let S be any element in $M_r(\mathcal{A})^*$ and let $(\Psi_k)_{k=1}^m$ be any element in $\prod_{k=1}^m X_k$. Restrict S to \mathcal{A} and consider the family $\{T_k : k = 1, ..., m\}$ in \mathcal{A}^* , found in Theorem 4.14 for the factorization of S so that

(21)
$$\langle \sum_{k=1}^{m} \Psi_k \cdot T_k, a \rangle = \langle S, a \rangle \text{ for every } a \in \mathcal{A}$$

Then by Hahn-Banach Theorem, regard this family $\{T_k : k = 1, ..., m\}$ in $M_r(\mathcal{A})^*$, and regard $\{\Psi_k : k = 1, ..., m\}$ in $M_r(\mathcal{A})^{**}$ (the unit ball if the elements a_{α}^k are bounded by 1), as explained in Notation 4.11. Let $(\varphi_i) \subset \mathcal{C} \cap \mathcal{A}$ be the net in \mathcal{C} with a left identity E as a weak*-limit in $M_r(\mathcal{A})^*$. Then using the fact that \mathcal{A} is a right ideal in $M_r(\mathcal{A})$, we apply (21) for the fourth of the following equalities

$$\begin{split} \langle \sum_{k=1}^{m} \Psi_k \cdot T_k, b \rangle &= \langle E(\sum_{k=1}^{m} \Psi_k \cdot T_k), b \rangle = \lim_i \langle \varphi_i(\sum_{k=1}^{m} \Psi_k \cdot T_k), b \rangle \\ &= \lim_i \langle \sum_{k=1}^{m} \Psi_k \cdot T_k, b\varphi_i \rangle = \lim_i \langle S, b\varphi_i \rangle \\ &= \lim_i \langle \varphi_i S, b \rangle = \langle ES, b \rangle = \langle S, b \rangle \end{split}$$

for every $b \in M_r(\mathcal{A})$, as required.

The η -factorization property is proved following the same steps as in the proof of Theorem 4.14.

Remark 4.18. Flipping definitions and conditions from left to right, the reader may check that $M_l(\mathcal{A})^*$ has the right η -factorization property when \mathcal{A} has a right $F\ell^1(\eta)$ -base for some infinite cardinal η .

5. The algebraic structure of \mathcal{A}^{**} , $\mathcal{LUC}(\mathcal{A})^*$ and $M_r(\mathcal{A})^{**}$

Various algebraic properties were proved in the algebras in harmonic analysis related to locally compact groups G and to some infinite, discrete semigroups Ssuch as $L^1(G)^{**}$, $\mathcal{LUC}(G)^*$, $\ell^{\infty}(S)^{**}$, $A(G)^{**}$ and $\mathcal{UC}_2(G)^*$.

Algebraic properties were also found in the Stone-Čech compactification βS of some discrete semigroups S and the \mathcal{LUC} -compactification G^{LUC} of a locally compact group G. See Notes 6.7 in next section. Most of these properties such as right cancellation, the number of topologically left invariant means or the number of left ideals in G^{LUC} and βS when S is a discrete semigroup are in fact related to the non-Arens regularity of the algebra. For instance when G is amenable, the number of topologically left invariant means $\mathcal{LUC}(G)^*$ or in $L^1(G)^{**}$ and the number of

disjoint left ideals in G^{LUC} are the same. For the support of a (topologically) left invariant, regarded as a Borel measure on G^{LUC} , is a left ideal in G^{LUC} . Conversely, if L is a left ideal in G^{LUC} , $x \in L$ and Ψ is a (topologically) left invariant mean, then Ψx is also (topologically) left invariant and regarded as a Borel measure on G^{LUC} , it has its support contained in L.

In fact, already in 1951, Arens himself showed that the semigroup algebra ℓ^1 with convolution is non-Arens regular. For this, he produced two distinct invariant means Φ and Ψ in $\ell^{1^{**}}$ (see [1]). This gave him immediately

$$\Phi \Psi = \Psi \neq \Phi = \Psi \Phi \quad \text{and} \quad \Phi \diamond \Psi = \Phi \neq \Psi = \Psi \diamond \Phi,$$

which means that

$$T_1 Z(\ell^{1^{**}}) \neq \ell^{1^{**}}$$
 and $T_2 Z(\ell^{1^{**}}) \neq \ell^{1^{**}}$

In 1957, Day used the same argument in [20] to show that $L^1(G)$ is non-Arens regular for many infinite discrete groups, including all Abelian ones. This same approach was followed by Civin and Yood in their seminal paper [12] where $L^1(G)$ was shown to be non-Arens regular for any infinite locally compact Abelian group. Their method relied again on Day's result when the group is discrete.

Once an infinite $F\ell^1(\eta)$ -base is available in \mathcal{A} , the theorems in this section unify all these results in \mathcal{A}^{**} , $\mathcal{LUC}(\mathcal{A})^*$ and $M_r(\mathcal{A})^{**}$ and many subsemigroups in these algebras.

We start with right cancellation. We say that a set Γ in an algebra is right cancellable if for every nonzero element a in the algebra , $ax \neq 0$ for some $x \in \Gamma$. When $\Gamma = \{x\}$ is right cancellable, we simply say that x is right cancellable.

Theorem 5.1. Let \mathcal{A} be a Banach algebra with a bai and an $F\ell^1(\eta)$ -base \mathcal{L} for some infinite cardinal η and some integer $m \in \mathbb{N}$. Let X be as in Notation 4.11 and S be any subsemigroup of $\mathcal{LUC}(\mathcal{A})^*$ with $X \subseteq S$, or any subsemigroup \mathcal{A}^{**} with $\widetilde{X} \subseteq S$.

- (i) If L is of type 1 or 2, then there are at least 2^{2ⁿ} many right cancellable elements in S.
- (ii) If L is of type 3, then there are at least 2^{2ⁿ} many right cancellable sets in S.

Proof. Let $\mathcal{L} := \bigcup_{\alpha < \eta} A_{\alpha} \{ a_{\alpha}^{k} : k = 1, ..., m \}$ be an $\mathbb{F}\ell^{1}(\eta)$ -base in \mathcal{A} . Let p and q be two different cofinal ultrafilters in $\beta\eta$ together with their corresponding elements

two different connal ultrafilters in $\beta\eta$ together with their corresponding elements $(\Psi_{pk})_{k=1}^m$ and $(\Psi_{qk})_{k=1}^m$ in $\prod_{k=1}^m X_k$. Pick two disjoint members I of p and J of q. For each k = 1, ..., m, Ψ_{pk} is the weak*-limit of a subnet of $(a_{\alpha}^k)_{\alpha \in I}$, which we denote for simplicity again by $(a_{\alpha}^k)_{\alpha \in I}$. In the same way, Ψ_{qk} is the weak*-limit of a subnet of $(a_{\alpha}^k)_{\alpha \in J}$. Then, for each $\alpha < \eta$ and k = 1, ..., m, we may take by Remark 4.13, an operator $P_k \in \mathcal{LUC}(\mathcal{A})$ such that

$$\langle P_k, a_{\alpha}^k \rangle = \begin{cases} 1, & \text{if } \alpha \in I \\ 0, & \text{elsewhere.} \end{cases}$$

Then, for each k = 1, ..., m, we have

$$\langle \Psi_{pk}, P_k \rangle = \lim_{\alpha \in I} \langle P_k, a^k_{\alpha} \rangle = 1$$
 while $\langle \Psi_{qk}, P_k \rangle = \lim_{\alpha \in J} \langle P_k, a^k_{\alpha} \rangle = 0.$

In other words, for each k = 1, ..., m, the elements Ψ_{pk} and Ψ_{qk} are different in X_k . Thus the map $\mathcal{C}(\eta) \to X_k : p \mapsto \Psi_{pk}$ is one-to-one for each k = 1, ..., m. Since the cardinality of $\mathcal{C}(\eta)$ is 2^{2^n} , we obtain the same number of points Ψ_{pk} in X_k for each k = 1, ..., m. This clearly implies that there are at least 2^{2^n} many elements $(\Psi_{pk})_{k=1}^m$ in $\prod_{k=1}^m X_k$. We claim first that each of the sets $\{\Psi_{pk} : k = 1, ..., m\}$ is right cancellable in S when S is a subsemigroup of $\mathcal{LUC}(\mathcal{A})^*$. So let Ψ be any

nonzero element in S and pick $P \in \mathcal{LUC}(\mathcal{A})$ with $\langle \Psi, P \rangle \neq 0$. Then by Theorem 4.16, there are operators $T_k \in \mathcal{LUC}(\mathcal{A}), k = 1, ..., m$, such that $\sum_{k=1}^{m} \Psi_{pk} \cdot T_k = P$. Consequently,

$$\langle \sum_{k=1}^{m} \Psi \Psi_{pk}, T_k \rangle = \langle \Psi, \sum_{k=1}^{m} \Psi_{pk} \cdot T_k \rangle = \langle \Psi, P \rangle \neq 0,$$

and so $\Psi \Psi_{pk}$ must be nonzero for at least one k = 1, ..., m, as wanted.

In the case when S is a subsemigroup of \mathcal{A}^{**} , we simply take for each element $\Psi_{pk} \in X_k$ a fixed element $\widetilde{\Psi_{pk}} \in \mathcal{A}^{**}$ such that $\pi(\widetilde{\Psi_{pk}}) = \Psi_{pk}$. Then for a nonzero element Ψ in S, pick $S \in \mathcal{A}^*$ such that $\langle \Psi, S \rangle \neq 0$ together with the *m* operators $T_k \in \mathcal{A}^*$ given by Theorem 4.14 such that

$$\sum_{k=1}^{m} \widetilde{\Psi_{pk}} \cdot T_k = \sum_{k=1}^{m} \Psi_{pk} \cdot T_k = S.$$

As before, this gives

$$\langle \sum_{k=1}^{m} \Psi \widetilde{\Psi_{pk}}, T_k \rangle = \langle \Psi, \sum_{k=1}^{m} \widetilde{\Psi_{pk}} \cdot T_k \rangle = \langle \Psi, S \rangle \neq 0,$$

and so again at least one of the terms $\Psi \widetilde{\Psi_{pk}}$ must be nonzero, showing that the set $\{\widetilde{\Psi_{pk}}: k = 1, ..., m\}$ is right cancellable in S for each $p \in \mathcal{C}(\eta)$.

This proves Statement (ii) when m is arbitrary, and Statement (i) when taking m = 1.

Theorem 5.2. Let \mathcal{A} be a Banach algebra with a bai and an $F\ell^1(\eta)$ -base \mathcal{L} for some infinite cardinal η and some integer $m \in \mathbb{N}$. Let X be as in Notation 4.11 and S be any non-trivial subalgebra of $\mathcal{LUC}(\mathcal{A})^*$ with $X \subseteq S$, or any non-trivial subalgebra of \mathcal{A}^{**} with $\widetilde{X} \subseteq S$. Then the dimension of any principal right ideal, and so of any nonzero right ideal, in S is at least $2^{2^{\eta}}$.

Proof. Let \mathcal{L} be an $\mathbb{F}\ell^1(\eta)$ -base in \mathcal{A} as in Theorem 5.1.

We prove the theorem when S is a subalgebra of $\mathcal{LUC}(\mathcal{A})^*$, the proof is similar for \mathcal{A}^{**} . Pick any nonzero element Ψ in a given nonzero right ideal of S. For each $p \in \mathcal{C}(\eta)$, let $(\Psi_{p1}, ..., \Psi_{pm})$ be the corresponding elements in $\prod_{k=1}^{m} X_k$ as in Theorem 5.1. Then we claim that the set

$$\{(\Psi\Psi_{p1},...,\Psi\Psi_{pm}): p \in \mathcal{C}(\eta)\}$$

is linearly independent in the product space $(\Psi S)^m$. Suppose otherwise that for some scalars c_n and distinct cofinal ultrafilters p_n , n = 1, ..., N, we have

$$\sum_{n=1}^{N} c_n(\Psi \Psi_{p_n 1},...,\Psi \Psi_{p_n m}) = 0.$$

Then

(22)
$$\sum_{n=1}^{N} c_n \Psi \Psi_{p_n k} = 0 \quad \text{for every} \quad k = 1, ..., m.$$

Let $P \in \mathcal{LUC}(\mathcal{A})$ such that $\langle \Psi, P \rangle = 1$. Pick pairwise disjoint members I_n of the ultrafilters $p_n, n = 1, ..., N$. For each i = 1, ..., N, pick the operators T_k in $\mathcal{LUC}(\mathcal{A})$, k = 1, ..., m, given by Theorem 4.16 as in Remark 4.15; that is,

$$\sum_{k=1} \Psi_{p_i k} \cdot T_k = P \quad \text{but} \quad \Psi_{p_n k} \cdot T_k = 0 \quad \text{for every} \quad k = 1, ..., m \quad \text{whenever} \quad n \neq i.$$

(The operators T_k depend on i, but this is not underlined since no confusion is caused.)

Then, for each i = 1, ..., N, we have by (22)

$$0 = \left\langle \sum_{k=1}^{m} \left(\sum_{n=1}^{N} c_n \Psi \Psi_{p_n k} \right), T_k \right\rangle = \sum_{k=1}^{m} \sum_{n=1}^{N} c_n \left\langle \Psi, \Psi_{p_n k} \cdot T_k \right\rangle$$
$$= \sum_{n=1}^{N} c_n \left\langle \Psi, \sum_{k=1}^{m} \Psi_{p_n k} \cdot T_k \right\rangle$$
$$= c_i \left\langle \Psi, P \right\rangle = c_i,$$

showing the linear independence of the set $\{(\Psi \Psi_{p1}, ..., \Psi \Psi_{pm}) : p \in \mathcal{C}(\eta)\}$, and so the dimension of the product space $(\Psi S)^m$ must be at least 2^{2^n} . Accordingly, the dimension of the coordinate space ΨS , as well as the right ideal containing Ψ , must be also at least 2^{2^n} . This completes the proof.

We deal now with the number and dimension of left ideals. As already observed in [45], nonzero left ideals in \mathcal{A}^{**} and $\mathcal{LUC}(\mathcal{A})^*$ may have finite dimensions. This is the case for example when \mathcal{A} is the Fourier algebra A(G) of a locally compact group G, or the group algebra $L^1(G)$ of an amenable locally compact group G. Topologically left invariant means generate, in this case, one-dimensional left ideals in both \mathcal{A}^{**} and $\mathcal{LUC}(\mathcal{A})^*$. When G is a nondiscrete, we find therefore $2^{2^{\eta}}$ many such left ideals in the case of the Fourier algebra, where η is the local weight of G(see [11] and [67]). When G is amenable and noncompact, we found $2^{2^{\eta}}$ many such left ideals in the case of the group algebra, where η is the compact covering of G(see [87] or [46]).

In what follows, we shall see that there is also an abundance of infinite-dimensional left ideals in the algebras when an $F\ell^1(\eta)$ -base is available in \mathcal{A} . In particular, we show that there are at least $2^{2^{\eta}}$ many left ideals each of dimension at least $2^{2^{\eta}}$ in both algebras \mathcal{A}^{**} and $\mathcal{LUC}(\mathcal{A})^*$. Under the conditions imposed on \mathcal{A} in Theorem 4.17, these theorems hold also for the multiplier algebra $M_r(\mathcal{A})$.

Theorem 5.3. Let \mathcal{A} be a Banach algebra with a bai and an $F\ell^1(\eta)$ -base \mathcal{L} for some infinite cardinal η and some integer $m \in \mathbb{N}$. Let X be as in Notation 4.11 and S be any subsemigroup or a subalgebra of $\mathcal{LUC}(\mathcal{A})^*$ with $X \subseteq S$, or any subsemigroup or a subalgebra of \mathcal{A}^{**} with $\widetilde{X} \subseteq S$.

- (i) If L is of type 1 or type 2 and S a subsemigroup, then there are at least 2^{2ⁿ} many disjoint principal left ideals in S. If S is a subalgebra, these left ideals have zero pairwise intersections.
- (ii) If \mathcal{L} is of type 3, then there are at least $2^{2^{\eta}}$ many principal left ideals in S.
- (iii) If S is subalgebra, then the dimension of each of these left ideals is at least $2^{2^{\eta}}$.

Proof. Let $\mathcal{L} := \bigcup_{\alpha < \eta} A_{\alpha} \{ a_{\alpha}^k : k = 1, ..., m \}$ be an $\mathbb{F}\ell^1(\eta)$ -base in \mathcal{A} . As in Theorem

5.1, the case m = 1 will correspond to both types of $F\ell^1(\eta)$ -base, 1 and 2.

We prove the theorem when S is a subsemigroup or a subalgebra of $\mathcal{LUC}(\mathcal{A})^*$, the proof is similar in the other case. Let p and q be two distinct ultrafilters in $\mathcal{C}(\eta)$) together with their corresponding elements Ψ_{pk} and Ψ_{qk} in X_k , k = 1, ..., m. When \mathcal{L} is of type 1 or 2, the corresponding elements in X will be taken as Ψ_p and Ψ_q .

When S is a subsemigroup, we consider the left ideals ${\mathcal P}$ and ${\mathcal Q}$ given, respectively, by the sets

$$S{\Psi_{pk}: k = 1, ..., m}$$
 and $S{\Psi_{qk}: k = 1, ..., m}$.

When S is a subalgebra, the left ideals \mathcal{P} and \mathcal{Q} are generated by these two sets. In both cases of whether S is a subsemigroup or a subalgebra, we prove that neither of these left ideals contains the other.

Pick any nonzero element Ψ in S together with an operator P in $\mathcal{LUC}(\mathcal{A})$ such that $\langle \Psi, P \rangle \neq 0$. By Theorem 4.16 and Remark 4.15, let T_k , k = 1, ..., m, be the operators in $\mathcal{LUC}(\mathcal{A})$ such that

$$\sum_{k=1}^{m} \Psi_{pk} \cdot T_k = P, \quad \text{but} \quad \Psi_{qk} \cdot T_k = 0 \quad \text{for every} \quad k = 1, ..., m.$$

So, for all $\Phi \in S$ and all k = 1, ..., m, we have

$$\langle \Phi \Psi_{qk}, T_k \rangle = \langle \Phi, \Psi_{qk} \cdot T_k \rangle = \langle \Phi, 0 \rangle = 0,$$

and accordingly,

$$\langle Q, T_k \rangle = 0$$
 for every $Q \in \mathfrak{Q}$ and $k = 1, ..., m$.

But since

$$\sum_{k=1}^{m} \langle \Psi \Psi_{pk}, T_k \rangle = \langle \Psi, \sum_{k=1}^{m} \Psi_{pk} \cdot T_k \rangle = \langle \Psi, P \rangle \neq 0,$$

we must have,

 $\langle \Psi \Psi_{pk}, T_k \rangle \neq 0$ for some k = 1, ..., m.

Therefore, $\Psi \Psi_{pk} \notin \Omega$, showing that \mathcal{P} is not contained in Ω . In the same way, we show that Ω is not contained in \mathcal{P} . If \mathcal{L} is of type 1 or 2, this shows that the principal left ideals $S\Psi_p$ and $S\Psi_q$ are disjoint when S is a subsemigroup, and have their intersection equal to zero when S is a subalgebra. If \mathcal{L} is of type 3, this shows that the left ideals \mathcal{P} and Ω are different. Since there are $2^{2^{\eta}}$ ultrafilters in $\mathcal{C}(\eta)$, the first two statements follow.

As for Statement (iii), suppose that S is a subalgebra of $\mathcal{LUC}(\mathcal{A})^*$. We show that the dimension of each of the left ideals of S found in the previous statements is at least $2^{2^{\eta}}$. Recall first that by Theorem 5.1, the cardinality of the set X is $2^{2^{\eta}}$. Moreover, arguing as in Theorem 5.2, it is quick to see that X is linearly independent. Therefore the dimension of S is at least equal to $2^{2^{\eta}}$ since $X \subseteq S$. Now let \mathcal{P} be any of the left ideals $S{\Psi_{pk}: k = 1, ..., m}$ of S, where p is a fixed cofinal ultrafilter in $\beta\eta$ and Ψ_{pk} are the corresponding elements in $X_k, k = 1, ..., m$. By Theorem 5.1, the map

$$\mathcal{S} \to \prod_{k=1}^m \mathcal{S}: \quad \Psi \mapsto \Psi(\Psi_{p1}, ..., \Psi_{pm})$$

is a linear isomorphism onto its image. Therefore, the dimension of the linear space $\prod_{k=1}^{m} \$\Psi_{pk}$ is at least $2^{2^{\eta}}$. Clearly, this implies that for at least one k = 1, ..., m, the linear space $\$\Psi_{pk}$ is of dimension at least $2^{2^{\eta}}$. Thus, the dimension of the left ideal generated by

$$S{\Psi_{pk}: k = 1, ..., m}$$

must be at least $2^{2^{\eta}}$.

Theorem 5.4. Let \mathcal{A} be a Banach algebra with a bai and an $F\ell^1(\eta)$ -base \mathcal{L} for some infinite cardinal η and some integer $m \in \mathbb{N}$. Suppose that $M_r(\mathcal{A})$ is a dual Banach algebra with predual algebra \mathbb{C} having a net in $\mathcal{A} \cap \mathbb{C}$ weak*-converging to a left identity in $M_r(\mathcal{A})^*$. Regard X as a subset of $M_r(\mathcal{A})^{**}$ as agreed in Notation 4.11 and let \mathbb{S} be any subsemigroup or a subalgebra of $M_r(\mathcal{A})^{**}$ with $X \subseteq \mathbb{S}$.

(i) If L is of type 1 or type 2, then there are at least 2^{2ⁿ} many right cancellable elements in S.

- (ii) If L is of type 3, then there are at least 2^{2^η} many right cancellable sets in S.
- (iii) If L is of type 1 or type 2 and S a subsemigroup, then there are at least 2^{2ⁿ} many disjoint principal left ideals in S. If S is a subalgebra, these left ideals have zero pairwise intersections.
- (iv) If \mathcal{L} is of type 3, then there are at least $2^{2^{\eta}}$ many left ideals in S.
- (v) If S is a non-trivial subalgebra of $M_r(\mathcal{A})^{**}$, then the dimension of each of these left ideals is at least 2^{2^n} .
- (vi) If S is a subalgebra of $M_r(\mathcal{A})^{**}$, then the dimension of any principal right ideal, and so of any nonzero right ideal, in S is at least $2^{2^{\eta}}$.

Proof. According to Theorem 4.17, under the condition imposed on \mathcal{A} , an $\mathbb{F}\ell^1(\eta)$ base in \mathcal{A} provides $M_r(\mathcal{A})^*$ with the η -factorization property. In particular, the 1-factorization together with Remark 4.15 (the ingredients necessary to prove Theorems 5.1, 5.2 and 5.3) are also available in this case. So we may proceed in the same way.

Remark 5.5. If $m \ge 2$, the set $\{\Psi_{pk} : k = 1, ...m\}$ is right cancellable in the subsemigroups S considered in the theorems above, but we do not know if each (or any of) the elements Ψ_{pk} , k = 1, ...m, is right cancellable in S.

The same problem concerns the left ideals generated by any of the elements Ψ_{pk} : k = 1, ...m. If m = 1 or \mathcal{L} is of type 1 or 2, then these principal left ideals have trivial intersections. But if $m \geq 2$, we are only able to show that neither one contains the other.

6. Application I: Group Algebra, semigroup Algebra and measure Algebra

A number of Banach algebras in harmonic analysis were proved in the past to be strongly Arens irregular and to carry a rich algebraic structure. In the rest of the paper, we see that $F\ell^1(\eta)$ -bases are indeed available in *almost* all these algebras, and so with the approach developed in the previous sections, we shall unify almost all what is known so far on the subjects since the introduction by Arens of his two products in 1951. Some further interesting new cases are also obtained.

6.1. Some definitions. Before we proceed with our first application, we summarize first our definitions and some properties. For a locally compact group G with a fixed left Haar measure, we will be considering two cardinal invariants of G. The *local weight* of G is the least cardinality of an open base at the identity of G. The *compact covering number* of G is the least cardinality of a compact covering of G.

The group algebra $L^1(G)$ is the Banach space of all equivalence classes of scalarvalued functions which are integrable with respect to the Haar measure under convolution, where as usual, two functions are equivalent if and only if they differ only on a set of Haar measure zero. By $L^{\infty}(G)$ we understand the Banach dual of $L^1(G)$. This is the C*-algebra of all essentially bounded locally measurable functions on G.

When G is an infinite discrete semigroup, we shall also use $L^1(G)$ and $L^{\infty}(G)$ instead of the usual $\ell^1(G)$ and $\ell^{\infty}(G)$, respectively.

The C^{*}-algebra $\mathcal{LUC}(G)$ is the space of all bounded *right uniformly continuous* functions on G. These are the bounded functions on G such that for every $\epsilon > 0$, there exists a neighbourhood U of e such that

$$|f(s) - f(t)| < \epsilon$$
 whenever $st^{-1} \in U$.

These are also the bounded functions on ${\cal G}$ which are left norm continuous in the sense that

$$s \mapsto {}_s f : G \to \mathcal{CB}(G)$$

is continuous, where $\mathcal{CB}(G)$ is the space of continuous bounded scalar valued functions equipped with its supremum norm and ${}_sf$ is the left translate of f by $s \in G$. In the literature, this C*-algebra is denoted by $C_{ru}(G)$, LC(G), $\mathcal{LC}(G)$ or $\mathcal{LUC}(G)$. We are using the latter notation. As known $\mathcal{LUC}(G)$ can be obtained as $L^1(G)*L^{\infty}(G)$, and so we have $\mathcal{LUC}(L^1(G)) = \mathcal{LUC}(G)$.

The C^{*}-subalgebra of $\mathcal{LUC}(G)$ consisting of functions vanishing at infinity will be denoted by $C_0(G)$.

The C*-subalgebra of $L^{\infty}(G)$ consisting of functions vanishing at infinity will be denoted by $L_0^{\infty}(G)$, that is,

$$L_0^{\infty}(G) = \{ f \in L^{\infty}(G) : \|f\|_{G \setminus K} \to 0 \text{ as } K \to G \},$$

where $||f||_{G\setminus K}$ is the essential supremum norm of f on $G\setminus K$.

The space M(G) is the usual measure algebra consisting of bounded Radon measures on G. The measure algebra M(G) is the dual space of $C_0(G)$, and by the well-known Wendel's theorem, M(G) is the multiplier algebra of $L^1(G)$, that is $M(L^1(G)) = M(G)$.

We shall consider the annihilator space of $L_0^{\infty}(G)$ in $L^1(G)^{**}$ given by

$$L_0^{\infty}(G)^{\perp} = \{ \Psi \in L^1(G)^{**} : \langle \Psi, f \rangle = 0 \text{ for every } f \in L_0^{\infty}(G) \}.$$

The space $C_0(G)^{\perp_{lu}}$ is the annihilator of $C_0(G)$ in $\mathcal{LUC}(G)^*$ when $C_0(G)$ is regarded as a subalgebra of $\mathcal{LUC}(G)$, i.e.,

$$C_0(G)^{\perp_{lu}} = \{ \Psi \in \mathcal{LUC}(G)^* : \langle \Psi, f \rangle = 0 \text{ for every } f \in C_0(G) \}.$$

The space $C_0(G)^{\perp_M}$ is the annihilator of $C_0(G)$ in $M(G)^{**}$ when $C_0(G)$ is regarded as a subalgebra of $M(G)^*$, i.e.,

$$C_0(G)^{\perp_M} = \{ \Psi \in M(G)^{**} : \langle \Psi, f \rangle = 0 \text{ for every } f \in C_0(G) \}.$$

When G is a discrete semigroup, we denote these annihilators simply by $c_0(G)^{\perp}$.

Recall that we have the following decompositions of $\mathcal{LUC}(G)^*$, $L^{\infty}(G)^*$ and $M(G)^{**}$ as Banach space direct sums

$$\begin{aligned} \mathcal{LUC}(G)^* &= M(G) \oplus C_0(G)^{\perp_{lu}}, \\ L^{\infty}((G)^* &= L_0^{\infty}(G)^* \oplus L_0^{\infty}(G)^{\perp}, \\ M(G)^{**} &= M(G) \oplus C_0(G)^{\perp_M}, \end{aligned}$$

where $C_0(G)^{\perp_{lu}}$, $L_0^{\infty}(G)^{\perp}$ and $C_0(G)^{\perp_M}$ are weak*-closed ideals in the corresponding algebra. See [16, 35, 54, 88].

We shall also be concerned with the \mathcal{LUC} -compactification G^{LUC} of a noncompact locally compact group G and the Stone-Čech compactification βG of an infinite discrete semigroup G. Both are compact right topological subsemigroups of $\mathcal{LUC}(G)^*$ and $\ell^{\infty}(G)^*$, respectively. With the Gelfand topology, they can be produced as the spectra (non-zero multiplicative linear functionals) of the C^{*}-algebras $\mathcal{LUC}(G)$ and $\ell^{\infty}(G)$, respectively. They are the largest semigroup compactifications of G with the joint-continuity property.

The canonical morphism $\epsilon\colon G\to G^{LUC}$ given by evaluations

$$\epsilon(s)(f) = f(s)$$
 for every $f \in \mathcal{LUC}(G)$ and $s \in G$

identifies G as a subgroup of G^{LUC} . The same morphism on βG , defined with $\ell^{\infty}(G)$ instead of $\mathcal{LUC}(G)$, identifies G as a subsemigroup of βG when G is a discrete semigroup. In both cases, G^* will denote the remainders $G^{LUC} \setminus G$ and $\beta G \setminus G$.

The reader is directed to [65] for the algebra in βG , and to our most recent investigation in [38] on G^{LUC} and other semigroup compactifications. It may be noted that the spectrum of $L^{\infty}(G)$ is not a semigroup unless G is compact or

discrete (see [89]), so this is going to be our concern only when G is discrete, i.e., the spectrum is βG .

6.2. $\mathbf{F}\ell^1(\eta)$ -bases. We start with the group algebra of a non-compact locally compact group. Variants of the construction below can be found in many articles dealing with the topological centre and the algebraic structure of $\mathcal{LUC}(G)^*$ and $L^1(G)^{**}$.

Theorem 6.1. Let G be a non-compact locally compact group with a compact covering η . Then the group algebra $L^1(G)$ has an $F\ell^1(\eta)$ -base of type 2.

Proof. Let λ be a fixed left Haar measure and Δ be the modular function on G. Let $\{K_{\alpha}\}_{\alpha < \eta}$ be a compact covering of G made of symmetric subsets of G with nonempty interiors and with $\lambda(K_{\alpha}) \leq 1$ for each α . Then put

$$V_{\alpha} = \bigcup_{\beta < \alpha} K_{\beta} \quad \text{for each} \quad \alpha < \eta,$$

and observe that $(V_{\alpha})_{\alpha < \eta}$ is an increasing family of symmetric subsets of G with nonempty interiors and $\kappa(V_{\alpha}) \leq \alpha$ for each $\alpha < \eta$. Now choose inductively elements $x_{\alpha} \in G$ such that

(23)
$$(V_{\alpha}x_{\alpha}) \cap (V_{\beta}x_{\beta}) = \emptyset$$
 when $\alpha \neq \beta$.

This is possible because, for every $\alpha < \eta$,

$$\kappa\left(V_{\alpha}(\bigcup_{\beta<\alpha}V_{\beta}x_{\beta})\right)\leq\alpha.$$

Let now, for each $\alpha < \eta$, A_{α} be the set of all continuous positive functions φ with compact supports contained in V_{α} and with $\|\varphi\|_1 = 1$.

We check that the conditions in Definition 4.8 are satisfied so that

(24)
$$\mathcal{L} = \bigcup_{\alpha < \eta} (A_{\alpha} * \delta_{x_{\alpha}})$$

is indeed an $F\ell^1(\eta)$ -base of type 2 for $L^1(G)$.

First note that the closed linear span of the set A_{η} is $L^{1}(G)$. Since the functions in $A_{\alpha} * \delta_{x_{\alpha}}$ are supported in $V_{\alpha}x_{\alpha}$, it is also clear that the intersection of the linear spans of each pair of sets $A_{\alpha} * \delta_{x_{\alpha}}$ is zero. Let then φ be any function in the linear span of \mathcal{L} of the form

$$\varphi = \sum_{n=1}^{p} \varphi_n * \delta_{x_{\alpha_n}},$$

where each φ_n (n = 1, ..., p) is in the linear span of A_{α_n} .

Then, for each n = 1, ..., p,

$$\varphi_n * \delta_{x_{\alpha_n}} = \Delta(x_{\alpha_n}^{-1})(\varphi_n)_{x_{\alpha_n}^{-1}}$$

(the right translate of φ_n by $x_{\alpha_n}^{-1}$), and so

(25)
$$\left\|\sum_{n=1}^{p}\varphi_{n}*\delta_{x_{\alpha_{n}}}\right\|_{1} = \int_{G}\left|\sum_{n=1}^{p}\varphi_{n}*\delta_{x_{\alpha_{n}}}\right|d\lambda = \int_{G}\sum_{n=1}^{p}\left|\varphi_{n}*\delta_{x_{\alpha_{n}}}\right|d\lambda$$
$$= \sum_{n=1}^{p}\left\|\varphi_{n}*\delta_{x_{\alpha_{n}}}\right\|_{1} = \sum_{n=1}^{p}\left\|\varphi_{n}\right\|_{1},$$

where the second equality is due to the fact that the supports of the functions $\varphi_n * \delta_{x_{\alpha_n}}$ are disjoint. The third equality shows that \mathcal{L} is selective (with $K_1 = 1$)

while the last equality shows the validity of Condition (F2) in Definition 4.8 with $K_2 = 1$. For, when p = 1, the identity (25) simply says that

$$\|\varphi * \delta_{x_{\alpha}}\|_{1} = \|\varphi\|_{1} \quad \text{for every} \quad \varphi \in \langle A_{\alpha} \rangle.$$

Note that Condition (f3) (and so Condition (F3) in Definition 4.9) is also valid here with m = 1, a_{α} and b_{α} may be simply taken as $\delta_{x_{\alpha}}$ and $\delta_{x_{\alpha}^{-1}}$, respectively, for every $\alpha < \eta$.

This ends the proof for $L^1(G)$.

The same construction gives $F\ell^1(\eta)$ -bases when G is a discrete semigroup which respects certain cancellation properties, where η is the cardinality of G. Recall first that a semigroup G is called *weakly cancellative* if the sets

$$s^{-1}t = \{u \in G : su = t\}$$
 and $ts^{-1} = \{u \in G : us = t\}$

are finite for every $s, t \in G$. The semigroup G is left (right) cancellative when $s^{-1}t$ (ts^{-1} , respectively) has at most one element. We shall use the notations

$$s^{-1}B = \{ u \in G : su \in B \}$$
 and $A^{-1}B = \bigcup_{s \in A} s^{-1}B$,

where $s \in G$ and $A, B \subseteq G$. The sets Bs^{-1} and BA^{-1} are defined similarly.

Theorem 6.2. Let G be an infinite discrete weakly cancellative, right cancellative semigroup with cardinality η . Then the semigroup algebra $\ell^1(G)$ has an $F\ell^1(\eta)$ -base of type 1.

Proof. Enumerate G as $G = \{s_{\alpha} : \alpha < \eta\}$, and let for each $\alpha < \eta$,

$$G_{\alpha} = \{ s_{\beta} : \beta < \alpha \}.$$

Then $\{G_{\alpha}\}_{\alpha < \eta}$ is an increasing cover of G with $|G_{\alpha}| = \alpha$ for every $\alpha < \eta$. Collect then by induction a set $\{x_{\alpha} : \alpha < \eta\}$ such that

(26)
$$G_{\alpha}x_{\alpha} \cap G_{\beta}x_{\beta} = \emptyset$$
 for every $\alpha \neq \beta$.

This is possible since G is weakly left cancellative, and so

$$|G_{\alpha}^{-1}G_{\beta}| \le \max\{\alpha, \beta\} < \eta.$$

Then, let for each $\alpha < \eta$, A_{α} consist of all point-mass measures δ_x with $x \in G_{\alpha}$, i.e., $A_{\alpha} = \{\delta_{s_{\beta}} : \beta < \alpha\}$. Now by construction, the linear spans of each pair of sets $A_{\alpha} * \delta_{x_{\alpha}}$ have zero intersection. Moreover, since G is right cancellative, $xx_{\alpha} \neq yx_{\alpha}$ for every $x \neq y$ in G and $\alpha < \eta$, and so the supports of the point measures $\delta_x * \delta_{x_{\alpha}}$ (that is $\{xx_{\alpha}\}$) are disjoint for every $\alpha < \eta$ and $x \in A_{\alpha}$. So the elements $\delta_x * \delta_{x_{\alpha}}$ in \mathcal{L} are all distinct for distinct pairs $(x, \alpha) \in G_{\alpha} \times \eta$. It is then easy to see that

(27)
$$\mathcal{L} = \bigcup_{\alpha < \eta} (A_{\alpha} * \delta_{x_{\alpha}})$$

is in fact an $\ell^1(\eta)$ -base in $\ell^1(G)$ (with K = 1), and so it is an $F\ell^1(\eta)$ -base of type 1 in $\ell^1(G)$.

6.3. Algebraic structure. Recall first the following facts.

(i) In LUC(G)*, non-trivial left ideals of finite dimensions exist if and only if G is amenable, they are generated by elements defined by mixing topological left invariant means (TIMs) together with coefficients of finite dimensional representations of G (see [32] and [43]). A TIM is a positive linear functional Ψ in LUC(G)* or L¹(G)** with ⟨Ψ, 1⟩ = 1 and

(28)
$$a\Psi = \langle 1, a \rangle \Psi = \left(\int_G a(x)d\lambda \right) \Psi \text{ for every } a \in L^1(G).$$

Since there is one TIM (the left Haar measure) when G is compact, and there are $2^{2^{\eta}}$ TIMs in $\mathcal{LUC}(G)^*$ when G is non-compact and amenable, at least max{ $|\hat{G}_F|, 2^{2^{\eta}}$ } many distinct left ideals of finite dimension exit in $\mathcal{LUC}(G)^*$ when G is infinite and amenable. Here η is the compact covering of G and \hat{G}_F is the set of all finite-dimensional representations of G.

- (ii) In $L^1(G)^{**}$, there are two types of left ideals of finite dimension (and the mixture of both). The first type exists whenever G is not discrete. They are generated by elements Ψ in $\mathcal{LUC}(G)^{\perp}$. Since $\mathcal{LUC}(G) = L^{\infty}(G) \cdot L^{1}(G) =$ $L^{1}(G) * L^{\infty}(G)$, these elements are easily seen to satisfy $L^{1}(G)^{**}\Psi = \{0\}$. Since these elements may be seen as the difference of two distinct right identities in $L^1(G)^{**}$, and since there are at least $2^{2^{\omega}}$ right identities in $L^{1}(G)^{**}$ ([73], see also [51, Remak 4.6.2(3)]), we see that the number of left ideals of finite dimension in $L^1(G)^{**}$ is at least $2^{2^{\omega}}$ for any non-discrete group. The second type of finite-dimensional left ideals (which are not generated by annihilators of $\mathcal{LUC}(G)$) exists if and only if G is amenable; they are generated as in $\mathcal{LUC}(G)^*$ by elements defined by mixing TIMs together with coefficients of finite dimensional representations of G (see [32] and [43]). Since the number of TIMs in $L^1(G)^{**}$ is 1 when G is compact and $2^{2^{\eta}}$ when G is non-compact and amenable, we see that the number of finitedimensional left ideals in $L^1(G)^{**}$ is at least equal to $\max\{2^{2^w}, |\widehat{G}_F|, 2^{2^\eta}\}$ for any infinite locally compact group. For more details, see [32] and [43].
- (iii) In $M(G)^{**}$, A TIM is a positive linear functional $\Psi \in M(G)^{**}$ satisfying the identity (28) for every $a \in M(G)$. TIMs exist in $M(G)^{**}$ if and only if G is amenable, see for example [79, Corollary 4.3], [18, Propositions 1.21 and 5.10], or Proposition 1.4 in our recent article [42]. So again, arguing as for $\mathcal{LUC}(G)^*$ in [32] or applying the general theorem proved in [43, Theorem 2.8], non-trivial left ideals of finite dimensions exist in $M(G)^{**}$ if and only if G is amenable, they are generated by elements defined by mixing topological left invariant means (TIMs) together with coefficients of finite dimensional representations of G. Since the number of TIMs in $M(G)^{**}$ is 1 when G is compact and at least 2^{2^n} when G is non-compact and amenable, we deduce that at least $\max\{|\hat{G}_F|, 2^{2^n}\}$ many distinct left ideals of finite dimension exit in $M(G)^{**}$ when G is infinite and amenable.

Remark 6.3. The number of TIMs in $\mathcal{LUC}(G)^*$ and $L^1(G)^{**}$ being $2^{2^{\eta}}$ when G is non-compact and amenable was proved in [87] (see also [46]), but can also be deduced as follows form the proof of Statements (ii) of Theorem 6.4 below. The number of TIMs in $M(G)^{**}$ is also included in our simple argument. This latter result seems to be new.

Proof. Let G be a non-compact locally compact group with compact covering η or an infinite discrete weakly cancellative, right cancellative semigroup with cardinality η . Let $\{x_{\alpha} : \alpha < \eta\}$ be the set constructed in Theorems 6.1 and 6.2 and X be as in Notation 4.11. If G is amenable and Φ is a nonzero TIM in $\mathcal{LUC}(G)^*$, $L^1(G)^{**}$ or $M(G)^{**}$, then $\Phi\Psi_1$ and $\Phi\Psi_2$ are easily checked to be TIMs, and Theorem 6.4 shows that $\Phi\Psi_1$ and $\Phi\Psi_2$ are distinct whenever Ψ_1 and Ψ_2 are distinct in X. So the cardinality of X, which is 2^{2^n} , equals that of the TIMs. \Box

Having $F\ell^1(\eta)$ -bases of type 2 and type 1 in the group algebra and the semigroup algebra, respectively, an application of the theorems in Section 3 yield immediately the following theorems on the algebra in the related semigroup compactifications and the Banach algebras with an Arens product. This include many results obtained in earlier papers, see Notes 6.7 below. The left ideals provided in the theorems have

infinite dimension and are of different type than those we have just discussed at the beginning of this subsection.

Theorem 6.4. Let G be a non-compact locally compact group with a compact covering η or an infinite discrete weakly cancellative right cancellative semigroup with cardinality η . Let S be any of the semigroups G^{LUC} , βG when G is an infinite weakly cancellative right cancellative discrete semigroup, G^* , or any of the algebras $\mathcal{LUC}(G)^*, L^1(G)^{**}, M(G)^{**}, L_0^{\infty}(G)^{\perp}, C_0(G)^{\perp_{lu}} \text{ or } C_0(G)^{\perp_M}.$ Then

- (i) there are at least 2^{2ⁿ} many right cancellable elements in S.
 (ii) there are at least 2^{2ⁿ} many principal left ideals in S. These left ideals are pairwise disjoint in the cases of G^{LUC}, βG and G^{*}, and have pairwise intersections equal to zero in the other cases.

Proof. Let $\{x_{\alpha} : \alpha < \eta\}$ be the subsets of G, needed for the $F\ell^{1}(\eta)$ -bases in $L^{1}(G)$ of a non-compact locally compact group and in $\ell^1(G)$ of an infinite discrete weakly cancellative right cancellative semigroup, constructed in Theorems 6.1 and 6.2, and let X and X be as in Notation 4.11. Then the sets X and X are contained in G^* and $L_0^{\infty}(G)^{\perp}$, respectively, and so are the elements Ψ_p and $\widetilde{\Psi}_p$, used in the proofs of Statement (i) of Theorems 5.1 and 5.3.

Therefore, a direct application of Theorems 5.1 (i) and 5.3 (i) gives the claimed properties when S is G^{LUC} , βG when G is discrete, G^* , $\mathcal{LUC}(G)^*$ or $C_0(G)^{\perp_{lu}}$ since G^* is contained in $\mathbb S$ in each case.

These theorems yield also the claimed properties when S is $L^1(G)^{**}$ or $L^{\infty}_0(G)^{\perp}$ since \widetilde{X} is contained in S in each case.

The measure algebra M(G) is a dual algebra with predual $C_0(G)$. If $(\varphi_i) \subset$ $C_{00}(G)$ is a bai for $C_0(G)$, then its weak*-limit in $M(G)^*$ is the identity 1 (the constant function 1) in $M(G)^*$ (remember that $C_0(G)$ is Arens regular). Since again the set X when regarded as a subset of $M(G)^{**}$ is clearly contained in $C_0(G)^{\perp_M}$, all the conditions are satisfied for Theorem 5.4 (i) and (iii) to hold in $M(G)^{**}$ as well. \square

In our next theorem, the dimension of left ideals and right ideals is provided when S is an algebra.

Theorem 6.5. Let G be a non-compact locally compact group with a compact covering η or an infinite discrete weakly cancellative right cancellative semigroup with cardinality η . Let S be any of the algebras $\mathcal{LUC}(G)^*$, $L^1(G)^{**}$, $M(G)^{**}$, $L_0^{\infty}(G)^{\perp}$, $C_0(G)^{\perp_{lu}}$ or $C_0(G)^{\perp_M}$. Then

- (i) the dimension of any principal right ideal, and so of any nonzero right ideal, in S is at least $2^{2^{\eta}}$.
- (ii) the dimension of each of the $2^{2^{\eta}}$ principal left ideals found in Theorem 6.4 in S is at least $2^{2^{\eta}}$.

Proof. Let again $\{x_{\alpha} : \alpha < \eta\}$, X and X be as in the previous theorem. Since $X \subseteq G^*$ and $\widetilde{X} \subseteq L_0^\infty(G)^{\perp}$, a direct application of Theorem 5.2 and Statement (iii) in Theorem 5.3 gives the claimed properties when S is any of the algebras $\mathcal{LUC}(G)^*$. $L^{1}(G)^{**}, L^{\infty}_{0}(G)^{\perp} \text{ or } C_{0}(G)^{\perp_{lu}}.$

For $M(G)^{**}$ and $C_0(G)^{\perp_M}$, we apply Theorem 5.4 since $X \subseteq C_0(G)^{\perp_M}$.

Remark 6.6. The number of right ideals is more difficult. Our approach in this paper does not apply when counting the number of right ideals in the semigroups and algebras. But at this point, we may direct the reader to [4], [65] and [113], where the number of right ideals in βG and G^{LUC} was computed in certain cases. We shall return for further developments on this problem in our next paper [42].

Notes 6.7. For the right cancellation property in βG when G is an infinite discrete weakly cancellative right cancellative semigroup, the reader is directed to [34], [36] and [65, Chapter 8]. For the right cancellation property in G^{LUC} when G is a noncompact locally compact group, the reader is directed to [30], [46] and [47]. For the number of left ideals in βG when G is an infinite discrete weakly cancellative right cancellative semigroup, the reader is directed to [10], [35], [34], [36], [48] and [65, Chapter 6]. For the number of left ideals in G^{LUC} when G is a non-compact locally compact group, the reader is directed to [36], [46], [47], [48] and [86].

For the right cancellation property in $\mathcal{LUC}(G)^*$ and $L^1(G)^{**}$, the reader is directed to [47]. For the number of left ideals in the algebras $\mathcal{LUC}(G)^*$ and $L^1(G)^{**}$ the reader is directed to [46] and [47].

6.4. Topological centres. We look now at the topological centres of $L^1(G)^{**}$, $\mathcal{LUC}(G)^*$ and $M(G)^{**}$ when G is a locally compact group, of $\ell^1(G)^{**}$ when G is weakly cancellative right cancellative semigroup, and many of their subsemigroups.

Recall that the group algebra of any locally compact group with a compact covering η has Mazur property of level $\eta.w$ (see [71, Corollary 3.3.(i)]) or [102, Theorem 3.4]).

The measure algebra of any locally compact group G has Mazur property of level |G|.w (see [71, Corollary 5.6(i)]), and $|G| = \eta \cdot 2^{w(G)}$ (see [14, Theorem 3.12(iii)] or [71, Corollary 5.5]). Recall also from [71, Corollary 5.6] that M(G) has Mazur property of level w (i.e., the classical Mazur property) if and only if |G| is non-measurable. For more information and references related to measurable cardinals, see [101, page 219].

With the $F\ell^1(\eta)$ -bases found in Theorems 6.1 and 6.2, Theorems 3.5 and 3.8 yield the following theorems, which summarizes some of the main results proved on topological centres in harmonic analysis in the last thirty years, and includes some new results as well.

Theorem 6.8. Let G be a locally compact group with compact covering η and local weight w(G). Then

- (i) the first topological centre of $L^1(G)^{**}$ is $L^1(G)$.
- (ii) the first topological centre of $\mathcal{LUC}(G)^*$ is M(G).
- (iii) the first topological centre of $L_0^{\infty}(G)^{\perp}$ is $\{0\}$.
- (iv) the first topological centre of $C_0(G)^{\perp_{lu}}$ is $\{0\}$.
- (v) the first topological centre of G^{LUC} is G.
- (vi) the first topological centre of G^* is empty.
- (vii) the first topological centre of $M(G)^{**}$ is M(G) whenever $\eta \ge 2^{w(G)}$ or |G| is non-measurable.
- (viii) the first topological centre of $C_0(G)^{\perp_M}$ is $\{0\}$ whenever $\eta \ge 2^{w(G)}$ or |G| is non-measurable.

Proof. When G is compact, the first statement was proved in [73], and it may also be deduced from [5, Theorem 2.2]. The five statements which follow are trivial when G is compact. So let us assume that G is non-compact, and let $\{x_{\alpha} : \alpha < \eta\} \subset G$ be as in Theorem 6.1, X and \widetilde{X} be as in Notation 4.11. Having an $F\ell^{1}(\eta)$ -base of type 2 in the group algebra, Theorem 4.14 and Theorem 4.17 provide $L^{\infty}(G)$ and $M(G)^{*}$ with the η -factorization property whenever G is a non-compact locally compact group.

Since the η -factorization property and the η -Mazur property are both available in $L^{\infty}(G)$ and $L^{1}(G)$, respectively, the first two statements follow directly from Theorems 3.5 (ii) and 3.8 (iii).

Now since $X \subseteq G^* \subseteq C_0(G)^{\perp_{lu}}$ in the case of $\mathcal{LUC}(G)^*$ and $\widetilde{X} \subseteq L_0^{\infty}(G)^{\perp}$ in the case of $L^1(G)^{**}$, Statements (iii)–(vi) follow also directly from 3.5 and 3.8 and

the facts that

$$L_0^{\infty}(G)^{\perp} \cap L^1(G) = C_0(G)^{\perp_{lu}} \cap M(G) = \{0\},\$$

$$G^{LUC} \cap M(G) = G \quad \text{and} \quad G^* \cap M(G) = \emptyset.$$

Finally, note that under the conditions imposed in Statements (vii) and (viii), M(G) has the required Mazur property, which is of level η when $\eta \geq 2^{w(G)}$ and of level w when |G| is non-measurable. Note also that in the latter case, $M(G)^*$ has in particular by Theorems 4.17 and 6.1 the w-factorization property (since up to η -many functionals in $M(G)^*$ may be factorized). Therefore Statement (vii) follows from Theorem 3.5.

Now since $C_0(G)^{\perp_M} \cap M(G) = \{0\}$ and $X \subseteq C_0(G)^{\perp_M}$ (when regarded as a subset of $M(G)^{**}$), the last statement is also a corollary of Theorem 3.5.

Having $F\ell^1(\eta)$ -bases of type 1 in the semigroup algebra, we also obtain in the same way the first topological centres when G is an infinite discrete weakly cancellative right cancellative semigroup.

Theorem 6.9. Let G be an infinite discrete weakly cancellative right cancellative semigroup. Then

- (i) the first topological centre of $\ell^1(G)^{**}$ is $\ell^1(G)$.
- (ii) the first topological centre of $c_0(G)^{\perp}$ is $\{0\}$.
- (iii) the first topological centre of βG is G.
- (iv) the first topological centre of G^* is empty.

Remark 6.10. The mirror theorems of Theorem 6.8 and 6.9 are also true; that is, if "first" is replaced by "second" and "right" replaced by "left" in these theorems, then the conclusions stay unchanged. To see this when G is not compact, one needs to return to Theorems 6.1 and 6.2 and define inductively the sets $\{x_{\alpha} : \alpha < \eta\} \subset G$ with the property

$$x_{\alpha}G_{\alpha} \cap x_{\beta}G_{\beta} = \emptyset$$
 for every $\alpha \neq \beta$

instead of (23) and (26). Then consider the family of functions given by

$$\mathcal{L} = \bigcup_{\alpha < \eta} (\delta_{x_{\alpha}} * A_{\alpha})$$

instead of (24) and (27). Precisely as in Theorems 6.1 and 6.2, this gives a right $F\ell^1(\eta)$ -base in $L^1(G)$. Following the steps of the proof of Theorem 4.14, we see then that $L^{\infty}(G)$ has the right η -factorization according to Remark 3.2. Then we are only left to apply Theorems 3.6 and 3.10.

When G is compact, we may use [5, Theorem 2.2(ii)] where Baker, Lau and Pym proved that $\mathcal{Z}_1(\mathcal{A}^{**}) = \mathcal{A}$ if \mathcal{A} is weakly sequentially complete, contains a bounded approximate identity and is an ideal in its second dual. Their proofs can also be applied to show that $\mathcal{Z}_2(\mathcal{A}^{**}) = \mathcal{A}$. So these algebras, in particular $L^1(G)$ when G is compact, are sAir by [5, Theorem 2.2(ii)].

Corollary 6.11. Let G be a locally compact group with compact covering η and local weight w(G), or an infinite discrete weakly cancellative right cancellative semigroup. Then

- (i) $L^1(G)$ is strongly Arens irregular.
- (ii) M(G) is strongly Arens irregular whenever $\eta \geq 2^{w(G)}$ or |G| is non-measurable.

Notes 6.12.

(i) The compact case in Statement (i) of Theorem 6.8 is not a consequence of our approach (see Final remarks 9.7(ii) below). This statement is due to Isik, Pym and Ülger ([73]). The general Statement in (i) is due to Lau and Losert ([83]). Statement (ii) is due to Lau ([80]). For non-compact groups, these results were also reached in [17], [48] and [100]. Statement (iii) and (iv) were proved in [48]. Statement (v) was proved originally by Lau and Pym in [90], then by Protasov and Pym in [106] and by Filali and Salmi in [48]. Statement (vi) was proved by Protasov and Pym in [106] and Filali and by Salmi in [48].

Statement (vii) is due to Neufang ([101].) The general statement that the topological centre of $M(G)^{**}$ is M(G) without any condition has been recently solved by Losert et al in [96]. It is worthwhile to note that a careful look at the constructions made in [96] may help to construct an $F\ell^1(|G|.w)$ -base in M(G). With Theorem 4.17 and Theorem 3.5, this would give another proof to the latter result. We hope to get back to this problem in the future. Statement (viii) is new.

(ii) In Theorem 6.9, Statement (i) was proved by Lau in [80] under a slightly weaker cancellation property of the semigroup G, and reached also in [48]. Statement (ii) was proved in [48]. Concerning Statements (iii) and (iv), Davenport and Hindman proved in [19], that the algebraic centre of βG coincides with that of G, and the algebraic centre of G^* is empty, when G is an infinite discrete cancellative semigroup. Under a weaker cancellation property of G, these two statements were proved again by Hindman and Strauss in [65, Theorem 6.54]. Statements (iii) and (iv) were proved in [48].

7. Application II: Weighted group, semigroup and measure algebras

Our approach applies also to the weighted (semi)group algebra and the weighted measure algebra when the weight is diagonally bounded.

7.1. Some definitions. Recall that a *weight* on a semigroup G (with a topology) is a continuous function $w: G \to (0, \infty)$ which is submultiplicative, that is,

$$w(st) \le w(s)w(t)$$
 for every $s, t \in G$.

If G is a group with identity e, we shall assume that w(e) = 1. The weight function w is called *diagonally bounded* on G if there exists K > 0 such that

 $w(s)w(t) \le Kw(st)$ for every $s, t \in G$.

For the analogues of the theorems proved in the previous section, we recall the following weighted spaces:

- $L^{\infty}(G, w^{-1})$ consists of all Borel measurable functions S on G such that $w^{-1}S \in L^{\infty}(G)$.
- $L_0^{\infty}(G, w^{-1})$ is the subspace of those functions $S \in L^{\infty}(G, w^{-1})$ such that $w^{-1}S \in L_0^{\infty}(G)$.
- $\mathcal{LUC}(G, w^{-1})$ is the subspace of those functions $S \in L^{\infty}(G, w^{-1})$ such that $w^{-1}S \in \mathcal{LUC}(G)$.
- $C_0(G, w^{-1})$ is the subspace of those functions $S \in \mathcal{LUC}(G, w^{-1})$ such that $w^{-1}S \in C_0(G)$.

The norm of S in these weighted spaces is given by $||w^{-1}S||$ so that the space and its weighted space are isometric.

• The weighted group algebra $L^1(G, w)$ consists of all Borel measurable functions a on G such that $wa \in L^1(G)$.

- The weighted semigroup algebra $\ell^1(G, w)$ is defined in the same way when G is a discrete semigroup.
- The weighted measure algebra M(G, w) consists of all regular compacted-Borel measures a on G such that $wa \in M(G)$.

The norm of a in these weighted spaces is given by ||wa||, so that the space and its weighted space are isometric. The definition of M(G, w) has been rectified recently by Stokke in [109] by adding the adjective "compacted".

As it is the case when w = 1, both $L^1(G, w)$ and M(G, w) are Banach algebras under convolution since w is submultiplicative, and so the Arens products may be defined on $L^1(G, w)^{**}$ and $M(G, w)^{**}$. Moreover, as known $L^1(G, w)$ is a closed two-sided ideal of M(G, w). The weighted Wendel's theorem is also true, so that M(G, w) is the multiplier algebra of $L^1(G, w)$, see [53, Lemma 2.3] and [112, Theorem 3.2].

The spaces $L^{\infty}(G, w^{-1})$ and $L^1(G, w)^*$ are identified by the usual dual pairing. And since

$$\mathcal{LUC}(G, w^{-1}) = L^{\infty}(G, w^{-1}) \cdot L^1(G, w)$$

(see [60] for the thorem and [16, Proposition 7.15] for the corrected proof), the actions of $\mathcal{LUC}(G, w^{-1})^*$ on $L^{\infty}(G, w^{-1})$ and $L^1(G, w)^{**}$ may be defined as previously.

The Banach space decompositions used in the previous section are true in the weighted situation, the proofs are as in [16, 35, 54, 88].

(29)

$$\mathcal{LUC}(G, w^{-1})^* = M(G, w) \oplus C_0(G, w^{-1})^{\perp_{lu}},$$

$$L^{\infty}((G, w^{-1})^* = L_0^{\infty}(G, w^{-1})^* \oplus L_0^{\infty}(G, w^{-1})^{\perp},$$

$$M(G, w)^{**} = M(G, w) \oplus C_0(G, w^{-1})^{\perp_M},$$

where $C_0(G, w^{-1})^{\perp_{lu}}$, $L_0^{\infty}(G, w^{-1})^{\perp}$ and $C_0(G, w^{-1})^{\perp_M}$ are weak*-closed ideals in the corresponding algebras. The proofs are easily adapted from the unweighted cases.

We shall not be concerned as in the previous section with the spectrum of $\mathcal{LUC}(G, w^{-1})$. This is not a subsemigroup of $\mathcal{LUC}(G, w^{-1})$ unless w is a homomorphism, in which case this spectrum is isomorphic to G^{LUC} as a semigroup compactification. For the same reason, the spectrum of $\ell^{\infty}(G, w^{-1})$ when G is an infinite discrete semigroup is not included in our theorem.

7.2. $\mathbf{F}\ell^1(\eta)$ -bases.

Theorem 7.1. Let G be a non-compact locally compact group with compact covering η and let w be a weight on G that is diagonally bounded. Then the weighted group algebra $L^1(G, w)$ has an $F\ell^1(\eta)$ -base of type 2.

Proof. Let $\{V_{\alpha}\}_{\alpha < \eta}$, $\{x_{\alpha}\}_{\alpha < \eta}$ and $\{A_{\alpha}\}_{\alpha < \eta}$ be as in Example 6.1 and let

$$\mathcal{L} = \bigcup_{\alpha < \eta} \left(\frac{A_{\alpha}}{w} \right) * \frac{\delta_{x_{\alpha}}}{w(x_{\alpha})}.$$

We claim that \mathcal{L} is an $F\ell^1(\eta)$ -base in $L^1(G, w)$. To see this, we only need to check that the analogue of (25) in Example 6.1 is true, the rest is clear and follows as in Example 6.1.

Note first that for every $\varphi \in L^1(G)$, we have

$$\frac{\varphi}{w} * \frac{\delta_{x_{\alpha}}}{w(x_{\alpha})} = \frac{1}{w(x_{\alpha})} \Delta(x_{\alpha}^{-1}) \left(\frac{\varphi}{w}\right)_{x_{\alpha}^{-1}} \quad \text{(the right translate of } \frac{\varphi}{w} \text{ by } x_{\alpha}^{-1}\text{)}.$$

Let now φ be a function in the closed linear span of \mathcal{L} . Then φ has the form

$$\varphi = \sum_{n=1}^{p} \frac{\varphi_n}{w} * \frac{\delta_{x_{\alpha_n}}}{w(x_{\alpha_n})},$$

where each φ_n (n = 1, ..., p) is in the closed linear span of A_{α_n} . Accordingly,

$$\begin{split} \left\| \sum_{n=1}^{p} \left(\frac{\varphi_n}{w} * \frac{\delta_{x_{\alpha_n}}}{w(x_{\alpha_n})} \right) \right\|_{1,w} &= \left\| \sum_{n=1}^{p} w \left(\frac{\varphi_n}{w} * \frac{\delta_{x_{\alpha_n}}}{w(x_{\alpha_n})} \right) \right\|_1 \\ &= \sum_{n=1}^{p} \int_G w \left| \left(\frac{\varphi_n}{w} * \frac{\delta_{x_{\alpha_n}}}{w(x_{\alpha_n})} \right) \right| d\lambda \\ &= \sum_{n=1}^{p} \left\| \frac{\varphi_n}{w} * \frac{\delta_{x_{\alpha_n}}}{w(x_{\alpha_n})} \right) \right\|_{1,w} \\ &= \sum_{n=1}^{p} \int_G \left| w \frac{1}{w(x_{\alpha_n})} \Delta(x_{\alpha_n}^{-1}) \left(\frac{\varphi_n}{w} \right)_{x_{\alpha_n}^{-1}} \right| d\lambda \\ &= \sum_{n=1}^{p} \int_G \left| \frac{w(yx_{\alpha_n})}{w(y)w(x_{\alpha_n})} \varphi_n(y) \right| d\lambda(y) \\ &\geq \sum_{n=1}^{p} \int_G \frac{1}{K} |\varphi_n(y)| d\lambda(y) = \frac{1}{K} \sum_{n=1}^{p} \left\| \frac{\varphi_n}{w} \right\|_{1,w} \end{split}$$

where as in Theorem 6.1, the second equality follows from the fact that the functions under the sum have disjoint supports. The third equality proves then that \mathcal{L} is selective. The inequality is due to the diagonal boundedness of w, and checks the validity of Condition (F2) in Definition 4.8. For, this inequality simply becomes

$$K \left\| \frac{\varphi}{w} * \frac{\delta_{x_{\alpha}}}{w(x_{\alpha})} \right\|_{1,w} \ge \left\| \frac{\varphi}{w} \right\|_{1,w}$$

when p = 1. As in the previous example, note that Condition (f3) (and so Condition (F3) in Definition 4.9) is also valid here with m = 1, a_{α} and b_{α} may be taken as $\frac{\delta_{x_{\alpha}}}{w(x_{\alpha})}$ and $w(x_{\alpha})\delta_{x_{\alpha}^{-1}}$, respectively, where $||a_{\alpha}||_{w} = 1$ and $||b_{\alpha}||_{w} = w(x_{\alpha})w(x_{\alpha}^{-1}) \leq K$ for every $\alpha < \eta$ since w is diagonally bounded by K.

Theorem 7.2. Let G be an infinite discrete weakly cancellative right cancellative semigroup with cardinality η and let w be a weight on G that is diagonally bounded. Then the weighted semigroup algebra $\ell^1(G, w)$ has an $F\ell^1(\eta)$ -base of type 1.

Proof. Proceed as in Theorems 6.2 and 7.1.

7.3. Algebraic structure. As in the previous section, having $F\ell^1(\eta)$ -bases, the theorems in Section 3 yield immediately the weighted analogues to Theorems 6.4 and 6.5. The proofs are precisely the same. Both analogues are summarized in the theorem below. The parts on the number of left ideals and the dimension of right ideals in $L^1(G, w)^{**}$ and $\mathcal{LUC}(G, w^{-1})^*$ appeared in [47]. The rest is new.

Theorem 7.3. Let G be a non-compact locally compact group with compact covering η or an infinite discrete weakly cancellative right cancellative semigroup with cardinality η , and let w be a diagonally bounded weight on G. Let S be any of the algebras $L^1(G, w)^{**}$, $\mathcal{LUC}(G, w^{-1})^*$, $M(G, w)^{**}$, $L_0^{\infty}(G, w^{-1})^{\perp}$, $C_0(G, w^{-1})^{\perp_{lu}}$ or $C_0(G, w^{-1})^{\perp_M}$. Then

- (i) there are at least $2^{2^{\eta}}$ many right cancellable elements in S.
- (ii) there are at least 2^{2ⁿ} many principal left ideals in S with pairwise intersections equal to zero, the dimension of each is at least 2^{2ⁿ}.

(iii) the dimension of any principal right ideal, and so of any nonzero right ideal, in S is at least $2^{2^{\eta}}$.

7.4. Topological centres. We turn again to topological centres. Now that we have $F\ell^1(\eta)$ -bases in $L^1(G, w)$ when G is a non-compact locally compact group, and of $\ell^1(G, w)$ when G is an infinite discret weakly cancellative, right cancellative semigroup, Theorems 4.14 and 4.17 provide $L^{\infty}(G, w^{-1})$ and $M(G, w)^*$ with the η -factorization property.

Since the Mazur property and its level are preserved under isomorphisms (as already observed by Neufang in [103, Remark 3.4] and Hu and Neufang in [71, Remark 1.5]), the weighted group algebra and the weighted measure algebra have each Mazur property of level $\eta.w$ and |G|.w, respectively. Theorems 4.14 and 4.17 together with Theorem 3.5 and 3.8 yield therefore, in our next theorem, the weighted analogues of Theorems 6.8 and 6.9.

Theorem 7.4. Let G be a locally compact group or a weakly cancellative right cancellative semigroup, and let w be a weight on G that is diagonally bounded. Then

- (i) the first topological centre of $L^1(G, w)^{**}$ is $L^1(G, w)$.
- (ii) the first topological centre of $\mathcal{LUC}(G, w^{-1})^*$ is M(G, w).
- (iii) the first topological centre of $M(G, w)^{**}$ is M(G, w) whenever $\eta \geq 2^{w(G)}$ or |G| is non-measurable.
- (iv) the first topological centre of $L_0^{\infty}(G, w^{-1})^{\perp}$ is $\{0\}$. (v) the first topological centre of $C_0(G, w^{-1})^{\perp_{lu}}$ is $\{0\}$.
- (vi) the first topological centre of $C_0(G, w^{-1})^{\perp_M}$ is $\{0\}$ whenever $\eta \geq 2^{w(G)}$ or |G| is non-measurable.

Proof. When G is compact, Statement (i) follows from [5, Theorem 2.2] since $L^{1}(G, w)$ is weakly sequentially complete, contains a bounded approximate identity and is an ideal in its second dual. Statements (ii), (iv) and (iv) are trivial when Gis compact. So let us assume that G is non-compact. Then again the proofs of the first three statements are exactly the same as those given for Theorem 6.8.

Since $X \subseteq C_0(G, w^{-1})^{\perp_{lu}}$ in the case of $\mathcal{LUC}(G, w^{-1})^*$, $\widetilde{X} \subseteq L_0^{\infty}(G, w^{-1})^{\perp}$ in the case of $L^1(G, w)^{**}$ and $X \subseteq C_0(G, w^{-1})^{\perp_M}$ in the case of $M(G, w)^{**}$, and since $L_0^{\infty}(G, w^{-1})^{\perp} \cap L^1(G, w) = C_0(G, w^{-1})^{\perp_{lu}} \cap M(G) = C_0(G, w^{-1})^{\perp_M} \cap M(G) = \{0\},\$

the last three statements of the theorem follow as in Theorem 6.8 as well.

As in Remark 6.10, the mirror theorem of Theorem 7.4 holds. In particular, we obtain the following corollary.

Corollary 7.5. Let G be a locally compact group with compact covering η and local weight w(G), or an infinite discrete weakly cancellative right cancellative semigroup with cardinality η , and let G have a diagonally bounded weight w. Then

- (i) $L^1(G, w)$ is strongly Arens irregular.
- (ii) M(G, w) is strongly Arens irregular whenever $\eta \geq 2^{w(G)}$ or |G| is nonmeasurable.

Notes 7.6. Statements (i) in Theorem 7.4 and Corollary 7.5 are due to Neufang in [103] for non-compact groups, Dales-Lau in [16] for locally compact compact groups with non-measurable compact covering, and Filali-Salmi in [49] for noncompact SIN-groups. Statement (ii) in Theorem 7.4 was also proved for locally compact compact groups in [103], [16] and [49]. Statement (iv) was proved in [49] when G is a non-compact SIN-group. Statement (v) in Theorem 7.4 was proved in [49]. The rest of the statements are new. Again, as noted earlier, a careful study of [96] may help to find an $F\ell^1(|G|.w)$ -base in M(G,w), which would yield Statements (iii) and (vi) and Corollary 7.5(ii) without any condition.

8. Application III: Fourier Algebra

This section deals with the Fourier algebra of compact non-metrizable groups, infinite products of compact groups and compact connected groups with infinite dual rank. The dual rank will be defined in time.

8.1. Some definitions and lemmas. Recall that the Fourier algebra is the collection of all functions h on G of the form $\overline{f} * \check{g}$ with $f, g \in L^2(G)$ and norm

$$||h|| = \inf \left\{ ||f||_2 ||g||_2 : h = \bar{f} * \check{g}, \ f, g \in L^2(G) \right\}.$$

The Banach dual of A(G) is the group von Neumann algebra VN(G) which is the closure in the weak operator topology of the linear span of $\{\lambda(x) : x \in G\}$ in $B(L^2(G))$, where λ is the left regular representation of G on $L^2(G)$ (see [29]).

A subspace of VN(G) which shall be of our interest is $UC_2(G)$. This is defined by

$$\mathfrak{UC}_2(G) = \overline{VN(G) \cdot A(G)}^{||.||} = \overline{\{T \in VN(G) : \operatorname{supp}(T) \text{ is compact}\}}^{||.||},$$

where the support $\operatorname{supp}(T)$ of T is in the sense of [29, page 227],

 $\operatorname{supp}(T) = \{ x \in G : u \in A(G) \text{ and } u \cdot T = 0 \implies u(x) = 0 \}.$

Recall that $VN(G) \cdot A(G)$ is automatically closed when G is amenable. This is usually called the space of uniformly continuous functionals on G, and is often denoted in the literature by $UC(\widehat{G})$. We are using the notation $\mathcal{UC}_2(G)$ to avoid possible confusion arising from the usage of the dual group \widehat{G} of G. The space $\mathcal{UC}_2(G)$ is in fact a C*-subalgebra of VN(G), proved by Granirer in [56, Proposition 2(a)]. This is obtained by regarding $\mathcal{UC}_2(G)$ as the space generated by operators in VN(G) with compact support and using [29, Proposition 4.8 (5°)]. In the subsequent paper [57, Theorem 3], Granirer proved that $\mathcal{UC}_2(G) = VN(G)$ if and only if G is compact.

Following [29], let P(G) be the space of continuous positive definite functions on G and B(G) be its linear span, equivalently, the space of all coefficient functions of continuous unitary representations of G. The space B(G) is a Banach algebra, called the Fourier-Stieltjes algebra, and if $C^*(G)$ is the group C^* -algebra of G, then B(G) is its Banach dual. The Fourier algebra A(G) is a closed ideal of B(G) and can be seen as the norm-closed linear span of $P(G) \cap C_{00}(G)$ in B(G), where $C_{00}(G)$ be the space of continuous functions with compact support.

As known B(G) is the multiplier algebra of A(G) if and only if G is amenable, see [105, page 209] for the sufficiency, [92] for the necessity, and [98] when G is discrete.

When G is a compact group, as it is the case in this section, B(G) = A(G) and so $UC_2(G) = VN(G) = B(G)^*$.

For more details on the Fourier algebra and the Fourier-Stieltjes algebra on locally compact groups, we direct the reader to the classical article [29] by Eymard and to the recent book [75] by Kaniuth and Lau.

When G is abelian and \widehat{G} is its dual group, then via the Fourier transform, A(G) identifies with $L^1(\widehat{G})$, VN(G) with $L^{\infty}(\widehat{G})$, $\mathcal{UC}_2(G)$ with $\mathcal{LUC}(\widehat{G})$, P(G) with the set of Fourier transforms of all probability measures on \widehat{G} and B(G) with $M(\widehat{G})$. So all the results stated for the group algebra and the measure algebra hold for the Fourier algebra and the Fourier-Stieltjes algebra, respectively, when G is abelian. However, unlike the case of $L^1(G)$, non-Arens regularity of A(G) has turned out to be more resistant in general, and the solution is still not complete.

We denote by $C^*(G)^{\perp}$ the annihilator space of $C^*(G)$ in $\mathcal{UC}_2(G)^*$ when $C^*(G)$ is regarded as a subalgebra of $\mathcal{UC}_2(G)$, i.e.,

 $C^*(G)^{\perp} = \{ \Psi \in \mathcal{UC}_2(G)^* : \langle \Psi, f \rangle = 0 \text{ for every } f \in C^*(G) \}.$

Recall also that we have the following decomposition of $\mathcal{UC}_2(G)^*$ as Banach space direct sum

$$\mathfrak{UC}_2(G)^* = B(G) \oplus C^*(G)^{\perp},$$

where $C^*(G)^{\perp}$ is a weak*-closed ideal in $\mathcal{UC}_2(G)^*$ (see for example [22, Corollary 3.5]).

For a locally compact group G, let $\mathcal{U}(G)$ be the set of all unitary representations of G on some Hilbert space \mathbb{H} , \widehat{G} be the dual object of G, that is the set of all equivalence classes of irreducible unitary representations of G on some Hilbert space \mathbb{H} , and $\mathcal{U}(\mathbb{H})$ be the space of all unitary operators on \mathbb{H} . If $\pi \in \mathcal{U}(G)$, \mathbb{H}_{π} will denote the representation space, i.e., $\pi: G \to \mathcal{U}(\mathbb{H}_{\pi})$. If $v \in \mathbb{H}_{\pi}$, we define the matrix coefficients $\pi^{uv}: G \to \mathbb{C}$ as usual by $\pi^{uv}(s) = \langle \pi(s)u, v \rangle$. The diagonal matrix coefficients will be denoted simply by π^{v} .

When the representation π has finite dimension n with a fixed base $\{u_1, ..., u_n\}$ say, we simply write π^{kl} , $1 \le k, l \le n$ for the matrix coefficients of π .

The \mathcal{Z} -**property** We shall prove first that $F\ell^1(\eta)$ -bases exist in the Fourier algebra when the group satisfies a certain property we shall call the \mathcal{Z} -property. We will see then that a large class of locally compact groups has this property.

Definition 8.1. Let G be a locally compact group. We say that G has a \mathbb{Z} -property of level η when its dual \widehat{G} has an upward directed cover $\{K_{\alpha}\}_{\alpha < \eta}^{1}$ together with a set of irreducible representations $\{\pi_{\alpha} : \alpha < \eta\}$ such that

$$K_{\alpha} \otimes \pi_{\alpha} \cap K_{\beta} \otimes \pi_{\beta} = \emptyset$$
 for every $\alpha < \beta < \eta$.

The following lemma was proved by Arsac for any locally compact group ([3, 3ème partie, B-C], or see [75, Section 2.8]). The lemma will be needed in each of our constructions of $F\ell^1(\eta)$ -bases in this section. When the group is compact (as it is in most of the cases dealt with in this section), the interested reader can easily reproduce the proof using Schur's orthogonality relations.

Lemma 8.2 (Arsac). Let G be a locally compact group, $\{\pi_{\alpha} : \alpha < \eta\}$ be a set of disjoint unitary representations on G and let $\pi = \bigoplus_{\alpha < \eta} \pi_{\alpha}$. Then

- (i) the closed linear spans of the coefficients of π_{α} have zero pairwise intersections.
- (ii) each φ is the closed linear span of the coefficients of π can be written uniquely as $\sum_{\alpha < \eta} \varphi_{\alpha}$, where each φ_{α} is in the closed linear span of the coefficients of π_{α} and

$$\|\varphi\|_{A(G)} = \sum_{\alpha < \eta} \|\varphi_{\alpha}\|_{A(G)}.$$

Lemma 8.3. Let G be a compact group with a \mathbb{Z} -property of level η , and let $\{\pi_{\alpha} : \alpha < \eta\}$ be the family of representations required for the \mathbb{Z} -property for the cover $\{K_{\alpha}\}_{\alpha < \eta}$ of \widehat{G} . For each $\alpha < \eta$, let

$$A_{\alpha} = \{ \sigma^{uv} : \sigma \in K_{\alpha}, u, v \in \mathbb{H}_{\sigma} \}.$$

¹The meaning of upward directed is: given α_1, α_2 there is $\alpha_3 > \alpha_1, \alpha_2$ such that $K_{\alpha_3} \supset K_{\alpha_1} \cup K_{\alpha_2}$.

(i) If for some constant $K_2 > 0$, for every $\alpha < \eta$, there exist $1 \le k, l \le \dim \pi_{\alpha}$ such that

(30)
$$\|\varphi\|_{A(G)} \le K_2 \|\varphi\pi_\alpha^{kl}\|_{A(G)} \quad \text{for every} \quad \varphi \in \langle A_\alpha \rangle,$$

then A(G) has an $F\ell^1(\eta)$ -base of type 2 with constants $K_1 = 1$ and $K_2 = K$.

(ii) If there are η -many representations π_{α} with a fixed dimension m, then A(G) has an $F\ell^{1}(\eta)$ -base of type 3 with constants $K_{1} = 1$ and $0 < K_{2} \leq m$.

Proof. For each $\alpha < \eta$, for simplicity and for keeping the notation used in Section 3, let for each $\alpha < \eta$, a_{α} be any coefficient of the representation π_{α} and let

$$\mathcal{L} = \bigcup_{\alpha < \eta} A_{\alpha} a_{\alpha}$$

We claim first that \mathcal{L} is a zooming $\ell^1(\eta)$ -selective base in A(G). For this, we need to check that the spans of the sets $A_{\alpha}a_{\alpha}$ have zero pairwise intersections as Conditions (i) and (ii) in Definition 4.6 are clearly valid.

If $\sigma_1 \in K_{\alpha}$ and $\sigma_2 \in K_{\beta}$ for $\alpha < \beta < \eta$, then

$$\sigma_1 \otimes \pi_\alpha \in K_\alpha \otimes \pi_\alpha$$
 and $\sigma_2 \otimes \pi_\beta \in K_\beta \otimes \pi_\beta$,

and so Condition (ii) of the \mathcal{Z} -property assures that

$$\sigma_1 \otimes \pi_\alpha \cap \sigma_2 \otimes \pi_\beta = \varnothing.$$

Therefore, by Lemma 8.2, the spans of the sets $A_{\alpha}a_{\alpha}$, $\alpha < \eta$, have zero pairwise intersections.

We still need check that Condition (10) in Definition 4.3 holds. Let $b \in \langle \mathcal{L} \rangle$. Then we may write b as

$$b = \sum_{n=1}^{p} \varphi_n a_{\alpha_n}$$
 with $\varphi_n \in \langle A_{\alpha_n} \rangle$, for $n = 1, ..., p$,

with $\alpha_1 < \alpha_2 < \ldots < \alpha_p$.

Since the sets of unitary representations $K_{\alpha} \otimes \pi_{\alpha}$ are disjoint, Lemma 8.2 implies that

$$\|b\| = \left\|\sum_{n=1}^{p} \varphi_n a_{\alpha_n}\right\| = \sum_{n=1}^{p} \|\varphi_n a_{\alpha_n}\|.$$

We conclude that \mathcal{L} is a zooming $F\ell^1(\eta)$ -selective base with constant $K_1 = 1$. In particular, this holds when $a_{\alpha} = \pi_{\alpha}^{kl}$, for each $\alpha < \eta$, where π_{α}^{kl} are the coefficients for which Condition (30) in the lemma holds.

The additional condition (30) makes $\bigcup_{\alpha < \eta} A_{\alpha} \pi_{\alpha}^{kl}$ an $F\ell^{1}(\eta)$ -base of type 2 with constant K_{2} , proving the first statement.

For the second statement, suppose that each representation in $\{\pi_{\alpha} : \alpha < \eta\}$ has dimension m and denote π_{α}^{1k} and $\overline{\pi_{\alpha}^{k1}}$ by a_{α}^{k} and b_{α}^{k} , respectively, where k = 1, ..., m. Then as in Statement (i),

$$\mathcal{L} = \bigcup_{\alpha < \eta} A_{\alpha} \{ a_{\alpha}^k : k = 1, ..., m \}$$

is a zooming $\ell^1(\eta)$ -selective base with constant $K_1 = 1$. Clearly,

$$\sum_{k=1}^m a_\alpha^k b_\alpha^k = 1 \text{ for each } \alpha < \eta,$$

giving Condition (f3) in Remark 4.12(ii), and so by Remark 4.12(ii), we have an $F\ell^1(\eta)$ -base of type 3 with constants

$$K_1 = 1$$
 and $K_2 = \sup_{\alpha} \|(\overline{\pi_{\alpha}^{k1}})_{k=1}^m\|_p = \sup_{\alpha} \|(\pi_{\alpha}^{1k})_{k=1}^m\|_p \le m.$

8.2. $\mathbf{F}\ell^1(\eta)$ -bases for compact non-metrizable group. We start with a lemma, which is needed for the construction of $F\ell^1(\eta)$ -bases of type 3 in A(G) when G is compact, non-metrizable with a local weight η having uncountable cofinality. The lemma is due to Hu [67, Proposition 4.3], it is a stronger version of an earlier result of Lau–Losert [84, Lemma 4.8] whose proof is based on Kakutani-Kodeira Theorem (see for example [63, 8.7]). With the $F\ell^1(\eta)$ -bases, we shall simplify and improve in Subsections 9.1, and 9.2 the main results on strong Arens irregularity of and algebraic structure proved in [44] and [45].

Lemma 8.4 (Hu). Let G be a σ -compact non-metrizable locally compact group with unit element e and local weight η , and let $\{O_{\alpha}: 0 \leq \alpha < \eta\}$ be an open base at e. Then there exists a decreasing family $(N_{\alpha})_{0 \leq \alpha \leq \eta}$ of normal subgroups of G (that is, $N_{\beta} \subset N_{\alpha}$ whenever $\alpha \leq \beta$) such that

- (i) $N_0 = G \text{ and } N_\eta = \{e\},\$
- (ii) $N_{\alpha+1} \subset N_{\alpha} \cap O_{\alpha}$ for all $0 \le \alpha < \eta$,
- (iii) N_{α} is compact for each $\alpha > 0$,
- (iv) $N_{\alpha}/N_{\alpha+1}$ is metrizable but $N_{\alpha+1} \neq N_{\alpha}$ for all $\alpha < \eta$,
- (v) $N_{\gamma} = \bigcap_{\alpha < \gamma} N_{\alpha}$ for every limit ordinal $\gamma < \eta$,
- (vi) the local weight of N_{α} is η for each $\alpha < \eta$.

Note that the condition in the following theorem on the cofinality of η being uncountable is equivalent to \hat{G} containing η many representations of a fixed dimension m. For, since each π_{α} has finite dimension, we may define $E_m = \{\alpha : \dim \pi_{\alpha} = m\}$. Then $\bigcup_{m=1}^{\infty} E_m = \{\alpha : \alpha < \eta\} = \eta$. So the condition that the cofinality of η being uncountable implies that the cardinality of E_m is η for at least one m ([74, Lemma 3.6]). The converse is clear.

Theorem 8.5. Let G be a compact non-metrizable group with a local weight η and suppose that η has uncountable cofinality. Then A(G) has an $F\ell^1(\eta)$ -base of type 3 with constants $K_1 = 1$ and $0 < K_2 \leq m$ for some $m \in \mathbb{N}$.

Proof. Let $\{N_{\alpha}\}$ be the family of normal subgroups of G given by Hu's Lemma. For each $\alpha < \eta$, put

$$K_{\alpha} = \{ \sigma \in G : N_{\alpha} \subseteq \ker \sigma \},\$$

and note that $K_{\alpha} \subseteq K_{\beta}$ if $\alpha \leq \beta < \eta$. We claim first that $\widehat{G} = \bigcup_{\alpha < \eta} K_{\alpha}$. Let $\sigma : G \to U(\mathbb{H}_{\sigma})$ be any representation in \widehat{G} . Let U be any neighbourhood of the identity in $U(\mathbb{H}_{\sigma})$. We can assume that U contains no nontrivial subgroups (see, e.g., [66, Corollary 2.40]). Then $\sigma^{-1}(U)$ is a neighbourhood of the identity in G and so it contains some basic neighbourood O_{α} of the identity in G. Thus there exists some $\alpha < \eta$ such that

$$N_{\alpha+1} \subseteq \sigma^{-1}(U).$$

Therefore, $\sigma(N_{\alpha+1})$ is a subgroup of $U(\mathbb{H}_{\sigma})$ contained in U. Our choice of U ensures then that $N_{\alpha+1} \subseteq \ker \sigma$, i.e., $\sigma \in K_{\alpha+1}$, as required.

Next we define representations $\pi_{\alpha} \in \widehat{G}$ such that

 $\sigma_1 \otimes \pi_\alpha \cap \sigma_2 \otimes \pi_\beta = \emptyset$ for every $\sigma_1 \in K_\alpha, \ \sigma_2 \in K_\beta, \ \beta < \alpha < \eta$

so that G has the required \mathcal{Z} -property of level η . To do so, start for each $\alpha < \eta$ with any non-trivial irreducible representation $\tilde{\pi}_{\alpha}$ on $N_{\alpha}/N_{\alpha+1}$ and extend it to an irreducible representation π^*_{α} on N_{α} , where $\pi^*_{\alpha} = \tilde{\pi}_{\alpha} \circ q_{\alpha}$ and $q_{\alpha} : N_{\alpha} \to N_{\alpha}/N_{\alpha+1}$ is the quotient mapping. This latter representation is then extended, as in [63, Theorem 27.46], to an irreducible representation π_{α} on G.

Let $\alpha > \beta$ and suppose otherwise that for some $\sigma_1 \in K_{\alpha}$ and $\sigma_2 \in K_{\beta}$, there is $\rho \in \widehat{G}$ such that

$$\rho \in (\sigma_1 \otimes \pi_\alpha) \cap (\sigma_2 \otimes \pi_\beta).$$

Then

$$\rho_{|N_{\alpha}} \in (\sigma_{1|N_{\alpha}} \otimes \pi_{\alpha|N_{\alpha}}) \cap (\sigma_{2|N_{\alpha}} \otimes \pi_{\beta|N_{\alpha}}).$$

Now $N_{\alpha} \subseteq N_{\beta+1} \subset N_{\beta}$ (since $\alpha > \beta$), so $\sigma_{1|N_{\alpha}} = \text{Id}$ and $\sigma_{2|N_{\alpha}} = \text{Id}$ by the definition of K_{α} . We also note that $\pi_{\beta|N_{\alpha}} = \text{Id}$ since by the definition of π_{β} , $\pi_{\beta|N_{\beta+1}} = \text{Id}$. Thus, $\rho_{|N_{\alpha}} = \text{Id}$, and consequently,

$$\mathrm{Id} \in \sigma_{1|N_{\alpha}} \otimes \pi_{\alpha|N_{\alpha}} = \mathrm{Id} \otimes \pi_{\alpha|N_{\alpha}} = \pi_{\alpha|N_{\alpha}} \oplus \ldots \oplus \pi_{\alpha|N_{\alpha}}$$

Now $\pi_{\alpha|N_{\alpha}}$ is not necessarily irreducible but it decomposes into a finite direct sum of irreducible ones; and the proof of Theorem 27.48 in [63] confirms that the original representation π_{α}^* of N_{α} appears in the decomposition. Thus $\mathrm{Id} = \pi_{\alpha}^*$, which goes against our choice of π_{α}^* . Therefore, $\sigma_1 \otimes \pi_{\alpha}$ and $\sigma_2 \otimes \pi_{\beta}$ must be disjoint for every $\sigma_1 \in K_{\alpha}, \sigma_2 \in K_{\beta}, \beta < \alpha < \eta$. Therefore, G has a Z-property of level η .

Now we go after Condition (F3) in Definition 4.9. Since each π_{α} has finite dimension, we may define $E_m = \{\alpha : \dim \pi_{\alpha} = m\}$ so that

$$\bigcup_{m=1}^{\infty} E_m = \{ \alpha \colon \alpha < \eta \} = \eta.$$

As observed before the statement of the theorem, by the condition that the cofinality of η being uncountable, we may take for some $m \in \mathbb{N}$, a subset E_m of η of cardinality η such that dim $\pi_{\alpha} = m$ for each $\alpha \in E_m$. Then by replacing the family of subgroups $\{N_{\alpha} : \alpha < \eta\}$ and its associated family of representations $\{\pi_{\alpha}^* : \alpha < \eta\}$ with those that are indexed by the set E_m , we obtain a family of irreducible unitary representations, which for simplicity we still denote by $\{\pi_{\alpha} : \alpha < \eta\}$, such that dim $\pi_{\alpha} = m$ for all $\alpha < \eta$. Since this new family of representations still provides Gwith a \mathcal{Z} -property of level η , we may proceed as in Lemma 8.3 (ii) to complete the proof. The $F\ell^1(\eta)$ -base so obtained has constants $K_1 = 1$ and $K_2 \leq m$.

8.3. $\mathbf{F}\ell^1(\eta)$ -bases for infinite products of compact groups. Our second task will be first with infinite products of compact groups. The case of countably infinite products of metrizable compact groups with a second countable locally compact amenable group was considered by Lau and Losert in [85]. We prove that infinite products of η many compact groups have the \mathcal{Z} -property of level η and satisfy Condition (30) in Lemma 8.3 (i), and so by this lemma the Fourier algebra of such groups contains an $\mathbf{F}\ell^1(\eta)$ -base of type 2. With the $\mathbf{F}\ell^1(\eta)$ -bases, we shall simplify and improve in Subsection 9.2 the main result on strong Arens irregularity proved in [85]. In addition to this, we shall also deduce in Subsection 9.1 some rich algebraic structures in the algebras.

We need first the following simple lemma.

Lemma 8.6. Let G be a locally compact group and suppose that G = LM for some subgroups L and M of G. Let $\pi : G \to U(\mathbb{H}_{\pi})$ be a representation of G with $\pi_{|M} = \mathrm{Id}$ (or $\pi_{|L} = \mathrm{Id}$). Then π is irreducible if and only if $\pi_{|L}$ ($\pi_{|M}$, respectively) is irreducible.

Proof. Let X be a closed subspace of \mathbb{H}_{π} . If X is invariant under π , then it is clearly so under $\pi_{|L}$ and $\pi_{|M}$. For the converse, suppose that $\pi_{|M} = \mathrm{Id}$ and X is invariant under $\pi_{|L}$. Then, writing each $s \in G$ as s = lm for some $l \in L$ and $m \in M$, we see that

 $\pi(s)(X) = \pi(lm)(X) = \pi(l)(X) \quad \text{for every} \quad s \in G,$

which shows that X is invariant under π . The proof is the same in the other situation.

Theorem 8.7. Let $\{G_{\alpha}\}_{\alpha < \eta}$ be an infinite family of non-trivial compact groups and let $G = \prod_{\alpha < \eta} G_{\alpha}$. Then A(G) has an $F\ell^{1}(\eta)$ -base of type 2 with constants $K_{1} = K_{2} = 1$.

Proof. First we prove first that \widehat{G} has a family of representations $\{\pi_{\alpha}\}_{\alpha < \eta}$ with the \mathcal{Z} -property. For each $\alpha < \eta$, let e_{α} be the identity in G_{α} . Let L_{α} and M_{α} be the subgroups given by

$$L_{\alpha} = \{e_{\gamma}\}_{\gamma < \alpha} \times \prod_{\gamma \ge \alpha} G_{\gamma} \quad \text{and} \quad M_{\alpha} = \prod_{\gamma < \alpha} G_{\gamma} \times \{e_{\gamma}\}_{\gamma \ge \alpha}.$$

Consider, for each $\alpha < \eta$, the set

$$K_{\alpha} = \{ \sigma \in \widehat{G} : L_{\alpha} \subseteq \ker \sigma \}.$$

Note that $K_{\alpha} \subseteq K_{\beta}$ if $\alpha \leq \beta < \eta$. We claim first that $\widehat{G} = \bigcup_{\alpha < \eta} K_{\alpha}$. Let

$$\sigma: \prod_{\alpha < \eta} G_{\alpha} \to U(\mathbb{H}_{\sigma})$$

be any representation in \widehat{G} . As in Theorem 8.5, let U be any neighbourhood of the identity in $U(\mathbb{H}_{\sigma})$ that contains no nontrivial subgroups. Then $\sigma^{-1}(U)$ is a neighbourhood of the identity in $\prod_{\alpha < \eta} G_{\alpha}$ and so it contains some basic neighbourhood of the identity in $\prod_{\alpha < \eta} G_{\alpha}$. Thus there exists some $\alpha < n$ such that

the identity in $\prod_{\alpha < \eta} G_{\alpha}$. Thus there exists some $\alpha < \eta$ such that

$$\{e_{\gamma}\}_{\gamma < \alpha} \times \prod_{\gamma \ge \alpha} G_{\gamma} \subseteq \sigma^{-1}(U),$$

that is, $L_{\alpha} \subseteq \sigma^{-1}(U)$. Therefore, $\sigma(L_{\alpha})$ is a subgroup of $U(\mathbb{H}_{\sigma})$ contained in U. Since U contains no nontrivial subgroups, our claim that $L_{\alpha} \subseteq \ker \sigma$ follows.

Denote now the projections of G onto G_{α} by pr_{α} , and take for each $\alpha < \eta$, a non-trivial representation $\pi^*_{\alpha} \in \widehat{G_{\alpha}}$.

Then, define for each $\alpha < \eta$, a representation π_{α} on G simply by $\pi_{\alpha}^* \circ pr_{\alpha}$, and consider the family $\{\pi_{\alpha} : \alpha < \eta\}$. To check that this family of cardinality η provides G with the \mathcal{Z} -property, suppose otherwise that for some $\alpha < \beta < \eta$, $\sigma_1 \in K_{\alpha}$ and $\sigma_2 \in K_{\beta}$, we have

$$\sigma_1 \otimes \pi_\alpha \cap \sigma_2 \otimes \pi_\beta \neq \emptyset,$$

and let $\rho \in \widehat{G}$ such that

$$\rho \leq \sigma_1 \otimes \pi_{\alpha} \quad \text{and} \quad \rho \leq \sigma_2 \otimes \pi_{\beta}.$$

Then, since $\alpha < \beta$, we see that $L_{\beta} \subseteq L_{\alpha}$, and so $L_{\beta} \subseteq \ker \sigma_1 \cap \ker \sigma_2$. Thus, $\sigma_{1|L_{\beta}}$ and $\sigma_{2|L_{\beta}}$ are identities. We also see by the definitions of L_{β} and π_{α} that $\pi_{\alpha|L_{\beta}}$ is the identity. Thus,

(31)
$$\begin{aligned} \rho|_{L_{\beta}} &\leq (\sigma_{1} \otimes \pi_{\alpha})|_{L_{\beta}} = \mathrm{Id} \oplus \cdots \oplus \mathrm{Id} \\ \rho|_{L_{\beta}} &\leq (\sigma_{2} \otimes \pi_{\beta})|_{L_{\beta}} = \pi_{\beta}|_{L_{\beta}} \oplus \ldots \oplus \pi_{\beta}|_{L_{\beta}} \end{aligned}$$

Now it is clear, that for each $\beta < \eta$, we have $G = L_{\beta}M_{\beta}$ and $\pi_{\beta|M_{\beta}} = \text{Id.}$ Therefore, by Lemma 8.6, each $\pi_{\beta|_{L_{\beta}}}$ is an irreducible representation of L_{β} . It follows from the equalities in (31) that $\pi_{\beta|_{L_{\beta}}}$ must be equal to the trivial one-dimensional representation 1. But this implies in turn that $\pi_{\beta} = 1$ which goes against our choice of π_{β} . Therefore,

 $\sigma_1 \otimes \pi_{\alpha} \cap \sigma_2 \otimes \pi_{\beta} = \emptyset$ for every $\alpha < \beta < \eta, \ \sigma_1 \in K_{\alpha}, \ \sigma_2 \in K_{\beta}$, as required for the \mathcal{Z} -property.

Next we check that Condition (30) in Lemma 8.3 holds. Recall that

$$A_{\alpha} = \{ \sigma^{uv} : \sigma \in K_{\alpha}, u, v \in \mathbb{H}_{\sigma} \}.$$

For each $\alpha < \eta$, let then $\varphi \in \langle A_{\alpha} \rangle$ and π_{α}^{k} be any diagonal coefficient of the

representation π_{α} (k may change with α). Then $\varphi = \sum_{n=1}^{p} z_n \sigma_n^{u_n v_n}$, where for n = 1, ..., p, z_n are scalars and $\sigma_n^{u_n v_n}$ are coefficients of the representations $\sigma_n \in K_{\alpha}$ so that $L_{\alpha} \subseteq \ker \sigma_n$.

Since $G = L_{\alpha}M_{\alpha}$, writing each $s \in G$ as s = hm with $h \in L_{\alpha}$ and $m \in M_{\alpha}$, we see that

$$\varphi(s)\pi_{\alpha}^{k}(s) = \varphi(hm)\pi_{\alpha}^{k}(hm) = \varphi(m)\pi_{\alpha}^{k}(h) = \varphi \otimes \pi_{\alpha}^{k}(m,h).$$

So if P is any operator in $VN(M_{\alpha})$ of norm 1 and I is the identity operator in $VN(L_{\alpha})$, then

 $\|\varphi\pi^k_{\alpha}\|_{A(G)} = \|\varphi \otimes \pi^k_{\alpha}\|_{A(L_{\alpha} \times M_{\alpha})} \ge |\langle\varphi \otimes \pi^k_{\alpha}, P \otimes I\rangle| = |\langle\varphi, P\rangle||\langle\pi^k_{\alpha}, I\rangle| = |\langle\varphi, P\rangle|.$ Thus, $\|\varphi \pi_{\alpha}^{k}\| \geq \|\varphi\|$, and so by Lemma 8.3, A(G) has an $\mathbb{F}\ell^{1}(\eta)$ -base of type 2 with constants $K_1 = K_2 = 1$.

Remark 8.8. We can actually prove a bit more in Theorem 8.7: For any $\alpha < \eta$ and $\sigma_1 \neq \sigma_2 \in K_{\alpha}$, the representations $\sigma_1 \otimes \pi_{\alpha}$ and $\sigma_2 \otimes \pi_{\alpha}$ are also disjoint. To prove this, suppose that for some $\alpha < \eta$, and some $\sigma_1, \sigma_2 \in K_{\alpha}$, the representations $\sigma_1 \otimes \pi_{\alpha}$ and $\sigma_2 \otimes \pi_{\alpha}$ contains a common irreducible representation ρ . Then since $\pi_{\alpha|M_{\alpha}} = \mathrm{Id}$, we obtain

$$\rho_{|M_{\alpha}} \leq (\sigma_1 \otimes \pi_{\alpha})_{|M_{\alpha}} \cap (\sigma_2 \otimes \pi_{\alpha})_{|M_{\alpha}} \\ = (\sigma_{1|M_{\alpha}} \oplus \sigma_{1|M_{\alpha}} \oplus \dots \oplus \sigma_{1|M_{\alpha}}) \cap (\sigma_{2|M_{\alpha}} \oplus \sigma_{2|M_{\alpha}} \oplus \dots \oplus \sigma_{2|M_{\alpha}}).$$

Therefore $(\sigma_{1|M_{\alpha}} \oplus \sigma_{1|M_{\alpha}} \oplus ... \oplus \sigma_{1|M_{\alpha}}) \cap (\sigma_{2|M_{\alpha}} \oplus \sigma_{2|M_{\alpha}} \oplus ... \oplus \sigma_{2|M_{\alpha}}) \neq \emptyset$. Now since $G = L_{\alpha}M_{\alpha}$ and $\sigma_{1|L_{\alpha}}$ and $\sigma_{2|L_{\alpha}}$ are identities, Lemma 8.6 implies that $\sigma_{1|M_{\alpha}}$ and $\sigma_{2|M_{\alpha}}$ are irreducible, and so $\sigma_{1|M_{\alpha}}$ and $\sigma_{2|M_{\alpha}}$ must be equal. Again, since $G = L_{\alpha}M_{\alpha}$, we must have $\sigma_1 = \sigma_2$, as required.

8.4. $\mathbf{F}\ell^1(\eta)$ -bases for compact connected groups. The results on infinite products of compact groups seen in the previous subsection can be transferred to compact connected groups G having a certain infinite "dimension". This will be our aim in the present subsection. The arguments are based on some structure theorems, where G is sandwiched between two groups having the form of the groups studied in the previous subsection (see [66, Theorems 9.25 2.26]).

8.4.1. The \mathcal{Z}' -property. We shall develop in this subsection the arguments used in the proof of Theorem 8.7 to include other compact groups having the \mathcal{Z} -property. For this we introduce another property of a family of representations of G called the \mathcal{Z}' -property of level η for some infinite cardinal η , as well as a new definition of a d-rank of a group. The \mathcal{Z}' -property is easier to handle than the \mathcal{Z} -property, while the d-rank of a group is a kind of a dimension of a group based on which the required representations with the \mathcal{Z}' -property are constructed. We shall see then that all compact connected groups with infinite d-rank have the \mathcal{Z}' -property.

Since \mathcal{Z}' -property of level η yields \mathcal{Z} -property of level η , and this type of groups satisfies Condition (30) in Lemma 8.3 (i), their Fourier algebras will contain an $F\ell^1(\eta)$ -base of type 2.

As with the previous cases, this will enable us to obtain in the last subsection 9.1 the strong Arens irregularity not only of A(G) when G belongs to this class of groups, but of $A(G \times H)$ where H is any locally compact amenable group. The centre of $\mathcal{UC}_2(G \times H)^*$ will be seen to be the Fourier-Stieltjes $B(G \times H)$. We shall also deduce some rich algebraic structures in the algebras related to these groups.

Definition 8.9. Let G be a locally compact group with its dual enumerated as $\widehat{G} = \{\sigma_{\alpha} : \alpha < \eta\}$. We say that G has the \mathcal{Z}' -property of level η when \widehat{G} contains a subset $\{\pi_{\alpha} : \alpha < \eta\}$ such that if

$$L_{\alpha} = \bigcap_{\beta < \alpha} (\ker \sigma_{\beta} \cap \ker \pi_{\beta}) \quad for \ every \quad \alpha < \eta.$$

then

$$G = L_{\alpha} \ker \pi_{\alpha}$$
 for every $\alpha < \eta$.

Notation 8.10. For a locally compact group G with a subgroup L, let

- (i) $A_L(G) = \{ \varphi \in A(G) : \varphi(nx) = \varphi(x) \text{ for every } x \in G, n \in L \}.$
- (ii) $R_L: A(G) \to A(L)$ be the restriction map.
- (iii) $Q_L: A(G/L) \to A_L(G)$ be given by $Q_L(u) = u \circ q_L$, when L is normal and $q_L: G \to G/L$ is the quotient homomorphism.

Lemma 8.11. Let G be a compact group such that G = LM, where L and M are normal subgroups of G with L compact and M closed. Then, for every $\varphi \in A_L(G)$, we have

$$||R_M(\varphi)||_{A(M)} = ||\varphi||_{A(G)}.$$

Proof. Consider the following diagram

$$\begin{array}{ccc} A(G/L) & \xrightarrow{Q_L} & A_L(G) \\ & & \downarrow^j & & \downarrow^{R_M} \\ A(M/(L \cap M)) & \xrightarrow{Q_{L \cap M}} & A_{L \cap M}(M). \end{array}$$

Then Q_L and $Q_{L\cap M}$ are isometric isomorphisms (see [75, Theorem 2.4.2], while the map j is the isometric isomorphism induced by the topological isomorphism $LM/L \to M/(L \cap M)$. Since the diagram commutes, R_M has to be an isometric isomorphism as well.

Lemma 8.12. Let G be a compact group with the \mathcal{Z}' -property of level η . Then A(G) contains an $F\ell^1(\eta)$ -base of type 2 with constants $K_1 = K_2 = 1$.

Proof. Let $\widehat{G} = \{\sigma_{\alpha} : \alpha < \eta\}$ with $\sigma_0 = 1$. Let $\{\pi_{\alpha} : \alpha < \eta\}$ be a family of irreducible representations having the \mathcal{Z}' -property for this enumeration of \widehat{G} , and let $\{L_{\alpha} : \alpha < \eta\}$ be, as in Definition 8.9, the family of closed subgroups of G directed downward,

$$L_{\alpha} = \bigcap_{\beta < \alpha} (\ker \sigma_{\beta} \cap \ker \pi_{\beta}) \quad \text{for every} \quad \alpha < \eta.$$

We see first that the family $\{\pi_{\alpha} : \alpha < \eta\}$ has the \mathcal{Z} - property. To begin with, define for each $\alpha < \eta$,

$$K_{\alpha} = \{ \sigma \in \widehat{G} \colon L_{\alpha} \subseteq \ker \sigma \}$$

Since $\sigma_{\alpha} \in K_{\alpha^+}$ for every $\alpha < \eta$, we see that $\widehat{G} = \bigcup_{\alpha < \eta} K_{\alpha}$ so that $\{K_{\alpha}\}_{\alpha < \eta}$ is an upward directed cover of \widehat{G} .

Next, let $\alpha_1 < \alpha_2 < \eta$ and $\sigma_i \in K_{\alpha_i}$, i = 1, 2, be given. Suppose that for some $\rho \in \widehat{G}$,

$$\rho \leq \sigma_1 \otimes \pi_{\alpha_1} \cap \sigma_2 \otimes \pi_{\alpha_2}.$$

Then, since $L_{\alpha_2} \subseteq \ker \sigma_1 \cap \ker \sigma_2$ and $\pi_{\alpha_1}(L_{\alpha_2}) = \operatorname{Id}$ (as $L_{\alpha_2} \subseteq \ker \pi_{\alpha_1}$ by definition), we find

$$\rho|_{L_{\alpha_{2}}} \leq (\sigma_{1} \otimes \pi_{\alpha_{1}})|_{L_{\alpha_{2}}} = \mathrm{Id} \oplus \cdots \oplus \mathrm{Id}$$
$$\rho|_{L_{\alpha_{2}}} \leq (\sigma_{2} \otimes \pi_{\alpha_{2}})|_{L_{\alpha_{2}}} = \pi_{\alpha_{2}}|_{L_{\alpha_{2}}} \oplus \ldots \oplus \pi_{\alpha_{2}}|_{L_{\alpha_{2}}}.$$

Now as before, Lemma 8.6 implies that $\pi_{\alpha_2}|_{_{L_{\alpha_2}}}$ is an irreducible representation of L_{α_2} since $G = L_{\alpha_2} \ker \pi_{\alpha_2}$. We deduce from the equalities above that $\pi_{\alpha_2}|_{_{L_{\alpha_2}}} = 1$, the trivial one-dimensional representation. But this implies that $\pi_{\alpha_2} = 1$ which goes against our construction. We thus deduce that Definition 8.1 holds, i.e., G has the \mathcal{Z} -property of level η .

Next we check that condition (30) in Lemma 8.3 is valid so that A(G) has an $F\ell^1(\eta)$ -base of type 2. Recall that for each $\alpha < \eta$,

$$A_{\alpha} = \{ \sigma^{uv} : \sigma \in K_{\alpha}, u, v \in \mathbb{H}_{\sigma} \},\$$

and as in the proof of Theorem 8.7, let $\varphi \in \langle A_{\alpha} \rangle$ and π_{α}^{k} be any diagonal coefficient of the representation π_{α} where k may change with α . Since φ is a linear combination of coefficients σ^{uv} where $\sigma \in K_{\alpha}$ (so that $L_{\alpha} \subseteq \ker \sigma$), we see that $\varphi \in A_{L_{\alpha}}(G)$. Now if M_{α} denotes $\ker(\pi_{\alpha})$, then $G = L_{\alpha}M_{\alpha}$ by the \mathcal{Z}' -property. We may apply therefore Lemma 8.11 to see that $\|\varphi\|_{A(G)} = \|R_{M_{\alpha}}(\varphi)\|_{A(M_{\alpha})}$. Choose then for every $\epsilon > 0$, $S_{\epsilon} \in VN(M_{\alpha})$ with $\|S_{\epsilon}\| \leq 1$ and

$$\|\varphi\|_{A(G)} = \|R_{M_{\alpha}}(\varphi)\|_{A(M_{\alpha})} \le |\langle S_{\epsilon}, R_{M_{\alpha}}(\varphi)\rangle| + \epsilon.$$

Since $M_{\alpha} = \ker \pi_{\alpha}$, we have $R_{M_{\alpha}}(\varphi \pi_{\alpha}^{k}) = R_{M_{\alpha}}(\varphi)$, whence

$$\|\varphi\|_{A(G)} \le |\langle S_{\epsilon}, R_{M_{\alpha}}(\varphi \pi_{\alpha}^{k})\rangle| + \epsilon = |\langle (R_{M_{\alpha}})^{*}(S_{\epsilon}), \varphi \pi_{\alpha}^{k}\rangle| + \epsilon.$$

Since $||R_{M_{\alpha}}^{*}(S_{\epsilon})|| \leq ||S_{\epsilon}|| \leq 1$ (see for instance [75, Theorem 2.4.1]), we conclude as required that $||\varphi||_{A(G)} \leq ||\varphi\pi_{\alpha}^{k}||_{A(G)}$. Therefore, A(G) has an $F\ell^{1}(\eta)$ -base of type 2 with constants $K_{1} = K_{2} = 1$.

Remarks 8.13.

- (i) Again, as in Remark 8.8, we can prove a bit more when G is as in Lemma 8.12. In fact, for every $\alpha < \eta$ and $\sigma_1, \sigma_2 \in K_{\alpha}$,
 - $\sigma_1 \otimes \pi_\alpha \cap \sigma_2 \otimes \pi_\alpha \neq \emptyset$ if and only if σ_1 and σ_2 are equivalent.

To prove this, suppose that

 $\rho \leq \sigma_1 \otimes \pi_\alpha \cap \sigma_2 \otimes \pi_\alpha$ for some $\rho \in \widehat{G}$. Then,

$$\rho\big|_{\ker \pi_{\alpha}} \leq \left(\sigma_{1} \otimes \pi_{\alpha}\right)\big|_{\ker \pi_{\alpha}} = \sigma_{1}\big|_{\ker \pi_{\alpha}} \oplus \ldots \oplus \sigma_{1}\big|_{\ker \pi_{\alpha}},$$
$$\rho\big|_{\ker \pi_{\alpha}} \leq \left(\sigma_{2} \otimes \pi_{\alpha}\right)\big|_{\ker \pi_{\alpha}} = \sigma_{2}\big|_{\ker \pi_{\alpha}} \oplus \ldots \oplus \sigma_{2}\big|_{\ker \pi_{\alpha}}.$$

Now, for i = 1, 2, $\sigma_i(L_\alpha) = \text{Id}$ and $G = L_\alpha \ker \pi_\alpha$. By Lemma 8.6, it follows that $\sigma_i|_{\ker \pi_\alpha}$ is an irreducible representation of $\ker \pi_\alpha$. The above equalities then imply that $\sigma_1|_{\ker \pi_\alpha}$ and $\sigma_2|_{\ker \pi_\alpha}$ are equivalent. Using again that $G = L_\alpha \ker \pi_\alpha$, we conclude that σ_1 and σ_2 are equivalent.

(ii) If $(G_{\alpha})_{\alpha < \eta}$ is an infinite family of non-trivial compact groups, then it is not difficult to verify that $G = \prod_{\alpha < \eta} G_{\alpha}$ has the \mathcal{Z}' -property of level η , and so A(G) contains an $\mathbb{F}\ell^{1}(\eta)$ -base of type 2 by Lemma 8.12 (as already

shown in the previous subsection). (η) -base of type 2 by Lemma 8.12 (as an shown in the previous subsection).

8.4.2. **d-rank of a compact group.** We proceed now to prove that the \mathcal{Z}' property is available for compact connected groups with an infinite dimension in
the sense of Definition 8.14 below. Since our task concerns elements picked from
the dual of G, we shall call our dimension of G the dual rank of G. When the
dual rank is infinite, we shall see that it coincides with the local weight η of the
group. So by Lemma 8.3, for these groups there are $F\ell^1(\eta)$ -bases of type 2 in A(G).
With these bases, as before, we deduce the strong Arens irregularity as well as the
algebraic structure in the algebras. Recall that when the group is abelian, A(G) is
strongly Arens irregular independently the dual rank of the group since we know

from Section 6 that $L^1(\widehat{G})$ is strongly Arens irregular for any locally compact group G.

Before we give the precise definition of a dual rank of a compact group, we recall the following facts.

The torsion-free rank $r_0(H)$ of an abelian discrete group H is the cardinality of a maximal independent subset of elements in H whose order is infinite. So if t(H) denotes the torsion subgroup of H, then $r_0(H)$ is the maximal number of independent elements in H/t(H) and $r_0(H) = r_0(H/t(H))$, see [63, A13]. Note that if the quotient H/t(H) is countable, then $r_0(H)$ may be either finite or countable. If the quotient H/t(H) is uncountable, then $r_0(H)$ is uncountable and in fact $r_0(H) =$ |H/t(H)|.

There are several notions of dimension of a topological space. The main ones coincide in some instances such as separable metrizable spaces or compact abelian groups, see the discussion at [66, 8.25] or [25, Chapter 7]. We take that given in [63, Definition 3.11]. With this definition, the dimension dim G of a compact abelian group G is the same as the torsion-free rank of its dual group \hat{G} , see [63, Theorem 24.28]. So we may define the dual rank of a compact abelian group G as

$$d - \operatorname{rank}(G) = \dim G = r_0(G).$$

When G is a connected compact abelian group, its dual group \widehat{G} is torsion free, see for example [63, Theorem 24.25], and so \widehat{G} contains a subgroup isomorphic to $\bigoplus_{\alpha < r_0(\widehat{G})} \mathbb{Z}$ and it is contained in its divisible hull $\bigoplus_{\alpha < r_0(\widehat{G})} \mathbb{Q}$ (the divisible hull being also torsion free, has the stated form, see for example the proof of [63, Theorem A. 14]). Hence, the following observation follows.

(32)
$$G \text{ is countable (i.e., } G \text{ is metrizable and } w(G) = w) \text{ if and only if} \\ r_0(\widehat{G}) \le w. \text{ Otherwise, } r_0(\widehat{G}) = |\widehat{G}| = w(G).$$

Hence for any connected compact abelian group, we have the three following cases

(33)
$$d - \operatorname{rank}(G) = r_0(\widehat{G}) \quad \text{when } w(G) = w \text{ and } r_0(\widehat{G}) \quad \text{is finite,} \\ d - \operatorname{rank}(G) = r_0(\widehat{G}) = |\widehat{G}| = w(G) \quad \text{when } r_0(\widehat{G}) = w = w(G), \\ d - \operatorname{rank}(G) = r_0(\widehat{G}) = |\widehat{G}| = w(G), \text{ when } w(G) > w,$$

where w(G) is the local weight of G.

We need also to recall the commutator G' of a topological group G. This is the subgroup of G generated by the commutators $sts^{-1}t^{-1}$ of all elements $s, t \in G$. The subgroup G' is normal but it is not in general closed in G. The quotient G/N of G by any normal subgroup N is abelian if and only if $G' \subseteq N$; in particular G/G' is abelian. A compact connected group is said to be semisimple when G = G'.

When G is a compact connected group, the commutator G' of G is closed and connected, [66, Theorem 9.2]. There is actually a family of simple simply connected compact Lie groups $\{S_j : j \in J\}$ such that G' is a quotient of the product $\prod_{j \in J} S_j$, [66, Theorem 9.19]. The cardinal card J is then an isomorphy invariant of G, denoted as $\aleph(G)$, see [66, Page 496 or 469].

Now every compact connected group G is a semidirect product of G' and the compact connected abelian group G/G' (see [66, Theorem 9.39]), and so G is homeomorphic to $G' \times G/G'$. So we may define the dual rank of G of a compact connected group as

$$d-\operatorname{rank}(G) = \aleph(G) + r_0(\overline{G}/\overline{G'}).$$

Finally, recall from with [63, Theorem 3.12] that the dimension dim G of a totally disconnected space is 0. Accordingly, since every compact group G is homeomorphic to $G_0 \times G/G_0$ ([66, Theorem 10.38]), where G_0 is the connected component of G,

we may define the dual rank of a compact group G as the dual rank of its connected component. Thus, we have

Definition 8.14. The dual rank of a compact group G is given by

 $d-\operatorname{rank}(G) = d-\operatorname{rank}(G_0) = \aleph(G_0) + r_0(\widehat{G_0}/\widehat{G_0}),$

where G_0 denotes the connected component of G and $\aleph(G_0)$ is the isomorphy invariant as defined above.

Remarks 8.15. We remark here some immediate consequences of the definition of the dual rank of a compact group G.

- (i) When G is abelian, G' is trivial, and so $d \operatorname{rank}(G) = r_0(\widehat{G_0})$.
- (ii) When G is connected, it follows from [66, Theorem 9.55 and Theorem 9.52] that $d-\operatorname{rank}(G)$ is finite if and only if dim G is finite. Indeed, these two theorems yield the following equivalent statements
 - (a) $d-\operatorname{rank}(G) = \aleph(G) + r_0(G/G') < \infty$.
 - (b) G' s a compact Lie group and G/G' is a finite dimensional compact abelian group.
 - (c) dim $G < \infty$.
- (iii) When G is connected the quotient group G/G' is a compact connected abelian group. Therefore, according to whether $r_0(\widehat{G/G'})$ is finite and so w(G/G') = w, or $r_0(\widehat{G/G'})$ is infinite, our previous observations in (33) imply, respectively, that

$$d-\operatorname{rank}(G) = \aleph(G) + r_0(\widehat{G}/\widehat{G'}) \quad \text{or} \\ d-\operatorname{rank}(G) = \aleph(G) + w(G/G').$$

For the following theorem, see [66, 9.24–9.26].

Theorem 8.16. Let G be a connected compact group, $Z_0(G)$ be the identity component of the centre Z(G) of G, and let $\Delta = Z_0(G) \cap G'$.

- (i) The subgroup Δ is totally disconnected and central, and $G = Z_0(G)G'$.
- (ii) There exist three continuous surjective morphisms

$$\pi_{G}: \prod_{\gamma < \aleph(G)} S_{\gamma} \to G',$$

$$\mu_{G}: Z_{0}(G) \times \prod_{\gamma < \aleph(G)} S_{\gamma} \to G \quad and$$

$$\nu_{G}: G \to \frac{Z_{0}(G)}{\Delta} \times \prod_{\gamma < \aleph(G)} \frac{S_{\gamma}}{Z_{0}(S_{\gamma})},$$

where each S_{γ} , $\gamma < \aleph(G)$, is a simple, simply connected compact Lie group such that

- (a) $\mu_G(z, \mathbf{s}) = z\pi_G(\mathbf{s})$, for each $z \in Z_0(G)$ and $\mathbf{s} \in \prod_{\gamma < \aleph(G)} S_{\gamma}$,
- (b) $\nu_G(g) = (z\Delta, (s_{\gamma}Z(S_{\gamma}))_{\gamma < \aleph(G)})$ for every $g \in G$, where $g = z\pi_G(\mathbf{s})$, $\mathbf{s} = (s_{\gamma})_{\gamma} \in$, is the decomposition of g seen as an element of $Z_0(G)G'$,
- (c) μ_G and ν_G have totally disconnected central kernels, and
- (d) $\nu_G \circ \mu_G = q$, where

$$q: Z_0(G) \times \prod_{\gamma < \aleph(G)} S_\gamma \to \frac{Z_0(G)}{\Delta} \times \prod_{\gamma < \aleph(G)} \frac{S_\gamma}{Z_0(S_\gamma)}$$

is given by $q(z, (s_{\gamma})_{\gamma < \aleph(G)}) = (z\Delta, (s_{\gamma}Z_0(S_{\gamma}))_{\gamma < \aleph(G)}).$

Lemma 8.17. Let G be an infinite compact group.

- (i) G has a finite dual rank if and only if its connected component G_0 has a finite dimension.
- (ii) If G has an infinite dual rank, then $d-\operatorname{rank}(G) = w(G_0)$.

Proof. Since $d-\operatorname{rank}(G) = d-\operatorname{rank}(G_0)$ by definition, Statement (i) is just a repetition of Remark 8.15(ii).

We prove the second assertion first for a connected compact group G; we shall be repeatedly looking at the equalities

(34)
$$w(G) = w(G') + w(G/G')$$
$$d-\operatorname{rank}(G) = \aleph(G) + r_0(\widehat{G/G'}).$$

We note first that, the existence of the surjective homomorphisms π_G and $\nu_{G'}$ in Lemma 8.16 shows that

(35)
$$\aleph(G) \le w(G') \le \max\{\aleph(G), \omega\}.$$

It follows that if $\aleph(G)$ is finite, then

(36)
$$\aleph(G) \le w(G') \le \omega$$
; otherwise, $\aleph(G) = w(G') \ge \omega$.

Assume now that G has infinite d-rank. Then $\aleph(G)$ and $r_0(\widehat{G/G'})$ cannot be both finite, and so to prove that d-rank(G) = w(G), we look at two separate cases. Case 1: If $r_0(\widehat{G/G'})$ is infinite, then by Remark 8.15 (iii),

$$d-\operatorname{rank}(G) = \aleph(G) + w(G/G').$$

If $\aleph(G) < w$, then by (36), $w(G') \leq \omega$, and so it is clear that

$$d-\operatorname{rank}(G) = w(G/G') + w(G') = w(G).$$

If $\aleph(G) \ge w$, then $\aleph(G) = w(G')$, and so

$$d-\operatorname{rank}(G) = w(G/G') + w(G') = w(G).$$

Case 2: If $r_0(G/G')$ is finite, then by the observation (32), $w(G/G') = \omega$. Since $\aleph(G)$ must be infinite, we have by (36), $\aleph(G) = w(G')$ and a further application of Remark 8.15 (iii) shows that

$$d-\operatorname{rank}(G) = \aleph(G) + r_0(\overline{G/G'}) = \aleph(G) = w(G') + w(G/G') = w(G),$$

as wanted.

Since $d-\operatorname{rank}(G) = d-\operatorname{rank}(G_0)$, the lemma is proved when G is an infinite compact group.

Lemma 8.18. Let G be a connected compact group and let L be a proper, closed, normal subgroup of G that is not totally disconnected. Then L has a normal supplement M, i.e., a proper closed normal subgroup M of G with G = LM.

Proof. We apply the structure theorem, Theorem 8.16.

Let L_0 be the connected component of L, and observe first that $\nu_G(L_0) \neq \{1\}$ since ker ν_G is totally disconnected. Let

$$p: \frac{Z_0(G)}{\Delta} \times \prod_{j \in J} \frac{S_j}{Z(S_j)} \to \prod_{j \in J} \frac{S_j}{Z(S_j)},$$
$$q: \frac{Z_0(G)}{\Delta} \times \prod_{j \in J} \frac{S_j}{Z(S_j)} \to \frac{Z_0(G)}{\Delta} \quad \text{and}$$
$$p_j: \frac{Z_0(G)}{\Delta} \times \prod_{j \in J} \frac{S_j}{Z(S_j)} \to \frac{S_j}{Z(S_j)} \quad \text{for } j \in J,$$

denote the natural projections. We consider two cases. (Remember $\aleph(G) = |J|$.)

Case I. $q(\nu_G(L_0))$ is trivial. Then $p(\nu_G(L_0))$ is not trivial. Note first that L_0 is also a normal subgroup of G (being the connected component of L). So $p(\nu_G(L_0))$ is a non-trivial normal subgroup of $\prod_{j \in J} \frac{S_j}{Z(S_j)}$. Since the factors $\frac{S_j}{Z(S_j)}$ are simple Lie groups and simple Lie groups contain no nontrivial connected normal subgroups, we conclude that there is some $j_0 \in J$ such that $p_{j_0}(\nu_G(L_0)) = \frac{S_{j_0}}{Z(S_{j_0})}$. Consider now the kernel of p_{j_0} . If 1_{j_0} is the identity in $\frac{S_{j_0}}{Z(S_{j_0})}$, then

$$\ker(p_{j_0}) = \frac{Z_0(G)}{\Delta} \times \prod_{j \in J, j \neq j_0} \frac{S_j}{Z(S_j)} \times \{1_{j_0}\}.$$

Then it is clear that ker (p_{j_0}) is a normal supplement of $\nu_G(L_0)$ in the group $\frac{Z_0(G)}{\Lambda} \times$ $\prod_{j \in J} \frac{S_j}{Z(S_j)}$. So for every $s \in G$ there exist $u \in \ker(p_{j_0})$ and $t \in L_0$ such that $\nu_G(s) = \nu_G(t)u$. Thus $u = \nu_G(t^{-1}s)$ and

$$s = t(t^{-1}s) \in L_0(\nu_G)^{-1}(\ker(p_{j_0})).$$

It follows that

$$M := (\nu_G)^{-1} \big(\ker(p_{j_0}) \big) = \ker \big(p_{j_0} \circ \nu_G \big)$$

is a normal supplement of L_0 , and so of L, in G.

Case II. $q(\nu_G(L_0))$ is nontrivial. Now $q(\nu_G(L_0))$ is a nontrivial subgroup of the compact abelian group $\frac{Z_0(G)}{\Delta}$. So let χ be any character of $\frac{Z_0(G)}{\Delta}$ that is not trivial on $q(\nu_G(L_0))$. We claim that $\frac{Z_0(G)}{\Delta} = q(\nu_G(L_0)) \ker \chi$, i.e., $\ker \chi$ is a normal supplement of $q(\nu_G(L_0))$ in $\frac{Z_0(G)}{\Delta}$. Since the circle group \mathbb{T} has no nontrivial connected subgroups and $q(\nu_G(L_0))$ is connected, we must have

$$\chi\left(q\left(\nu_G(L_0)\right)\right) = \chi\left(\frac{Z_0(G)}{\Delta}\right) = \mathbb{T}.$$

We can therefore find for each element $x \in \frac{Z_0(G)}{\Delta}$, an element $s \in L_0$ with $\chi(q(\nu_G(s))) = \chi(x)$. In other words,

$$x = q(\nu_G(s))((q(\nu_G(s)))^{-1}x)$$
 with $(q(\nu_G(s)))^{-1}x \in \ker \chi_s$

which is what is required.

Once we have

$$\frac{Z_0(G)}{\Delta} = q(\nu_G(L_0)) \ker \chi$$

and observe that

$$(q(\nu_G(L_0))\ker\chi) \times \prod_{j\in J} \frac{S_j}{Z(S_j)} \subseteq (\nu_G(L_0)(\ker\chi \times \prod_{j\in J} \frac{S_j}{Z(S_j)}),$$

it is easy to see that

$$(\ker \chi) \times \prod_{j \in J} \frac{S_j}{Z(S_j)}$$

is a normal supplement of $\nu_G(L_0)$ in $\frac{Z_0(G)}{\Delta} \times \prod_{j \in J} \frac{S_j}{Z(S_j)}$. Accordingly, as argued in the previous case,

$$M = (\nu_G)^{-1} \left((\ker \chi) \times \prod_{j \in J} \frac{S_j}{Z(S_j)} \right)$$

is a normal supplement of L_0 , and so of L in G, as wanted.

Proof. If $r_0(\widehat{G})$ is infinite, we can find an infinite set $\{\chi_n : n \in \mathbb{N}\} \subset \widehat{G}$ consisting of independent characters with $\chi_n^k \neq 1$ for any $n \in \mathbb{N}$ and $k \in \mathbb{Z} \setminus \{0\}$. Since $\widehat{G/H}$ is isomorphic to $H^{\perp} = \{\chi \in \widehat{G} : \chi(H) = \{1\}\}$ and $r_0(\widehat{G/H})$ is finite, we see that $\langle \chi_n \rangle \cap H^{\perp} = \{1\}$ for all but finitely many $n \in \mathbb{N}$. Pick $n_0 \in \mathbb{N}$ with $\langle \chi_{n_0} \rangle \cap H^{\perp} = \{1\}$. Then $\chi_{n_0}^k \notin H^{\perp}$ for any $k \in \mathbb{Z} \setminus \{0\}$, hence $\chi_{n_0}|_{H^k} \neq 1$ for every $k \in \mathbb{Z} \setminus \{0\}$. Having a character of infinite order, H cannot be totally disconnected (see [63, Theorem 24.26 or Corollary 24.18]). We conclude that $r_0(\widehat{G})$ must be finite. \Box

Lemma 8.20. Let G be a compact connected group and let H be a totally disconnected normal subgroup of G. Then

- (i) w(G/H) = w(G).
- (ii) If G/H has a finite dual rank, then so has G.

Proof. The first assertion follows from [66, Proposition 12.26].

We divide the proof of the second assertion into two cases:

Case I: G is semisimple, i.e., G' = G. In this case, the dual rank of G is simply $d-\operatorname{rank}(G) = \aleph(G)$. Let $\widetilde{G} = \prod_{j \in J} S_j$ be the simply connected semisimple compact group whose quotient is G by the map $\pi_G : \widetilde{G} \to G' = G$. Then G/H is also a quotient of $\prod_{j \in J} S_j$ by the map $q \circ \pi_G$ where $q : G \to G/H$ is the quotient map. Since H is totally disconnected, we see that $\ker(q \circ \pi_G)$ is totally disconnected. So, by the uniqueness theorem proved in [66, Theorem 9.19] (i), we see the family of simple Lie groups $\{S_j : j \in J\}$ defining $\aleph(G)$ is the same as the one defining $\aleph(G/H)$. Therefore $\aleph(G) = \aleph(G/H)$, and so the claim is clear.

Case II: G is an arbitrary compact connected group. In this case

$$d-\operatorname{rank}(G) = \aleph(G) + r_0(\widehat{G}/\widehat{G'})$$

Then G = Z(G)G', since $G = Z_0(G)G'$ by [66, Theorem 9.24]. Since every totally disconnected normal subgroup of a connected group is central (see for instance [66, Proposition A4.27 or Lemma 6.13]), we see that H is central, i.e., $H \subset Z(G)$. Therefore we can write G/H as

$$G/H = (Z(G)G')/H = (Z(G)/H)(G'H/H).$$

It follows that each of the groups (Z(G)/H) and (G'H/H) has a finite dual rank, since the group G/H has a finite dual rank by assumption. On one hand, using Lemma 8.19, we see that Z(G) has a finite dual rank, i.e., $r_0(\widehat{Z(G)})$ is finite. And because G/G' is a quotient of Z(G) since it is topologically isomorphic to $Z(G)/(G' \cap Z(G))$ by the second isomorphism theorem, we deduce that $r_0(\widehat{G/G'}) \leq r_0(\widehat{Z(G)})$, and so $r_0(\widehat{G/G'})$ is finite.

On the other hand, as G'H/H is topologically isomorphic to $G'/(H \cap G')$, it follows that $G'/(H \cap G')$ has a finite dual rank, and so Case I implies that G' has a finite dual rank, i.e., $\aleph(G)$ is finite. Accordingly, $d-\operatorname{rank}(G) = \aleph(G) + r_0(\widehat{G/G'})$ is finite, as wanted.

Lemma 8.21. Every compact connected group with an infinite dual rank η has the \mathcal{Z}' -property of level η .

Proof. By Lemma 8.17. $d - \operatorname{rank}(G) = w(G)$.

Let $\widehat{G} = \{\sigma_{\alpha} : \alpha < w(G)\}$ be any enumeration of \widehat{G} with $\sigma_0 = 1$, the trivial representation. We construct our sequence $\{\pi_{\alpha} : \alpha < w(G)\}$ inductively. Choose π_0 to be any nontrivial unitary representation of G and let $L_1 = \ker \pi_0$.

Since the quotient group $G/\ker \pi_0$ (i.e., the image im (π_0) of π_0 by the first isomorphism theorem) is a subgroup of a unitary group, and unitary groups are,

topologically, subspaces of \mathbb{C}^n , $n \in \mathbb{N}$, Lemma 8.17 shows that $G/ \ker \pi_0$ must have a finite dual rank.

The second assertion of Lemma 8.20 implies then that ker π_0 cannot be totally disconnected, otherwise G will have a finite dual rank. Lemma 8.18 gives then a proper normal subgroup M_1 of G such that $G = \ker \pi_0 M_1$. We then choose $\pi_1 \in \widehat{G}$ such that $M_1 \subset \ker \pi_1$. Thus, $G = L_1 \ker \pi_1$.

Assume now that, for some $\beta < w(G)$, $\{\pi_{\alpha} : \alpha < \beta\}$ have already been chosen with the \mathcal{Z}' -property and define $L_{\beta} = \bigcap_{\alpha < \beta} (\ker \sigma_{\alpha} \cap \ker \pi_{\alpha}).$

If we denote $(\ker \sigma_{\alpha} \cap \ker \pi_{\alpha})$ by H_{α} , i.e., $L_{\beta} = \bigcap_{\alpha < \beta} H_{\alpha}$, then the natural map

$$gL_{\beta} \mapsto (gH_{\alpha})_{\alpha < \beta}$$

shows there is a copy of G/L_{β} in $\prod_{\alpha < \beta} G/H_{\alpha}$, and so, by the first isomorphism theorem, we may see the quotient group G/L_{β} isomorphic to a subgroup of

$$\prod_{\alpha < \beta} \left(G/(\ker \sigma_{\alpha}) \times G/(\ker \pi_{\alpha}) \right) = \prod_{\alpha < \beta} \left(\operatorname{im} \left(\sigma_{\alpha} \right) \times \operatorname{im} \left(\pi_{\alpha} \right) \right).$$

If $\beta < \omega$, then as above Lemma 8.17 shows that G/L_{β} has finite dual rank, and so the second assertion of Lemma 8.20 shows that L_{β} cannot be totally disconnected. If $\beta \ge \omega$, we have $w(G/L_{\beta}) \le \beta < w(G)$ since each factor im $(\sigma_{\alpha}) \times \operatorname{im}(\pi_{\alpha})$ under the product space is a subspace of some \mathbb{C}^n . Then, the first assertion of Lemma 8.20 shows again that L_{β} cannot be totally disconnected.

So in both cases, applying again Lemma 8.18, we find a proper closed normal subgroup M_{β} of G with $G = L_{\beta}M_{\beta}$. The representation π_{β} is then chosen such that $M_{\beta} \subset \ker \pi_{\beta}$. Obviously, $G = L_{\beta} \ker \pi_{\beta}$ and the inductive process is concluded. \Box

Theorem 8.22. Let G be a compact connected group with an infinite dual rank and a local weight η . Then A(G) has an $F\ell^1(\eta)$ -base of type 2 with constants $K_1 = K_2 = 1$.

Proof. This follows directly from Lemmas 8.12 and 8.21.

Having $F\ell^1(\eta)$ -bases in A(G) for the class of compact groups studied in this section, we may apply as previously Sections 5 and 3 to obtain the centres of the Banach algebras $A(G)^{**}$, $\mathcal{UC}_2(G)^*$ and $C^*(G)^{\perp}$ and their algebraic structure. We shall however postpone the statements of these results for next section. The factorization of $VN(G \times H)$, where H is any amenable locally compact group, proved in Thereforem 9.1, enables us to extend the results and state them for a larger class of groups.

9. Factorizations of $VN(G \times H)$ and applications

Throughout the section, we assume that G is as earlier, i.e., a compact group whose local weight η has uncountable cofinality, an infinite product $\prod_{\alpha < \eta} G_{\alpha}$ of non-trivial compact groups, or a compact connected group with infinite d-rank η , and let H be an amenable locally compact group. The Fourier algebra $A(G \times H)$ may not have an $\mathbb{F}\ell^1(\eta)$ -base, but we shall prove factorization theorems for $VN(G \times H)$ and $\mathcal{UC}_2(G \times H)$ using the factorization theorems available for VN(G) and $\mathcal{UC}_2(G)$. This yields information on the algebraic structure of $A(G \times H)^{**}$ and $\mathcal{UC}_2(G \times H)^*$ as well as the strong Arens irregularity of $A(G \times H)$ and the centre of $\mathcal{UC}_2(G \times H)^*$.

If $a \in B(G)$, we define $u \in B(G \times H)$ by u(s,t) = a(s); that is, $u = a \otimes 1_H$, where 1_H is the function with the constant value 1 on H. Also if $T \in VN(G)$, we consider $T \otimes I \in VN(G \times H)$ where I is the identity operator on $L^2(H)$. So

$$T \otimes I(f \otimes h) = (Tf) \otimes h$$
 for $f \in L^2(G)$ and $h \in L^2(H)$.

It is then very quick to verify the following identities, for any locally compact groups G and H,

(37)
$$(u \otimes v) \cdot (S \otimes P) = (u \cdot S) \otimes (v \cdot P)$$

for every $u \in B(G), v \in B(H), S \in VN(G)$ and $P \in VN(H)$. So

(38)
$$(\Psi \otimes v) \cdot (S \otimes P) = (\Psi \cdot S) \otimes (v \cdot P)$$

for every $\Psi \in \mathfrak{UC}_2(G)^*$, $v \in B(H), S \in VN(G)$ and $P \in VN(H)$.

Let $\mathcal{L} = \bigcup_{\alpha < \eta} A_{\alpha} \{ a_{\alpha}^{k} : k = 1, ..., m \}$ be the $\mathbb{F}\ell^{1}(\eta)$ -base in A(G) obtained in Theorem 8.5, 8.7 or 8.22. So in the first case, \mathcal{L} is of type 3, $m \in \mathbb{N}$ is the dimension of the representations π_{α} picked for the \mathcal{Z} -property and $a_{\alpha}^{k} = \pi_{\alpha}^{1k}, k = 1, ..., m$, is the first row of the matrix coefficients of the representation π_{α} for each $\alpha < \eta$. In the second and third cases, \mathcal{L} is of type 2, m = 1 and $a_{\alpha} = \pi_{\alpha}^{k}$ is any diagonal coefficient of the representation π_{α} for each $\alpha < \eta$ (k may depend on α).

Recall also from Section 4 the sets

$$X_k = \{\Psi_{pk} : p \in \mathcal{C}(\eta)\}$$
 and let $X = \bigcup_{k=1}^m X_k$,

where Ψ_{pk} are the cluster points of $\{a_{\alpha}^k : \alpha < \eta, k = 1, ..., m\}$ which correspond to the cofinal ultrafilters p in η , they are in the closed unit ball of $\mathcal{UC}_2(G)^*$.

For each $p \in \mathcal{C}(\eta)$ and k = 1, ..., m, denote by $\Psi_{pk} \otimes 1_H$ the corresponding weak*-cluster point of $\{a_{\alpha}^k \otimes 1_H : \alpha < \eta\}$. Let, for each $k = 1, ..., m, X_k \otimes 1_H$ be the set formed by these cluster points in the unit ball of $\mathcal{UC}_2(G \otimes H)^*$. For an arbitrary element in $\prod_{k=1}^m X_k \otimes 1_H$, we will not refer to the ultrafilters to which it corresponds, and write it simply as $\Psi \otimes 1_H$.

Theorem 9.1. Let G be a compact group whose local weight η has uncountable cofinality, a compact connected group with infinite d-rank η , or an infinite product $\prod_{\alpha < \eta} G_{\alpha}$ of non-trivial compact groups G_{α} , and let H be an amenable locally compact group. Then

- (i) $VN(G \times H)$ has the uniform η -factorization through $\prod_{k=1}^{m} X_k \otimes 1_H$.
- (ii) $\operatorname{UC}_2(G \times H)$ has the 1-factorization through $\prod_{k=1}^m X_k \otimes \mathbb{1}_H$.

Proof. (i) We prove the 1-factorization of $VN(G \times H)$ through $\prod_{k=1}^{m} X_k \otimes 1_H$ when G has the cofinality of its local weight uncountable. The other two cases follow in the same way with m = 1. The case when $Q \in VN(G \times H)$ has the form $S \otimes P$ with $S \in VN(G)$ and $P \in VN(H)$ is straightforward. Pick by Theorem 4.14, any $\Psi = (\Psi_k)_{k=1}^m \in \prod_{k=1}^m X_k$ and $T = (T_k)_{k=1}^m \in (VN(G))^m$ such that

(39)
$$S = \sum_{k=1}^{m} \Psi_k \cdot T_k = \Psi \cdot T_k$$

Then, using (38), we find

$$(\Psi \otimes 1_H) \cdot (T \otimes P) = \sum_{k=1}^m (\Psi_k \otimes 1_H) \cdot (T_k \otimes P) = \sum_{k=1}^m (\Psi_k \cdot T_k) \otimes P = S \otimes P = Q,$$

as required.

Let now $Q \in VN(G \times H)$ be arbitrary and see it by [24], as a weak*-limit of a bounded net $(Q_F)_{F \in \mathbb{N}^{\leq w}}$, where $\mathbb{N}^{\leq w}$ is the set of all finite subsets of \mathbb{N} and

$$Q_F = \sum_{i \in F} S_i \otimes P_i$$
, with $S_i \in VN(G)$, $P_i \in VN(H)$ for each $i \in F$.

Let $\mathcal{A} = \mathcal{A}(G)$ and \mathcal{A}_{α} be as defined in Lemma 8.3. For each $i \in F$, define S^i on $\mathcal{L} = \bigcup_{\alpha < \eta} \mathcal{A}_{\alpha} \mathbf{a}_{\alpha}$ by

(40)
$$\langle S^i, a\mathbf{a}_{\alpha} \rangle = \langle S_i, a \rangle$$
, for every $a \in A_{\alpha}$ and $\alpha < \eta$,

where $\mathbf{a}_{\alpha} = (a_{\alpha}^k)_{k=1}^m = (\pi_{\alpha}^{1k})_{k=1}^m$. Let for each $F \in \mathbb{N}^{\leq w}$,

$$R_F = \sum_{i \in F} S^i \otimes P_i$$

be defined on $\mathcal{L} \otimes A(H)$. Let $x \in \langle \mathcal{L} \rangle \otimes A(H)$ and write it as $x = (\sum_{n=1}^{p} a_n \mathbf{a}_{\alpha_n}) \otimes b$, where $a_n \in \langle A_{\alpha_n} \rangle$ and $\alpha_1 < \alpha_2 < ... < \alpha_p$. Accordingly,

$$\begin{aligned} |\langle R_F, x \rangle| &= |\langle \sum_{i \in F} \sum_{n=1}^p \langle S^i, a_n \mathbf{a}_{\alpha_n} \rangle \langle P_i, b \rangle| = |\langle \sum_{i \in F} \sum_{n=1}^p \langle S_i, a_n \rangle \langle P_i, b \rangle| \\ &= |\langle \sum_{i \in F} S_i \otimes P_i, \sum_{n=1}^p a_n \otimes b \rangle| \\ \leq \|\sum_{i \in F} S_i \otimes P_i\| \|\sum_{n=1}^p a_n \otimes b\| = \|Q_F\| \|\sum_{n=1}^p a_n\| \|b\| \\ &\leq \|Q_F\| \sum_{n=1}^p \|a_n\| \|b\| \leq K_2 \|Q_F\| \sum_{n=1}^p \|a_n \mathbf{a}_{\alpha_n}\| \|b\| \\ &\leq K_1 K_2 \|Q_F\| \sum_{n=1}^p a_n \mathbf{a}_{\alpha_n}\| \|b\| = K_1 K_2 \|Q_F\| \sum_{n=1}^p a_n \mathbf{a}_{\alpha_n} \otimes b\| \\ &= K_1 K_2 \|Q_F\| \|x\|, \end{aligned}$$

where by Lemma 8.3 since A(G) has the \mathbb{Z} -property of level η , $K_1 = 1$ is the selective constant and $K_2 == \sup_{\alpha} ||(\pi_{\alpha}^{1k})_{k=1}^m|| \leq m$ is the $\mathbb{F}\ell^1(\eta)$ -base constant. (In the two other cases, $K_1 = K_2 = 1$.) In other words, R_F is a bounded operator (by $||Q_F||$) on the subspace $\langle \mathcal{L} \rangle \otimes A(H)$ of $A(G \times H)$. We may therefore extend R_F to an operator in $VN(G \times H)$ with $||R_F|| \leq ||Q_F||$. We denote this extension again by R_F and consider the net $(R_F)_{F \in \mathbb{N}^{\leq w}}$. This is a bounded net since $(Q_F)_{F \in \mathbb{N}^{\leq w}}$ is bounded, and so it weak*-clusters at some $R \in VN(G \times H)$. We claim that

$$(\Psi \otimes 1_H) \cdot R = Q.$$

If $a \in A(G)$ and $b \in A(H)$, Then, $a \in A_{\beta}$ for some β and so for every $\alpha \ge \beta$, we find

$$\langle R_F, (a\mathbf{a}_{\alpha}) \otimes b \rangle = \langle \sum_{i \in F} S^i \otimes P_i, (a\mathbf{a}_{\alpha}) \otimes b \rangle$$

$$= \langle \sum_{i \in F} S^i, (a\mathbf{a}_{\alpha}) \rangle \langle P_i, b \rangle$$

$$= \langle \sum_{i \in F} S_i, a \rangle \langle P_i, b \rangle = \langle \sum_{i \in F} S_i \otimes P_i, a \otimes b \rangle$$

$$= \langle Q_F, a \otimes b \rangle.$$

Therefore,

$$\langle R, (a\mathbf{a}_{\alpha}) \otimes b \rangle = \lim_{F} \langle R_F, (a\mathbf{a}_{\alpha}) \otimes b \rangle = \lim_{F} \langle Q_F, a \otimes b \rangle = \langle Q, a \otimes b \rangle.$$

Accordingly,

$$\langle (\Psi \otimes 1_H) \cdot R, a \otimes b \rangle = \lim_{\gamma} \langle (\mathbf{a}_{h(\gamma)} \otimes 1_H) \cdot R, a \otimes b \rangle$$

=
$$\lim_{\gamma} \langle R, (a \otimes b) (\mathbf{a}_{h(\gamma)} \otimes 1_H) \rangle$$

=
$$\lim_{\gamma} \langle R, (a\mathbf{a}_{h(\gamma)}) \otimes b) \rangle = \langle Q, a \otimes b \rangle$$

as wanted.

To prove that $VN(G \times H)$ has the uniform η -factorization, we may argue in two ways either as in Theorem 4.14 or as in Remark 4.15(ii). We choose the shorter route of the remark. Let $\{Q_{\lambda}\}_{\lambda < \eta}$ be any bounded family of operators in $VN(G \times H)$, and regard each Q_{λ} as a weak^{*}-limit of a net $(Q_{\lambda F})_{F \in \mathbb{N}^{<w}}$ made as earlier of simple operators

$$Q_{\lambda F} = \sum_{i \in F} S_{\lambda i} \times P_{\lambda i}, \quad F \in \mathbb{N}^{< w}$$

Partition η into η many subsets I_{λ} such that each I_{λ} is cofinal in η . Take for each $\lambda < \eta$ and each k = 1, ..., m, an element $\Psi_{k\lambda} \otimes 1_H$ in $X_k \otimes 1_H$ corresponding to some ultrafilter $p \in \mathcal{C}(I_{\lambda})$, i.e., an element which comes as a weak*-cluster point of $\{a_{\alpha}^k \otimes 1_H : \alpha \in I_{\lambda}\}$, where $a_{\alpha}^k = \pi_{\alpha}^{1k}$ are picked as in Theorem 8.5 or $a_{\alpha}^k = \pi_{\alpha}^k$ are picked as in Theorems 8.7 and 8.22 depending on the type of our base. The elements $\Psi_{k\lambda} \otimes 1_H$ are in the unit ball of $\mathcal{UC}_2(G \otimes H)^*$ and belong to $C^*(G \times H)^{\perp}$. Put $\Psi_{\lambda} \otimes 1_H = (\Psi_{k\lambda} \otimes 1_H)_{k=1}^m$. ($\Psi_{k\lambda}$ and Ψ_{pk} are not to be confused, one corresponds to the cofinal ultrafilter p on η while the other corresponds to the cofinal subset I_{λ} of η .) For each $\lambda < \eta$, $F \in \mathbb{N}^w$ and $i \in F$, instead of (40) define $S^{\lambda i}$ on $\mathcal{L} = \bigcup_{\alpha < \eta} A_{\alpha} \mathbf{a}_{\alpha}$ by

(42)
$$\langle S^{\lambda i}, a\mathbf{a}_{\alpha} \rangle = \begin{cases} \langle S_{\lambda i}, a \rangle, & \text{if } \alpha \in I_{\lambda} \\ 0 & \text{if } \alpha \notin I_{\lambda}, \end{cases}$$

and let

$$R_{\lambda F} = \sum_{i \in F} S^{\lambda i} \otimes P_{\lambda i}.$$

Then the argument in (41) shows that $R_{\lambda F}$ is a bounded operator on $\langle \mathcal{L} \rangle \otimes A(H)$ and so may be extended to an operator in $VN(G \times H)$. The bounded net $(R_{\lambda F})_{F \in \mathbb{N}^{\leq w}}$ so obtained, weak*-clusters at some $R_{\lambda} \in VN(G \times H)$. The operator R_{λ} satisfies

$$\langle R_{\lambda}, (aa_{\alpha}) \otimes b \rangle = \begin{cases} \langle Q_{\lambda}, a \otimes b \rangle & \text{if } \alpha \in I_{\lambda} \\ 0 & \text{if } \alpha \notin I_{\lambda}. \end{cases}$$

Now let $R = \sum_{\lambda < \eta} R_{\lambda}$ be first defined on $\mathcal{L} \otimes A(H)$ and continue as in Remark 4.15(ii).

(ii) Since $G \times H$ is amenable, $A(G \times H)$ has a bai. So we may proceed precisely as in Theorem 4.16 for the 1-factorization of $\mathcal{UC}_2(G \times H)$ since now that we have the 1-factorization of $VN(G \times H)$.

Remark 9.2. Due to the availability of the $F\ell^1(\eta)$ -bases, together with the factorization theorems, Remark 4.15 was also necessary for the algebraic structure reached in the previous sections. Next we check that this remark is also valid here.

So let our group have the form $G \times H$ as in Theorem 9.1, and Q be in $VN(G \times H)$ or in $\mathcal{UC}_2(G \times H)$ and I be any cofinal subset of η . Then there exists $R \in VN(G \times H)$ or $\mathcal{UC}_2(G \times H)$, respectively such that

(43)
$$\Psi_p \cdot R = \begin{cases} Q, & \text{if } I \in p \\ 0, & \text{otherwise.} \end{cases}$$

To see this, suppose that Q be in $VN(G \times H)$ and let p be any ultrafilter having I as a member and Ψ_p be any element in $\prod_{k=1}^m X_k \otimes 1$. Let $(\sum_{i \in F} S_i \otimes P_i)_{F \in \mathbb{N}^w}$ be the net in $VN(G \times H)$ with Q as a weak*-limit. As in (42), define for each $F \in \mathbb{N}^w$ and $i \in F$, S^i on \mathcal{L} by

$$\langle S^i, a\mathbf{a}_{\alpha} \rangle = \begin{cases} \langle S_i, a \rangle, & \text{if } \alpha \in I \\ 0 & \text{if } \alpha \notin I, \end{cases}$$

and let

$$R_F = \sum_{i \in F} S^i \otimes P_i.$$

Proceed then as in the proof of Theorem 9.1 to see that $(R_F)_F$ is bounded, and so has a weak*-cluster operator $R \in VN(G \times H)$ having the property required in (43)

If $Q = S \cdot u \in \mathcal{UC}_2(G \times H)$ with $S \in VN(G \times H)$ and $u \in A(G \times H)$ then the factorization of S by Ψ_p and R as in (43) yields the same type of factorization of Q by Ψ_p and $R \cdot u \in \mathfrak{UC}_2(G \times H)$.

9.1. Algebraic structure. As it is with the group algebra in Section 6.3, nontrivial left ideals of finite dimensions exist in $A(G)^{**}$ and $\mathcal{UC}_2(G)^*$ for any locally compact group (not necessarily amenable in this case), they are generated by the topological invariant means (TIMs), by the annihilators of $\mathcal{UC}_2(G)$ when G is not compact, or the mixture of both. Since there are 2^{2^n} TIMs in $A(G)^{**}$ and $\mathcal{UC}_2(G)^*$ when G is non-discrete (see [11, Theorem 3.3] for metrizable G, and [67, Theorem 5.9] or [41, Corollary 5.4] for nonmetrizable G), at least this many distinct left ideals of finite dimension exists in $A(G)^{**}$ and $\mathcal{UC}_2(G)^*$ when G is non-discrete. For more details, see [43]. But as we shall soon see, there are also plenty of distinct left ideals of infinite dimension when the group is as in Theorem 9.1.

The part on $A(G \times H)^{**}$ and $\mathcal{UC}_2(G \times H)^*$ in Theorem 9.3 made up the main results in [44]. However, our proofs here are much more simplified.

Theorem 9.3. Let G be a compact group such that $\eta = w(G)$ has uncountable cofinality and H be an amenable locally compact group. Let S be any of the algebras $A(G \times H)^{**}$, $\mathfrak{UC}_2(G \times H)^*$ or $C^*(G \times H)^{\perp}$. Then

- (i) there are at least 2^{2ⁿ} many right cancellable sets in S.
 (ii) there are at least 2^{2ⁿ} many left ideals in S with dimension at least 2^{2ⁿ}.
- (iii) the dimension of any principal right ideal, and so of any nonzero right ideal, in S is at least $2^{2^{\eta}}$.

Proof. Since X is contained in $C^*(G \times H)^{\perp}$ and since we have the 1-factorization of $VN(G \times H)$ and $\mathcal{UC}_2(G \times H)$ by Theorem 9.1, using the proofs of Theorems 5.1(ii), 5.2 and 5.3(ii) with Remarks 4.13 and 9.2 in mind gives the claimed properties. \Box

Theorem 9.4. Let G be a compact group which is either connected with infinite d-rank η or is an infinite product $\prod_{\alpha < \eta} G_{\alpha}$ of non-trivial compact groups, and H be an amenable locally compact group. Let S be any of the algebras $A(G \times H)^{**}$, $\mathfrak{UC}_2(G \times H)^*$ or $C^*(G \times H)^{\perp}$. Then

- (i) there are at least $2^{2^{\eta}}$ many right cancellable elements in S.
- (ii) there are at least $2^{2^{\eta}}$ many principal left ideals in S with trivial pairwise intersections and with dimension at least $2^{2^{\eta}}$.
- (iii) the dimension of any principal right ideal, and so of any nonzero right ideal, in S is at least $2^{2^{\eta}}$.

Proof. This follows with the same proof as above using the proofs of Theorems 5.1(i), 5.2 and 5.3(i) and Remarks 4.13 and 9.2. \square

9.2. Topological centres. Statements (i) and (ii) in the following theorem were proved in [44, Corollary 4.7] for the case when G is compact non-metrizable with the cofinality of w(G) uncountable. The condition that $\omega(H) \leq w(G)$, when H is not necessarily compact, was missing in [44, Corollary 4.7]. This condition guarantees that $A(G \times H)$ has Mazur property of level w(G), required for Theorems 3.5 and 3.8 to apply. This is now corrected in the following theorem. In Statements (i) and (ii), we also simplify and improve the main result proved by Lau and Losert in [85], which was proved with $\eta = w$, each G_{α} is a non-trivial metrizable compact group and H is a second countable locally compact amenable group.

The rest of the results are new.

Theorem 9.5. Let G be a compact group with local weight $\eta = w(G)$ and H be an amenable locally compact group with $w(H) \leq \eta$. Then in the following cases

- (i) G is non-metrizable such that η has uncountable cofinality,
- (ii) G is an infinite product $\prod_{\alpha < \eta} G_{\alpha}$ of non-trivial compact groups such that $\sup \omega(G_{\alpha}) \leq \eta$,
- (iii) G is connected with infinite dual rank η ,
 - (i) the topological centre of $A(G \times H)^{**}$ is $A(G \times H)$.
- (ii) the topological centre of $UC_2(G \times H)^*$ is $B(G \times H)$.
- (iii) the topological centre of $C^*(G \times H)^{\perp}$ is $\{0\}$.

Proof. Note first that the local weight of G is η in each of the three cases. And as already mentioned the condition $w(H) \leq w(G)$ guarantees that $A(G \times H)$ has Mazur property of level η . Since Theorems 8.5, 8.7 and 8.22 provide $F\ell^1(\eta)$ -bases in A(G) for each of these groups G, we have the η -factorization of $A(G \times H)^*$ by Theorem 9.1. Therefore, Statements (i) and (ii) follow by Theorems 3.5(ii) and 3.8(iii), respectively.

Statement (iii) follows from Theorem 3.8(ii) since $X \subseteq C^*(G \times H)^{\perp}$ and $C^*(G \times H) \cap B(G \times H) = \{0\}$.

Corollary 9.6. Let G be a compact group as in Theorem 9.5. Then $A(G \times H)$ is strongly Arens irregular.

- Final remarks 9.7. (i) For simplicity let H be trivial. Then Theorem 9.5 leaves out the case when $w(G) = \sup \omega(G_{\alpha})$ has countable cofinality and $w(G) > \eta$. In fact, We still do not know about the interesting case when G is compact, non-metrizable with countable cofinality. We also do not know what happens when η is finite and w(G) is countable as it is the case if G = SU(3), which is still undecided. On the other hand, A(G) is strongly Arens irregular when G = SU(2) as claimed in [95].
 - (ii) Note that in all the examples studied previously, for the cardinality of the Fℓ¹-bases to be large, it is essential in each case that either κ(G), |G| or w(G) to be as large. So our approach (as well as most of the articles cited above except [73], [84] and [96]) cannot be applied to give the strong Arens irregularity of L¹(G) and M(G) when G is compact, and that of A(G) and B(G) when G is discrete and amenable. In the case of the group algebra, one needs to start first with the topological centre of LUC(G)* being M(G) (since LUC(G) = CB(G)), then deduce that the topological centre of L[∞](G)* is L¹(G) as it can be seen in the proofs provided in [73] and [83, Theorem 1].

One encounters the same problem with the Fourier algebra of an amenable discrete group. This is proved in [84, Theorem 6.5 (i)]. By analogy to the group algebra, one needs to notice first that $\mathcal{UC}_2(G)^*$ is B(G) (since

 $\mathfrak{UC}_2(G) = C^*(G)$, then use [84, Theorem 6.4] to deduce that the topological centre of $VN(G)^*$ is A(G). The approach of [84, Theorem 6.4] is not straightforward.

Another approach can be found [5, Theorem 2.2], where a weakly sequentially complete Banach algebra, containing a bounded approximate identity and is an ideal in its second dual, as it is the situation here with $L^1(G)$ and A(G), is always strongly Arens irregular. See also our recent paper [28].

The same remark in these situations concerns the theorems proved in Section 5 on the algebraic structure of these algebras. Our approach does not cover $L^1(G)^{**}$, M(G) and $M(G)^{**}$ when G is compact, neither $A(G)^{**}$, B(G), $B_{\rho}(G)$ and $B(G)^{**}$ when G is discrete, and cannot lead to any algebraic structure in these algebras. We hope to return to this situation in the future.

(iii) The other two known groups for which the centres of $A(G)^{**}$ and $\mathcal{UC}_2(G)^*$ are A(G) and B(G), respectively, are any locally compact, second countable group G having the closure of its commutator subgroup $\overline{[G,G]}$ not open, and the special unitary group SU(2), see [84] and [95]. We have not been able to construct $F\ell_1$ -bases in A(G) with these two groups, and we believe a further refinement of our approach is necessary to deal with these two groups. We hope to return to this situation in the future.

ACKNOWLEDGEMENT. This paper was written during several visits of the first author to the University of Jaume I in Castellón in his off-teaching periods. The work was partially supported several times by the University of Jaume I. This support is gratefully acknowledged. The work would never have been completed (or even started) without such a support.

He would also like to thank Jorge Galindo for his hospitality and all the folks at the department of mathematics in Castellón.

References

- 1. R. Arens, The adjoint of a bilinear operation, Proc. Amer. Math. Soc. 2 (1951), 839-848.
- 2. R. Arens, Operations induced in function classes, Monatsh. Math. 55 (1951), 1–19.
- G. Arsac, Sur l'espace de Banach engendré par les coefficients d'une représentation unitaire, Publ. Dép. Math. (Lyon) 13 (1976), no. 2, 1–101.
- J.W. Baker and P. Milnes, . The ideal structure of the Stone-Čech compactification of a group, Math. Proc. Cambridge Philos. Soc. 82 (1977), no. 3, 401–409.
- J.W. Baker, A. T. -M. Lau and J. S. Pym, Module homomorphisms and topological centres associated with weakly sequentially complete Banach algebras, J. Funct. Anal. 158 (1998), 186–208.
- J. F. Berglund, H. D. Junghenn and P. Milnes, Analysis on Semigroups: Function Spaces, Compactifications, Representations, Wiley, New York (1989).
- 7. F. F. Bonsall and J. Duncan, Complete normed algebras, Springer-Verlag, Berlin, 1973.
- A. Bouziad and M. Filali, On the size of quotients of function spaces on a topological group, Studia Math. 202 (2011), no. 3, 243–259.
- T. Budak, N. Işık, and J. Pym, Minimal determinants of topological centres for some algebras associated with locally compact groups, Bull. Lond. Math. Soc. 43 (2011), 495–506.
- C. Chou, On the size of the set of left invariant means on a semigroup, Proc. Amer. Math. Soc. 23 (1969), 199–205.
- C. Chou, Topological invariant means on the von Neumann algebra VN(G), Trans. Amer. Math. Soc. 273 (1982), no. 1, 207–229.
- P. Civin and B. Yood, The second conjugate space of a Banach algebra as an algebra, Pacific J. Math. 11 (1961), 847–870.
- W. W. Comfort, A survey of cardinal invariants, General Topology and Appl. 1 (1971), no. 2, 163–199.
- W. W. Comfort, *Topological groups*, In: Handbook of Set-Theoretic Topology, North-Holland, Amsterdam, 1984, pp. 1143–1263.

- 15. H. G. Dales, *Banach algebras and automatic continuity*, The Clarendon Press, Oxford University Press, New York, 2000.
- H. G. Dales and A. T.-M. Lau, The second duals of Beurling algebras, Memoires Amer. Math. Soc.177(836) (2005).
- H. G. Dales, A. T.-M. Lau, D. Strauss, Banach algebras on semigroups and on their compactifications, Mem. Amer. Math. Soc. 205 (2010), vi+165 pp.
- H. G. Dales, A. T.-M. Lau, D. Strauss, Second duals of measure algebras, Dissertationes Math. 481 (2012), 1–121.
- D. E. Davenport and N. Hindman, A proof of van Douwen's right ideal theorem, Proc. Amer. Math. Soc. 113 (1991), no. 2, 573–580
- 20. M. M. Day, Amenable semigroups, Illinois J. Math. 1 (1957), 509-44.
- A. Derighetti, Convolution operators on groups, Lecture Notes of the Unione Matematica Italiana, 11. Springer, Heidelberg; UMI, Bologna, 2011.
- 22. A. Derighetti, M. Filali, M. Monfared, On the ideal structure of some Banach algebras related to convolution operators on $L^p(G)$, J. Funct. Anal.215 (2004), no.2, 341–365.
- J. Duncan and S. A. R. Hosseiniun, The second dual of a Banach algebra, Proc. Roy. Soc. Edinburgh Sect. A 84 (1979), 309–325.
- E. G. Effros and Z-J. Ruan, On approximation properties for operator spaces, Internat. J. Math. 1 (1990), 163–187.
- R. Engelking, *General Topology*, Translated from the Polish by the author, Second edition, Sigma Series in Pure Mathematics 6, Heldermann Verlag, Berlin, 1989.
- M. Eshaghi Gordji and M. Filali, Arens regularity of module actions, Studia Math. 181 (2007), no. 3, 237–254.
- 27. R. Esmailvandi, M. Filali and J. Galindo, Arens regularity of ideals of the group algebra of a compact Abelian group, Proceedings of the Royal Society of Edinburgh, Published online by Cambridge University Press: 27 October 2023.
- R. Esmailvandi, M. Filali and J. Galindo, Special subsets of discrete groups and Arens regularity of ideals in Fourier and group algebras, preprint, 2024.
- P. Eymard, L'algèbre de Fourier d'un groupe localement compact, Bull. Soc. Math. France 92 (1964), 181–236.
- M. Filali, The uniform compactification of a locally compact abelian group, Math. Proc. Cambridge Philos. Soc. 108 (1990), no. 3, 527–538.
- M. Filali, The ideal structure of some Banach algebras, Math. Proc. Cambridge Philos. Soc. 111 (1992), no. 3, 567–576.
- M. Filali, Finite-dimensional left ideals in some algebras associated with a locally compact group. Proc. Amer. Math. Soc. 127 (1999), no. 8, 2325–2333
- M. Filali, Finite dimensional right ideals in algebras associated to a locally compact group, Proc. Amer. Math. Soc. 127 (1999) 1729–1734.
- 34. M. Filali, On a conjecture of Chou and applications, Topology Proc. 24 (1999) 139-147.
- M. Filali, On the ideal structure of some algebras with an Arens product, Recent progress in functional analysis (Valencia, 2000), 289–297, North-Holland Math. Stud., 189, North-Holland, Amsterdam, 2001
- M. Filali, t-Sets and some algebraic properties in βS and ℓ_∞(S)*), Semigroup Forum 65, (2002) 285–300.
- M. Filali and J. Galindo, Extreme Non-Arens Regularity of the Group Algebra, Forum Math. 30, No. 5, 1193–1208 (2018).
- M. Filali and J. Galindo, Algebraic structure of semigroup compactifications: Pym's and Veech's theorems and strongly prime points, J. Math. Anal. Appl. 456 (2017), no. 1, 117–150.
- 39. M. Filali and J. Galindo, ℓ¹ -bases in Banach algebras and Arens irregularities in harmonic analysis, Banach algebras and applications. Proceedings of the international conference held at the University of Oulu, July 3–11, 2017, De Gruyter Proceedings in Mathematics. Berlin: De Gruyter, 37 p. (2020).
- M. Filali and J. Galindo, On the extreme non-Arens regularity of Banach algebras, J. Lond. Math. Soc. (2) 104 (2021), no. 4, 1840–1860.
- M. Filali and J. Galindo, Orthogonal l₁-sets and extreme non-Arens regularity of preduals of von Neumann algebras, J. Math. Anal. Appl. 512 (2022), no. 1, Paper No. 126137, 21 pp.
- 42. M. Filali and J. Galindo, Number of right ideals in the second dual of algebras in harmonic analysis, preprint 2024.
- M. Filali and M. S. Monfared, Finite-dimensional left ideals in the duals of introverted spaces. Proc. Amer. Math. Soc. 139 (2011), no. 10, 3645–3656
- 44. M. Filali, M. Neufang and M. Monfared, Weak factorizations of operators in the group von Neumann algebras of certain amenable groups and applications, Math. Ann. 351 (2011), no. 4, 935–961.

- M. Filali, M. Neufang and M. Monfared, On ideals in the bidual of the Fourier algebra and related algebras, J. Funct. Anal. 258(2010), no. 9, 3117–3133.
- M. Filali and J. S. Pym, Right cancellation in LUC-compactification of a locally compact group, Bull. London. Math. Soc. 35 (2003), 128–134.
- M. Filali and P. Salmi, One-sided ideals and right cancellation in the second dual of the group algebra and similar algebras, J. London Math. Soc. 75 (2007), 35–46.
- J. Funct. Anal. 250 (2007), 144–166.
- <u>Topological centres of the weighted convolution algebras</u>, J. Funct. Anal. 278, No. 11, 22 p. (2020).
- 50. _____, Right cancellation, factorisation and right isometries, preprint 2015.
- 51. M. Filali and A. I. Singh, Recent developments on Arens regularity and ideal structure of the second dual of a group algebra and some related topological algebras, General topological algebras (Tartu, 1999), Math. Stud. (Tartu), vol. 1, Est. Math. Soc., 2001, pp. 95–124.
- M. Filali and T. Vedenjuoksu, Extreme non-Arens regularity of semigroup algebras, Topology Proc. 33 (2009), 185–196.
- 53. F. Ghahramani, Weighted group algebra as an ideal in its second dual space, Proc. Amer. Math. Soc. 90 (1984), no. 1, 71–76.
- F. Ghahramani and J. Laali, Amenability and topological centres of the second duals of Banach algebras, Bull. Austral. Math. Soc. 65 (2002), no. 2, 191–197.
- 55. F. Ghahramani, J. P. McClure and M. Meng, On asymmetry of topological centers of the second duals of Banach algebras, Proc. Amer. Math. Soc. 126 (1998), no. 6, 1765–1768.
- 56. E. E. Granirer, Density theorems for some linear subspaces and some C*-subalgebras of VN(G), Symposia Mathematica, Vol. XXII (Convegno sull'Analisi Armonica e Spazi di Funzioni su Gruppi Localmente Compatti, INDAM, Rome, 1976), pp. 61–70. Academic Press, London, 1977.
- 57. E. E. Granirer, On group representations whose C*-algebra is an ideal in its von Neumann algebra, Ann. Inst. Fourier (Grenoble) 29 (1979), no. 4, v, 37–52.
- 58. E. E. Granirer, Day points for quotients of the Fourier algebra A(G), extreme nonergodicity of their duals and extreme non-Arens regularity, Illinois J. Math. 40 (1996), no. 3, 402–419.
- 59. F. P. Greenleaf, *Invariant means on topological groups*, Van Nostrand Math. Studies, Van Nostrand Reinhold, 1969.
- N. Groenbaek, Amenability of weighted convolution algebras on locally compact groups, Trans. Amer. Math. Soc. 319 (1990), no. 2, 765–775
- M. Grosser, Bidualräume und Vervollständigungen von Banach-moduln, Lecture notes in Mathematics, Vol. 717, Springer-Verlag, Berlin, 1979.
- M. Grosser and V. Losert, The norm strict bidual of a Banach algebra and the dual of C_u(G), Manuscripta Math. 45 (1984), 127–146.
- E. Hewitt and K. A. Ross, Abstract harmonic analysis I, II, Springer-Verlag, Berlin, 1963, 1970.
- 64. C. Herz, *Harmonic synthesis for subgroups*, Ann. Inst. Fourier (Grenoble) 23 (1973), no. 3, 91–123.
- N. Hindman and D. Strauss, Algebra in the Stone-Čech Compactification, de Gruyter Exp. Math. 27, Walter de Gruyter, Berlin, (1998).
- 66. K. H. Hofmann and S. A. Morris, *The structure of compact groups. A primer for the student a handbook for the expert.*, 2nd edition. Walter de Gruyter & Co., Berlin, 2006.
- 67. Z. Hu, On the set of topologically invariant means on the von Neumann algebra VN(G), Illinois J. Math. **39** (1995), 463–490.
- 68. _____, Extreme non-Arens regularity of quotients of the Fourier algebra A(G), Colloq. Math. **72** (1997), 237–249.
- 69. _____, Inductive extreme non-Arens regularity of the Fourier algebra A(G), Studia Math. **151** (2002), 247–264.
- 70. _____, Open subgroups and the centre problem for the Fourier algebra, Proc. Amer. Math. Soc. 134 (2006), 3085–3095.
- Z. Hu and M. Neufang, Decomposability of von Neumann algebras and the Mazur property of higher level, Canad. J. Math. 58 (2006), 768–795.
- Z. Hu, M, M. Sangani and T. Traynor, On character amenable Banach algebras, Studia Math. 193 (2009), no. 1, 53–78.
- N. Işik, J. Pym, and A. Ülger, The second dual of the group algebra of a compact group, J. London Math. Soc. 35 (1987), 135–158.
- 74. T. Jech, Set Theory, Academic Press, New York, 1978.
- E. Kaniuth and A. T.-M. Lau, Fourier algebras and amenability, Banach algebras and their applications, 181-192, Contemp. Math., 363, Amer. Math. Soc., Providence, RI, 2004.

- E. Kaniuth, A. Lau, and J. Pym, On φ-amenability of Banach algebras, Math. Proc. Cambridge Philos. Soc. 144 (2008), 85–96.
- 77. E. Kaniuth, A. Lau, and J. Pym, On character amenability of Banach algebras, J. Math. Anal. Appl. 344 (2008), 942–955.
- A. T.-M. Lau, Uniformly continuous functionals on the Fourier algebra of any locally compact group, Trans. Amer. Math. Soc. 251 (1979), 39–59.
- 79. A. T.-M. Lau, Analysis on a class of Banach algebras with applications to harmonic analysis on locally compact groups and semigroups, Fund. Math. 118 (1983), no. 3, 161–175
- A. T.-M. Lau, Continuity of Arens multiplication on the dual space of bounded uniformly continuous functions on locally compact groups and topological semigroups, Math. Proc. Cambridge Phil. Soc. 99(1986), 273–283.
- A. T.-M. Lau, Uniformly continuous functionals on Banach algebras, Colloquium Math. 51(1987), 195-205.
- A. T.-M. Lau, Fourier and Fourier-Stieltjes algebras of a locally compact group and amenability, Topological vector spaces, algebras and related areas (Hamilton, ON, 1994), 79-92, Pitman Res. Notes Math. Ser., 316, Longman Sci. Tech., Harlow, 1994.
- A. T.-M. Lau and V. Losert, On the second conjugate algebra of L¹(G) of a locally compact group, J. London Math. Soc. 37 (1988), 464–470.
- The C*-algbera generated by the operators with compact support on a locally compact group, J. Funct. Anal. 112 (1993), 1–30.
- The center of the second conjugate algebra of the Fourier algebra for infinite products of groups, Math. Proc. Camb. Phil. Soc. 138 (2005), 27–39.
- Lau, A. T.; Milnes, P.; Pym, J. S. Locally compact groups, invariant means and the centres of compactifications. J. London Math. Soc. (2) 56 (1997), no. 1, 77–90.
- A. T.-M. Lau and A. L. T. Paterson, The exact cardinality of the set of topological left invariant means on an amenable locally compact group, Proc. Amer. Math. Soc. 98 (1986) 75–80.
- A. T. Lau and J. Pym, Concerning the second dual of the group algebra of a locally compact group, J. London Math. Soc. (2) 41 (1990) 445–460.
- 89. A. T. Lau, A. R. Medghalchi and J. S. Pym, On the spectrum of $L^{\infty}(G)$, J. London Math. Soc. (2) 48 (1993), no. 1, 152–166.
- A. T.-M. Lau and J. S. Pym, The topological centre of a compactification of a locally compact group, Math. Z. 219 (1995), no. 4, 567–79.
- A. T.-M. Lau and A. Ülger, Topological centers of certain dual algebras, Trans. Amer. Math. Soc. 348 (1996), 1191–1212.
- V. Losert, Properties of the Fourier algebra that are equivalent to the amenability, Proc. Amer. Math. Soc. 92 (1984), 347–354.
- V. Losert, On tensor products of Fourier algebras, Arch. Math. (Basel) 43 (1984), no. 4, 370–372.
- 94. _____, Properties of the Fourier algebra that are equivalent to amenability, Proc. Amer. Math. Soc. 92 (1984), 347–354.
- Further results on the centre of the bidual of Fourier algebras, Lectures in Harmonic Analysis, Operator Algebras and representations, CIRM 3-08.11.2008.
- V. Losert, The centre of the bidual of Fourier algebras (discrete groups), Bull. Lond. Math. Soc. 48 (2016), no. 6, 968–976.
- M. S. Monfared, *Character amenability of Banach algebras*, Math. Proc. Cambridge Philos. Soc. 144 (2008), no. 3, 697–706.
- C. Nebbia, Multipliers and asymptotic behaviour of the Fourier algebra of nonamenable groups, Proc. Amer. Math. Soc. 84 (1982), 549-554.
- M. Neufang, Solution to a conjecture by Hofmeier-Wittstock, J. Funct. Anal. 217 (2004), 171–180.
- 100. _____, A unified approach to the topological centre problem for certain Banach algebras arising in abstract harmonic analysis, Arch. Math. 82 (2004), 164–171.
- 101. _____, On a conjecture by Ghahramani-Lau and related problems concerning topological centres, J. Funct. Anal. 224 (2005), 217–229.
- 102. _____, On Mazur's property and property (X), J. Operator Theory 60 (2008), no. 2, 301–316.
- 103. _____, On the topological centre problem for weighted convolution algebras and semigroup compactifications Proc. Amer. Math. Soc. 136 (2008), no. 5, 1831–1839.
- 104. _____, On one-sided strong Arens irregularity, Arch. Math. (Basel) 92 (2009), no. 5, 519–524.
- 105. J. Pier, Amenable locally compact groups, Wiley-Interscience, New York, 1984.

- 106. I. V. Protasov and J. S. Pym, Continuity of multiplication in the largest compactification of a locally compact group, Bull. London Math. Soc. 33 (2001), no. 3, 279–282.
- 107. H.P. Rosenthal, A characterization of Banach spaces containing $\ell^1,$ Proc. Nat. Acad. Sci. U.S.A., 71 (1974), 2411-2413.
- S. Sherman, The second adjoint of a C*-algebra, Proc. International Congress Mathematicians, Cambridge, Mass., Volume 1, (1950), 470.
- 109. R. Stokke, On Beurling measure algebras, Comment. Math. Univ. Carol. (to appear).
- 110. Z. Takeda, Conjugate spaces of operator algebras, Proc. Japan Academy, 30 (1954), 90–95.
- 111. E. K. van Douwen, The number of cofinal ultrafilters, Topol. Appl. 39 (1991), 61-63.
- 112. S. Zadeh, Isometric isomorphisms of Beurling algebras, J. Math. Anal. Appl. 438 (2016), no. 1, 1–13.
- 113. Y. Zelenyuk, The number of minimal right ideals of βG, Proc. Amer. Math. Soc. 137 (2009), no. 7, 2483–2488.

MAHMOUD FILALI, DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF OULU, OULU, FINLAND.

E-MAIL: mfilali@cc.oulu.fi

Jorge Galindo, Instituto Universitario de Matemáticas y Aplicaciones (IMAC), Universidad Jaume I, E-12071, Castellón, Spain. E-mail: jgalindo@mat.uji.es