Orthospectrum and simple orthospectrum rigidity: finiteness and genericity.

Nolwenn Le Quellec

2024

Abstract. We study the orthospectrum and the simple orthospectrum of compact hyperbolic surfaces with geodesic boundary. We show that there are finitely many hyperbolic surfaces sharing the same simple orthospectrum and finitely many hyperbolic surfaces sharing the same orthospectrum. Then, we show that generic surfaces are determined by their orthospectrum and by their simple orthospectrum. We conclude with the example of the one-holed torus which is determined by its simple orthospectrum.

1 Introduction

In a by now famous paper [\[Kac66\]](#page-29-0), Kac asked "Can one hear the shape of a drum?". The question can be formalized mathematically using the fact that the sound made by a drum is linked to the frequencies at which a drumhead vibrates, that is, the spectrum of the Laplacian. Kac's question then becomes: "Does the spectrum of the Laplacian of a planar domain determines the domain itself?". This question was then also asked for general Riemannian manifolds. In the case of closed hyperbolic surfaces, Huber and Selberg (see [\[Bus10,](#page-29-1) Chapter 7.1]) showed that the Laplacian spectrum determines and is determined by the length spectrum, which is the set of lengths of closed geodesics counted with multiplicities. This shifted the question to "Does the length spectrum of a hyperbolic surface determine the metric up to isometry?". In 1978, Vignéras provided the first example of isospectral non-isometric closed hyperbolic surfaces [\[Vig78\]](#page-30-0) showing that the answer to the question is negative, and in 1985, Sunada gave a general criterion to construct isospectral non-isometric manifolds [\[Sun85\]](#page-30-1). On the other hand, McKean proved that there is a finite number of isospectral non-isometric hyperbolic surfaces [\[McK72\]](#page-30-2). In 1979, Wolpert showed that a generic hyperbolic surface is determined by its length spectrum [\[Wol79\]](#page-30-3). In 1985, Haas proved that the one-holed hyperbolic torus with a fixed boundary length is determined by the length spectrum [\[Haa85\]](#page-29-2), and later Buser and Semmler removed the condition on the boundary length [\[BS88\]](#page-29-3). In the case of the simple spectrum (where only the simple closed geodesics are considered) the question is still open: there are no known examples of non-isometric hyperbolic surfaces with the same simple length spectrum. There is, however, a result by Baik, Choi and Kim showing that generic hyperbolic surfaces are also determined by their simple length spectrum [\[BCK20\]](#page-29-4).

In 1993, Basmajian introduced the orthospectrum of hyperbolic surfaces with boundary, defined as the set of lengths of geodesic arcs orthogonal to the boundary (called or-

thogeodesics) counted with multiplicities [\[Bas93\]](#page-29-5). This object, analogous to the length spectrum, aims at taking the boundary of a surface more into account. In particular, Basmajian gave a formula to compute the boundary length of a hyperbolic surface from its orthospectrum [\[Bas93\]](#page-29-5). Motivated by Kac's question, Masai and McShane considered the analogous problem for the orthospectrum: "Does the orthospectrum of a hyperbolic surface determine the metric up to isometry?". They showed that there is a finite number of non-isometric hyperbolic surfaces with one boundary component sharing the same orthospectrum [\[MM23\]](#page-30-4). In the same paper, they showed that the one-holed torus is determined its orthospectrum. They also gave a general construction for isospectral non-isometric hyperbolic surfaces, thus answering the question in the negative.

In this paper, we investigate further Masai and McShane's question, both for the orthospectrum and the simple orthospectrum. The simple orthospectrum being the multiset of lengths of simple orthogeodesics, counted with multiplicities. Note that even if the orthospectrum and simple orthospectrum seems related, there no know way to deduced one from the other. Denoting by $\mathcal{O}(X)$ and $\mathcal{O}_S(X)$ the orthospectrum and the simple orthospectrum of a hyperbolic surface X , our main results are the following.

Theorem [3.1.](#page-11-0) Let S_g^b be a compact, genus g surface of negative Euler characteristic with b boundary components. Let X be a hyperbolic structure on S_g^b with geodesic boundary. Then, up to isometry, there is a finite number of hyperbolic structures Y on S_g^b such that

$$
\mathcal{O}_S(Y)=\mathcal{O}_S(X).
$$

Similarly, up to isometry, there is also a finite number of hyperbolic structures Y on S_g^b such that

$$
\mathcal{O}(Y) = \mathcal{O}(X).
$$

The proof is inspired by Masai and McShane's proof but diverges in the way we control the systole of the surfaces (*i.e.*, the length of the shortest closed geodesic). Our proof also works for the orthospectrum, thus extending Masai and McShane's result to hyperbolic surfaces with any finite number of boundary components. We also establish an analogue of Wolpert's result both for the orthospectrum and the simple orthospectrum.

Theorem [4.1.](#page-19-0) Generic surfaces in $\text{Teich}(S_g^b)$ are determined, up to isometry, by their orthospectrum. Similarly, generic surfaces in Teich (S_g^b) are also determined, up to isometry, by their simple orthospectrum.

We took inspiration in Wolpert's proof of the generic determination by the length spectrum. The main difference is the use of a different type of coordinates for the Teichmüller space. The same proof works both for the orthospectrum and the simple orthospectrum.

Finally, we show:

Theorem [5.2.](#page-25-0) Let T and T' be two hyperbolic structures with geodesic boundary on the one-holed torus. Then T and T' are isometric if and only if $\mathcal{O}_S(T) = \mathcal{O}_S(T')$.

This paper is organized as follows. We recall in Section [2](#page-2-0) properties of hyperbolic surfaces and geodesics that will be needed in the rest of the paper. In Section [3,](#page-11-1) we prove Theorem [3.1.](#page-11-0) The proof is divided in several steps corresponding each to a subsection. Using a theorem from Section [3,](#page-11-1) we prove Theorem [4.1](#page-19-0) in Section [4.](#page-19-1) Finally in Section [5,](#page-24-0) we show Theorem [5.2.](#page-25-0)

Acknowledgments. We appreciate the support and help of Federica Fanoni and Stéphane Sabourau, both PhD advisors of the author, in particular for their help in the redaction.

2 Preliminaries

2.1 Curves and arcs on surfaces.

Throughout this paper, S_g^b will denote an orientable, compact, genus g surface of negative Euler characteristic with $b > 0$ boundary components. Moreover, arcs are always defined with endpoints on the boundary and closed curves are always considered primitive.

Definition 2.1. A closed curve on S_g^b is *essential* if it is not homotopic to a boundary component or to a point. It is simple if it has no self-intersection.

Definition 2.2. A pair of pants is a surface homeomorphic to S_0^3 . A pants decomposition of S_g^b is a maximal collection of pairwise non-homotopic, disjoint, essential simple closed curves on S_g^b .

Remark 2.3. The cardinality of a pants decomposition is $3g + b - 3$ and a pants decomposition cuts S_g^b into $2g + b - 2$ pairs of pants [\[Abi80\]](#page-29-6).

When we study surfaces with boundary, it can be useful to change the perspective from the closed curve/closed geodesic point of view to the arc/orthogeodesic point of view.

Definition 2.4. An arc on S_g^b is *essential* if it is not homotopic relatively to the boundary, into a boundary component. It is simple if it has no self-intersection.

An *orthogeodesic* of a compact hyperbolic surface is the shortest geodesic representative of the homotopy class relative to the boundary of an essential arc.

Remark 2.5. Endpoints of an orthogeodesic are orthogonals to the boundary [\[Bas93\]](#page-29-5).

Definition 2.6. A hexagon decomposition of a surface S_g^b is a maximal collection of pairwise non-homotopic and disjoint simple essential arcs on S_g^b .

Remark 2.7. There always exists a hexagon decomposition on a negative Euler characteristic surface. The cardinality of a hexagon decomposition is $6g + 3b - 6$ and the hexagon decomposition cuts S_g^b into $4g + 2b - 4$ hexagons. See [\[Ush99\]](#page-30-5).

When we endow S_g^b with a hyperbolic metric, and cut the surface along the orthogeodesic representatives of a hexagon decomposition, the surface decomposes into right-angled hexagons.

Figure 1: Example of a hexagon decomposition of the one-holed torus.

There is a link between pants and hexagon decomposition through the concept of double of surfaces.

Definition 2.8. Let S be a surface of genus q with $b > 0$ boundary components. The *double of* S is the surface obtained by gluing two copies of S along their corresponding boundary components. When S is endowed with a hyperbolic metric with geodesic boundary components, the double of S is endowed with the hyperbolic metric which coincides with the hyperbolic metric of S on each copy of S and such that, the reflection R between the two copies of S along their boundary is an isometry.

Let a be an arc of S. We call the *double of a* the union of the two copies of a on the double of S.

Remark 2.9. The double of S_g^b is the surface S_{2g+b-1} of genus $2g + b - 1$ with no boundary.

Lemma 2.10. Let H be a hexagon decomposition of S_g^b . Denote by S_{2g+b-1} the double of S_g^b and by R the reflection associated to it. Then, $\mathcal{H} \cup R(\mathcal{H})$ is a pants decomposition on S_{2g+b-1} .

Proof. The double of every arc of a hexagon decomposition forms a simple closed curve of S_{2g+b-1} . By construction, $\mathcal{H} \cup R(\mathcal{H})$ is a set of $6g + 3b - 6$ pairwise non-homotopic, disjoint, essential, simple, closed curves on S_{2q+b-1} . Therefore, it is a maximal collection of such curves, and hence a pants decomposition. \Box

Now, let us define the geometric intersection number between closed curves and arcs.

Definition 2.11. Let α and β be closed curves or arcs on a surface S. The geometric intersection number $i(\alpha, \beta)$ between α and β is the minimum number of intersection counted with multiplicities between curves/arcs in their homotopy class (relative to the boundary).

Note that, if α and β are different closed geodesics or orthogeodesics $i(\alpha, \beta) = |\alpha \cap \beta|$.

2.2 Teichmüller space and Moduli space.

The hyperbolic surfaces studied in this paper either live in the Teichmüller space or the moduli space of S_g^b . Moreover, we assume they all have a geodesic boundary.

Definition 2.12. The *Teichmüller space* $\text{Teich}(S_g^b)$ of S_g^b is the set of homotopy classes of hyperbolic structures on S_g^b .

Definition 2.13. The mapping class group $\mathcal{MCG}(S_g^b)$ of S_g^b is the quotient of the group of orientation-preserving homeomorphisms of S_g^b by the subgroup of homeomorphisms isotopic to the identity.

$$
\mathcal{MCG}(S_g^b) = \text{Homeo}^+(S_g^b) / \text{Homeo}_0(S_g^b).
$$

Definition 2.14. The moduli space $\mathcal{M}(S_g^b)$ is the space of isometry classes of hyperbolic surfaces homeomorphic to S_g^b .

The moduli space is the quotient space

$$
\mathcal{M}(S_g^b) = \mathrm{Teich}(S_g^b) / \mathcal{MCG}(S_g^b)
$$

and Teichmüller space is the universal cover of $\mathcal{M}(S_g^b)$.

The following function on Teich (S_g^b) will be used throughout this paper.

Definition 2.15. For any essential closed curve/arc α on S_g^b and any hyperbolic surface $\chi \in \text{Teich}(S_g^b)$, the *length function* $\ell_{\chi}(\alpha)$ is the length of the shortest curve/arc in the homotopy class of α (relative to the boundary if α is an arc) on χ . If C is a collection of curves and arcs, we define $\ell_{\chi}(\mathcal{C}) = \sum_{\alpha \in \mathcal{C}} \ell_{\chi}(\alpha)$.

We may write $\ell_X(\alpha)$ with $X \in \mathcal{M}(S_g^b)$ instead of its lift $\chi \in \text{Teich}(S_g^b)$ when there is no ambiguity on the marking of χ . When there is no ambiguity about the surface, we may just write $\ell(\alpha)$. With the length function and hexagon decomposition, we can now define a system of coordinates on the Tecihmüller space. A right-angled hexagon is determined up to isometry by the lengths of three pairwise non-consecutive sides [\[Rat19,](#page-30-6) Theorem 3.5.14]. Thus, if we fix a hexagon decomposition on S_g^b , the lengths of the orthogeodesic representatives of the hexagon decomposition on S_g^b determine a hyperbolic surface in the Teichmüller space. Furthermore, Ushijima showed in [\[Ush99,](#page-30-5) Theorem 4.1] the following theorem:

Theorem 2.16. Given a hexagon decomposition $\mathcal{H} = {\alpha_1, ..., \alpha_{6g+3b-6}}$ on S_g^b , the map

$$
\varphi_{\mathcal{H}} : \mathrm{Teich}(S_g^b) \to \mathbb{R}_+^{6g+3b-6}
$$

defined by

$$
\varphi_{\mathcal{H}}(\chi) = (\ell_{\chi}(\alpha_1), ..., \ell_{\chi}(\alpha_{6g+3b-6}))
$$

is a homeomorphism.

Definition 2.17. For a fixed hexagon decomposition $\mathcal{H} = {\alpha_1, ..., \alpha_{6q+3b-6}}$, we call $\varphi_{\mathcal{H}}(\cdot) = (\ell(\alpha_1), ..., \ell(\alpha_{6q+3b-6}))$ Ushijima coordinates function and denote by

$$
\chi^{\omega} = \varphi_{\mathcal{H}}^{-1}(\omega)
$$

the surface in Teich (S_g^b) associated to $\omega \in \mathbb{R}^{6g+3b-6}_+$.

Teichmüller space of S_g^b admits a real analytic structure and the Fenchel-Nielsen coordinates are real analytic (independently from the pants decomposition) [\[Abi80\]](#page-29-6). By identifying $\text{Teich}(S_g^b)$ with a real analytic subvariety of $\text{Teich}(S_{2g+b-1})$ we obtain that this structure is compatible with the one we get via Ushijima's coordinates: fix a hexagon decomposition $\mathcal{H} = \{\alpha_1, ..., \alpha_{6g+3b-6}\}\$ on S_g^b , by Lemma [2.10](#page-3-0) this hexagon decomposition induce a pants decomposition on S_{2g+b-1} , the double of S_g^b , we denote it $\mathcal{P} = \{\alpha'_1, ..., \alpha'_{6g+3b-6}\}\$ where α'_i is the double of α_i . We denote by $T(S_g^b)$ the subset of surfaces in Teich (S_{2g+b-1}) that are double of surfaces in Teich (S_g^b) , which is identified with Teich (S_g^b) . We have

$$
T(S_g^b) = \{ \chi \in \text{Teich}(S_{2g+b-1}) \mid t_{\alpha'_i}(\chi) = 0, 1 \leq i \leq 6g + 3b - 6 \}
$$

with $t_{\alpha_i'}(.)$ the twists parameters in the Fenchel-Nielsen coordinates. The lengths parameters are real analytic on $T(S_g^b)$ and by identifying Teich (S_g^b) with $T(S_g^b)$ Ushijima's coordinates $(\ell(\alpha_1), ..., \ell(\alpha_{6g+3b-6})) = (\frac{\ell(\alpha'_1)}{2})$ $\frac{\alpha'_1}{2}, ..., \frac{\ell(\alpha'_{6g+3b-6})}{2}$ $\frac{+3b-6}{2}$ are real analytic. This does not depend on the hexagon decomposition. Through the same process with a pants decomposition on S_{2g+b-1} induced by a pants decomposition on S_g^b , we show that Fenchel-Nielsen coordinates are real analytic for the same analytic structure as Ushijima's coordinates.

Lemma 2.18. For any arc β , the function

$$
\ell_{\bullet}(\beta) : \text{Teich}(S_g^b) \to \mathbb{R}^+ \to \ell_{\chi}(\beta)
$$

is real analytic on Teich (S_g^b) .

Proof. Let β be an arc on S_g^b and β' be the double of β on S_{2g+b-1} . By [\[Bus10,](#page-29-1) Lemma 10.2.3 (see also [\[MP08,](#page-30-7) Proposition 2.4]), we know that for any closed curve β' the function $\ell(\beta')$ is real analytic on Teich(S_{2g+b-1}) and thus on $T(S^g_b)$ $\binom{g}{b}$. By identifying $T(S_g^b)$ and Teich (S_g^b) , we get that $\ell(\beta) = \frac{\ell(\beta')}{2}$ $\frac{\beta'}{2}$ is real analytic on Teich (S_g^b) . \Box

Definition 2.19. Given two surfaces χ , $\Upsilon \in \text{Teich}(S_g^b)$ and $K \geq 1$, a homeomorphism $\phi: \chi \to \Upsilon$ is said to be K-quasiconformal if its distributional derivatives are locally in L_2 and

$$
|\phi_{\overline{z}}| \leqslant \frac{K-1}{K+1} |\phi_z|
$$
 a.e.

For any $\chi, \Upsilon \in \text{Teich}(S_g^b)$, there exists a unique K-quasiconformal map $\phi_{\chi, \Upsilon}$ between χ and Υ with K minimal (see [\[Abi80\]](#page-29-6) for more details). The Teichmüller metric d_{Teich} is defined by $d_{\text{Teich}}(\chi, \Upsilon) = \frac{1}{2} \log(K)$; see [\[FM12\]](#page-29-7).

Now, let us state an extension of Wolpert's lemma [\[FM12,](#page-29-7) 12.3.2] to any closed curve, simple or not, due to Buser [\[Bus10,](#page-29-1) Theorem 6.4.3].

Lemma 2.20 (Wolpert's lemma). Let $\phi : \chi_1 \to \chi_2$ be a K-quasiconformal homeomorphism between two hyperbolic surfaces χ_1 and χ_2 . For any isotopy class c of a closed curve in χ_1 , the following inequalities hold:

$$
\frac{\ell_{\chi_1}(c)}{K} \leq \ell_{\chi_2}(\phi(c)) \leqslant K\ell_{\chi_1}(c).
$$

The previous lemma applied to the double of the surface implies the following.

Lemma 2.21 (Orthogeodesic Wolpert's lemma). Let $\phi : \chi_1 \to \chi_2$ be a K-quasiconformal homeomorphism between two hyperbolic surfaces χ_1 and χ_2 . For any isotopy class c of arc in χ_1 , the following inequalities hold:

$$
\frac{\ell_{\chi_1}(c)}{K} \leq \ell_{\chi_2}(\phi(c)) \leqslant K\ell_{\chi_1}(c).
$$

By definition of the Teichmüller metric, for any $\chi_1, \chi_2 \in \text{Teich}(S_g^b)$ with $d_{\text{Teich}}(\chi_1, \chi_2) = \log(K)/2$, there is a K-quasiconformal homeomorphism between χ_1 and χ_2 . By Lemma [2.21,](#page-6-0) for any $\chi_1, \chi_2 \in \text{Teich}(S_g^b)$ with $d_{\text{Teich}}(\chi_1, \chi_2) \leq \log(K)/2$ and any isotopy class c of simple arcs on S_g^b , we have $\frac{\ell_{\chi_1}(c)}{K} \leq \ell_{\chi_2}(c) \leq K\ell_{\chi_1}(c)$.

Then, we state a corollary of Lemma [2.21.](#page-6-0)

Corollary 2.22. Let $\mathcal{Q} \subset \text{Teich}(S_g^b)$ be a compact subset. There exists a constant $q \geqslant 1$ which depends only on Q such that

$$
\frac{\ell_\chi(\beta)}{q} \leqslant \ell_{\chi'}(\beta) \leqslant q\ell_\chi(\beta)
$$

for any $\chi, \chi' \in \mathcal{Q}$ and any orthogeodesic β .

Proof. Let $q > 0$ such that diam $\mathcal{Q} = \frac{1}{2}$ $\frac{1}{2} \log(q)$. Then for any $\chi, \chi' \in \mathcal{Q}$, we have $d_{\text{Teich}}(\chi, \chi') \leq \log(q)/2$. By Lemma [2.21,](#page-6-0) we have

$$
\frac{\ell_\chi(\beta)}{q} \leqslant \ell_{\chi'}(\beta) \leqslant q\ell_\chi(\beta).
$$

 \Box

Definition 2.23. The *systole* $sys(X)$ of a hyperbolic surface X is the shortest length of an essential closed curve on X . Note that the systole is realized by the length of a simple closed geodesic, unless X is a pair of pants.

Similarly to the systole, we also define the orthosystole.

Definition 2.24. The *orthosystole* $\cos(X)$ of a hyperbolic surface X is the shortest length of an orthogeodesic on X.

Finally, let us define an interesting subset of the moduli space.

Definition 2.25. Let $\gamma_0, ..., \gamma_{b-1}$, be the boundary component of S_g^b . We define the set

$$
\mathcal{M}_{A,\varepsilon}(S_g^b) = \{ X \in \mathcal{M}(S_g^b) \mid \text{sys}(X) \geq \varepsilon \text{ and } A \geq \ell_X(\gamma_0), \dots, \ell_X(\gamma_{b-1}) \geq \varepsilon \}
$$

The following result by Parlier [\[Par23\]](#page-30-8) will help us define a property of $\mathcal{M}_{A,\varepsilon}(S_g^b)$.

Theorem 2.26. Let X be a finite area hyperbolic surface, possibly with geodesic boundary ∂X . Then X admits a pants decomposition where each curve is of length at most

$$
B = \max{\{\ell(\partial X), \text{area}(X)\}}.
$$

Theorem 2.27. For all $A \geq \varepsilon > 0$, $\mathcal{M}_{A,\varepsilon}(S_g^b)$ is compact.

This result was proven by Mumford [\[Mum71\]](#page-30-9) in the closed case. In [\[LSZ15\]](#page-30-10), a different version of the theorem is stated. Here, we use Farb and Margalit's proof in [\[FM12,](#page-29-7) Chap. 12] in the case $b = 0$ and adapt it to the case $b > 0$ to prove Theorem [2.27.](#page-7-0) With Corollary [3.3,](#page-12-0) we obtain an equivalence of Theorem [2.27](#page-7-0) with [\[LSZ15,](#page-30-10) Theorem 4.1].

Proof. Since $\mathcal{M}(S_g^b)$ inherits the Teichmüller metric from Teich (S_g^b) , we just need to show that $\mathcal{M}_{A,\varepsilon}(S_g^b)$ is sequentially compact. Let (X_i) be a sequence in $\mathcal{M}_{A,\varepsilon}(S_g^b)$ and $\chi_i \in \text{Teich}(S_g^b)$ a lift of X_i for all i.

To show that a subsequence of (X_i) converges in $\mathcal{M}_{\varepsilon}(S_g^b)$, we show that for a fixed choice of Fenchel-Nielsen coordinates, we can choose lifts χ_i of X_i to Teich (S_g^b) inside a rectangular compact set of the Euclidean space $\mathbb{R}^{3g-3+2b}_{>0} \times \mathbb{R}^{3g-3+b}$.

By Theorem [2.26,](#page-7-1) there is a pants decomposition \mathcal{P}_i of S_g^b such that $\ell_{\chi_i}(\gamma) \in [\varepsilon, B]$ for all $\gamma \in \mathcal{P}_i$, where $B = \max(bA, 2\pi(2g + b - 1)).$

Since there are a finitely many topological types of pants decompositions of S_g^b , we can choose a sequence (f_i) in $\mathcal{MCG}(S_g^b)$ such that, up to passing to a subsequence, $f_i(\mathcal{P}_i) = \mathcal{P}_1$. The hyperbolic structure $\Upsilon_i = f_i \cdot \chi_i$ is also a lift of X_i , whose length parameters in the Fenchel-Nielsen coordinates with respect to \mathcal{P}_1 are between ε and B.

Since the Dehn twists on the curves in \mathcal{P}_1 change the twist parameters by 2π , there is a product h_i of Dehn twists on the curves of \mathcal{P}_1 such that the twist parameters of $h_i \Upsilon_i$ are between 0 and 2π . Thus, the lifts $h_i \Upsilon_i$ of X_i are all inside a compact set. Therefore, there exists a converging subsequence which projects to a converging subsequence of (X_i) . \Box

Then we state a corollary of Theorem 2.[27](#page-7-0).

Corollary 2.28. Let $A \geq \varepsilon > 0$. There exists a compact subset $\mathcal{Q}(A, \varepsilon) \subset \text{Teich}(S_g^b)$ such that for each surface $\chi \in \text{Teich}(S_b^b)$ with $\text{sys}(\chi) \geq \varepsilon$ and boundary component lengths between ε and A, there exists an isometric surface $\chi' \in \mathcal{Q}(A,\varepsilon)$.

Proof. By Theorem [2.27,](#page-7-0) the set $\mathcal{M}_{A,\varepsilon}(S_g^b)$ of hyperbolic surfaces X, up to isometry, with $sys(X) \geq \varepsilon$ and boundary component length between ε and A is compact. We set $\mathcal{Q}(A,\varepsilon)$ a compact lift of $\mathcal{M}_{A,\varepsilon}(S_g^b)$ in the Teichmüller space. By definition, any surface $\chi' \in \text{Teich}(S_g^b)$ with $\text{sys}(\chi') \geqslant \varepsilon$ and boundary component length between ε and A is sent by the cover on a surface $X' \in \mathcal{M}_{A,\varepsilon}(S_g^b)$. By construction, there is a lift χ of X' in $\mathcal{Q}(A,\varepsilon)$ and χ is isometric to χ' . \Box

2.3 Orthospectrum.

Let us introduce the orthospectrum, an object analogous to the length spectrum and first defined by Basmajian in [\[Bas93\]](#page-29-5).

Definition 2.29. The *orthospectrum* of a hyperbolic surface X is the multiset $\mathcal{O}(X)$ of lengths of orthogeodesics on X , counted with multiplicities.

Along with its definition, Basmajian showed the following in [\[Bas93\]](#page-29-5).

Theorem 2.30. The orthospectrum is discrete.

We also define the simple orthospectrum, which is the focus of Theorem [1.](#page-0-0)

Definition 2.31. The *simple orthospectrum* of a hyperbolic surface X is the multiset $\mathcal{O}_S(X)$ of lengths of simple orthogeodesics on X counted with multiplicities.

Among the properties of the orthospectrum, we highlight the following one by [\[Bas93\]](#page-29-5), which shows that the boundary length of a hyperbolic surface is determined by its orthospectrum.

Theorem 2.32 (Basmajian's Identity.). Let X be a compact hyperbolic surface with geodesic boundary. Then,

$$
\ell(\partial X) = \sum_{\ell \in \mathcal{O}(X)} B(\ell)
$$

where $B(\ell) = 2 \sinh^{-1} \left(\frac{1}{\sinh \ell} \right)$ $\frac{1}{\sinh(\ell)}$. Note that B is a positive decreasing function.

For the simple orthospectrum, the theorem implies that

$$
\ell(\partial X) > \sum_{\ell \in \mathcal{O}_S(X)} B(\ell).
$$

Thus, X has a boundary component of length greater than $\frac{\sum_{\ell \in \mathcal{O}_S(X)} B(\ell)}{h}$ $\frac{b^{(X)D(x)}}{b}$.

2.4 Hyperbolic geometry.

Let us state several properties on geodesics and orthogeodesics using the length function. The following result is shown in [\[Bus10,](#page-29-1) Theorem 4.2.1].

Theorem 2.33. Let X be a hyperbolic surface. Then every non-simple closed geodesic on X has length greater than 1.

By doubling the surface, we deduce

Corollary 2.34. Let X be a hyperbolic surface. Then every non-simple orthogeodesic on X has length greater than $\frac{1}{2}$.

We also have:

Lemma 2.35 (Half-collar lemma). Let Y be a pair of pants with boundary geodesics $\gamma_1, \gamma_2, \gamma_3$. The sets

$$
\mathcal{C}^*[\gamma_i] = \{ p \in Y \mid \sinh(\text{dist}(p, \gamma_i)) \sinh(\frac{1}{2}\ell(\gamma_i)) \leq 1 \}
$$

for $i = 1, 2, 3$ are pairwise disjoint and each of them is homeomorphic to a cylinder.

Lemma 2.36 (Collar lemma). Let α and β be two distinct closed geodesics on a hyperbolic surface X such that $\alpha \cap \beta \neq \emptyset$. If β is simple, then

$$
\sinh\left(\frac{\ell(\alpha)}{2}\right)\sinh\left(\frac{\ell(\beta)}{2}\right) > 1.
$$

The proofs of Lemmas [2.35](#page-9-0) and [2.36](#page-9-1) can be found in [\[Bus10\]](#page-29-1). We can state a version of the last one with orthogeodesics instead of closed geodesics, which can be deduced by doubling the surface.

Lemma 2.37 (OrthoCollar lemma). Let α and β be two distinct orthogeodesics on a hyperbolic surface X such that $\alpha \cap \beta \neq \emptyset$. If β is simple, then

$$
\sinh(\ell(\alpha))\sinh(\ell(\beta)) > 1.
$$

Now, let us state hyperbolic trigonometry formulas that we will need in the different proofs of this paper.

Lemma 2.38. For any right-angled hexagon with consecutive sides β , c, α , b , γ , a , we have

$$
\cosh(c) = \sinh(a)\sinh(b)\cosh(\gamma) - \cosh(a)\cosh(b). \tag{1}
$$

For every trirectangle with sides labelled as in Figure [2,](#page-9-2) the following relation is true:

$$
\cos(\varphi) = \tanh(\sigma)\tanh(\tau).
$$

Figure 2: Right-angled hexagon and trirectangle

The proofs can be found in [\[Bus10\]](#page-29-1).

As already mentioned, any right-angled hexagon is determined by the lengths of three pairwise disjoint sides. A right-angled octagon can be obtained by gluing two rightangled hexagons along one side. Thus, any right-angled octagon is determined by the length of four pairwise disjoint sides and the length of one orthogonal arc between opposite sides. In the following lemma, we will see how to compute the length of the orthogonal between the last two other sides.

Lemma 2.39 (Right-angled octagon). We define the function $f_{\text{octa}} : \mathbb{R}^5_+ \to \mathbb{R}_+$ given by

$$
f_{\text{octa}}(a, x, y, z, t) = -\cosh(x)\cosh(t) + \sinh(x)\sinh(t) \times
$$

$$
\cosh\left[\cosh^{-1}\left(\frac{\cosh(y) + \cosh(x)\cosh(a)}{\sinh(x)\sinh(a)}\right) + \cosh^{-1}\left(\frac{\cosh(y) + \cosh(t)\cosh(a)}{\sinh(t)\sinh(a)}\right)\right].
$$

For any right-angled octagon with four disjoint sides $\delta_1, \delta_2, \delta_3, \delta_4$ and two orthogonal arcs α and β joining two opposite sides different from the δ_i as in Figure [3,](#page-10-0) we have:

$$
\cosh(\beta) = f_{\text{octa}}(\alpha, \delta_1, \delta_2, \delta_3, \delta_4).
$$

Proof. We can decompose the octagon into two right-angled hexagons with three disjoint sides $(\beta, \delta_4, \delta_1)$ and $(\beta, \delta_2, \delta_3)$. We can also decompose it into two other rightangled hexagons with three disjoint sides $(\alpha, \delta_1, \delta_2)$ and $(\alpha, \delta_3, \delta_4)$. The arc α decomposes the side between δ_1 and δ_4 into two segments x and x' such that x is a side of the hexagon $(\alpha, \delta_1, \delta_2)$ and x' is a side of the hexagon $(\alpha, \delta_3, \delta_4)$ as in Figure [3.](#page-10-0)

Figure 3: Right-angled octagon

We then apply [\(1\)](#page-9-3) from Lemma [2.38](#page-9-4) to the hexagons $(\alpha, \delta_1, \delta_2)$, $(\alpha, \delta_3, \delta_4)$ and $(\beta, \delta_4, \delta_1)$:

$$
\cosh(x) = \frac{\cosh(\delta_2) + \cosh(\delta_1)\cosh(\alpha)}{\sinh(\delta_1)\sinh(\alpha)}
$$

$$
\cosh(x') = \frac{\cosh(\delta_3) + \cosh(\delta_4)\cosh(\alpha)}{\sinh(\delta_4)\sinh(\alpha)}
$$

$$
\cosh(\beta) = \sinh(\delta_1)\sinh(\delta_4)\cosh(x + x') - \cosh(\delta_1)\cosh(\delta_4).
$$

We use the first two equalities to compute $x + x'$ and we use the expression in the last equality to conclude. \Box

In [\[MM23,](#page-30-4) Lemma 3.2] , Masai and McShane prove the following result.

Lemma 2.40. Let P be a pair of pants with boundary geodesics α, β, γ such that $\ell(\beta) \leq \ell(\alpha)$. Let τ be the unique simple orthogeodesic with both endpoints on γ as in Figure [4.](#page-11-2) Then, we have

Figure 4

3 Finite characterization

We recall that S_g^b is a genus g surface of negative Euler characteristic with b boundary components. We label the boundary components $\gamma_0, \gamma_1, ..., \gamma_{b-1}$. We pick a hyperbolic structure X on S_g^b and we define

$$
I(X) = \{ Y \in \mathcal{M}(S_g^b) \text{ such that } \mathcal{O}_S(Y) = \mathcal{O}_S(X) \}.
$$

In this section, we are going to show [Theorem](#page-1-0) [3.1.](#page-11-0) We restate it as follows.

Theorem 3.1. Let X be a hyperbolic structure on S_g^b with geodesic boundary. Then, $I(X)$ is finite.

In the first step of the proof, we establish an upper bound on the length of every boundary component of a hyperbolic surface in $I(X)$. In the second step, we obtain a lower bound on the systole and the length of every boundary component of a hyperbolic surface from its simple orthospectrum. By Theorem [2.27,](#page-7-0) we deduce that $I(X)$ lies in a compact set of the moduli space. Finally, with the help of the previous steps and the discreteness of the orthospectrum (Theorem [2.30\)](#page-8-0), we deduce that $I(X)$ is not only included in a compact set, but it is compact and discrete, and therefore finite.

Remark 3.2. The idea of the proof comes from Masai and McShane's article [\[MM23\]](#page-30-4), where they show an analogous result for the orthospectrum of surfaces with a single boundary component. Our proof also works for the orthospectrum of surfaces with a finite number of component, so it recovers and extends their result.

3.1 Step one: Upper bounds

In this section, we show that there exists $A = A(\mathcal{O}_S(X))$ such that $\ell_Y(\gamma_i) \leq A$ for all $i \in \{0, ..., b-1\}$ and $Y \in I(X)$.

Let us state a corollary of [\[Bav05,](#page-29-8) Théorème 1].

Corollary 3.3. Let t be the orthosystole of X . Then

$$
\sinh(2t)\sinh\left(\frac{\ell(\partial X)}{24g+12b-24}\right) \leqslant \frac{1}{2}.
$$

The orthosystole is attained by the length of a simple orthogeodesic, meaning it is the smallest length in $\mathcal{O}_S(X)$. This gives us an upper bound

$$
A = (24g + 12b - 24)\sinh^{-1}\left(\frac{1}{2\sinh(2t)}\right)
$$

on $\ell(\partial X)$, and in particular on the length of any boundary component γ_i of X, which depends only on $\mathcal{O}_S(X)$, g and b.

3.2 Step two: Compactness

In this section, we will show the following intermediate result.

Theorem 3.4. Let $\varepsilon_1 \geq \varepsilon_2 > 0$. Then the set

$$
\mathcal{OM}_{[\varepsilon_1,\varepsilon_2]}(S_g^b) = \{ X \in \mathcal{M}(S_g^b) \mid \varepsilon_1 \leqslant \text{osys}(X) \leqslant \varepsilon_2 \}
$$

is included in a compact. In particular, there exist $A \geq \varepsilon > 0$ such that

$$
\mathcal{OM}_{[\varepsilon_1,\varepsilon_2]}(S_g^b) \subset \mathcal{M}_{A,\varepsilon}(S_g^b).
$$

Proof. By Corollary [3.3,](#page-12-0) we have

$$
A = (24g + 12b - 24)\sinh^{-1}\left(\frac{1}{2\sinh(2\varepsilon_1)}\right) \ge \ell(\partial X)
$$

Thus, there is an upper bound on the length of every boundary component of any surface in $\mathcal{OM}_{[\varepsilon_1,\varepsilon_2]}(S_g^b)$.

To show that $\mathcal{OM}_{[\varepsilon_1,\varepsilon_2]}(S_g^b) \subset \mathcal{M}_{A,\varepsilon}(S_g^b)$, we still need a lower bound on the systole and on the length of the boundary components of any surface in $\mathcal{OM}_{[\varepsilon_1,\varepsilon_2]}(S_g^b)$. By contradiction, let us suppose that there is an infinite family of hyperbolic surfaces $(X_n)_{n\in\mathbb{N}}\in\mathcal{OM}_{[\varepsilon_1,\varepsilon_2]}(S_g^b)$ which leaves every compact: $\min(\mathrm{sys}(X_n),\ell_{X_n}(\gamma_0),...,\ell_{X_n}(\gamma_{b-1}))\to 0$ as n goes to infinity.

From Basmajian's Identity [2.32,](#page-8-1) each surface X_n has a boundary component of length greater than $\frac{\sum_{\ell \in \mathcal{O}_S(X)} B(\ell)}{h}$ $\frac{(X)^{D(x)}}{b}$. The function B is decreasing (See Theorem [2.32\)](#page-8-1) so in particular, each surface X_n has a boundary component of length greater than $\frac{B(\varepsilon_2)}{b}$.

For each X_n , we choose a lift $\chi_n \in \text{Teich}(S_g^b)$ such that γ_0 is such a boundary component. From Theorem [2.26,](#page-7-1) for every $n \in \mathbb{N}$, the surface χ_n admits a pant decomposition \mathcal{P}_n such that any curve in it has length at most

$$
B = \max(A, 2\pi(2g + b - 1)).
$$

There is a finite number of topological types of pants decomposition, so without loss of generality, we can take a subsequence $(\chi_n)_{n\in\mathbb{N}}$ such that for every $n, \mathcal{P}_n = \mathcal{P}$. If there is an essential closed curve σ on S_g^b such that $\ell_{\chi_n}(\sigma) \to 0$ when n goes to infinity,

then σ is homotopic to a curve in \mathcal{P} . Indeed, let us suppose it is not the case. Then, σ intersects a curve α in P . Hence, by the collar Lemma [2.36,](#page-9-1) $\ell_{\chi_n}(\alpha) \to \infty$, contradicting our assumption on P . As a consequence, to find a lower bound on the systole of any $Y \in \mathcal{OM}_{[\varepsilon_1,\varepsilon_2]}(S_g^b)$, we only need to find a lower bound on the lengths of the curves in P , which is independent from *n*.

In the following, we show how to obtain a lower bound on the length of the curves in P and on the length of the boundary components. We construct a rooted graph G as follows. Each vertex corresponds to a pair of pants in χ_n given by the pants decomposition P. The root corresponds to the pair of pants, denoted by P_0 , with γ_0 as one of its boundary component. A pair of vertices is joined by an edge each time the corresponding pairs of pants have a boundary component in common.

Figure 5: Example of a rooted graph G.

We choose a spanning rooted tree T in G . We show by induction on the number of vertices that we have a lower bound on the length of each boundary component of each pair of pants corresponding to the vertices.

Base case P_0 :

One of the boundary components of P_0 is γ_0 and we label the other two by α_0 and β_0 . We consider the simple orthogeodesics τ_{α_0} and τ_0 of P_0 with endpoints on α_0 and γ_0 as in Figure [6.](#page-13-0)

Figure 6: Pair of pants P_0 .

We already have a positive lower bound on the length of γ_0 and since $\ell_{\chi_n}(\alpha_0), \ell_{\chi_n}(\beta_0) \leqslant B$, Lemma [2.40](#page-11-3) gives us an upper bound on $\ell_{\chi_n}(\tau_0)$. Our goal is to find an upper bound for $\ell_{\chi_n}(\tau_{\alpha_0})$ independent from *n* and then use Lemma [2.35](#page-9-0) to find a lower bound for $\ell_{\chi_n}(\alpha_0)$ independent from *n*. We start by cutting P_0 into two symmetric rightangled hexagons and we look at one of them; see Figure [7.](#page-14-0)

Figure 7: Right-angled hexagon obtained by cutting P_0 .

The three altitudes of a right-angled hexagon are concurrent (see [\[Bus10,](#page-29-1) Theorem 2.4.3]) and we call Q the point of intersection of the hexagon altitudes. Then we label its half-altitudes $t_a^1, t_a^2, t_0^1, t_0^2, t'$ and t'' as in Figure [7.](#page-14-0) We note that $t_0^1+t_0^2=\frac{\ell_{\chi_n}(\tau_0)}{2}$ 2 and $t_a^1 + t_a^2 = \frac{\ell_{\chi_n}(\tau_{\alpha_0})}{2}$ $\frac{(\tau_{\alpha_0})}{2}$. Note that since t_a^2 is the distance between Q and the edge of the hexagon intersected by τ_{α_0} (because τ_{α_0} intersects the edge at a right angle), it is shorter than the path following τ_0 from Q to γ_0 then following γ_0 to the edge intersected by τ_{α_0} . In other words, $t_a^2 < t_0^2 + \frac{\ell_{\chi_n}(\gamma_0)}{2} \leq \frac{\ell_{\chi_n}(\tau_0)}{2} + \frac{\ell_{\chi_n}(\gamma_0)}{2}$ $\frac{1}{2}$. To bound $\ell_{\chi_n}(\tau_{\alpha_0})$, we still need to bound t_a^1 . Lemma [2.38](#page-9-4) gives us several relations between the angles φ, ψ, θ and the lengths $t_a^1, t_a^2, t_0^1, t_0^2, t', t''$. Namely,

$$
\cos(\varphi) = \tanh(t_a^2)\tanh(t_0^2) = \tanh(t_0^1)\tanh(t_a^1),\tag{2}
$$

$$
\cos(\psi) = \tanh(t_a^2)\tanh(t'') = \tanh(t_a)\tanh(t'),\tag{3}
$$

$$
\cos(\theta) = \tanh(t_0^2)\tanh(t') = \tanh(t'')\tanh(t_0^1). \tag{4}
$$

Using [\(2\)](#page-14-1), we obtain

$$
\frac{\tanh(t_a^1)}{\tanh(t_a^2)} = \frac{\tanh(t_0^2)}{\tanh(t_0^1)}.
$$

That is,

$$
\frac{\tanh\left(\frac{\ell_{\chi_n}(\tau_{\alpha_0})}{2} - t_a^2\right)}{\tanh(t_a^2)} = \frac{\tanh\left(\frac{\ell_{\chi_n}(\tau_0)}{2} - t_0^1\right)}{\tanh(t_0^1)}
$$

We recognize on the left-hand side the function $g(x) = \frac{\tanh(x - \frac{x}{y})}{\tanh(\frac{x}{y})}$ $\frac{\tanh(x-\frac{x}{J})}{\tanh(\frac{x}{J})}$ with $J=\frac{\ell_{\chi_n}(\tau_{\alpha_0})}{2t_a^2}$ $\frac{a^{(7\alpha_0)}}{2t_a^2}$ and $x = \frac{\ell_{\chi_n}(\tau_{\alpha_0})}{2}.$

If $J \leq 2$, then $t_a^1 \leq t_a^2$ and $\ell_{\chi_n}(\tau_{\alpha_0}) = 2(t_a^1 + t_a^2) \leq 4t_a^2 \leq 2(\ell_{\chi_n}(\tau_0) + \ell_{\chi_n}(\gamma_0)).$ If $J > 2$, we study the function g and see that $\sup g = \lim_{x \to 0} g(x) = J - 1$.

To bound $\ell_{\chi_n}(\tau_{\alpha_0})$, we want an upper bound on $\frac{\tanh\left(\frac{\ell_{\chi_n}(\tau_0)}{2}-t_0^1\right)}{\tanh(t_0)}$ $\frac{(1 - 2)^{-1}}{\tanh(t_0)}$, that is, a lower bound on t_0^1 . Suppose by contradiction that t_0^1 converges to 0. Since $|\tanh(x)| < 1$ for all x , we derive from the relations (2) and (4) that

$$
\tanh(t_0^1)\tanh(t_a^1) = \tanh(t_a^2)\tanh(t_0^2) \to 0\tag{5}
$$

$$
\tanh(t'')\tanh(t_0^1) = \tanh(t_0^2)\tanh(t') \to 0\tag{6}
$$

We cannot have $t_0^2 \to 0$ because $2(t_0^2 + t_0^1) = \ell_{\chi_n}(\tau_0) \geq \min(\mathcal{O}_S(X)) > 0$. If instead, we have $t_a^2 \to 0, t' \to 0$, then by Lemma [2.35,](#page-9-0) the lengths of the sides of the hexagon with extremities on γ_0 go to infinity. Thus, $t_0^2 \to \infty$ which is not possible because we have an upper bound on $\ell_{\chi_n}(\tau_0) = 2(t_0^1 + t_0^2)$. Hence a contradiction.

Therefore, there exists $\varepsilon'_0 > 0$ such that $t_0^1 > \varepsilon'_0$. Hence,

$$
\frac{\tanh(\frac{\ell_{\chi_n}(\tau_0)}{2}-\varepsilon_0')}{\tanh(\varepsilon_0')}+1\geqslant \sup\limits_{\frac{\ell_{\chi_n}(\tau_{\alpha_0})}{2}}\left(\frac{\tanh(\frac{\ell_{\chi_n}(\tau_{\alpha_0})}{2}-t_a^2)}{\tanh(t_a^2)}\right)+1=J
$$

and we have an upper bound on $2Jt_a^2 \geq \ell_{\chi_n}(\tau_{\alpha_0})$. Since both endpoints of τ_{α_0} lie in α_0 , we have $\ell_{\chi_n}(\tau_{\alpha_0}) \geq 2d$, where d is the width of the half-collar of α_0 . By Proposi-tion [2.35,](#page-9-0) if $\ell_{\chi_n}(\alpha_0) \to 0$, the length of τ_{α_0} goes to infinity, which contradicts the fact that $\ell_{\chi_n}(\tau_{\alpha_0})$ is bounded. By symmetry, we can also show that $\ell_{\chi_n}(\beta_0)$ is bounded away from zero. We denote by ε_0 the minimum between the positive lower bound on $\ell_{\chi_n}(\alpha_0)$, $\ell_{\chi_n}(\beta_0)$ and $\ell_{\chi_n}(\gamma_0)$. Observe that ε_0 only depends on the simple orthospectrum and the topology of X .

Induction step:

Now let us choose another vertex of T . We have a unique path in T going from the root to our vertex. Let us label the pair of pants corresponding to the vertices on the path by $P_0, ..., P_l, P_{l+1}$ with P_0 the pair of pants corresponding to the root.

Figure 8

Suppose we already have a lower bound on the length of the boundary components of the pair of pants from P_0 to P_l . We show that there is a lower bound on the length of the boundary components α_{l+1} and β_{l+1} of P_{l+1} . We call χ'_n the sub-surface of χ_n composed of the pairs of pants P_0, \ldots, P_{l+1} as in Figure [9.](#page-16-0)

Figure 9: The sub-surface χ_n .

We define P'_l to be the pair of pants embedded in χ'_n with γ_0 and α_l as boundary components. Let a be a shortest path between γ_0 and α_l . The third boundary component of P'_l is homotopic to the piecewise geodesic obtained by following a, going around α_l , following a in the other direction and then going around γ_0 . In the same way, we define P'_{l+1} with γ_0 and α_{l+1} as boundary components. Then, we let τ_l be the unique simple orthogeodesic of P'_l with both endpoints on γ_0 , and τ_{l+1} be the unique simple orthogeodesic of P'_{l+1} with both endpoints on γ_0 . These two orthogeodesics are also simple orthogeodesics of X_n . We let τ_{α_l} and $\tau_{\alpha_{l+1}}$ be the unique simple orthogeodesics of P'_l and P'_{l+1} with both endpoints on α_l and α_{l+1} . Finally, we let τ' be the unique simple orthogeodesic of P_{l+1} with both endpoints on α_l . See Figure [11](#page-17-0) and [10.](#page-17-1)

Figure 10: The pair of pants P'_l and its orthogeodesic embedded in χ'_n .

Figure 11: The pair of pants P'_{l+1} and its orthogeodesic embedded in χ'_n .

By induction, we have an upper bound on $\ell_{\chi_n}(\tau_l)$ and $\ell_{\chi_n}(\tau_{\alpha_l})$, and Lemma [2.40](#page-11-3) gives us an upper bound on $\ell_{\chi_n}(\tau')$. We construct a piecewise geodesic σ homotopic to τ_{l+1} as follows: we follow τ_l from one of its endpoints on γ_0 until we meet τ_{α_l} , then we follow τ_{α_l} along the shortest path toward α_l , we follow α_l until we meet an endpoint of τ' , we follow τ' , then α_l again until we meet the other endpoint of τ_{α_l} , we follow τ_{α_l} until we meet τ_l , and finally, we follow τ_l until we meet γ_0 again (see Figure [12\)](#page-18-0).

Figure 12: The piecewise geodesic σ on χ'_n .

The length of σ yields an upper bound on $\ell(\tau_{l+1})$:

$$
\ell_{\chi_n}(\tau_{l+1}) \leqslant \ell_{\chi_n}(\sigma) \leqslant \ell_{\chi_n}(\tau_l) + \ell_{\chi_n}(\tau_{\alpha_l}) + 2\ell_{\chi_n}(\alpha_l) + \ell_{\chi_n}(\tau').
$$

Now that we have an upper bound on $\ell_{\chi_n}(\tau_{l+1})$, we can apply to P'_{l+1} what we did in the base case P_0 . As a result, we obtain an upper bound on $\ell_{\chi_n}(\tau_{\alpha_{l+1}})$ and a positive lower bound on $\ell_{\chi_n}(\alpha_{l+1})$. By symmetry, we also have a lower bound on $\ell_{\chi_n}(\beta_{l+1})$. We observe that the lower bounds on $\ell_{\chi_n}(\alpha_{l+1})$ and $\ell_{\chi_n}(\beta_{l+1})$ depends only on the simple orthospectrum and the topology of X.

Let $\varepsilon = \varepsilon(\varepsilon_1, \varepsilon_2, b, g)$ be the minimum between the positive lower bounds on the length of the boundary component of the vertices of T. Then, ε is a positive lower bound independent from n on the systole and on the length of every boundary component of χ_n . By Corollary [3.3,](#page-12-0) we obtain that $\mathcal{OM}_{[\epsilon_1,\epsilon_2]}(S_g^b) \subset \mathcal{M}_{A,\epsilon}(S_g^b)$ which is compact by Theorem [2.27.](#page-7-0) \Box

If two surfaces share the same simple orthospectrum, then they share the same orthosystole and we can deduce from Theorem [3.4](#page-12-1) the following corollary:

Corollary 3.5. The set $I(X)$ lies in a compact set of $\mathcal{M}(S_g^b)$. In particular, there exist $A \geqslant \varepsilon > 0$ such that

$$
I(X) \subset \mathcal{M}_{A,\varepsilon}(S_g^b).
$$

If they share the same orthospectrum, they also share the same orthosystole thus we have the same result for the orthospectrum.

3.3 Step three: Discreteness

We can now prove the main theorem of this section.

Proof of Theorem [3.1.](#page-11-0) First, let us fix a hexagon decomposition \mathcal{H} on S_g^b and set $\varphi_{\mathcal{H}}$ as in Theorem [2.16.](#page-4-0)

Because I(X) is included in a compact subset of $\mathcal{M}(S_g^b)$ (by Corollary [3.5\)](#page-18-1), there is a lift $\tilde{I}(X)$ of $I(X)$ in Teich (S_g^b) which is also included in a compact C. We will show that $\tilde{I}(X)$ is finite, and so is $I(X)$.

Since $\tilde{I}(X)$ is included in a compact, for any sequence $\chi_n \in \tilde{I}(X)$, there is a subsequence such that $\chi_n \to \chi \in C$. The map $\varphi_{\mathcal{H}}$ is continuous, so $\varphi_{\mathcal{H}}(\chi_n) \to \varphi_{\mathcal{H}}(\chi)$. We recall that for every n, we have $\mathcal{O}_S(\chi_n) = \mathcal{O}_S(X)$ and that by Theorem [2.30](#page-8-0) the orthospectrum is discrete.

As for all $n, \varphi_{\mathcal{H}}(\chi_n) \subset \mathcal{O}_S(X)$, and applying Wolpert's lemma [2.21,](#page-6-0) we deduce that, there exists $N \in \mathbb{N}^*$ such that for all $n', m' > N$, $\varphi_{\mathcal{H}}(\chi_{n'}) = \varphi_{\mathcal{H}}(\chi_{m'})$. Thus, for all $n', m' > N$, $\chi_{n'} = \chi_{m'} = \chi$ and $\chi \in \tilde{I}(X)$. So $\tilde{I}(X)$ is compact and discrete, thus finite, and so is $I(X)$. \Box

4 Generic characterization

In this section, we are going further in our characterization of the rigidity of the orthospectrum and the simple orthospectrum. We prove the following theorem:

Theorem 4.1. Let \mathcal{V}_{g}^{b} be the subset of all $\chi \in \text{Teich}(S_{g}^{b})$ for which there exists $\Upsilon \in \text{Teich}(S_g^b)$ non-isometric to χ such that $\mathcal{O}(\chi) = \mathcal{O}(\Upsilon)$. Similarly, let \mathcal{W}_g^b be the subset of all $\chi \in \text{Teich}(S_g^b)$ for which there exists $\Upsilon \in \text{Teich}(S_g^b)$ non-isometric to χ such that $\mathcal{O}_S(\chi) = \mathcal{O}_S(\Upsilon)$.

Then, \mathcal{V}_g^b and \mathcal{W}_g^b are proper local real analytic subvarieties of Teich(S_g^b). In particular, they are negligible set.

This result is a version of Wolpert's Theorem [\[Wol79\]](#page-30-3) for the orthospectrum instead of the length spectrum. We adapt the proof given by Buser in [\[Bus10,](#page-29-1) Chap. 10] to our case. Doing so, we no longer require non-simple curves, making the theorem true both for the orthospectrum and the simple orthospectrum. As in Buser's proof, we also show an intermediate theorem before proving the main theorem.

Let $\mathcal{T}_{g}^{b}([\varepsilon_{1}, \varepsilon_{2}]) \subset \text{Teich}(S_{g}^{b})$ denote the set of all compact hyperbolic surfaces of genus g, with b boundary component and orthosystole between ε_1 and ε_2 in Teich (S_g^b) .

Theorem 4.2. Fix $\varepsilon_1, \varepsilon_2 > 0$ with $\varepsilon_1 \leq \varepsilon_2$. Then, there exists a real number $t = t(g, b, \varepsilon_1, \varepsilon_2)$ such that for $\chi, \Upsilon \in \mathcal{T}_{g}^b([\varepsilon_1, \varepsilon_2])$, we have $\mathcal{O}(\chi) = \mathcal{O}(\Upsilon)$ if and only if

$$
\mathcal{O}(\chi) \cap [0, t] = \mathcal{O}(\Upsilon) \cap [0, t]
$$

and $\mathcal{O}_S(\chi) = \mathcal{O}_S(\Upsilon)$ if and only if

$$
\mathcal{O}_S(\chi) \cap [0,t] = \mathcal{O}_S(\Upsilon) \cap [0,t].
$$

We first set up all the context and notation, then we prove Theorem [4.2](#page-19-2) and finally Theorem [4.1.](#page-19-0)

4.1 Set up and prerequisites

Fix a hexagon decomposition $\mathcal{H} = \{\alpha_1, ..., \alpha_{6g+3b-6}\}\$ on S_g^b . Let $\chi_0 \in \text{Teich}(S_g^b)$ be given by $(\ell_{\chi_0}(\alpha_1), ..., \ell_{\chi_0}(\alpha_{6g+3b-6})) = (1, ..., 1)$. This surface is going to be a point of

reference. Recall that; see Definition [2.17,](#page-5-0) for any $\omega = (\ell_1, ..., \ell_{6g+3b-6}) \in \mathbb{R}^{6g+3b-6}_+$, we set $\chi^{\omega} := \varphi_{\mathcal{H}}^{-1}(\omega) \in \text{Teich}(S_g^b)$. Then we choose a quasi-conformal homeomorphism ϕ^{ω} between χ_0 and χ^{ω} . For each orthogeodesic τ on χ_0 , we denote by $\tau(\chi^{\omega})$ the unique orthogeodesic in the free homotopy class of the orthogeodesic $\phi^{\omega} \circ (\tau)$ on χ^{ω} . For any finite or infinite ordered set Λ of orthogeodesics τ_1, τ_2, \ldots on χ_0 , we define the sequences

$$
\Lambda(\chi) = (\tau_1(\chi), \tau_2(\chi), \ldots)
$$

$$
\ell_{\chi}(\Lambda) = (\ell_{\chi}(\tau_1), \ell_{\chi}(\tau_2), \ldots).
$$

We set $\Pi = (\beta_1, \beta_2, \dots)$ to be the sequence of all orthogeodesics on χ_0 , arranged so that $\ell_{\chi_0}(\beta_1) \leq \ell_{\chi_0}(\beta_2) \leq \dots$ and set $\Pi_k = (\beta_1, \dots, \beta_k)$. Then, set $\Pi'_k = (\beta'_1, \beta'_2, \dots)$ the sequence of all simple orthogeodesics on χ_0 , arranged so that $\ell_{\chi_0}(\beta_1') \leq \ell_{\chi_0}(\beta_2') \leq \ldots$ and set $\Pi'_k = (\beta'_1, ..., \beta'_k)$. Note that $\ell_{\chi_0}(\Pi) = \mathcal{O}(\chi_0)$ and $\ell_{\chi_0}(\Pi') = \mathcal{O}_S(\chi_0)$. Finally, let $\delta_1, ..., \delta_{6g+3b-6}$ be the unique collection of simple orthogeodesics on χ_0 such that $i(\alpha_i, \delta_j) = \delta_{ij}$ for all $1 \leq i, j \leq 6g + 3b - 6$, and set $\Sigma = \mathcal{H} \cup \{\delta_1, ..., \delta_{6g + 3b - 6}\}.$

We fix $\varepsilon_2 \geqslant \varepsilon_1 > 0$ and we choose $A \geqslant \varepsilon > 0$ as in Theorem [3.4.](#page-12-1) Let $\mathcal{Q}(A, \varepsilon)$ be as in Corollary [2.28.](#page-7-2) We choose an open neighborhood $U \subset \text{Teich}(S_g^b)$ with compact closure which contains $\mathcal{Q}(A,\varepsilon)$. By Corollary [2.22,](#page-6-1) there exist $\varepsilon_2' \geqslant \varepsilon_1' > 0$ such that the orthosystole of any surface $\chi \in U$ lies between ε_1' and ε_2' . By Theorem [3.4](#page-12-1) and Corollary [2.28,](#page-7-2) there exist $A_U \geq \varepsilon_U > 0$ such that there is a compact subset $\mathcal{Q}(A_U, \varepsilon_U)$ (as in Corollary [2.28\)](#page-7-2) of Teich (S_g^b) containing U. By definition, if $\mathcal{O}(\chi) = \mathcal{O}(\chi')$ (or if $\mathcal{O}_S(\chi) = \mathcal{O}_S(\chi')$ for $\chi \in U$ and $\chi' \in \text{Teich}(S_g^b)$, then χ and χ' have the same orthosystole which lies between ε_1 and ε_2 , and therefore both have a systole greater than ε_U and boundary length between ε_U and A_U . Thus, there exists a surface isometric to χ' in $\mathcal{Q}(A_U, \varepsilon_U)$ and we may assume without loss of generality that $\chi' \in \mathcal{Q}(A_U, \varepsilon_U)$.

Let $D = \{ \chi^{\omega} \in \text{Teich}(S_g^b) \mid \frac{1}{10} \leq \ell_{\chi^{\omega}}(\alpha_1), \dots, \ell_{\chi^{\omega}}(\alpha_{6g+3b-6}) \leq 10 \}$ and let C, C_1 be compact sets whose interiors $\check{C}, \check{C}_1 \subset \text{Teich}(S_g^b)$ are connected and

$$
(U\cup\mathcal{Q}(A_U,\varepsilon_U)\cup D)\subset\mathring{C}\subset C\subset\mathring{C}_1.
$$

By Corollary [2.22,](#page-6-1) there exists $q \geq 1$ such that

$$
\frac{\ell_{\chi}(\beta)}{q} \leq \ell_{\chi'}(\beta) \leq q\ell_{\chi}(\beta)
$$
\n(7)

for any $\chi, \chi' \in C_1$ and any $\beta \in \Pi$. This q will remain fixed during the proof.

Lemma 4.3. For any $k \in \mathbb{N}$, there exists an integer $k^* \geq k$, which depends only on k and C_1 , with the following property. If $\chi', \chi'' \in C_1$ and if $\rho : \Pi_k \to \Pi$ is an injection such that

$$
\ell_{\chi'}(\Pi_k) = \ell_{\chi''}(\rho(\Pi_k))
$$

then $\rho(\Pi_k) \subset \Pi_{k^*}$. The same is true with Π', Π'_k and Π'_{k^*} instead of Π, Π_k and Π_{k^*} .

Proof. Let $c_1 = \max{\{\ell_\chi(\beta) \mid \chi \in C_1, \beta \in \Pi_k\}}$, and let $k^* \geq k$ be such that, on the base surface χ_0 , we have $\ell_{\chi_0}(\beta_j) > qc_1$ for all $j > k^*$. By [\(7\)](#page-20-0) and since χ_0 is in $D \subset C_1$, we have $\ell_{\chi}(\beta_j) \geqslant \frac{\ell_{\chi_0}(\beta_j)}{q}$ $\frac{(\beta_j)}{q} > c_1$ for all $j > k^*$ and $\chi \in C_1$. Since we have $\ell_{\chi''}(\rho(\beta_i)) = \ell_{\chi'}(\beta_i) \leqslant c_1$ for $i \leqslant k$ by definition of c_1 , it follows that $\rho(\beta_i) \in \Pi_{k^*}$. For the simple orthospectrum, just replace β_i with β'_i and Π , Π_k and Π_{k^*} with Π' , Π'_k and $\Pi'_{k^*}.$ \mathbb{R}^n

4.2 The first lengths of the orthospectrum

Let us proceed to the proof of Theorem [4.2.](#page-19-2)

Proof of Theorem [4.2.](#page-19-2) We define for each $k \in \mathbb{N}$ the following sets:

$$
V_k^1 = \{ (\chi, \chi') \in \mathring{C}_1 \times \mathring{C}_1 \mid \ell_\chi(\Pi_k) \subset \mathcal{O}(\chi') \text{ and } \ell_{\chi'}(\Pi_k) \subset \mathcal{O}(\chi) \}
$$

\n
$$
V_k = V_k^1 \cap (\mathring{C} \times \mathring{C})
$$

\n
$$
W_k^1 = \{ (\chi, \chi') \in \mathring{C}_1 \times \mathring{C}_1 \mid \ell_\chi(\Pi'_k) \subset \mathcal{O}_S(\chi') \text{ and } \ell_{\chi'}(\Pi'_k) \subset \mathcal{O}_S(\chi) \}
$$

\n
$$
W_k = W_k^1 \cap (\mathring{C} \times \mathring{C}).
$$

Let $k^*, k'^* \in \mathbb{N}$ be as in Lemma [4.3.](#page-20-1) Given any two pair of injections $\rho_1, \rho_2 : \Pi_k \to \Pi_{k^*}$ and $\rho'_1, \rho'_2 : \Pi'_k \to \Pi'_{k'^*},$ we set

$$
V[\rho_1, \rho_2] = \{ (\chi, \chi') \in \mathring{C}_1 \times \mathring{C}_1 \mid \ell_{\chi}(\Pi_k) = \ell_{\chi'}(\rho_1(\Pi_k)) \text{ and } \ell_{\chi'}(\Pi_k) = \ell_{\chi}(\rho_2(\Pi_k)) \}
$$

$$
W[\rho'_1, \rho'_2] = \{ (\chi, \chi') \in \mathring{C}_1 \times \mathring{C}_1 \mid \ell_{\chi}(\Pi'_k) = \ell_{\chi'}(\rho'_1(\Pi'_k)) \text{ and } \ell_{\chi'}(\Pi'_k) = \ell_{\chi}(\rho'_2(\Pi'_k)) \}.
$$

By Lemma [2.18,](#page-5-1) the spaces $V[\rho_1, \rho_2]$ and $W[\rho'_1, \rho'_2]$ are real analytic subvarieties of $C_1 \times C_1$. Then, thanks to Lemma [4.3,](#page-20-1) we have

$$
V_k^1 = \bigcup_{(\rho_1, \rho_2)} V[\rho_1, \rho_2]
$$

$$
W_k^1 = \bigcup_{(\rho'_1, \rho'_2)} W[\rho'_1, \rho'_2].
$$

Since there are finitely many pairs (ρ_1, ρ_2) and (ρ'_1, ρ'_2) , the unions V_k^1 and W_k^1 are also real analytic.

Next, we need the following result:

Lemma 4.4. There exists $K \in \mathbb{N}$ such that $V_{K+j} = V_K$ and $W_{K+j} = W_K$ for all $j \ge 1$.

Proof. Teichmüller space is a real analytic space and C_1 is a compact subset of Teich (S_g^b) . Thus, by [\[Fri67,](#page-29-9) Théorème I.9], the ring of real analytic functions on $C_1 \times C_1$ is Noetherian. Moreover, any real analytic subvariety V is associated with the ideal $I(V)$ of real analytic functions vanishing on the subvariety. For two real analytic subvarieties V_1, V_2 , the inclusion $V_1 \subset V_2$ is equivalent to $I(V_2) \subset I(V_1)$ (see [\[Bru52\]](#page-29-10) for more details). By definition, increasing sequences of ideals of a Noetherian ring are stationary, so decreasing sequences of subvarieties are stationary. Therefore, there is $K \in \mathbb{N}$ such that $V_{K+j} = V_K$ and $W_{K+j} = W_K$ for all $j \geq 1$. \Box

Finally, we define

$$
t(A, \varepsilon) = \max\{\ell_{\chi}(\beta) \mid \chi \in \mathcal{Q}(A, \varepsilon), \beta \in \Pi_K\}
$$

$$
t'(A, \varepsilon) = \max\{\ell_{\chi}(\beta) \mid \chi \in \mathcal{Q}(A, \varepsilon), \beta \in \Pi_K'\}.
$$

If the orthosystole of χ and χ' is between ε_1 and ε_2 , then without loss of generality $\chi, \chi' \in \mathcal{Q}(A, \varepsilon)$. If in addition $\mathcal{O}(\chi) \cap [0, t(A, \varepsilon)] = \mathcal{O}(\chi') \cap [0, t(A, \varepsilon)]$, then $\ell_{\chi}(\Pi_K) \subset \mathcal{O}(\chi')$ and $\ell_{\chi'}(\Pi_K) \subset \mathcal{O}(\chi)$. Thus, $(\chi, \chi') \in V_K = V_{K+j}$ for all $j \geq 1$. In conclusion, $\mathcal{O}(\chi) = \mathcal{O}(\chi')$.

The same argument shows that if $\mathcal{O}_S(\chi) \cap [0, t'(A, \varepsilon)] = \mathcal{O}_S(\chi') \cap [0, t'(A, \varepsilon)]$ then $\mathcal{O}_S(\chi) = \mathcal{O}_S(\chi').$ \Box

4.3 Generic surfaces are determined by their (simple) orthospectrum

We define $U \subset \text{Teich}(S_g^b), D, C$ and C_1 as in Section [4.1.](#page-19-3) To avoid repetition, we prove Theorem [4.1](#page-19-0) only for the orthospectrum. For the simple orthospectrum, the proof is the same with Π′ instead of Π. Note that some verifications we perform about the simplicity of curves are not necessary when proving the theorem for the simple orthospectrum.

Proof of Theorem [4.1.](#page-19-0) We fix K as in Lemma [4.4](#page-21-0) and large enough so that $\Sigma \subset \Pi_K$. Then, with the notations of Lemma [4.3,](#page-20-1) we fix $M = K^*$ and $N = M^*$. Now, let $\rho: \Pi_M \to \Pi_N$ be any injection such that $\Pi_K \subset \rho(\Pi_M)$. We define

$$
V_{\rho} = \{ \chi \in \mathring{C} \mid \text{ there exists } \chi^{\rho} \in \text{Teich}(S_g^b) \text{ such that } \ell_{\chi}(\Pi_M) = \ell_{\chi^{\rho}}(\rho(\Pi_M)) \}.
$$

For $\chi \in V_\rho$, the surface χ^ρ is unique and $\mathcal{O}(\chi) = \mathcal{O}(\chi^\rho)$: indeed, since $\mathcal{H} \subset \Sigma \subset \Pi_K \subset \rho(\Pi_M)$, the vector $\ell_{\chi^{\rho}}(\mathcal{H})=\ell_{\chi}(\rho^{-1}(\mathcal{H}))$ represents Ushijima's coordinates of χ^{ρ} . Then, we have $\ell_{\chi}(\Pi_K) \subset \ell_{\chi}(\rho(\Pi_M)) = \ell_{\chi}(\Pi_M) \subset \mathcal{O}(\chi)$. Conversely, since $\Pi_K \subset \Pi_M$, we also have $\ell_\chi(\Pi_K) \subset \ell_\chi(\Pi_M) = \ell_{\chi^{\rho}}(\rho(\Pi_M)) \subset \mathcal{O}(\chi^{\rho})$ so $\mathcal{O}(\chi) = \mathcal{O}(\chi^{\rho})$ by Lemma [4.4.](#page-21-0)

Now, we define the real analytic mapping m_{ρ} : Teich $(S_g^b) \to$ Teich (S_g^b) given by $m_{\rho}(\chi^{\omega}) = \chi^{\ell_{\chi}\omega(\rho^{-1}(\mathcal{H}))}$. If χ^{ρ} exists, we have $m_{\rho}(\chi) = \chi^{\rho}$. Thus,

$$
V_{\rho} = \{ \chi \in \mathring{C} \mid \ell_{\chi}(\Pi_M) = \ell_{m_{\rho}(\chi)}(\rho(\Pi_M)) \},
$$

is a real analytic subvariety of \check{C} . We define

$$
V_{\rho}^* = \{ \chi \in V_{\rho} \mid \chi^{\rho} \text{ is isometric to } \chi \}.
$$

A surface $\chi \in V_\rho$ is isometric to $\chi^\rho = m_\rho(\chi)$ if and only if χ and χ^ρ are in the same $MCG(S_g^b)$ orbit, that is, if and only if there is an injection $r: \mathcal{H} \to \Pi$, induced by a homeomorphism of the base surface χ_0 , such that

$$
\ell_{\chi}(\mathcal{H}) = \ell_{m_{\rho}(\chi)}(r(\mathcal{H})).
$$

By Lemma [4.3,](#page-20-1) we have $r(\mathcal{H}) \subset \Pi_M$. This shows that the set R^* of all such possible injections r is finite. This implies that

$$
V_{\rho}^* = \bigcup_{r \in R^*} \{ \chi \in V_{\rho} \mid \ell_{\chi}(\mathcal{H}) = \ell_{m_{\rho}(\chi)}(r(\mathcal{H})) \}
$$

is a real analytic subvariety of V_o .

Let R be the set of injective maps $\rho : \Pi_M \to \Pi_N$ satisfying $\Pi_K \subset \rho(\Pi_M)$. Note that this set is finite. Define

$$
V = \bigcup_{\rho \in \mathcal{R}} (V_{\rho} \setminus V_{\rho}^*).
$$

The sets V_{ρ}, V_{ρ}^* are real analytic subvarieties so V is a real analytic subvariety of \dot{C} . By construction, we have $V \cap \overset{\circ}{C} \subset \mathcal{V}_g^b \cap \overset{\circ}{C}$, hence $V \cap U \subset \mathcal{V}_g^b \cap U$. Conversely, if $\chi \in \mathcal{V}_g^b$ then there exists χ' not isometric to χ such that $\mathcal{O}(\chi) = \mathcal{O}(\chi')$. This implies that $\chi, \chi' \in \mathcal{Q}(A_U, \varepsilon_U)$. Furthermore, if $\chi, \chi' \in \mathcal{Q}(A_U, \varepsilon_U)$ have the same orthospectrum, then there exists a bijection $\rho : \Pi \to \Pi$ satisfying $\ell_{\chi}(\Pi) = \ell_{\chi'}(\rho(\Pi))$. By Lemma [4.3,](#page-20-1) we have $\rho^{-1}(\Pi_K) \subset \Pi_M$ and $\rho(\Pi_M) \subset \Pi_N$. In other words, there exists an injection $\rho : \Pi_M \to \Pi_N$ such that

$$
\Pi_K \subset \rho(\Pi_M)
$$

$$
\ell_\chi(\Pi_M) = \ell_{\chi'}(\rho(\Pi_M)).
$$

By definition of V, we have $\chi \in V \cap U$. In conclusion, $V \cap U = V_g^b \cap U$ for any neighborhood U. We still need to show that $\mathcal{V}_g^b \cap U = V \cap U \neq U$, *i.e.*, that $\dim V < \dim \text{Teich}(S_g^b)$. Since V_ρ is a real analytic subvariety of $\overset{\circ}{C}$ and $\overset{\circ}{C}$ is connected by definition, we either have $V_{\rho} = \dot{C}$ or else dim $V_{\rho} < \dim \dot{C} = \dim \text{Teich}(S_g^b)$. We want to show that if $V_{\rho} = \dot{C}$ then $V_{\rho} = V_{\rho}^*$. If this is true, for all $\rho \in \mathcal{R}$ $\dim(V_\rho \setminus V_\rho^*) < \dim \mathrm{Teich}(S_g^b)$, thus $\dim V < \dim \mathrm{Teich}(S_g^b)$.

So suppose $V_\rho = \mathring{C}$, then there is a map $m_\rho : \mathring{C} \to \mathrm{Teich}(S_g^b)$ such that $\ell_{m_\rho(\chi)}(\rho(\Sigma)) =$ $\ell_{\mathcal{X}}(\Sigma)$ for all $\chi \in \mathring{C}$. In the following, we abbreviate $\tilde{\beta} := \rho(\beta)$ for any $\beta \in \Sigma$, and $\tilde{\chi} := m(\chi)$ for any $\chi \in \tilde{C}$.

Step 1: $\rho(\mathcal{H})$ is a hexagon decomposition.

Set $\omega = (\frac{1}{4}, ..., \frac{1}{4})$ $\frac{1}{4}$). Then $\ell_{\tilde{\chi}^{\omega}}(\tilde{\alpha}_i) = \frac{1}{4}$ for $1 \leqslant i \leqslant 6g + 3b - 6$. By Theorem [2.33,](#page-8-2) we deduce that the orthogeodesics $\tilde{\alpha}_i$ are simple. The fact that they are pairwise disjoint fol-lows from Lemma [2.37.](#page-9-5) If $\tilde{\alpha}_i$ and $\tilde{\alpha}_j$ intersect each other, then $\sinh(\ell_{\tilde{\chi}^{\omega}}(\tilde{\alpha}_i))\sinh(\ell_{\tilde{\chi}^{\omega}}(\tilde{\alpha}_j)) > 1$. This is impossible since $\sinh(\frac{1}{4}) \sinh(\frac{1}{4}) < 1$. Therefore, ρ sends H to another hexagon decomposition $\mathcal H$ of χ_0 .

Step 2: Understand the relative position of the $\tilde{\alpha}_i$.

We want to show that if $\alpha_{i_1}, \alpha_{i_2}, \alpha_{i_3}, \alpha_{i_4}$ are orthogeodesics delimiting an octagon of orthogonals α_j and δ_j on χ_0 , then $\tilde{\alpha}_{i_1}, \tilde{\alpha}_{i_2}, \tilde{\alpha}_{i_3}, \tilde{\alpha}_{i_4}$ are also orthogeodesics delimiting an octagon of orthogonals $\tilde{\alpha}_j$ and $\tilde{\delta}_j$. Indeed, fix χ and $1 \leqslant j \leqslant 6g + 3b - 6$ such that $\ell_{\chi}(\alpha_i) = \frac{1}{4}$ and $\ell_{\chi}(\alpha_j) = 6$ for all $1 \leq i \neq j \leq 6g + 3b - 6$. By Lemma [2.39,](#page-10-1) we have

$$
\cosh(\ell_{\chi}(\delta_j)) = f_{\text{octa}}(\ell_{\chi}(\alpha_j), \ell_{\chi}(\alpha_{i_1}), \ell_{\chi}(\alpha_{i_2}), \ell_{\chi}(\alpha_{i_3}), \ell_{\chi}(\alpha_{i_4}))
$$

and $\ell_{\tilde{\chi}}(\tilde{\delta}_j) = \ell_{\chi}(\delta_j) < \frac{1}{2}$ $\frac{1}{2}$. As before, by Theorem [2.33](#page-8-2) and Lemma [2.37,](#page-9-5) the curve $\tilde{\delta}_j$ is simple and disjoint from $\tilde{\alpha}_i$ for any $1 \leq i \neq j \leq 6g+3b-6$. It follows that $i(\tilde{\delta}_j, \tilde{\alpha}_j) = 1$ and the arcs $\tilde{\delta}_j$, $\tilde{\alpha}_j$ lie in an octagon delimited by four orthogeodesics among the $\tilde{\alpha}_i$ for $i \neq j$. Since $\cosh(\ell_{\tilde{\chi}}(\tilde{\delta}_j)) = f_{\text{octa}}(\ell_{\tilde{\chi}}(\tilde{\alpha}_j), \ell_{\tilde{\chi}}(\tilde{\alpha}_{i_1}), \ell_{\tilde{\chi}}(\tilde{\alpha}_{i_2}), \ell_{\tilde{\chi}}(\tilde{\alpha}_{i_3}), \ell_{\tilde{\chi}}(\tilde{\alpha}_{i_4}))$ we see that if we vary the length of one $\tilde{\alpha}_l \in \tilde{\mathcal{H}}$ and fix the other ones, the length of $\tilde{\delta}_i$ only depends on the lengths of $\tilde{\alpha}_{i_1}, \tilde{\alpha}_{i_2}, \tilde{\alpha}_{i_3}, \tilde{\alpha}_{i_4}$ and $\tilde{\alpha}_j$. This means that the orthogeodesics delimiting the octagon containing $\tilde{\delta}_j$ and $\tilde{\alpha}_j$ are $\tilde{\alpha}_{i_1}, \tilde{\alpha}_{i_2}, \tilde{\alpha}_{i_3}, \tilde{\alpha}_{i_4}$.

Moreover, the non-symmetry of f_{octa} also gives an indication about how the orthogeodesics delimiting the octagon are placed. Indeed, if α , δ_1 , δ_2 , δ_3 and δ_4 are as in Figure [3](#page-10-0) then we obtain different values of β when we exchange δ_1 and δ_2 or when we exchange δ_1 and δ_4 . Thus, if we choose χ such that $\alpha_{i_1}, \alpha_{i_2}, \alpha_{i_3}, \alpha_{i_4}, \alpha_j$ and δ_j all have different lengths. Then, we know that if $(\alpha_{i_1}, \alpha_{i_2}, \alpha_j)$ delimits a hexagon, then $(\tilde{\alpha}_{i_1}, \tilde{\alpha}_{i_2}, \tilde{\alpha}_j)$ also delimits an hexagon. Thus, the two surfaces are isometric because they are obtained by gluing isometric hexagons with the same pattern. In conclusion, if $V_{\rho} = \mathring{C}$, then $V_{\rho} = V_{\rho}^*$ and we have $V_{\rho} \setminus V_{\rho}^* = \emptyset$. \Box

5 Rigidity results

In [\[MM23\]](#page-30-4), Masai and McShane prove orthospectrum rigidity for the one-holed torus. In the case of the simple orthospectrum, the same proof does not work as it relies on computing the length of the boundary using Basmajian's identity. However in this section, we prove simple orthospectrum rigidity for the one-holed torus with a different proof, which relies on Ushijima's coordinates instead of Fenchel-Nielsen coordinates.

The first result we need to prove rigidity is the following:

Proposition 5.1. Let X be a compact hyperbolic surface with geodesic boundary. Then the first two lengths in $\mathcal{O}_S(X)$ are the lengths of two disjoint orthogeodesics.

Proof. Let τ_1 and τ_2 be two orthogeodesics realizing the first two lengths of the simple orthospectrum with $\ell(\tau_1) \leq \ell(\tau_2)$. Let us suppose that $i(\tau_1, \tau_2) > 0$. The idea is to get a contradiction by constructing a new orthogeodesic τ shorter than τ_2 .

We construct a piecewise geodesic path σ as follows. Start at an endpoint of τ_1 such that the length between this endpoint and the first intersection point p between τ_1 and τ_2 is less than $\frac{\ell(\tau_1)}{2}$. Then follow τ_1 until p, and finally follow τ_2 until its closest endpoint. We obtain $\ell(\sigma) \leq \frac{\ell(\tau_2)}{2} + \frac{\ell(\tau_1)}{2} \leq \ell(\tau_2)$. Note that σ is essential, otherwise, together with an arc of the boundary of X , we get a hyperbolic triangle with two right angles, which is impossible.

Figure 13: Construction of σ .

Note that the orthogeodesic τ homotopic to σ is simple. Since the two arcs forming σ meet at some angle different from π , the geodesic representative τ is strictly shorter than σ and $\ell(\tau) < \ell(\sigma) \leq \ell(\tau_2)$. Moreover, by construction $i(\tau, \tau_1) < i(\tau_1, \tau_2)$. Suppose then that $\tau = \tau_1$. In this case, the segment of τ_2 that σ follows and the segment of τ_1 that σ does not follow are homotopic and form with a segment of ∂X a hyperbolic triangle with two right angles. This is impossible, so $\tau \neq \tau_1$.

Thus, the simple orthogeodesic τ is different from τ_1 and shorter than τ_2 , which is a contradiction. \Box With this result at hand, we can prove the desired rigidity statement.

Theorem 5.2. Let T and T' be two hyperbolic structures with geodesic boundary on the one-holed torus. Then T and T' are isometric if and only if $\mathcal{O}_S(T) = \mathcal{O}_S(T')$.

Proof. A hexagon decomposition of a one-holed torus is formed by three arcs. Our goal is to find a hexagon decomposition where the three orthogeodesics have length $t_1 \leq t_2 \leq t_3$, which are the first three lengths of the simple orthospectrum.

By Proposition [5.1,](#page-24-1) the first two lengths t_1 and t_2 of $\mathcal{O}_S(T) = \mathcal{O}_S(T')$ correspond to two disjoint simple orthogeodesics τ_1 and τ_2 on T (respectively τ'_1 and τ'_2 on T'). To visualize this situation, we give τ_1 and τ_2 an orientation, cut the one-holed torus along τ_1 and τ_2 and obtain a right-angled octagon as in Figure [14.](#page-25-1)

There are exactly two simple orthogeodesics, τ_3 and $\tilde{\tau}_3$, disjoint from τ_1 and τ_2 , each of which joins two opposite sides of the octagon corresponding to arcs in ∂X . Assume that $\ell(\tau_3) \leq \ell(\tilde{\tau}_3)$. Our goal is to prove that any simple orthogeodesic which is not disjoint from τ_1 or τ_2 (or both) is longer than τ_3 .

Figure 14: τ_3 and $\tilde{\tau}_3$ on the octagon

Let τ be a simple orthogeodesic intersecting τ_1 or τ_2 (or both). Without loss of generality, we can assume that τ has its endpoints on the same sides b_1 and b_2 as τ_3 . Indeed, if τ has its endpoints on the opposite sides, we just replace τ_3 by $\tilde{\tau}_3$ in the proof and show that $\ell(\tau_3) \leq \ell(\tilde{\tau}_3) < \ell(\tau)$. Orient τ from its endpoint on b_1 to the endpoint on b_2 . Assume that τ first intersects τ_1 before possibly intersecting τ_2 (the other case being analogous). Let n be the number of time that τ intersects τ_1 before possibly intersecting τ_2 . We prove that $\ell(\tau_3) < \ell(\tau)$ by induction on n.

Base case $n = 1$: The orthogeodesic τ intersects τ_1 exactly once before intersecting τ_2 .

We denote by p_1 the first point of intersection between τ and τ_1 , and by p_2 the last one. Since τ is simple, p_2 does not lie between p_1 and b_2 . In other words, we have $d(p_2, b_1) + d(p_1, b_2) \leq \ell(\tau_1)$. We label by x the segment of τ between p_2 and b_2 , and by y the segment of τ between p_1 and b_1 (as in Figure [15\)](#page-26-0).

Figure 15: Example of case $n = 1$.

We construct two arcs σ and $\tilde{\sigma}$ homotopic to τ_3 as follows: σ is the union of x with the segment of τ_1 between p_2 and b_1 , and $\tilde{\sigma}$ is the union of y with the segment of τ_1 between p_1 and b_2 , as in Figure [16.](#page-26-1) We have

$$
length(\sigma) = d(p_2, b_1) + length(x)
$$

$$
length(\tilde{\sigma}) = d(p_1, b_2) + length(y).
$$

Figure 16: Construction of σ and $\tilde{\sigma}$ for case $n = 1$.

We obtain

 $2\ell(\tau_3) < \text{length}(\sigma) + \text{length}(\tilde{\sigma}) \leq d(p_2, b_1) + d(p_1, b_2) + \text{length}(x) + \text{length}(y) \leq \ell(\tau_1) + \ell(\tau).$ Since $\ell(\tau_1) \leq \ell(\tau)$, we conclude that $\ell(\tau_3) < \ell(\tau)$.

Induction step: Suppose the result is true for any simple orthogeodesic which intersects τ_1 at most *n* times before intersecting τ_2 . Let τ be a simple orthogeodesic which intersects τ_1 $n+1$ times before intersecting τ_2 .

We denote by $p_1, p_2, ..., p_n, p_{n+1}$ the first $n+1$ intersection points of τ and τ_1 . These points cut τ_1 into $n+2$ segments.

Figure 17: Several example of τ .

Let us construct two arcs σ and $\tilde{\sigma}$ as follows. For σ , we start from the edge b_2 and we follow τ until it intersects τ_1 , then we follow τ_1 until p_n , then we follow τ between p_n and p_{n-1} , then τ_1 between p_{n-1} and p_{n-2} and so on until we reach p_1 . If $n+1$ is even, we close up σ by following τ until the edge b_1 ; if $n+1$ is odd, we follow τ_1 until the edge b_1 . We construct $\tilde{\sigma}$ in a similar way, using segments of τ and τ_1 between the intersection points p_i that we did not already use. We start from the side b_2 and follow τ_1 until p_{n+1} , then we follow τ until p_n , then τ_1 until p_{n-1} and we repeat the process until we reach p_1 . Then, if $n + 1$ is even, we follow τ_1 until b_1 ; if $n + 1$ is odd, we follow τ until b_1 .

Figure 18: Several examples of the construction of σ and $\tilde{\sigma}$ in Case 2.

Since σ and $\tilde{\sigma}$ use different segments of τ and τ_1 , we have $\ell(\sigma) + \ell(\tilde{\sigma}) \leq \ell(\tau_1) + \ell(\tau)$. By induction, we know that $\ell(\tau_3) < \ell(\sigma)$ and $\ell(\tau_3) < \ell(\tilde{\sigma})$. Indeed, when $n+1$ is even, the arc σ is homotopic to τ in the induction step $\frac{n+1}{2}$ for $i(\tau, \tau_2) = 0$ and the arc $\tilde{\sigma}$ is homotopic to τ in the induction step $(\frac{n+1}{2} - 1)$ for $i(\tau, \tau_2) = 0$. When $n+1$ is odd, the arcs σ and $\tilde{\sigma}$ are homotopic to τ in the induction step $\frac{n}{2}$ for $i(\tau, \tau_2) = 0$. To conclude, we have $2\ell(\tau_3) < \ell(\sigma) + \ell(\tilde{\sigma}) \leq \ell(\tau) + \ell(\tau_1)$. Thus, $\ell(\tau_3) < \ell(\tau)$.

So the first three lengths of $\mathcal{O}_S(T)$ and $\mathcal{O}_S(T')$ are realized by disjoint orthogeodesics τ_1, τ_2 and τ_3 on T and τ'_1, τ'_2 and τ'_3 on T'. The set $\{\tau_1, \tau_2, \tau_3\}$ is a hexagon decomposition of T and the set $\{\tau'_1, \tau'_2, \tau'_3\}$ is also a hexagon decomposition of T'. Cutting T and T' along their respective hexagon decomposition, we obtain two isometric sets of two hexagons. We have only two hexagons per set and they are isometric, so there is no ambiguity as to how to glue them back into T and T' , which are then isometric. \Box

Finally, in [\[MM23\]](#page-30-4), Masai and McShane also gave an example of two non-isometric hyperbolic surfaces with the same orthospectrum. Their proof does not provide such

an example in the case of the simple orthospectrum. Indeed, they used the fact that if we have a regular d-cover $\pi : X \to X$ of a hyperbolic surface X with boundary, then any orthogeodesic on X is covered by exactly d orthogeodesics on X [\[MM23,](#page-30-4) Lemma 6.1. It is then possible to compute the orthospectrum of X from $\mathcal{O}(X)$ and the degree of the cover. They construct two non-isometric regular degree d cover of the same hyperbolic surface, which then have the same orthospectrum. To use the same argument for the simple orthospectrum, we would need to control which non-simple orthogeodesics on X have simple lift to X . So, similarly to the simple spectrum case, the question of whether the simple orthospectrum determine the surface is still open.

References

- [Abi80] William Abikoff. The real analytic theory of Teichmüller space, volume 820 of Lecture Notes in Mathematics. Springer, Berlin, 1980.
- [Bas93] Ara Basmajian. The orthogonal spectrum of a hyperbolic manifold. Amer. J. Math., 115(5):1139–1159, 1993.
- [Bav05] Christophe Bavard. Anneaux extrémaux dans les surfaces de Riemann. Manuscripta Math., 117(3):265–271, 2005.
- [BCK20] Hyungryul Baik, Inhyeok Choi, and Dongryul M. Kim. Simple length spectra as moduli for hyperbolic surfaces and rigidity of length identities. $arXiv$ eprints, December 2020.
- [Bru52] François Bruhat. Sous-variétés analytiques ; variétés principales. Séminaire Henri Cartan, 4:1–11, 1951-1952. talk:12.
- [BS88] Peter Buser and Klaus-Dieter Semmler. The geometry and spectrum of the one-holed torus. Comment. Math. Helv., 63(2):259–274, 1988.
- [Bus10] Peter Buser. Geometry and spectra of compact Riemann surfaces. Modern Birkhäuser Classics. Birkhäuser Boston, Ltd., Boston, MA, 2010. Reprint of the 1992 edition.
- [FM12] Benson Farb and Dan Margalit. A primer on mapping class groups, volume 49 of Princeton Mathematical Series. Princeton University Press, Princeton, NJ, 2012.
- [Fri67] Jacques Frisch. Points de platitude d'un morphisme d'espaces analytiques complexes. Invent. Math., 4:118–138, 1967.
- [Haa85] Andrew Haas. Length spectra as moduli for hyperbolic surfaces. Duke Math. J., 52(4):923–934, 1985.
- [Kac66] Mark Kac. Can one hear the shape of a drum? Amer. Math. Monthly, 73(4):1–23, 1966.
- [LSZ15] Lixin Liu, Weixu Su, and Youliang Zhong. On metrics defined by length spectra on Teichmüller spaces of surfaces with boundary. Ann. Acad. Sci. Fenn. Math., 40(2):617–644, 2015.
- [McK72] Henry P. McKean. Selberg's trace formula as applied to a compact Riemann surface. Comm. Pure Appl. Math., 25:225–246, 1972.
- [MM23] Hidetoshi Masai and Greg McShane. On systoles and ortho spectrum rigidity. Math. Ann., 385(1-2):939–959, 2023.
- [MP08] Greg McShane and Hugo Parlier. Multiplicities of simple closed geodesics and hypersurfaces in Teichmüller space. Geom. Topol., 12(4):1883–1919, 2008.
- [Mum71] David Mumford. A remark on Mahler's compactness theorem. Proc. Amer. Math. Soc., 28:289–294, 1971.
- [Par23] Hugo Parlier. A shorter note on shorter pants. arXiv e-prints, April 2023.
- [Rat19] John G. Ratcliffe. Foundations of hyperbolic manifolds, volume 149 of Graduate Texts in Mathematics. Springer, Cham, third edition, 2019.
- [Sun85] Toshikazu Sunada. Riemannian coverings and isospectral manifolds. Ann. of Math. (2), 121(1):169–186, 1985.
- [Ush99] Akira Ushijima. A canonical cellular decomposition of the Teichmüller space of compact surfaces with boundary. Comm. Math. Phys., 201(2):305–326, 1999.
- [Vig78] Marie-France Vignéras. Exemples de sous-groupes discrets non conjugués de PSL(2, R) qui ont même fonction zéta de Selberg. C. R. Acad. Sci. Paris Sér. A-B, 287(2):A47–A49, 1978.
- [Wol79] Scott Wolpert. The length spectra as moduli for compact Riemann surfaces. Ann. of Math. (2), 109(2):323–351, 1979.

Univ Gustave Eiffel, Univ Paris Est Creteil, CNRS, LAMA UMR8050 F-77447 Marne-la-Vallée, France Email address: nolwenn.le-quellec@univ-eiffel.fr