## A REFINEMENT OF THE COHERENCE CONJECTURE OF PAPPAS AND RAPOPORT

#### JIUZU HONG AND HUANHUAN YU

ABSTRACT. The coherence conjecture of Pappas and Rapoport, proved by Zhu, asserts the equality of dimensions for the global sections of a line bundle over a spherical Schubert variety in the affine Grassmannian and those of another line bundle over a certain union of Schubert varieties in a partial affine flag variety.

In this paper, we upgrade this equality of dimensions to an isomorphism of representations. This has interesting consequences in the setting of affine Demazure modules. Zhu's proof of coherence conjcture and our comparison theorem are established by introducing a parahoric Bruhat-Tits group scheme  $\mathcal G$  over the affine line that is ramified at 0. We further strengthen this comparison theorem by equipping any line bundle on the global affine Grassmannian of  $\mathcal G$  with a unique equivariant structure of the global jet group scheme of  $\mathcal G$ .

## **CONTENTS**

1. Introduction	2
2. Parahoric group schemes revisited	4
2.1. Notations	4
2.2. Parahoric group schemes from the equivariant point of view	6
2.3. Parahoric group associated to a facet	8
2.4. A generation theorem	10
3. A refinement of the coherence conjecture (after Zhu)	12
3.1. Statement of theorem	12
3.2. Equivariance of line bundles on partial affine flag varieties	15
3.3. Equivariance of line bundles on ramified global affine Grassmannians	19
3.4. Proof of Theorem 3.5	22
4. Application of the coherence conjecture to Affine Demazure modules	24
4.1. Affine Kac-Moody algebras	24
4.2. The coherence conjecture and affine Demazure modules	25
4.3. Inclusion relation among Demazure modules	28
5. Canonical $L^+\mathcal{G}$ -equivariance on line bundles over $Gr_{\mathcal{G}}$	31
5.1. Picard group of $Gr_{\mathcal{G}}$	31
5.2. Canonical equivariance of line bundles on $Gr_{\mathcal{G}}$	32
5.3. General case	36
Appendix A. Examples of Theorem 4.5	36
References	39

1

### 1. Introduction

The coherence conjecture of Pappas and Rapoport [PR] has important applications in local models of Shimura varieties, and it was proved by Zhu in [Zh1]. Despite its significance in arithmetic, its representation-theoretic consequences have not yet been fully explored. In this paper, we will refine Zhu's theorem by upgrading the dimension equality to an isomorphism of representations, which will be further enhanced in the context of line bundles on ramified global affine Grassmannians. Additionally, we will discuss its applications to affine Demazure modules.

Let G be a simple algebraic group G over an algebraically closed field k of characteristic p. Let  $Gr_G$  denote the affine Grassmannian of G. Let L denote the line bundle on  $Gr_G$  of central charge 1. For each dominant coweight  $\mu$  of G, let  $Gr_{G,\mu}$  denote the associated spherical Schubert variety in  $Gr_G$ . Let G((z)) denote the loop group. Let  $\mathcal{P}_{\vartheta}$  be the parahoric subgroup in G((z)) associated to a rational point  $\vartheta$  in the fundamental alcove of G((z)). Let Y be set of vertices of the minimal facet containing  $\vartheta$ . We then attach a partial affine flag variety  $Fl_Y = G((z))/\mathcal{P}_{\vartheta}$ . For any dominant coweight  $\mu$  of G, we can attach a subvariety  $\mathcal{A}_Y(\mu)$  in  $Fl_Y$ ,

(1) 
$$\mathcal{A}_{Y}(\mu) := \bigcup_{w \in W} \overline{\mathrm{Fl}}_{Y,w(\mu)},$$

where W is the Weyl group of G and  $\overline{\operatorname{Fl}}_{Y,w(\mu)}$  is the Iwahori Schubert variety associated to coweight  $w(\mu)$  as an element in the extended affine Weyl group of W. The variety  $\mathcal{A}_Y(\mu)$  was originally defined in terms of admissible set, see Lemma 3.4. The following is the coherence conjecture of Pappas and Rapoport [PR], which is extended and proved by Zhu [Zh1].

**Theorem 1.1.** For any ample line bundle  $\mathcal{L}$  on  $\operatorname{Fl}_Y$  of central charge c, we have the following dimension equality

$$\dim H^0(\overline{\mathrm{Gr}}_{G,\mu}, L^c) = \dim H^0(\mathcal{A}_Y(\mu), \mathcal{L}),$$

where  $H^0(\cdot)$  denote the space of global sections of line bundles.

In fact, Theorem 1.1 also holds in the twisted case, where on the left-hand side remains unchanged and on the right-hand side  $Fl_Y$  is a partial affine flag variety of twisted type. For simplicity of illustration, in this introduction we only focus on the untwisted situation. All of our results presented in the introduction are proved in full generality, with the exception of certain constraints on the characteristic of k.

In the proof of Theorem 1.1, Zhu considered a parahoric Bruhat-Tits group scheme  $\mathcal{G}$  over  $\mathbb{A}^1$  that is ramified at 0. The main geometric result proved by Zhu is that the global Schubert variety  $\overline{\mathrm{Gr}}_{\mathcal{G},\mu}$  of  $\mathcal{G}$ , which is flat over  $\mathbb{A}^1$ , has the generic fiber  $\overline{\mathrm{Gr}}_{\mathcal{G},\mu}$  and the special fiber  $\mathcal{A}_Y(\mu)$  over 0. Furthermore, Zhu constructed a line bundle on the global affine Grassmannian  $\mathrm{Gr}_{\mathcal{G}}$  which specializes to powers of  $L^c$  and  $\mathcal{L}$  at generic and special fibers. Using Frobenius splitting technique, he derived the equality in Theorem 1.1 comparing the global sections of  $L^c$  and  $\mathcal{L}$ .

Any ample line bundle  $\mathcal{L}$  on  $\operatorname{Fl}_Y$  corresponds to an affine dominant weight  $\Lambda$ , and is denoted by  $\mathcal{L}(\Lambda)$ . Let G' be the simply-connected cover of G, and let T' denote the

preimage of T in G'. Let  $\lambda$  be the restriction of  $\Lambda$  to T', and it is a dominant weight of G'. Let  $G'_{\vartheta}$  be the centralizer of a multiple of  $\vartheta$  in G', defined in (15). The image of  $G'_{\vartheta}$  in G is denoted by  $G_{\vartheta}$ . Then, there is a unique G'(O)-equivariant structure on  $\mathscr{L}(\Lambda)$ . The following theorem is the untwisted case of Theorem 3.5 in Section 3.

**Theorem 1.2.** Let  $\mathcal{L}(\Lambda)$  be the ample line bundle on  $\operatorname{Fl}_Y$ . When  $\operatorname{char}(k) = 0$ , there is an isomorphism of representations of  $G'_{\mathfrak{P}}$ :

$$H^0(\overline{\mathrm{Gr}}_{G,\mu},L^c)\simeq H^0(\mathcal{A}_Y(\mu),\mathcal{L}(\Lambda))\otimes \mathrm{k}_\lambda,$$

where the action of  $G'_{\theta}$  on the left side (resp. right side) arises from the G'(O)-equivariant structure on  $L^c$  (resp.  $\mathcal{L}(\Lambda)$ ). In general, this is an isomorphism of T'-representations when char(k) satisfies a very mild condition.

We now explain the main idea of the proof and for simplicity we assume G is simplyconnected and char(k) = 0. Using the method of Faltings [Fa], we construct in Section 3.2 a  $\mathcal{P}_{\vartheta}^-$ -equivariant line bundle  $\mathscr{L}(\Lambda)$  on  $\mathrm{Fl}_Y$ , where  $\mathcal{P}_{\vartheta}^-$  denotes the opposite parahoric subgroup in G((z)) relative to  $\mathcal{P}_{\vartheta}$ . This descends to a line bundle  $\mathcal{L}_{\text{Bun}_G}$  on the moduli stack  $\operatorname{Bun}_{\mathcal{G}}$  of  $\mathcal{G}$ -bundles over  $\mathbb{P}^1$ , where  $\mathcal{G}$  is the parahoric Bruhat-Tits group scheme over  $\mathbb{P}^1$  with parahoric subgroup  $\mathcal{P}_{\vartheta}$  at 0 and the parahoric subgroup  $\mathcal{P}_{-\vartheta}$  at  $\infty$ . By pulling-back, it gives rise to a line bundle  $\mathcal L$  on the global affine Grassmannian  $\mathrm{Gr}_{\mathcal G}$  of  $\mathcal{G}$ . The line bundle  $\mathcal{L}$  specializes to the line bundle  $\mathcal{L}(\Lambda)$  on Fl<sub>y</sub> at 0, and specializes to  $L^c$  on  $Gr_G$  at a generic point in  $\mathbb{P}^1$ . In Theorem 3.21, we equip the line bundle  $\mathcal{L}$  with a natural  $G_{\vartheta} \times \mathbb{P}^1$ -equivariant structure, where  $G_{\vartheta} \times \mathbb{P}^1$  is a subgroup scheme of the global jet group scheme  $L^+\mathcal{G}$ . In the proof, we make crucial use of an equivariant realization of  $\mathcal{G}$ , which relies on the local result stated in Theorem 2.4. Note that  $L^+\mathcal{G}$  specializes to  $\mathcal{P}_{\vartheta}$  at 0 and to G(O) at a generic point. In particular,  $G_{\vartheta}$  is a subgroup of  $\mathcal{P}_{\vartheta}$ . In the statement of Theorem 1.2, we regard  $G_{\theta}$  as a subgroup of G(O), and the discrepancy of  $k_{\lambda}$  arises from comparing equivariant structures of G(O) and  $\mathcal{P}_{\vartheta}$  on  $\mathcal{L}(\Lambda)$ , see Corollary 3.15.

In Section 2, we revisit the theory of parahoric subgroups in the loop groups and their scheme structures. In Theorem 2.4, we extend the equivariant construction of parahoric group schemes from the simply-connected case in [DH1] to general simple algebraic groups. In Proposition 2.8, we describe a canonical Levi subgroup in any parahoric subgroup. Finally, in Theorem 2.10, we prove a generation theorem for parahoric subgroups. These results will play a crucial role in Sections 3 and 5.

In Section 4, we explore a consequence of Theorem 1.2 in the context of affine Demazure modules. By a theorem of Kumar and Mathieu, the dual of  $H^0(\overline{\mathrm{Gr}}_{G,\mu}, L^c)$  is an affine Demazure module  $D(c,\mu)$ , see Theorem 4.6. Furthermore, with additional work (primarily using a result of Littelmann [Li]), the dual of  $H^0(\mathcal{A}_Y(\mu), \mathcal{L}(\Lambda))$  is a summation  $\sum_{w \in W} D(\Lambda, w(\mu))$  of Demazure modules in the integrable highest weight module  $V(\Lambda)$ . Consequently, Theorem 1.2 can be reformulated as the following purely algebraic statement, which is the untwisted case of Theorem 4.5.

**Theorem 1.3.** For any dominant coweight  $\mu \in X_*(T)$ , there is an isomorphism of  $\mathfrak{g}_{\vartheta}$ -modules

$$D(c,\mu) \otimes \mathbf{k}_{\lambda} \simeq \sum_{\substack{w \in W \\ 3}} D(\Lambda, w(\mu)),$$

where  $g_{\vartheta}$  denotes the Lie algebra of  $G_{\vartheta}$  and  $k_{\lambda}$  is the 1-dimensional representation of  $g_{\vartheta}$ .

Demazure modules have been extensively studied in the literature, including their character formula, cf. [Ku2, Ma]. Theorem 1.3 reveals a surprising relationship between different Demazure modules, one that has not been previously noticed. In the Appendix A, we present explicit examples to illustrate this phenomenon. In Section 4.3, we also discuss the inclusion relation among Demazure modules  $\{D(\Lambda, w(\mu)) | w \in W\}$ , as well as the closure relation between Schubert varieties  $\overline{Fl}_{Xw(\mu)}$  contained in  $\mathcal{A}_Y(\mu)$ .

In Section 5.1, we classify all line bundles on the global affine Grassmannian  $Gr_{\mathcal{G}}$ , where  $\mathcal{G}$  is the parahoric Bruhat-Tits group scheme over  $\mathbb{A}^1$  ramified at 0. In fact, any line bundle  $\mathcal{L}$  on  $Gr_{\mathcal{G}}$  is uniquely determined by its restriction  $\mathcal{L}_0$  on the fiber  $Gr_{\mathcal{G},0} \simeq Fl_Y$  over 0, and thus it is parametrized by an affine dominant weight  $\Lambda$ , see Proposition 5.1. The following theorem is the untwisted case of Theorem 5.2, assuming G is simply-connected. A version of this theorem is proved in Theorem 5.6 for any simple algebraic group G.

**Theorem 1.4.** When char(k) = 0, any line bundle  $\mathcal{L}$  on  $Gr_{\mathcal{G}}$  has a unique  $L^+\mathcal{G}$ -equivariant structure, where  $L^+\mathcal{G}$  is the global jet group scheme of  $\mathcal{G}$ .

This theorem can be viewed as an enhancement of Theorem 1.2 or, more precisely, Theorem 3.21, as we can regard  $G_{\vartheta} \times \mathbb{A}^1$  as a closed subgroup scheme of  $L^+\mathcal{G}$ . When  $\mathcal{G}$  is special parahoric at  $0 \in \mathbb{A}^1$ , this result was essentially proved in a previous work [HY] of the authors. We expect that Theorem 1.4 will have applications in the equivariant K-theory of partial affine flag varieties, which will be the focus of a future research project.

We now outline the strategy for proving Theorem 1.4. By Lemma 5.3 we first reduce the question to global Schubert varieties. For any line bundle  $\mathcal{L}$  on  $Gr_{\mathcal{G}}$ , let  $\widehat{L^+\mathcal{G}}$  be the central extension of  $L^+\mathcal{G}$  by the constant group scheme  $\mathbb{G}_{m,\mathbb{A}^1}$  arising from  $\mathcal{L}$ . Using the equivariant method, we show that the central extension  $\widehat{L^+\mathcal{G}}$  splits uniquely over the jet group scheme of any global root subgroup. The splitting over  $T_{\mathbb{A}^1}$  is already established in Theorem 3.21. Then, using the generation theorem in Theorem 2.10, these splittings together yield a splitting of  $\widehat{L^+\mathcal{G}} \to L^+\mathcal{G}$ , thereby proving Theorem 1.4.

**Acknowledgement**: We would like to thank S. Kato, S. Kumar, S. Nie and X. Zhu for helpful discussions. We also thank P. Littelmann for valuable email correspondence. J. Hong was partially supported by NSF grant DMS-2001365 and Simons grant MPS-TSM-00007468.

#### 2. Parahoric group schemes revisited

In this section, we review and extend some results in [DH1] on the equivariant construction of parahoric group schemes over the ring of formal power series. We describe a natural Levi subgroup in any parahoric group. We also prove a generation theorem for parahoric subgroups. The results in this section will be used in later sections.

2.1. **Notations.** Let k be an algebraically closed field with characteristic p. Let G be a connected simple algebraic group over k. We fix a Borel subgroup  $B \subseteq G$  and a maximal

torus  $T \subseteq B$ . Denote by  $X_*(T)$  the lattice of cocharacters of T. Let  $G_{ad}$  be the adjoint quotient of G and let  $T_{ad}$  be the image of T in  $G_{ad}$ . Similarly, let  $X_*(T_{ad})$  denote the lattice of cocharacters of  $T_{ad}$ .

Let  $\tau$  be a diagram automorphism of G preserving B and T with order r. Note that  $\tau$  can be trivial, and r can only be either 1, 2, or 3. Throughout this paper, we always assume  $p \neq r$ . Let  $\mathcal{K} := \mathrm{k}((z))$  and  $O := \mathrm{k}[[z]]$  be the ring of formal Laurent series and formal power series respectively. We fix an r-th primitive unit root  $\epsilon$  and define  $\tau(z) := \epsilon^{-1}z$ . Set  $\bar{\mathcal{K}} := \mathrm{k}((z^r))$ , and  $\bar{O} := \mathrm{k}[[z^r]]$ .

Let  $G^{\tau,\circ}$  (resp.  $B^{\tau,\circ}$ ,  $T^{\tau,\circ}$ ) denote the neutral component of the  $\tau$  fixed point subgroup of G (resp. B, T). Then,  $B^{\tau,\circ}$  is a Borel subgroup of  $G^{\tau,\circ}$  and  $T^{\tau,\circ}$  is a maximal torus. In the following sections,  $\Box^{\circ}$  always stands for the neutral component of  $\Box$ . Note that when G is simply-connected or adjoint,  $G^{\tau,\circ} = G^{\tau}$  and  $T^{\tau,\circ} = T^{\tau}$ .

Let  $I_{\tau}$  (resp. I) be the set of vertices of the Dynkin diagram of  $G^{\tau}$  (resp. G). Then,  $I_{\tau}$  can be identified with the set of  $\tau$ -orbits in I. Let  $X_N^{(r)}$  be the affine Dynkin diagram associated to the twisted loop group  $G(\mathcal{K})^{\tau}$ . By adjoining an affine node o,  $\hat{I}_{\tau} := I_{\tau} \sqcup \{o\}$  can be naturally regarded as the set of vertices of  $X_N^{(r)}$ . We remark that o corresponds to the vertex 0 if  $X_N^{(r)} \neq A_{2\ell}^{(2)}$  or the vertex  $\ell$  if  $X_N^{(r)} = A_{2\ell}^{(2)}$  in [Kac, Chapter 4]. Let  $(a_i)_{i \in \hat{I}_{\tau}}$  be the Kac labels in the table in [Kac, Chapter 4] and let  $(\check{a}_i)_{i \in \hat{I}_{\tau}}$  be the dual

Let  $(a_i)_{i \in \hat{I}_{\tau}}$  be the Kac labels in the table in [Kac, Chapter 4] and let  $(\check{a}_i)_{i \in \hat{I}_{\tau}}$  be the dual Kac labels. Note that  $\check{a}_o = 2$  if  $X_N^{(r)} = A_{2\ell}^{(2)}$ , and  $\check{a}_o = 1$  otherwise; moreover,  $a_o = 1$  for any cases. Let  $\{\alpha_i \mid i \in \hat{I}_{\tau}\}$  be the set of simple affine roots and let  $\{\check{\alpha}_i \mid i \in \hat{I}_{\tau}\}$  be the set of simple affine coroots. Set

(2) 
$$\theta_0 := \sum_{i \in I_\tau} a_i \alpha_i,$$

We remark that  $\theta_0$  agrees with the definition in [Kac, Proposition 8.3], which can be described as in (40). Let  $\{\omega_i | i \in I_\tau\}$  be the set of fundamental weights of the simply connected cover  $(G^{\tau,\circ})'$  of  $G^{\tau,\circ}$ .

Following [Kac, §6.2], we can associate the following vector space

$$\mathfrak{h}^{ au}_{\mathbb{Q}} := igoplus_{i \in I_{ au}} \mathbb{Q} \check{lpha}_i, \quad \hat{\mathfrak{h}}^{ au}_{\mathbb{Q}} := igoplus_{i \in \hat{I}_{ au}} \mathbb{Q} \check{lpha}_i \oplus \mathbb{Q} d,$$

where d is the scaling element. We can also define the central element as  $K = \sum_{i \in \hat{I}_{\tau}} \check{a}_i \check{\alpha}_i$ . For each  $i \in \hat{I}_{\tau}$ , let  $\Lambda_i \in (\hat{\mathfrak{h}}_{\mathbb{Q}}^{\tau})^*$  be the associated affine fundamental weight, i.e.  $\langle \Lambda_i, \check{\alpha}_j \rangle = \delta_{ij}$ , and  $\langle \Lambda_i, d \rangle = 0$ . Then  $(\hat{\mathfrak{h}}_{\mathbb{Q}}^{\tau})^*$  admits the following decomposition

(3) 
$$(\hat{\mathfrak{h}}_{\mathbb{Q}}^{\tau})^* = (\mathfrak{h}_{\mathbb{Q}}^{\tau})^* \oplus \mathbb{Q}\Lambda_o \oplus \mathbb{Q}\delta,$$

where  $\delta = \sum_{i \in \hat{I}_{\tau}} a_i \alpha_i$ . Let  $(\cdot|\cdot)$  be the normalized invariant form on  $\hat{\mathfrak{h}}_{\mathbb{Q}}^{\tau}$  defined as in [Kac, §6.2]. This bilinear form induced a map  $\nu : \hat{\mathfrak{h}}_{\mathbb{Q}}^{\tau} \to (\hat{\mathfrak{h}}_{\mathbb{Q}}^{\tau})^*$ . For any  $i \in \hat{I}_{\tau}$ , we have

(4) 
$$\nu(\check{\alpha}_i) = \frac{a_i}{\check{a}_i}\alpha_i, \quad \nu(K) = \delta.$$

Note that we have the following natural identifications:  $\mathfrak{h}^{\tau}_{\mathbb{Q}} \simeq X_*(T)^{\tau}_{\mathbb{Q}} \simeq X_*(T_{\mathrm{ad}})^{\tau}_{\mathbb{Q}}$ . Moreover,  $\nu$  restricts to a map  $\nu : \mathfrak{h}^{\tau}_{\mathbb{Q}} \simeq (\mathfrak{h}^{\tau}_{\mathbb{Q}})^*$ .

2.2. Parahoric group schemes from the equivariant point of view. Let  $\mathscr{G}_{\bar{K}}$  be the quasi-split group scheme  $\operatorname{Res}_{\mathcal{K}/\bar{K}}(G_{\mathcal{K}})^{\tau}$  over  $\bar{\mathcal{K}}$ , i.e. the subgroup scheme of  $\tau$ -invariants of the Weil restriction  $\operatorname{Res}_{\mathcal{K}/\bar{K}}(G_{\mathcal{K}})$ . Note that any tamely-ramified quasi-split simple group scheme over  $\bar{\mathcal{K}}$  can be realized from such construction.

Let S denote the group scheme  $(T^{\tau,\circ})_{\bar{K}}$ , which is a maximal  $\bar{K}$ -split torus of  $\mathscr{G}_{\bar{K}}$ . Then, the relative root system  $\mathcal{R}$  for  $(\mathscr{G}_{\bar{K}},S)$  can be identified with the quotient of the root system R of (G,T) by the  $\tau$ -action. We remark that the root system  $R(G^{\tau,\circ},T^{\tau,\circ})$  of  $(G^{\tau,\circ},T^{\tau,\circ})$  is naturally a subsystem of  $\mathcal{R}$ . In fact,  $R(G^{\tau,\circ},T^{\tau,\circ})=\mathcal{R}$  if and only if  $(G,r)\neq (A_{2\ell},2)$ .

Let  $\Delta_{\alpha} \subseteq R$  denote the set of preimages of  $\alpha \in R$  via the projection map  $R \to R$ . For each  $\alpha \in R$ , we denote by  $\mathcal{K}_{\alpha}$  the subfield of  $\mathcal{K}$  which is fixed by the stabilizer of any  $\tilde{\alpha} \in \Delta_{\alpha}$  under the action of the cyclic group  $\langle \tau \rangle$ . Let  $d_{\alpha}$  be the cardinality of the set  $\Delta_{\alpha}$ . Then,  $\mathcal{K}_{\alpha} = k((z^{r/d_{\alpha}}))$ .

We now recall the root subgroup  $\mathcal{U}_{\alpha}$  of  $\mathcal{G}_{\bar{K}}$  for  $\alpha \in \mathcal{R}$ . When  $2\alpha, \frac{1}{2}\alpha \notin \mathcal{R}$ , the root subgroup  $\mathcal{U}_{\alpha}$  is given by

 $\Big(\prod_{\tilde{\alpha}\in\Lambda_{\alpha}}U_{\tilde{\alpha}}(\mathcal{K}_{\alpha})\Big)^{\tau},$ 

where  $U_{\tilde{\alpha}}$  denote the root subgroup of G associated to  $\tilde{\alpha}$ . Since the  $\tau$ -action on  $\Delta_{\alpha}$  is transitive, the root subgroup homomorphism  $x_{\tilde{\alpha}}: \mathbb{G}_a \to U_{\tilde{\alpha}}$  induces a Chevalley-Steinberg pinning  $\{x_{\alpha}: \mathcal{K}_{\alpha} \to \mathcal{U}_{\alpha} \mid \alpha \in \mathcal{R}\}$  of  $\mathcal{G}_{\tilde{\mathcal{K}}}$ , cf. [BT, 4.1.5]. Let  $\nu$  be the valuation of  $\mathcal{K}$  with  $\nu(z) = 1$ . Via the pinning  $x_{\alpha}$ , the restriction of  $\nu$  on  $\mathcal{K}_{\alpha}$  induces a valuation  $\nu$  on  $\mathcal{U}_{\alpha}$  (by abuse of notation). For every  $\ell \in \mathbb{R}$ , we define  $\mathcal{U}_{\alpha,\ell} := \nu^{-1}[\ell, \infty] \subseteq \mathcal{U}_{\alpha}$ .

When  $2\alpha \in \mathcal{R}$ , we have r = 2,  $\Delta_{\alpha} = \{\tilde{\alpha}, \tau(\tilde{\alpha})\}$  is the set of two adjacent roots in R, and  $\Delta_{2\alpha} = \{\tilde{\alpha} + \tau(\tilde{\alpha})\}$ . In this case, we have

$$\mathscr{U}_{\alpha} = (U_{\tilde{\alpha}}(\mathcal{K})U_{\tilde{\alpha}+\tau(\tilde{\alpha})}(\mathcal{K})U_{\tau(\tilde{\alpha})}(\mathcal{K}))^{\tau}, \quad \mathscr{U}_{2\alpha} = U_{\tilde{\alpha}+\tau(\tilde{\alpha})}(\bar{\mathcal{K}}).$$

There are natural valuations defined on  $\mathcal{U}_{\alpha}$  and  $\mathcal{U}_{2\alpha}$ , and we can accordingly define subgroups  $\mathcal{U}_{\alpha,\ell}$  and  $\mathcal{U}_{2\alpha,\ell}$  for every  $\ell \in \mathbb{R}$ , cf. [BT, 4.1.9]. By abuse of notation, we will use the notation  $\nu$  uniformly for the valuation on the root subgroup  $\mathcal{U}_{\alpha}$  for any  $\alpha \in \mathcal{R}$ .

**Definition 2.1.** Following [DH1, Definition 2.1.2] and references therein, for any  $\vartheta \in X_*(S)_{\mathbb{R}} \simeq X_*(T)_{\mathbb{R}}^{\tau}$ , we define the parahoric subgroup  $\mathcal{P}_{\vartheta}$  to be the subgroup of  $G(\mathcal{K})^{\tau}$  generated by  $\{\mathscr{U}_{\alpha, \neg \langle \vartheta, \alpha \rangle}\}_{\alpha \in \mathcal{R}}$  and  $T(O)^{\tau, \circ}$  (the neutral component of  $T(O)^{\tau}$ ). Then, there exists a unique affine smooth group scheme  $\mathscr{G}_{\vartheta}$  over  $\bar{O}$ , which extends  $\mathscr{G}_{\bar{\mathcal{K}}}$  and satisfies that  $\mathscr{G}_{\vartheta}(\bar{O}) = \mathcal{P}_{\vartheta}$ .

Similarly, we define a subgroup  $\mathcal{P}^-_{\vartheta}$  of  $G[z,z^{-1}]^{\tau}$  as follows. Let  $\nu_{\infty}$  be the valuation of  $k[z,z^{-1}]$  with  $\nu_{\infty}(z^{-1})=1$ . When  $2\alpha,\frac{1}{2}\alpha\notin\mathcal{R}$ , via the pinning  $x_{\alpha}$ , the restriction of  $\nu_{\infty}$  on  $k[z,z^{-1}]\cap\mathcal{K}_{\alpha}$  induces a valuation  $\nu_{\infty}$  on  $\mathcal{U}_{\alpha}\cap G[z,z^{-1}]^{\tau}$ . For every  $\ell\in\mathbb{R}$ , we define  $\mathcal{U}^-_{\alpha,\ell}:=\nu_{\infty}^{-1}[\ell,\infty]\subseteq\mathcal{U}_{\alpha}\cap G[z,z^{-1}]^{\tau}$ . When  $2\alpha\in\mathcal{R}$ , we can define subgroups  $\mathcal{U}^-_{\alpha,\ell}$  and  $\mathcal{U}^-_{2\alpha,\ell}$  for every  $\ell\in\mathbb{R}$  in a similar way as explained before. Then,  $\mathcal{P}^-_{\vartheta}$  is the subgroup of  $G[z,z^{-1}]^{\tau}$  generated by  $\{\mathcal{U}^-_{\alpha(\vartheta,\alpha)}\}_{\alpha\in\mathcal{R}}$  and  $T^{\tau,\circ}$ .

Remark 2.2. The valuation on k((z)) used in our definition is different from the one used in [DH1], and the parahoric subgroup  $\mathcal{P}_{\vartheta}$  defined in this paper coincides with the parahoric subgroup  $\mathcal{P}_{r\vartheta}$  defined in loc. cit.

Fix a point  $\vartheta \in X_*(T)^\tau_{\mathbb{Q}}$ . Let m be the minimal positive multiple of r such that  $\frac{m}{r}\vartheta \in X_*(T_{\mathrm{ad}})^\tau$ . Identify the lattice  $X_*(T_{\mathrm{ad}})^\tau$  with  $\bigoplus_{i\in I_\tau}\mathbb{Z}\cdot\check{\omega}_i\subseteq\mathfrak{h}^\tau_{\mathbb{Q}}$ , where  $\check{\omega}_i$  represents the fundamental coweight of  $G^\tau$  associated to  $i\in I_\tau$ . Then, we can write

$$\frac{m}{r}\vartheta=\sum_{i\in I_{\tau}}s_{i}\check{\omega}_{i},$$

where  $s_i$  are integers. Suppose that  $p \nmid m$ . Fix a primitive m-th root of unity  $\epsilon_m$ . We associate an element  $(\epsilon_m)^{m\vartheta/r} \in T_{\mathrm{ad}}$ , and we define an automorphism of G

(5) 
$$\sigma := \tau \circ \mathrm{Ad}_{(\epsilon_m)^{m\theta/r}}.$$

## **Lemma 2.3.** The order of $\sigma$ is m.

*Proof.* Clearly, we have  $\sigma^m = \text{id}$ . Suppose  $\operatorname{ord}(\sigma) = m'$ . Then,  $m' \mid m$ . Recall that  $I_{\tau}$  can be identified with the set of  $\tau$ -orbits in I. For each  $i \in I_{\tau}$ , let  $\eta_i$  be the corresponding  $\tau$ -orbit in I. Pick  $j \in \eta_i$  and let  $x_{\tilde{\alpha}_j} : \mathbb{G}_a \to G$  be the root subgroup associated to the simple root  $\tilde{\alpha}_j$  of G. Then,  $\tilde{\alpha}_j|_{T^{\tau,\circ}} = \alpha_i$  and we have

$$x_{\tilde{\alpha}_i}(u) = \sigma^{m'}(x_{\tilde{\alpha}_i}(u)) = x_{\tau^{m'}\tilde{\alpha}_i}((\epsilon_m)^{m'\langle h, \alpha_i \rangle}u) = x_{\tau^{m'}\tilde{\alpha}_i}((\epsilon_m)^{m's_i}u), \quad u \in \mathbb{R}.$$

Hence, m' is a multiple of r, and  $m's_i$  is a multiple of m for all  $i \in I_\tau$ . Therefore,  $m' \cdot \gcd\{s_i \mid i \in I_\tau\}$  is a multiple of m. From the definition of  $s_i$ , we have  $\gcd\{s_i \mid i \in I_\tau\}$  and m are coprime. Thus,  $m \mid m'$  which implies  $m = m' = \operatorname{ord}(\sigma)$ .

Set  $t = z^{r/m}$ ,  $O_t := k[[t]]$ , and  $\mathcal{K}_t := k((t))$ . Define an action of  $\sigma$  on t by  $\sigma(t) := (\epsilon_m)^{-1}t$ . Then,  $\bar{O} = k[[z^r]] = k[[t^m]]$  is the set of  $\sigma$ -fixed elements in  $O_t$ . Consider the group scheme  $\mathcal{H} := \operatorname{Res}_{O_t/\bar{O}}(G_{O_t})^{\sigma}$  over  $\bar{O}$ , which is the subgroup scheme of  $\sigma$ -invariants of the Weil restriction  $\operatorname{Res}_{O_t/\bar{O}}(G_{O_t})$ . We define the neutral component  $\mathcal{H}^{\circ}$  of  $\mathcal{H}$  to be the union of the neutral components  $(\mathcal{H}_a)^{\circ}$  of fibers at every  $a \in \operatorname{Spec}(\bar{O})$ , cf. [BT, 1.1.12]. Then,  $\mathcal{H}^{\circ}$  is an open subscheme of  $\mathcal{H}$ , denoted by  $\mathscr{G}_{\sigma}$ , i.e.

(6) 
$$\mathscr{G}_{\sigma} := \operatorname{Res}_{O/\bar{O}}(G_{O_{\ell}})^{\sigma,\circ}.$$

We can regard  $G(O_t)^{\sigma}$  as a pro-algebraic group over k. Let  $G(O_t)^{\sigma,\circ}$  be the connected component of  $G(O_t)^{\sigma}$  as a pro-algebraic group.

The following theorem generalizes [DH1, Theorem 4.1.2].

**Theorem 2.4.** When  $p \nmid m$ , there is a unique isomorphism of group schemes over  $\bar{O}$ :

(7) 
$$\phi_0: \mathcal{G}_{\vartheta} \simeq \mathcal{G}_{\sigma}.$$

such that over its  $\bar{O}$ -points,  $\phi_0: \mathcal{P}_{\vartheta} \simeq G(O_t)^{\sigma,\circ}$  is the restriction of the isomorphism  $\mathrm{Ad}_{z^{\vartheta}}: G(\mathcal{K})^{\tau} \simeq G(\mathcal{K}_t)^{\sigma}$ .

*Proof.* When G is simply-connected, the theorem is proved in [DH1, Theorem 4.1.2]. For general simple algebraic group G, we can adapt the same argument, except that we need to check  $\mathscr{G}_{\sigma}(\bar{O}) = G(O_t)^{\sigma,\circ}$ . Since  $\mathscr{G}_{\sigma}$  is the open subscheme of  $\operatorname{Res}_{O_t/\bar{O}}(G_{O_t})^{\sigma}$  throwing away the non-neutral components of the special fiber

$$\operatorname{Res}_{O_t/\bar{O}}(G_{O_t})^{\sigma}|_{0} = G(\mathbf{k}[t]/(t^m))^{\sigma},$$

 $\mathscr{G}_{\sigma}(\bar{O})$  consists of the points  $x \in G(k[[t]])^{\sigma}$  such that its image  $\bar{x}_m \in G(k[t]/(t^m))^{\sigma}$  is contained in the neutral component  $G(k[t]/(t^m))^{\sigma,\circ}$ , equivalently the image  $\bar{x}$  of x in  $G^{\sigma}$  is contained in  $G^{\sigma,\circ}$ . Thus,  $\mathscr{G}_{\sigma}(\bar{O}) = G(O_t)^{\sigma,\circ}$ .

Remark 2.5. The parahoric subgroup is only determined by the minimal facet of the alcove containing  $\vartheta$ . In fact, under the condition that  $p \neq r$  and that p is a good prime, we can always find a rational interior point  $\vartheta$  of this facet such that  $p \nmid m$ , including the case when the facet is a vertex, see Lemma 2.6.

Replacing z by  $z^{-1}$  and let  $v_{\infty}$  be the valuation on  $k((z^{-1}))$  with  $v_{\infty}(z^{-1})=1$ , we can similarly define a parahoric group scheme  $\mathscr{G}_{-\vartheta,\infty}$  over  $\bar{O}_{\infty}$  associated with  $-\vartheta$ , where  $\bar{O}_{\infty}:=k[[z^{-r}]]$ . Set  $O_{t,\infty}:=k[[t^{-1}]]$  and define a parahoric group scheme  $\mathscr{G}_{\sigma,\infty}:=\mathrm{Res}_{O_{t,\infty}/\bar{O}_{\infty}}(G_{O_{t,\infty}})^{\sigma,\circ}$ . Again, we have an isomorphism parahoric group schemes

(8) 
$$\phi_{\infty}: \mathcal{G}_{-\vartheta,\infty} \simeq \mathcal{G}_{\sigma,\infty}.$$

such that over its  $\bar{O}_{\infty}$ -points,  $\phi_{\infty}: \mathscr{G}_{-\vartheta,\infty}(\bar{O}_{\infty}) \simeq G[[t^{-1}]]^{\sigma,\circ}$  is the restriction of the isomorphism

(9) 
$$Ad_{(z^{-1})^{-\theta}} = Ad_{z^{\theta}} : G((z^{-1}))^{\tau} \simeq G((t^{-1}))^{\sigma}.$$

2.3. **Parahoric group associated to a facet.** Let A be the fundamental alcove in  $X_*(T)^\tau_\mathbb{R}$  defined by inequalities  $\langle x, \alpha_i \rangle \geq 0$  for each  $i \in I_\tau$ , and  $\langle x, \theta_0 \rangle \leq 1$ . For convenience, set  $\check{\omega}_o = 0$ . Then, A is a simplex with vertices  $\{\frac{\check{\omega}_i}{a_i} \mid i \in \hat{I}_\tau\}$ . In this section, we fix a nonempty subset  $Y \subseteq \hat{I}_\tau$  and a facet  $F_Y$  of A spanned by the set of vertices  $\{\frac{\check{\omega}_i}{a_i} \mid i \in Y\}$ . We associate a positive integer

$$(10) a_Y := r \cdot \gcd\{a_i \mid i \in Y\}.$$

**Lemma 2.6.** For any  $p \nmid a_Y$ , there is a rational interior point  $\vartheta \in F_Y$  such that  $p \nmid m$ , where m is the minimal positive multiple of r such that  $\frac{m}{r}\vartheta \in X_*(T_{ad})^{\tau}$ .

*Proof.* Set  $n_Y := \sum_{i \in Y} a_i$  and  $x_Y := \frac{1}{n_Y} \sum_{i \in Y} \check{\omega}_i$ . For each  $k \in Y$ , set  $n_k := a_k + n_Y$  and  $x_k := \frac{1}{n_k} (\check{\omega}_k + \sum_{i \in Y} \check{\omega}_i)$ . Then,  $x_Y$  and  $x_k$  are rational interior points of  $F_Y$  for any  $k \in Y$ . We will show that some point among  $\{x_Y, x_k \mid k \in Y\}$  satisfies the lemma.

Let  $m_Y$  and  $m_k$  be minimal positive multiple of r such that  $\frac{m_Y}{r}x_Y \in X_*(T_{\mathrm{ad}})^{\mathsf{T}}$  and  $\frac{m_k}{r}x_k \in X_*(T_{\mathrm{ad}})^{\mathsf{T}}$  respectively. Clearly,  $\frac{m_Y}{r} \mid n_Y$  and  $\frac{m_k}{r} \mid n_k$ . We claim  $p \nmid \frac{m_Y}{r}$  or  $p \nmid \frac{m_k}{r}$  for some  $k \in Y$ . Suppose that  $p \mid \frac{m_Y}{r}$  and  $p \mid \frac{m_k}{r}$  for all  $k \in Y$ . Then,  $p \mid n_Y$  and  $p \mid n_k$  for all  $k \in Y$ . Hence,  $p \mid a_k$  for all  $k \in Y$ . This contradicts with the assumption that  $p \nmid a_Y$ . Therefore,  $p \nmid \frac{m_Y}{r}$  or  $p \nmid \frac{m_k}{r}$  for some  $k \in Y$ . Note that the assumption  $p \nmid a_Y$  implies that  $p \nmid r$ . Since p is a prime, it follows that  $p \nmid m_Y$  or  $p \nmid m_k$  for some  $k \in Y$ .

From now on, we always assume  $p \nmid a_Y$ . Moreover, we fix a rational interior point  $\vartheta \in F_Y$  so that  $p \nmid m$  (the existence follows from Lemma 2.6).

**Definition 2.7.** Set  $\mathcal{P}_Y := \mathcal{P}_{\vartheta}$  and  $\mathcal{G}_Y := \mathcal{G}_{\vartheta}$ . In fact,  $\mathcal{P}_Y$  is the parahoric subgroup corresponding to the facet  $F_Y$ . We call it the parahoric subgroup associated to Y. When  $Y = \hat{I}_{\tau}$ ,  $\mathcal{P}_Y$  is the Iwahori subgroup, denoted by I; when  $Y = \{o\}$ ,  $\mathcal{P}_Y = G(O)^{\tau,\circ}$ . Similarly, we set  $\mathcal{P}_Y^- := \mathcal{P}_{\vartheta}^-$ . When  $Y = \hat{I}_{\tau}$ , we denote  $\mathcal{P}_Y^-$  by  $I^-$ .

Set  $h := \frac{m}{r} \vartheta \in X_*(T_{ad})^{\tau}$ . We can write  $h = \sum_{i \in I_{\tau}} s_i \check{\omega}_i$  for non-negative integers  $s_i, i \in I_{\tau}$ . Since  $\langle \vartheta, \theta_0 \rangle \leq 1$ , we have  $\sum_{i \in I_{\tau}} a_i s_i \leq \frac{m}{r}$ . Set

$$s_o := \frac{m}{r} - \sum_{\substack{i \in I_\tau \ g}} a_i s_i \in \mathbb{Z}^{\geq 0}.$$

Since  $\vartheta$  is an interior point of  $F_Y$ , we must have  $Y = \{i \in \hat{I}_\tau \mid s_i \neq 0\}$ . Define an automorphism  $\sigma := \tau \circ \operatorname{Ad}_{(\epsilon_m)^{m\vartheta/r}}$  of G, whose order is m, see Lemma 2.3. By Theorem 2.4, we have an isomorphism of parahoric group schemes

$$\phi: \mathcal{G}_{Y} \simeq \mathcal{G}_{\sigma}.$$

such that over its  $\bar{O}$ -points,  $\phi: \mathcal{P}_Y \simeq G(O_t)^{\sigma,\circ}$  is the restriction of the isomorphism  $\mathrm{Ad}_{t^h}: G(\mathcal{K})^{\tau} \simeq G(\mathcal{K}_t)^{\sigma}$ . We define  $M_Y = \mathrm{Ad}_{t^{-h}}(G^{\sigma,\circ})$  as a subgroup of  $\mathcal{P}_Y$ . Clearly,  $M_Y$  is a Levi subgroup of  $\mathcal{P}_Y$ .

We describe a pinning of  $G^{\tau,\circ}$  as follows, cf. [HS, Lemma 2.2]. Let  $\{\tilde{\alpha}_j | j \in I\} \subseteq R$  denote the set of simple roots of G with respect to the pair (B,T). We fix a pinning  $\{x_{\pm \tilde{\alpha}_j} : \mathbb{G}_a \to G | j \in I\}$  of G which is preserved by  $\tau$ , i.e.  $\tau(x_{\pm \tilde{\alpha}_j}(a)) = x_{\pm \tilde{\alpha}_{\tau(j)}}(a)$  for any  $a \in k$ . For each  $i \in I_{\tau}$ , let  $\eta_i$  denote the preimage of i via the orbit map  $I \to I_{\tau}$ .

If the vertices in  $\eta_i$  are not adjacent, set

(12) 
$$x_{\alpha_i}(a) = \prod_{j \in \eta_i} x_{\tilde{\alpha}_j}(a), \quad x_{-\alpha_i}(a) = \prod_{j \in \eta_i} x_{-\tilde{\alpha}_j}(a), \quad \forall a \in \mathbf{k}.$$

If the vertices in  $\eta_i$  are adjacent, we can write  $\eta_i = \{j, \tau(j)\}$ . Set

$$(13) x_{\alpha_i}(a) = x_{\tilde{\alpha}_j}(a)x_{\tilde{\alpha}_{\tau(j)}}(2a)x_{\tilde{\alpha}_j}(a), x_{-\alpha_i}(a) = x_{-\tilde{\alpha}_j}(\frac{a}{2})x_{-\tilde{\alpha}_{\tau(j)}}(a)x_{-\tilde{\alpha}_j}(\frac{a}{2}), \forall a \in \mathbf{k}.$$

Let  $\theta^0$  be the root of G described in [Kac, Section 8.3]. Then,

$$\theta^0|_{T^{\tau,\circ}} = \theta_0.$$

When  $(G,r)=(A_{2\ell},2)$ , from [Kac, Section 8.3]  $\theta^0$  is the highest root of G, we choose any root subgroups  $x_{\pm\theta^0}$  associated to  $\pm\theta^0$ . When r>1 and  $(G,r)\neq (A_{2\ell},2)$ ,  $\tau(\theta^0)\neq \theta^0$ . In this case, we can choose root subgroups  $\{x_{\pm\tau^k(\theta^0)}\mid k=0,\cdots,r-1\}$  so that they are preserved by  $\tau$ .

For any  $u \in \mathcal{K}$ , we set

$$x_{\pm\theta_0}(u) := \begin{cases} x_{\pm\theta_0}(u) & \text{when } r = 1 \\ x_{\pm\theta^0}(u) & \text{when } (G,r) = (A_{2\ell},2) \\ \prod_{k=0}^{r-1} x_{\pm\tau^k(\theta^0)}(\tau^k(u)) & \text{when } r \neq 1 \text{ and } (G,r) \neq (A_{2\ell},2) \end{cases}.$$

For any  $a \in k$ , we define

$$x_{\pm \alpha_o}(a) := x_{\mp \theta_0}(az^{\pm 1}) \in G(\mathcal{K})^{\tau}.$$

**Proposition 2.8.** With the same setting as above, we have

(1) The group  $G^{\sigma,\circ}$  is generated by root subgroups  $\{x_{\pm\alpha} \mid \alpha \in \Pi_{\sigma}\}$  and  $T^{\tau,\circ}$ , where  $\Pi_{\sigma}$  is the following set of roots

$$\Pi_{\sigma} = \begin{cases} \{ \alpha_i \mid i \notin Y \}, & if \ o \in Y; \\ \{ -\theta_0, \alpha_i \mid i \notin Y \cup \{o\} \}, & if \ o \notin Y \end{cases}.$$

(2) The group  $M_Y$  is generated by  $\{x_{\pm\alpha_i}(a) | i \notin Y, a \in k\}$  and  $T^{\tau,\circ}$ .

*Proof.* By Lemma 2.3 and the assumption that  $p \nmid m$ , p does not divide the order of  $\sigma$ . In view of [St, Theorem 8.1],  $G^{\sigma,\circ}$  is a connected reductive algebraic group. Moreover, by [Bor, 9.1],  $Lie(G^{\sigma,\circ}) = \mathfrak{g}^{\sigma}$ . When char(k) = 0, the root system of  $\mathfrak{g}^{\sigma}$  is generated by

 $\Pi_{\sigma}$  as a set of simple roots, cf. [HK1, Section 2]. Clearly, when  $p \nmid m$ ,  $\mathfrak{g}^{\sigma}$  has the same root system as in the case of characteristic zero (with respect to the action of  $T^{\tau,\circ}$ ). Thus, the root system of  $\mathfrak{g}^{\sigma}$  (also  $G^{\sigma,\circ}$ ) is generated by  $\Pi_{\sigma}$  as a set of simple roots. By [Sp, Corollary 8.2.10],  $G^{\sigma,\circ}$  is generated by the root subgroups  $\{x_{\pm \alpha} \mid \alpha \in \Pi_{\sigma}\}$  and  $T^{\tau,\circ}$ . This proves part (1).

For any  $i \notin Y \cup \{o\}$ , by definition  $\langle h, \alpha_i \rangle = 0$ . Then,

$$\operatorname{Ad}_{t^{-h}}(x_{\pm \alpha_i}(a)) = x_{\pm \alpha_i}(a), \text{ for any } a \in \mathbf{k}.$$

If  $o \notin Y$ , then  $s_o = 0$ , equivalently  $\langle h, \theta_0 \rangle = \sum_{i \in I_\tau} a_i s_i = m/r$ . Then,

$$\operatorname{Ad}_{t^{-h}}(x_{\pm\theta_0}(a)) = x_{\pm\theta_0}(at^{\langle -h, \pm\theta_0 \rangle}) = x_{\pm\theta_0}(at^{\mp m/r}) = x_{\mp\alpha_0}(a).$$

Thus, part (2) follows from part (1) and the definition of  $M_Y$ .

Note that  $h \in X_*(T_{\mathrm{ad}})^{\tau}$  can be viewed as a cocharacter  $h : \mathbb{G}_m \to T_{\mathrm{ad}}^{\tau}$ . We define  $G_{\vartheta}$  to be the centralizer of h in  $G^{\tau,\circ}$ , i.e.

(15) 
$$G_{\vartheta} = \{ g \in G^{\tau,\circ} \mid \operatorname{Ad}_{h(s)}(g) = g, \text{ for any } s \in \mathbb{G}_m \}.$$

Here, we use the notation  $G_{\vartheta}$ , as this group only depends on  $\vartheta$ . In fact, it only depends on Y.

**Corollary 2.9.** Set  $Y' = Y \cup \{o\}$ . We have  $G_{\vartheta} = M_{Y'}$ . In particular, if  $o \in Y$ , then  $G_{\vartheta} = M_{Y'}$ ; if  $o \notin Y$ , then  $G_{\vartheta} \subseteq M_{Y}$ .

*Proof.* First of all, we have  $G_{\vartheta} = \operatorname{Ad}_{t^h}(G_{\vartheta}) \subseteq G^{\sigma,\circ}$ . Hence,  $G_{\vartheta} = \operatorname{Ad}_{t^{-h}}(G_{\vartheta}) \subseteq M_Y$ .

When  $o \in Y$ ,  $M_Y$  is generated by  $\{x_{\pm \alpha_i}(a) \mid i \notin Y, a \in k\}$  and  $T^{\tau,\circ}$ , which is contained in  $G_{\vartheta}$ . Therefore,  $G_{\vartheta} = M_Y = M_{Y'}$ .

When  $o \notin Y$ , i.e.  $s_o = 0$ , we have  $\langle \vartheta, \theta_0 \rangle = 1$ . Let  $q > \frac{m}{r}$  be a prime number so that  $\frac{1}{q}h \notin X_*(T_{ad})^{\mathsf{T}}$  and  $q \neq p$ . Set  $\vartheta' := \frac{1}{q}h$ . We claim that q is the minimal positive integer so that  $q\vartheta' \in X_*(T_{ad})^{\mathsf{T}}$ . Suppose  $a\vartheta' \in X_*(T_{ad})^{\mathsf{T}}$  for some integer 0 < a < q. Pick integers u, v so that uq + va = 1. Then,  $\vartheta' = (uq + va)\vartheta' = uh + va\vartheta' \in X_*(T_{ad})^{\mathsf{T}}$ , which is a contradiction. Hence, q is the minimal positive integer so that  $q\vartheta' \in X_*(T_{ad})^{\mathsf{T}}$ . Therefore, m' := qr is the minimal positive multiple of r so that  $\frac{m'}{r}\vartheta' \in X_*(T_{ad})^{\mathsf{T}}$ . Since  $h' := \frac{m'}{r}\vartheta' = h$ , we conclude that  $G_{\vartheta'} = G_{\vartheta}$ . Note that  $\langle \vartheta', \theta_0 \rangle = \frac{1}{q} \cdot \frac{m}{r} < 1$ . The element  $\vartheta'$  is a rational interior point of  $F_{Y'}$ . By the first case, we have  $G_{\vartheta'} = M_{Y'}$  (the condition  $p \nmid m'$  clearly holds). Therefore,  $G_{\vartheta} = M_{Y'}$ . In this case, we have  $M_{Y'} \subsetneq M_Y$ , see Proposition 2.8.

2.4. **A generation theorem.** The following theorem will be crucially used in the proof of Proposition 3.12 and also in the proof of Theorem 5.2.

**Theorem 2.10.** Following the same notations as in Definition 2.1, we further assume that  $p \nmid |\pi_1(G)|$ . For any point  $\vartheta \in X_*(T)^{\tau}_{\mathbb{Q}}$ , the parahoric subgroup  $\mathcal{P}_{\vartheta}$  is generated by  $T^{\tau,\circ}$  and  $\{\mathscr{U}_{\alpha,\neg(\vartheta,\alpha)}\}_{\alpha\in\mathbb{R}}$ .

*Proof.* We first consider the case when G is simply-connected. It suffices to show that the kernel  $K := \ker(T(O)^{\tau} \to T^{\tau})$  can be generated by  $\{\mathcal{U}_{\alpha, \neg(\vartheta, \alpha)}\}_{\alpha \in \mathcal{R}}$ . Let  $\{\tilde{\alpha}_j \in R \mid j \in I\}$ 

be the set of simple roots of G. There is an isomorphism  $\prod_{j \in I} \mathbb{G}_m \xrightarrow{11\tilde{\alpha}_j^{\vee}} T$ . Set

$$O_j = \begin{cases} \bar{O} & \text{if } \tau(j) = j, \\ O & \text{if } \tau(j) \neq j. \end{cases}$$

For any  $j \in I$  and  $u \in O_j^{\times}$ , we define the following element in  $T(O)^{\tau}$ ,

$$h_j(u) = \begin{cases} \tilde{\alpha}_j^{\vee}(u) & \text{if } \tau(j) = j, \\ \tilde{\alpha}_j^{\vee}(u) \cdots \tilde{\alpha}_{\tau^{r-1}(j)}^{\vee}(\tau^{r-1}(u)) & \text{if } \tau(j) \neq j. \end{cases}$$

Then, the kernel *K* is generated by

$$\{h_i(u) \mid j \in I, u = 1 + u' \in O_i^{\times}, \text{val}(u') > 0\}.$$

For any  $j \in I$ , let  $\alpha_j$  be the image of  $\tilde{\alpha}_j$  under the map  $R \to \mathcal{R}$ . We claim that  $h_j(u)$  belongs to the subgroup generated by  $\{\mathscr{U}_{\alpha, -\langle \vartheta, \alpha \rangle}, \mathscr{U}_{-\alpha, -\langle \vartheta, -\alpha \rangle} \mid \alpha \text{ is a multiple of } \alpha_j\}$ .

**Case 1:** When j and  $\tau(j)$  are not adjacent or  $\tau(j) = j$ , we only need to show that  $\tilde{\alpha}_j^{\vee}(u)$  can be write as a product of  $x_{\pm \tilde{\alpha}_j}(v)$  with  $v \in O_j$  and  $\operatorname{val}(v) \ge -\langle \vartheta, \pm \alpha_j \rangle$ . This reduces to the computation on  $\operatorname{SL}_2$ :

$$\begin{pmatrix} u & 0 \\ 0 & u^{-1} \end{pmatrix} = \begin{pmatrix} 1 & -z^{-n} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -u'z^n & 1 \end{pmatrix} \begin{pmatrix} 1 & u^{-1}z^{-n} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ uu'z^n & 1 \end{pmatrix}.$$

Hence,

$$\tilde{\alpha}_j^\vee(u) = x_{\tilde{\alpha}_j}(-z^{-n})x_{-\tilde{\alpha}_j}(-u'z^n)x_{\tilde{\alpha}_j}(u^{-1}z^{-n})x_{-\tilde{\alpha}_j}(uu'z^n).$$

We take *n* to be the integer so that  $n \le \langle \vartheta, \alpha_i \rangle < n + 1$ . Then,

$$\operatorname{val}(uu'z^n) = \operatorname{val}(-u'z^n) \ge n+1 > -\langle \vartheta, -\alpha_i \rangle, \quad \operatorname{val}(u^{-1}z^{-n}) = -n \ge -\langle \vartheta, \alpha_i \rangle.$$

It follows that  $h_j(u)$  belongs to the subgroup generated by  $\{\mathscr{U}_{\alpha_j,-\langle\vartheta,\alpha_j\rangle},\mathscr{U}_{-\alpha_j,-\langle\vartheta,-\alpha_j\rangle}\}$ .

**Case 2:** When j and  $\tau(j)$  are adjacent, we first recall that  $\mathscr{U}_{\alpha_j, -\langle \vartheta, \alpha_j \rangle}$  (resp.  $\mathscr{U}_{-\alpha_j, -\langle \vartheta, -\alpha_j \rangle}$ ) is isomorphic to the following subgroup of  $SL_3(O)$ , cf. [BT],

$$(16) \quad \left\{ \begin{pmatrix} 1 & w & v \\ 0 & 1 & \tau(w) \\ 0 & 0 & 1 \end{pmatrix} \middle| v + \tau(v) = w\tau(w), \operatorname{val}(w) \ge -\langle \vartheta, \alpha_j \rangle \right\} \quad (\text{resp.} \begin{pmatrix} 1 & 0 & 0 \\ w & 1 & 0 \\ v & \tau(w) & 1 \end{pmatrix}).$$

and  $\mathscr{U}_{2\alpha_j,-\langle\vartheta,2\alpha_j\rangle}$  (resp.  $\mathscr{U}_{-2\alpha_j,-\langle\vartheta,-2\alpha_j\rangle}$ ) is isomorphic to the following subgroup of  $SL_3(O)$ ,

(17) 
$$\left\{ \begin{pmatrix} 1 & 0 & v \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \middle| v + \tau(v) = 0, \text{val}(v) \ge -\langle \vartheta, 2\alpha_j \rangle \right\} \quad (\text{resp.} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ v & 0 & 1 \end{pmatrix}).$$

This reduces our claim to the computation on SL<sub>3</sub>. Note that u = 1 + u' can be written as  $u = (1 + u_1)(1 + u_2)$ , where  $u_1 = (u' + \tau(u'))/2$  is even,  $val(u_1) > 0$ , and  $u_2 = \frac{u' - \tau(u')}{2 + 2u_1}$  is odd. Then, we can write  $h_j(u) = h_j(1 + u_1)h_j(1 + u_2)$ . Therefore, it suffices to consider the following two cases.

Case 2.1: For any  $u = 1 + u' \in O^{\times}$  with  $\tau(u') = u'$  and val(u') > 0, we have

$$(18) \quad \begin{pmatrix} u & 0 & 0 \\ 0 & u^{-1} & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & \tau(u) & 0 \\ 0 & 0 & \tau(u)^{-1} \end{pmatrix} = \begin{pmatrix} 1 & 0 & -z^{1-2n} \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -u'z^{2n-1} & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & z^{1-2n}u^{-1} \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ uu'z^{2n-1} & 0 & 1 \end{pmatrix}.$$

Choose n to be the integer so that  $2n-1 \le \langle \vartheta, 2\alpha_i \rangle < 2n+1$ . Since val $(u') \ge 2$ , we have

$$\operatorname{val}(uu'z^{2n-1}) = \operatorname{val}(-u'z^{2n-1}) \geq 2n+1 > -\langle \vartheta, -2\alpha_j \rangle, \quad \operatorname{val}(z^{1-2n}u^{-1}) = 1-2n \geq -\langle \vartheta, 2\alpha_j \rangle.$$

Thus, the matrices on the right side of decomposition (18) belong to (17). Hence,  $h_j(u) = \tilde{\alpha}_j^{\vee}(u)\tilde{\alpha}_{\tau(j)}^{\vee}(\tau(u))$  belongs to the subgroup generated by  $\{\mathscr{U}_{2\alpha_j,-\langle\vartheta,-2\alpha_j\rangle},\mathscr{U}_{-2\alpha_j,-\langle\vartheta,-2\alpha_j\rangle}\}$ .

Case 2.2: For any  $u = 1 + u' \in O^{\times}$  with  $\tau(u') = -u'$ , we have  $\tau(u) = 1 - u'$  and

$$\begin{pmatrix} u & 0 & 0 \\ 0 & u^{-1} & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & \tau(u) & 0 \\ 0 & 0 & \tau(u)^{-1} \end{pmatrix}$$

$$\stackrel{\bigoplus}{=} \begin{pmatrix} 1 & -\sqrt{2}z^{-n} & (-1)^n z^{-2n} \\ 0 & 1 & -\sqrt{2}(-z)^{-n} \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ (-1)^n z^{2n} u' & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & \sqrt{2}z^{-n} u^{-1} & (-1)^n z^{-2n} \tau(u)^{-1} \\ 0 & 1 & \sqrt{2}(-z)^{-n} \tau(u)^{-1} \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ \sqrt{2}z^n u' u \tau(u)^{-1} & 1 & 0 \\ (-1)^{n+1} z^{2n} u u' & -\sqrt{2}(-z)^n u' u^{-1} \tau(u) & 1 \end{pmatrix} .$$

$$\stackrel{\bigoplus}{=} \begin{pmatrix} 1 & \sqrt{2}z^{-n} u' u \tau(u)^{-1} & (-1)^{n+1} z^{-2n} u u' \\ 0 & 1 & -\sqrt{2}(-z)^{-n} u' u^{-1} \tau(u) \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ \sqrt{2}z^n u^{-1} & 1 & 0 \\ (-1)^n z^{2n} \tau(u)^{-1} & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & (-1)^n z^{-2n} u' \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & (-1)^n z^{-2n} u' \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & (-1)^n z^{-2n} u' \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & (-1)^n z^{-2n} u' \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & (-1)^n z^{-2n} u' \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & (-1)^n z^{-2n} u' \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & (-1)^n z^{-2n} u' \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & (-1)^n z^{-2n} u' \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & (-1)^n z^{-2n} u' \\ 0 & 1 & 0 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 & (-1)^n z^{-2n} u' \\ 0 & 1 & 0 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 & (-1)^n z^{-2n} u' \\ 0 & 1 & 0 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 & (-1)^n z^{-2n} u' \\ 0 & 1 & 0 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 & (-1)^n z^{-2n} u' \\ 0 & 1 & 0 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 & (-1)^n z^{-2n} u' \\ 0 & 1 & 0 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 & (-1)^n z^{-2n} u' \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 & (-1)^n z^{-2n} u' \\ 0 & 1 & 0 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 & (-1)^n z^{-2n} u' \\ 0 & 1 & 0 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 & (-1)^n z^{-2n} u' \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 & (-1)^n z^{-2n} u' \\ 0 & 1 & 0 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 & (-1)^n z^{-2n} u' \\ 0 & 1 & 0 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 & (-1)^n z^{-2n} u' \\ 0 & 1 & 0 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 & (-1)^n z^{-2n} u' \\ 0 & 1 & 0 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 & (-1)^n z^{-2n} u' \\ 0 & 1 & 0 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 & (-1)^n z^{-2n} u' \\ 0 & 1 & 0 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 & (-1)^n z^{-2n} u' \\ 0 & 1 & 0 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 & (-1)^n z^{-2n} u' \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 & (-1)^n z^{-2n} u' \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 & (-1)^$$

When  $2n \leq \langle \vartheta, 2\alpha_j \rangle < 2n+1$  for some integer n, the matrices in decomposition ① belong to (16) and (17). When  $2n-1 \leq \langle \vartheta, 2\alpha_j \rangle < 2n$  for some integer n, the matrices in decomposition ② belong to (16) and (17). Hence,  $h_j(u)$  belongs to the subgroup generated by  $\{\mathscr{U}_{\alpha_j, -\langle \vartheta, \alpha_j \rangle}, \mathscr{U}_{-\alpha_j, -\langle \vartheta, -\alpha_j \rangle}, \mathscr{U}_{2\alpha_j, -\langle \vartheta, 2\alpha_j \rangle}, \mathscr{U}_{-2\alpha_j, -\langle \vartheta, -2\alpha_j \rangle}\}$ . This proves our claim. Therefore, the kernel K is generated by  $\{\mathscr{U}_{\alpha, -\langle \vartheta, \alpha \rangle}\}_{\alpha \in \mathcal{R}}$ . This proves the lemma for simply-connected group G.

For general case, let  $\pi: G' \to G$  be the simply-connected cover of G. The morphism  $\pi$  is étale since  $p \nmid \pi_1(G)$ . Let T' be the preimage of T in G'. Furthermore,  $\pi: T' \to T$  is étale. Denote  $O_k := k[z]/(z^k)$  and  $\mathbb{D}_k := \operatorname{Spec}(O_k)$ . Let  $i: \operatorname{Spec}(k) \to \mathbb{D}_k$  be the natural morphism. By [SP, Tag 02HF],  $\pi$  is formally étale, i.e. for any  $g: \mathbb{D}_k \to T$  so that  $g \circ i: \operatorname{Spec}(k) \to T$  is the identity  $e \in T(k)$ , there is a unique lifting  $\tilde{g}: \mathbb{D}_k \to T'$  so that  $\tilde{g} \circ i$  is the identity  $e' \in T'(k)$ . Thus,  $K \simeq K'$ , where  $K := \ker(T(O) \to T)$  and  $K' := \ker(T'(O) \to T')$ . Hence,

$$\ker(T(O)^{\tau} \to T^{\tau}) = K^{\tau} \simeq K'^{\tau} = \ker(T'(O)^{\tau} \to T'^{\tau}).$$

By the proof above in the simply-connected case,  $K'^{\tau}$  is generated by  $\{\mathscr{U}'_{\alpha,-\langle\vartheta,\alpha\rangle}\}_{\alpha\in\mathcal{R}}$ . Moreover,  $\mathscr{U}_{\alpha,-\langle\vartheta,\alpha\rangle} \simeq \mathscr{U}'_{\alpha,-\langle\vartheta,\alpha\rangle}$  for any  $\alpha\in\mathcal{R}$ . It follows that  $K^{\tau}$  is generated by  $\{\mathscr{U}_{\alpha,-\langle\vartheta,\alpha\rangle}\}_{\alpha\in\mathcal{R}}$ . Now, the lemma follows from the definition of  $\mathscr{P}_{\vartheta}$ .

## 3. A REFINEMENT OF THE COHERENCE CONJECTURE (AFTER ZHU)

3.1. **Statement of theorem.** In this section, we follow the same notations and assumption as in Section 2. We emphasize again that the automorphism  $\tau$  on G can be trivial.

Let W be the Weyl group of G with respect to the maximal torus T. Then, the subgroup  $W^{\tau}$  of  $\tau$ -fixed points is naturally isomorphic to the Weyl group of  $G^{\tau,\circ}$  with respect to  $T^{\tau,\circ}$ . Let  $X_*(T)_{\tau}$  be the lattice of  $\tau$ -coinvariants of  $X_*(T)$ .

**Definition 3.1.** Let  $\tilde{W}_{\rm aff}:=X_*(T)_{\tau}\rtimes W^{\tau}$  be the extended affine Weyl group. The elements in  $\tilde{W}_{\rm aff}$  are written as  $\varrho^{\bar{\mu}}w$  for any  $\bar{\mu}\in X_*(T)_{\tau}$  and  $w\in W^{\tau}$ , where  $\varrho^{\bar{\mu}}$  represents the translation on the apartment of  $(\mathcal{G}_{\bar{K}},S)$  by  $-\bar{\mu}$ .

Let G' be the simply-connected cover of G with the maximal torus T' mapping to T. Then,  $W_{\rm aff}:=X_*(T')_{\tau}\rtimes W^{\tau}$  is an affine Weyl group, cf. [DH1, A.1]. Let  $\Omega$  denote the group of length zero elements in  $\tilde{W}_{\rm aff}$ . Then,  $\Omega\simeq X_*(T)_{\tau}/X_*(T')_{\tau}$  and  $\tilde{W}_{\rm aff}=W_{\rm aff}\rtimes\Omega$ . Moreover, the group  $\Omega$  acts on  $\hat{I}_{\tau}$ .

Let  $w_0$  be the longest element in  $W^{\tau}$ . Any element in  $\Omega$  can be written as  $\varsigma_{\kappa} := \varrho^{-\kappa} w_{\kappa} w_0$ , where  $\kappa$  is 0 or a miniscule dominant element in  $X_*(T)_{\tau}$  (cf. [BH2, Section 2.3]), and  $w_{\kappa}$  is the longest word in the stabilizer  $W_{\kappa}^{\tau}$  of  $\kappa$  in  $W^{\tau}$ , cf. [Bou, Chapter 6, Section 2.3]

For any nonempty subset  $Y \subset \hat{I}_{\tau}$ , let  $\mathcal{P}_{Y} \subseteq G(\mathcal{K})^{\tau}$  be the parahoric subgroup associated to Y, cf. Definition 2.7. Let  $\mathcal{G}_{Y}$  denote the corresponding parahoric group scheme over  $\bar{O}$ . When  $Y = \hat{I}_{\tau}$ ,  $\mathcal{P}_{Y}$  is the Iwahori subgroup of  $G(\mathcal{K})^{\tau}$ , denoted by I.

**Definition 3.2.** We define the partial affine flag variety

$$\operatorname{Fl}_Y := L\mathscr{G}_Y/L^+\mathscr{G}_Y,$$

where  $L\mathscr{G}$  (resp.  $L^+\mathscr{G}$ ) is the loop group scheme (resp. jet group scheme) of  $\mathscr{G}$ . Then,  $\mathrm{Fl}_Y(k) = G(\mathcal{K})^\tau/\mathcal{P}_Y$ . In particular, when  $\tau$  is trivial and  $Y = \{o\}$ ,  $\mathrm{Fl}_Y$  is the affine Grassmannian of G, denoted by  $\mathrm{Gr}_G$ . Let  $e_0$  denote the base point in  $\mathrm{Fl}_Y$  and  $\mathrm{Gr}_G$ 

Set  $\bar{Y} =: \hat{I}_{\tau} \setminus Y$ , and let  $W_{\bar{Y}}$  denote the parabolic subgroup of  $W_{\text{aff}}$  generated by the set of simple reflections  $\{r_i \mid i \in \bar{Y}\}$ .

**Definition 3.3.** For any  $w \in \widetilde{W}_{aff}/W_{\bar{Y}}$ , define the Schubert cell  $\mathrm{Fl}_{Y,w} := I\dot{w}e_0$ , and the Schubert variety  $\overline{\mathrm{Fl}}_{Y,w}$  is the closure of  $\mathrm{Fl}_{Y,w}$ , where  $e_0$  is the based point in  $\mathrm{Fl}_Y$  and  $\dot{w}$  is a lift of w in  $G(\mathcal{K})^{\tau}$ . When  $Y = \hat{I}_{\tau}$ , we simply use the notations  $\mathrm{Fl}_w$  and  $\overline{\mathrm{Fl}}_w$ .

For any dominant coweight  $\mu \in X_*(T)^+$ , we define the affine Schubert variety  $\overline{\text{Gr}}_{G,\mu}$  to the closure of the orbit  $G(O) \cdot e_{\mu}$  in  $\text{Gr}_G$ , where  $e_{\mu} := z^{\mu}e_0$ .

Given any  $\bar{\mu} \in X_*(T)_\tau$ , we simply denote by  $\mathrm{Fl}_{Y,\bar{\mu}}$  (resp.  $\overline{\mathrm{Fl}}_{Y,\bar{\mu}}$ ) the Schubert cell  $\mathrm{Fl}_{Y,\varrho^{\bar{\mu}}W_{\bar{\gamma}}}$  (resp.  $\overline{\mathrm{Fl}}_{Y,\varrho^{\bar{\mu}}W_{\bar{\nu}}}$ ). In fact, a lift of  $\varrho^{\bar{\mu}}$  can be taken to be the following element in  $T(\mathcal{K})^\tau$ ,

(19) 
$$n^{\mu} := \prod_{i=0}^{r-1} \tau^i(z^{\mu}) \in T(\mathcal{K})^{\tau},$$

where  $\mu$  is a lifting of  $\bar{\mu}$  in  $X_*(T)$ . Then,  $\mathrm{Fl}_{Y,\bar{\mu}}$  is exactly the orbit  $\mathcal{I} \cdot e_{\bar{\mu}}$  in  $\mathrm{Fl}_Y$ , where  $e_{\bar{\mu}} := n^{\mu}e_0$ . We define the following union of Schubert varieties in  $\mathrm{Fl}_Y$ ,

(20) 
$$\mathcal{A}_{Y}(\bar{\mu}) := \bigcup_{w \in W^{\tau}} \overline{\mathrm{Fl}}_{Y,w(\bar{\mu})}.$$

We now recall the admissible set of affine Weyl group, cf. [Zh1, 2.1.6]. For  $\mu \in X_*(T)$ , define

$$Adm(\mu) := \{ w \in \tilde{W}_{aff} \mid w \leq \rho^{\eta} \text{ for some } \eta \in W^{\tau} \bar{\mu} \}.$$

Let  $\mathrm{Adm}^Y(\mu)$  denote the subset  $W_{\bar{Y}}\mathrm{Adm}(\mu)W_{\bar{Y}} \subset \tilde{W}_{\mathrm{aff}}$ . For any  $w \in \tilde{W}_{\mathrm{aff}}$ , let  ${}_{Y}\mathrm{Fl}_{Y,w}$  denote the  $\mathcal{P}_{Y}$ -orbit through  $\dot{w}$  in  $\mathrm{Fl}_{Y}$  and let  ${}_{Y}\overline{\mathrm{Fl}}_{Y,w}$  be the closure of  ${}_{Y}\mathrm{Fl}_{Y,w}$  in  $\mathrm{Fl}_{Y}$ . In the following proposition, we show that  $\mathcal{A}_{Y}(\bar{\mu})$  defined in (20) agrees with the definition in [Zh1, 2.2.2] in terms of admissible set.

**Lemma 3.4.** For any  $\mu \in X_*(T)$ , let  $\bar{\mu}$  be the image of  $\mu$  in  $X_*(T)_{\tau}$ . We have

$$\mathcal{A}_{Y}(\bar{\mu}) = \bigcup_{w \in \mathrm{Adm}^{Y}(\mu)} {}_{Y}\overline{\mathrm{Fl}}_{Y,w}.$$

*Proof.* Denote the right side by  $X_Y(\mu)$ . For any  $w \in W^{\tau}$ , by definition, we have  $\operatorname{Fl}_{Y,w(\bar{\mu})} \subseteq {}_Y\operatorname{Fl}_{Y,o^{w(\bar{\mu})}}$ . Hence,  $\overline{\operatorname{Fl}}_{Y,w(\bar{\mu})} \subseteq {}_Y\overline{\operatorname{Fl}}_{Y,o^{w(\bar{\mu})}} \subseteq X_Y(\mu)$ . This shows that  $\mathcal{A}_Y(\bar{\mu}) \subseteq X_Y(\mu)$ .

Conversely, for any  $w \in \operatorname{Adm}^{Y}(\underline{\mu})$ , we write  $w = w_1 w' w_2$ , where  $w_1, w_2 \in W_{\bar{Y}}$  and  $w' \leq \varrho^{\eta}$  for some  $\eta \in W^{\tau}\bar{\mu}$ . Then,  ${}_{Y}\overline{\operatorname{Fl}}_{Y,w} = {}_{Y}\overline{\operatorname{Fl}}_{Y,w'} \subseteq {}_{Y}\overline{\operatorname{Fl}}_{Y,\varrho^{\eta}}$ . We have

(21) 
$$\mathcal{P}_{Y}\varrho^{\eta}\mathcal{P}_{Y} = \bigcup_{v \in W_{\bar{v}}} I_{y}I_{\varrho}^{\eta}\mathcal{P}_{Y} \stackrel{(*)}{\subseteq} \bigcup_{v \in W_{\bar{v}}} I_{y}\varrho^{\eta}\mathcal{P}_{Y} = \bigcup_{v \in W_{\bar{v}}} I_{y}\varrho^{\eta}y^{-1}\mathcal{P}_{Y}.$$

The inclusion (\*) follows from the fact that  $IyI = Is_{i_1}I \cdots Is_{i_k}I$  for reduced word  $y = s_{i_1} \cdots s_{i_k}$  and  $Is_iII\varrho^{\eta}I \subseteq I\varrho^{\eta}I \cup Is_i\varrho^{\eta}I$  for any simple reflection  $s_i \in W_{\text{aff}}$ , cf. [Ku2, Theorem 5.1.3(d)]. For any  $y \in W_{\bar{Y}}$ , we can write  $y = y'\varrho^{\gamma}$  for some  $y' \in W^{\tau}$ ,  $\gamma \in X_*(T)_{\tau}$ . Then,  $y\varrho^{\eta}y^{-1} = y'\varrho^{\eta}y'^{-1} = \varrho^{y'(\eta)}$  and we have

$$Iy\varrho^{\eta}y^{-1}\mathcal{P}_Y=I\varrho^{y'(\eta)}\mathcal{P}_Y\subseteq\bigcup_{v\in W^\tau}I\varrho^{v(\eta)}\mathcal{P}_Y.$$

Hence,  $\mathcal{P}_{Y}\varrho^{\eta}\mathcal{P}_{Y}\subseteq\bigcup_{\nu\in W^{\tau}}I\varrho^{\nu(\eta)}\mathcal{P}_{Y}$ . Therefore,  ${}_{Y}\overline{\mathrm{Fl}}_{Y,w}\subseteq{}_{Y}\overline{\mathrm{Fl}}_{Y,\varrho^{\eta}}\subseteq\bigcup_{\nu\in W^{\tau}}{}_{Y}\overline{\mathrm{Fl}}_{Y,\varrho^{\nu(\eta)}}=\mathcal{A}_{Y}(\bar{\mu})$ . This shows that  $X_{Y}(\mu)\subseteq\mathcal{A}_{Y}(\bar{\mu})$ .

For any  $\varsigma_{\kappa} \in \Omega$ , let  $\operatorname{Fl}_{Y}^{\kappa}$  denote the connected component of  $\operatorname{Fl}_{Y}$  containing  $\varsigma_{\kappa}$  (equivalently  $\varrho^{-\kappa}$ ). Then,  $\operatorname{Fl}_{Y}^{\kappa}$  is a homogeneous space of  $G'(\mathcal{K})^{\tau}$ , and

(22) 
$$\operatorname{Fl}_{Y}^{\kappa} \simeq G'(\mathcal{K})^{\tau}/\mathcal{P}_{Y_{\kappa}}' =: \operatorname{Fl}_{Y_{\kappa}}',$$

where  $\mathcal{P}'_{Y_{\kappa}}$  is the parahoric subgroup of  $G'(\mathcal{K})^{\tau}$  associated to  $Y_{\kappa} := \{\varsigma_{\kappa}(i) \mid i \in Y\}$ . By [PR, Section 10], we have the following isomorphism,

(23) 
$$\operatorname{Pic}(\operatorname{Fl}_{Y}^{\kappa}) \simeq \bigoplus_{i \in Y_{\kappa}} \mathbb{Z}\Lambda_{i}.$$

Given any  $\bar{\mu} \in X_*(T)_\tau$ , let  $\varsigma_{\kappa} \in \Omega$  be the unique element such that  $\varrho^{\bar{\mu}} \varsigma_{\kappa}^{-1} \in W_{\text{aff}}$  (equivalently  $\bar{\mu} + \kappa \in X_*(T')_\tau$ ). Then, for any  $w \in W^\tau$ , we have  $w(\bar{\mu}) + \kappa \in X_*(T')_\tau$ , equivalently  $\varrho^{w(\bar{\mu})} \varsigma_{\kappa}^{-1} \in W_{\text{aff}}$ . Hence,  $\operatorname{Fl}_{Y,w(\bar{\mu})} \subseteq \operatorname{Fl}_Y^{\kappa}$  for any  $w \in W^\tau$ . Let  $\dot{\varsigma}_{\kappa}$  be a lifting of  $\varsigma_{\kappa}$  in  $G(\mathcal{K})^\tau$ . Consider the translation

$$\dot{\varsigma}_{\kappa}: \mathrm{Fl}_{\nu}^{\circ} \to \mathrm{Fl}_{\nu}^{\kappa}$$

given by  $x \mapsto \dot{\varsigma}_{\kappa} x$ . For any  $w \in W^{\tau}$ , the orbit  $I(\dot{\varsigma}_{\kappa})^{-1} n^{w(\bar{\mu})} e_0$  maps to the orbit  $In^{w(\bar{\mu})} e_0$ . Thus, we have an isomorphism  $\overline{Fl}_{Y,\varsigma_{\nu}^{-1}\rho^{w(\bar{\mu})}} \simeq \overline{Fl}_{Y,w(\bar{\mu})}$  under the translation map.

Let  $\tilde{\varsigma}_{\kappa}$  be a lifting of  $\dot{\varsigma}_{\kappa}$  in  $G'(\mathcal{K})^{\tau}$ . Consider the conjugation map  $\mathrm{Ad}_{\tilde{\varsigma}_{\kappa}}: G'(\mathcal{K})^{\tau} \to G'(\mathcal{K})^{\tau}$  given by  $x \mapsto \tilde{\varsigma}_{\kappa} x(\tilde{\varsigma}_{\kappa})^{-1}$ . It induces an isomorphism  $\mathrm{Ad}_{\tilde{\varsigma}_{\kappa}}: \mathrm{Fl}'_{Y} \to \mathrm{Fl}'_{Y_{\kappa}}$ . Then, (22) is the composition of the following isomorphisms

$$\operatorname{Fl}_Y^{\kappa} \xrightarrow{(\dot{\varsigma}_{\kappa})^{-1}} \operatorname{Fl}_Y^{\circ} \simeq \operatorname{Fl}_Y' \xrightarrow{\operatorname{Ad}_{\bar{\varsigma}_{\kappa}}} \operatorname{Fl}_{Y_{\kappa}}'.$$

Under these isomorphisms, we have  $\overline{\mathrm{Fl}}_{Y,w(\bar{\mu})} \simeq \overline{\mathrm{Fl}}_{Y,\varsigma_{\kappa}^{-1}\varrho^{w(\bar{\mu})}} \simeq \overline{\mathrm{Fl}}_{Y,\varsigma_{\kappa}^{-1}\varrho^{w(\bar{\mu})}} \simeq \overline{\mathrm{Fl}}_{Y_{\kappa},\varrho^{w(\bar{\mu})}\varsigma_{\kappa}^{-1}}.$  Therefore, (22) restricts to an isomorphism

(25) 
$$\overline{\mathrm{Fl}}_{Y,w(\bar{\mu})} \simeq \overline{\mathrm{Fl}}'_{Y_{\kappa},\mathcal{O}^{w(\bar{\mu})}S_{\kappa}^{-1}}.$$

Hence, the variety  $\mathcal{A}_Y(\bar{\mu})$  maps to  $\bigcup_{w \in W^{\tau}} \overline{\mathrm{Fl}}'_{Y_{\kappa},\mathcal{Q}^{w(\bar{\mu})}S_{\kappa}^{-1}}$  under the map (22).

Let  $\mathcal{L}(\Lambda)$  be an ample line bundle on  $\operatorname{Fl}_Y^{\kappa}$  associated to  $\Lambda = \sum_{i \in Y_{\kappa}} n_i \Lambda_i$ , i.e.  $n_i > 0$  for any  $i \in Y_{\kappa}$ . Let  $c := \Lambda(K)$  be the level of  $\Lambda$  and let  $\lambda$  be the restriction of  $\Lambda$  to  $(T^{\tau,\circ})'$ .

Suppose that  $p \nmid a_{Y_{\kappa}}$ , where  $a_{Y_{\kappa}}$  is defined in (10) for the subset  $Y_{\kappa} \subset \hat{I}_{\tau}$ . By Lemma 2.6, we can choose an interior point  $\vartheta$  in the facet associated to  $Y_{\kappa}$  so that  $p \nmid m$ . Let  $G_{\vartheta}$  be the subgroup of  $G^{\tau,\circ}$  associated to  $\vartheta$  defined in (15) of Section 2.3. Let  $G'_{\vartheta}$  (resp.  $(T^{\tau,\circ})'$ ) be the pre-image of  $G_{\vartheta}$  (resp.  $T^{\tau,\circ}$ ) via the simply-connected cover map  $(G^{\tau,\circ})' \to G^{\tau,\circ}$ .

In Theorem 3.15, we will show that the level one line bundle L on  $Gr_G$  has a unique G'(O)-equivariant structure and the line bundle  $\mathcal{L}(\Lambda)$  on  $Fl_Y^{\kappa}$  has a unique  $(G(O)^{\tau,\circ})'$ -equivariant structure. These equivariant structures induce  $G'_{\vartheta}$ -equivariant structures on L and  $\mathcal{L}(\Lambda)$  via the composition of natural morphisms

$$G'_{\vartheta} \hookrightarrow (G^{\tau,\circ})' \to G' \hookrightarrow G'(O)$$

and the composition map  $G'_{\vartheta} \hookrightarrow (G^{\tau,\circ})' \to (G(O)^{\tau,\circ})'$  respectively.

We now state our main result of this section, which will be proved in the rest part.

**Theorem 3.5.** Let  $\mu \in X_*(T)^+$ . Let  $\mathcal{L}(\Lambda)$  be an ample line bundle on  $\mathrm{Fl}_Y^{\kappa}$  of central charge c > 0 such that  $\overline{\mathrm{Fl}}_{Y,\bar{\mu}}$  is contained in  $\mathrm{Fl}_Y^{\kappa}$  (equivalently  $\varrho^{\bar{\mu}} \varsigma_{\kappa}^{-1} \in W_{\mathrm{aff}}$ ).

(1) Assume  $p \nmid a_{Y_{\kappa}}$ , we have an isomorphism of  $(T^{\tau,\circ})'$ -representations:

$$H^0(\mathcal{A}_Y(\bar{\mu}), \mathcal{L}(\Lambda)) \simeq H^0(\overline{\mathrm{Gr}}_{G,\mu}, L^c) \otimes \mathrm{k}_{-\lambda},$$

where  $k_{-\lambda}$  is the 1-dimensional representation of  $(T^{\tau,\circ})'$ .

(2) Assume p = 0, we have an isomorphism of  $G'_{\vartheta}$ -representations:

$$H^0(\mathcal{A}_Y(\bar{\mu}), \mathcal{L}(\Lambda)) \simeq H^0(\overline{\mathrm{Gr}}_{G,\mu}, L^c) \otimes \mathrm{k}_{-\lambda}.$$

3.2. Equivariance of line bundles on partial affine flag varieties. In this subsection, we assume G is simply-connected.

As in Section 2.3, we fix a non-empty subset  $Y \subseteq \hat{I}_{\tau}$  and assume  $p \nmid a_Y$ . Then, we can choose an interior point  $\vartheta$  in the facet associated to Y so that  $p \nmid m$ , where m is the minimal positive multiple of r such that  $h := \frac{m}{r}\vartheta \in X_*(T_{ad})^{\tau}$ . Define an automorphism  $\sigma := \tau \circ \operatorname{Ad}_{(\epsilon_m)^{m\vartheta/r}}$  of G, whose order is m, see Lemma 2.3.

Recall the groups  $\mathcal{P}_{Y}^{-}$  and  $\mathcal{I}^{-}$  in Definition 2.7 and the parahoric group scheme  $\mathscr{G}_{-\vartheta,\infty}$  over  $\bar{\mathcal{O}}_{\infty}$  defined in Section 2.2.

**Proposition 3.6.** The isomorphism  $\mathrm{Ad}_{t^h}: G((z^{-1}))^{\tau} \simeq G((t^{-1}))^{\sigma}$  in equation (9) restricts to an isomorphism

(26) 
$$\operatorname{Ad}_{t^{h}}: \mathcal{P}_{Y}^{-} \simeq G[t^{-1}]^{\sigma}.$$

As a consequence,  $\mathcal{P}_{Y}^{-}$  has an integral ind-group scheme structure and  $\mathcal{P}_{Y}^{-} = \mathscr{G}_{-\vartheta,\infty}(\bar{O}_{\infty}) \cap G[z,z^{-1}]^{\mathsf{T}}$ .

*Proof.* It is clear that  $L^-\mathscr{G}_\sigma:=G[t^{-1}]^\sigma$  admits a natural ind-group scheme structure. Define  $L^{--}\mathscr{G}_\sigma$  to be

$$L^{--}\mathscr{G}_{\sigma} := \ker(\operatorname{ev}_{\infty} : L^{-}\mathscr{G}_{\sigma} \to G^{\sigma}).$$

By Theorem 2.4 and [PR, Theorem 0.1, Theorem 0.2], the twisted affine Grassmannian  $Gr_{\mathscr{G}_{\sigma}} := G((t))^{\sigma}/G[[t]]^{\sigma}$  is reduced and irreducible. Moreover, by [HR, Lemma 3.1] the morphism  $L^{--}\mathscr{G}_{\sigma} \to Gr_{\mathscr{G}_{\sigma}}$  given by  $g \mapsto g \cdot e$  is an open embedding. Thus,  $L^{--}\mathscr{G}_{\sigma}$  is integral. Since G is simply-connected,  $G^{\sigma}$  is connected. Therefore,  $G[t^{-1}]^{\sigma}$  is integral.

By definition of  $\mathcal{P}_{Y}^{-}$ ,  $\operatorname{Ad}_{t^{h}}(\mathcal{P}_{Y}^{-})$  is contained in  $G[t^{-1}]^{\sigma}$ . Note that,  $B[t^{-1}]^{\sigma}U^{-}[t^{-1}]^{\sigma}$  is open in  $G[t^{-1}]^{\sigma}$ , and  $B[t^{-1}]^{\sigma}$  and  $U^{-}[t^{-1}]^{\sigma}$  are both contained in  $\operatorname{Ad}_{t^{h}}(\mathcal{P}_{Y}^{-})$ . Thus,  $\operatorname{Ad}_{t^{h}}(\mathcal{P}_{Y}^{-}) = G[t^{-1}]^{\sigma}$ . The group  $\mathcal{P}_{Y}^{-}$  naturally inherits an integral ind-group scheme structure from  $G[t^{-1}]^{\sigma}$ .

Remark 3.7. In the above proposition, the ind-group scheme structure on  $\mathcal{P}_{Y}^{-}$  relies on the condition  $p \nmid a_{Y}$ . It is possible that this condition can be removed. However, we don't know an appropriate reference.

Recall that  $M_Y = \operatorname{Ad}_{t^{-h}}(G^{\sigma})$ . As a corollary of Theorem 2.4 and Proposition 3.6, we have

**Corollary 3.8.** 
$$M_Y$$
 is a reductive algebraic group and  $M_Y = \mathcal{P}_Y \cap \mathcal{P}_Y^-$ .

Consider the full affine flag variety  $\mathrm{Fl}=G(\mathcal{K})^{\mathrm{T}}/I$  and the natural morphism  $p_Y:\mathrm{Fl}\to\mathrm{Fl}_Y.$  Let  $\mathbb{P}^1_j$  denote the image of the morphism  $\mathbb{P}^1\simeq\mathcal{P}_{\hat{I}_{\mathrm{T}}\setminus\{j\}}/I\to\mathrm{Fl}$ . For each  $i\in Y$ , let  $\mathscr{L}_i$  denote the line bundle  $\mathscr{L}(\Lambda_i)$  on  $\mathrm{Fl}_Y.$  Then, we have

$$\deg\left(p_Y^*(\mathscr{L}_i)\big|_{\mathbb{P}^1_i}\right) = \delta_{i,j}$$

for any  $i \in Y$  and  $j \in \hat{I}_{\tau}$ , cf. [PR, Section 10].

Let  $e_0$  denote the base point of  $Fl_Y$ . We have the following decompositions

(27) 
$$\operatorname{Fl}_{Y} = \bigsqcup_{w \in W_{\operatorname{aff}}/W_{\bar{Y}}} I \dot{w} e_{0}, \quad \operatorname{Fl}_{Y} = \bigsqcup_{w \in W_{\operatorname{aff}}/W_{\bar{Y}}} I^{-} \dot{w} e_{0},$$

where  $\dot{w}$  is a lift of w in  $G(\mathcal{K})^{\tau}$ . In fact,  $\operatorname{Fl}_{Y}$  can be identified with the Kac-Moody flag variety  $\operatorname{Fl}_{Y}^{\operatorname{KM}}$  in the sense of [Ku1] and [Ma], cf. [PR, §9.f]. Thus, the decompositions in (27) arise from the decompositions for  $\operatorname{Fl}_{Y}^{\operatorname{KM}}$  in [Ku1] and [Ma].

For any  $w \in W_{\text{aff}}$ , let  $Fl_Y^w$  denote the  $I^-$ -orbit  $I^-we_0$  in  $Fl_Y$  and let  $\overline{Fl}_Y^w$  denote the closure of  $Fl_Y^w$ . When  $Y = \hat{I}_\tau$ , we simply use the notations  $Fl_Y^w$  and  $\overline{Fl}_Y^w$ .

**Lemma 3.9.**  $T^-ye_0 \subseteq \overline{Fl}_Y^w$  if and only if  $w_{\min} \leq y_{\min}$ , where  $w_{\min}$  represents the minimal element in the left coset  $wW_{\bar{Y}}$  (similarly for  $y_{\min}$ ).

*Proof.* When char(k)=0, it follows from [Ku2, Prop.7.1.21]. For general characteristic, it follows from [Kat, Theorem 1.26, Lemma 2.9]. □

**Lemma 3.10.**  $\overline{\operatorname{Fl}}_{Y}^{w} \cap \operatorname{Fl}_{Y,y} \neq \emptyset$  if and only if  $w_{\min} \leq y_{\min}$ .

*Proof.* Assume  $w_{\min} \leq y_{\min}$ . By Lemma 3.9, we have  $I^-ye_0 \subseteq \overline{\operatorname{Fl}}_Y^w$ . Hence,  $\overline{\operatorname{Fl}}_Y^w \cap \operatorname{Fl}_{Y,y} \supseteq I^-ye_0 \cap \operatorname{Fl}_{Y,y} \ni ye_0$ . Therefore,  $\overline{\operatorname{Fl}}_Y^w \cap \operatorname{Fl}_{Y,y} \neq \emptyset$ . Conversely, assume  $\overline{\operatorname{Fl}}_Y^w \cap \operatorname{Fl}_{Y,y} \neq \emptyset$ ,

then  $\overline{\mathrm{Fl}}_{Y}^{w} \cap \overline{\mathrm{Fl}}_{Y,y} \neq \emptyset$ . Hence, the  $T^{\tau}$ -fixed point of the projective variety  $\overline{\mathrm{Fl}}_{Y}^{w} \cap \overline{\mathrm{Fl}}_{Y,y}$  is nonempty. Note that the  $T^{\tau}$ -fixed points of  $\mathrm{Fl}_{Y}$  are taking the form  $xe_{0}$ , where  $x \in W_{\mathrm{aff}}$ .

Define  ${}^{\bar{Y}}W^{\bar{Y}}$  to be the set of minimal length coset representatives of  $W_{\bar{Y}} \setminus W_{\text{aff}} / W_{\bar{Y}}$ . Let  $R \subseteq {}^{\bar{Y}}W^{\bar{Y}}$  be a finite subset so that  $w \in R$  and  $w' \leq w$  imply  $w' \in R$ . Set

$$S := W_{\bar{Y}} \cdot R \cdot W_{\bar{Y}} \subseteq W_{\text{aff}}.$$

Then,  $w \in S$  and  $w' \leq w$  implies  $w' \in S$ .

We define the following open subset of  $Fl_Y$ ,

$$V^S := \bigcup_{w \in S} w \mathcal{N}^- e_0,$$

where  $\mathcal{N}^-$  is the unipotent radical of  $\mathcal{I}^-$ .

**Lemma 3.11.** We have  $V^S = \bigcup_{w \in S} \mathcal{I}^- w e_0$ . As a consequence,  $V^S$  is  $\mathcal{P}_Y^-$ -stable.

*Proof.* We write  $\mathcal{N}^-V^S = \bigcup_{v \in S'} \mathcal{N}^-ye_0$  for some  $S' \subseteq W_{\text{aff}}$ . Clearly,  $S \subseteq S'$ . Note that

$$\mathcal{N}^-V^S = \bigcup_{w \in S} \mathcal{N}^-w\mathcal{N}^-e_0 = \bigcup_{w \in S} \mathcal{N}^-(\mathcal{N}^+ \cap w\mathcal{N}^-w^{-1})we_0 = \bigcup_{w \in S} \mathcal{N}^-Iwe_0,$$

where  $\mathcal{N}^+$  is the unipotent radical of I. Then, for each  $y \in S'$ , there exists  $w \in S$  such that  $\mathcal{N}^-ye_0 \cap \mathcal{N}^-Iwe_0 \neq \emptyset$ . This implies that  $\mathcal{N}^-ye_0 \cap Iwe_0 \neq \emptyset$ . By Lemma 3.10, we get  $y_{\min} \leq w_{\min}$ . Since S is saturated and stable under right action of  $W_{\bar{Y}}$ , we must have  $y \in S$ . Therefore,  $\mathcal{N}^-V^S = \bigcup_{y \in S'} \mathcal{N}^-ye_0 \subseteq \bigcup_{y \in S} \mathcal{N}^-ye_0$ . It follows that  $V^S \subseteq \mathcal{N}^-V^S \subseteq \bigcup_{y \in S} \mathcal{N}^-ye_0$ . Conversely, we have

$$\mathcal{N}^- y e_0 = (\mathcal{N}^- \cap y \mathcal{N}^- y^{-1}) y e_0 \subseteq y \mathcal{N}^- e_0.$$

Hence,  $\bigcup_{y \in S} \mathcal{N}^- y e_0 \subseteq V^S$ . Therefore,  $V^S = \bigcup_{y \in S} \mathcal{N}^- y e_0 = \bigcup_{y \in S} \mathcal{I}^- y e_0$ .

Set  $\overline{\mathrm{Fl}}_{Y}^{w}(S) := \overline{\mathrm{Fl}}_{Y}^{w} \cap V^{S}$ . In particular, when  $Y = \hat{I}_{\tau}$ , we simply denote it by  $\overline{\mathrm{Fl}}^{w}(S)$ . For each  $i \in Y$ , let  $\mathcal{L}_{i}$  denote the line bundle  $\mathcal{L}(\Lambda_{i})$  on  $\mathrm{Fl}_{Y}$ .

**Proposition 3.12.** For any  $i \in Y$ , the line bundle  $\mathcal{L}_i$  has a unique  $\mathcal{P}_Y^-$ -equivariant structure with  $M_Y$  acting trivially on the fiber  $\mathcal{L}_i|_{e_0}$  at the base point. Moreover, the line bundle  $\mathcal{L}_i$  has a unique  $\mathcal{P}_Y$ -equivariant structure with  $M_Y$  acting trivially on  $\mathcal{L}_i|_{e_0}$ .

*Proof.* Following the idea of [Fa, Theorem 7], we shall construct a  $\mathcal{P}_{Y}^{-}$ -equivariant line bundle  $L_{i}$  on Fl<sub>Y</sub>, and then show that  $L_{i}$  is isomorphic to  $\mathcal{L}_{i}$ .

Given any  $i \in Y$ , choose R as above so that  $s_i \in R$ . Then, we have  $W_{\bar{Y}}s_i \subseteq S$ . Set  $D := \overline{\mathrm{Fl}}_Y^{s_i}(S)$ . Then,  $\overline{\mathrm{Fl}}_Y^w(S) \subseteq D$  for any  $w \in W_{\bar{Y}}s_i$  since  $w \geq s_i$ . Thus, D is  $\mathcal{P}_Y^-$ -stable.

For each  $w \in S$ , we define a subgroup  $\mathcal{N}_w^- := \mathcal{N}^- \cap w \mathcal{N}^- w^{-1}$ . Consider the subgroup  $\mathcal{N}^-(n) \subseteq \mathcal{N}^-$  whose elements are identity modulo  $z^{-n}$ . Choose n big enough so that  $\mathcal{N}^-(n) \subseteq \mathcal{N}_w^-$  for all  $w \in S$ . By the same argument of [Fa, Lemma 6],  $\mathcal{N}^-(n) \setminus V^S$  is covered by finite many open subsets, which are finite dimensional affine spaces of same dimension. Thus,  $\mathcal{N}^-(n) \setminus V^S$  is a smooth variety.

Recall that D is  $\mathcal{P}_{Y}^{-}$ -stable and of codimension 1 in  $V^{S}$ . Thus,  $\mathcal{N}^{-}(n)\backslash D$  is a  $\mathcal{N}^{-}(n)\backslash \mathcal{P}_{Y}^{-}$ -stable divisor of  $\mathcal{N}^{-}(n)\backslash V^{S}$ . This defines an  $\mathcal{N}^{-}(n)\backslash \mathcal{P}_{Y}^{-}$ -equivariant line bundle  $L_{i,S}$  on

 $\mathcal{N}^-(n)\backslash V^S$  and  $\mathcal{N}^-(n)\backslash M_Y$  acts trivially on the canonical section  $1\in\Gamma(\mathcal{N}^-(n)\backslash V^S,L_{i,S})$ . Thus, the action of  $\mathcal{N}^-(n)\backslash M_Y$  on  $L_{i,S}|_{e_0}$  is trivial. By taking S arbitrarily large, we get a line bundle on  $\mathrm{Fl}_Y$ , denoted by  $L_i$ . From construction, this line bundle  $L_i$  has an  $\mathcal{P}^-_Y$ -equivariant structure such that  $M_Y$  acts on  $L_i|_{e_0}$  trivially.

Consider the natural morphism  $p_Y: \operatorname{Fl} \to \operatorname{Fl}_Y$ . We have  $p_Y^{-1}(\overline{\operatorname{Fl}}_Y^{s_i}(S)) = \overline{\operatorname{Fl}}^{s_i}(S)$ . Thus, the degree of the restriction of the pull-back  $p_Y^*(L_i)|_{\mathbb{P}^1_j}$  equals the degree of the divisor  $\overline{\operatorname{Fl}}^{s_i}(S) \cap \mathbb{P}^1_j$  in  $\mathbb{P}^1_j$ , and equals to the intersection number  $\#\{\overline{\operatorname{Fl}}^{s_i}(S) \cap \mathbb{P}^1_j\}$  since  $\overline{\operatorname{Fl}}^{s_i}(S) \cap \mathbb{P}^1_j$  intersects transversely. By Lemma 3.10,  $\overline{\operatorname{Fl}}^{s_i} \cap \mathbb{P}^1_j$  is empty if  $j \neq i$ . When j = i, from the proof of Lemma 3.10, we have  $\overline{\operatorname{Fl}}^{s_i} \cap \mathbb{P}^1_i = I^-s_ie_0 \cap Is_ie_0 \subsetneq \mathbb{P}^1_i$ . Hence, every point in  $\overline{\operatorname{Fl}}^{s_i} \cap \mathbb{P}^1_i$  is fixed by  $T^\tau$ . Thus,  $\overline{\operatorname{Fl}}^{s_i} \cap \mathbb{P}^1_i = \{s_ie_0\}$ . Therefore,  $\deg(p_Y^*(L_i)|_{\mathbb{P}^1_j})$  equals 1 if j = i and equals 0 if  $j \neq i$ . This shows that  $L_i \simeq \mathscr{L}_i$  as line bundles on  $\operatorname{Fl}_Y$ . Therefore,  $\mathscr{L}_i$  has an  $\mathscr{P}^-_Y$ -equivariant structure such that  $M_Y$  acts on  $\mathscr{L}_i|_{e_0}$  trivially. This equivariant structure is unique since the character of  $M_Y$  uniquely extends to a character of  $\mathscr{P}^-_Y$ . This proves the first statement.

We now prove the second statement. Let  $K_n$  be the n-th congruence subgroup in  $G(O)^{\tau}$ , i.e. whose elements are identity modulo  $z^n$ . Let  $\mathcal{P}_{Y,n}$  be the quotient group  $\mathcal{P}_Y/K_n$ . For any  $w \in W_{\text{aff}}$ , we consider the  $\mathcal{P}_Y$ -Schubert variety  ${}_YFl_{Y,w}$ . It is well-known that when n is sufficiently large, the action of  $\mathcal{P}_Y$  factors through  $\mathcal{P}_{Y,n}$ . Consider the restriction of  $\mathscr{L}_i$  to  $_Y\overline{\mathrm{Fl}}_{Y,w}$ , which will still be denoted by  $\mathscr{L}_i$ . Let  $\widehat{\mathscr{P}_{Y,n}}$  be the  $\mathbb{G}_{\mathrm{m}}$ -central extension of  $\mathcal{P}_{Y,n}$  associated to the line bundle  $\mathcal{L}_i$  on  ${}_Y\overline{\mathrm{Fl}}_{Y,w}$ . Note that  $\mathcal{P}_{Y,n}$  is the semiproduct of  $M_Y$  with a unipotent subgroup  $U_{Y,n}$ . By the first part of the proposition, we have a splitting  $M_Y \to \mathcal{P}_{Yn}$  over  $M_Y$ . On the other hand, by [CT, Corollary 5.7], the group  $Pic(U_{Y,n})$  classifies the isomorphism classes of extensions of  $U_{Y,n}$  by  $\mathbb{G}_m$ . Moreover, by [Po, Theorem1], the Picard group  $Pic(U_{Y,n})$  of unipotent group is trivial. Hence,  $U_{Y,n}$ only has trivial central extension by  $\mathbb{G}_m$ . Thus, we have a unique splitting  $U_{Y,n} \to \widehat{\mathcal{P}_{Y,n}}$ over  $U_{Y,n}$ . These splittings uniquely glue to a splitting  $\mathcal{P}_{Y,n} \to \widehat{\mathcal{P}_{Y,n}}$ . Therefore, the line bundle  $\mathcal{L}_i$  on  $_Y\overline{\mathrm{Fl}}_{Y,w}$  is  $\mathcal{P}_{Y,n}$ -equivariant. Taking a limit of  $_Y\overline{\mathrm{Fl}}_{Y,w}$  with respect to the Bruhat order on  $W_{\text{aff}}$ , we conclude that the line bundle  $\mathcal{L}_i$  on Fl<sub>Y</sub> is  $\mathcal{P}_Y$ -equivariant. From our argument, the induced  $M_Y$ -action on  $\mathcal{L}_i|_{e_0}$  is trivial. Finally, the equivariant structure with the desired property is unique since the character of  $M_Y$  uniquely extends to a character of  $\mathcal{P}_{Y}$ .

Let  $(G^{\tau})'$  denote the simply connected cover of  $G^{\tau}$  and set  $(G(O)^{\tau})' := G(O)^{\tau} \times_{G^{\tau}} (G^{\tau})'$ . Note that when G is simply-connected,  $G^{\tau}$  is connected but not necessarily simply-connected.

**Proposition 3.13.** For any  $i \in Y$ , the line bundle  $\mathcal{L}_i$  has a unique  $(G(O)^{\tau})'$ -equivariant structure. Under this equivariant structure,  $(T^{\tau})'$  acts on  $\mathcal{L}_i|_{e_0}$  via the weight  $-\omega_i$  if  $i \neq o$  and trivially if i = o.

*Proof.* Similar to the argument in Proposition 3.12, we consider any  $G(O)^{\tau}$ -Schubert variety  $X_w$  in Fl<sub>Y</sub>. Then, the action of  $G(O)^{\tau}$  factors through a group  $G(O_n)^{\tau}$ , where  $O_n = k[z]/(z^n)$ . Let  $(G(O_n)^{\tau})' := G(O_n)^{\tau} \times_{G^{\tau}} (G^{\tau})'$ . Note that  $(G(O_n)^{\tau})'$  is the semiproduct of  $(G^{\tau})'$  with the kernel  $J_n$  of the projection  $(G(O_n)^{\tau})' \to (G^{\tau})'$ . In view of [Po, Theorem1],

 $\operatorname{Pic}((G(O_n)^{\tau})') = \operatorname{Pic}((G^{\tau})') = 1$ . By [Po, Theorem1] again, any  $\mathbb{G}_m$ -central extension of  $(G(O_n)^{\tau})'$  is trivial. Hence, the restriction of the line bundle  $\mathcal{L}_i$  on  $X_w$  has a unique  $(G(O_n)^{\tau})'$ -equivariant structure, thus a unique  $(G(O)^{\tau})'$ -equivariant structure. Taking limit of  $X_w$ , we get a unique  $(G(O)^{\tau})'$ -equivariant structure on  $\mathcal{L}_i$  over  $\operatorname{Fl}_Y$ .

On the other hand, we have  $\deg\left(p_Y^*(\mathscr{L}_i)\big|_{\mathbb{P}^1_j}\right) = \delta_{i,j}$  for any  $i \in Y$  and  $j \in \hat{I}_\tau$ , cf. [PR, Section 10]. Hence,  $p_Y^*(\mathscr{L}_i)\big|_{\mathbb{P}^1_i} \simeq \mathscr{O}_{\mathbb{P}^1_i}(1)$  and  $p_Y^*(\mathscr{L}_i)\big|_{\mathbb{P}^1_j} \simeq \mathscr{O}_{\mathbb{P}^1_j}$  for  $j \neq i$ . Let  $\alpha_j$  be the simple root of  $G^\tau$  associated to  $j \in I_\tau$  and  $\{x_{\pm \alpha_j}\}$  be the root subgroups associated to  $\pm \alpha_i$ , cf. (12). Let  $G_{\alpha_j}$  be the subgroup of the simply-connected cover  $(G^\tau)'$  of  $G^\tau$  generated by  $\{x_{\pm \alpha_j}(a) \mid a \in k\}$ . The group  $G_{\alpha_j} \simeq \mathrm{SL}_2$  acts on  $\mathbb{P}^1_j$  transitively and has a unique equivariant structure on  $p_Y^*(\mathscr{L}_i)\big|_{\mathbb{P}^1_j}$  determined by its degree. In fact,  $a \in \mathbb{G}_m \subset G_{\alpha_j}$  acts on  $(p_Y^*(\mathscr{L}_i)\big|_{\mathbb{P}^1_j})\big|_{e_0} = \mathscr{L}_i\big|_{e_0}$  by  $a^{-\delta_{ij}}$ . This concludes the proof of the proposition.  $\Box$ 

*Remark* 3.14. The induced actions of  $(T^{\tau})'$  on  $\mathcal{L}|_{e_0}$  are different in Proposition 3.12 and 3.13, as we regard  $T^{\tau}$  and  $(T^{\tau})'$  as subgroups in  $\mathcal{P}_Y$  and  $(G(O)^{\tau})'$  respectively.

Let  $\mathcal{L}(\Lambda)$  be an ample line bundle on Fl<sub>Y</sub> associated to  $\Lambda = \sum_{i \in Y} n_i \Lambda_i$ , i.e.  $n_i > 0$  for any  $i \in Y$ . Let  $\lambda$  be the restriction of  $\Lambda$  to  $(T^{\tau})'$ .

Recall Lemma 2.6, when  $p \nmid a_Y$ , we can choose an interior point  $\vartheta$  in the facet associated to Y so that  $p \nmid m$ . Define a subgroup  $G_{\vartheta}$  of  $G^{\tau}$  as in (15), which is a subgroup of  $M_Y$ . Let  $G'_{\vartheta}$  be the pre-image of  $G_{\vartheta}$  via the map  $(G^{\tau})' \to G^{\tau}$ . Recall Corollary 2.9,  $G_{\vartheta}$  is a subgroup of  $M_Y$ . Thus, we have a natural morphism  $G'_{\vartheta} \to M_Y$ .

Applying Proposition 3.12 and Proposition 3.13, we get

**Corollary 3.15.** The line bundle  $\mathcal{L}(\Lambda)$  has a unique  $\mathcal{P}_Y$ -equivariant structure with  $M_Y$  acting on  $\mathcal{L}(\Lambda)|_{e_0}$  trivially and a unique  $(G(O)^{\tau})'$ -equivariant structure. Moreover, the  $G'_{\vartheta}$ -actions on  $\mathcal{L}(\Lambda)$  induced from these two equivariant structures differ by a weight  $\lambda$ .

*Proof.* By Proposition 3.12, the line bundle  $\mathcal{L}(\Lambda)$  has a unique  $\mathcal{P}_Y$ -equivariant structure with  $M_Y$  acting on  $\mathcal{L}(\Lambda)|_{e_0}$  trivially. This induces a  $G'_{\vartheta}$ -action on  $\mathcal{L}(\Lambda)$  via the natural morphism  $G'_{\vartheta} \to M_Y$  described above. On the other hand, by Proposition 3.13, the line bundle  $\mathcal{L}(\Lambda)$  has a unique  $(G(O)^{\tau})'$ -equivariant structure on  $\mathcal{L}(\Lambda)$ . Under the second equivariant structure,  $G'_{\vartheta}$  acts on  $\mathcal{L}(\Lambda)|_{e_0}$  via the weight  $-\sum_{i\in Y} n_i \omega_i = -\lambda$ , cf. Proposition 3.13. Therefore, these two  $G'_{\vartheta}$ -actions differ by a weight  $\lambda$ .

# 3.3. Equivariance of line bundles on ramified global affine Grassmannians. In this section, we assume that G is simply-connected.

Recall Section 2.3, we fix an interior point  $\vartheta$  in the facet  $F_Y$  of A associated to  $Y \subseteq \hat{I}_{\tau}$  so that  $p \nmid m$ . Define parahoric group schemes  $\mathscr{G}_{\vartheta}$  over  $\bar{O}$  and  $\mathscr{G}_{-\vartheta,\infty}$  over  $\bar{O}_{\infty}$  as in Section 2.2. Let  $\mathbb{A}^1 = \operatorname{Spec}(\Bbbk[z])$  be the affine line with a  $\tau$ -action given by  $\tau(z) = \epsilon^{-1}z$ . Let  $C := \mathbb{A}^1 \cup \{\infty\}$  be the projective line. Let  $\bar{\mathbb{A}}^1$  and  $\bar{C}$  be the quotient of  $\mathbb{A}^1$  and C by the  $\tau$ -action. We can naturally glue  $\mathscr{G}_{\vartheta}$ ,  $\operatorname{Res}_{\mathbb{A}^1 - 0/\bar{\mathbb{A}}^1 - 0}(G_{\mathbb{A}^1 - 0})^{\tau}$ , and  $\mathscr{G}_{-\vartheta,\infty}$ , and get a parahoric Bruhat-tits group scheme G over  $\bar{C}$ .

By definition, we have the following property:

(28) 
$$\mathcal{G}|_{\mathbb{D}_{a}} \simeq \begin{cases} G_{\mathbb{D}_{a}} & \text{if } a \neq 0, \infty \\ \mathcal{G}_{\vartheta} & \text{if } a = 0 \\ \mathcal{G}_{-\vartheta} & \text{if } a = \infty \end{cases}$$

where  $\mathbb{D}_a$  denotes the formal disk around  $a \in \bar{C}$ .

Similar to the local description in Theorem 2.4, we shall give another description of the parahoric Bruhat-tits group scheme from the equivariant point of view. Let  $\tilde{\mathbb{A}}^1 = \operatorname{Spec}(\mathbb{k}[t])$  be the affine line with a  $\sigma$ -action given by  $\sigma(t) := (\epsilon_m)^{-1}t$ , where  $\epsilon_m$  is a fixed primitive m-th root of unity. Let  $\tilde{C} := \tilde{\mathbb{A}}^1 \cup \{\infty\}$  be the projective line. Let  $\tilde{C} \to C$  be the map given by  $a \mapsto a^{m/r}$  for any  $a \in \tilde{C}$ . Then,  $\bar{C} = C/\langle \tau \rangle = \tilde{C}/\langle \sigma \rangle$ .

Define the following group scheme over  $\bar{C}$ ,

$$\mathcal{G}_{\sigma} := \operatorname{Res}_{\tilde{C}/\bar{C}}(G \times \tilde{C})^{\sigma},$$

where  $\sigma$  acts on  $\tilde{C}$  and  $\sigma$  acts on G defined in (5). Clearly, we have  $\mathcal{G}_{\sigma}|_{\bar{\mathbb{D}}_0} \simeq \mathcal{G}_{\sigma}$  and  $\mathcal{G}_{\sigma}|_{\bar{\mathbb{D}}_{\infty}} \simeq \mathcal{G}_{\sigma,\infty}$ , where  $\mathcal{G}_{\sigma}$  and  $\mathcal{G}_{\sigma,\infty}$  are defined in Section 2.2, and  $\bar{\mathbb{D}}_0 = \operatorname{Spec}(\bar{O})$  denotes the formal disks around  $0 \in \bar{C}$ .

**Lemma 3.16.** The map  $Ad_{\tau^{\theta}}$  induces an isomorphism of group schemes over  $\bar{\mathbb{A}}^1 - 0$ ,

(29) 
$$\phi_{\tilde{\mathbb{A}}^{1}-0} : \operatorname{Res}_{\mathbb{A}^{1}-0/\tilde{\mathbb{A}}^{1}-0}(G_{\mathbb{A}^{1}-0})^{\tau} \simeq \operatorname{Res}_{\tilde{\mathbb{A}}^{1}-0/\tilde{\mathbb{A}}^{1}-0}(G_{\tilde{\mathbb{A}}^{1}-0})^{\sigma}.$$

*Proof.* The proof is identical to the proof of [DH1, Proposition 4.1.1].

In view of Theorem 2.4 and Lemma 3.16, the isomorphisms  $\phi_0$ ,  $\phi_\infty$  (defined in Section 2.2), and  $\phi_{\bar{\mathbb{A}}^1-0}$  can be glued to an isomorphism of group schemes over  $\bar{C}$ ,

$$\phi: \mathcal{G} \simeq \mathcal{G}_{\sigma}.$$

**Definition 3.17.** The global affine Grassmannian  $Gr_{\mathcal{G},\bar{C}}$  and the jet group scheme  $L^+\mathcal{G}$  over  $\bar{C}$  are defined as follows, for any k-algebra R,

$$\operatorname{Gr}_{\mathcal{G}}(R) := \left\{ (a, \mathcal{F}, \beta) \,\middle|\, a \in \bar{C}(R), \mathcal{F} \text{ a } \mathcal{G}\text{-torsor over } \bar{C}_R, \beta : \mathcal{F}|_{\bar{C}_R \setminus \Delta_a} \simeq \mathring{\mathcal{F}}|_{\bar{C}_R \setminus \Delta_a} \right\}$$

and

(31) 
$$L^{+}\mathcal{G}(R) := \left\{ (a, \eta) \mid a \in \bar{C}(R), \eta \in \mathcal{G}(\hat{\Delta}_{a}) \right\},$$

where  $\mathring{\mathcal{F}}$  is the trivial  $\mathcal{G}$ -torsor on  $\bar{C}_R$ , and  $\hat{\Delta}_a$  is the completion of  $\bar{C}_R$  along the graph  $\Delta_a$  of  $a: \operatorname{Spec}(R) \to \bar{C}$ .

For any  $a \in \bar{C}$ , let  $Gr_{\mathcal{G},a}$  and  $L^+\mathcal{G}_a$  denote the fiber of  $Gr_{\mathcal{G},C}$  and  $L^+\mathcal{G}_C$  at a respectively.

The global Grassmannian  $Gr_{\mathcal{G}}$  over  $\bar{C}$  has the following property,

$$\operatorname{Gr}_{\mathcal{G},a} \simeq \begin{cases} \operatorname{Fl}_{Y} & \text{when } a = 0 \\ \operatorname{Gr}_{G} & \text{when } a \neq 0, \infty. \end{cases}$$

**Proposition 3.18.** Let  $\operatorname{Bun}_{\mathcal{G}}$  be the moduli stack of  $\mathcal{G}$ -torsors on  $\bar{\mathcal{C}}$ . We have

$$\operatorname{Bun}_{\mathcal{G}} \simeq [\mathcal{P}_{\mathcal{Y}} \backslash \operatorname{Fl}_{\mathcal{Y}}].$$

Moreover, the line bundle  $\mathcal{L}(\Lambda)$  descends to a line bundle  $\mathcal{L}_{Bun_G}$  on  $Bun_G$ .

*Proof.* We define the following ind-group scheme  $L^-\mathcal{G}$ , for any k-algebra R,

(32) 
$$L^{-}\mathcal{G}(R) = \{(a,g) \mid a \in \bar{C}(R), g \in \mathcal{G}(\bar{C}_R \setminus \Delta_a)\}.$$

Taking fiber at  $0 \in \bar{C}$ , we have  $L^-\mathcal{G}_0 \simeq \mathcal{P}_Y^-$ , see Proposition 3.6. By uniformization theorem (cf. [He]), we have  $\operatorname{Bun}_{\mathcal{G}} \simeq [\mathcal{P}_Y^- \backslash \operatorname{Fl}_Y]$ . The second part of the statement follows from the  $\mathcal{P}_Y^-$ -equivariance of the line bundle  $\mathscr{L}(\Lambda)$  on  $\operatorname{Fl}_Y$  (Theorem 3.15).

Remark 3.19. For general parahoric Bruhat-Tits group scheme  $\mathcal{G}$  over a smooth projective curve, line bundles on  $\operatorname{Bun}_{\mathcal{G}}$  have been further investigated in [DH2] by Damiolini and the first author.

We denote by  $\mathcal{L} := \operatorname{pr}^*(\mathcal{L}_{\operatorname{Bun}_{\mathcal{G}}})$  the pullback of  $\mathcal{L}_{\operatorname{Bun}_{\mathcal{G}}}$  via the natural projection pr :  $\operatorname{Gr}_{\mathcal{G}} \to \operatorname{Bun}_{\mathcal{G}}$ . For each  $a \in \overline{C}$ , let  $\mathcal{L}_a$  denote the restriction of  $\mathcal{L}$  to the fiber  $\operatorname{Gr}_{\mathcal{G},a}$  of  $\operatorname{Gr}_{\mathcal{G}}$  at  $a \in \overline{C}$ .

**Proposition 3.20.** We have an isomorphism  $\mathcal{L}_0 \simeq \mathcal{L}(\Lambda)$  of line bundles on  $\operatorname{Gr}_{\mathcal{G},0} \simeq \operatorname{Fl}_Y$ . Moreover, for any  $a \neq 0, \infty$  in  $\bar{C}$ , we have an isomorphism  $\mathcal{L}_a \simeq L^c$  of line bundles on  $\operatorname{Gr}_{\mathcal{G},a} \simeq \operatorname{Gr}_{\mathcal{G}}$ , where L denotes the level one line bundle on  $\operatorname{Gr}_{\mathcal{G}}$ .

*Proof.* The first statement follows from the construction of  $\mathcal{L}$ . For the second statement, applying [Zh1, Proposition 4.1], the central charge of the restriction  $\mathcal{L}_a$  is constant, hence is equal to c. Since the central charge map  $\operatorname{Pic}(\operatorname{Gr}_G) \to \mathbb{Z}$  is an isomorphism, we conclude that  $\mathcal{L}_a \simeq L^c$ .

Recall the  $\mathcal{P}_Y$ -equivariant structure of the line bundle  $\mathcal{L}(\Lambda)$  on  $\mathrm{Fl}_Y$  and the G(O)-equivariant structure of line bundle L on  $\mathrm{Gr}_G$ , cf. Theorem 3.15 and [Fa, Theorem 7]. Via the embeddings  $G_{\vartheta} \to \mathcal{P}_Y$  and  $G_{\vartheta} \to G(O)$ , they induce  $G_{\vartheta}$ -equivariant structures on line bundles  $\mathcal{L}(\Lambda)$  and L with  $G_{\vartheta}$  acting on  $\mathcal{L}|_{e_0}$  and  $L|_{e_0}$  trivially.

**Theorem 3.21.** The line bundle  $\mathcal{L}$  over  $\operatorname{Gr}_{\mathcal{G}}$  has an  $G_{\vartheta} \times \bar{C}$ -equivariant structure such that the induced  $G_{\vartheta}$ -equivariant structure on  $\mathcal{L}_a$  is the same as the  $G_{\vartheta}$ -equivariant structure on  $\mathcal{L}(\Lambda)$  (when  $a=0,\infty$ ) or  $L^c$  (when  $a\neq 0,\infty$ ) defined above.

*Proof.* Recall the ind-group scheme  $L^-\mathcal{G}$  defined in Proposition 3.18. There is an isomorphism  $[L^-\mathcal{G}\backslash \mathrm{Gr}_{\mathcal{G}}]\simeq \bar{C}\times \mathrm{Bun}_{\mathcal{G}}$ . Thus, the pullback  $\mathrm{pr}^*(\mathcal{L}_{\mathrm{Bun}_{\mathcal{G}}})$  is naturally equipped with a  $L^-\mathcal{G}$ -equivariant structure.

For the parahoric Bruhat-Tits group scheme  $\mathcal{G}_{\sigma}$ , we can similarly define the ind-group scheme  $L^-\mathcal{G}_{\sigma}$  over  $\bar{C}$  as in (32). Note that, for any commutative ring R over k,  $\mathcal{G}_{\sigma}(\bar{C}_R)$  can be naturally identified with the set of  $\sigma$ -equivariant morphisms from  $C_R \setminus \bigcup_{i=0}^{m-1} \Delta_{\sigma^i \bar{a}}$  to G, where  $\tilde{a}$  is a preimage of  $\bar{a}$  under the map  $\tilde{C}(R) \to \bar{C}(R)$  and  $\Delta_{\sigma^i \bar{a}}$  is the graph of  $\sigma^i \tilde{a}$  in  $C_R$ . There is a natural embedding  $G^{\sigma} \hookrightarrow \mathcal{G}_{\sigma}(\bar{C}_R \setminus \Delta_{\bar{a}})$  giving by sending  $g \in G^{\sigma}$  to the constant morphism landing at g. This induces an embedding  $G^{\sigma} \times \bar{C} \hookrightarrow L^-\mathcal{G}_{\sigma}$  of group schemes over  $\bar{C}$ . Moreover, the map  $\phi$  in (30) induces an isomorphism  $L^-\mathcal{G} \simeq L^-\mathcal{G}_{\sigma}$ . Hence, we have the following composition of morphisms

(33) 
$$G_{\vartheta} \times \bar{C} \subseteq G^{\sigma} \times \bar{C} \hookrightarrow L^{-}G_{\sigma} \simeq L^{-}G.$$

Restricting to the fibers at  $0 \in \bar{C}$ , the composition map (33) is given by

(34) 
$$G_{\vartheta} \subseteq G^{\sigma} \hookrightarrow G[t^{-1}] \xrightarrow{\operatorname{Ad}_{r^{-h}}} \mathcal{P}_{Y}^{-},$$

where the isomophism  $\mathrm{Ad}_{t^{-h}}$  is the restriction of  $\mathrm{Ad}_{t^{-h}}: G((t^{-1}))^{\sigma} \simeq G((z^{-1}))^{\tau}$ , see Proposition 3.6. Since  $\mathrm{Ad}_{t^{-h}}$  restricts to the identity map on  $G_{\vartheta}$ , the composition map in (34) is exactly the natural embedding  $G_{\vartheta} \subseteq \mathcal{P}_{V}^{-}$ .

We now consider restricting the composition map (33) to the fibers at  $\bar{a} \in \bar{C} \setminus \{0, \infty\}$ . Let a be a lifting of  $\bar{a}$  in C, and let  $\tilde{a}$  be a lifting of a in  $\tilde{C}$ . Then, the restriction of the composition map (33) at  $\bar{a}$  given by

$$G_{\vartheta} \subseteq G^{\sigma} \hookrightarrow G[t_{\tilde{a}}^{-1}] \xrightarrow{\operatorname{Ad}_{r^{-h}}} G[z_{a}^{-1}],$$

where the isomorphism  $\operatorname{Ad}_{r^{-h}}$  is the restriction of  $\operatorname{Ad}_{r^{-h}}: G((t)) \simeq G((z))$ , and  $t_{\tilde{a}} = t - \tilde{a}, z_a = z - a$ . By the definition of  $G_{\vartheta}$  in (15),  $\operatorname{Ad}_{r^{-h}}$  restricts to the identity map on  $G_{\vartheta}$ . Thus, this composition is the natural embedding  $G_{\vartheta} \subseteq G[z_a^{-1}]$ .

In summary, the natural  $L^-\mathcal{G}$ -equivariant structure on  $\operatorname{pr}^*(\mathcal{L}_{\operatorname{Bun}_{\mathcal{G}}})$  induces a  $G_{\vartheta} \times \bar{C}$  equivariant structure on it via the embedding  $G_{\vartheta} \times \bar{C} \hookrightarrow L^-\mathcal{G}$  in (33). Restricting to the fiber at  $0 \in \bar{C}$ , the induced  $G_{\vartheta}$ -equivariant structure on  $\mathcal{L}_0$  is the same as the one induced from  $\mathcal{P}_Y^-$ -equivariant structure via the embedding  $G_{\vartheta} \subseteq \mathcal{P}_Y^-$ . In view of Theorem 3.12, this is the same  $G_{\vartheta}$ -equivariant structure as the one induced from  $\mathcal{P}_Y$ -equivariant structure via the embedding  $G_{\vartheta} \subseteq \mathcal{P}_Y$ . Moreover, restricting to the fiber at  $\bar{a} \in \bar{C} \setminus \{0\}$ , the induced  $G_{\vartheta}$ -equivariant structure on  $\mathcal{L}_{\bar{a}}$  is the same as the one induced from  $G[z_a^{-1}]$ -equivariant structure via the embedding  $G_{\vartheta} \subseteq G[z_a^{-1}]$ . In view of [Fa, Theorem 7], this is the same  $G_{\vartheta}$ -equivariant structure as the one induced from  $G[[z_a]]$ -equivariant structure via the embedding  $G_{\vartheta} \subseteq G[[z_a]]$ .

3.4. **Proof of Theorem 3.5.** When G is simply-connected, Theorem 3.5 follows from Theorem 3.21 and Corollary 3.15. We now assume G is a general simple algebraic group.

For any  $\mu \in X_*(T)$ , let  $\mathrm{Fl}_Y^{\kappa}$  be the component of  $\mathrm{Fl}_Y$  containing  $\mathrm{Fl}_{Y,\bar{\mu}}$  (equivalently  $\varrho^{\bar{\mu}}\varsigma_{\kappa}^{-1} \in W_{\mathrm{aff}}$ ). Recall that we have an isomorphism  $\mathrm{Fl}_Y^{\kappa} \simeq \mathrm{Fl}_{Y_{\kappa}}^{\prime}$  defined in (22). Let  $\mathscr{L}(\Lambda)$  be the ample line bundle on  $\mathrm{Fl}_Y^{\kappa}$  associated to  $\Lambda = \sum_{i \in Y_{\kappa}} n_i \Lambda_i$ , i.e.  $n_i > 0$  for any  $i \in Y_{\kappa}$ .

Assume  $p \nmid a_{Y_{\kappa}}$ . By Lemma 2.6, we can choose an interior point  $\vartheta$  in the facet associated to  $Y_{\kappa}$  so that  $p \nmid m$ . Let G' denote the simply-connected cover of G. Let  $G_{\vartheta}$  be the subgroup of  $G^{\tau,\circ}$  associated to  $\vartheta$  defined in (15) of Section 2.3. Let  $G'_{\vartheta}$  (resp.  $(T^{\tau,\circ})'$ ) be the pre-image of  $G_{\vartheta}$  (resp.  $T^{\tau,\circ}$ ) via the simply-connected cover map  $(G^{\tau,\circ})' \to G^{\tau,\circ}$ .

Let  $\mathscr{G}_{\vartheta}$  (resp.  $\mathscr{G}_{-\vartheta,\infty}$ ) be parahoric group schemes in  $G((z))^{\tau}$  (resp.  $G((z^{-1}))^{\tau}$ ) over O (resp.  $\bar{O}_{\infty}$ ) defined in Section 2.2. We can naturally glue  $\mathscr{G}_{\vartheta}$ ,  $\operatorname{Res}_{\mathbb{A}^1-0/\bar{\mathbb{A}}^1-0}(G_{\mathbb{A}^1-0})^{\tau}$ , and  $\mathscr{G}_{-\vartheta,\infty}$ , and get a parahoric Bruhat-tits group scheme G over  $\bar{C}$ . Consider the following global affine Grassmannian over C,

$$Gr_{G,C} := Gr_G \times_{\bar{C}} C.$$

For each  $a \in C$ , we denote  $Gr_{\mathcal{G},a}$  the fiber of  $Gr_{\mathcal{G},C}$  at a. Then,  $Gr_{\mathcal{G},0} \simeq Fl_Y$  and  $Gr_{\mathcal{G},a} \simeq Gr_G$  for  $a \neq 0, \infty$ . Similarly, we set  $L\mathcal{G}_C := L\mathcal{G} \times_{\bar{C}} C$ .

Note that  $Gr_{\mathcal{G},\mathbb{A}^1-0} \simeq Gr_G \times (\mathbb{A}^1-0)$ , whose connected components are parametrized by  $X_*(T)/X_*(T')$ . Let M denote the set consisting of 0 and miniscule dominant coweights of G. There is a natural bijection  $M \simeq X_*(T)/X_*(T')$ . For any  $\xi \in M$ , let  $Gr_{\mathcal{G},\mathbb{A}^1-0}[\xi]$  denote the component containing  $e_{-\xi} \times (\mathbb{A}^1-0)$ , where  $e_{-\xi} \in Gr_G$ . Let  $Gr_{\mathcal{G},\mathcal{C}}[\xi]$  be the

closure of  $Gr_{G,\mathbb{A}^1-0}[\xi]$  in  $Gr_{G,\mathbb{C}}$ . When  $\xi=0$ , we call  $Gr_{G,\mathbb{C}}[\xi]$  the neutral component and denote it by  $Gr_{G,C}[o]$ .

The cover  $G' \to G$  induces a natural morphism  $G' \to G$ . By a similar proof of [HY, Lemma 5.16], there is an isomorphism  $Gr_{\mathcal{G}',\mathcal{C}} \simeq Gr_{\mathcal{G},\mathcal{C}}[o]$ . Recall Theorem 3.18, which states that the line bundle  $\mathcal{L}(\varsigma_{\kappa}^{-1}\Lambda)$  on  $\mathrm{Fl}_Y'$  descends to a line bundle on  $\mathrm{Bun}_{\mathscr{G}'}$ . Pulling back via the composition of morphisms  $Gr_{G',C} \to Gr_{G'} \to Bun_{G'}$ , we get a line bundle  $\mathcal{L}_o$  on  $Gr_{\mathcal{G}',\mathcal{C}}$ , which can be regarded as a line bundle on  $Gr_{\mathcal{G},\mathcal{C}}[o]$ .

For any dominant coweight  $\mu \in X_*(T)$ , let  $s_\mu : C \to L\mathcal{G}_C$  be the section constructed in [Zh1, Section 3.3]. It induces a section  $C \to Gr_{\mathcal{G},C}$ , which is still denoted by  $s_{\mu}$  if there is no confusion. For any  $a \in C - 0$ ,  $s_{\mu}(a) = e_{\mu} \in Gr_G$  under the identification  $Gr_{\mathcal{G},a} \simeq Gr_{\mathcal{G}}$ ; and  $s_{\mu}(0) = e_{\bar{\mu}} \in Gr_{\mathcal{G},0} \simeq Fl_{Y}$ .

For any miniscule dominant element  $\xi \in X_*(T)$ , let  $T_{\xi} : \operatorname{Gr}_{\mathcal{G},C}[o] \simeq \operatorname{Gr}_{\mathcal{G},C}[\xi]$  be the translation map by multiplying  $s_{-\xi}$ . Let  $\mathcal{L}_{\xi} := (T_{\xi})_*(\mathcal{L}_o)$  denote the push-forward of  $\mathcal{L}_o$ .

**Lemma 3.22.** The line bundle  $\mathcal{L}_{\xi}$  on  $Gr_{\mathcal{G},C}[\xi]$  has a  $T'^{\tau} \times C$ -equivariant structure. Moreover, the  $T'^{\tau}$ -equivariant structure on the fiber  $(\mathcal{L}_{\xi})_a$  comes from the embedding  $T'^{\tau} \to \mathcal{P}'_{Y_{-}}$  for a = 0 and  $T'^{\tau} \to G'[z_a]$  for  $a \neq 0$ .

*Proof.* By Theorem 3.21, the line bundle  $\mathcal{L}_o$  on  $Gr_{\mathcal{G},C}[o]$  is  $T'^{\tau} \times C$ -equivariant. Note that a  $T'^{\tau} \times C$ -equivariant structure on  $\mathcal{L}_o$  is equivalent to a  $\mathrm{Ad}_{s_{-\xi}}(T'^{\tau} \times C)$ -equivariant structure on  $\mathcal{L}_{\xi}$ . Moreover,  $\mathrm{Ad}_{s_{-\xi}}(T'^{\tau} \times C) = T'^{\tau} \times C$ . Therefore, the line bundle  $\mathcal{L}_{\xi}$  on  $Gr_{G,C}[\xi]$  has a  $T'^{\tau} \times C$ -equivariant structure.

**Definition 3.23.** Given any dominant  $\mu \in X_*(T)$ , define the global Schubert variety  $Gr_{G,C}^{\mu}$ to be the minimal  $L^+\mathcal{G}$ -stable irreducible closed subvariety of  $Gr_{\mathcal{G},\mathcal{C}}$  that contains  $s_\mu$ . Let  $Gr_{\mathcal{G},C}[\xi]$  be the component of  $Gr_{\mathcal{G},C}$  containing  $\overline{Gr}_{\mathcal{G},C}^{\mu}$  (equivalently  $\mu + \xi \in X_*(T')$ ). The line bundle  $\mathcal{L}_{\xi}$  restricts to a line bundle on  $\overline{\mathrm{Gr}}_{GC}^{\mu}$ , which will be denoted by  $\mathcal{L}$ .

We are now ready to conclude Theorem 3.5.

By [Zh1, Theorem 3], the global Schubert variety  $\overline{\operatorname{Gr}}_{G,C}^{\mu}$  is flat over C, and for any  $a \in C$ , we have

(35) 
$$\overline{\mathrm{Gr}}_{\mathcal{G},a}^{\mu} \simeq \begin{cases} \mathcal{A}_{Y}(\bar{\mu}) & \text{if } a = 0\\ \overline{\mathrm{Gr}_{G,\mu}} & \text{if } a \neq 0, \infty \end{cases}.$$

Combining with Lemma 3.22, we conclude that the space  $H^0(\overline{\mathrm{Gr}}_{\mathcal{G},C}^{\mu},\mathcal{L})$  is a  $T'^{\tau}$ -module and flat over C. By the complete reducibility of  $T'^{\tau}$ , we have an  $T'^{\tau}$ -isomorphism

(36) 
$$H^{0}(\overline{\mathrm{Gr}}_{\mathcal{G},0}^{\mu},\mathcal{L}_{0}) \simeq H^{0}(\overline{\mathrm{Gr}}_{\mathcal{G},a}^{\mu},\mathcal{L}_{a}).$$

Note that over  $0 \in C$  the translation map  $T_{\xi} : Gr_{\mathcal{G}} \to Gr_{\mathcal{G}}$  is the map  $Fl_Y \to Fl_Y$  given by  $x \mapsto n^{-\xi} \cdot x$ . This restricts to an isomorphism  $n^{-\xi} : \operatorname{Fl}_{Y}^{\circ} \simeq \operatorname{Fl}_{Y}^{\kappa}$ . Thus, we have

$$\mathcal{L}_0 = (\mathcal{L}_{\xi})|_{\mathsf{Fl}_{\nu}^{\kappa}} = (n^{-\xi})_*((\mathcal{L}_{\varrho})|_{\mathsf{Fl}_{\nu}^{\circ}}) = (n^{-\xi})_*(\mathcal{L}(\varsigma_{\nu}^{-1}(\Lambda))) = \mathcal{L}(\Lambda).$$

Therefore, (36) can be rewritten as the following isomorphsim of  $T^{\prime\tau}$ -modules

(37) 
$$H^{0}(\mathcal{A}_{Y}(\bar{\mu}), \mathcal{L}(\Lambda)) \simeq H^{0}(\overline{\mathrm{Gr}}_{G,\mu}, L^{c}).$$

In view of Lemma 3.22, the  $T'^{\tau}$ -action on the fiber  $\mathcal{L}_a$  comes from the embedding  $T'^{\tau} \to \mathcal{P}'_{Y_{\kappa}}$  for a=0 and from the embedding  $T'^{\tau} \to G'[z_a]$  for  $a \neq 0$ . Consider the natural map  $(T^{\tau,\circ})' \to T'^{\tau}$ . This induces  $(T^{\tau,\circ})'$ -module structures on both sides of (37).

On the other hand, by Corollary 3.15, the line bundle  $\mathcal{L}(\Lambda)$  has equivariant structures of  $\mathcal{P}'_{Y_{\kappa}}$  and  $(G'(O)^{\tau})' \simeq (G(O)^{\tau,\circ})'$ . Moreover,  $(T^{\tau,\circ})'$ -actions on  $\mathcal{L}(\Lambda)_{e_0}$  via the morphisms  $(T^{\tau,\circ})' \to T'^{\tau} \hookrightarrow \mathcal{P}'_{Y_{\kappa}}$  and  $(T^{\tau,\circ})' \hookrightarrow (G^{\tau,\circ})' \to (G(O)^{\tau,\circ})'$  differ by a weight  $\lambda$ . Thus, (37) can be rewritten as the following isomorphsim of  $(T^{\tau,\circ})'$ -modules

(38) 
$$H^{0}(\mathcal{A}_{Y}(\bar{\mu}), \mathcal{L}(\Lambda)) \otimes \mathbf{k}_{\lambda} \simeq H^{0}(\overline{\operatorname{Gr}}_{G}^{\mu}, L^{c}),$$

where the  $(T^{\tau,\circ})'$ -action on the left side comes from the unique  $(G'(O)^{\tau})'$ -equivariant structure on  $\mathcal{L}(\Lambda)$  and  $(T^{\tau,\circ})'$ -action on the right side comes from the unique G'(O)-equivariant structure on L. This completes the proof of Part (1) of Theorem 3.5.

We now assume char(k) = 0. As mentioned in Section 3.1, both sides of (38) are representations of  $G'_{\vartheta}$ . Since two sides of (38) have the same  $(T^{\tau,\circ})'$ -character, by standard representation theory of reductive group we can conclude that (38) are isomorphic as  $G'_{\vartheta}$ -representations.

### 4. Application of the coherence conjecture to Affine Demazure modules

4.1. **Affine Kac-Moody algebras.** Let k be an algebraically closed field with characteristic 0. Let g be a simple Lie algebra over k. We fix a Borel subalgebra  $\mathfrak{b} \subseteq \mathfrak{g}$  and a Cartan subalgebra  $\mathfrak{h} \subseteq \mathfrak{b}$ . Let  $(\cdot, \cdot)$  be the invariant (symmetric, nondegenerate) bilinear form on g normalized so that the induced form on the dual space  $\mathfrak{h}^*$  satisfies  $(\theta, \theta) = 2$  for the highest root  $\theta$  of g.

Let  $\tau$  be a diagram automorphism of g of order r preserving b and b, which is allowed to be trivial. We fix an r-th primitive root of unity  $\epsilon$ . Let  $g(t) := g \otimes_k k(t)$  be the loop algebra and let  $g(t)^{\tau}$  denote the  $\tau$ -fixed point subalgebra with respect to the extended  $\tau$ -action by  $\tau(t) := \epsilon^{-1}t$ . Then, the (twisted) affine Lie algebra  $\tilde{L}(g, \tau)$  associated to g and  $\tau$  is defined as follows,

(39) 
$$\tilde{L}(\mathfrak{g},\tau) := \mathfrak{g}((t))^{\tau} \oplus kK,$$

where K denotes the canonical central element and the Lie bracket on  $\tilde{L}(\mathfrak{g},\tau)$  is given by

$$[x(f) + zK, x'(f') + z'K] := [x, x'](ff') + r^{-1}Res_{t=0}((df)f')(x, x')K.$$

Let  $\hat{L}(\mathfrak{g}, \tau) := \tilde{L}(\mathfrak{g}, \tau) \oplus kd$  be the affine Kac-Moody algebra, where d is the scaling element. When  $\tau$  is trivial, we simply denote  $\hat{L}(\mathfrak{g}, \tau)$  by  $\hat{L}(\mathfrak{g})$ . Set  $\hat{\mathfrak{h}}^{\tau} := \mathfrak{h}^{\tau} \oplus kK \oplus kd$ , which is the Cartan subalgebra of  $\hat{L}(\mathfrak{g}, \tau)$ . Let  $(\hat{\mathfrak{h}}^{\tau})^*$  denote the linear dual of  $\hat{\mathfrak{h}}^{\tau}$ .

Let  $X_N^{(r)}$  be the affine Dynkin diagram associated to  $\hat{L}(\mathfrak{g}, \tau)$ , and we similarly have the notations  $I_{\tau}$ ,  $\hat{I}_{\tau}$ , and  $\{\alpha_i, \check{\alpha}_i | i \in \hat{I}_{\tau}\}$ , as defined in Section 2.1. Recall the weight  $\theta_0$  of  $\mathfrak{g}^{\tau}$ 

defined in (2). We have another description of  $\theta_0$  and its dual  $\check{\theta}_0$  as follows,

(40) 
$$\theta_0 = \begin{cases} \text{highest root of } \mathfrak{g}, & \text{if } r = 1 \\ \text{highest short root of } \mathfrak{g}^{\tau}, & \text{if } r > 1 \text{ and } (\mathfrak{g}, r) \neq (A_{2n}, 2) \\ 2 \cdot \text{highest short root of } \mathfrak{g}^{\tau}, & \text{if } (\mathfrak{g}, r) = (A_{2n}, 2). \end{cases}$$

(41) 
$$\check{\theta}_0 = \begin{cases} \text{highest short coroot of } \mathfrak{g} & \text{if } r = 1\\ \text{highest coroot of } \mathfrak{g}^{\tau} & \text{if } r > 1 \text{ and } (\mathfrak{g}, r) \neq (A_{2n}, 2)\\ \frac{1}{2} \cdot \text{highest coroot of } \mathfrak{g}^{\tau} & \text{if } (\mathfrak{g}, r) = (A_{2n}, 2) \end{cases}$$

Recall the normalized symmetric bilinear form  $(\cdot | \cdot)$  on  $\hat{\mathfrak{h}}^{\tau}$  in [Kac, §6.1]. This bilinear form induces a map  $\nu : \hat{\mathfrak{h}}^{\tau} \to (\hat{\mathfrak{h}}^{\tau})^*$ , which satisfies the property in (4). By [Kac, (6.2.2) and Proposition 5.1], we have

(42) 
$$(\theta_0 \mid \theta_0) = 2a_o \check{a}_o = 2\check{a}_o; \quad \nu(\check{\theta}_0) = \frac{\theta_0}{a_o \check{a}_o} = \frac{\theta_0}{\check{a}_o}$$

Let  $\check{P}$  (resp.  $\check{Q}$ ) denote the coweight (resp. coroot) lattice of g. Let  $\check{P}_{\tau}$  (resp.  $\check{Q}_{\tau}$ ) denote the set of  $\tau$ -coinvariants of  $\check{P}$  (resp.  $\check{Q}$ ). We have the following well-defined map

$$\check{P}_{\tau} \to (\mathfrak{h}^{\tau})^*, \quad \mu \mapsto \iota(\tilde{\mu})|_{\mathfrak{h}^{\tau}},$$

where  $\tilde{\mu}$  is a lifting of  $\mu$  in  $\check{P}$ , and  $\iota : \mathfrak{h} \to \mathfrak{h}^*$  is induced from the normalized Killing form  $(\cdot, \cdot)$  on g. We still denote this map by  $\iota$  if there is no confusion.

Let  $A: \mathfrak{h} \to \mathfrak{h}^{\tau}$  be the map defined by  $A(h) := \sum_{i=0}^{r-1} \tau^i(h)$ , for any  $h \in \mathfrak{h}$ . This map descends to a map  $\check{P}_{\tau} \to \mathfrak{h}^{\tau}$ , which is still denoted by A if there is no confusion.

**Lemma 4.1.** For any  $\mu \in \check{P}_{\tau}$ , we have  $\nu(A(\mu)) = \iota(\mu)$ .

*Proof.* For any  $h \in \mathfrak{h}^{\tau}$ , by [Kac, Corollary 6.4], we have

$$\langle h, \nu(A(\mu)) \rangle := \langle h, \nu(A(\tilde{\mu})) \rangle = (h \mid A(\tilde{\mu})) = \frac{1}{r}(h, A(\tilde{\mu})) = (h, \tilde{\mu}) = \langle h, \iota(\tilde{\mu}) \rangle =: \langle h, \iota(\mu) \rangle,$$
 where  $\tilde{\mu} \in \check{P}$  is a lifting of  $\mu$ . Therefore,  $\nu(A(\mu)) = \iota(\mu)$ .

4.2. The coherence conjecture and affine Demazure modules. Let G be the adjoint group associated to g. We fix a torus T and a Borel subgroup B corresponding to the Lie subalgebras  $\mathfrak{h} \subseteq \mathfrak{b}$ .

We follow the notations in Section 3.1. Let  $W^{\tau}$  be the Weyl group of  $\mathfrak{g}^{\tau}$ . We can identify  $\check{P}$  (resp.  $\check{Q}$ ) with the set  $X_*(T)$  (resp.  $X_*(T')$ ). Recall the affine Weyl group  $W_{\rm aff} = \check{Q}_{\tau} \rtimes W^{\tau}$  and the extended affine Weyl group  $\widetilde{W}_{\rm aff} = X_*(T)_{\tau} \rtimes W^{\tau}$ . The elements in  $\widetilde{W}_{\rm aff}$  are written as  $\varrho^{\mu}w$  for  $\mu \in X_*(T)_{\tau}$  and  $w \in W^{\tau}$ .

We regard  $W_{\text{aff}}$  as the Weyl group of  $\hat{L}(g, \tau)$ , whose action on  $(\hat{h}^{\tau})^*$  is given by

(43) 
$$\varrho^{\mu}w(x) = w(x) - \langle w(x), K \rangle \iota(\mu) + \left( (w(x) | \iota(\mu)) - \frac{1}{2} (\iota(\mu) | \iota(\mu)) \langle w(x), K \rangle \right) \delta$$

for any  $\varrho^{\mu}w \in \tilde{W}_{aff}$ ,  $x \in (\hat{\mathfrak{h}}^{\tau})^*$ , cf. [Kac, (6.5.2)].

For any dominant integrable weight  $\Lambda$  of  $\hat{L}(\mathfrak{g}, \tau)$ , we denote by  $V(\Lambda)$  the irreducible highest weight integrable representation of  $\hat{L}(\mathfrak{g}, \tau)$ . For any  $\bar{\mu} \in X_*(T)_\tau$ , there exists a unique element  $\varsigma_{\kappa} \in \Omega$  such that  $\varrho^{\bar{\mu}} \varsigma_{\kappa}^{-1} \in W_{\mathrm{aff}}$ . Let  $v_{\varrho^{\bar{\mu}} \varsigma_{\kappa}^{-1}(\Lambda)} \in V(\Lambda)$  be an extremal weight vector of weight  $\varrho^{\bar{\mu}} \varsigma_{\kappa}^{-1}(\Lambda)$ .

**Definition 4.2.** For any  $\bar{\mu} \in X_*(T)_\tau$  and dominant weight  $\Lambda$  of  $\hat{L}(\mathfrak{g}, \tau)$ , we define the affine Demazure module  $D^{\tau}(\Lambda, \bar{\mu})$  as follows,

$$D^{\tau}(\Lambda, \bar{\mu}) := \mathrm{U}(\hat{\mathfrak{b}}^{\tau}) \cdot \nu_{\rho^{\bar{\mu}} \varsigma_{\kappa}^{-1}(\Lambda)} \subseteq V(\Lambda),$$

where  $\hat{\mathfrak{b}}^{\tau} := \mathfrak{b}^{\tau} \oplus (t\mathfrak{g}[[t]])^{\tau} \oplus kK \oplus kd$  is a Borel subalgebra of  $\hat{L}(\mathfrak{g}, \tau)$ .

When  $\tau$  is trivial, for any coweight  $\mu \in X_*(T)$  and dominant weight  $\Lambda$  of  $\hat{L}(\mathfrak{g})$ , the Demazure module will be denoted by  $D(\Lambda, \mu)$ . In particular, when  $\mu$  is dominant,  $D(c\Lambda_{\kappa}, \mu)$  is  $\mathfrak{g}[t]$ -stable, where  $\Lambda_{\kappa} = \Lambda_o + \iota(\kappa)$ . We also denote  $D(c\Lambda_{\kappa}, \mu)$  by  $D(c, \mu)$ .

Remark 4.3. The  $\mathfrak{g}[t]$ -stable Demazure module  $D(c, \mu)$  has a lowest weight vector whose  $\mathfrak{h}$  weight is  $-c\iota(\mu)$ , see [BH2], [HY].

Let c be the level of the dominant weight  $\Lambda$ , i.e.  $c = \langle \Lambda, K \rangle$ . Let  $\lambda$  be the restriction of  $\Lambda$  to  $\mathfrak{h}^{\tau}$ . Consider the element  $\vartheta = \frac{1}{c} \nu^{-1}(\lambda) \in \mathfrak{h}^{\tau}$ . Define  $\mathfrak{g}_{\lambda} := Z_{\mathfrak{g}^{\tau}}(\vartheta)$  to be the centralizer of  $\vartheta$  in  $\mathfrak{g}^{\tau}$ . We write  $\Lambda = \sum_{i \in \hat{I}_{\tau}} n_i \Lambda_i$ . For any  $\bar{\mu} \in X_*(T)_{\tau}$ , there is a unique element  $\mathcal{G}_{\kappa} \in \Omega$  so that  $\varrho^{\bar{\mu}} \mathcal{G}_{\kappa}^{-1} \in W_{\text{aff}}$ . Set

$$Y_{\kappa} := \{ i \in \hat{I}_{\tau} \mid n_i \neq 0 \}, \quad Y := \varsigma_{\kappa}^{-1}(Y_{\kappa}).$$

**Lemma 4.4.** The element  $\vartheta$  is an rational interior point in the facet  $F_{Y_{\kappa}}$  associated to  $Y_{\kappa}$ . Further more, we have  $\text{Lie}(G_{\vartheta}) = \mathfrak{g}_{\lambda}$ .

*Proof.* By direct computation, we have

$$\vartheta = \frac{1}{c} v^{-1}(\lambda) = \sum_{i \in I_{\tau}} \frac{n_i}{c} v^{-1}(\omega_i) = \sum_{i \in I_{\tau}} \frac{n_i}{c} \cdot \frac{\check{\alpha}_i}{a_i} \check{\omega}_i = \sum_{i \in Y_{\kappa}} \frac{n_i}{c} \cdot \frac{\check{\alpha}_i}{a_i} \check{\omega}_i,$$

where by convention  $\check{\omega}_o = 0$ . Since  $\sum_{i \in Y_\kappa} \frac{n_i \check{u}_i}{c} = 1$ ,  $\vartheta$  is an rational interior point in the facet  $F_{Y_\kappa}$  spanned by  $\{\frac{\check{\omega}_i}{a_i} \mid i \in Y_\kappa\}$ . By (15), we have

$$\operatorname{Lie}(G_{\vartheta}) = Z_{\mathfrak{q}^{\tau}}(\frac{m}{r}\vartheta) = \mathfrak{g}_{\lambda},$$

where m is the minimal positive multiple of r so that  $\frac{m}{r}\vartheta \in X_*(T_{ad})^{\tau}$ .

Now, we state the main theorem of this section.

**Theorem 4.5.** For any dominant coweight  $\mu \in X_*(T)$ , there is an isomorphism of  $\mathfrak{g}_{\lambda}$ -modules

$$D(c,\mu) \otimes \mathbf{k}_{\lambda} \simeq \sum_{w \in W^{\mathsf{T}}} D^{\mathsf{T}}(\Lambda, w(\bar{\mu})),$$

where  $k_{\lambda}$  is the 1-dimensional representation of  $g_{\lambda}$ .

Consider the partial affine flag variety  $\operatorname{Fl}_Y$  corresponding to Y. Recall (22), we have an isomorphism  $\operatorname{Fl}_Y^{\kappa} \simeq \operatorname{Fl}_{Y_{\kappa}}'$  of homogeneous spaces of  $G'((z))^{\tau}$ . Moreover,  $\overline{\operatorname{Fl}}_{Y_{\kappa}}$  is contained in  $\operatorname{Fl}_Y^{\kappa}$ . Let  $\mathscr{L}'(\Lambda)$  be the ample line bundle on  $\operatorname{Fl}_{Y_{\kappa}}'$  associated to  $\Lambda$ . Pulling back via the isomorphism (22), we get an line bundle on  $\operatorname{Fl}_Y^{\kappa}$ , denoted by  $\mathscr{L}(\Lambda)$ .

By the affine Borel-Weil theorem (due to Kumar and Mathieu), we have

**Theorem 4.6.** (1) For any  $\bar{\mu} \in X_*(T)_{\tau}$ , there is a isomorphism of  $\hat{\mathfrak{b}}^{\tau}$ -modules

$$H^0(\overline{\mathrm{Fl}}_{Y,\bar{\mu}},\mathcal{L}(\Lambda))^{\vee} \simeq D^{\tau}(\Lambda,\bar{\mu}).$$

(2) For any dominant coweight  $\mu \in X_*(T)$ , there is an isomorphism of  $\mathfrak{g}[t]$ -modules

$$H^0(\overline{\operatorname{Gr}}_{G,\mu}, L^c)^{\vee} \simeq D(c,\mu),$$

where L denotes the level one line bundle on  $Gr_G$ .

*Proof.* (1) Recall that the isomorphism (22) restricts to an isomorphism  $\overline{Fl}_{Y,\bar{\mu}} \simeq \overline{Fl}'_{Y_{\kappa},\mathcal{Q}^{\bar{\mu}}\mathcal{S}_{\kappa}^{-1}}$ , see (25). By [Ku2, Theorem 8.2.2 (a)] and [PR, §9f], the space

$$H^{0}(\overline{\mathrm{Fl}}_{Y,\bar{\mu}},\mathcal{L}(\Lambda))^{\vee} \simeq H^{0}(\overline{\mathrm{Fl}}'_{Y_{\nu},o^{\bar{\mu}}\varsigma^{-1}_{\nu}},\mathcal{L}'(\Lambda))^{\vee}$$

is isomorphic to the affine Demazure module  $D^{\tau}(\Lambda, \bar{\mu})$ .

(2) Let  $\tau$  be tivial and take  $Y = \{o\}$ . Then,  $\operatorname{Fl}_Y$  is the affine Grassmannian  $\operatorname{Gr}_G$ . Moreover, for any dominant coweight  $\mu \in X_*(T)$ , we have

$$\overline{\mathrm{Gr}}_{G,\mu} = {}_{Y}\overline{\mathrm{Fl}}_{Y,\varrho^{\mu}} = \overline{\mathrm{Fl}}_{Y,\mu},$$

where the first equality is by definition and the second equality follows from Corollary 4.17. Let  $Gr_G^{\kappa} := Fl_Y^{\kappa}$  be the connected component containing  $\overline{Gr}_{G,\mu}$  (equivalently  $\varrho^{\mu} \varsigma_{\kappa}^{-1} \in W_{\text{aff}}$ ). Note that the restriction of L to  $Gr_G^{\kappa}$  is the line bundle  $\mathscr{L}(\Lambda_{\kappa})$  on  $Fl_Y^{\kappa}$ . By part (1), we get

$$H^0(\overline{\operatorname{Gr}}_{G,\mu}, L^c)^{\vee} = H^0(\overline{\operatorname{Fl}}_{Y,\mu}, \mathscr{L}(c\Lambda_{\kappa}))^{\vee} \simeq D(c,\mu),$$

where  $D(c, \mu) := D(c\Lambda_{\kappa}, \mu)$ .

Recall the following variety defined in (20),

$$\mathcal{A}_Y(\bar{\mu}) := \bigcup_{w \in W^{\tau}} \overline{\mathrm{Fl}}_{Y,w(\bar{\mu})} \subseteq \mathrm{Fl}_Y^{\kappa}.$$

**Theorem 4.7.** Let  $\bar{\mu} \in X_*(T)_{\tau}$ .

(1) The following restriction map is surjective

$$H^0(\mathrm{Fl}_{\mathrm{v}}^{\kappa}, \mathcal{L}(\Lambda)) \twoheadrightarrow H^0(\mathcal{A}_{\mathrm{v}}(\bar{\mu}), \mathcal{L}(\Lambda)).$$

(2)  $H^i(\mathcal{A}_V(\bar{\mu}), \mathcal{L}(\Lambda)) = 0$  for  $i \geq 1$ .

*Proof.* Note that any Schubert variety  $\overline{Fl}_{Y,\bar{\mu}}$  is normal (cf. [PR, Theorem 0.3]), and by [PR, §9.h] and [Ku2, Theorem 8.2.2],  $H^i(\overline{Fl}_{Y,\bar{\mu}}, \mathcal{L}(\Lambda)) = 0$  when  $i \geq 1$ . For the union  $\mathcal{H}_Y(\bar{\mu})$ , this theorem follows from the same argument as in [LLM, Theorem 6.1, Corollary 6.2].

Note that the above theorem also holds when k is of positive characteristic p and  $p \neq r$ . This follows from the fact that  $\mathcal{A}_Y(\bar{\mu})$  is Frobenius splitting, cf. [PR].

**Proof of Theorem 4.5.** We first note that there is an injective map

$$H^0(\mathcal{A}_Y(\bar{\mu}), \mathcal{L}(\Lambda)) \hookrightarrow \bigoplus_{w \in W^{\tau}} H^0(\overline{\mathrm{Fl}}_{Y,w(\bar{\mu})}, \mathcal{L}(\Lambda)).$$

Composing with the map in part (1) of Theorem 4.7 and taking dual, we get

$$\bigoplus_{w \in W_{\mathcal{T}}} H^0(\overline{\mathrm{Fl}}_{Y,w(\bar{\mu})}, \mathcal{L}(\Lambda))^{\vee} \twoheadrightarrow H^0(\mathcal{R}_Y(\bar{\mu}), \mathcal{L}(\Lambda))^{\vee} \hookrightarrow H^0(\mathrm{Fl}_Y^{\kappa}, \mathcal{L}(\Lambda))^{\vee}.$$

Clearly, the image of the composition map is

$$\sum_{w \in W^{\mathsf{T}}} \mathrm{Im} \Big( H^{0}(\overline{\mathrm{Fl}}_{Y,w(\bar{\mu})}, \mathscr{L}(\Lambda))^{\vee} \hookrightarrow H^{0}(\mathrm{Fl}_{Y}^{\kappa}, \mathscr{L}(\Lambda))^{\vee} \Big).$$

We regard  $H^0(\overline{\mathrm{Fl}}_{\mathrm{Y}_W(\overline{\mu})}, \mathcal{L}(\Lambda))^{\vee}$  as a subspace of  $H^0(\mathrm{Fl}_{\mathrm{Y}}^{\kappa}, \mathcal{L}(\Lambda))^{\vee}$ , cf [Ku2, Theorem 8.2.2 (d)]. Then,

$$H^0(\mathcal{A}_Y(\bar{\mu}),\mathcal{L}(\Lambda))^\vee\simeq \sum_{w\in W^\tau}H^0(\overline{\mathrm{Fl}}_{Y,w(\bar{\mu})},\mathcal{L}(\Lambda))^\vee\subseteq H^0(\mathrm{Fl}_Y^\kappa,\mathcal{L}(\Lambda))^\vee.$$

Now, the theorem follows from Theorem 3.5 and Theorem 4.6.

4.3. Inclusion relation among Demazure modules. Recall the element  $\check{\theta}_0 \in \mathfrak{h}^{\tau}$  defined in (41) and the map  $A: X_*(T)_{\tau} \to \mathfrak{h}^{\tau}$  in Section 4.1. Let  $\check{M}$  be the lattice spanned by  $W^{\tau} \cdot \check{\theta}_0$ . Then, A restricts to an isomorphism  $\check{Q}_{\tau} \simeq \check{M}$ , see the proof in [DH1, Proposition A.1.8]. Let  $\mu_0 \in \check{Q}_{\tau}$  be the element such that  $A(\mu_0) = \check{\theta}_0 \in \check{M}$ . In fact, we have the following description of  $\mu_0$ . Recall the root  $\theta^0$  of g introduced in Section 2.3 (also in [Kac, §8.3]). We have  $\theta^0|_{\mathfrak{h}^r} = \theta_0$ , see (14). Let  $\check{\theta}^0$  denote the coroot of  $\theta^0$ . Define the following element in  $\check{Q}$ ,

$$\mu^{0} := \begin{cases} \check{\theta}_{0}, & \text{when } r = 1\\ \check{\theta}^{0}, & \text{when } r \neq 1 \text{ and } X_{N}^{(r)} \neq A_{2\ell}^{(2)}\\ \check{\alpha}_{1} + \dots + \check{\alpha}_{\ell}, & \text{when } X_{N}^{(r)} = A_{2\ell}^{(2)}, \end{cases}$$

where we label the vertices of  $g = A_{2\ell}$  by  $1, \ldots, 2\ell$ .

**Lemma 4.8.** Let  $\overline{\mu^0}$  denote the image of  $\mu^0$  under the map  $X_*(T) \to X_*(T)_\tau$ . Then,

$$\mu_0 = \overline{\mu^0}$$
.

*Proof.* Since  $A: \check{Q}_{\tau} \simeq \check{M}$  is an isomorphism, it suffices to show that  $A(\overline{\mu^0}) = \check{\theta}_0$ .

When r=1, the map A is the identity map. Hence,  $A(\overline{\mu^0})=\overline{\mu^0}=\mu^0=\check{\theta}_0$ . When r>1 and  $X_N^{(r)}\neq A_{2\ell}^{(2)}$ , since g is simply-laced, we have  $(\theta^0,\theta^0)=2$  under the normalized Killing form on g. Note that  $\langle \check{\theta}^0,\theta^0\rangle=2$ . We get  $\iota(\check{\theta}^0)=\theta^0$  and

$$\nu(A(\overline{\mu^0})) = \iota(\overline{\mu^0}) = \iota(\mu^0)|_{b^{\tau}} = \iota(\check{\theta}^0)|_{b^{\tau}} = \theta^0|_{b^{\tau}} = \theta_0,$$

where the first equality follows from Lemma 4.1 and the second equality is by definition. Therefore,  $A(\mu^0) = \nu^{-1}(\theta_0) = \check{\theta}_0$ , where the second equality follows from (42).

When  $X_N^{(r)} = A_{2\ell}^{(2)}$ , the simple coroots of  $\mathfrak{g}^{\tau}$  are  $\check{\beta}_i = \check{\alpha}_i + \check{\alpha}_{2\ell+1-i}$ ,  $(1 \leq i < \ell)$ , and  $\check{\beta}_{\ell} = 2(\check{\alpha}_{\ell} + \check{\alpha}_{\ell+1})$ , see [Kac, §8.3 Case 5]. Hence,

$$A(\overline{\mu^0}) = \mu^0 + \tau(\mu^0) = \check{\alpha}_1 + \dots + \check{\alpha}_{2\ell} = \frac{1}{2} \cdot (2\check{\beta}_1 + \dots + 2\check{\beta}_{\ell-1} + \check{\beta}_{\ell}) = \check{\theta}_0.$$

Remark 4.9. By [BH2], [Zh2], we can regard  $X_*(T)_\tau$  the weight lattice of the reductive group  $(G^{\vee})^{\tau}$  where  $G^{\vee}$  is the Langlands dual group of G and  $\tau$  is the diagram automorphism on  $G^{\vee}$ . Then,  $\mu_0$  is the highest short root of  $(G^{\vee})^{\tau}$ .

Recall the affine Weyl group  $W_{\text{aff}} = \check{Q}_{\tau} \rtimes W^{\tau}$ . For any  $i \in I_{\tau}$ , let  $r_i \in W^{\tau}$  denote the simple reflection associated to i. Set

$$r_o := \varrho^{-\mu_0} r_{\theta_0} \in W_{\text{aff}}.$$

Then,  $W_{\text{aff}}$  is generated by the simple reflections  $\{r_i | i \in \hat{I}_{\tau}\}$ . By (43), the action of  $r_o$  on  $(\hat{\mathfrak{h}}^{\tau})^*$  is given by  $r_o(x) = x - \langle \check{\alpha}_o, x \rangle \alpha_o$ .

As in Section 3.1, we fix a non-empty subset  $Y \subseteq \hat{I}_{\tau}$ . Set  $\bar{Y} = \hat{I}_{\tau} \setminus Y$ . Recall the partial affine flag variety  $\mathrm{Fl}_Y$  and the parabolic subgroup  $W_{\bar{Y}}$  of  $W_{\mathrm{aff}}$  generated by the simple reflections  $r_i$ ,  $i \in \bar{Y}$ . Let  $\leq$  be the Bruhat order on  $W_{\mathrm{aff}}$ .

**Definition 4.10.** For each element  $w \in W_{\text{aff}}$ , the coset  $wW_{\bar{Y}}$  has a unique element with minimal length, denoted by  $w_{\min}$ . For any  $w, v \in W_{\text{aff}}$ , we define  $w \leq_Y v$  if  $w_{\min} \leq v_{\min}$ .

Let  $\Lambda = \sum_{i \in Y} n_i \Lambda_i$  be a dominant integrable weight of  $\hat{L}(\mathfrak{g}, \tau)$  with  $n_i > 0$ . For any  $\mu \in \check{Q}_{\tau}$ , recall the affine Schubert variety  $\overline{\mathrm{Fl}}_{Y,\mu}$  and the affine Demazure module  $D^{\tau}(\Lambda, \mu)$ . The following result is well-known, cf. [BH1, Proposition 3.4 and 3.7]

**Proposition 4.11.** Given any  $\mu_1, \mu_2 \in \check{Q}_{\tau}$ , the following are equivalent:

- (1)  $\varrho^{\mu_1} \leq_Y \varrho^{\mu_2}$ ,
- (2)  $D^{\tau}(\Lambda, \mu_1) \subseteq D^{\tau}(\Lambda, \mu_2)$ ,
- $(3) \ \overline{\mathrm{Fl}}_{Y,\mu_1} \subseteq \overline{\mathrm{Fl}}_{Y,\mu_2}.$

**Definition 4.12.** For any  $w \in W_{\text{aff}}$ , let  $\bar{w} \in W^{\tau}$  denote its image under the composition map  $W_{\text{aff}} \to \check{Q}_{\tau} \backslash W_{\text{aff}} \simeq W^{\tau}$ . For example,  $\bar{r}_o = r_{\theta_0}$  and  $\bar{r}_i = r_i$  for  $i \neq o$ .

**Lemma 4.13.** Let  $\mu \in \check{Q}_{\tau}$ .

- (1) For any  $w \in W_{\text{aff}}$ , we have  $w\varrho^{\mu}w^{-1} = \varrho^{\bar{w}(\mu)}$ .
- (2) Recall the map  $\iota: \check{Q}_{\tau} \to (\mathfrak{h}^{\tau})^*$ . We have

$$r_i(\iota(\mu)) = \begin{cases} \iota(\bar{r}_i(\mu)) & \text{if } i \neq o, \\ \iota(\bar{r}_o(\mu)) + \langle \check{\theta}_0, \iota(\mu) \rangle \delta & \text{if } i = o. \end{cases}$$

*Proof.* For any  $w \in W_{\text{aff}}$ , we have  $w = \varrho^{\eta} \bar{w}$  for some  $\eta \in \check{Q}_{\tau}$ . Then,

$$w\varrho^\mu w^{-1} = \varrho^\eta \bar w \varrho^\mu \bar w^{-1} \varrho^{-\eta} = \varrho^\eta \varrho^{\bar w(\mu)} \varrho^{-\eta} = \varrho^{\bar w(\mu)}.$$

This proves part (1). Part (2) follows from the equation (43) and the fact that  $w(\iota(\mu)) = \iota(w(\mu))$  for any  $w \in W^{\tau}$ .

**Lemma 4.14.** Given any  $\mu \in \check{Q}_{\tau}$  and  $i \in \bar{Y}$ , we have  $r_i \varrho^{\mu} \leq_Y \varrho^{\mu}$  if and only if  $\langle \check{\alpha}_i, \iota(\mu) \rangle \geq 0$ . Furthermore,  $(r_i \varrho^{\mu})_{\min} = (\varrho^{\mu})_{\min}$  if and only if  $\langle \check{\alpha}_i, \iota(\mu) \rangle = 0$ , where  $\check{\alpha}_i$  is the simple coroot associated to i.

*Proof.* We first assume that  $\langle \check{\alpha}_i, \iota(\mu) \rangle \geq 0$ . Then,  $(\alpha_i | \iota(\mu)) \geq 0$ . By (43), we have  $\varrho^{\mu}(\alpha_i) = \alpha_i + (\alpha_i | \iota(\mu))\delta$ , which is a positive affine root. By [Kac, Lemma 3.11 (a)],  $\ell(\varrho^{\mu}r_i) > \ell(\varrho^{\mu})$ . Since we always have  $\ell(\varrho^{\mu}r_i) = \ell(\varrho^{\mu}) \pm 1$ , it follows that  $\ell(\varrho^{\mu}r_i) = \ell(\varrho^{\mu}) + 1$ . Note that  $\ell(r_i\varrho^{\mu}r_i) = \ell(\varrho^{\bar{r}_i\mu}) = \ell(\varrho^{\mu}) = \ell(\varrho^{\mu}) = \ell(\varrho^{\mu}r_i) - 1$ . Thus,  $r_i\varrho^{\mu}r_i < \varrho^{\mu}r_i$ . By [Ku2, Lemma 1.3.18], we get  $(r_i\varrho^{\mu}r_i)_{\min} \leq (\varrho^{\mu}r_i)_{\min}$ . Thus,  $(r_i\varrho^{\mu})_{\min} \leq (\varrho^{\mu})_{\min}$ , i.e.  $r_i\varrho^{\mu} \leq r_i$ 

Conversely, assume that  $r_i\varrho^{\mu} \leq_Y \varrho^{\mu}$ . Suppose that  $(\iota(\mu) \mid \alpha_i) < 0$ . Then  $\varrho^{\mu}(\alpha_i) = \alpha_i + (\alpha_i \mid \iota(\mu))\delta$  is a negative affine root. By [Kac, Lemma 3.11 (a)],  $\ell(\varrho^{\mu}r_i) < \ell(\varrho^{\mu})$ .

Therefore, we have  $\ell(\varrho^{\mu}r_i) = \ell(\varrho^{\mu}) - 1$ , and hence  $(\varrho^{\mu})_{\min} \leq (r_i\varrho^{\mu})_{\min}$ . Combining with our assumption that  $r_i\varrho^{\mu} \leq_Y \varrho^{\mu}$ , we must have  $(\varrho^{\mu})_{\min} = (r_i\varrho^{\mu})_{\min}$ . Then,  $\varrho^{\mu} = r_i\varrho^{\mu}w$  for some  $w \in W_{\bar{Y}}$ . By part (1) of Lemma 4.13,  $\varrho^{\mu} = \varrho^{\bar{r}_i(\mu)}w'$  for some  $w' \in W_{\bar{Y}}$ . Thus,  $\varrho^{\mu-\bar{r}_i(\mu)} \in W_{\bar{Y}}$ . Note that  $W_{\bar{Y}}$  is a finite group and every element has finite order. It follows that  $\mu - \bar{r}_i(\mu) = 0$ . By part (2) of Lemma 4.13, we conclude that  $(\iota(\mu) - r_i(\iota(\mu)) | \alpha_i) = 0$ . Hence,  $\langle \check{\alpha}_i, \iota(\mu) \rangle = 0$ . This contradicts with our assumption that  $(\iota(\mu) | \alpha_i) < 0$ . Therefore,  $(\iota(\mu) | \alpha_i) \geq 0$ , equivalently  $\langle \check{\alpha}_i, \iota(\mu) \rangle \geq 0$ . This conclude the proof of the first statement in the Lemma.

From above argument, we have shown that  $(\varrho^{\mu})_{\min} = (r_i \varrho^{\mu})_{\min}$  implies  $\langle \check{\alpha}_i, \iota(\mu) \rangle = 0$ . Conversely, if  $\langle \check{\alpha}_i, \iota(\mu) \rangle = 0$ , then  $r_i(\iota(\mu)) = \iota(\mu)$ . By Lemma 4.13, we get  $\iota(\bar{r}_i(\mu)) = \iota(\mu)$ . Hence,  $\bar{r}_i(\mu) = \mu$ . It follows that  $r_i \varrho^{\mu} r_i = \varrho^{\bar{r}_i(\mu)} = \varrho^{\mu}$ . Hence,  $(r_i \varrho^{\mu})_{\min} = (\varrho^{\mu})_{\min}$ .

We remark that  $\langle \check{\alpha}_o, \iota(\mu) \rangle = \langle -\check{\theta}_0, \iota(\mu) \rangle$  for any  $\mu \in \check{Q}_{\tau}$ .

**Definition 4.15.** Define  $\beta_i := \alpha_i$  for  $i \neq o$  and  $\beta_o := -\theta_0$ . Then,  $\{\beta_i\}_{i \in \bar{Y}}$  form a base of a root system, denoted by  $\Phi_{\bar{Y}}$ , where  $\bar{Y} := \hat{I}_{\tau} \setminus Y$ . Let  $\Phi_{\bar{Y}}^+$  and  $\Phi_{\bar{Y}}^-$  denote the sets of positive roots and negative roots of  $\Phi_{\bar{Y}}$  with respect to the base  $\{\beta_i\}_{i \in \bar{Y}}$ .

**Proposition 4.16.** Let  $\mu \in \check{Q}_{\tau}$  and  $w \in W_{\bar{Y}}$ . If  $\langle \check{\beta}, \iota(\mu) \rangle \geq 0$  for any  $\beta \in \Phi_{\bar{Y}}^+$  with  $\bar{w}(\beta) \in \Phi_{\bar{Y}}^-$ , then we have

- (1)  $\varrho^{\bar{w}(\mu)} \leq_Y \varrho^{\mu}$ ,
- (2)  $D^{\tau}(\Lambda, \bar{w}(\mu)) \subseteq D^{\tau}(\Lambda, \mu)$ ,
- (3)  $\overline{\mathrm{Fl}}_{Y,\bar{w}(\mu)} \subseteq \overline{\mathrm{Fl}}_{Y,\mu}$ .

*Proof.* By Proposition 4.11, it suffices to prove part (1). Let  $W_{\bar{Y},f}$  be a subgroup of  $W^{\tau}$  defined as follows,

$$W_{\bar{Y},f} := \{ \bar{w} \mid w \in W_{\bar{Y}} \}.$$

Then,  $W_{\bar{Y},f}$  is generated by  $\{\bar{r}_i | i \in \bar{Y}\}$ . Note that for any  $i \in \bar{Y}$  and  $\beta \in \Phi_{\bar{Y}}$ , we have  $\bar{r}_i(\beta) = \beta - \langle \check{\beta}_i, \beta \rangle \beta_i$ . Hence,  $W_{\bar{Y},f}$  is the Weyl group of  $\Phi_{\bar{Y}}$  with simple generators  $\{\bar{r}_i | i \notin Y\}$ . Let  $\bar{w} = \bar{r}_{i_n} \cdots \bar{r}_{i_1}$  be a reduced word in  $W_{\bar{Y},f}$ , where  $i_1, \dots, i_n \in \bar{Y}$ . Set  $\bar{w}_0 = e$  and  $\bar{w}_k := \bar{r}_{i_k} \bar{w}_{k-1}$  for  $1 \le k \le n$ . Then, for any k, we have

$$\bar{w}\bar{w}_{k-1}^{-1}(\beta_{i_k})=\bar{r}_{i_n}\cdots\bar{r}_{i_k}(\beta_{i_k})\in\Phi_{\bar{Y}}^-.$$

By assumption,  $\langle \bar{w}_{k-1}^{-1}(\check{\beta}_{i_k}), \iota(\mu) \rangle \geq 0$ . By Lemma 4.14, we get  $r_{i_k}\varrho^{\bar{w}_{k-1}\mu} \leq_Y \varrho^{\bar{w}_{k-1}\mu}$ . Multiplying  $r_{i_k}$  on the right, we have  $\varrho^{\bar{w}_k\mu} \leq_Y \varrho^{\bar{w}_{k-1}\mu}$ . It follows that

$$\varrho^{\bar{w}(\mu)} = \varrho^{\bar{w}_n(\mu)} \leq_Y \cdots \leq_Y \varrho^{\bar{w}_1(\mu)} \leq_Y \varrho^{\mu}.$$

We say  $\mu \in \check{Q}_{\tau}$  is  $\bar{Y}$ -dominant if  $\langle \check{\beta}, \iota(\mu) \rangle \geq 0$  for any  $\beta \in \Phi_{\bar{Y}}^+$ . For any  $\mu \in \check{Q}_{\tau}$ , let  $S_{Y,\mu}$  be the following subset of  $\check{Q}_{\tau}$ ,

$$S_{Y,\mu} := \{ \eta \in W^{\tau} \mu \mid \eta \text{ is } \bar{Y}\text{-dominant} \}.$$

Corollary 4.17. Let  $\mu \in \check{Q}_{\tau}$ .

(1) Assume that  $\mu$  is  $\overline{Y}$ -dominant. For any  $w \in W_{\overline{Y}}$ , we have  $\overline{Fl}_{Y,\overline{w}(\mu)} \subseteq \overline{Fl}_{Y,\mu}$ . Moreover,  ${}_{Y}\overline{Fl}_{Y,\rho^{\mu}} = \overline{Fl}_{Y,\mu}$ .

(2) The union in (20) can be simplified as follows,

$$\mathcal{A}_{Y}(\mu) = \bigcup_{\eta \in S_{Y,\mu}} \overline{\mathrm{Fl}}_{Y,\eta}.$$

(3) Theorem 4.5 can be restated as follows. There is an isomorphism of  $g_{\lambda}$ -modules

$$D(c,\mu) \otimes \mathbf{k}_{\lambda} \simeq \sum_{\eta \in S_{Y,\mu}} D^{\tau}(\Lambda,\eta),$$

where  $k_{\lambda}$  is the 1-dimensional representation of  $g_{\lambda}$ .

*Proof.* Part (1) follows from Proposition 4.16, and the fact that  ${}_Y\overline{Fl}_{Y,\mathcal{Q}^\mu}=\bigcup_{w\in W_{\bar{Y}}}\overline{Fl}_{Y,\bar{w}(\mu)}$ , see (21) in Lemma 3.4. For part (2), it suffices to show that the left hand side is contained in the right hand side. Since  $W_{\bar{Y},f}$  is the Weyl group of  $\Phi_{\bar{Y}}$ , for any  $\eta\in \check{Q}_{\tau}$ , there exists a  $\bar{w}\in W_{\bar{Y},f}$  such that  $\bar{w}(\eta)$  is Y-dominant. By part (1), we have  $\overline{Fl}_{Y,\eta}\subseteq \overline{Fl}_{Y,\bar{w}(\eta)}$ . Hence,

$$\mathcal{A}_{Y}(\mu) = \bigcup_{\eta \in W^{\tau}(\mu)} \overline{\mathrm{Fl}}_{Y,\eta} \subseteq \bigcup_{\eta \in S_{Y,\mu}} \overline{\mathrm{Fl}}_{Y,\eta}.$$

This proves part (2). The third part can be proved in the same way.

5. Canonical  $L^+\mathcal{G}$ -equivariance on line bundles over  $\mathrm{Gr}_{\mathcal{G}}$ 

In this section, we always assume char(k) = 0.

5.1. **Picard group of**  $Gr_{\mathcal{G}}$ . Let G be a simply-connected simple algebraic group over  $\mathbb{R}$ . As in Section 3, we fix a non-empty subset Y of  $\hat{I}_{\tau}$ . Let  $\mathcal{G}_{Y}$  be the parahoric group scheme associated to Y, and let G be the parahoric Bruhat-Tits group scheme over  $\mathbb{R}^1$  glued from  $\mathcal{G}_{Y}$  and  $Res_{\mathbb{A}^1-0/\bar{\mathbb{A}}^1-0}(G_{\mathbb{A}^1-0})^{\tau}$ . Let  $Gr_{\mathcal{G}}$  be the global affine Grassmannian of G over  $\mathbb{A}^1$  and let  $L^+G$  be the jet group scheme of G over  $\mathbb{A}^1$ , see Definition 3.17.

Let X be a projective scheme flat over k-scheme S. Let  $e: S \to X$  be a section. A rigidified line bundle on X with respect to e is a pair  $(L, \eta)$  of a line bundle L on X and an isomorphism  $\eta: \mathscr{O}_S \simeq e^*L$ . Let  $\operatorname{Pic}^e(X)$  denote the group of isomorphism classes of these rigidified line bundles on X, which is called the rigidified Picard group of X along e. If  $S = \mathbb{A}^n$ , then  $\operatorname{Pic}^e(X)$  is isomorphic to usual Picard group  $\operatorname{Pic}(X)$  of X, since any line bundle on  $\mathbb{A}^n$  is trivializable.

Let  $e : \bar{\mathbb{A}}^1 \to \operatorname{Gr}_{\mathcal{G}}$  the section of base points, and let  $\operatorname{Pic}^e(\operatorname{Gr}_{\mathcal{G}})$  be the rigidified Picard group of  $\operatorname{Gr}_{\mathcal{G}}$  along e. Note that  $\operatorname{Gr}_{\mathcal{G}}|_0 \simeq \operatorname{Fl}_Y$ . We have the following restriction map,

$$\mathfrak{R}: \operatorname{Pic}^{e}(\operatorname{Gr}_{\mathcal{G}}) \to \operatorname{Pic}(\operatorname{Fl}_{Y}).$$
$$[(\mathcal{L}, \eta)] \mapsto [\mathcal{L}_{0}]$$

This map is well-defined since  $(\mathcal{L}, \eta) \simeq (\mathcal{L}', \eta')$  implies  $\mathcal{L} \simeq \mathcal{L}'$ .

**Proposition 5.1.** The restriction map  $\Re$  is an isomorphism.

*Proof.* By Proposition 3.18, any line bundle  $\mathcal{L}(\Lambda)$  on  $\mathrm{Fl}_Y$  descends to a line bundle  $\mathcal{L}_{\mathrm{Bun}_{\mathcal{G}}}$  on  $\mathrm{Bun}_{\mathcal{G}}$ . Let  $\mathcal{L}(\Lambda)$  be the pullback of  $\mathcal{L}_{\mathrm{Bun}_{\mathcal{G}}}$  via morphism  $\mathrm{pr}:\mathrm{Gr}_{\mathcal{G}}\to\mathrm{Bun}_{\mathcal{G}}$ . Then,  $\mathcal{L}(\Lambda)$  is a line bundle on  $\mathrm{Gr}_{\mathcal{G}}$  and  $\mathcal{L}(\Lambda)_0\simeq\mathcal{L}(\Lambda)$ . Note that every line bundle on  $\mathrm{Gr}_{\mathcal{G}}$  is equipped with a rigidification automatically. It follows that the restriction map is surjective.

Let  $\mathcal{L}$  be a line bundle on  $Gr_{\mathcal{G}}$ , which is automatically equipped with a rigidification. Assume the restriction  $\mathcal{L}_0$  is trivial, we shall show  $\mathcal{L}$  is also trivial. Let  $\pi: \operatorname{Gr}_{\mathcal{G}} \to \bar{\mathbb{A}}^1$ denote the projection map. For any  $a \in \bar{\mathbb{A}}^1$ , by [Zh1, Proposition 4.1], the central charge of the restriction  $\mathcal{L}_a$  is equal to the central charge of  $\mathcal{L}_0$ , hence is equal to 0. For any  $a \neq 0$ , the central charge map  $Pic(Gr_{\mathcal{G},a}) \rightarrow \mathbb{Z}$  is an isomorphism, cf. [Fa, Corollary 12]. Therefore,  $\mathcal{L}_a$  is also trivial for any  $a \neq 0$ . Since  $Gr_{\mathcal{G}}$  is projective over  $\bar{\mathbb{A}}^1$ , we have dim  $(H^0(Gr_{\mathcal{G},a}, \mathcal{L}_a)) = 1$  for any  $a \in \bar{\mathbb{A}}^1$ . It follows that  $\pi_*(\mathcal{L})$  is a locally free sheaf of rank 1 on  $\mathbb{A}^1$ . Thus,  $\pi_*(\mathcal{L}) \simeq \mathscr{O}_{\bar{\mathbb{A}}^1}$ . By adjunction, we have a morphism  $\pi^*\pi_*(\mathcal{L}) \to \mathcal{L}$ . Set  $X := \operatorname{Gr}_{\mathcal{G}}$ . For any  $x \in X$ , the map  $(\pi^*\pi_*\mathcal{L})|_{x} = H^0(\operatorname{Gr}_{\mathcal{G},a}, \mathcal{L}_a) \to \mathcal{L}|_{x}$ is an isomorphism, since  $\mathcal{L}_a := \mathcal{L}|_{Gr_{\mathcal{G}_a}}$  is trivial on  $Gr_{\mathcal{G}_{a}}$  and  $Gr_{\mathcal{G}_{a}}$  is ind-projective, where  $a = \pi(x)$ . It follows that  $\mathcal{L} \simeq \pi^* \pi_*(\mathcal{L}) \simeq \pi^* \mathcal{O}_{\mathbb{A}^1}$  is trivial. Therefore, the restriciton map  $\Re$  is also injective.

For any affine weight  $\Lambda = \sum_{i \in \hat{I}_{\tau}} n_i \Lambda_i$ , by this proposition we can uniquely associate to a line bundle  $\mathcal{L}(\Lambda)$  on  $Gr_{\mathcal{G}}$  such that  $\mathcal{L}(\Lambda)_0 \simeq \mathcal{L}(\Lambda)$  on  $Fl_Y$ .

5.2. Canonical equivariance of line bundles on  $Gr_G$ . Assume G is simply connected. Let  $\vartheta$  be a rational interior point of the facet associated to Y. Let  $\sigma$  be the automorphism of G defined as in (5), which is of order m.

Set  $O_t := k[[t]]$  and  $O := k[[t^m]]$ . Let  $\sigma$  act on  $O_t$  by  $\sigma(t) := (\epsilon_m)^{-1}t$  and  $\sigma|_k = \mathrm{Id}_k$ . As in (6), we define the following group scheme over  $\bar{O}$ ,

(44) 
$$\mathscr{G}_{\sigma} := \operatorname{Res}_{\mathcal{O}_{\sigma}/\bar{\mathcal{O}}}(G_{\mathcal{O}_{\sigma}})^{\sigma}.$$

Set  $\mathcal{K}_t := k(t)$  and  $\mathcal{K} := k(t)$ , where  $t = t^{m/r}$ . Recall in Section 2.2, we can attach a parahoric subgroup  $\mathcal{P}_{\vartheta} \subseteq G(\mathcal{K})^{\tau}$  and a parahoric group scheme  $\mathscr{G}_{\vartheta}$  over  $\bar{O}$  associated to  $\vartheta$ , so that  $\mathscr{G}_{\vartheta}(\bar{O}) = \mathcal{P}_{\vartheta}$ . Furthermore, by Theorem 2.4 there is an isomorphism of group schemes  $\phi_0: \mathscr{G}_{\vartheta} \simeq \mathscr{G}_{\sigma}$ , such that over its  $\bar{O}$ -points,  $\mathscr{P}_{\vartheta} \simeq G(O_t)^{\sigma}$  is the restriction of the isomorphism  $\mathrm{Ad}_{\tau^{\theta}}: G(\mathcal{K})^{\tau} \simeq G(\mathcal{K}_{t})^{\sigma}$ . As in Section 3.3, we have an action of  $\sigma$  on  $\tilde{\mathbb{A}}^{1}$ , such that  $\bar{\mathbb{A}}^1 = \mathbb{A}^1/\langle \tau \rangle = \tilde{\mathbb{A}}^1/\langle \sigma \rangle$ . Similarly, we consider the following group scheme over  $\bar{\mathbb{A}}^1$ ,

$$\mathcal{G}_{\sigma} := \operatorname{Res}_{\tilde{\mathbb{A}}^1/\bar{\mathbb{A}}^1} (G \times \tilde{\mathbb{A}}^1)^{\sigma}.$$

Then, we have  $\mathcal{G}_{\sigma}|_{\bar{\mathbb{D}}_0} \simeq \mathcal{G}_{\sigma}$  and  $\mathcal{G}_{\sigma}|_{\bar{\mathbb{D}}_{\infty}} \simeq \mathcal{G}_{\sigma,\infty}$ , where  $\bar{\mathbb{D}}_0 = \operatorname{Spec}(\bar{O})$ . Let  $\mathcal{G}$  be the parahoric BT group scheme over  $\bar{\mathbb{A}}^1$  by glueing  $\mathcal{G}_Y$  and  $\operatorname{Res}_{\mathbb{A}^1 - 0/\bar{\mathbb{A}}^1 - 0}(G_{\mathbb{A}^1 - 0})^{\tau}$ . By (30), there is an isomorphism  $\phi: \mathcal{G} \simeq \mathcal{G}_{\sigma}$ . This induces an isomorphism of jet group schemes  $L^+\mathcal{G} \simeq L^+\mathcal{G}_{\sigma}$  over  $\mathbb{A}^1$ .

For any relative root  $\alpha \in \mathcal{R}$ , we can attach a root subgroup  $\mathscr{U}_{\alpha}^{\dagger}$  of  $\operatorname{Res}_{\mathcal{K}/\bar{\mathcal{K}}}(G_{\mathcal{K}})^{\sigma}$  and a natural valuation on  $\mathcal{U}_{\alpha}^{\dagger}$  as in Section 2.2. By [DH1, Theorem 4.1.2], we have

$$\mathrm{Ad}_{z^{\vartheta}}(\mathscr{U}_{\alpha,-\langle\vartheta,\alpha\rangle})=\mathscr{U}_{\alpha,0}^{\dagger}.$$

We now define global root subgroups of  $\mathcal{G}_{\sigma}$ . For each root  $\tilde{\alpha} \in R$  of G, let  $U_{\tilde{\alpha}}$  be the root subgroup of G and let  $U_{\tilde{\alpha},\tilde{\mathbb{A}}^1}$  denote the constant group scheme  $U_{\tilde{\alpha}}$  over  $\tilde{\mathbb{A}}^1$ . Set

$$\mathcal{U}_{\tilde{\alpha}}^{\dagger} := \begin{cases} \operatorname{Res}_{\tilde{\mathbb{A}}^{1}/\tilde{\mathbb{A}}^{1}}(U_{\tilde{\alpha},\tilde{\mathbb{A}}^{1}})^{\sigma} & \text{if } \tau(\tilde{\alpha}) = \tilde{\alpha} \\ \operatorname{Res}_{\tilde{\mathbb{A}}^{1}/\tilde{\mathbb{A}}^{1}}(\prod_{i=0}^{m-1} U_{\sigma^{i}(\tilde{\alpha}),\tilde{\mathbb{A}}^{1}})^{\sigma} & \text{if } \tau(\tilde{\alpha}) \text{ and } \tilde{\alpha} \text{ are not adjacent} \\ \operatorname{Res}_{\tilde{\mathbb{A}}^{1}/\tilde{\mathbb{A}}^{1}}(\prod_{i=0}^{\frac{m}{r}-1} U_{\sigma^{2i}(\tilde{\alpha}),\tilde{\mathbb{A}}^{1}}U_{\sigma^{2i}(\tilde{\alpha})+\sigma^{2i+1}(\tilde{\alpha}),\tilde{\mathbb{A}}^{1}}U_{\sigma^{2i+1}(\tilde{\alpha}),\tilde{\mathbb{A}}^{1}})^{\sigma} & \text{if } \tau(\tilde{\alpha}) \text{ and } \tilde{\alpha} \text{ are adjacent.} \end{cases}$$

Let  $L^+\mathcal{U}_{\bar{\alpha}}^{\dagger}$  be the jet group scheme of  $\mathcal{U}_{\bar{\alpha}}^{\dagger}$  as in (31). Let  $\alpha$  be the image of  $\bar{\alpha}$  under the natural map  $R \to \mathcal{R}$ . For any  $x \in \bar{\mathbb{A}}^1$ , we have

$$L^{+}\mathcal{U}_{\tilde{\alpha},x}^{\dagger} \simeq \begin{cases} \mathscr{U}_{\alpha,0}^{\dagger}, & \text{if } x = 0 \\ U_{\tilde{\alpha}}(O_{t}), & \text{if } x \neq 0 \end{cases}.$$

Via the isomorphism (30), the root subgroup  $\mathcal{U}_{\tilde{\alpha}}^{\dagger}$  of  $\mathcal{G}_{\sigma}$  corresponds to a root subgroup of  $\mathcal{G}$ , which will be denoted by  $\mathcal{U}_{\tilde{\alpha}}$ . For any  $x \in \mathbb{A}^1$ , we have

(45) 
$$L^{+}\mathcal{U}_{\tilde{\alpha},x} \simeq \begin{cases} \mathscr{U}_{\alpha,-\langle \vartheta,\alpha\rangle}, & \text{if } x = 0 \\ U_{\tilde{\alpha}}(O), & \text{if } x \neq 0 \end{cases}.$$

By the argument in Proposition 5.1, any line bundle  $\mathcal{L}$  on  $Gr_{\mathcal{G}}$  is isomorphic to  $\mathcal{L}(\Lambda)$  for a unique affine weight  $\Lambda$ . By Theorem 3.21,  $\mathcal{L}(\Lambda)$  admits a  $T^{\tau} \times \bar{\mathbb{A}}^1$ -equivariant structure. In the following theorem, we extend this  $T^{\tau} \times \bar{\mathbb{A}}^1$ -equivariance to a canonical equivariant structure of  $L^+\mathcal{G}$ .

**Theorem 5.2.** Assume that G is simiply-connected. Let  $\mathcal{L}$  be a line bundle on  $Gr_{\mathcal{G}}$ . There is a unique  $L^+\mathcal{G}$ -equivariant structure on  $\mathcal{L}$ .

We first need a lemma to reduce the proof of our theorem to global Schubert varieties. Recall the global affine Grassmannian  $\mathrm{Gr}_{\mathcal{G},\mathbb{A}^1}:=\mathrm{Gr}_{\mathcal{G}}\times_{\bar{\mathbb{A}}^1}\mathbb{A}^1$  over  $\mathbb{A}^1$ , considered in Section 3.4. There is a section  $s_\mu:\mathbb{A}^1\to\mathrm{Gr}_{\mathcal{G},\mathbb{A}^1}$  for any  $\mu\in X_*(T)^+$ . When  $\mu$  is  $\tau$ -invariant i.e.  $\tau(\mu)=\mu$ , the section  $s_\mu$  descends to a section  $\bar{s}_\mu:\bar{\mathbb{A}}^1\to\mathrm{Gr}_{\mathcal{G}}$ , see [Zh1, Remark 3.2]. Let  $(X_*(T)^+)^\tau$  denote the set of  $\tau$ -invariant elements of  $X_*(T)^+$ . For any  $\mu\in (X_*(T)^+)^\tau$ , we define the global Schubert variety  $\overline{\mathrm{Gr}}_{\mathcal{G}}^\mu$  to be the minimal  $L^+\mathcal{G}$ -stable irreducible closed subvariety of  $\mathrm{Gr}_{\mathcal{G}}$  containing  $\bar{s}_\mu$ .

**Lemma 5.3.** The global affine Grassmannian  $Gr_{\mathcal{G}}$  is a direct limit of global Schubert varieties with respect to the natural order  $\langle on(X_*(T)^+)^{\tau}, i.e. Gr_{\mathcal{G}} = \varinjlim_{\mu} \overline{Gr}_{\mathcal{G}}^{\mu}$ .

*Proof.* Consider the ind-closed embedding  $j: \varinjlim_{\mu} \overline{\operatorname{Gr}}_{\mathcal{G}}^{\mu} \to \operatorname{Gr}_{\mathcal{G}}$  over  $\bar{\mathbb{A}}^1$ , where  $\mu \in$ 

 $(X_*(T)^+)^{\tau}$ . Note that  $\operatorname{Gr}_{\mathcal{G},\mathbb{A}^1-0} \simeq \operatorname{Gr}_{G,\mathbb{A}^1-0}$  and it restricts to an isomorphism  $\overline{\operatorname{Gr}}_{\mathcal{G},\mathbb{A}^1-0}^{\mu} \simeq \overline{\operatorname{Gr}}_{G,\mathbb{A}^1-0}^{\mu}$  for any  $\mu \in (X_*(T)^+)^{\tau}$ . Since  $\varinjlim_{\mu} \overline{\operatorname{Gr}}_{G,\mathbb{A}^1-0}^{\mu} \simeq \operatorname{Gr}_{G,\mathbb{A}^1-0}$ , it induces an isomorphism

 $\overline{\operatorname{Gr}}_{\mathcal{G},\mathbb{A}^1-0}^{\mu} \simeq \operatorname{Gr}_{\mathcal{G},\mathbb{A}^1-0}$ . As  $\mathbb{A}^1-0 \to \overline{\mathbb{A}}^1-0$  is étale  $\mathbb{Z}/r\mathbb{Z}$ -cover, by descent theory this isomorphism implies  $j|_{\bar{\mathbb{A}}^1-0}: \varinjlim_{\mu} \overline{\operatorname{Gr}}_{\mathcal{G},\bar{\mathbb{A}}^1-0}^{\mu} \to \operatorname{Gr}_{\mathcal{G},\bar{\mathbb{A}}^1-0}$  is an isomorphism.

We now show that the restrictions  $j|_0$  is also an isomorphism. Recall that  $\operatorname{Gr}_{\mathcal{G},0} \simeq \operatorname{Fl}_Y$ 

We now show that the restrictions  $j|_0$  is also an isomorphism. Recall that  $\operatorname{Gr}_{\mathcal{G},0} \simeq \operatorname{Fl}_Y$  and  $\overline{\operatorname{Gr}}_{\mathcal{G},0}^{\mu} \simeq \mathcal{A}^Y(\bar{\mu})$  for any  $\mu \in (X_*(T)^+)^{\tau}$ , see (35). Thus, the restriction  $j|_0$  is the ind-closed embedding  $\varinjlim_{\mu} \mathcal{A}^Y(\bar{\mu}) \to \operatorname{Fl}_Y$ , where  $\mu \in (X_*(T)^+)^{\tau}$ . Given any  $w \in W_{\operatorname{aff}}$ , we shall show that  $w \leq_Y \varrho^{\bar{\mu}}$  for some  $\mu \in (X_*(T)^+)^{\tau}$ .

Let  $\mu \in (X_*(T)^+)^{\tau}$  be an element such that  $\bar{\mu} \neq 0$ . Since the element  $\varrho^{\bar{\mu}}$  has infinite order, any reduced word expression of  $\varrho^{\bar{\mu}}$  must contain all simple reflections  $s_i$  of  $W_{\text{aff}}$ .

Therefore,  $s_i < \varrho^{\bar{\mu}}$  for any  $i \in \hat{I}_{\tau}$ . Let  $w = s_{i_1} \cdots s_{i_k}$  be a reduced word in  $W_{\text{aff}}$ . Since  $\ell(\varrho^{k\bar{\mu}}) = k\ell(\varrho^{\bar{\mu}})$ , the element  $\varrho^{k\bar{\mu}}$  has a reduced word expression which contains  $s_{i_1} \cdots s_{i_k}$ as a subword. Thus,  $w < \varrho^{k\bar{\mu}}$ . By [Ku2, Lemma 1.3.18], we get  $w \leq_Y \varrho^{k\bar{\mu}}$ . Thus,  $\overline{\mathrm{Fl}}_{Y,w} \subseteq \overline{\mathrm{Fl}}_{Y,k\bar{\mu}} \subseteq \mathcal{A}^Y(k\bar{\mu})$ . Since  $\mathrm{Fl}_Y$  is the union of Schubert varieties  $\bigcup_{w \in W_{\mathrm{aff}}} \overline{\mathrm{Fl}}_{Y,w}$ , it follows that  $j|_0: \varinjlim \mathcal{A}^{\gamma}(\bar{\mu}) \simeq \operatorname{Fl}_{\gamma}$ . Finally, by a Theorem of W. Reeves (cf. [HY,

Theorem 4.23]), j is an isomorphism.

The following lemma will also be used in the proof of Theorem 5.2.

**Lemma 5.4.** Let H be a connected smooth affine group scheme over a smooth curve X over k, such that for any point  $x \in X$  the fiber  $H_x$  unipotent. Suppose we are given a central extension  $1 \to \mathbb{G}_{m,X} \to \hat{H} \xrightarrow{p} H \to 1$  over X, where  $\mathbb{G}_{m,X} := \mathbb{G}_m \times X$  is the constant group scheme over X. If the projection p splits over an open subset U of X, then p has a unique splitting over X.

*Proof.* The uniqueness is obvious, as every fiber  $H_x$  is unipotent. For the existence, we can assume  $X \setminus U$  only consists of a point  $x_0$ . Let  $\beta$  be a splitting of p over U. Let  $\overline{H}$  be the schematic closure of  $\text{Im}(\beta)$  (we remove the non-identity components of  $\bar{H}_{x_0}$  if they occur). By [Ha, III, Proposition 9.8],  $\bar{H}$  is a flat group scheme over X. By Cartier's Theorem (cf. [DG, II, $\S6,1.1$ ]) and [Ha, III, Theorem 10.2],  $\bar{H}$  is smooth over X. In particular, the fibers  $\bar{H}_{x_0}$  and  $H_{x_0}$  have the same dimension. By assumption, H is generically unipotent over X. Then, in view of [VD, Theorem 1.1],  $\bar{H}_{x_0}$  is unipotent. Thus,  $\bar{H}_{x_0}$  does not contain any non-trivial element in  $\mathbb{G}_{\mathrm{m}}$ . It follows that  $p: \mathrm{Lie}(\bar{H}_{x_0}) \to \mathrm{Lie}(H_{x_0})$  is an isomorphism. Applying [HY, Lemma 5.13],  $p: \bar{H}_{x_0} \to H_{x_0}$  is an isomorphism. By assumption,  $p: H \to H$  is an isomorphism over U. Resorting to [HY, Lemma 5.12],  $p: \bar{H} \to H$  is an isomorphism over X. This gives rise to a splitting of p over X.

**Proof of Theorem 5.2.** As in [HY, Section 5.2], we can define the smooth group scheme  $L_k^+\mathcal{G}$  over  $\bar{\mathbb{A}}^1$  such that  $L^+\mathcal{G} = \lim_{k \to \infty} L_k^+\mathcal{G}$  and  $L_k^+\mathcal{G}_a(k) \simeq \mathcal{G}_a(k[z_a]/(z_a^k))$  for any  $a \in \bar{\mathbb{A}}^1$ , where  $z_a$  is the local coordinate at a. For each root  $\tilde{\alpha}$  of G, recall that we have defined  $\mathcal{U}_{\tilde{\alpha}}$  and the jet group scheme  $L^+\mathcal{U}_{\tilde{\alpha}}$ . For each k, we can similarly define  $L^+_{k}\mathcal{U}_{\tilde{\alpha}}$ , which is a unipotent subgroup of  $L_k^+\mathcal{G}$ . Then,  $L^+\mathcal{U}_{\tilde{\alpha}} = \lim_{k \to \infty} L_k^+\mathcal{U}_{\tilde{\alpha}}$ .

To prove Theorem 5.2, it suffices, by Lemma 5.3, to show that for any  $\mu \in (X_*(T)^+)^{\tau}$ the line bundle  $\mathcal{L}|_{\overline{\mathrm{Gr}}_{\mathcal{G}}^{\mu}}$  on  $\overline{\mathrm{Gr}}_{\mathcal{G}}^{\mu}$  is  $L^{+}\mathcal{G}$ -equivariant. We fix an arbitrary  $\mu \in (X_{*}(T)^{+})^{\tau}$  and simply denote the restriction  $\mathcal{L}|_{\overrightarrow{Gr}_{G}^{\mu}}$  by  $\mathcal{L}$ . By the same argument as in [HY, Lemma 5.11], the action of  $L^+\mathcal{G}$  on  $\overline{\mathrm{Gr}}^\mu_\mathcal{G}$  factors through the action of  $L^+_k\mathcal{G}$  for large enough integer k. In the following, we will only work with large enough k.

As in [HY, Section 5.2], via the line bundle  $\mathcal{L}$ , we can define a central extension  $L_k^+\mathcal{G}$ of  $L_k^+\mathcal{G}$  by  $\mathbb{G}_{m,\bar{\mathbb{A}}^1}$ . Then, an  $L^+\mathcal{G}$ -equivariant structure on  $\mathcal{L}$  is equivalent to a splitting of the following extension,

$$(46) 1 \to \mathbb{G}_{m,\bar{\mathbb{A}}^1} \to \widehat{L_k^+ \mathcal{G}} \xrightarrow{p} L_k^+ \mathcal{G} \to 1.$$

Note that  $\overline{\mathrm{Gr}}_{\mathcal{G},\mathbb{A}^1-0}^{\mu} \simeq \overline{\mathrm{Gr}}_{G}^{\mu} \times (\mathbb{A}^1-0)$  and  $L_k^+\mathcal{G}_{\mathbb{A}^1-0} \simeq L_k^+G \times (\mathbb{A}^1-0)$ . By base change to  $\mathbb{A}^1-0$  via the morphism  $\mathbb{A}^1-0 \to \bar{\mathbb{A}}^1$ , it is clear that there is a unique splitting

 $\tilde{\beta}: L_k^+\mathcal{G}_{\mathbb{A}^1-0} \to \widehat{L_k^+\mathcal{G}}_{\mathbb{A}^1-0}$  of the extension (46) over  $\mathbb{A}^1 - 0$ . By uniqueness,  $\tilde{\beta}$  is  $\tau$ -equivariant. Thus, it descends to a splitting  $\beta: L_k^+\mathcal{G}_{\bar{\mathbb{A}}^1-0} \to \widehat{L_k^+\mathcal{G}}_{\bar{\mathbb{A}}^1-0}$  of (46) over  $\bar{\mathbb{A}}^1 - 0$ .

Let  $\overline{L_k^+\mathcal{G}}$  denote the closure of  $\operatorname{Im}(\beta)$  in  $\widehat{L_k^+\mathcal{G}}$  (and remove the non-identity components if they occur). By the same argument as in the proof of Lemma 5.4,  $\overline{L_k^+\mathcal{G}}$  is smooth group scheme over  $\overline{\mathbb{A}}^1$ . We claim that the morphism

$$(47) p: \overline{L_k^+ \mathcal{G}} \to L_k^+ \mathcal{G}.$$

is an isomorphism. Clearly, this is an isomorphism over  $\bar{\mathbb{A}}^1 - 0$ . By [HY, Lemma 5.12], it suffices to show that over  $0 \in \bar{\mathbb{A}}^1$  it is also an isomorphism,

$$(48) \overline{L_k^+ \mathcal{G}_0} \to L_k^+ \mathcal{G}_0 =: \mathcal{P}_{Y,k}.$$

By Proposition 3.12, the line bundle  $\mathcal{L}_0 = \mathcal{L}(\Lambda)$  on  $Gr_{\mathcal{G},0} = Fl_Y$  is  $\mathcal{P}_Y$ -equivariant, which induces a  $\mathcal{P}_{Y,k}$ -equivariant structure on  $\mathcal{L}|_{\overline{Gr}_{\mathcal{G},0}^{\mu}}$ . This further induces a splitting  $\beta_0 : \mathcal{P}_{Y,k} \to \widehat{L_k^+} \mathcal{G}_0$ . We shall show that  $Im(\beta_0) \subseteq \overline{L_k^+} \mathcal{G}_0$ .

By Lemma 5.4, we have a unique splitting  $\beta_{\tilde{\alpha}}: L_k^+\mathcal{U}_{\tilde{\alpha}} \to \widehat{L_k^+\mathcal{U}}_{\tilde{\alpha}} \subseteq \widehat{L_k^+\mathcal{G}}$ . By the uniqueness, we have  $\beta_{\tilde{\alpha}}(L_k^+\mathcal{U}_{\tilde{\alpha},\bar{\mathbb{A}}^1-0}) = \beta(L_k^+\mathcal{U}_{\tilde{\alpha},\bar{\mathbb{A}}^1-0})$ . From the proof of Lemma 5.4, we have  $\operatorname{Im}(\beta_{\tilde{\alpha}}) \subset \overline{L_k^+\mathcal{G}}$ . On the other hand,  $\beta_{\tilde{\alpha}}|_0$  is a splitting over  $L_k^+\mathcal{U}_{\tilde{\alpha},0}$ . By the uniqueness of splitting it must agree with  $\beta_0|_{L_k^+\mathcal{U}_{\tilde{\alpha},0}}$ . Therefore, we have

$$\beta_0(L_k^+\mathcal{U}_{\tilde{\alpha},0}) = \operatorname{Im}(\beta_{\tilde{\alpha}}|_0) = \operatorname{Im}(\beta_{\tilde{\alpha}})_0 \subseteq \overline{L_k^+\mathcal{G}}_0.$$

By Theorem 3.21, we have a splitting  $\zeta: T_{\bar{\mathbb{A}}^1}^{\tau} \to \widehat{L_k^+\mathcal{G}}$ . Moreover, the restriction  $\zeta|_0$  agrees with  $\beta_0|_{T^{\tau}}$  and the restriction  $\zeta|_{\bar{\mathbb{A}}^1-0}$  agrees with  $\beta$ . Then,

$$\beta_0(T^{\tau}) = \operatorname{Im}(\zeta|_0) \subseteq \overline{L_k^+ \mathcal{G}}_0.$$

By Theorem 2.10 and the isomorphism (45),  $L^+\mathcal{G}_0 = \mathcal{P}_Y$  is generated by  $T^\tau$  and  $\{L^+\mathcal{U}_{\tilde{\alpha},0}\}_{\tilde{\alpha}\in R}$ . Hence,  $\mathcal{P}_{Y,k} = L_k^+\mathcal{G}_0$  is generated by  $T^\tau$  and  $\{L_k^+\mathcal{U}_{\tilde{\alpha},0}\}_{\tilde{\alpha}\in R}$ . We conclude that  $\mathrm{Im}(\beta_0)\subseteq \overline{L_k^+\mathcal{G}_0}$ . In particular, it follows that  $p_0:\overline{L_k^+\mathcal{G}_0}\to L_k^+\mathcal{G}_0$  is surjective.

Note that  $L_k^+\mathcal{G}$  and  $\overline{L_k^+\mathcal{G}}$  are both smooth group schemes over  $\overline{\mathbb{A}}^1$  and isomorphic over  $\overline{\mathbb{A}}^1 - 0$ . Their fibers at 0 have the same dimension. Since  $\overline{L_k^+\mathcal{G}}_0$  is irreducible, it follows that  $\text{Im}(\beta_0) = \overline{L_k^+\mathcal{G}}_0$  and that (48) is an isomorphism. This proves our claim that (47) is an isomorphism. Then, the inverse map gives a splitting of the extension (46).

Finally, we show the uniqueness. Note that any splitting  $\beta: L_k^+\mathcal{G} \to \widehat{L_k^+\mathcal{G}}$  is uniquely determined by its restriction over  $\bar{\mathbb{A}}^1 - 0$ . It is clear that the splitting over  $\bar{\mathbb{A}}^1 - 0$  is unique, as there is no nontrivial character of  $L^+\mathcal{G}_a \simeq L^+G$  for any  $a \in \bar{\mathbb{A}}^1 - 0$ .

*Remark* 5.5. In our proof of Theorem 5.2, we crucially make use of Cartier theorem, which replies on the characteristic 0 condition. However, we expect that Theorem 5.2 also holds when  $p \nmid a_Y$ , where  $a_Y$  is defined in (10). Similarly, we expect Theorem 5.6 holds under a mild condition on the characteristic p.

5.3. **General case.** In this section, we drop the condition that G is simply-connected, and allow G to be a general simple algebraic group over k. As in Section 3.3, we fix a non-empty subset Y of  $\hat{I}_{\tau}$ , and let G be the parahoric Bruhat-Tits group scheme over  $\bar{\mathbb{A}}^1$  by gluing  $\mathcal{G}_Y$  and  $\operatorname{Res}_{\mathbb{A}^1-0/\bar{\mathbb{A}}^1-0}(G_{\mathbb{A}^1-0})^{\tau}$ . Let  $\operatorname{Gr}_{G}$  be the global affine Grassmannian of G over  $\bar{\mathbb{A}}^1$ . Set  $\operatorname{Gr}_{G,\mathbb{A}^1}:=\operatorname{Gr}_{G}\times_{\bar{\mathbb{A}}^1}\mathbb{A}^1$ . Let G' be the simply connected cover of G. We can similarly attach a parahoric Bruhat-Tits group scheme G' over  $\bar{\mathbb{A}}^1$ . Let  $L^+G'$  be the jet group scheme of G' over  $\bar{\mathbb{A}}^1$ . Set  $L^+G'_{\mathbb{A}^1}:=L^+G'\times_{\bar{\mathbb{A}}^1}\mathbb{A}^1$ .

As in Section 3.4, let M be the set of zero or miniscule coweights of G. For any  $\xi \in M$ , let  $Gr_{\mathcal{G},\mathbb{A}^1}[\xi]$  be the  $\xi$ -component of  $Gr_{\mathcal{G}}$ . Note that  $Gr_{\mathcal{G},\mathbb{A}^1}[\xi]|_0 \simeq Fl_Y^{\kappa}$ , where  $\kappa$  has the same image as  $\xi$  in  $X_*(T)_{\tau}/X_*(T')_{\tau}$ . Moreover, for any  $\mu \in X_*(T)_{\tau}$ , the global Schubert variety  $\overline{Gr}_{\mathcal{G},\mathbb{A}^1}^{\mu}$  is contained in  $Gr_{\mathcal{G},\mathbb{A}^1}[\xi]$  if and only if  $\mu + \xi \in X_*(T')$ , see Definition 3.23. Recall the section  $s_{-\xi}: \mathbb{A}^1 \to L\mathcal{G} \times_{\bar{\mathbb{A}}^1} \mathbb{A}^1$  which satisfies  $s_{-\xi}(0) = n^{-\xi} \in T(\mathcal{K})^{\tau}$ , where  $n^{-\xi}$  is defined in (19). The translation by  $s_{-\xi}$  gives rise to a morphism  $T_{\xi}: Gr_{\mathcal{G},\mathbb{A}^1}[o] \simeq Gr_{\mathcal{G},\mathbb{A}^1}[\xi]$ .

**Theorem 5.6.** Any line bundle  $\mathcal{L}$  on  $Gr_{\mathcal{G},\mathbb{A}^1}[\xi]$  is uniquely determined by its restriction  $\mathcal{L}_0$  on  $Fl_Y^{\kappa}$ . Moreover, there is a unique  $L^+\mathcal{G}'_{\mathbb{A}^1}$ -equivariant structure on  $\mathcal{L}$ .

*Proof.* The pull-back  $T_{\xi}^*\mathcal{L}$  is a line bundle on the neutral component  $\mathrm{Gr}_{\mathcal{G},\mathbb{A}^1}[o] \simeq \mathrm{Gr}_{\mathcal{G}',\mathbb{A}^1}$ . By Proposition 5.1,  $T_{\xi}^*\mathcal{L}$  is uniquely determined by its restriction on the neutral component  $\mathrm{Fl}_Y^\circ$  of  $\mathrm{Fl}_Y$ . This translates to the first part of our theorem, as  $(T_{\xi}^*\mathcal{L})|_{\mathrm{Fl}_Y^\circ} \simeq (n^{-\xi})^*(\mathcal{L}|_{\mathrm{Fl}_Y^\circ}) = (n^{-\xi})^*(\mathcal{L}_0)$ .

We consider the action of  $L^+\mathcal{G}'_{\mathbb{A}^1}$  on  $Gr_{\mathcal{G},\mathbb{A}^1}[\xi]$ . Similar to Lemma 5.3, we have

$$\mathrm{Gr}_{\mathcal{G},\mathbb{A}^1}[\xi] = \varinjlim_{\mu + \xi \in X_*(T')} \overline{\mathrm{Gr}}_{\mathcal{G},\mathbb{A}^1}^\mu.$$

By the same argument as in Section 5.2, there is a unique  $L^+\mathcal{G}'_{\mathbb{A}^1}$ -equivariant structure on  $\mathcal{L}|_{\overline{\mathrm{Gr}}^\mu_{\mathcal{G},\mathbb{A}^1}}$  for each  $\overline{\mathrm{Gr}}^\mu_{\mathcal{G},\mathbb{A}^1}$ , hence on  $\mathcal{L}$  over  $\mathrm{Gr}_{\mathcal{G},\mathbb{A}^1}[\xi]$ .

Remark 5.7. In Theorem 5.6, we need to work with  $Gr_{\mathcal{G},\mathbb{A}^1} = Gr_{\mathcal{G}} \times_{\bar{\mathbb{A}}^1} \mathbb{A}^1$ , as for general  $\xi \in M$  one can only construct a section  $s_{\xi} : \mathbb{A}^1 \to L\mathcal{G}_{\mathbb{A}^1}$  after the base change by  $\mathbb{A}^1 \to \bar{\mathbb{A}}^1$ .

## APPENDIX A. EXAMPLES OF THEOREM 4.5

**Example A.1.** Let  $g = sl_2$  and let  $\tau$  be trivial. Then,  $\hat{L}(g)$  is the untwisted affine algebra of type  $A_1^{(1)}$ . We label the vertex of the Dynkin diagram of g by 1. Then,  $\hat{I}_{\tau} = \{o, 1\}$ . Let  $\alpha := \alpha_1$  and  $\check{\alpha} := \check{\alpha}_1$  be the simple root and coroot of g. Let  $\omega := \omega_1$  and  $\check{\omega} := \check{\omega}_1$  denote the fundamental weight and coweight. There are two level one dominant integrable weights  $\Lambda_o$  and  $\Lambda_1 = \Lambda_o + \omega$  of  $\hat{L}(g)$ .

(1) We take  $\Lambda = \Lambda_1$  and  $\mu = \check{\alpha}$  in Theorem 4.5. In this case, c = 1 and  $\lambda = \omega$ . We have the following isomorphsim of  $\mathfrak{h}$ -modules,

$$D(1,\check{\alpha}) \otimes \mathbf{k}_{\omega} \simeq D(\Lambda_1, -\check{\alpha}) + D(\Lambda_1, \check{\alpha}).$$

Note that  $Y = \{1\}$  and  $-\check{\alpha}$  is  $\bar{Y}$ -dominant. By Corollary 4.17, the above isomorphism can be simplified to the isomorphism  $D(1, \check{\alpha}) \otimes k_{\omega} \simeq D(\Lambda_1, -\check{\alpha})$ . In fact,

by (43), we have

$$\rho^{\pm\check{\alpha}}(\Lambda_{\alpha}) = \Lambda_{\alpha} \mp \alpha - \delta; \quad \rho^{\check{\alpha}}(\Lambda_{1}) = \Lambda_{1} - \alpha, \quad \rho^{-\check{\alpha}}(\Lambda_{1}) = \Lambda_{1} + \alpha - 2\delta.$$

The modules  $D(1, \check{\alpha}) := D(\Lambda_o, \check{\alpha})$  and  $D(\Lambda_1, -\check{\alpha})$  are 4-dimensional  $\hat{\mathfrak{b}}$ -modules as shown in Figure 1a and 1b.




Figure 1

(2) We take  $\Lambda = 2\Lambda_o + \omega$  and  $\mu = \check{\omega}$  in Theorem 4.5. In this case, c = 2,  $\lambda = \omega$ . We get an isomorphism of  $\mathfrak{h}$ -modules

$$D(2,\check{\omega}) \otimes \mathbf{k}_{\omega} \simeq D(2\Lambda_{\alpha} + \omega, -\check{\omega}) + D(2\Lambda_{\alpha} + \omega, \check{\omega}).$$

In fact, by (43), we have

$$\varrho^{\check{\omega}}(\Lambda) = \Lambda - \alpha, \quad \varrho^{-\check{\omega}}(\Lambda) = \Lambda + \alpha - \delta.$$

The modules  $D(2\Lambda_o + \omega, -\check{\omega})$  and  $D(2\Lambda_o + \omega, \check{\omega})$  are 2-dimensional spaces as shown in Figure 2b labeled by red and blue colors respectively. They intersect at a 1-dimensional subspace. Hence, the sum of two spaces  $D(2\Lambda_o + \omega, -\check{\omega}) + D(2\Lambda_o + \omega, \check{\omega})$  is a 3-dimensional  $\hat{\mathfrak{b}}$ -module.

On the other hand, we have  $\Omega = \{id, \varsigma_{\check{\omega}}\}$ , where  $\varsigma_{\check{\omega}} = \varrho^{-\check{\omega}} s_{\alpha}$ . Note that  $\varsigma_{\check{\omega}}^{-1} = \varsigma_{\check{\omega}}$ . Then,  $\varrho^{\check{\omega}} \varsigma_{\check{\omega}}^{-1} = s_{\alpha} \in W_{\text{aff}}$  and we have

$$\varrho^{\check{\omega}}\varsigma_{\check{\omega}}^{-1}(2\Lambda_1) = s_{\alpha}(2\Lambda_1) = 2\Lambda_1 - 2\alpha.$$

Thus,  $D(2, \check{\omega})$  is a 3-dimensional  $\mathfrak{g}[t]$ -module shown in Figure 2a.




Figure 2

(3) We take  $\Lambda = 2\Lambda_o + \omega$  and  $\mu = \check{\alpha}$  in Theorem 4.5. Then,  $c = 2, \lambda = \omega$ . We get an isomorphism of  $\mathfrak{h}$ -modules

$$D(2, \check{\alpha}) \otimes k_{\omega} \simeq D(2\Lambda_o + \omega, -\check{\alpha}) + D(2\Lambda_o + \omega, \check{\alpha}).$$

In fact, by (43), we have

$$\varrho^{\check{\alpha}}(2\Lambda_{\varrho}) = 2\Lambda_{\varrho} - 2\alpha - 2\delta; \quad \varrho^{\check{\alpha}}(\Lambda) = \Lambda - 2\alpha - \delta, \quad \varrho^{-\check{\alpha}}(\Lambda) = \Lambda + 2\alpha - 3\delta.$$

The module  $D(2,\check{\alpha}) := D(2\Lambda_o,\check{\alpha})$  is a 9-dimensional g[t]-module as shown in Figure 3a. Moreover, the modules  $D(2\Lambda_o + \omega, -\check{\alpha})$  and  $D(2\Lambda_o + \omega, \check{\alpha})$  are 6dimensional spaces shown in Figure 3b labeled by red and blue colors respectively. They intersect at a 3-dimensional subspace. Hence, the sum of two spaces  $D(2\Lambda_o + \omega, -\check{\alpha}) + D(2\Lambda_o + \omega, \check{\alpha})$  is a 9-dimensional  $\hat{b}$ -module.




Figure 3

**Example A.2.** Let  $g = sl_3$ ,  $G = PGL_3$ , and let  $\hat{L}(g)$  be the untwisted affine algebra of type  $A_2^{(1)}$ . We label the vertices of the Dynkin diagram of g by 1, 2. Then,  $\hat{I} = \{0, 1, 2\}$ . Let  $\alpha_1, \alpha_2$  and  $\check{\alpha}_1, \check{\alpha}_2$  denote the simple roots and coroots of g respectively. Let  $\check{\omega}_1, \check{\omega}_2$ denote the fundamental coweights of g. Then,  $\Omega = \{id, \varsigma_{\check{\omega}_1}, \varsigma_{\check{\omega}_2}\}$ , where  $\varsigma_{\check{\omega}_1} = \varrho^{-\check{\omega}_1} s_{\alpha_1} s_{\alpha_2}$ and  $\varsigma_{\check{\omega}_2} = \varrho^{-\check{\omega}_2} s_{\alpha_2} s_{\alpha_1}$ . Consider the dominant weight  $2\Lambda_1$ , where  $\Lambda_1$  is the fundamental weight of  $\hat{L}(\mathfrak{g})$  associated to  $1 \in \hat{I}$ . Note that  $\varsigma_{\check{\omega}_1}^{-1} = \varsigma_{\check{\omega}_2}$ . We have  $\varrho^{\check{\omega}_2} \varsigma_{\check{\omega}_1}^{-1} = s_{\alpha_2} s_{\alpha_1} \in W_{\text{aff}}$ and

$$\varrho^{\check{\omega}_2}\varsigma_{\check{\omega}_1}^{-1}(2\mathbf{\Lambda}_1) = s_{\alpha_1}s_{\alpha_2}(2\mathbf{\Lambda}_1) = 2\mathbf{\Lambda}_1 - 2\alpha_1 - 2\alpha_2.$$

Thus,  $D(2, \check{\omega}_2) := D(2\Lambda_1, \check{\omega}_2)$  has only one layer at degree 0 as shown in Figure 4a.

Let  $\tau$  be the nontrivial diagram automorphism on g. Let  $\hat{L}(g,\tau)$  be the twisted affine algebra of type  $A_2^{(2)}$  and  $g^{\tau} \simeq sl_2$ . We label the vertex of the Dynkin diagram of  $g^{\tau}$  by 1. Then,  $\hat{I}_{\tau} = \{o, 1\}$ . Let  $\beta := \alpha_1|_{\mathfrak{h}^{\tau}}$  be the restriction of the simple root  $\alpha_1$  of g to  $\mathfrak{h}^{\tau}$ . Then,  $\beta$ is the simple root of  $g^{\tau}$ . There are two level two dominant weights  $\Lambda_o$  and  $2\Lambda_1 = \Lambda_o + \beta$ of  $\hat{L}(g,\tau)$ , where  $\Lambda_i$  is the fundamental dominant weight of  $\hat{L}(g,\tau)$  associated to  $i \in \hat{I}$ . Consider the fundamental coweight  $\check{\omega}_2 \in X_*(T)$ . Let  $\overline{\check{\omega}}_2$  denote its image in  $X_*(T)_{\tau}$ . Then,  $\iota(\overline{\check{\omega}}_2) = \beta$ , see Section 4.1. By (43), we have

$$\varrho^{\pm \overline{\omega}_2}(\Lambda_o) = \Lambda_o \mp 2\beta - \delta; \quad \varrho^{\overline{\omega}_2}(2\Lambda_1) = 2\Lambda_1 - 2\beta, \quad \varrho^{-\overline{\omega}_2}(2\Lambda_1) = 2\Lambda_1 + 2\beta - 2\delta.$$

The twisted Demazure modules  $D^{\tau}(\Lambda_0, \overline{\check{\omega}}_2)$  and  $D^{\tau}(2\Lambda_1, -\overline{\check{\omega}}_2)$  are 6-dimensional as shown in Figure 4.

Now, we have following examples of Theorem 4.5:

(1) We take  $\Lambda = \Lambda_o$  and  $\mu = \check{\omega}_2$ . Then,  $c = 2, \lambda = 0$ . We get the following isomorphism of  $g^{\tau}$ -modules,

$$D(2, \check{\omega}_2) \simeq D^{\tau}(\Lambda_o, \overline{\check{\omega}}_2) + D^{\tau}(\Lambda_o, -\overline{\check{\omega}}_2).$$

- Note that  $Y = \{o\}$  and  $\overline{\check{\omega}}_2$  is  $\bar{Y}$ -dominant. By Corollary 4.17, it can be simplified as  $D(2, \check{\omega}_2) \simeq D^{\tau}(\Lambda_o, \overline{\check{\omega}}_2)$ .
- (2) We take  $\Lambda = 2\Lambda_1$  and  $\mu = \check{\omega}_2$ . Then,  $c = 2, \lambda = \beta$ . We get the following isomorphism of  $\mathfrak{h}^{\tau}$ -modules,

$$D(2, \check{\omega}_2) \otimes \mathbf{k}_{\beta} \simeq D^{\tau}(2\Lambda_1, \overline{\check{\omega}}_2) + D^{\tau}(2\Lambda_1, -\overline{\check{\omega}}_2).$$

Note that  $Y = \{1\}$  and  $-\overline{\check{\omega}}_2$  is  $\overline{Y}$ -dominant. By Corollary 4.17, it can be simplified as  $D(2, \check{\omega}_2) \otimes k_{\beta} \simeq D^{\tau}(2\Lambda_1, -\overline{\check{\omega}}_2)$ .




FIGURE 4

#### REFERENCES

- [BH1] M. Besson and J. Hong, A combinatorial study of affine Schubert varieties in affine Grassmannian. Transformation Groups volume 27, pages 1189–1221 (2022).
- [BH2] M. Besson and J. Hong, Smooth locus of twisted affine Schubert varieties and twisted affine Demazure modules. arXiv:2010.11357.
- [Bor] A. Borel, Linear Algebraic Groups, 2nd. Enl. Ed., Springer-Verlag, New York, 1991.
- [Bou] N. Bourbaki *Lie Groups and Lie Algebras*. Chapters 4-6, Springer-Verlag Berlin Heidelberg 2002
- [BT] F. Bruhat and J. Tits, *Groupes réductifs sur un corps local. II. Schémas en groupes. Existence d'une donnée radicielle valuée.* Inst. Hautes Efudes Sci. Publ. Math., (60):197–376, 1984.
- [CT] J.-L. Colliot-Thélène, *Résolutions flasques des groupes linéaires connexes*. Journal für die reine und angewandte Mathematik, vol. 2008, no. 618, 2008, pp. 77-133.
- [DG] M. Demazure and P. Gabriel, *Groupes Algébriques I.* Masson/North Holland, 1970.
- [DH1] C. Damiolini and J. Hong, Local types of  $(\Gamma, G)$ -bundles and parahoric group scheme. Proceedings of the London Mathematical Society, Volume127, Issue2, August 2023, Pages 261-294.
- [DH2] C. Damiolini and J. Hong, *Line bundles on the moduli stack of parahoric bundles*. arXiv:2412.08826.
- [Fa] G. Faltings, *Algebraic loop groups and moduli spaces of bundles*. J. Eur. Math. Soc. (JEMS) 5 (2003), no. 1, 41-68.
- [Ha] R. Hartshorne, *Algebraic geometry*. Graduate Texts in Mathematics, No. 52. Springer-Verlag, New York-Heidelberg, 1977.
- [He] J. Heinloth, Uniformization of G-bundles. Math. Annalen, 347(3): 499-528, 2010.

- [HK1] J. Hong and S. Kumar, *Conformal blocks for Galois covers of algebraic curves*. Compos.Math.159,2191–2259 (2023).
- [HK2] J. Hong and S. Kumar, *Twisted conformal blocks and their dimension*. Math. Z. 306, 76 (2024).
- [HR] T. J. Haines and T. Richarz, The test function conjecture for parahoric local models. J. Amer. Math. Soc. 34 (2021), 135-218.
- [HS] J. Hong and L. Shen *Tensor invariants, saturation problems and Dynkin automorphisms*. Advances in Mathematics, Volume 285, Pages 629-657, 2015.
- [HY] J. Hong and H. Yu, Beilinson-Drinfeld Schubert varieties of parahoric group schemes and twisted global Demazure modules. arXiv:2209.07347. To appear in Selecta Mathematica.
- [Kac] V. Kac, *Infinite-dimensional Lie algebras*. *Third edition*. Cambridge University Press, Cambridge, 1990.
- [Kat] S. Kato, Frobenius Splitting of Thick Flag Manifolds of Kac–Moody Algebras, International Mathematics Research Notices, Volume 2020, Issue 17, September 2020, Pages 5401–5427.
- [Ku1] S. Kumar, *Demazure character formula in arbitrary Kac-Moody setting*. Invent. Math. 89 (1987), 395–423.
- [Ku2] S. Kumar, *Kac-Moody Groups, their Flag Varieties and Representation Theory*. Progress in Mathematics Vol. 214, Birkhauser (2002).
- [LLM] V. Lakshmibai, P. Littelmann, and P. Magyar, *Standard Monomial Theory and applications* In Representation theories and algebraic geometry, pp. 319-364. Dordrecht: Springer Netherlands, 1998.
- [Li] P. Littelmann, Contracting Modules and Standard Monomial Theory for Symmetrizable Kac-Moody Algebras J. Amer. Math. Soc. 11 (1998), 551-567.
- [Ma] O. Mathieu, Formules de caractères pour les algèbres de Kac-Moody générales Astérisque 159-160 (1988).
- [Po] V. L. Popov, *Picard group of connected affine algebraic group*. Russian Mathematical Surveys, 2023, Volume 78, Issue 4, Pages 794-796. arXiv:2302.13374.
- [PR] G. Pappas and M. Rapoport, *Twisted loop groups and their affine flag varieties*. Adv. Math. 219 (2008), no. 1, 118-198.
- [Sp] T.A. Springer, *Linear algebraic groups. Second edition*. Progress in Mathematics, 9. Birkhäuser Boston, Inc., Boston, MA, 1998.
- [SP] The Stacks project authors, *The Stacks project*. https://stacks.math.columbia.edu
- [St] R. Steinberg, *Endomorphisms of Linear Algebraic Groups*, Memoirs of the American Mathematical Society, vol. 80. American Mathematical Society, Providence (1968).
- [VD] B. Veisfeiler and I. Dolgacev, *Unipotent group schemes over integral rings*. Izv. Akad. Nauk SSSR Ser. Mat. Tom 38 (1974), No. 4.
- [Zh1] X. Zhu, On the coherence conjecture of Pappas and Rapoport. Ann. of Math. (2) 180 (2014), no. 1, 1-85.
- [Zh2] X. Zhu, *The geometric Satake correspondence for ramified groups*. Ann. Sci. Éc. Norm. Supér. (4) 48 (2015), no. 2, 409-451.

- [Zh3] X. Zhu, An introduction to affine Grassmannians and the geometric Satake equivalence. Geometry of moduli spaces and representation theory, 59-154, IAS/Park City Math. Ser., 24, Amer. Math. Soc., Providence, RI, 2017.
- $(J.\,Hong)\,\,Department\,\,of\,\,Mathematics,\,\,University\,\,of\,\,North\,\,Carolina\,\,at\,\,Chapel\,\,Hill,\,\,Chapel\,\,Hill,\,\,NC\,\,27599-3250,\,\,U.S.\,A.$

Email address: jiuzu@email.unc.edu

(H. Yu) Beijing International Center for Mathematical Research, Peking University, No. 5 Yiheyuan Road Haidian District, Beijing, P.R.China 100871

Email address: huanhuan.yu@outlook.com