

CORSIKA 8 with PYTHIA 8: Simulating Vertical Proton Showers

C. Gaudu¹, M. Reininghaus² and F. Riehn^{3,4*} for the CORSIKA 8 Collaboration[†]

¹*Bergische Universität Wuppertal,*

Gaußstraße 20, 42119 Wuppertal, Germany

²*Karlsruhe Institute of Technology, Institute of Experimental Particle Physics,
Hermann-von-Helmholtz-Platz 1, 76344 Eggenstein-Leopoldshafen, Germany*

³*Universidade de Santiago de Compostela,*

Instituto Galego de Física de Altas Enerxías,

Rúa de Xoaquín Díaz de Rábago, E-15782 Santiago de Compostela, Spain

⁴*Laboratório de Instrumentação e Física Experimental de Partículas,
Avenida Professor Gama Pinto 2, 1649-003 Lisboa, Portugal*

Abstract. The field of air shower physics, dedicated to understanding the development of cosmic-ray interactions with the Earth’s atmosphere, faces a significant challenge regarding the muon content of air showers observed by the Pierre Auger Observatory, and numerous other observatories. Thorough comparisons between extensive air shower (EAS) measurements and simulations are imperative for determining the primary energy and mass of ultra-high energy cosmic rays. Current simulations employing state-of-the-art hadronic interaction models reveal a muon deficit compared to experimental measurements, commonly known as the “Muon Puzzle”. The primary cause of this deficit lies in the uncertainties surrounding high-energy hadronic interactions.

In this contribution, we discuss the integration of a new hadronic interaction model, PYTHIA 8, into the effort to resolve the Muon Puzzle. While the PYTHIA 8 model is well-tailored in the context of Large Hadron Collider (LHC) experiments, its application in air shower studies remained limited until now. However, recent advancements, particularly in the ANGANTYR model of PYTHIA 8, offer promising enhancements in describing hadron-nucleus interactions, thereby motivating its potential application in air shower simulations.

We present results from EAS simulations conducted using CORSIKA 8, wherein PYTHIA 8 is employed to model hadronic interactions.

Key words: PYTHIA 8 – Muon Puzzle – hadronic interactions – CORSIKA 8 – air showers

*Now at

*Technische Universität Dortmund, August-Schmidt-Straße 4, 44221 Dortmund, Germany
Ruhr-Universität Bochum, Universitätsstraße 150, 44801 Bochum, Germany*

[†]FAL: gitlab.iap.kit.edu/AirShowerPhysics/corsika/-/wikis/Current-CORSIKA-8-author-list

Introduction

To gain insights into the “Muon puzzle” – a persistent muon deficit observed in air shower simulations compared to measurements, e.g. from the Pierre Auger Observatory – several studies took place: from ad hoc modification of cross-section, multiplicity, and elasticity of hadronic interactions models (Ulrich et al. (2011); Vicha (2023b,a); Ebr et al. (2023)), or by altering directly particle production, to performing a multi-parameter fit of model predictions against Auger data (Riehn et al. (2024)). The purpose of this proceeding is to introduce another hadronic interaction model, PYTHIA 8 (Bierlich et al. (2022)), into the landscape of air shower simulations. Unlike the models most commonly used in this kind of study, namely EPOS-LHC, SIBYLL 2.3d and QGSJET-II.04 (Pierog et al. (2015); Riehn et al. (2020); Ostapchenko (2011)), PYTHIA 8 was not built with the constraints of air showers physics in mind.

In this work, we present the first results from the *ongoing implementation* of PYTHIA 8 in CORSIKA 8 (Engel et al. (2019); Alameddine et al. (2023, 2025)) for vertical proton-induced showers. We discuss the longitudinal shower and energy loss profiles, as well as the lateral distributions and energy spectra of the particles at ground.

1. PYTHIA 8

PYTHIA was conceived as a general purpose particle interaction model, optimized for describing LHC measurements, i.e. e^+e^- , pp , $\bar{p}p$, pPb , $PbPb$ interactions. The nuclear interactions are handled by the ANGANTYR model (Bierlich et al. (2018), Bierlich et al. (2021)) in PYTHIA 8, bridging the gap between high energy and heavy ion phenomenology.

In Fig.1, the cross-sections for proton-proton and proton-air interactions, from ~ 100 GeV to 100 EeV are displayed¹. PYTHIA 8 is overall compatible with the commonly high energy models used in air shower physics: EPOS-LHC, SIBYLL 2.3d and QGSJET-II.04. In the proton-proton case, all hadronic interaction models yield an extremely similar description, except for the highest energies where PYTHIA diverges, while in proton-air interactions, this divergence starts at the TeV scale already, establishing PYTHIA 8 as the model with the highest cross-sections.

¹Further collisions systems, such as πp and π -air, can be found in Gaudu (2024).




Figure 1: Lines: Inelastic cross-sections as a function of the momentum in the laboratory frame of the projectile for proton-proton (left) and proton-air (right) collisions, assuming the composition of air as 80% ^{14}N to 20% ^{16}O mix. Markers: inelastic cross-section for proton-proton collisions from Abreu et al. (2012).

2. CORSIKA 8 + PYTHIA 8

The implementation of PYTHIA in CORSIKA 8 follows in the footsteps of previous studies by Reininghaus (2024) and Reininghaus et al. (2023), but now using the ANGANTYR model in PYTHIA 8.312 and its opportunities as discussed in Sjöstrand and Utheim (2022) and Gaudu (2024). A new feature allows for a single instance of PYTHIA to run, from which the beam system can be switched and the collision energy can vary on an event-by-event basis.

The plan is for all collisions that includes a nucleus to be treated by ANGANTYR², while any other collisions are handled by PYTHIA. A comprehensive set of total and partial cross-sections for atmospheric collisions was tabulated to simulate interactions using PYTHIA within CORSIKA 8. This includes pions, kaons, protons, neutrons, other long-lived mesons and baryons as projectiles, as well as protons, ^{12}C , ^{14}N , ^{16}O and ^{50}Ar as targets.

For this work, 300 vertical air showers that were induced by protons with a primary energy of 10^{17} eV were simulated using four different high-energy

²Using nuclear projectiles handled via ANGANTYR within C8 are a work-in-progress.

interaction models. The following energy thresholds for particle tracking were used: e^\pm/γ particles at 10 MeV, hadrons and muons at 300 MeV. Electromagnetic particles with an energy below 10^{-6} times the primary energy were combined statistically to reduce runtime; this thinning is detailed in Alameddine et al. (2023). For technical reasons, the energy threshold to transition from the lower energy model FLUKA (The FLUKA Collaboration et al. (2024)) to PYTHIA 8 is currently set at 4 TeV, while the CORSIKA 8 default value is ~ 80 GeV when using the other high energy models.

2.1. SHOWER DEVELOPMENT

The so-called longitudinal shower profile tracks the number of secondary particles as a function of atmospheric depth. It provides insights into the development of the shower as it propagates through the atmosphere.




Figure 2: Average longitudinal shower profiles for vertical proton-induced 10^{17} eV air showers for various particle types: photons (upper left), electrons/positrons (upper right), hadrons (lower left) and muons (lower right), using several high-energy hadronic interaction models and FLUKA as low-energy interaction model.

The longitudinal profile of PYTHIA 8 showers for several particle types (photons, electrons/positrons, hadrons and muons) are displayed in Fig. 2. These profiles are compatible to those from Alameddine et al. (2023) and (cite C8@ECRS2024) when previously comparing CORSIKA 8 to CORSIKA 7 outputs, with interaction models EPOS-LHC, SIBYLL 2.3d and QGSJET-II.04. All models display a *crossing* behaviour relative to EPOS-LHC, with higher muon content predicted in the early stages of shower development, and similar or lower content at later stages. PYTHIA 8 and QGSJET-II.04 predict up to 50% more muons at the early stages but end up with about 10% fewer muons by $X \sim 1000 \text{ g cm}^{-2}$ compared to EPOS-LHC.




Figure 3: Average total energy loss distributions for the same showers as in Fig. 2.

In Fig. 3 the total energy loss profile is shown, from which the depth of shower maximum, X_{max} , is inferred and displayed in Fig. 4. The PYTHIA 8 prediction for the average depth of shower maximum is found to be slightly below the values for SIBYLL 2.3d and EPOS-LHC, while being in good agreement with QGSJET-II.04.




Figure 4: Average depth of the shower maximum for the same showers as in Fig. 2.

It has been discussed in a recent study by Abdul Halim et al. (2024) that a deeper X_{\max} would lead to a larger number of muons at ground, making the description of PYTHIA 8 and QGSJET-II.04 the least attractive toward a resolution of the Muon Puzzle in this scenario.

2.2. PARTICLES AT GROUND

The lateral distribution function shows the particle number density as a function of distance from the shower impact point at a specific altitude. Unlike the longitudinal profile, it represents the shower at a particular stage of its development. As shown in Fig. 5, PYTHIA 8 predicts fewer electrons, positrons, and muons from near the shower core up to ~ 1 km, at sea level, in comparison to the other hadronic interaction models.




Figure 5: Average lateral distributions of electrons/positrons (left) and muons (right) at sea level for the same showers as displayed in Fig. 2.

Another observable related to the secondaries reaching the ground is the average energy spectrum of charged leptons depicted in Fig. 6. It seems that PYTHIA produces fewer electrons and positrons reaching the ground compared to the other interaction models. On the other hand, it offers a similar description for secondary muons at the ground as QGSJET-II.04, up to the TeV scale, with a slight offset towards higher energy muons. Above ~ 10 TeV, fluctuations from the limited size of the set of simulated showers are dominant for all models.




Figure 6: Average energy spectra of electrons/positrons (left) and muons (right) at sea level for the same showers as displayed in Fig. 2. High-energy interactions are modeled by PYTHIA 8 above 4 TeV and by other high-energy models above 80 GeV, with FLUKA handling interactions below these thresholds.

Conclusion

In summary, this work shows that full air shower simulations can now be performed using PYTHIA 8 within the CORSIKA 8 framework. Focusing on vertical proton-induced showers at 10^{17} eV, we found that PYTHIA 8 predicts shower observables that are in agreement, for the most part, with those from other hadronic interaction models. The longitudinal profiles are generally consistent across the models. While PYTHIA and EPOS-LHC exhibit some differences in the depth of shower maximum, PYTHIA remains compatible with both SIBYLL 2.3d and QGSJET-II.04. The lateral distributions and energy spectra at the ground exhibit distinct behaviours but remain broadly comparable. This highlights the potential of PYTHIA 8 as a viable tool for air shower simulations.

The strength of PYTHIA resides in its tuning opportunities, allowing in a friendly and well-documented manner its users to modify internal parameters of the model with a strong impact on air shower observables. Therefore, once the investigations of vertical showers with nuclear primaries and inclined air showers for both proton and nuclear primaries are finalized, a tuning study of PYTHIA will be the next logical step towards refining PYTHIA in relation to the Muon Puzzle. A forward physics tune of PYTHIA was previously presented by Fieg et al. (2024), while ongoing efforts for a tune using fixed-target measurement are discussed by Gaudu (2024).

Acknowledgements

C. Gaudu acknowledges funding from the German Research Foundation (DFG, Deutsche Forschungsgemeinschaft) via the Collaborative Research Center SFB1491: Cosmic Interacting Matters – from Source to Signal (F4) – project no. 445052434.

F. Riehn received funding from the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 101065027.

References

- Abdul Halim, A. et al. (2024). Testing hadronic-model predictions of depth of maximum of air-shower profiles and ground-particle signals using hybrid data of the Pierre Auger Observatory. *Phys. Rev. D*, 109(10):102001.
- Abreu, P. et al. (2012). Measurement of the proton-air cross-section at $\sqrt{s} = 57$ TeV with the Pierre Auger Observatory. *Phys. Rev. Lett.*, 109:062002.
- Alameddine, J., Albrecht, J., Ammerman-Yebra, J., Arrabito, L., Alves, A., Baack, D., Coleman, A., Dembinski, H., Elsässer, D., Engel, R., Faure, A., Ferrari, A., Gaudu, C., Glaser, C., Gottowik, M., Heck, D., Huege, T., Kampert, K., Karastathis, N., Nellen, L., Pierog, T., Prechelt, R., Reininghaus, M., Rhode, W., Riehn, F., Sackel, M., Sampathkumar, P., Sandrock, A., Soedingrekso, J., and Ulrich, R. (2025). Simulating radio emission from particle cascades with corsika 8. *Astroparticle Physics*, 166:103072.
- Alameddine, J.-M. et al. (2023). The particle-shower simulation code CORSIKA 8. *PoS, ICRC2023*:310.
- Bierlich, C. et al. (2022). A comprehensive guide to the physics and usage of PYTHIA 8.3. *SciPost Phys. Codeb.*, 2022:8.
- Bierlich, C., Gustafson, G., Lönnblad, L., and Shah, H. (2018). The Angantyr model for Heavy-Ion Collisions in PYTHIA8. *JHEP*, 10:134.
- Bierlich, C., Sjöstrand, T., and Utheim, M. (2021). Hadronic rescattering in pA and AA collisions. *Eur. Phys. J. A*, 57(7):227.
- Ebr, J., Blažek, J., Vícha, J., Pierog, T., Santos, E., Trávníček, P., and Denner, N. (2023). Impact of modified characteristics of hadronic interactions on cosmic-ray observables for proton and nuclear primaries. *PoS, ICRC2023*:245.
- Engel, R., Heck, D., Huege, T., Pierog, T., Reininghaus, M., Riehn, F., Ulrich, R., Unger, M., and Veberič, D. (2019). Towards a Next Generation of CORSIKA: A Framework for the Simulation of Particle Cascades in Astroparticle Physics. *Comput. Softw. Big Sci.*, 3(1):2.

- Fieg, M., Kling, F., Schulz, H., and Sjöstrand, T. (2024). Tuning pythia for forward physics experiments. *Phys. Rev. D*, 109(1):016010.
- Gaudu, C. (2024). Pythia 8 and Air Shower Simulations: A Tuning Perspective. In *22nd International Symposium on Very High Energy Cosmic Ray Interactions*.
- Ostapchenko, S. (2011). Monte Carlo treatment of hadronic interactions in enhanced Pomeron scheme: I. QGSJET-II model. *Phys. Rev. D*, 83:014018.
- Pierog, T., Karpenko, I., Katzy, J. M., Yatsenko, E., and Werner, K. (2015). EPOS LHC: Test of collective hadronization with data measured at the CERN Large Hadron Collider. *Phys. Rev. C*, 92(3):034906.
- Reininghaus, M. (2024). Air showers and hadronic interactions with CORSIKA 8. *SciPost Phys. Proc.*, 15:019.
- Reininghaus, M., Sjöstrand, T., and Utheim, M. (2023). Pythia 8 as hadronic interaction model in air shower simulations. *EPJ Web Conf.*, 283:05010.
- Riehn, F., Engel, R., Fedynitch, A., Gaisser, T. K., and Stanev, T. (2020). Hadronic interaction model Sibyll 2.3d and extensive air showers. *Phys. Rev. D*, 102(6):063002.
- Riehn, F., Fedynitch, A., and Engel, R. (2024). Sibyll*. *Astropart. Phys.*, 160:102964.
- Sjöstrand, T. and Utheim, M. (2022). Hadron interactions for arbitrary energies and species, with applications to cosmic rays. *Eur. Phys. J. C*, 82(1):21.
- The FLUKA Collaboration, Ballarini, Francesca, Batkov, Konstantin, Battistoni, Giuseppe, Bisogni, Maria Giuseppina, Böhlen, Till T., Campanella, Mauro, Carante, Mario P., Chen, Daiyuan, De Gregorio, Angelica, Degtiarenko, Pavel V., De la Torre Luque, Pedro, dos Santos Augusto, Ricardo, Engel, Ralph, Fassò, Alberto, Fedynitch, Anatoli, Ferrari, Alfredo, Ferrari, Anna, Franciosini, Gaia, Kraan, Aafke Christine, Lascaud, Julie, Li, Wenxin, Liu, Juntao, Liu, Zhiyi, Magro, Giuseppe, Mairani, Andrea, Mattei, Ilaria, Mazziotta, Mario N., Morone, Maria C., Müller, Stefan E., Muraro, Silvia, Ortega, Pablo G., Parodi, Katia, Patera, Vincenzo, Pinsky, Lawrence S., Ramos, Ricardo L., Ranft, Johannes, Rosso, Valeria, Sala, Paola R., Santana Leitner, Mario, Sportelli, Giancarlo, Tessonier, Thomas, Ytre-Hauge, Kristian S., and Zana, Lorenzo (2024). The fluka code: Overview and new developments. *EPJ Nuclear Sci. Technol.*, 10:16.
- Ulrich, R., Engel, R., and Unger, M. (2011). Hadronic Multiparticle Production at Ultra-High Energies and Extensive Air Showers. *Phys. Rev. D*, 83:054026.
- Vícha, J. (2023a). Probing hadronic interaction models with the hybrid data of the Pierre Auger Observatory. *SciPost Phys. Proc.*, 13:026.
- Vícha, J. (2023b). Testing Model Predictions of Depth of Air-Shower Maximum and Signals in Surface Detectors using Hybrid Data of the Pierre Auger Observatory. *EPJ Web Conf.*, 283:02012.