

Status of the CORSIKA 8 particle-shower simulation code

C. Gaudu¹ for the CORSIKA 8 Collaboration*

¹*Bergische Universität Wuppertal,
Gaußstraße 20, 42119 Wuppertal, Germany,*

Abstract. For over two decades, CORSIKA 7 and its previous versions have been the leading Monte Carlo code for simulating extensive air showers. However, its monolithic Fortran-based software design and hand-optimized code has created challenges for maintenance, adaptation to new computing paradigms, and extensions for more complex simulations. Addressing these limitations, the CORSIKA 8 project represents a comprehensive rewrite of CORSIKA 7, seeing its core functionality re-envisioned in a modern and modular C++ framework.

CORSIKA 8 has now reached a “physics-complete” state, offering a robust platform that encourages expert development for specialized applications. It supports high-energy hadronic interactions using models such as SIBYLL 2.3d, QGSJET-II.04, EPOS-LHC, and PYTHIA 8.3, alongside the treatment of electromagnetic cascades with PROPOSAL 7.6.2. Key highlights are the support for multiple interaction media, including cross-media particle showers, and an enhanced calculation of radio emissions from particle showers.

This contribution provides an overview of the current functionalities, showcases validation results of its simulations, and discusses future development plans.

Key words: CORSIKA 8 – status – Monte-Carlo – particle showers simulation code

1. Introduction

The astroparticle physics community significantly depends on Monte Carlo simulations to analyze particle showers occurring in air and various other media. For over two decades, the CORSIKA code (Heck et al. (1998)) has served as the standard for air shower simulations, initially developed for the KASCADE experiment. Currently maintained by KIT, CORSIKA 7.75 faces challenges due to its monolithic Fortran architecture and the retirement of key developers, complicating ongoing maintenance efforts. Additionally, forthcoming experiments necessitate a level of flexibility that the hand-optimized CORSIKA 7 (C7) code cannot accommodate in the long run.

*FAL: gitlab.iap.kit.edu/AirShowerPhysics/corsika/-/wikis/Current-CORSIKA-8-author-list

Consequently, in 2018, the development of CORSIKA 8 (C8) was initiated, which entails a complete rewrite of the core functionalities of C7 within a modular C++-based framework. The design philosophy behind C8 was previously discussed in Alameddine et al. (2023c), alongside with the general structure of the code. By the time of ICRC 2023, the CORSIKA 8 code was deemed “physics-complete” and since then, the development was focused primarily on enhancing the user experience.

This work reviews the most recent results in terms of electromagnetic and hadronic cascades, radio emission calculations and discuss the usage of CORSIKA 8 by the physics community.

2. Electromagnetic cascades

Electromagnetic cascades in CORSIKA 8 are treated using the PROPOSAL code (Alameddine et al. (2020)) at current version 7.6.2. In Sandrock et al. (2023), an overview of the simulation of electromagnetic cascades within CORSIKA 8 is presented, with an emphasis on electron-induced showers. It includes a comparative analysis of the implemented physics and simulation outcomes with those produced by EGS4, as is the case in CORSIKA 7¹.




Figure 1: Longitudinal profile of electromagnetic particles for 100 TeV photon-induced showers, observed at a height of 5.8 km above sea level, in CORSIKA 7 and CORSIKA 8.

¹It is important to note that although CORSIKA 7 serves as a benchmark for comparison, certain physical processes have been enhanced and differ in CORSIKA 8; e.g. C8 includes the process of triplet production, which is not accounted for by EGS4 in C7. A 1:1 agreement with C7 is neither expected nor intended.

In this section, we display results of the electromagnetic component of *photon*-induced air showers. The longitudinal profiles in Fig. 1 shows a consistency between CORSIKA 7 and CORSIKA 8 on the few-percent level, except for very low particle numbers at the earliest stage of the shower development.




Figure 2: Lateral profile (left) and energy distribution on the observation plane (right) of electromagnetic particles for 100 TeV photon-induced showers, observed at a height of 5.8 km above sea level, in CORSIKA 7 and CORSIKA 8.

The lateral distributions and energy spectra are shown in Fig. 2. The lateral distributions find themselves in agreement at the 10% level for e^\pm , while the codes agree below the 5% level for photons, except for very close distances to the shower core, at the cm scale, where both C8 distributions diverge from the C7 description. The energy spectra for C8 and C7 are in agreement typically on a 5% level, with some more pronounced differences at very low energies.

At extremely high energies and also at large densities, the Landau-Pomeranchuk-Migdal effect changes the behavior of electromagnetic showers, slowing down the development of the cascade and shifting the shower maximum to larger depth, as discussed in Sandrock et al. (2023).

3. Hadronic cascades

CORSIKA 8 employs a modular framework that integrates state-of-the-art physics models to simulate hadronic processes with precision and adaptability. It supports a versatile suite of high-energy interaction models, includ-

ing EPOS-LHC (Pierog et al. (2015)), SIBYLL 2.3d (Riehn et al. (2020)), QGSJET-II.04 (Ostapchenko (2011)), and an experimental integration of PYTHIA 8.3 (Bierlich et al. (2022)), catering to diverse physics scenarios. Most recent efforts on the integration of the PYTHIA 8.3 hadronic interaction model are discussed in Gaudu (2024) and (cite C8+P8@ECRS2024), while in parallel, ongoing work to interface the recently updated models, EPOS LHC-R (Pierog and Werner (2023)) and QGSJET-III (Ostapchenko (2024)) is taking place.

The model used in CORSIKA 8 for low-energy interactions is FLUKA (The FLUKA Collaboration et al. (2024)). The transition threshold between the high- and low-energy models is fixed at ~ 80 GeV as default, but can be modified. Decay processes are supported through SIBYLL 2.3d and PYTHIA 8.




Figure 3: Average lateral distributions of electrons/positrons (left) and muons (right) at ground for 300 vertical proton-induced 10^{17} eV air showers, using several high-energy hadronic interaction models and FLUKA as low-energy interaction model, for CORSIKA 8 and CORSIKA 7. From Alameddine et al. (2023c).

In Fig. 3, comparisons of the lateral distributions of electrons/positrons and muons at the ground are displayed for proton-induced showers with the interaction models EPOS-LHC, SIBYLL 2.3d, QGSJET-II.04. Reaching agreement at this level with completely independent codes is no small accomplishment, though further investigation is needed to understand the differences in the lateral distribution of electrons and positrons in hadron-induced air showers between C7 and C8. The muon content at ground agrees at the 5–10% level near the shower axis, while discrepancies at larger distances remain to be explored.

4. Radio-emission calculations

The radio-emission calculations have significantly influenced the development of CORSIKA 8. This is especially true given the radio-detection community’s requirement for a flexible framework that can accommodate complex simulation scenarios. For instance, in-ice radio detection experiments necessitate consideration of the radio emission from air showers as they propagate from air into ice, a background that is presently simulated by piecing together several simulation codes (De Kockere et al. (2024)). The implementation of the radio process in CORSIKA 8 is discussed in Alameddine et al. (2023a) and most recently by Alameddine et al. (2025).

The best method to examine the agreement for all observer positions between CORSIKA 7 (C7), CORSIKA 8 (C8) and ZHAIRES (Zas et al. (1992)) is with the energy fluence, which is the energy deposited by radio waves per unit area in a given frequency band, predicted at the ground. Displayed in Fig. 4 is the 2D fluence map for different signal polarizations of the electric field in the 30 MHz to 80 MHz, and 50 MHz to 350 MHz frequency bands in the shower plane, for small step sizes (MaxRad \sim 0.001 rad), for C8 with “CoREAS” formalism, C8 with “ZHS” formalism, and C7 with CoREAS extension.

The emission footprints from CORSIKA 8 with both formalism, and CORSIKA 7, demonstrate an agreement in terms of shape and symmetry of the energy fluence for all of the electric field components. In Alameddine et al. (2025), it is shown that as the particle track length in the simulation is decreased, the “CoREAS” and “ZHS” formalisms agree to within 2% in the 30–80 MHz band and to within 1% in the 50–350 MHz band. For small particle tracks, the agreement in terms of radiation energy, which is the area integral over the energy fluence, is on the level of better than 10% for the 30-80 MHz band.

An interesting feature of the vertical polarization is the presence of a *blip* near the shower axis predicted by the “ZHS” formalism in both C8 and ZHAIRES², a feature that is not observed in the calculations employing the “CoREAS” formalism. From a practical point of view, however, the polarization along the shower axis holds little relevance, as the fluence in that region is negligible.

²This *blip* can be spotted for C8(ZHS) in the bottom row of Fig. 4, and in Alameddine et al. (2025) for C8(ZHS) and ZHAIRES.




Figure 4: Fluence maps for 30-80 MHz (left) and 50-350 MHz (right) radio emission from a vertical 10^{17} eV iron-induced air shower simulated with both the endpoint (CoREAS) and ZHS formalisms in CORSIKA 8 compared with CoREAS from CORSIKA 7, for smaller step size (MaxRad ~ 0.001 rad). From top to bottom, the rows show: total energy fluence, fluence in the $\vec{v} \times (\vec{v} \times \vec{B})$ (north-south), $\vec{v} \times \vec{B}$ (east-west) and \vec{v} (vertical) polarizations. The x/y -axes show core distances $\vec{v} \times \vec{B}$ and $\vec{v} \times (\vec{v} \times \vec{B})$ directions. The color scales are identical within a given row.

5. New functionalities

Certain functionalities introduced in CORSIKA 8 cannot be independently achieved using its predecessor, CORSIKA 7. It is now feasible to trace the ancestry of particles beyond the immediate mother and grandmother generations, enabled by the newly introduced air shower genealogy feature, as outlined in Reininghaus et al. (2021). Furthermore, the thinning algorithm for electromagnetic cascades has been enhanced by incorporating elements of the Hillas algorithm, previously utilized in CORSIKA 7, alongside a statistical thinning approach. More details about this updated thinning algorithm are discussed in Alameddine et al. (2023c).




Figure 5: Longitudinal profile of a vertical 100 PeV proton *cross-media* shower, starting at the top of the atmosphere, and intersecting an ice layer located at an altitude of 2.4 km above sea level. Data from Alameddine et al. (2023b).

A most-awaited implementation for the in-ice radio-detection community is the ability to simulate showers propagating through multiple media *within a single framework*, now made possible by the cross-media shower feature in CORSIKA 8. Combining this novel feature with radio emission calculations will allow the study of complex geometries relevant for radio neutrino detectors. Recent advancements in simulating in-ice radio emissions from air showers using CORSIKA 8 are presented in Coleman et al. (2024). This study shows, within the single framework of C8, that air-shower cores generate coherent in-ice radio emissions similar to those generated by neutrino-induced in-ice showers of equivalent energy. This represents an important step forward for the in-ice radio community, expanding the range of available tools.

Conclusion

There were many improvements over the last two years, and we now stand at a stage where the CORSIKA 8 code is considered “physics-complete”. This means the implementation of the physical processes taking place during the development of particle showers is done.

Recent enhancements include the implementation of the low-energy interaction model FLUKA, ongoing efforts to integrate the high-energy interaction model PYTHIA 8.3, consideration of photo-hadronic interactions and the Landau-Pomeranchuk-Migdal (LPM) effect, refinement of the electromagnetic cascade thinning algorithm, a fully integrated Cherenkov light calculation using GPU parallelization (Albrecht et al. (2023)), and a realistic simulation of radio emission in air accompanied by a proof-of-concept CPU parallelization (Alves et al. (2023)). Extensive validation was done against CORSIKA 7 and other codes; what remains is understanding differences at the 5-10% level, polishing user-level and technical aspects, such as refining simulation steering, enhancing performance, and finalizing the documentation. An expert release of CORSIKA 8 is expected by the end of this year.

A future implementation of interaction models QGSJET-III and EPOS LHC-R to CORSIKA 8 is shortly expected. Moreover, additional features are anticipated as user requests are communicated to the development team. Users interested in testing CORSIKA 8 for specific use cases are encouraged to contact the development team through the channels provided on the Corsika 8 GitLab page³.

Acknowledgements

C. Gaudu acknowledges funding from the German Research Foundation (DFG, Deutsche Forschungsgemeinschaft) via the Collaborative Research Center SFB1491: Cosmic Interacting Matters – from Source to Signal (F4) – project no. 445052434.

³gitlab.iap.kit.edu/AirShowerPhysics/corsika

References

- Alameddine, J., Albrecht, J., Ammerman-Yebra, J., Arrabito, L., Alves, A., Baack, D., Coleman, A., Dembinski, H., Elsässer, D., Engel, R., Faure, A., Ferrari, A., Gaudu, C., Glaser, C., Gottowik, M., Heck, D., Huege, T., Kampert, K., Karastathis, N., Nellen, L., Pierog, T., Prechelt, R., Reininghaus, M., Rhode, W., Riehn, F., Sackel, M., Sampathkumar, P., Sandrock, A., Soedingrekso, J., and Ulrich, R. (2025). Simulating radio emission from particle cascades with corsika 8. *Astroparticle Physics*, 166:103072.
- Alameddine, J.-M. et al. (2023a). Simulating radio emission from air showers with CORSIKA 8. *PoS, ICRC2023*:425.
- Alameddine, J.-M. et al. (2023b). Simulations of cross media showers with CORSIKA 8. *PoS, ICRC2023*:442.
- Alameddine, J.-M. et al. (2023c). The particle-shower simulation code CORSIKA 8. *PoS, ICRC2023*:310.
- Alameddine, J.-M., Soedingrekso, J., Sandrock, A., Sackel, M., and Rhode, W. (2020). Proposal: A library to propagate leptons and high energy photons. *Journal of Physics: Conference Series*, 1690(1):012021.
- Albrecht, J. et al. (2023). Comparison and efficiency of GPU accelerated optical light propagation in CORSIKA 8. *PoS, ICRC2023*:417.
- Alves, A. A. et al. (2023). Parallel processing of radio signals and detector arrays in CORSIKA 8. *PoS, ICRC2023*:469.
- Bierlich, C. et al. (2022). A comprehensive guide to the physics and usage of PYTHIA 8.3. *SciPost Phys. Codeb.*, 2022:8.
- Coleman, A., Glaser, C., Rice-Smith, R., Barwick, S., and Besson, D. (2024). In-ice Askaryan Emission from Air Showers: Implications for Radio Neutrino Detectors.
- De Kockere, S., Van den Broeck, D., Latif, U. A., de Vries, K. D., van Eijndhoven, N., Huege, T., and Buitink, S. (2024). Simulation of radio signals from cosmic-ray cascades in air and ice as observed by in-ice Askaryan radio detectors. *Phys. Rev. D*, 110(2):023010.
- Gaudu, C. (2024). Pythia 8 and Air Shower Simulations: A Tuning Perspective. In *22nd International Symposium on Very High Energy Cosmic Ray Interactions*.
- Heck, D., Knapp, J., Capdevielle, J. N., Schatz, G., and Thouw, T. (1998). CORSIKA: A Monte Carlo code to simulate extensive air showers.
- Ostapchenko, S. (2011). Monte Carlo treatment of hadronic interactions in enhanced Pomeron scheme: I. QGSJET-II model. *Phys. Rev. D*, 83:014018.
- Ostapchenko, S. (2024). QGSJET-III: predictions for extensive air shower characteristics and the corresponding uncertainties.

- Pierog, T., Karpenko, I., Katzy, J. M., Yatsenko, E., and Werner, K. (2015). EPOS LHC: Test of collective hadronization with data measured at the CERN Large Hadron Collider. *Phys. Rev. C*, 92(3):034906.
- Pierog, T. and Werner, K. (2023). EPOS LHC-R : up-to-date hadronic model for EAS simulations. *PoS, ICRC2023*:230.
- Reininghaus, M., Ulrich, R., and Pierog, T. (2021). Air shower genealogy for muon production. *PoS, ICRC2021*:463.
- Riehn, F., Engel, R., Fedynitch, A., Gaisser, T. K., and Stanev, T. (2020). Hadronic interaction model Sibyll 2.3d and extensive air showers. *Phys. Rev. D*, 102(6):063002.
- Sandrock, A., Alameddine, J.-M., and Riehn, F. (2023). Validation of Electromagnetic Showers in CORSIKA 8. *PoS, ICRC2023*:393.
- The FLUKA Collaboration, Ballarini, Francesca, Batkov, Konstantin, Battistoni, Giuseppe, Bisogni, Maria Giuseppina, Böhlen, Till T., Campanella, Mauro, Carante, Mario P., Chen, Daiyuan, De Gregorio, Angelica, Degtiarenko, Pavel V., De la Torre Luque, Pedro, dos Santos Augusto, Ricardo, Engel, Ralph, Fassò, Alberto, Fedynitch, Anatoli, Ferrari, Alfredo, Ferrari, Anna, Franciosini, Gaia, Kraan, Aafke Christine, Lascaud, Julie, Li, Wenxin, Liu, Juntao, Liu, Zhiyi, Magro, Giuseppe, Mairani, Andrea, Mattei, Ilaria, Mazziotta, Mario N., Morone, Maria C., Müller, Stefan E., Muraro, Silvia, Ortega, Pablo G., Parodi, Katia, Patera, Vincenzo, Pinsky, Lawrence S., Ramos, Ricardo L., Ranft, Johannes, Rosso, Valeria, Sala, Paola R., Santana Leitner, Mario, Sportelli, Giancarlo, Tessonier, Thomas, Ytre-Hauge, Kristian S., and Zana, Lorenzo (2024). The fluka code: Overview and new developments. *EPJ Nuclear Sci. Technol.*, 10:16.
- Zas, E., Halzen, F., and Stanev, T. (1992). Electromagnetic pulses from high-energy showers: Implications for neutrino detection. *Phys. Rev. D*, 45:362–376.