LIMIT POINTS OF UNIFORM ARITHMETIC BASS NOTES

WILL HIDE AND BRAM PETRI

ABSTRACT. We prove that the set of limit points of the set of all spectral gaps of closed arithmetic hyperbolic surfaces equals $[0, \frac{1}{4}]$.



FIGURE 1. A tree cover of a surface of genus two.

1. INTRODUCTION

The spectral gap $\lambda_1(X)$ of a closed orientable hyperbolic surface X is the minimal non-zero eigenvalue of its Laplacian. A natural question, that fits into a larger context of the study of the bass note spectra of locally uniform geometries [Sar23], is which numbers can appear as spectral gaps of hyperbolic surfaces. It can be derived in various ways from various combinations of [Che75, Gag80, Jen81, Bav96, SU13, Bon22, KMP24, FBP23] that the **bass note spectrum** of the set of closed orientable hyperbolic surfaces satisfies:

 $Bass(hyp_2^c) := \{ \lambda_1(X) : X \text{ a closed orientable hyperbolic surface} \} = (0, \Lambda_2],$

where $\Lambda_2 = \sup \{ \lambda_1(X) : X \text{ a closed hyperbolic surface of genus 2} \}$. The latter conjecturally equals the spectral gap of the Bolza surface (see Section 2.1 for a description).

If we restrict to arithmetic surfaces, the set of spectral gaps we obtain becomes countable. Recently, Magee [Mag24a] proved that the spectral gaps – defined as the infimum of the non-zero spectrum in this setting – of non-compact arithmetic surfaces form a dense set in $[0, \frac{1}{4}]$. He also asked whether the same is true for closed arithmetic surfaces. The goal of this article is to answer this question in the affirmative:

Theorem 1. The set $\{\lambda_1(X) : X \text{ a closed arithmetic surface}\} \cap [0, \frac{1}{4}]$ is dense in $[0, \frac{1}{4}]$.

Date: December 20, 2024.

In fact, we show that, like in the non-compact case, it suffices to consider only finite index torsion free subgroups of a single uniform arithmetic group. In our case this is the (2,3,8)-triangle group.

Similarly, the same method can be used to prove that if $X = \Gamma \backslash \mathbb{H}^2$ is a closed orientable surface then

$$\left\{\lambda_1(\Gamma' \setminus \mathbb{H}^2): \ \Gamma' < \Gamma, \ [\Gamma:\Gamma'] < \infty\right\} \bigcap [0,\Lambda]$$

is dense in $[0, \Lambda]$, where $\Lambda = \min \{ \lambda_1(\Gamma' \setminus \mathbb{H}^2) : \Gamma' < \Gamma, [\Gamma : \Gamma'] = 2 \}.$

Finally, we also obtain a different proof of [LM23, Theorem 1.8], the fact that there exists a sequence of finite sheeted covers $Y_n \to X$ of a given closed orientable arithmetic hyperbolic surface X, whose genera tend to infinity and whose spectral gap tends to $\frac{1}{4}$.

1.1. Strategy of proof. Like Magee's proof in the non-uniform case, we build our arithmetic surfaces as covers of degree two of a certain type of random covers of a fixed base surface. In our case, this base surface is the Bolza surface X_B . As such, the overarching strategy of our proof is similar to that of Magee. However, several of the key ingredients of his proof are not present in our setting, so in what follows we will describe the similarities and the differences.

The first thing that makes our case different, is that the fundamental group $\Gamma_B = \pi_1(X_B)$ of the X_B is not free, which makes random finite degree covers harder to control (see [MP23, MNP22] for the geometry of uniform random covers of a closed hyperbolic surface). We remedy this by considering random covers $Y^{(n)} \to X_B$ of degree *n* whose mondromy factors through an epimorphism from our surface group $\Gamma_B \to F_2$ to a free group F_2 on two generators. This allows us to use the results of Bordenave–Collins [BC19] on strong convergence of permutation representations combined with results by Hide–Magee [LM23, Appendix] on the convergence of the spectral gaps of the associated covers. In [LM23], Louder–Magee also did this. However, we need more control over the geometry of the associated cover, so we use a different argument.

We need to control the following things:

- (1) In order to obtain near optimal spectral gaps, we need to understand the spectral gap of the cover of the Bolza surface associated to the map $\Gamma_B \to F_2$ (see Figure 1). We tackle this using Cheeger's inequality. Using Mirzakhani's result on random pants decompositions of hyperbolic surfaces [Mir16] combined with some further geometric arguments, we show that by varying the map to the free group (see Section 4 and Proposition 6.1) we can obtain a spectral gap as close to $\frac{1}{4}$ as we like.
- (2) To reach intermediate spectral gaps, our aim is to use the covers of degree two of our random cover $Y^{(n)} \to X_B$. To make this work, we need there to be at least one cover among these degree two covers that has a large spectral gap. We obtain such a cover by restricting a non-constant homomorphism $\Gamma_B \to F_2 \to \mathbb{Z}/2\mathbb{Z}$ to the fundamental group of $Y^{(n)}$, seen as a subgroup of Γ_B . The fact that this indeed yields two sheeted covers of $Y^{(n)}$ with large spectral gaps relies on a strong convergence argument combined with the fact that all connected two sheeted covers

 $\widehat{Y} \to X_B$ satisfy $\lambda_1(\widehat{Y}) > \frac{1}{4}$. The latter fact we prove (in Section 3) using the linear programming methods developed in [FBP23, FBGMPP23].

(3) Like Magee, we then need to show that, when n is large enough, "small" modifications of the degree two covers of $Y^{(n)}$ do not influence the spectral gap too much. These modifications consist of changing the cover along the lifts of a simple closed curve in $Y^{(n)}$ and will be called switches (see Section 6.5). In Magee's paper, logarithmic tangle freeness is used to prove that similar moves don't influence the spectral gap too much. Unfortunately, our covers are, by construction, much less tangle free: the infinite cover associated to the map $\Gamma_B \to F_2$ contains many short curves. As a result, the injectivity radius of $Y^{(n)}$ is uniformly bounded.

We will however show, again relying on Mirzakhani's results on random pants decompositions and also using Nica's work on cycles of random permutations coming from word maps [Nic94], that we can find maps $\Gamma_B \to F_2$ such that, with positive probability, the curves in $Y^{(n)}$ along which we switch, admit uniformly wide collars (see Section 6.6). This suffices to make the argument, based on the L^{∞} -bounds on eigenfunctions by Gilmore–Le Masson–Sahlsten–Thomas [GLMST21], work. This implies the density result because any two covers of degree two of $Y^{(n)}$ can be connected by a finite sequence of switches.

1.2. Further notes and references. The study of spectral gaps of hyperbolic surfaces has a long history, which we will not attempt to summarize here. We refer to [Bus10, Ber16] for introductory texts. The shape of the bass note spectrum has been studied in the case of compact hyperbolic 2-orbifolds [Bon22, KMP24], non-compact arithmetic surfaces [Mag24a] and regular graphs [AW23, DM24]. In the case of orbifolds, Kravchuk–Mazac–Pal give a complete description of the set

 $\{\lambda_1(O) : O \text{ a compact orientable hyperbolic 2-orbifold}\}.$

This description is conjectural, but a large part of it is proven (see [KMP24, Conjecture 4.2]).

The case of arithmetic surfaces is very different. There are only finitely many arithmetic hyperbolic surfaces of any given genus (up to isometry). As such, the theorem above can be stated as saying that the set of limit points of the set of spectral gaps of closed hyperbolic surfaces equals $[0, \frac{1}{4}]$. There are however isolated points outside of this interval, for instance corresponding to the Bolza surface and the Klein quartic. Sarnak [Sar23] conjectured that there should be infinitely many such isolated points.

Acknowledgements. We are very grateful to Maxime Fortier Bourque for his part in developing the code that we use in Section 3. We also thank the SageMath [The21], Arb [Joh17] and GAP [GAP22] developers for making the calculations using that same code possible.

We thank the CIRM for hosting the workshop Random hyperbolic surfaces and random graphs, part of the Jean-Morlet Chair semester Random Matrices, Representation Theory and Quantum Information, during which this project started.

WILL HIDE AND BRAM PETRI

Finally, we thank Michael Magee and Frédéric Naud for useful conversations surrounding this project.

Funding. BP was partially supported by the grant ANR-23-CE40-0020-02 "MOST".

2. The model

In this section we formally describe our model of random surfaces, which is a variation of the models appearing in [BCP21, LM23, Mat24].

2.1. The Bolza surface. The model we will describe samples a certain type of covers of a closed hyperbolic surface. The model makes sense for any closed base surface, but because of the application we have in mind, we will assume the base surface is the Bolza surface. We will use this subsection to describe some of its properties.

The quickest way to define the Bolza surface – that we shall denote X_B – is to say that it's a Riemann surface of genus 2 with 48 automorphisms. Indeed, it turns out that this is the maximal possible number of automorphisms of a Riemann surface of genus 2 and there is a unique such surface. More concretely, X_B can be obtained by taking a regular octagon O with interior angles $\pi/4$ in the hyperbolic plane \mathbb{H}^2 and identifying opposite sides. We will also fix a Fuchsian group Γ_B such that $X_B = \Gamma_B \setminus \mathbb{H}^2$ and we will moreover fix an identification of Γ_B with $\pi_1(X_B)$.

The surface X_B is also a (2,3,8)-triangle surface. That is, its fundamental group is a normal subgroup (of index 48) in the orientable triangle group

$$\Theta(2,3,8) = \langle x, y, z | x^2, y^3, z^8, xyz \rangle$$

whose generators act on \mathbb{H}^2 by the order 2, 3 and 8 rotations in the vertices of a hyperbolic triangle whose interior angles are $\pi/2$, $\pi/3$ an $\pi/8$ respectively. The quotient group is described explicitly as item T2.1 in [Con15]. Since the (2,3,8)-triangle group is arithmetic [Tak77] (see also [MR03, Section 13.3]), X_B is arithmetic.

Very good numerical estimates of the spectral gap of X_B have been obtained in [SU13]. That is,

$$\lambda_1(X_B) = 3.838887 \pm 10^{-6}.$$

The best known upper bound on the maximal spectral gap of a hyperbolic surface of genus 2 equals 3.8388976481 [Bon22, KMP24].

2.2. Handlebody attachments, maps to free groups and pants decompositions. A handlebody H is a three-manifold that can be obtained by attaching 1-handles to a closed 3-ball. Alternatively it can be seen as the closed regular neighborhood of a finite graph in \mathbb{R}^3 . In particular, its fundamental group is a free group and its boundary is a closed orientable surface. The genus of this surface will be called the **genus** of H. This genus coincides with the rank of the fundamental group of H. The inclusion map $\iota : \partial H \hookrightarrow H$ induces an epimorphism

$$\iota_*: \pi_1(\partial H) \longrightarrow F_g,$$

where g is the genus of H. In fact, we obtain a large family of such maps, because we can precompose the inclusion by a homeomorphism $\phi : \partial H \to \partial H$.

Any such homeomorphism yields an F_q -cover

$$T_{\phi} \to \partial H$$

corresponding to the subgroup ker($(\iota \circ \phi)_*$). The surface T_{ϕ} is what is sometimes called a Cantor tree: a surface homemorphic to $\mathbb{S}^2 - C$, where C is a Cantor set. As such, we will call the F_q -covers constructed by the procedure above **tree covers**.

Finally, we observe that if \mathcal{P} is a pants decomposition of ∂H consisting entirely of **meridians** – simple closed curves that bound disks in H – then $\phi^{-1}(\mathcal{P})$ lifts to a pants decomposition of T_{ϕ} . For a picture of this set-up, see Figure 1.

2.3. Random covers. We will now once and for all fix a handlebody attachment ι : $X_B \hookrightarrow H$ of the Bolza surface X_B and a pants decomposition $\mathcal{P}_0 = (\alpha_1, \alpha_2, \alpha_3)$ of the X_B that consists entirely of meridians.

Given a homeomorphism $\phi : X_B \to X_B$, we obtain, as in the previous section, a homomorphism $(\iota \circ \phi)_* : \Gamma_B \to F_2$ and a tree cover $T_\phi \to X_B$. The pants decomposition of T_ϕ that we obtain by lifting $\phi^{-1}(\mathcal{P}_0)$ will be called the **canonical pants decomposition** of T_ϕ in what follows.

Now, given a homomorphism $\rho \in \text{Hom}(F_2, \mathfrak{S}_n)$, where \mathfrak{S}_n denotes the symmetric group on *n* letters, we obtain a subgroup

$$H^{(n)} = \operatorname{Stab}_{\rho}(\{1\}) \quad < \quad F_2,$$

but also a homomorphism

$$\rho \circ (\iota \circ \phi)_* : \Gamma_B \longrightarrow \mathfrak{S}_n$$

and hence a subgroup

$$\Lambda_{\phi,\rho} = \operatorname{Stab}_{\rho \circ (\iota \circ \phi)_*}(\{1\}) \quad < \quad \Gamma_B$$

of index at most n and thus a cover

$$Y_{\phi,\rho} \longrightarrow X_B$$

of degree at most n.

Our random surface $Y_{\phi}^{(n)}$ will be obtained by taking the permutation representation $\rho \in \text{Hom}(F_2, \mathfrak{S}_n)$ uniformly at random. It follows from results by Dixon [Dix69] that, as $n \to \infty$, the probability that the cover $Y_{\phi}^{(n)}$ is of degree exactly *n* tends to 1.

3. The bass notes of degree two covers of the Bolza surface

In this section we will explain how we prove, with the help of a computer, that all the degree two covers $Y \to X_B$ of the Bolza surface satisfy $\lambda_1(Y) > \frac{1}{4}$. To do so, we will employ the linear programming methods from [FBP23, FBGMPP23].

In fact, just like in [Mag24a], this can be proved by verifying the spectral gap of a single surface Y_{17} obtained as the $H_1(X_B, \mathbb{Z}/2\mathbb{Z})$ -cover of X_B . Indeed, any homomorphism $\Gamma_B \to \mathbb{Z}/2\mathbb{Z}$ factors through $H_1(X_B, \mathbb{Z}/2\mathbb{Z})$ and as such any degree two cover of X_B is intermediate to the cover $Y_{17} \to X_B$.

3.1. Identifying the cover. First of all, we need a description of Y_{17} . Since

$$|\mathrm{H}_1(X_B, \mathbb{Z}/2\mathbb{Z})| = 16,$$

the genus of Y_{17} equals 17 (whence the notation). Moreover, by construction, the fundamental group of Y_{17} is a characteristic¹ subgroup of Γ_B . This implies that it's a normal subgroup of the (2,3,8)-triangle group $\Theta(2,3,8)$. In other words, Y_{17} is a (2,3,8)-triangle surface of genus 17. It turns out there is only one such surface, which appears as item T17.2 in Conder's list [Con15]. In particular, Y_{17} corresponds to the quotient

$$G = \langle x, y, z | x^2, y^3, z^8, xyz, z^2yxz^2yxzy^{-1}z^{-1}xzy^{-1}z^{-1}x \rangle$$

of $\Theta(2, 3, 8)$.

3.2. Excluding small eigenvalues. In order to prove that Y_{17} has no small eigenvalues, we use the method from [FBGMPP23] that we will very briefly describe here. For more details, we refer to loc. cit. and [CP23].

First of all, the Deck group G of the orbifold cover $Y_{17} \rightarrow S_{2,3,8}$, where $S_{2,3,8} = \Theta(2,3,8) \setminus \mathbb{H}^2$ denotes the orientable (2,3,8)-triangle orbifold, acts on functions on Y_{17} . This action commutes with the Laplacian, so we can split the eigenspaces of the Laplacian into G-representations.

Our goal is now to exclude small eigenvalues from these representations. To do so, we use a criterion based on the twisted Selberg trace formula corresponding to every such representation.

We start with some set up. We will write $\operatorname{spec}(Y_{17})$ for the spectrum of the Laplacian on Y_{17} , as a multi-set. Moreover, $\operatorname{Irr}(G)$ will denote the finite set of real *G*-representations $\rho: G \to \operatorname{GL}(V^{\rho})$. The splitting of the eigenspaces into *G*-representations described above yields a splitting of the spectrum

$$\operatorname{spec}(Y_{17}) = \bigcup_{\rho \in \operatorname{Irr}(G)} \dim(\rho) \cdot \operatorname{spec}(Y_{17}, \rho),$$

where multiplication of a multi-set by a positive integer means multiplying the multiplicities by that integer and spec (Y_{17}, ρ) is the spectrum of the Laplacian acting on functions $\mathbb{H}^2 \to V^{\rho}$ that are equivariant with respect to the action of $\Theta(2,3,8)$ on the left and $\rho(G)$ on the right. This spectrum can also be interpreted in terms of the associated bundle.

For d > 0, we define $f_d : \mathbb{R} \to \mathbb{R}$ by

$$f_d(x) = \left(\frac{1}{2d}\chi_{[-d,d]}\right)^{*4}(x) = \begin{cases} \frac{1}{12d} \left(4 - \frac{3}{2d^2}x^2 + \frac{3}{8d^3} |x|^3\right) & \text{if } 0 \le |x| \le 2d, \\ \frac{1}{12d} \left(2 - \frac{|x|}{2d}\right)^3 & \text{if } 2d \le |x| \le 4d \text{ and} \\ 0 & \text{otherwise.} \end{cases}$$

Here $\chi_{[-d,d]}$ denotes the characteristic function of the interval [-d,d] and "*" denotes the convolution product.

¹Recall that a characteristic subgroup of Γ_B is a subgroup that is preserved by all elements of Aut(Γ_B). This is a stronger assumption than being a normal subgroup.

The Fourier transform $\widehat{f}_d : \mathbb{C} \to \mathbb{C}$ of f is given by

$$\widehat{f}_d(y) = \int_{-\infty}^{\infty} f_d(x) e^{-iy \cdot x} dx = \frac{\sin(d \cdot y)^4}{(d \cdot y)^4}, \quad y \in \mathbb{C}.$$

Given $\rho \in \operatorname{Irr}(G)$, we define

(3.1)
$$\mathcal{G}_{d}(\rho) = -\frac{\dim(V^{\rho})}{96} \int_{-\infty}^{\infty} \frac{f'_{d}(x)}{\sinh(x/2)} dx + \sum_{[\gamma]\in\mathcal{E}(\Gamma)} \frac{\operatorname{tr}\left(\widetilde{\rho}(\gamma)\right)}{m(\gamma)} \int_{0}^{\infty} \frac{\cosh(x/2)}{\cosh(x) - 1 + 2\sin(\theta(\gamma))^{2}} f_{d}(x) dx + \sum_{[\gamma]\in\mathcal{P}(\Gamma)} \ell(\gamma) \sum_{n\geq 1} \frac{\operatorname{tr}\left(\widetilde{\rho}(\gamma^{n})\right)}{2\sinh(n\ell(\gamma)/2)} f_{d}(n\ell(\gamma)),$$

where

- $\tilde{\rho}$ is the composition of $\rho: G \to \operatorname{GL}(V^{\rho})$ with the quotient map $\Theta(2,3,8) \to G$,
- $\mathcal{E}(\Gamma)$ denotes the set of conjugacy classes of elliptic elements in $\Theta(2,3,8)$,
- for an elliptic element $\gamma \in \Theta(2,3,8)$, $m(\gamma)$ denotes its order and $\theta(\gamma)$ denotes half the angle of rotation, i.e., is such that γ is conjugate to

$$\begin{bmatrix} \cos(\theta(\gamma)) & \sin(\theta(\gamma)) \\ -\sin(\theta(\gamma)) & \cos(\theta(\gamma)) \end{bmatrix} \in \mathrm{PSL}(2,\mathbb{R}),$$

- $\mathcal{P}(\Gamma)$ denotes the set of conjugacy classes of primitive hyperbolic elements in $\Theta(2,3,8)$,
- for a hyperbolic element $\gamma \in \Theta(2,3,8)$, $\ell(\gamma)$ denotes its translation length on \mathbb{H}^2 .

We observe that, because f_d has finite support, the sum and the integrals appearing on the right hand side of (3.1) are all finite.

Given these definitions, the criterion we use is the following:

Proposition 3.1 ([FBGMPP23, Proposition 3.1]). Let $\rho \in Irr(G)$ and let $\lambda > 0$. Suppose that

$$\widehat{f}_d\left(\sqrt{\lambda-\frac{1}{4}}\right) > \begin{cases} \mathcal{G}_d(\rho) & \text{if } \rho \text{ is non-trivial} \\ \mathcal{G}_d(\rho) - \widehat{f}(i/2) & \text{if } \rho \text{ is trivial,} \end{cases}$$

for some d > 0. Then $\lambda \notin \operatorname{spec}(Y_{17}, \rho)$.

We prove the following lower bound on the spectral gap of Y_{17} :

Proposition 3.2. The criterion from Proposition 3.1 hold for the parameter $d = \frac{3}{4}$ for all $\rho \in Irr(G)$ and for all $\lambda \leq 0.2501$. In particular,

$$\lambda_1(Y_{17}) \ge 0.2501.$$

Proof. The final proof of this proposition is performed using SageMath and its interface to GAP. Here we will discuss the main ingredients.

WILL HIDE AND BRAM PETRI

- (1) We compute the character table of G, using GAP.
- (2) Now we observe that when $d = \frac{3}{4}$, the support $\operatorname{supp}(f_d) = [-3, 3]$. As such, we need to know all the hyperbolic conjugacy classes of translation length at most 3 and all the elliptic conjugacy classes of $\Theta(2, 3, 8)$ in order to compute $\mathcal{G}_d(\rho)$. The latter are the just the conjugacy classes of the generators x, y and z and their powers. The hyperbolic conjugacy classes we need were determined in [FBGMPP23].
- (3) The integrals appearing in $\mathcal{G}_d(\rho)$ are treated using interval arithmetic as implemented in the Arb package.

The claim in our proposition is verified in the Jupyter notebook computation_spectral_gap.ipynb that is attached to the arXiv version of this article as an ancillary file.

Remark 3.3. Because 0.2501 suffices for what follows, we have made no effort to optimize the bound in the proposition above.

4. Bass notes of planar surfaces of infinite area

Given a hyperbolic surface X, we define its **Cheeger constant** by

$$h(X) = \inf \left\{ \begin{array}{l} \frac{\ell(\partial A)}{\operatorname{area}(A)} : \begin{array}{l} A \subset X \text{ compact} \\ \partial A \text{ rectifiable} \end{array} \right\}.$$

The **systole** – the length of the shortest closed geodesic on X – of a hyperbolic surface X will be denoted sys(X). We call a surface **planar** if it has genus 0. Note that such a surface either has cusps or infinite area.

We claim:

Lemma 4.1. Let X be a planar orientable hyperbolic surface without cusps. Then

$$h(X) \ge 1 - \frac{2\pi}{\operatorname{sys}(X) + 2\pi}.$$

Proof. The case in which the fundamental group of X is elementary – which in our setting implies that X is either the hyperbolic plane or an annulus $\langle g \rangle \backslash \mathbb{H}^2$ with $g \in PSL(2, \mathbb{R})$ hyperbolic – is covered by the results of Adams–Morgan [AM99, Theorem 2.2, Lemma 2.3].

As such, we may assume the fundamental group of X is non-elementary. We define

$$H(X) = \inf \left\{ \begin{array}{l} \ell(\partial A) \\ \overline{\operatorname{area}(A)} \end{array} : \begin{array}{l} A \subset X \text{ compact} \\ \partial A \text{ consists of simple closed geodesics} \end{array} \right\}$$

Clearly $h(X) \leq H(X)$. It turns out that

$$h(X) \ge \frac{H(X)}{H(X) + 1}$$

see $[Mat05, Theorem 7]^2$ and [Mir13, Proposition 4.7].

²Note that Matsuzaki uses the inverse convention for the Cheeger constant.

Because X is planar, any candidate subsurface A for H(X) is necessarily planar as well. Suppose A is an *n*-holed sphere. This means that

$$\operatorname{area}(A) = 2\pi \cdot (n-2) \quad \text{and} \quad \ell(\partial A) \ge n \cdot \operatorname{sys}(X).$$

So, $H(X) \ge \frac{n}{n-2} \frac{\operatorname{sys}(X)}{2\pi} \ge \frac{\operatorname{sys}(X)}{2\pi}$ and hence

$$h(X) \ge 1 - \frac{2\pi}{\operatorname{sys}(X) + 2\pi}$$

Using Cheeger's inequality [Che70] (see also [Cha84, Section IV.3]), we obtain:

Proposition 4.2. Let X be a planar hyperbolic surface without cusps. Then

$$\lambda_1(X) \ge \frac{1}{4} \cdot \left(1 - \frac{2\pi}{\operatorname{sys}(X) + 2\pi}\right)^2.$$

5. Eigenfunctions near pants curves

In this section, we describe the effect of flattening an eigenfunction near a pants curve on the Rayleigh quotient of that function.

5.1. **Delocalization.** One of the inputs we will need, is a delocalization result by Gilmore–Le Masson–Sahlsten–Thomas [GLMST21, Theorem 1.5]:

Theorem 5.1 (Gilmore–Le Masson–Sahlsten–Thomas). Suppose that $X = \Gamma \setminus \mathbb{H}^2$ is a closed hyperbolic surface of genus $g \geq 2$. Then there exists a universal constant A > 0 such that

$$|f(z)|^2 \leq A \cdot e^{-t \cdot \sqrt{\frac{1}{4} - \lambda}} \cdot |\{\gamma \in \Gamma : d_{\mathbb{H}^2}(z, \gamma \cdot z) \leq t\}|$$

for all t > 0, $\lambda \in (0, \frac{1}{4})$ and all Laplacian eigenfunctions f of eigenvalue λ with $||f||_{L^2} = 1$.

5.2. Flattening an eigenfunction near a pants curve. Given the result above, we need a bound on the Rayleigh quotient of a flattened eigenfunction in terms of the L^{∞} -norm of the eigenfunction. Our main technical input is the Ismagilov–Morgan–Simo–Sigal localization formula [CFKS87, Theorem 3.2]:

Theorem 5.2 (Ismagilov–Morgan–Simo–Sigal). Let M be a Riemannian manifold and suppose $\{J_i\}_{i \in \mathcal{I}}$ is a family of smooth functions $J_i : M \to [0, 1]$ such that

- (1) $\sum_{i \in \mathcal{I}} J_i^2 \equiv 1$,
- (2) on any compact subset $K \subset M$, only finitely many J_i are non-zero, and

(3)
$$\sup_{x \in M} \sum_{i \in \mathcal{I}} |\nabla J_i(x)|^2 < \infty.$$

Then

$$\Delta = \sum_{i \in \mathcal{I}} J_i \Delta J_i - \sum_{i \in \mathcal{I}} |\nabla J_i|^2.$$

Given a simple closed geodesic γ in a hyperbolic surface X and w > 0, we will write

$$C_w(\gamma) = \{ x \in X : d(x, \gamma) \le w \}$$

for the **collar** of width w around γ . If $C_w(\gamma)$ is isometric to a cylinder $[-w, w] \times \mathbb{S}^1$ with the Riemannian metric

$$ds^2 = d\rho^2 + \ell^2 \cosh^2(\rho) dt^2,$$

where ℓ is the length of γ , we will call the this collar **standard**. The collar lemma (see [Kee74] and [Bus10, Chaper 4]) states that the collar of width $w(\ell) = \operatorname{arcsinh}\left(\frac{1}{\sinh(\ell/2)}\right)$ is always standard. Note that $w(\ell) \sim e^{-\ell/2}$ as $\ell \to \infty$. For our arguments, we will need to work with (long) geodesics that admit standard collars of uniform width.

We will use the following bound, which is the analogue of [Mag24a, Proposition 8.1.] in our setting:

Proposition 5.3. For all w > 0 there exists a constant $\varepsilon > 0$ such that the following statement holds. Let X be a closed hyperbolic surface and let $\gamma_1, \ldots, \gamma_k \subset X$ be closed geodesics of length ℓ such that the collars of width w > 0 around $\gamma_1, \ldots, \gamma_k$ are standard and disjoint from each other. Moreover, let $f : X \to \mathbb{R}$ be an L^2 -normalized Laplacian eigenfunction of eigenvalue $\lambda > 0$. Suppose furthermore that³

$$||f||_{\infty}^2 \cdot \ell < \varepsilon.$$

Then there exists a smooth function $f': X \to \mathbb{R}$ such that

$$f'|_{C_{w/2}(\gamma)} \equiv 0, \quad f'|_{X-C_w(\gamma)} = f|_{X-C_w(\gamma)} \quad and \quad \frac{\langle \Delta f', f' \rangle}{\langle f', f' \rangle} - \lambda \le B \cdot ||f||_{\infty}^2 \cdot \ell \cdot k,$$

where B = B(w) > 0 is a constant depending only on w and ε .

Proof. Once and for all, fix a symmetric smooth function $J_0: \mathbb{R} \to [0,1]$ such that

$$J_0(x) = \begin{cases} 1 & \text{if } |x| \ge 2, \\ 0 & \text{if } |x| \le 1. \end{cases}$$

Using this function, we define $J: C_w(\gamma) \to \mathbb{R}$ by

$$J(x) = J_0\left(\frac{2 \cdot d(x, \gamma)}{w}\right)$$

and define $f': X \to \mathbb{R}$ by

$$f'(x) = \begin{cases} J(x) \cdot f(x) & \text{if } x \in C_w(\gamma) \\ f(x) & \text{otherwise.} \end{cases}$$

f' satsifies the first two thirds of the claim, so we need to verify the last third.

We will work with one curve (i.e. k = 1), because we assume disjointness, the argument in the general case is the same, except that the final errors multiply by k.

 $^{^{3}}$ This is a technical condition we set to make the bound come out nicer. The regime we will be interested in is when this quantity tends to 0 in any event.

We now first estimate the numerator of the Rayleigh quotient. Using Theorem 5.2, we obtain:

$$\begin{split} \langle \Delta f', f' \rangle - \langle \Delta f, f \rangle &= \int_{C_w(\gamma)} (\Delta J f) \cdot J f - (\Delta f) \cdot f \\ &= \int_{C_w(\gamma)} \left(|\nabla J|^2 + \left| \nabla (\sqrt{1 - J^2}) \right|^2 \right) \cdot f^2 \\ &- \int_{C_w(\gamma)} \Delta (\sqrt{1 - J^2} f) \cdot \sqrt{1 - J^2} f \\ &\leq \int_{C_w(\gamma)} \left(|\nabla J|^2 + \left| \nabla (\sqrt{1 - J^2}) \right|^2 \right) \cdot f^2 \\ &\leq ||f||_{\infty}^2 \cdot \int_{C_w(\gamma)} \left(|\nabla J|^2 + \left| \nabla (\sqrt{1 - J^2}) \right|^2 \right) \end{split}$$

Here we have used the fact that Δ is a positive operator in order to get from the second to the third line.

Now, using the fact $C_w(\gamma)$ is standard and that the function J depends on the distance to the core curve only, we get that

$$\begin{split} \int_{C_w(\gamma)} \left(|\nabla J|^2 + \left| \nabla (\sqrt{1 - J^2}) \right|^2 \right) \\ &= \ell \cdot \int_{-w}^w \left[\left(\frac{d}{d\rho} J_0\left(\frac{\rho}{w}\right) \right)^2 + \left(\frac{d}{d\rho} \sqrt{1 - J_0\left(\frac{\rho}{w}\right)^2} \right)^2 \right] \cdot \cosh(\rho) d\rho \\ &= \frac{\ell}{w} \int_{-1}^1 \left[\left(\frac{dJ_0(y)}{dy} \right)^2 + \left(\frac{d}{dy} \sqrt{1 - J_0(y)^2} \right)^2 \right] \cdot \cosh(w \cdot y) dy \\ &\leq A_1 \cdot \frac{\ell \cdot \cosh(w)}{w} \end{split}$$

where $A_1 > 0$ is a constant depending on our choice of J_0 alone. The division by w in the beforelast line comes from the combination of the chain rule and the substitution rule. All in all, we obtain

$$\langle \Delta f', f' \rangle \leq \langle \Delta f, f \rangle + A_1 \cdot ||f||_{\infty}^2 \cdot \ell.$$

Likewise, for the denominator, we have:

$$\langle f', f' \rangle = \langle f, f \rangle - \int_{C_{\gamma}} (1 - J^2) f^2 \ge 1 - ||f||_{\infty}^2 \cdot \int_{C_{\gamma}} (1 - J^2) \ge 1 - A_2 \cdot ||f||_{\infty}^2 \cdot \ell \cdot w \cdot \cosh(w)$$

for some constant $A_2 > 0$ depending on J_0 only.

This means that, setting $E = A_3 \cdot ||f||_{\infty}^2 \cdot \ell$ (where $A_3 > 0$ is large enough to absorb the previous constants and the factors depending on w),

$$\frac{\langle \Delta f', f' \rangle}{\langle f', f' \rangle} \le \frac{\langle \Delta f, f \rangle + E}{\langle f, f \rangle - E} = \frac{\lambda + E}{1 - E} \le \lambda + A_4 \cdot E,$$

WILL HIDE AND BRAM PETRI

for some constant $A_4 > 0$, where we've used the geometric series around E = 0 and our assumption that $||f||_{\infty}^2 \cdot \ell < \varepsilon$. Here we can choose any $\varepsilon \in (0, 1/A_3)$.

6. Building good surfaces

The goal of this section is to prove the main result of this article: we can produce sequences of covers of the Bolza surface whose spectral gaps are dense in $[0, \frac{1}{4}]$. We will use a variant of the switching technique from [Mag24a], based on the model of random surfaces described in Section 2.

6.1. The order of constants. There are many constants involved in our construction and even though they're uniform, the order in which they're fixed matters. So we will record this order here:

- C1 We fix a constant $\eta > 0$. Our goal will be to build a set of covers whose spectral gaps are η -dense in $[0, \frac{1}{4}]$.
- **C2** We also once and for all fix a parameter w > 0 (independent of η). This number appears as a collar width in Proposition 5.3 and determines another parameter $\varepsilon > 0$ and a third number B = B(w) that both appear in that same proposition.
- C3 Given all the parameters above, we fix any $\ell > 4w$ such that

$$\frac{1}{4} \cdot \left(1 - \frac{2\pi}{\ell + 2\pi}\right)^2 > \frac{1}{4} - \frac{1}{2} \cdot \eta \quad \text{and} \quad \ell \cdot e^{-\ell \cdot \sqrt{\eta}} < \min\{\varepsilon/A, \eta/B\},$$

where B = B(w) is the constant that appears in Proposition 5.3 and A the constant that appears in Theorem 5.1.

6.2. Building a good tree cover. From hereon out the parameters chosen in the previous subsection will be considered fixed.

We will now first prove that we can find good tree covers:

Proposition 6.1. The Bolza surface admits a tree cover $T_{\phi} \to X_B$, coming from a homeomorphism $\phi : X_B \to X_B$ such that the following conditions hold:

- (1) $\operatorname{sys}(T_{\phi}) > \ell$,
- (2) the collars of width w around all geodesics in the canonical pants decomposition of T_{ϕ} are all standard, and
- (3) $\lambda_1(T_{\phi}) > \frac{1}{4} \frac{1}{2} \cdot \eta.$

Proof. First of all, Mirakhani [Mir16, Theorem 1.2] determined the asymptotic distribution of the normalized length vector of a uniform random pants decomposition

$$\mathcal{P}_L \in \left\{ \mathcal{P} \in \mathrm{MCG}(X_B) \cdot \mathcal{P}_0 : \sum_{\alpha \in \mathcal{P}} \ell_\alpha(X_B) \le L \right\},\$$

where $MCG(X_B)$ denotes the mapping class group of X_B . The normalized length vector of a pants decomposition is the vector

$$\overline{\ell}(\mathcal{P}_L) = \left(\ell_{\alpha}(X_B) \middle/ \sum_{\beta \in \mathcal{P}_L} \ell_{\beta}(X_B) \right)_{\alpha \in \mathcal{P}_L} \in \sigma_3 = \left\{ x \in \mathbb{R}^3 : \begin{array}{c} x_i \ge 0, \ i = 1, 2, 3\\ \text{and } x_1 + x_2 + x_3 = 1 \end{array} \right\}$$

Mirzakhani proved that, as $L \to \infty$, this random vector converges in distribution to a random vector in σ_3 distributed according to a measure that can be explicitly computed. This was generalized to other multi-curves by Arana-Herrera [AH22] and Liu [Liu22].

For us, the only thing that matters is that the limit measure is absolutely continuous with respect to the Lebesgue measure with a Radon–Nikodym derivative that is strictly positive in the interior of σ_3 . This implies that for all L > 0 and all $\delta > 0$, we can find a homeomorphism $\phi : X_B \to X_B$ such that the pants decomposition $\phi^{-1}(\mathcal{P}_0) = (\beta_1, \beta_2, \beta_3)$ satifies

$$\ell(\beta_i) \ge L$$
 and $1 - \delta < \frac{\ell(\beta_i)}{\ell(\beta_i)} < 1 + \delta$ for all $i, j \in \{1, 2, 3\}$

We claim that the tree cover $T_{\phi} \longrightarrow X_B$ has the properties we require, as soon as $L > \ell/(1-\delta)$.

We start with item 1: the systole. First observe that, if the systole corresponds to a pants curve in the canonical pants decomposition of T_{ϕ} , then it automatically satisfies the inequality above. If it does not, the geodesic realizing it contains at least two arcs that each run between two points on one of the boundary components of one of the pairs of pants in the pants decomposition. Indeed, the systole is necessarily simple, so it crosses at least two pairs of pants. As such the path it traces in the tree dual to the canonical pants decomposition has at least two leaves. So the length of the systole is at least twice the length of such an arc. All that remains is to find a lower bound on this length.

To do this, consider the arc α in Figure 2. The arc α cuts the boundary component



FIGURE 2. A pair of pants

 β_1 into two arcs, the shortest one of which has length at most $\frac{1+\delta}{2}L$. If we suppose this shortest arc lies on the side of β_2 , we obtain

$$\ell(\alpha) + \frac{1+\delta}{2}L \ge \ell(\beta_2) \ge L,$$

which gives us $\ell(\alpha) \geq \frac{1-\varepsilon}{2}L$ and hence

$$\operatorname{sys}(T_{\phi}) \ge (1-\delta) \cdot L > \ell.$$

We prove item 2 using essentially the same argument. Indeed, if a collar of width u around one of the pants curves is not standard, then this yields a simple arc of length at

most 2u between two points on the corresponding pants curve. This arc must be essential relative to its endpoints and by the same argument as above, such an arc has length at least $\ell/2$. So, the collars of width $\ell/4 > w$ (by assumption C3) are standard.

Finally, item 3 follows from the second part of the same assumption C3, combined with Proposition 4.2.

6.3. Strong convergence. To obtain covers of X_B whose spectral gap is nearly optimal we will rely on the techniques developed by Hide–Magee [HM23, LM23] based on strong convergence of permutation representations.

Let Γ be a finitely generated group and suppose $(\mathcal{H}_n)_n$ is a sequence of Hilbert spaces. Given a sequence of unitary representations $\rho_n : \Gamma \to U(\mathcal{H}_n)$, where $U(V \mathcal{H}_n)$ denotes the unitary group of \mathcal{H}_n , we say that this sequence **strongly converges** to $\rho_\infty : \Gamma \to U(\mathcal{H}_\infty)$ if

$$\lim_{n \to \infty} ||\rho_n(z)|| = ||\rho_\infty(z)||$$

for all $z \in \mathbb{C}[\Gamma]$.

We will need the following result which is a version of the results in [Mag24b], specialized to our set-up:

Theorem 6.2. Let $\Gamma < PSL(2, \mathbb{R})$ be a uniform lattice and let Λ be a group. Moreover suppose that

- $p: \Gamma \to \Lambda$ is an epimorphism and write $K = \ker(p) \triangleleft \Gamma$,
- $\left(\rho_n \in \operatorname{Hom}(\Lambda, \mathfrak{S}_n)\right)_{n \ge 1}$ is a sequence of permutation representations such that $\operatorname{std}_n \circ \rho_n \xrightarrow{\operatorname{strong}} \rho_{\operatorname{reg}}, \quad as \ n \to \infty$

where $\operatorname{std}_n : \mathfrak{S}_n \to U(\mathbb{C}^{n-1})$ denotes the (n-1)-dimensional irreducible representation and $\rho_{\operatorname{reg}} : \Lambda \to \mathcal{B}(\ell^2(\Lambda))$ denotes the left regular representation, and

• write $\Gamma_n = \operatorname{Stab}_{\rho_n \circ p}(\{1\}) < \Gamma$

Then

$$\lim_{n \to \infty} \lambda_1(\Gamma_n \backslash \mathbb{H}^2) = \min \left\{ \lambda_1(K \backslash \mathbb{H}^2), \lambda_1(\Gamma \backslash \mathbb{H}^2) \right\}.$$

Proof sketch. The precise statement is not yet available in the literature although it can be derived using small modifications of the arguments in [HM23, LM23, MT23]. A representation theoretic argument due to Magee [Mag24b] is perhaps the fastest way to obtain it. We will very briefly describe how this works here.

The fact that std $\circ \rho_n \xrightarrow{\text{strong}} \rho_{\text{reg}}$, implies that

$$\operatorname{std}_n \circ \rho_n \circ p \xrightarrow{\operatorname{strong}} \rho_{\operatorname{reg}} \circ p : \Gamma \to \mathcal{B}(\ell^2(\Lambda)).$$

Magee [Mag24b, Theorem 3.1] proves that this implies that:

$$\operatorname{Ind}_{\Gamma}^{\operatorname{PSL}(2,\mathbb{R})}\left(\operatorname{std}_{n}\circ\rho_{n}\circ p\right)\xrightarrow{\operatorname{strong}}\operatorname{Ind}_{\Gamma}^{\operatorname{PSL}(2,\mathbb{R})}\left(\rho_{\operatorname{reg}}\circ p\right).$$

By [Mag24b, Proposition 2.9], this implies our statement.

Strongly convergent sequences of representations are notoriously hard to find. However, the following theorem by Bordenave–Collins [BC19] gives us what we need:

Theorem 6.3 (Bordenave–Collins). Let F be a finitely generated non-abelian free group and let $\rho_n \in \text{Hom}(F, \mathfrak{S}_n)$ be chosen uniformly at random. Then

$$\operatorname{std}_n \circ \rho_n \xrightarrow{strong} \rho_{\operatorname{reg}}, \quad as \ n \to \infty$$

in probability.

6.4. Covers of degree two and Hamming distance. Using good tree covers, combined with the existence of strongly convergent sequences of permutation representations of F_2 and Theorem 6.2 above, we can already use our construction to build sequences of finite covers of X_B whose spectral gaps are arbitrarily close to $\frac{1}{4}$. As such, we need to explain how to get intermediate spectral gaps, which is what the remainder of the text is devoted to.

From hereon out, we will suppose that $\phi : X_B \to X_B$ is a homeomorphism such that T_{ϕ} satisfies the result from Proposition 6.1. Recall that the sequence $\left(Y_{\phi}^{(n)} \to X_B\right)_{n\geq 1}$ denotes the associated sequence of random covers (see Section 2).

In order to obtain finite covers of X_B whose spectral gaps are η -dense, we will use a variation of the switching argument of Magee. Magee phrases the argument in terms of bundles over $Y_{\phi}^{(n)}$ coming from homomorphisms $\pi_1(Y_{\phi}^{(n)}) \to \mathbb{Z}/2\mathbb{Z}$. We will phrase it entirely in terms of covers of degree two of $Y_{\phi}^{(n)}$, which is an equivalent point of view.

First of all, we will restrict to covers that factor through the handle body attachment. That is, the groups defined in Section 2.3 fit in the following commutative diagram:

$$\begin{split} \Lambda_{\phi}^{(n)} & \longleftrightarrow & \Gamma_B \\ & \downarrow^{(\iota \circ \phi)_*} & \downarrow^{(\iota \circ \phi)_*} \\ & H^{(n)} & \longleftrightarrow & F_2 \end{split}$$

which allows us to define the set

$$\mathcal{H}_{\phi}^{(n)} = \left\{ \rho \circ (\iota \circ \phi)_* : \rho \in \operatorname{Hom}(H^{(n)}, \mathbb{Z}/2\mathbb{Z}) \right\}.$$

Given a free generating set $S = (s_1, \ldots, s_k)$ of $H^{(n)}$, we can identify the set of maps $\mathcal{H}_{\phi}^{(n)}$ with $(\mathbb{Z}/2\mathbb{Z})^k$ through the map that associates $(\rho(s_1), \ldots, \rho(s_k))$ to $\rho \circ (i \circ \phi)_* \in \mathcal{H}_{\phi}^{(n)}$. This in turn allows us to define a **Hamming distance** $d_S : \mathcal{H}_{\phi}^{(n)} \times \mathcal{H}_{\phi}^{(n)} \to \mathbb{N}$ by

$$d_S\left(\rho_1\circ(\iota\circ\phi)_*,\ \rho_2\circ(\iota\circ\phi)_*\right) = \left|\left\{j\ :\ \rho_1(s_j)\neq\rho_2(s_j)\right\}\right|.$$

We note that every element $\rho \in \mathcal{H}_{\phi}^{(n)}$ corresponds to a cover of $Y_{\phi}^{(n)}$. If the image of ρ is trivial, we will let this cover be the cover of $Y_{\phi}^{(n)}$ by two disjoint copies of itself. This way, all the corresponding covers are of degree two.

6.5. Switching. In this section we will describe how two covers of degree two of $Y_{\phi}^{(n)}$ whose Hamming distance is 1 differ from each other.



FIGURE 3. A local picture of a switch. The surface below represents a neighborhood of one of the prefered pants curves $\alpha_i^{(n)}$ of $Y_{\phi}^{(n)}$. The surfaces up top (one on the left and one on the right) are the neighborhoods of the lifts of $\alpha_i^{(n)}$ in two distinct covers of degree two. The shading is there to indicate that the curves that are involved are meridians.

To make this precise, we need to first decribe a free generating set of $H^{(n)}$. One way to do this is to identify F_2 with the fundamental group of a graph $W^{(1)}$. We will let $W^{(1)}$ be a wedge of two circles. Because the pants curves in \mathcal{P}_0 are meridians, the two edges of $W^{(1)}$ are dual to two of the pants curves in the pants decomposition \mathcal{P}_0 we've fixed. If we change the handlebody attachment by a homeomorphism $\phi : X_B \to X_B$, this changes the identification. That is, we can think of the edges of $W^{(1)}$ as corresponding to two of the three pants curves in $\phi^{-1}(\mathcal{P}_0)$. $H^{(n)}$ is the fundamental group of a graph $W^{(n)}$ that covers $W^{(1)}$. The edges in this graph are dual to the canonical pants decomposition of $Y^{(n)}_{\phi}$.

In order to obtain a prefered free generating set of $H^{(n)}$, we pick an arbitrary spanning tree in $W^{(n)}$ and contract it. The edges that do not appear in the tree yield a generating set $S^{(n)} = (s_1, \ldots, s_k)$ of $H^{(n)}$. Because of the correspondance between edges and pants curves described above, each of these generators is dual to a pants curve in the canonical pants decomposition of $Y^{(n)}_{\phi}$. We will denote these prefered pants curves by $(\alpha_1^{(n)}, \ldots, \alpha_k^{(n)})$. Given a map $\rho: H^{(n)} \to \mathbb{Z}/2\mathbb{Z}$, the corresponding cover of degree two of $Y_{\phi}^{(n)}$ can be built as follows:

- Write $Z_{\phi}^{(n)} = \overline{Y_{\phi}^{(n)} \bigcup_{i=1}^{k} \alpha_i}$, where \overline{X} denotes the completion of a hyperbolic surface X, so $Z_{\phi}^{(n)}$ is a surface with boundary,
- take two disjoint copies Z_1 and Z_2 of $Z_{\phi}^{(n)}$, and
- for each pants curve α_i :
 - if $\rho(s_i) = [0]$, glue the two boundary components of Z_1 corresponding to α_i back together and do the same for those in Z_2 ,
 - if $\rho(s_i) = [1]$, glue the "left" boundary component of Z_1 corresponding to α_i to the "right" boundary component of Z_2 and, likewise, glue the corresponding right boundary component of Z_1 to the left boundary component of Z_2 .

All these gluings above should be performed with the correct twist, so as to guarantee that the resulting surface covers $Y_{\phi}^{(n)}$.

It follows from this desciption that, if

$$\mathbf{d}_{S^{(n)}}\left(\rho_1 \circ (i \circ \phi)_*, \ \rho_2 \circ (i \circ \phi)_*\right) = 1,$$

then the two corresponding covers can be obtained from each other by opening up the lifts corresponding to one of the prefered pants curves $\alpha_i^{(n)} \subset Y_{\phi}^{(n)}$ and gluing them back together with the opposite pattern. We will call this operation a **switch** – this same terminology is used in graph theory for what happens at the level of the 2-covers of the graph $W^{(n)}$. Figure 3 shows a local picture of this move.

6.6. Finishing the proof. We now have everything we need in place to finish the proof of our main theorem. Indeed, Theorem 1 is implied directly by the following result:

Theorem 6.4. For all $n \in \mathbb{N}$ large enough, the probability that the set

$$\left\{ \, \lambda_1(\widehat{Y}) \, : \, \widehat{Y} \, \, is \, \, a \, \, connected \, \, cover \, \, of \, Y_{\phi}^{(n)} \, \, of \, \, degree \, \, two \,
ight\}$$

is η -dense in $[0, \frac{1}{4}]$, is strictly positive.

Proof. First of all, we use Proposition 6.1 to choose a homeomorphism $\phi : X_B \to X_B$ that gives us a tree cover with the properties in that same proposition.

It now follows from Theorems 6.2 and 6.3 that $\lambda_1(Y_{\phi}^{(n)}) > \frac{1}{4} - \eta$ with high probability. In fact, using the same argument as in [Mag24a, Section 4],

$$\theta \otimes \rho_n \xrightarrow{\text{strong}} \rho_{\text{reg}}$$

in probability as $n \to \infty$, for any $\theta \in \text{Hom}(F_2, \mathbb{Z}/2\mathbb{Z})$. For us this means that, again using Theorem 6.2, we obtain covers of degree two, corresponding to some element in $\mathcal{H}_{\phi}^{(n)}$, of

 $Y_{\phi}^{(n)}$ whose spectral gap is at least

$$\lambda_1(T_\phi) - \eta/2 > \frac{1}{4} - \eta$$

for n large enough.

The set $\mathcal{H}_{\phi}^{(n)}$ also contains the trivial homorphism, the corresponding degree two cover of which has $\lambda_1(\hat{Y}) = 0$.

We can go between these covers using a finite number of switches, so as soon as we prove that a single switch doesn't change the spectral gap by more than η , we are done.

In order to do this, we need to make one further assumption on the surfaces $Y^{(n)}$: the preferred pants curves in $Y^{(n)}$ need to have collar width at least w and the systole of $Y^{(n)}$ needs to be at least ℓ .

Both of these statements are true in the tree cover T_{ϕ} . This means, that if they're not true in $Y^{(n)}$, there needs to be a closed geodesic $\gamma \subset X_B$ of uniformly bounded length $L = L(w, \ell)$ such that

(6.1)
$$\iota_*(\gamma) \neq e \quad \text{but} \quad \rho_n \circ (\iota \circ \phi)_*(\gamma) \cdot 1 = 1.$$

It follows from Nica's work [Nic94] that, with asymptotically strictly positive probability, $\rho_n(g)$ is fixed point free for all g in any fixed finite set of non-trivial conjugacy classes in F_2 . The statement we use here is the following. First of all, let $K_1, \ldots, K_m \subset F_2$ be a finite collection of distinct non-trivial conjugacy classes, none of which contain a power of a non-trivial element of F_2 . Moreover, let (c_1, \ldots, c_k) be a finite set of cycle lengths. Then the vector of random variables $\left(X_{c_i}(K_j)\right)_{1\leq i\leq m, 1\leq j\leq k}$, where $X_{c_i}(K_j)$ denotes the number of c_i -cycles in the image of an element of K_j under a uniformly random homomorphism $\rho \in \text{Hom}(F_2, \mathfrak{S}_n)$ converges in distribution to a vector of independent random variables. Furthermore, the limit of $X_{c_i}(K_j)$ is Poisson $(1/c_i)$ -distributed. This statement is not literally contained in Nica's paper, but it can be proven using similar methods. Versions of this statement are also available in [LP10, Section 4], [BP23, Theorem 1.4], [PZ24, Corollary 1.7 and Theorem 1.14]. The only consequence we need is that, there exists a uniform a > 0such that for all n large enough:

$$\mathbb{P}\left(X_{c_i}(K_j)=0 \text{ for } i=1,\ldots,k \text{ and } j=1,\ldots,m\right) > a.$$

Also observe that if $g = u^k \in F_2$ and if, under a permutation representation of F_2 , u has no d-cycles for any divisor d of k, then g has no fixed points.

Since Γ_B contains a finite number of conjugacy classes of elements of bounded translation length, we can avoid the existence of a closed geodesic $\gamma \subset X_B$ as in (6.1) with strictly positive probability. Moreover, the existence of a 2-cover with spectral gap above $\frac{1}{4} - \eta$ has probability tending to 1 as $n \to \infty$, so this property is preserved when we restrict to covers without short geodesics that don't lift to T_{ϕ} . In other words, the probability that $Y_{\phi}^{(n)}$ has systole at least ℓ , all the curves in the canonical pants decomposition have standard collars of width w and $Y_{\phi}^{(n)}$ has a 2-cover with spectral gap at least $\frac{1}{4} - \eta$, is asymptotically strictly positive. So now suppose \widehat{Y}_1 and \widehat{Y}_2 are two covers of degree two of $Y_{\phi}^{(n)}$ that differ by a switch move. Moreover, let f be a $\lambda_1(\widehat{Y}_1)$ -eigenfunction on \widehat{Y}_1 and let α_i be the curve in which we need to switch to obtain \widehat{Y}_2 . Using Proposition 5.3, there exists a smooth function f' that restricts to the 0 function on both lifts of α_i and

$$\frac{\langle \Delta f', f' \rangle}{\langle f', f' \rangle} \le \lambda_1(\widehat{Y}_2) + \eta_2$$

Here we've used both the assumption on the collar widths and the systole of $Y_{\phi}^{(n)}$ and the relations between our constants (Section 6.1). Because f' cancels on the lifts of α_i , it defines a function on the cover \hat{Y}_2 and hence

$$\lambda_1(\widehat{Y}_2) \le \lambda_1(\widehat{Y}_1) + \eta_2$$

by the variational characterization of λ_1 , and we are done.

References

- [AH22] Francisco Arana-Herrera. Counting hyperbolic multigeodesics with respect to the lengths of individual components and asymptotics of Weil-Petersson volumes. Geom. Topol., 26(3):1291–1347, 2022.
 [AM99] Colin Adams and Frank Morgan. Isoperimetric curves on hyperbolic surfaces. Proc. Amer.
- [AM99] Colin Adams and Frank Morgan. Isoperimetric curves on hyperbolic surfaces. Proc. Amer. Math. Soc., 127(5):1347–1356, 1999.
- [AW23] Noga Alon and Fan Wei. The limit points of the top and bottom eigenvalues of regular graphs. Preprint, arXiv:2304.01281, 2023.
- [Bav96] C. Bavard. Disques extrémaux et surfaces modulaires. Ann. Fac. Sci. Toulouse Math. (6), 5(2):191–202, 1996.
- [BC19] Charles Bordenave and Benoît Collins. Eigenvalues of random lifts and polynomials of random permutation matrices. Ann. of Math. (2), 190(3):811–875, 2019.
- [BCP21] Thomas Budzinski, Nicolas Curien, and Bram Petri. On the minimal diameter of closed hyperbolic surfaces. *Duke Math. J.*, 170(2):365–377, 2021.
- [Ber16] Nicolas Bergeron. *The spectrum of hyperbolic surfaces*. Universitext. Springer, Cham; EDP Sciences, Les Ulis, 2016. Appendix C by Valentin Blomer and Farrell Brumley, Translated from the 2011 French original by Brumley [2857626].
- [Bon22] J. Bonifacio. Bootstrapping closed hyperbolic surfaces. J. High Energy Phys., 93:Paper No. 093, 18, 2022.
- [BP23] Elizabeth Baker and Bram Petri. Statistics of finite degree covers of torus knot complements. Ann. H. Lebesgue, 6:1213–1257, 2023.
- [Bus10] Peter Buser. Geometry and spectra of compact Riemann surfaces. Modern Birkhäuser Classics. Birkhäuser Boston, Ltd., Boston, MA, 2010. Reprint of the 1992 edition.
- [CFKS87] H. L. Cycon, R. G. Froese, W. Kirsch, and B. Simon. Schrödinger operators with application to quantum mechanics and global geometry. Texts and Monographs in Physics. Springer-Verlag, Berlin, study edition, 1987.
- [Cha84] Isaac Chavel. *Eigenvalues in Riemannian geometry*, volume 115 of *Pure and Applied Mathematics*. Academic Press, Inc., Orlando, FL, 1984. Including a chapter by Burton Randol, With an appendix by Jozef Dodziuk.
- [Che70] Jeff Cheeger. A lower bound for the smallest eigenvalue of the Laplacian. Problems in analysis (Papers dedicated to Salomon Bochner, 1969), pages 195–199, 1970.
- [Che75] S. Y. Cheng. Eigenvalue comparison theorems and its geometric applications. *Math. Z.*, 143(3):289–297, 1975.
- [Con15] M. D. E. Conder. Quotients of triangle groups acting on surfaces of genus 2 to 101. math.auckland.ac.nz/~conder/TriangleGroupQuotients101.txt, 2015.

20	WILL HIDE AND BRAM PETRI
[CP23]	G. Cornelissen and N. Peyerimhoff. Twisted isospectrality, homological wideness, and isometry—a sample of algebraic methods in isospectrality. SpringerBriefs in Mathematics. Springer, Cham. [2023] ©2023.
[Dix69]	J. D. Dixon. The probability of generating the symmetric group. <i>Math. Z.</i> , 110:199–205, 1969.
[DM24]	Dingding Dong and Theo McKenzie. Arbitrary spectral edge of regular graphs. Preprint, arXiv:2412.09570, 2024.
[FBGMPP23]	Maxime Fortier Bourque, Émile Gruda-Mediavilla, Bram Petri, and Mathieu Pineault. Two counterexamples to a conjecture of Colin de Verdière on multiplicity. Preprint, arXiv:2312.03504, 2023.
[FBP23]	Maxime Fortier Bourque and Bram Petri. Linear programming bounds for hyperbolic surfaces. Preprint, arXiv:2302.02540, 2023.
[Gag80]	M. E. Gage. Upper bounds for the first eigenvalue of the Laplace-Beltrami operator. <i>Indiana</i> Univ. Math. J., 29(6):897–912, 1980.
[GAP22] [GLMST21]	The GAP Group. GAP – Groups, Algorithms, and Programming, Version 4.12.2, 2022. Clifford Gilmore, Etienne Le Masson, Tuomas Sahlsten, and Joe Thomas. Short geodesic loops and L^p norms of eigenfunctions on large genus random surfaces. Geom. Funct. Anal., 31(1):62-110, 2021.
[HM23]	Will Hide and Michael Magee. Near optimal spectral gaps for hyperbolic surfaces. Ann. of Math. (2), 198(2):791–824, 2023.
[Jen81]	F. Jenni. Ueber das spektrum des Laplace-operators auf einer schar konmpakter Rie- mannscher flächen. PhD thesis, University of Basel, 1981.
[Joh17]	F. Johansson. Arb: efficient arbitrary-precision midpoint-radius interval arithmetic. <i>IEEE Transactions on Computers</i> , 66:1281–1292, 2017.
[Kee74]	Linda Keen. Collars on Riemann surfaces. In <i>Discontinuous groups and Riemann surfaces</i> (<i>Proc. Conf., Univ. Maryland, College Park, Md., 1973</i>), volume No. 79 of <i>Ann. of Math.</i> <i>Stud.</i> , pages 263–268. Princeton Univ. Press, Princeton, NJ, 1974.
[KMP24]	Petr Kravchuk, Dalimil Mazáč, and Sridip Pal. Automorphic spectra and the conformal bootstrap. Commun. Am. Math. Soc., 4:1–63, 2024.
[Liu22]	Mingkun Liu. Length statistics of random multicurves on closed hyperbolic surfaces. <i>Groups Geom. Dyn.</i> , 16(2):437–459, 2022.
[LM23]	Larsen Louder and Michael Magee. Strongly convergent unitary representations of limit groups. Preprint, arXiv:2210.08953. With an appendix by Will Hide and Michael Magee, 2023.
[LP10]	Nati Linial and Doron Puder. Word maps and spectra of random graph lifts. <i>Random Structures Algorithms</i> , 37(1):100–135, 2010.
[Mag24a]	Michael Magee. The limit points of the bass notes of arithmetic hyperbolic surfaces. Preprint, arXiv:2403.00928, 2024.
[Mag24b]	Michael Magee. Strong convergence of unitary and permutation representations of discrete groups, to appear, 2024+.
[Mat05]	Katsuhiko Matsuzaki. Isoperimetric constants for conservative Fuchsian groups. Kodai Math. J., 28(2):292–300, 2005.
[Mat24]	Joffrey Mathien. Diameter of a new model of random hyperbolic surfaces. Preprint, arXiv:2403.01925, 2024.
[Mir13]	Maryam Mirzakhani. Growth of Weil-Petersson volumes and random hyperbolic surfaces of large genus I Differential Geom 94(2):267–300 2013
[Mir16]	Maryam Mirzakhani. Counting mapping class group orbits on hyperbolic surfaces. Preprint, arXiv:1601.03342, 2016
[MNP22]	Michael Magee, Frédéric Naud, and Doron Puder. A random cover of a compact hyperbolic surface has relative spectral gap $\frac{3}{2} = c_{c} C_{com} E_{unct} Angl. 32(3):505.661, 2022$
[MP23]	Surface has relative spectral gap $\frac{1}{16} - \varepsilon$. Geom. Funct. Anal., 52(5):555–601, 2022. Michael Magee and Doron Puder. The asymptotic statistics of random covering surfaces. Forum Math. Pi, 11:Paper No. e15, 51, 2023.

- [MR03] Colin Maclachlan and Alan W. Reid. The arithmetic of hyperbolic 3-manifolds, volume 219 of Graduate Texts in Mathematics. Springer-Verlag, New York, 2003.
 [MT23] Michael Magee and Joe Thomas. Strongly convergent unitary representations of right-angled artin groups. Preprint, arXiv:2308.00863, 2023.
- [Nic94] Alexandru Nica. On the number of cycles of given length of a free word in several random permutations. *Random Structures Algorithms*, 5(5):703–730, 1994.
- [PZ24] Doron Puder and Tomer Zimhoni. Local statistics of random permutations from free products. Int. Math. Res. Not. IMRN, (5):4242–4300, 2024.
- [Sar23] Peter Sarnak. Spectra of locally uniform geometries. Chern Lectures, U.C. Berkeley. Available at: http://publications.ias.edu/sarnak/paper/2728, 2023.
- [SU13] A. Strohmaier and V. Uski. An algorithm for the computation of eigenvalues, spectral zeta functions and zeta-determinants on hyperbolic surfaces. *Comm. Math. Phys.*, 317:827–869, 2013.
- [Tak77] Kisao Takeuchi. Arithmetic triangle groups. J. Math. Soc. Japan, 29(1):91–106, 1977.
- [The21] The Sage Developers. SageMath, the Sage Mathematics Software System (Version 9.3), 2021. https://www.sagemath.org.

(Will Hide) MATHEMATICAL INSTITUTE, UNIVERSITY OF OXFORD, ANDREW WILES BUILDING, OX2 6GG OXFORD, UNITED KINGDOM

Email address: william.hide@maths.ox.ac.uk

(Bram Petri) Institut de Mathématiques de Jussieu-Paris Rive Gauche and Institut universitaire de France ; Sorbonne Université and Université Paris Cité, CNRS, IMJ-PRG, F-75005 Paris, France

Email address: bram.petri@imj-prg.fr