INDUCED ARITHMETIC REMOVAL FOR PARTITION-REGULAR PATTERNS OF COMPLEXITY 1

V. GLADKOVA

ABSTRACT. In 2019, Fox, Tidor and Zhao [7] proved an induced arithmetic removal lemma for linear patterns of complexity 1 in vector spaces over a fixed finite field. With no further assumptions on the pattern, this induced removal lemma cannot guarantee a fully patternfree recolouring of the space, as some 'non-generic' instances must necessarily remain. On the other hand, Bhattacharyya et al. [3] showed that in the case of translation-invariant patterns, it is possible to obtain recolourings that eliminate the given pattern completely, with no exceptions left behind. This paper demonstrates that such complete removal can be achieved for all partition-regular arithmetic patterns of complexity 1.

1. INTRODUCTION

The triangle removal lemma for graphs and, more generally, arbitrary subgraph removal lemmas [5] are a well-known application of Szemerédi's regularity lemma [20]. These removal results state that if a graph G contains few copies of a graph H as a subgraph, then G can be made H-free by removing only a small proportion of its edges. The same holds for H occurring as an induced subgraph, as proved by Alon, Fischer, Krivelevich, and Szegedy [1] via a strong variant of the regularity lemma, which produces two nested vertex partitions with certain desirable properties.

Analogous induced removal lemmas in the setting of vector spaces over finite fields assert that if \mathbb{F}_p^n is *r*-coloured and a given arithmetic pattern occurs with small density under this colouring, then it is possible to recolour a small proportion of the space to eliminate (nearly) all instances of the pattern. Here, an arithmetic pattern is a solution to a given system of equations with a specified assignment of colours. The following definition specifies a pattern using a collection of linear forms but this can be readily translated to the language of systems of equations by standard methods, as laid out in Section 4.

Definition 1.1 (Arithmetic patterns). An arithmetic pattern \mathcal{H} in l variables is a tuple $(\mathcal{L}, \mathcal{X})$, where \mathcal{L} is a collection of linear forms $\{L_1, \ldots, L_m\}$ in l variables over \mathbb{F}_p , and $\mathcal{X} = \{\chi : [m] \to [r]\}$ is a collection of r-colourings.

Given an r-colouring ϕ of \mathbb{F}_p^n , an instance of \mathcal{H} under ϕ is an l-tuple $\underline{x} \in (\mathbb{F}_p^n)^l$ such that for some $\chi \in \mathcal{X}$, $\phi(L_j(\underline{x})) = \chi(j)$ for each $1 \leq j \leq m$. A colouring ϕ is said to be \mathcal{H} -free if there are no instances of \mathcal{H} under ϕ .

Note. Unlike in previous work, this definition of an arithmetic pattern incorporates multiple colourings at once. In this way, a monochromatic 3-term arithmetic progression can be expressed as a single pattern instead of a union of r separate patterns. For example, when r = 2, this is given by $(\{x, x + d, x + 2d\}, \{\chi_1, \chi_2\})$, where $\chi_i : [3] \rightarrow [2]$ is the constant colouring $\chi_i \equiv i$.

Induced arithmetic removal lemmas were established in the work of [4, 3, 7], and [21], but, unlike in the graph-theoretic setting, the results of both [7] and [21] only hold with the caveat that a small number of 'non-generic' instances might remain behind. In the case of complexity 1 patterns, Fox, Tidor and Zhao [7] proved the following, with $\Lambda_{\mathcal{H}}(\phi)$ denoting the density of the instances of \mathcal{H} under a colouring ϕ (this is defined formally in Definition 2.1).

Theorem 1.2 (Induced removal for complexity 1 patterns [7]). Fix $\epsilon > 0$, an integer r > 0, and an arithmetic pattern \mathcal{H} of complexity 1. There exists a $\delta = \delta(\epsilon, r, \mathcal{H})$ satisfying the following. If $\phi : \mathbb{F}_p^n \to [r]$ is an r-colouring of \mathbb{F}_p^n such that $\Lambda_{\mathcal{H}}(\phi) \leq \delta$, then ϕ can be made \mathcal{H} -free on $\mathbb{F}_p^n \setminus \{0\}$ by recolouring at most an ϵ -proportion of \mathbb{F}_p^n .

This result cannot be extended to recolour all of \mathbb{F}_p^n when the arithmetic pattern in question is not partition-regular (see Example 1.4). The definition of partition regularity in the context of vector spaces over finite fields follows the work of Bergelson, Deuber, and Hindman [2].

Definition 1.3 (Partition regularity). An $l \times m$ matrix A with coefficients in \mathbb{F}_p is said to be partition-regular if the following holds. For any $r \in \mathbb{N}$, there exists N = N(r, p, m) such that, if $n \geq N$, $A\underline{x} = 0$ has a monochromatic solution in $\mathbb{F}_p^n \setminus \{0\}$ under any r-colouring of the space.

Example 1.4. Consider a matrix A over \mathbb{F}_p that is not partition-regular. Then for any $r, n \in \mathbb{N}$, there is an r-colouring of \mathbb{F}_p^n such that the only monochromatic solution to $A\underline{x} = 0$ is 0. However, this remains a monochromatic solution no matter how the space is recoloured. (A special case of this was given in [7, Non-example 1.3].)

As a consequence, any induced arithmetic removal lemma for general arithmetic patterns must necessarily make exceptions for instances of a certain form. However, for partitionregular patterns (including translation-invariant patterns), an obstacle such as in Example 1.4 does not apply: indeed, any colouring of \mathbb{F}_p^n must have many monochromatic instances of such patterns (see Theorem 4.2), whereas the induced removal lemma only applies when the number of instances is small to begin with.

In fact, the induced removal lemma of Bhattacharyya et al. [3] for translation-invariant patterns guarantees the removal of *all* instances. Such complete removal is possible due to the additional flexibility afforded by being able to translate instances of the pattern without breaking their structure (see the discussion in Section 3).

The main result of this chapter demonstrates that it is possible to arrange for similar flexibility and therefore achieve complete removal in the case of partition-regular patterns of complexity 1.

Theorem 1.5 (Induced removal for partition-regular patterns). Fix $\epsilon > 0$, an integer r > 0, and a partition-regular pattern \mathcal{H} of complexity 1. There exists a $\delta = \delta(\epsilon, r, \mathcal{H})$ satisfying the following. If $\phi : \mathbb{F}_p^n \to [r]$ is an r-colouring of \mathbb{F}_p^n such that $\Lambda_{\mathcal{H}}(\phi) \leq \delta$, then ϕ can be made \mathcal{H} -free on the whole of \mathbb{F}_p^n by recolouring at most an ϵ -proportion of \mathbb{F}_p^n .

The following example serves as an illustration of this theorem.

Example 1.6. Let \mathcal{H} be the arithmetic pattern encoding rainbow solutions to x + y + z - w = 0 under ϕ , noting that this is a partition-regular pattern of complexity 1 which is not translation-invariant when p > 2. For a given $\epsilon > 0$, let $\phi : \mathbb{F}_p^n \to [4]$ be any surjective 4-colouring such that $0 \in \phi^{-1}(1)$ and $|\phi^{-1}(1)| \leq \delta |\mathbb{F}_p^n|^4$, where $\delta = \delta_{1.5}(\epsilon, 4, \mathcal{H})$. Since one of the colour classes is small, there are at most $\delta |\mathbb{F}_p^n|^3$ instances of \mathcal{H} under ϕ . Then Theorem 1.5 implies that it is possible to eliminate all rainbow solutions to x + y + z - w = 0 by recolouring at most an ϵ -proportion of \mathbb{F}_p^n .

Of course, in this example, this is easy to see without resorting to Theorem 1.5, as we can simply replace all of $\phi^{-1}(1)$ with another colour. However, note that the induced removal lemma of Fox, Tidor and Zhao [7] (Theorem 1.2) applied to this same setting would only eliminate rainbow solutions for which each of x, y, z, w is non-zero.

Additionally, an easy corollary is provided here as another instance of the kind of results implied by Theorem 1.5. Here $A\Delta H$ denotes the symmetric difference of A and H, so that (iii) says that A is ϵ -close to being a subspace.

Corollary 1.7. Fix $\epsilon > 0$, and let A be a subset of \mathbb{F}_p^n . There exists a $\delta = \delta(\epsilon)$ such that one of the following holds:

- (i) $|A| \leq \epsilon |\mathbb{F}_p^n|;$
- (*ii*) $|(A+A) \cap A^c| > \delta |\mathbb{F}_p^n|;$
- (iii) there is a subspace $H \leq \mathbb{F}_p^n$ such that $|A\Delta H| \leq \epsilon |\mathbb{F}_p^n|$.

Proof. Let $\phi : \mathbb{F}_p^n \to \{0, 1\}$ be the 2-colouring $\phi = \mathbb{1}_A$, and let \mathcal{H} be the arithmetic pattern corresponding to the solutions of x + y = z for which x, y have colour 1 and z has colour 0. This is a partition-regular pattern of complexity 1, so Theorem 1.5 applies. If $|(A+A) \cap A^c| \leq \delta |\mathbb{F}_p^n|$ where $\delta = \delta_{1.5}(\epsilon, 2, \mathcal{H})$, then there are at most $\delta |\mathbb{F}_p^n|^2$ such solutions to x + y = z under ϕ . Therefore, by Theorem 1.5, there is a 2-colouring $\psi : \mathbb{F}_p^n \to \{0, 1\}$ that differs from ϕ in at most $\epsilon |\mathbb{F}_p^n|$ places such that ψ has no solutions to x + y = z satisfying $\psi(x) = \psi(y) = 1$ and $\psi(z) = 0$.

In particular, letting $A' = \psi^{-1}(1)$, this implies that $A' + A' \subseteq A'$. Hence A' is either a subspace or an empty set. Moreover, $|A\Delta A'| \leq \epsilon |\mathbb{F}_p^n|$, as $A\Delta A'$ is precisely equal to the set of $x \in \mathbb{F}_p^n$ such that $\phi(x) \neq \psi(x)$. Therefore either (i) or (iii) holds, as required. \Box

The proof of Theorem 1.5 follows the same lines as existing induced removal lemmas but utilises a more general subcoset selection scheme, reminiscent of the argument for the induced graph removal lemma [1]. The proposed approach is sketched out in Section 3 before being formally applied in Section 5.

Acknowledgments. The author would like to thank Julia Wolf for many discussions and advice given in the course of this research, as well as Sean Prendiville and Julian Sahasrabudhe for their helpful comments on an earlier version of this paper.

2. Preliminaries

This section sets out some essential definitions and concepts used in the rest of the chapter, beginning with the formal definition of the density of a pattern.

Definition 2.1 (Pattern density). Given a system of linear forms $\mathcal{L} = \{L_1, \ldots, L_m\}$ and functions $f_1, \ldots, f_m : \mathbb{F}_p^n \to [-1, 1]$, define the operator

$$\Lambda_{\mathcal{L}}(f_1,\ldots,f_m) = \mathop{\mathbb{E}}_{\underline{x} \in (\mathbb{F}_p^n)^l} f_1(L_1(\underline{x})) \ldots f_m(L_m(\underline{x})).$$

For an r-colouring ϕ , the density of $\mathcal{H} = (\mathcal{L}, \mathcal{X})$ under ϕ is given by

$$\Lambda_{\mathcal{H}}(\phi) = \sum_{\chi \in \mathcal{X}} \Lambda_{\mathcal{L}}(\mathbb{1}_{\phi^{-1}(\chi(1))}, \dots, \mathbb{1}_{\phi^{-1}(\chi(m))}).$$

It is easy to check that $\Lambda_{\mathcal{L}}$ satisfies the telescoping identity

(1)
$$\Lambda_{\mathcal{L}}(f_1,\ldots,f_m) - \Lambda_{\mathcal{L}}(g_1,\ldots,g_m) = \sum_{i=1}^m \Lambda_{\mathcal{L}}(h_1^{(i)},\ldots,h_m^{(i)})$$

where $h_j^{(i)}$ is equal to f_j when j < i, $f_i - g_i$ when j = i, and g_j otherwise.

An integral tool in proving removal lemmas is an arithmetic analogue of Szemerédi's regularity lemma. Given a function $f : \mathbb{F}_p^n \to [0, 1]$, this arithmetic regularity lemma [13] provides a partition of \mathbb{F}_p^n into cosets of a 'large' subspace such that f behaves 'pseudorandomly' on almost all cosets, with the measure of pseudorandomness given by Fourier uniformity. In the following definition, $\mathcal{P}(H)$ denotes the partition of \mathbb{F}_p^n into cosets of a subspace H.

Definition 2.2 (Fourier uniformity). Let H be a subspace of \mathbb{F}_p^n . Given a function $F : \mathbb{F}_p^n \to \mathbb{C}$ and elements $c, r \in \mathbb{F}_p^n$, the Fourier transform of F on H + c at r is defined as

$$\widetilde{F}|_{H+c}(r) = \mathbb{E}_{x \in H+c} F(x) e_p(r^T x),$$

where $e_p(\cdot)$ denotes $\exp(2\pi i \cdot p)$.

A function $f : \mathbb{F}_p^n \to \mathbb{C}$ is said to be ϵ -uniform on H + c if $|\widehat{F}|_{H+c}(r)| \leq \epsilon$ for all $r \in \mathbb{F}_p^n$, where $F = f - \mathbb{E}_{x \in H+c} f(x)$. When f is ϵ -uniform on H + c, the latter is referred to as an ϵ -regular coset for f. The partition $\mathcal{P}(H)$ is ϵ -regular for f if for all but an ϵ -proportion of $c \in \mathbb{F}_p^n$, H + c is ϵ -regular for f.

The precise statement of the arithmetic regularity lemma, due to Green [13], is then as follows.

Theorem 2.3 (Arithmetic regularity lemma [13]). Fix $\epsilon > 0$, an integer r > 0, and a subspace $H_0 \leq \mathbb{F}_p^n$. There exists $C_{arl} = C_{arl}(\epsilon, r)$ such that for any functions f_1, \ldots, f_r : $\mathbb{F}_p^n \rightarrow [0,1]$, there is a subspace $H \leq H_0$ of codimension at most $C_{arl}(\epsilon, r)$ in H_0 such that $\mathcal{P}(H)$ is ϵ -regular for f_1, \ldots, f_r .

Fourier uniformity allows us to count the number of instances for linear systems of complexity 1. Specifically, the complexity of a linear system here refers to the true complexity of Gowers and Wolf [11], originally defined in terms of Gowers uniformity norms $||f||_{U^{s+1}}$ (see [9]). While the latter will not be defined here, it is a well-known fact that functions with small U^2 -norm are precisely those that have small Fourier transforms: in fact, it is easy to show that $||f||_{U^2}^4 = ||\hat{f}||_4^4$ [10, Lemma 2.4], from which the last part of the definition below follows.

Definition 2.4 (True complexity). Let $\mathcal{L} = \{L_1, \ldots, L_m\}$ be a system of linear forms. The true complexity of \mathcal{L} is the least positive integer s (if it exists) with the following property. For every $\delta > 0$, there is an $\epsilon(\delta) > 0$ such that for any functions $F_1, \ldots, F_m : \mathbb{F}_p^n \to [-1, 1]$ satisfying $\min_i ||F_i||_{U^{s+1}} \leq \epsilon(\delta)$,

$$\left|\Lambda_{\mathcal{L}}(F_1,\ldots,F_m)\right|\leq \delta.$$

In particular, \mathcal{L} has true complexity 1 if for every $\delta > 0$, there is an $\epsilon_{count}(\delta) > 0$ such that for any $F_1, \ldots, F_m : \mathbb{F}_p^n \to [-1, 1]$ with $\min_i \max_{\eta \in \mathbb{F}_p^n} |\widehat{F}_i(\eta)| \leq \epsilon_{count}(\delta)$,

$$\left|\Lambda_{\mathcal{L}}(F_1,\ldots,F_m)\right|\leq \delta.$$

Example 2.5. The linear system $\{x, x + d, x + 2d\}$, corresponding to 3-term arithmetic progressions, has true complexity 1 (for example, see [15, Proposition 1.8]). On the other hand, $\{x, x + d, x + 2d, x + 3d\}$ does not. There is a beautifully simple criterion that may be used to verify this fact. Gowers and Wolf [12, Theorem 6.1] proved that for sufficiently large p, the true complexity of a linear system $\{L_1, \ldots, L_m\}$ over \mathbb{F}_p^n is equal to the smallest s such that the set $\{L_1^{s+1}, \ldots, L_m^{s+1}\}$ is linearly independent. It is easy to check that while x - 2(x+d) + (x+2d) = 0, the set $\{x^2, (x+d)^2, (x+2d)^2\}$ is linearly independent, so the true complexity of the 3-term arithmetic progression is 1. On the other hand, $\{x^2, (x+d)^2, (x+2d)^2, (x+3d)^2\}$ satisfies the linear dependence $x^2 - 3(x+d)^2 + 3(x+2d)^2 - (x+3d)^2 = 0$, meaning that the true complexity of the 4-term arithmetic progression is at least 2.

Together with the telescoping identity (1) and an ϵ -regular partition provided by Theorem 2.3, Definition 2.4 can be used to count instances of a linear system \mathcal{L} that satisfy $L_i(\underline{x}) \in$

 $H + c_i$ for given cosets $H + c_1, \ldots, H + c_m$. In fact, if each $H + c_i$ is regular for f_i , then there are many such instances of \mathcal{L} , so long as the cosets $H + c_1, \ldots, H + c_m$ 'align' in the right way. This is a necessary condition as, for example, there cannot be any instances of a 3-term arithmetic progression x_1, x_2, x_3 with each $x_i \in H + c_i$ unless $H + c_1, H + c_2, H + c_3$ themselves form an arithmetic progression. The following definition arises as a special case of [21, Definition 3.17].

Definition 2.6 (Coset consistency). Let $H \leq \mathbb{F}_p^n$ be a subspace and let $\mathcal{L} = (L_1, \ldots, L_m)$ be a linear system of complexity 1 in l variables. Given $c_1, \ldots, c_m \in \mathbb{F}_p^n$, the cosets $H + c_1, \ldots, H + c_m$ are said to be consistent with \mathcal{L} if there exists $\underline{x} \in (\mathbb{F}_p^n)^l$ such that $L_i(\underline{x}) \in H + c_i$ for each $i \in [m]$.

If the cosets $H + c_1, \ldots, H + c_m$ are in fact consistent with \mathcal{L} , then the expected number of instances of \mathcal{L} satisfying $L_i(\underline{x}) \in H + c_i$ is approximately the same as one would expect in a random setting. This result, known as a counting lemma, appears in various forms in the literature, often stated in the more general setting of higher-order Fourier analysis (see, for instance, [3, Theorem 3.10], [21, Theorem 3.19], and the discussion following [15, Lemma 1.3]) or for linear systems of a particular form (such as [13, Proposition 6.2]). The statement below pertains to general linear systems of complexity 1, with the proof given for completeness in Appendix A.

Lemma 2.7 (Counting Lemma). Fix $\delta > 0$, $\epsilon \leq \epsilon_{count}(\delta)$, and a linear system $\mathcal{L} = (L_1, \ldots, L_m)$ of complexity 1. Let $H \leq \mathbb{F}_p^n$ be a subspace of codimension d. Then for any functions $f_1, \ldots, f_m : \mathbb{F}_p^n \to [-1, 1]$ and any $c_1, \ldots, c_m \in \mathbb{F}_p^n$, the following holds. If the cosets $H + c_1, \ldots, H + c_m$ are consistent with \mathcal{L} and are ϵ -regular for f_1, \ldots, f_m , then

$$\left|\Lambda_{\mathcal{L}}(f_1\mathbb{1}_{H+c_1},\ldots,f_m\mathbb{1}_{H+c_m}) - p^{-d(m-\operatorname{rank}(\mathcal{L}))}\prod_{i=1}^m \alpha_i\right| \le p^{-d(m-\operatorname{rank}(\mathcal{L}))}m\delta$$

where α_i denotes the average of f_i on $H + c_i$.

3. Overview of the argument

It is instructive to consider first the proof a non-induced arithmetic removal lemma. The latter states that if a given set doesn't have too many instances of a linear system \mathcal{L} , then all such instances can be eliminated by removing at most an ϵ -proportion of \mathbb{F}_p^n from the set.

Theorem 3.1 (Arithmetic removal lemma [13]). Fix $\epsilon > 0$ and a linear system \mathcal{L} of complexity 1. There exists $\delta = \delta(\epsilon, \mathcal{L})$ satisfying the following. If $A \subseteq \mathbb{F}_p^n$ is a set such that $\Lambda_{\mathcal{L}}(\mathbb{1}_A) \leq \delta$, then A can be made \mathcal{L} -free on the whole of \mathbb{F}_p^n by removing at most $\epsilon |\mathbb{F}_p^n|$ elements from A.

The proof of Theorem 3.1 begins with an application of the arithmetic regularity lemma (Theorem 2.3) to $\mathbb{1}_A$, which gives a partition of \mathbb{F}_p^n into cosets of a subspace such that all

but an ϵ -proportion of the cosets are regular for $\mathbb{1}_A$. The next step is to remove from A all elements that lie either in a non-regular coset or in a coset on which A has low density. Call the resulting set $A' \subseteq A$.





FIGURE 1. The sets A (left) and A' (right) depicted as hashed areas. Blue squares correspond to regular cosets after an application of the arithmetic regularity lemma.

A crucial property of A' is that if a coset contains at least one element of A', then A must both be Fourier-uniform and have high density on this coset. As a result, if there were a single instance of \mathcal{L} in A', we would be able to deduce that there are *many* instances of \mathcal{L} in A via the counting lemma (Lemma 2.7), as depicted in the figure below.



FIGURE 2. An instance of \mathcal{L} in A' would imply many instances of \mathcal{L} in A as a consequence of Lemma 2.7.

A set A may equivalently be viewed as a 2-colouring of \mathbb{F}_p^n given by $\mathbb{1}_A$, and the modified set A' as an \mathcal{L} -free recolouring. In this way, Theorem 3.1 is a special case of the induced removal lemma, where the pattern in question is $\mathcal{H} = (\mathcal{L}, \{\chi\})$ for the constant colouring $\chi \equiv 1$. Removing an element from A, then, corresponds to recolouring it with the colour 0. Note that this never creates new instances of \mathcal{H} , which gives an easy way of avoiding non-regular cosets. On the other hand, if we had $\chi(1) = 0$ and $\chi(i) = 1$ everywhere else, the same would no longer hold.

This necessitates a different approach for proving induced removal lemmas. The existing proofs [4, 3, 7, 21] all follow roughly the same strategy, involving two nested partitions rather than one.

- (1) Use a strong version of the arithmetic regularity lemma ([7, Theorem 5.4]) to find subspaces $H_2 \leq H_1$ such that
 - (i) nearly all cosets of H_2 are regular for the given colouring ϕ (that is, for the indicator functions of all of its colour classes);
 - (ii) the colour densities on almost all cosets of H_2 are a good approximation for the densities on the corresponding coset of H_1 .
- (2) Inside each coset of H_1 , select a coset of H_2 to act as its representative in such a way that
 - (i) each of the chosen subcosets is regular for ϕ ;
 - (ii) the colour densities on (almost all) chosen subcosets approximate the densities on the cosets they are representing;
 - (iii) the chosen subcosets preserve consistency with the given pattern, i.e. if a collection of cosets of H_1 is consistent with the pattern, their subcoset representatives should be as well.

The resulting collection of subcosets can be thought of as a 'regular model' for H_1 (see [7, Section 2]): while it is impossible to guarantee that every coset of H_1 is regular for ϕ [14], we can pass to the chosen regular subcosets without losing too much information.

(3) Define a recolouring ψ as follows. For each coset of H_1 , its subcoset representative determines whether a colour is 'removed'. Specifically, if a colour occurs with low density on the subcoset, then it is replaced with some high-density colour on the whole of the coset. This recolours only a small proportion of all elements since the colour densities on the subcoset are approximately the same as on the coset it is representing.

The resulting colouring ψ has the property that if a colour occurs at least *once* in a coset of H_1 under ψ , then it occurs with *high density* on its subcoset representative under the original colouring ϕ . Now, if there is an instance of the pattern under ψ , we can obtain a contradiction as in Figure 2 using the counting lemma on the subcoset representatives. This works since the representatives were chosen to preserve consistency (see Step 2).

In order to make sure that the chosen subcosets preserve consistency, Fox, Tidor and Zhao [7] require that these subcosets themselves form a subspace. Of course, this leaves no choice for the subcoset representing the zero coset H_1 , as it must be H_2 itself. This presents an issue since it is impossible to guarantee that H_2 is regular [14]. As a result, their argument



*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*

*
</tr

(A) Subcoset representatives outlined in orange. Blue subcosets are regular for ϕ , and the density is well-approximated on subcosets with a star.

(B) Each non-zero coset is recoloured according to its chosen subcoset; the zero coset is handled separately with a Ramsey-type argument.

FIGURE 3. The strategy of Fox, Tidor, and Zhao [7] for the induced arithmetic removal of complexity-1 patterns, depicted for a 2-colouring. Instances of the pattern which contain 0 may remain.

requires the zero coset to be treated separately from the rest of the space with a Ramseytheoretic patching argument (Figure 3b). It is this last step that leads to a small number of pattern instances remaining after the recolouring in Theorem 1.2.

This obstacle does not arise in the translation-invariant setting [3], since the subspace of subcosets can be translated as desired without breaking consistency with the pattern. This added flexibility allows Bhattacharyya et al. [3] to ensure that all chosen subcosets are regular.

The main contribution of this paper is a modified subcoset selection strategy which allows to extend similar flexibility to the wider class of partition-regular patterns. The idea is to choose multiple subcosets inside each coset of H_1 , so that the recolouring of each coset is governed by a group of representatives. In fact, this is reminiscent of the proof of the induced graph removal lemma by Alon, Fischer, Krivelevich and Szegedy (see the use of [1, Corollary 3.4] in the proof of [1, Theorem 5.1]). The technical details of this modification and the proof of Theorem 1.5 are contained in Section 5.

4. LINEAR SYSTEMS AND PARTITION REGULARITY

Recall the well-known fact due to Rado [18] (extended to general abelian groups by Deuber [6]) that a matrix A is partition-regular if and only if A satisfies Rado's column conditions.

Definition 4.1 (Rado's column conditions). Let A be an $l \times m$ matrix over \mathbb{F}_p . Then A is said satisfy Rado's column conditions if there are integers $0 < k_1 < \ldots < k_t = m$ and an ordering of the column vectors $\underline{c}_1, \ldots, \underline{c}_m$ such that

(1)
$$\sum_{i=1}^{k_1} \underline{c}_i = 0;$$

(2) for each $1 < i \leq t$, $\sum_{j=k_{i-1}+1}^{k_i} \underline{c}_j$ is in the span of $\underline{c}_1, \ldots, \underline{c}_{k_{i-1}}$.

As outlined at the start of the chapter, the proof of Theorem 1.5 relies on being able to choose subcosets with certain nice properties. Partition regularity facilitates this because we can define an auxiliary colouring on the set of all subcosets in a way that encodes these properties, and know that there are many monochromatic configurations of subcosets that are consistent with a given system \mathcal{L} (see Lemma 5.3 for details). Specifically, we will be using the following quantitative form of Rado's theorem over finite fields due to Serra and Vena [19, Theorem 2.1].

Theorem 4.2 (Rado's theorem over finite fields [19]). Fix integers r, n > 0, and let A be an $l \times m$ matrix over \mathbb{F}_p that satisfies Rado's column conditions. Then there exist constants $c_{rado} = c_{rado}(p, r, m)$ and $n_{rado} = n_{rado}(p, r, m)$ such that for all $n \ge n_{rado}$ and for every r-colouring of \mathbb{F}_p^n , the system $A\underline{x} = 0$ has at least $c_{rado} |\mathbb{F}_p^n|^{m-l}$ monochromatic solutions.

Of course, in order to apply Theorem 4.2 to arithmetic patterns, we need a way of turning a collection of linear forms into matrices. Such a translation is standard and outlined in this section for completeness.

Let $\mathcal{L} = \{L_1, \ldots, L_m\}$ be a system of linear forms in l variables over \mathbb{F}_p so that

$$L_i(\underline{x}) = c_1^{(i)} x_1 + \ldots + c_l^{(i)} x_l.$$

Define an $m \times l$ matrix $\mathcal{M}(\mathcal{L}) = (c_j^{(i)})$, where the *i*th row contains the coefficients of L_i . Then Im(\mathcal{M}) precisely corresponds to all values that the linear system \mathcal{L} can take.

Moreover, it is easy to see that the linear dependencies between L_1, \ldots, L_m are given by vectors $\underline{y} \in \mathbb{F}_p^m$ satisfying $\underline{y}\mathcal{M} = 0$, or, equivalently, $\underline{y} \in \ker(\mathcal{M}^T)$. On the other hand, these dependencies are precisely the vectors in $\operatorname{Im}(\mathcal{M})^{\perp}$, as the following lemma shows.

Lemma 4.3. Let A be any matrix. Then $\operatorname{Im}(A)^{\perp} = \ker(A^T)$.

Proof. Take $v \in \text{Im}(A)^{\perp}$. Then for all x, $v^T A x = 0$ and so $v^T A = 0$. Hence $A^T v = 0$, i.e. $v \in \text{ker}(A^T)$. For the other direction, take $v \in \text{ker}(A^T)$. Then for all $x \in \mathbb{F}_p^n$, $v^T A x = (A^T v)^T x = 0$. Hence $v \in \text{Im}(A)^{\perp}$.

Now let $\{v_1, \ldots, v_l\}$ be a basis of ker (\mathcal{M}^T) and define $\mathcal{K}(\mathcal{L})$ to be the matrix with rows given by $\{v_1^T, \ldots, v_l^T\}$. While the definition of $\mathcal{K}(\mathcal{L})$ depends on the choice of $\{v_1, \ldots, v_l\}$, the properties we are interested in are invariant under a change of basis, so this choice is of no import.

Proposition 4.4. The solutions to $\mathcal{K}(\mathcal{L})\underline{z} = 0$ are precisely the values taken by \mathcal{L} as \underline{x} ranges over \mathbb{F}_p^l . That is, for each such \underline{z} there is an $\underline{x} \in \mathbb{F}_p^l$ such that

$$\underline{z} = (L_1(\underline{x}), \dots, L_m(\underline{x})).$$

Proof. The definition of $\mathcal{K}(\mathcal{L})$ implies that $\operatorname{Im}(\mathcal{K}^T) = \ker(\mathcal{M}^T)$. Then, by Lemma 4.3, $\ker(\mathcal{K}) = \operatorname{Im}(\mathcal{K}^T)^{\perp} = \ker(\mathcal{M}^T)^{\perp} = (\operatorname{Im}(\mathcal{M})^{\perp})^{\perp} = \operatorname{Im}(\mathcal{M})$, which implies the result. \Box

In line with Proposition 4.4, we will say that a linear system \mathcal{L} is partition-regular if and only if $\mathcal{K}(\mathcal{L})$ satisfies Rado's column conditions.

5. Proof of Theorem 1.5

This section contains the core of the proof and the main original contribution of the chapter. Whereas previous methods [7], [3] depended on choosing a single suitable subcoset within each coset of the coarser partition (Figure 3a), the proof of Theorem 1.5 requires an additional refining step which allows us to choose *several* subcosets at once. In fact, we will use the subcoset selection of [7] (Figure 4a) as an intermediate stage, then find the final subcosets inside each of the intermediate ones (Figure 4b). The following is [7, Proposition 3.2].

Proposition 5.1 (Intermediate subcoset selection). Fix $\epsilon, \zeta > 0$ and integers r, d > 0. There exist $n_{reg} = n(\epsilon, \zeta, r, d)$ and $C_{reg}(\epsilon, \zeta, r, d)$ such that for any $n > n_{reg}$ the following holds. Given a subspace $H_0 \leq \mathbb{F}_p^n$ of codimension d and functions $f_1, \ldots, f_r : \mathbb{F}_p^n \to [0, 1]$, there are subspaces $H_2 \leq H_1 \leq H_0$ of codimensions D_2 and D_1 respectively, as well as a choice of complement U satisfying $U \oplus H_1 = \mathbb{F}_p^n$ such that

- (i) $D_1 \leq D_2 \leq C_{reg}(\epsilon, \zeta, r, N, d);$
- (ii) for all $u \in U \setminus \{0\}$, $H_2 + u$ is ϵ -regular for f_1, \ldots, f_r ;
- (iii) for all but at most a ζ -proportion of $u \in U$ and all $i \in [r]$,

$$\left| \mathop{\mathbb{E}}_{x \in H_2 + u} f_i(x) - \mathop{\mathbb{E}}_{x \in H_1 + u} f_i(x) \right| < \zeta.$$

In Figure 4a below, the chosen cosets are those given by H + u for $u \in U$, with the regular cosets coloured in blue. The star inside a coset is used to depict property (iii) of Proposition 5.1.





(A) A selection of cosets of H_2 inside those of H_1 , for $H_2 \leq H_1$, as produced in [7].

(B) A selection of cosets of H_3 inside previously chosen cosets of H_2 , for $H_3 \leq H_2$.

It is not hard to show that the property of being ϵ -regular is inherited by subcosets and, moreover, provides us with a way of inheriting property (iii) as well.

Proposition 5.2 (Regularity properties). Fix $\epsilon > 0$. Let $f : \mathbb{F}_p^n \to [0,1]$, and let $H_2 \leq H_1$ be subspaces of \mathbb{F}_p^n with $\operatorname{codim}_{H_1}H_2 = d$. Suppose that a coset C_1 of H_1 is ϵ -regular for f. Then for any coset C_2 of H_2 such that $C_2 \subseteq C_1$,

- (i) C_2 is $p^d \epsilon$ -regular for f;
- (*ii*) $\left| \mathbb{E}_{x \in C_2} f(x) \mathbb{E}_{x \in C_1} f(x) \right| \le p^d \epsilon.$

Proof. Write $\alpha_i = \mathbb{E}_{x \in C_i} f(x)$ and $C_i = H_i + z_i$ for i = 1, 2, where $z_1, z_2 \in \mathbb{F}_p^n$. Since C_1 is ϵ -regular for f, we know that for any $r \in \mathbb{F}_p^n$,

(2)
$$\left| \underset{x \in C_1}{\mathbb{E}} (f(x) - \alpha_1) e_p(r^T x) \right| \le \epsilon$$

As $\operatorname{codim}_{H_1}H_2 = d$, there exist some $h_1, \ldots, h_d \in H_1$ such that $H_2 = H_1 \cap \langle h_1, \ldots, h_d \rangle^{\perp}$. The indicator function $\mathbb{1}_{H_2}$ can then be written as

(3)
$$\mathbb{1}_{H_2}(x) = \mathbb{1}_{H_1}(x) \prod_{i=1}^d \mathbb{E}_{a_i \in \mathbb{F}_p} e_p(a_i h_i^T x) = \mathbb{1}_{H_1}(x) \mathbb{E}_{a \in \mathbb{F}_p^d} e_p(R_a^T x).$$

where $R_a = \sum_{i=1}^d a_i h_i$. Now fix some $r \in \mathbb{F}_p^n$. If $r \in H_2^{\perp}$, then $\widehat{f|_{C_2}}(r) = 0$ trivially. Otherwise,

$$\widehat{f|_{C_2}}(r) = \frac{|C_1|}{|C_2|} \mathop{\mathbb{E}}_{x \in C_1} (f(x) - \alpha_2) \mathbb{1}_{H_2} (x - z_2) e_p(r^T x) = p^d \mathop{\mathbb{E}}_{a \in \mathbb{F}_p^d} e_p(-R_a^T z_2) \Big[\mathop{\mathbb{E}}_{x \in C_1} (f(x) - \alpha_2) e_p((r + R_a)^T x) \Big].$$

Then using (2) and the triangle inequality gives

$$\left|\widehat{f}|_{C_2}(r)\right| \le p^d \epsilon + \left| \underset{x \in C_1}{\mathbb{E}} (\alpha_1 - \alpha_2) e_p((r + R_a)^T x) \right|.$$

But the last term in this equation is 0 since $(r + R_a)^T x \neq 0$ on C_1 when $r \notin H_2^{\perp}$. This is because $H_1^{\perp} \leq H_2^{\perp}$ so $r \notin H_1^{\perp}$ and $R_a \in H_1$ by definition, which implies $r + R_a \notin H_1^{\perp}$. Hence part (i) of the proposition holds.

Part (ii) is proved similarly. From equation (3),

$$\mathop{\mathbb{E}}_{x \in C_2} f(x) - \alpha_1 = p^d \mathop{\mathbb{E}}_{a \in \mathbb{F}_p^d} e_p(-R_a^T z_2) \Big[\mathop{\mathbb{E}}_{x \in C_1} (f(x) - \alpha_1) e_p(R_a^T x) \Big],$$

which implies, by (2) and the triangle inequality, that $|\alpha_2 - \alpha_1| \leq p^d \epsilon$, as required.

Thanks to Proposition 5.2, if $\operatorname{codim}_{H_2}H_3$ is sufficiently small, any cosets of H_3 picked inside the regular cosets of H_2 (as in Figure 4b) will still be regular, and f will still have approximately the same density on them as on the coset of H_1 containing them. Since non-zero cosets of H_2 can be made regular by Proposition 5.1, it is enough to make a 'good' selection of subcosets inside the zero coset of H_2 . This is the purpose of the following lemma.

Lemma 5.3 (Selecting multiple subcosets). Fix $\epsilon, \delta > 0$, an integer r > 0, and an $l \times m$ partition-regular matrix A over \mathbb{F}_p . There exist $n_{zreg} = n(\epsilon, \delta, r, m)$ and $C_{zreg}(\epsilon, \delta, r, m)$ such

that for any $n > n_{zreg}$ the following holds. Given functions $f_1, \ldots, f_r : \mathbb{F}_p^n \to [0, 1]$, there is a subspace $H \leq \mathbb{F}_p^n$ of codimension $D \leq C_{zreg}$, as well as $z_1, \ldots, z_m \in \mathbb{F}_p^n$ such that

- (*i*) $A(z_1, \ldots, z_m)^T = 0;$
- (ii) for all $j \in [m]$, $H + z_j$ is ϵ -regular for f_1, \ldots, f_r ;
- (iii) for each $i \in [r]$, $\mathbb{E}_{x \in H+z_j} f_i(x) < \delta$ for all $j \in [m]$ or $\mathbb{E}_{x \in H+z_j} f_i(x) \ge \delta$ for all $j \in [m]$.

Proof. Let H_0 be an arbitrary subspace of codimension $n_{rado}(p, 2^r, m)$ and define $\epsilon' = \min(\epsilon, c_{rado}(p, 2^r, m)/2)$, where n_{rado} and c_{rado} are as in Theorem 4.2. Apply the arithmetic regularity lemma (Theorem 2.3) to f_1, \ldots, f_r inside H_0 with parameter ϵ' to obtain a subspace $H \leq H_0$ such that

- *H* has codimension *D* satisfying $n_{rado}(p, 2^r, m) \leq D \leq C_{arl}(\epsilon', r) + n_{rado}(p, 2^r, m);$
- all but an ϵ' -proportion of the cosets of H are ϵ -regular for f_1, \ldots, f_r .

Let U be such that $U \oplus H = \mathbb{F}_p^n$, and define a colouring $\psi : U \to \mathcal{P}([r])$ by setting $\psi(u) = \{i \in [r] : \mathbb{E}_{x \in H+u} f_i(x) \geq \delta\}$, so that $\psi(u)$ corresponds to the set of those f_i that have high density on H + u. Apply Theorem 4.2 to ψ in U (which is isomorphic to \mathbb{F}_p^D) to find at least $c_{rado}(p, 2^r, m) |\mathbb{F}_p^D|^{m-l}$ monochromatic solutions to $A\underline{z} = 0$ under ψ . Note that by definition of ψ , any such solution satisfies property (iii) of the lemma.

(b)	(r ,b,g)	(r,g)	(r,g)	(b)	(r,b,g)	(r,g)	(r,g)
(b)	()	()	(r,g)	(b)	()	()	(r,g)
(r,g)	(r,g)	(r,b,g)	(b)	(r,g)	(r,g)	(r,b,g)	(b)
()	(r,b,g)	(r)	(b)	()	(r,b,g)	(r)	(b)

FIGURE 5. An auxiliary colouring ψ arising in the proof of Lemma 5.3 from a 3-colouring $\phi : \mathbb{F}_p^n \to \{r, b, g\}$. Due to a large proportion of regular cosets (shown in blue), we can find a monochromatic solution consisting entirely of regular cosets.

On the other hand, the number of solutions for which property (ii) does not hold is at most $\epsilon' |\mathbb{F}_p^D|^{m-l} \leq \frac{1}{2} c_{rado}(p, 2^r, m) |\mathbb{F}_p^D|^{m-l}$, which is strictly less than the total number of monochromatic solutions. As such, there is at least one solution (z_1, \ldots, z_m) that satisfies all properties.

All results stated so far have referred to 1-bounded functions whereas our aim is to say something about an *r*-colouring $\phi : \mathbb{F}_p^n \to [r]$. Of course, it is easy to turn ϕ into a collection of *r* functions by letting $\phi_i : \mathbb{F}_p^n \to [0, 1]$ be the indicator function of colour *i*, i.e. $\phi_i = \mathbb{1}_{\phi^{-1}(i)}$.

We can then combine the other results in this section to define an appropriate recolouring for Theorem 1.5.

Lemma 5.4 (Recolouring). Fix $\epsilon > 0$, an integer r > 0, and an $l \times m$ partition-regular matrix A over \mathbb{F}_p . There exist $n_{rcl} = n_{rcl}(\epsilon, r, m)$ and $C_{rcl} = C_{rcl}(\epsilon, r, m)$ such that for any $n > n_{rcl}$ the following holds. Given an r-colouring $\phi : \mathbb{F}_p^n \to [r]$, there exist subspaces $H_3 \leq H_2 \leq H_1$ of codimensions D_3, D_2, D_1 respectively, a choice of complement U satisfying $U \oplus H_1 = \mathbb{F}_p^n$, and $z_1, \ldots, z_m \in H_2$ such that

- (i) $\log_p(\frac{\epsilon}{4r}) \leq D_1, D_2, D_3 \leq C_{rcl};$
- (*ii*) $A(z_1, \ldots, z_m)^T = 0;$
- (iii) for all $u \in U$ and $j \in [m]$, $H_3 + u + z_j$ is ϵ -regular for ϕ_1, \ldots, ϕ_r .

Additionally there is an r-colouring $\phi' : \mathbb{F}_p^n \to [r]$ such that ϕ' differs from ϕ in at most $\epsilon |\mathbb{F}_p^n|$ places and satisfies the following property.

(iv) For any $u \in U$, if $\phi'(y) = i$ for some $y \in H_1 + u$, then $\mathbb{E}_{x \in H_3 + u + z_j} \phi_i(x) \ge \epsilon/8r$ for all $j \in [m]$.

Proof. Define $D_0 = \lceil \log_p(\epsilon/4r) \rceil$, $C_{zreg} = C_{zreg}(\epsilon, \epsilon/8r, m)$, $\epsilon' = \epsilon p^{-C_{zreg}}/8r$ and $\zeta = \epsilon/4r$, where C_{zreg} is as in Lemma 5.3. Set $C_{rcl} = C_{reg}(\epsilon', \zeta, D_0) + C_{zreg}$ and

$$n_{rcl} = \max\left(n_{reg}(\epsilon', \zeta, D_0), n_{zreg}(\epsilon, \zeta, m) + C_{reg}(\epsilon', \zeta, D_0)\right),$$

with n_{reg} and C_{reg} as in Proposition 5.1. Let H_0 be an arbitrary subspace of codimension D_0 , and apply Proposition 5.1 to ϕ_1, \ldots, ϕ_r and H_0 with parameters ϵ' and ζ . The result is subspaces $H_2 \leq H_1 \leq H_0$ of codimensions D_2 and D_1 respectively, as well as a complement U satisfying $U \oplus H_1 = \mathbb{F}_n^n$ such that the following holds:

- (a) $\log_p(\epsilon/4r) \leq D_1, D_2 \leq C_{reg}(\epsilon', \zeta, D_0);$
- (b) for all $u \in U \setminus \{0\}$, $H_2 + u$ is ϵ' -regular for ϕ_1, \ldots, ϕ_r ;
- (c) for all but at most a ζ -proportion of $u \in U$ and all $i \in [r]$,

$$\mathbb{E}_{x\in H_2+u}\phi_i(x) - \mathbb{E}_{x\in H_1+u}\phi_i(x)\big| < \zeta.$$

Now apply Lemma 5.3 to H_2 with parameters ϵ and $\delta = \epsilon/8r$ to obtain a subspace $H_3 \leq H_2$ of codimension $D_3 = C_{zreg} + D_2 \leq C_{rcl}$ in \mathbb{F}_p^n , as well as $z_1, \ldots, z_m \in H_2$ such that

- (d) $A(z_1, \ldots, z_m)^T = 0;$
- (e) for all $j \in [m]$, $H_3 + z_j$ is ϵ -regular for ϕ_1, \ldots, ϕ_r ;
- (f) for all $i \in [r]$, either all $j \in [m]$ satisfy $\mathbb{E}_{x \in H_3 + z_j} \phi_i(x) < \delta$ or all $j \in [m]$ satisfy $\mathbb{E}_{x \in H_3 + z_j} \phi_i(x) \ge \delta$.

Property (b) and Proposition 5.2 imply that for all $u \in U \setminus \{0\}$ and $j \in [m]$, the coset $H_3 + u + z_j$ is $\epsilon' p^{C_{zreg}}$ -regular for ϕ_1, \ldots, ϕ_r . By the choice of parameters, $\epsilon' p^{C_{zreg}} \leq \epsilon/8r$ which together with (e) gives part (iii) of the lemma. All that remains is to define a suitable recolouring ϕ' .

Fix $u \in U$. If $u \neq 0$, let c be a colour with density at least 1/r on $H_2 + u$.

- If $|\mathbb{E}_{x \in H_2+u} \phi_i(x) \mathbb{E}_{x \in H_1+u} \phi_i(x)| > \zeta$ for some $i \in [r]$, recolour all of $H_1 + u$ with c; note that this happens for at most a ζ -proportion of all possible $u \in U \setminus \{0\}$.
- Otherwise, for each colour $i \in [r]$ with density less than $\epsilon/4r$ on $H_2 + u$, recolour all occurrences of i in $H_1 + u$ with c; note that this changes at most a $\epsilon/4r + \zeta \leq \epsilon/2r$ proportion of $H_1 + u$.

These steps affect at most a $\zeta + r\epsilon/2r \leq 3\epsilon/4$ proportion of the whole space.

For u = 0, let c be a colour that has density at least 1/r on $H_3 + z_1$. Then by property (f), c has density at least $\delta = \epsilon/8r$ on each $H_3 + z_j$. Recolour all of H_1 with c and note that this means ϕ' satisfies property (iv) when u = 0.

In total, the recolouring ϕ' defined by following the steps above differs from ϕ in at most $(3\epsilon/4 + p^{-D_1})|\mathbb{F}_p^n| \leq \epsilon |\mathbb{F}_p^n|$ places.

To show that property (iv) holds for all u, fix $u \in U \setminus \{0\}$ and some $y \in H_1+u$. By definition of ϕ' , if $\phi'(y) = i$, then $\mathbb{E}_{x \in H_2+u} \phi_i(x) \ge \epsilon/4r$. On the other hand, $H_3 + u + z_j \subseteq H_2 + u$ for any $j \in [m]$, and $H_2 + u$ is ϵ' -uniform for ϕ_i ; then Proposition 5.2 implies

$$\mathop{\mathbb{E}}_{x \in H_3 + u + z_j} \phi_i(x) \ge \mathop{\mathbb{E}}_{x \in H_2 + u} \phi_i(x) - \epsilon' p^{C_{zreg}} \ge \frac{\epsilon}{4r} - \frac{\epsilon}{8r} = \frac{\epsilon}{8r}$$

as required. Together with the earlier observation that property (iv) holds for u = 0, this completes the proof.

Lemma 5.4 contains all of the ingredients necessary to prove Theorem 1.5, restated as Theorem 5.5 below. To complete the proof, we must show that the recolouring provided by Lemma 5.4 has no instances of the given partition-regular pattern.

Theorem 5.5. Fix $\epsilon > 0$, and let $\mathcal{H} = (\mathcal{L}, \mathcal{X})$ be a partition-regular pattern of complexity 1 such that \mathcal{L} consists of m linear forms. There exists a $\delta = \delta(\epsilon, m)$ with the following property. If $\phi : \mathbb{F}_p^n \to [r]$ is an r-colouring of \mathbb{F}_p^n such that $\Lambda_{\mathcal{H}}(\phi) \leq \delta$, then there is a recolouring $\phi' : \mathbb{F}_p^n \to [r]$ that differs from ϕ in at most $\epsilon |\mathbb{F}_p^n|$ places such that $\Lambda_{\mathcal{H}}(\phi') = 0$.

Proof. Let $A = \mathcal{K}(\mathcal{L})$ be an $l \times m$ partition-regular matrix so that $\underline{y} \in (\mathbb{F}_p^n)^m$ is an instance of \mathcal{L} if and only if $A\underline{y} = 0$. Set $\delta' = (\epsilon/8r)^m/2m$ and $\epsilon' = \min(\epsilon, \epsilon_{count}(\delta'))$, where $\epsilon_{count}(\delta')$ is as in Definition 2.4. Apply Lemma 5.4 with ϵ' to obtain ϕ' , three subspaces $H_3 \leq H_2 \leq H_1$ of codimensions $D_3, D_2, D_1 \leq C_{rcl}(\epsilon', r, m)$, a complement U such that $U \oplus H_1 = \mathbb{F}_p^n$, and $z_1, \ldots, z_m \in H_2$ satisfying $A\underline{z} = 0$.

Suppose there exists an instance of \mathcal{H} under ϕ' , i.e. $y_1, \ldots, y_m \in \mathbb{F}_p^n$ such that $A(y_1, \ldots, y_m)^T = 0$ and $\chi \in \mathcal{X}$ such that $\phi'(y_i) = \chi(i)$ for all $i \in [m]$. Let $u_1, \ldots, u_m \in U$ be such that $y_i \in H_1 + u_i$.

Claim. The cosets $H_3 + u_1 + z_1, \ldots, H_3 + u_m + z_m$ are consistent with \mathcal{L} .

Proof of Claim. For each *i*, write $y_i = h_i + u_i$ for some $h_1, \ldots, h_m \in H_1$. We know that $A(y_1, \ldots, y_m)^T = 0$ so

$$A(u_1,\ldots,u_m)^T = -A(h_1,\ldots,h_m)^T.$$

Since H_1 and U are subspaces, $A\underline{h} \in H_1$ and $A\underline{u} \in U$, meaning that $A\underline{u} \in H_1 \cap U = \{0\}$. Therefore (u_1, \ldots, u_m) is itself a solution to A. Since (z_1, \ldots, z_m) is also a solution, so is $(u_1 + z_1, \ldots, u_m + z_m)$, which completes the proof of the claim.

As a shorthand, write B_i for the coset $H_3 + u_i + z_i$. By Lemma 5.4(iii), $\phi_{\chi(i)}$ is ϵ' -uniform on B_i for all $i \in [m]$, and B_1, \ldots, B_m are consistent with \mathcal{L} by the claim. Then the counting lemma (Lemma 2.7) applies to give

$$\left|\Lambda_{\mathcal{H}}(\phi_{\chi_1}\mathbb{1}_{B_1},\ldots,\phi_{\chi_m}\mathbb{1}_{B_m})-p^{-D_3(m-\operatorname{\mathbf{rank}}(\mathcal{L}))}\prod_{i=1}^m\alpha_i\right|\leq p^{-D_3(m-\operatorname{\mathbf{rank}}(\mathcal{L}))}m\delta',$$

where $\alpha_i = \mathbb{E}_{x \in B_i} \phi_{\chi(i)}(x) \ge \epsilon/8r$ by Lemma 5.4(iv).

Finally, note that $\Lambda_{\mathcal{H}}(\phi) \geq \Lambda_{\mathcal{H}}(\phi_{\chi_1} \mathbb{1}_{B_1}, \ldots, \phi_{\chi_m} \mathbb{1}_{B_m})$, so that, in particular, the density of \mathcal{H} under ϕ is at least

$$\Lambda_{\mathcal{H}}(\phi_{\chi_{1}}\mathbb{1}_{B_{1}},\ldots,\phi_{\chi_{m}}\mathbb{1}_{B_{m}}) \geq p^{-D_{3}(m-\operatorname{rank}(\mathcal{L}))} \left(\prod_{i=1}^{m} \alpha_{i} - m\delta'\right)$$
$$\geq p^{-mC_{rcl}(\epsilon',r,m)} \left((\epsilon/8r)^{m} - \frac{1}{2}(\epsilon/8r)^{m}\right)$$
$$\geq \frac{1}{2} \left(8rp^{C_{rcl}(\epsilon',r,m)}\epsilon^{-1}\right)^{-m},$$

which contradicts the assumption of the theorem when $\delta < \left(8rp^{C_{rcl}(\epsilon',r,m)}\epsilon^{-1}\right)^{-m}/2$.

6. DISCUSSION

The largest contribution to the bound on δ in Theorem 5.5 is $C_{rcl}(\epsilon', r, m)$ arising from the use of Lemma 5.4. In turn, the proof of Lemma 5.4 shows that $C_{rcl}(\epsilon', r, m)$ has the form $C_{rcl} = C_{reg}(\epsilon' p^{-C_{zreg}}/8r, \epsilon'/4r, \lceil \log_p(\epsilon/4r) \rceil) + C_{zreg}$, where $C_{zreg} = C_{zreg}(\epsilon', \epsilon'/4r, m)$ comes from Lemma 5.3 and C_{reg} from Proposition 5.1.

Lemma 5.3 follows from a single application of the arithmetic regularity lemma (Theorem 2.3) and thus has the same growth in ϵ' as C_{arl} in Theorem 2.3. This is known to be of tower-type in general, with $\mathbf{twr}(n)$ defined recursively by $\mathbf{twr}(1) = 2$, $\mathbf{twr}(n) = 2^{\mathbf{twr}(n-1)}$. The lower bound of tower-type on $C_{arl}(\epsilon, 1)$ was first proved by Green in [13] as $\mathbf{twr}(\lceil \log \epsilon^{-1} \rceil)$ and later strengthened by Hosseini et al. in [16] to $\mathbf{twr}(\lceil \epsilon^{-1} \rceil)$. As such, C_{zreg} is $\mathbf{twr}(\lceil 1/\epsilon' \rceil)$ at best.

Proposition 5.1 follows from an iterated version of the arithmetic regularity lemma, known as the strong arithmetic regularity lemma ([7, Theorem 5.4]). As a consequence of the iteration, one might expect wowzer-type growth in its bounds, where wowzer is an iterated tower defined by $\mathbf{wwz}(1) = 2$, $\mathbf{wwz}(n) = \mathbf{twr}(\mathbf{wwz}(n-1))$. Indeed, upcoming work of the author [8] confirms that wowzer-type growth in the strong arithmetic regularity lemma is unavoidable. This leads to wowzer-type bounds on C_{reg} when the strong arithmetic regularity lemma is used.

However, Fox, Tidor and Zhao found an alternative proof of Proposition 5.1 that bypasses the strong regularity lemma, resulting in better, tower-type bounds [7, Section 5.2]. As a result, $C_{reg}(\epsilon' p^{-C_{zreg}}/8r, \epsilon/4r, \lceil \log_p(\epsilon/4r) \rceil)$ grows like a double tower in $1/\epsilon'$.

Finally, ϵ' was chosen to be at most $\epsilon_{count}(O(\epsilon^m))$ where ϵ_{count} is as in the definition of true complexity (Definition 2.4). In recent work, Manners [17, Theorem 1.1.5] showed that $\epsilon_{count}(\delta)$ can be taken as polynomial in δ .

Theorem 6.1 (Polynomial bounds for true complexity [17]). Let \mathcal{L} be a linear system of finite complexity. Then for all $\delta > 0$, $\epsilon_{2.4}(\delta)$ and $\epsilon_{count}(\delta)$ in Definition 2.4 may be taken as $\delta^{O_{\mathcal{L}}(1)}$.

As a result, $\epsilon' = \epsilon^{O(1)}$, meaning that C_{rcl} , and therefore δ , has growth of the order of $\mathbf{twr}(\mathbf{twr}(\epsilon^{-O(1)}))$.

It is not hard to modify the proof of Theorem 5.5 in order to remove any finite number of partition-regular patterns $\mathcal{H}_1 = (\mathcal{L}_1, \mathcal{X}_1), \ldots, \mathcal{H}_t = (\mathcal{L}_t, \mathcal{X}_t)$ at once (indeed, the induced removal lemmas of [3] and [7] are both able to remove multiple arithmetic patterns). This can be achieved by defining a single pattern encompassing all of $\mathcal{H}_1, \ldots, \mathcal{H}_t$. Let A_1, \ldots, A_t denote the matrices $A_i = \mathcal{M}(\mathcal{L}_i)$, each of dimensions $l_i \times m_i$. By taking $m = \max_i(m_i)$ and $l = \max_i(l_i)$, as well as adding zero rows or columns where appropriate, we may think of these as having the same dimension $l \times m$ – note that adding zero rows or columns does not affect partition regularity.

Let A^* be a $tl \times tm$ block-diagonal matrix composed of A_1, \ldots, A_t , meaning that A^* has the form

$$A^* = \begin{pmatrix} A_1 & \underline{0} & \dots & \underline{0} \\ \underline{0} & A_2 & \dots & \underline{0} \\ & & \dots & \\ \underline{0} & \underline{0} & \dots & A_t \end{pmatrix}.$$

Observe that A^* inherits Rado's column conditions (Definition 4.1) from A_1, \ldots, A_t via taking unions of the corresponding sets of columns. Thus, A^* is itself partition-regular. Moreover, a solution to $A^*\underline{z} = 0$ is a vector $\underline{z} = (\underline{z}_1, \ldots, \underline{z}_t) \in (\mathbb{F}_p^n)^{mt}$ such that $A_i\underline{z}_i = 0$, i.e. \underline{z}_i is an instance of \mathcal{L}_i .

Additionally, given colourings $\chi_1, \ldots, \chi_t : [m] \to [r]$, let $(\chi_1, \ldots, \chi_t) : [tm] \to [r]$ denote the joint colouring obtained by setting $(\chi_1, \ldots, \chi_t)(a) = \chi_{t'}(j)$ where $1 \le t' \le t$ and $1 \le j \le m$ are the unique integers satisfying a = (t'-1)m + j.

Let $\Psi = \{\psi : [m] \to [r]\}$ be the set of all possible r-colourings of m and define \mathcal{X}^* by

$$\mathcal{X}^* = \{ (\chi_1, \dots, \chi_t) \in \Psi^t : \exists i \text{ s.t. } \chi_i \in \mathcal{X}_i \}.$$

Consider a new pattern $\mathcal{H}^* = (\mathcal{L}^*, \mathcal{X}^*)$. As noted, \mathcal{H}^* is still partition-regular. Moreover, by definition, $\underline{z} = (\underline{z}_1, \ldots, \underline{z}_t) \in (\mathbb{F}_p^n)^{mt}$ is an instance of \mathcal{H}^* if and only if each \underline{z}_i is an instance of \mathcal{L}_i and at least one of \underline{z}_i is an instance of \mathcal{H}_i . Therefore, if $\Lambda(\mathcal{H}_i)(\phi) \leq \delta$ for each i, then $\Lambda(\mathcal{H}^*)(\phi) \leq t\delta$. In particular, Theorem 5.5 applies to \mathcal{H}^* to give a \mathcal{H}^* -recolouring. In fact, such a recolouring must also be \mathcal{H}_i -free for every $i \in [t]$, since for any instance $\underline{z}_i \in (\mathbb{F}_p^n)^m$ of $\mathcal{H}_i, (\underline{0}, \ldots, \underline{z}_i, \ldots, \underline{0})$ with \underline{z}_i in the ith position is an instance of \mathcal{H}^* .

In fact, given this, it is possible to remove an infinite number of partition-regular patterns by reducing to the case of finitely many patterns, as was done, for instance, in [7, Section 6] or [3, Section 5.2].

Finally, it is worth noting that all arguments in Section 5 can be readily translated to apply to patterns of higher complexities, with the exception of a suitable version of Lemma 5.3. In the higher-order setting, the types of partitions one obtains from the arithmetic regularity lemma are the level sets of polynomials (in the complexity 1 case, these polynomials are linear and thus result in cosets of a subspace). In order for such level sets to form a (nearly) equipartition, which is hard to forgo while counting, the polynomials need to be chosen carefully of high enough 'rank' relative to each other (see [15] for an excellent introduction to the quadratic case of this). To make the proof work, one would need a version of Lemma 5.3 that can be applied to the zero level set of the intermediate partition (the equivalent of $\mathcal{P}(H_2)$) in such a way that the polynomials of the new partition (the equivalent of $\mathcal{P}(H_3)$) interact well with the previously-obtained polynomials in terms of rank. However, this is hard to arrange without introducing a circular dependence between the various parameters in Lemma 5.4. It is the belief of the author that Theorem 1.5 holds for partition-regular patterns of all finite complexities (and perhaps infinite complexity, as is the case in the induced removal lemma of Tidor and Zhao [21]), but the proof of this might require a novel way of handling the rank of partition refinements, or a different approach entirely.

References

- N. Alon, E. Fischer, M. Krivelevich, and M. Szegedy. "Efficient testing of large graphs". In: *Combinatorica* 20.4 (2000), pp. 451–476.
- [2] V. Bergelson, W. A. Deuber, and N. Hindman. "Rado's Theorem for finite fields". In: Sets, graphs and numbers. Vol. 60. Colloq. Math. Soc. János Bolyai. János Bolyai Math. Soc., Budapest, 1992, pp. 77–88.
- [3] A. Bhattacharyya, E. Fischer, H. Hatami, P. Hatami, and S. Lovett. "Every locally characterized affine-invariant property is testable". In: *Proceedings of the 2013 ACM* Symposium on Theory of Computing. ACM, New York, 2013, pp. 429–435.
- [4] A. Bhattacharyya, E. Grigorescu, and A. Shapira. "A unified framework for testing linear-invariant properties". In: *Random Structures Algorithms* 46.2 (2015), pp. 232– 260.

REFERENCES

- [5] D. Conlon and J. Fox. "Graph removal lemmas". In: Surveys in combinatorics 2013.
 Vol. 409. London Math. Soc. Lecture Note. Cambridge Univ. Press, 2013, pp. 1–49.
- [6] W. Deuber. "Partition theorems for abelian groups". In: J. Combinatorial Theory Ser. A 19 (1975), pp. 95–108.
- [7] J. Fox, J. Tidor, and Y. Zhao. "Induced arithmetic removal: complexity 1 patterns over finite fields". In: Israel J. Math. 248.1 (2022), pp. 1–38.
- [8] V. Gladkova. "A lower bound for the strong arithmetic regularity lemma". forthcoming.
- [9] W. T. Gowers. "A new proof of Szemerédi's theorem". In: Geom. Funct. Anal. 11.3 (2001), pp. 465–588.
- [10] W. T. Gowers. "Decompositions, approximate structure, transference, and the Hahn-Banach theorem". In: *Bull. Lond. Math. Soc.* 42.4 (2010), pp. 573–606.
- [11] W. T. Gowers and J. Wolf. "The true complexity of a system of linear equations". In: Proc. Lond. Math. Soc. (3) 100.1 (2010), pp. 155–176.
- [12] W. T. Gowers and J. Wolf. "Linear forms and higher-degree uniformity for functions on \mathbb{F}_p^{n} ". In: *Geom. Funct. Anal.* 21.1 (2011), pp. 36–69.
- B. Green. "A Szemerédi-type regularity lemma in abelian groups, with applications". In: Geom. Funct. Anal. 15.2 (2005), pp. 340–376.
- [14] B. Green and T. Sanders. "Fourier uniformity on subspaces". arXiv:1510.08739. 2015.
- B. Green. "Montréal notes on quadratic Fourier analysis". In: Additive combinatorics.
 Vol. 43. CRM Proc. Lecture Notes. Amer. Math. Soc., Providence, RI, 2007, pp. 69–102.
- [16] K. Hosseini, S. Lovett, G. Moshkovitz, and A. Shapira. "An improved lower bound for arithmetic regularity". In: *Math. Proc. Cambridge Philos. Soc.* 161.2 (2016), pp. 193– 197.
- [17] F. Manners. "True complexity and iterated Cauchy–Schwarz". arXiv:2109.05731. 2021.
- [18] R. Rado. "Studien zur Kombinatorik". In: Math. Z. 36.1 (1933), pp. 424–470.
- [19] O. Serra and L. Vena. "On the number of monochromatic solutions of integer linear systems on abelian groups". In: European J. Combin. 35 (2014), pp. 459–473.
- [20] E. Szemerédi. "Regular partitions of graphs". In: Problèmes combinatoires et théorie des graphes. Vol. 260. Colloq. Internat. CNRS. CNRS, 1978, pp. 399–401.
- [21] J. Tidor and Y. Zhao. "Testing linear-invariant properties". In: SIAM J. Comput. 51.4 (2022), pp. 1230–1279.

Appendix A. Proof of the counting Lemma

This appendix details the proof of the counting lemma stated in the introduction. Recall that ϵ_{count} was defined alongside true complexity in Definition 2.4.

Lemma 2.7. Fix $\delta > 0$, $\epsilon \leq \epsilon_{count}(\delta)$, and a linear system $\mathcal{L} = (L_1, \ldots, L_m)$ of complexity 1. Let $H \leq \mathbb{F}_p^n$ be a subspace of codimension d. For any functions $f_1, \ldots, f_m : \mathbb{F}_p^n \to [-1, 1]$

REFERENCES

and any $c_1, \ldots, c_m \in \mathbb{F}_p^n$, if cosets $H + c_1, \ldots, H + c_m$ are consistent with \mathcal{L} and ϵ -regular for f_1, \ldots, f_m , then

$$\left|\Lambda_{\mathcal{L}}(f_1\mathbb{1}_{H+c_1},\ldots,f_m\mathbb{1}_{H+c_m}) - p^{-d(m-\operatorname{rank}(\mathcal{L}))}\prod_{i=1}^m \alpha_i\right| \le p^{-d(m-\operatorname{rank}(\mathcal{L}))}m\delta_{\mathcal{L}}$$

We will first prove a short technical result that allows the density of the instances of \mathcal{L} to be rewritten in terms of the matrix $\mathcal{K}(\mathcal{L})$ (see Section 4).

Proposition A.1. Let $\mathcal{L} = (L_1, \ldots, L_m)$ be a linear system in l variables. If $A = \mathcal{K}(\mathcal{L})$, then for any functions $h_1, \ldots, h_m : \mathbb{F}_p^n \to [-1, 1], \Lambda_{\mathcal{L}}(h_1, \ldots, h_m) = \mathbb{E}_{\underline{z} \in \ker(A)} \prod_{i=1}^m h_i(z_i)$.

Proof. Recall from Section 4 that $M = \mathcal{M}(\mathcal{L})$ is the $m \times l$ matrix containing the coefficients of L_i in the *i*th row, so that for any $\underline{x} \in (\mathbb{F}_p^n)^l$, $M\underline{x} = (L_1(\underline{x}), \ldots, L_m(\underline{x}))$. This means that $\mathrm{Im}(M)$ is precisely the same as the set of instances of \mathcal{L} . In particular,

$$\sum_{\underline{x}\in(\mathbb{F}_p^n)^l}\prod_{i=1}^m h_i(L_i(\underline{x})) = \sum_{\underline{z}\in\mathrm{Im}(M)}|\ker(M)|\prod_{i=1}^m h_i(z_i).$$

Moreover, $\ker(A) = \operatorname{Im}(M)$ by Proposition 4.4 and $|\mathbb{F}_p^n|^l = |\operatorname{Im}(M)| |\ker(M)|$ by standard linear algebra, so

$$\Lambda_{\mathcal{L}}(h_1,\ldots,h_m) = \mathop{\mathbb{E}}_{\underline{x}\in(\mathbb{F}_p^n)^l} \prod_{i=1}^m h_i(L_i(\underline{x})) = \mathop{\mathbb{E}}_{\underline{z}\in\ker(A)} \prod_{i=1}^m h_i(z_i).$$

Proof of Lemma 2.7. Let $A = \mathcal{K}(\mathcal{L})$ and write ker^H(A) for the restriction of ker(A) to H^m . Since $H + c_1, \ldots, H + c_m$ are consistent with \mathcal{L} , we may assume without loss of generality that (c_1, \ldots, c_m) is itself an instance of \mathcal{L} , i.e. $A\underline{c} = 0$. Then for any $h_1, \ldots, h_m : \mathbb{F}_p^n \to [-1, 1]$,

$$\mathbb{E}_{\underline{z}\in\ker(A)}\prod_{i=1}^{m}h_i(z_i)\mathbb{1}_{H+c_i}(z_i) = \mathbb{E}_{\underline{z}\in\ker(A)}\prod_{i=1}^{m}h_i(z_i)\mathbb{1}_H(z_i-c_i) = \mathbb{E}_{\underline{z}\in\ker^H(A)}\prod_{i=1}^{m}h_i(z_i+c_i)$$

In particular, $\Lambda_{\mathcal{L}}(\mathbb{1}_{H+c_1}, \dots, \mathbb{1}_{H+c_m}) = |\ker^H(A)|/|\mathbb{F}_p^n| = p^{-d(m-\operatorname{rank}(\mathcal{L}))}$. Writing $\Lambda_{\mathcal{L}}^H$ for the density of \mathcal{L} -instances in H, we additionally have

(4)
$$\Lambda_{\mathcal{L}}(f_1 \mathbb{1}_{H+c_1}, \dots, f_m \mathbb{1}_{H+c_m}) = p^{-d(m-\operatorname{rank}(\mathcal{L}))} \Lambda_{\mathcal{L}}^H(g_1, \dots, g_m),$$

where $g_i: H \to [-1, 1]$ is the function given by $g_i(x) = f_i(x + c_i)$.

On the other hand, using $\prod_{i=1}^{m} \alpha_i = \Lambda_{\mathcal{L}}^H(\alpha_1, \ldots, \alpha_m)$ in the telescoping identity (1) gives

(5)
$$\Lambda_{\mathcal{L}}^{H}(g_{1},\ldots,g_{m}) - \prod_{i=1}^{m} \alpha_{i} = \sum_{i=1}^{m} \Lambda_{\mathcal{L}}^{H}(h_{1}^{(i)},\ldots,h_{m}^{(i)}),$$

where for each *i*, at least one $1 \leq j \leq m$ satisfies $h_j^{(i)} = g_j - \alpha_j$. Crucially, all g_1, \ldots, g_m are ϵ -uniform on *H* by assumption, so that by Definition 2.4, $\left|\Lambda_{\mathcal{L}}^H(h_1^{(i)}, \ldots, h_m^{(i)})\right| \leq \delta$. As a result, (5) can be bounded by $\left|\Lambda_{\mathcal{L}}^H(g_1, \ldots, g_m) - \prod_{i=1}^m \alpha_i\right| \leq m\delta$, which together with (4) completes the proof.