# Complete background cosmology of parity-even quadratic metric-affine gravity

T. Dyer,<sup>1,2,\*</sup> W. Barker,<sup>3</sup> and D. Iosifidis<sup>4,†</sup>

<sup>1</sup>Astrophysics Group, Cavendish Laboratory, JJ Thomson Avenue, Cambridge CB3 0HE, UK

<sup>2</sup>Kavli Institute for Cosmology, Madingley Road, Cambridge CB3 0HA, UK

<sup>3</sup>Central European Institute for Cosmolgy and Fundamental Physics,

Institute of Physics of the Czech Academy of Sciences, Na Slovance 1999/2, 182 00 Prague 8, Czechia

<sup>4</sup>Laboratory of Theoretical Physics, Institute of Physics,

University of Tartu, W. Ostwaldi 1, 50411 Tartu, Estonia.

The cosmology of metric-affine gravity is studied for the general, parity preserving action quadratic in curvature, torsion and non-metricity. The model contains 27 a priori independent couplings in addition to the Einstein constant. Linear and higher order relations between the quadratic operators in a Friedmann–Lemaître–Robertson–Walker spacetime are obtained, along with the modified Friedmann, torsion and non-metricity equations. Extra parameter constraints lead to two special branches of the model. Firstly, a branch is found in which the Riemannian spatial curvature (thought to be slightly closed or flat in the  $\Lambda$ CDM model of our Universe) is entirely screened from all the field equations, regardless of its true value. Secondly, an integrable branch is found the need to eliminate higher-spin particles as well as ghosts and tachyons motivates further parameter constraints in each case. The most general model is also found which reproduces the exact Friedmann equations of general relativity. The full set of equations describing closed, open or flat cosmologies, for general parity-even quadratic metric-affine gravity, is made available for *SymPy*, *Mathematica* and *Maple* platforms.

### Contents

I	Introduction	1
Π	Theoretical development	2
	A Metric-affine geometry	3
	B Background cosmology	3
	C The Einstein–Hilbert case	4
III	Structure of the field equations	5
	A Reduced couplings for cosmology	5
	B Minisuperspace computation	6
	C Conditions for projective invariance	7
IV	Special models and their particles	8
	A Gravity with K-screening	8
	B Integrable gravity and its Maxwell limit	9
V	Concluding remarks	11
	References	12
A	Quadratic and cubic relations	15
B	The field equations in full	16
С	Particle spectrographs	18

#### I. Introduction

Reformulation of gravity — Metric-affine gravity (MAG) [1-8] is a natural generalisation of Einstein's general relativity (GR) [9]. MAG assumes that the connection, uniquely defined in GR as the Levi-Civita connection, is an independent dynamical variable in addition to the metric. A completely general affine connection endows the spacetime with non-Riemannian geometric properties beyond the Riemannian curvature of GR, namely torsion and non-metricity, see Fig. 1 for an illustration. MAG can be motivated as a natural extension of the theory of elastic continua in three dimensions (which describes deformations beyond curvature) to a four-dimensional spacetime [1, 10-12]. The conjugate source of the affine connection, the hypermomentum, is equivalently the source for torsion and non-metricity [3, 10, 12–14]. In this framework, not only does gravitation represent a 'metrical elasticity' of space [15], but it also possesses torsional and non-metric degrees of freedom (d.o.f) generated by the microstructure of matter, for which GR has no equivalent. Since its inception [16], MAG has had many applications and has received interest in a range of areas (see reviews [1, 8]). In particular today, the phenomenology of torsion and non-metricity couplings in the standard model of particle physics is an area of active focus [17–19].

**Motivation for replacing GR** — While GR is an extremely successful physical theory of gravitation and has passed many experimental and observational tests to high precision, in isolation it fails to account for certain phenomena. Chief among these are the strongly accelerated expansion of the early Universe [20, 21], and the enhancement of gravity across a range of astrophysical and cosmological scales [22]. Accordingly, a separate inflationary mechanism is sought after, along with a feebly-interacting dust-like ingredient in the concordance model of cosmology: the dark-energy-cold-dark-

<sup>\*</sup> td455@cantab.ac.uk

<sup>&</sup>lt;sup>†</sup> damianos.iosifidis@ut.ee




FIG. 1. Left: the effect of the curvature in Eq. (6) – the rotation of a vector after parallel transport in a closed loop. Centre: the effect of torsion in Eq. (4) – non-closure of infinitesimal parallelograms formed from parallely-transported vectors. Right: the effect of non-metricity in Eq. (5) – the change in vector norm under parallel transport.

matter ( $\Lambda$ CDM) paradigm [23, 24]. Despite being the 'best fit' model, there have been some suggestions in recent years that  $\Lambda$ CDM may itself be in tension with observations [25– 27]. Underpinning the  $\Lambda$ CDM model are the Friedmann equations, which describe the background expansion of the Universe when GR is coupled to matter-like sources, along with the theory of perturbations around this background. By modifying GR, so the Friedmann and perturbation equations are themselves modified: there is some hope that the resulting phenomenology will lead to a less ad hoc cosmological model than  $\Lambda$ CDM, or one with fewer tensions.

A very general geometry — Non-Riemannian models offer a natural alternative to GR. For two very special cases, GR itself can be equivalently formulated with vanishing curvature using either only torsion, or only non-metricity (all other quantities in Fig. 1 vanishing). These are called the teleparallel and symmetric-teleparallel equivalents of GR (TEGR [28] and STEGR).<sup>1</sup> Together with Riemannian GR, these three formulations are frequently referred to as the 'geometrical trinity' of gravity [30]. However, the geometry does not have to be thus restricted. An admixture of geometries is found in Weyl spacetime [31, 32], which admits part of the general non-metricity tensor in addition to curvature. A more substantial extension is to Riemann–Cartan spacetime [33–39], where the full torsion is instead admitted: models with this geometry may be thought of as examples of Poincaré gauge theory (PGT), where the gauge-theoretic structure is motivated by graivtational coupling to spinorial matter [40-42]. By blending the Weyl and Riemann-Cartan spacetimes, models of dynamical Weyl-Cartan spacetime may be thought of as examples of Weyl gauge theory (WGT) [43]. At the most general level, however, MAG encompasses all these non-Riemannian geometries, and specialises to them depending on properties of the connection.

*A very general Lagrangian* — The choice of geometry is not the only consideration when constructing a non-Riemannian model: one must also select the operators which appear in the action. Quadratic gravity (QG) is the class of the-

ories in Riemannian geometry that model gravity with an action up to quadratic order in curvature [44]. It is closely related to quadratic MAG (OMAG), in which operators quadratic in torsion and non-metricity are also included [4]. Classically, QMAG is a straightforward extension of QG, being indistinguishable from Riemannian quadratic gravity coupled (albeit in a highly non-minimal way) to torsion and non-metricity tensors, which can be thought of as extra matter fields. QG is a generic expectation of GR as a low-energy effective field theory [45–48]: the same may possibly be true of QMAG, though investigations into the non-Riemannian quantum theory are still relatively nascent [7, 49–52].<sup>2</sup> Famously, unitarity is not a general feature of the QG theory-space, according to the standard definition of ghost modes [44]. The particle spectrum of OMAG is much richer than that of OG: without careful tuning the linear spectrum is also blighted by ghosts and tachyons [7, 53–98], though these are not generally the same species which spoil QG. Moreover, there is a particular danger in QMAG that the linear spectrum is not a faithful representation of the general degrees of freedom that are propagating: some modes may become strongly coupled near Minkowski spacetime [5, 58, 95, 99–122].

In this work — We obtain background cosmology of QMAG, with the arbitrary but pragmatic restriction that parity-violating operators are excluded. Indeed, parity is not a good quantum number, but given a particular non-Riemannian geometry it is common to first explore the quadratic theory in the parity-preserving case, following up the investigation with the parity-violating extension. Cosmology has been extensively studied in various kinds of non-Riemannian geometry, see [3, 123-128] (also [60] and references therein). Recent investigations into the full MAG geometry show a diverse collection of phenomena [128, 129], particularly when including the hypermomentum as a source [123]. Despite these advances, the full QMAG action remains unexplored. The rest of this work is structured as follows. In Section II we set out the MAG formalism and discuss our methods. In Section III we obtain the cosmological equations and discuss their structure. In Section IV we motivate two special cases of QMAG by their cosmology, and use linearised particle spectroscopy near Minkowski spacetime to further constrain them. Conclusions follow in Section V, along with technical appendices. We work in natural units  $c \equiv \hbar \equiv 1$ . Further conventions are introduced as needed.

### II. Theoretical development

Having motivated MAG in Section I, we will define our conventions for the 64 extra kinematic d.o.f implied by metricaffine geometry (relative to the ten d.o.f present in the Riemannian case) in Section II A. In Section II B will then understand

<sup>&</sup>lt;sup>1</sup> There is also the generalized teleparallel gravity with both torsion and nonmetricity [29].

<sup>&</sup>lt;sup>2</sup> An immediate objection to QMAG being quantum in origin is that the low-energy effective theory of a dynamical connection should be a priori spanned by a much larger basis of operators than those which can be formed from the squares of curvature, torsion and non-metricity.

how the restriction to an isotropic and homogeneous cosmology eliminates all but five of these.

## A. Metric-affine geometry

Our notation mainly follows [3]. A non-Riemannian spacetime is a differentiable manifold equipped with a symmetric metric  $g_{\mu\nu}$  of signature (-, +, +, +) and an independent affine connection  $\Gamma^{\lambda}_{\mu\nu}$ , where the covariant derivative  $\nabla_{\mu}$  of some tensor field  $X^{\mu}_{\nu}$  is

$$\nabla_{\alpha}X^{\mu}_{\ \nu} \equiv \partial_{\alpha}X^{\mu}_{\ \nu} - \Gamma^{\lambda}_{\ \nu\alpha}X^{\mu}_{\ \lambda} + \Gamma^{\mu}_{\ \lambda\alpha}X^{\lambda}_{\ \nu}.$$
 (1)

Since the difference of two connections is always a tensor, the deviation of the affine connection from the unique, symmetric and metric-compatible Levi–Civita connection  $\tilde{\Gamma}^{\lambda}_{\mu\nu}$  gives the  $4 \times 4 \times 4 = 64$  new d.o.f (i.e. the components of an asymmetric rank-three four-tensor) of the distortion tensor

$$N^{\lambda}_{\ \mu\nu} \equiv \Gamma^{\lambda}_{\ \mu\nu} - \tilde{\Gamma}^{\lambda}_{\ \mu\nu}, \tag{2}$$

where  $\tilde{\Gamma}^{\lambda}_{\mu\nu}$  is explicitly given by the Christoffel formula

$$\tilde{\Gamma}^{\lambda}_{\ \mu\nu} \equiv \frac{1}{2} g^{\alpha\lambda} \left( \partial_{\mu} g_{\mu\nu} + \partial_{\nu} g_{\alpha\mu} - \partial_{\alpha} g_{\mu\nu} \right). \tag{3}$$

From this point onwards, quantities with a tilde over them will denote Riemannian parts derived from  $\tilde{\Gamma}^{\lambda}_{\mu\nu}$  unless stated otherwise. It is possible to derive two non-Riemannian quantities from the affine connection: the torsion and the non-metricity. The torsion tensor is defined to be the antisymmetric part of the connection, with  $4 \times 6 = 24$  d.o.f

$$T^{\lambda}_{\mu\nu} \equiv \Gamma^{\lambda}_{[\mu\nu]} \equiv N^{\lambda}_{[\mu\nu]}. \tag{4}$$

The torsion in Eq. (4) can be said to measure the non-closure of infinitesimal parallelograms formed from parallel transport of tangents along two intersecting paths (see the central diagram in Fig. 1). The non-metricity tensor with  $4 \times 10 = 40$  d.o.f is defined as the degree to which the metricity condition is violated by the independent connection (see rightmost diagram in Fig. 1)

$$Q_{\alpha\mu\nu} \equiv -\nabla_{\alpha}g_{\mu\nu} \equiv 2N_{(\mu\nu)\alpha}.$$
 (5)

The second equality of Eq. (5) follows from expanding the covariant derivative into the connection using Eq. (1), and then expanding the connection using Eq. (2). The Levi–Civita connection in Eq. (3) uniquely cancels all but the non-Riemannian contributions. This post-Riemannian decomposition (PRD) — whereby the original Palatini variables  $\left\{g_{\mu\nu}, \Gamma^{\lambda}_{\mu\nu}\right\}$  are superseded by  $\left\{g_{\mu\nu}, N^{\lambda}_{\ \mu\nu}\right\}$  or  $\left\{g_{\mu\nu}, T^{\lambda}_{\ \mu\nu}, Q_{\lambda\mu\nu}\right\}$  through a simple field reparameterisation — will be used throughout our investigations.

Curvature is defined as the failure of covariant derivatives to commute (see leftmost diagram in Fig. 1). The non-Riemannian curvature tensor is

$$R^{\mu}_{\ \nu\rho\sigma} \equiv \partial_{\rho}\Gamma^{\mu}_{\ \nu\sigma} - \partial_{\sigma}\Gamma^{\mu}_{\ \nu\rho} + \Gamma^{\tau}_{\ \nu\sigma}\Gamma^{\mu}_{\ \tau\rho} - \Gamma^{\tau}_{\ \nu\rho}\Gamma^{\mu}_{\ \tau\sigma}.$$
 (6)

Note that, unlike in the Riemannian case, the non-Riemannian curvature is only antisymmetric in its last two indices. By plugging Eq. (2) into Eq. (6) the post-Riemannian decomposition of the curvature tensor is found to be

$$R^{\mu}_{\nu\rho\sigma} \equiv \tilde{R}^{\mu}_{\nu\rho\sigma} + \tilde{\nabla}_{\rho} N^{\mu}_{\nu\sigma} - \tilde{\nabla}_{\sigma} N^{\mu}_{\nu\rho} + N^{\tau}_{\nu\sigma} N^{\mu}_{\tau\rho} - N^{\tau}_{\nu\rho} N^{\mu}_{\tau\sigma}.$$
(7)

Without using the metric, two independent contractions of the curvature can be defined

$$R_{\mu\nu} \equiv R^{\alpha}_{\ \mu\alpha\nu}, \quad \hat{R}_{\mu\nu} \equiv R^{\alpha}_{\ \alpha\mu\nu}. \tag{8}$$

The former is the Ricci tensor and the latter is the homothetic curvature. A third tensor — the co-Ricci — is defined as  $\check{R}^{\mu}{}_{\rho} \equiv R^{\mu}{}_{\nu\sigma\rho}g^{\nu\sigma}$ . The Ricci scalar remains uniquely defined, since

$$R \equiv R_{\mu\nu}g^{\mu\nu} \equiv -\check{R}_{\mu\nu}g^{\mu\nu} \equiv -R_{\mu\ \alpha\nu}g^{\mu\nu}, \qquad (9)$$

meanwhile the homothetic curvature in Eq. (8) is traceless due to its antisymmetry  $\hat{R}_{\mu\nu}g^{\mu\nu} \equiv 0$ . Note that the notation and index conventions here are different to [4]. The latter two traces of the Riemann tensor do not exist in Riemannian geometry due to the index symmetry  $\tilde{R}_{\alpha\beta\mu\nu} \equiv -\tilde{R}_{\beta\alpha\mu\nu}$ . Consequently, we have  $\hat{R}_{\mu\nu} \equiv \tilde{R}^{\alpha}_{\ \alpha\mu\nu} \equiv 0$  and  $\check{R}_{\mu\nu} \equiv \tilde{R}^{\ \alpha}_{\ \alpha\nu} \equiv -\tilde{R}_{\mu\nu}$  in the Riemannian limit.

It is also useful to define non-Riemannian four-vectors using the torsion and non-metricity. Based on Eqs. (4) and (5) the torsion forms one vector and the non-metricity forms two vectors:

$$t_{\mu} \equiv T^{\lambda}_{\ \mu\lambda}, \quad Q_{\alpha} \equiv Q_{\alpha\mu\nu}g^{\mu\nu}, \quad q_{\alpha} \equiv Q_{\mu\nu\alpha}g^{\mu\nu}.$$
 (10)

Note that Eq. (10) omits the axial vector  $\epsilon_{\mu\alpha\beta\gamma}T^{\alpha\beta\gamma}$ , which will not need a separate notation in this work. The non-metricity does not form an axial vector because of its index symmetries. Having defined the basic geometric ingredients, their explicit forms in a cosmological setting can be introduced.

#### B. Background cosmology

We will now introduce the Friedmann–Lemaître–Robertson– Walker (FLRW) metric in the context of MAG, giving the forms of non-Riemannian ingredients from Section II A in an FLRW coordinate basis derived from the assumed symmetries of the spacetime. The line element in a spatially flat FLRW spacetime is given by

$$ds^{2} = -dt^{2} + a^{2} \left[ dr^{2} + r^{2} \left( d\theta^{2} + \sin^{2}(\theta) d\phi^{2} \right) \right], \qquad (11)$$

in spatially co-moving spherical polar coordinates, where the dimensionless scale factor  $a \equiv a(t)$  is normalised to unity at the current epoch. In Sections II C and III, a minisuperspace method will be used to extract the background dynamics, and for this a 'lapse-like' function  $b \equiv b(t)$  will be introduced, alongside *a*. This is done so that the action can then be varied

with respect to a and b separately, setting b to unity afterwards as a choice of time gauge. With the inclusion of the 'lapse-like' function, Eq. (11) becomes

$$ds^{2} = -b^{2}dt^{2} + a^{2} \left[ dr^{2} + r^{2} \left( d\theta^{2} + \sin^{2}(\theta) d\phi^{2} \right) \right].$$
(12)

The co-moving observer in a homogeneous, isotropic Universe has four-velocity  $u_{\mu}u^{\mu} \equiv -1$ . Based on this velocity, a projection tensor can be defined as

$$P_{\mu\nu} \equiv g_{\mu\nu} + u_{\mu}u_{\nu}. \tag{13}$$

The conditions of homogeneity and isotropy imply that, of the 64 d.o.f in Eq. (2), only five remain. We parameterise these using the 'scalars'  $\{X, Y, Z, V, W\}$ , which depend only on cosmic time t so that  $X \equiv X(t)$ , etc. Using Eq. (13) these scalars are distributed according to

$$N_{\lambda\mu\nu} \equiv X P_{\mu\nu} u_{\lambda} + Y P_{\nu\lambda} u_{\mu} + Z P_{\mu\lambda} u_{\nu} + V u_{\lambda} u_{\mu} u_{\nu} + W \epsilon_{\lambda\mu\nu\kappa} u^{\kappa}.$$
(14)

Equally, with Eqs. (4) and (5) we see that Eq. (14) becomes

$$Q_{\alpha\mu\nu} \equiv 2Zu_{\alpha}P_{\mu\nu} + 2(X+Y)P_{\alpha(\mu}u_{\nu)} + 2Vu_{\alpha}u_{\mu}u_{\nu}, \quad (15a)$$

$$T_{\mu\nu\nu} \equiv (Y+Z)u_{\mu}P_{\nu} + W\epsilon_{\mu\nu\alpha\rho}u^{\rho}.$$
 (15b)

The restrictions in Eqs. (14), (15a) and (15b) may be formally derived from the vanishing of the Lie derivative — see e.g. [3, 126, 130] - however they are also self-evident. Indeed, it follows in cosmology that only the spin-zero parts of the tensor field  $N^{\lambda}_{\mu\nu}$  may contribute to the dynamics. As is well known, an asymmetric tensor of third rank contains five spin-zero modes, one of negative parity (W in this notation) and all the others of positive parity. Thus, Eqs. (14), (15a) and (15b) constitute the spin-zero part of the spin-parity decomposition of the distortion tensor. This decomposition is used for, example, in the canonical analysis, where the analogue of  $u^{\mu}$  is provided by the normal to the space-like hypersurfaces. As will be shown in Appendix C, this decomposition is also used in the study of the particle spectrum, where the analogue of  $u^{\mu}$  is provided by  $n^{\mu} \equiv k^{\mu}/k$  for  $k^2 \equiv k_{\mu}k^{\mu}$ and  $k^{\mu}$  the four-momentum of the (massive) particles. The same principle applies to the metric: it is well known that a symmetric tensor of second rank contains two scalars, both of positive parity. This is the reason why we need the two variables  $\{a, b\}$  in Eq. (12) for a lossless minisuperspace reduction. As a final comment before using these definitions, it will be useful for us to define the following combination of distortion scalars

$$U \equiv V + Z, \tag{16}$$

for various results in Appendix B.

## C. The Einstein-Hilbert case

Having restricted to cosmology in Section II B we will compute the background cosmological equations for the minimal case of MAG, in which the Lagrangian density comprises only the Einstein–Hilbert term

$$S = \frac{1}{2\kappa} \int d^4 x \sqrt{-g} R.$$
 (17)

The first method to obtain the Friedmann equations is covariantly, via field equations. We will work with the PRD and separately vary Eq. (17) with respect to the distortion and the metric

$$\delta S = \frac{1}{2\kappa} \int d^4 x \sqrt{-g} \, \delta g^{\mu\nu} \bigg[ \tilde{R}_{\mu\nu} - \frac{1}{2} g_{\mu\nu} \tilde{R} + \left( N^{\sigma}_{\lambda\sigma} N^{\lambda}_{\ \mu\nu} - N^{\sigma}_{\ \lambda\nu} N^{\lambda}_{\ \mu\sigma} \right) - \frac{1}{2} g_{\mu\nu} g^{\alpha\beta} \left( N^{\sigma}_{\ \lambda\sigma} N^{\lambda}_{\ \alpha\beta} - N^{\sigma}_{\ \lambda\beta} N^{\lambda}_{\ \alpha\sigma} \right) \bigg] + \sqrt{-g} \, \delta N^{\sigma}_{\ \kappa\gamma} \left[ \delta^{\gamma}_{\ \sigma} N^{\kappa\alpha}_{\ \alpha} + g^{\kappa\gamma} N^{\alpha}_{\ \sigma\alpha} - N^{\kappa\gamma}_{\ \sigma} - N^{\gamma}_{\ \sigma}^{\kappa} \right].$$
(18)

On shell we take  $\delta S = 0$  so that Eq. (18) yields the field equations

$$\tilde{R}_{\mu\nu} - \frac{1}{2}g_{\mu\nu}\tilde{R} + D_{\mu\nu} - \frac{1}{2}g_{\mu\nu}D^{\alpha}_{\ \alpha} = 0,$$
(19a)

$$\delta^{\gamma}{}_{\sigma}N^{\kappa\alpha}{}_{\alpha} + g^{\kappa\gamma}N^{\alpha}{}_{\sigma\alpha} - N^{\kappa\gamma}{}_{\sigma} - N^{\gamma}{}_{\sigma}{}^{\kappa} = 0, \qquad (19b)$$

where  $D_{\mu\nu} \equiv 2 N^{\lambda}_{\mu[\nu]} N^{\sigma}_{\lambda|\sigma]}$ . Referring back to the line element in Eq. (11) and the cosmological form of the distortion Eq. (14), the FLRW component forms of these equations simplify to give

$$H = W = X = Y = V + Z = 0,$$
 (20)

where  $H \equiv H(t)$  is the Hubble parameter defined as  $H \equiv \dot{a}/a$ , and an over-dot will always denote differentiation with respect to the cosmic time t, so that  $\dot{a} \equiv da/dt$ . Recall that the 'lapse-like' function introduced in Eq. (12) is set to unity throughout. Note that Eq. (20) agrees with a well known result in MAG: an action linear in the Ricci scalar should return the same dynamics as normal GR in a vacuum. Note however that there is one d.o.f between V and Z (namely U) which is left unconstrained by Eq. (20). We will show in Section III C that this is a result of the invariance of Eq. (17) under the projective transformation

$$\Gamma^{\lambda}_{\ \mu\nu} \mapsto \Gamma^{\lambda}_{\ \mu\nu} + \delta^{\lambda}_{\mu} \xi_{\nu}, \qquad (21)$$

where  $\xi_v \equiv \xi_v(x)$  is an arbitrary local four-vector gauge generator [4, 131].

The second method of calculating the Friedmann equations is to use the minisuperspace<sup>3</sup> method (or Weyl method). This

<sup>&</sup>lt;sup>3</sup> The term '*minisuperspace*' comes from quantum cosmology, when considering the full gravitational path integral. Instead of integrating over all components of the metric  $g_{\mu\nu} \equiv g_{\mu\nu}(x)$ , an approximation can be made by assuming the off-diagonal components to vanish, and and the remaining components to be functions only of *t*, thereby reducing the number of fields to two – the 'lapse-like' and 'shift-like' functions. Including the 'lapse-like' function is what allows for time reparameterisation invariance, and its variation yields a constraint equation.

method can be adapted to classical cosmology [132, 133], it involves imposing Eqs. (12), (14), (15a) and (15b) first, before taking variational derivatives. The three-dimensional spacelike hypersurfaces are integrated out of the original action with an appropriate cut-off to give a fiducial volume. In general, the imposition of constraints on the fields prior to variations is not guaranteed to lead to the correct equations of motion, since the two operations do not commute. However, for Bianchi class A cosmological models, the method holds as guaranteed by Palais' principle of symmetric criticality [134–137] and since the metric here fulfills the requirements<sup>4</sup> we can proceed to obtain the correct equations of motion. The Ricci scalar in a metric-affine FLRW spacetime is

$$R = 3 \left[ 2 \left( \frac{\ddot{a}}{ab^2} + \left( \frac{\dot{a}}{ab} \right)^2 - \frac{\dot{a}\dot{b}}{ab^3} \right) + 3 \frac{\dot{a}}{ab} (X - Y) + \frac{1}{b} (\dot{X} - \dot{Y}) + (X + Y)(V + Z) - 2XY - 2W^2 \right].$$
(22)

Now, using also  $\sqrt{-g} = a^3 b$ , we can use Eq. (22) to write the Lagrangian density in Eq. (17) as

$$S \propto \int dt \left[ a^{3}b \left( (X+Y)(Z+V) - 2XY - 2W^{2} \right) - 2\frac{a\dot{a}^{2}}{b} + \frac{d}{dt} \left[ 2\frac{a^{2}\dot{a}}{b} + a^{3}(X-Y) \right] \right], \quad (23)$$

where the proportionality holds up to constants and the fiducial volume. Note that in Eq. (23) the last term can be dropped, since this is a total derivative. Variation with respect to the minisuperspace field variables  $\{a, b, X, Y, Z, V, W\}$  gives us seven (or six independent) equations, respectively

$$6H^{2} + 4\dot{H} = 6W^{2} + 6XY - 3(X + Y)(V + Z),$$
  

$$2H^{2} = 2W^{2} + 2XY - (X + Y)(V + Z),$$
  

$$V + Z = 2Y, \quad V + Z = 2X, \quad X + Y = 0,$$
  

$$X + Y = 0, \quad W = 0.$$
(24)

Collectively, Eq. (24) simplify to yield precisely Eq. (20), showing that the covariant and minisuperspace methods are consistent.

### III. Structure of the field equations

As discussed in Section I, it is interesting to consider the most general action quadratic in curvature, torsion, and nonmetricity, i.e. QMAG. The QMAG action is given by<sup>5</sup>

$$S = -\frac{1}{2} \int d^4x \sqrt{-g} \left[ -a_0 R + R^{\rho\sigma\mu\nu} \left( c_1 R_{\rho\sigma\mu\nu} + c_2 R_{\sigma\rho\mu\nu} \right) \right] \right]$$

$$+ c_{3}R_{\mu\nu\rho\sigma} + c_{4}R_{\mu\sigma\rho\nu} + c_{5}R_{\mu\rho\sigma\nu} + c_{6}R_{\rho\mu\sigma\nu} + c_{7}R^{\mu\nu}R_{\mu\nu} + c_{8}R^{\mu\nu}R_{\nu\mu} + c_{9}\check{R}^{\mu\nu}\check{R}_{\mu\nu} + c_{10}\check{R}^{\mu\nu}\check{R}_{\nu\mu} + c_{11}R^{\mu\nu}\check{R}_{\mu\nu} + c_{12}R^{\mu\nu}\check{R}_{\nu\mu} + c_{13}\hat{R}^{\mu\nu}\hat{R}_{\mu\nu} - c_{14}R^{\mu\nu}\hat{R}_{\mu\nu} - c_{15}\hat{R}^{\mu\nu}\check{R}_{\mu\nu} + c_{16}R^{2} + 4a_{1}T^{\mu\nu\rho}T_{\mu\nu\rho} + 4a_{2}T^{\mu\nu\rho}T_{\nu\mu\rho} + 4a_{3}t^{\mu}t_{\mu} + a_{4}Q^{\mu\nu\rho}Q_{\mu\nu\rho} + a_{5}Q^{\mu\nu\rho}Q_{\nu\mu\rho} + a_{6}Q^{\mu}Q_{\mu} + a_{7}q^{\mu}q_{\mu} + a_{8}Q^{\mu}q_{\mu} + 2a_{9}Q^{\mu\nu\rho}T_{\nu\rho\mu} + 2\left(a_{10}Q^{\mu} + a_{11}q^{\mu}\right)t_{\mu}\right].$$
(25)

This action has many parameters and is very cumbersome; therefore, the goal of Section III A will be to make it as hygienic as possible to work with.

### A. Reduced couplings for cosmology

As in Section II C, the action in Eq. (25) can be reduced by using the FLRW forms of tensors which were introduced in Section II B. The large symmetry of the cosmological ansatz restricts the forms of the operators in the action, causing some to vanish and others to be related to each other. Accordingly, we will next obtain the *reduced* couplings in Eq. (30). Before quoting these results, however, we will discuss the formal procedure for obtaining them.

We define the operators of dimensions four and two as being respectively  $\mathscr{C}_i \equiv \delta S / \delta c_i$  and  $\mathscr{A}_i \equiv \delta S / a_i$  for all relevant *i*. The operators are hexic polynomials in  $\{x_1, \dots, x_{13}\} \equiv \{X, Y, Z, V, W, \dot{X}, \dot{Y}, \dot{W}, a^{-1}, \dot{a}, \ddot{a}, b^{-1}, \dot{b}\}$  with rational coefficients. For QMAG, the problem of extracting from the  $\{c_1, \dots, c_{16}\}$  and  $\{a_0, \dots, a_{11}\}$  some minimal set of couplings relevant to the background cosmology is evidently related to finding all non-trivial solutions for  $\{\chi_1, \dots, \chi_{16}\}$ and  $\{\alpha_0, \dots, \alpha_{11}\}$  to the equations

$$\sum_{i=1}^{16} \chi_i \mathscr{C}_i \equiv 0, \quad \sum_{i=0}^{11} \alpha_i \mathscr{A}_i \equiv 0.$$
 (26)

The problem in Eq. (26) is linear: its non-linear generalisation in commutative algebra is the so-called *ideal of relations* for each of the algebraically dependent sets  $\{\mathscr{C}_1, \ldots, \mathscr{C}_{16}\}$ and  $\{\mathscr{A}_0, \ldots, \mathscr{A}_{11}\}$  [138]. This latter problem is not relevant to QMAG, having instead applications to quartic or hexic MAG and beyond.<sup>6</sup> Nonetheless, the full ideal of relations can be obtained with surprisingly little effort using computer algebra techniques, and so we use two such techniques to solve Eq. (26).

A Gröbner basis technique was used for the  $\{\mathscr{A}_0, \dots, \mathscr{A}_{11}\}$ . We introduce new indeterminates  $\{\mathbf{a}_0, \dots, \mathbf{a}_{11}\}$ , and ascribe

<sup>&</sup>lt;sup>4</sup> The metric ansatz in Eq. (12) meets the conditions in [135, 136].

<sup>&</sup>lt;sup>5</sup> Note that the couplings  $\{c_1, \ldots, c_{16}\}$  and  $\{a_0, \ldots, a_{11}\}$  are carefully tuned so as to be identical to those in [4]. Since our conventions for the geometry itself, as set out in Section II A, follow [3] instead of [4], there are apparent sign changes in front two terms.

<sup>&</sup>lt;sup>6</sup> Moreover, quartic and hexic MAG are additionally spanned by many more operators which are not complete squares of the QMAG operators, in addition to cubic operators and quintic operators.

a monomial elimination order such that for any *i* and *j* we have  $\mathbf{x}_i > \mathbf{a}_j$ . With this order, we compute the Gröbner basis for the ideal  $\langle \mathbf{a}_0 - \mathscr{A}_0, \dots, \mathbf{a}_{11} - \mathscr{A}_{11} \rangle$ , which is a subset of the polynomial ring  $\mathbb{Q} [\mathbf{x}_1, \dots, \mathbf{x}_{13}, \mathbf{a}_0, \dots, \mathbf{a}_{11}]$  over the rationals. By dropping all elements of this basis which refer explicitly to the  $\{\mathbf{x}_1, \dots, \mathbf{x}_{13}\}$ , so the ideal of relations is obtained. Those elements of the ideal of relations which are linear in the  $\{\mathbf{a}_0, \dots, \mathbf{a}_{11}\}$  evidently solve for the  $\{\alpha_0, \dots, \alpha_{11}\}$  in Eq. (26), once the identifications  $\mathbf{a}_i \equiv \mathscr{A}_i$  are made for all *i*. Using the *Mathematica* function GroebnerBasis, we find the three relations

$$\begin{aligned} \mathcal{A}_{1} + \mathcal{A}_{2} - \mathcal{A}_{3} &\equiv 0, \quad 3\mathcal{A}_{9} + \mathcal{A}_{10} - \mathcal{A}_{11} &\equiv 0, \\ 3\mathcal{A}_{4} - 3\mathcal{A}_{5} - \mathcal{A}_{6} - \mathcal{A}_{7} + 2\mathcal{A}_{8} &\equiv 0. \end{aligned}$$
(27)

Consequently, of the 11 original non-Riemannian operators, it follows from Eq. (27) that only eight are independent. These relations are consistent with the results in [128]. The non-linear extension also reveals a fourth, quartic relation: this is shown in Eq. (A1).

The GroebnerBasis implementation was not powerful enough for the  $\{\mathscr{C}_1, \ldots, \mathscr{C}_{16}\}$ . An alternative method for finding the ideal of relations is to find the kernel of the ring map  $F : \mathbb{Q}[\mathscr{C}_1, \ldots, \mathscr{C}_{16}] \mapsto \mathbb{Q}[x_1, \ldots, x_{13}]$ , i.e. those elements which map to zero, since these are the polynomial functions of the  $\{\mathscr{C}_1, \ldots, \mathscr{C}_{16}\}$  which identically vanish. An algorithm implemented in *Macaulay2* [139] finds

$$\mathcal{C}_{16} - \mathcal{C}_1 - 2\mathcal{C}_2 - \mathcal{C}_6 - 4\mathcal{C}_5 - \mathcal{C}_4 + 6\mathcal{C}_{12} + \mathcal{C}_3 \equiv 0,$$

$$\begin{aligned} \mathscr{C}_8 &-\mathscr{C}_1 - \mathscr{C}_2 - \mathscr{C}_6 - 2\mathscr{C}_5 - \mathscr{C}_4 + 2\mathscr{C}_{12} + \mathscr{C}_{10} \equiv 0, \\ \mathscr{C}_7 - \mathscr{C}_8 &\equiv \mathscr{C}_9 - \mathscr{C}_{10} \equiv \mathscr{C}_{11} - \mathscr{C}_{12} \equiv 0, \\ \mathscr{C}_{13} &\equiv \mathscr{C}_{14} \equiv \mathscr{C}_{15} \equiv 0. \end{aligned}$$
(28)

This implementation finds in addition to Eq. (28) two extra relations in Eqs. (A2) and (A3) which apply to quartic MAG, and one invariant in Eq. (A4) which applies to hexic MAG. The final three relations in Eq. (28) follow by construction because the homothetic curvature  $\hat{R}_{\mu\nu} \equiv 0$  is defined to be the Maxwell operator for the Weyl vector, and the Maxwell tensor always vanishes in cosmology. Similarly, the third, fourth and fifth relations in Eq. (28) are equivalent to

$$R^{[\bar{\mu}\bar{\nu}]}R_{[\bar{\mu}\bar{\nu}]} \equiv \check{R}^{[\bar{\mu}\bar{\nu}]}\check{R}_{[\bar{\mu}\bar{\nu}]} \equiv R^{[\bar{\mu}\bar{\nu}]}\check{R}_{[\bar{\mu}\bar{\nu}]} \equiv 0, \qquad (29)$$

which is to be expected for the same reason. The conditions in Eq. (28) imply that of the original 16 curvature operators, only eight are independent. This is also consistent with the results in [128]. In summary, of the total 28 operators entering Eq. (25), we expect only 17 to be linearly independent. The couplings entering in Eq. (25) are variables that form the basis of a 28-dimensional space, and their linear combinations in front of the monomials are vectors in that space. This linear system can be reduced via standard Gaussian elimination to yield a set of 17 linearly independent coefficients: we relabel these as  $\{a_0, d_1, \dots, d_{16}\}$ . Thus the Einstein–Hilbert parameter  $a_0$  already plays a unique role in the cosmology; the relevant formulae for the other new couplings are:

$$\begin{aligned} d_1 &\equiv -6 \left( 2c_1 + 2c_{10} - 2c_{11} - 2c_{12} + 6c_{16} - 2c_2 + 2c_3 + c_4 - c_5 + c_6 + 2 \left( c_7 + c_8 + c_9 \right) \right), \\ d_2 &\equiv 12 \left( c_{10} - c_{11} - c_{12} + 6c_{16} + c_7 + c_8 + c_9 \right), \\ d_3 &\equiv -6 \left( 4c_1 + 4c_{10} - 5c_{11} - 5c_{12} + 2 \left( 9c_{16} - 2c_2 + 2c_3 + c_4 - c_5 + c_6 + 3 \left( c_7 + c_8 \right) + 2c_9 \right) \right), \\ d_4 &\equiv -3 \left( 6c_1 + 7c_{10} - 6c_{11} - 6c_{12} + 27c_{16} - 4c_2 + 5c_3 + 3c_4 - 2c_5 + 3c_6 + 9 \left( c_7 + c_8 \right) + 7c_9 \right), \\ d_5 &\equiv 6 \left( 14c_1 + 4c_{10} - 4c_{11} - 4c_{12} + 12c_{16} - 14c_2 + 12c_3 - 7c_4 + 7c_5 - 7c_6 + 4 \left( c_7 + c_8 + c_9 \right) \right), \\ d_6 &\equiv 6 \left( 12c_1 + 4c_{10} - 5c_{11} - 5c_{12} + 18c_{16} - 12c_2 + 8c_3 - 2c_4 + 4c_5 - 6c_6 + 6 \left( c_7 + c_8 \right) + 4c_9 \right), \\ d_7 &\equiv -6 \left( 12c_1 + 6c_{10} - 5c_{11} - 5c_{12} + 18c_{16} - 12c_2 + 8c_3 - 6c_4 + 4c_5 - 2c_6 + 4 \left( c_7 + c_8 \right) + 6c_9 \right), \\ d_8 &\equiv 24 \left( c_1 - c_6 \right), \quad d_9 &\equiv 3 \left( 2a_4 + a_5 + 3a_7 \right), \quad d_{10} &\equiv 3 \left( 3a_{11} + 4a_4 + 2a_5 + 6a_7 + a_9 \right), \\ d_{11} &\equiv 3 \left( 2a_1 + 3a_{11} + a_2 + 3a_3 + 2a_4 + a_5 + 3a_7 + a_9 \right), \\ d_{12} &\equiv -6 \left( 2a_7 + a_8 \right), \quad d_{13} &\equiv -6 \left( a_{10} + a_{11} + 2a_7 + a_8 \right), \\ d_{14} &\equiv 4 \left( a_4 + a_5 + a_6 + a_7 + a_8 \right), \quad d_{15} &\equiv -3 \left( 3a_{11} - 4a_5 - 6a_8 + a_9 \right), \quad d_{16} &\equiv -24 \left( a_1 - a_2 \right). \end{aligned}$$

Note in Eq. (30) that the couplings  $\{d_1, \ldots, d_8\}$  are of mass dimension zero whilst the  $\{d_9, \ldots, d_{16}\}$  are of mass dimension two.

### B. Minisuperspace computation

The minisuperspace method outlined in Section II C is now used to calculate the modified Friedmann equations corre-

sponding to the full model in Eq. (25). As found in Section III A, the minisuperspace Lagrangian density corresponding to Eq. (25) can be expressed entirely in terms of the reduced couplings in Eq. (30). This action is then varied with respect to the seven fields  $\{a, b, X, Y, Z, V, W\}$ , subsequently setting the 'lapse-like' function to unity and so restricting to FLRW coordinates by passing from Eq. (12) to Eq. (11). To our knowledge, the full equations were not yet obtained elsewhere. They are, however, very cumbersome, and we defer

them to Appendix B.

The *b*-equation is a constraint which we take to be the first modified Friedmann equation, in the sense that it always allows one to solve for the term proportional to  $a_0H^2$ . The first Friedmann equation in Eq. (B1) has mass dimension four, and contains besides the  $a_0H^2$  term a whole host of other terms. Terms linear in W or its velocities are generally absent, since these are pseudoscalar in nature, whilst Eq. (25) is parity-preserving.

The second modified Friedmann equation should allow one to solve for the term  $a_0\dot{H}$ ; it is constructed from the equations of motion for *a*, *b*, *X* and *Y* according to

$$\frac{\delta S}{\delta a} - 3\frac{\delta S}{\delta b} + \frac{d}{dt} \left(\frac{\delta S}{\delta X} - \frac{\delta S}{\delta Y}\right) = 0.$$
(31)

The reason for the final term in Eq. (31) is to remove from the *a*-equation in Eq. (B2) the triple-derivative structure

$$\frac{\delta S}{\delta a} \supset 2d_1 \ddot{H} + \left(3d_1 - d_3 - \frac{d_2}{2}\right) \ddot{X} + \left(d_1 - d_3 - \frac{d_2}{2}\right) \ddot{Y}. \tag{32}$$

As an aside, an alternative route to removing the higherderivative terms in Eq. (32) is to set the coefficients in Eq. (32) to zero, resulting in a special case of QMAG. Indeed, the resulting conditions  $d_1 = 0$  and  $d_2 + 2d_3 = 0$  greatly simplify the whole system, though we do not assume them moving forward. In GR, the second Friedmann equation can be derived from the first by taking its derivative and subtracting from it the product of itself with the Hubble parameter. In the metricaffine case, this procedure generalises to the identity

$$H\frac{\delta S}{\delta a} \equiv \left(\frac{d}{dt} + 3H\right)\frac{\delta S}{\delta b} - \dot{X}\frac{\delta S}{\delta X} - \dot{Y}\frac{\delta S}{\delta Y} - \dot{Z}\frac{\delta S}{\delta Z} - \dot{V}\frac{\delta S}{\delta V} - \dot{W}\frac{\delta S}{\delta W},$$
(33)

which holds both on- and off-shell. The reasoning behind Eq. (33) is as follows. In the case of GR, Eq. (33) holds because of the projection of the Riemannian Bianchi identity  $2u^{\nu}\tilde{\nabla}^{\mu}\tilde{R}_{\mu\nu} \equiv u^{\nu}\tilde{\nabla}_{\nu}\tilde{R}$  along the time direction. When the theory is extended from GR to Eq. (25), the Lagrangian density acquires (i) terms quadratic in  $\tilde{R}^{\mu}_{\ \nu\rho\sigma}$ , (ii) a dynamical theory of  $N^{\lambda}_{\ \mu\nu}$  minimally coupled to the metric, and (iii) nonminimal couplings between  $N^{\lambda}_{\mu\nu}$  and  $\tilde{R}^{\mu}_{\nu\rho\sigma}$  (i.e., the crossterms from the PRD of operators quadratic in  $R^{\mu}_{\nu\rho\sigma}$ ). Focussing just on the type-(i) terms, it is well known that the Bianchi identity extends to QG [140] - a result which is consistent with the LHS and the first term on the RHS of Eq. (33). To understand the origin of the remaining terms on the RHS of Eq. (33), we consider the type-(ii) operators. These extend QG with an effective matter sector, such that the Bianchi identity must be extended by the four-divergence of the effective stress-energy tensor. As expected, when  $N_{\mu\nu}^{\lambda}$  is restricted as in Eq. (14), it can be shown that the extra terms in Eq. (33) derive from applying the operator  $u^{\nu} \tilde{\nabla}^{\mu} \delta / \delta g^{\mu\nu}$  to the action of a general theory of  $N^{\lambda}_{\mu\nu}$  minimally coupled to the metric, and then assuming FLRW-type background values for all fields. This also explains why Eq. (33) is invariant under linear redfinitions of the distortion scalars, i.e. it doesn't matter

how we normalise the  $\{X, Y, Z, V, W\}$ . The type-(iii) terms are similarly consistent with Eq. (33).

As a calibration of the full system of equations we compared against [127, 128], where an action quadratic in torsion and non-metricity was considered and the vacuum dynamics found to reduce to GR. In terms of the reduced couplings in Eq. (30), this model corresponds to setting all the  $\{d_1, \ldots, d_8\}$  to zero. After simplifying, the result is

$$H = W = X = Y = V = Z = 0,$$
 (34)

which differs from Eq. (20) by the requirement that V and Z both vanish. This is an expected consequence of the broken projective invariance of the model in [127, 128].

## C. Conditions for projective invariance

It remains to explain the claim, regarding Eq. (20), that projective-invariant models exclusively propagate the linear combination V+Z, rather than V and Z separately. The transformation of the affine connection in Eq. (21) can be equivalently viewed as a transformation of the distortion

$$N^{\lambda}_{\ \mu\nu} \mapsto N^{\lambda}_{\ \mu\nu} + \delta^{\lambda}_{\mu}\xi_{\nu}, \qquad (35)$$

where  $\xi_v \equiv \xi_v(x)$  is an arbitrary local vector gauge generator. By combining Eqs. (13) and (14), the distortion in Eq. (2) is restricted by cosmology to

$$N^{\lambda}_{\mu\nu} = (X + V + Y + Z) u^{\lambda} u_{\mu} u_{\nu} + X g_{\mu\nu} u^{\lambda} + Z \delta^{\lambda}_{\mu} u_{\nu} + Y \delta^{\lambda}_{\nu} u_{\mu} + W \epsilon^{\lambda}_{\mu\nu\kappa} u^{\kappa}.$$
(36)

The only projective transformation which is consistent with isotropy and homogeneity is  $\xi_v = \lambda u_v$ , where  $\lambda \equiv \lambda(t)$ . By combining Eqs. (35) and (36) it follows that the scalars transform according to

$$Z \mapsto Z - \lambda, \quad V \mapsto V + \lambda,$$
 (37)

and from Eq. (37) the invariance of V + Z follows immediately. As a consequence, projective invariance in cosmology requires the minisuperspace Lagrangian just to depend on the combination V + Z. Equivalently, the condition

$$\frac{\delta S}{\delta V} - \frac{\delta S}{\delta Z} = 0, \tag{38}$$

must hold also off-shell, not just on-shell, for a model invariant under the projective transformation. Calculating Eq. (38) explicitly for the QMAG model and couplings in Eqs. (25) and (30) yields

$$\begin{pmatrix} d_{10} + 2d_{12} - d_{13} - 8d_{14} + d_{15} + 2d_9 \end{pmatrix} V + \begin{pmatrix} -d_{12} + d_{15} \end{pmatrix} X + \begin{pmatrix} -2d_{11} + 3d_{12} - 4d_{13} + d_{15} + 2d_9 \end{pmatrix} Y + \begin{pmatrix} -3d_{10} + 2d_{11} \\ -8d_{12} + 7d_{13} + 24d_{14} - 5d_{15} - 8d_9 \end{pmatrix} Z = 0.$$
(39)

If Eq. (39) holds off-shell as well as on-shell, then the coefficients of each scalar must vanish separately. When reduced as far as possible, and translated back into the couplings in Eq. (25) using Eq. (30), the conditions are

$$2a_1 - 4a_{10} - a_{11} + a_2 + 3a_3 + a_9 = 0,$$
  

$$4a_4 - 3a_{10} + 16a_6 + 2a_8 + a_9 = 0,$$
  

$$4a_5 - 3a_{11} + 4a_7 + 8a_8 - a_9 = 0.$$
(40)

It is noted that Eq. (40) are precisely the constraints on the mass-dimension-two parameters obtained in [4, 131]. In [4], three constraints on the mass-dimension-zero couplings are obtained, which beyond Eq. (40) are additionally needed for full projective invariance of Eq. (25) in completely general spacetimes. By contrast, the mass-dimension-zero couplings are entirely unconstrained by invariance under our cosmological projective transformation in Eq. (37): all the quadratic curvature operators automatically have this restricted version of the symmetry. In fact, this is to be expected. The restricted transformation is a special case of the scenario  $\xi_{\mu} = \partial_{\mu} \phi$ where a conservative field — equivalently the local scalar  $\phi \equiv$  $\phi(x)$  — acts as the vector generator. It has been shown already in [141] that any OMAG constructed entirely from curvature operators supports this restricted symmetry. In the case of cosmology, the scalar generator 'rolls' in the direction of time according to  $\phi \equiv \phi(t)$ , and in this sense it may be termed a khronon field [142].

#### IV. Special models and their particles

The field equations obtained in Section III still lack any predictive power, due to the abundance of unconstrained parameters. We will now identify some special cases of the model in Eq. (25) whose cosmological dynamics have some interesting features. Regardless of how compelling these features might be, this is evidently a very superficial analysis, not extending, for example, to the perturbation theory. This is a severe limitation, because almost all metric-affine models one can write down are expected to be inconsistent as physical theories, regardless of how well they compare against observations. The most pernicious inconsistencies cannot be detected at the level of the background cosmology, having instead to do with unitarity violation, or strong coupling.

As a first step towards consistency, we compute for each special case the linear particle spectra near Minkowski spacetime. If the models admit a perturbative, weak gravity regime at all, then the elimination of higher-spin particles along with ghosts and tachyons is a necessary (but not sufficient) condition for consistency. The computations are performed with the *PSALTer* software [143], and the resulting particle spectrographs are presented in Appendix C.

### A. Gravity with K-screening

As mentioned already in Section IIB, generally accepted bounds on the  $\Lambda$ CDM model [23] are not strong enough to guarantee that the spatially flat line element in Eq. (12) describes the background metric of our Universe. In general, some curvature scale K should be assumed, according to

$$ds^{2} = -dt^{2} + a^{2} \left[ \frac{dr^{2}}{1 - Kr^{2}} + r^{2} \left( d\theta^{2} + \sin^{2}(\theta) d\phi^{2} \right) \right].$$
(41)

All the methods and conclusions of Section III apply equally well to the spatially curved line element in Eq. (41), though the expressions are more cumbersome and the full field equations are confined to the supplemental materials [144]. An obvious question to ask, given this full set of equations, is whether it is possible to completely screen K from the background dynamics. This so-called 'K-screening' effect certainly does not apply in GR, so that K is in principle an important observable. Recent observational probes of  $\Lambda$ CDM are found to be in moderate tension over K, and their exclusion suggests Kto be a small positive scale, or an effectively negative energy density weighing in at a few percent of the combined baryonic and dark-sector energy budget [25]. If this is true, then the Riemannian geometry of the observable Universe is very slightly closed. In any case, the smallness of this curvature scale (the base model of  $\Lambda$ CDM actually assumes K = 0) is a key motivation for inflation in the early Universe. Just at the background level, therefore, the possibility of a K-screening effect seems like a promising starting point for modifying GR so as to alleviate cosmological tensions, or formulate a more holistic replacement for  $\Lambda$ CDM.

There is, in fact, a clear precedent for K-screening leading to a compelling non-Riemannian model. Thorough analyses of the particle spectra of parity-preserving, quadratic PGT (i.e. the  $Q_{\mu\nu\sigma} \equiv 0$  limit of MAG) identified certain special cases which (i) are free of ghosts and tachyons, (ii) propagate light d.o.f which might plausibly correspond to graviton polarisations, and (iii) are power-counting renormalisable [80, 86]. The presence of a graviton-like particle is very significant in these models, because the PGT K-screening conditions eliminate the Einstein-Hilbert term. A study of the background cosmology later revealed that these promising models are precisely those which are K-screened [88]. Moreover, these models affect the background dynamics in other ways, suggesting a possible mechanism for alleviating yet another tension within  $\Lambda CDM$  — the so-called *Hubble tension* — regarding the current value of H [88, 145, 146].

For the full MAG model in Eq. (25), and referring to the reduced couplings in Eq. (30), we find that *K*-screening occurs (i.e. *K* is eliminated from all the equations) when the following conditions are satisfied:

$$a_0 = d_1 = d_2 = d_3 = 0. \tag{42}$$

Thus, in MAG as with PGT, the Einstein–Hilbert term is excluded by Eq. (42). We will make no attempt here to connect Eq. (42) with the exact solutions proposed in [88], but instead proceed directly to the particle analysis. The constraints in Eq. (42) are so-far quite minimal, and even when  $a_1$  and  $a_4$  are the only mass-dimension-two parameters present, the preliminary analysis in Appendix C suggests a rich spectrum of massive modes. By imposing further, minimal constraints in an algorithmic manner, it is possible to ensure that the squares

of the masses of all the particles are given by rational functions of the remaining Lagrangian couplings. The resulting model propagates only a massive  $0^-$  particle (see unitarity conditions in Eq. (C6)), without any massleess d.o.f. This is somehow disappointing, but hardly surprising. Indeed, the conditions Eqs. (C1) to (C4), (C5a) and (C5b), when imposed on Eq. (25), reduce the quadratic curvature sector to the square of the Holst pseudoscalar  $\epsilon_{\mu}{}^{\nu\rho\sigma}R^{\mu}{}_{\nu\rho\sigma}$ . It is well known (see e.g. [19]) that this operator propagates a single massive pseudoscalar particle (without strong coupling). Moreover, it does not involve the Riemannian curvature at all: this is consistent with the expectation of K-sceening. Our analysis shows only that K-screening in QMAG can lead to wholly self-consistent theoretical models, it does not (yet) point to a phenomenologically viable model. In particlular, a replacement for the graviton is still needed. This should presumably be derived from the  $2^+$  or  $2^-$  sectors of the theory, which still contain a priori many massive modes once the K-screening condition in Eq. (42) is imposed. A thorough investigation of this matter is left to future work, as is an analysis of the rather large array of source constraints in the minimal Holst-squared model in Fig. 5.

### B. Integrable gravity and its Maxwell limit

Finally, we provide one alternative branch of MAG to that proposed in Section IV A, motivated by an exact solution to the cosmological equations of motion. Our starting point will be to remove odd powers of H from the equations, wherever they are not coupled to other fields. This leads to the conditions

$$d_6 = d_7, \quad d_5 = d_4 = d_3 = d_2 = d_1 = 0.$$
 (43)

Whilst Eq. (43) are somewhat arbitrary, they greatly simplify the field equations, which can be written down compactly as:

$$\begin{aligned} 6a_0H^2 + d_{14}V^2 - (6a_0 - d_{16})W^2 + (3a_0 + d_{12})VX + d_9X^2 \\ + d_8W^2X^2 + (3a_0 + d_{13})VY - (6a_0 - d_{10})XY \\ - 2(d_7 - d_8)W^2XY + d_{11}Y^2 - (2d_7 - d_8)W^2Y^2 \\ + (d_{10} + 2d_{12} - d_{13} - 6d_{14} + d_{15} + 2d_9)VZ \\ + (3a_0 + d_{15})XZ + (3a_0 - 2d_{11} + 3d_{12} - 3d_{13} \\ + d_{15} + 2d_9)YZ - (d_{10} - d_{11} + 3d_{12} - 3d_{13} - 9d_{14} \\ + 2d_{15} + 3d_9)Z^2 = 0, \end{aligned}$$
(44a)

$$12a_{0}\dot{H} - d_{7}W\left(5\dot{W}\left(X+Y\right) + W\left(\dot{X}+\dot{Y}\right)\right) = 0, \quad (44b)$$

$$\begin{aligned} (3a_0 + d_{12})V + d_7 H W^2 + 2d_9 X + 2d_8 W^2 X \\ + (d_{10} - 6a_0)Y + 2(d_8 - d_7)W^2 Y + (3a_0 + d_{15})Z \\ - d_7 W \dot{W} &= 0, \end{aligned} \tag{44c}$$

$$(3a_0 + d_{13})V + d_7HW^2 + (d_{10} - 6a_0)X + 2(d_8 - d_7)W^2X + 2d_{11}Y + 2(d_8 - 2d_7)W^2Y$$

$$+ (3a_0 - 2d_{11} + 3d_{12} - 3d_{13} + d_{15} + 2d_9)Z - d_7W\dot{W} = 0,$$
(44d)

$$\begin{aligned} (d_{10} + 2d_{12} - d_{13} - 6d_{14} + d_{15} + 2d_9)V \\ &+ (3a_0 + d_{15})X + (3a_0 - 2d_{11} + 3d_{12} - 3d_{13} \\ &+ d_{15} + 2d_9)Y - 2(d_{10} - d_{11} + 3d_{12} - 3d_{13} - 9d_{14} \\ &+ 2d_{15} + 3d_9)Z = 0, \end{aligned} \tag{44e}$$

$$2d_{14}V + (3a_0 + d_{12})X + (3a_0 + d_{13})Y + (d_{10} + 2d_{12}) - d_{13} - 6d_{14} + d_{15} + 2d_9)Z = 0,$$
(44f)

$$2(d_{16} - 6a_0)W + 5d_7HW (X + Y) + 2d_8WX^2 + 4(d_8 - d_7)WXY + 2(d_8 - 2d_7)WY^2 + d_7W (\dot{X} + \dot{Y}) = 0.$$
(44g)

It is important to note that if W = 0 then everything trivializes as, in this case, it follows that X = Y = Z = V = H = 0as can be easily verified. We will therefore look for solutions with  $W \neq 0$ , so that Eq. (44g) becomes

$$2(d_{16} - 6a_0) + 5d_7 H (X + Y) + 2d_8 X^2 + 4(d_8 - d_7) XY + 2(d_8 - 2d_7) Y^2 + d_7 (\dot{X} + \dot{Y}) = 0.$$
(45)

**The case**  $d_7 = 0$  — The parameter  $d_7$  heavily influences the behaviour of the solutions. For  $d_7 = 0$ , and after some simple algebra, it follows that

$$H = H_0 \equiv \text{const.}, \quad Y \propto V \propto Z \propto X = X_0 \equiv \text{const.},$$
$$W = W_0 \equiv \pm \sqrt{\frac{\sigma_1 X_0^2 - 6a_0 H_0^2}{\sigma_2 X_0^2 + \sigma_3}},$$
(46)

where the proportionality factors as well as the  $\{\sigma_i\}$  are some long expressions of the coefficients  $a_0$  and  $\{d_i\}$ . Therefore, for  $d_7 = 0$  one can only have de Sitter expansion and constant distortion.

**The case**  $d_7 \neq 0$  — Now let us focus on the non-trivial  $d_7 \neq 0$  case. Subtracting Eq. (44c) and Eq. (44d) we get

$$(d_{12} - d_{13})V + (2d_9 - d_{10} + 6a_0)X + (d_{10} - 6a_0 - 2d_{11})Y + (2d_{11} - 3d_{12} + 3d_{13} - 2d_9)Z + 2d_7W^2(X + Y) = 0.$$
(47)

When the system in Eqs. (44e) and (44f) is not degenerate we can solve for V and Z in terms of X and Y, viz.

$$V = k_1 X + k_2 Y \quad Z = k_3 X + k_4 Y, \tag{48}$$

where again the  $\{k_i\}$  are some linear combinations of the parameters of the theory. We may then substitute Eq. (48) back into Eq. (47) and solve for  $W^2$ , to yield

$$W^{2} = \frac{k_{5}X + k_{6}Y}{2d_{7}(X+Y)} = \frac{k_{6}}{2d_{7}} + \frac{(k_{5} - k_{6})X}{2d_{7}(X+Y)}.$$
 (49)

The case  $k_5 \neq k_6$  — Note that choosing the theory parameters such that  $k_6 = k_5$  implies that W = const.. Let us firstly consider the case  $k_6 \neq k_5$ . On substituting Eq. (48) into Eq. (44a) we obtain

$$6a_0H^2 + k_7X^2 + k_8XY + k_9Y^2 + W^2 \Big[ d_{16} - 6a_0 + d_8(X+Y)^2 - 2d_7Y(X+Y) \Big] = 0.$$
(50)

Then, using Eq. (49) we may eliminate  $W^2$  from Eq. (50) and solve for *H* in terms of *X* and *Y* according to

$$6a_0H^2 - \frac{6a_0 - d_{16}}{2d_7} \left[ k_6 + (k_5 - k_6)\frac{X}{X + Y} \right] - \left[ \frac{d_8}{2d_7} k_5 - k_7 \right] X^2 - \left[ \frac{d_8}{2d_7} k_6 - k_6 - k_9 \right] Y^2 - \left[ \frac{d_8}{2d_7} (k_5 + k_6) - k_5 - k_8 \right] XY = 0.$$
(51)

We have therefore expressed everything in terms of X and Y. These can be inverted in terms of H and W as the dynamical variables or any other pick of two out of four. Then, by eliminating the rest of the variables, Eq. (44b) combined with either of Eq. (44c) or Eq. (44d) gives us a system of coupled second order ordinary differential equations for, say, the variables H and W. The resulting system is, in general, somewhat complicated. However, we do get some quite simple expressions if  $d_{16} = 6a_0$  and in Eq. (51) remaining terms on the RHS form a perfect square  $(X + Y)^2$ . For such a configuration, taking the square root of the aforementioned equation, yields

$$H = \lambda_0 (X + Y), \tag{52}$$

provided the following conditions hold:

$$k_5 + k_8 = k_6 + k_7 + k_9, \quad \frac{d_8(k_7 - k_8 + k_9)}{k_6 - k_7 + k_9} + 2d_7 = 0.$$
 (53)

Combining this with Eq. (49) we deduce

$$X = \frac{W^2 - \lambda_1}{\lambda_0 \lambda_2} H.$$
 (54)

Then eliminating X and Y, equations Eq. (44b) and Eq. (44c) read

$$12a_0\lambda_0\dot{H} = d_7 \left(5W\dot{W}H + W^2\dot{H}\right),$$
 (55)

$$\lambda_3 H + \lambda_4 H W^2 + \lambda_5 W^4 H = \lambda_0 d_7 W \dot{W}.$$
 (56)

Now, Eq. (55) directly integrates to

$$H = C_1 \left( 12a_0\lambda_0 - d_7 W^2 \right)^{-\frac{5}{2}}.$$
 (57)

Substituting this in Eq. (56) and changing variables as

$$u = 12a_0\lambda_0 - d_7W^2, (58)$$

we obtain the differential equation

$$\dot{u} = -\frac{2C_1}{\lambda_0} \frac{(\tilde{\lambda}_1 + \tilde{\lambda}_4 u + \tilde{\lambda}_5 u^2)}{u^{\frac{5}{2}}}.$$
(59)

This is then integrated straightforwardly but its exact form depends crucially on the values of the parameters. For instance, if  $\tilde{\lambda}_1 = 0 = \tilde{\lambda}_5$ , it follows that  $u \propto t^{2/5}$  and consequently  $H \propto \frac{1}{t}$ , yielding a power-law behaviour for the scale factor.

**The case**  $k_5 = k_6$  — Let us now focus on the  $k_6 = k_5$  case. From Eq. (49) we find

$$W^2 = W_0^2 \equiv \frac{k_5}{2d_7}.$$
 (60)

First, note that under the conditions

$$6a_0 = d_{16}, \quad k_6 = k_5, \quad k_7 + k_9 = k_8, \quad k_5 + 2k_9 = k_8,$$
(61)

then the RHS of Eq. (51) is once again a perfect square and combining Eq. (51) with Eqs. (44b) and (60) we get an H = const. de-Sitter solution (even with  $d_7 \neq 0$ ).

Returning to the general case of  $k_6 = k_5$ , with Eq. (60) we find that Eq. (44b) directly integrates to

$$H = \frac{k_5}{24a_0}(X+Y) + \mathscr{C}_0,$$
(62)

with  $\mathscr{C}_0$  the integration constant. Then, for W = const. and taking also into account Eq. (48), we find that Eq. (44c) implies

$$k_{10}X + k_{11}Y = k_5H. ag{63}$$

Combining Eq. (63) with Eq. (62), we get

$$X = \lambda_1 H + X_0, \quad Y = \lambda_2 H + Y_0. \tag{64}$$

From Eq. (64) it follows that Eq. (48) can also be expressed as

$$V = \lambda_3 H + V_0, \quad Z = \lambda_4 H + Z_0.$$
 (65)

Geometrically it follows that in the configuration space of the cosmology  $\{H, X, Y, V, Z, W\}$  the  $k_6 = 0$  solutions are represented by lines lying on the W = const. hyperplane. Of course this is not the end of the story, if we plug the above results back to Eq. (45) we obtain the first order differential equation<sup>7</sup>

$$\dot{H} = \mu_2 H^2 + \mu_1 H + \mu_0. \tag{66}$$

Obviously, Eq. (66) is separable and can be readily integrated. However, the form of the solutions depends on the values  $\{\mu_i\}$  which are given in terms of the parameters of the theory and the integration constant. For  $\mu_2 = 0$  and  $\mu_1 \neq 0$  the solution reads

$$H = \frac{1}{\mu_1} \Big( C_1 e^{\mu_1 t} - \mu_0 \Big), \quad a = C_2 e^{\frac{1}{\mu_1} \Big( C_1 e^{\mu_1 t} - \mu_0 t \Big)}, \quad (67)$$

<sup>7</sup> Eq. (66) also known as a Riccati equation.

i.e. a super-exponential expansion. For  $\mu_2 \neq 0$  we can define  $\gamma \equiv \mu_1^2/4\mu_2 - \mu_0$  and shift  $\tilde{H} \equiv H + \mu_1/2\mu_2$ , then Eq. (66) is written as

$$\frac{\mathrm{d}\tilde{H}}{\tilde{H}^2 + \gamma/\mu_2} = \mu_2 \mathrm{d}t,\tag{68}$$

and of course the form of the solution to Eq. (68) depends on the ratio  $\gamma/\mu_2$ . For  $\gamma/\mu_2 < 0$  and defining  $\omega_0^2 \equiv |\gamma/\mu_2|$  we have

$$H = \omega_0 \frac{1 + C_1 e^{2\mu_2 \omega_0 t}}{1 - C_1 e^{2\mu_2 \omega_0 t}}, \quad a = \frac{C_2 e^{\omega_0 t}}{\left(1 - C_1 e^{2\mu_2 \omega_0 t}\right)^{1/\mu_2}}.$$
 (69)

Notice that for late times Eq. (69) always implies de Sitter or anti-de Sitter expansion depending on whether  $\mu_2 \omega_0$  is positive or negative. When  $\gamma/\mu_2 > 0$  the solution to Eq. (68) is

$$H = -\frac{\mu_1}{2\mu_2} + \omega_0 \tan\left(\mu_2 \omega_0 t + C_1\right).$$
 (70)

The integrable model contains too many unconstrained parameters for a thorough analysis of its particle spectrum. A severe restriction on the model set out in Eq. (43) is the case where all of the  $\{d_1, \ldots, d_8\}$  are set to zero. Whilst the Lagrangian density may still contain quadratic curvature operators in this case, they will not affect the cosmology at the background level. Since quadratic curvature operators endow the distortion with kinetic terms, it is interesting to consider the particle spectrum of such a model. The analogue in Riemannian cosmology would be Einstein-Maxwell theory: the Maxwell operator does not affect cosmology at the background level<sup>8</sup>, yet it still contributes two massless photon polarizations to the spectrum. For simplicity, we take the case wheere the  $\{a_1, \ldots, a_{11}\}$  are all taken to be zero, so that  $a_0$  is the only non-vanishing coupling of mass dimension two. As shown in Appendix C, this leads to a unitary model when some further constraints are imposed (concretely, sequential imposition of Eqs. (C7) to (C10) on Eq. (25)). According to the resulting particle spectrograph in Fig. 8, this unitarity is guaranteed by Eq. (C11), which suggests that  $c_{13} > 0$  is the noghost condition on the new species. By comparing Eq. (C11) with Eq. (25) we see that  $c_{13}$  modulates the square of the Homothetic curvature. We may therefore conjecture that, despite the presence of many other kinetic operators, the propagating part of the model is effectively Einstein-Maxwell theory, where the Weyl vector is massless. We note also in Fig. 8 the presence of *five* source constraints. This is expected: there are four gauge generators associated with diffeomeorphisms and one associated with the integrable (i.e. conservative) Weyl symmetry discussed in Section III C, which is common to all QMAG models built from the curvature alone [141].

## 11

### V. Concluding remarks

*Main findings* — The results of this work are as follows:

- Based on the general (i.e. spatially curved) Friedmann– Lemaître–Robertson–Walker line element in Eq. (41), we have derived the equations describing the background dynamics of the general parity-preserving quadratic metric-affine gravity model in Eq. (25). The full field equations are confined to the supplemental materials [144], where versions have been prepared for direct import into *SymPy*, *Mathematica* and *Maple* platforms. The simplified equations for the spatially flat case are on the edge of becoming human-readable, and so are formatted in Section III and Appendix B.
- Only the reduced set of 16 parameters in Eq. (30) are relevant to the background dynamics, down from the 28 Lagrangian couplings which parameterise Eq. (25).
- In terms of the reduced parameters in Eq. (30), the conditions in Eq. (42) completely eliminate the spatial curvature *K* from the field equations: this phenomenon is termed '*K*-screening'.
- As an alternative to *K*-screening, the conditions in Eq. (43) lead to simple equations which may be integrated analytically. This model always transitions to a de Sitter or anti-de Sitter epoch in the late Universe.

*Further work* — The following should be investigated:

- A straightforward extension of this analysis to parityviolating quadratic metric-affine gravity.
- The use of *K*-screening to construct models which alleviate current tensions in the dark-energy-cold-dark-matter (ACDM) model. Any serious attempt at solving these problems requires a complete understanding of the cosmological perturbation theory.
- The degree if any to which the parameter constraints identified in this work may lead to consistent models, i.e. those which are unitary and weakly coupled. The relevant tools in this endeavour include nonlinear Hamiltonian analysis and functional renormalisation group methods.

**Final note of caution** — We reiterate that the last point of further work is especially crucial: the analyses conducted here are superficial when compared to those which are needed to demonstrate consistency. Despite the colourful dynamics of the background cosmology, metric-affine gravity in general has a poor record in this regard, and to date only a very few special cases (mostly minimal extensions to general relativity itself) were shown to be consistent. It is perhaps telling that, starting with constraints motivated by the background cosmology, a systematic search for consistent weak-field particle spectra leads back to two such theories: the square-Holst theory (see Fig. 5) and the square-homothetic theory (see Fig. 8). It is conjectured in [19] that these *are* the only self-consistent kinetic operators in QMAG, though the point remains to be proven.

<sup>&</sup>lt;sup>8</sup> Of course, we know that radiative d.o.f actually dominate the background dynamics in the early Universe, but this is an average effect of locally anisotropic and inhomogeneous field configurations.

## Acknowledgments

We are grateful for correspondence with Will Handley and Tobias Mistele. This work is based in part on a Master's thesis by TD, which was itself improved by feedback from the assessor, Paul Rimmer.

This work used the DiRAC Data Intensive service (CSD3 www.csd3.cam.ac.uk) at the University of Cambridge, managed by the University of Cambridge University Information Services on behalf of the STFC DiRAC HPC Facility (www.dirac.ac.uk). The DiRAC component of CSD3 at Cam-

- F. W. Hehl, J. McCrea, E. W. Mielke, and Y. Ne'eman, Metricaffine gauge theory of gravity: field equations, noether identities, world spinors, and breaking of dilation invariance, Physics Reports 258, 10.1016/0370-1573(94)00111-f (1995).
- [2] D. Iosifidis, Metric-Affine Gravity and Cosmology/Aspects of Torsion and non-Metricity in Gravity Theories, Ph.D. thesis (2019), arXiv:1902.09643 [gr-qc].
- [3] D. Iosifidis, Cosmological hyperfluids, torsion and non-metricity, The European Physical Journal C 80, 10.1140/epjc/s10052-020-08634-z (2020).
- [4] R. Percacci and E. Sezgin, New class of ghost- and tachyon-free metric affine gravities, Physical Review D 101, 10.1103/physrevd.101.084040 (2020).
- [5] A. Jiménez Cano, Metric-affine Gauge theories of gravity. Foundations and new insights, Ph.D. thesis, Granada U., Theor. Phys. Astrophys. (2021), arXiv:2201.12847 [gr-qc].
- [6] O. A. Belarbi and A. Meziane, Overview and perspectives on metric-affine gravity, J. Phys. Conf. Ser. 1766, 012007 (2021).
- [7] A. Baldazzi, O. Melichev, and R. Percacci, Metric-Affine Gravity as an effective field theory, Annals Phys. 438, 168757 (2022), arXiv:2112.10193 [gr-qc].
- [8] M. Blagojevic and F. W. Hehl, Gauge Theories of Gravitation, (2012), arXiv:1210.3775 [gr-qc].
- [9] A. Einstein, Die Feldgleichungen der Gravitation, Sitzungsber. Preuss. Akad. Wiss 18, 844 (1915).
- [10] F. W. Hehl, G. D. Kerlick, and P. von der Heyde, On hypermomentum in general relativity i. the notion of hypermomentum, Zeitschrift für Naturforschung A 31, 111 (1976).
- [11] F. W. Hehl, G. D. Kerlick, and P. von der Heyde, On hypermomentum in general relativity ii. the geometry of spacetime, Zeitschrift für Naturforschung A 31, 524 (1976).
- [12] F. W. Hehl, G. D. Kerlick, and P. von der Heyde, On hypermomentum in general relativity iii. coupling hypermomentum to geometry, Zeitschrift für Naturforschung A 31, 823 (1976).
- [13] D. Iosifidis, Non-Riemannian cosmology: The role of shear hypermomentum, Int. J. Geom. Meth. Mod. Phys. 18, 2150129 (2021), arXiv:2010.00875 [gr-qc].
- [14] D. Iosifidis and T. S. Koivisto, Hyperhydrodynamics: relativistic viscous fluids from hypermomentum, JCAP 05, 001, arXiv:2312.06780 [gr-qc].
- [15] A. D. Sakharov, Vacuum quantum fluctuations in curved space and the theory of gravitation, Soviet Physics Uspekhi 34, 394 (1991).
- [16] F. Hehl, G. Kerlick, and P. Von der Heyde, On a new metric affine theory of gravitation, Physics Letters B 63, 446 (1976).
- [17] C. Rigouzzo and S. Zell, Coupling metric-affine gravity to the standard model and dark matter fermions, Phys. Rev. D 108,

bridge was funded by BEIS, UKRI and STFC capital funding and STFC operations grants. DiRAC is part of the UKRI Digital Research Infrastructure.

This work also used the Newton server, access to which was provisioned by Will Handley using an ERC grant.

WB is grateful for the support of Marie Skłodowska-Curie Actions and the Institute of Physics of the Czech Academy of Sciences.

The work of DI was supported by the Estonian Research Council via the Center of Excellence '*Foundations of the Uni*verse' (TK202U4).

124067 (2023), arXiv:2306.13134 [gr-qc].

- [18] W. Barker and C. Marzo, Particle spectra of general ricci-type palatini or metric-affine theories (2024), arXiv:2402.07641 [hep-th].
- [19] W. Barker and S. Zell, Consistent particle physics in metricaffine gravity from extended projective symmetry, (2024), arXiv:2402.14917 [hep-th].
- [20] A. A. Starobinsky, Spectrum of relict gravitational radiation and the early state of the universe, JETP Lett. 30, 682 (1979).
- [21] A. H. Guth, The Inflationary Universe: A Possible Solution to the Horizon and Flatness Problems, Phys. Rev. D 23, 347 (1981).
- [22] F. Zwicky, Die Rotverschiebung von extragalaktischen Nebeln, Helv. Phys. Acta 6, 110 (1933).
- [23] N. Aghanim *et al.* (Planck), Planck 2018 results. VI. Cosmological parameters, Astron. Astrophys. **641**, A6 (2020), [Erratum: Astron.Astrophys. 652, C4 (2021)], arXiv:1807.06209 [astro-ph.CO].
- [24] D. Scott, The standard model of cosmology: A skeptic's guide, Proc. Int. Sch. Phys. Fermi 200, 133 (2020), arXiv:1804.01318 [astro-ph.CO].
- [25] W. Handley, Curvature tension: evidence for a closed universe, Phys. Rev. D 103, L041301 (2021), arXiv:1908.09139 [astroph.CO].
- [26] E. Di Valentino, O. Mena, S. Pan, L. Visinelli, W. Yang, A. Melchiorri, D. F. Mota, A. G. Riess, and J. Silk, In the realm of the Hubble tension—a review of solutions, Class. Quant. Grav. 38, 153001 (2021), arXiv:2103.01183 [astro-ph.CO].
- [27] L. Perivolaropoulos and F. Skara, Challenges for ACDM: An update, New Astron. Rev. 95, 101659 (2022), arXiv:2105.05208 [astro-ph.CO].
- [28] S. Bahamonde, K. F. Dialektopoulos, C. Escamilla-Rivera, G. Farrugia, V. Gakis, M. Hendry, M. Hohmann, J. Levi Said, J. Mifsud, and E. Di Valentino, Teleparallel gravity: from theory to cosmology, Rept. Prog. Phys. 86, 026901 (2023), arXiv:2106.13793 [gr-qc].
- [29] J. Beltrán Jiménez, L. Heisenberg, D. Iosifidis, A. Jiménez-Cano, and T. S. Koivisto, General teleparallel quadratic gravity, Phys. Lett. B 805, 135422 (2020), arXiv:1909.09045 [grqc].
- [30] J. B. Jimenez, L. Heisenberg, and T. S. Koivisto, The geometrical trinity of gravity (2019), arXiv:1903.06830 [hep-th].
- [31] H. Weyl, Gravitation and electricity, Sitzungsber. Preuss. Akad. Wiss. **26**, 465 (1918).
- [32] H. Weyl, Space-Time-Matter (Dover Publications, 1922).
- [33] É. Cartan, Sur les variétés à connexion affine et la théorie de la relativité généralisée (première partie), in Annales sci-

entifiques de l'École normale supérieure, Vol. 40 (Gauthier-Villars, 1923) pp. 325-412.

- [34] A. S. Eddington, *The Mathematical Theory of Relativity* (Cambridge University Press, 1923).
- [35] É. Cartan, Sur les variétés à connexion affine, et la théorie de la relativité généralisée (première partie)(suite), in Annales scientifiques de l'École Normale Supérieure, Vol. 41 (Gauthier-Villars, 1924) pp. 1–25.
- [36] É. Cartan, Sur les variétés à connexion affine, et la théorie de la relativité généralisée (deuxième partie), in Annales scientifiques de l'École normale supérieure, Vol. 42 (Gauthier-Villars, 1925) pp. 17–88.
- [37] A. Einstein, Einheitliche Feldtheorie von Gravitation und Elektrizität, Sitzungsber. Preuss. Akad. Wiss 22, 414 (1925).
- [38] A. Einstein, Riemanngeometrie mit Aufrechterhaltung des Begriffes des Fern-Parallelismus, Sitzungsber. Preuss. Akad. Wiss 17, 217 (1928).
- [39] A. Einstein, Neue Möglichkeit für eine einheitliche Feldtheorie von Gravitation und Elektrizität, Sitzungsber. Preuss. Akad. Wiss 18, 224 (1928).
- [40] R. Utiyama, Invariant theoretical interpretation of interaction, Phys. Rev. 101, 1597 (1956).
- [41] D. W. Sciama, On the analogy between charge and spin in general relativity, in *Recent developments in general relativity* (Pergamon Press, Oxford, 1962) p. 415.
- [42] T. W. B. Kibble, Lorentz invariance and the gravitational field, J. Math. Phys. 2, 212 (1961).
- [43] M. Blagojevic, Gravitation and gauge symmetries (2002).
- [44] A. Salvio, Quadratic Gravity, Front. in Phys. 6, 77 (2018), arXiv:1804.09944 [hep-th].
- [45] A. Salam and J. Strathdee, Remarks on high-energy stability and renormalizability of gravity theory, Phys. Rev. D 18, 4480 (1978).
- [46] K. S. Stelle, Renormalization of higher-derivative quantum gravity, Phys. Rev. D 16, 953 (1977).
- [47] M. B. Einhorn and D. R. T. Jones, Renormalizable, asymptotically free gravity without ghosts or tachyons, Phys. Rev. D 96, 124025 (2017), arXiv:1710.03795 [hep-th].
- [48] J. F. Donoghue, General relativity as an effective field theory: The leading quantum corrections, Phys. Rev. D 50, 3874 (1994), arXiv:gr-qc/9405057.
- [49] C. Pagani and R. Percacci, Quantum gravity with torsion and non-metricity, Class. Quant. Grav. 32, 195019 (2015), arXiv:1506.02882 [gr-qc].
- [50] R. Percacci, Towards Metric-Affine Quantum Gravity, Int. J. Geom. Meth. Mod. Phys. 17, 2040003 (2020), arXiv:2003.09486 [gr-qc].
- [51] O. Melichev and R. Percacci, On the renormalization of Poincaré gauge theories, JHEP 03, 133, arXiv:2307.02336 [hep-th].
- [52] O. Melichev, On the renormalization of Metric-Affine Gravity theories, (2024), arXiv:2406.14146 [hep-th].
- [53] E. Sezgin, Class of Ghost Free Gravity Lagrangians With Massive or Massless Propagating Torsion, Phys. Rev. D 24, 1677 (1981).
- [54] M. Blagojevic and I. A. Nikolic, Hamiltonian dynamics of Poincare gauge theory: General structure in the time gauge, Phys. Rev. D 28, 2455 (1983).
- [55] M. Blagojevic and M. Vasilic, EXTRA GAUGE SYMME-TRIES IN A WEAK FIELD APPROXIMATION OF AN R + T\*\*2 + R\*\*2 THEORY OF GRAVITY, Phys. Rev. D 35, 3748 (1987).
- [56] R. Kuhfuss and J. Nitsch, Propagating Modes in Gauge Field Theories of Gravity, Gen. Rel. Grav. 18, 1207 (1986).

- [57] H.-j. Yo and J. M. Nester, Hamiltonian analysis of Poincare gauge theory scalar modes, Int. J. Mod. Phys. D 8, 459 (1999), arXiv:gr-qc/9902032.
- [58] H.-J. Yo and J. M. Nester, Hamiltonian analysis of Poincare gauge theory: Higher spin modes, Int. J. Mod. Phys. D 11, 747 (2002), arXiv:gr-qc/0112030.
- [59] M. Blagojevic, Gravitation and gauge symmetries (CRC Press, Bristol and Philadelphia, 2002).
- [60] D. Puetzfeld, Status of non-Riemannian cosmology, New Astron. Rev. 49, 59 (2005), arXiv:gr-qc/0404119.
- [61] H.-J. Yo and J. M. Nester, Dynamic Scalar Torsion and an Oscillating Universe, Mod. Phys. Lett. A 22, 2057 (2007), arXiv:astro-ph/0612738.
- [62] K.-F. Shie, J. M. Nester, and H.-J. Yo, Torsion Cosmology and the Accelerating Universe, Phys. Rev. D 78, 023522 (2008), arXiv:0805.3834 [gr-qc].
- [63] V. P. Nair, S. Randjbar-Daemi, and V. Rubakov, Massive Spin-2 fields of Geometric Origin in Curved Spacetimes, Phys. Rev. D 80, 104031 (2009), arXiv:0811.3781 [hep-th].
- [64] V. Nikiforova, S. Randjbar-Daemi, and V. Rubakov, Infrared Modified Gravity with Dynamical Torsion, Phys. Rev. D 80, 124050 (2009), arXiv:0905.3732 [hep-th].
- [65] H. Chen, F.-H. Ho, J. M. Nester, C.-H. Wang, and H.-J. Yo, Cosmological dynamics with propagating Lorentz connection modes of spin zero, JCAP 10, 027, arXiv:0908.3323 [gr-qc].
- [66] W.-T. Ni, Searches for the role of spin and polarization in gravity, Rept. Prog. Phys. 73, 056901 (2010), arXiv:0912.5057 [grqc].
- [67] P. Baekler, F. W. Hehl, and J. M. Nester, Poincare gauge theory of gravity: Friedman cosmology with even and odd parity modes. Analytic part, Phys. Rev. D 83, 024001 (2011), arXiv:1009.5112 [gr-qc].
- [68] F.-H. Ho and J. M. Nester, Poincaré gauge theory with even and odd parity dynamic connection modes: isotropic Bianchi cosmological models, J. Phys. Conf. Ser. **330**, 012005 (2011), arXiv:1105.5001 [gr-qc].
- [69] F.-H. Ho and J. M. Nester, Poincaré Gauge Theory With Coupled Even And Odd Parity Dynamic Spin-0 Modes: Dynamic Equations For Isotropic Bianchi Cosmologies, Annalen Phys. 524, 97 (2012), arXiv:1106.0711 [gr-qc].
- [70] Y. C. Ong, K. Izumi, J. M. Nester, and P. Chen, Problems with Propagation and Time Evolution in f(T) Gravity, Phys. Rev. D 88, 024019 (2013), arXiv:1303.0993 [gr-qc].
- [71] D. Puetzfeld and Y. N. Obukhov, Prospects of detecting spacetime torsion, Int. J. Mod. Phys. D 23, 1442004 (2014), arXiv:1405.4137 [gr-qc].
- [72] G. K. Karananas, The particle spectrum of parity-violating Poincaré gravitational theory, Class. Quant. Grav. 32, 055012 (2015), arXiv:1411.5613 [gr-qc].
- [73] W.-T. Ni, Searches for the role of spin and polarization in gravity: a five-year update, Int. J. Mod. Phys. Conf. Ser. 40, 1660010 (2016), arXiv:1501.07696 [hep-ph].
- [74] F.-H. Ho, H. Chen, J. M. Nester, and H.-J. Yo, General Poincaré Gauge Theory Cosmology, Chin. J. Phys. 53, 110109 (2015), arXiv:1512.01202 [gr-qc].
- [75] G. K. Karananas, Poincaré, Scale and Conformal Symmetries Gauge Perspective and Cosmological Ramifications, Ph.D. thesis, Ecole Polytechnique, Lausanne (2016), arXiv:1608.08451 [hep-th].
- [76] Y. N. Obukhov, Gravitational waves in Poincaré gauge gravity theory, Phys. Rev. D 95, 084028 (2017), arXiv:1702.05185 [gr-qc].
- [77] M. Blagojević, B. Cvetković, and Y. N. Obukhov, Generalized plane waves in Poincaré gauge theory of gravity, Phys. Rev. D

96, 064031 (2017), arXiv:1708.08766 [gr-qc].

- [78] M. Blagojević and B. Cvetković, General Poincaré gauge theory: Hamiltonian structure and particle spectrum, Phys. Rev. D 98, 024014 (2018), arXiv:1804.05556 [gr-qc].
- [79] H.-H. Tseng, Gravitational Theories with Torsion, Ph.D. thesis, Taiwan, Natl. Tsing Hua U. (2018), arXiv:1812.00314 [grqc].
- [80] Y.-C. Lin, M. P. Hobson, and A. N. Lasenby, Ghost and tachyon free Poincaré gauge theories: A systematic approach, Phys. Rev. D 99, 064001 (2019), arXiv:1812.02675 [gr-qc].
- [81] J. Beltrán Jiménez and A. Delhom, Ghosts in metric-affine higher order curvature gravity, Eur. Phys. J. C 79, 656 (2019), arXiv:1901.08988 [gr-qc].
- [82] H. Zhang and L. Xu, Late-time acceleration and inflation in a Poincaré gauge cosmological model, JCAP 09, 050, arXiv:1904.03545 [gr-qc].
- [83] K. Aoki and K. Shimada, Scalar-metric-affine theories: Can we get ghost-free theories from symmetry?, Phys. Rev. D 100, 044037 (2019), arXiv:1904.10175 [hep-th].
- [84] H. Zhang and L. Xu, Inflation in the parity-conserving Poincaré gauge cosmology, JCAP 10, 003, arXiv:1906.04340 [gr-qc].
- [85] J. Beltrán Jiménez and F. J. Maldonado Torralba, Revisiting the stability of quadratic Poincaré gauge gravity, Eur. Phys. J. C 80, 611 (2020), arXiv:1910.07506 [gr-qc].
- [86] Y.-C. Lin, M. P. Hobson, and A. N. Lasenby, Power-counting renormalizable, ghost-and-tachyon-free Poincaré gauge theories, Phys. Rev. D 101, 064038 (2020), arXiv:1910.14197 [grqc].
- [87] R. Percacci and E. Sezgin, New class of ghost- and tachyonfree metric affine gravities, Phys. Rev. D 101, 084040 (2020), arXiv:1912.01023 [hep-th].
- [88] W. E. V. Barker, A. N. Lasenby, M. P. Hobson, and W. J. Handley, Systematic study of background cosmology in unitary Poincaré gauge theories with application to emergent dark radiation and  $H_0$  tension, Phys. Rev. D **102**, 024048 (2020), arXiv:2003.02690 [gr-qc].
- [89] J. Beltrán Jiménez and A. Delhom, Instabilities in metric-affine theories of gravity with higher order curvature terms, Eur. Phys. J. C 80, 585 (2020), arXiv:2004.11357 [gr-qc].
- [90] F. J. Maldonado Torralba, New effective theories of gravitation and their phenomenological consequences, Ph.D. thesis, Cape Town U., Dept. Math. (2020), arXiv:2101.11523 [gr-qc].
- [91] W. E. V. Barker, A. N. Lasenby, M. P. Hobson, and W. J. Handley, Nonlinear Hamiltonian analysis of new quadratic torsion theories: Cases with curvature-free constraints, Phys. Rev. D 104, 084036 (2021), arXiv:2101.02645 [gr-qc].
- [92] C. Marzo, Ghost and tachyon free propagation up to spin 3 in Lorentz invariant field theories, Phys. Rev. D 105, 065017 (2022), arXiv:2108.11982 [hep-ph].
- [93] C. Marzo, Radiatively stable ghost and tachyon freedom in metric affine gravity, Phys. Rev. D 106, 024045 (2022), arXiv:2110.14788 [hep-th].
- [94] A. de la Cruz Dombriz, F. J. Maldonado Torralba, and D. F. Mota, Dark matter candidate from torsion, Phys. Lett. B 834, 137488 (2022), arXiv:2112.03957 [gr-qc].
- [95] J. Annala and S. Rasanen, Stability of non-degenerate Riccitype Palatini theories, JCAP 04, 014, [Erratum: JCAP 08, E02 (2023)], arXiv:2212.09820 [gr-qc].
- [96] Y. Mikura, V. Naso, and R. Percacci, Some simple theories of gravity with propagating torsion, Phys. Rev. D 109, 104071 (2024), arXiv:2312.10249 [gr-qc].
- [97] Y. Mikura and R. Percacci, Some simple theories of gravity with propagating nonmetricity, (2024), arXiv:2401.10097 [gr-

qc].

- [98] W. Barker and C. Marzo, Particle spectra of general Ricci-type Palatini or metric-affine theories, Phys. Rev. D 109, 104017 (2024), arXiv:2402.07641 [hep-th].
- [99] C. Møller, Conservation Law and Absolute Parallelism in General Relativity, K. Dan. Vidensk. Selsk. Mat. Fys. Skr. 1, 1 (1961).
- [100] C. Pellegrini and J. Plebanski, Tetrad Fields and Gravitational Fields, K. Dan. Vidensk. Selsk. Mat. Fys. Skr. 2, 1 (1963).
- [101] K. Hayashi and T. Nakano, Extended translation invariance and associated gauge fields, Prog. Theor. Phys. 38, 491 (1967).
- [102] Y. M. Cho, Einstein Lagrangian as the Translational Yang-Mills Lagrangian, Phys. Rev. D 14, 2521 (1976).
- [103] K. Hayashi and T. Shirafuji, New general relativity., Phys. Rev. D 19, 3524 (1979), [Addendum: Phys.Rev.D 24, 3312–3314 (1982)].
- [104] A. Dimakis, The Initial Value Problem of the Poincare Gauge Theory in Vacuum. 1: Second Order Formalism, Ann. Inst. H. Poincare Phys. Theor. **51**, 371 (1989).
- [105] A. Dimakis, THE INITIAL VALUE PROBLEM OF THE POINCARE GAUGE THEORY IN VACUUM. 1: FIRST OR-DER FORMALISM, Ann. Inst. H. Poincare Phys. Theor. 51, 389 (1989).
- [106] J. Lemke, Shock waves in the Poincare gauge theory of gravitation, Phys. Lett. A 143, 13 (1990).
- [107] R. D. Hecht, J. Lemke, and R. P. Wallner, Tachyonic torsion shock waves in Poincare gauge theory, Phys. Lett. A 151, 12 (1990).
- [108] R. D. Hecht, J. Lemke, and R. P. Wallner, Can Poincare gauge theory be saved?, Phys. Rev. D 44, 2442 (1991).
- [109] N. Afshordi, D. J. H. Chung, and G. Geshnizjani, Cuscuton: A Causal Field Theory with an Infinite Speed of Sound, Phys. Rev. D 75, 083513 (2007), arXiv:hep-th/0609150.
- [110] J. Magueijo, Bimetric varying speed of light theories and primordial fluctuations, Phys. Rev. D 79, 043525 (2009), arXiv:0807.1689 [gr-qc].
- [111] C. Charmousis and A. Padilla, The Instability of Vacua in Gauss-Bonnet Gravity, JHEP **12**, 038, arXiv:0807.2864 [hep-th].
- [112] C. Charmousis, G. Niz, A. Padilla, and P. M. Saffin, Strong coupling in Horava gravity, JHEP 08, 070, arXiv:0905.2579 [hep-th].
- [113] A. Papazoglou and T. P. Sotiriou, Strong coupling in extended Horava-Lifshitz gravity, Phys. Lett. B 685, 197 (2010), arXiv:0911.1299 [hep-th].
- [114] D. Baumann, L. Senatore, and M. Zaldarriaga, Scale-Invariance and the Strong Coupling Problem, JCAP 05, 004, arXiv:1101.3320 [hep-th].
- [115] G. D'Amico, C. de Rham, S. Dubovsky, G. Gabadadze, D. Pirtskhalava, and A. J. Tolley, Massive Cosmologies, Phys. Rev. D 84, 124046 (2011), arXiv:1108.5231 [hep-th].
- [116] A. E. Gumrukcuoglu, C. Lin, and S. Mukohyama, Anisotropic Friedmann-Robertson-Walker universe from nonlinear massive gravity, Phys. Lett. B 717, 295 (2012), arXiv:1206.2723 [hep-th].
- [117] A. Wang, Hořava gravity at a Lifshitz point: A progress report, Int. J. Mod. Phys. D 26, 1730014 (2017), arXiv:1701.06087 [gr-qc].
- [118] C. Mazuet, S. Mukohyama, and M. S. Volkov, Anisotropic deformations of spatially open cosmology in massive gravity theory, JCAP 04, 039, arXiv:1702.04205 [hep-th].
- [119] J. Beltrán Jiménez and A. Jiménez-Cano, On the strong coupling of Einsteinian Cubic Gravity and its generalisations, JCAP 01, 069, arXiv:2009.08197 [gr-qc].

- [120] W. E. V. Barker, Supercomputers against strong coupling in gravity with curvature and torsion, Eur. Phys. J. C 83, 228 (2023), arXiv:2206.00658 [gr-qc].
- [121] A. Delhom, A. Jiménez-Cano, and F. J. Maldonado Torralba, Instabilities in Field Theory: A Primer with Applications in Modified Gravity (Springer, 2022) arXiv:2207.13431 [gr-qc].
- [122] W. Barker and S. Zell, Einstein-Proca theory from the Einstein-Cartan formulation, Phys. Rev. D 109, 024007 (2024), arXiv:2306.14953 [hep-th].
- [123] D. Iosifidis and K. Pallikaris, Into the mag-verse or: Cosmology of the complete quadratic metric-affine gravity (2024), arXiv:2404.19498 [gr-qc].
- [124] K. Shimada, K. Aoki, and K.-i. Maeda, Metric-affine Gravity and Inflation, Phys. Rev. D 99, 104020 (2019), arXiv:1812.03420 [gr-qc].
- [125] I. D. Gialamas and K. Tamvakis, Inflation in metric-affine quadratic gravity, JCAP 03, 042, arXiv:2212.09896 [gr-qc].
- [126] A. V. Minkevich and A. S. Garkun, Isotropic cosmology in metric-affine gauge theory of gravity (1998), arXiv:grqc/9805007 [gr-qc].
- [127] Y. N. Obukhov, E. J. Vlachynsky, W. Esser, and F. W. Hehl, Irreducible decompositions in metric-affine gravity models, (1997), arXiv:gr-qc/9705039 [gr-qc].
- [128] D. Iosifidis and L. Ravera, Cosmology of quadratic metricaffine gravity, Physical Review D 105, 10.1103/physrevd.105.024007 (2022).
- [129] D. Iosifidis, The full quadratic metric-affine gravity (including parity odd terms): exact solutions for the affine-connection, Class. Quant. Grav. **39**, 095002 (2022), arXiv:2112.09154 [grqc].
- [130] M. Tsamparlis, Cosmological principle and torsion, Physics Letters A 75, 27 (1979).
- [131] D. Iosifidis, Linear Transformations on Affine-Connections, Class. Quant. Grav. 37, 085010 (2020), arXiv:1911.04535 [grqc].
- [132] G. N. Remmen and S. M. Carroll, Attractor Solutions in Scalar-Field Cosmology, Phys. Rev. D 88, 083518 (2013), arXiv:1309.2611 [gr-qc].
- [133] J. B. Achour, Proper time reparametrization in cosmology: Möbius symmetry and Kodama charges, JCAP 12 (12), 005, arXiv:2103.10700 [gr-qc].
- [134] S. Deser and B. Tekin, Shortcuts to high symmetry solutions in gravitational theories, Class. Quant. Grav. 20, 4877 (2003), arXiv:gr-qc/0306114.

- [135] M. A. H. Maccallum and A. H. Taub, Variational principles and spatially-homogeneous universes, including rotation, Commun. Math. Phys. 25, 173 (1972).
- [136] M. Fels and C. Torre, The principle of symmetric criticality in general relativity, Classical and Quantum Gravity 19 (2001).
- [137] S. Hawking, On the Rotation of the Universe, Monthly Notices of the Royal Astronomical Society 142, 129 (1969).
- [138] D. A. Cox, J. B. Little, and D. O'Shea, *Ideals, varieties, and algorithms : an introduction to computational algebraic geometry and commutative algebra*, 1st ed., Undergraduate texts in mathematics (Springer New York, 1992).
- [139] D. R. Grayson and M. E. Stillman, Macaulay2, a software system for research in algebraic geometry, Available at http: //www2.macaulay2.com.
- [140] S. Kolekar, On the Bianchi identity in generalized theories of gravity, Gen. Rel. Grav. 54, 92 (2022).
- [141] D. Iosifidis and T. Koivisto, Scale transformations in metricaffine geometry, Universe 5, 82 (2019), arXiv:1810.12276 [grqc].
- [142] T. Koivisto, M. Hohmann, and T. Złośnik, The General Linear Cartan Khronon, Universe 5, 168 (2019), arXiv:1905.02967 [gr-qc].
- [143] W. Barker, Particle Spectrum for Any Tensor Lagrangian (PSALTer), download: github.com/wevbarker/PSALTer.
- [144] T. Dyer, D. Iosifidis, and W. Barker, *Supplemental materials* at www.github.com/wevbarker/SupplementalMaterials-2411.
- [145] W. E. V. Barker, A. N. Lasenby, M. P. Hobson, and W. J. Handley, Mapping Poincaré gauge cosmology to Horndeski theory for emergent dark energy, Phys. Rev. D 102, 084002 (2020), arXiv:2006.03581 [gr-qc].
- [146] C. Rew and W. E. V. Barker, The effective inflationary potential of constant-torsion emergent gravity, (2023), arXiv:2302.07250 [gr-qc].
- [147] W. Barker, Particle spectra of gravity based on internal symmetry of quantum fields, (2023), arXiv:2311.11790 [hep-th].
- [148] W. Barker, C. Marzo, and C. Rigouzzo, PSALTer: Particle Spectrum for Any Tensor Lagrangian, (2024), arXiv:2406.09500 [hep-th].
- [149] W. Barker, M. Hobson, A. Lasenby, Y.-C. Lin, and Z. Wei, Every Poincaré gauge theory is conformal: a compelling case for dynamical vector torsion, (2024), arXiv:2406.12826 [hepth].

### A. Quadratic and cubic relations

**Distortion invariants** — In this appendix we provide additional algebraic identities obeyed by the quadratic invariants in Eq. (25) in a cosmological setting. These take the form of identically vanishing sums of squares or cubes of the quadratic invariants formed from the curvature, torsion and non-metricity tensors. The means for obtaining these relations is discussed in Section III A. There is precisely one relation of fourth order in the non-metricity, i.e. quadratic in the quadratic non-metricity invariants

$$\mathscr{A}_6 \,\mathscr{A}_7 \, - \,\mathscr{A}_8^{\ 2} = 0. \tag{A1}$$

*Curvature invariants* — Meanwhile, there are two relations of fourth order in the curvature, and one sixth order relation

$$\begin{aligned} & \mathcal{C}_{1}^{2} - 2\mathcal{C}_{1}\mathcal{C}_{3} - \mathcal{C}_{1}\mathcal{C}_{4} - 2\mathcal{C}_{1}\mathcal{C}_{5} - \mathcal{C}_{1}\mathcal{C}_{6} - \mathcal{C}_{2}^{2} - 2\mathcal{C}_{2}\mathcal{C}_{3} + \mathcal{C}_{2}\mathcal{C}_{4} + 2\mathcal{C}_{2}\mathcal{C}_{5} + \mathcal{C}_{2}\mathcal{C}_{6} + 2\mathcal{C}_{3}\mathcal{C}_{4} + 4\mathcal{C}_{3}\mathcal{C}_{5} + 2\mathcal{C}_{3}\mathcal{C}_{6} \\ & - 2\mathcal{C}_{4}^{2} + 4\mathcal{C}_{4}\mathcal{C}_{6} - 2\mathcal{C}_{6}^{2} = 0, \end{aligned}$$

$$\begin{aligned} & (A2) \\ & \mathcal{C}_{1}\mathcal{C}_{3} - 4\mathcal{C}_{1}\mathcal{C}_{10} - 6\mathcal{C}_{1}\mathcal{C}_{12} + 3\mathcal{C}_{1}\mathcal{C}_{2} + 3\mathcal{C}_{1}\mathcal{C}_{4} + 8\mathcal{C}_{1}\mathcal{C}_{5} + 3\mathcal{C}_{1}\mathcal{C}_{6} + 4\mathcal{C}_{10}^{2} + 8\mathcal{C}_{10}\mathcal{C}_{12} - 4\mathcal{C}_{10}\mathcal{C}_{2} - 4\mathcal{C}_{10}\mathcal{C}_{4} - 8\mathcal{C}_{10}\mathcal{C}_{5} \end{aligned}$$

$$- 4\mathscr{C}_{10}\mathscr{C}_{6} + 4\mathscr{C}_{12}^{2} - 6\mathscr{C}_{12}\mathscr{C}_{2} - 6\mathscr{C}_{12}\mathscr{C}_{4} - 12\mathscr{C}_{12}\mathscr{C}_{5} - 6\mathscr{C}_{12}\mathscr{C}_{6} + 3\mathscr{C}_{2}^{2} + \mathscr{C}_{2}\mathscr{C}_{3} + 2\mathscr{C}_{2}\mathscr{C}_{4} + 6\mathscr{C}_{2}\mathscr{C}_{5} + 2\mathscr{C}_{2}\mathscr{C}_{6} - 3\mathscr{C}_{3}\mathscr{C}_{4} \\ - 6\mathscr{C}_{3}\mathscr{C}_{5} - 3\mathscr{C}_{3}\mathscr{C}_{6} + 3\mathscr{C}_{4}^{2} + 6\mathscr{C}_{4}\mathscr{C}_{5} - 2\mathscr{C}_{4}\mathscr{C}_{6} + 8\mathscr{C}_{5}^{2} + 6\mathscr{C}_{5}\mathscr{C}_{6} + 3\mathscr{C}_{6}^{2} = 0,$$
(A3)
$$8\mathscr{C}_{1}\mathscr{C}_{10}\mathscr{C}_{5} - 3\mathscr{C}_{1}\mathscr{C}_{10}^{2} - 6\mathscr{C}_{1}\mathscr{C}_{10}\mathscr{C}_{12} + 4\mathscr{C}_{1}\mathscr{C}_{10}\mathscr{C}_{4} + 4\mathscr{C}_{1}\mathscr{C}_{10}\mathscr{C}_{6} - 3\mathscr{C}_{1}\mathscr{C}_{12}^{2} + 6\mathscr{C}_{1}\mathscr{C}_{12}\mathscr{C}_{4} + 12\mathscr{C}_{1}\mathscr{C}_{12}\mathscr{C}_{5} + 6\mathscr{C}_{1}\mathscr{C}_{12}\mathscr{C}_{6} \\ + 2\mathscr{C}_{1}\mathscr{C}_{3}\mathscr{C}_{4} + 4\mathscr{C}_{1}\mathscr{C}_{3}\mathscr{C}_{5} + 2\mathscr{C}_{10}^{2}\mathscr{C}_{6} - 3\mathscr{C}_{1}\mathscr{C}_{4}^{2} - 8\mathscr{C}_{1}\mathscr{C}_{4}\mathscr{C}_{5} - 10\mathscr{C}_{1}\mathscr{C}_{5}^{2} - 8\mathscr{C}_{1}\mathscr{C}_{5} - 3\mathscr{C}_{1}\mathscr{C}_{6}^{2} + 3\mathscr{C}_{10}^{2}\mathscr{C}_{2} + 6\mathscr{C}_{10}^{2}\mathscr{C}_{3} \\ + 2\mathscr{C}_{1}^{2}\mathscr{C}_{3}\mathscr{C}_{4} + 4\mathscr{C}_{10}^{2}\mathscr{C}_{5} + 2\mathscr{C}_{10}^{2}\mathscr{C}_{6} - 3\mathscr{C}_{1}\mathscr{C}_{4}^{2} - 3\mathscr{C}_{1}\mathscr{C}_{6}^{2} - 3\mathscr{C}_{1}\mathscr{C}_{6}^{2} + 3\mathscr{C}_{1}\mathscr{C}_{2} + 4\mathscr{C}_{1}\mathscr{C}_{5} - 8\mathscr{C}_{1}\mathscr{C}_{5}^{2} - 8\mathscr{C}_{1}\mathscr{C}_{5}^{2} - 3\mathscr{C}_{1}\mathscr{C}_{6}^{2} - 3\mathscr{C}_{1}\mathscr{C}_{6}^{2} + 3\mathscr{C}_{1}\mathscr{C}_{6} - 3\mathscr{C}_{1}\mathscr{C}_{6}^{2} - 3\mathscr{C}_{1}\mathscr{C}_{6}^{2} + 3\mathscr{C}_{1}\mathscr{C}_{2} + 6\mathscr{C}_{1}\mathscr{C}_{2}^{2} + 12\mathscr{C}_{1}\mathscr{C}_{1}\mathscr{C}_{4} + 4\mathscr{C}_{1}\mathscr{C}_{1}\mathscr{C}_{5} - 3\mathscr{C}_{1}\mathscr{C}_{6}^{2} - 3\mathscr{C}_{1}\mathscr{C}_{6}^{2} - 3\mathscr{C}_{1}\mathscr{C}_{6}^{2} - 3\mathscr{C}_{1}\mathscr{C}_{6}^{2} - 3\mathscr{C}_{1}\mathscr{C}_{1}^{2} - 3\mathscr{C}_{1}^{2} - 3\mathscr{C}_{$$

## B. The field equations in full

Friedmann equations — In this appendix we display the background field equations of Eq. (25) subject only to the condition K = 0 of vanishing spatial curvature, corresponding to Eq. (11) — the full equations, with general  $K \neq 0$  corresponding to Eq. (41), are made available in the supplemental materials [144]. The means by which these equations are obtained is described in Section III B. The first and second modified Friedmann equations are respectively:

$$\begin{aligned} 6a_0H^2 + d_1\left(2H\ddot{H} + 6H^2\dot{H} - \dot{H}^2\right) + \left(\mathcal{A}_{ij}N_i\dot{N}_j + \ddot{\Xi}\right)H + \left(\frac{1}{2}\mathcal{B}_{ij}N_iN_j + f_1(\dot{N})\right)H^2 + 3f_2(N)H^3 \\ &- \dot{\Xi}\dot{H} + \frac{1}{2}\mathcal{C}_{ij}\dot{N}_i\dot{N}_j + \frac{1}{2}\mathcal{G}_{ij}N_iN_j + F_1(N^4) = 0, \end{aligned} \tag{B1} \\ 12a_0\dot{H} + 2d_1\left(2\dot{H}^2 - H\ddot{H}\right) + \left(\mathcal{D}_{ij}N_i\dot{N}_j - \ddot{\Xi}\right)H + \left(\frac{1}{2}\mathcal{M}_{ij}N_iN_j + 4\dot{\Xi}\right)\dot{H} - f_2(\dot{N})H^2 + f_2(N)H\dot{H} \\ &- \Xi\ddot{H} + \mathcal{K}_{ij}N_i\ddot{N}_j + \mathcal{J}_{ij}\dot{N}_i\dot{N}_j + f_3(\dot{N}) + F_2(N^2,\dot{N}) = 0. \end{aligned} \tag{B2}$$

/

In Eqs. (B1) and (B2) we define the following quantities:

$$N_i \in \{X, Y, V, Z, W\}, \quad \Xi \equiv \alpha X + \beta Y, \quad U \equiv V + Z,$$
  
$$\alpha \equiv 3d_1 - d_3 - \frac{d_2}{2}, \quad \beta \equiv d_1 - d_3 - \frac{d_2}{2}, \quad \gamma \equiv d_3 - 2d_4, \quad \delta \equiv 2d_4,$$
  
$$f_1(\dot{N}) = (3\alpha + \beta) \dot{X} + (\alpha + 3\beta) \dot{Y} + (\alpha + \beta) \dot{U}, \quad f_2(N) = \beta X + \alpha Y + (\alpha + \beta)U,$$

$$f_2(\dot{N}) = \beta \dot{X} + \alpha \dot{Y} + (\alpha + \beta) \dot{U}, \quad f_3(\dot{N}) = \left(6a_0 + 2d_9 - d_{10}\right) \dot{X} + \left(-6a_0 + d_{10} - 2d_{11}\right) \dot{Y} + \left(d_{12} - d_{13}\right) \dot{U},$$

$$\mathcal{A}_{ij} = \begin{pmatrix} \gamma & \alpha + \gamma & \alpha & \alpha & 0 \\ \gamma - \beta & \gamma & -\beta - \beta & 0 \\ 2(\delta + \beta + 2\gamma) & 2(\delta + \beta + 2\gamma) & 0 & 0 & 0 \\ 2(\delta + \beta + 2\gamma) & 2(\delta + \beta + 2\gamma) & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & d_2 - 2d_5 + d_6 - d_7 \end{pmatrix},$$

$$\mathcal{B}_{ij} = \begin{pmatrix} -\delta & 3\alpha - 4\beta - \delta & 3\alpha + 2\beta + 3\gamma + \delta & 3\alpha + 2\beta + 3\gamma + \delta & 0 \\ 3\alpha - 4\beta - \delta & -\alpha - \beta - \delta & 3\gamma - \alpha - 2\beta + \delta & 3\gamma - \alpha - 2\beta + \delta \\ 3\alpha + 2\beta + 3\gamma + \delta & 3\gamma - \alpha - 2\beta + \delta & 5\beta - \alpha + 8\gamma + 4\delta & 5\beta - \alpha + 8\gamma + 4\delta & 0 \\ 3\alpha + 2\beta + 3\gamma + \delta & 3\gamma - \alpha - 2\beta + \delta & 5\beta - \alpha + 8\gamma + 4\delta & 5\beta - \alpha + 8\gamma + 4\delta \\ 0 & 0 & 0 & 0 & 2d_2 - d_5 \end{pmatrix},$$

$$G_{ij} = \begin{pmatrix} 2d_9 & d_{10} - 6a_0 & 3a_0 + d_{12} & 3a_0 + d_{15} & 0 \\ d_{10} - 6a_0 & 2d_{11} & 3a_0 + d_{13} & 3a_0 - 2d_{11} + 3d_{12} - 3d_{13} + d_{15} + 2d_9 & 0 \\ 3a_0 + d_{12} & 3a_0 + d_{13} & d_{14} & d_{10} + 2d_{12} - d_{13} - 6d_{14} + d_{15} + d_9 & 0 \\ 3a_0 + d_{15} & 3a_0 - 2d_{11} + 3d_{12} - 3d_{13} + d_{15} + 2d_9 & d_{10} + 2d_{12} - d_{13} - 6d_{14} + d_{15} + d_9 & 0 \\ 0 & 0 & 0 & 0 & 2(d_{16} - 6a_0) \end{pmatrix},$$

$$\mathcal{D}_{ij} = \begin{pmatrix} -\gamma & -2\alpha - \beta - \gamma & 2(-\alpha + \beta + 2\gamma + \delta) & 2(-\alpha + \beta + 2\gamma + \delta) & 0\\ \alpha + 2\beta - \gamma & -\gamma & 4(\beta + \gamma) + 2\delta & 4(\beta + \gamma) + 2\delta & 0\\ \alpha - 4(\beta + 2\gamma + \delta) & -5\beta - 4(2\gamma + \delta) & 0 & 0 & 0\\ \alpha - 4(\beta + 2\gamma + \delta) & -5\beta - 4(2\gamma + \delta) & 0 & 0 & 0\\ 0 & 0 & 0 & 0 & 2d_5 - d_2 - d_6 + d_7 \end{pmatrix},$$

$$\mathcal{M}_{ij} = \begin{pmatrix} 2\gamma & \alpha - \beta + 2\gamma & \alpha + 2(\beta + 2\gamma + \delta) & \alpha + 2(\beta + 2\gamma + \delta) & 0\\ \alpha - \beta + 2\gamma & 2\gamma & \beta + 4\gamma + 2\delta & \beta + 4\gamma + 2\delta & 0\\ \alpha + 2(\beta + 2\gamma + \delta) & \beta + 4\gamma + 2\delta & 0 & 0 & 0\\ \alpha + 2(\beta + 2\gamma + \delta) & \beta + 4\gamma + 2\delta & 0 & 0 & 0\\ 0 & 0 & 0 & 0 & d_2 - 2d_5 + d_6 - d_7 \end{pmatrix},$$

$$\begin{split} F_1(N^4) &= \frac{1}{2} \left( \alpha - \beta \right) W^4 - \left( 2\beta + \gamma + \delta \right) UYW^2 + \left( \alpha - \beta - \gamma - \delta \right) UXW^2 - \frac{1}{2} \left( 2\beta + 2\gamma + \delta \right) U^2Y^2 - \left( 2\beta + \gamma + \delta \right) UXY^2 \\ &+ \frac{1}{2} \left( \alpha - \beta \right) X^2Y^2 + \left( \alpha - \beta - \gamma - \delta \right) UX^2Y + \frac{1}{2} \left( \alpha - \beta - 2\gamma - \delta \right) U^2X^2 + \left( \beta + 2\gamma + \delta \right) U^2XY \\ &+ \frac{1}{3} \left( 4d_5 - 7d_6 + d_7 + 6d_8 + 6\alpha - 8 \left( \beta + \gamma + \delta \right) \right) XYW^2 + \left( \alpha + \beta - d_6 - d_7 + d_8 \right) Y^2W^2 + d_8X^2W^2, \end{split}$$

$$\begin{split} F_2(N^2,\dot{N}) &= (-\alpha + \beta + 2\gamma + \delta)X^2\dot{U} - (2\beta + 2\gamma + \delta)Y^2\dot{U} - (2\beta + \gamma + \delta)Y^2\dot{X} + (\alpha - \beta - \gamma - \delta)XY\dot{X} + (\alpha - \beta - 2\gamma - \delta)UX\dot{X} \\ &+ 3(\beta + 2\gamma + \delta)UY\dot{X} - 3(\beta + 2\gamma + \delta)UX\dot{Y} + (-\alpha + \beta + \gamma + \delta)X^2\dot{Y} + (2\beta + \gamma + \delta)XY\dot{Y} + (2\beta + 2\gamma + \delta)UY\dot{Y} \\ &+ \frac{1}{3}\left(-3\alpha + \beta + \gamma + \delta + 4d_5 - 4d_6 + d_7\right)W\left(X\dot{W} - W\dot{X}\right) + \frac{1}{3}\left(4\beta + \gamma + \delta + 4d_5 - d_6 + 4d_7\right)W\left(W\dot{Y} - Y\dot{W}\right). \end{split}$$

**Distortion equations** — The remaining five equations of motion correspond to variations of the distortion fields:  $\alpha \ddot{X} + \beta \ddot{Y} + (\alpha X + \beta Y) \dot{H} + 2d_1 (4H\dot{H} + \ddot{H}) - (6a_0 + 2d_9 - d_{10}) X + (6a_0 - d_{10} + 2d_{11}) Y + (d_{13} - d_{12}) U + 2(d_9 - d_{11}) U + 2(d_9$  $+2d_{12}-2d_{13})Z + \left[ (3\alpha - 9\beta - 6(\gamma + \delta) - d_6 + d_7)W^2 + (5\beta - \alpha + 8\gamma + 4\delta)U^2 + (5\alpha + 2(\beta + \gamma))UX - (2\alpha + 5\beta - 2\gamma)UY + (2\alpha + 5\beta - 2\gamma)UY +$  $-(\gamma+\delta)X^{2} + (5\alpha - 7\beta - 2(\gamma+\delta))XY + (-\alpha - \beta - \gamma - \delta)Y^{2}\Big]H + \Big[(4\alpha + \beta)\dot{X} + (\alpha + 4\beta)\dot{Y} + (\alpha + \beta)\dot{U}\Big]H$  $+4\left[(\alpha+\beta)U+\beta X+\alpha Y\right]H^2+(\beta-\alpha)XY^2+(2(\beta+2\gamma+\delta)-\alpha)U^2X-(3\beta+4\gamma+2\delta)U^2Y+(\alpha-\beta)X^2Y-(\alpha+\beta)UW^2$  $+\frac{1}{3}\left(6\alpha - 8(\beta + \gamma + \delta) + 4d_5 - 7d_6 + d_7\right)W^2X + \frac{1}{3}\left(14\beta + 8(\gamma + \delta) - 4d_5 + d_6 - 7d_7\right)W^2Y + (\alpha - \beta - \gamma - \delta)UX^2$  $-2(\alpha + \beta)UXY + (2\beta + \gamma + \delta)UY^2 + 2(\beta + 2\gamma + \delta)U\dot{Y} - \beta Y\dot{U} + 2(\beta + 2\gamma + \delta)U\dot{X} + \alpha X\dot{U} + (\alpha - \beta - \gamma - \delta)X\dot{X} - (2\beta + \beta)UXY + (2\beta + \gamma + \delta)UY^2 + 2(\beta + 2\gamma + \delta)U\dot{Y} - \beta Y\dot{U} + 2(\beta + 2\gamma + \delta)U\dot{X} + \alpha X\dot{U} + (\alpha - \beta - \gamma - \delta)X\dot{X} - (2\beta + \beta)U\dot{X} + \alpha X\dot{U} + (\alpha - \beta - \gamma - \delta)X\dot{X} - (2\beta + \beta)U\dot{X} + \alpha X\dot{U} + (\alpha - \beta - \gamma - \delta)X\dot{X} - (2\beta + \beta)U\dot{X} + \alpha X\dot{U} + (\alpha - \beta - \gamma - \delta)X\dot{X} - (2\beta + \beta)U\dot{X} + \alpha X\dot{U} + (\alpha - \beta - \gamma - \delta)X\dot{X} - (2\beta + \beta)U\dot{X} + \alpha X\dot{U} + (\alpha - \beta - \gamma - \delta)X\dot{X} - (2\beta + \beta)U\dot{X} + \alpha X\dot{U} + (\alpha - \beta - \gamma - \delta)X\dot{X} - (2\beta + \beta)U\dot{X} + \alpha X\dot{U} + (\alpha - \beta - \gamma - \delta)X\dot{X} - (2\beta + \beta)U\dot{X} + \alpha X\dot{U} + (\alpha - \beta - \gamma - \delta)X\dot{X} - (2\beta + \beta)U\dot{X} + \alpha X\dot{U} + (\alpha - \beta - \gamma - \delta)X\dot{X} - (2\beta + \beta)U\dot{X} + \alpha X\dot{U} + (\alpha - \beta - \gamma - \delta)X\dot{X} - (2\beta + \beta)U\dot{X} + \alpha X\dot{U} + (\alpha - \beta - \gamma - \delta)X\dot{X} - (2\beta + \beta)U\dot{X} + \alpha X\dot{U} + (\alpha - \beta - \gamma - \delta)X\dot{X} - (2\beta + \beta)U\dot{X} + \alpha X\dot{U} + (\alpha - \beta - \gamma - \delta)X\dot{X} - (2\beta + \beta)U\dot{X} + \alpha X\dot{U} + (\alpha - \beta - \gamma - \delta)X\dot{X} - (2\beta + \beta)U\dot{X} + \alpha X\dot{U} + (\alpha - \beta - \gamma - \delta)X\dot{X} - (2\beta + \beta)U\dot{X} + (\alpha - \beta - \gamma - \delta)X\dot{X} + (\alpha - \beta - \delta)X\dot{X} + (\alpha - \beta - \gamma - \delta)X\dot{X} + (\alpha - \beta - \delta)X\dot{X} + (\alpha$  $+\gamma+\delta )Y\dot{X} + (\alpha-\beta-\gamma-\delta)X\dot{Y} - (2\beta+\gamma+\delta)Y\dot{Y} - \frac{1}{3}\left(-3\alpha+5\beta+2(\gamma+\delta)+8d_5-5d_6+5d_7\right)W\dot{W} = 0$ (B3)  $(\beta+2\gamma+\delta)\ddot{X} + (2(\beta+\gamma)+\delta)\ddot{Y} + \left[(\alpha+\gamma)X + \gamma Y + 2(\beta+2\gamma+\delta)U\right]\dot{H} + (\alpha-3\beta)H\dot{H} - \beta\ddot{H} + \left(d_{10} - 6a_0\right)X + 2d_{11}Y\dot{H} + (\alpha-3\beta)H\dot{H} + \beta\dot{H} + \left(d_{10} - 6a_0\right)X + 2d_{11}Y\dot{H} + (\alpha-3\beta)H\dot{H} + \beta\dot{H} + \beta\dot{H}$  $+ \left(3 a_0 + d_{13}\right) U + \left(2 d_9 - 2 d_{11} + 3 d_{12} - 4 d_{13} + d_{15}\right) Z + \left[\left(d_7 - 3 (2\beta + \gamma + \delta)\right) W^2 + (3\beta + 4\gamma + 2\delta) U^2 + (2\alpha + 5\beta + 6\gamma + 6\gamma + \delta) U^2 + (2\beta + 4\gamma + 2\delta) U^2 + (2\beta + 4\gamma +$  $+3\delta)UX + (-6\beta - 4\gamma - 3\delta)UY + (-\gamma - \delta)X^{2} + (2\alpha - 4\beta - \gamma - \delta)XY \Big]H + \Big[(\alpha + 4\beta + 6\gamma + 3\delta)\dot{X} + 3(2(\beta + \gamma) + \delta)\dot{Y} + (-\beta - 4\gamma - 3\delta)\dot{X} + 3(2(\beta + \gamma) + \delta)\dot{Y} \Big]$  $+ (3\beta + 4\gamma + 2\delta)\dot{U}\Big]H + \Big[(\alpha + 8\beta + 9\gamma + 5\delta)U + (4\beta + 3\gamma + \delta)X + (\alpha + \beta + 3\gamma + \delta)Y\Big]H^2 + \frac{1}{2}(6\alpha - 8(\beta + \gamma + \delta) + 4d_5)H^2 + \frac{1}{2}(6\alpha - 8(\beta + \gamma + \delta) + \frac{1}{2}(6\alpha - 8(\beta + \gamma + \delta) + 4d_5)H^2 + \frac{1}{2}(6\alpha - 8(\beta + \gamma + \delta) + \frac{1}{2}(6\alpha - 8(\beta + \beta + \delta) + \frac{1}{2}($  $-7d_{6} + d_{7} + 6d_{8}W^{2}X + 2(\alpha + \beta - d_{6} - d_{7} + d_{8}W^{2}Y + (\beta + 2\gamma + \delta)U^{2}X - (2(\beta + \gamma) + \delta)U^{2}Y - (2\beta + \gamma + \delta)UW^{2}$  $+ (\alpha - \beta - \gamma - \delta)UX^{2} + 2(\beta + 2\gamma + \delta)U\dot{X} - 2(2\beta + \gamma + \delta)UXY + (\alpha - \beta)X^{2}Y + (\beta + 2\gamma + \delta)X\dot{U} - (2(\beta + \gamma) + \delta)Y\dot{U}$  $+ (\alpha - \beta - \gamma - \delta)X\dot{X} - (2\beta + \gamma + \delta)Y\dot{X} - \frac{1}{2}(4\beta + \gamma + \delta + 4d_5 - d_6 + 4d_7)W\dot{W} = 0,$ (B4)  $(\alpha + \beta)H\dot{H} + (3a_0 + d_{12})X + (3a_0 + d_{13})Y + 2d_{14}U + (2d_9 + d_{10} + 2d_{12} - d_{13} - 8d_{14} + d_{15})Z + [(\alpha - 5\beta - 4(2\gamma + \delta))U]$  $+\left(-2\beta-3\gamma-\delta\right)X+(\alpha-\beta-3\gamma-\delta)Y\Big]H^2+\Big[2(\alpha-2(\beta+2\gamma+\delta))UX+(6\beta+8\gamma+4\delta)UY+(\alpha+\beta)W^2-\gamma X^2+(\alpha+\beta)W^2-\gamma X^2+(\alpha+\beta)W$  $+2(\alpha+\beta)XY+\gamma Y^{2}+(\alpha-2(\beta+2\gamma+\delta))\dot{X}+(-3\beta-4\gamma-2\delta)\dot{Y}\Big]H+(\alpha X-\beta Y)\dot{H}+(\alpha-\beta-2\gamma-\delta)UX^{2}$  $+2(\beta+2\gamma+\delta)UXY - (2(\beta+\gamma)+\delta)UY^2 + (\alpha-\beta-\gamma-\delta)W^2X - (2\beta+\gamma+\delta)W^2Y + (\alpha-\beta-\gamma-\delta)X^2Y$  $-(2\beta + \gamma + \delta)XY^{2} + (\alpha - \beta - 2\gamma - \delta)X\dot{X} + (\beta + 2\gamma + \delta)Y\dot{X} - (\beta + 2\gamma + \delta)X\dot{Y} + (2(\beta + \gamma) + \delta)Y\dot{Y} = 0,$ (B5)  $(d_{12} - d_{15}) X + (-2d_9 + 2d_{11} - 3d_{12} + 4d_{13} - d_{15}) Y - (2d_9 + d_{10} + 2d_{12} - d_{13} - 8d_{14} + d_{15}) V$ +  $(8d_9 + 3d_{10} - 2d_{11} + 8d_{12} - 7d_{13} - 24d_{14} + 5d_{15}) Z = 0$ , (B6)  $2(-6a_{0} + d_{16})W + (-4d_{5} + 3d_{6} - 3d_{7} - \alpha + 3\beta + 2(\gamma + \delta))H^{2}W + 2(\alpha - \beta)W^{3} + 2(\alpha - \beta - \gamma - \delta)UWX + 2d_{8}WX^{2}$  $-2(2\beta + \gamma + \delta)UWY + \frac{2}{3}(4d_5 - 7d_6 + d_7 + 6d_8 + 6\alpha - 8(\beta + \gamma + \delta))WXY + 2(-d_6 - d_7 + d_8 + \alpha + \beta)WY^2$  $+\left(d_{6}-2d_{5}-d_{7}+\alpha-3\beta-2\left(\gamma+\delta\right)\right)W\dot{H}+\left[2\left(\alpha+\beta\right)UW-\left(4d_{5}-6d_{6}+d_{7}+3\alpha-5\left(\beta+\gamma+\delta\right)\right)WX$  $+ \left(4d_{5} - d_{6} + 6d_{7} - 8\beta - 5(\gamma + \delta)\right)WY + 2\left(d_{5} - d_{6} + d_{7} - 2(\beta + \gamma + \delta)\right)\dot{W} H - \frac{1}{3}\left(4d_{5} - 4d_{6} + d_{7} - 3\alpha + \beta + \gamma + \delta\right)W\dot{X}$  $+\frac{1}{2}\left(4d_{5}-d_{6}+4d_{7}+4\beta+\gamma+\delta\right)W\dot{Y}+\frac{1}{2}\left(d_{5}-d_{6}+d_{7}-2\left(\beta+\gamma+\delta\right)\right)\ddot{W}=0.$ (**B**7) Specifically Eq. (B3) is  $\delta S/\delta Y - \delta S/\delta X = 0$ , Eq. (B4) is  $\delta S/\delta Y = 0$ , Eq. (B5) is  $\delta S/\delta V = 0$ , Eq. (B6) is  $\delta S/\delta V - \delta S/\delta Z = 0$ ,

and Eq. (B7) is  $\delta S / \delta W = 0$ .

#### C. Particle spectrographs

**Conventions** — The particle spectra are obtained using the *Particle Spectrum for Any Tensor Lagrangian (PSALTer)* software [19, 98, 147–149]. Near Minkowski spacetime, we take the metric perturbation to be  $h_{\mu\nu} \equiv g_{\mu\nu} - \eta_{\mu\nu}$ . In MAG, we take the connection  $\Gamma_{\mu\nu}^{\ \rho}$  to be inherently perturbative. Conjugate to  $h_{\mu\nu}$  is the linearised stress-energy tensor  $T^{\mu\nu}$ , and conjugate to  $\Gamma_{\mu\nu}^{\ \rho}$  is the linearised hypermomentum  $\Delta^{\mu\nu}_{\ \rho}$ . Under projection with the unit-timelike vector  $n_{\mu}n^{\mu} \equiv 1$ , where  $n^{\mu} \equiv k^{\mu}/k$  for  $k^2 \equiv k^{\mu}k_{\mu}$  and  $k^{\mu}$  the massive four-momentum, the various SO(3) irreducible parts of these quantities are presented in Figs. 2 and 3. They

have spin-parity  $(J^P)$  labels to identify them. Duplicate  $J^P$  states can arise, and these are distinguished by extra labels '#1', '#2', etc. The analyses in Figs. 4 to 8 were performed across 64 AMD<sup>®</sup> *Ryzen Threadripper* CPUs. Further details are provided in the supplemental materials [144].

Fundamental fiel	J Symmetries	Decomposition into SO(3) irrep(s)	Source
h <sub>αβ</sub>	Symmetry[2, $h^{\bullet 1 \bullet 2}$ , { $\bullet 1 \rightarrow -a$ , $\bullet 2 \rightarrow -b$ }, StrongGenSet[{1, 2}, GenSet[(1,2)]]	$\frac{1}{3} \eta_{\alpha\beta} h_{0}^{\theta 1} + h_{2}^{\pm 1}{}_{\alpha\beta} + h_{1}^{\pm 1}{}_{\beta} \eta_{\alpha} + h_{1}^{\pm 1}{}_{\alpha} \eta_{\beta} - \frac{1}{3} h_{0}^{\theta 1} \eta_{\alpha} \eta_{\beta} + h_{0}^{\theta 2} \eta_{\alpha} \eta_{\beta}$	τ <sub>αβ</sub>
SO(3) irrep	Symmetries	Expansion in terms of the fundamental field	Source SO(3) irrep
h <sub>0+</sub> <sup>#1</sup>	Symmetry[0, $h_{0^+}^{\pm 1}$ , {}, StrongGenSet[{}, GenSet[]]]	$h^{\alpha}_{\alpha} = h_{\alpha\beta} n^{\alpha} n^{\beta}$	τ <sub>0+</sub> #1
h <sub>0</sub> <sup>#2</sup>	Symmetry[0, $h_{0^+}^{\#2}$ , {}, StrongGenSet[{}, GenSet[]]]	$h_{\alpha\beta} n^{\alpha} n^{\beta}$	τ <sub>0+</sub> #2
$h_{2^{+} \alpha \beta}^{\#1}$	Symmetry[2, $h_{2^+}^{\#1 \bullet 1 \bullet 2}$ , $\{\bullet 1 \rightarrow -a, \bullet 2 \rightarrow -b\}$ , StrongGenSet[ $\{1, 2\}$ , GenSet[ $(1, 2)$ ]]]	$h_{\alpha\beta} - \frac{1}{3} \eta_{\alpha\beta} h_{\chi}^{\chi} + \frac{1}{3} h_{\chi}^{\chi} n_{\alpha} n_{\beta} - h_{\beta\chi} n_{\alpha} n^{\chi} - h_{\alpha\chi} n_{\beta} n^{\chi} + \frac{1}{3} \eta_{\alpha\beta} h_{\chi\sigma} n^{\chi} n^{\delta} + \frac{2}{3} h_{\chi\sigma} n_{\alpha} n_{\beta} n^{\chi} n^{\delta}$	$\tau_{2^{+} \alpha \beta}^{\#1}$
$h_1^{\#1}{}_{\alpha}$	Symmetry[1, $h_1^{\#1\bullet1}$ , $\{\bullet1 \rightarrow -a\}$ , StrongGenSet[{}, GenSet[]]]	$h_{\alpha\beta} \ n^{\beta} - h_{\beta\chi} \ n_{\alpha} \ n^{\beta} \ n^{\chi}$	$\tau_1^{\#1}{}_{\alpha}$

FIG. 2. Kinematic structure of the metric perturbation  $h_{\mu\nu}$ . These definitions are used in Figs. 4 to 8. See [148] for further notational details. This is a vector graphic: all details are visible under magnification.

*Gravity with K-screening* — We will first study the model considered in Section IV A. The conditions in Eq. (42) may be imposed with the rules

$$a_0 \mapsto 0, \quad c_6 \mapsto -2c_1 + 6c_{16} + 2c_2 - 2c_3 - c_4 + c_5, \quad c_8 \mapsto \frac{c_{11}}{2} + \frac{c_{12}}{2} - 3c_{16} - c_7, \quad c_9 \mapsto -c_{10} + \frac{c_{11}}{2} + \frac{c_{12}}{2} - 3c_{16}.$$
 (C1)

To simplify the formulae for the masses of the various propagating particles, we consider the case where only  $a_1$  and  $a_4$  are non-zero. By comparing with Eq. (25), we see that these parameters modulate the 'index-for-index squares' of the torsion and non-metricity, respectively. Thus, they are somewhat analagous to the Kretschmann scalar for curvature, and we take them to be representative of the mass-dimension-two parameters that may be included in the model. Accordingly, we impose the conditions

$$a_2 \mapsto 0, \quad a_3 \mapsto 0, \quad a_5 \mapsto 0, \quad a_6 \mapsto 0, \quad a_7 \mapsto 0, \quad a_8 \mapsto 0, \quad a_9 \mapsto 0, \quad a_{10} \mapsto 0, \quad a_{11} \mapsto 0. \tag{C2}$$

When Eqs. (C1) and (C2) are applied to Eq. (25), the resulting partial particle spectrum is shown in Fig. 4. The abundance of unconstrained parameters makes the computation of the full spectrum difficult, and the 1<sup>-</sup> sector is entirely neglected at this stage. Despite this, there is enough information about the remaining massive  $J^P$  sectors to begin constraining the model. Beginning with the 2<sup>-</sup> sector, we see that the denominator of the saturated propagator (the pseudo-determinant of the wave operator matrix) is a quadratic in  $k^2$  – we thus expect *two* massive poles. Generically, the expressions for the square masses in terms of the remaining Lagrangian couplings will contain radicals owing to the quadratic formula, and these are awkward points about which to compute pole residues when determining the unitarity of the model. To proceed, we impose further constraints so as to remvoe the coefficient of  $k^4$ . This coefficient is a quadratic form in the mass-dimension-zero couplings, which is problematic because it is computationally preferable to impose *linear* constraints among the couplings. An impartial (i.e. parameterisation-independent) approach is to diagonalise the quadratic form as a weighted sum of squares, and insist that each term vanish separately. This approach yields the constraints

$$c_4 \mapsto 2c_2, \quad c_5 \mapsto 4c_1, \quad c_{16} \mapsto -\frac{c_1}{3} + \frac{c_2}{3} + \frac{c_3}{3},$$
 (C3)

and, in fact, these eliminate not only the  $k^4$  coefficient but also the coefficient of  $k^2$ : there is accordingly no massive 2<sup>-</sup> particle from this point forward. After imposing Eq. (C3), the determinants for the remaining  $J^P$  sectors are naturally much simplified. Proceeding to the 1<sup>+</sup> sector, a similar quartic system is attempted to be reduced to a single massive pole using the same method, yielding the constraints

$$c_2 \mapsto -c_1 + c_{10} - c_{11} + c_7, \quad c_3 \mapsto 2c_1 - c_{10} + c_7, \quad c_{12} \mapsto 2c_{10} - 3c_{11} + 6c_7, \tag{C4}$$

but once again this removes the 1<sup>+</sup> particle entirely. Imposing Eqs. (C3) and (C4) on Fig. 4 suggests that one massive particle will still arise from each of the 0<sup>+</sup>, 0<sup>-</sup> 2<sup>+</sup> and 3<sup>-</sup> sectors. However, the 1<sup>-</sup> sector was not yet considered at all. When Eqs. (C1) to (C4) are applied to Eq. (25), the partial particle spectrum for the 1<sup>-</sup> sector is simple enough to be obtainable, but still too large to present (see the supplemental materials [144]). Commensurate with the large size of its wave operator and saturated propagator matrices, the 1<sup>-</sup> sector contains a *quartic* in  $k^2$ , suggesting that there will be *four* massive poles. It is easiest to begin 'half-way through' by eliminating the  $k^4$  coefficient once again. This coefficient is a sum of quadratic forms in the mass-dimension-zero couplings, one multiplied by  $a_1a_4^3$  and the other by  $a_1^2a_4^2$ . Eliminating them both independently yields, respectively

$$c_{11} \mapsto 2c_7, \quad c_{14} \mapsto 3c_{10} + 2c_{13}, \quad c_{15} \mapsto 3c_{10} + 2c_{13},$$
(C5a)

$$c_7 \mapsto 0, \quad c_{10} \mapsto 0, \quad c_{13} \mapsto 0.$$
 (C5b)

As was the experience with Eqs. (C3) and (C4), the whole of the  $1^-$  sector is actually rendered non-propagating by Eqs. (C5a) and (C5b), although this was not our intention. When Eqs. (C5a) and (C5b) are applied sequentially on Fig. 5, all the massive

Fundamental fi	eld Symmetries	Decomposition into SO(3) irrep(s)	Source
Γαβχ	Symmetry[3, r●1≠3#3, {●1→-a, ●2→-b, ●3→-c}, StrongGenSet[{}, GenSet[]]]	$\begin{split} & \frac{1}{2} \left( \sigma_{xx} + \frac{1}{2}^{-1} s + \frac{1}{2} - \sigma_{yx} + \frac{1}{2}^{-1} s^{-2} - \sigma_{xx} + \frac{1}{2} - \frac{1}{2} s^{-2} - \sigma_{xx} - \frac{1}{2} -$	$\Delta_{a\beta\chi}$
SO(3) irrep	Symmetries	Expansion in terms of the fundamental field	Source SO(3) irrep
Γ <sub>0+</sub> <sup>#1</sup>	Symmetry[0, F <sup>#1</sup> <sub>0+</sub> , {}, StrongGenSet[{}, GenSet[]]]	$\cdot \frac{1}{2} \Gamma^{\alpha}{}_{\alpha}{}^{\beta} n_{\beta} + \frac{1}{2} \Gamma^{\alpha\beta}{}_{\alpha} n_{\beta}$	∆ <sub>0</sub> <sup>#1</sup>
F <sub>0</sub> <sup>#2</sup>	Symmetry[0, Γ <sup>#2</sup> <sub>0+</sub> , {}, StrongGenSet[{}, GenSet[]]]	$\Gamma^{\alpha\beta\chi} n_{\alpha} n_{\beta} n_{\chi}$	Δ# <sup>2</sup> 0+
Γ#3	Symmetry[0, Γ <sup>#3</sup> , {}, StrongGenSet[{}, GenSet[]]]	$\Gamma^{\alpha\beta}{}_{\beta} n_{\alpha} + \Gamma^{\alpha}{}_{\alpha}{}^{\beta} n_{\beta} + \Gamma^{\alpha\beta}{}_{\alpha} n_{\beta} \cdot \Im \Gamma^{\alpha\beta\chi} n_{\alpha} n_{\beta} n_{\chi}$	$\Delta_{0^{+}}^{#3}$
F <sup>#4</sup>	Symmetry[0, Γ <sub>0</sub> <sup>#4</sup> , {}, StrongGenSet[{}, GenSet[]]]	$\Gamma^{\alpha\beta}{}_{\beta} n_{\alpha} - \frac{1}{2} \Gamma^{\alpha}{}_{\alpha}{}^{\beta} n_{\beta} - \frac{1}{2} \Gamma^{\alpha\beta}{}_{\alpha} n_{\beta}$	$\Delta_{0^{+}}^{\#4}$
F <sub>0</sub> #1	Symmetry[0, Γ <sub>0</sub> <sup>#1</sup> , {}, StrongGenSet[{}, GenSet[]]]	$\epsilon \eta_{a\beta\chi\delta} \Gamma^{a\beta\chi} n^{\delta}$	Δ#1
Γ <sup>#1</sup> <sub>1<sup>+</sup> αβ</sub>	Symmetry[2, $\Gamma_{1^+}^{\#_1 \bullet 1 \bullet 2}$ , $\{\bullet 1 \rightarrow -a, \bullet 2 \rightarrow -b\}$ , StrongGenSet[{1, 2}, GenSet[-(1,2)]]]	$\frac{1}{4}\Gamma_{\alpha\beta}^{\ \ x}n_{\chi} - \frac{1}{4}\Gamma_{\alpha\beta}^{\ \ x}n_{\chi} - \frac{1}{4}\Gamma_{\beta\alpha}^{\ \ x}n_{\chi} + \frac{1}{4}\Gamma_{\beta\alpha}^{\ \ \chi}n_{\chi} - \frac{1}{4}\Gamma_{\alpha\beta}^{\ \ \chi}n_{\alpha} - \frac{1}{4}\Gamma_{\alpha\beta}^{\ \ \chi}n_{\alpha} - \frac{1}{4}\Gamma_{\alpha\beta}^{\ \ \chi}n_{\alpha} - \frac{1}{4}\Gamma_{\alpha\beta}^{\ \ \chi}n_{\alpha} + \frac{1}{4}\Gamma_{\alpha\beta}^{\ \ \chi}n_{\alpha} - \frac{1}{4}\Gamma_{\alpha\beta}^$	$\Delta_{1^{+} \alpha \beta}^{\#1}$
Γ <sup>#2</sup> <sub>1<sup>+</sup> αβ</sub>	Symmetry[2, $\Gamma_{1+}^{\#2 \bullet 1 \bullet 2}$ , { $\bullet 1 \rightarrow -a, \bullet 2 \rightarrow -b$ }, StrongGenSet[{1, 2}, GenSet[-(1,2)]]]	$\frac{1}{2} \Gamma_{\alpha\beta}^{\chi} n_{\chi} - \frac{1}{2} \Gamma_{\beta\alpha}^{\chi} n_{\chi} + \frac{1}{2} \Gamma_{\beta}^{\chi} \delta n_{\alpha} n_{\chi} n_{\delta} - \frac{1}{2} \Gamma_{\alpha}^{\chi} \delta n_{\alpha} n_{\chi} n_{\delta} - \frac{1}{2} \Gamma_{\alpha}^{\chi} \delta n_{\beta} n_{\chi} n_{\delta} + \frac{1}{2} \Gamma_{\alpha}^{\chi} \delta n_{\delta} n_$	Δ <sup>#2</sup> <sub>1</sub> + <sub>αβ</sub>
Γ#3 1+ αβ	Symmetry[2, $\Gamma_1^{g_2^{a} \rightarrow b^2}$ , { $\bullet$ 1 $\rightarrow$ -a, $\bullet$ 2 $\rightarrow$ -b}, StrongGenSet[{1, 2}, GenSet[-(1,2)]]	$ \frac{1}{2} \Gamma_{ab}^{-5} n_{a}^{-\frac{1}{2}} \Gamma_{ab}^{-5} n_{x}^{-\frac{1}{2}} \Gamma_{\mu}^{-5} n_{x}^{-\frac{1}{2}} \Gamma_{\mu}^{-5} n_{x}^{-\frac{1}{2}} \Gamma_{\mu}^{-5} n_{x}^{-\frac{1}{2}} \Gamma_{\mu}^{-5} n_{x}^{-\frac{1}{2}} \Gamma_{\mu}^{-5} n_{x}^{-\frac{1}{2}} n_{x}^{-\frac{1}{2}} \Gamma_{\mu}^{-5} n_{\mu}^{-\frac{1}{2}} n_{\mu}^{-\frac{1}{2}} \Gamma_{\mu}^{-\frac{1}{2}} n_{\mu}^{-\frac{1}{2}} \Gamma_{\mu}^{-\frac{1}{2}} n_{\mu}^{-\frac{1}{2}} n_{\mu}^{$	Δ#3 1+ αβ
Γ1***	Symmetry[1, $\Gamma_1^{\pm 1 \bullet 1}$ , { $\bullet 1 \rightarrow -a$ }, StrongGenSet[{}, GenSet[]]]	$-\frac{1}{2}\Gamma^{\beta}{}_{\alpha\beta}+\frac{1}{2}\Gamma^{\beta}{}_{\beta\alpha}{}_{\alpha}{}-\frac{1}{2}\Gamma^{\beta}{}_{\beta}{}^{\chi}n_{\alpha}n_{\chi}+\frac{1}{2}\Gamma^{\beta\chi}{}_{\beta}n_{\alpha}n_{\chi}+\frac{1}{2}\Gamma^{\beta}{}_{\alpha}{}^{\chi}n_{\beta}n_{\chi}{}-\frac{1}{2}\Gamma^{\beta\chi}{}_{\alpha}n_{\beta}n_{\chi}$	Δ <sup>#1</sup> <sub>α</sub>
Γ <sub>1</sub> <sup>#2</sup> α	Symmetry[1, Γ <sup>+2•1</sup> , {●1→-a}, StrongGenSet[{}, GenSet[]]]	$\frac{1}{2}\Gamma_a^{\beta} x \eta_B \eta_z - \frac{1}{2}\Gamma_a^{\beta} \eta_B \eta_z$	Δ <sup>#2</sup> <sub>α</sub>
Γ <sup>#3</sup> <sub>1α</sub>	Symmetry[1, $\Gamma_{1}^{\#3}\bullet1$ , $\{\bullet1\rightarrow -a\}$ , StrongGenSet[{}, GenSet[]]]	$\frac{z}{\Gamma_{\alpha}\beta^{\chi}} \frac{r}{n_{\beta}} \frac{r}{n_{\gamma}} + \frac{r}{\Gamma_{\alpha}} \frac{r}{n_{\beta}} \frac{r}{n_{\gamma}} + \frac{r}{\Gamma_{\alpha}} \frac{r}{n_{\beta}} \frac{r}{n_{\gamma}} + \frac{r}{3} \frac{r}{\Gamma_{\alpha}} \frac{r}{n_{\beta}} \frac{r}{n_{\beta$	Δ <sup>#3</sup> <sub>1</sub>
Γ <sub>1</sub> <sup>#4</sup>	Symmetry[1, $\Gamma_1^{\texttt{ide1}}$ , $\{\bullet 1 \rightarrow -a\}$ , StrongGenSet[ $\{\}$ , GenSet[]]]	$ \Gamma_{\alpha\beta}^{\ \beta} + \Gamma_{\alpha\beta}^{\beta} + \Gamma_{\beta\alpha}^{\beta} - \Gamma_{\beta\alpha}^{\beta\chi} n_{\alpha} n_{\beta} - \Gamma_{\beta\chi}^{\beta\chi} n_{\alpha} n_{\chi} - \Gamma_{\beta\chi}^{\beta\chi} n_{\alpha} n_{\chi} - \Gamma_{\alpha\beta\chi}^{\beta\chi} n_{\beta} n_{\chi} - \Gamma_{\alpha\beta\chi}^{\beta\chi} n_{\beta} n_{\chi} - \Gamma_{\alpha\beta\chi}^{\beta\chi} n_{\beta} n_{\chi} - \Gamma_{\beta\chi}^{\beta\chi} n_{\gamma} - \Gamma_{\gamma\chi}^{\beta\chi} n_{\gamma} - \Gamma$	Δ <sup>#4</sup> <sub>1</sub>
Γ <sup>#5</sup> <sub>1</sub>	Symmetry[1, $\Gamma_{1}^{\bullet 5 \bullet 1}$ , $\{\bullet 1 \rightarrow -a\}$ , StrongGenSet[{}, GenSet[]]]	$ \Gamma_{\alpha}^{\ \beta \chi} \ n_{\beta} \ n_{\chi} - \frac{1}{2} \ \Gamma_{\beta}^{\ \alpha} \ n_{\beta} \ n_{\chi} - \frac{1}{2} \ \Gamma_{\beta}^{\ \alpha} \ n_{\beta} \ n_{\chi} $	Δ <sup>#5</sup> <sub>α</sub>
Γ1#6	Symmetry[1, $\Gamma_{1}^{\#6\bullet1}$ , $\{\bullet1 \rightarrow -a\}$ , StrongGenSet[{}, GenSet[]]]	$\Gamma_{\alpha\beta}^{\ \beta} = \frac{1}{2} \Gamma_{\alpha\beta}^{\beta} - \frac{1}{2} \Gamma_{\beta\alpha}^{\beta} - \frac{\Gamma^{\beta\chi}}{2} \Gamma_{\alpha}^{\alpha} n_{\beta} + \frac{1}{2} \Gamma_{\beta}^{\beta\chi} n_{\alpha} n_{\chi} + \frac{1}{2} \Gamma^{\beta\chi}_{\ \beta} n_{\alpha} n_{\chi} - \Gamma_{\alpha}^{\ \beta\chi} n_{\beta} n_{\chi} + \frac{1}{2} \Gamma^{\beta\chi}_{\ \alpha} n_{\chi} + \frac{1}{2} \Gamma^{\beta\chi}_{\ \alpha} n_{\chi} + \frac{1}{2} \Gamma^{\beta\chi}_{\ \alpha} + \frac{1}{2} \Gamma^{\beta\chi}_{\ \alpha} n_{\chi} + \frac$	Δ <sup>#6</sup> <sub>1 α</sub>
Γ <sup>#1</sup> <sub>2<sup>+</sup> αβ</sub>	Symmetry[2, $\Gamma_2^{g_1 \bullet 1 \bullet 2}$ , $\{\bullet 1 \rightarrow \cdot a, \bullet 2 \rightarrow \cdot b\}$ , StrongGenSet[{1, 2}, GenSet[(1.2)]]]	$ \frac{1}{4} \Gamma_{\alpha\beta}^{\ \ \alpha} n_{\beta} \frac{1}{h_{\alpha}} \frac{1}{h_{\alpha}$	$\Delta_{2^+ \alpha\beta}^{\#1}$
Γ <sub>2+ αβ</sub>	Symmetry(2, $\Gamma_2^{\phi_2 \bullet i \bullet 2}$ , { $\bullet 1 \rightarrow \cdot a$ , $\bullet 2 \rightarrow \cdot b$ }, StrongGenSet{{1, 2}, GenSet{(1,2)]}	$ \begin{array}{c} \frac{1}{2} \Gamma_{\alpha\beta}^{\ \ \ } n_{\alpha} + \frac{1}{2} \Gamma_{\beta}^{\ \ \ \ } n_{\alpha} + \frac{1}{2} \Gamma_{\beta}^{\ \ \ \ } n_{\alpha} + \frac{1}{2} \Gamma_{\alpha\beta}^{\ \ \ \ } n_{\alpha} + \frac{1}{2} \Gamma_{\beta\alpha}^{\ \ \ \ \ } n_{\alpha} + \frac{1}{2} \Gamma_{\beta\alpha}^{\ \ \ \ \ } n_{\alpha} + \frac{1}{2} \Gamma_{\beta\alpha}^{\ \ \ \ \ } n_{\alpha} + \frac{1}{2} \Gamma_{\beta\alpha}^{\ \ \ \ \ \ } n_{\alpha} + \frac{1}{2} \Gamma_{\beta\alpha}^{\ \ \ \ \ } n_{\alpha} + \frac{1}{2} \Gamma_{\beta\alpha}^{\ \ \ \ \ \ } n_{\alpha} + \frac{1}{2} \Gamma_{\beta\alpha}^{\ \ \ \ \ \ \ } n_{\alpha} + \frac{1}{2} \Gamma_{\beta\alpha}^{\ \ \ \ \ \ \ } n_{\alpha} + \frac{1}{2} \Gamma_{\beta\alpha}^{\ \ \ \ \ \ \ } n_{\alpha} + \frac{1}{2} \Gamma_{\beta\alpha}^{\ \ \ \ \ \ \ \ \ } n_{\alpha} + \frac{1}{2} \Gamma_{\beta\alpha}^{\ \ \ \ \ \ \ \ \ \ \ } n_{\alpha} + \frac{1}{2} \Gamma_{\beta\alpha}^{\ \ \ \ \ \ \ \ \ \ } n_{\alpha} + \frac{1}{2} \Gamma_{\beta\alpha}^{\ \ \ \ \ \ \ \ \ \ \ \ \ \ } n_{\alpha} + \frac{1}{2} \Gamma_{\beta\alpha}^{\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ $	$\Delta_2^{\#2}{}_{\alpha\beta}$
Γ <sub>2+αβ</sub>	Symmetry(2, $\Gamma_{2}^{\#3} \bullet i \bullet^2$ , { $\bullet 1 \rightarrow \cdot a$ , $\bullet 2 \rightarrow \cdot b$ }, StrongGenSet[{1, 2}, GenSet[(1, 2)]]	$ \frac{1}{4} \begin{bmatrix} r_{ab}^{ s} & n_{c}^{-\frac{1}{4}} \end{bmatrix} \begin{bmatrix} r_{ab}^{ s} & n_{c}^{-\frac{1}{4}} \end{bmatrix} \begin{bmatrix} r_{ba}^{ s} & n_{c} + \frac{1}{2} \end{bmatrix} \begin{bmatrix} r_{ba}^{ s} & n_{c} + n_{c} + \frac{1}{2} \end{bmatrix} \begin{bmatrix} r_{ba}^{ s} & n_{c} + n_{c} + \frac{1}{2} \end{bmatrix} \begin{bmatrix} r_{ba}^{ s} & n_{c} + n_{c} + \frac{1}{2} \end{bmatrix} \begin{bmatrix} r_{ba}^{ s} & n_{c} + n_{c} + \frac{1}{2} \end{bmatrix} \begin{bmatrix} r_{ba}^{ s} & n_{c} + n_{c} + \frac{1}{2} \end{bmatrix} \begin{bmatrix} r_{ba}^{ s} & n_{c} + n_{c} + \frac{1}{2} \end{bmatrix} \begin{bmatrix} r_{ba}^{ s} & n_{c} + n_{c} + \frac{1}{2} \end{bmatrix} \begin{bmatrix} r_{ba}^{ s} & n_{c} + n_{c} + \frac{1}{2} \end{bmatrix} \begin{bmatrix} r_{ba}^{ s} & n_{c} + n_{c} + \frac{1}{2} \end{bmatrix} \begin{bmatrix} r_{ba}^{ s} & n_{c} + n_{c} + \frac{1}{2} \end{bmatrix} \begin{bmatrix} r_{ba}^{ s} & n_{c} + n_{c} + \frac{1}{2} \end{bmatrix} \begin{bmatrix} r_{ba}^{ s} & n_{c} + n_{c} + \frac{1}{2} \end{bmatrix} \begin{bmatrix} r_{ba}^{ s} & n_{c} + n_{c} + \frac{1}{2} \end{bmatrix} \begin{bmatrix} r_{ba}^{ s} & n_{c} + n_{c} + \frac{1}{2} \end{bmatrix} \begin{bmatrix} r_{ba}^{ s} & n_{c} + n_{c} + \frac{1}{2} \end{bmatrix} \begin{bmatrix} r_{ba}^{ s} & n_{c} + n_{c} + \frac{1}{2} \end{bmatrix} \begin{bmatrix} r_{ba}^{ s} & r_{b} + n_{c} + \frac{1}{2} \end{bmatrix} \begin{bmatrix} r_{ba}^{ s} & r_{b} + n_{c} + \frac{1}{2} \end{bmatrix} \begin{bmatrix} r_{ba}^{ s} & r_{b} + n_{c} + \frac{1}{2} \end{bmatrix} \begin{bmatrix} r_{ba}^{ s} & r_{b} + n_{c} + \frac{1}{2} \end{bmatrix} \begin{bmatrix} r_{ba}^{ s} & r_{b} + n_{c} + \frac{1}{2} \end{bmatrix} \begin{bmatrix} r_{ba}^{ s} & r_{b} + n_{c} + \frac{1}{2} \end{bmatrix} \begin{bmatrix} r_{ba}^{ s} & r_{b} + n_{c} + \frac{1}{2} \end{bmatrix} \begin{bmatrix} r_{ba}^{ s} & r_{b} + n_{c} + \frac{1}{2} \end{bmatrix} \begin{bmatrix} r_{ba}^{ s} & r_{b} + n_{c} + \frac{1}{2} \end{bmatrix} \begin{bmatrix} r_{ba}^{ s} & r_{b} + n_{c} + \frac{1}{2} \end{bmatrix} \begin{bmatrix} r_{ba}^{ s} & r_{b} + n_{c} + \frac{1}{2} \end{bmatrix} \begin{bmatrix} r_{ba}^{ s} & r_{b} + n_{c} + \frac{1}{2} \end{bmatrix} \begin{bmatrix} r_{ba}^{ s} & r_{b} +$	$\Delta_{2^+ \alpha\beta}^{\#3}$
Γ <sub>2</sub> <sup>#1</sup> αβχ	Symmetry[3, $\Gamma_2^{\pm 1 \oplus 3 \oplus 3 + 3}$ , $\{ \bullet 1 \rightarrow -a, \bullet 2 \rightarrow -b, \bullet 3 \rightarrow -c \}$ , StrongGenSet[{1, 2}, GenSet[-(1, 2)]]	$ \frac{1}{16} \begin{bmatrix} r_{\alpha\beta} + \frac{1}{8} & \Gamma_{\alpha\beta\beta} + \frac{1}{8} & \Gamma_{\beta\alpha} + \frac{1}{8} & \Gamma_{\beta\alpha} + \frac{1}{4} & \Gamma_{\alpha\beta\beta} - \frac{1}{4} & \Gamma_{\beta\beta\alpha} - \frac{1}{4} & \Gamma_{\beta\beta} $	$\Delta_{2}^{\#1}{}_{aB\chi}$
$\Gamma_{2}^{\#2}a_{\beta\chi}$	Symmetry[3, $\Gamma_2^{g_2 \bullet i_2 \bullet 3}$ , $\{\bullet 1 \rightarrow \cdot a, \bullet 2 \rightarrow \cdot b, \bullet 3 \rightarrow \cdot c\}$ , StrongGenSet[{1, 2}, GenSet[-(1,2)]]]	$ \begin{array}{c} \frac{1}{2} \int_{-\infty}^{\infty} g_{+}^{-1} \frac{1}{2} \int_{-\infty}^{\infty} g_{+}^{-1} \int_{-\infty}^{\infty} g_{+}^{-1} \frac{1}{2} \int_{-\infty}^{\infty} g_{+} \frac{1}{2} \int_{-\infty}^{$	$\Delta_2^{\#2}{}_{abx}$
Γ <sup>#1</sup> <sub>3</sub> , <sub>αβχ</sub>	Symmetry[3, r <sup>g1●10,20,3</sup> , {●1→-a, ●2→-b, ●3→-c}, StrongGenSet[{1, 2, 3}, GenSet[(1,2), (2,3)]]]	$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	Δ <sup>#1</sup> <sub>3</sub> , <sub>ηρχ</sub>

FIG. 3. Kinematic structure of the affine connection  $\Gamma_{\mu\nu}^{\ \rho}$ . These definitions are used in Figs. 4 to 8. See [148] for further notational details. This is a vector graphic: all details are visible under magnification.

poles vanish except for that arising from the  $0^-$  sector: the system is now simple enough that the full spectral analysis may be performed and presented. When Eqs. (C1) to (C4), (C5a) and (C5b) are applied to Eq. (25), the resulting particle spectrograph is shown in Fig. 5. The  $0^-$  sector contains a single massive pole, as expected. The outcome of the massless analysis is very much harder to anticipate from the computations above. In this case, the software tells us that there are *no massless particles whatever*. The paucity of propagating d.o.f in the final model means that unitarity is easy to achieve, indeed the no-tachyon and no-ghost conditions are respectively

$$a_1 < 0 < c_1. \tag{C6}$$

As explained in Section IV A, the model we have constructed is simply the square of the Holst operator, with added mass terms.




FIG. 4. Partial particle spectrograph of Eq. (25) after the sequential imposition of Eqs. (C1) and (C2). For brevity, the analysis is halted once the matrix representations of the wave operators and saturated propagators have been computed. Neither the wave operator nor the saturated propagator matrix for the spin-parity  $1^-$  sector are attempted to be computed at this stage, since they are too cumbersome. All quantities are defined in Figs. 2 and 3. See [148] for further notational details. This is a vector graphic: all details are visible under magnification.




FIG. 5. Complete particle spectrograph of Eq. (25) after the sequential imposition of Eqs. (C1) to (C4), (C5a) and (C5b), to be compared with Fig. 4. The saturated propagator matrix for the spin-parity  $1^-$  sector has been omitted due to its large size (though it is still computed). All quantities are defined in Figs. 2 and 3. See [148] for further notational details. This is a vector graphic: all details are visible under magnification.

Integrable gravity and its Maxwell limit — We now turn to the model considered in Section IV B. The condition that all the  $\{d_1, \ldots, d_8\}$  vanish may be imposed with the replacements

$$c_{5} \mapsto 4c_{6} + 2c_{7} + 2c_{8}, \quad c_{4} \mapsto c_{6}, \quad c_{3} \mapsto -c_{6} - c_{7} - c_{8}, \quad c_{2} \mapsto 2c_{6} + c_{7} + c_{8}, \quad c_{16} \mapsto -c_{6} - c_{7} - c_{8}, \\ c_{11} \mapsto -c_{12} - 6c_{6} - 4c_{7} - 4c_{8}, \quad c_{10} \mapsto c_{7} + c_{8} - c_{9}, \quad c_{1} \mapsto c_{6},$$
(C7)

and the condition that the  $\{a_1, \dots, a_{11}\}$  vanish may be imposed with the replacements

$$a_1 \mapsto 0, \quad a_2 \mapsto 0, \quad a_3 \mapsto 0, \quad a_4 \mapsto 0, \quad a_5 \mapsto 0, \quad a_6 \mapsto 0, \quad a_7 \mapsto 0, \quad a_8 \mapsto 0, \quad a_9 \mapsto 0, \quad a_{10} \mapsto 0, \quad a_{11} \mapsto 0.$$
(C8)

When Eqs. (C7) and (C8) are applied to the model in Eq. (25), the resulting partial particle spectrum is shown in Fig. 6. As with Fig. 4, in the first instance, the analysis is performed only up to the computation of the denominators of the propagators for each spin-parity sector. It is clear from Fig. 6 that there will be multiple massive d.o.f, because these denominators contain various roots. Out of necessity, we will eliminate the  $3^-$  and two  $2^-$  d.o.f by sending their square masses to infinite magnitude, and as a means of simplifying the model we do the same for the  $1^+$  mode. Neither of the  $0^+$  or  $0^-$  modes contains a massive pole;

the  $1^-$  mode contains a single massive pole, but the expression for its mass is still slightly cumbersome. The relevant masses are made to diverge by imposing the further replacements

$$c_6 \mapsto 0, \quad c_8 \mapsto -c_7, \quad c_{12} \mapsto c_7 + c_9. \tag{C9}$$

The complete particle spectrograph resulting from the sequential imposition of Eqs. (C7) to (C9) on Eq. (25) is shown in Fig. 7. This full analysis confirms the massive  $1^-$  mode, along with four massless polarisations whose presence was not previously obvious. Two of these polarisations may be associated with the graviton, since their no-ghost conditions depend only on the  $a_0$ -coupling. The other two polarisations presumably have their origin in the massless limit of some higher-spin mode, whose mass was removed by the conditions in Eq. (C8). The expression for the mass of the  $1^-$  mode is now simpler, but it turns out to have a cumbersome pole residue which still hinders the unitarity analysis: to remove this d.o.f entirely we once again take the limit of infinite mass with the final condition

$$c_{15} \mapsto -c_{14}. \tag{C10}$$

The complete particle spectrograph resulting from the sequential imposition of Eqs. (C7) to (C10) on Eq. (25) is shown in Fig. 8. The massless spectra are the same as in Fig. 7, though the extra massless pole residue is somwhat simplified. The total conditions for unitarity can be readily computed as

$$a_0 < 0 < c_{13}.$$
 (C11)

As explained in Section IV B, the model leading to Eq. (C11) can be understood as the square of the homothetic curvature.




FIG. 6. Partial particle spectrograph of Eq. (25) after the sequential imposition of Eqs. (C7) and (C8). For brevity, the analysis is halted once the matrix representations of the wave operators and saturated propagators have been computed. The saturated propagator matrix for the spin-parity  $1^-$  sector has been omitted due to its large size (though it is still computed). All quantities are defined in Figs. 2 and 3. See [148] for further notational details. This is a vector graphic: all details are visible under magnification.






FIG. 7. Complete particle spectrograph of Eq. (25) after the sequential imposition of Eqs. (C7) to (C9), to be compared with Fig. 6. The saturated propagator matrix for the spin-parity  $1^-$  sector has been omitted due to its large size (though it is still computed). All quantities are defined in Figs. 2 and 3. See [148] for further notational details. This is a vector graphic: all details are visible under magnification.




FIG. 8. Complete particle spectrograph of Eq. (25) after the sequential imposition of Eqs. (C7) to (C10), to be compared with Fig. 7. The saturated propagator matrix for the spin-parity  $1^-$  sector has been omitted due to its large size (though it is still computed). All quantities are defined in Figs. 2 and 3. See [148] for further notational details. This is a vector graphic: all details are visible under magnification.