

Non-equatorial scalar rings supported by rapidly spinning Gauss-Bonnet black holes

Shahar Hod

The Ruppin Academic Center, Emeq Hefer 40250, Israel

and

The Hadassah Institute, Jerusalem 91010, Israel

(Dated: December 23, 2024)

Black-hole spacetimes that possess stationary equatorial matter rings are known to exist in general relativity. We here reveal the existence of black-hole spacetimes that support *non*-equatorial matter rings. In particular, it is proved that rapidly-rotating Kerr black holes in the dimensionless large-spin regime $\bar{a} > \bar{a}_{\text{crit}} = \sqrt{\{7 + \sqrt{7} \cos[\frac{1}{3} \arctan(3\sqrt{3})] - \sqrt{21} \sin[\frac{1}{3} \arctan(3\sqrt{3})]\}/12} \simeq 0.78$ can support a pair of non-equatorial massive scalar rings which are negatively coupled to the Gauss-Bonnet curvature invariant of the spinning spacetime (here $\bar{a} \equiv J/M^2$ is the dimensionless angular momentum of the central supporting black hole). We explicitly prove that these non-equatorial scalar rings are characterized by the dimensionless functional relation $-\frac{57+28\sqrt{21} \cos[\frac{1}{3} \arctan(\frac{1}{3\sqrt{3}})]}{8(1+\sqrt{1-\bar{a}^2})^6} \cdot \frac{\bar{\eta}}{\bar{\mu}^2} \rightarrow 1^+$ in the large-mass $\bar{\mu} \equiv M\mu \gg 1$ regime (here $\{\bar{\eta} < 0, \mu\}$ are respectively the non-trivial coupling parameter of the composed Einstein-Gauss-Bonnet-massive-scalar field theory and the proper mass of the supported non-minimally coupled scalar field).

I. INTRODUCTION

The physically important no-hair conjecture [1, 2] for black holes has motivated the mathematical development of elegant non-existence theorems that established the intriguing fact that bound-state static scalar field configurations with minimal [3–6] (as well as with non-minimal [7, 8]) coupling to the spatially-dependent Ricci curvature scalar R cannot be supported in the external regions of asymptotically flat black-hole spacetimes.

However, recent studies [9–12] have explicitly demonstrated that the no-hair conjecture can be violated by asymptotically flat non-vacuum black holes with spatially regular horizons in generalized Einstein-Gauss-Bonnet-scalar field theories that contain a non-trivial interaction term $f(\phi)\mathcal{G}$ between the scalar field ϕ of the theory and the Gauss-Bonnet invariant \mathcal{G} of the curved spacetime [13–20].

The physically intriguing observation [9–12] (see also [21–27]) that black holes in composed Einstein-Gauss-Bonnet-scalar field theories can be spontaneously scalarized is a direct consequence of the fact that the generalized Klein-Gordon differential equation that determines the spatio-temporal functional behavior of the non-minimally coupled scalar fields in the supporting curved black-hole spacetime contains an effective mass term of the form $-\bar{\eta}G$ [here $\bar{\eta}$ is the non-trivial Gauss-Bonnet-scalar-field dimensionless coupling parameter of the theory, see Eq. (8) below] which, depending on the physical parameters of the central supporting black hole, may become negative (thus representing an attractive black-hole-field interaction potential) in the highly curved near-horizon region.

Interestingly, it has been demonstrated numerically [21, 22] and analytically [28] that composed Einstein-Gauss-Bonnet-scalar field theories are characterized by an *existence-line* which marks the sharp boundary between the familiar (scalarless) Kerr black-hole solutions of the Einstein field equations and composed spinning-black-hole-nonminimally-coupled-scalar-field hairy configurations. In particular, it has been demonstrated [21, 22, 28] that the existence-line of the black-hole-field system describes the spin-dependent unique values $\bar{\eta} = \bar{\eta}(\bar{a})$ of the non-trivial Gauss-Bonnet-scalar-field coupling parameter that allow spinning Kerr black holes to support spatially regular bound-state linearized configurations (linearized ‘scalar clouds’ [29, 30]) of the non-minimally coupled matter fields (here $\bar{a} \equiv J/M^2$ is the dimensionless spin parameter of the supporting black hole).

As nicely discussed in [9–12, 21–27], one finds that the critical existence-line of the black-hole-field system is universal in the sense that different Einstein-Gauss-Bonnet-scalar field theories, which are characterized by different non-linear scalar coupling functions $\{f(\phi)\}$ that share the same weak-field functional expansion $f(\phi) = 1 + \bar{\eta}\phi^2/2 + O(\phi^4)$, are characterized by the same critical boundary (critical existence-line) $\bar{\eta} = \bar{\eta}(\bar{a})$ that separates bald (scalarless) Kerr black-hole spacetimes from non-trivial spinning-black-hole-scalar-field hairy spacetime configurations.

Intriguingly, it has recently been proved [22] that, in Einstein-Gauss-Bonnet-scalar field theories in which the scalar field is *negatively* coupled ($\bar{\eta} < 0$) to the Gauss-Bonnet curvature invariant of the spacetime, the spontaneous scalarization phenomenon of spinning Kerr black holes is restricted to the super-critical regime [22, 31, 32]

$$\bar{a} > \frac{1}{2} \tag{1}$$

of the black-hole dimensionless spin parameter. In addition, it has been revealed that, in the $\bar{\eta} < 0$ regime of the non-minimal Gauss-Bonnet-scalar-field coupling parameter, spontaneously scalarized spinning black holes are

characterized by the near-critical *large-coupling* behavior [22, 28]

$$-\bar{\eta} \rightarrow \infty \quad \text{for} \quad \bar{a} \rightarrow \left(\frac{1}{2}\right)_{\text{crit}}^+ . \quad (2)$$

The main goal of the present paper is to explore, using *analytical* techniques, the physical and mathematical properties of composed spinning-Kerr-black-hole-massive-scalar-field cloudy configurations in the negative large-coupling regime. In particular, below we shall explicitly prove that the addition of proper masses to the non-trivially coupled scalar fields allows rapidly-rotating Kerr black holes [see Eq. (18) below] to support a pair of *non-equatorial* scalar rings which are negatively coupled to the Gauss-Bonnet invariant of the curved spacetime. Interestingly, we shall prove that the composed Kerr-black-hole-non-equatorial-scalar-rings cloudy configurations are characterized by the double *large-coupling-large-mass* behavior

$$-\bar{\eta} \rightarrow \infty \quad \text{and} \quad \bar{\mu} \rightarrow \infty \quad (3)$$

with the finite dimensionless ratio

$$-\frac{\bar{\eta}}{\bar{\mu}^2} \rightarrow \text{finite value} , \quad (4)$$

where $\bar{\mu} \equiv M\mu$ is the dimensionless proper mass of the supported non-minimally coupled scalar field.

II. DESCRIPTION OF THE SYSTEM

We study the physical and mathematical properties of ‘cloudy’ black-hole spacetimes which are composed of central spinning Kerr black holes that support spatially regular linearized bound-state configurations of massive scalar fields with a non-trivial (non-minimal) direct coupling to the Gauss-Bonnet invariant,

$$\mathcal{G} \equiv R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma} - 4R_{\mu\nu}R^{\mu\nu} + R^2 , \quad (5)$$

of the curved black-hole spacetime.

The action of the Einstein-Gauss-Bonnet-nonminimally-coupled-massive-scalar field theory is given by the integral expression [11, 12, 33]

$$S = \frac{1}{2} \int d^4x \sqrt{-g} \left[R - \frac{1}{2} \nabla_\alpha \phi \nabla^\alpha \phi - \frac{1}{2} \mu^2 \phi^2 + f(\phi) \mathcal{G} \right] . \quad (6)$$

Following [11, 12, 14], we shall assume that the non-minimal coupling function of the massive scalar field is characterized by the leading order compact functional behavior

$$f(\phi) = \frac{1}{2} \eta \phi^2 , \quad (7)$$

which guarantees that the familiar (scalarless) spinning Kerr black-hole spacetime [see Eq. (9) below] is a valid solution of the coupled Einstein-scalar field equations in the $\phi \rightarrow 0$ limit [11, 12, 14]. The strength of the direct non-minimal interaction between the supported massive scalar field configurations and the Gauss-Bonnet invariant (5) of the curved spacetime is controlled by the dimensionless physical parameter

$$\bar{\eta} \equiv \frac{\eta}{M^2} . \quad (8)$$

The weak-field coupling function (7) of the composed Einstein-Gauss-Bonnet-nonminimally-coupled-massive-scalar field theory (6) may be characterized by either positive [21] or negative [22] values of the non-minimal expansion parameter $\bar{\eta}$.

The supporting Kerr black-hole spacetime of mass M and angular momentum $J \equiv Ma$ [34] is described, using the familiar Boyer-Lindquist coordinates, by the curved line element [35, 36]

$$ds^2 = -\frac{\Delta}{\rho^2} (dt - a \sin^2 \theta d\phi)^2 + \frac{\rho^2}{\Delta} dr^2 + \rho^2 d\theta^2 + \frac{\sin^2 \theta}{\rho^2} [adt - (r^2 + a^2)d\phi]^2 . \quad (9)$$

The spatially-dependent metric functions in (9) are given by the functional expressions

$$\Delta \equiv r^2 - 2Mr + a^2 \quad ; \quad \rho^2 \equiv r^2 + a^2 \cos^2 \theta . \quad (10)$$

The black-hole horizon radii,

$$r_{\pm} = M \pm (M^2 - a^2)^{1/2}, \quad (11)$$

are determined by the radial roots of the metric function $\Delta(r)$.

The supported massive scalar field configurations that we shall analyze in the present paper are assumed to be directly coupled [see the action (6)] to the spatially-dependent Gauss-Bonnet invariant (5) which, in the spinning curved Kerr black-hole spacetime (9) can be expressed in the form [22]

$$\mathcal{G}_{\text{Kerr}}(r, \theta) = \frac{48M^2}{(r^2 + a^2 \cos^2 \theta)^6} \cdot (r^6 - 15a^2 r^4 \cos^2 \theta + 15a^4 r^2 \cos^4 \theta - a^6 \cos^6 \theta). \quad (12)$$

The action (6), which characterizes the composed Einstein-Gauss-Bonnet-nonminimally-coupled-massive-scalar field theory, yields the generalized Klein-Gordon equation [11, 12, 14, 37, 38]

$$\nabla^\nu \nabla_\nu \phi = \mu_{\text{eff}}^2 \phi, \quad (13)$$

where the effective scalar-field-Gauss-Bonnet mass term, whose presence in (13) reflects the direct coupling between the scalar field and the Gauss-Bonnet curvature invariant (5), is given by the spatially-dependent ($\{r, \theta\}$)-dependent functional expression

$$\mu_{\text{eff}}^2(r, \theta; M, a) = \mu^2 - \eta \cdot \mathcal{G}_{\text{Kerr}}(r, \theta). \quad (14)$$

Intriguingly, taking cognizance of Eq. (12), one finds that, depending on the values of the physical parameters $\{M, a, \eta, \mu\}$ that characterize the composed black-hole-field system, the effective mass term (14) may become negative in the exterior region of the black-hole spacetime. In the next section we shall explicitly prove that this physically interesting property of the spatially-dependent effective mass term (14) allows the existence of composed black-hole-massive-scalar-field bound-state configurations in the non-trivial Einstein-Gauss-Bonnet-nonminimally-coupled-massive-scalar field theory (6).

III. NON-EQUATORIAL MASSIVE SCALAR RINGS SUPPORTED BY RAPIDLY-SPINNING KERR BLACK HOLES

In the present section we shall analyze the physical and mathematical properties of cloudy spinning-Kerr-black-hole-massive-scalar-field configurations that mark the *onset* of the spontaneous scalarization phenomenon in the composed Einstein-Gauss-Bonnet-nonminimally-coupled-massive-scalar field theory (6). In particular, we shall reveal the physically interesting fact that for negative values,

$$\bar{\eta} < 0, \quad (15)$$

of the non-trivial dimensionless coupling parameter of the theory, rapidly-rotating black holes can support *non-equatorial* thin matter rings which are made of the non-minimally coupled massive scalar fields.

To this end, we first point out that one deduces from Eq. (12) that, depending on the relative magnitudes of the physical parameters η and μ that characterize the non-trivial field theory (6), the functional expression (14) for the effective mass term of the scalar field may become negative (thus representing an *attractive* black-hole-field binding potential well) in the exterior region of the spinning Kerr black-hole spacetime. This physically interesting property of the composed Einstein-Gauss-Bonnet-nonminimally-coupled-massive-scalar field theory allows the central black hole to support bound-state configurations of the non-minimally coupled massive scalar fields [21, 31, 37, 39].

In particular, in the dimensionless large-mass regime

$$\bar{\mu} \equiv M\mu \gg 1 \quad (16)$$

of the composed black-hole-field system [or equivalently, in the regime $-\bar{\eta} \gg 1$ of large negative values of the non-minimal coupling parameter, see Eq. (22) below], the *onset* of the spontaneous scalarization phenomenon in the non-trivial field theory (6) is marked by the critical functional behavior [31, 39, 40]

$$\min\{\mu_{\text{eff}}^2(r, \theta; M, a)\} \rightarrow 0^- \quad (17)$$

of the spatially-dependent effective mass term that characterizes the generalized Klein-Gordon equation (13) of the non-minimally coupled scalar field.

Performing a functional analysis of the two-dimensional ($\{r, \theta\}$ -dependent) expression (12) one finds that, for rapidly-rotating Kerr black holes in the dimensionless spin regime

$$\bar{a} \equiv \left(\frac{a}{M}\right) > \left(\frac{a}{M}\right)_{\text{crit}} = \sqrt{\frac{7 + \sqrt{7} \cos \left[\frac{1}{3} \arctan(3\sqrt{3})\right] - \sqrt{21} \sin \left[\frac{1}{3} \arctan(3\sqrt{3})\right]}{12}}, \quad (18)$$

the Kerr Gauss-Bonnet curvature invariant is characterized by the extremum relation

$$\min \left\{ M^4 \cdot \mathcal{G}_{\text{Kerr}}(r \in [r_+, \infty], \cos^2 \theta; \bar{a}) \right\} = -\frac{57 + 28\sqrt{21} \cos \left[\frac{1}{3} \arctan\left(\frac{1}{3\sqrt{3}}\right)\right]}{8(1 + \sqrt{1 - \bar{a}^2})^6}, \quad (19)$$

where the spin-dependent minimum (19) is characterized by the polar angular relation [41]

$$(\cos^2 \theta)_{\min} = \frac{(1 + \sqrt{1 - \bar{a}^2})^2}{\bar{a}^2} \cdot \left\{ 7 - 4\sqrt{7} \sin \left[\frac{1}{3} \arctan\left(\frac{1}{3\sqrt{3}}\right)\right] - 4\sqrt{\frac{7}{3}} \cos \left[\frac{1}{3} \arctan\left(\frac{1}{3\sqrt{3}}\right)\right] \right\} < 1 \quad (20)$$

with

$$r_{\min} \rightarrow r_+(M, a). \quad (21)$$

It is interesting to note that the analytically derived expression (20) for $(\cos^2 \theta)_{\min}$ is a monotonically decreasing function of the black-hole dimensionless spin parameter \bar{a} in its regime of validity (18). The data presented in Table I clearly demonstrates the simple *monotonic* functional behavior of the spin-dependent angular function $(\cos^2 \theta)_{\min}$ [see Eq. (20)] in the super-critical regime (18). In particular, the value of $|\theta_{\min}(\bar{a})|$ increases monotonically from $\theta_{\min}(\bar{a} = \bar{a}_{\text{crit}}) = 0^\circ$ to $|\theta_{\min}(\bar{a} = 1)| = 61.21^\circ$.

\bar{a}	0.79	0.82	0.85	0.88	0.91	0.94	0.97	1.00
$[(\cos^2 \theta)_{\min}](\bar{a})$	0.967	0.853	0.748	0.652	0.560	0.472	0.381	0.232

TABLE I: Non-equatorial scalar rings supported by rapidly spinning Gauss-Bonnet black holes. We display, for various values of the black-hole dimensionless spin parameter \bar{a} in the super-critical regime (18), the values of $[(\cos^2 \theta)_{\min}](\bar{a})$ as calculated from the analytically derived formula (20). The data presented demonstrates the simple *monotonic* functional behavior of the angular function (20). In particular, one finds that the value of $|\theta_{\min}|$ increases monotonically from $\theta_{\min}(\bar{a} = \bar{a}_{\text{crit}}) = 0^\circ$ to $|\theta_{\min}(\bar{a} = 1)| = 61.21^\circ$.

From Eqs. (18) and (19) one deduces that, in the dimensionless large-mass (or equivalently, large-coupling $-\bar{\eta} \gg 1$) regime (16) with the critical relation [42]

$$-\frac{57 + 28\sqrt{21} \cos \left[\frac{1}{3} \arctan\left(\frac{1}{3\sqrt{3}}\right)\right]}{8(1 + \sqrt{1 - \bar{a}^2})^6} \cdot \frac{\bar{\eta}}{\bar{\mu}^2} \rightarrow 1^+, \quad (22)$$

the effective mass term (14) that appears in the scalar Klein-Gordon equation (13) of the composed black-hole-field system becomes *negative* (thus representing an attractive black-hole-field potential well) in a pair of narrow *non-equatorial* rings which are characterized by the polar angular relation (20).

This physically intriguing property of the composed Einstein-Gauss-Bonnet-nonminimally-coupled-massive-scalar field theory (6) implies that, in the large-mass (large-coupling) regime (16), rotating Kerr black holes in the dimensionless large-spin regime (18) can support pairs of non-equatorial thin matter rings [see Eqs. (20) and (21)] which are made of massive scalar fields that are negatively coupled to the Gauss-Bonnet curvature invariant of the black-hole spacetime.

In particular, the analytically derived functional expression (22) determines the spin-dependent critical existence-line $\bar{\eta} = \bar{\eta}(\bar{\mu}, \bar{a})$ that characterizes the onset of the spontaneous scalarization phenomenon in the composed Einstein-Gauss-Bonnet-nonminimally-coupled-massive-scalar field theory (6) in the dimensionless large-coupling-large-mass ($-\bar{\eta} \gg 1, \bar{\mu} \gg 1$) regime.

IV. THE CLASSICALLY ALLOWED REGION OF THE NON-EQUATORIAL NEAR-CRITICAL MASSIVE SCALAR RINGS

In the present section we shall use analytical techniques in order to determine the classically allowed spatial region of the supported non-minimally coupled massive scalar rings in the near-critical regime (22). To this end, it proves

useful to use the dimensionless physical parameters $\{\epsilon, x, y\}$ which are defined by the following near-critical relations [see Eqs. (20), (21), and (22)]

$$\frac{\bar{\eta}}{\bar{\mu}^2} = \left(\frac{\bar{\eta}}{\bar{\mu}^2}\right)_{\text{crit}} \cdot (1 + \epsilon) \quad ; \quad \left(\frac{\bar{\eta}}{\bar{\mu}^2}\right)_{\text{crit}} = -\frac{8(1 + \sqrt{1 - \bar{a}^2})^6}{57 + 28\sqrt{21} \cos\left[\frac{1}{3} \arctan\left(\frac{1}{3\sqrt{3}}\right)\right]} \quad ; \quad 0 \leq \epsilon \ll 1, \quad (23)$$

$$\cos^2 \theta = (\cos^2 \theta)_{\text{min}} \cdot (1 + x) \quad ; \quad x \ll 1, \quad (24)$$

and

$$r = r_+ \cdot (1 + y) \quad ; \quad 0 \leq y \ll 1. \quad (25)$$

Substituting the functional relations (23), (24), and (25) into the expression (14) for the spatially-dependent [see (12)] effective mass term that characterizes the composed spinning-Kerr-black-hole-nonminimally-coupled-massive-scalar-field system, one finds that the black-hole-field effective interaction potential is negative (thus representing an *attractive* black-hole-field binding potential) in a pair of narrow non-equatorial rings which are characterized by the dimensionless functional relation

$$\gamma x^2 + 6y \leq \epsilon, \quad (26)$$

where

$$\gamma \equiv -\frac{6\beta^2(\beta^3 - 27\beta^2 + 63\beta - 21)}{(1 + \beta)^2(\beta^3 - 15\beta^2 + 15\beta - 1)} > 0 \quad (27)$$

with

$$\beta \equiv 7 - 4\sqrt{7} \sin\left[\frac{1}{3} \arctan\left(\frac{1}{3\sqrt{3}}\right)\right] - 4\sqrt{\frac{7}{3}} \cos\left[\frac{1}{3} \arctan\left(\frac{1}{3\sqrt{3}}\right)\right]. \quad (28)$$

The functional relation (26) with the definitions (24) and (25) determine, in the near-critical regime (23), the classically allowed spatial region for the spontaneous scalarization phenomenon of negatively-coupled massive scalar fields in rapidly-spinning Kerr black-hole spacetimes [see (18)].

Interestingly, from Eqs. (24), (25), and (26) one deduces that, for a given value of the dimensionless near-critical parameter ϵ [with $\epsilon \ll 1$, see Eq. (23)], the maximum effective angular width of the supported non-equatorial scalar rings is given by the functional relation [43]

$$[\Delta(\cos^2 \theta)]_{\text{ring}} = \frac{(1 + \sqrt{1 - \bar{a}^2})^2}{\bar{a}^2} \cdot \sqrt{-\frac{2(1 + \beta)^2(\beta^3 - 15\beta^2 + 15\beta - 1)}{3(\beta^3 - 27\beta^2 + 63\beta - 21)}} \cdot \sqrt{\epsilon} \quad ; \quad \epsilon \ll 1, \quad (29)$$

and the maximum effective radial width of the supported scalar rings is given by the relation [44]

$$(\Delta r)_{\text{ring}} = \frac{1}{6} \cdot r_+ \cdot \epsilon \quad ; \quad \epsilon \ll 1. \quad (30)$$

V. SUMMARY AND DISCUSSION

The recently published physically interesting studies [9–12] (see also [21–27]) of the composed Einstein-Gauss-Bonnet-scalar field theory have revealed that black holes with spatially regular horizons can support hairy matter configurations which are made of scalar fields with a non-trivial (non-minimal) coupling to the spatially dependent Gauss-Bonnet invariant \mathcal{G} of the curved spacetime.

Intriguingly, it has been demonstrated [21, 22] that, for spinning black holes in Einstein-Gauss-Bonnet-scalar field theories, the sharp boundary between the familiar (scalarless) Kerr black-hole spacetime and spontaneously scalarized hairy black-hole spacetimes is determined by the presence of cloudy black-hole-field configurations, Kerr black holes that support linearized bound-state configurations of the spatially regular non-minimally coupled scalar fields. In particular, the composed black-hole-field cloudy configurations determine a universal [that is, independent of the non-linear functional behavior of the scalar coupling function $f(\phi)$, see (6)] critical existence-line $\bar{\eta}_{\text{crit}} = \bar{\eta}_{\text{crit}}(\bar{a})$ that

describes the non-trivial spin-dependent functional behavior of the non-minimal coupling parameter that characterizes the composed Einstein-Gauss-Bonnet-scalar field theory.

In the present paper we have used *analytical* techniques in order to explore the onset of the negative-coupling ($\bar{\eta} < 0$) spontaneous scalarization phenomenon of spinning Kerr black-hole spacetimes in the large-coupling-large-mass regime

$$\{-\bar{\eta}, \bar{\mu}\} \rightarrow \infty \quad \text{with} \quad -\frac{\bar{\eta}}{\bar{\mu}^2} \rightarrow \text{finite value} . \quad (31)$$

The main results derived in this paper and their physical implications are as follows:

(1) We have revealed the physically interesting fact that in the Einstein-Gauss-Bonnet-nonminimally-coupled-massive-scalar field theory (6), rapidly-rotating Kerr black holes in the large-spin regime

$$\bar{a} > \bar{a}_{\text{crit}} \quad (32)$$

with [see Eq. (18)]

$$\bar{a}_{\text{crit}} = \sqrt{\frac{7 + \sqrt{7} \cos \left[\frac{1}{3} \arctan(3\sqrt{3}) \right] - \sqrt{21} \sin \left[\frac{1}{3} \arctan(3\sqrt{3}) \right]}{12}} \simeq 0.78 \quad (33)$$

can support a pair of *non*-equatorial thin matter rings.

(2) It has been shown that the supported non-minimally coupled massive scalar rings are characterized by the analytically derived dimensionless angular relation [see Eqs. (11) and (22)]

$$\left(\frac{a}{r_+}\right)^2 \cdot (\cos^2 \theta)_{\text{ring}} = 7 - 4\sqrt{7} \sin \left[\frac{1}{3} \arctan\left(\frac{1}{3\sqrt{3}}\right) \right] - 4\sqrt{\frac{7}{3}} \cos \left[\frac{1}{3} \arctan\left(\frac{1}{3\sqrt{3}}\right) \right] . \quad (34)$$

(3) We have proved that the negatively-coupled non-equatorial scalar rings are characterized by the large-mass-large-coupling critical ratio [see Eqs. (11) and (22)]

$$\left(-\frac{\bar{\eta}}{\bar{\mu}^2}\right)_{\text{crit}} \rightarrow \left\{ \frac{8}{57 + 28\sqrt{21} \cos \left[\frac{1}{3} \arctan\left(\frac{1}{3\sqrt{3}}\right) \right]} \cdot \left(\frac{r_+}{M}\right)^6 \right\}^+ . \quad (35)$$

It is worth emphasizing again that the physical significance of the analytically derived spin-dependent dimensionless functional relation (35), which characterizes the composed Kerr-linearized-massive-scalar-field cloudy configurations, stems from the fact that this critical existence-line determines the sharp boundary between bald Kerr black holes and spontaneously scalarized hairy spinning black holes in the Einstein-Gauss-Bonnet-nonminimally-coupled-massive-scalar field theory (6).

(4) It has been shown that, in the near-critical regime

$$\frac{\bar{\eta}}{\bar{\mu}^2} = \left(\frac{\bar{\eta}}{\bar{\mu}^2}\right)_{\text{crit}} \cdot (1 + \epsilon) \quad ; \quad 0 \leq \epsilon \ll 1 \quad (36)$$

of the composed Einstein-Gauss-Bonnet-nonminimally-coupled-massive-scalar field theory (6), the maximum effective spatial widths of the supported non-equatorial scalar rings are characterized by the small- ϵ dimensionless functional relations [see Eqs. (29) and (30)]

$$\frac{\bar{a}^2}{(1 + \sqrt{1 - \bar{a}^2})^2} \cdot [\Delta(\cos^2 \theta)]_{\text{ring}} = \sqrt{-\frac{2(1 + \beta)^2(\beta^3 - 15\beta^2 + 15\beta - 1)}{3(\beta^3 - 27\beta^2 + 63\beta - 21)}}} \cdot \sqrt{\epsilon} \quad ; \quad \frac{(\Delta r)_{\text{ring}}}{r_+} = \frac{1}{6} \cdot \epsilon . \quad (37)$$

ACKNOWLEDGMENTS

This research is supported by the Carmel Science Foundation. I would like to thank Yael Oren, Arbel M. Ongo, Ayelet B. Lata, and Alona B. Tea for helpful discussions.

[1] R. Ruffini and J. A. Wheeler, *Physics Today* **24**, 30 (1971).

- [2] J. D. Bekenstein, Contribution to the 2nd International Sakharov Conference on Physics, 216-219 [e-Print: gr-qc/9605059].
- [3] J. D. Bekenstein, Phys. Rev. D **5**, 1239 (1972).
- [4] T. P. Sotiriou, Class. Quant. Grav. **32**, 214002 (2015).
- [5] C. A. R. Herdeiro and E. Radu, Int. J. Mod. Phys. D **24**, 1542014 (2015).
- [6] T. P. Sotiriou and V. Faraoni, Phys. Rev. Lett. **108**, 081103 (2012).
- [7] A. E. Mayo and J. D. Bekenstein and, Phys. Rev. D **54**, 5059 (1996).
- [8] S. Hod, Phys. Lett. B **771**, 521 (2017) [arXiv:1911.08371]; S. Hod, Phys. Rev. D **96**, 124037 (2017) [arXiv:2002.05903].
- [9] T. P. Sotiriou and S.-Y. Zhou, Phys. Rev. Lett. **112**, 251102 (2014); T. P. Sotiriou and S.-Y. Zhou, Phys. Rev. D **90**, 124063 (2014).
- [10] T. P. Sotiriou, Lect. Notes Phys. **892**, 3 (2015) [arXiv:1404.2955].
- [11] D. D. Doneva and S. S. Yazadjiev, Phys. Rev. Lett. **120**, 131103 (2018).
- [12] H. O. Silva, J. Sakstein, L. Gualtieri, T. P. Sotiriou, and E. Berti, Phys. Rev. Lett. **120**, 131104 (2018).
- [13] See [14–20] for the physically related case of charged black hole in composed Einstein-Maxwell-scalar field theories that can support spatially regular bound-state configurations of static scalar fields which are directly (non-minimally) coupled to the spatially-dependent Maxwell electromagnetic invariant.
- [14] C. A. R. Herdeiro, E. Radu, N. Sanchis-Gual, and J. A. Font, Phys. Rev. Lett. **121**, 101102 (2018).
- [15] P. G. S. Fernandes, C. A. R. Herdeiro, A. M. Pombo, E. Radu, and N. Sanchis-Gual, Class. Quant. Grav. **36**, 134002 (2019) [arXiv:1902.05079].
- [16] S. Hod, Phys. Lett. B **798**, 135025 (2019) [arXiv:2002.01948].
- [17] S. Hod, Phys. Rev. D **101**, 104025 (2020) [arXiv:2005.10268].
- [18] M. Khodadi, A. Allahyari, S. Vagnozzi, and D. F. Mota, JCAP **09**, 026 (2020) [arXiv:2005.05992].
- [19] S. Hod, The Euro. Phys. Jour. C **80**, 1150 (2020).
- [20] S. Hod, Jour. of High Energy Phys. **08**, 272 (2022) [arXiv:2206.12074].
- [21] P. V. P. Cunha, C. A. R. Herdeiro, and E. Radu, Phys. Rev. Lett. **123**, 011101 (2019).
- [22] A. Dima, E. Barausse, N. Franchini, and T. P. Sotiriou, Phys. Rev. Lett. **125**, 231101 (2020) [arXiv:2006.03095].
- [23] S. Hod, Phys. Rev. D **100**, 064039 (2019) [arXiv:1912.07630].
- [24] S. Hod, The Euro. Phys. Jour. C **79**, 966 (2019).
- [25] D. D. Doneva, L. G. Collodel, C. J. Krüger, and S. S. Yazadjiev, Phys. Rev. D **102**, 104027 (2020).
- [26] C. A. R. Herdeiro, E. Radu, H. O. Silva, T. P. Sotiriou, and N. Yunes, Phys. Rev. Lett. **126**, 011103 (2021) [arXiv:2009.03904].
- [27] E. Berti, L. G. Collodel, B. Kleihaus, and J. Kunz, Phys. Rev. Lett. **126**, 011104 (2021).
- [28] S. Hod, Phys. Rev. D **105**, 024074 (2022).
- [29] S. Hod, Phys. Rev. D **86**, 104026 (2012) [arXiv:1211.3202]; S. Hod, The Euro. Phys. Journal C **73**, 2378 (2013) [arXiv:1311.5298]; S. Hod, Phys. Rev. D **90**, 024051 (2014) [arXiv:1406.1179].
- [30] C. A. R. Herdeiro and E. Radu, Phys. Rev. Lett. **112**, 221101 (2014).
- [31] S. Hod, Phys. Rev. D **102**, 084060 (2020) [arXiv:2006.09399].
- [32] It is worth noting that, as explicitly proved in [31], the critical black-hole spin (1) corresponds to the regime $l \gg 1$. In particular, one finds [31] that for smaller l values the critical rotation parameter that allows for the existence of negatively coupled scalar clouds is larger than the critical value (1).
- [33] We use natural units in which $G = c = \hbar = 1$. Note that in these units the mass μ of the non-minimally coupled scalar field has the dimensions of length⁻¹. The black-hole-scalar-field composed mass parameter $\bar{\mu} \equiv M\mu$ is dimensionless.
- [34] We shall assume, without loss of generality, the relation $\bar{a} \equiv J/M^2 > 0$ for the angular momentum parameter of the supporting Kerr black hole.
- [35] C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation*, (W. H. Freeman, San Francisco, 1973).
- [36] S. Chandrasekhar, *The Mathematical Theory of Black Holes*, (Oxford University Press, New York, 1983).
- [37] D. D. Doneva, L. G. Collodel, C. J. Krüger, S. S. Yazadjiev, The Eur. Phys. Jour. C **80**, 1205 (2020) [arXiv:2009.03774].
- [38] It is worth emphasizing that the fact that the non-minimally coupled scalar field has a finite mass allows one to obtain non-equatorial scalar rings whose widths can be made arbitrarily thin. In particular, we shall explicitly prove below that, by adjusting properly the mass term of the non-minimally coupled scalar field [see Eq. (22) below], the classically allowed region of the composed black-hole-field effective potential term (14) [the region in which the effective mass term (14) is negative and therefore represents a binding potential well] can be made arbitrarily narrow, thus allowing for the existence of arbitrarily thin non-equatorial scalar rings which are characterized by the property (17).
- [39] C. A. R. Herdeiro, A. M. Pombo, and E. Radu, Universe **7**, 483 (2021) [arXiv:2111.06442].
- [40] S. Hod, Jour. of High Energy Phys. **02**, 039 (2022) [arXiv:2201.03503].
- [41] Note that the inequality in (20) is valid in the super-critical regime (18).
- [42] Note that the critical dimensionless ratio (22) corresponds to the fundamental composed black-hole-field configuration. The composed black-hole-field system also has excited bound states which are characterized by the near-critical relation (23) with $\epsilon > 0$.
- [43] Note that the angular width is defined as the angular interval in which the effective mass term (14) in the scalar equation (13) is negative and therefore represents an attractive binding potential well of the composed black-hole-field system.
- [44] Note that the radial width is defined as the radial interval in which the effective mass term (14) is negative and therefore represents an attractive black-hole-field binding potential well.