# Weak Hopf non-invertible symmetry-protected topological spin liquid and lattice realization of (1+1)D symmetry topological field theory

# Zhian Jia<sup>0</sup><sup>*a,b,\**</sup>

<sup>a</sup> Centre for Quantum Technologies, National University of Singapore, SG 117543, Singapore <sup>b</sup> Department of Physics, National University of Singapore, SG 117543, Singapore \*Email: giannjia@foxmail.com

ABSTRACT: We propose weak Hopf symmetry as a general framework to explore (1 + 1)D topological phases that exhibit non-invertible symmetries. Inspired by the Symmetry Topological Field Theory (SymTFT) description of quantum phases with non-invertible symmetry, we construct a lattice model by introducing two distinct topological boundary conditions for a weak Hopf lattice gauge theory. One boundary encodes the topological symmetry information, while the other incorporates the non-topological dynamics. The resulting model is termed the cluster ladder model. We demonstrate that the cluster state model is a special case of this broader class of lattice models exhibiting weak Hopf symmetry  $H \times \hat{H}$ , where H is a weak Hopf algebra and  $\hat{H}$  is its dual weak Hopf algebra. On a closed manifold, the symmetry reduces to  $Cocom(H) \times Cocom(\hat{H})$ , corresponding to the cocommutative subalgebras of  $H \times \hat{H}$ . An essential weak Hopf sub-symmetry is  $Cocom(H) \times \text{Rep}(H)$ , which, in the finite group case, reduces to the familiar symmetry  $G \times \text{Rep}(G)$ . To exactly solve the lattice model, we introduce a weak Hopf tensor network. Furthermore, we demonstrate how to construct the lattice realization of an arbitrary fusion category symmetry S via combining Tannaka–Krein reconstruction or weak Hopf tube algebra and the cluster ladder model.

KEYWORDS: Anyon, Quantum Symmetry, Topological State of Matter, Topological Field Theory, Symmetry-Protected Topological (SPT) Phase

## Contents

1	Intr	oduction	1				
2	War	m-up example: finite-group quantum cluster ladder	4				
	2.1	Abelian cluster ladder model	7				
	2.2	S3 cluster ladder model	8				
3	Weak Hopf non-invertible symmetry						
	3.1	Weak Hopf algebra symmetry and weak Hopf comodule algebra symmetry	10				
		3.1.1 Hopf symmetry	10				
		3.1.2 Weak Hopf symmetries	13				
	3.2	Fusion category symmetry as dual weak Hopf symmetry	16				
	3.3	Weak Hopf symmetric single qudit system	20				
4	Pair	ing-based weak Hopf tensor network states	22				
5	Symmetry TFT for weak Hopf lattice gauge theory						
	5.1	Weak Hopf lattice gauge theory	24				
	5.2	Algebraic theory of SymTFT for weak Hopf lattice gauge theory	27				
6	Wea	k Hopf cluster ladder model I: smooth-rough construction	28				
	6.1	Weak Hopf cluster state model	28				
	6.2	Weak Hopf cluster state	29				
	6.3	Symmetry of the model	30				
7	Weak Hopf cluster ladder model II: general construction						
	7.1	Weak Hopf cluster ladder model	34				
	7.2	Ground sate degeneracy	37				
	7.3	Solving the model via tensor network	38				
8	Disc	cussion	40				

# 1 Introduction

The symmetry-protected topological (SPT) phases [1–11] have garnered significant attention in recent years. The ground state of an SPT phase is short-range entangled, and its topological order is characterized as an invertible topological order [12]. This implies that, on any closed manifold, SPT phases have a unique ground state. Distinct SPT phases with a given symmetry cannot be smoothly deformed into each other without undergoing a phase transition, provided the deformation preserves the symmetry. However, if the symmetry is completely broken during the deformation, the SPT states can all be smoothly deformed into a product state. The SPT phases are of particular interest due to their boundary behavior, the boundary of a SPT state is either gapless or degenerate. The boundary effective theory of a non-trivial SPT state always exhibits a pure gauge anomaly or a mixed gauge-gravity anomaly with respect to the symmetry group [13]. The classification of *d*-dimensional bosonic SPT phases with an internal symmetry group *G* is described by the group cohomology  $H^{d+1}(G, U(1))$  [13, 14]<sup>1</sup>. However, it is known that some bosonic SPT phases cannot be captured by this cohomology-based framework [15–17]. This limitation has motivated a broader classification approach based on cobordism theory, in which SPT phases are characterized by the cobordism group  $\text{Hom}(\Omega_{d+1}^G, U(1))$ . Here,  $\Omega_{d+1}^G$  denotes the bordism group of (d + 1)-dimensional manifolds with a *G*-structure [18–21]. For fermionic SPT phases, a complete classification of non-interacting cases has been achieved via K-theory [22, 23]. In the case of interacting fermionic SPT phases, alternative frameworks based on group supercohomology [24] and cobordism theory [25] have been proposed.

On the other hand, significant advancements have been made in the study of generalized symmetries, which encompass new and novel forms of symmetries. Notable examples include high-form symmetries [26–29], non-invertible symmetries [29–36], fusion category symmetries [37–46], algebraic higher symmetries [47], and Hopf and weak Hopf symmetries [48–54]. Category theory provides a general framework for us to investigate these generalized symmetries. The SPT phases protected by these generalized symmetries have attracted considerable attention [35, 47, 55–64].

The concept of categorical symmetry can be understood through a holographic perspective [47, 64, 65]: an *n*-dimensional categorical symmetry corresponds to a topological order in one higher dimension. This approach, now referred to as symmetry topological field theory (SymTFT) or topological holography, provides a comprehensive framework for understanding non-invertible or categorical symmetries in both gapped and gapless phases [35, 47, 56–64, 66, 67].

Consider an *n*-dimensional topological order, the (n - 1)-dimensional defects can be regarded as 1-morphisms, (n-2)-dimensional defects between (n-1)-dimensional defects can be regarded as 2morphisms, and so on. Thus, the corresponding topological order is described by a fusion *n*-category [47, 68-70]. We can combine all d-dimensional topological orders to form a category  $TO_d$ . For this category, we need to include domain walls between different topological orders and the stacking operation of topological orders to ensure that **TO**<sub>d</sub> is closed. From Hamiltonian point of view, consider the moduli space  $\mathfrak{M}_{nd}$  formed by all nd gapped liquid Hamiltonians. An elements in  $\pi_0(\mathfrak{M}_{nd})$  is a topological order [71–73]. Two topological orders are considered equivalent if they can be smoothly deformed into each other through a continuous path of Hamiltonians without closing the energy gap. These equivalence classes of topological orders are described first fundamental group  $\pi_1(\mathfrak{M})$ . From algebraic point of view, two topological orders are equivalent if they can be connected by some invertible domain wall, the invertible domain walls are classified by  $\pi_1(\mathfrak{M})$ . The (n-1)-dimensional defects can fuse but not braid. If we exclude these defects, we obtain a braided fusion (n-1)-category of excitations. This braided fusion category is the looping of the *n*-fusion category of topological order [47, 70]. A special type of topological order is called an invertible topological order C, which is invertible under the stacking operation. Specifically, there exists a topological order  $\mathcal D$  such that the stacking of  $\mathcal C$  and  $\mathcal D$  yields the trivial topological order:  $\mathbb{C} \boxtimes \mathbb{D} \simeq \mathbb{J}$ . The SPT phases possess exclusively invertible topological order, which guarantees the unique ground state of the SPT phase on any closed manifold.

<sup>&</sup>lt;sup>1</sup>In this work, *d* denotes the spatial dimension, while *D* represents the spacetime dimension.

For an *n*-dimensional system, its categorical symmetry is characterized by a fusion *n*-category S. From a topological holography perspective, this fusion *n*-symmetry S corresponds to an (n+1)-dimensional topological order, which is described by the Drinfeld center Z(S). Using the SymTFT sandwich construction [35, 47, 56–64, 66, 67], this relationship can be understood as follows: the symmetry  $S = B_{sym}$  represents a topological boundary condition of the topological field theory Z(S). Additionally, there may exist another boundary,  $\mathcal{B}_{phys}$ , of Z(S), which encodes the non-topological aspects of the system. The boundary  $\mathcal{B}_{phys}$  can be either gapped or gapless. For an n = 1-dimensional system, the categorical symmetry is described by a fusion category S (fusion 1-category). The symmetry is considered anomaly-free if there exists a 1-dimensional S-SPT phase; otherwise, it is referred to as anomalous. For an anomaly-free S, the S-SPT phases are classified by fiber functors  $F : S \rightarrow \text{Vect}$ , which are tensor functors mapping S to the category of finite-dimensional vector spaces [42].

An interesting and challenging question is whether it is possible to construct a lattice model that realizes a given S-SPT phase [45, 57, 62, 74–87]. Despite the numerous examples of S-symmetric phases, a systematic approach for constructing such models remains elusive. In this work, we address this question by exploring it within the framework of weak Hopf symmetry.

For a (1 + 1)D system, the categorical symmetry is described by a (multi)fusion category S. It has been proven that any (multi)fusion category can be realized as the representation category of a weak Hopf algebra [88–90]. Motivated by this result, we introduce weak Hopf symmetries and use them to construct lattice models by generalizing the cluster state model.

The cluster state model is a widely studied example of lattice models exhibiting SPT phases. For the standard  $\mathbb{Z}_2$  cluster state, the Hamiltonian takes the form

$$H_{\text{cluster}} = -\sum_{i} Z_{i-1} X_i Z_{i+1},$$
 (1.1)

which is known to host  $\mathbb{Z}_2 \times \mathbb{Z}_2$  SPT order [91]. By applying Hadamard gates to all even vertices of the lattice, one can map this system to the Calderbank-Shor-Steane (CSS)-type cluster state [54, 78, 92]. The stabilizers for the transformed state are of the CSS-type, specifically  $X_{i-1}X_iX_{i+1}$  for odd sites ( $i \in 2\mathbb{N} + 1$ ) and  $Z_{i-1}Z_iZ_{i+1}$  for even sites ( $i \in 2\mathbb{N}$ ). This leads to a modified Hamiltonian of the form

$$H = -\sum_{i \in 2\mathbb{N}} Z_{i-1} Z_i Z_{i+1} - \sum_{i \in 2\mathbb{N}+1} X_{i-1} X_i X_{i+1}.$$
 (1.2)

Recent studies have revealed that the cluster state model exhibits non-invertible global symmetries. In particular, the symmetry can be described by the fusion category  $\text{Rep}(D_8)$ , where  $D_8$  is the dihedral group of order 8 [85].

Since the qubit-based cluster state corresponds to systems valued in the Abelian group  $\mathbb{Z}_2$ , a natural generalization is to consider systems defined over arbitrary finite groups G [92]. This generalization involves introducing higher-dimensional analogs of Pauli operators, similar to those found in Kitaev's quantum double model [93, 94]. The SPT order and non-invertible symmetries of such finite-group cluster states were recently analyzed in Ref. [78]. Using tensor network techniques, it was demonstrated that the symmetry of a G-valued cluster state is described by  $G \times \text{Rep}(G)$ , where Rep(G) is the representation category of G. Additionally, it was shown in Ref. [54] that the cluster state model can be understood as a specific realization of a quantum double ladder model.

In this work, we systematically construct lattice models that realize an S-symmetric phase. To achieve this, we introduce a general framework for weak Hopf symmetric phases and demonstrate how such phases can be realized via SymTFT. We also show that the (multi)fusion symmetry can be interpreted as a dual weak Hopf symmetry. Building on the observation that a cluster state SPT phase can be viewed as a thin quantum double model with one smooth and one rough boundary [54], we introduce the weak Hopf cluster state model. Inspired by this construction, we further propose a generalized model, termed the weak Hopf cluster ladder model, where the two topological boundaries are chosen to be distinct and do not allow any non-trivial anyonic tunneling channels between them. A crucial observation is that the symmetry of the (1+1)D cluster ladder model is given by  $H \times \hat{H}$ , where H represents a weak Hopf symmetry and  $\hat{H}$  is its dual weak Hopf symmetry. Notably, the  $\mathbb{Z}_2 \times \mathbb{Z}_2$  symmetry of the  $\mathbb{Z}_2$ -cluster state, the  $G \times \text{Rep}(G)$  symmetry of the G-cluster state model, and the  $\text{Cocom}(H) \times \text{Rep}(H)$  symmetry are all special cases of this general framework. Also note that our model is quite different from that based on anyonic chains [57, 62, 79–81], where the Hilbert space is constructed from the fusion tree and thus does not have a well-defined local space or tensor product structure <sup>2</sup>.

The paper is organized as follows. In Section 2, we use finite groups as examples to illustrate the key ideas of our lattice model. For the  $\mathbb{Z}_p$  group with p a prime number, there exists a single type of cluster ladder model, which corresponds to the  $\mathbb{Z}_p$  cluster state model. In contrast, for the  $S_3$  group, two distinct types of cluster ladder models that realize SPT phases arise: one corresponds to the cluster state model, while the other exhibits topological boundary conditions that differ from the conventional smooth and rough boundaries. In Section 3, we provide a detailed discussion of weak Hopf symmetries, introducing the necessary mathematical formalism and results. We begin with the case of invertible symmetries, which correspond to group algebra Hopf symmetries, and then extend the analysis to general weak Hopf symmetries and comodule algebra symmetries. Furthermore, we demonstrate how fusion category symmetries can be interpreted as dual weak Hopf symmetries. Finally, we explore the implications of weak Hopf symmetries for single-qudit systems. Section 4 introduces weak Hopf tensor network states, which serve as a powerful tool for analyzing and solving our lattice model. In Section 6, we present the weak Hopf generalization of the cluster state model. The model is solved using weak Hopf tensor network states. We explore the weak Hopf symmetry of the model on both closed and open manifolds, demonstrating that there is a  $H \times \hat{H}$  symmetry for open manifolds, while for closed manifolds, the symmetry is given by  $Cocom(H) \times Cocom(H)$ . Section 7 extends the cluster state model to incorporate general boundary conditions. The lattice model is constructed using comodule algebras and can also be solved via weak Hopf tensor network states. Finally, Section 8 provides a discussion of the results and outlines several open problems for future research.

## 2 Warm-up example: finite-group quantum cluster ladder

In Ref. [54], it is demonstrated that the cluster state model is indeed a quantum double ladder model with rough and smooth boundaries. The aim of this paper is to generalize the construction presented therein to arbitrary weak Hopf algebras, encompassing all possible non-invertible symmetries in (1+1)D. Before delving into the cluster ladder model realization of non-invertible symmetry for the most general weak Hopf case, let us first consider a warm-up example of the finite-group cluster ladder model, focusing specifically on the  $\mathbb{Z}_p$  and  $S_3$  quantum cluster ladder models. The finite-group cluster state model is also discussed in Refs. [78, 92]. The macroscopic theory of the corresponding SymTFT is given in Ref. [56].

<sup>&</sup>lt;sup>2</sup>While a tensor product structure can be assigned to each edge, followed by a projection to enforce the two labels on the edge to be identical, this approach for introducing tensor product structure is not so natural.

Here, we emphasize the comodule algebra approach and discuss it in a more general setting via the socalled quantum cluster ladder model.

The group-based Pauli X operators are defined as

$$\vec{X}_g|h\rangle = |gh\rangle, \quad \overleftarrow{X}_g|h\rangle = |hg^{-1}\rangle.$$
 (2.1)

For some irreducible representation  $\Gamma \in \operatorname{Rep}(G)$ , the group-based Pauli Z operators are defined by

$$Z_{\Gamma} = \sum_{g \in G} \Gamma(g) \otimes |g\rangle \langle g|, \quad Z_{\Gamma}^{\ddagger} = \sum_{g \in G} \Gamma(g^{-1})|g\rangle \langle g|.$$
(2.2)

If we take trace of the representation matrix, we obtain

$$\vec{Z}_{\chi\Gamma} = \sum_{g \in G} \chi_{\Gamma}(g) |g\rangle \langle g|, \quad \overleftarrow{Z}_{\chi\Gamma} = \sum_{g \in G} \chi_{\Gamma}(g^{-1}) |g\rangle \langle g|, \quad (2.3)$$

where  $\chi_{\Gamma}$  is character of  $\Gamma \in \text{Rep}(G)$ .

For a quantum double model D(G) of finite group G, the topological boundary conditions are determined by a pair  $(K, \omega)$  with K a subgroup of G and  $\omega$  a two-cocycle [93, 95–97]. The topological boundary conditions are equivalently classified by the comodule algebras over Hopf algebra  $\mathbb{C}[G]$  [51, 52]. Here for simplicity, we will only consider the case with trivial 2-cocycle. The 1d cluster ladder is defined as the following lattice (with periodic boundary condition at two ends):



For the symmetry boundary (blue edges), we place  $K = \mathbb{C}[M]$  on each edge. For the physical boundary (red edges), we place  $J = \mathbb{C}[N]$  on each edge. In the bulk (black edges), we assign  $W = \mathbb{C}[G]$  to each edge. This means that the total Hilbert space is  $\mathcal{H}_{tot} = \mathcal{H}_{bk} \otimes \mathcal{H}_{sym} \otimes \mathcal{H}_{phys}$  with  $\mathcal{H} = \bigotimes_j \mathbb{C}[G]$ ,  $\mathcal{H}_{sym} = \bigotimes_k \mathbb{C}[M]$ , and  $\mathcal{H}_{phys} = \bigotimes_l \mathbb{C}[N]$ .

The SymTFT bulk operator is the face operator which is defined as

$$\mathbb{B}_f = \sum_{\Gamma \in \operatorname{Irr}(G)} \frac{d_{\Gamma}}{|G|} \mathbb{B}_f^{\chi_{\Gamma}}$$
(2.5)

where

$$\mathbb{B}_{f}^{\chi_{\Gamma}} | x_{3} \underbrace{\begin{array}{c} x_{2} \\ \bullet \\ x_{4} \end{array}}_{x_{4}} x_{1} \rangle = \chi_{\Gamma}(x_{1}^{-1}x_{2}x_{3}x_{4}^{-1}) | x_{3} \underbrace{\begin{array}{c} x_{2} \\ \bullet \\ x_{4} \end{array}}_{x_{4}} x_{1} \rangle.$$
(2.6)

The symmetry boundary vertex operator is defined as

$$\mathbb{A}_{\nu_s}\Big|\underset{x_3}{\overset{x_2}{\longrightarrow}}x_1\Big\rangle = \frac{1}{|M|} \sum_{g \in M} \Big|\underset{gx_3}{\overset{x_2g^{-1}}{\longrightarrow}}x_1g^{-1}\Big\rangle.$$
(2.7)

The physical boundary vertex operator is defined similarly. Both the face and vertex operators are equipped with a counterclockwise ordering (around the face and around the vertex). Alternatively, we can also define the model with a clockwise ordering. The Hamiltonian is of the form

$$\mathbb{H}_{G}^{M,N} = \mathbb{H}_{bk} + \mathbb{H}_{sym}^{M} + \mathbb{H}_{phys}^{N}$$
(2.8)

where  $\mathbb{H}_{bk} = \sum_{f} \mathbb{B}_{f}$ ,  $\mathbb{H}_{sym} = \sum_{v_s} \mathbb{A}_{v_s}$  and  $\mathbb{H}_{phys} = \sum_{v_p} \mathbb{A}_{v_p}$ . This model realize the SymTFT with fusion category symmetry give by the boundary excitations over the symmetry boundary.

The topological charges of the 2d bulk are labeled by the irreducible representations of the quantum double D(G) [51, 52, 93, 95–97]. These representations are classified by two components: the conjugacy class [g] of G, determining the magnetic charge, and the irreducible representations  $\pi$  of the centralizer  $C_G([g])$ , labeling the electric charge. For symmetry boundaries  $K = \mathbb{C}[M]$ , the boundary excitation is described by the fusion category

$$\mathcal{B}_{\text{sym}} = {}_{K}^{\mathbb{C}[G]} \mathsf{Mod}_{K}, \tag{2.9}$$

which is the category of  $\mathbb{C}[G]$ -covariant K|K-bimodules [51, 52, 98]. The boundary excitations are equivalently classified by pairs  $(S, \pi)$ , where  $S \in M \setminus G/M$  is a double coset and  $\pi$  is an irreducible representation of  $M \cap rMr^{-1}$ , with  $r \in S$  [96]. For physical boundaries  $J = \mathbb{C}[N]$ , the excitations can be characterized in a similar manner. Since we have chosen both boundaries to be topological boundaries, the cluster ladder model exhibits the following symmetry:

$$\mathcal{B}_{\text{sym}} \times \mathcal{B}_{\text{phys}} = {}_{K}^{\mathbb{C}[G]} \operatorname{\mathsf{Mod}}_{K} \times {}_{J} \operatorname{\mathsf{Mod}}_{J}^{\mathbb{C}[G]}, \qquad (2.10)$$

where  $\mathbb{C}[G]$  is placed on the left and right, respectively, to emphasize that K is a left  $\mathbb{C}[G]$ -comodule algebra and J is a right  $\mathbb{C}[G]$ -comodule algebra.

The CSS-type cluster state model is a special case of the quantum cluster ladder model, featuring one rough boundary and one smooth boundary [54]. For this case, M = G and  $N = \{1\}$ . The symmetry boundary excitation becomes  $\mathcal{B}_{sym} = \text{Rep}(G)$ , while the physical boundary excitation becomes  $\text{Vect}_G$ , the category of *G*-graded vector spaces. Notice that for the rough boundary edge, the boundary space reduces to  $\mathbb{C}$ , allowing us to effectively remove it. Additionally,  $\mathbb{A}_{v_p}$  becomes the identity operator, so we can also remove the boundary vertex operator. The lattice then simplifies to:



and the corresponding Hamiltonian becomes

$$\mathbb{H}_{G}^{G,\{1\}} = -\sum_{f} \mathbb{B}_{f}^{G} - \sum_{\nu_{s}:\text{sym}} \mathbb{A}_{\nu_{s}}^{G}.$$
(2.12)

The symmetry of the model is

$$\mathcal{B}_{\text{sym}} \times \mathcal{B}_{\text{phys}} = \text{Rep}(G) \times \text{Vect}_G.$$
(2.13)

Notice that choosing different configurations of orientations for each edge results in distinct cluster state models, and choosing a clockwise local ordering also yields a different model. However, all these models realize the same phase. The models discussed in Refs. [54, 78] are examples of the clockwise model. In this paper, we follow the conventions in Refs. [51, 52, 93] by using the counterclockwise local ordering. This will be elaborated in more detail later when discussing weak Hopf cluster ladder models.

## 2.1 Abelian cluster ladder model

Consider the  $\mathbb{Z}_p$  quantum double model, where p is a prime number.<sup>3</sup> The topological charges are given by irreducible representations of  $D(\mathbb{Z}_p)$ :

$$a_{[g],\pi_h} = e^g \otimes m^h =: \varepsilon^{g,h}, g, h \in \mathbb{Z}_p.$$
(2.14)

Since  $\mathbb{Z}_p$  is Abelian, [g] consists of a single element  $\{g\}$  and  $C_{\mathbb{Z}_p}([g]) = \mathbb{Z}_p$ . The irreducible representations of the Abelian group are also labeled by elements  $h \in \mathbb{Z}_p$ . The fusion rule reads

$$\varepsilon^{g,h} \otimes \varepsilon^{k,l} = (e^g \otimes m^h) \otimes (e^k \otimes m^l) = e^{g+k} \otimes m^{h+l} = \varepsilon^{g+k,h+l}, \tag{2.15}$$

namely, they form  $\mathbb{Z}_p \times \mathbb{Z}_p$  fusion algebra. In a good choice of gauge, the fusion F-symbols are trivial (i.e., they equal to 1 when allowed by fusion), and the braiding R-symbols read

$$R_{\varepsilon^{g,h},\varepsilon^{k,l}} = \exp(\frac{2\pi i}{p}gl).$$
(2.16)

There are two possible topological boundary conditions of the  $D(\mathbb{Z}_p)$  quantum double model:

- *Rough boundary*: This corresponds to the trivial subgroup  $M_r = \{0\}^4$ . The Lagrangian algebra is  $\mathcal{A}_r = \mathbf{1} \oplus e \oplus \cdots \oplus e^{p-1}$ . The boundary excitation is given by the unitary fusion category  $\mathsf{Vect}_{\mathbb{Z}_p}$ .
- Smooth boundary: This corresponds to the subgroup  $N_s = \mathbb{Z}_p$ . The Lagrangian algebra is  $\mathcal{A}_s = \mathbf{1} \oplus e \oplus \cdots \oplus e^{p-1}$ . The boundary excitation is given by  $\operatorname{Rep}(\mathbb{Z}_p)$ .

In both cases, the associated 2-cocycles are trivial. The triviality of the 2-cocycle can be understood as follows: the case of  $H^2(\mathbb{Z}_1, \mathbb{C}^{\times}) = 0$  is straightforwardly trivial. For  $H^2(\mathbb{Z}_p, \mathbb{C}^{\times})$ , we use the relation  $H^n(G, \mathbb{Z}) \simeq H^{n-1}(G, \mathbb{Q}/\mathbb{Z}) \simeq H^{n-1}(G, \mathbb{C}^{\times})$ , for all  $n \ge 2$ , along with the fact that  $H^3(\mathbb{Z}_p, \mathbb{Z}) = 0$ .

For the Abelian group  $\mathbb{Z}_p$ , the Weyl-Heisenberg X operator is given by

$$X_g = \sum_{h \in G} |h + g\rangle \langle h|,$$

where  $X_g = \overrightarrow{X}_g$  and  $X_g^{\dagger} = \overleftarrow{X}_g$ . Recall that all irreducible representations (irreps) of  $\mathbb{Z}_p$  are onedimensional, and  $\chi_{\Gamma}$  is a complex character function, with  $\Gamma_k$  labeled by a group element  $k \in \mathbb{Z}_p$ . In this way,  $\chi_k(h) = \exp(2\pi i k h/p)$ . The Weyl-Heisenberg Z operator is defined as

$$Z_k = \sum_{h \in G} \chi_k(h) |h\rangle \langle h|,$$

where  $Z_k = \overrightarrow{Z}_{\Gamma_k}$  and  $Z_k^{\dagger} = \overleftarrow{Z}_{\Gamma_k}$ .

The face operator in this case can be simplified as  $\mathbb{B}_{f}^{\mathcal{X}\Gamma} = \overrightarrow{Z}_{k} \otimes \overleftarrow{Z}_{k} \otimes \overleftarrow{Z}_{k}$ :

$$\mathbb{B}_{f}^{\chi_{\Gamma}} \Big| \begin{array}{c} x_{1} \\ \vdots \\ x_{2} \end{array} \Big\rangle = \chi_{\Gamma}(x_{1}x_{2}^{-1}x_{3}^{-1}) \Big| \begin{array}{c} x_{1} \\ \vdots \\ x_{2} \end{array} \Big\rangle.$$
(2.17)

<sup>&</sup>lt;sup>3</sup>We consider this case since there are only two possible topological boundary conditions; the discussion can be generalized to a general Abelian group straightforwardly.

<sup>&</sup>lt;sup>4</sup>Here we use 0 to denote the identity of the group.

$\otimes$	A	B	С	D	Ε	F	G	Н							
A	A	B	С	D	Е	F	G	Н							
B	B	A	С	Е	D	F	G	Н							
С	С	C	$A \oplus B \oplus C$	$D \oplus E$	$D \oplus E$	$G \oplus H$	$F\oplus H$	$F \oplus G$							
ת	D	F	$D \oplus E$	$A \oplus C \oplus F$	$B \oplus C \oplus F$	$D \oplus E$	$D \oplus E$	$D \oplus E$							
		L		$\oplus G \oplus H$	$\oplus G \oplus H$										
F	E	מ	$D \oplus F$	$B \oplus C \oplus F$	$A \oplus C \oplus F$	$D \oplus F$	$D \oplus F$ $D \oplus F$	$D \oplus F$							
											$D \oplus L$	$\oplus G \oplus H$	$\oplus G \oplus H$	$D \oplus L$	$D \oplus L$
F	F	F	$G \oplus H$	$D \oplus E$	$D \oplus E$	$A \oplus B \oplus F$	$H \oplus C$	$G \oplus C$							
G	G	G	$F \oplus H$	$D \oplus E$	$D \oplus E$	$H \oplus C$	$A\oplus B\oplus G$	$F \oplus C$							
Η	Η	H	$F \oplus G$	$D \oplus E$	$D \oplus E$	$G\oplus C$	$F \oplus C$	$A \oplus B \oplus H$							

**Table 1**. Fusion rules of  $D(S_3)$  quantum double model.

Thus, the face operator is given by

$$\mathbb{B}_f = \frac{1}{p} \sum_{k \in \mathbb{Z}_p} \overrightarrow{Z}_k \otimes \overleftarrow{Z}_k \otimes \overleftarrow{Z}_k.$$

The symmetry boundary operator becomes

$$\mathbb{A}_{v_s} = \frac{1}{p} \sum_{g \in \mathbb{Z}_p} \overleftarrow{X}_g \otimes \overleftarrow{X}_g \otimes \overrightarrow{X}_g$$

This model is nothing but the cluster state model [54, 78, 92]. The symmetry of the model is  $\operatorname{Rep}(\mathbb{Z}_p) \times \operatorname{Vect}_{\mathbb{Z}_p}$ , with the fusion algebra  $\operatorname{Gr}(\operatorname{Rep}(\mathbb{Z}_p)) = \operatorname{Gr}(\operatorname{Vect}_{\mathbb{Z}_p}) = \mathbb{Z}_p$  (we use Gr to denote Grothendieck group), which gives the usual  $\mathbb{Z}_p \times \mathbb{Z}_p$  cluster state model.

## 2.2 S<sub>3</sub> cluster ladder model

For non-Abelian group  $S_3 = \langle \sigma, \tau : \sigma^2 = \tau^3 = 1, \sigma\tau = \tau^{-1}\sigma \rangle$  (we take  $\sigma = (12)$  and  $\tau = (123)$ ), the corresponding quantum double model  $D(S_3)$  has eight topological charges described in the following table:

	A	В	С	D	Ε	F	G	H
conjugacy class [g]	[1]	[1]	[1]	$[\sigma]$	$[\sigma]$	$[\tau]$	$[\tau]$	[ au]
irrep of the centralizer	1	sign	π	$\omega_2^0$	$\omega_2^1$	$\omega_3^0$	$\omega_3^1$	$\omega_3^2$

Notice  $C_{S_3}([1]) = S_3$ , we use **1** to denote the trivial irrep, sign to denote the sign representation, and  $\pi$  to denote the two-dimensional representation of  $S_3$ . For  $C_{S_3}([\sigma]) = \mathbb{Z}_2$ , we use  $\omega_2^0$  and  $\omega_2^1$  to denote the two irreducible representations (irreps) of  $\mathbb{Z}_2$  ( $\omega_2$  is the second root of unity). For  $C_{S_3}([\tau]) = \mathbb{Z}_3$ , we use  $\omega_3^k$ , where k = 0, 1, 2, to denote its three irreps ( $\omega_3$  is the third root of unity). Their fusion rules are given in Table 1 [95, 99].

The *S* and *T* matrices of  $D(S_3)$  topological phase are of the form:

$$S = \frac{1}{6} \begin{bmatrix} 1 & 1 & 2 & 3 & 3 & 2 & 2 & 2 \\ 1 & 1 & 2 & -3 & -3 & 2 & 2 & 2 \\ 2 & 2 & 4 & 0 & 0 & -2 & -2 & -2 \\ 3 & -3 & 0 & 3 & -3 & 0 & 0 & 0 \\ 3 & -3 & 0 & -3 & 3 & 0 & 0 & 0 \\ 2 & 2 & -2 & 0 & 0 & 4 & -2 & -2 \\ 2 & 2 & -2 & 0 & 0 & -2 & -2 & 4 \\ 2 & 2 & -2 & 0 & 0 & -2 & 4 & -2 \end{bmatrix},$$

$$T = \operatorname{diag}(1, 1, 1, \omega_2^0, \omega_1^1, \omega_3^1, \omega_3^1, \omega_3^1, \omega_3^2).$$
(2.19)

Notice that the first row of the S-matrix represents the quantum dimensions of all topological charges, while the T-matrix encodes the topological spins of these charges. The fusion rule  $N_{XY}^Z$  can be determined from the S-matrix using the Verlinde formula:

$$N_{XY}^{Z} = \sum_{U \in Irr(D(G))} \frac{S_{XU} S_{YU} S_{ZU}^{*}}{S_{1U}},$$
(2.20)

where the summation runs over all irreducible representations U, and **1** denotes the trivial charge. For more details of modular data, see Refs. [100–102].

There are four types of topological boundary conditions [96]:

- Rough boundary: This corresponds to the trivial subgroup  $M_r = \{1\}$ . The Lagrangian algebra is  $\mathcal{A}_r = A \oplus B \oplus 2C$ . The boundary excitation is given by  $\operatorname{Vect}_{S_3}$ .
- Smooth boundary: This corresponds to the trivial subgroup  $M_s = S_3$ . The Lagrangian algebra is  $\mathcal{A}_s = A \oplus D \oplus F$ . The boundary excitation is given by  $\mathsf{Rep}(S_3)$ .
- $\mathbb{Z}_2$  boundary: This corresponds to the subgroup  $\mathbb{Z}_2$  of  $S_3$ . The Lagrangian algebra is  $\mathcal{A}_2 = A \oplus C \oplus D$ .
- $\mathbb{Z}_3$  boundary: This corresponds to the subgroup  $\mathbb{Z}_3$  of  $S_3$ . The Lagrangian algebra is  $\mathcal{A}_3 = A \oplus B \oplus 2F$ .

For the cluster ladder lattice, if there is no anyonic tunneling channel between the two boundaries, the ground state degeneracy of the model is 1. Such a model is a candidate for a non-invertible SPT phase. This includes the following models:

- 1. *The smooth and rough boundary pair:*  $\mathbb{H}[\mathcal{A}_s, \mathcal{A}_r]$ . This corresponds to the  $S_3$  cluster state model. The symmetry is given by  $\operatorname{Rep}(S_3) \times \operatorname{Vect}_{S_3}$ .
- 2. The  $\mathbb{Z}_2$  and  $\mathbb{Z}_3$  boundary pair:  $\mathbb{H}[\mathcal{A}_2, \mathcal{A}_3]$ . This model is entirely different from the cluster state model. Its symmetry is given by  $\operatorname{Rep}(D(S_3))_{\mathcal{A}_2} \times \operatorname{Rep}(D(S_3))_{\mathcal{A}_3}$ , where we use the notation  $\operatorname{Rep}(D(S_3))_{\mathcal{A}}$  to denote the category of  $\mathcal{A}$ -modules in  $\operatorname{Rep}(D(S_3))$ . Equivalently, the symmetry can be described as  ${\mathbb{C}}^{[S_3]}_K \operatorname{Mod}_K \times {}_J \operatorname{Mod}_J^{\mathbb{C}}^{[S_3]}$ , where  $K = \mathbb{C}[\mathbb{Z}_2]$  and  $J = \mathbb{C}[\mathbb{Z}_3]$ .

The lattice model with  $\text{Rep}(S_3)$  symmetry has been discussed carefully based on the anyonic chain in Ref. [57]. Our model here can be regarded as an alternative lattice realization.

For other pairings, we also provide the corresponding lattice Hamiltonian, e.g.,  $\mathbb{H}[\mathcal{A}_r, \mathcal{A}_2], \mathbb{H}[\mathcal{A}_s, \mathcal{A}_3]$ , etc. Due to the existence of anyonic tunneling channels between the two boundaries, the symmetry is either fully or partially broken.

## 3 Weak Hopf non-invertible symmetry

From a modern perspective, symmetries can be characterized by the algebraic structure of topological defects [26, 35, 40, 41, 47, 55–65]. Symmetries associated with non-invertible topological defects are known as non-invertible symmetries. For (1+1)D system, the symmetry is described by a fusion 1-category, viz., the usual fusion category, thus it's also called fusion category symmetry [40–42]. Since any unitary (multi)fusion category is equivalent to the representation category Rep(H) for some weak Hopf algebra H, it is natural for us to investigate the symmetry in the framework of Hopf and weak Hopf algebras [48, 49, 51–54, 93, 103, 104]. These symmetries, characterized by quantum algebras such as Hopf, quasi-Hopf and weak Hopf algebras, as well as module and comodule algebras over (weak, quasi-) Hopf algebras, constitute a significant class of non-invertible symmetries [29–35].

As we will see, in many cases, by fusion category symmetry  $\mathcal{C} \simeq \operatorname{Rep}(H)$ , people usually refer to the fusion algebra symmetry, whose algebraic structure is the fusion ring (Grothendieck ring)  $\operatorname{Gr}(\mathcal{C})$  of the category  $\mathcal{C}$  (or more generally,  $R(\mathcal{C}) = \operatorname{Gr}(\mathcal{C}) \otimes_{\mathbb{Z}} \mathbb{C}$ ). This fusion ring can be naturally embedded into the dual weak Hopf algebra  $\hat{H}$ . This suggests that weak Hopf symmetry provides a comprehensive framework for understanding non-invertible symmetries in (1+1)D systems.

#### 3.1 Weak Hopf algebra symmetry and weak Hopf comodule algebra symmetry

Here, we will review the definitions of various quantum algebras used in this work and discuss the symmetries characterized by these algebras. For further details on quantum algebras, see, e.g., Refs. [105–108]. Notice that, unless otherwise specified, all algebras and categories in this work are over the complex number field  $\mathbb{C}$ .

#### 3.1.1 Hopf symmetry

A complex Hopf algebra is a complex vector space H equipped with the five structure linear morphisms: multiplication  $\mu : H \otimes H \to H$ , unit  $\eta : \mathbb{C} \to H$ , comultiplication  $\Delta : H \to H \otimes H$ , counit  $\varepsilon : H \to \mathbb{C}$ , and antipode  $S : H \to H$ . These maps satisfy the following conditions:

- 1.  $(H, \mu, \eta)$  is an algebra, meaning  $\mu \circ (\mu \otimes id) = \mu \circ (id \otimes \mu)$  and  $\mu \circ (\eta \otimes id) = id = \mu \circ (id \otimes \eta)$ .
- 2.  $(H, \Delta, \varepsilon)$  is a coalgebra, meaning  $(\Delta \otimes id) \circ \Delta = (id \otimes \Delta) \circ \Delta$  and  $(\varepsilon \otimes id) \circ \Delta = id = (id \otimes \varepsilon) \circ \Delta$ .
- 3.  $(H, \mu, \eta, \Delta, \varepsilon)$  forms a bialgebra, where  $\Delta$  and  $\varepsilon$  are algebra homomorphisms, and equivalently,  $\mu$  and  $\eta$  are coalgebra homomorphisms.
- 4. The antipode *S* satisfies  $\mu \circ (S \otimes id) \circ \Delta = \eta \circ \varepsilon = \mu \circ (id \otimes S) \circ \Delta$ .

We will use the notation  $\mu(x \otimes y) = xy$  and  $\eta(1) = 1_H$  for multiplication and unit. We will also adopt the Sweedler's notation  $\Delta(u) = \sum_{(u)} u^{(1)} \otimes u^{(2)} = \sum_i u_i^{(1)} \otimes u_i^{(2)}$ . The comultiplication law ensures that  $(\Delta \otimes id) \circ \Delta(u) = (id \otimes \Delta) \circ \Delta(u) = \sum_{(u)} u^{(1)} \otimes u^{(2)} \otimes u^{(3)}$ . In general, we define  $\Delta_1 = \Delta$  and  $\Delta_n = (id \otimes \cdots \otimes id \otimes \Delta) \circ \Delta_{n-1}$ , then  $\Delta_n(u) = \sum_{(u)} u^{(1)} \otimes \cdots \otimes u^{(n+1)}$ . From the definition above, we have the following useful identity:

$$\sum_{(x)} \varepsilon(x^{(1)}) x^{(2)} = x = \sum_{(x)} x^{(1)} \varepsilon(x^{(2)}).$$
(3.1)

The swap operation is defined as  $\tau(x \otimes y) = y \otimes x$ , by which one denotes  $\mu^{op} = \mu \circ \tau$  and  $\Delta^{op} = \tau \circ \Delta$ . The opposite Hopf algebra  $H^{op}$  is defined as  $(H, \mu^{op}, \eta, \Delta, \varepsilon, S^{-1})$ , and the coopposite weak Hopf algebra  $H^{cop}$  is defined as  $(H, \mu, \eta, \Delta^{op}, \varepsilon, S^{-1})$ . For more details about our notation and convention, see [51, 52, 54].

**Definition 1** (Hopf symmetry). A quantum system  $(\mathbb{H}, \mathcal{H})$  is said to exhibit Hopf symmetry if and only if the Hilbert space  $\mathcal{H}$  supports a representation of a Hopf algebra, given by  $W_{\bullet} : H \to \text{End}(\mathcal{H})$ , and satisfies the following conditions: (i)  $[W_h, \mathbb{H}] = 0$  for all  $h \in H$ ; (ii) all eigenspaces  $\mathcal{V}$  of  $\mathbb{H}$  are invariant under the H-action. This symmetry will also be referred to as Hamiltonian symmetry to distinguish it from state symmetry.

In certain cases (especially when discussing weak Hopf symmetry), it is useful to discuss symmetry within the vacuum space. Specifically, this refers to an operator  $W_h$  that commutes with the Hamiltonian  $\mathbb{H}$  only within the ground state space, i.e.,

$$[W_h, \mathbb{H}]_{\mathcal{V}_{GS}} = 0, \quad \forall h \in H.$$

The ground state symmetry is generally larger than the Hamiltonian symmetry defined above.

All group symmetries are special cases of Hopf symmetries. A key example of a non-group Hopf algebra symmetry is quantum double symmetry D(G). Consider a (2 + 1)D discrete gauge theory, such as Gauge-Higgs theory, where a Higgs field breaks the continuous gauge group U to a finite group G [109, 110]. In the low-energy or long-distance limit, the theory becomes topological, with excitations characterized by fusion and braiding data. These data are captured by the representation category Rep(D(G)), corresponding to the quantum double of G. This formalism is generalized in [48], where the unbroken symmetry is described by a Hopf algebra H, of which the group algebra is a special case. Hopf algebraic symmetry has broad applications in quantum gravity [105, 111, 112], conformal field theory (CFT) [113], topological quantum field theory (TQFT) [103, 114], and the quantum Hall effect [115]. The Hopf quantum double model provides a general framework for realizing Hopf symmetry [49–51, 93, 103, 104, 114, 116–118], while certain string-net models exhibit Hopf symmetry [53, 119], among other examples.

**Invertible Hopf symmetry** A special type of Hopf symmetry is the conventional invertible group symmetry, where  $H = \mathbb{C}[G]$ . The group elements form a basis, with multiplication induced by group multiplication. The comultiplication and counit are defined as  $\Delta(g) = g \otimes g$  and  $\varepsilon(g) = 1$  for all basis elements  $g \in G$ . The antipode is given by  $S(g) = g^{-1}$  for each  $g \in G$ , and the \*-operation (to be defined later for more general weak Hopf algebras) is specified by  $g^* = g^{-1}$ .

Recall that for invertible symmetry G,  $U_g^{\dagger} = U_{g^*} = U_{g^{-1}}$ .  $U_g$ 's commute with Hamiltonian  $\mathbb{H}$  for all  $g \in G$ . The total Hilbert space supports a representation of G, and all eigenspace of  $\mathbb{H}$  is invariant under the G-action. When regarded as a group algebra symmetry  $\mathbb{C}[G]$ , we say  $(\mathbb{H}, \mathcal{H})$  possesses a  $\mathbb{C}[G]$ 

symmetry means that (i)  $[U_x, \mathbb{H}] = 0$  for all  $x \in \mathbb{C}[G]$ ; (ii) any eigenspace  $\mathcal{V}$  of  $\mathbb{H}$  is invariant under the *G*-action, note that this a result of condition (i).

A natural question now arises: under what circumstances is a Hopf symmetry invertible? We provide the following formal definition:

**Definition 2.** A (weak) Hopf symmetry H is called invertible if there exists a group G such that  $H \cong \mathbb{C}[G]$  as Hopf algebras.

Notice that the elements in *G* are all invertible, but the elements in  $\mathbb{C}[G]$  are generally not invertible. Determining whether a given Hopf symmetry is invertible is generally a challenging task. It is clear that group algebra symmetries must be cocommutative. For pointed cocommutative Hopf symmetries<sup>5</sup>, a useful result is the Cartier-Konstant-Milnor-Moore theorem [102, 121–123].

For a (weak) Hopf algebra H, an element  $g \in H$  is called a group-like element if and only if  $\Delta(g) = g \otimes g$  and  $\varepsilon(g) = 1$ . The set of all group-like elements of H is denoted by  $\mathcal{G}(H)$ ; it forms a group when the antipode S is invertible. If  $x \in H$  satisfies  $\Delta(x) = x \otimes 1 + 1 \otimes x$ , we call x a primitive element. If x is primitive, then we have  $\varepsilon(x) = 0$ . The set of all primitive element of H is denoted as  $\mathcal{P}(H)$ , and  $\mathcal{P}(H)$  is a Lie algebra with [x, y] := xy - yx. For group algebra  $\mathbb{C}[G]$ , it is clear that  $\mathcal{P}(\mathbb{C}[G]) = \{0\}$  and  $\mathcal{G}(\mathbb{C}[G]) = G$ .

The Cartier-Konstant-Milnor-Moore theorem [102, 121-123] claims that, for pointed cocommutative Hopf algebras H, we have

$$H \cong U(\mathcal{P}(H)) \# \mathbb{C}[\mathcal{G}(H)], \tag{3.2}$$

where  $\mathcal{P}(H)$  is the set of primitive elements of H,  $U(\mathcal{P}(H))$  is the universal enveloping algebra of  $\mathcal{P}(H)$ , and  $\mathcal{G}(H)$  is the set of group elements in H; '#' represents smash product (see, e.g., [107]).

Since  $\mathcal{P}(H) = 0$  for a finite-dimensional complex Hopf algebra H [124, Exercise 4.2.16], a direct consequence of the Cartier–Kostant–Milnor–Moore theorem is that for finite-dimensional pointed cocommutative Hopf algebras, we have  $H \cong \mathbb{C}[\mathcal{G}(H)]$ . Therefore, all finite-dimensional pointed cocommutative Hopf symmetries are invertible symmetries.

**Comodule algebra symmetry** As shown in Refs. [51–53], (weak) Hopf symmetry characterizes 2d topologically ordered phases, while 1d gapped boundaries exhibit comodule algebra symmetry (or equivalently, module algebra symmetry). These symmetries are fundamental to our construction of non-invertible SPT lattice models. We now review their definitions.

Let *H* be a Hopf algebra, and let  $(K, \mu_K, \eta_K)$  be an algebra. If *K* is a left *H*-comodule with coaction  $\beta_K : K \to H \otimes K^6$ , satisfying the conditions:

$$\beta_K(xy) = \beta_K(x)\beta_K(y), \quad \beta_K(1_K) = 1_H \otimes 1_K, \tag{3.3}$$

then *K* is called a *left H-comodule algebra*. A *right H-comodule algebra* can be defined analogously. For left and right comodules, we adopt Sweedler's notation for left and right coactions as follows:

$$\beta(x) = \sum_{(x)} x^{[-1]} \otimes x^{[0]}, \beta(x) = \sum_{(x)} x^{[0]} \otimes x^{[1]}.$$
(3.4)

<sup>&</sup>lt;sup>5</sup> A Hopf algebra H over a field  $\mathbb{F}$  is called pointed if all its simple left or right comodules are one-dimensional. See Ref. [120] for more details.

<sup>&</sup>lt;sup>6</sup>This means that  $(id \otimes \beta) \circ \beta = (\Delta \otimes id) \circ \beta$  and  $(\varepsilon \otimes id) \circ \beta = id$ .

Let H be a Hopf algebra and M an algebra. If M is a left H-module and satisfies the following conditions:

$$h \triangleright (xy) = \sum_{(h)} (h^{(1)} \triangleright x) (h^{(2)} \triangleright y), \quad h \triangleright 1_M = \varepsilon(h) 1_M, \tag{3.5}$$

for all  $h \in H$  and  $x, y \in M$ , then M is called a *left H-module algebra*. A *right H-module algebra* is defined similarly.

From a macroscopic perspective, the description of the 1d gapped boundary of the quantum double model in terms of comodule algebra and module algebra is equivalent. Henceforth, following the convention in Refs. [51–53], we adopt the comodule algebra description.

## 3.1.2 Weak Hopf symmetries

It is well-known that not all fusion categories can be regarded as the representation category of a Hopf algebra. Typical examples include the Haagerup category and  $\text{Vect}_G^{\omega}$  (the category of *G*-graded vector spaces with a non-trivial 3-cocycle  $\omega$ ). To represent an arbitrary fusion category as a representation category, it is necessary to introduce the concept of weak Hopf algebras [88–90].

A complex weak Hopf algebra [125] is a complex vector space equipped with the same structural morphisms as a Hopf algebra, but with modified compatibility conditions:

- 1. The triple  $(H, \mu, \eta)$  is an algebra, and the triple  $(H, \Delta, \varepsilon)$  forms a coalgebra.
- 2. The quintuple  $(H, \mu, \eta, \Delta, \varepsilon)$  is a weak bialgebra, meaning that the algebra structure and coalgebra structure are compatible in the following way:

(i) *Multiplicativity of comultiplication:* The comultiplication map satisfies  $\Delta(xy) = \Delta(x) \cdot \Delta(y)$ , explicitly,

$$\sum_{(x \cdot y)} (x \cdot y)^{(1)} \otimes (x \cdot y)^{(2)} = \sum_{(x), (y)} x^{(1)} \cdot y^{(1)} \otimes x^{(2)} \cdot y^{(2)}.$$
 (3.6)

This property holds for Hopf algebras as well.

(ii) Weak comultiplicativity of the unit: The comultiplication of the identity element satisfies

$$(\Delta \otimes \mathrm{id}) \circ \Delta(1_H) = (\Delta(1_H) \otimes 1_H) \cdot (1_H \otimes \Delta(1_H)) = (1_H \otimes \Delta(1_H)) \cdot (\Delta(1_H) \otimes 1_H).$$
(3.7)

This indicates that the comultiplication of the identity is not factorizable but instead takes the form of a linear combination:

$$\Delta(1_H) = \sum_{(1_H)} 1_H^{(1)} \otimes 1_H^{(2)}.$$
(3.8)

In the case of Hopf algebras, we have the stronger condition

$$\Delta(1_H) = 1_H \otimes 1_H. \tag{3.9}$$

(iii) Weak multiplicativity of the counit: For all  $x, y, z \in H$ , the counit satisfies

$$\varepsilon(x \cdot y \cdot z) = \sum_{(y)} \varepsilon(x \cdot y^{(1)}) \varepsilon(y^{(2)} \cdot z) = \sum_{(y)} \varepsilon(x \cdot y^{(2)}) \varepsilon(y^{(1)} \cdot z).$$
(3.10)

For Hopf algebras, since  $\Delta(1_H) = 1_H \otimes 1_H$ , setting y = 1 in the above equation gives the stronger constraint  $\varepsilon(xz) = \varepsilon(x)\varepsilon(z)$ .

- 3. The antipode *S* plays the role of taking the inverse of elements, and its axioms differ significantly from those of Hopf algebras. It satisfies the following properties:
  - (i) *Left counit:* For all *x* in *H*, we have

$$\sum_{(x)} x^{(1)} S(x^{(2)}) = \varepsilon_L(x) := \sum_{(1_H)} \varepsilon(1_H^{(1)} x) 1_H^{(2)}.$$
(3.11)

For Hopf algebras, we have the stronger relation  $\sum_{(x)} x^{(1)} S(x^{(2)}) = \varepsilon_L(x) = \varepsilon(x) \mathbf{1}_H$ .

(ii) *Right counit:* For all *x* in *H*, we have

$$\sum_{(x)} S(x^{(1)}) x^{(2)} = \varepsilon_R(x) := \sum_{(1_H)} 1_H^{(1)} \varepsilon(x 1_H^{(2)}).$$
(3.12)

For Hopf algebra, we have the stronger relation  $\sum_{(x)} S(x^{(1)})x^{(2)} = \varepsilon_R(x) = \varepsilon(x)\mathbf{1}_H$ . (iii) *Antipode decomposition:* For all x in H, we have

$$\sum_{(x)} S(x^{(1)}) x^{(2)} S(x^{(3)}) = S(x).$$
(3.13)

This condition also holds for Hopf algebras.

Notice that the images of the left and right counit maps are denoted as  $H_L = \varepsilon_L(H)$  and  $H_R = \varepsilon_R(H)$ . They serve as the tensor unit (vacuum charge or vacuum sector) of the representation category Rep(H) of H. In general, the tensor unit is not simple, and the representation category Rep(H) is a multifusion category. When H is connected [90] (also called pure in Ref. [125]), Rep(H) is a fusion category. If the dual weak Hopf algebra  $\hat{H}$  is connected, we call H coconnected [126]. In this work, we primarily consider connected and coconnected weak Hopf algebras, although all discussions can be straightforwardly generalized.

We will also assume that the weak Hopf algebra is a  $C^*$ -weak Hopf algebra. A complex weak Hopf algebra H is called simple (or indecomposable) if its underlying algebra  $(H, \mu, \eta)$  has no nontrivial subalgebras, and it is called semisimple if its underlying algebra can be written as a direct sum of simple algebras. A \*-weak Hopf algebra (H, \*) is a weak Hopf algebra H equipped with a  $C^*$ -structure  $* : H \to H$  such that  $\Delta$  is a \*-homomorphism. That is, for all  $x, y \in H$  and  $c \in \mathbb{C}$ ,

$$(x^*)^* = x, \quad (x+y)^* = x^* + y^*, \quad (xy)^* = y^* x^*, \quad (cx)^* = \bar{c}x^*, \quad \Delta(x)^* = \Delta(x^*).$$
 (3.14)

We also have

$$S(x^*) = S^{-1}(x)^*.$$
(3.15)

A  $C^*$ -weak Hopf algebra (H, \*) is one for which there exists a fully faithful \*-representation  $\rho : H \to \mathbf{B}(\mathcal{H})$  for some operator space over a Hilbert space  $\mathcal{H}$ .

The weak Hopf morphism between two weak Hopf algebras  $H_1$  and  $H_2$  is a map  $f : H_1 \to H_2$ which is both an algebra and coalgebra homomorphism preserving unit and counit and which intertwines the antipodes,  $f \circ S_1 = S_2 \circ f$ . The category of all finite dimensional weak Hopf algebras will be denoted as WHopfAlg<sub>C</sub>.

A weak Kac algebra is a weak Hopf algebra such that  $S^2 = \text{id} [127, 128]$ . All finite-dimensional semisimple Hopf algebras are weak Kac algebras due to the Larson-Radford theorem (See, e.g., Ref. [108]).

**Definition 3** (Weak Hopf symmetry). A Hamiltonian  $\mathbb{H} : \mathcal{H} \to \mathcal{H}$  is said to possess weak Hopf symmetry H if  $\mathcal{H}$  supports a representation  $W_{\bullet} : H \to \text{End}(\mathcal{H})$  of the weak Hopf algebra H, such that  $[W_h, \mathbb{H}] = 0$ for all  $h \in H$ . In some contexts, we consider symmetries that act only on the ground states, where  $[\mathbb{H}, W_h] = 0$ holds (at least) within the ground state space  $\mathcal{V}_{GS}$ . In such cases, the symmetry of the ground state space must be larger than that of the Hamiltonian, i.e.,  $\text{Sym}_{GS} \supseteq \text{Sym}_{\mathbb{H}}$ . If the ground state space is non-degenerate, meaning dim  $\mathcal{V}_{GS} = 1$ , the ground state must be an eigenstate of the symmetry operator  $W_h$ . When the weak Hopf algebra is a weak Kac algebra, we refer to the corresponding symmetry as weak Kac symmetry.

To summarize, the hierarchy of symmetries is as follows:

Invertible symmetry  $\subset$  Hopf symmetry  $\subset$  Weak Kac symmetry  $\subset$  Weak Hopf symmetry. (3.16)

For weak Hopf symmetry, the notion of invariance under the symmetry action differs from that of groups and Hopf algebras [52]. A state  $|\psi\rangle \in \mathcal{H}$  is called left-invariant under the action if and only if

$$W_h|\psi\rangle = W_{\varepsilon_L(h)}|\psi\rangle. \tag{3.17}$$

The notion of right-invariance is defined analogously. For a Hopf algebra, Eq. (3.17) reduces to

$$W_h|\psi\rangle = \varepsilon(h)|\psi\rangle, \quad \forall h \in H,$$
(3.18)

which means that  $|\psi\rangle$  is an eigenstate of  $W_h$  with eigenvalue  $\varepsilon(h)$ . For a group algebra  $\mathbb{C}[G]$ , where  $\varepsilon(g) = 1$  for all  $g \in G$ , this further simplifies to the usual definition of invariance for group symmetries.

Weak Hopf comodule algebra symmetry For the topological boundary condition of 2d weak Hopf lattice gauge theory, we require the concept of a comodule algebra over weak Hopf algebras [52]. The axioms for the comodule algebra need to be adjusted compared to the standard Hopf algebra case.

**Definition 4.** Let H be a weak Hopf algebra, an algebra K is called a left H-comodule algebra [129] if there exists a comodule map  $\beta : K \to H \otimes K$  such that

$$\beta(xy) = \beta(x)\beta(y), \qquad (3.19)$$

and

$$(1_H \otimes x)\beta(1_K) = (\varepsilon_R \otimes \mathrm{id}_K) \circ \beta(x), \tag{3.20}$$

for all  $x, y \in K$ . We adopt Sweedler's notation  $\beta(x) = \sum_{[x]} x^{[-1]} \otimes x^{[0]}, \beta_2(x) = \sum_{[x]} x^{[-2]} \otimes x^{[-1]} \otimes x^{[0]}, \text{ etc.}^7$  The right H-comodule algebra can be defined similarly.

*Remark* 3.1. In literature, there are several equivalent expressions of Eq. (3.20) (written more explicitly and omitting the sum symbol  $1_K^{[-1]} \otimes x 1_K^{[0]} = \varepsilon_R(x^{[-1]}) \otimes x^{[0]}$ ):

- 1.  $\beta(1_K) \in H_R \otimes K;$
- 2.  $\beta(1_K) = (\varepsilon_R \otimes id) \circ \beta(1_K);$

<sup>&</sup>lt;sup>7</sup>We use square brackets to denote the comodule structure to distinguish it from the comultiplication of the weak Hopf algebra.

3.  $1_{H}^{(1)} \otimes 1_{H}^{(2)} 1_{K}^{[-1]} \otimes 1_{K}^{[0]} = 1_{H}^{(1)} \otimes 1_{K}^{[-1]} 1_{H}^{(2)} \otimes 1_{K}^{[0]} = 1_{K}^{[-2]} \otimes 1_{K}^{[-1]} \otimes 1_{K}^{[0]}$ , where we have omitted the sum symbol;

For the convenience of the reader, we provide the proof here.

The equivalence of 1 and 2 is obvious since  $\varepsilon_R^2 = \varepsilon_R$ . For "1  $\Rightarrow$  3", note that  $1^{(1)} \otimes 1^{(2)} \in H_R \otimes H_L$ and hg = gh for all  $h \in H_L$  and  $g \in H_R$ . Using the fact that for all  $h \in H_R$  we have

$$\sum_{(1)} 1^{(1)} \otimes h 1^{(2)} = \Delta(h), \tag{3.21}$$

we arrive at the required conclusion.

For "3  $\Rightarrow$  Eq. (3.20)", note that for  $x \in K$ , we have

$$\beta(x1) = x^{[-1]} 1^{[-1]} \otimes x^{[0]} 1^{[0]} = x^{[-1]} \otimes x^{[0]}.$$
(3.22)

Then

$$\varepsilon_{R}(x^{[-1]}) \otimes x^{[0]} = 1_{H}^{(1)} \varepsilon(x^{[-1]} 1_{H}^{(2)}) \otimes x^{[0]}$$
  

$$= 1_{H}^{(1)} \varepsilon(x^{[-1]} 1_{K}^{[-1]} 1_{H}^{(2)}) \otimes x^{[0]} 1_{K}^{[0]}$$
  

$$= 1_{K}^{[-2]} \varepsilon(x^{[-1]} 1_{K}^{[-1]}) \otimes x^{[0]} 1_{K}^{[0]}$$
  

$$= 1_{K}^{[-1]} \otimes x 1_{K}^{[0]}.$$
(3.23)

Finally, "Eq.  $(3.20) \Rightarrow 1$ " is obvious.

Note that H is a comodule algebra over itself with the comodule map given by the comultiplication. Since  $\Delta$  is an algebra map and

$$\Delta(1_H) \in H_R \otimes H_L, \tag{3.24}$$

the comodule algebra conditions are satisfied. The topological boundary condition can also be characterized via the module algebra. An algebra A is called a *left* H-module algebra if A is a left H-module such that  $h \triangleright (xy) = \sum_{(h)} (h^{(1)} \triangleright x) (h^{(2)} \triangleright y)$  and  $h \triangleright 1_A = \varepsilon_L(h) \triangleright 1_A$  for all  $h \in H$  and  $x \in A$ . The *right* H-module algebra can be defined similarly.

## 3.2 Fusion category symmetry as dual weak Hopf symmetry

Let us now consider how to understand (multi)fusion category symmetry in the framework of weak Hopf symmetries. By definition, a fusion category symmetry is a fusion category  $\mathcal{C}$  that acts on a quantum system  $\mathcal{T}$ , like lattice system or quantum field theory, as a symmetry. Treating the symmetry action as an algebra, the symmetry algebra for fusion category  $\mathcal{C}$  is the Grothendieck ring Gr( $\mathcal{C}$ ). By definition, Gr( $\mathcal{C}$ ) is generated by the equivalence classes of simple objects in  $\mathcal{C}$  as a free  $\mathbb{Z}$ -module. The multiplication is defined by the fusion rule:

$$[a] \cdot [b] := [a \otimes b] = \sum_{c \in \operatorname{Irr}(\mathbb{C})} N^c_{ab}[c], \quad N^c_{ab} \in \mathbb{Z}_{\ge 0}.$$
(3.25)

This is also called the fusion ring, and it is a  $\mathbb{Z}_+$ -ring [102]. When  $\mathcal{C}$  is a multifusion category, the unit of the ring is a nonnegative linear combination of the simple objects that constitute the tensor unit **1** of  $\mathcal{C}$ .

When C is a fusion category, [1] is one of the basis elements. Hereinafter, we will primarily focus on the fusion category case, as the generalization to multifusion categories is straightforward.

The simple topological symmetry defect lines or symmetry operators  $W_{[a]}$ 's are labeled by basis elements  $[a] \in Gr(\mathcal{C})$ , which is isomorphism class of simple objects in  $\mathcal{C}$ . These symmetry defect line obey the fusion algebra:

$$W_{[a]}W_{[b]} = W_{[a\otimes b]} = W_{\sum_{c} N_{ab}^{c}[c]}, N_{ab}^{c} \in \mathbb{Z}_{\geq 0}.$$
(3.26)

To make the fusion algebra a  $\mathbb{C}$ -algebra, we introduce the character algebra  $R(\mathbb{C}) := \operatorname{Gr}(\mathbb{C}) \otimes_{\mathbb{Z}} \mathbb{C}$  [126]. The fusion category symmetry is thus equivalently described by the character algebra  $R(\mathbb{C})$ , with the symmetry operator  $W_{\alpha}$  defined for  $\alpha \in R(\mathbb{C})$ .

Recall that a weak Hopf algebra H is called connected if and only if the trivial representation 1 of H is a simple object. In this case, the representation category  $\operatorname{Rep}(H)$  is a unitary fusion category. On the other hand, for any fusion category  $\mathbb{C}$ , there exists a (not unique) weak Hopf algebra H such that  $\operatorname{Rep}(H) \simeq \mathbb{C}$ as tensor categories [88–90, 125]. Two weak Hopf algebras H and H' are called Morita equivalent if and only if  $\operatorname{Rep}(H) \simeq \operatorname{Rep}(H')$  as tensor categories. Thus for a given fusion category symmetry, we can determine the weak Hopf symmetry H up to Morita equivalence.

The weak Hopf algebra associated with a given fusion category can be reconstructed in the following two ways:

- 1. Tannaka-Krein reconstruction from weak fiber functor. Given any fusion category  $\mathcal{C}$ , there exists an algebra A over  $\mathbb{C}$  such that we can endow  $\mathcal{D}$  with an exact faithful monoidal functor  $F : \mathcal{C} \to {}_A \operatorname{Mod}_A$  (known as the weak fiber functor over fusion category <sup>8</sup>). The algebra  $H = \operatorname{End}(F)$ , consisting of natural transformations from F to itself. It can be proved that H forms a weak Hopf symmetry, and the representation category  $\operatorname{Rep}(H)$  is equivalent to  $\mathcal{C}$  as unitary fusion categories [88, 89].
- 2. Boundary tube algebra. An intuitive approach involves the concept of boundary tube algebra [53, 119, 130] as applied to the string-net model. For a given bulk fusion category  $\mathcal{C}$ , we consider a gapped boundary described by a  $\mathcal{C}$ -module category  $\mathcal{C}\mathcal{M}$  and construct the corresponding boundary tube algebra **Tube**( $\mathcal{C}\mathcal{M}$ ), which is a  $C^*$  weak Hopf algebra. Setting  $\mathcal{M} = \mathcal{C}$ , namely, considering the smooth boundary, we have the boundary tube algebra as follows:

$$\left\{ \begin{array}{ccc}
g \\
f \\
e \\
v \\
c
\end{array} : a, c, e, f, g \in \operatorname{Irr}(\mathcal{C}), \mu, v \in \operatorname{Hom}_{\mathcal{C}} \\
e \\
v \\
c
\end{array} \right\}.$$
(3.27)

The fusion category  $\mathcal{C} = \operatorname{Fun}_{\mathcal{C}}(\mathcal{C}, \mathcal{C})$  (regarded as boundary phase) can be embedded into the representation category of this tube algebra  $\operatorname{Rep}(\operatorname{Tube}(_{\mathcal{C}}\mathcal{C}))$ . Thus, the fusion category  $\mathcal{C}^*_{\mathcal{M}}$  can be realized via the weak Hopf algebra  $\operatorname{Tube}(_{\mathcal{C}}\mathcal{C})$ .

<sup>&</sup>lt;sup>8</sup>Notice that in Ref. [53], we refer to this functor as a fiber functor. We emphasize that this is not the same as a fiber functor  $F : \mathcal{C} \to \text{Vect}$ , which does not always exist for general fusion categories.

Before we discuss how to treat fusion category symmetry as a dual weak Hopf symmetry, let's recall some basics about the representations of weak Hopf algebra. The  $H_L$  is the tensor unit with H-action given by

$$x \triangleright a = \varepsilon_L(xh) = x^{(1)}aS(x^{(2)}), \forall x \in H, \forall a \in H_L.$$
(3.28)

For two representations V, W, the action of H on  $V \otimes_{\mathbb{C}} W$  is defined as

$$h \triangleright (v \otimes w) := \sum_{(h)} h^{(1)} v \otimes h^{(2)} w, \qquad (3.29)$$

the tensor product of V and W is defined as

$$V \otimes W = \{ x \in V \otimes_{\mathbb{C}} W | \Delta(1) \triangleright x = x \}.$$
(3.30)

The dual representation of *V* is  $\hat{V} := \text{Hom}(V, \mathbb{C})$  equipped with the action  $(h \triangleright f)(x) := f(S(h) \triangleright x)$ .

Recall that, an element l (or r) is called a left (or right) integral of H if it satisfies  $xl = \varepsilon_L(x)l$  (or  $rx = r\varepsilon_R(x)$ ). A left (or right) integral l (or r) is considered left (or right) normalized if  $\varepsilon_L(l) = 1_H$  (or  $\varepsilon_R(r) = 1_H$ ). When an element h functions as both a left and right integral, it is referred to as a two-sided integral. A Haar integral in H is a two-sided integral that is normalized on both sides. For a  $C^*$  weak Hopf algebra H, there always exist a unique Haar integral.

**Lemma 1** (Schur's lemma for weak Hopf algebra). Let  $\Gamma : H \to \text{End}(V)$  and  $\Phi : H \to \text{End}(W)$  be two irreducible representations of a weak Hopf algebra H, and let  $f : V \to W$  be a linear map such that  $\Phi(x) \circ f = f \circ \Gamma(x)$  for all  $x \in H$ . If  $\Gamma$  and  $\Phi$  are not isomorphic, then f = 0. When  $\Gamma = \Phi$ , f is a scalar multiple of the identity map on V, i.e.,  $f \propto \text{id}_V$ .

*Proof.* The proof follows the same reasoning as in Ref. [54].

For Haar integral  $\lambda \in H$  and arbitrary  $x \in H$ , we have the following [125, 126]

$$\sum_{(\lambda)} x S(\lambda^{(1)}) \otimes \lambda^{(2)} = \sum_{(\lambda)} S(\lambda^{(1)}) \otimes \lambda^{(2)} x.$$
(3.31)

Then for a linear map  $f: V \to W$ , we define

$$F = \sum_{(\lambda)} \Phi(S(\lambda^{(1)})) \circ f \circ \Gamma(\lambda^{(2)}).$$
(3.32)

It's clear from Eq. (3.31) that

$$\Phi(x) \circ F = F \circ \Gamma(x), \quad \forall x \in H.$$
(3.33)

**Lemma 2.** (1) For any  $f : V \to W$ , and define F as in Eq. (3.32), we have: (i) If  $\Gamma$  and  $\Phi$  are not isomorphic, F = 0; (ii) when  $\Gamma = \Phi$ , we have

$$F = \frac{1}{d_{\Gamma}} \operatorname{Tr}(f) \operatorname{id}_{V}.$$
(3.34)

(2) From the above result, we obtain the orthogonality relation for irreps of H:

$$\sum_{(\lambda)} \Phi_{ij}(S(\lambda^{(1)})) \Gamma_{kl}(\lambda^{(2)}) = \frac{\delta_{\Phi,\Gamma} \delta_{il} \delta_{jk}}{d_{\Gamma}} = \sum_{(\lambda)} \Phi_{ij}(S(\lambda^{(2)})) \Gamma_{kl}(\lambda^{(1)}).$$
(3.35)

where we have used the cocommutativity of  $\lambda$ .

*Proof.* (1) Note that (i) is a direct result of Proposition 1. For (ii), we have

$$\operatorname{Tr} F = \sum_{(\lambda)} \operatorname{Tr}[\Gamma(S(\lambda^{(1)})) \circ f \circ \Gamma(\lambda^{(2)})] = \operatorname{Tr}[\Gamma(\varepsilon_L(\lambda))f] = \operatorname{Tr} f, \qquad (3.36)$$

which implies Eq. (3.34), notice that we have used  $\varepsilon_L(\lambda) = 1_H$ .

(2) From (1), the matrix elements  $F_{il}$  is of the form

$$\sum_{(\lambda)} \sum_{jk} \Phi_{ij}(\mathcal{S}(\lambda^{(1)})) f_{jk} \Gamma_{kl}(\lambda^{(2)}) = \frac{\delta_{\Phi,\Gamma}}{d_{\Gamma}} \delta_{il} \sum_{jk} f_{jk} \delta_{jk}.$$
(3.37)

This implies the required result by taking  $f_{jk} = \delta_{jj'} \delta_{kk'}$ .

Suppose we have identified a weak Hopf algebra H that realizes the fusion category symmetry  $\mathcal{C} \simeq \operatorname{Rep}(H)$ . In this case, the simple topological defect lines are labeled by the irreducible representations  $\Gamma \in \operatorname{Irr}(H)$  of the weak Hopf algebra H. The fusion ring of  $\operatorname{Rep}(H)$  can equivalently be described using the irreducible characters  $\chi_{\Gamma}$ , where  $\Gamma \in \operatorname{Irr}(H)$ :

$$\chi_{\Gamma} \cdot \chi_{\Phi} = \chi_{\Gamma \otimes \Phi} = \sum_{\Psi \in \operatorname{Irr}(H)} N_{\Gamma \Phi}^{\Psi} \chi_{\Psi}.$$
(3.38)

Here, the product on the left-hand side is defined by  $\chi_{\Gamma} \cdot \chi_{\Phi}(x) = \sum_{(x)} \chi_{\Gamma}(x^{(1)}) \chi_{\Phi}(x^{(2)})$ , which corresponds to the multiplication in the dual weak Hopf algebra  $\hat{H}$ . Recall that for a  $C^*$  weak Hopf algebra H, its dual space  $\hat{H} := \text{Hom}(H, \mathbb{C})$  acquires a canonical  $C^*$  weak Hopf algebra structure induced by the canonical pairing  $\langle \bullet, \bullet \rangle : \hat{H} \times H \to \mathbb{C}$ , defined by  $\langle \varphi, h \rangle := \varphi(h)^9$ . The structure morphisms are given by:

$$\langle \hat{\mu}(\varphi \otimes \psi), x \rangle = \langle \varphi \otimes \psi, \Delta(x) \rangle,$$
 (3.39)

$$\langle \hat{\eta}(1), x \rangle = \varepsilon(x), \text{ i.e., } \hat{1} = \varepsilon,$$
 (3.40)

$$\langle \hat{\Delta}(\varphi), x \otimes y \rangle = \langle \varphi, \mu(x \otimes y) \rangle,$$
 (3.41)

$$\hat{\varepsilon}(\varphi) = \langle \varphi, \eta(1) \rangle,$$
 (3.42)

$$\langle \hat{S}(\varphi), x \rangle = \langle \varphi, S(x) \rangle.$$
 (3.43)

The star operation on  $\hat{H}$  is defined as

$$\langle \varphi^*, x \rangle = \overline{\langle \varphi, S(x)^* \rangle}.$$
 (3.44)

Since irreducible characters form a basis of  $\operatorname{Gr}(\operatorname{Rep}(H))$ , and each  $\chi_{\Gamma} \in \hat{H}$ , we can embed  $\operatorname{Gr}(\operatorname{Rep}(H))$ into  $\hat{H}$ . We can consider  $R(\operatorname{Rep}(H)) = \operatorname{Gr}(\operatorname{Rep}(H)) \otimes_{\mathbb{Z}} \mathbb{C}$ , this is a subalgebra of  $\hat{H}$ . This means that any fusion category symmetry  $\mathbb{C}$  can be treated as a subalgebra in some dual weak Hopf algebra  $\hat{H}$ . To summarize, we have

**Theorem 1.** For (1+1)D phase, the fusion category symmetry C can be equivalently characterized by some dual weak Hopf algebra  $\hat{H}$ , where H is the weak Hopf algebra that satisfies  $\operatorname{Rep}(H) \simeq C$  as fusion categories.

<sup>&</sup>lt;sup>9</sup>In this work, we will interchangeably use  $\hat{H}$  and  $H^{\vee}$  do denote the dual space of H.

The above discussion suggests that weak Hopf symmetry is the most general symmetry for (1 + 1)dimensional systems, with fusion category symmetry as a special case. As seen in the previous subsection, both group symmetry and Hopf symmetry are special cases of weak Hopf symmetry. Furthermore, invertible group symmetry can be viewed as a fusion category symmetry by categorifying the group. For a finite group *G*, we introduce the discrete category <u>*G*</u>, whose objects are the group elements, and define the fusion as

$$g \otimes h := gh. \tag{3.45}$$

The fusion category symmetry  $\underline{G}$  corresponds exactly to group symmetry. Notice that  $\underline{G}$  has the same fusion rule as Vect<sub>G</sub>. Recall that Vect<sub>G</sub>  $\simeq$  Rep( $\mathbb{C}[G]^{\vee}$ ), where H is chosen as  $\mathbb{C}[G]$ . Thus, the fusion category symmetry  $\underline{G}$  is equivalent to dual weak Hopf symmetry  $\hat{H} = \mathbb{C}[G]^{\vee\vee} \cong \mathbb{C}[G]$ , which aligns perfectly with the structure of invertible symmetry.

Moreover, when considering general (not necessarily connected) weak Hopf symmetries, this framework also includes multifusion category symmetries as a special case.

Weak Hopf qudit	$\mathcal{H} = H$					
Regular action	$ \begin{split} \overrightarrow{X}_{g} h\rangle &=  gh\rangle, \ \overleftarrow{X}_{g} h\rangle =  hS^{-1}(g)\rangle, \\ \widetilde{\overrightarrow{X}}_{g} h\rangle &=  S^{-1}(g)h\rangle, \ \overleftarrow{\overleftarrow{X}}_{g} h\rangle =  hg\rangle \end{split} $					
Dual action	$\vec{Z}_{\psi} h\rangle =  \psi \rightarrow h\rangle = \sum_{(h)} \psi(h^{(2)}) h^{(1)}\rangle$ $\vec{Z}_{\psi} h\rangle =  h \leftarrow \hat{S}(\psi)\rangle = \sum_{(h)} \psi(S(h^{(1)})) h^{(2)}\rangle$ $\vec{Z}_{\psi} h\rangle =  \hat{S}(\psi) \rightarrow h\rangle = \sum_{(h)} \psi(S(h^{(2)})) h^{(1)}\rangle,$ $\vec{Z}_{\psi} h\rangle =  h \leftarrow \psi\rangle = \sum_{(h)} \psi(h^{(1)}) h^{(2)}\rangle$					

## 3.3 Weak Hopf symmetric single qudit system

Table 2. Summary of quantum operations for weak Hopf qudit system.

Now let us consider a simplest example of weak Hopf symmetric system, a  $C^*$  weak Hopf qudit. By a weak Hopf qudit we mean a complex weak Hopf algebra H with inner product given by

$$\langle a, b \rangle = \Lambda(a^*b),$$
 (3.46)

where  $\Lambda$  is the Haar integral of  $\hat{H}$  (also called Haar measure of H). We will hereinafter denote Haar integral of  $\hat{H}$  as  $\lambda$  and Haar integral of  $\hat{H}$  as  $\Lambda$ . For the specail case of group algebra  $H = \mathbb{C}[G]$ , the Haar integral is of the form

$$\lambda = \frac{1}{|G|} \sum_{g \in G} g, \quad \Lambda = \delta_{1_G}(\cdot).$$
(3.47)

The inner product for this case it the usual  $\langle g, h \rangle = \delta_{1_G}(g^*h) = \delta_{1_G,g^{-1}h} = \delta_{g,h}$  for all  $g, h \in G^{10}$ .

<sup>&</sup>lt;sup>10</sup>Recall that the involution of  $\mathbb{C}[G]$  is defined by  $g^* = g^{-1}$  for all  $g \in G$ , extended antilinearly over the entire space.

The regular action of a weak Hopf algebra on itself can be viewed as a generalization of Pauli X-type operators. For the left action  $H \curvearrowright H$ , we define:

$$\vec{X}_g|h\rangle = |gh\rangle, \quad \overleftarrow{X}_g|h\rangle = |hS^{-1}(g)\rangle.$$
 (3.48)

For the right action  $H \curvearrowleft H$ , we define:

$$\tilde{\vec{X}}_g|h\rangle = |S^{-1}(g)h\rangle, \quad \tilde{\vec{X}}_g|h\rangle = |hg\rangle.$$
(3.49)

The use of  $S^{-1}$  in the above definitions ensures the commutative local stabilizer in weak Hopf lattice gauge theory [52] <sup>11</sup>. Additionally, note that for a  $C^*$  Hopf algebra,  $S^{-1} = S$ .

There are also canonical actions of the dual weak Hopf algebra  $\hat{H}$  on the Hopf qudit H, defined using Sweedler's notation as follows:

$$\varphi \rightharpoonup x := \sum_{(x)} x^{(1)} \langle \varphi, x^{(2)} \rangle, \quad x \leftarrow \varphi := \sum_{(x)} \langle \varphi, x^{(1)} \rangle x^{(2)},$$
 (3.50)

for all  $\varphi \in \hat{H}$  and  $x \in H$ . The operators corresponding the  $\hat{H}$  action on weak Hopf qudit H can be regarded as generalized Pauli Z-type operators. For left action  $\hat{H} \frown H$ , we define:

$$\vec{Z}_{\psi}|h\rangle = |\psi \rightarrow h\rangle = \sum_{(h)} \psi(h^{(2)})|h^{(1)}\rangle, \qquad (3.51)$$

$$\overleftarrow{Z}_{\psi}|h\rangle = |h \leftarrow \widehat{S}(\psi)\rangle = \sum_{(h)} \psi(S(h^{(1)}))|h^{(2)}\rangle.$$
(3.52)

For right action  $H \curvearrowleft \hat{H}$ , we define

$$\vec{Z}_{\psi}|h\rangle = |\hat{S}(\psi) \rightarrow h\rangle = \sum_{(h)} \psi(S(h^{(2)}))|h^{(1)}\rangle, \qquad (3.53)$$

$$\tilde{\tilde{Z}}_{\psi}|h\rangle = |h \leftarrow \psi\rangle = \sum_{(h)} \psi(h^{(1)})|h^{(2)}\rangle.$$
(3.54)

Since the character  $\chi_{\Gamma}$  of irreducible representations  $\Gamma \in \operatorname{Irr}(H)$  lies in  $\hat{H}$ , we can define  $\vec{Z}_{\Gamma}, \vec{Z}_{\Gamma}, \vec{Z}_{\Gamma}, \vec{Z}_{\Gamma}, \vec{Z}_{\Gamma}$  by replacing  $\psi$  with  $\chi_{\Gamma}$  in the expressions given above.

It is convenient to introduce representation-matrix-valued Z-type operators:

$$Z_{\Gamma} = \sum_{(h)} \Gamma(h^{(2)}) | h^{(1)} \rangle, \quad Z_{\Gamma}^{\ddagger} = \sum_{(h)} \Gamma(S(h^{(1)})) | h^{(2)} \rangle.$$
(3.55)

Similarly, we define

$$\tilde{Z}_{\Gamma} = \sum_{(h)} \Gamma(h^{(1)}) |h^{(2)}\rangle, \quad \tilde{Z}_{\Gamma}^{\ddagger} = \sum_{(h)} \Gamma(S(h^{(2)})) |h^{(1)}\rangle.$$
(3.56)

Using Tr' to denote the trace over the representation space, we obtain

$$\vec{Z}_{\Gamma} = \operatorname{Tr}' Z_{\Gamma}, \quad \overleftarrow{Z}_{\Gamma} = \operatorname{Tr}' Z_{\Gamma}^{\ddagger}, \quad \overleftarrow{Z}_{\Gamma} = \operatorname{Tr}' \tilde{Z}_{\Gamma}, \quad \vec{Z}_{\Gamma} = \operatorname{Tr}' \tilde{Z}_{\Gamma}^{\ddagger}. \quad (3.57)$$

<sup>&</sup>lt;sup>11</sup>Alternatively, we can choose to include  $S^{-1}$  in the weak Hopf Pauli Z operators, in which case there is no need to use  $S^{-1}$  in the definition of the weak Hopf Pauli X operators.

*Remark* 3.2. Following the notation of Kitaev [49, 51, 52, 93], we have  $L_{+} = \vec{X}$ ,  $L_{-} = \vec{X}$ ,  $T_{+} = \vec{Z}$ , and  $T_{-} = \vec{Z}$ . The notations here primarily follows Refs. [54, 78, 94], with slight modifications to accommodate the weak Hopf algebra case.

The simplest example of a Hamiltonian with weak Hopf symmetry can be constructed from the center of *H*. Let  $\gamma \in \text{Center}(H)$  (the center of *H* is defined as  $\text{Center}(H) = \{h \in H | hg = gh, \forall g \in H\}$ ), and define  $\mathbb{H} = \vec{X}_{\gamma}$ . Then,  $[\vec{X}_h, \mathbb{H}] = 0$  for all  $h \in H$ . Similarly, a model exhibiting comodule algebra symmetry can be constructed.

#### 4 Pairing-based weak Hopf tensor network states

The key feature of Hopf qudits is their comultiplication. In Ref. [54], we used the Hopf tensor network as a basic tool to solve the Hopf cluster state model. For the more general weak Hopf cluster ladder model, we need to apply a generalization [49, 51, 52, 54, 104] of the Hopf tensor network, which is based on the pairing of weak Hopf algebras and dual weak Hopf algebras (Eqs. (3.39)-(3.43)).

**Definition 5.** A pairing  $\langle \bullet, \bullet \rangle : M \otimes N \to \mathbb{C}$  between two weak Hopf algebras M, N is a bilinear map satisfying

$$\langle hg, a \rangle = \sum_{(a)} \langle h, a^{(1)} \rangle \langle g, a^{(2)} \rangle, \tag{4.1}$$

$$\langle h, ab \rangle = \sum_{(h)} \langle h^{(1)}, a \rangle \langle h^{(2)}, b \rangle, \tag{4.2}$$

$$\langle 1_M, a \rangle = \varepsilon_N(a),$$
 (4.3)

$$\langle h, 1_N \rangle = \varepsilon_M(h),$$
 (4.4)

$$\langle S_M(h), a \rangle = \langle h, S_N(a) \rangle.$$
 (4.5)

Note that the pairing between  $\hat{H}$  and H is a specific instance of the pairing defined above. Using this, we introduce a generalized pairing-based weak Hopf (PWH) tensor network with advantageous properties, suitable for solving the lattice model discussed later. While our primary focus is the pairing between  $\hat{H}$  and H, generalization to other pairings is straightforward.

We will draw elements  $h \in H$  and  $\psi \in \hat{H}$  and their pairing  $\langle \psi, h \rangle = \psi(h) \in \mathbb{C}$  as

$$\frac{1}{h}, \frac{1}{h}, \frac{1}{h}.$$
(4.6)

The diagram for *H* is read upwards, while the diagram for  $\hat{H}$  is read downwards. The tensor product  $h \otimes g$  (or  $\psi \otimes \phi$ ) is represented by concatenating two diagrams and swapping of elements is represented as

$$\tau(h \otimes g) = \bigvee_{\psi} (\psi \otimes \phi) = \bigvee_{\psi} (\psi \otimes \phi), \quad (4.7)$$

where  $\tau = \tau^{-1}$  denotes the swapping map in Vect<sub>C</sub>, allowing us to avoid specifying upper or lower crossings. The multiplication in *H* and  $\hat{H}$  are represented as

$$h \cdot g = \underbrace{\bigwedge_{h}}_{g}, \quad \psi \cdot \phi = \underbrace{\bigvee_{h}}_{g}. \tag{4.8}$$

The comultiplication can be represented as

$$\Delta(h) = \bigvee_{\psi^{(1)}}^{h^{(1)}} h^{(2)}, \quad \Delta(\psi) = \bigvee_{\psi^{(1)}}^{\psi^{(2)}} \psi^{(2)}.$$
(4.9)

The counit  $\varepsilon$  of H is an element of  $\hat{H}$  and it is the unit element:  $1_{\hat{H}} = \varepsilon$ . The unit  $1_H \in H$  is the counit of  $\hat{H}$  in the sense that  $\hat{\varepsilon}(\psi) = \psi(1_H)$ . Both of them are removable under the respective multiplication map. The antipodes of H and  $\hat{H}$  are represented as

$$S(h) = \underbrace{\stackrel{\bullet}{\overset{\bullet}{\overset{\bullet}{\overset{\bullet}{\overset{\bullet}}}}}_{h}, \quad \hat{S}(\psi) = \underbrace{\stackrel{\bullet}{\overset{\bullet}{\overset{\bullet}{\overset{\bullet}}}}_{\bullet}. \tag{4.10}$$

The pairing diagram introduced above has the convenient property of satisfying the pairing conditions (Eqs. (3.39)-(3.43)). Eq. (3.39) can be represented as

When read from top to bottom, this represents the left-hand side of Eq. (3.39), and when read from bottom to top, it represents the right-hand side of Eq. (3.39). Similarly, Eqs. (3.41) and (3.43) can be represented as

$$(4.12)$$

The tensor network representation presented here is more compact and concise than that in Ref.[54]. It can be effectively utilized to solve the cluster ladder model (see Sections6 and 7 for details).

## 5 Symmetry TFT for weak Hopf lattice gauge theory

The Symmetry TFT, also known as topological holography, provides a general framework for understanding non-invertible or categorical symmetries in both gapped and gapless phases [35, 47, 56–64]. For an *n*-dimensional quantum field theory  $QFT_n$  with fusion category symmetry S, the corresponding Symmetry TFT  $\mathcal{Z}(S)$  has the advantage of separating the topological data of the fusion category symmetry



Figure 1. An illustration of a quantum double lattice.

from the non-topological dynamics of the theory. The symmetry S is encoded in the symmetry boundary  $\mathcal{B}_{sym}^{S}$ , while the non-topological data resides on the physical boundary  $\mathcal{B}_{phys}$ . Upon compactification of the (n+1)-dimensional Symmetry TFT, we obtain an *n*-dimensional QFT<sub>n</sub> with non-invertible symmetry S.

## 5.1 Weak Hopf lattice gauge theory

The weak Hopf lattice gauge theory, also referred to as the weak Hopf quantum double model, is comprehensively examined in Refs. [52, 131] as a lattice-based realization of the weak Hopf generalization of BF-theory or Dijkgraaf-Witten theory [132]. For a weak Hopf algebra H, the representation category Rep(H) is typically a multifusion category, establishing an equivalence between the weak Hopf lattice gauge theory and the multifusion string-net model [53], a lattice representation of the Turaev-Viro-Barrett-Westbury topological quantum field theory (TQFT). Although the input data Rep(H) forms a multifusion category, the associated topological excitations are described by a modular tensor category,  $\mathcal{Z}(\text{Rep}(H)) \simeq \text{Rep}(D(H))$  as braided fusion category. Here, D(H) represents the quantum double of H, while  $\mathcal{Z}(\text{Rep}(H))$  denotes the Drinfeld center of Rep(H) [53].

Since quantum double plays a crucial role in our construction, let us briefly review its definition here. Consider a weak Hopf algebra H. The tensor product  $\hat{H}^{cop} \otimes H$  is equipped with an algebra structure where multiplication is given by [133–135]

$$(\varphi \otimes h)(\psi \otimes g) = \sum_{(h)} \sum_{(\psi)} \varphi \psi^{(2)} \otimes h^{(2)} g \langle \psi^{(1)}, S^{-1}(h^{(3)}) \rangle \langle \psi^{(3)}, h^{(1)} \rangle,$$
(5.1)

where we use the pairing between  $\hat{H}$  and H and the comultiplication of  $\psi$  is considered within  $\hat{H}$ . It is useful to rewrite the multiplication using the so-called straightening relation:

$$h \cdot \psi = \psi^{(2)} \otimes g^{(2)} \langle \psi^{(1)}, S^{-1}(h^{(3)}) \rangle \langle \psi^{(3)}, h^{(1)} \rangle.$$
(5.2)

The unit of this algebra is given by  $\varepsilon \otimes 1_H$ . The linear span *M* of the elements

$$\varphi \otimes xh - \varphi(x \to \varepsilon) \otimes h, \quad x \in H_L,$$
(5.3)

$$\varphi \otimes yh - \varphi(\varepsilon \leftarrow y) \otimes h, \quad y \in H_R,$$
(5.4)

constitutes a two-sided ideal of  $\hat{H}^{cop} \otimes H$ . We denote the quotient algebra  $(\hat{H}^{cop} \otimes H)/M$  as D(H), with equivalence classes in D(H) represented by  $[\varphi \otimes h]$  for  $\varphi \otimes h \in \hat{H}^{cop} \otimes H$ .

**Definition 6** (Quantum double). The quantum double of a weak Hopf algebra H, denoted by D(H), is a weak Hopf algebra equipped with the following structure:

(1) The multiplication operation is defined by

$$[\varphi \otimes h][\psi \otimes g] = \sum_{(\psi),(h)} [\varphi \psi^{(2)} \otimes h^{(2)}g] \langle \psi^{(1)}, S^{-1}(h^{(3)}) \rangle \langle \psi^{(3)}, h^{(1)} \rangle.$$

(2) The unit element is

 $[\varepsilon \otimes 1_H].$ 

(3) The comultiplication is given by

$$\Delta([\varphi \otimes h]) = \sum_{(\varphi),(h)} [\varphi^{(2)} \otimes h^{(1)}] \otimes [\varphi^{(1)} \otimes h^{(2)}].$$

(4) The counit is defined as

$$\varepsilon([\varphi \otimes h]) = \langle \varphi, \varepsilon_R(S^{-1}(h)) \rangle$$

(5) The antipode is given by

$$S([\varphi \otimes h]) = \sum_{(\varphi),(h)} [\hat{S}^{-1}(\varphi^{(2)}) \otimes S(h^{(2)})] \langle \varphi^{(1)}, h^{(3)} \rangle \langle \varphi^{(3)}, S^{-1}(h^{(1)}) \rangle.$$

Weak Hopf quantum double lattice model Consider a 2d manifold  $\Sigma$ , a lattice on it is a cellulation  $C(\Sigma) = C^0(\Sigma) \cup C^1(\Sigma) \cup C^2(\Sigma)$  with  $C^0(\Sigma)$  denoting the set of vertices,  $C^1(\Sigma)$  denoting the set of edges, and  $C^2(\Sigma)$  denoting the set of faces. The bulk input data is a weak Hopf algebra H. For a directed lattice on 2d closed surface, we assign each edge a weak Hopf qudit  $\mathcal{H}_e = H$ , the total Hilbert space is  $\mathcal{H}_{tot} = \bigotimes_{e \in C^1(\Sigma)} \mathcal{H}_e$ . All edges of the lattice are oriented, to construct the local stabilizers, we adopt the following convention:

$$\overleftarrow{X} \bullet \overrightarrow{Z} \bullet \overrightarrow{X}$$
 (5.5)

With this convention, consider the following configuration:

$$j_5 \xrightarrow{j_4} j_1 j_2 \\ j_6 j_6$$
 (5.6)

Here, s = (v, f) is an edge-face link, and we assume a counterclockwise order around each face and vertex, with the starting point at link *s*.

The corresponding vertex and face operators are defined based on comultiplication as follows:

$$\mathbb{A}^{h}(s) = \sum_{(h)} \overleftarrow{X}_{h^{(1)}}(j_{4}) \otimes \overrightarrow{X}_{h^{(2)}}(j_{5}) \otimes \overrightarrow{X}_{h^{(3)}}(j_{6}) \otimes \overleftarrow{X}_{h^{(4)}}(j_{1}),$$

$$\mathbb{B}^{\varphi}(s) = \sum_{(\varphi)} \overleftarrow{Z}_{\varphi^{(1)}}(j_{1}) \otimes \overleftarrow{Z}_{\varphi^{(2)}}(j_{2}) \otimes \overrightarrow{Z}_{\varphi^{(3)}}(j_{3}) \otimes \overrightarrow{Z}_{\varphi^{(4)}}(j_{4}),$$
(5.7)

where  $h \in H$  and  $\varphi \in \hat{H}$ . At the link *s*, we have a representation of quantum double D(H):

$$[\varphi \otimes h] \mapsto \mathbb{B}^{\varphi}(s) \mathbb{A}^{h}(s), \tag{5.8}$$

where  $[\varphi \otimes h]$  is an element in quantum double, see [52] for the proof.

For any site *s*, the operators  $\mathbb{A}^{h}(s)$  and  $\mathbb{B}^{\varphi}(s)$  obey the following commutation relations [52]:

$$\mathbb{A}^{h}(s)\mathbb{B}^{\varphi}(s) = \sum_{(h),(\varphi)} \mathbb{B}^{\varphi^{(2)}}(s)\mathbb{A}^{h^{(2)}}(s)\langle\varphi^{(1)}, S^{-1}(h^{(3)})\rangle\langle\varphi^{(3)}, h^{(1)}\rangle,$$
(5.9)

$$\mathbb{B}^{\varphi}(s)\mathbb{A}^{xh}(s) = \mathbb{B}^{\varphi(x \to \varepsilon)}(s)\mathbb{A}^{h}(s), \mathbb{B}^{\varphi}(s)\mathbb{A}^{yh}(s) = \mathbb{B}^{\varphi(\varepsilon \leftarrow y)}(s)\mathbb{A}^{h}(s), \tag{5.10}$$

for any  $x \in H_L = \varepsilon_L(H)$  and  $y \in H_R = \varepsilon_R(H)$ . From this, we obtain that if *h* is taken as the Haar integral  $\lambda$  and  $\varphi$  is taken as the Haar measure  $\Lambda$ , then the corresponding local operators are projectors that commute with each other. Additionally, the definition of the operator does not depend on the initial link *s* but only on the vertex and faces. Therefore, we define the operators as  $\mathbb{A}_v = \mathbb{A}^{\lambda}(s)$  and  $\mathbb{B}_f = \mathbb{B}^{\Lambda}(s)$ . The Hamiltonian is defined as

$$\mathbb{H} = -\sum_{\nu} \mathbb{A}_{\nu} - \sum_{f} \mathbb{B}_{f}.$$
(5.11)

The topological excitations of the model are characterized by the representation category of quantum double D(H).

**Topological boundary condition of weak Hopf lattice gauge theory** The topological boundary condition of weak Hopf lattice gauge theory can be characterized in several different ways. From anyon condensation perspective, we have the following (see Refs. [51, 52, 96, 136–139]):

- Lagrangian algebra  $\mathcal{A}$ . The topological boundary condition is characterized by a Lagrangian algebra in the bulk phase  $\operatorname{Rep}(D(H))$ , the boundary phase is given by anyon condensation  $\mathcal{B} = \operatorname{Rep}(D(H))_{\mathcal{A}}$ , the category of  $\mathcal{A}$ -modules in  $\operatorname{Rep}(D(H))$ .
- Frobenius algebra  $\mathcal{F}$ . For this case, the topological boundary condition is characterized by a Frobenius algebra in the bulk phase  $\operatorname{Rep}(D(H))$ . The anyon condensation is realized in two steps, we first take the quotient category  $\operatorname{Rep}(D(H))/\mathcal{F}$ , then we take idempotent complete to obtain the boundary phase  $\mathcal{B} = \operatorname{IC}(\operatorname{Rep}(D(H))/\mathcal{F})$ .

Since  $\operatorname{Rep}(D(H))$  is non-chiral topological phase, the boundary can also be characterized at the level a weak Hopf gauge symmetry W (or equivalently at the level of (multi)fusion category  $\operatorname{Rep}(H)$ ).

- In the framework of quantum double lattice model, the bulk gauge symmetry is the weak Hopf algebra *W*, the boundary gauge symmetry is a comodule algebra *K* over *H*.
- In the framework of string-net model, the bulk input (multi)fusion category is Rep(H), the boundary is determined by a Rep(H)-module category  $\mathcal{M}$ . For a given comodule algebra H, the category of K-modules  $\text{Mod}_K$  is of module category over Rep(H).



**Figure 2**. Depiction of the SymTFT sandwich: the symmetry boundary is drawn in cyan, while the physical boundary is drawn in magenta.

## 5.2 Algebraic theory of SymTFT for weak Hopf lattice gauge theory

The SymTFT is a sandwich structure as shown in Figure 2, the sandwich manifold defined as  $\mathbb{M}^{1,1} \times [0,1]$  where  $\mathbb{M}^{1,1}$  represents the (1+1) D manifold on which our system lies on. The sandwich manifold  $\mathbb{M}^{1,1} \times [0,1]$  is a (2+1)D manifold that has two boundaries. The basic idea is we put the non-invertible symmetry on one of the boundary, and put our physical system on the other boundary, then after doing compactification over the interval [0,1], we obtain a (1+1)D system with non-invertible symmetry.

Consider a (1+1)D phase  $\mathcal{T}$  with fusion category symmetry given by the fusion category  $\mathcal{B}_{sym}$ , whose physical information is captured by  $\mathcal{B}_{phys}$ . The SymTFT for this phase is a triple ( $\mathcal{B}_{sym}$ ,  $\mathcal{B}_{phys}$ ,  $\mathcal{Z}(\mathcal{B}_{sym})$ ), where  $\mathcal{Z}(\mathcal{B}_{sym})$  is the Drinfeld center of  $\mathcal{B}_{sym}$ , a unitary modular tensor category characterizing a (2+1)D topological phase. The symmetry boundary  $\mathcal{B}_{sym}$  must be a gapped topological boundary condition of  $\mathcal{Z}(\mathcal{B}_{sym})$ . The physical boundary  $\mathcal{B}_{phys}$  is also a boundary condition for the bulk TQFT, but it is not necessarily topological or even gapped. If  $\mathcal{T}$  is gapless,  $\mathcal{B}_{phys}$  corresponds to a gapless boundary condition. Conversely, if  $\mathcal{T}$  is gapped,  $\mathcal{B}_{phys}$  represents a gapped topological boundary condition.

For this section, we will assume that  $\mathcal{T}$  is gapped. Therefore, both  $\mathcal{B}_{sym}$  and  $\mathcal{B}_{phys}$  are chosen as gapped boundary conditions of the TQFT  $\mathcal{Z}(\mathcal{B}_{sym})$ . Now, we set the (2 + 1)D bulk theory as the weak Hopf lattice gauge theory  $\mathcal{Z}(\mathcal{B}_{sym}) \simeq_{\otimes, br} \text{Rep}(D(H)) \simeq_{\otimes, br} \mathcal{Z}(\text{Rep}(H))$ , where the weak Hopf gauge symmetry is given by H. Note that this does not imply that  $\mathcal{B}_{sym}$  is monoidally equivalent to Rep(H); it only means that they are weak Morita equivalent (i.e., their Drinfeld centers are equivalent as unitary modular tensor categories). All topological boundary conditions of a weak Hopf lattice gauge theory are weak Morita equivalent, allowing us to choose  $\mathcal{B}_{sym}$  as an arbitrary topological gapped boundary condition. For weak Hopf gauge symmetry H, the corresponding SymTFT is a sandwich that consists of the following data:

- For the (2+1)D bulk, we put a weak Hopf lattice gauge theory whose topological excitation is given by the representation category Rep(D(H)) of the quantum double D(H).
- The symmetry boundary is a topological boundary condition of bulk lattice gauge phase Rep(D(H)), this can be characterized by a comodule algebra K over weak Hopf algebra H. We denote the boundary vacuum charge as  $\mathcal{A}_K$ , which is a Lagrangian algebra in Rep(D(H)).
- The physical boundary condition  $\mathcal{B}_{phys}$  is also chosen as a topological boundary condition, thus it is also characterized by a comodule algebra J over H. We denote the boundary vacuum charge as

 $\mathcal{A}_J$ , which is also a Lagrangian algebra in  $\operatorname{Rep}(D(H))$ .

- The local operators of are generated by Wilson lines that can end at both boundaries. In weak Hopf lattice gauge theory, the Wilson lines are ribbon operators that connect two boundaries [50–52, 103, 114], representing the transport of vacuum charge from one boundary to the other.
- The symmetry action on local operators is realized by half-braiding. In weak Hopf lattice gauge theory, this is characterized by boundary closed ribbons [50–52, 103, 114], which are ribbons with both ends anchored on the symmetry boundary. Since the symmetry boundary closed ribbon intersects with the local operator ribbon, this intersection can be used to realize the half-braiding.

## 6 Weak Hopf cluster ladder model I: smooth-rough construction

In this section, we consider a special case of cluster ladder model, for which one boundary is chosen as smooth boundary and the other is chosen as rough boundary. The main result can be summarized as follows:

**Theorem 2** (Informal summary). Any SymTFT with (multi)fusion category symmetry S can be realized via a cluster state model based on the smooth-rough construction. We first use Tannaka-Krein reconstruction (or tube algebra approach [53, 119]) to obtain a weak Hopf gauge symmetry  $H_S$  such that  $S \simeq \text{Rep}(H_S)$ . Then, we construct the cluster state model based on this weak Hopf symmetry  $\mathbb{H}_{\text{cluster}}[H_S]$ , which realizes the S-SymTFT.

## 6.1 Weak Hopf cluster state model

As shown in Ref. [54] for Hopf case (and finite group case), the CSS-type cluster state model can be regarded as the quantum double model that has one smooth boundary and one rough boundary. Inspired by the construction there, we define weak Hopf cluster state model as a weak Hopf quantum lattice gauge theory with one smooth boundary and one rough boundary.

**Definition** 7 (Weak Hopf cluster state model). *The weak Hopf cluster state model is a weak Hopf lattice gauge theory with one rough boundary and one smooth boundary.* 

For bulk weak Hopf gauge symmetry H, the smooth boundary corresponds to comodule algebra  $K_s = H$ , namely, choose boundary gauge symmetry as H itself. The corresponding boundary excitation is  $\mathcal{B}_s \cong \operatorname{Rep}(H)$ . The rough boundary corresponds to the trivial comodule algebra  $K_r = \mathbb{C}$ , the rough boundary excitation  $\mathcal{B}_r = \operatorname{Rep}(\hat{H})$ .

Starting from a weak Hopf quantum double model with one smooth boundary and one rough boundary, and after coarse-graining the bulk, we ultimately obtain the following ladder lattice:



There are two types of stabilizers: one is the smooth boundary vertex operator  $\mathbb{A}_{v_s}$ , and the other is the face operator  $\mathbb{B}_f$  (the rough boundary vertex operators are trivial and thus omitted). The Hamiltonian reads

$$\mathbb{H}_{\text{cluster}}[H, K_s, K_r] = -\sum_{\nu_s} \mathbb{A}_{\nu_s} - \sum_f \mathbb{B}_f.$$
(6.2)

Recall that  $\mathbb{A}_{v_s} = \mathbb{A}_{v_s}^{\lambda}$  and  $\mathbb{B}_f = \mathbb{B}_f^{\Lambda}$ , where  $\lambda$  and  $\Lambda$  represent the Haar integral and measure of the weak Hopf algebra H. Consequently, these local stabilizers commute with one another.

Since for arbitrary (multi)fusion category symmetry S, we can find a corresponding weak Hopf algebra  $H_S$  to realize it, the weak Hopf cluster state model provides a lattice realization of all S-symmetric theories. Moreover, since there is no anyonic tunneling channel between the smooth and rough boundaries, this model realizes an S-SPT phase.

*Remark* 6.1. In Ref. [54], we adopt a convention different from the one used here, where the local ordering around each vertex and face is clockwise. Consequently, the vertex operator and face operator must be modified accordingly. Different conventions lead to different lattice models, but they result in the same quantum phase. Additionally, when constructing the lattice model, it is necessary to set an initial configuration of orientations for all edges. Different choices of this configuration also yield distinct lattice models; however, their quantum phases remain identical. In the case of  $\mathbb{C}[\mathbb{Z}_2]$ , all these different models are the same due to the fact that the inverse of 1 is itself, and they correspond to the CSS-type cluster state model [92]. For a general Abelian group, varying the local ordering around vertices and faces does not alter the lattice model, but different initial configurations still result in distinct models. In contrast, for a non-Abelian group, both the local ordering and the initial configurations lead to different models [54, 78].

*Remark* 6.2 (String-net realization). In the multifusion string-net framework [53, 119, 140], the input data is a unitary multifusion category  $\mathcal{C} = \operatorname{Rep}(H)$  for a weak Hopf gauge symmetry H. The smooth boundary is a C-module category  $\mathcal{M}_s = \mathcal{C}$ , with boundary excitations given by  $\mathcal{B}_s = \operatorname{Fun}_{\mathbb{C}}(\mathcal{M}_s, \mathcal{M}_s) = \mathcal{C}$ , where  $\operatorname{Fun}_{\mathbb{C}}(\mathcal{C}, \mathbb{C})$  is the category of all C-module functors from  $\mathcal{M}_s$  to itself. The rough boundary is a C-module category  $\mathcal{M} = \operatorname{Vect}_{\mathbb{C}}$ , with  $\operatorname{Vect}_{\mathbb{C}}$  as the category of all finite-dimensional complex vector spaces. Rough boundary excitations are given by  $\mathcal{B}_r = \operatorname{Fun}_{\mathbb{C}}(\operatorname{Vect}_{\mathbb{C}}, \operatorname{Vect}_{\mathbb{C}}) = \operatorname{Rep}(\hat{H})$ . Using the Kitaev-Kong construction, we can also obtain a lattice model for the cluster state model.

## 6.2 Weak Hopf cluster state

The cluster state model can be solved using the weak Hopf tensor network introduced in Section 4. Assuming periodic boundary conditions for the cluster model, we need to consider two types of edges: the bottom smooth boundary edge (blue in Eq. 6.1) and the bulk edge (black in Eq. 6.1).

For boundary edge, the corresponding local tensor is the comultiplication of Haar integral  $\lambda$ :

$$\Delta(\lambda) = \bigwedge_{\lambda^{(1)}}^{\lambda^{(2)}} := \bigwedge_{\lambda^{(1)}}^{\lambda^{(1)}} A^{(2)} .$$
(6.3)

Similarly, for bulk edge, the local tensor corresponds to  $\Delta_2(\lambda)$ :

$$\Delta(\lambda) = \begin{array}{c} \lambda^{(3)} & \lambda \\ & \lambda^{(2)} \\ & \lambda^{(1)} \end{array} := \begin{array}{c} \lambda^{(1)} & \lambda^{(2)} & \lambda^{(3)} \\ & & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & & \\ & & & \\ & & & \\ & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ &$$

Since Haar integral  $\lambda$  is cocommutative, we use a simplified notation in which the comultiplication components (1), (2), . . . can be cyclically permuted. For example,  $\lambda^{(1)} \otimes \lambda^{(2)} \otimes \lambda^{(3)} = \lambda^{(3)} \otimes \lambda^{(1)} \otimes \lambda^{(2)}$ .

For face, a local tensor corresponds to the Haar measure  $\Lambda \in \hat{H}$  is needed to glue the edge tensors together:

$$(\mathrm{id}\otimes\mathrm{id}\otimes\hat{S})\circ\hat{\Delta}(\Lambda) = \bigwedge_{\Lambda^{(1)}}^{\hat{S}(\Lambda^{(3)})} - \bigwedge_{\Lambda^{(2)}}^{\Lambda} = \bigwedge_{\Lambda^{(1)}}^{\Lambda^{(2)}} = \bigwedge_{\Lambda^{(1)}}^{\Lambda^{(2)}} .$$
(6.5)

The antipode  $\hat{S}$  is introduced because the direction of the lattice is chosen as in Eq. (6.1). If a different lattice configuration is chosen, the antipode may be placed on different legs of the tensor.

The cluster state is defined as the following tensor network state:



where we have drawn the free leg (corresponding to physical degrees of freedom) in red. The cluster state is the ground state of the cluster state model defined in Eq. (6.2)

## 6.3 Symmetry of the model

The correspondence between the cluster state model and the quantum double model can also be extended to their symmetry operators. We propose that the symmetry of the cluster state model is governed by the ribbon operators. Ribbon operators are fundamental tools for studying topological excitations. In this subsection, following the works of [50–52, 93, 96, 97], we recall the construction of ribbon operators. For more details, see [51, 52].

A direct triangle  $\tau = (s_0, s_1, e)$  consists of two adjacent face-vertex links,  $s_0$  and  $s_1$ , connected by a directed edge e. Similarly, a dual triangle  $\tilde{\tau} = (s_0, s_1, \tilde{e})$  consists of two adjacent links  $s_0$  and  $s_1$  connected by a dual edge  $\tilde{e}$  (by "dual edge" we mean an edge in dual lattice, we will draw dual edge as dashed line, the orientation of dual edge is obtained by rotating the direct edge  $\pi/2$  clockwise). The triangle's direction is defined as the direction from  $s_0$  to  $s_1$ , which may or may not align with the edge direction. A triangle is left-handed if its edge lies on the left when traversing in its positive direction, and right-handed otherwise. Left-handed and right-handed direct triangles are denoted by  $\tau_L$  and  $\tau_R$ , respectively, with dual counterparts  $\tilde{\tau}_L$  and  $\tilde{\tau}_R$ . A ribbon  $\rho$  is an ordered sequence of triangles  $\tau_1, \ldots, \tau_n$  with a consistent direction, such that  $\partial_1 \tau_j = \partial_0 \tau_{j+1}$  for all j, and with no self-intersections. A ribbon is closed if it forms a loop, i.e.,  $\partial_1 \tau_n = \partial_0 \tau_1$ . Direct and dual triangles in a directed ribbon must have opposite chiralities. Ribbons, denoted by  $\rho_B$ , consist of left-handed direct triangles and right-handed dual triangles. Type-A ribbons, denoted by  $\rho_B$ , consist of right-handed direct triangles and right-handed dual triangles.

The ribbon operator is constructed recursively, starting with the definition of the triangle operator and extending it through a recursive relation. For triangles, we have (for  $h, x \in H$  and  $\varphi \in \hat{H}$ )

$$\sum_{s_1 \dots s_0}^{x} F^{h,\varphi}(\tau_R)|x\rangle = \varepsilon(h)\overleftarrow{Z}_{\varphi}|x\rangle = \varepsilon(h)|x \leftarrow \widehat{S}(\varphi)\rangle, \tag{6.7}$$

$$F^{h,\varphi}(\tilde{\tau}_R)|x\rangle = \hat{\varepsilon}(\varphi)\overleftarrow{X}_h|x\rangle = \hat{\varepsilon}(\varphi)|x \triangleleft S(h)\rangle, \tag{6.9}$$

$$s_1 \qquad s_0 \qquad F^{h,\varphi}(\tilde{\tau}_R)|x\rangle = \hat{\varepsilon}(\varphi) \overrightarrow{X}_h |x\rangle = \hat{\varepsilon}(\varphi)|h \triangleright x\rangle.$$
(6.10)

$$F^{h,\varphi}(\tilde{\tau}_L)|x\rangle = \hat{\varepsilon}(\varphi)\tilde{X}_h|x\rangle = \hat{\varepsilon}(\varphi)|x \triangleleft h\rangle.$$
(6.11)

$$F^{h,\varphi}(\tilde{\tau}_L)|x\rangle = \hat{\varepsilon}(\varphi)\tilde{\vec{X}}_h|x\rangle = \hat{\varepsilon}(\varphi)|S(h) \triangleright x\rangle.$$
(6.12)

$$x = s_0 \qquad F^{h,\varphi}(\tau_L)|x\rangle = \varepsilon(h)\tilde{Z}_{\varphi}|x\rangle = \varepsilon(h)|x \leftarrow \varphi\rangle, \tag{6.13}$$

For a type-B ribbon  $\rho = \rho_B$  and an element  $h \otimes \varphi \in H \otimes \hat{H}$ , the ribbon operator is defined as  $F^{h,\varphi}(\rho) = F^{h\otimes\varphi}(\rho_B)$ . The operator acts non-trivially only on the edges of  $\rho_B$ . Consider the decomposition  $\rho = \rho_B = \rho_1 \cup \rho_2$ , where both  $\rho_1$  and  $\rho_2$  are oriented in the same direction as  $\rho$ , and  $\partial_1 \rho_1 = \partial_0 \rho_2$ . For  $h \otimes \varphi \in H^{\text{op}} \otimes \hat{H}$ , the ribbon operator on the composite ribbon is defined as

$$F^{h,\varphi}(\rho) = \sum_{(h\otimes\varphi)} F^{(h\otimes\varphi)^{(1)}}(\rho_1) F^{(h\otimes\varphi)^{(2)}}(\rho_2) = \sum_k \sum_{(k),(h)} F^{h^{(1)},\hat{k}}(\rho_1) F^{S(k^{(3)})h^{(2)}k^{(1)},\varphi(k^{(2)}\bullet)}(\rho_2).$$
(6.15)

where  $\{k\}$  is an orthogonal basis of H with  $\{\hat{k}\}$  its dual basis and the comultiplication is taken in the dual weak Hopf algebra  $D(H)^{\vee}$ . This definition is independent of the decomposition  $\rho = \rho_1 \cup \rho_2$ . A similar construction applies for type-A ribbons.

**Symmetry for closed 1d ladder** A 1d closed cluster lattice can also be regarded as a bicycle-wheel quantum double lattice, which is a specific cellulation of a disk. We assign a weak Hopf qudit to each edge, corresponding to the sites of the cluster lattice. The boundary edges correspond to even sites, while the internal edges correspond to odd sites. For example, we have



The solid lattice represents the quantum double lattice, while the dotted lattice represents the cluster lattice, where the qudits are placed on the vertices. In the context of the cluster state model, qudits are typically placed on the vertices of the cluster lattice. To distinguish these from the vertices of the quantum double lattice, we refer to them as sites. Even sites are depicted as blue squares, and odd sites as red dots (see also Figure 5 of Ref. [54] for an illustration of how the 1d cluster lattice is folded into a 2d configuration). Each bicycle-wheel quantum double lattice corresponds to a 1d cluster lattice, and every 1d cluster lattice can be transformed into a bicycle-wheel quantum double lattice. For convenience, we will use these two lattices interchangeably.

The symmetry operators can be easily understood from the perspective of the bicycle-wheel lattice. In this framework, the rough boundary corresponds to a vertex in the bicycle-wheel lattice. Consider a closed ribbon  $\sigma$  surrounding the vertex *w*, consisting exclusively of dual triangles (illustrated in gray as follows):



The corresponding ribbon operator is simply the vertex operator

$$F^{h,\hat{1}}(\sigma) = \mathbb{A}^h_w, \tag{6.17}$$

as there is no direct triangle. Thus, the operator is independent of the unit  $\hat{1} \in \hat{H}$  (we retain this  $\hat{1}$  to emphasize that this is a special case of a more general situation, which will be discussed later). In the finite group case, this operator reduces to the Pauli X-string operator. Note that this symmetry operator acts only on the red edges, which correspond to the odd sites of the cluster lattice.

It is clear  $[\mathbb{A}_{w}^{h}, \mathbb{A}_{v}] = 0$  for all vertex operators  $A_{v}$  on the smooth boundary (depicted as the blue circle). This is because, for the two endpoints of an edge, one operator acts as  $\overrightarrow{X}$  and the other as  $\overleftarrow{X}$ , which commute with each other. Using a result from [131, Corollary 1], if *h* is cocommutative, then  $[\mathbb{A}_{w}^{h}, \mathbb{B}_{f}] = 0$  for all face operators. This implies that the symmetry arising from the rough boundary is

$$\operatorname{Sym}_{\operatorname{rough}} = \operatorname{Cocom}(H),$$
 (6.18)

namely, the set of all cocommutative elements in H.  $h \in \text{Cocom}(H)$  For  $H = \mathbb{C}[G]$ , it follows that  $\text{Cocom}(\mathbb{C}[G]) = \mathbb{C}[G]$ , and thus the symmetry is  $\mathbb{C}[G]$ . This conclusion also aligns with a macroscopic perspective, where we assert that the rough boundary symmetry is  $\text{Rep}(\hat{H})$ , which can be embedded into H. Note that all characters are cocommutative, as  $\chi_{\Gamma}(fg) = \chi_{\Gamma}(gf)$  for all  $\Gamma \in \text{Rep}(\hat{H})$ . Here, the comultiplication of  $\chi_{\Gamma}$  is defined in Definition 5.

For the smooth boundary, we can further deform the bicycle-wheel lattice into a cone as follows:



On the base face, we also have a ribbon  $\xi$  consisting only of direct triangles surrounding the face  $f_s$ . The corresponding symmetry ribbon operator is simply the face operator

$$F^{1,\varphi}(\xi) = \mathbb{B}^{\varphi}_{f_{\tau}},\tag{6.19}$$

as there are no dual triangles. Thus, the operator does not depend on elements in H.

The symmetry operator  $\mathbb{B}_{f_s}^{\varphi}$  commutes with all face operators of the cluster state model because, on the two sides of an edge, the operators are chosen as  $\vec{Z}$  and  $\vec{Z}$  respectively, which always commute. Moreover, if  $\varphi$  is cocommutative, a result from Ref. [131, Corollary 1] shows that  $\mathbb{B}_{f_s}^{\varphi}$  also commutes with all vertex operators of the cluster state model.

This implies that the symmetry arising from the smooth boundary is

$$\operatorname{Sym}_{\operatorname{smooth}} = \operatorname{Cocom}(\hat{H}),$$
 (6.20)

namely, the subalgebra of all cocommutative elements in  $\hat{H}$ . From the macroscopic analysis, all characters in Rep(H) are cocommutative elements in  $\hat{H}$ , which matches well with the result obtained from the lattice here.

The symmetry operators arising from the rough and smooth boundaries commute with each other, as they act non-trivially on different weak Hopf qudits. This is analogous to the  $\mathbb{Z}_2$  cluster state (as well as finite-group or Hopf cluster states), where one symmetry operator acts on odd sites and the other acts on even sites.

**Theorem 3.** The weak Hopf cluster state model on a closed manifold exhibits the symmetry

$$\operatorname{Sym}_{\operatorname{closed cluster}} = \operatorname{Cocom}(H) \times \operatorname{Cocom}(\hat{H}),$$
 (6.21)

which is a subset of  $H \times \hat{H}$ . Moreover, note that Gr(Rep(H)), or more generally,  $R(Rep(H)) = Gr(Rep(H)) \otimes_{\mathbb{Z}} \mathbb{C}$ , is a subalgebra of  $Cocom(\hat{H})$ . Consequently, this symmetry contains a weak Hopf sub-symmetry  $Cocom(H) \times Rep(H)$ , where we use the notation Rep(H) to refer to R(Rep(H)), as is standard in the physical literature.

**Symmetry for open 1d ladder** For an open 1d manifold, the corresponding weak Hopf cluster state model exhibits a larger weak Hopf symmetry:  $H \times \hat{H}$ . Since the symmetry ribbons on the rough and smooth boundaries are not required to be closed ribbons, the elements assigned to the ribbon operators

are no longer constrained to be cocommutative. Nonetheless, the symmetry ribbon operators still commute with the vertex and face operators in the cluster state model [52, 131].

In the case of a finite group, this is analogous to the group symmetry and dual group symmetry  $G \times \hat{G}$ [47, 64]. Using our notation,  $\operatorname{Cocom}(\mathbb{C}[G]) = \mathbb{C}[G]$  and  $\operatorname{Cocom}(\mathbb{C}[G]^{\vee}) = \operatorname{Rep}(G)$ , indicating that both the closed ladder and open ladder share the same symmetry, given by  $\mathbb{C}[G] \times \operatorname{Rep}(G)$ .

When acting on the ground state of the cluster state model, the symmetry ribbon operators create edge modes at the two ends of the open 1D lattice. Consider a more general ribbon operator as illustrated below:



In this case, the ribbon operator  $F^{h,\varphi}(\rho)$  is obtained. A result in [52, Proposition 3] guarantees that  $F^{h,\varphi}(\rho)$  commutes with all vertex and face operators of the cluster state model. At the two ends of the boundary, it creates edge modes.

Theorem 4. The weak Hopf cluster state model on an open manifold exhibits the weak Hopf symmetry

$$\operatorname{Sym}_{\operatorname{open cluster}} = H \times \hat{H}.$$
(6.22)

This symmetry contains a weak Hopf sub-symmetry  $H \times \text{Rep}(H)$ , where we use the notation Rep(H) to refer to R(Rep(H)), with the usual abuse of notation.

### 7 Weak Hopf cluster ladder model II: general construction

In this section, we extend the cluster state model to the more general cluster ladder model, where the topological boundary conditions on the two boundaries can be chosen arbitrarily.

## 7.1 Weak Hopf cluster ladder model

For the most general case, consider a sandwich manifold  $\Sigma \times [0, 1]$ . For the bulk  $\Sigma \times (0, 1)$ , we assign a weak Hopf algebra. On the symmetry boundary  $\Sigma \times \{0\}$ , we assign a weak Hopf comodule algebra K. Depending on the orientation of the boundary, we can choose either left or right comodule algebras as input data [51, 52]. Similarly, on the physical boundary  $\Sigma \times \{1\}$ , we assign a weak Hopf comodule algebra J.

The compactification over the interval [0, 1] can be realized by considering an ultra-thin sandwich lattice, which is a ladder lattice denoted by  $\mathbb{M}^1$ :



The edges for the symmetry boundary are drawn in blue, the edges for the physical boundary are drawn in red, and the bulk edges are drawn in black. Note that in the above lattice, the bulk is positioned on the left side of the symmetry boundary and on the right side of the physical boundary when traversing along the positive direction of the boundary. Consequently, the corresponding comodule algebras K and J must be chosen as a left H-comodule algebra and a right H-comodule algebra, respectively.

We will adopt Sweedler's notation for the left *H*-comodule algebra *K* as  $\beta_K(x) = \sum_{[x]} x^{[-1]} \otimes x^{[0]}$ , where  $x^{[-1]} \in H$  and  $x^{[0]} \in K$ . Similarly, for the right *H*-comodule algebra *J*, we write  $\beta_J(x) = \sum_{[x]} x^{[0]} \otimes x^{[1]}$ , where  $x^{[0]} \in J$  and  $x^{[1]} \in H$ .

For the symmetry boundary, we assign comodule algebra  $\mathcal{H}_{e_s} = K$  on each boundary edge. To build the boundary local stabilizers, we need the notion of *symmetric separability idempotent* of an algebra K(see [51, 52, 117, 141]). By definition, it is an element  $\zeta = \sum_{\langle \zeta \rangle} \zeta^{\langle 1 \rangle} \otimes \zeta^{\langle 2 \rangle} \in K \otimes K$  such that

- 1.  $\sum_{\langle \zeta \rangle} x \zeta^{\langle 1 \rangle} \otimes \zeta^{\langle 2 \rangle} = \sum_{\langle \zeta \rangle} \zeta^{\langle 1 \rangle} \otimes \zeta^{\langle 2 \rangle} x$  for all  $x \in K$ ;
- 2.  $\sum_{\langle \zeta \rangle} \zeta^{\langle 1 \rangle} \zeta^{\langle 2 \rangle} = 1;$
- 3.  $\sum_{\langle \zeta \rangle} \zeta^{\langle 1 \rangle} \otimes \zeta^{\langle 2 \rangle} = \sum_{\langle \zeta \rangle} \zeta^{\langle 2 \rangle} \otimes \zeta^{\langle 1 \rangle}.$

We have used " $\sum_{\langle \zeta \rangle}$ " to represent that  $\zeta$  is a linear combination of elements in  $K \otimes K$ , in order to make the notation consistent with that for comultiplication. It can be shown that  $\zeta$  is an idempotent element of the enveloping algebra  $K \otimes K^{\text{op}}$ . If K is a weak Kac algebra <sup>12</sup> with Haar integral  $\lambda_K$ , then it is straightforward to verify that  $\zeta = \sum_{(\lambda_K)} \lambda_K^{(1)} \otimes S(\lambda_K^{(2)})$  serves as a symmetric separability idempotent. The existence and uniqueness of the symmetric separability idempotent for a finite-dimensional semisimple algebra over an algebraically closed field of characteristic zero are established in [141, Corollary 3.1].

For the symmetry boundary, as depicted in Eq. (7.1), with the bulk on the left-hand side of the boundary, we define the following operator for  $z \otimes w \in K \otimes K^{\text{op}}$ :

Now we define the boundary operator as

$$\mathbb{A}_{\nu_{s}}^{K} = \mathbb{A}_{\nu_{s}}^{\zeta_{K}} = \sum_{\langle \zeta_{K} \rangle} \mathbb{A}_{\nu_{s}}^{\zeta_{K}^{(1)} \otimes \zeta_{K}^{\langle 2 \rangle}}$$
(7.3)

where  $\zeta_K$  is the symmetric separability idempotent of *K*. The Hamiltonian for the symmetry boundary is

$$\mathbb{H}_{\text{sym}} = -\sum_{\nu_s:\text{sym}} \mathbb{A}_{\nu_s}^K.$$
(7.4)

*Remark* 7.1. When K is a weak Kac subalgebra of H, it is also a left H-comodule algebra. In this case, the symmetric separability idempotent of K is given by

$$\zeta_{K} = \sum_{(\lambda_{K})} \lambda_{K}^{(1)} \otimes S(\lambda_{K}^{(2)}) = \sum_{(\lambda_{K})} \lambda_{K}^{(2)} \otimes S(\lambda_{K}^{(1)}) = \sum_{(\lambda_{K})} S^{-1}(\lambda_{K}^{(2)}) \otimes \lambda_{K}^{(1)} = \sum_{(\lambda_{K})} S^{-1}(\lambda_{K}^{(1)}) \otimes \lambda_{K}^{(2)}$$

where  $\lambda_K$  denotes the Haar integral of K. Here, we have used the properties  $S(\lambda_K) = S^{-1}(\lambda_K) = \lambda_K$ , the cocommutativity of  $\lambda_K$ , and the fact that the antipode  $S : K \to K$  acts as an anti-homomorphism

<sup>&</sup>lt;sup>12</sup>Recall that weak Kac algebra is a weak Hopf algebra such that  $S^2 = id$ , see Refs. [127, 128]. All finite dimensional  $C^*$  Hopf algebras are weak Kac algebras.

on the coalgebra. In this case, the boundary vertex operator in Eq. (7.2) becomes:

$$\begin{array}{ccc}
h \\
x \\ y \\
\end{array} \xrightarrow{K} A_{\nu_s}^{K} = \overrightarrow{X}_{\lambda_K^{(1)}} \otimes \overleftarrow{X}_{\lambda_K^{(2)}} \otimes \overleftarrow{X}_{\lambda_K^{(3)}}, \\
A_{\nu_s}^{K} | x, y, h \rangle = |\lambda_K^{(1)} x, y S^{-1}(\lambda_K^{(2)}), h S^{-1}(\lambda_K^{(3)}) \rangle.
\end{array}$$
(7.5)

The cocommutativity of  $\lambda$  implies that the operator  $A_{v_s}^K$  is independent of the specific starting link  $s = (v_s, f)$  chosen at the vertex  $v_s$ . Consequently,  $A_{v_s}^K$  depends solely on the vertex  $v_s$  itself.

For the physical boundary as drawn in Eq. (7.1), the bulk is on the right-hand side of the boundary, and J is a right H-comodule algebra. We define the following operator for  $z \otimes w \in J \otimes J^{\text{op}}$ :

$$x \quad y \\ h \qquad A_{v_p}^{z \otimes w} |x, h, y\rangle = \sum_{[z]} |w^{[0]}x, w^{[1]}h, yz\rangle.$$
(7.6)

The boundary operator is defined similarly as that for symmetry boundary

$$\mathbb{A}_{\nu_p}^J = \mathbb{A}_{\nu_p}^{\zeta_J} = \sum_{\langle \zeta_J \rangle} \mathbb{A}_{\nu_p}^{\zeta_J^{(1)} \otimes \zeta_J^{(2)}}$$
(7.7)

where  $\zeta_J$  is the symmetric separability idempotent of J. The Hamiltonian for the symmetry boundary is

$$\mathbb{H}_{\text{phys}} = -\sum_{\nu_p:\text{phys}} \mathbb{A}^J_{\nu_s}.$$
(7.8)

*Remark* 7.2. When *J* is chosen as a weak Kac subalgebra of *H*, it is also a right *H*-comodule algebra. In this case, the symmetric separability idempotent of *J* is given by  $\zeta_J = \sum_{(\lambda_J)} \lambda_J^{(1)} \otimes S(\lambda_J^{(2)})$  with  $\lambda_J$  denotes the Haar integral of *J*. In this case, the boundary vertex operator becomes:

$$A_{\nu_{p}}^{J} = \overrightarrow{X}_{\lambda_{J}^{(1)}} \otimes \overrightarrow{X}_{\lambda_{J}^{(2)}} \otimes \overleftarrow{X}_{\lambda_{J}^{(3)}},$$

$$A_{\nu_{p}}^{J} | x, h, y \rangle = |\lambda_{J}^{(1)} x, \lambda_{J}^{(2)} h, y S^{-1}(\lambda_{J}^{(3)}) \rangle.$$
(7.9)

The cocommutativity of  $\lambda$  implies that the operator  $A_{v_p}^J$  is independent of the specific starting link  $s = (v_p, f)$  chosen at the vertex  $v_s$ . Consequently,  $A_{v_p}^J$  depends solely on the vertex  $v_s$  itself.

The bulk stabilizers are face operators  $\mathbb{B}_{f}^{H}$ , whose expression is the same as that in Eq. (5.7) but with a different interpretation. For the symmetry boundary K, we need to introduce

$$\overleftarrow{Z}_{\psi}^{K}|y\rangle = \sum_{[y]} \psi(S(y^{[-1]}))|y^{[0]}\rangle, \quad y \in K,$$

where there is no direct pairing between K and  $\hat{H}$ ; instead, we use the pairing between H and  $\hat{H}$ , with  $y^{[-1]} \in H$ . For the physical boundary, we similarly introduce  $Z_{\psi}^{J}$  as

$$\vec{Z}_{\psi}^{J}|x\rangle = \sum_{[x]} \psi(x^{[1]})|x^{[0]}\rangle, \quad x \in J.$$

Using these two edge operators, we can define face operator as

$$g \xrightarrow{x} h \qquad \mathbb{B}_{f}^{\psi} = \overrightarrow{Z}_{\psi^{(1)}} \otimes \overleftarrow{Z}_{\psi^{(2)}}^{K} \otimes \overleftarrow{Z}_{\psi^{(3)}} \otimes \overrightarrow{Z}_{\psi^{(4)}}^{J} \\ \mathbb{B}_{f}^{\psi}|g, y, h, x\rangle = \sum \psi(g^{(2)}S(y^{[-1]})S(h^{(1)})x^{[1]})|g^{(1)}, y^{[0]}, h^{(2)}, x^{[0]}\rangle$$
(7.10)

By setting  $\psi$  as Haar measure  $\Lambda \in \hat{H}$ , we obtain face stabilizer operator  $\mathbb{B}_{f}^{H} = \mathbb{B}_{f}^{\Lambda}$ , the corresponding bulk Hamiltonian reads

$$\mathbb{H}_{bk} = -\sum_{f} \mathbb{B}_{f}^{H}.$$
(7.11)

The total weak Hopf cluster ladder Hamiltonian is of the form

$$\mathbb{H}[H, K, J] = \mathbb{H}_{bk} + \mathbb{H}_{sym} + \mathbb{H}_{phys}$$
(7.12)

which realize the SymTFT for a given weak Hopf gauge symmetry *H*.

The bulk phase is given by  $\operatorname{Rep}(D(H)) \simeq \mathbb{Z}(\operatorname{Rep}(H))$  as we have discussed before. The symmetry boundary is equivalently characterized by the category  $_K \operatorname{Mod}$  of all left K-modules, and it can be checked that this forms a left module category over  $\operatorname{Rep}(H)$ . The symmetry is thus given by the category of end-ofunctor category

$$S = \mathcal{B}_{sym} \simeq \operatorname{Fun}_{\operatorname{Rep}(H)}({}_{K}\operatorname{Mod}, {}_{K}\operatorname{Mod}).$$
 (7.13)

For the physical boundary, since J is a right H-comodule algebra, the boundary is equivalently characterized by Mod<sub>J</sub>, which is a right module category over Rep(H). The physical boundary excitation is given by the category of endofunctor category

$$\mathcal{B}_{phys} = \mathsf{Fun}_{\mathsf{Rep}(H)}(\mathsf{Mod}_J, \mathsf{Mod}_J). \tag{7.14}$$

*Remark* 7.3 (String-net realization). From a macroscopic point of view, the bulk of the ladder is characterized by the multifusion category  $\operatorname{Rep}(H)$ . The symmetry boundary is characterized by the category of left *K*-modules  $\mathcal{M}_K = {}_K \operatorname{Mod}$ , which is a left module category over  $\operatorname{Rep}(H)$ . Similarly, the physical boundary is given by the module category  $\mathcal{M}_J = \operatorname{Mod}_J$  over  $\operatorname{Rep}(H)$ . The category  $\mathcal{C} = \operatorname{Rep}(H)$  can be treated as the bulk input data, while  $\mathcal{M}_K$  and  $\mathcal{M}_J$  serve as two boundary input data. We can also construct a string-net lattice model that realizes the corresponding cluster ladder model.

## 7.2 Ground sate degeneracy

On a ladder lattice with periodic boundary condition, the ground state degeneracy can be calculated using a macroscopic argument. The bulk topological phase is given by the Drinfeld center  $\text{Bulk}(H) = \mathcal{Z}(\text{Rep}(H))$  of Rep(H), which is braided monoidal equivalent to the representation category of charge symmetry, Drinfeld double D(H) of gauge symmetry H. The bulk phase is a unitary modular tensor category, from the anyon condensation point of view, the topological boundary is determined by Lagrangian algebras in Bulk(H). For each boundary labeled by H comodule algebra K, there is a corresponding Lagrangian algebra  $\mathcal{A}_K$  in Bulk(H).

Before discussing the ground state degeneracy of the weak Hopf ladder model, let us first consider a more general case. Take a closed sphere  $S^2$  and punch *n* holes in it. For each hole, assign a topological

boundary condition denoted by  $K_i$ , where i = 1, ..., n. For example, when n = 3 we have the following manifold:



Then for each boundary there is a Lagrangian algebra  $A_{K_i}$ , which can be regarded as a charge in the bulk. The sphere with *n* gapped holes is equivalent to a sphere with *n* charges. In this case the ground state degeneracy is just the number of independent fusion channel between these charges and the vacuum charge **1** of the bulk phase:

$$GSD = \dim \operatorname{Hom}(\mathbf{1}, \mathcal{A}_{K_1} \otimes \cdots \otimes \mathcal{A}_{K_n}). \tag{7.15}$$

If we replace the sphere with a more complicated g-genus surface and punch n holes, the ground state degeneracy can also be derived from the Moore-Seiberg formula [142, 143]. Consider an orientable g-genus surface  $\mathbb{M}_{g,n}$  with n holes, the anyons for there holes are still  $\mathcal{A}_{K_1}, \dots, \mathcal{A}_{K_k}$ . This surface can be decomposed into surfaces of pants, which are then glued together. It is important to note that this decomposition is generally not unique, but the associativity of anyon fusion ensures that the final ground state degeneracy remains consistent:

$$\operatorname{GSD}[\mathbb{M}_{g,n}, \mathcal{A}_{K_1}, \cdots, \mathcal{A}_{K_k}] = \sum_{Y \in \operatorname{Irr}(\operatorname{Rep}(D(H)))} \left( \prod_{j=1}^k S_{\mathcal{A}_{K_j}, Y} \right) S_{\mathbf{1}, Y}^{2-2g-n},$$
(7.16)

where *S* is the S-matrix for the bulk phase.

The weak Hopf ladder model corresponds to the case where the manifold is a cylinder, which can be thought of as a sphere with two punctures. In this model, the boundary conditions are labeled as follows: the symmetry boundary is associated with  $A_K$ , and the physical boundary is associated with  $A_J$ . By applying the Eq. (7.15) or Eq. (7.16), one can determine the ground state degeneracy for this configuration.

#### 7.3 Solving the model via tensor network

The cluster ladder model can be solved using the weak Hopf tensor network introduced in Section 4. There will be four types of local tensors we need to construct the cluster ladder ground state based on a tensor network construction:

(1) For symmetry boundary edge, the corresponding local tensor is the comultiplication of Haar integral  $\lambda_K$  of weak Kac subalgebra  $K \leq H$ :

$$\Delta(\lambda_K) = \begin{matrix} \lambda_K^{(2)} \\ \vdots \\ \lambda_K^{(1)} \\ \lambda_K^{(1)} \end{matrix} := \begin{matrix} \lambda_K^{(1)} & \lambda_K^{(2)} \\ \vdots \\ \vdots \\ \lambda_K^{(1)} \end{matrix} .$$
(7.17)

The leg of the tensor that will be contracted in the tensor network is  $\lambda_{K}^{(2)}$ .

(2) For physical boundary edge, the corresponding local tensor is the comultiplication of Haar integral  $\lambda_J$  of weak Kac subalgebra  $J \leq H$ :

$$\Delta(\lambda_J) = \begin{array}{c} \lambda_J^{(1)} \\ \lambda_J \\ \\ \\ \lambda_I^{(2)} \end{array} := \begin{array}{c} \lambda_J^{(1)} \\ \lambda_J^{(2)} \\ \\ \lambda_J^{(2)} \end{array} .$$
(7.18)

Notice that we have put  $\lambda_J^{(1)}$  at the top; the leg of the tensor that will be contracted in the tensor network is  $\lambda_J^{(2)}$ .

(3) For the bulk edge, the local tensor corresponds to  $\Delta_2(\lambda)$ :

$$\Delta_2(\lambda) = \underbrace{\lambda^{(3)}}_{\lambda^{(1)}} \stackrel{\lambda}{\longrightarrow} \lambda^{(2)} := \underbrace{\lambda^{(1)}}_{\lambda^{(2)}} \stackrel{\lambda^{(2)}}{\longrightarrow} \frac{\lambda^{(3)}}{\lambda^{(3)}}.$$
(7.19)

The legs of the tensor that will be contracted in the tensor network are  $\lambda^{(2)}, \lambda^{(3)}$ .

(4) For face, a local tensor corresponds to the Haar measure  $\Lambda \in \hat{H}$  is needed to glue the edge tensors together:

$$(\mathrm{id}\otimes\mathrm{id}\otimes\hat{S}\otimes\hat{S})\circ\hat{\Delta}(\Lambda) = \hat{S}(\Lambda^{(4)}) \stackrel{\bullet}{\bullet} \stackrel{\Lambda}{\wedge} _{\Lambda^{(2)}} = \bigwedge_{\Lambda^{(1)}}^{\Lambda^{(2)}} \hat{S}(\Lambda^{(3)}) \hat{S}(\Lambda^{(4)}) .$$
(7.20)

The antipode  $\hat{S}$  is introduced because the direction of the lattice is chosen as in Eq. (6.1). If a different lattice configuration is chosen, the antipode may be placed on different legs of the tensor.

**Theorem 5.** The cluster ladder state  $|\Psi\rangle_{\text{cluster}}$  is defined as the following tensor network state:



where we have drawn the free leg (corresponding to physical degrees of freedom) in red. The cluster state  $|\Psi\rangle_{\text{cluster}}$  is the ground state of the cluster ladder model defined in Eq. (7.12), namely,

$$\mathbb{A}_{\nu_s}^K |\Psi\rangle_{\text{cluster}} = \mathbb{A}_{\nu_p}^J |\Psi\rangle_{\text{cluster}} = |\Psi\rangle_{\text{cluster}} = \mathbb{B}_f^H |\Psi\rangle_{\text{cluster}}, \tag{7.22}$$

for all  $v_s, v_p, f$ .

*Proof.* This can be proved using the techniques developed in Ref. [52, Section 7] and the property of pairing,  $\lambda^2 = \lambda$ ,  $\Lambda^2 = \Lambda$ , along with the cyclic property of the comultiplication of the Haar integral and Haar measure.

#### 8 Discussion

In this paper, we propose a general framework for (1 + 1)D lattice models exhibiting non-invertible symmetries. We demonstrate that these models can be analyzed using weak Hopf algebras. The weak Hopf cluster state model and its generalizations are discussed in detail, and we show that the underlying symmetry takes the form  $H \times \hat{H}$ . Thus, dual symmetry plays a crucial role. In the group case, non-invertibility primarily arises due to the existence of dual symmetry. For the general Hopf and weak Hopf cases, non-invertibility emerges from both components of the  $H \times \hat{H}$  symmetry.

Despite the progress made, several important open problems remain. A key question is the classification of all (1 + 1)D phases with weak Hopf symmetry. We conjecture that they can be classified using weak Hopf cohomology theory. Another challenge is extending these results to higher dimensions. For 2D and beyond, additional structures must be introduced to fully characterize the symmetry. Based on Tannaka–Krein duality, there should exist an algebraic structure corresponding to the fusion *n*-category. Identifying this structure and constructing the corresponding lattice model will be a central problem. We plan to address these questions in future work.

#### Acknowledgments

I would like to extend my sincere gratitude to Dagomir Kaszlikowski for his support. I also thank Liang Kong and Sheng Tan for discussions on SymTFT during my visit at BIMSA. This work is supported by the National Research Foundation in Singapore, the A\*STAR under its CQT Bridging Grant and CQT-Return of PIs EOM YR1-10 Funding.

## References

- F. D. M. Haldane, "Nonlinear field theory of large-spin Heisenberg antiferromagnets: Semiclassically quantized solitons of the one-dimensional easy-axis néel state," Phys. Rev. Lett. 50, 1153 (1983).
- [2] F. D. M. Haldane, "Continuum dynamics of the 1-d Heisenberg antiferromagnet: Identification with the o (3) nonlinear sigma model," Physics letters a 93, 464 (1983).
- [3] I. Affleck, T. Kennedy, E. H. Lieb, and H. Tasaki, "Rigorous results on valence-bond ground states in antiferromagnets," Phys. Rev. Lett. 59, 799 (1987).
- [4] Z.-C. Gu and X.-G. Wen, "Tensor-entanglement-filtering renormalization approach and symmetry-protected topological order," Phys. Rev. B 80, 155131 (2009), arXiv:0903.1069 [cond-mat.str-el].
- [5] X. Chen, Z.-C. Gu, and X.-G. Wen, "Classification of gapped symmetric phases in one-dimensional spin systems," Phys. Rev. B 83, 035107 (2011), arXiv:1008.3745 [cond-mat.str-el].
- [6] F. Pollmann, E. Berg, A. M. Turner, and M. Oshikawa, "Symmetry protection of topological phases in one-dimensional quantum spin systems," Phys. Rev. B 85, 075125 (2012), arXiv:0909.4059 [cond-mat.str-el].

- [7] X. Chen, Z.-X. Liu, and X.-G. Wen, "Two-dimensional symmetry-protected topological orders and their protected gapless edge excitations," Phys. Rev. B 84, 235141 (2011), arXiv:1106.4752 [cond-mat.str-el].
- [8] F. Pollmann and A. M. Turner, "Detection of symmetry-protected topological phases in one dimension," Phys. Rev. B 86, 125441 (2012), arXiv:1204.0704 [cond-mat.str-el].
- [9] N. Schuch, D. Pérez-García, and I. Cirac, "Classifying quantum phases using matrix product states and projected entangled pair states," Phys. Rev. B 84, 165139 (2011), arXiv:1010.3732 [cond-mat.str-el].
- [10] X. Chen, Z.-C. Gu, Z.-X. Liu, and X.-G. Wen, "Symmetry-protected topological orders in interacting bosonic systems," Science 338, 1604 (2012), arXiv:1301.0861 [cond-mat.str-el].
- [11] B. Zeng, X. Chen, D.-L. Zhou, and X.-G. Wen, "Quantum information meets quantum matter-from quantum entanglement to topological phase in many-body systems," arXiv preprint arXiv:1508.02595 (2015).
- [12] X.-G. Wen, "Colloquium: Zoo of quantum-topological phases of matter," Rev. Mod. Phys. 89, 041004 (2017), arXiv:1610.03911 [cond-mat.str-el].
- [13] X.-G. Wen, "Classifying gauge anomalies through symmetry-protected trivial orders and classifying gravitational anomalies through topological orders," Phys. Rev. D 88, 045013 (2013), arXiv:1303.1803 [hep-th].
- [14] X. Chen, Z.-C. Gu, Z.-X. Liu, and X.-G. Wen, "Symmetry protected topological orders and the group cohomology of their symmetry group," Phys. Rev. B 87, 155114 (2013), arXiv:1106.4772 [cond-mat.str-el]
- [15] A. Vishwanath and T. Senthil, "Physics of three-dimensional bosonic topological insulators: Surface-deconfined criticality and quantized magnetoelectric effect," Phys. Rev. X 3, 011016 (2013), arXiv:1209.3058 [cond-mat.str-el].
- [16] C. Wang and T. Senthil, "Boson topological insulators: A window into highly entangled quantum phases," Phys. Rev. B 87, 235122 (2013), arXiv:1302.6234 [cond-mat.str-el].
- [17] F. J. Burnell, X. Chen, L. Fidkowski, and A. Vishwanath, "Exactly soluble model of a three-dimensional symmetry-protected topological phase of bosons with surface topological order," Phys. Rev. B 90, 245122 (2014), arXiv:1302.7072 [cond-mat.str-el].
- [18] A. Kapustin, "Symmetry protected topological phases, anomalies, and cobordisms: Beyond group cohomology," (2014), arXiv:1403.1467 [cond-mat.str-el].
- [19] A. Kapustin, "Bosonic topological insulators and paramagnets: a view from cobordisms," (2014), arXiv:1404.6659 [cond-mat.str-el].
- [20] D. S. Freed and M. J. Hopkins, "Reflection positivity and invertible topological phases," Geometry & Topology 25, 1165 (2021), arXiv:1604.06527 [hep-th].
- [21] K. Yonekura, "On the cobordism classification of symmetry protected topological phases," Communications in Mathematical Physics 368, 1121 (2019), arXiv:1803.10796 [hep-th].
- [22] A. Kitaev, "Periodic table for topological insulators and superconductors," in *AIP conference proceedings*, Vol. 1134 (American Institute of Physics, 2009) pp. 22–30, arXiv:0901.2686 [cond-mat.mes-hall].
- [23] S. Ryu, A. P. Schnyder, A. Furusaki, and A. W. Ludwig, "Topological insulators and superconductors: tenfold way and dimensional hierarchy," New Journal of Physics 12, 065010 (2010), arXiv:0912.2157 [cond-mat.mes-hall].
- [24] Z.-C. Gu and X.-G. Wen, "Symmetry-protected topological orders for interacting fermions: Fermionic

topological nonlinear  $\sigma$  models and a special group supercohomology theory," Phys. Rev. B **90**, 115141 (2014), arXiv:1201.2648 [cond-mat.str-el].

- [25] A. Kapustin, R. Thorngren, A. Turzillo, and Z. Wang, "Fermionic symmetry protected topological phases and cobordisms," Journal of High Energy Physics 2015, 1 (2015), arXiv:1406.7329 [cond-mat.str-el].
- [26] D. Gaiotto, A. Kapustin, N. Seiberg, and B. Willett, "Generalized global symmetries," Journal of High Energy Physics 2015, 1 (2015), arXiv:1412.5148 [hep-th].
- [27] A. Kapustin and R. Thorngren, "Higher symmetry and gapped phases of gauge theories," Algebra, Geometry, and Physics in the 21st Century: Kontsevich Festschrift, 177 (2017).
- [28] P. R. Gomes, "An introduction to higher-form symmetries," SciPost Physics Lecture Notes, 074 (2023), arXiv:2303.01817 [hep-th].
- [29] L. Bhardwaj, L. E. Bottini, L. Fraser-Taliente, L. Gladden, D. S. Gould, A. Platschorre, and H. Tillim, "Lectures on generalized symmetries," Physics Reports **1051**, 1 (2024), arXiv:2307.07547 [hep-th].
- [30] C. Cordova, T. T. Dumitrescu, K. Intriligator, and S.-H. Shao, "Snowmass white paper: Generalized symmetries in quantum field theory and beyond," (2022), arXiv:2205.09545 [hep-th].
- [31] T. D. Brennan and S. Hong, "Introduction to generalized global symmetries in QFT and particle physics," (2023), arXiv:2306.00912 [hep-ph].
- [32] J. McGreevy, "Generalized symmetries in condensed matter," Annual Review of Condensed Matter Physics 14, 57 (2023), arXiv:2204.03045 [cond-mat.str-el].
- [33] R. Luo, Q.-R. Wang, and Y.-N. Wang, "Lecture notes on generalized symmetries and applications," Physics Reports 1065, 1 (2024), arXiv:2307.09215 [hep-th].
- [34] S.-H. Shao, "What's done cannot be undone: TASI lectures on non-invertible symmetries," (2024), arXiv:2308.00747 [hep-th].
- [35] S. Schäfer-Nameki, "ICTP lecture on (non-)invertible generalized symmetries," Physics Reports 1063, 1 (2024), arXiv:2305.18296 [hep-th].
- [36] C. Delcamp and A. Tiwari, "Higher categorical symmetries and gauging in two-dimensional spin systems," SciPost Physics 16, 110 (2024), arXiv:2301.01259 [hep-th].
- [37] J. Fröhlich, J. Fuchs, I. Runkel, and C. Schweigert, "Kramers-Wannier duality from conformal defects," Phys. Rev. Lett. 93, 070601 (2004), arXiv:cond-mat/0404051 [cond-mat.stat-mech].
- [38] J. Fuchs, M. R. Gaberdiel, I. Runkel, and C. Schweigert, "Topological defects for the free boson CFT," Journal of Physics A: Mathematical and Theoretical **40**, 11403 (2007), arXiv:0705.3129 [hep-th].
- [39] J. Fröhlich, J. Fuchs, I. Runkel, and C. Schweigert, "Defect lines, dualities and generalised orbifolds," in XVIth International Congress On Mathematical Physics: (With DVD-ROM) (World Scientific, 2010) pp. 608–613, arXiv:0909.5013 [math-ph].
- [40] L. Bhardwaj and Y. Tachikawa, "On finite symmetries and their gauging in two dimensions," Journal of High Energy Physics 2018, 1 (2018), arXiv:1704.02330 [hep-th].
- [41] C.-M. Chang, Y.-H. Lin, S.-H. Shao, Y. Wang, and X. Yin, "Topological defect lines and renormalization group flows in two dimensions," Journal of High Energy Physics 2019, 1 (2019), arXiv:1802.04445 [hep-th].
- [42] R. Thorngren and Y. Wang, "Fusion category symmetry. part i. anomaly in-flow and gapped phases," Journal of High Energy Physics 2024, 1 (2024), arXiv:1912.02817 [hep-th].

- [43] R. Thorngren and Y. Wang, "Fusion category symmetry ii: Categoriosities at c = 1 and beyond," (2021), arXiv:2106.12577 [hep-th].
- [44] Z. Komargodski, K. Ohmori, K. Roumpedakis, and S. Seifnashri, "Symmetries and strings of adjoint QCD2," Journal of High Energy Physics 2021, 1 (2021), arXiv:2008.07567 [hep-th].
- [45] K. Inamura, "On lattice models of gapped phases with fusion category symmetries," Journal of High Energy Physics 2022, 1 (2022), arXiv:2110.12882 [cond-mat.str-el].
- [46] K. Inamura, "Fermionization of fusion category symmetries in 1+1 dimensions," Journal of High Energy Physics 2023, 1 (2023), arXiv:2206.13159 [cond-mat.str-el].
- [47] L. Kong, T. Lan, X.-G. Wen, Z.-H. Zhang, and H. Zheng, "Algebraic higher symmetry and categorical symmetry: A holographic and entanglement view of symmetry," Phys. Rev. Res. 2, 043086 (2020), arXiv:2005.14178 [cond-mat.str-el].
- [48] A. F. Bais, B. J. Schroers, and J. K. Slingerland, "Hopf symmetry breaking and confinement in (2+1)-dimensional gauge theory," Journal of High Energy Physics 2003, 068 (2003), arXiv:hep-th/0205114 [hep-th].
- [49] O. Buerschaper, J. M. Mombelli, M. Christandl, and M. Aguado, "A hierarchy of topological tensor network states," Journal of Mathematical Physics 54, 012201 (2013), arXiv:1007.5283 [cond-mat.str-el].
- [50] B. Yan, P. Chen, and S. Cui, "Ribbon operators in the generalized Kitaev quantum double model based on Hopf algebras," Journal of Physics A: Mathematical and Theoretical (2022), arXiv:2105.08202 [cond-mat.str-el].
- [51] Z. Jia, D. Kaszlikowski, and S. Tan, "Boundary and domain wall theories of 2d generalized quantum double model," Journal of High Energy Physics **2023**, 1 (2023), arXiv:2207.03970 [quant-ph].
- [52] Z. Jia, S. Tan, D. Kaszlikowski, and L. Chang, "On weak Hopf symmetry and weak Hopf quantum double model," Communications in Mathematical Physics 402, 3045 (2023), arXiv:2302.08131 [hep-th].
- [53] Z. Jia, S. Tan, and D. Kaszlikowski, "Weak Hopf symmetry and tube algebra of the generalized multifusion string-net model," Journal of High Enegy Physics **07**, 207 (2024), arXiv:2403.04446 [hep-th].
- [54] Z. Jia, "Generalized cluster states from hopf algebras: non-invertible symmetry and hopf tensor network representation," Journal of High Energy Physics **2024**, 147 (2024), arXiv:2405.09277 [quant-ph].
- [55] K. Inamura, "Topological field theories and symmetry protected topological phases with fusion category symmetries," Journal of High Energy Physics **2021**, 1 (2021), arXiv:2103.15588 [cond-mat.str-el].
- [56] S.-J. Huang and M. Cheng, "Topological holography, quantum criticality, and boundary states," (2023), arXiv:2310.16878 [cond-mat.str-el].
- [57] L. Bhardwaj, L. E. Bottini, S. Schafer-Nameki, and A. Tiwari, "Lattice models for phases and transitions with non-invertible symmetries," (2024), arXiv:2405.05964 [cond-mat.str-el].
- [58] D. S. Freed, G. W. Moore, and C. Teleman, "Topological symmetry in quantum field theory," (2024), arXiv:2209.07471 [hep-th].
- [59] D. Gaiotto and J. Kulp, "Orbifold groupoids," Journal of High Energy Physics 2021, 1 (2021), arXiv:2008.05960 [hep-th].
- [60] L. Bhardwaj and S. Schafer-Nameki, "Generalized charges, part ii: Non-invertible symmetries and the symmetry TFT," (2023), arXiv:2305.17159 [hep-th].
- [61] F. Apruzzi, F. Bonetti, I. García Etxebarria, S. S. Hosseini, and S. Schäfer-Nameki, "Symmetry TFTs from string theory," Communications in mathematical physics **402**, 895 (2023), arXiv:2112.02092 [hep-th].

- [62] L. Bhardwaj, L. E. Bottini, D. Pajer, and S. Schafer-Nameki, "Gapped phases with non-invertible symmetries: (1+1)d," (2024), arXiv:2310.03784 [hep-th].
- [63] C. Zhang and C. Córdova, "Anomalies of (1 + 1)-dimensional categorical symmetries," Phys. Rev. B 110, 035155 (2024).
- [64] W. Ji and X.-G. Wen, "Categorical symmetry and noninvertible anomaly in symmetry-breaking and topological phase transitions," Phys. Rev. Res. 2, 033417 (2020).
- [65] A. Chatterjee, W. Ji, and X.-G. Wen, "Emergent generalized symmetry and maximal symmetry-topological-order," (2024), arXiv:2212.14432 [cond-mat.str-el].
- [66] L. Bhardwaj, L. E. Bottini, D. Pajer, and S. Schafer-Nameki, "The club sandwich: Gapless phases and phase transitions with non-invertible symmetries," (2024), arXiv:2312.17322 [hep-th].
- [67] L. Bhardwaj, D. Pajer, S. Schafer-Nameki, and A. Warman, "Hasse diagrams for gapless spt and SSB phases with non-invertible symmetries," (2024), arXiv:2403.00905 [cond-mat.str-el].
- [68] L. Kong and X.-G. Wen, "Braided fusion categories, gravitational anomalies, and the mathematical framework for topological orders in any dimensions," arXiv preprint arXiv:1405.5858 (2014).
- [69] L. Kong, X.-G. Wen, and H. Zheng, "Boundary-bulk relation in topological orders," Nuclear Physics B 922, 62 (2017), arXiv:1702.00673 [cond-mat.str-el].
- [70] T. Johnson-Freyd, "On the classification of topological orders," Communications in Mathematical Physics , 1 (2022), arXiv:2003.06663 [math.CT].
- [71] B. Zeng, X. Chen, D.-L. Zhou, and X.-G. Wen, Quantum Information Meets Quantum Matter–From Quantum Entanglement to Topological Phase in Many-Body Systems (Springer, 2019) arXiv:1508.02595 [cond-mat.str-el].
- [72] B. Swingle and J. McGreevy, "Renormalization group constructions of topological quantum liquids and beyond," Phys. Rev. B 93, 045127 (2016), arXiv:1407.8203 [cond-mat.str-el].
- [73] P.-S. Hsin and Z. Wang, "On topology of the moduli space of gapped Hamiltonians for topological phases," Journal of Mathematical Physics 64, 041901 (2023), arXiv:2211.16535 [cond-mat.str-el].
- [74] D. Aasen, R. S. Mong, and P. Fendley, "Topological defects on the lattice: I. The Ising model," Journal of Physics A: Mathematical and Theoretical 49, 354001 (2016), arXiv:1601.07185 [cond-mat.stat-mech].
- [75] D. Aasen, P. Fendley, and R. S. Mong, "Topological defects on the lattice: dualities and degeneracies," arXiv preprint arXiv:2008.08598 (2020).
- [76] K. Inamura and K. Ohmori, "Fusion surface models: 2+1d lattice models from fusion 2-categories," SciPost Phys. 16, 143 (2024), arXiv:2305.05774 [cond-mat.str-el].
- [77] L. Eck and P. Fendley, "Generalizations of kitaev's honeycomb model from braided fusion categories," (2024), arXiv:2408.04006 [cond-mat.str-el].
- [78] C. Fechisin, N. Tantivasadakarn, and V. V. Albert, "Non-invertible symmetry-protected topological order in a group-based cluster state," (2023), arXiv:2312.09272 [cond-mat.str-el].
- [79] A. Feiguin, S. Trebst, A. W. W. Ludwig, M. Troyer, A. Kitaev, Z. Wang, and M. H. Freedman, "Interacting anyons in topological quantum liquids: The golden chain," Phys. Rev. Lett. 98, 160409 (2007), arXiv:cond-mat/0612341 [cond-mat.str-el].
- [80] S. Trebst, M. Troyer, Z. Wang, and A. W. Ludwig, "A short introduction to fibonacci anyon models," Progress of Theoretical Physics Supplement 176, 384 (2008).

- [81] M. Buican and A. Gromov, "Anyonic chains, topological defects, and conformal field theory," Communications in Mathematical Physics 356, 1017 (2017), arXiv:1701.02800 [hep-th].
- [82] L. Lootens, C. Delcamp, G. Ortiz, and F. Verstraete, "Dualities in one-dimensional quantum lattice models: Symmetric hamiltonians and matrix product operator intertwiners," PRX Quantum 4, 020357 (2023), arXiv:2112.09091 [quant-ph].
- [83] L. Lootens, C. Delcamp, and F. Verstraete, "Dualities in one-dimensional quantum lattice models: Topological sectors," PRX Quantum 5, 010338 (2024), arXiv:2211.03777 [quant-ph].
- [84] N. Seiberg, S. Seifnashri, and S.-H. Shao, "Non-invertible symmetries and LSM-type constraints on a tensor product Hilbert space," SciPost Phys. 16, 154 (2024), arXiv:2401.12281 [cond-mat.str-el].
- [85] S. Seifnashri and S.-H. Shao, "Cluster state as a noninvertible symmetry-protected topological phase," Phys. Rev. Lett. 133, 116601 (2024), arXiv:2404.01369 [cond-mat.str-el].
- [86] K. Inamura and S. Ohyama, "1+1d SPT phases with fusion category symmetry: interface modes and non-abelian thouless pump," (2024), arXiv:2408.15960 [cond-mat.str-el].
- [87] C. Cordova, D. B. Costa, and P.-S. Hsin, "Non-invertible symmetries in finite group gauge theory," (2024), arXiv:2407.07964 [cond-mat.str-el].
- [88] K. Szlachányi, "Finite quantum groupoids and inclusions of finite type," in Mathematical physics in mathematics and physics: quantum and operator algebraic aspects, Fields Institute Communications, Vol. 30 (American Mathematical Soc., 2001) p. 393–407, arXiv:math/0011036 [math.QA].
- [89] V. Ostrik, "Module categories, weak Hopf algebras and modular invariants," Transformation Groups 8, 177 (2003), arXiv:math/0111139 [math.QA].
- [90] P. Etingof, D. Nikshych, and V. Ostrik, "On fusion categories," Annals of Mathematics , 581 (2005), arXiv:math/0203060 [math.QA].
- [91] W. Son, L. Amico, and V. Vedral, "Topological order in 1d cluster state protected by symmetry," Quantum Information Processing **11**, 1961 (2012), arXiv:1111.7173 [quant-ph].
- [92] C. G. Brell, "Generalized cluster states based on finite groups," New Journal of Physics 17, 023029 (2015), arXiv:1408.6237 [quant-ph].
- [93] A. Kitaev, "Fault-tolerant quantum computation by anyons," Annals of Physics **303**, 2 (2003), arXiv:quant-ph/9707021 [quant-ph].
- [94] V. V. Albert, D. Aasen, W. Xu, W. Ji, J. Alicea, and J. Preskill, "Spin chains, defects, and quantum wires for the quantum-double edge," (2021), arXiv:2111.12096 [cond-mat.str-el].
- [95] S. Beigi, P. W. Shor, and D. Whalen, "The quantum double model with boundary: Condensations and symmetries," Communications in Mathematical Physics **306**, 663 (2011), arXiv:1006.5479 [quant-ph].
- [96] I. Cong, M. Cheng, and Z. Wang, "Hamiltonian and algebraic theories of gapped boundaries in topological phases of matter," Communications in Mathematical Physics 355, 645 (2017), arXiv:1707.04564 [cond-mat.str-el].
- [97] H. Bombin and M. A. Martin-Delgado, "Family of non-Abelian Kitaev models on a lattice: Topological condensation and confinement," Phys. Rev. B 78, 115421 (2008), arXiv:0712.0190 [cond-mat.str-el].
- [98] N. Andruskiewitsch and J. M. Mombelli, "On module categories over finite-dimensional Hopf algebras," Journal of Algebra 314, 383 (2007), arXiv:math/0608781 [math.QA].
- [99] S. X. Cui, S.-M. Hong, and Z. Wang, "Universal quantum computation with weakly integral anyons," Quantum Information Processing 14, 2687 (2015), arXiv:1401.7096 [quant-ph].

- [100] B. Bakalov and A. A. Kirillov, Lectures on tensor categories and modular functors, Vol. 21 (American Mathematical Soc., 2001).
- [101] Z. Wang, Topological quantum computation, CBMS No. 112 (American Mathematical Soc., 2010).
- [102] P. Etingof, S. Gelaki, D. Nikshych, and V. Ostrik, *Tensor categories*, Vol. 205 (American Mathematical Soc., 2016) pp. xvi+343.
- [103] C. Meusburger, "Kitaev lattice models as a Hopf algebra gauge theory," Communications in Mathematical Physics 353, 413 (2017), arXiv:1607.01144 [math.QA].
- [104] F. Girelli, P. K. Osei, and A. Osumanu, "Semidual Kitaev lattice model and tensor network representation," Journal of High Energy Physics 2021, 1 (2021), arXiv:1709.00522 [math.QA].
- [105] S. Majid, Foundations of quantum group theory (Cambridge university press, Cambridge, 2000) pp. x+607.
- [106] E. Abe, *Hopf algebras*, Cambridge Tracts in Mathematics, Vol. 74 (Cambridge University Press, 2004) pp. xii+284.
- [107] S. Montgomery, Hopf algebras and their actions on rings, 82 (American Mathematical Soc., 1993).
- [108] C. Kassel, Quantum groups, Graduate Texts in Mathematics, Vol. 155 (Springer-Verlag, New York, 1995) pp. xii+531.
- [109] M. d. W. Propitius and F. A. Bais, "Discrete gauge theories," (1995), arXiv:hep-th/9511201 [hep-th].
- [110] F. Alexander Bais, P. van Driel, and M. de Wild Propitius, "Quantum symmetries in discrete gauge theories," Physics Letters B 280, 63 (1992), arXiv:hep-th/9203046 [hep-th].
- [111] F. Bais, N. Muller, and B. Schroers, "Quantum group symmetry and particle scattering in (2+1)-dimensional quantum gravity," Nuclear Physics B 640, 3 (2002), arXiv:hep-th/0205021 [hep-th].
- [112] C. Delcamp, B. Dittrich, and A. Riello, "Fusion basis for lattice gauge theory and loop quantum gravity," Journal of High Energy Physics 2017, 1 (2017), arXiv:1607.08881 [hep-th].
- [113] J. Fuchs, Affine Lie algebras and quantum groups: An Introduction, with applications in conformal field theory (Cambridge university press, 1995).
- [114] C. Meusburger and D. K. Wise, "Hopf algebra gauge theory on a ribbon graph," Reviews in Mathematical Physics, 2150016 (2021), arXiv:1512.03966 [math.QA].
- [115] J. Slingerland and F. Bais, "Quantum groups and non-Abelian braiding in quantum Hall systems," Nuclear Physics B 612, 229 (2001), arXiv:cond-mat/0104035 [cond-mat.mes-hall].
- [116] O. Buerschaper, M. Christandl, L. Kong, and M. Aguado, "Electric-magnetic duality of lattice systems with topological order," Nuclear Physics B 876, 619 (2013), arXiv:1006.5823 [cond-mat.str-el].
- [117] V. Koppen, "Defects in Kitaev models and bicomodule algebras," arXiv preprint arXiv:2001.10578 (2020).
- [118] T. Voß, Defects and symmetries in Hopf algebra lattice models, Ph.D. thesis, Friedrich-Alexander-Universität Erlangen-Nürnberg (FAU) (2021).
- [119] A. Kitaev and L. Kong, "Models for gapped boundaries and domain walls," Communications in Mathematical Physics 313, 351 (2012), arXiv:1104.5047 [cond-mat.str-el].
- [120] N. Andruskiewitsch and H.-J. Schneider, "On the classification of finite-dimensional pointed hopf algebras," Annals of Mathematics, 375 (2010).
- [121] J. W. Milnor and J. C. Moore, "On the structure of Hopf algebras," Annals of Mathematics 81, 211 (1965).

- [122] P. Cartier, "Groupes algébriques et groupes formels," in *Colloq. Théorie des Groupes Algébriques* (1962) pp. 87–111.
- [123] M. Sweedler, *Hopf Algebras*, Mathematics lecture note series (W.A. Benjamin, 1969).
- [124] S. Dascalescu, C. Nastasescu, and S. Raianu, Hopf algebra: An introduction (CRC Press, 2000).
- [125] G. Böhm, F. Nill, and K. Szlachányi, "Weak Hopf algebras: I. Integral theory and C\*-structure," Journal of Algebra 221, 385 (1999), arXiv:math/9805116 [math.QA].
- [126] D. Nikshych, "Semisimple weak Hopf algebras," Journal of Algebra 275, 639 (2004), arXiv:math/0304098 [math.QA].
- [127] T. Yamanouchi, "Duality for generalized Kac algebras and a characterization of finite groupoid algebras," Journal of Algebra 163, 9 (1994).
- [128] G. Böhm and K. Szlachányi, "Weak Hopf algebras: II. representation theory, dimensions, and the markov trace," Journal of Algebra 233, 156 (2000).
- [129] G. Böhm, "Doi-hopf modules over weak Hopf algebras," Communications in Algebra 28, 4687 (2000), arXiv:math/9905027 [math.QA].
- [130] J. C. Bridgeman, L. Lootens, and F. Verstraete, "Invertible bimodule categories and generalized Schur orthogonality," Communications in Mathematical Physics 402, 2691 (2023), arXiv:2211.01947 [math.QA].
- [131] L. Chang, "Kitaev models based on unitary quantum groupoids," Journal of Mathematical Physics 55, 041703 (2014), arXiv:1309.4181 [math.QA].
- [132] R. Dijkgraaf and E. Witten, "Topological gauge theories and group cohomology," Communications in Mathematical Physics 129, 393 (1990).
- [133] V. G. Drinfel'd, "Quantum groups," Journal of Soviet mathematics 41, 898 (1988).
- [134] S. Majid, "Physics for algebraists: Non-commutative and non-cocommutative Hopf algebras by a bicrossproduct construction," Journal of Algebra 130, 17 (1990).
- [135] S. Majid, "Some remarks on the quantum double," Czechoslovak Journal of Physics 44, 1059 (1994), arXiv:hep-th/9409056 [hep-th].
- [136] L. Kong, "Anyon condensation and tensor categories," Nuclear Physics B 886, 436 (2014), arXiv:1307.8244 [cond-mat.str-el].
- [137] Z. Jia, D. Kaszlikowski, and S. Tan, "Electric-magnetic duality and Z<sub>2</sub> symmetry enriched cyclic Abelian lattice gauge theory," Journal of Physics A: Mathematical and Theoretical 57, 255203 (2024), arXiv:2201.12361 [quant-ph].
- [138] F. J. Burnell, "Anyon condensation and its applications," Annual Review of Condensed Matter Physics 9, 307 (2018), arXiv:1706.04940 [cond-mat.str-el].
- [139] S. Eliëns, "Anyon condensation," MSc Thesis, University of Amsterdam, 2010, available at https://www.cs.vu.nl/ eliens/download/thesis-sebas.pdf (2010).
- [140] M. A. Levin and X.-G. Wen, "String-net condensation: A physical mechanism for topological phases," Phys. Rev. B 71, 045110 (2005), arXiv:cond-mat/0404617 [cond-mat.str-el].
- [141] M. Aguiar, "A note on strongly separable algebras," Bol. Acad. Nac. Cienc. (Córdoba) 65, 51 (2000).
- [142] G. Moore and N. Seiberg, "Classical and quantum conformal field theory," Communications in Mathematical Physics 123, 177 (1989).

[143] A. Ritz-Zwilling, J.-N. Fuchs, S. H. Simon, and J. Vidal, "Topological and nontopological degeneracies in generalized string-net models," (2023), arXiv:2309.00343 [cond-mat.other].