# COMPLEX STRUCTURES ON PRODUCT MANIFOLDS

LEONARDO BILIOTTI AND ALESSANDRO MINUZZO

ABSTRACT. Let  $M_i$ , for i = 1, 2, be a Kähler manifold, and let G be a Lie group acting on  $M_i$  by Kähler isometries. Suppose that the action admits a momentum map  $\mu_i$  and let  $N_i := \mu_i^{-1}(0)$ be a regular level set. When the action of G on  $N_i$  is proper and free, the Meyer-Marsden-Weinstein quotient  $P_i := N_i/G$  is a Kähler manifold and  $\pi_i : N_i \to P_i$  is a principal fiber bundle with base  $P_i$  and characteristic fiber G. In this paper, we define an almost complex structure for the manifold  $N_1 \times N_2$  and give necessary and sufficient conditions for its integrability. In the integrable case, we find explicit holomorphic charts for  $N_1 \times N_2$ . As applications, we consider a non integrable almost-complex structure on the product of two complex Stiefel manifolds and the infinite Calabi-Eckmann manifolds  $\mathbb{S}^{2n+1} \times S(\mathcal{H})$ , for  $n \geq 1$ , where  $S(\mathcal{H})$  denotes the unit sphere of an infinite dimensional Hilbert space  $\mathcal{H}$ .

#### 1. INTRODUCTION

In a classical paper by E. Calabi and B. Eckmann [1] it is shown that the compact manifold  $\mathbb{S}^{2n+1} \times \mathbb{S}^{2m+1}$  admits a complex structure coming from the natural U(1)-action on  $\mathbb{C}^n \times \mathbb{C}^m$ , but does not admit a Kähler structure for  $n \geq 1$ , as an immediate cohomological computation shows. For example, the second de Rham cohomology group  $H^2_{dR}(\mathbb{S}^{2n+1} \times \mathbb{S}^{2m+1}) = \{0\}$ , precisely when  $n \geq 1$  or  $m \geq 1$ .

We generalize the idea of their construction by considering two Kähler manifolds  $(M_i, h_i, J_i, \omega_i)$ , i = 1, 2, where  $h_i$  is the Riemannian metric,  $J_i$  the complex structure,  $\omega_i$  the compatible symplectic structure and letting G be a Lie group acting on each  $M_i$  by Kähler isometries. This means that for all  $g \in G$  we have

$$g^*h_i = h_i, \qquad g_* \circ J_i = J_i \circ g_*, \qquad g^*\omega_i = \omega_i$$

where by  $g^*$  and  $g_*$  we denote respectively the pull-back and the push-farward, associated to the diffeomorphism induced by g. We suppose that each action admits a momentum map  $\mu_i: M_i \to \mathfrak{g}^*$ , where  $\mathfrak{g} := \operatorname{Lie}(G)$ , and let  $N_i$  be a regular level set of it. We further suppose that we can reduce  $N_i$  to its orbit space  $P_i$  by taking the quotients by the isotropy subgroup  $F_i$  of the momentum, relative to the co-adjoint action of G on  $\mathfrak{g}^*$ . This is the Meyer–Marsden–Weinstein quotient, and is known to be a Kähler manifold  $(P_i, \tilde{h}_i, \tilde{J}_i, \tilde{\omega}_i)$ , where  $h_i|_{N_i} = \pi_i^* \tilde{h}_i, \omega_i|_{N_i} = \pi_i^* \tilde{\omega}_i$ and  $\pi_i \circ J_i = \tilde{J}_i \circ \pi_i$ . Moreover, the natural projection  $\pi_i: N_i \to P_i$  is a principal fiber bundle with base  $P_i$  and characteristic fiber  $F_i$ . For the definition of the momentum map and its use in the Meyer–Marsden–Weinstein reduction, we refer to [2].

There is a natural decomposition of the tangent space of  $N_i$  as we now describe. For further details see [3]. The projection defines a distribution inside  $TN_i$ , the so called *vertical distribution*, given by the kernel of its differential. At each point the vertical distribution is isomorphic (not

<sup>2000</sup> Mathematics Subject Classification: Primary 53C55, 57S20

Key words: group action, momentum map, reduction, complex structures.

The first author was partially supported by PRIN 2017 "Real and Complex Manifolds: Topology, Geometry and holomorphic dynamics" and "National Group for Algebraic and Geometric Structures, and their Applications" (GNSAGA - INDAM).

canonically) to the Lie algebra of the isotropy subgroup,  $\mathfrak{f}_i := \operatorname{Lie}(F_i)$ , because we are in the hypothesis that the action of  $F_i$  on  $N_i$  is free. We denote the complementary subspace at  $p_i \in N_i$  with respect to  $T_{p_i}N_i$  as  $H_{p_i}$ , i.e.  $T_{p_i}N_i \simeq H_{p_i} \oplus \mathfrak{f}_i$ . We can therefore define a distribution  $H_i := \bigcup_{p_i \in N_i} H_{p_i} \subset TN_i$ , the horizontal distribution, and we also have that for all  $g \in G$  that  $g_*H_i = H_i$ , because we are considering an isometric action. This tells us that  $H_i \subset TN_i$  is a principal connection for the principal fiber bundle  $\pi_i : N_i \to P_i$ .

principal connection for the principal fiber bundle  $\pi_i : N_i \to P_i$ . In the following, we will consider the case in which  $N_i := \mu_i^{-1}(0)$  is a regular level set of the momentum map  $\mu_i$ , so that  $F_1 \simeq G \simeq F_2$ . Moreover, we will use the splitting

(1) 
$$T_{(p_1,p_2)}(N_1 \times N_2) \simeq (H_{p_1} \oplus \mathfrak{g}) \oplus (H_{p_2} \oplus \mathfrak{g}) .$$

**Definition 1.1.** We define the following endomorphism J of  $T(N_1 \times N_2)$  as

$$J := (J_1 \oplus T_1) \oplus (J_2 \oplus T_2) ,$$

where we have used the splitting (1), and we have set for  $\xi \in \mathfrak{g}$ 

$$T_1(\xi_{N_1}) := \xi_{N_2} \qquad T_2(\xi_{N_2}) := -\xi_{N_1} ,$$

which  $\xi_{N_i}$  the infinitesimal generator of the action of G on  $N_i$ , relative to  $\xi$ .

Our first result is the following.

**Theorem 1.2.** The endomorphism J defines an almost complex structure on  $N_1 \times N_2$ , i.e.  $J^2 = -id_{T(N_1 \times N_2)}$ . Furthermore, J is integrable, i.e. complex, if and only if G is abelian.

By the celebrated Newlander–Nirenberg Theorem [5], the integrability of the almost-complex structure J on  $N_1 \times N_2$  is equivalent to the vanishing of the Nijenhuis tensor associated to J. This tensor is defined as

$$N_J(X,Y) := [JX, JY] - [X,Y] - J[JX,Y] - [X,JY],$$

where X, Y are smooth vector fields on  $N_1 \times N_2$ .

We then study the action by the abelian group  $(\mathbb{C}, +)$  defined as

(2)  $\Psi: \mathbb{C} \times (N_1 \times N_2) \to (N_1 \times N_2), \qquad (a+ib, (x,y)) \mapsto (e^{2\pi i (a+b)} \cdot x, e^{2\pi i (a-b)} \cdot y) ,$ 

where the dot denotes the action on the two factors. We prove the following proposition, telling us that the map  $\Psi$  is holomorphic.

- **Proposition 1.3.** (1) The map  $\Psi_{(x,y)} : \mathbb{C} \to (N_1 \times N_2), \ \Psi_{(x,y)}(a+ib) := \Psi(a+ib, (x,y))$ is holomorphic with respect to the complex structure of  $\mathbb{C}$  and the complex structure J of definition 1.1;
  - (2) The map  $\Psi_{a+ib}: (N_1 \times N_2) \to (N_1 \times N_2), \ \Psi_{a+ib}(x, y) := \Psi(a+ib, (x, y))$  is holomorphic with respect to the complex structure J of definition 1.1.

Let now  $\Lambda := \{a + ib : a, b \in \mathbb{Z}/2\}$  be the lattice of  $\mathbb{C}$  with cell  $[0, \frac{1}{2}] \times [0, \frac{i}{2}]$ . Then  $\mathbb{C}/\Lambda \simeq \mathbb{T}^2 = \mathbb{S}^1 \times \mathbb{S}^1$  is the standard one dimensional complex torus. This is a compact abelian group and the quotient action  $\Psi$  by  $\mathbb{T}^2$  is free. Without using theorem 1.2, we are able to prove the following

**Theorem 1.4.** Given the action (2), the natural projection  $\pi : (N_1 \times N_2) \to (P_1 \times P_2)$  is a principal holomorphic bundle with characteristic fiber  $\mathbb{T}^2$ . Moreover, the construction can be easily generalized to the case of an action of  $\mathbb{C}^n / \Lambda_n = \mathbb{T}^{2n}$ .

### 2. PROOFS OF THE RESULTS

*Proof.* of **Theorem 1.2.** We start by proving that J is an almost complex structure. Consider a basis element for  $T_{(p_1,p_2)}(N_1 \times N_2) \simeq (H_{p_1} \oplus \mathfrak{g}) \bigoplus (H_{p_2} \oplus \mathfrak{g})$  denoted as  $(v_1,\xi_1,v_2,\xi_2)$ , where  $v_i \in H_{p_i}$  and  $\xi_i \in \mathfrak{g}$  are nonzero elements. By definition 1.1, and since  $J_i$  is a complex structure on  $H_{p_i}$ , we have that

$$J^{2}(v_{1},\xi_{1},v_{2},\xi_{2}) = J(J_{1}v_{1},\xi_{2},J_{2}v_{2},-\xi_{1}) = (J_{1}^{2}v_{1},-\xi_{1},J_{2}^{2}v_{2},-\xi_{2}) = -(v_{1},\xi_{1},v_{2},\xi_{2})$$

To prove integrability, we show the vanishing of the Nijenhuis tensor of J, denoted as  $N_J$ . We do this for each case comprised in the splitting (1). If  $X_i$  is a horizontal vector field the integrability follows directly from the integrability of the  $J_i$ . Let  $Y_1 := \xi_{N_1}, Y_2 := \eta_{N_2}$  be vertical vector fields, we have

$$\begin{split} N_J(Y_1,Y_2) &= [JY_1,JY_2] - [Y_1,Y_2] - J[JY_1,Y_2] - J[Y_1,JY_2] = -J[JY_1,Y_2] - J[Y_1,JY_2] = \\ &= -J([\xi_{N_2},\eta_{N_2}] - [\xi_{N_1},\eta_{N_1}]) = -J([\xi,\eta]_{N_1} - [\xi,\eta]_{N_2}) \;. \end{split}$$

Therefore,  $N_J(Y_1, Y_2) = 0$  if and only if G is abelian. By skew-symmetry and linearity we are left to check that  $N_J(X_1, Y_i) = 0$ .

$$N_J(X_1, Y_1) = -[X_1, Y_1] - J[JX_1, Y_1] = -[X_1, Y_1] - J^2[X_1, Y_1] = [X_1, Y_1] - [X_1, Y_1] = 0.$$

Moreover,

$$N_J(X_1, Y_2) = [JX_1, JY_2] - J[X_1, JY_2] = J[X_1, JY_2] - J[X_1, JY_2] = 0$$

In the last two equations we used the fact that for  $X_i$  horizontal and  $Y_i = \xi_{N_i}$  vertical we have

$$[JX_i, Y_i] = [J_iX_i, Y_i] = J_i[X_i, Y_i] = J[X_i, Y_i]$$

Indeed  $J_i \circ g_* = g_* \circ J_i$  is equivalent to  $\mathcal{L}_{\xi_{N_i}} J_i = 0$  for all  $\xi \in \mathfrak{g}$ , and hence

$$Y_i, J_i X_i] = \mathcal{L}_{Y_i}(J_i X_i) = (\mathcal{L}_{Y_i} J_i)(X_i) + J_i(\mathcal{L}_{Y_i} X_i) = J_i(\mathcal{L}_{Y_i} X_i) = J_i[Y_i, X_i].$$

### *Proof.* of **Proposition 1.3.**

(1) Consider the curve  $t \mapsto (a + t + ib) \in \mathbb{C}$ , so that  $\frac{d}{dt}(a + t + ib)|_{t=0} = 1 \in T_{(a+ib)}\mathbb{C}$ . Therefore,

$$\left(\mathrm{d}\Psi_{(x,y)}\right)_{a+ib}(1) = \frac{d}{dt}\Psi_{(x,y)}(a+t+ib)\big|_{t=0} = (1_{N_1}, 1_{N_2}) \; .$$

For the tangent vector i consider the curve  $t \mapsto (a + i(t + b)) \in \mathbb{C}$ , so that  $\frac{d}{dt}(a + i(t + b))|_{t=0} = i \in T_{(a+ib)}\mathbb{C}$ . Now,

$$\left(\mathrm{d}\Psi_{(x,y)}\right)_{a+ib}(i) = \frac{d}{dt}\Psi_{(x,y)}(a+i(t+b))\big|_{t=0} = (i_{N_1}, i_{N_2}) = (1_{N_2}, -1_{N_1}) = J(1_{N_1}, 1_{N_2}) \ .$$

The result follows observing that  $i = i \cdot 1$  is the action of the complex structure of  $\mathbb{C}$  on the vector 1.

(2) Using the decomposition of  $TN_i \simeq H_i \oplus \mathfrak{g} \simeq H_i \oplus \mathbb{R}^2$  and since  $\Psi$  is holomorphic we have  $(\Psi_{a+ib})_* \circ J_i = J_i \circ (\Psi_{a+ib})_*$ . For the vertical vectors we have, since they are invariant under  $(\Psi_{a+ib})_*$ 

$$J((\Psi_{a+ib})_*\xi_{N_1},(\Psi_{a+ib})_*\eta_{N_2})) = ((\Psi_{a+ib})_*\eta_{N_1},-(\Psi_{a+ib})_*\xi_{N_2}) = (\eta_{N_1},-\xi_{N_2}) + ((\Psi_{a+ib})_*\eta_{N_2}) = (\eta_{N_1},-\xi_{N_2}) + ((\Psi_{a+ib})_*\eta_{N_2}) = (\eta_{N_1},-\xi_{N_2}) + ((\Psi_{a+ib})_*\eta_{N_2}) = ((\Psi_{a+ib})_*\eta_{N_2}) + ((\Psi_{a+ib})_*\eta_{N_$$

and also,

$$(\Psi_{a+ib})_* J(\xi_{N_1}, \eta_{N_2}) = (\Psi_{a+ib})_* (\eta_{N_1}, -\xi_{N_2}) = (\eta_{N_1}, -\xi_{N_2}) .$$

Remark 2.1. We observe that, in general, given  $A \in GL(n, \mathbb{Z})$ , the action  $\Psi : \mathbb{C} \times (N_1 \times N_2) \to (N_1 \times N_2)$  defined as  $\Psi(a+ib, (x, y)) := (e^{2\pi i (A_{11}a+A_{21}b)}.x, e^{2\pi i (A_{12}a+A_{22}b)}.y)$  is holomorphic with respect to J, and induces a free holomorphic action of  $\mathbb{C}/\Lambda_p$ , where  $\Lambda_p := \{c+id : c, d \in \mathbb{Z}/p\}$ , and  $p = |\det(A)|$ .

*Proof.* of **Theorem 1.4** We find holomorphic charts of  $N_1 \times N_2$  and the holomorphic trivializations of  $\pi : (N_1 \times N_2) \to (P_1 \times P_2)$  for the case of an  $\mathbb{S}^1$  action.

**Holomorphic charts for**  $N_1 \times N_2$ . Let  $U_1 \subseteq P_1$  and  $U_2 \subseteq P_2$  open neighborhoods of  $x_1 \in U_1$ and  $x_2 \in U_2$  respectively. Then if  $\pi_i : N_i \to P_i$ , there are diffeomorphisms

$$\phi_i: \pi_i^{-1}(U_i) \to U_i \times \mathbb{S}^1$$

Let us denote with  $\Sigma_i := \phi^{-1}(U_i \times \{1\})$ . Then we have

$$T_{(x_1,x_2)}(N_1 \times N_2) \simeq T_{(x_1,x_2)}(\Sigma_1 \times \Sigma_2) \oplus T_{(x_1,x_2)}G.(x_1,x_2)$$
.

Up to shrinking  $\Sigma_i$  we can assume the previous relation to hold everywhere on  $\Sigma_1 \times \Sigma_2$ . Let  $\Phi: \Sigma_1 \times \Sigma_2 \times G \to N_1 \times N_2$  be the map

$$\Phi(s_1, s_2, g) := g(s_1, s_2)$$

This map is holomorphic thanks to proposition 1.3 and a local diffeomorphism.

**Lemma 2.2.** The map  $\Phi$  defined above is injective and hence a biholomorphism.

*Proof.* Suppose by contradiction that  $\Phi$  is not injective. Then pick

$$\{(s_1^n, s_2^n)\}_{n \in \mathbb{N}} \subset \Sigma_1 \times \Sigma_2 \supset \{(\tilde{s}_1^n, \tilde{s}_2^n)\}_{n \in \mathbb{N}}$$

such that  $(s_1^n, s_2^n) \neq (\tilde{s}_1^n, \tilde{s}_2^n)$ ,  $g_n$  for all  $n \in \mathbb{N}$  and

$$(s_1^n, s_2^n) \xrightarrow{n \to \infty} (s_1, s_2) \xleftarrow{n \to \infty} (\tilde{s}_1^n, \tilde{s}_2^n)$$
.

We can then find  $\{g_n\}_{n\in\mathbb{N}}\subset G\supset \{\tilde{g}_n\}_{n\in\mathbb{N}}$  such that

$$g_n.(s_1^n, s_2^n) = \tilde{g}_n.(\tilde{s}_1^n, \tilde{s}_2^n)$$

Therefore if  $\theta_n := \tilde{g}_n^{-1} g_n$  we have

$$\Phi(s_1^n, s_2^n, \theta_n) = \Phi(\tilde{s}_1^n, \tilde{s}_2^n, 1) = (\tilde{s}_1^n, \tilde{s}_2^n)$$

We conclude that  $\Phi(s_1^n, s_2^n, \theta_n) \xrightarrow{n \to \infty} (s_1, s_2)$  and by properness of the action  $\theta_n \xrightarrow{n \to \infty} 1$ . But this means that  $\Phi$  is not injective in a neighborhood of  $(s_1, s_2, 1)$ , which is a contradiction.

**Transition functions.** Let  $\tilde{U}_i \subseteq P_i$  with  $\tilde{U}_i \cap U_i \neq \emptyset$  and diffeomorphism

$$\tilde{\phi}_i : \pi_i^{-1}(\tilde{U}_i) \to \tilde{U}_i \times \mathbb{S}^1$$

and denote  $\tilde{\Sigma}_i := \tilde{\phi}_i^{-1}(\tilde{U}_i \times \{1\})$ . Again we have a biholomorphism with the image  $\tilde{\Phi} : \tilde{\Sigma}_1 \times \tilde{\Sigma}_2 \times G \to N_1 \times N_2$ . Let us use the following notation for simplicity  $\Sigma := \Sigma_1 \times \Sigma_2$ ,  $\tilde{\Sigma} := \tilde{\Sigma}_1 \times \Sigma_2$ ,  $U := U_1 \times U_2$ ,  $\tilde{U} := \tilde{U}_1 \times \tilde{U}_2$ ,

$$\begin{split} \Phi^{-1} &: \Phi(\Sigma \times G) \to \Sigma \times G, \ n \mapsto (\Pi^{-1}(\pi(n)), \Theta(n)) \ , \\ \tilde{\Phi}^{-1} &: \tilde{\Phi}(\tilde{\Sigma} \times G) \to \tilde{\Sigma} \times G, \ n \mapsto (\tilde{\Pi}^{-1}(\pi(n)), \tilde{\Theta}(n)) \ , \end{split}$$

Where  $\Pi := \pi|_{\Sigma}$  and  $\tilde{\Pi} := \pi|_{\tilde{\Sigma}}$  are smooth invertible maps. Then we can define trivializations  $H: U \times G \to \pi^{-1}(U)$  as

$$H(p,g) := \Phi(\Pi^{-1}(p),g)$$
.

Similarly,

$$\tilde{H}(p,g):=\tilde{\Phi}(\tilde{\Pi}^{-1}(p),g)\ .$$
 We have  $H^{-1}(n)=(\pi(n),\Theta(n)).$  Now,

 $(H^{-1} \circ \tilde{H})(p,g) = H^{-1}(\tilde{\Phi}(\tilde{\Pi}^{-1}(p),g) = (\pi(\tilde{\Phi}(\tilde{\Pi}^{-1}(p),g)), \Theta(\tilde{\Phi}(\tilde{\Pi}^{-1}(p),g))) = (p,\Theta(\tilde{\Phi}(\tilde{\Pi}^{-1}(p),g))) .$ Let h(p) such that  $\tilde{\Phi}(\tilde{\Pi}^{-1}(p),e) = h(p).\Phi(\Pi^{-1}(p),e)$ , where e = (1,1). Then, by equivariance,

$$(H^{-1} \circ H)(p,g) = (p,\Theta(\Phi(\Pi^{-1}(p),g))) = (p,\Theta(g.\Phi(\Pi^{-1}(p),e))) = = (p,\Theta(gh(p).\Phi(\Pi^{-1}(p),e))) = (p,\Theta(\Phi(\Pi^{-1}(p),gh(p)))) = (p,gh(p)) .$$

In summary, we have proved that the transition functions of the principal fiber bundle  $\pi$ :  $(N_1 \times N_2) \rightarrow (P_1 \times P_2)$  are holomorphic. Hence  $N_1 \times N_2$  is a complex manifold, since the map  $\Phi$  is holomorphic with respect to J. It also follows that the natural complex structure induced by the complex atlas is in fact J. Moreover, if we realize  $P_1 \times P_2$  as Meyer–Marsden– Weinstein reduction by an action of  $\mathbb{S}^1 \times \overset{k-\text{times}}{\cdot \cdot \cdot \cdot} \times \mathbb{S}^1$ , then the same argument proves that  $\pi : (N_1 \times N_2) \rightarrow (P_1 \times P_2)$  is a  $(\mathbb{T}^2 \times \overset{k-\text{times}}{\cdot \cdot \cdot \cdot} \times \mathbb{T}^2)$ -principal bundle. The transition functions are then holomorphic functions and then, as before  $N_1 \times N_2$  admits a complex atlas and J is precisely the complex structure induced by the atlas.  $\Box$ 

#### 3. Applications

3.1. Complex Stiefel manifolds. Consider the action of the unitary group in complex dimension k, U(k), on the space of linear maps  $\operatorname{Hom}(\mathbb{C}^k, \mathbb{C}^n)$ , for  $n \ge k$ .

$$\Psi: U(k) \times \operatorname{Hom}(\mathbb{C}^k, \mathbb{C}^n) \to \operatorname{Hom}(\mathbb{C}^k, \mathbb{C}^n)$$
,

$$\Psi(u,A) := A \circ u^{\dagger} .$$

This is a proper left action<sup>1</sup>. The action is also isometric with respect to the Hermitian metric metric on  $\operatorname{Hom}(\mathbb{C}^k, \mathbb{C}^n)$  given by

$$h(A, B) := \operatorname{Tr}(A \circ B^{\dagger})$$

The symplectic form associated,  $\omega(A, B) := \mathfrak{Im}(\operatorname{Tr}(A \circ B^{\dagger}))$ , is exact and equal to the differential of the Liouville 1-form  $\theta|_A(B) = \mathfrak{Im}(\operatorname{Tr}(A \circ B^{\dagger}))$ . The infinitesimal generator of the action at  $A \in \operatorname{Hom}(\mathbb{C}^k, \mathbb{C}^n)$  for  $\xi \in \mathfrak{u}(k)$  is given by  $\frac{d}{dt}(A \circ \exp t\xi)|_{t=0} = A \circ \xi$  and then we have

$$\langle \mu(A)|\xi\rangle = \theta_A(A\circ\xi) = \Im\mathfrak{m}(\operatorname{Tr}(A\circ\xi^{\dagger}\circ A^{\dagger})) = -\Im\mathfrak{m}(\operatorname{Tr}(A^{\dagger}\circ A\circ\xi)) = i\operatorname{Tr}(A^{\dagger}\circ A\circ\xi) \ .$$

By duality then,

$$\langle \mu(A)|\xi \rangle = \operatorname{Tr}(\mu(A) \circ \xi^{\dagger}) = -\operatorname{Tr}(\mu(A) \circ \xi) = i\operatorname{Tr}(A^{\dagger} \circ A \circ \xi) ,$$

and hence

$$\mu(A) = i(A^{\dagger} \circ A) \in \mathfrak{u}(k)^{*}$$
 .

This momentum map is  $Ad_{U(k)}$ -equivariant, indeed

$$\mu(A \circ u^{\dagger}) = i(u \circ A^{\dagger} \circ A \circ u^{\dagger}) = u \circ \mu(A) \circ u^{\dagger} .$$

<sup>&</sup>lt;sup>1</sup>The action is proper since U(k) is compact, and is a left action since  $\Psi(u_2u_1, A) = A \circ (u_2u_1)^{\dagger} = A \circ u_1^{\dagger} \circ u_2^{\dagger} = \Psi(u_2, \Psi(u_1, A)).$ 

Notice that the level set of  $\mu$  relative to the element  $i\mathbb{I} \in \mathfrak{u}(k)^*$  is  $\mu^{-1}(i\mathbb{I}) = \{A \in \operatorname{Hom}(\mathbb{C}^k, \mathbb{C}^n) : A^{\dagger} \circ A = \mathbb{I}\}$  is equal to the set of unitary k-frames in  $\mathbb{C}^n$ , the so called Stiefel manifold  $V_k(\mathbb{C}^n)$ , and that the isotropy subgroup of any element, which is the whole U(k), acts freely on it. We also know that the quotient space is given by the complex Grassmannian  $\operatorname{Gr}_{\mathbb{C}}(k,n)$ , which is Kähler. Thanks to our construction, the product  $V_k(\mathbb{C}^n) \times V_k(\mathbb{C}^n)$  carries a non integrable almost complex structure. We can also consider a p-dimensional torus in U(k),  $p \leq k$ , and obtain a free action by  $\mathbb{C}^p/\Lambda$  on  $V_k(\mathbb{C}^n) \times V_k(\mathbb{C}^n)$ , which is then foliated by tori. In particular, if p = 1, the orbits are J-holomorphic curves (see [4]).

3.2. Infinite Calabi-Eckmann manifolds. Consider an Hilbert space  $\mathcal{H}$  and its unit sphere  $S(\mathcal{H})$ . Theorem 1.4 tells us that the product  $\mathbb{S}^{2n+1} \times S(\mathcal{H})$  is holomorphic with complex structure J and then we have the following.

**Theorem 3.1.** There cannot be a *J*-invariant symplectic structure  $\omega$  on  $\mathbb{S}^{2n+1} \times S(\mathcal{H})$  such that  $\omega(\cdot, J \cdot)$  is Riemannian. In other words  $\mathbb{S}^{2n+1} \times S(\mathcal{H})$  cannot be Kähler with respect to the complex structure *J*.

*Proof.* If such a symplectic form  $\omega$  does exist, the complex submanifolds  $j : \mathbb{S}^{2n+1} \times \mathbb{S}^{2m+1} \hookrightarrow \mathbb{S}^{2n+1} \times S(\mathcal{H})$  would be Kähler with respect to  $j^*\omega$ . This is a contradiction since  $\mathbb{S}^{2n+1} \times \mathbb{S}^{2m+1}$  cannot be Kähler, e.g. as we have already recalled, for cohomological reasons, when  $n \geq 1$  or  $m \geq 1$ .

## References

- CALABI E., ECKMANN B. A Class of Compact, Complex Manifolds Which are not Algebraic, Annals of Mathematics 58 (1953), 494.
- [2] CANNAS DA SILVA A., Lectures on Symplectic Geometry, Lecture Notes in Math. 1764, Springer-Verlag, Berlin, 2001.
- [3] KOBAYASHI, S., NOMIZU, N., Foundations of Differential Geometry Vol. I & II, Interscience Publisher, J. Wiles & Sons, New-York, (1963).
- [4] MCDUFF, D., SALAMON, D. Introduction to Symplectic Topology OXFORD UNIVERSITY PRESS, OX-FORD, (2017).
- [5] NEWLANDER, A., NIRENBERG, L., Complex analytic coordinates in almost complex manifolds, ANN. OF MATH. 65 (1957), 391-404.

DIPARTIMENTO DI SCIENZE MATEMATICHE, FISICHE E INFORMATICHE, UNIVERSITÀ DI PARMA (ITALY) Email address: leonardo.biliotti@unipr.it Email address: alessandro.minuzzo@unipr.it