Rainbow Arborescence Conjecture

Kristóf Bérczi^a

Tamás Király^b

Yutaro Yamaguchi^c

Yu Yokoi^d

Abstract

The famous Ryser–Brualdi–Stein conjecture asserts that every $n \times n$ Latin square contains a partial transversal of size n-1. Since its appearance, the conjecture has attracted significant interest, leading to several generalizations. One of the most notable extensions is to matroid intersection given by Aharoni, Kotlar, and Ziv, focusing on the existence of a common independent transversal of the common independent sets of two matroids. In this paper, we study a special case of this setting, the *Rainbow Arborescence Conjecture*, which states that any graph on n vertices formed by the union of n-1 spanning arborescences contains an arborescence using exactly one arc from each. We prove that the computational problem of testing the existence of such an arborescence with a fixed root is NP-complete, verify the conjecture in several cases, and explore relaxed versions of the problem.

Keywords: Latin squares, Matroid intersection, Rainbow arborescences, Transversals

1 Introduction

A Latin square is an $n \times n$ matrix where each row/column contains all distinct numbers, 1 through n. A partial transversal is a subset of entries in the matrix that are distinct numbers in distinct rows and columns. Ryser, Brualdi, and Stein [5,16] conjectured that every $n \times n$ Latin square contains a partial transversal of size n - 1. Woolbright [18] and independently Brouwer, de Vries, and Wieringa [4] showed that a partial transversal of size $n - \sqrt{n}$ always exists. Shor and Hatami [11,15] improved this bound to $n - O(\log^2 n)$. Cameron and Wanless [6] further relaxed the problem, showing that every Latin square contains a subset of entries such that they are in distinct rows and columns and no number appears more than twice.

A matroidal generalization was proposed by Chappell [7] and later by Kotlar and Ziv [12]. A matroidal Latin square is an $n \times n$ matrix whose elements are members of the ground set of a matroid, where each row/column forms an independent set. An independent partial transversal is an independent set comprising entries from distinct rows and columns. As a matroidal analogue of the Ryser-Brualdi-Stein conjecture, Kotlar and Ziv [12] conjectured that every $n \times n$ matroidal Latin square has an independent partial transversal of size n - 1, and proved a lower bound $\lfloor 2n/3 \rfloor$ for the maximum size of such a set.

Aharoni, Kotlar, and Ziv [2,3] further generalized the conjecture to matroid intersection as follows. If M and N are matroids on the same ground set, then for any n pairwise disjoint common independent sets of size n, there exists a common independent set of size n - 1 intersecting each of the n sets in at most one element. They also established a lower bound $n - \sqrt{n}$ for the size of such a set, extending the results

^aMTA-ELTE Matroid Optimization Research Group and HUN-REN-ELTE Egerváry Research Group, Department of Operations Research, Eötvös Loránd University, Budapest, Hungary. Email: kristof.berczi@ttk.elte.hu.

^bHUN-REN-ELTE Egerváry Research Group, Department of Operations Research, Eötvös Loránd University, Budapest, Hungary. Email: tamas.kiraly@ttk.elte.hu.

^cDepartment of Information and Physical Sciences, Graduate School of Information Science and Technology, Osaka University, Osaka, Japan. Email: yutaro.yamaguchi@ist.osaka-u.ac.jp.

^dDepartment of Mathematical and Computing Science, School of Computing, Institute of Science Tokyo, Tokyo, Japan. Email: yokoi@comp.isct.ac.jp.

of [4,18], and improving the lower bound in [12]. Additionally, Kotlar and Ziv [13] extended the work of Drisko [9] and Chappell [7] and showed that for any 2n - 1 disjoint common independent sets of size n, such a common independent set of size n always exists. For a comprehensive overview of results on Latin squares, we recommend the survey by Wanless [17].

It should be noted that the problem proposed in [2,3] can also be rephrased as a statement about the maximum common independent set of three matroids. While finding a maximum common independent set in the intersection of three matroids is oracle-hard, the question naturally arises whether reasonable lower bounds can be obtained using existing approximation methods. Unfortunately, neither the technique of Aharoni and Berger [1] nor the iterative refinement approach by Linhares, Olver, Swamy, and Zenklusen [14] suffices to establish the strong lower bound proposed in the conjecture.

Our results. We study the arborescence analogue of the Aharoni–Kotlar–Ziv conjecture and explore stronger results. An *arborescence* is a connected digraph in which exactly one vertex has no incoming arc and every other vertex has exactly one incoming arc. Similar to bipartite matchings (corresponding to Latin squares), arborescences (on the same vertex set) are well-known as one of the most fundamental graph structures that are naturally represented as matroid intersection.

Throughout the paper, G is a digraph on the vertex set V of size $n (\geq 2)$ that is obtained as the disjoint union of $k (\geq 1)$ arborescences A_1, A_2, \ldots, A_k on V, where each index $i \in [k]$ is called a *color*. Note that G may have parallel arcs of different colors. A subgraph of G is often identified with its arc set. A subgraph B of G is called *rainbow* if $|B \cap A_i| \leq 1$ for all $i \in [k]$. An arborescence in G is said to be *spanning* if its vertex set is V. Yokoi [8, Open Problem: Rainbow Arborescence Problem] proposed the following conjecture.

Conjecture 1.1. Let G be the disjoint union of n-1 spanning arborescences $A_1, A_2, \ldots, A_{n-1}$. Then, G has a rainbow spanning arborescence B.

We here remark on a small but possibly significant difference from the conjectures discussed above. In Conjecture 1.1, since |B| = n - 1, all the colors are used exactly once; that is, B is a transversal rather than a (proper) partial transversal. On the other hand, the -1 term in the previous conjectures cannot be dismissed, even in the case of bipartite matchings. Indeed, it is not difficult to construct a bipartite graph that is the union of n perfect matchings, yet there is no matching of size n that uses exactly one edge from each of them (e.g., consider the complete bipartite graph $K_{2,2}$). Somewhat surprisingly, no similar obstacle is known for the problem when specialized to arborescences.

Each of the following conjectures is equivalent to Conjecture 1.1, and may be easier when $k \neq n-1$. Note that the conjectures are completely the same when k = n - 1, and Conjecture 1.1 implies Conjectures 1.2 and 1.3 for $k \neq n-1$ by easy observations.

Conjecture 1.2. Let $k \ge n-1$, and G be the disjoint union of k spanning arborescences A_1, A_2, \ldots, A_k . Then, G has a rainbow spanning arborescence B.

Conjecture 1.3. Let $k \le n-1$, and G be the disjoint union of n-1 spanning arborescences $A_1, A_2, \ldots, A_{n-1}$. Then, G has a rainbow arborescence B of size k.

Our contribution has three main components. In Section 2, we show that deciding whether a rainbow arborescence with a fixed root exists is NP-hard. In Section 3, we verify Conjecture 1.1 in various settings. Finally, relaxations of the problem are considered in Section 4.

Notation and Terminology For an arborescence, the unique vertex having no incoming arc is called the *root*; we also say that an arborescence is *rooted at* r if its root is r. For each non-root vertex, its *parent* is the tail of the unique incoming arc. A vertex is called a *leaf* if it has no outgoing arc. A *path* is an arborescence having exactly one leaf, and a *star* is one in which every non-root vertex is a leaf. A *branching* is a digraph such that every connected component is an arborescence.

For a digraph S and a vertex v, we denote by $\delta_S^-(v)$ and $\delta_S^+(v)$ the sets of arcs in S incoming to v and outgoing from v, respectively.

2 Complexity with fixed root

Observe that if all colors have the same root r, then a rainbow spanning arborescence rooted at r exists and one can find one by a simple greedy approach, while there exists no spanning arborescence rooted at a different vertex, since r has no incoming arc in G. In the general setting, a natural approach to find a rainbow spanning arborescence would be to fix a root, start with an empty arc set, and augment it by adding an arc of each color one-by-one. It, however, turns out that finding a rainbow arborescence rooted at a fixed vertex is hard. More precisely, let ROOTED-RA denote the problem where the input is a graph G that is the disjoint union of n-1 spanning arborescences $A_1, A_2, \ldots, A_{n-1}$ (with arbitrary roots) along with a target root vertex r, and the goal is to decide if G contains a rainbow spanning arborescence rooted at r.

Theorem 2.1. ROOTED-RA is NP-complete even if exactly two vertices are roots of the input arborescences.

Proof. The problem is clearly in NP. We prove hardness by reduction from the 3-dimensional matching problem, denoted by 3DM. In 3DM, we are given a tripartite 3-uniform hypergraph $\mathcal{H} = (X, Y, Z; \mathcal{E})$ where |X| = |Y| = |Z| = p, $|\mathcal{E}| = q$, and $\mathcal{E} \subseteq X \times Y \times Z$, and the goal is to decide whether \mathcal{H} has a perfect matching. This problem is known to be NP-complete [10, (SP2)].

Let $\mathcal{H} = (X, Y, Z; \mathcal{E})$ be an instance of 3DM. We construct an instance of ROOTED-RA as follows. Let D_{st} denote the digraph obtained by taking two vertices s and t, and adding a directed s-t path $P_H = (e_{H,x}, e_{H,y}, e_{H,z}, f_H)$ of length 4 for each hyperedge H = (x, y, z), where all the paths are openly disjoint; thus, D_{st} consists of 3q + 2 vertices and 4q arcs. We take p copies of D_{st} and apply the notational convention that in the jth copy, we add j as an upper index to the notation. We connect these p digraphs D_{st}^j by identifying t^j with s^{j+1} for $j \in [p-1]$. We also add an extra vertex t.

We assign color c_v to all the arcs of the form $e_{H,v}^j$ for each $v \in X \cup Y \cup Z$. Furthermore, we assign a unique color $c_{H,j}$ to every arc of the form f_H^j . Note that at this point, every color class forms a branching. We extend these branchings to spanning arborescences as follows. For each $v \in X \cup Y \cup Z$, we add an arc of color c_v from t_p to every other vertex with no incoming arc of color c_v . That is, the root of the arborescence of color c_v is t_p . For each $H \in \mathcal{E}$ and $j \in [p]$, we add an arc of color $c_{H,j}$ from t to every other vertex with no incoming arc of color $c_{H,j}$. That is, the root of the arborescence of color $c_{H,j}$ is t. Finally, we add 2pq - 2p + 1 stars rooted at t_p of distinct colors. Let G = (V, E) be the digraph thus obtained. It is easy to check from definition that |V| = 3pq + p + 2 and E is the union of 3pq + p + 1 spanning arborescences. We claim that G has a rainbow spanning arborescence rooted at s_1 if and only if \mathcal{H} has a perfect matching.

Assume first that G has a rainbow spanning arborescence rooted at s_1 and \mathcal{H} has no perfect matching. By construction, the arborescence must contain a rainbow s_1-t_p path. This path is the union of s_j-t_j paths for $j \in [p]$, each corresponding to some hyperedge $H_j \in \mathcal{E}$. Since the s_1-t_p path is rainbow, the hyperedges H_1, H_2, \ldots, H_p must be disjoint and hence is a perfect matching in \mathcal{H} , a contradiction.

Now suppose that \mathcal{H} has a perfect matching, consisting of hyperedges $H_1, H_2, \ldots, H_p \in \mathcal{E}$. The union of the paths $P_{H_j}^j$ is a rainbow s_1-t_p path. Using one of the stars rooted at t_p , we can extend this path to a rainbow s_1-t path. Finally, using the remaining stars and the colors $c_{H,j}$ rooted at t, we can finish the construction to obtain a rainbow spanning arborescence rooted at s_1 .

3 Verifying special cases

As a step toward verifying Conjecture 1.1, we show that the statement is true under some further assumptions: when each A_i is a path or a star (Section 3.2), when there are at most two vertices that are roots of multiple arborescences (Section 3.3), and when the underlying graph is a tree (Section 3.4).

3.1 Preparations

For the proofs, we need some technical lemmas. We first observe that we can get rid of stars among the arborescences A_i . However, since the reduction may change the structure of other arborescences, we should be careful when we use this lemma in an argument based on induction.

Lemma 3.1. Let G be the disjoint union of n-1 spanning arborescences $A_1, A_2, \ldots, A_{n-1}$ on V, and suppose that A_{n-1} is a star rooted at r. For each $i \in [n-2]$, we define an arborescence A'_i on V - r as follows.

- If the root of A_i is r, then choose any vertex $r' \in V r$ with $(r, r') \in \delta^+_{A_i}(r)$ and let $A'_i \coloneqq \{(r', v) \mid (r, v) \in \delta^+_{A_i}(r), v \neq r'\} \cup (A_i \setminus \delta^+_{A_i}(r)).$
- Otherwise, let r' be the parent of r on A_i and $A'_i := \{ (r', v) \mid (r, v) \in \delta^+_{A_i}(r) \} \cup (A_i \setminus (\delta^+_{A_i}(r) \cup \{ (r', r) \})).$

In particular, if A_i is a star, then this is true for A'_i . If there exists a rainbow arborescence on V - r with respect to $(A'_1, A'_2, \ldots, A'_{n-2})$, then this is true on V with respect to $(A_1, A_2, \ldots, A_{n-1})$.

Proof. Let B' be a rainbow arborescence on V - r with respect to $(A'_1, A'_2, \ldots, A'_{n-2})$, and define a subgraph \hat{B} as follows. For each $i \in [n-2]$, let $\{e'_i\} = B' \cap A'_i$. If $e'_i \in A_i$ as it is, then \hat{B} contains the arc e'_i . Otherwise, by definition, $e'_i = (r', v)$ for some $v \in V - r$, where r' is the root of A'_i , and it corresponds to two arcs in A_i as follows.

- If the root of A_i is r, then e'_i corresponds to $(r, r'), (r, v) \in A_i$.
- Otherwise, e'_i corresponds to $(r', r), (r, v) \in A_i$.

In either case, \hat{B} contains both of the two corresponding arcs.

Suppose that |B| = n - 2. In this case, B = B', and we obtain a desired arborescence on V just by adding an arc in the star A_{n-1} from r to the root of B'.

Otherwise, we have $|\hat{B}| \ge n-1$. Then, $|\hat{B} \cap A_i| = 2$ for at least one index $i \in [n-2]$, and $\hat{B} \cap A_i$ contains an arc (r, v) such that $e'_i = (r', v)$. Let B be the subgraph of G obtained from \hat{B} by removing the other arc for each of such indices i((r, r') or (r', r) depending on the root of A_i). Then, B is disjoint only from A_{n-1} and consists of two arborescences, one of whose root is r. Thus, by adding an arc in the star A_{n-1} from rto the other root, we obtain a rainbow spanning arborescence.

For each vertex $v \in V$, let $\rho(v) := \{i \in [n-1] \mid \text{the root of } A_i \text{ is } v\}$. A vertex v is called a *multi-root* if $|\rho(v)| \ge 2$, and a *non-root* if $|\rho(v)| = 0$. We then observe the following sufficient condition for the existence of a rainbow spanning arborescence.

Lemma 3.2. Let G be the disjoint union of n-1 spanning arborescences $A_1, A_2, \ldots, A_{n-1}$ on V. If G has a rainbow arborescence \tilde{B} on $\tilde{V} \subseteq V$ such that \tilde{V} contains all the multi-roots, then G has a rainbow spanning arborescence.

Proof. We show that we can augment (\tilde{B}, \tilde{V}) to (\tilde{B}', \tilde{V}') with $\tilde{V}' = \tilde{V} + v$ for some $v \in V \setminus \tilde{V}$ unless $\tilde{V} = V$. Suppose that there exists a color $i \in [n-1]$ with $|\tilde{B} \cap A_i| = 0$ and $r_i \in \tilde{V}$, where r_i is the root of A_i . Then, A_i has an arc e = (u, v) with $u \in \tilde{V}$ and $v \notin \tilde{V}$, and hence we just define $\tilde{B}' \coloneqq \tilde{B} + e$ and $\tilde{V} \coloneqq \tilde{V} + v$. Thus, we may assume that $|\tilde{B} \cap A_i| = 1$ for any color i with $r_i \in \tilde{V}$. This implies $\sum_{v \in \tilde{V}} \rho(v) \leq |\tilde{B}| = |\tilde{V}| - 1$. Since $\sum_{v \in V} \rho(v) = |V| - 1$ and $\rho(v) \leq 1$ for every $v \in V \setminus \tilde{V}$, we also have $\sum_{v \in \tilde{V}} \rho(v) \geq |\tilde{V}| - 1$. Thus, the equality holds, and $|\tilde{B} \cap A_i| = 1$ if and only if $r_i \in \tilde{V}$.

Let \tilde{r} be the root of \tilde{B} . Pick an unused color i, and let $e = (v, \tilde{r})$ be the unique arc in $\delta_{A_i}^-(\tilde{r})$. If $v \notin \tilde{V}$, then we define $\tilde{B}' := \tilde{B} + e$ and $\tilde{V}' := \tilde{V} + v$. Otherwise, let f = (u, v) be the unique arc in $\delta_{\tilde{B}}^-(v)$ and we first update \tilde{B} to $\tilde{B} + e - f$. After this update, \tilde{B} is still a rainbow arborescence on \tilde{V} , whose root is v, and satisfies $|\tilde{B} \cap A_j| = 0$ and $r_j \in \tilde{V}$ for the color j with $f \in A_j$. Thus, A_j has an arc e' = (u', v') with $u' \in \tilde{V}$ and $v' \notin \tilde{V}$, and we can define $\tilde{B}' := \tilde{B} + e'$ and $\tilde{V}' := \tilde{V} + v'$.

3.2 Each arborescence being a path or a star

We first verify the conjecture when each arborescence is a path.

Theorem 3.3. Conjecture 1.1 is true when each A_i is a path.

Proof. Since each A_i is a path, it has the unique leaf. We first claim the existence of a pair of a subgraph S and a vertex $r \in V$ satisfying the following three conditions:

- (1) $|S \cap A_i| = 1$ for each $i \in [n-1]$,
- (2) $|\delta_S^-(v)| = 1$ for each $v \in V \setminus \{r\}$ and $|\delta_S^-(r)| = 0$, and
- (3) for each $i \in [n-1]$, the unique arc in $S \cap A_i$ lies between r and the leaf on the path A_i .

Indeed, such (S, r) can be found by the following procedure. Initially, let S be empty. Then, for each i = 1, 2, ..., n-1, do the following. Let v_i be the vertex on A_i closest to the leaf among those that have no incoming arcs in S and add to S the arc of A_i entering v_i . Then, the size of the resulting S is n-1. Let r be the unique vertex that has no incoming arc in S. We see that these S and r satisfy conditions (1)-(3).

Among the pairs (S, r) satisfying conditions (1)–(3), take the one minimizing $\sum_{i \in [n-1]} |A_i(r, S)|$, where $|A_i(r, S)|$ is the length of the path on A_i from r to the head of the unique arc in $S \cap A_i$. We show that S is an arborescence, which completes the proof by condition (1).

Suppose, to the contrary, that S is not so. Since we have conditions (1) and (2), S is not connected, and at least one connected component must be a cycle. That is, there is a vertex set $\{u^1, u^2, \ldots, u^k\}$ such that $(u^j, u^{j+1}) \in S$ for $j \in [k]$, where we let $u^{k+1} = u^1$. Note that r is not contained in this cycle as we have $|\delta_S^-(r)| = 0$. Let S' be a subgraph obtained from S by replacing each arc (u^j, u^{j+1}) in the cycle with the arc of the same color that enters u^j . Such an arc does exist because $u^j \neq r$ and we have condition (3). This S' satisfies conditions (1)–(3) with the same r while $\sum_{i \in [n-1]} |A_i(r, S')| < \sum_{i \in [n-1]} |A_i(r, S)|$, which contradicts the choice of S.

By combining Theorem 3.3 with Lemma 3.1, we get the following corollary.

Corollary 3.4. Conjecture 1.1 is true when each A_i is either a path or a star.

Proof. We prove this by induction on n. If all A_i are paths, just apply Theorem 3.3. Otherwise, without loss of generality, we may assume that A_{n-1} is a star. Then, by Lemma 3.1, we can reduce the instance by removing the star and its root. By the induction hypothesis, it suffices to show that the reduction given in Lemma 3.1 preserves the condition that an arborescence is a path.

Suppose that A_i $(i \in [n-2])$ is a path. If the root of A_i is r, then $A'_i = A_i \setminus \{(r, v)\}$, where v is the next vertex of r on the path A_i ; thus, this is indeed a path. Otherwise, $A'_i = A_i \setminus \{(r', r)\}$ if r is the leaf, and $A'_i = (A_i \setminus \{(r', r), (r, v)\}) \cup \{(r', v)\}$ otherwise, where r' is the parent of r and v is the next vertex of r on the path A_i ; thus, this is also a path.

3.3 At most two multi-roots

Our next result shows that the conjecture also holds if the number of multi-roots is small. An *in-arborescence* is a connected digraph which becomes an arborescence by flipping the direction of all the arcs, i.e., in which exactly one vertex has no outgoing arc and every other vertex has exactly one outgoing arc.

Theorem 3.5. Conjecture 1.1 is true when there exist at most two multi-roots.

Proof. When there exists at most one multi-root, the claim follows from Lemma 3.2 by taking $\tilde{B} = \emptyset$ and $\tilde{V} = \{r\}$ for a vertex $r \in V$ with $\rho(r) = \max_{v \in V} \rho(v)$. The remaining case is when there exist exactly two multi-roots, say x_1 and x_2 . By Lemma 3.2 again, it suffices to show that G has a rainbow arborescence \tilde{B} on some \tilde{V} with $\{x_1, x_2\} \subseteq \tilde{V}$.

In order to obtain such (\tilde{B}, \tilde{V}) , we show that G has a pair of arc-disjoint in-arborescences \tilde{B}_1 and \tilde{B}_2 on \tilde{V}_1 and \tilde{V}_2 , respectively, satisfying the following three conditions:

- (1) \tilde{B}_j is rooted at x_j for each $j \in [2]$,
- (2) $\tilde{B}_1 \cup \tilde{B}_2$ is rainbow, and
- (3) $\tilde{V}_1 \cap \tilde{V}_2 \neq \emptyset$.

It is easy to see that $\tilde{B}_1 \cup \tilde{B}_2$ includes a desired rainbow arborescence \tilde{B} as it contains arc-disjoint paths from each $\tilde{r} \in \tilde{V}_1 \cap \tilde{V}_2$ to x_1 and x_2 .

Observe that if $|\tilde{B}_1 \cup \tilde{B}_2| = n - 1$ in addition to conditions (1)–(2), then condition (3) is automatically satisfied by the pigeonhole principle (as $|\tilde{V}_1| + |\tilde{V}_2| = |\tilde{B}_1| + |\tilde{B}_2| + 2 = n + 1 > |V|$). Also, $\tilde{B}_1 = \tilde{B}_2 = \emptyset$, $\tilde{V}_1 = \{x_1\}$, and $\tilde{V}_2 = \{x_2\}$ satisfy conditions (1)–(2). Thus, it suffices to show that we can augment at least one of the pairs $(\tilde{B}_1, \tilde{V}_1)$ and $(\tilde{B}_2, \tilde{V}_2)$ satisfying conditions (1)–(2) as long as $\tilde{V}_1 \cap \tilde{V}_2 = \emptyset$.

Suppose that $V_1 \cap V_2 = \emptyset$. Then, there exists a color $i \in [n-1]$ such that $(B_1 \cup B_2) \cap A_i = \emptyset$. As $\tilde{V}_1 \cap \tilde{V}_2 = \emptyset$, at least one \tilde{V}_j does not contain the root of A_i . Thus, A_i contains some arc $e = (u, v) \in A_i$ with $u \notin \tilde{V}_j$ and $v \in \tilde{V}_j$, and we can indeed augment $(\tilde{B}_j, \tilde{V}_j)$ to $(\tilde{B}_j + e, \tilde{V}_j + u)$.

As a corollary, we get that the conjecture is true for graphs on a small number of vertices.

Corollary 3.6. Conjecture 1.1 is true when $n \leq 6$.

Proof. When $n \le 6$, there are at most five colors. Thus, there are at most two multi-roots, and Theorem 3.5 completes the proof.

3.4 Underlying graph is a tree

Finally, we show that the conjecture is true if the underlying graph is a tree, i.e., each arborescence is obtained by orienting the edges of the same tree.

Theorem 3.7. Conjecture 1.1 is true when the underlying graph of G is a tree.

Proof. By the assumption of the theorem, each A_i is obtained as an orientation of some tree on V. We prove the statement by induction on n. The base case is n = 2, and it is trivial. Suppose that $n \ge 3$. Let $v \in V$ be a leaf of A_{n-1} and $e = (u, v) \in A_{n-1}$ be its incoming arc. For each $i \in [n-2]$, let A'_i be the arborescence on V - v obtained from A_i by removing the unique arc incident to v. By induction hypothesis, there exists an arborescence B' on V - v such that $|B' \cap A'_i| = 1$ for all $i \in [n-2]$. Thus, we obtain a desired rainbow arborescence B = B' + e on V.

4 Relaxations

In this section, we consider relaxations of Conjecture 1.1 in two directions. First, we relax the number of input arborescences (Section 4.1), and then we relax the size of the rainbow arborescence to be found (Section 4.2).

4.1 Relaxing the number of input arborescences

The result of Kotlar and Ziv [13], when specialized to arborescences, implies that if G is the disjoint union of 2n-3 spanning arborescences, then there exists a rainbow spanning arborescence. We improve this bound to 2n-4 using a simpler proof.

Theorem 4.1. Conjecture 1.2 is true when $k \ge 2n-3$. Furthermore, if $n \ge 3$, Conjecture 1.2 is true when $k \ge 2n-4$.

Proof. We prove the first claim by induction on n. The base case is n = 2, and it is trivial. Suppose that $n \geq 3$. Let \tilde{B} be an inclusion-wise maximal rainbow arborescence in G and \tilde{V} be its vertex set; that is, \tilde{B} is a rainbow arborescence on \tilde{V} such that for any color $i \in [k]$ with $\tilde{B} \cap A_i = \emptyset$ and any arc $(u, v) \in A_i$, we have $u \notin \tilde{V}$ or $v \in \tilde{V}$. If $\tilde{V} = V$, then we have obtained a desired arborescence. Otherwise, for every color $i \in [k]$ with $\tilde{B} \cap A_i = \emptyset$, the root of A_i must be in $V' \coloneqq V \setminus \tilde{V}$, and the restriction A'_i of A_i to $V \setminus \tilde{V}$ is also an arborescence. The number of such colors is

$$k' \coloneqq k - (|V| - 1) \ge 2n - |V| - 2.$$

As $\tilde{V} \neq \emptyset$, we obtain $k' \ge 2(n - |\tilde{V}|) - 1 > 2|V'| - 3$. Thus, by induction hypothesis if $|V'| \ge 2$ and trivially if |V'| = 1, there exists a rainbow arborescence B' on V' only using these k' colors. Also, since $k' \ge n - 1$ as $|\tilde{V}| \le n - 1$, starting with an arborescence B' on V', we can obtain a rainbow arborescence B on V by adding outgoing arcs of unused ones among the k' colors one-by-one (as they have roots in V').

Finally, we verify the second claim. Suppose that $n \ge 3$ and k = 2n - 4 (note that $k \ge n - 1$). If we fail the induction proof above, then we have k' < n - 1, which implies $|\tilde{V}| = n - 1$ and k' = n - 2. In this case, let r be the unique vertex in V', and then we can construct a rainbow arborescence B' rooted at r with |B'| = k' = n - 2 by adding outgoing arcs of the k' colors one-by-one. If we can add one more arc of an unused color, then we obtain a desired arborescence. Otherwise, all the k - k' = n - 2 unused colors have the same root, which is a unique vertex not spanned by B'. Thus, G has at most two multi-roots, and Theorem 3.5 completes the proof.

As a corollary, we get a counterpart of the result of Cameron and Wanless [6], but for arborescences.

Corollary 4.2. Let $n \ge 3$ and G be the disjoint union of n-2 spanning arborescences $A_1, A_2, \ldots, A_{n-2}$ on V. Then, G has a spanning arborescence using at most two arcs from each A_i .

Proof. For each arborescence A_i , make its copy A'_i . Then, by Theorem 4.1 (with k = 2n - 4), there exists a spanning arborescence B' on V using at most one arc from each A_i and A'_i . Thus, one can obtain a desired spanning arborescence B from B' by replacing each copy arc from A'_i with its original in A_i .

4.2 Relaxing the size of output arborescence

The result of Kotlar and Ziv [13] implies that if G is the disjoint union of n-1 spanning arborescences, then it has a rainbow branching of size $\lfloor n/2 \rfloor$. We prove a strengthening of this observation by showing that the branching in fact can be chosen to be an arborescence.

Theorem 4.3. Conjecture 1.3 is true when $k \leq \lfloor n/2 \rfloor$.

Proof. By Corollary 3.6, we assume that $n \ge 7$. As in the proof of Theorem 4.1, let \tilde{B} be an inclusion-wise maximal rainbow arborescence in G and \tilde{V} be its vertex set. If $|\tilde{B}| \ge \lfloor n/2 \rfloor$, then we have obtained a desired arborescence. We show that otherwise we can augment (\tilde{B}, \tilde{V}) by exchanging several arcs.

Let \tilde{r} be the root of \tilde{B} , and J be the set of unused colors. Since \tilde{B} is inclusion-wise maximal, for every color $i \in J$, the root of A_i is not in \tilde{V} , and the restriction of A_i to $V \setminus \tilde{V}$ is also an arborescence. Thus, each A_i includes a path P_i satisfying the following three conditions:

- the root is not in \tilde{V} ,
- all other vertices are in \tilde{V} , and
- the leaf is \tilde{r} .

Claim 4.4. The disjoint union $\bigcup_{i \in I} P_i$ contains a rainbow path P satisfying the same three conditions.

Proof. It suffices to show that $\bigcup_{i \in J} P_i$ has a rainbow in-arborescence B' on $V' \not\subseteq \tilde{V}$ rooted at \tilde{r} , which must contain a desired rainbow path P.

We prove this constructively. Initially, let $B' = \emptyset$ and $V' = \{\tilde{r}\}$. For each color $i \in J$ (in any order), we do the following update. If $V' \not\subseteq \tilde{V}$ at some point, then we have obtained a desired in-arborescence B'. Otherwise, P_i has an arc e = (u, v) such that $u \notin V'$ and $v \in V'$ (as P_i starts outside of $\tilde{V} \supseteq V'$ and ends at $\tilde{r} \in V'$); we add e to B' and u to V'. By the assumptions, we have

$$|\tilde{V}| = |\tilde{B}| + 1 \le \lfloor n/2 \rfloor \le n - 1 - (\lfloor n/2 \rfloor - 1) \le n - 1 - |\tilde{B}| = |J|.$$

This implies $|V'| = |J| + 1 > |\tilde{V}|$ at the end, and hence $V' \not\subseteq \tilde{V}$.

Let P be a path obtained by Claim 4.4, and let Q be the set of arcs $e \in \tilde{B}$ such that the head of e is in P. Then, |Q| = |P| - 1 as \tilde{r} is the root of the arborescence \tilde{B} , and we can indeed augment \tilde{B} to $(\tilde{B} \cup P) \setminus Q$. This completes the proof of the theorem.

Acknowledgments. We are grateful to Shin-ichi Tanigawa for the initial discussion on the problem, which contains a key observation leading to Theorem 3.5.

Yutaro Yamaguchi was supported by JSPS KAKENHI Grant Numbers 20K19743 and 20H00605, and by Start-up Program in Graduate School of Information Science and Technology, Osaka University. Yu Yokoi was supported by JST PRESTO Grant Number JPMJPR212B and by JST ERATO Grant Number JPMJER2301. The research was supported by the Lendület Programme of the Hungarian Academy of Sciences – grant number LP2021-1/2021, by Dynasnet European Research Council Synergy project – grant number ERC-2018-SYG 810115, by the Ministry of Innovation and Technology of Hungary – grant number ELTE TKP 2021-NKTA-62, and by JST CRONOS Japan Grant Number JPMJCS24K2.

References

- R. Aharoni and E. Berger. The intersection of a matroid and a simplicial complex. Transactions of the American Mathematical Society, 358(11):4895–4917, 2006.
- [2] R. Aharoni, D. Kotlar, and R. Ziv. On independent transversals in matroidal Latin rectangles. In The Seventh European Conference on Combinatorics, Graph Theory and Applications: EuroComb 2013, pages 549–550. Springer, 2013.
- [3] R. Aharoni, D. Kotlar, and R. Ziv. Rainbow sets in the intersection of two matroids. Journal of Combinatorial Theory, Series B, 118:129–136, 2016.
- [4] A. Brouwer, A. de Vries, and R. Wieringa. A lower bound for the length of partial transversals in a Latin square. *Nieuw Archief voor Wiskunde*, 26(2):330–332, 1978.
- [5] R. A. Brualdi and H. J. Ryser. Combinatorial Matrix Theory. Cambridge University Press, 1991.
- [6] P. J. Cameron and I. M. Wanless. Covering radius for sets of permutations. Discrete Mathematics, 293(1-3):91–109, 2005.
- [7] G. G. Chappell. A matroid generalization of a result on row-Latin rectangles. Journal of Combinatorial Theory, Series A, 88(2):235-245, 1999.
- [8] J. De Loera, S. Iwata, and M. Skutella. Combinatorial optimization. Oberwolfach Reports, 15(4):2969– 3023, 2019.
- [9] A. A. Drisko. Proof of the Alon-Tarsi conjecture for $n = 2^r p$. The Electronic Journal of Combinatorics, 5(1):R28, 1998.
- [10] M. R. Garey and D. S. Johnson. Computers and Intractability. Freeman San Francisco, 1979.
- [11] P. Hatami and P. W. Shor. A lower bound for the length of a partial transversal in a Latin square. Journal of Combinatorial Theory, Series A, 115(7):1103–1113, 2008.
- [12] D. Kotlar and R. Ziv. On the length of a partial independent transversal in a matroidal Latin square. The Electronic Journal of Combinatorics, 19(2):P12, 2012.
- [13] D. Kotlar and R. Ziv. Rainbow sets in the intersection of two matroids: A generalization of results of Drisko and Chappell. Discrete Mathematics, 338(5):695–697, 2015.
- [14] A. Linhares, N. Olver, C. Swamy, and R. Zenklusen. Approximate multi-matroid intersection via iterative refinement. *Mathematical Programming*, 183:397–418, 2020.
- [15] P. W. Shor. A lower bound for the length of a partial transversal in a Latin square. Journal of Combinatorial Theory, Series A, 33(1):1–8, 1982.

- [16] S. Stein. Transversals of Latin squares and their generalizations. Pacific Journal of Mathematics, 61(2):567–575, 1975.
- [17] I. M. Wanless. Transversals in Latin squares: a survey. Surveys in Combinatorics, 392:403–437, 2011.
- [18] D. E. Woolbright. An $n \times n$ Latin square has a transversal with at least $n \sqrt{n}$ distinct symbols. Journal of Combinatorial Theory, Series A, 24(2):235–237, 1978.