
















## IXPE detection of highly polarized X-rays from the magnetar 1E 1841–045

MICHELA RIGOSELLI <sup>1,2</sup> ROBERTO TAVERNA <sup>3</sup> SANDRO MEREGHETTI <sup>2</sup> ROBERTO TUROLLA <sup>3,4</sup>  
GIAN LUCA ISRAEL <sup>5</sup> SILVIA ZANE <sup>4</sup> LORENZO MARRA <sup>3</sup> FABIO MULERI <sup>6</sup> ALICE BORGHESE <sup>7</sup>  
FRANCESCO COTI ZELATI <sup>8,9</sup> DAVIDE DE GRANDIS <sup>8,9</sup> MATTEO IMBROGNO <sup>10,5</sup> RUTH M.E. KELLY <sup>4</sup>  
PAOLO ESPOSITO <sup>11</sup> AND NANDA REA <sup>8,9</sup>

<sup>1</sup>INAF, Osservatorio Astronomico di Brera, via Brera 28, I-20121 Milano, Italy

<sup>2</sup>INAF, Istituto di Astrofisica Spaziale e Fisica Cosmica di Milano, via Corti 12, I-20133 Milano, Italy

<sup>3</sup>Dipartimento di Fisica e Astronomia, Università degli Studi di Padova, Via Marzolo 8, I-35131 Padova, Italy

<sup>4</sup>Mullard Space Science Laboratory, University College London, Holmbury St Mary, Dorking, Surrey RH5 6NT, UK

<sup>5</sup>INAF, Osservatorio Astronomico di Roma, via Frascati 33, I-00078 Monteporzio Catone, Italy

<sup>6</sup>INAF, Istituto di Astrofisica e Planetologia Spaziali, Via del Fosso del Cavaliere 100, I-00133 Roma, Italy

<sup>7</sup>European Space Agency (ESA), European Space Astronomy Centre (ESAC), Camino Bajo del Castillo s/n, 28692 Villanueva de la Cañada, Madrid, Spain

<sup>8</sup>Institute of Space Sciences (ICE, CSIC), Campus UAB, Carrer de Can Magrans s/n, E-08193 Barcelona, Spain

<sup>9</sup>Institut d'Estudis Espacials de Catalunya (IEEC), 08860 Castelldefels (Barcelona), Spain

<sup>10</sup>Dipartimento di Fisica, Università degli Studi di Roma Tor Vergata, Via della Ricerca Scientifica 1, I-00133, Rome, Italy

<sup>11</sup>Scuola Universitaria Superiore IUSS Pavia, Palazzo del Broletto, piazza della Vittoria 15, I-27100 Pavia, Italy

### ABSTRACT

The Imaging X-ray Polarimetry Explorer (IXPE) observed for the first time highly polarized X-ray emission from the magnetar 1E 1841–045, targeted after a burst-active phase in August 2024. To date, IXPE has observed four other magnetars during quiescent periods, highlighting substantially different polarization properties. 1E 1841–045 exhibits a high, energy-dependent polarization degree, which increases monotonically from  $\approx 15\%$  at 2–3 keV up to  $\approx 55\%$  at 5.5–8 keV, while the polarization angle, aligned with the celestial North, remains fairly constant. The broadband spectrum (2–79 keV) obtained by combining simultaneous IXPE and NuSTAR data is well modeled by a blackbody and two power-law components. The unabsorbed 2–8 keV flux ( $\approx 2 \times 10^{-11}$  erg cm<sup>−2</sup> s<sup>−1</sup>) is about 10% higher than that obtained from archival XMM-Newton and NuSTAR observations. The polarization of the soft, thermal component does not exceed  $\approx 25\%$ , and may be produced by a condensed surface or a bombarded atmosphere. The intermediate power law is polarized at around 30%, consistent with predictions for resonant Compton scattering in the star magnetosphere; while, the hard power law exhibits a polarization degree exceeding 65%, pointing to a synchrotron/curvature origin.

**Keywords:** stars: magnetars — techniques: polarimetry — X-rays: stars

### 1. INTRODUCTION

Magnetars are ultra-magnetized neutron stars whose high-energy emission is mainly powered by their own magnetic energy (Duncan & Thompson 1992; Thompson & Duncan 1993). The presence of magnetic fields up to three orders of magnitude stronger than those typically found in ordinary young neutron stars is at

the basis of their enhanced X-ray emission and extreme variability phenomena on different timescales, ranging from short (tens of ms) bursts to months- or even years-long outbursts (see, e.g., Turolla et al. 2015; Kaspi & Beloborodov 2017; Esposito et al. 2021, and references therein).

Such a strong magnetic field ( $B \sim 10^{13}$ – $10^{15}$  G) makes magnetars extremely interesting targets for polarimetric observations in the X-ray range, which have been recently made possible by the NASA-ASI Imaging X-ray Polarimetry Explorer (IXPE; Weisskopf et al. 2022). Radiation in a strongly magnetized medium propagates

Corresponding author: Michela Rigoselli  
michela.rigoselli@inaf.it

in the ordinary (O) and extraordinary (X) normal modes (Gnedin & Pavlov 1974). The opacity for X-mode photons is strongly suppressed, below the electron cyclotron energy, relative to that of the O-mode ones (see Harding & Lai 2006, and references therein). The X-ray emission from magnetar sources is, then, expected to be substantially polarized (up to  $\approx 80\%$ ; Fernández & Davis 2011; Taverna et al. 2014, 2020; Caiazzo et al. 2022).

The IXPE satellite observed four magnetars during the first two years of operations. Polarization was clearly measured in the 2–8 keV band for 4U 0142+61 (Taverna et al. 2022), 1RXSJ170849.0–400910 (hereafter 1RXSJ1708 for short; Zane et al. 2023) and 1E 2259+586 (Heyl et al. 2024), while an unexpectedly low flux prevented significant detection in SGR 1806–20 (Turolla et al. 2023). In the two brightest sources the polarization properties turned out to strongly depend on energy. In 1RXSJ1708, the degree of polarization increases monotonically from  $\approx 20\%$  to  $\approx 80\%$  at a constant polarization angle making it the most polarized source detected by IXPE so far, while in 4U 0142+61 it is  $\approx 15\%$  at 2–3 keV, is consistent with zero around 4–5 keV, where the polarization angle swings by  $90^\circ$ , and then reaches  $\approx 35\%$  at 6–8 keV.

All the IXPE observations described above were targeted on bright magnetars during periods of quiescence (see also Taverna & Turolla 2024 for a review of all the results). On the other hand, most magnetars exhibit periods of activity during which they emit frequent short bursts of hard X-rays accompanied by flux and spectral changes and/or anomalies in the timing properties (glitches, spin-down rate variations, pulse profile changes). Since these phenomena are probably associated with magnetospheric reconfigurations, it is interesting to explore the polarimetric properties during or after periods of activity.

The recent reactivation of the magnetar 1E 1841–045 offered this possibility. The source lies at the center of the shell-type, 4'-diameter supernova remnant (SNR) Kes 73, at an estimated distance of  $D \approx 8.5$  kpc (Tian & Leahy 2008; Kumar & Safi-Harb 2010; Olausen & Kaspi 2014). Its X-ray luminosity  $L_X \sim 5 \times 10^{35}$  erg s $^{-1}$ , largely in excess of the rotational energy loss rate, and the detection of pulsations suggested a magnetar interpretation (Vasisht & Gotthelf 1997). This was confirmed by the subsequent detection of short bursts at different epochs (Morii et al. 2003; Wachter et al. 2004). Very recently, after more than 9 years of inactivity (An et al. 2013; Barthelmy et al. 2015), 1E 1841–045 emitted a series of hard X-ray bursts, beginning on 2024 August 20th (Swift Team 2024a,b; Fermi GBM Team 2024; GECAM team 2024; SVOM/GRM Team 2024). Soft X-ray ob-

servations of the source, performed by Swift-XRT and NICER (NICER Team 2024) about one day after the first burst, revealed an enhancement of  $\approx 25\%$  of the 0.5–10 keV flux with respect to the quiescent level.

In this *Letter* we report on an IXPE target of opportunity (ToO) observation of 1E 1841–045 performed soon after the burst-active phase, along with NuSTAR and archival Chandra, XMM-Newton and NuSTAR campaigns. Observations are detailed in §2 and the results of timing, spectral, and polarimetric analyses are presented in §3. Discussion follows in §4. In a companion paper (Stewart et al. 2024) an independent analysis of the same IXPE datasets is presented. The results and theoretical interpretation are largely consistent, except for the discussion on the nature of the soft X-ray component, where the two papers present distinct interpretive elements.

## 2. OBSERVATIONS AND DATA ANALYSIS

### 2.1. IXPE

Following reports of bursting activity and rebrightening of 1E 1841–045, an IXPE ToO pointing was requested (PI G. Younes). The observation, divided into two segments, started on 2024 September 28 01:32:39 UTC and ended on 2024 October 10 05:12:53 UTC, for a total of  $\approx 290$  ks of on source time for each of the three detector units (DUs).

We retrieved the level 1 and level 2 photon lists from the IXPE archive<sup>1</sup> and performed background rejection (see e.g. Di Marco et al. 2023), excising a  $\approx 100$  s interval, which includes solar flares. Source counts were extracted from a circular region with radius  $48''$ , centered on the position of 1E 1841–045. We checked that by taking a source extraction radius of  $30''$  (Weisskopf et al. 2022) the overall counts decrease (especially at higher energies), implying that some source events are lost without a definite improvement of the signal-to-noise (S/N) ratio. We extracted background counts from a concentric annulus with inner and outer radii  $r_{\text{int}}^{\text{bkg}} = 60''$  and  $r_{\text{ext}}^{\text{bkg}} = 240''$  solely for the polarization analysis (detailed in §3.2.1 and §3.2.2), that is, subtracting the SNR contribution from the source counts as normal background. For the spectral analysis, instead, we selected  $r_{\text{int}}^{\text{bkg}} = 150''$ , to include the SNR contribution as well (see §3.2.3).

No significant variation in the IXPE count rates is visible throughout the entire observation, so in the following we analyze the dataset obtained by joining the two segments. Given that background counts are always

<sup>1</sup> <https://heasarc.gsfc.nasa.gov/docs/ixpe/archive/>

**Table 1.** Log of X-ray observations and measured spin periods used in this work

Observatory	Instrument/Mode	ObsID	Start time (UTC)	End time (UTC)	Exposure (ks)	$P$ (s)
Chandra	ACIS/TE	729	2000-07-23 20:55:29	2000-07-24 05:46:55	29.26	–
Chandra	ACIS/TE	6732	2006-07-30 06:54:37	2006-07-30 14:30:26	24.86	–
Chandra	ACIS/TE	16950	2015-06-04 20:42:20	2015-06-05 05:16:21	28.69	–
Chandra	ACIS/TE	17668	2015-07-07 11:21:02	2015-07-07 17:32:53	20.89	–
Chandra	ACIS/TE	17692	2015-07-08 22:57:29	2015-07-09 05:54:24	23.26	–
Chandra	ACIS/TE	17693	2015-07-09 22:00:45	2015-07-10 04:58:34	22.77	–
XMM-Newton	EPIC/FF	0783080101	2017-03-31 05:54:27	2017-04-01 04:41:49	71.45	11.79693(2)
NuSTAR	–	30001025012	2013-09-21 11:26:07	2013-09-23 22:51:07	100.5	11.792407(8)
NuSTAR	–	91001330002	2024-08-29 04:21:09	2024-08-30 07:11:09	50.46	11.80646(3)
NuSTAR	–	91001335002	2024-09-28 23:36:09	2024-09-30 07:36:09	55.36	11.80654(2)
IXPE	–	03250499	2024-09-28 01:32:39	2024-10-10 05:12:53	292.5	11.80659(1)

subdominant with respect to those of the source within the IXPE band, we opted for an unweighted analysis of the level 2 photon lists, using version 20240701-v013 of the response files, provided in the online calibration database<sup>2</sup>. We checked that a weighted analysis does not produce significant differences.

## 2.2. Chandra

To estimate the contribution of the Kes 73 SNR to the magnetar spectra obtained with other instruments, we analyzed the archival data obtained with Chandra, which, thanks to its unmatched angular resolution, is the best X-ray telescope to disentangle the magnetar and SNR emissions. We used all the ACIS observations in Timed-Exposure (TE) mode listed in Table 1 (see also DOI: 10.25574/cdc.322), with the SNR located at the aim point of observation. We reprocessed the data with the tool `chandra_repro` of the Chandra Interactive Analysis of Observation software (CIAO; Fruscione et al. 2006), version 4.16, using the calibration database CALDB 4.11.1. The mosaic of the Chandra-ACIS images in the 0.5–8 keV energy band is shown in Figure 1.

For each of the six observations, we extracted three spectra from annuli centered on the magnetar position and with inner radius  $r_{\text{int}}^{\text{SNR}} = 6''$ , to avoid contamination from the magnetar, and outer radii  $r_{\text{ext}}^{\text{SNR}} = 15''$ ,  $48''$ , and  $60''$  (see Figure 1). These values were chosen to match the extraction regions used for XMM-Newton-EPIC (§2.3), IXPE (§2.1) and NuSTAR (§2.4), respectively. We extracted background counts from nearby boxes of size  $400'' \times 150''$ .

For each set of extraction regions, we simultaneously fit the spectra of the six observations with the sum of two non-equilibrium collisional ionization plasma models, absorbed by the interstellar medium (TBABS\*(VPSHOCK+VPSHOCK), see also Kumar et al. 2014). The parameters of the thermal component are the two plasma temperatures,  $kT_s$  and  $kT_h$ , and the two ionization timescales,  $\tau_s$  and  $\tau_h$ .


The fit with all the element abundances fixed at solar values (Anders & Grevesse 1989) gives, for the three regions, best fits with  $\chi^2/\text{dof} = 495.45/475$  ( $6'' - 15''$ ),  $1733.25/1254$  ( $6'' - 48''$ ) and  $2111.99/1414$  ( $6'' - 60''$ ). We then allowed the abundances of Mg, Si, and S — which are responsible for the main features observed at energies  $\approx 1.25$ ,  $\approx 1.8$  and  $\approx 2.4$  keV, respectively — to vary in the hotter plasma component. This led to a significant improvement in the  $6'' - 48''$  and  $6'' - 60''$  regions (F-test probability  $< 10^{-2}$ ), while the fit of the  $6'' - 15''$  region did not change significantly (F-test probability = 0.38).

These models (Figure A1 and the best fit parameters in Table A1), properly rescaled to account for the excision of the inner  $6''$  circle, were then added as fixed components in the spectral fits of the data from XMM-Newton, IXPE and NuSTAR (§3.1.1 and §3.1.2).

## 2.3. XMM-Newton

We analyzed the latest and deepest XMM-Newton observation of 1E1841–045 (see Table 1). The analysis of this dataset is presented in this *Letter* for the first time. The European Photon Imaging Cameras (EPIC) instrument was operated in Prime Full Window (pn camera, Strüder et al. 2001) and Prime Partial Window

<sup>2</sup> <https://heasarc.gsfc.nasa.gov/docs/ixpe/caldb>



**Figure 1.** Chandra mosaic of SNR Kes 73. The solid white circles mark the three annular extraction regions for the SNR spectrum, at  $r_{\text{int}}^{\text{SNR}} = 6''$  (innermost circle, common to all three) and  $r_{\text{ext}}^{\text{SNR}} = 15''$ ,  $48''$  and  $60''$  (outer circles). The dashed magenta circles mark the extraction region used for the background of XMM-Newton, NuSTAR and IXPE data:  $r_{\text{int}}^{\text{bkg}} = 150''$  and  $r_{\text{ext}}^{\text{bkg}} = 240''$ .

(MOS1/2 cameras, [Turner et al. 2001](#)) with the thick filter.

Data reduction was performed using the `eproc` and `emproc` pipelines of version 18 of the Science Analysis System (SAS). The extraction regions for the source and background are shown in Figure 1. We selected single- and multiple-pixel events (`pattern`  $\leq 4$  for the EPIC-pn and  $\leq 12$  for the EPIC-MOS).

#### 2.4. NuSTAR

NuSTAR ([Harrison et al. 2013](#)) observed 1E 1841–045 twice, following the recent bursting activity: the first observation started on August 29 and lasted 50 ks; the second started on September 28 and lasted 56 ks. Furthermore, to compare the properties of broadband emission before and after the bursting episode, we analyzed a long observation (100 ks) made in September 2013 (see Table 1). Data were processed using the `nupipeline` tool with default screening parameters, and spectra were extracted throughout the energy bandpass using `nuproducts`. The extraction regions for the source and background are shown in Figure 1.

### 3. RESULTS

We converted the times of arrival of the photons to the Solar System barycenter using the Jet Propulsion Laboratory Development Ephemeris JPL DE430, with source coordinates R.A. =  $18^{\text{h}}41^{\text{m}}19.34^{\text{s}}$ , Dec. =  $-04^{\circ}56'11.16''$ . Spectral and spectro-

polarimetric analyses were performed using `XSPEC` ([Arnaud 1996](#), version 12.11.0). Absorption from the interstellar medium was accounted for using the TBABS models with cross sections and abundances from [Anders & Grevesse \(1989\)](#). The  $\chi^2$  statistics was used to assess the goodness of the fits, and all reported errors are at  $1\sigma$  confidence level (cl hereafter), unless otherwise specified.

#### 3.1. Spectral analysis

##### 3.1.1. Pre bursting activity

We simultaneously fit the XMM-Newton-EPIC and NuSTAR spectra of 1E 1841–045 obtained before the August 2024 bursting phase. The emission component of SNR Kes 73 was included as explained in §2.2, and we added a constant to account for the calibration uncertainties between the three EPIC cameras and the two NuSTAR modules. We found that these normalization parameters for NuSTAR are 15%–20% higher than those of XMM-Newton for all the tested models.


We fitted to the spectra a (absorbed) two-component model, either two power laws (PL+PL), or a blackbody and a power law (BB+PL). In both cases, the fit was poor, with  $\chi^2/\text{dof} = 3319.90/2671$  and  $3331.42/2671$ , respectively, with a structured modulation in the residuals, suggesting the presence of an additional component. We then added a BB as a third component, obtaining  $\chi^2/\text{dof} = 2847.25/2669$  for the BB+PL+PL model and  $\chi^2/\text{dof} = 2919.53/2669$  for the BB+BB+PL model (see the two panels in Figure 2 and Table 2). The addition of the third component yields in both cases an F-test probability  $< 10^{-16}$ . The total unabsorbed fluxes in the 2–8 keV energy range turned out to be  $F_{\text{BB+PL+PL}} = (2.11 \pm 0.01) \times 10^{-11} \text{ erg cm}^{-2} \text{ s}^{-1}$  and  $F_{\text{BB+BB+PL}} = (2.03 \pm 0.01) \times 10^{-11} \text{ erg cm}^{-2} \text{ s}^{-1}$ .

##### 3.1.2. Post bursting activity

We fit the simultaneous IXPE and NuSTAR spectra in the 2–79 keV energy range. A cross-calibration constant was introduced to account for the different response of the three IXPE DUs and the two NuSTAR modules; we found that these normalization parameters for NuSTAR are  $\approx 30\%$  higher than those of IXPE for all the tested models. The contribution of SNR Kes 73 was included as explained in §2.2.

We started with absorbed PL+PL and BB+PL models and obtained best fits with  $\chi^2/\text{dof} = 904.11/883$  and  $972.07/883$ , respectively. The PL+PL fit is formally acceptable, but yields a very high column absorption  $N_{\text{H}} = (4.4 \pm 0.2) \times 10^{22} \text{ cm}^{-2}$  and an unreasonably steep photon index of  $\Gamma_{\text{soft}} = 5.00 \pm 0.14$ , probably mimicking the presence of a thermal component. If we fix the





**Figure 2.** Best fit and residuals to the XMM-Newton (EPIC-pn in black, MOS1 and MOS2 in red and green respectively) and NuSTAR (FPMA and FPMB detectors in blue and cyan respectively) pre-burst data. The solid lines show the single components of the models, while the dashed lines show the SNR contribution as computed in §2.2, which was kept fixed in all fits. The plotted spectra were rebinned to improve visualization and single components are only displayed in the cases of the EPIC-pn and FPMA detectors for clarity.

absorption to the value used by [Stewart et al. \(2024\)](#),  $N_{\text{H}} = 2.6 \times 10^{22} \text{ cm}^{-2}$ , we obtain  $\Gamma_{\text{soft}} = 3.72 \pm 0.04$  and  $\Gamma_{\text{hard}} = 1.21 \pm 0.02$ , which are quite in agreement with what found by [Stewart et al. \(2024\)](#). However, this fit is marginally acceptable ( $\chi^2/\text{dof} = 1007.90/884$ ) and the addition of the thermal component is statistically required (F-test probability  $\approx 2 \times 10^{-21}$ , see below).

As a consequence, following the approach discussed in 3.1.1, we considered a three-model spectral decomposition for the IXPE and NuSTAR data. Given the reduced energy band of IXPE and the lower number of source counts, we froze the value of the column density to those obtained in §3.1.1. We found that both the BB+PL+PL and BB+BB+PL decompositions provide an acceptable fit ( $\chi^2/\text{dof} = 904.54/882$  and  $900.77/882$ , respectively) with an even distribution of the residuals. The data, together with the best-fit models, are shown in Figure 3, and the corresponding best-fit parameters are reported in Table 2. The unabsorbed fluxes in the 2–8 keV energy range are  $F_{\text{BB+PL+PL}} = (2.31 \pm 0.02) \times 10^{-11} \text{ erg cm}^{-2} \text{ s}^{-1}$  and  $F_{\text{BB+BB+PL}} = (2.22 \pm 0.02) \times 10^{-11} \text{ erg cm}^{-2} \text{ s}^{-1}$ .

### 3.2. Polarization analysis


#### 3.2.1. Energy-resolved polarimetric analysis

The phase-averaged Stokes parameters  $I$ ,  $Q$  and  $U$  integrated over the 2–8 keV band were obtained from the photon lists for each of the IXPE DUs using the *ixpeobssim* package ([Baldini et al. 2022](#)). The results


for the normalized Stokes parameters  $Q/I$  and  $U/I$  are shown in Figure 4. The detection obtained from the sum of the three DUs is highly significant ( $\gtrsim 8\sigma$ ), yielding a polarization degree  $\text{PD} = \sqrt{Q^2 + U^2}/I = 25.9 \pm 3.1\%$ , well above the minimum detectable polarization at 99% cl (MDP<sub>99</sub>, see [Weisskopf et al. 2010](#)) of 9.5%, and a polarization angle  $\text{PA} = \arctan(U/Q)/2 = 1.1 \pm 3.5^\circ$ , measured east of celestial north.

We then divided the 2–8 keV energy band into four energy bins and extracted the values of PD and PA in each of them. Polarization properties were derived using both the *pcube* algorithm within the *ixpeobssim* suite and by fitting simultaneously the  $I$ ,  $Q$  and  $U$  spectra with *XSPEC* in each energy interval. In the latter case, we first fit the IXPE spectrum (Stokes  $I$ ) in the 2–8 keV band using a simple BB+PL model with  $N_{\text{H}}$  frozen at the value listed in Table 2 ( $\chi^2/\text{dof} = 230.95/216$ ). Then, we fit simultaneously the spectra of  $I$ ,  $U$ , and  $Q$  in each energy interval, keeping the parameters of BB and PL at their best-fit values found previously, and convolving the spectral model with a constant polarization (POLCONST).

The results, reported in Table 3, show an overall agreement (within  $1\sigma$ ) between the parameters obtained using the two methods, confirming a posteriori that the method used to extract PD and PA with *XSPEC* is essentially model independent. The measurements are above MDP<sub>99</sub> in each energy bin, with significance  $> 3\sigma$ . The most probable values of PD and PA, together with the




**Figure 3.** Best fit and residuals to the IXPE (DU1, DU2 and DU3 in black, red and green respectively) and NuSTAR (FPMA and FPMB detectors in blue and cyan respectively) 2024 data. Line code is as in Figure 2.



**Figure 4.** The normalized Stokes parameters  $Q/I$  and  $U/I$  measured by the IXPE DU1 (cyan), DU2 (orange) and DU3 (green) along with the combined measurements from the three DUs (black);  $1\sigma$  error bars are also shown. Dashed circles represent the loci of constant PD while straight lines those of constant PA. The  $\text{MDP}_{99}$  relative to the combined measurement is marked by the filled circle.

corresponding confidence contours at 68% and 99% cl, are also plotted in Figure 5. The degree of polarization increases monotonically with energy, from  $\approx 15\%$  at 2–3 keV up to more than 50% at higher energies (5.5–8 keV), while the polarization angle remains fairly constant, with only a slight deviation in the bin at the highest energy. However, the polarization direction is



**Figure 5.** Polar plot of the energy-dependent PD and PA obtained using XSPEC (see text for details). Filled and empty contours correspond to the 68% and 99% confidence regions, respectively, around the most probable values reported in Table 3.

compatible with that of the celestial north in the entire IXPE band within  $1\sigma$ .

### 3.2.2. Phase-resolved polarimetric analysis

To perform a phase-dependent polarimetric analysis, we measured the spin period of 1E1841–045 at the epoch of the IXPE observation. Using only the IXPE data, we determined the best period by searching around the frequency 0.085 Hz reported in Dib & Kaspi (2014) and using the  $Z_n^2$ -search technique included in the HENDRICS suite v.7.0 (Bachetti 2018). We found that the most likely value for the frequency (using  $n = 3$  harmonics,  $\max Z_3^2 = 850$ ) is  $f = 0.08469848(8)$  Hz, corresponding to a period  $P = 11.80659(1)$  s, at the MJD

**Table 2.** Fit parameters of 1E 1841–045 spectra

	BB+PL+PL		BB+BB+PL	
	PRE	POST	PRE	POST
$N_{\text{H}}$ ( $10^{22} \text{ cm}^{-2}$ )	$2.20^{+0.03}_{-0.03}$	2.20*	$2.009^{+0.015}_{-0.014}$	2.009*
$kT_{\text{BB1}}$ (keV)	$0.491^{+0.004}_{-0.004}$	$0.501^{+0.007}_{-0.007}$	$0.465^{+0.007}_{-0.008}$	$0.524^{+0.008}_{-0.008}$
$R_{\text{BB1}}$ (km)	$3.94^{+0.12}_{-0.13}$	$4.62^{+0.23}_{-0.23}$	$5.58^{+0.18}_{-0.15}$	$4.65^{+0.16}_{-0.15}$
$F_{\text{BB1}}^{2-8}$ ( $10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$ )	$5.25^{+0.24}_{-0.25}$	$8.2^{+0.5}_{-0.6}$	$7.6^{+0.3}_{-0.4}$	$10.6^{+0.3}_{-0.3}$
$kT_{\text{BB2}}$ (keV)	—	—	$1.04^{+0.05}_{-0.05}$	$1.4^{+0.1}_{-0.1}$
$R_{\text{BB2}}$ (km)	—	—	$0.56^{+0.07}_{-0.06}$	$0.24^{+0.04}_{-0.03}$
$F_{\text{BB2}}^{2-8}$ ( $10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$ )	—	—	$4.3^{+0.2}_{-0.2}$	$2.6^{+0.3}_{-0.3}$
$\Gamma_{\text{PL2}}$	$2.5^{+0.1}_{-0.1}$	$2.4^{+0.2}_{-0.3}$	—	—
$F_{\text{PL2}}^{2-8}$ ( $10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$ )	$12.2^{+0.4}_{-0.4}$	$9.3^{+0.8}_{-0.9}$	—	—
$\Gamma_{\text{PL3}}$	$0.93^{+0.06}_{-0.07}$	$1.1^{+0.06}_{-0.07}$	$1.34^{+0.02}_{-0.02}$	$1.30^{+0.03}_{-0.03}$
$F_{\text{PL3}}^{2-8}$ ( $10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$ )	$3.3^{+0.6}_{-0.5}$	$5.7^{+1.1}_{-1.7}$	$8.4^{+0.3}_{-0.3}$	$9.0^{+0.4}_{-0.4}$
$F_{\text{TOT}}^{2-8}$ ( $10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$ )	$21.1^{+0.1}_{-0.1}$	$23.1^{+0.2}_{-0.2}$	$20.3^{+0.1}_{-0.1}$	$22.2^{+0.2}_{-0.2}$
$F_{\text{TOT}}^{8-79}$ ( $10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$ )	$54.3^{+0.7}_{-0.7}$	$59^{+1}_{-1}$	$50.0^{+0.7}_{-0.7}$	$58^{+1}_{-1}$
$K_{\text{pn}}$ or $K_{\text{DU1}}$	1*	1*	1*	1*
$K_{\text{MOS1}}$ or $K_{\text{DU2}}$	$1.139 \pm 0.005$	$0.972 \pm 0.011$	$1.139 \pm 0.005$	$0.971 \pm 0.011$
$K_{\text{MOS2}}$ or $K_{\text{DU3}}$	$1.061 \pm 0.005$	$0.978 \pm 0.011$	$1.061 \pm 0.005$	$0.978 \pm 0.011$
$K_{\text{FMFA}}$	$1.185 \pm 0.007$	$1.328 \pm 0.017$	$1.183 \pm 0.007$	$1.326 \pm 0.017$
$K_{\text{FMFB}}$	$1.147 \pm 0.007$	$1.314 \pm 0.017$	$1.143 \pm 0.007$	$1.312 \pm 0.017$
$\chi^2/\text{dof}$	2847.25/2669	904.54/882	2919.53/2669	900.77/882

NOTE— The parameters marked with an asterisk were kept fixed in the fit. Blackbody radii are calculated assuming a distance of 8.5 kpc (see [Tian & Leahy 2008](#)). The fluxes, corrected for the absorption, are in the specified energy range.  $K$  are the cross-calibration constants between the XMM-Newton-EPIC cameras and the NuSTAR FPM detectors (PRE), and between the IXPE detector units and the NuSTAR FPM detectors (POST).

epoch 60587.140813793325, while the value of the frequency derivative was not constrained.

The pulse profile shown in the left panel of Figure 6 is obtained folding the data at our best spin period and dividing the counts into 20 equally spaced bins. The pulse shape and the pulsed fraction<sup>3</sup> do not appear to change significantly with energy, with the latter being  $(21.2 \pm 1.5)\%$  in the 2–4 keV band and only slightly

higher (but still compatible within  $1\sigma$ ) in the 4–8 keV range,  $(27.6 \pm 7.3)\%$ .


For our phase-dependent polarimetric analysis we consider four phase intervals, corresponding to the peak, decline, dip, and rise of the pulse (shown by the dotted vertical lines in the left panel of Figure 6). This choice allows us to characterize the polarization properties in the relevant parts of the light curve and, at the same time, ensures a significant detection ( $\text{PD} > \text{MDP}_{99}$ ) in each bin in the 2–8 keV range. However, the relatively short exposure time and the consequent choice of rather wide phase bins necessarily limits the inferences that can be drawn from such an analysis. The results for

<sup>3</sup> Defined as  $(\text{max} - \text{min})/(\text{max} + \text{min})$ , where max and min are the maximum and the minimum count rates, respectively, of the pulse profile.

**Table 3.** Energy-resolved polarization degree and angle

	2.0–3.0 keV	3.0–4.0 keV	4.0–5.5 keV	5.5–8.0 keV	2.0–8.0 keV
pcube PD (%)	$16.1 \pm 3.5$	$28.0 \pm 4.0$	$37.9 \pm 6.1$	$54.2 \pm 15.2$	$25.8 \pm 3.1$
pcube PA (deg)	$4.6 \pm 6.3$	$2.1 \pm 4.1$	$2.0 \pm 4.6$	$-6.7 \pm 8.0$	$1.1 \pm 3.5$
pcube PD S/N	$4.6\sigma$	$7.0\sigma$	$6.2\sigma$	$3.6\sigma$	$8.3\sigma$
pcube MDP <sub>99</sub> (%)	10.7	12.1	18.5	45.9	9.5
XSPEC PD (%)	$14.5 \pm 3.8$	$27.9 \pm 4.1$	$43.1 \pm 6.6$	$52.9 \pm 12.1$	$25.3 \pm 2.4$
XSPEC PA (deg)	$3.3 \pm 7.5$	$1.7 \pm 4.2$	$4.1 \pm 4.4$	$-12.9 \pm 6.7$	$2.3 \pm 2.7$
XSPEC PD S/N	$3.8\sigma$	$6.8\sigma$	$6.5\sigma$	$4.4\sigma$	$10.5\sigma$

NOTE—Values are extracted combining the contributions of the three IXPE DUs, using the `ixpeobssim` and `XSPEC` suites (see text for details). Signal-to-noise (S/N) ratios are obtained dividing the most probable value by the corresponding  $1\sigma$  error.



**Figure 6.** IXPE count rate (left panel), polarization degree (central panel) and polarization angle (right panel) as a function of the rotational phase; two cycles are shown for clarity. Results are presented for the 2–8 (blue), 2–4 (orange) and 4–8 keV (green) energy ranges. Polarization degree and angle are shown for the four phase intervals marked by vertical dotted lines in the left panel. In all the bins where PD is below MDP<sub>99</sub>, a  $3\sigma$  upper limit (marked by a downward arrow) is shown for PD and the errors on PA are extended over the entire  $[-90^\circ, 90^\circ]$  range (double-headed arrows).

PD and PA, obtained using the `ixpeobssim-pcube` algorithm, are shown in Figure 6 (center and right panels) and reported in Table 4. In the 2–8 keV range, the polarization degree varies between  $\approx 20\%$  and  $\approx 36\%$ . A fit with a simple sinusoidal profile of both counts and PD is acceptable around or better than  $3\sigma$  cl and suggests that the two oscillations are in-phase within  $1\sigma$  errors; fitting PD with a constant still produces an acceptable result, albeit only at  $2\sigma$  cl.

We also analyzed the phase-dependent behavior of the polarization in the 2–4 and 4–8 keV energy ranges. At lower energies, the degree of polarization is above MDP<sub>99</sub> in three phase bins and slightly below around the pulse minimum. If we assume the most probable value at the dip, PD still follows a sinusoidal pattern

that is in phase with the light curve; a fit with a constant is ruled out at 96% cl. At higher energies (4–8 keV), the polarization signal is significant in only one out of four phase intervals, excluding any conclusions.

Concerning the variation of PA with phase, both the fits with a constant and a sinusoid in the 2–8 keV range are acceptable at better than  $3\sigma$  cl, while those in the 2–4 keV range are both rejected. We then tried a fit in the 2–8 keV band using a rotating vector model (RVM, see Radhakrishnan & Cooke 1969). This is motivated by the fact that, if vacuum birefringence is indeed at work in magnetars, it is expected to produce phase-dependent variations in the polarization angle that match the RVM predictions – a behavior already observed in a few sources (see Taverna et al. 2022; Zane et al. 2023; Heyl



**Table 4.** Phase-resolved polarization degree and angle.

Phase bin	Energy range (keV)	PD (%)	PA (deg)	PD S/N
0.0 – 0.15	2.0 – 8.0	$26.1 \pm 7.1$	$-8.5 \pm 7.8$	$3.7\sigma$
	2.0 – 4.0	$19.0 \pm 6.3$	$-14.3 \pm 9.4$	$3.0\sigma$
	4.0 – 8.0	$< 98.7$	-	-
0.15 – 0.4	2.0 – 8.0	$35.8 \pm 6.0$	$-0.9 \pm 4.8$	$6.0\sigma$
	2.0 – 4.0	$29.4 \pm 5.3$	$3.7 \pm 5.1$	$5.6\sigma$
	4.0 – 8.0	$56.1 \pm 13.6$	$-7.5 \pm 6.9$	$4.1\sigma$
0.4 – 0.8	2.0 – 8.0	$18.5 \pm 5.5$	$0.2 \pm 8.4$	$3.4\sigma$
	2.0 – 4.0	$< 25.7$	-	-
	4.0 – 8.0	$< 76.2$	-	-
0.8 – 1.0	2.0 – 8.0	$29.0 \pm 6.8$	$12.2 \pm 6.7$	$4.3\sigma$
	2.0 – 4.0	$28.9 \pm 5.7$	$16.5 \pm 5.6$	$5.1\sigma$
	4.0 – 8.0	$< 80.1$	-	-


NOTE—  $3\sigma$  upper limits are provided when  $PD < MDP_{99}$ . S/N ratios are obtained dividing the most probable value by the corresponding  $1\sigma$  error.

et al. 2024). However, the two parameters of the model (the inclinations of the line-of-sight and magnetic axis with respect to the rotation axis) turned out to be totally unconstrained because of the limited statistics. In this respect, the analysis of the phase-dependent PA does not provide useful information on the source geometry, and the only conclusion that can be reached is that PA is not compatible with a constant in the soft energy band.

### 3.2.3. Spectro-polarimetric analysis

As a final step, we explored the polarization properties of the individual components entering the spectral models discussed in §3.1, i.e. TBABS  $\times$  (SNR + BBODYRAD + BBODYRAD + POWERLAW) (hereafter model 1) and TBABS  $\times$  (SNR + BBODYRAD + POWERLAW + POWERLAW) (hereafter model 2), where SNR stands for the SNR spectral model discussed in §2.2. We froze all spectral parameters at the values reported in Table 2 (POST) and convolved each spectral component with a constant polarization model (POLCONST), simultaneously fitting the  $I$ ,  $Q$  and  $U$  spectra. Although the contribution of the SNR to the flux is sizable at 2–4 keV (see Figure 3 and Table A1), its polarization is expected to be small, as discussed at the end of the section. Consequently, we froze the SNR polarization degree to zero.

The hard PL accounts for almost all the counts above  $\approx 5$  keV in the case of model 1 (see Figure 3, right panel). Therefore, we fixed its polarization to the value measured with IXPE in the 5.5–8 keV range (see Table 3) and allowed the polarization of the soft and hard BBs



**Figure 7.** Polarization of the single spectral components (see text for details). Filled and empty contours correspond to the 68% and 99% confidence regions, respectively, around the most probable values; crosses mark the values at which PD and PA were frozen for the specified component.

to vary. As shown in Figure 7 (panel A), the polarization of the softer thermal component is approximately 20% and is constrained at 99% cl, while that of the hot blackbody is unconstrained, and we can only conclude that the two thermal components are polarized in the same direction at a confidence level of 68%.

On the other hand, in model 2 no single component dominates the spectrum in any energy range (see Figure 3, left panel). So, none of them can be associated a priori with the polarization measured by IXPE in a given energy interval. Since a fit with all parameters left free is particularly poor, with all the parameters uncon-

strained, we decided to run different tests, allowing the fit to compute the polarization of two components and freezing that of the third one.

First, we checked that, by freezing only the degree of polarization of one of the three components and leaving all other polarization parameters free to vary, the polarization of the other two components is unconstrained at 68%. Then, for the component with frozen polarization, we fixed both PD and PA. In particular, we decided to take  $PA = 0^\circ$ , in agreement, within  $1\sigma$  cl, with the IXPE measurement in all the energy bins (see Figure 5).

The high polarization degree obtained in the 2–8 keV energy interval suggests that, in the case of model 1, the hard PL component may be due to synchrotron radiation. Hence, for model 2 we fixed the PD of the hard PL, taking 75% as representative of synchrotron emission (see e.g. Rybicki & Lightman 1979). Note that this is not the PD measured in the 5.5–8 keV bin, since both PLs contribute in this energy range, allowing the soft PL to be less polarized. The result is reported in panel B of Figure 7. Although the polarization of both components turns out to be unconstrained at 99% cl, the derived values at 68% cl are  $\approx 20\%$  for the BB and  $\approx 30\%$  for the soft PL, while PA is compatible with zero, in agreement with observations. The latter value of PD suggests that the soft PL may be produced by resonant Compton scattering (RCS, which predicts  $PD \approx 33\%$ ; Taverna et al. 2020). We then froze this parameter at 33%, imposing again  $PA = 0^\circ$ , and leaving the polarization of the BB and the hard PL free to vary. The result is reported in Figure 7 (panel C). In this case, the degree of polarization of the hard PL is well constrained at  $PD \approx 65\%$  and its polarization angle is  $PA \approx -6^\circ$  (and compatible with  $0^\circ$  within  $1\sigma$ ). This suggests that, under the assumption that the polarization of each spectral component is constant across the IXPE band, the data can be reproduced by a model comprising a soft BB with  $PD \approx 20\%$ , a mildly polarized soft PL ( $PD \approx 30\%$ ) and a highly polarized hard PL, all with a polarization direction compatible with  $0^\circ$ .

Finally, we checked that by extracting the IXPE counts in an annular region centered on the source, with inner and outer radii  $30''$  and  $150''$ , respectively, from which the background was subtracted as described above, the `pcube` PD turns out to be below  $MDP_{99} \approx 20\%$ , both in the 2–8 keV and in the 2–4 keV energy ranges. This supports the assumption that SNR emission exhibits relatively low polarization, consistent with other SNRs observed by IXPE, where the average degree of polarization does not exceed  $\approx 10\%$  at most (see e.g. Slane et al. 2024). As a further test, we performed the fit with model 2 under the same conditions discussed

above, but considering the polarization angle frozen at  $+90^\circ$  or  $-90^\circ$ . This resulted in a  $> 100\%$  polarization of at least one of the two free components, and thus reasonably rules out the possibility that the emission associated with a specific spectral component is polarized perpendicularly to that associated with the other two.

#### 4. DISCUSSION


In this work, we investigated the spectro-polarimetric properties of 1E 1841–045 after it entered a period of bursting activity. We found that the emission over the whole IXPE energy band is strongly polarized, with  $PD = (25.9 \pm 3.1)\%$  and  $PA = (1.1 \pm 3.5)^\circ$  at a significance greater than  $8\sigma$ . The degree of polarization increases monotonically with energy, and its variation with the rotational phase broadly follows the pulse profile. The polarization angle remains fairly constant in both energy and phase, with the exception of the 2–4 keV energy range, where a phase modulation in PA is detected.

The broadband (2–79 keV) spectrum, obtained joining IXPE and simultaneous NuSTAR data, can be reproduced equally well by a BB+PL+PL or BB+BB+PL model. The 2–8 keV fluxes are about 10% higher than those obtained with the same spectral decompositions from archival XMM-Newton and NuSTAR data collected before the bursting phase. The increase in flux is more pronounced (about 15%) at higher energies, between 8 and 79 keV<sup>4</sup>.

There is no substantial evolution of the spectral parameters with respect to the pre-burst state, although the temperature (radius) of the hotter BB increased (decreased) after the onset of the bursting activity. No evidence of a further hot and bright thermal component associated with a large heat deposition in the crust was found, as instead observed in full-fledged outbursts (see e.g. Coti Zelati et al. 2018).

However, a significant evolution is seen in the pulse profile. We found the best-folding spin period in each of the XMM-Newton and NuSTAR observations (see Table 1) and we obtained the pulse profiles shown in Figure 8. Before the burst, the 2017 EPIC-pn and the 2013 NuSTAR data (first and second panels in Figure 8) consistently show a double-peaked profile, with  $PF = (23.6 \pm 0.9)\%$  at 2–8 keV and  $(25.2 \pm 1.6)\%$  at

<sup>4</sup> Due to the  $\approx 30\%$  cross-correlation uncertainty between IXPE and NuSTAR (see Table 2 and Stewart et al. 2024), the reported flux increments depend on our choice to consider IXPE as the reference instrument; the different choice operated by Stewart et al. (2024) leads to a flux variation of 40%.



**Figure 8.** Background-subtracted EPIC-pn pulse profiles in the 2–8 (blue), 2–4 (orange) and 4–8 keV (green) energy bands (left panel). The three rightmost panels show the NuSTAR pulse profiles in the 2–79 (red), 2–8 (blue) and 8–79 keV (black) energy bands at three different epochs.

8–79 keV. In the NuSTAR data obtained a few days after the onset of the active burst phase, a dip appeared in the rise toward the primary peak (third panel, see also [Younes et al. 2024](#)) that partially recovered one month later (fourth panel). The pulse profile measured by IXPE (see Figure 6, left panel) reveals the presence of two plateaux, just before and after the main peak, which are reminiscent of the secondary peak and the small dip seen in the NuSTAR 2024 data (fourth panel in Figure 8). These features are not well resolved in the IXPE light curve, likely due to the lower number of events collected in the relatively short exposure time. The pulsed fraction attains a value quite similar to that of the other instruments, with  $PF = (21.5 \pm 2.2)\%$  in the 2–8 keV band, without significant changes with energy.

Like other magnetars, 1E1841–045 is known to exhibit significant hard X-ray emission. These high-energy tails were originally detected by INTEGRAL ([Kuiper et al. 2004, 2006](#); [Mereghetti et al. 2005](#); [Molkov et al. 2005](#); [Götz et al. 2006](#)), can extend up to a few hundreds of keV, and are highly phase dependent ([den Hartog et al. 2008a,b](#); [Vogel et al. 2014](#); [Yang et al. 2016](#)). Their physical origin is still poorly understood and different interpretations have been proposed, ranging from thermal bremsstrahlung in the surface layers heated by returning currents, to synchrotron emission from pairs in the magnetosphere ([Thompson & Beloborodov 2005](#)), to Compton resonant upscattering from relativistic charges ([Baring et al. 2005](#); [Baring & Harding 2008](#); [Beloborodov 2013](#); [Wadiasingh et al. 2018](#)). Thanks to NuSTAR, we were able to characterize this emission component after the beginning of the burst-active phase. We find that the hard power law contributes significantly to the large polarization of the source at higher energies (5–8 keV).

Attempts to perform a spectro-polarimetric analysis of the IXPE data alone by associating a constant polarization model with each spectral component and leaving all parameters free to vary resulted in a largely un-

constrained fit. However, fixing the polarization parameters of one of the spectral components produces a reasonably constrained fit, and the hard PL turns out to be polarized at more than 65%, regardless of which component is actually frozen. This large polarization is difficult to reconcile with resonant Compton scattering models, which predict  $PD \approx 30\%–40\%$  because of the 1/3 ratio of the X- and O-mode photon scattering cross sections ([Fernández & Davis 2011](#); [Taverna et al. 2014](#); [Wadiasingh et al. 2018](#), see also [Stewart et al. 2024](#)). However, the polarization degree and the power-law spectral distribution may be compatible with a synchrotron/curvature origin for the high-energy tail. In fact, if one assumes that synchrotron radiation is produced by a power-law distribution of electrons, the photon index  $\Gamma_{\text{hard}} = \Gamma_{\text{PL}_3} \sim 1$  (see Table 2 POST) implies a degree of polarization of  $\approx 75\%$  ([Rybicki & Lightman 1979](#)). Interestingly, by fixing the polarization degree of the hard PL at 75%, the soft PL turned out to be polarized at around 30%, as expected if it is produced by resonant Compton scattering in the star magnetosphere (see [Taverna et al. 2014, 2020](#)). Moreover, the polarization of the soft thermal component is  $\approx 25\%$ , close to that observed in other magnetars ([Taverna et al. 2022](#); [Zane et al. 2023](#); [Heyl et al. 2024](#)). Such polarization can hardly come from a strongly passive, magnetized atmosphere (for  $B \gtrsim 10^{14}$  G, also allowing for mode conversion; [Kelly et al. 2024b](#)), while it may be produced by a condensed surface or a bombarded atmosphere ([Taverna et al. 2020](#); [Kelly et al. 2024a](#)).

Although the presence of a soft, thermal, mildly polarized component and a hard, strongly polarized power law appears to be well established, the nature of emission at intermediate energies ( $\approx 3–6$  keV) is not so clear. It could be either a softer power law, due to the RCS of the seed thermal photons, or a second, hotter thermal component. In the latter case, the relatively large polarization ( $\approx 30\%$ ) is incompatible with emission from a

condensed surface and suggests that part of the star is covered by an atmosphere, as proposed for 1RXS J1708 (Zane et al. 2023).

Because of vacuum birefringence, the polarization angle in magnetars is actually fixed at the polarization limiting radius ( $r_{\text{pl}} \approx 80 (B/10^{14} \text{ G})^{2/5} (E/1 \text{ keV})^{1/5} R_{\text{NS}}$ ; Heyl & Shaviv 2002), at least when radiation is produced inside  $r_{\text{pl}}$ . As a consequence, radiation polarized in a given direction (wrt the local  $B$ -field) at emission is characterized by the same (phase-averaged) PA at infinity, which is related to the projection of the dipole axis on the plane of the sky (Taverna et al. 2015). This implies that the observed PA for the two normal modes must differ by  $90^\circ$ . In particular, if PA is counted from the projection of the star rotation axis in the plane of the sky, it is  $\text{PA} = 0^\circ/90^\circ$  for O-/X-mode photons.

Since in 1E 1841–045 the polarization direction changes little with energy, the three spectral components should be polarized in the same mode if they come from below  $r_{\text{pl}}$ . Although there is no way to tell which is the dominant polarization mode from the data, we note that both atmospheric emission (if the preferred model is BB+BB+PL) and RCS (in case of the BB+PL+PL decomposition) result in an intermediate component mostly polarized in the X-mode. This lends support to the soft BB being polarized in the X-mode, too. This is compatible with the emission of a magnetic condensate which can be polarized in the X- or O-mode, depending on the emission/viewing geometry; the former was invoked for the thermal component in 1RXS J1708. Intriguingly, the polarization direction of the hard PL photons seems to be the same.

Synchrotron emission is polarized perpendicular to the magnetic field projection in the plane of the sky and is not, in general, associated with the X- and O-modes. However, if synchrotron radiation comes from below  $r_{\text{pl}}$ , the strong  $B$ -field forces also synchrotron photons to propagate in the two normal modes, likely in the X-mode for the case at hand. A possibility is that synchrotron emission is due to a population of relativistic pairs produced (similarly to what suggested by Beloborodov 2013) at the base of a localized twisted bundle, where the magnetic field direction is expected to change little. In this picture, the phase-dependent polarization angle should oscillate according to the rotating vector model, with the amplitude depending on the geometry of the source. Unfortunately, as discussed above, not enough IXPE counts have been collected to reconstruct the behavior of PA above 4 keV.

The general behavior of the polarization in 1E 1841–045 (monotonic increase with energy of PD at a constant polarization angle) is reminiscent of that

observed in 1RXS J1708. The similarity between the two sources goes deeper, since both exhibit a double power-law tail with comparable spectral indices ( $\Gamma_{\text{soft}} \sim 2.5$ – $2.7$  and  $\Gamma_{\text{hard}} \sim 1$ ; see den Hartog et al. 2008a, for the X-ray spectrum of 1RXS J1708). Zane et al. (2023) proposed two possible scenarios for 1RXS J1708: either two thermal components, one with low polarization and the other with high polarization, or one thermal component polarized at  $\approx 20\%$  and a power law, with  $\Gamma \sim 3$ , polarized at  $\approx 70\%$ . According to the latter interpretation, the same scenario discussed here for 1E 1841–045 may also apply to 1RXS J1708, although the lack of simultaneous high-energy observations prevented in their case the complete characterization of the spectro-polarimetric properties at the upper end of the IXPE energy range.

## 5. CONCLUSIONS


We have reported the first X-ray polarimetric measurements of the magnetar 1E 1841–045 obtained with the IXPE satellite, together with a detailed analysis of data from Chandra, XMM-Newton and NuSTAR. These complementary data have been essential to properly take into account the non-negligible contribution of the Kes 73 SNR (20% of the 2–8 keV flux in the IXPE source extraction region) and to place the observed broadband flux and spectral properties of 1E 1841–045 in the context of the previous history of this persistent magnetar. In fact, the IXPE observation was obtained 40 days after the period of bursting activity and the increase in persistent flux that occurred in August 2024.

The polarization properties found for 1E 1841–045 are compared to those of the other four magnetars observed with IXPE in Figure 9. Similarly to 1RXS J1708, the linear polarization increases with the energy from  $\approx 15\%$  to  $\approx 55\%$ , without any change in the polarization angle.

We found that a three-component spectral model (BB+PL+PL or BB+BB+PL) is required to fit the simultaneous IXPE and NuSTAR data, resulting in a flux of  $8 \times 10^{-11} \text{ erg cm}^{-2} \text{ s}^{-1}$  between 2–79 keV. The analysis of the archival XMM-Newton and NuSTAR data obtained before the source activation favors a BB+PL+PL model with flux of  $7.5 \times 10^{-11} \text{ erg cm}^{-2} \text{ s}^{-1}$  (2–79 keV, see also An et al. 2013). Although only a moderate increase of the flux was found between the pre- and the post-burst phases, the pulse profile experienced a noticeable evolution, especially above 8 keV where the hard power law dominates the flux.

High-energy tails are pretty ubiquitous in the magnetar population, but their origin is still poorly understood. Our spectro-polarimetric analysis showed that in 1E 1841–045 this component is polarized at more





**Figure 9.** Energy dependence of the linear polarization degree (left panel) and polarization angle (right panel) of the five magnetars observed with IXPE. Thin lines indicate results for the total energy range (in case of significant polarization detection in the 2–8 keV range). See details in Taverna et al. (2022); Zane et al. (2023); Turolla et al. (2023); Heyl et al. (2024).

than 65% and can be well interpreted in terms of synchrotron/curvature emission. The intermediate power law component has a degree of polarization of 30%, consistent with the predictions for resonant Compton scattering in the magnetosphere, while the moderate polarization of the soft, thermal component ( $\approx 25\%$ ) may be produced by a condensed surface or a bombarded atmosphere.

Future polarimetric missions in hard X-rays (like PHEMTO and ASTROMEV; Laurent et al. 2021; De Angelis et al. 2021) will prove key in assessing the synchrotron/curvature origin of magnetar hard tails and shed light on the nature of the other spectral components.

1 We acknowledge financial support from INAF Funda-  
 2 mental Research Grants through the “Magnetars” Large  
 3 Program (PI S.Mereghetti). The work of RTa, RTu,  
 4 LM, and FM is partially supported by the PRIN grant  
 5 2022LWPEXW of the Italian Ministry of University and  
 6 Research (MUR). FCZ acknowledges support from a  
 7 Ramón y Cajal fellowship (grant agreement RYC2021-  
 8 030888-I). RK acknowledges support from the Science  
 9 and Technology Facilities Council (STFC) for funding  
 10 through a Ph.D. studentship. We thank the anonymous  
 11 referee for her/his helpful comments.

*Facilities:* IXPE, XMM-Newton (EPIC), Chandra (ACIS), NuSTAR.

*Software:* Astropy (Astropy Collaboration et al. 2013, 2018), `ixpeobssim` (<https://ixpeobssim.readthedocs.io/en>, Baldini et al. 2022), SAS (<https://www.cosmos.esa.int/web/xmm-newton/sas>), XSPEC (Arnaud 1996), CIAO (Fruscione et al. 2006).

## APPENDIX

### A. SNR KES 73


We report here the results of the spectral fit to the Chandra data of SNR Kes 73 (see §2.2 for details). The best fit parameters are listed in Table A1; the spectrum together with the best-fit model is shown in Figure A1.



**Table A1.** Fit parameters of SNR Kes 73 spectra

	SNR 6'' – 15''	SNR 6'' – 48''	SNR 6'' – 60''
$N_{\text{H}}$ ( $10^{22} \text{ cm}^{-2}$ )	$2.31^{+0.06}_{-0.05}$	$2.26^{+0.06}_{-0.04}$	$2.32^{+0.02}_{-0.03}$
$kT_s$ (keV)	$0.61^{+0.05}_{-0.04}$	$0.52^{+0.05}_{-0.03}$	$0.49^{+0.02}_{-0.01}$
$\tau_s$ ( $10^{10} \text{ cm}^{-3} \text{ s}$ )	$0.15^{+0.07}_{-0.08}$	$0.12^{+0.02}_{-0.02}$	$0.14^{+0.02}_{-0.01}$
$F_s^{2-8}$ ( $10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$ )	$0.25^{+0.02}_{-0.02}$	$1.21^{+0.14}_{-0.06}$	$1.72^{+0.06}_{-0.08}$
$kT_h$ (keV)	$3.2^{+0.2}_{-0.2}$	$2.04^{+0.04}_{-0.12}$	$1.79^{+0.04}_{-0.02}$
$\tau_h$ ( $10^{10} \text{ cm}^{-3} \text{ s}$ )	$3.9^{+0.4}_{-0.4}$	$4.3^{+0.3}_{-0.2}$	$4.9^{+0.2}_{-0.2}$
$F_h^{2-8}$ ( $10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$ )	$0.68^{+0.03}_{-0.03}$	$3.67^{+0.08}_{-0.12}$	$5.67^{+0.18}_{-0.04}$
Mg	1*	$1.38^{+0.06}_{-0.10}$	$1.35^{+0.05}_{-0.05}$
Si	1*	$1.58^{+0.05}_{-0.07}$	$1.61^{+0.05}_{-0.05}$
S	1*	$2.01^{+0.08}_{-0.11}$	$2.05^{+0.08}_{-0.08}$
$K$	1.19048	1.01587	1.01010
$F_{\text{TOT}}^{2-8}$ ( $10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$ )	$1.11^{+0.02}_{-0.02}$	$4.98^{+0.04}_{-0.03}$	$7.47^{+0.05}_{-0.04}$
$\chi^2/\text{dof}$	495.45/475	1537.00/1251	1813.83/1411

NOTE—The parameters marked with an asterisk are kept fixed in the fit. All elemental abundances are in solar units as given by [Anders & Grevesse \(1989\)](#). The fluxes, corrected for absorption, are in the 2–8 keV range;  $F_s$  and  $F_h$  are computed in the annular regions of radii 6'' – 15'', 6'' – 48'' and 6'' – 60'', while  $F_{\text{TOT}}$  is corrected by the corresponding scaling factor  $K$  to account for the area lost in the inner circles. Errors are at  $1\sigma$  cl level.



**Figure A1.** Best fit and residuals to the Chandra spectra of the SNR Kes 73. The three panels show the results for three annular extraction regions centered at (but excluding) 1E1841–045, as detailed in §2.2. The six observations, displayed in black, red, green, blue, cyan and magenta in increasing order of date, were jointly fit with two VPSHOCK components, shown as a solid (hot component) and dashed (soft component) black lines.

## REFERENCES

- An, H., Hascoët, R., Kaspi, V. M., et al. 2013, *ApJ*, 779, 163, doi: [10.1088/0004-637X/779/2/163](https://doi.org/10.1088/0004-637X/779/2/163)
- Anders, E., & Grevesse, N. 1989, *GeoCoA*, 53, 197, doi: [10.1016/0016-7037\(89\)90286-X](https://doi.org/10.1016/0016-7037(89)90286-X)
- Arnaud, K. A. 1996, in *Astronomical Society of the Pacific Conference Series*, Vol. 101, *Astronomical Data Analysis Software and Systems V*, ed. G. H. Jacoby & J. Barnes,

- Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, *A&A*, 558, A33, doi: [10.1051/0004-6361/201322068](https://doi.org/10.1051/0004-6361/201322068)
- Astropy Collaboration, Price-Whelan, A. M., Sipőcz, B. M., et al. 2018, *AJ*, 156, 123, doi: [10.3847/1538-3881/aabc4f](https://doi.org/10.3847/1538-3881/aabc4f)
- Bachetti, M. 2018, *Astrophysics Source Code Library*, ascl:1805.019
- Baldini, L., Bucciantini, N., Di Lalla, N., et al. 2022, *SoftwareX*, 19, 101194, doi: [10.1016/j.softx.2022.101194](https://doi.org/10.1016/j.softx.2022.101194)
- Baring, M. G., Gonthier, P. L., & Harding, A. K. 2005, *ApJ*, 630, 430, doi: [10.1086/431895](https://doi.org/10.1086/431895)
- Baring, M. G., & Harding, A. K. 2008, in *American Institute of Physics Conference Series*, Vol. 968, *Astrophysics of Compact Objects*, ed. Y.-F. Yuan, X.-D. Li, & D. Lai (AIP), 93–100, doi: [10.1063/1.2840459](https://doi.org/10.1063/1.2840459)
- Barthelmy, S. D., Kennea, J. A., Marshall, F. E., Maselli, A., & Sbarufatti, B. 2015, *GRB Coordinates Network*, 18024, 1
- Beloborodov, A. M. 2013, *ApJ*, 762, 13, doi: [10.1088/0004-637X/762/1/13](https://doi.org/10.1088/0004-637X/762/1/13)
- Caiazzo, I., González-Caniulef, D., Heyl, J., & Fernández, R. 2022, *MNRAS*, 514, 5024, doi: [10.1093/mnras/stac1571](https://doi.org/10.1093/mnras/stac1571)
- Coti Zelati, F., Rea, N., Pons, J. A., Campana, S., & Esposito, P. 2018, *MNRAS*, 474, 961, doi: [10.1093/mnras/stx2679](https://doi.org/10.1093/mnras/stx2679)
- De Angelis, A., Tatischeff, V., Argan, A., et al. 2021, *Experimental Astronomy*, 51, 1225, doi: [10.1007/s10686-021-09706-y](https://doi.org/10.1007/s10686-021-09706-y)
- den Hartog, P. R., Kuiper, L., & Hermsen, W. 2008a, *A&A*, 489, 263, doi: [10.1051/0004-6361:200809772](https://doi.org/10.1051/0004-6361:200809772)
- den Hartog, P. R., Kuiper, L., Hermsen, W., et al. 2008b, *A&A*, 489, 245, doi: [10.1051/0004-6361:200809390](https://doi.org/10.1051/0004-6361:200809390)
- Di Marco, A., Soffitta, P., Costa, E., et al. 2023, *AJ*, 165, 143, doi: [10.3847/1538-3881/acba0f](https://doi.org/10.3847/1538-3881/acba0f)
- Dib, R., & Kaspi, V. M. 2014, *ApJ*, 784, 37, doi: [10.1088/0004-637X/784/1/37](https://doi.org/10.1088/0004-637X/784/1/37)
- Duncan, R. C., & Thompson, C. 1992, *ApJL*, 392, L9, doi: [10.1086/186413](https://doi.org/10.1086/186413)
- Esposito, P., Rea, N., & Israel, G. L. 2021, in *Astrophysics and Space Science Library*, Vol. 461, *Timing Neutron Stars: Pulsations, Oscillations and Explosions*, ed. T. M. Belloni, M. Méndez, & C. Zhang, 97–142, doi: [10.1007/978-3-662-62110-3\\_3](https://doi.org/10.1007/978-3-662-62110-3_3)
- Fermi GBM Team. 2024, *GRB Coordinates Network*, 37234, 1
- Fernández, R., & Davis, S. W. 2011, *ApJ*, 730, 131, doi: [10.1088/0004-637X/730/2/131](https://doi.org/10.1088/0004-637X/730/2/131)
- Fruscione, A., McDowell, J. C., Allen, G. E., et al. 2006, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 6270, *Observatory Operations: Strategies, Processes, and Systems*, ed. D. R. Silva & R. E. Doxsey, 62701V, doi: [10.1117/12.671760](https://doi.org/10.1117/12.671760)
- GECAM team. 2024, *GRB Coordinates Network*, 37240, 1
- Gnedin, Y. N., & Pavlov, G. G. 1974, *Soviet Journal of Experimental and Theoretical Physics*, 38, 903
- Götz, D., Mereghetti, S., Tiengo, A., & Esposito, P. 2006, *A&A*, 449, L31, doi: [10.1051/0004-6361:20064870](https://doi.org/10.1051/0004-6361:20064870)
- Harding, A. K., & Lai, D. 2006, *Reports on Progress in Physics*, 69, 2631, doi: [10.1088/0034-4885/69/9/R03](https://doi.org/10.1088/0034-4885/69/9/R03)
- Harrison, F. A., Craig, W. W., Christensen, F. E., et al. 2013, *ApJ*, 770, 103, doi: [10.1088/0004-637X/770/2/103](https://doi.org/10.1088/0004-637X/770/2/103)
- Heyl, J., Taverna, R., Turolla, R., et al. 2024, *MNRAS*, 527, 12219, doi: [10.1093/mnras/stad3680](https://doi.org/10.1093/mnras/stad3680)
- Heyl, J. S., & Shaviv, N. J. 2002, *PhRvD*, 66, 023002, doi: [10.1103/PhysRevD.66.023002](https://doi.org/10.1103/PhysRevD.66.023002)
- Kaspi, V. M., & Beloborodov, A. M. 2017, *ARA&A*, 55, 261, doi: [10.1146/annurev-astro-081915-023329](https://doi.org/10.1146/annurev-astro-081915-023329)
- Kelly, R. M. E., González-Caniulef, D., Zane, S., Turolla, R., & Taverna, R. 2024a, *MNRAS*, 534, 1355, doi: [10.1093/mnras/stae2163](https://doi.org/10.1093/mnras/stae2163)
- Kelly, R. M. E., Zane, S., Turolla, R., & Taverna, R. 2024b, *MNRAS*, 528, 3927, doi: [10.1093/mnras/stae159](https://doi.org/10.1093/mnras/stae159)
- Kuiper, L., Hermsen, W., den Hartog, P. R., & Collmar, W. 2006, *ApJ*, 645, 556, doi: [10.1086/504317](https://doi.org/10.1086/504317)
- Kuiper, L., Hermsen, W., & Mendez, M. 2004, *ApJ*, 613, 1173, doi: [10.1086/423129](https://doi.org/10.1086/423129)
- Kumar, H. S., & Safi-Harb, S. 2010, *ApJL*, 725, L191, doi: [10.1088/2041-8205/725/2/L191](https://doi.org/10.1088/2041-8205/725/2/L191)
- Kumar, H. S., Safi-Harb, S., Slane, P. O., & Gotthelf, E. V. 2014, *ApJ*, 781, 41, doi: [10.1088/0004-637X/781/1/41](https://doi.org/10.1088/0004-637X/781/1/41)
- Laurent, P., Acero, F., Beckmann, V., et al. 2021, *Experimental Astronomy*, 51, 1143, doi: [10.1007/s10686-021-09723-x](https://doi.org/10.1007/s10686-021-09723-x)
- Mereghetti, S., Götz, D., Mirabel, I. F., & Hurley, K. 2005, *A&A*, 433, L9, doi: [10.1051/0004-6361:200500088](https://doi.org/10.1051/0004-6361:200500088)
- Molkov, S., Hurley, K., Sunyaev, R., et al. 2005, *A&A*, 433, L13, doi: [10.1051/0004-6361:200500087](https://doi.org/10.1051/0004-6361:200500087)
- Morii, M., Sato, R., Kataoka, J., & Kawai, N. 2003, *PASJ*, 55, L45, doi: [10.1093/pasj/55.3.L45](https://doi.org/10.1093/pasj/55.3.L45)
- NICER Team. 2024, *The Astronomer’s Telegram*, 16789, 1
- Olausen, S. A., & Kaspi, V. M. 2014, *ApJS*, 212, 6, doi: [10.1088/0067-0049/212/1/6](https://doi.org/10.1088/0067-0049/212/1/6)
- Radhakrishnan, V., & Cooke, D. J. 1969, *Astrophys. Lett.*, 3, 225
- Rybicki, G. B., & Lightman, A. P. 1979, *Radiative processes in astrophysics* (Wiley-Interscience)

- Slane, P., Ferrazzoli, R., Zhou, P., & Vink, J. 2024, *Galaxies*, 12, 59, doi: [10.3390/galaxies12050059](https://doi.org/10.3390/galaxies12050059)
- Stewart, R., Younes, G., Harding, A., et al. 2024, arXiv e-prints, arXiv:2412.16036, doi: [10.48550/arXiv.2412.16036](https://doi.org/10.48550/arXiv.2412.16036)
- Strüder, L., Briel, U., Dennerl, K., et al. 2001, *A&A*, 365, L18, doi: [10.1051/0004-6361:20000066](https://doi.org/10.1051/0004-6361:20000066)
- SVOM/GRM Team. 2024, GRB Coordinates Network, 37297, 1
- Swift Team. 2024a, GRB Coordinates Network, 37211, 1
- . 2024b, GRB Coordinates Network, 37222, 1
- Taverna, R., Muleri, F., Turolla, R., et al. 2014, *MNRAS*, 438, 1686, doi: [10.1093/mnras/stt2310](https://doi.org/10.1093/mnras/stt2310)
- Taverna, R., & Turolla, R. 2024, *Galaxies*, 12, 6, doi: [10.3390/galaxies12010006](https://doi.org/10.3390/galaxies12010006)
- Taverna, R., Turolla, R., Gonzalez Caniulef, D., et al. 2015, *MNRAS*, 454, 3254, doi: [10.1093/mnras/stv2168](https://doi.org/10.1093/mnras/stv2168)
- Taverna, R., Turolla, R., Suleimanov, V., Potekhin, A. Y., & Zane, S. 2020, *MNRAS*, 492, 5057, doi: [10.1093/mnras/staa204](https://doi.org/10.1093/mnras/staa204)
- Taverna, R., Turolla, R., Muleri, F., et al. 2022, *Science*, 378, 646, doi: [10.1126/science.add0080](https://doi.org/10.1126/science.add0080)
- Thompson, C., & Beloborodov, A. M. 2005, *ApJ*, 634, 565, doi: [10.1086/432245](https://doi.org/10.1086/432245)
- Thompson, C., & Duncan, R. C. 1993, *ApJ*, 408, 194, doi: [10.1086/172580](https://doi.org/10.1086/172580)
- Tian, W. W., & Leahy, D. A. 2008, *ApJ*, 677, 292, doi: [10.1086/529120](https://doi.org/10.1086/529120)
- Turner, M. J. L., Abbey, A., Arnaud, M., et al. 2001, *A&A*, 365, L27, doi: [10.1051/0004-6361:20000087](https://doi.org/10.1051/0004-6361:20000087)
- Turolla, R., Zane, S., & Watts, A. L. 2015, *Reports on Progress in Physics*, 78, 116901, doi: [10.1088/0034-4885/78/11/116901](https://doi.org/10.1088/0034-4885/78/11/116901)
- Turolla, R., Taverna, R., Israel, G. L., et al. 2023, *ApJ*, 954, 88, doi: [10.3847/1538-4357/aced05](https://doi.org/10.3847/1538-4357/aced05)
- Vasisht, G., & Gotthelf, E. V. 1997, *ApJL*, 486, L129, doi: [10.1086/310843](https://doi.org/10.1086/310843)
- Vogel, J. K., Hascoët, R., Kaspi, V. M., et al. 2014, *ApJ*, 789, 75, doi: [10.1088/0004-637X/789/1/75](https://doi.org/10.1088/0004-637X/789/1/75)
- Wachter, S., Patel, S. K., Kouveliotou, C., et al. 2004, *ApJ*, 615, 887, doi: [10.1086/424704](https://doi.org/10.1086/424704)
- Wadiasingh, Z., Baring, M. G., Gonthier, P. L., & Harding, A. K. 2018, *ApJ*, 854, 98, doi: [10.3847/1538-4357/aaa460](https://doi.org/10.3847/1538-4357/aaa460)
- Weisskopf, M. C., Guainazzi, M., Jahoda, K., et al. 2010, *ApJ*, 713, 912, doi: [10.1088/0004-637X/713/2/912](https://doi.org/10.1088/0004-637X/713/2/912)
- Weisskopf, M. C., Soffitta, P., Baldini, L., et al. 2022, *Journal of Astronomical Telescopes, Instruments, and Systems*, 8, 1, doi: [10.1117/1.JATIS.8.2.026002](https://doi.org/10.1117/1.JATIS.8.2.026002)
- Yang, C., Archibald, R. F., Vogel, J. K., et al. 2016, *ApJ*, 831, 80, doi: [10.3847/0004-637X/831/1/80](https://doi.org/10.3847/0004-637X/831/1/80)
- Younes, G., Hu, C. P., Enoto, T., et al. 2024, *The Astronomer's Telegram*, 16802, 1
- Zane, S., Taverna, R., González-Caniulef, D., et al. 2023, *ApJL*, 944, L27, doi: [10.3847/2041-8213/acb703](https://doi.org/10.3847/2041-8213/acb703)