

# TYPE II SINGULARITIES OF LAGRANGIAN MEAN CURVATURE FLOW WITH ZERO MASLOV CLASS

XIANG LI<sup>1</sup>, YONG LUO<sup>2</sup>, AND JUN SUN<sup>3</sup>

ABSTRACT. In this paper, we will prove some rigidity theorems for blow up limits to Type II singularities of Lagrangian mean curvature flow with zero Maslov class or almost calibrated Lagrangian mean curvature flows, especially for Lagrangian translating solitons in any dimension. These theorems generalized previous corresponding results from two dimensional case to arbitrarily dimensional case.

Keywords. Lagrangian mean curvature flow, translating solitons, zero Maslov class

## 1. INTRODUCTION

The existence of Special Lagrangians in a Calabi-Yau manifold is an interesting problem in differential geometry and mirror symmetry. The Special Lagrangians not only are minimal submanifolds, but also are calibrated submanifolds of Calabi-Yau manifolds ([HL82]). Therefore, the Special Lagrangians are area-minimizing in their homology class. In a compact Kähler-Einstein surface, Schoen-Wolfson ([SW01]) have studied the minimization problem and have shown the existence of a surface which minimizes area among Lagrangian competitors in each Lagrangian homology class, by variational methods. In addition to using variational methods to find minimal surfaces, another important idea is to use the mean curvature flow, which is the negative gradient flow for the area functional.

Smoczyk ([Smo96]) proved that “Lagrangian” condition is preserved by mean curvature flow, in which case the flow is called the Lagrangian mean curvature flow. It also follows from the maximum principle that “zero Maslov class” property or “almost calibrated” condition is preserved by the Lagrangian mean curvature flow, in which case the flow is called Lagrangian mean curvature flow with zero Maslov class, or almost calibrated Lagrangian mean curvature flow.

By establishing new monotonicity formula, Chen-Li ([CL04]) and Wang ([Wan01]) independently proved that there is no finite time Type I singularity for almost calibrated Lagrangian mean curvature flow. Neves ([Nev07]) proved the same conclusion for Lagrangian mean curvature flow with zero Maslov class. Furthermore, Neves constructed examples of Lagrangian mean curvature flow with zero Maslov class which develops finite time singularity at finite time ([Nev07], [Nev13]). Therefore, it is important to study blow up flows for Type II singularities of the Lagrangian mean curvature flow, which is a nonflat complete eternal solution to the Lagrangian mean curvature in  $\mathbb{C}^n$  with bounded second fundamental form, at most Euclidean area growth, and bounded Lagrangian angle (or almost calibrated).

In  $\mathbb{C}^2$ , Han-Li-Sun ([HLS11]) have studied the properties of the general limit flow  $\Sigma_s^\infty$  of the almost calibrated Lagrangian mean curvature flow. In particular, they provided an estimate for the norm of the mean curvature vector in terms of the oscillation of the Lagrangian angle for the limit flow. The first result in this paper is to generalize this result for Lagrangian mean curvature flow with zero Maslov class in any dimension  $n$ . Specifically, we have:

**Theorem 1.** *Suppose that  $\Sigma_t$  ( $t \in (-\infty, 0]$ ) is a complete proper Lagrangian mean curvature flow in  $\mathbb{C}^n$  with bounded Lagrangian angle. Assume further that  $\sup_{t \in (-\infty, 0]} \sup_{\Sigma_t} |A|^2 = 1$ . Then we have*

$$h^2 = \sup_{t \in (-\infty, 0]} \sup_{\Sigma_t} |H|^2 \leq \left( \sup_{t \in (-\infty, 0]} \sup_{\Sigma_t} \theta - \inf_{t \in (-\infty, 0]} \inf_{\Sigma_t} \theta \right)^2.$$

In the same paper, Han-Li-Sun ([HLS11]) also proved that an eternal solution to the almost calibrated Lagrangian mean curvature flow in  $\mathbb{C}^2$  which is flat at all time must be flat planes, hence cannot arise as blow up limit of almost calibrated Lagrangian mean curvature flow. By the Gauss equation, we can see that a surface is flat (i.e. the Gauss curvature vanishes everywhere) if and only if  $|H|^2 = |A|^2$  on  $\Sigma$ . Recently, Li-Sun ([LS24]) have shown that a stronger result that any almost calibrated Lagrangian eternal mean curvature flow in  $\mathbb{C}^2$  with  $\cos \theta \geq \delta > 0$  and  $|H|^2 \geq \varepsilon |A|^2$  with  $\varepsilon > 1 - \delta$  must be flat planes. Our next theorem generalizes it to arbitrary dimension  $n$ .

**Theorem 2.** *Any proper almost calibrated lagrangian eternal mean curvature flow in  $\mathbb{C}^n$  with  $\cos \theta \geq \delta > 0$  and  $|H|^2 \geq \varepsilon |A|^2$  with  $\varepsilon > 1 - \delta$  must be flat planes.*

As a consequence, we also have the following improvement in ([HLS11]), ([HS10]) and ([LS24]).

**Corollary 3.** *Any proper eternal mean curvature flow in  $\mathbb{C}^n$  with scalar curvature  $R_{\Sigma_t} \geq -(1 - \varepsilon)|A|^2$  for  $\varepsilon > 1 - \delta$  cannot arise as blow up flow of almost calibrated Lagrangian mean curvature flow.*

An important type of eternal solutions to the Lagrangian mean curvature flow are Lagrangian translating solitons. There are some nonexistence results and rigidity theorems on Lagrangian translating solitons (e.g., [HL09], [HS10], [NT13], [Qiu24], [Sun13, Sun14, Sun15], etc.).

Recall that a surface  $\Sigma^n$  in  $\mathbb{R}^{n+k}$  is called a translating soliton (or translator) of the mean curvature flow, if it satisfies

$$T^\perp = H,$$

where  $H$  is the mean curvature vector of  $\Sigma^n$  in  $\mathbb{R}^{n+k}$  and  $T$  is a constant vector in  $\mathbb{R}^{n+k}$ . Let  $V$  be the tangent part of  $T$ . Then we have

$$T = V + H.$$

In [Sun14], Sun proved that the infimum of  $|H|$  on any complete almost-calibrated Lagrangian translating soliton in  $\mathbb{C}^2$  with  $\cos \theta \geq \delta > 0$  and quadratic area growth must be zero. Our next result is to improve this result for complete Lagrangian translating soliton with zero Maslov class of any dimension:

**Theorem 4.** *Suppose  $\Sigma$  is a complete Lagrangian translating soliton with zero Maslov class in  $\mathbb{C}^n$ . Then:*

$$\inf_{\Sigma} |H|^2 = 0.$$

It is well known that a translating soliton can be viewed as a critical point of the functional

$$\mathcal{L}(\Sigma) = \int_{\Sigma} e^{\langle \mathbf{T}, \mathbf{x} \rangle} d\mu,$$

where  $\mathbf{x}$  is the position vector in  $\mathbb{R}^{n+k}$ , and  $d\mu$  is the volume form on  $\Sigma$  induced from the Euclidean space  $\mathbb{R}^{n+k}$ . For our convenience, we denote  $d\tilde{\mu} = e^{\langle \mathbf{T}, \mathbf{x} \rangle} d\mu$ .

It is easy to see that a complete translating soliton cannot be compact. Inspired by Xin ([Xin15]) and Sun's ([Sun15]) work, we have

**Theorem 5.** *Suppose  $\Sigma$  is a complete Lagrangian translating soliton with zero Maslov class in  $\mathbb{C}^n$  and mean curvature vector  $H \in L^1(d\tilde{\mu})$ . Then  $\Sigma$  must be a minimal Lagrangian submanifold.*

**Remark 6.** In Theorem 5 and Theorem 9 below, we can only prove that the translating solitons with suitable restrictions are minimal instead of flat plane. Actually, when  $n = 2$ , it is easy to see that a minimal Lagrangian translating soliton in  $\mathbb{C}^2$  must be a plane (see (7)). However, when  $n \geq 3$ , a minimal Lagrangian translating soliton in  $\mathbb{C}^n$  need not be a flat plane ([NT13]). For instance, if  $\Sigma$  is any nonflat Special Lagrangian in  $\mathbb{C}^2$ , then  $\tilde{\Sigma} := \mathbb{R} \times \Sigma \subset \mathbb{C}^3$  is minimal Lagrangian translating soliton but nonflat.

**Corollary 7.** *Any translating soliton with mean curvature vector  $H \in L^1(d\tilde{\mu})$  in  $\mathbb{C}^2$  cannot arise as blow up limit of the Lagrangian mean curvature flow with zero Maslov class.*

In order to state the next theorem, we give the definition of weighted polynomial area growth:

**Definition 8.** We say a submanifold  $\Sigma^n$  in  $\mathbb{R}^{n+k}$  has weighted polynomial area growth, if there is a constant  $D_0 > 0$  and  $d > 0$ , such that

$$(1) \quad \tilde{\mu}(\Sigma \cap B(r)) := \int_{\Sigma \cap B(r)} e^{\langle \mathbf{T}, \mathbf{x} \rangle} d\mu \leq D_0 r^d,$$

for any  $r \geq 1$  holds, where  $B(r)$  is the ball of radius  $r$  in  $\mathbb{R}^{n+k}$ .

Then we have

**Theorem 9.** *Suppose  $\Sigma$  is a complete Lagrangian translating soliton with zero Maslov class in  $\mathbb{C}^n$  and weighted polynomial area growth. Then  $\Sigma$  must be a minimal submanifold.*

As a corollary, we get that

**Corollary 10.** *Any translating soliton with weighted polynomial area growth in  $\mathbb{C}^2$  cannot arise as blow up limit of the Lagrangian mean curvature flow with zero Maslov class.*

By the monotonicity formula to the mean curvature flow, we know that the blowup limit of mean curvature flow must have polynomial area growth. However it is not clear whether the blowup limit has weighted polynomial area growth.

In the end of this section let us emphasize several significance of results and innovations of methodology in this paper. Firstly, all the results proved in this paper are true for any dimension instead of dimension  $n = 2$ . Actually, the dimension 2 is a crucial assumption in [HLS11] and [LS24]. For  $n = 2$ , they can use the weighted monotonicity formula to estimate the maximum of the mean curvature vector. These arguments are invalidated for  $n > 2$ . To overcome this difficulty, instead of deriving the estimate for the mean curvature vector, we choose a new cutoff functions to obtain the desired decay. Secondly, all the previous results are proved for almost calibrated Lagrangian translating solitons, where they used  $\frac{1}{\cos \theta}$  to construct the test function. In this paper, we choose to look for cutoff functions depending on  $\theta$  itself so that most of our results are true for Lagrangian translating solitons with zero Maslov class.

The subsequent sections are organized as follows: in Section 2, we provide some preliminary materials and in Section 3 and Section 4, we prove the main theorems.

## 2. NOTATIONS AND PRELIMINARIES

Let  $M^n$  be a Calabi-Yau manifold of complex dimensional  $n$  with a Kähler form  $\omega$ , a complex structure  $J$ , a Kähler metric  $g$  and a parallel holomorphic  $(n, 0)$ -form  $\Omega$  of unit length.

An immersed submanifold  $\Sigma$  in  $M$  is Lagrangian if  $\omega|_{\Sigma} = 0$ . The induced volume form  $d\mu_{\Sigma}$  on a Lagrangian submanifold  $\Sigma$  from the Ricci-flat metric  $g$  is related to  $\Omega$  by (see [HL82])

$$\Omega|_{\Sigma} = e^{i\theta} d\mu_{\Sigma},$$

where  $\theta$  is a multivalued function called the Lagrangian angle. When the Lagrangian angle is a single valued function the Lagrangian is called *zero-Maslov class* and if  $\cos \theta \geq \delta > 0$  for some positive  $\delta$ , then  $\Sigma$  is called *almost calibrated*. If  $\theta \equiv \text{constant}$ , then  $\Sigma$  is a *Special Lagrangian*.

Let  $F_0 : \Sigma^n \rightarrow M^{n+k}$  be a smooth immersion of a compact manifold  $\Sigma$ . The mean curvature flow with initial condition  $F_0$  is a smooth family of immersions  $F : \Sigma^n \times [0, T) \rightarrow M^{n+k}$  satisfying

$$\begin{cases} \frac{\partial}{\partial t} F(x, t) = H(x, t), \\ F(x, 0) = F_0(x), \end{cases}$$

where  $H(x, t) = H^{\alpha} \nu_{\alpha} = h_{i\bar{j}}^{\alpha} \nu_{\alpha}$  is the mean curvature vector of the submanifold  $\Sigma_t = F(\Sigma \cdot t)$  at  $x$ . The mean curvature flow is the negative gradient flow of the area functional. If the flow exists globally and converges at infinity, then the limit must be a minimal submanifold.

For a Lagrangian submanifold  $\Sigma$ , we have the following relation between mean curvature vector and the Lagrangian angle ([TY02]):

$$H = J\nabla\theta.$$

It then follows that

$$(2) \quad |\nabla\theta| = |H|.$$

Smoczyk ([Smo96]) proved that if the initial submanifold is Lagrangian, then along the mean curvature flow, it remain Lagrangian for positive time. In this case, the flow is called “Lagrangian mean curvature flow”. Smoczyk ([Smo99]) also derived the following equation for the Lagrangian angle along the mean curvature flow:

$$(3) \quad \left(\frac{\partial}{\partial t} - \Delta\right)\theta = 0$$

and thus

$$(4) \quad \left(\frac{\partial}{\partial t} - \Delta\right)\cos\theta = |H|^2\cos\theta.$$

As a result, if the initial surface is almost calibrated, i.e.,  $\cos\theta(\cdot, 0)$  has a positive lower bound, then by applying the parabolic maximum principle to this evolution equation, one concludes that  $\cos\theta$  remains positive as long as the mean curvature flow has a smooth solution, i.e., almost calibrated condition is preserved by mean curvature flow, in which case the flow is called “almost calibrated Lagrangian mean curvature flow”. (3) also implies that “zero Maslov class” property is preserved along the Lagrangian mean curvature flow. In this case the flow is called “Lagrangian mean curvature flow with zero Maslov class”.

From (3), it is easy to see that on Lagrangian translating solitons, we have:

$$(5) \quad \Delta\theta + \langle V, \nabla\theta \rangle = 0.$$

This implies that

$$(6) \quad -\Delta\cos\theta = |H|^2\cos\theta + V \cdot \nabla\cos\theta,$$

which will be used later.

For a two dimensional translating soliton in  $\mathbb{R}^4$ , Han-Li ([HL09]) computed the following identities on translating solitons:

$$(7) \quad |A|^2 = |H|^2 + 2\frac{|\nabla H|^2}{|V|^2} + \frac{V \cdot \nabla|H|^2}{|V|^2}.$$

This implies that two dimensional translating soliton in  $\mathbb{R}^4$  is a flat plane if and only if it is minimal.

Finally, we also have the following well-known evolution equation for the mean curvature vector along mean curvature flow in  $\mathbb{R}^{n+k}$  which will be used later (see, for example, ([AB10])):

$$(8) \quad \left(\frac{\partial}{\partial t} - \Delta\right)|H|^2 = -2|\nabla^\perp H|^2 + 2\sum_{i,j,\alpha,\beta} H^\alpha H^\beta h_{ij}^\alpha h_{ij}^\beta.$$

### 3. RIGIDITY THEOREMS FOR BLOW UP FLOWS OF LAGRANGIAN MEAN CURVATURE FLOWS

In this section, we will prove the first two main theorems for blow up flows of Lagrangian mean curvature flows.

**Proof of Theorem 1:** Without loss of generality, we may assume

$$\inf_{t \in (-\infty, 0]} \inf_{\Sigma_t} \theta = 0.$$

If  $h = 0$  or  $\eta := \sup_{t \in (-\infty, 0]} \sup_{\Sigma_t} \theta = 0$ , it is evident that the result holds. Now we assume that  $h > 0$ ,  $\eta > 0$ .

We introduce a new function  $f(X, t) = |H|^2 + p\theta^2$ , where  $p > 1$ ,  $t \in [-T, 0]$ . Using the evolution equation (3) for  $\theta$  and the evolution equation (8) for  $|H|^2$ , we get

$$\begin{aligned} (\Delta - \frac{\partial}{\partial t})f &\geq 2|\nabla^\perp H|^2 - 2|H|^2|A|^2 + 2p|\nabla\theta|^2 \\ &\geq 2(p-1)|H|^2. \end{aligned}$$

Here, we have used the assumption  $\sup_{t \in (-\infty, 0]} \sup_{\Sigma_t} |A|^2 = 1$  and the fact (2).

Let  $\psi(x)$  be a  $C^2$  function on  $[0, \infty)$  such that

$$\psi(x) = \begin{cases} 1, & \text{if } 0 \leq x \leq \frac{1}{2}, \\ 0, & \text{if } x \geq 1, \end{cases}$$

$$0 \leq \psi(x) \leq 1, \quad \psi'(x) \leq 0, \quad |\psi''(x)| \leq C \quad \text{and} \quad \frac{|\psi'(x)|^2}{\psi(x)} \leq C,$$

where  $C$  is an absolute constant.

We construct a cutoff function  $g(X, t) = \psi(\frac{|X|^2}{R^2})\psi(\frac{t}{T})$ . A straightforward computation shows that

$$\begin{aligned} (\Delta - \frac{\partial}{\partial t})g &= 4\psi''(\frac{|X|^2}{R^2})\psi(\frac{t}{T})\frac{\langle X, \nabla X \rangle^2}{R^4} + \psi(\frac{t}{T})\frac{2n\psi'(\frac{|X|^2}{R^2})}{R^2} - \frac{1}{T}\psi'(\frac{t}{T})\psi(\frac{|X|^2}{R^2}) \\ &\geq -\frac{C}{R^2} - \frac{C}{T} \end{aligned}$$

and

$$\frac{|\nabla g|^2}{g} \leq \frac{C}{R^2}.$$

Set  $\phi = gf$ . Since  $\phi|_{\partial(\bar{B}_R \times [-T, 0])} = 0$ , then  $\phi$  must achieve its maximum at some points  $(x_i, t_i) \in B_R \times [-T, 0]$ . By the maximum value principle, at  $(x_i, t_i)$ , we obtain that

$$\frac{\partial}{\partial t}\phi(x_i, t_i) \geq 0,$$

$$\nabla\phi(x_i, t_i) = 0,$$

$$\Delta\phi(x_i, t_i) \leq 0,$$

i.e.,

$$(\Delta - \frac{\partial}{\partial t})\phi = (\Delta - \frac{\partial}{\partial t})gf \leq 0$$

and

$$\nabla g = -\frac{g}{f}\nabla f.$$

Hence, we get

$$\begin{aligned}
0 &\geq (\Delta - \frac{\partial}{\partial t})gf \\
&= f(\Delta - \frac{\partial}{\partial t})g + g(\Delta - \frac{\partial}{\partial t})f + 2\nabla g \nabla f \\
&\geq -\frac{C}{R^2}f - \frac{C}{T}f + 2(p-1)g|H|^2 - \frac{|\nabla g|^2}{g}f \\
&\geq -\frac{C}{R^2}f - \frac{C}{T}f + 2(p-1)g|H|^2.
\end{aligned}$$

Since  $p > 1$  and  $f$  is bounded, we get

$$g|H|^2(x_i, t_i) \leq \frac{C}{(p-1)R^2} + \frac{C}{(p-1)T}.$$

Therefore,

$$\sup_{B_{\frac{R}{2}} \times [-T, 0]} f(x_i, t_i) \leq \frac{C}{(p-1)R^2} + \frac{C}{(p-1)T} + p \sup_{B_R \times [-T, 0]} \theta^2.$$

Letting  $R \rightarrow \infty$  and  $T \rightarrow \infty$ , we obtain

$$h^2 \leq p\eta^2.$$

Letting  $p \rightarrow 1$ , we get the desired inequality.  $\square$

**Proof of Theorem 2:** We first prove that the eternal flow is minimal under the assumption that

$$(9) \quad |H|^2 \geq \varepsilon|A|^2,$$

where  $\varepsilon > 1 - \delta$ . We prove it by contradiction.

As explained earlier, we suppose on the contrary that the eternal solution is not minimal. Without loss of generality, we may assume that  $|H|(0, 0) = 1$ . Let  $f$  be a positive increasing function which is to be determined later. We compute  $(\Delta - \frac{\partial}{\partial t})(|H|^2 f(\frac{1}{\cos \theta}))$ . First note that by (4),

$$\begin{aligned}
(10) \quad (\Delta - \frac{\partial}{\partial t}) \frac{1}{\cos \theta} &= -\frac{(\Delta - \frac{\partial}{\partial t}) \cos \theta}{\cos^2 \theta} + 2 \cos \theta |\nabla \frac{1}{\cos \theta}|^2 \\
&= \frac{|H|^2}{\cos \theta} + 2 \cos \theta |\nabla \frac{1}{\cos \theta}|^2.
\end{aligned}$$

On the other hand, the equality (8) combining with Cauchy-Schwarz inequality yields

$$\begin{aligned}
(11) \quad (\Delta - \frac{\partial}{\partial t})|H|^2 &= 2|\nabla^\perp H|^2 - 2 \sum_{i,j,\alpha,\beta} H^\alpha H^\beta h_{ij}^\alpha h_{ij}^\beta \\
&\geq 2|\nabla^\perp H|^2 - 2|A|^2|H|^2.
\end{aligned}$$

We calculate

$$\begin{aligned}
(\Delta - \frac{\partial}{\partial t})(|H|^2 f(\frac{1}{\cos \theta})) &= (\Delta - \frac{\partial}{\partial t})|H|^2 f(\frac{1}{\cos \theta}) + |H|^2 (\Delta - \frac{\partial}{\partial t})(f(\frac{1}{\cos \theta})) \\
&\quad + 2\nabla |H|^2 \cdot \nabla f(\frac{1}{\cos \theta}) \\
&\geq f(\frac{1}{\cos \theta})(2|\nabla^\perp H|^2 - 2|A|^2|H|^2)
\end{aligned}$$

$$\begin{aligned}
(12) \quad & + |H|^2 \left( f' \frac{|H|^2}{\cos \theta} + 2f' \cos \theta \left| \nabla \frac{1}{\cos \theta} \right|^2 + f'' \left| \nabla \frac{1}{\cos \theta} \right|^2 \right) \\
& + 2 \frac{\nabla(f|H|^2) - |H|^2 \nabla f}{f} \cdot \nabla f \left( \frac{1}{\cos \theta} \right) \\
& = |H|^2 f \left( 2 \frac{|\nabla^\perp H|^2}{|H|^2} - 2|A|^2 + \frac{f'}{f} \frac{|H|^2}{\cos \theta} \right) \\
& + |H|^2 \left( f'' - 2 \frac{(f')^2}{f} + 2f' \cos \theta \right) \left| \nabla \frac{1}{\cos \theta} \right|^2 \\
& + 2|H|^2 \frac{\nabla(f|H|^2)}{f|H|^2} \cdot \nabla f \left( \frac{1}{\cos \theta} \right).
\end{aligned}$$

Set  $\phi = f|H|^2$ . At the point where  $\phi \neq 0$ , it is easy to see that

$$\nabla \phi = f \nabla |H|^2 + |H|^2 f' \nabla \frac{1}{\cos \theta},$$

i.e.,

$$(13) \quad \nabla \frac{1}{\cos \theta} = \frac{f}{f'} \left( \frac{\nabla \phi}{\phi} - \frac{\nabla |H|^2}{|H|^2} \right).$$

Plugging (13) into (12), we obtain

$$\begin{aligned}
(14) \quad & \left( \Delta - \frac{\partial}{\partial t} \right) \phi \geq \phi \left( 2 \frac{|\nabla^\perp H|^2}{|H|^2} - 2|A|^2 + \frac{f'}{f} \frac{|H|^2}{\cos \theta} \right) \\
& + \frac{\phi f}{(f')^2} \left( f'' - 2 \frac{(f')^2}{f} + 2f' \cos \theta \right) \left( \frac{|\nabla |H|^2|^2}{|H|^4} - 2 \frac{\nabla |H|^2}{|H|^2} \cdot \frac{\nabla \phi}{\phi} + \frac{|\nabla \phi|^2}{\phi^2} \right) \\
& + 2|H|^2 f' \frac{\nabla \phi}{\phi} \nabla \frac{1}{\cos \theta} \\
& = \phi \left( \frac{f'}{f} \frac{|H|^2}{\cos \theta} - 2|A|^2 \right) + \phi \left( 2 \frac{|\nabla^\perp H|^2}{|H|^2} + 4 \frac{f f''}{(f')^2} \frac{|\nabla |H|^2|^2}{|H|^2} - 8 \frac{|\nabla |H|^2|^2}{|H|^2} \right. \\
& \left. + 8 \frac{f}{f'} \cos \theta \frac{|\nabla |H|^2|^2}{|H|^2} \right) + \phi \left( \frac{f f''}{(f')^2} + 2 \frac{f}{f'} \cos \theta - 2 \right) \left( \frac{|\nabla \phi|^2}{\phi^2} - 2 \frac{\nabla |H|^2}{|H|^2} \cdot \frac{\nabla \phi}{\phi} \right) \\
& + 2|H|^2 f' \frac{\nabla \phi}{\phi} \nabla \frac{1}{\cos \theta} \\
& \geq \phi \left( \frac{f'}{f} \frac{|H|^2}{\cos \theta} - 2|A|^2 \right) + \phi \left( 4 \frac{f f''}{(f')^2} + 8 \frac{f}{f'} \cos \theta - 6 \right) \frac{|\nabla |H|^2|^2}{|H|^2} \\
& + \phi \left( \frac{f f''}{(f')^2} + 2 \frac{f}{f'} \cos \theta - 2 \right) \left( \frac{|\nabla \phi|^2}{\phi^2} - 2 \frac{\nabla |H|^2}{|H|^2} \cdot \frac{\nabla \phi}{\phi} \right) \\
& + 2|H|^2 f' \frac{\nabla \phi}{\phi} \nabla \frac{1}{\cos \theta} \\
& = \phi \left( \frac{f'}{f} \frac{|H|^2}{\cos \theta} - 2|A|^2 \right) + \phi \left( 4 \frac{f f''}{(f')^2} + 8 \frac{f}{f'} \cos \theta - 6 \right) \frac{|\nabla |H|^2|^2}{|H|^2} \\
& - \phi \left( \frac{f f''}{(f')^2} + 2 \frac{f}{f'} \cos \theta - 2 \right) \left( \frac{|\nabla \phi|^2}{\phi^2} - 2 \frac{f'}{f} \nabla \frac{1}{\cos \theta} \cdot \frac{\nabla \phi}{\phi} \right) \\
& + 2|H|^2 f' \frac{\nabla \phi}{\phi} \nabla \frac{1}{\cos \theta},
\end{aligned}$$

where we used the Kato's inequality

$$|\nabla|H||^2 \leq |\nabla^\perp H|^2.$$

Set  $\frac{f'}{f} = g$ . We choose  $g$  such that for  $x \in [1, \frac{1}{\delta}]$

$$(15) \quad \begin{cases} \varepsilon x/g > 2, \\ -4g' + 8g/x - 2 = 0, \end{cases}$$

where  $\varepsilon$  is a fixed constant. Note that  $x$  in this place represents the rang of Lagrangian angle. Let  $g(x) = \frac{x}{2}c(x)$ , then  $c(x)$  need satisfy

$$\begin{cases} 0 < c(x) < \varepsilon, \\ -xc' = 1 - c. \end{cases}$$

We choose  $c(x) = 1 - \eta x$  by solving the last equation. Then we have that

$$1 - \frac{\eta}{\delta} \leq c(x) \leq 1 - \eta.$$

Therefore, the system (15) has a solution if and only if  $1 - \varepsilon < \eta < \delta$ . Hence, our assumption that  $\varepsilon > 1 - \delta$  implies that we can choose

$$\eta = \frac{1 - \varepsilon + \delta}{2}.$$

Wth this choice of  $\eta$ , we have

$$g = \frac{x}{2}(1 - \eta x)$$

and

$$f(x) = \frac{x^2}{(1 - \eta x)^2}, \quad x \in [1, \frac{1}{\delta}].$$

It is evident that for  $x \in [1, \frac{1}{\delta}]$ ,

$$\frac{1}{(1 - \eta)^2} \leq f(x) \leq \frac{1}{(\delta - \eta)^2}.$$

On the other hand, by our assumption(9), we know from (14) that

$$(16) \quad \begin{aligned} (\Delta - \frac{\partial}{\partial t})\phi &\geq 2\phi|A|^2\left(\frac{\varepsilon}{1 - \frac{\eta}{\cos\theta}} - 1\right) + \frac{\phi}{2}\left(\frac{|\nabla\phi|^2}{\phi^2} - 2\frac{f'}{f}\nabla\frac{1}{\cos\theta} \cdot \frac{\nabla\phi}{\phi}\right) \\ &\quad + 2|H|^2f'\frac{\nabla\phi}{\phi} \cdot \nabla\frac{1}{\cos\theta} \\ &\geq \frac{2(\varepsilon + \eta - 1)}{1 - \eta}\phi|A|^2 + \frac{|\nabla\phi|^2}{2\phi} - \left(\phi\frac{f'}{f}\nabla\frac{1}{\cos\theta} - 2|H|^2f'\nabla\frac{1}{\cos\theta}\right) \cdot \frac{\nabla\phi}{\phi} \\ &\geq \frac{\varepsilon + \delta - 1}{1 + \varepsilon - \delta}\phi|H|^2 - \mathbf{b} \cdot \frac{\nabla\phi}{\phi}, \end{aligned}$$

where  $\mathbf{b} = \phi\frac{f'}{f}\nabla\frac{1}{\cos\theta} - 2|H|^2f'\nabla\frac{1}{\cos\theta}$ .

We claim that  $\mathbf{b}$  is bounded on  $\Sigma$ . Indeed, the definition of  $f$  guarantees that  $f'$  and  $\frac{f'}{f}$  are uniformly bounded by a constant depending only on  $\delta$ . Therefore we only need to control  $\frac{\nabla\cos\theta}{\cos^2\theta} = \frac{-\sin\theta}{\cos^2\theta}\nabla\theta$ . But by our assumption,  $\cos\theta$  is bounded below by  $\delta$ . Thus, (2) gives the upper bound of  $\frac{\nabla\cos\theta}{\cos^2\theta}$  because of the boundedness of  $|A|^2$ .

Assume  $|H|^2$  is not identically zero, we may assume  $\phi(0, 0) = 1 > 0$ .

Let  $\psi(x)$  be a  $C^2$  function on  $[0, \infty)$  such that

$$\psi(x) = \begin{cases} 1, & \text{if } 0 \leq x \leq \frac{1}{2}, \\ 0, & \text{if } x \geq 1, \end{cases}$$

$$0 \leq \psi(x) \leq 1, \quad \psi'(x) \leq 0, \quad |\psi''(x)| \leq C \quad \text{and} \quad \frac{|\psi'(x)|^2}{\psi(x)} \leq C,$$

where  $C$  is an absolute constant.

We construct a cutoff function  $h(X, t) = \psi(\frac{|X|^2}{R^2})\psi(\frac{t}{T})$ .

Now we consider the function  $F$  on  $B_R(0) \times [s_i, 0]$  defined by

$$F(x, t) = h\phi(x, t).$$

Using the fact that  $|\nabla X|^2 = n$ , a straightforward computation shows that

$$\begin{aligned} (\Delta - \frac{\partial}{\partial t})h &= 4\psi''(\frac{|X|^2}{R^2})\psi(\frac{t}{T})\frac{\langle X, \nabla X \rangle^2}{R^4} + \psi(\frac{t}{T})\frac{2n\psi'(\frac{|X|^2}{R^2})}{R^2} - \frac{1}{T}\psi'(\frac{t}{T})\psi(\frac{|X|^2}{R^2}) \\ &\geq -\frac{C}{R^2} - \frac{C}{T} \end{aligned}$$

$$(17) \quad \frac{|\nabla h|^2}{h} \leq \frac{C}{R^2}.$$

Since  $F|_{\partial(\overline{B_R}) \times [s_i, 0]} = 0$ , then  $F$  must achieve its maximum at some  $(y, t) \in B_R(0) \times [s_i, 0]$ .

By the maximum principle, at  $(y, t)$ , we obtain that

$$\begin{aligned} \frac{\partial}{\partial t}F(y, t) &\geq 0, \\ \nabla F(y, t) &= 0, \\ \Delta F(y, t) &\leq 0. \end{aligned}$$

Hence,

$$\begin{aligned} (\Delta - \frac{\partial}{\partial t})F &= (\Delta - \frac{\partial}{\partial t})h\phi \leq 0, \\ \nabla h &= -\frac{h}{\phi}\nabla\phi. \end{aligned}$$

Using (16) and (17), we obtain

$$\begin{aligned} (18) \quad 0 &\geq (\Delta - \frac{\partial}{\partial t})(h\phi) \\ &= \phi(\Delta - \frac{\partial}{\partial t})h + h(\Delta - \frac{\partial}{\partial t})\phi + 2\nabla h \cdot \nabla\phi \\ &\geq -\frac{C}{R^2}\phi - \frac{C}{T}\phi + \frac{\varepsilon + \delta - 1}{1 + \varepsilon - \delta}\phi|H|^2h - \mathbf{b} \cdot \frac{\nabla\phi}{\phi}h + 2\nabla h \cdot (-\frac{\phi}{h})\nabla h \\ &= -\frac{C}{R^2}\phi - \frac{C}{T}\phi + \frac{\varepsilon + \delta - 1}{1 + \varepsilon - \delta}\phi|H|^2h + \mathbf{b} \cdot \nabla h - 2\frac{\phi}{h}|\nabla h|^2 \\ &\geq -\frac{C}{R^2}\phi - \frac{C}{T}\phi + \frac{\varepsilon + \delta - 1}{1 + \varepsilon - \delta}\phi|H|^2h - |\mathbf{b}|\frac{C}{R} - 2\frac{C}{R^2}\phi \\ &\geq \frac{\varepsilon + \delta - 1}{1 + \varepsilon - \delta}|H|^2 - \frac{C}{R} - \frac{C}{R^2} - \frac{C}{T}, \end{aligned}$$

where  $C$  is constant which depend only on  $\delta$  and the bound of  $|A|^2$ . By (18), it implies that

$$\frac{\varepsilon + \delta - 1}{1 + \varepsilon - \delta} |H|^2(y, t) \leq \frac{C}{R} + \frac{C}{R^2} + \frac{C}{T},$$

which gives us the desired contradiction by letting  $R \rightarrow \infty$  and  $T \rightarrow \infty$ . This proves that the eternal solution satisfies

$$|H|^2 \equiv 0,$$

which implies that eternal mean curvature flow  $\tilde{\Sigma}$  is a minimal surface. Then by our assumption (9), we know  $|A| = 0$ . This finishes the proof of the theorem.  $\square$

#### 4. RIGIDITY THEOREMS FOR LAGRANGIAN TRANSLATING SOLITON WITH ZERO MASLOV CLASS

In this section, we will prove the rigidity theorems for Lagrangian translating soliton with zero Maslov class .

**Proof of Theorem 4:** We prove it by contradiction. Suppose there is a complete Lagrangian translating soliton  $\Sigma$  in  $\mathbb{C}^n$  with  $-D_1 \leq \theta \leq D_2$ ,  $|T| = 1$  and

$$(19) \quad \inf_{\Sigma} |H|^2 := a > 0,$$

where  $a \in (0, \frac{1}{2})$ . Set  $u = \theta^2$ , then by (5), we can easily see that

$$(20) \quad \Delta u = 2|H|^2 - \langle V \cdot \nabla u \rangle,$$

where  $\Delta$  and  $\nabla$  are the Laplacian and gradient operator on  $\Sigma$  with respect to the induced metric, respectively.

Let  $\phi$  be any cutoff function and  $p$  be a positive number to be determined later. Multiplying both sides of (20) by  $\phi^2 u^p$  and integrating by parts yields

$$\begin{aligned} & 2 \int_{\Sigma} \phi^2 u^p |H|^2 d\mu - \int_{\Sigma} \phi^2 u^p \langle V \cdot \nabla u \rangle d\mu \\ &= \int_{\Sigma} \phi^2 u^p \Delta u d\mu = -p \int_{\Sigma} \phi^2 u^{p-1} |\nabla u|^2 d\mu - 2 \int_{\Sigma} \phi u^p \nabla \phi \cdot \nabla u d\mu \end{aligned}$$

Rearranging this equality and using Young's inequality, we obtain

$$\begin{aligned} & 2 \int_{\Sigma} \phi^2 u^p |H|^2 d\mu + p \int_{\Sigma} \phi^2 u^{p-1} |\nabla u|^2 d\mu \\ &= \int_{\Sigma} \phi^2 u^p \langle V \cdot \nabla u \rangle d\mu - 2 \int_{\Sigma} \phi u^p \nabla \phi \cdot \nabla u d\mu \\ &\leq \int_{\Sigma} \phi^2 u^p |V| |\nabla u| d\mu + 2 \int_{\Sigma} \phi u^p |\nabla \phi| |\nabla u| d\mu \\ &\leq \varepsilon \int_{\Sigma} \phi^2 u^{p+1} |V|^2 d\mu + \frac{1}{4\varepsilon} \int_{\Sigma} \phi^2 u^{p-1} |\nabla u|^2 d\mu \\ &\quad + 2 \int_{\Sigma} \phi u^p |\nabla \phi| |\nabla u| d\mu, \end{aligned}$$

which implies that

$$(21) \quad \begin{aligned} & \int_{\Sigma} \phi^2 u^p (2|H|^2 - \varepsilon u|V|^2) d\mu + (p - \frac{1}{4\varepsilon}) \int_{\Sigma} \phi^2 u^{p-1} |\nabla u|^2 d\mu \\ & \leq 2 \int_{\Sigma} \phi u^p |\nabla \phi| |\nabla u| d\mu. \end{aligned}$$

From  $1 = |T|^2 = |H|^2 + |V|^2$ ,  $-D_1 \leq \theta \leq D_2$  and (19), we get that

$$2|H|^2 - \varepsilon u|V|^2 \geq 2a - \varepsilon D^2(1-a) \geq 2a - \varepsilon D^2,$$

where  $D = \max\{D_1, D_2\}$ . We first choose  $\varepsilon = \frac{a}{D^2}$ , then

$$2|H|^2 - \varepsilon D^2 |V|^2 \geq a.$$

Then we obtain from (21) that

$$\begin{aligned} & a \int_{\Sigma} \phi^2 u^p d\mu + (p - \frac{D^2}{4a}) \int_{\Sigma} \phi^2 u^{p-1} |\nabla u|^2 d\mu \\ & \leq 2 \int_{\Sigma} \phi u^p |\nabla \phi| |\nabla u| d\mu. \end{aligned}$$

Next, we choose  $p = \frac{D^2}{4a} + 1 := P_0$ , then

$$(22) \quad a \int_{\Sigma} \phi^2 u^{P_0} d\mu + \int_{\Sigma} \phi^2 u^{P_0-1} |\nabla u|^2 d\mu \leq 2 \int_{\Sigma} \phi u^{P_0} |\nabla \phi| |\nabla u| d\mu.$$

Now we take a smooth function  $\phi(x) : \mathbb{R} \rightarrow \mathbb{R}$  supported on  $[-R, R]$  which has the property:  $\phi = 1$  on  $[-\frac{R}{2}, \frac{R}{2}]$  and there is a constant  $C_b > 0$  such that  $|\nabla \phi| \leq C_b \phi^b \frac{1}{R}$  for  $0 < b < 1$ .

From Huisken's monotonicity formula, we see that the blow-up limit of Lagrangian mean curvature flow always has polynomial area growth (e.g., [Col12]). By using Young's inequality and the property of  $\phi$ , we can estimate  $\int_{\Sigma} \phi u^{P_0} |\nabla \phi| |\nabla u| d\mu$  as follows:

$$(23) \quad \begin{aligned} \int_{\Sigma} \phi u^{P_0} |\nabla \phi| |\nabla u| d\mu & \leq \frac{C_b}{R} \int_{\Sigma} \phi^{1+b} u^{P_0} |\nabla u| d\mu \\ & \leq 2D \frac{C_b}{R} \int_{\Sigma} \phi^{1+b} u^{\frac{P_0(1+b)}{2}} u^{\frac{P_0(1-b)}{2}} d\mu \\ & \leq \frac{a}{4} \int_{\Sigma} \phi^2 u^{P_0} d\mu + \frac{C}{R^{\frac{2}{1-b}}} \int_{\Sigma \cap \text{supp} \phi} u^{P_0} d\mu \\ & \leq \frac{a}{4} \int_{\Sigma} \phi^2 u^{P_0} d\mu + CD^{2P_0} R^{n-\frac{2}{1-b}}, \end{aligned}$$

where we used the fact  $|\nabla u| = |\nabla \theta|^2 = 2|\theta| |\nabla \theta| \leq 2D$ . Finally we choose  $b > 0$  such that  $n - \frac{2}{1-b} < 0$ , i.e.,  $b > 1 - \frac{2}{n}$ . Using (23) in (22), we have

$$\frac{a}{2} \int_{\Sigma} \phi^2 d\mu \leq C(a, D, n) R^{n-\frac{2}{1-b}}.$$

This gives the desired contradiction as  $R \rightarrow \infty$ .  $\square$

**Proof of Theorem 5:** Let  $\phi$  be any cutoff function and  $p$  be a positive number to be determined later. Multiplying both sides of (20) by  $\phi^2 u e^{(T, \mathbf{x})}$  and integrating

by parts yields

$$\begin{aligned}
2 \int_{\Sigma} \phi^2 u |H|^2 e^{\langle \mathbf{T}, \mathbf{x} \rangle} d\mu &= \int_{\Sigma} \phi^2 u (\Delta u + \langle \mathbf{V}, \nabla u \rangle) e^{\langle \mathbf{T}, \mathbf{x} \rangle} d\mu \\
&= \int_{\Sigma} \phi^2 u \operatorname{div}_{\Sigma}(e^{\langle \mathbf{T}, \mathbf{x} \rangle} \nabla u) d\mu \\
&= - \int_{\Sigma} \langle \nabla(\phi^2 u), \nabla u \rangle e^{\langle \mathbf{T}, \mathbf{x} \rangle} d\mu \\
&= - \int_{\Sigma} \phi^2 |\nabla u|^2 e^{\langle \mathbf{T}, \mathbf{x} \rangle} d\mu \\
&\quad - 2 \int_{\Sigma} \phi u \langle \nabla \phi, \nabla u \rangle e^{\langle \mathbf{T}, \mathbf{x} \rangle} d\mu
\end{aligned}$$

which implies that

$$(24) \quad \int_{\Sigma} \phi^2 (2u|H|^2 + |\nabla u|^2) e^{\langle \mathbf{T}, \mathbf{x} \rangle} d\mu \leq 2 \int_{\Sigma} \phi u |\nabla \phi| |\nabla u| e^{\langle \mathbf{T}, \mathbf{x} \rangle} d\mu.$$

Since  $-D_1 \leq \theta \leq D_2$ , we see that  $\theta \leq D := \max\{D_1, D_2\}$ . Furthermore, from (2), we have

$$(25) \quad |\nabla u| = |\nabla \theta^2| = 2|\theta| |\nabla \theta| \leq 2D|H|.$$

Therefore, by (24), we have

$$\begin{aligned}
(26) \quad \int_{\Sigma} \phi^2 (2u|H|^2 + |\nabla u|^2) e^{\langle \mathbf{T}, \mathbf{x} \rangle} d\mu &\leq C(D) \int_{\Sigma} \phi |\nabla \phi| |H| e^{\langle \mathbf{T}, \mathbf{x} \rangle} d\mu \\
&= C(D) \int_{\Sigma} \phi |\nabla \phi| |H| d\tilde{\mu}.
\end{aligned}$$

Now for any fixed  $R > 0$ , we take the cutoff function  $\phi = \phi_R$  such that  $\phi \in C_0^\infty(B(2R))$ ,  $\phi = 1$  on  $B(R)$ , and  $|\nabla \phi| \leq |D\phi| \leq \frac{C}{R}$ . Here,  $B(R)$  is the ball of radius in  $R^n$ ,  $D\phi$  is the gradient of  $\phi$  with respect to the Euclidean metric in  $R^n$ , and  $C$  is an absolute constant. Taking  $\phi = \phi_R$  in (26) yields

$$\begin{aligned}
(27) \quad \int_{\Sigma} \phi^2 (2u|H|^2 + |\nabla u|^2) e^{\langle \mathbf{T}, \mathbf{x} \rangle} d\mu &\leq C(D) \int_{\Sigma} \phi |\nabla \phi| |H| d\tilde{\mu} \\
&\leq \frac{C(D)}{R} \int_{\Sigma} |H| d\tilde{\mu}.
\end{aligned}$$

By our assumption,  $\int_{\Sigma} |H| d\tilde{\mu} < \infty$ . Letting  $R \rightarrow \infty$  in (27), we finally obtain that  $H \equiv 0$  on  $\Sigma$ . This means that  $\Sigma$  is a minimal submanifold.

When  $n = 2$ , by (7), we see that  $A \equiv 0$  on  $\Sigma$ . Thus it must be a flat plane.  $\square$

**Proof of Theorem 9:**

Let  $\phi$  be any cutoff function and  $p$  be a positive number to be determined later. Multiplying both sides of (20) by  $\phi^2 u e^{\langle \mathbf{T}, \mathbf{x} \rangle}$  and integrating by parts yields

$$\begin{aligned} 2 \int_{\Sigma} \phi^2 u |H|^2 e^{\langle \mathbf{T}, \mathbf{x} \rangle} d\mu &= \int_{\Sigma} \phi^2 u (\Delta u + \langle V, \nabla u \rangle) e^{\langle \mathbf{T}, \mathbf{x} \rangle} d\mu \\ &= \int_{\Sigma} \phi^2 u \operatorname{div}_{\Sigma}(e^{\langle \mathbf{T}, \mathbf{x} \rangle} \nabla u) d\mu \\ &= - \int_{\Sigma} \langle \nabla(\phi^2 u), \nabla u \rangle e^{\langle \mathbf{T}, \mathbf{x} \rangle} d\mu \\ &= - \int_{\Sigma} \phi^2 |\nabla u|^2 e^{\langle \mathbf{T}, \mathbf{x} \rangle} d\mu \\ &\quad - 2 \int_{\Sigma} \phi u \langle \nabla \phi, \nabla u \rangle e^{\langle \mathbf{T}, \mathbf{x} \rangle} d\mu \end{aligned}$$

which implies that

$$(28) \quad \int_{\Sigma} \phi^2 (2u|H|^2 + |\nabla u|^2) e^{\langle \mathbf{T}, \mathbf{x} \rangle} d\mu \leq 2 \int_{\Sigma} \phi u |\nabla \phi| |\nabla u| e^{\langle \mathbf{T}, \mathbf{x} \rangle} d\mu.$$

Now we take a smooth function  $\phi(x) : \mathbb{R} \rightarrow \mathbb{R}$  supported on  $[-R, R]$  which has the property:  $\phi = 1$  on  $[-\frac{R}{2}, \frac{R}{2}]$  and there is a constant  $C_b > 0$  such that  $|\nabla \phi| \leq C_b \phi^b \frac{1}{R}$  for  $0 < b < 1$ .

By using Young's inequality, the bound of  $u = \theta^2$  and the property of  $\phi$ , we can estimate (28) as follows:

$$\begin{aligned} \int_{\Sigma} \phi^2 (2u|H|^2 + |\nabla u|^2) e^{\langle \mathbf{T}, \mathbf{x} \rangle} d\mu &\leq C \int_{\Sigma} \phi^{1+b} u |\nabla u| \frac{1}{R} e^{\langle \mathbf{T}, \mathbf{x} \rangle} d\mu \\ &\leq \tilde{C} \frac{1}{R} \int_{\Sigma} \phi^{1+b} |\nabla u| e^{\langle \mathbf{T}, \mathbf{x} \rangle} d\mu \\ &\leq \int_{\Sigma} \phi^2 |\nabla u|^2 e^{\langle \mathbf{T}, \mathbf{x} \rangle} d\mu + \int_{\Sigma \cap \operatorname{supp} \phi} \frac{C}{R^{\frac{2}{1-b}}} e^{\langle \mathbf{T}, \mathbf{x} \rangle} d\mu. \end{aligned}$$

Finally we choose  $b > 0$  such that  $n - \frac{2}{1-b} < 0$ , i.e.,  $b > 1 - \frac{2}{n}$ . Then

$$\int_{\Sigma} \phi^2 u |H|^2 e^{\langle \mathbf{T}, \mathbf{x} \rangle} d\mu \leq \frac{C}{R^{\frac{2}{1-b}}} \tilde{\mu}(\Sigma \cap B_{2R}(0)).$$

Letting  $R \rightarrow \infty$  in (4), we obtain that  $|H| \equiv 0$  on  $\Sigma$ . This means that  $\Sigma$  is a minimal submanifold.

When  $n = 2$ , by (7), we see that  $A \equiv 0$  on  $\Sigma$ . Thus it must be a flat plane.  $\square$

**Acknowledgement.** The second author was supported by the National Natural Science Foundation of China (Grant No. 12271069) and the third author was supported by the National Natural Science Foundation of China (Grant No. 12071352, 12271039). The authors would like to thank Professor Hongbing Qiu for stimulating discussions on this subject.

**Author Contributions** All authors have contributed equally.

**Data availability** No datasets were generated or analysed during the current study.

## Declarations

**Conflict of interest** Not applicable.

## REFERENCES

- [AB10] Ben Andrews and Charles Baker, *Mean curvature flow of pinched submanifolds to spheres*, Journal of Differential Geometry **85** (2010), no. 3, 357–396. ↑5
- [CL04] Jingyi Chen and Jiayu Li, *Singularity of mean curvature flow of Lagrangian submanifolds*, Inventiones mathematicae **156** (2004), no. 1, 25–51. ↑1
- [Col12] Tobias H and Minicozzi Colding William P, *Generic mean curvature flow I; generic singularities*, Annals of mathematics (2012), 755–833. ↑12
- [HL09] Xiaoli Han and Jiayu Li, *Translating solitons to symplectic and Lagrangian mean curvature flows*, International Journal of Mathematics **20** (2009), no. 04, 443–458. ↑2, 5
- [HLS11] Xiaoli Han, Jiayu Li, and Jun Sun, *The second type singularities of symplectic and Lagrangian mean curvature flows*, Chinese Annals of Mathematics, Series B **32** (2011), no. 2, 223–240. ↑2, 4
- [HS10] Xiaoli Han and Jun Sun, *Translating solitons to symplectic mean curvature flows*, Annals of Global Analysis and Geometry **38** (2010), no. 2, 161–169. ↑2
- [HL82] Reese Harvey and H Blaine Lawson, *Calibrated geometries*, Acta Mathematica **148** (1982), 47–157. ↑1, 4
- [HS99] Gerhard Huisken and Carlo Sinestrari, *Mean curvature flow singularities for mean convex surfaces*, Calculus of Variations and Partial Differential Equations **8** (1999), no. 1, 1–14. ↑
- [LS24] Xiang Li and Jun Sun, *Eternal solutions to almost calibrated Lagrangian and symplectic mean curvature flows*, Journal of Mathematical Analysis and Applications **531** (2024), no. 1. ↑2, 4
- [Nev07] André Neves, *Singularities of Lagrangian Mean Curvature Flow: Zero-Maslov class case*, Inventiones mathematicae **168** (2007), no. 3, 449–484. ↑1
- [Nev13] ———, *Finite time singularities for Lagrangian mean curvature flow*, Annals of Mathematics. Second Series **177** (2013), no. 3, 1029–1076. ↑1
- [NT13] André Neves and Gang Tian, *Translating solutions to Lagrangian mean curvature flow*, Transactions of the American Mathematical Society **365** (2013), no. 11, 5655–5680. ↑2, 3
- [Qiu24] Hongbing Qiu, *Rigidity theorems of Lagrangian and symplectic translating solitons*, preprint (2024). ↑2
- [SW01] Richard Schoen and Jon Wolfson, *Minimizing area among Lagrangian surfaces: the mapping problem*, Journal of Differential Geometry **58** (2001), no. 1, 1–86. ↑1
- [Smo96] Knut Smoczyk, *A canonical way to deform a Lagrangian submanifold*, arXiv preprint dg-ga/9605005 (1996). ↑1, 5
- [Smo99] ———, *Harnack inequality for the Lagrangian mean curvature flow*, Calculus of Variations and Partial Differential Equations **8** (1999), no. 3, 247–258. ↑5
- [Sun13] Jun Sun, *A gap theorem for translating solitons to Lagrangian mean curvature flow*, Differential Geometry and its Applications **31** (2013), no. 5, 568–576. ↑2
- [Sun14] ———, *Mean curvature decay in symplectic and lagrangian translating solitons*, Geometriae Dedicata **172** (2014), no. 1, 207–215. ↑2
- [Sun15] ———, *Rigidity results on Lagrangian and symplectic translating solitons*, Communications in Mathematics and Statistics **3** (2015), no. 1, 63–68. ↑2, 3
- [TY02] Richard P. Thomas and Shing-Tung Yau, *Special Lagrangians, stable bundles and mean curvature flow*, Communications in Analysis and Geometry **10** (2002), no. 5, 1075–1113. ↑4
- [Wan01] Mu-Tao Wang, *Mean curvature flow of surfaces in Einstein four-manifolds*, Journal of Differential Geometry **57** (2001), no. 2, 301–338. ↑1
- [Xin15] Yuanlong Xin, *Translating solitons of the mean curvature flow*, Calculus of Variations and Partial Differential Equations **54** (2015), no. 2, 1995–2016. ↑3

<sup>1</sup>SCHOOL OF MATHEMATICAL SCIENCES, JIANGSU UNIVERSITY, ZHENJIANG 212013, CHINA  
*Email address:* [lihsiang@mail.ustc.edu.cn](mailto:lihsiang@mail.ustc.edu.cn)

<sup>2</sup>MATHEMATICAL SCIENCE RESEARCH CENTER, CHONGQING UNIVERSITY OF TECHNOLOGY, CHONGQING  
400054, CHINA  
*Email address:* [yongluo-math@cqut.edu.cn](mailto:yongluo-math@cqut.edu.cn)

<sup>3</sup>SCHOOL OF MATHEMATICS AND STATISTICS, WUHAN UNIVERSITY, WUHAN 430072, CHINA  
*Email address:* [sunjun@whu.edu.cn](mailto:sunjun@whu.edu.cn)