The homotopy theory of cyclotomic spectra 10 years later

Andrew J. Blumberg Michael A. Mandell

Author address:

DEPARTMENT OF MATHEMATICS, COLUMBIA UNIVERSITY, NEW YORK, NY 10027

 $Email \ address: \verb"blumberg@math.columbia.edu"$

Department of Mathematics, Indiana University, Bloomington, IN47405

Email address: mmandell@iu.edu

Contents

Chapter I. Introduction	7
Chapter II. Model Categories and Geometric Fixed Points on Equivariant Commutative Ring Spectra (Theorem E)	15
 V-constrained F-model categories and their variants The multiplicative geometric fixed point functor on variant model 	15
structures and Theorem E	21
Chapter III. The Circle Group and Cyclotomic Spectra (Theorems B, C, D)	27
1. The circle group and Theorem C	27
2. Pre-cyclotomic spectra and Theorems D and B	33
3. Construction of \mathcal{A}^{cyc} and proof of Lemma 1.4	39
4. The positive convenient Σ -model structure and Theorem 1.3	46
Chapter IV. The Multiplicative tom Dieck Splitting (Theorem A)	51
1. Construction of the multiplicative splitting	51
2. The factors in the multiplicative splitting	55
Chapter V. Generalized Orbit Desuspension Spectra: Main Properties	59
1. Generalized orbit desuspension spectra	59
2. Geometric fixed points and smash products	62
3. Geometric fixed points of symmetric powers	65
Chapter VI. Generalized Orbit Desuspension Spectra: Proofs	69
1. The quotient fixed point lemma and Theorems V.2.4, V.2.5, and V.2.7	69
2. Reduction of Theorems V.2.6 and V.2.9	76
3. Proof of Theorem 2.2	79
4. Proof of Theorem V.2.8	87
5. Proof of Theorem V.3.1	93
6. The diagonal fixed point lemma and Theorem V.3.5	96
7. Indexed smash powers, HHR norms, geometric fixed points, and the	
HHR diagonal	99
Bibliography	107

Abstract

This paper studies the foundations of the geometric fixed point functor in multiplicative equivariant stable homotopy theory. We introduce a new class of equivariant orthogonal spectra called generalized orbit desuspension spectra and analyze their homotopical behavior with respect to the geometric fixed point functor and especially the interaction with smash products and symmetric powers. This analysis leads to several new foundational results, including the construction of the derived functor of geometric fixed points on equivariant commutative ring orthogonal spectra and its comparison to the derived functor on the underlying equivariant orthogonal spectra. In addition this theory provides foundations for the category of commutative ring pre-cyclotomic spectra and a formula for the derived space of maps in this category (a formula needed in the authors' paper with Yuan on relative TC). Finally, we prove a new multiplicative tom Dieck splitting for equivariant commutative ring spectra.

²⁰²⁰ Mathematics Subject Classification. Primary 55P91. Secondary 19D55.

Key words and phrases. Topological cyclic homology, cyclotomic spectrum, geometric fixed points.

The first author was supported in part by NSF grants DMS-2052970, DMS-2104420.

The second author was supported in part by NSF grants DMS-2052846, DMS-2104348.

CHAPTER I

Introduction

The last 50 years of work on equivariant stable homotopy theory has led to spectacular accomplishments both internally (e.g., [8, 13]) and externally (e.g., [17, 11, 12, 6, 14, 2, 15]. However, despite monumental work codifying the theory by Lewis-May-Steinberger [20] (later extended by Mandell-May [22]), the foundations of the theory, particularly the treatment of multiplicative phenomena, remain unsettled. Notably, the spectacular recent advances of Hill-Hopkins-Ravenel [15] depended on a new foundational construction, the multiplicative norm, and this has led to an array of new computations and insights (e.g., see [16, 3]).

A particularly glaring gap in the current foundations is the interaction between geometric fixed points and the homotopy theory of equivariant E_{∞} ring spectra. In particular, the following basic questions do not have good answers.

- (i) Is there a derived functor of geometric fixed points on equivariant commutative ring orthogonal spectra?
- (ii) If so, does it agree with the derived functor on the underlying equivariant orthogonal spectra?

This issue was cleverly sidestepped in Hill-Hopkins-Ravenel, and arose again in our work [4] on the homotopy theory of cyclotomic spectra. Specifically, lack of control on the derived functor of geometric fixed points in the multiplicative setting has prevented the detailed study of commutative ring objects in cyclotomic spectra. For work along these lines, the existence of derived functors is rarely enough and what is needed is some kind of additional homotopical control, for example, a model structure where the geometric fixed points have good properties.

The purpose of this paper is to return to these foundations and fill in the gaps. In particular for a compact Lie group G and a closed subgroup H < G, we construct a closed model structure \mathcal{M} (that depends on H) on the category of G-equivariant commutative ring orthogonal spectra with the following properties, positively answering the questions above.

- (i) The point-set geometric fixed point functor Φ^H preserves weak equivalences between \mathcal{M} -cofibrant objects.
- (ii) If R is \mathcal{M} -cofibrant and $X \to R$ is a cofibrant approximation in the standard model structure on orthogonal spectra, then the induced map $\Phi^H X \to \Phi^H R$ is a weak equivalence.

Here (ii) implies (i) and is established in Theorem E below. The \mathcal{M} -cofibrant objects also have the following further properties.

(iii) If R is \mathcal{M} -cofibrant, then $\Phi^H R$ is flat for the smash product of WHequivariant orthogonal spectra and in particular flat for the smash product of non-equivariant orthogonal spectra. (Proposition II.2.6)

(iv) If R is \mathcal{M} -cofibrant and X is any G-equivariant orthogonal spectrum, then the lax monoidal structure map

$$\Phi^H R \wedge \Phi^H X \longrightarrow \Phi^H (R \wedge X)$$

is an isomorphism. (Proposition II.2.7)

We also mention here the following foundational property of the smash product, which is not directly related to the \mathcal{M} model structure and should have been observed earlier; it is essential to our work with Yuan in [5] but does not seem to be well-known.

(v) If M, N are R-modules with M cofibrant (in the usual model structure), then the lax symmetric monoidal structure map

 $\Phi^H M \wedge_{\Phi^H R} \Phi^H N \longrightarrow \Phi^H (M \wedge_R N)$

is an isomorphism. (Proposition II.2.8)

We expect these new foundations to have a wide variety of applications; we already deploy them here in a number of interesting directions. A particularly striking new structural result about multiplicative equivariant stable homotopy theory establishes a previously unnoticed multiplicative version of the tom Dieck splitting. The tom Dieck splitting, a foundational result in equivariant stable homotopy theory, is usually stated for suspension spectra, but it has a version for genuine *G*-equivariant spectra promoted from non-equivariant spectra, whether or not suspension spectra. In the multiplicative context, we consider the left derived functor ϵ^* that makes a non-equivariant commutative ring orthogonal spectrum into a *G*-equivariant commutative ring orthogonal spectrum. We prove the following theorem in Chapter IV.

THEOREM A (Multiplicative tom Dieck Splitting). Let G be a finite group and let ϵ^* denote the left derived functor from the homotopy category of nonequivariant commutative ring orthogonal spectra to the homotopy category of genuine G-equivariant commutative ring orthogonal spectra. There exists a natural weak equivalence

$$\Phi^G(\epsilon^* R) \simeq R \otimes (\prod BWH)$$

where \otimes denotes the left derived tensor (topologically indexed colimit) in the category of commutative ring orthogonal spectra and the disjoint union is over conjugacy classes of subgroups H < G.

Our original motivation for this paper was to provide the technical foundations for our work [5] with Yuan on relative cyclotomic structures. That project requires us to construct the homotopy theory of commutative ring objects in cyclotomic spectra and to compute the derived mapping spaces. The existence of a model structure presenting the correct (∞ , 1)-category of commutative ring objects in cyclotomic spectra (stated as Theorem E below) is an immediate consequence of our foundational work, and using this we prove the following characterization of the derived mapping spaces which provides the commutative ring analogue of the main theorem on mapping spectra in [4, 5.12].

THEOREM B. If A and B are commutative ring p-pre-cyclotomic spectra where the underlying \mathbb{T} -equivariant commutative ring orthogonal spectra of A and B are cofibrant and fibrant, respectively, in the model structure of Theorem C.(ii) below,

8

then the derived space of maps of commutative ring p-pre-cyclotomic spectra from A to B is represented by the homotopy equalizer

$$\mathcal{C}om^{\mathbb{T}}(A,B) \rightrightarrows \mathcal{C}om^{\mathbb{T}}(\Phi A,B)$$

where $Com^{\mathbb{T}}$ denotes the point-set mapping space in \mathbb{T} -equivariant commutative ring orthogonal spectra. Here one map is induced by the cyclotomic structure map $\Phi A \to A$ and the other map is induced by applying Φ and post-composing with the cyclotomic structure map $\Phi B \to B$.

The foundation for all the theorems above is the identification and study of a new class of G-equivariant orthogonal spectra that behave well with respect to smash product and the point-set geometric fixed point functors. We call these spectra "generalized orbit desuspension spectra". A generalized orbit desuspension spectrum is a certain kind of equivariant orthogonal spectrum $J\eta$ constructed from an equivariant vector bundle η . (See Terminology V.1.1 and Construction V.1.2) for specifics on the hypotheses on η and the construction $J\eta$.) The orbit desuspension spectra $G/H_+ \wedge F_V S^0$ and the HHR complete cells $G \wedge_K F_W S^0$ are examples that come from the trivial G-equivariant vector bundles $G/H \times V \rightarrow G/H$ and the extended G-equivariant vector bundles $G \times_K W \to G/K$, respectively. The multiplicative homotopy theory of G-equivariant orthogonal spectra works because orbit desuspension spectra are flat for the smash product of G-equivariant orthogonal spectra (smashing with one preserves weak equivalences between arbitrary Gequivariant orthogonal spectra). The point-set geometric fixed point functors are useful because they are homotopically correct on the orbit desuspension spectra and they commute up to isomorphism with the smash product of an orbit desuspension spectrum and an arbitrary G-equivariant orthogonal spectrum (the latter being the main technical result of [4, App. A] that makes the homotopy theory of pre-cyclotomic spectra work). Under mild hypotheses, generalized orbit desuspension spectra share these properties. Details of the theory take up the bulk of the paper, starting in Chapter V.

We now move on to a more detailed exposition of the homotopical results of the paper, specifically the construction of the model structures, with precise statements of some the most important results. In this exposition and most of the rest of the paper, we use the following terminology:

TERMINOLOGY.

- "spectrum" means orthogonal spectrum; for a compact Lie group G, "G-spectrum" means G-equivariant orthogonal spectrum. Often these refer to point set categories, but when homotopy theory is implied, we understand the complete universe theory unless explicitly indicated otherwise.
- We abbreviate pre-cyclotomic spectrum to "pc spectrum" and *p*-precyclotomic spectrum to "*p*pc spectrum". These concepts are defined in [4], and we give only minimal review here.
- For spectra and G-spectra, we prepend "ring" or "commutative ring" to indicate the categories of associative monoids or commutative monoids for the smash product (e.g., we say "ring G-spectra" for associative ring G-equivariant orthogonal spectra). Similar conventions apply to module categories.

• For ppc spectra and pc spectra, we review ring, commutative ring, and module categories in Section III.2.

We start the detailed exposition with our results on commutative ring ppc spectra. In [4], we introduced a homotopy theory for cyclotomic spectra and ppc spectra in order to justify a construction of derived mapping spectra and to prove corepresentability of TC. (Corepresentability of TC(-;p), now often treated as a definition rather than a theorem, was first conjectured in [18, 7.9] and proved in [4, 1.4].) The definition of the category of ppc spectra depends on a version of the point-set multiplicative geometric C_p fixed point functor Φ^{C_p} . This functor goes from \mathbb{T} -spectra to \mathbb{T}/C_p -spectra, but using the pth root isomorphism $\rho \colon \mathbb{T} \cong \mathbb{T}/C_p$, we get a version that is an endofunctor functor on \mathbb{T} -spectra, which we denote as Φ :

$$\Phi := \rho^* \Phi^{C_p} \colon \mathcal{S}^{\mathbb{T}} \longrightarrow \mathcal{S}^{\mathbb{T}}.$$

A key fact that made the mapping spectra analyzable is that the endofunctor Φ takes cofibrant T-spectra to cofibrant T-spectra (in the standard stable model structures and appropriate family variants of [22, IV§6]). The same does not hold in the setting of commutative ring T-spectra: for a commutative ring T-spectrum R, ΦR has the canonical structure of a commutative ring T-spectrum, but even when R is cofibrant as a commutative ring T-spectrum, ΦR may not be. (More generally, Φ does not preserve cofibrant objects in the model structures introduced in [27], [7, 3.3.12].) We prove the following theorem in Section III.1:

THEOREM C. Let \mathcal{F} be either the family of \mathcal{F}_{fin} of all finite subgroups of \mathbb{T} , or the family \mathcal{F}_p of finite subgroups of \mathbb{T} of order a power of p. There exists a symmetric monoidal topological model structure $\mathcal{M}_{\mathcal{F}}$ on \mathbb{T} -spectra which has weak equivalences the genuine \mathcal{F} -equivalences, is enriched over the standard model structure on non-equivariant spectra (in the sense of [21, 3.3]), and which satisfies the following properties:

- (i) The endofunctor Φ preserves cofibrations and acyclic cofibrations.
- (ii) The category of commutative ring T-spectra admits a topological model structure with fibrations and weak equivalences created by the forgetful functor to T-spectra.
- (iii) The endofunctor Φ preserves cofibrant objects in the model structure in (ii).
- (iv) If R is cofibrant in the model structure of (ii), then it is cofibrant in the undercategory of S in the model structure $\mathcal{M}_{\mathcal{F}}$, i.e., the unit $\mathbb{S} \to R$ is a cofibration in $\mathcal{M}_{\mathcal{F}}$.

Moreover, the forgetful functor from \mathbb{T} -spectra to non-equivariant spectra takes cofibrations and fibrations in this model structure to cofibrations and fibrations in the positive convenient Σ -model structure of Definition III.4.1.

The positive convenient Σ -model structure is the orthogonal spectrum analogue of Shipley's "S-model structure" on symmetric spectra from [26] and a variant of Stolz's "S-model structure" from [27, §1.3.3] or [7, 3.3.12] for the trivial group. See Section III.4 for details.

By the K. Brown lemma, (i) implies that Φ preserves weak equivalences between cofibrant objects; combining with (iv), we see that for R a cofibrant \mathbb{T} -equivariant commutative ring spectrum and $X \to R$ a cofibrant approximation in the model structure $\mathcal{M}_{\mathcal{F}}$, the map $\Phi X \to \Phi R$ is a weak equivalence.

10

The model structures on commutative ring T-spectra in the theorem lead to corresponding model structures on modules and algebras.

COROLLARY. Let R be a commutative ring \mathbb{T} -spectra. For the model structure $\mathcal{M}_{\mathcal{F}}$ constructed in the proof of Theorem C, the categories of R-modules and Ralgebras admit model structures with the fibrations and weak equivalences created in $\mathcal{M}_{\mathcal{F}}$. Moreover, if R is cofibrant in the model structure of (ii), then cofibrant R-modules are cofibrant in the model structure $\mathcal{M}_{\mathcal{F}}$ and cofibrant R-algebras are cofibrant in the model structure $\mathcal{M}_{\mathcal{F}}$ on the undercategory of S.

Recall that a ppc spectrum consists of a \mathbb{T} -spectrum X together with a cyclotomic structure map

$$\Phi X \longrightarrow X$$

and a map of ppc spectra is a map of spectra which makes the diagram of cyclotomic structure maps commute. As shown in [4, 5.3], the category of ppc spectra have a model structure with fibrations and weak equivalences created by the forgetful functor to the standard model structure on \mathbb{T} -spectra with the \mathcal{F}_p -equivalences. Using the model structure of Theorem C, we get a new model structure on ppcspectra and a compatible model structure on the category of commutative ring ppcspectra.

THEOREM D. There exists a topological model structure on the category of ppc spectra with weak equivalences the genuine \mathcal{F}_p -equivalences and fibrations created in the model category of \mathbb{T} -spectra of Theorem C. Moreover, it is enriched over the standard model structure on non-equivariant spectra, and it satisfies the following additional properties:

- (i) The forgetful functor to T-spectra sends cofibrations to cofibrations in the model structure of Theorem C.
- (ii) The smash product ∧ extends to a symmetric monoidal product on the subcategory of cofibrant ppc spectra in this model structure.
- (iii) The category of commutative ring ppc spectra admits a topological model structure where fibrations and weak equivalences are created by the forgetful functor to ppc spectra, and the forgetful functor to ppc spectra under S preserves cofibrant objects.

Because of the issues with Φ , categories of modules and algebras in ppc spectra are hard to study in general, but when the ring ppc spectrum R has underlying ppcspectrum cofibrant in the undercategory of \mathbb{S} , the usual arguments can be adapted to work. As a particular case, we have the following addendum to the previous theorem for module and algebra model structures.

ADDENDUM D. Let R be a cofibrant commutative ring ppc spectrum.

- (i) The category of R-module ppc spectra admits a model structure where fibrations and weak equivalences are created by the forgetful functor to ppc spectra, and the forgetful functor preserves cofibrant objects. The model structure is enriched over the standard model structure on nonequivariant spectra. Moreover, ∧_R extends to a symmetric monoidal product on the subcategory of cofibrant R-module ppc spectra.
- (ii) The category of R-algebra ppc spectra admits a model structure where fibrations and weak equivalences are created by the forgetful functor to

ppc spectra, and the forgetful functor to the undercategory of S preserves cofibrant objects.

(iii) The category of commutative R-algebra ppc spectra admits a model structure where fibrations and weak equivalences are created by the forgetful functor to ppc spectra, and the forgetful functor to the undercategory of S preserves cofibrant objects.

The particulars of Theorems C and D allow us to extend the main result on mapping spaces of [4] to the multiplicative context. We stated the case for commutative R-algebras as Theorem B above; see Section III.2 for the other cases and for other variants of cyclotomic spectra.

The fact that the geometric fixed points Φ^{C_p} admits a refinement into an endofunctor Φ is a special feature of the circle group \mathbb{T} . As we mentioned above, we have theorems for the homotopical properties of geometric H fixed points of commutative ring G-spectra more generally, but they are not as clean to state. The easiest statement that looks useful is the following. It is proved at the end of Section II.2; see Theorem II.2.5 for another theorem along these lines.

THEOREM E. Let G be a compact Lie group and \mathcal{H} a finite set of closed subgroups of G. There exists a model structure on the category of commutative ring Gspectra with the genuine weak equivalences, having the following property: when R is cofibrant in this model structure, $X \to R$ a cofibrant approximation in the (genuine) standard or positive stable model structure on G-spectra, and $H \in \mathcal{H}$, the induced map on the point-set multiplicative geometric fixed point functor $\Phi^H X \to \Phi^H R$ is a weak equivalence. In particular, for $H \in \mathcal{H}$, Φ^H preserves weak equivalences between cofibrant objects in this model structure.

Theorem E establishes the existence of a left derived functor for geometric fixed points, viewed as a functor from commutative ring G-spectra to commutative ring WH-spectra and a natural isomorphism from this derived functor composed with the forgetful functor to the genuine WH-stable category and the derived geometric fixed point functor on the underlying G-spectrum.

Technical conventions. In addition to the terminology for spectra and precyclotomic spectra set forth on page 9 above, we also have the following technical conventions.

We use the phrase "orthogonal G-representation" to mean a finite dimensional real inner product space with action of G by isometries, rather than the isomorphism class of such an object (as the phrase is often used in algebra). For an orthogonal G-representation V, we write [V] for its isomorphism class.

For orthogonal G-representations V, W, we write V < W to denote that V is a G-stable vector subspace of W, not necessarily a proper subspace. We write [V] < [W] to indicate that V is isomorphic to a G-stable vector subspace of W (not necessarily a proper subspace).

We write H < G to indicate that H is a closed subgroup of G, not necessarily a proper subgroup.

In Chapters V and VI, Γ denotes a compact Lie group. Wherever it appears in this paper, the phrase " Γ -space" means space with continuous Γ action, and not Γ -space in the sense of Segal.

Acknowledgments. This project was motivated by the authors' work with Allen Yuan on relative cyclotomic spectra; we would not have discovered the results here without their necessity in that work. We would like to thank our collaborators on [1], Vigleik Angeltveit, Teena Gerhardt, Mike Hill, and Tyler Lawson for many discussions of related topics. The first author also like to particularly thank the Mikes Hill and Hopkins for years of illuminating discussions about geometric fixed points and equivariant commutative ring spectra.

CHAPTER II

Model Categories and Geometric Fixed Points on Equivariant Commutative Ring Spectra (Theorem E)

This chapter proves Theorem E from the introduction, which constructs the left derived functor for geometric H fixed points on commutative ring G-spectra and shows that it agrees with the corresponding functor on the underlying G-spectra. This requires the construction of some new model structures, and the first section is devoted to a review of some general facts for constructing cofibrantly generated model structures on G-spectra that are Quillen equivalent to the standard model structure.

1. V-constrained F-model categories and their variants

The purpose of this section is to specialize some standard results on creating new model structures from old ones in the context of the standard model structure on equivariant orthogonal spectra. We claim no novelty on these results and in fact we regard the techniques as well-known enough to omit all proofs.

Let G be a compact Lie group. The standard stable model structure on Gspectra has its weak equivalences defined to be the maps that induce isomorphisms on homotopy groups π^H_* for all closed subgroups H < G; its cofibrations are the retracts of relative cell complexes where the cells are of the form

$$G/H_+ \wedge F_V S^0 \wedge (D^n, \partial D^n)_+$$

where V is an orthogonal G-representation, and F_V denotes the functor from based G-spaces to G-spectra that is left adjoint to the Vth space functor. Hill-Hopkins-Ravenel introduced the *complete model structure* that is better behaved with respect to induction along a closed subgroup. It has the same weak equivalences but more cofibrations; a cofibration is a retract of a relative cell complex where the cells are of the form

$$G_+ \wedge_H \wedge F_V S^0 \wedge (D^n, \partial D^n)_+$$

where now V is allowed to be an orthogonal H-representation.

There are variants of the standard and complete stable model structures in the context of a family \mathcal{F} of subgroups of G. A family \mathcal{F} is a non-empty set of closed subgroups of G with the property that when $H \in \mathcal{F}$ so is every conjugate and every closed subgroup of H. Given a family \mathcal{F} , the \mathcal{F} -equivalences are the maps that induce isomorphisms on homotopy groups π^H_* for H in \mathcal{F} . The standard and complete \mathcal{F} -model structures have cofibrations the retracts of relative cell complexes with the cells displayed above (for the standard and complete stable model structures, respectively) but with the restriction that the subgroup H be in \mathcal{F} . From the very beginning, dealing with commutative ring orthogonal spectra necessitated the creation of additional model structures, the so-called *positive* model structures, which put a constraint on the orthogonal G-representations V used in the definition of cells for defining cofibrations. We have positive variants of all the model structures mentioned above where we require V to contain a positive dimensional trivial representation. In later sections, we will see that asking for even stricter constraints on V will lead to homotopical results on the geometric fixed points of commutative ring orthogonal spectra. The following definition gives the framework for the types of constraints we consider.

DEFINITION 1.1. A family representation constraint $(\mathcal{F}, \mathcal{V})$ consists of a family \mathcal{F} of subgroups of G and a set of isomorphism classes of orthogonal H-representations $\mathcal{V}(H)$ for each $H \in \mathcal{F}$ such that:

- (i) If $[V], [W] \in \mathcal{V}(H)$, then $[V] \oplus [W] \in \mathcal{V}(H)$.
- (ii) For any orthogonal *H*-representation *W*, there exists $[V] \in \mathcal{V}(H)$ with [W] < [V].
- (iii) If $H' = gHg^{-1}$, then $[V] \mapsto [gV]$ gives a bijection from $\mathcal{V}(H)$ to $\mathcal{V}(H')$.
- (iv) If K < H, then the isomorphism classes of underlying K-representations of the elements of $\mathcal{V}(H)$ lie in $\mathcal{V}(K)$.

We say that \mathcal{V} is *standard compatible* if it has the property that for every orthogonal *G*-representation *W* and every $H \in \mathcal{F}$, there exists an orthogonal *G*representation *V* with W < V and the underlying isomorphism class of orthogonal *H*-representations in $\mathcal{V}(H)$.

We have the following basic examples.

EXAMPLE 1.2. For any family, we have a constraint \mathcal{V}_0 defined by taking $\mathcal{V}_0(H)$ to be all isomorphism classes of orthogonal *H*-representations. We have a constraint \mathcal{V}_+ defined by taking $\mathcal{V}_+(H)$ to be all isomorphism classes [V] such that $V^H \neq 0$ (i.e., [V] contains a copy of the trivial representation). For a family consisting of finite groups, we have a constraint \mathcal{V}_{reg} defined by taking $\mathcal{V}_{\text{reg}}(H)$ to consist of sums of regular representations. The constraints \mathcal{V}_0 and \mathcal{V}_+ are standard compatible and \mathcal{V}_{reg} is standard compatible in the case when *G* is finite or $G = \mathbb{T}$.

We also have the following example that plays an important role in Theorem C.

EXAMPLE 1.3. Recall that a permutation representation of H is an orthogonal H-representation isomorphic to one of the form $\mathbb{R}\langle X \rangle$, where X is a finite H-set; equivalently, it is an orthogonal H-representation that has an orthonormal basis on which the H action restricts. We have a constraint \mathcal{V}_{π} defined by taking $\mathcal{V}(H)$ to consist of the isomorphism classes of permutation representations. In the work on pc spectra, we need a family constraint that combines \mathcal{V}_{reg} and \mathcal{V}_{π} . Let $\mathcal{V}_{\pi!}$ be the family constraint where $[V] \in \mathcal{V}_{\pi!}(H)$ when $V \cong \mathbb{R}\langle X \rangle$ for an H-set X that has at least one free H-orbit. We call these representations the semiregular permutation representations. The constraints \mathcal{V}_{π} and $\mathcal{V}_{\pi!}$ are standard compatible when G is finite or $G = \mathbb{T}$.

A family representation constraint specifies a constraint on the allowed cell types as follows.

NOTATION 1.4. Given a family representation constraint $(\mathcal{F}, \mathcal{V})$, let $\mathcal{A}^{\natural}(\mathcal{F}, \mathcal{V})$ denote the set of *G*-spectra $G/H_+ \wedge F_V S^0$ for $H \in \mathcal{F}$ and *V* an orthogonal *G* action

16

on \mathbb{R}^n (for some $n \geq 0$) such that its underlying orthogonal *H*-representation satisfies $[V] \in \mathcal{V}(H)$; let $\mathcal{A}^{\circ}(\mathcal{F}, \mathcal{V})$ denote the set of *G*-spectra $G_+ \wedge_H F_V S^0$ for $H \in \mathcal{F}$ and *V* an orthogonal *H* action on \mathbb{R}^n (for some $n \geq 0$) that satisfies $[V] \in \mathcal{V}(H)$.

The set $\mathcal{A}^{\natural}(\mathcal{F}, \mathcal{V})$ is only useful for us when \mathcal{V} is standard compatible. With $\mathcal{A} = \mathcal{A}^{\natural}(\mathcal{F}, \mathcal{V})$ and $\mathcal{A} = \mathcal{A}^{\circ}(\mathcal{F}, \mathcal{V})$ in the following definition, we get the standard and complete cells (respectively) with subgroup and representation constraints specified by \mathcal{F} and \mathcal{V} .

NOTATION 1.5. For any subset \mathcal{A} of G-spectra, let $I_{\mathcal{A}}$ be the set of maps of the form

$$A \wedge (\partial D^n_+ \longrightarrow D^n_+)$$

for $A \in \mathcal{A}$ and $n \ge 0$ (with ∂D^0 the empty set).

As a technical hypothesis, we will always assume that the orthogonal spectra in \mathcal{A} are spacewise non-degenerately based. This hypothesis implies that the maps in $I_{\mathcal{A}}$ and the resulting cofibrations defined below are spacewise unbased Hurewicz cofibrations. While this plays no role in the (omitted) proofs of the model category structures below, it is indispensable in other contexts where Lillig's theorem is used, for example, checking that geometric realization preserves levelwise weak equivalences.

DEFINITION 1.6. Let \mathcal{A} be a set of spacewise non-degenerately based G-spectra. A relative \mathcal{A} -cell complex is a map $X \to Y$ such that $Y \cong \operatorname{colim} X_n$ where $X_0 = X$, and inductively X_{n+1} is formed as a pushout from X_n over a wedge of maps in $I_{\mathcal{A}}$ (with the isomorphism to the colimit inducing the given map $X \to Y$). A regular $I_{\mathcal{A}}$ -cofibration is a map $X \to Y$ given by transfinite composition of pushouts along maps in $I_{\mathcal{A}}$: there exists an ordinal λ , G-spectra X_α for $\alpha \leq \lambda$, and compatible maps $X_\alpha \to X_\beta$ for $\alpha < \beta \leq \lambda$ such that $X_0 = X$, $X_\lambda \cong Y$ (compatibly with the given map $X \to Y$), for every $\alpha < \lambda$, $X_{\alpha+1}$ is formed by pushout along a map in I_{α} , and for every limit ordinal $\beta \leq \lambda$, $X_{\beta} = \operatorname{colim}_{\alpha < \beta} X_{\alpha}$.

It is easy to see by transfinite induction that for a regular $I_{\mathcal{A}}$ -cofibration $X \to X_{\lambda}$, all the maps $X_{\alpha} \to X_{\beta}$ in the composition system are Hurewicz cofibrations of equivariant orthogonal spectra. That observation also implies that for any Gspectrum Z, when λ is a cardinal and greater than the sum of the cardinalities of the spaces $Z(\mathbb{R}^n)$, then any map $Z \to Y$ factors through some X_{α} with $\alpha < \lambda$. This allows use of the small object argument in this context.

A relative \mathcal{A} -cell complex is in particular a regular $I_{\mathcal{A}}$ -cofibration (for example by well ordering the set of maps in each pushout). Under certain "smallness" hypotheses, the reverse is also true.

DEFINITION 1.7. A G-spectrum A is topologically point-set small if the set of maps out of A commutes with transfinite compositions when the constituent maps are spacewise closed inclusions.

The orbit desuspension spectrum F_VG/H_+ is point-set topologically small since maps out of it represent the functor $X \mapsto X(V)^H$. By adjunction it follows that Ais point-set topologically small if and only if the function spectrum functor F(A, -)commutes with transfinite compositions when the constituent maps are spacewise closed inclusions. Adjunction again implies that the smash product of topologically point-set small G-spectra is topologically point-set small. The relevance of this concept is the following proposition which follows from a transfinite induction argument (and the observation above that the maps in the composition system for a regular I_A -cofibration are Hurewicz cofibrations and so in particular spacewise closed inclusions).

PROPOSITION 1.8. Let \mathcal{A} be as in Definition 1.6. If every object A in \mathcal{A} is topologically point-set small then every regular $I_{\mathcal{A}}$ -cofibration is a relative \mathcal{A} -cell complex.

Standard techniques then give the following theorems, simultaneously generalizing the standard, positive, and complete stable and \mathcal{F} -model structures to more general family representation constraints.

THEOREM 1.9 (\mathcal{V} -constrained \mathcal{F} -model structure). For any compact Lie group G and any family representation constraint $(\mathcal{F}, \mathcal{V})$ with \mathcal{V} standard compatible, the category of G-spectra admits a symmetric monoidal topological model structure where:

- (i) The cofibrations are the retracts of relative $\mathcal{A}^{\natural}(\mathcal{F}, \mathcal{V})$ -cell complexes.
- (ii) The fibrations are the maps X → Y such that for every H ∈ F and every orthogonal G-representation V whose underlying H-representation satisfies [V] ∈ [V(H)]:
 - The map of non-equivariant spaces $X(V)^H \to Y(V)^H$ is a Serre fibration, and
 - For every orthogonal G-representation W with $[V \oplus W] \in \mathcal{V}(H)$, the square of non-equivariant spaces

$$\begin{array}{c} X(V)^H \longrightarrow (\Omega^W X(V \oplus W))^H \\ \downarrow \qquad \qquad \downarrow \\ Y(V)^H \longrightarrow (\Omega^W Y(V \oplus W))^H \end{array}$$

is homotopy cartesian.

(iii) The weak equivalences are the \mathcal{F} -equivalences.

THEOREM 1.10 (\mathcal{V} -constrained complete \mathcal{F} -model structure). For any compact Lie group G and any family representation constraint (\mathcal{F}, \mathcal{V}), the category of G-spectra admits a symmetric monoidal topological model structure where:

- (i) The cofibrations are the retracts of $\mathcal{A}^{\circ}(\mathcal{F}, \mathcal{V})$ -cell complexes.
- (ii) The fibrations are the maps $X \to Y$ such that for every $H \in \mathcal{F}$ and every orthogonal H-representation V that satisfies $[V] \in \mathcal{V}(H)$:
 - The map of non-equivariant spaces $X(V)^H \to Y(V)^H$ is a Serre fibration, and
 - For every orthogonal H-representation W with $[V \oplus W] \in \mathcal{V}(H)$, the square of non-equivariant spaces

$$\begin{array}{c} X(V)^H \longrightarrow (\Omega^W X(V \oplus W))^H \\ \downarrow \qquad \qquad \downarrow \\ Y(V)^H \longrightarrow (\Omega^W Y(V \oplus W))^H \end{array}$$

is homotopy cartesian.

18

(iii) The weak equivalences are the \mathcal{F} -equivalences.

See [25, 2.1] for a definition of monoidal model structure and e.g., [23, 5.12] for a definition of topological model structure.

In the case when $\mathcal{V} = \mathcal{V}_0$, we get the standard stable model structure and complete model structure with the family of all subgroups or the corresponding \mathcal{F} -model structures for an arbitrary family. When $\mathcal{V} = \mathcal{V}_+$, we get the corresponding positive model structures.

For the theorems in the introduction, we need to consider more general classes of cells \mathcal{A} . (For the following theorem, we note that $\mathcal{A}^{\natural}(\mathcal{F}, \mathcal{V}) \subset \mathcal{A}^{\circ}(\mathcal{F}, \mathcal{V})$. A version holds for constraints \mathcal{V} that are not standard compatible where we replace $\mathcal{A}^{\natural}(\mathcal{F}, \mathcal{V})$ with $\mathcal{A}^{\circ}(\mathcal{F}, \mathcal{V})$ and replace " \mathcal{V} -constrained \mathcal{F} -model structure" with " \mathcal{V} constrained complete \mathcal{F} -model structure".)

THEOREM 1.11 (Variant \mathcal{V} -constrained \mathcal{F} -model structures). Let G be a compact Lie group, and $(\mathcal{F}, \mathcal{V})$ a family representation constraint with \mathcal{V} standard compatible. Let \mathcal{A} be a set of spacewise non-degenerately based G-spectra with $\mathcal{A}^{\natural}(\mathcal{F}, \mathcal{V}) \subset \mathcal{A}$. Then the category of G-spectra admits a topological model structure where:

- (i) The cofibrations are the $I_{\mathcal{A}}$ -regular cofibrations.
- (ii) The fibrations are the maps $X \to Y$ such that for every $A \in \mathcal{A}$:
 - The induced map of mapping spaces $\mathcal{S}^G(A, X) \to \mathcal{S}^G(A, Y)$ is a Serre fibration, and
 - For every cofibrant object à in the V-constrained F-model structure and F-equivalence à → A, the square of mapping spaces

$$\begin{array}{c} \mathcal{S}^G(A,X) \longrightarrow \mathcal{S}^G(\tilde{A},X) \\ \downarrow \qquad \qquad \downarrow \\ \mathcal{S}^G(A,Y) \longrightarrow \mathcal{S}^G(\tilde{A},Y) \end{array}$$

is homotopy cartesian.

(iii) The weak equivalences are the \mathcal{F} -equivalences.

Moreover:

- If every object in A is topologically point-set small, then the cofibrations are the retracts of the relative A-cell complexes.
- If all objects A of A are flat in the sense that A∧(−) preserves weak equivalences, then all cofibrant objects are flat and this model structure satisfies the unit axiom and monoid axiom of Schwede-Shipley [25, 3.2,3.3].
- If for all A, B ∈ A, A ∧ B is cofibrant in this model structure, then this model structure satisfies the pushout-product axiom of Schwede-Shipley [25, 3.4].

We also have the following alternative identification of the fibrations. (It mainly changes the quantifier on the \mathcal{F} -equivalences $\tilde{A} \to A$ in the last bullet point.)

PROPOSITION 1.12. For \mathcal{A} , \mathcal{F} , \mathcal{V} in the previous theorem, the fibrations in the model structure there are precisely the maps $X \to Y$ that satisfy all of the following conditions.

- $X \to Y$ is a fibration in the \mathcal{V} -constrained \mathcal{F} -model structure.
- For every $A \in \mathcal{A}$, the map $\mathcal{S}^G(A, X) \to \mathcal{S}^G(A, Y)$ is a Serre fibration.

 For every A ∈ A, there exists a cofibrant object à in the V-constrained F-model structure and an F-equivalence à → A such that the square

$$\begin{array}{c} \mathcal{S}^{G}(A,X) \longrightarrow \mathcal{S}^{G}(\tilde{A},X) \\ \downarrow \\ \mathcal{S}^{G}(A,Y) \longrightarrow \mathcal{S}^{G}(\tilde{A},Y) \end{array}$$

is homotopy cartesian.

When the model structure satisfies the unit axiom, monoid axiom, and pushoutproduct axiom, we get corresponding model structures on categories of modules and algebras by [25]. For commutative rings and commutative algebras we need additional hypotheses.

THEOREM 1.13. Let G be a compact Lie group, and $(\mathcal{F}, \mathcal{V})$ a family representation constraint with \mathcal{V} standard compatible. Let \mathcal{A} be a set of spacewise nondegenerately based G-spectra with $\mathcal{A}^{\natural}(\mathcal{F}, \mathcal{V}) \subset \mathcal{A}$. Assume that the smash product of any two elements of \mathcal{A} is cofibrant in the \mathcal{A} -variant \mathcal{V} -constrained \mathcal{F} -model structure and assume that for every $A \in \mathcal{A}$, and every $n \geq 1$:

- (i) the nth symmetric power $A^{(n)}/\Sigma_n$ is flat for the smash product of G-spectra, and
- (ii) for every cofibrant approximation à → A in the V-constrained F-model structure, the induced map on nth symmetric powers

$$\tilde{A}^{(n)}/\Sigma_n \longrightarrow A^{(n)}/\Sigma_n$$

is an \mathcal{F} -equivalence.

Then the forgetful functor from commutative ring G-spectra to G-spectra with the A-variant V-constrained \mathcal{F} -model structure creates fibrations and weak equivalences in a model structure. Moreover, if \mathcal{A} is closed up to retract under nth symmetric powers for all $n \geq 1$, then commutative ring G-spectra that are cofibrant in this model structure are cofibrant in the undercategory of S in the \mathcal{A} -variant V-constrained \mathcal{F} -model structure on G-spectra.

We note that in the previous theorem, $G/H_+ \wedge \mathbb{S}$ does not satisfy condition (ii) and so cannot be an element of \mathcal{A} . In particular, $\mathcal{V}(H)$ must contain only positive dimensional isomorphism classes for every $H \in \mathcal{F}$.

The hypotheses in Theorem 1.13 include the extra hypotheses of Theorem 1.11 that imply the unit, monoid, and pushout product axioms, and in particular give model structures on categories of R-modules and R-algebras. Since a commutative R-algebra is just a map out of R in the commutative ring category, the theorem also constructs model structures on categories of commutative R-algebras.

Several theorems in this section assert symmetric monoidal model structures (in some cases under extra hypotheses). When the \mathcal{A} -variant \mathcal{V} -constrained \mathcal{F} -model structure is symmetric monoidal, then the model is enriched over itself in the sense of [21, 3.3], and *a fortiori* enriched over the \mathcal{V} -constrained \mathcal{F} -model category of Gspectra. We caution that it is not always the case that the \mathcal{A} -variant \mathcal{V} -constrained \mathcal{F} -model category of G-spectra is enriched over a model structure on the category of non-equivariant spectra: this depends on whether the objects in \mathcal{A} smashed with the non-equivariant desuspension spectra $\mathcal{F}_{\mathbb{R}^n} S^0$ are cofibrant in the \mathcal{A} -variant \mathcal{V} constrained \mathcal{F} -model structure. (This does not hold for example when G is finite or $G = \mathbb{T}$ and $A = \mathcal{A}^{\natural}(\mathcal{F}, \mathcal{V}_{reg})$ for any family \mathcal{F} containing a non-trivial subgroup.) Using the formulation of enrichment in terms of the Pushout-Tensor Axiom of [21, 3.4], this is clearly a necessary condition and given the characterization of the cofibrations, it is easy to see that it also sufficient.

PROPOSITION 1.14. With notation as in Theorem 1.11, assume that for every $n \geq 1$ and every $A \in \mathcal{A}$, the G-spectrum $A \wedge F_{\mathbb{R}^n} S^0$ is cofibrant in the \mathcal{A} -variant \mathcal{V} -constrained \mathcal{F} -model structure; then this model structure is enriched over the standard model structure on non-equivariant spectra in the sense of [21, 3.3].

2. The multiplicative geometric fixed point functor on variant model structures and Theorem E

The purpose of this section is to provide enough examples of \mathcal{A} -variant \mathcal{V} constrained model structures to be able to construct the left derived functor of the
multiplicative geometric fixed point functor Φ^H on the category of commutative
ring *G*-spectra. As a consequence, we prove Theorem E from the introduction;
at the end of the section, we review the individual definitions and results to give
a self-contained description of a model structure with the properties asserted in
Theorem E.

Some of the proofs in this section involve the properties of the generalized orbit desuspension spectra studied in Chapter V–VI. The background needed on the theory of generalized orbit desuspension spectra is Proposition V.1.4, which relates the symmetric power construction to the theory and Theorem V.2.4, which under a mild faithfulness hypothesis (Definition V.2.2) relates the geometric fixed point construction to the theory. The faithfulness hypothesis of Definition V.3.3 is also needed in places. With this background, the arguments below rely on specific references to results in Chapter V.

We begin with the following basic result.

THEOREM 2.1. Let G be a compact Lie group, \mathcal{F}, \mathcal{V} a family representation constraint with \mathcal{V} standard compatible, and \mathcal{A} a set of spacewise non-degenerately based G-spectra with $\mathcal{A}^{\natural}(\mathcal{F}, \mathcal{V}) \subset \mathcal{A}$. Fix a closed subgroup H < G and a family \mathcal{F}_{WH} of closed subgroups of WH. Assume that for every $A \in \mathcal{A}$ and every cofibrant approximation $\tilde{A} \to A$ in the \mathcal{V} -constrained \mathcal{F} -model structure, the induced map $\Phi^H \tilde{A} \to \Phi^H A$ is an \mathcal{F}_{WH} -equivalence. Then Φ^H sends \mathcal{F} -equivalences between cofibrant objects in the \mathcal{A} -variant \mathcal{V} -constrained \mathcal{F} -model structure to \mathcal{F}_{WH} equivalences of WH-spectra. Moreover, for any cofibrant object X in the \mathcal{A} -variant \mathcal{V} -constrained \mathcal{F} -model structure and any cofibrant approximation $\tilde{X} \to X$ in the \mathcal{V} -constrained \mathcal{F} -model structure, the induced map $\Phi^H \tilde{X} \to \Phi^H X$ is an \mathcal{F}_{WH} equivalence.

PROOF. Let $\mathcal{A}^{\Phi H}$ be the set of WH-spectra

$$\mathcal{A}^{\Phi H} = \{ \Phi^H A \mid A \in \mathcal{A} \}.$$

By transfinite induction, Φ^H sends an I_A -regular cofibration to the corresponding $\mathcal{I}_{A^{\Phi H}}$ -regular cofibration. Standard arguments then prove the statements. \Box

COROLLARY 2.2. Let G be a compact Lie group, and \mathcal{F}, \mathcal{V} a family representation constraint with \mathcal{V} standard compatible. Let \mathcal{A} be a set of spacewise nondegenerately based G-spectra satisfying the hypotheses of Theorem 1.13 that is closed under nth symmetric powers for all $n \geq 1$. Fix a closed subgroup H < G and a family \mathcal{F}_{WH} of closed subgroups of WH. Assume that for every $A \in \mathcal{A}$ and every cofibrant approximation $\tilde{A} \to A$ in the \mathcal{V} -constrained \mathcal{F} -model structure, the induced map $\Phi^H \tilde{A} \to \Phi^H A$ is an \mathcal{F}_{WH} -equivalence. Then Φ^H sends \mathcal{F} -equivalences between cofibrant objects in the \mathcal{A} -variant \mathcal{V} -constrained \mathcal{F} -model structure on commutative ring G-spectra to \mathcal{F}_{WH} -equivalences of commutative ring WH-spectra. Moreover, for any commutative ring G-spectrum R that is cofibrant in the \mathcal{A} -variant \mathcal{V} -constrained \mathcal{F} -model structure and any cofibrant approximation $\tilde{X} \to R$ in the \mathcal{V} -constrained \mathcal{F} -model structure on G-spectra, the induced map $\Phi^H \tilde{X} \to \Phi^H R$ is an \mathcal{F}_{WH} -equivalence.

PROOF. By Theorem 1.13, if R is cofibrant in the A-variant \mathcal{V} -constrained \mathcal{F} model structure of G-equivariant commutative ring orthogonal spectra, the unit
map $\mathbb{S} \to R$ is a cofibration of the underlying G-spectra in the A-variant \mathcal{V} constrained \mathcal{F} -model structure. The statements then follow from the previous
theorem.

To give the previous theorems substance, we need examples where the hypotheses hold. Here is one such example.

THEOREM 2.3. Let G be a compact Lie group. Let \mathcal{F} be any family of closed subgroups of G, H any closed subgroup of G, and \mathcal{F}_{WH} any family of closed subgroups of WH. Let \mathcal{V} be a family representation constraint for \mathcal{F} that is standard compatible with the property that for every $K \in \mathcal{F}$, every $[V] \in \mathcal{V}(K)$ satisfies the following conditions:

- (1) V contains a copy of the trivial representation, and
- (2) If K < H and contains the identity component of H then V contains a copy of the regular representation of $\pi_0 K$.

Let \mathcal{A} be the closure of $\mathcal{A}^{\natural}(\mathcal{F}, \mathcal{V})$ or $\mathcal{A}^{\circ}(\mathcal{F}, \mathcal{V})$ under nth symmetric power $(n \geq 1)$ and (finite, non-empty) smash products. Then \mathcal{F} , V, H, \mathcal{F}_{WH} , and \mathcal{A} satisfy the hypotheses of Corollary 2.2.

PROOF. We need to check the following properties hold for every $A \in \mathcal{A}$:

- (i) For every $n \ge 1$, $A^{(n)}/\Sigma_n$ is flat for the smash product of G-spectra.
- (ii) For every $n \ge 1$ and every cofibrant approximation $\tilde{A} \to A$ in the \mathcal{V} constrained \mathcal{F} -model structure, $\tilde{A}^{(n)}/\Sigma_n \to A^{(n)}/\Sigma_n$ is an \mathcal{F} -equivalence.
- (iii) For every cofibrant approximation $\tilde{A} \to A$ in the \mathcal{V} -constrained \mathcal{F} -model structure, $\Phi^H \tilde{A} \to \Phi^H A$ is an \mathcal{F}_{WH} -equivalence.

We use the theory of Chapter V, and particularly the theorems in Section V.2, with K the trivial group and $G = \Gamma = \Gamma/K$. We note that by induction, every $A \in \mathcal{A}$ can be expressed as a generalized orbit desuspension spectrum $J\eta$ for some (G, Q) vector bundle η of the form

$$\eta_1^{n_1} \times \cdots \times \eta_r^{n_r}$$

where $Q < \Sigma_{n_1} \times \cdots \times \Sigma_{n_r}$ and each η_i is of the form

$$G \times_K V \longrightarrow G/K$$

for $K \in \mathcal{F}$ and $[V] \in \mathcal{V}(H)$. To apply the results of Section V.2, we need to check that η is *Q*-faithful and for some results that $\eta(H)$ (in the notation of Definition V.2.10) is *Q*-faithful. For this, consider the bundle

$$\theta = \eta_1 \amalg \cdots \amalg \eta_r$$

and the bundle $\operatorname{Sym}_Q^n \theta$ (in the notation of Proposition V.1.4) where $n = n_1 + \cdots + n_r$ and we regard Q as a subgroup of Σ_n via block sum. Then η is obtained from $\operatorname{Sym}_Q^n \theta$ by the restriction (pullback) along the inclusion of a $(G \times Q^{\operatorname{op}})$ -stable subset of the base; likewise $\eta(H)$ is obtained from $\operatorname{Sym}_Q^n \theta(H)$ by the restriction along the inclusion of a $(G \times Q^{\operatorname{op}})$ -stable subset of the base. Thus (by Proposition V.2.3.(ii)), to see that η and $\eta(H)$ are Q-faithful, it is enough to show that $\operatorname{Sym}_Q^n \theta$ and $\operatorname{Sym}_Q^n \theta(H)$ are Q-faithful. Proposition V.3.2 (and Proposition V.2.3.(i)) shows that $\operatorname{Sym}_Q^n \theta$ is Q-faithful. Proposition V.3.6.(b) applies to θ (with $\Lambda = H$) because of the requirement on regular representations in \mathcal{V} and so by Theorem V.3.5, θ is inheritably faithful for Φ^H . This implies that $\operatorname{Sym}^n \theta(H)$ is Σ_n -faithful, which implies that $\operatorname{Sym}_Q^n \theta(H)$ is Q-faithful (Proposition V.3.4).

By Theorem V.2.9, we then have that A is flat for all $A \in \mathcal{A}$, and since \mathcal{A} is closed under symmetric power, property (i) holds. Property (ii) follows by Theorem V.2.8. Property (iii) holds by Theorem V.2.7.

Such a representation constraint always exists: for example, the Peter-Weyl theorem implies that for any orthogonal H-representation V there exists an orthogonal G-representation W whose underlying H-representation contains a copy of V. Applying this with V the regular representation of $\pi_0 H$ and taking $\mathcal{V}(K)$ (for all $K \in \mathcal{F}$) to be the set of isomorphism classes of orthogonal K-representations that contain a copy of the underlying K-representation of W gives a representation constraint satisfying the hypothesis of the theorem.

The previous theorem is general enough that it establishes the existence of derived functor of Φ^H on commutative ring *G*-spectra and a comparison with the derived functor of Φ^H on *G*-spectra.

COROLLARY 2.4. For any compact Lie group G and any closed subgroup H, the geometric H fixed point functor from commutative ring G-spectra to commutative ring WH-spectra has a left derived functor, which lifts the derived geometric H fixed point functor from G-spectra to commutative ring WH-spectra.

The previous result only treats the geometric fixed points of a single subgroup (or a finite set of subgroups) at a time. We also have the following variant, which handles all finite subgroups at the same time. The proof is identical, replacing part (b) of Proposition V.3.6 with part (a) (applied with Q the trivial group).

THEOREM 2.5. Let G be a compact Lie group. Let \mathcal{F} be any family of finite subgroups, and let \mathcal{V} be a family representation constraint for \mathcal{F} that is standard compatible and that has the property that for every subgroup $K \in \mathcal{F}$, every $[V] \in \mathcal{V}(K)$ contains the regular representation of K. Let \mathcal{A} be the closure of $\mathcal{A}^{\circ}(\mathcal{F}, \mathcal{V})$ under nth symmetric power $(n \geq 1)$ and smash product. Then \mathcal{F} , V, and \mathcal{A} satisfy the hypotheses of Corollary 2.2 for any finite subgroup H of G and any family \mathcal{F}_{WH} of closed subgroups of WH.

In both Theorems 2.3 and 2.5, the cofibrant objects are built from cells of the form $A \wedge (D^n, \partial D^n)_+$ where A is a generalized orbit desuspension spectrum $J\eta$ for some (G, Q) vector bundle (for some finite group Q). Using the additional properties of generalized orbit desuspension spectra from Chapter V, we see that the cofibrant objects in these model structures have additional good properties, which we catalog below. In the statements below, H denotes the particular subgroup in Theorem 2.3 or a subgroup in \mathcal{F} in Theorem 2.5. In each case all that is needed is that η and $\eta(H)$ are Q-faithful and have base spaces isomorphic to $(G \times Q^{\text{op}})$ -cell complexes. Theorem V.2.9 implies:

PROPOSITION 2.6. Suppose X is cofibrant or cofibrant under S in the model structure of Theorem 2.3 or 2.5. Then X is flat for the smash product of G-spectra and $\Phi^H X$ is flat for the smash product of WH-spectra.

Theorem V.2.6 implies:

PROPOSITION 2.7. Suppose X is cofibrant or cofibrant under S in the model structure of Theorem 2.3 or 2.5. Then for any G-spectrum Y, the canonical map

$$\Phi^H X \wedge \Phi^H Y \longrightarrow \Phi^H (X \wedge Y)$$

 $is \ an \ isomorphism.$

The previous proposition and a cell by cell argument also gives the following result.

PROPOSITION 2.8. Let R be an associative ring G-spectrum. Let M and N be left and right R-modules (respectively) and assume one of M, N is cofibrant either in the standard model structure, the model structure of Theorem 2.3, or the model structure of Theorem 2.5 for its respective category of R-modules. Then the canonical map

$$\Phi^H M \wedge_{\Phi^H R} \Phi^H N \longrightarrow \Phi^H (M \wedge_R N)$$

is an isomorphism.

Model structures for Theorem E. Theorem 2.3 combined with Theorem 1.13 gives a model structure that works for an arbitrary finite set \mathcal{H} of closed subgroups. The following is a step-by-step guide through the definitions and results above to see this. (We now assume that \mathcal{H} is non-empty, as otherwise there is nothing to show.)

Let W be an orthogonal G-representation with the property that for every $H \in \mathcal{H}$, the restriction of W as an orthogonal H-representation contains a copy of the regular representation of $\pi_0 H$. Let \mathcal{A}_W^{\natural} be the set of spectra of the form

$$F_V(G/K)_+$$

for K a closed subgroup of G and V an orthogonal G-representation containing W. Let $\mathcal{A}_W^{(\wedge)}$ denote the closure of \mathcal{A}_W^{\natural} under (finite, non-empty) smash products and symmetric powers: an element of $\mathcal{A}_W^{(\wedge)}$ is in particular a spectrum of the form

$$((F_{V_1}(G/K_1)_+)^{(q_1)} \wedge \cdots \wedge (F_{V_r}(G/K_r)_+)^{(q_r)})/Q$$

where $r \geq 1$, $q_i \geq 1$, each V_i contains W, and Q is a subgroup of the product of symmetric groups $\Sigma_{q_1} \times \cdots \times \Sigma_{q_r}$. The objects in \mathcal{A}_W^{\natural} are topologically point-set small (Definition 1.7) because $F_V(G/K)_+$ is and the class of topologically point-set small objects is closed under smash products and quotients.

We get two model structures on G-spectra with $\mathcal{A} = \mathcal{A}_W^{\natural}$ and $\mathcal{A} = \mathcal{A}_W^{(\wedge)}$, resp., where:

- The cofibrations are the maps $X \to Y$ that are retracts of relative \mathcal{A} -cell complexes (see Definition 1.6).
- The weak equivalences are the standard weak equivalences

In the case of \mathcal{A}_W^{\natural} , a map $X \to Y$ is a fibration if and only if for every orthogonal G-representation V containing W, the map $X(V) \to Y(V)$ is a G-equivariant Serre fibration and for every orthogonal G-representation U containing V (which contains W) the diagram

$$\begin{array}{c} X(V) \longrightarrow \Omega^{U-V} X(U) \\ \downarrow \qquad \qquad \downarrow \\ Y(V) \longrightarrow \Omega^{U-V} Y(U) \end{array}$$

is homotopy cartesian. In the case of $\mathcal{A}_W^{(\wedge)}$, a map $X \to Y$ is a fibration if and only if:

- $X \to Y$ is a fibration in the model structure for \mathcal{A}_W^{\natural} ,
- For every $A \in \mathcal{A}_W^{(\wedge)}$, the map $\mathcal{S}^G(A, X) \to \mathcal{S}^G(A, Y)$ is a Serre fibration, and
- For every $A \in \mathcal{A}_W^{(\wedge)}$, there exists a cofibrant object \tilde{A} in the model structure for \mathcal{A}_W^{\natural} and a weak equivalence $\tilde{A} \to A$ such that the diagram

$$\begin{array}{c} \mathcal{S}^G(A,X) \longrightarrow \mathcal{S}^G(\tilde{A},X) \\ & \downarrow \\ \mathcal{S}^G(A,Y) \longrightarrow \mathcal{S}^G(\tilde{A},Y) \end{array}$$

is homotopy cartesian.

See for example Theorem 1.9 for \mathcal{A}_W^{\natural} and Theorem 1.11 with Proposition 1.12 for $\mathcal{A}_W^{(\wedge)}$.

The identity gives a zigzag of Quillen equivalences of model structures relating these two model structures and the standard (or positive) stable model structure on *G*-spectra: the identity is a Quillen left adjoint from the model structure for \mathcal{A}_W^{\natural} to the standard (or positive) model structure and to the model structure for \mathcal{A}_W^{\natural} .

Theorem 2.3 implies that every $A \in \mathcal{A}_W^{(\wedge)}$ satisfies the following properties:

- For any $n \ge 1$, $A^{(n)}/\Sigma_n$ is flat for the smash product of G-spectra
- For any $n \ge 1$ and any cofibrant approximation $\tilde{A} \to A$ in the model category for \mathcal{A}_W^{\natural} , the induced map $\tilde{A}^{(n)}/\Sigma_n \to A^{(n)}/\Sigma_n$ is a weak equivalence.

(While the statement of Theorem 2.3 is peripatetic in this regard, this assertion is also made in the first few lines of its proof.) Since $\mathcal{A}_W^{\natural} \subset \mathcal{A}_W^{(\wedge)}$, Theorem 1.13 applies to show that the forgetful functor from commutative ring *G*-spectra to *G*spectra creates a model structure using either the model structure for \mathcal{A}_W^{\natural} or the model structure for $\mathcal{A}_W^{(\wedge)}$; moreover, using the latter model structure, cofibrant commutative ring *G*-spectra are cofibrant in the undercategory of S in *G*-spectra.

Now assume that R is a cofibrant commutative ring G-spectrum in the model structure for $\mathcal{A}_W^{(\wedge)}$ (which holds in particular if R a cofibrant commutative ring Gspectrum in the model structure for \mathcal{A}_W^{\natural}). We need to show that for any cofibrant approximation $X \to R$ in the standard model structure on G-spectra, the induced map $\Phi^H X \to \Phi^H R$ is a weak equivalence for every $H \in \mathcal{H}$. Since Φ^H preserves weak equivalences between cofibrant objects in the standard model structure, it suffices to show that there exists a cofibrant approximation with this property. Let $X \to R$ be a cofibrant approximation of R in the undercategory of S in the model structure for \mathcal{A}_W^{\natural} on G-spectra; then $X \to R$ is a cofibrant approximation in the standard model structure on the (absolute) category of G-spectra. The inclusions of S in X and R are Hurewicz cofibrations and Φ^H preserves Hurewicz cofibrations and point-set quotients by Hurewicz cofibrations, so we get a map of long exact sequences of homotopy groups

$$\begin{array}{c} \cdots \longrightarrow \pi_q^K(\Phi^H \mathbb{S}) \longrightarrow \pi_q^K(\Phi^H X) \longrightarrow \pi_q^K(\Phi^H(X/\mathbb{S})) \longrightarrow \pi_{q-1}^K(\Phi^H \mathbb{S}) \longrightarrow \cdots \\ = \downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow \qquad \qquad = \downarrow \\ \cdots \longrightarrow \pi_q^K(\Phi^H \mathbb{S}) \longrightarrow \pi_q^K(\Phi^H R) \longrightarrow \pi_q^K(\Phi^H(R/\mathbb{S})) \longrightarrow \pi_{q-1}^K(\Phi^H \mathbb{S}) \longrightarrow \cdots \end{array}$$

for every K < WH (for example, by [22, III.3.5.(iv),(vi)]). The map $X/\mathbb{S} \to Y/\mathbb{S}$ is a weak equivalence of cofibrant objects in the model structure for $\mathcal{A}_W^{(\wedge)}$ on *G*-spectra and so induces a weak equivalence on Φ^H by Corollary 2.2, which holds by Theorem 2.3. It follows that $\Phi^H X \to \Phi^H R$ is a weak equivalence.

CHAPTER III

The Circle Group and Cyclotomic Spectra (Theorems B, C, D)

This chapter treats the case when the compact Lie group is the circle group \mathbb{T} of unit complex numbers. In this case, we have significantly better results on geometric fixed point functors than we have for general compact Lie groups. We study this in Section 1, where we prove Theorem C. We then apply the results on the circle group to study ppc and pc spectra in Section 2, where we prove Theorems B and D. The construction of the model structure used in these sections is stated in terms of a black box set of \mathbb{T} -spectra \mathcal{A}^{cyc} (and its subset \mathcal{A}_p^{cyc}); Section 3 opens this box, filling in the details and proving the equivariant properties of the \mathbb{T} -spectra in \mathcal{A}^{cyc} needed to prove the geometric fixed point results. Section 4 fills in the properties of \mathcal{A}^{cyc} needed to compare the model structure on \mathbb{T} -spectra to the positive convenient Σ -model structure on non-equivariant spectra.

1. The circle group and Theorem C

The purpose of this section is to begin the proof of Theorem C from the introduction. Here we have broken it into two parts: Theorem 1.2 below generalizes the model structure statement of Theorem C and its properties that involve just the equivariant categories, while Theorem 1.3 restates the strong relationship asserted between the model structures on the equivariant and non-equivariant categories. As in the statement of Theorem C, we let \mathcal{F}_{fin} denote the family of finite subgroups of T and \mathcal{F}_p denote the family of finite *p*-subgroups of T. In contrast, we use slightly different notation for the geometric fixed point endofunctor denoted Φ in the introduction in order to consider the construction for more general subgroups of T.

NOTATION 1.1. Define endofunctors Φ_n , $n \ge 1$ by

$$\Phi_n := \rho_n^* \Phi^{C_n}.$$

Here we first take the geometric fixed points Φ^{C_n} to \mathbb{T}/C_n -spectra and then use the *n*th root isomorphism $\rho_n \colon \mathbb{T} \cong \mathbb{T}/C_n$ to convert back to \mathbb{T} -spectra.

To establish the model structure, we use the general result of Theorem II.1.13, applied with the family \mathcal{F}_{fin} or \mathcal{F}_p and the representation constraint $\mathcal{V}_{\pi!}$ from Example II.1.2. Namely, for $C \in \mathcal{F}_{\text{fin}}$ (or \mathcal{F}_p), $\mathcal{V}_{\pi!}(C)$ consists of the semiregular permutation representations, that is, those orthogonal *C*-representations which admit an orthonormal basis to which the *C* action restricts, and which has at least one free orbit. The set of model cells that we take for \mathcal{A} in Theorem II.1.13 is denoted \mathcal{A}^{cyc} (in the case of \mathcal{F}_{fin}) and $\mathcal{A}_p^{\text{cyc}}$ (in the case of \mathcal{F}_p). The \mathbb{T} -spectra in these sets are complicated to specify, but have very nice properties, as stated in Lemma 1.4. We postpone the details of the construction of these \mathbb{T} -spectra to Section 3, as the list of properties suffices for what we need here.

THEOREM 1.2. The sets \mathcal{A}^{cyc} and \mathcal{A}^{cyc}_p of \mathbb{T} -spectra in Definition 3.3 below satisfy the hypotheses of Theorem II.1.13 for $\mathcal{F} = \mathcal{F}_{fin}$, $\mathcal{V} = \mathcal{V}_{\pi!}$ and for $\mathcal{F} = \mathcal{F}_p$, $\mathcal{V} = \mathcal{V}_{\pi!}$, respectively. In particular, the forgetful functor from commutative ring \mathbb{T} -spectra to \mathbb{T} -spectra creates model structures for both the \mathcal{A}^{cyc} -variant $\mathcal{V}_{\pi!}$ -constrained \mathcal{F}_{fin} -model structure and the \mathcal{A}^{cyc}_p -variant $\mathcal{V}_{\pi!}$ -constrained \mathcal{F}_p -model structure. These model structures are enriched over the standard model structure in non-equivariant spectra. Moreover in these variant model structures:

- (i) For A^{cyc}, the endofunctors Φ_n preserve cofibrations and acyclic cofibrations of both T-spectra and commutative ring T-spectra. For A^{cyc}_p, the endofunctors Φ_{pⁿ} preserve cofibrations and acyclic cofibrations of both T-spectra and commutative ring T-spectra.
- (ii) If R is a cofibrant commutative ring T-spectrum, then S → R is a cofibration of T-spectra.

The remainder of Theorem C is the relationship to non-equivariant spectra. We state the following result here but prove it in Section 4.

THEOREM 1.3. The forgetful functor from \mathbb{T} -spectra to non-equivariant spectra takes cofibrant objects in the \mathcal{A}^{cyc} -variant $\mathcal{V}_{\pi^{\dagger}}$ -constrained \mathcal{F}_{fin} -model structure to cofibrant objects in the positive convenient Σ -model structure of Definition 4.1. The free functor from non-equivariant spectra to \mathbb{T} -spectra takes cofibrant objects in the positive convenient Σ -model structure on spectra of Definition 4.1 to cofibrant objects in the \mathcal{A}_p^{cyc} -variant $\mathcal{V}_{\pi^{\dagger}}$ -constrained \mathcal{F}_p -model structure on \mathbb{T} -spectra.

The hypotheses of Theorem II.1.13 imply in particular the last two bullet point hypotheses of Theorem II.1.11, which give the unit, monoid, and pushout-product axioms of [25, 3.2–4], which then give model structures on module and algebra categories. When R is a commutative ring T-spectrum whose underlying T-spectrum is cofibrant in the undercategory of S, then cofibrant R-algebras are cofibrant as T-spectra in the undercategory of S and cofibrant R-modules are cofibrant as T-spectra in these model structures.

The definition of \mathcal{A}^{cyc} and \mathcal{A}_{p}^{cyc} is in terms of specific generalized orbit desuspension spectra described in detail in Section 3. What we need for the proof of the theorem is given by the following lemma, proved in Section 3.

LEMMA 1.4. Let $A \in \mathcal{A}^{\text{cyc}}, k \geq 1$.

- (i) A is $i^* J^{\Gamma/K} \theta_A$, for some Γ/K -compatible Q-faithful positive dimensional (Γ, Q) vector bundle θ_A , where $K \lhd \Gamma$ is a finite normal subgroup, i is an isomorphism $i: \mathbb{T} \cong \Gamma/K$, and the vector bundle θ_A has base space a finite product of orbits of Γ . Moreover, for every $K < \Lambda \lhd \Gamma$, the (Γ, Q) vector bundle $\theta_A(\Lambda|K)$ is Q-faithful.
- (ii) If $B \in \mathcal{A}^{cyc}$, then $A \wedge B$ is isomorphic to an element of \mathcal{A}^{cyc} ; if $A, B \in \mathcal{A}_{p}^{cyc}$, the $A \wedge B$ is isomorphic to an element of \mathcal{A}_{p}^{cyc} .
- (iii) $\Phi_m A$ is isomorphic to an element of \mathcal{A}^{cyc} ; if $A \in \mathcal{A}_p^{cyc}$ and m is a power of p, then $\Phi_m A$ is isomorphic to an element of \mathcal{A}_p^{cyc} .
- (iv) $A^{(k)}/\Sigma_k$ is isomorphic to an element of \mathcal{A}^{cyc} ; if $A \in \mathcal{A}_p^{cyc}$, then $A^{(k)}/\Sigma_k$ is isomorphic to an element of \mathcal{A}_p^{cyc} .

- (v) There exists a cofibrant object \tilde{A} in the complete positive model structure on \mathbb{T} -spectra and a weak equivalence $\tilde{A} \to A$ such that the induced map $\tilde{A}^{(k)}/\Sigma_k \to A^{(k)}/\Sigma_k$ is a weak equivalence.
- (vi) For any $n \ge 1$, $A \wedge F_{\mathbb{R}^n} S^0$ is isomorphic to an object of \mathcal{A}^{cyc} , which is in \mathcal{A}_p^{cyc} if A is.

As a consequence of the lemma, objects of \mathcal{A}^{cyc} behave well with respect to smash product and the functors Φ_n . For other statements, it is useful to provide a label for the spectra with the properties listed below.

DEFINITION 1.5. Let \mathcal{E}_{Φ} be the class of \mathbb{T} -spectra A that satisfy all of the following properties:

- (i) For every $m \ge 1$, $\Phi_m A$ is flat for the smash product of T-spectra.
- (ii) For every $m \geq 1$ and every cofibrant approximation $\tilde{A} \to A$ in the $\mathcal{V}_{\pi!}$ -constrained \mathcal{F}_{fin} -model structure, the map $\Phi_m \tilde{A} \to \Phi_m A$ is a \mathcal{F}_{fin} -equivalence.
- (iii) For every $m, n \ge 1$, the canonical map $\Phi_{mn}A \to \Phi_n \Phi_m A$ of [4, 2.6] is an isomorphism.
- (iv) For every $m, n \ge 1$ and every \mathbb{T} -spectrum X, the canonical map

$$\Phi_n(\Phi_m A) \land \Phi_n X \longrightarrow \Phi_n(\Phi_m A \land X)$$

is an isomorphism.

It is clear that \mathcal{E}_{Φ} is closed under Φ_m , retracts, smash products, pushouts along Hurewicz cofibrations, and transfinite compositions of Hurewicz cofibrations. We note that $\mathbb{S} \in \mathcal{E}_{\Phi}$ and more generally, the cofibrant objects in the standard stable and complete model structures on \mathbb{T} -spectra are in \mathcal{E}_{Φ} .

PROPOSITION 1.6. $\mathcal{A}^{cyc} \subset \mathcal{E}_{\Phi}$.

PROOF. This is simplified by remembering that if $A \in \mathcal{A}^{cyc}$, then $\Phi_n A$ is isomorphic to an object of \mathcal{A}^{cyc} by Lemma 1.4.(ii). Applying Lemma 1.4.(i), we see that objects of \mathcal{A}^{cyc} have property (i) by Theorem V.2.9, have property (ii) by Theorem V.2.7, have property (iii) by Theorem V.2.5, and have property (iv) by Theorem V.2.6.

Before beginning the proof of Theorem 1.2, we state some auxiliary results that simplify some aspects of the proof and are also needed in other contexts. The first allows us to use relative cell complexes in place of regular cofibrations. It is an easy consequence of the fact that in the equivariant vector bundles used to construct objects of \mathcal{A}^{cyc} , the base spaces are compact.

PROPOSITION 1.7. Objects in \mathcal{A}^{cyc} are topologically point-set small.

The other result we need regards the interaction of the balanced smash product with the endofunctors Φ_n . In the statement, the undercategory case is designed to handle cofibrations of commutative ring orthogonal spectra: if $R \to A$ is a cofibration of commutative ring orthogonal spectra in one of the model categories of Theorem 1.2, then $R \to A$ is a cofibrant object in the undercategory of R in the category of (symmetric) R-modules.

PROPOSITION 1.8. Let R be a associative ring \mathbb{T} -spectrum. Let M be a right R-module and N a left R-module. If either M or N is cofibrant in the respective

R-module model structure (from Theorem 1.2) or in the undercategory of R in the respective R-module model structure, then for any $m \ge 1$, the canonical map

$$\Phi_m M \wedge_{\Phi_m R} \Phi_m N \longrightarrow \Phi_m (M \wedge_R N)$$

is an isomorphism.

PROOF. Without loss of generality, we can assume that the cofibrant object is a cell complex. The R-module cells are then in particular of the form

$$R \wedge A \wedge (D^n, \partial D^n)_+$$

for $A \in \mathcal{A}^{\text{cyc}}$, with the left (for M) or right (for N) R-module structure on R. Working cell by cell, the result follows from property (iv) defining \mathcal{E}_{Φ} for the objects in \mathcal{A}^{cyc} .

We are now ready for the proof of Theorem 1.2.

PROOF OF THEOREM 1.2. First, to check the hypotheses of Theorem II.1.13, we observe the following:

- Smash product of objects of \mathcal{A}^{cyc} or \mathcal{A}^{cyc}_p are isomorphic to objects of \mathcal{A}^{cyc} or \mathcal{A}^{cyc}_p (Lemma 1.4.(ii)) and are therefore cofibrant in their variant model structures.
- The symmetric power of any object of \mathcal{A}^{cyc} is flat for the smash product by Lemma 1.4.(iv) and Proposition 1.6 (q.v. Definition 1.5.(i)).
- Any \mathcal{F}_{fin} or \mathcal{F}_p -equivalence $\tilde{A} \to A$ from a positive complete cofibrant object to an object of \mathcal{A}^{cyc} induces an \mathcal{F}_{fin} or \mathcal{F}_p -equivalence on symmetric powers by Lemma 1.4.(v).

The last bullet point holds because the symmetric power functor preserves \mathcal{F}_{fin} - and \mathcal{F}_p -equivalences between cofibrant objects in the positive complete model structure (e.g., by applying [15, B.111,B.116] and restricting the group from \mathbb{T} to C_m or C_{p^k}). Objects that are cofibrant in the $\mathcal{V}_{\pi!}$ -constrained \mathcal{F}_{fin} -model structure and in the $\mathcal{V}_{\pi!}$ -constrained \mathcal{F}_p -model structure are in particular cofibrant in the positive complete model structure, so the last bullet point in particular implies hypothesis (ii) of Theorem II.1.13. We therefore get $\mathcal{A}_p^{\text{cyc}}$ and $\mathcal{A}_p^{\text{cyc}}$ variant model structures on \mathbb{T} -equivariant commutative ring spectra.

To check the enrichment statement, we apply Lemma 1.4.(vi) and Proposition II.1.14.

To check statement (ii), that cofibrant commutative ring T-spectra have underlying T-spectra cofibrant in the undercategory of S, we need to see that \mathcal{A}_{p}^{cyc} and \mathcal{A}_{p}^{cyc} are closed up to isomorphism under symmetric powers; this is Lemma 1.4.(iv).

To check statement (i) in the case of \mathbb{T} -spectra, we use the fact that a cofibration is a retract of a relative cell complex, with cells of the form

 $A \wedge (D^n, \partial D^n)_+$

for A in \mathcal{A}^{cyc} or $\mathcal{A}_p^{\text{cyc}}$. Because \mathcal{A}^{cyc} is closed up to isomorphism under Φ_n , Φ_n preserves cell attachments of this form (with A replaced by $\Phi_n A$). Likewise $\mathcal{A}_p^{\text{cyc}}$ is closed up to isomorphism under Φ_{p^n} . The endofunctors Φ_n preserve Hurewicz cofibrations, and pushouts along Hurewicz cofibrations give Mayer-Vietoris sequences of homotopy groups in \mathbb{T} -spectra, so Φ_n and Φ_{p^n} also preserve acyclic cofibrations in the \mathcal{A}^{cyc} -variant $\mathcal{V}_{\pi!}$ -constrained \mathcal{F}_{fin} -model structure and $\mathcal{A}_p^{\text{cyc}}$ -variant $\mathcal{V}_{\pi!}$ -constrained \mathcal{F}_p -model structure on \mathbb{T} -spectra, respectively.

To check statement (i) in the case of commutative ring \mathbb{T} -spectra, the main new tool is Theorem V.3.1. The usual retract and colimit argument reduces the statement to showing that applying Φ_n to a cell attachment in this model structure

$$R \longrightarrow R \wedge_{\mathbb{P}(A \wedge \partial D^k_+)} (\mathbb{P}(A \wedge D^k_+))$$

gives a cofibration in this model structure. (Here \mathbb{P} denotes the free functor from \mathbb{T} -spectra to commutative ring \mathbb{T} -spectra.) By Proposition 1.8, we have

$$\Phi_n(R \wedge_{\mathbb{P}(A \wedge \partial D^k_{\pm})} (\mathbb{P}(A \wedge D^k_{\pm}))) \cong (\Phi_n R) \wedge_{\Phi_n(\mathbb{P}(A \wedge \partial D^k_{\pm}))} \Phi_n(\mathbb{P}(A \wedge D^k_{\pm})),$$

so it suffices to check that

$$\Phi_n(\mathbb{P}(A \wedge \partial D^k_+)) \longrightarrow \Phi_n(\mathbb{P}(A \wedge D^k_+))$$

is a cofibration when $A \in \mathcal{A}^{\text{cyc}}$. By Theorem V.3.1, this map is the retract of a finite smash product of maps of the form

$$\mathbb{P}(X \wedge (\partial D^k)^q_+) \longrightarrow \mathbb{P}(X \wedge (D^k)^q_+)$$

where X is a retract of $A^{(q)}/\Sigma_q$. Since $A^{(q)}/\Sigma_q$ is a isomorphic to an object of \mathcal{A}^{cyc} , X is a retract of an object of \mathcal{A}^{cyc} , and the map is a cofibration in the \mathcal{A}^{cyc} -variant model structure for commutative ring T-spectra. Any acyclic cofibration is a transfinite composition of maps, each of which is formed as a pushout along an acyclic cofibration of cofibrant objects. Because as just shown Φ_n preserves cofibrations and because it preserves weak equivalences between cofibrant objects, it follows that Φ_n preserves acyclic cofibrations in the \mathcal{A}^{cyc} -variant model structure for commutative ring T-spectra. The argument for the case of $\mathcal{A}^{\text{cyc}}_p$ is entirely similar.

Model structures for Theorem C. Theorem 1.2 combined with Theorem II.1.13 gives a model structure that satisfies the conclusions of Theorem C. The following is a step-by-step guide through the definitions and results above to see this. Theorem C gives two sets of statements, one for \mathcal{F}_{fin} and one for \mathcal{F}_p ; let \mathcal{F} denote \mathcal{F}_{fin} or \mathcal{F}_p in the respective cases, and let \mathcal{A} denote the set \mathcal{A}^{cyc} or $\mathcal{A}_p^{\text{cyc}}$ of Definition 3.3 in the respective cases. Write \mathcal{A}° for the set of T-spectra of the form

$$\mathbb{T}_+ \wedge_C F_V S^0$$


where $C \in \mathcal{F}$ and V is an orthogonal C-representation whose isomorphism class is in $\mathcal{V}_{\pi!}(C)$: V is non-zero and has an orthonormal basis to which the C action restricts. Finally, let \mathcal{A}^{\natural} be the set of \mathbb{T} -spectra of the form $\mathbb{T}/C_+ \wedge F_V S^0$ where $C \in \mathcal{F}$ and V is an orthogonal \mathbb{T} -representation whose underlying isomorphism class of orthogonal C-representation is in $\mathcal{V}_{\pi!}(C)$. (We will more concisely write this condition as "V is in $\mathcal{V}_{\pi!}(C)$ " in what follows.)

We then get model structures on \mathbb{T} -spectra for $\mathcal{A}, \mathcal{A}^{\circ}$, and \mathcal{A}^{\natural} , where:

- The cofibrations are the maps $X \to Y$ that are retracts of relative \mathcal{A} -,
 - \mathcal{A}° , and \mathcal{A}^{\natural} -cell complexes, respectively (see Definition II.1.6).
- The weak equivalences are the \mathcal{F} -equivalences

In the case of \mathcal{A}^{\natural} (or \mathcal{A}°), a map $X \to Y$ is a fibration if and only if for every $C \in \mathcal{F}$ and every orthogonal \mathbb{T} -representation (or *C*-representation, respectively) V in $\mathcal{V}_{\pi!}(C)$, the map $X(V) \to Y(V)$ is a *C*-equivariant Serre fibration (a Serre fibration on all fixed point spaces) and for every orthogonal \mathbb{T} -representation (or

C-representation, respectively) W in $\mathcal{V}_{\pi!}(C)$ containing such a V, the diagram of C-spaces



is homotopy cartesian. In the case of \mathcal{A} , a map $X \to Y$ is a fibration if and only if:

- $X \to Y$ is a fibration in the model structure for \mathcal{A}^{\natural} ,
- For every $A \in \mathcal{A}$, the map $\mathcal{S}^G(A, X) \to \mathcal{S}^G(A, Y)$ is a Serre fibration, and
- For every $A \in \mathcal{A}$, there exists a cofibrant object \tilde{A} in the model structure for \mathcal{A}^{\natural} and a weak equivalence $\tilde{A} \to A$ such that the diagram

$$\begin{array}{c} \mathcal{S}^{G}(A,X) \longrightarrow \mathcal{S}^{G}(\tilde{A},X) \\ \downarrow \\ \mathcal{S}^{G}(A,Y) \longrightarrow \mathcal{S}^{G}(\tilde{A},Y) \end{array}$$

is homotopy cartesian.

(The fibrations in the model structure in the case of \mathcal{A} can alternatively be characterized in terms of \mathcal{A}° , just replacing \mathcal{A}^{\natural} by \mathcal{A}° in the statements.)

These model structures follow for example from Theorems II.1.9 and II.1.10 in the case of \mathcal{A}^{\ddagger} and \mathcal{A}° (respectively) and Theorem II.1.11 with Proposition II.1.12 and Proposition 1.7 in the case of \mathcal{A} .

The identity gives a zigzag of Quillen equivalences of model structures relating these two model structures and the standard (or positive) complete stable model structure on T-spectra: the identity is a Quillen left adjoint from the model structure for \mathcal{A}^{\natural} to the standard (or positive) model structure and to the model structure for \mathcal{A} and the identity is also a Quillen left adjoint from the model structure for \mathcal{A}° to the complete (or positive complete) model structure and to the model structure for \mathcal{A} . Writing the functors in the direction of Quillen left adjoints, we have the following zigzags of Quillen equivalences of model structures on the category of T-spectra with underlying functor the identity.

(1.9) $\begin{array}{c} \text{standard} \longleftarrow \text{positive} \longleftarrow \text{model}(\mathcal{A}^{\natural}) \longrightarrow \text{model}(\mathcal{A})\\ \text{complete} \longleftarrow \text{pos. complete} \leftarrow \text{model}(\mathcal{A}^{\circ}) \longrightarrow \text{model}(\mathcal{A}) \end{array}$

Theorem 1.2 now proves Theorem C: Theorem 1.2 asserts that Φ_n preserves cofibrations and acyclic cofibrations in the model structure for \mathcal{A} on \mathbb{T} -spectra, which is statement (i) of Theorem C.

Theorem 1.2 asserts that the hypotheses of Theorem II.1.13 hold which implies that the forgetful functor creates a topological model structure on commutative ring \mathbb{T} -spectra. This is statement (ii) of Theorem C.

Theorem 1.2 asserts that Φ_n preserves cofibrations in the model structure for \mathcal{A} on the category of commutative ring \mathbb{T} -spectra, and this in particular implies statement (iii) of Theorem C.

Theorem 1.2 asserts that in the model structures for \mathcal{A} , the forgetful functor takes the cofibrant commutative ring \mathbb{T} -spectra to cofibrant objects in the undercategory of \mathbb{S} in \mathbb{T} -spectra. This is statement (iv) of Theorem C.

The final part of Theorem C, relating the model structures on \mathbb{T} -spectra to the positive convenient Σ -model on non-equivariant spectra is proved as follows. Since cofibrations both equivariantly and non-equivariantly are formed as sequential colimits of pushouts along certain maps between certain cofibrant objects, Theorem 1.3 (and the fact that $\mathcal{A}_p^{\text{cyc}} \subset \mathcal{A}^{\text{cyc}}$) implies that both the forgetful functor and free functor preserve cofibrations. Since the free functor preserves all weak equivalences, by adjunction, the forgetful functor also preserves fibrations.

2. Pre-cyclotomic spectra and Theorems D and B

The purpose of this section is to prove Theorem B and Theorem D (and its addendum) using the results of the previous section.

The model structure on ppc spectra alluded to in the statement of Theorem D is the one arising out of Theorem 1.2 above. Here is the precise statement both in this case and in the case of pc spectra. The proof given in [4] for the model structure considered there extends to prove this case.

THEOREM 2.1. The category of ppc spectra (respectively, pc spectra) admits a topological model structure with fibrations and weak equivalences created by the forgetful functor to \mathbb{T} -equivariant spectra with the $\mathcal{A}_p^{\text{cyc}}$ -variant $\mathcal{V}_{\pi^{\text{l}}}$ -constrained \mathcal{F}_p model structure (resp., the \mathcal{A}^{cyc} -variant $\mathcal{V}_{\pi^{\text{l}}}$ -constrained \mathcal{F}_{fin} -model structure). This model structure is enriched over the standard model structure on non-equivariant spectra.

The generating cofibrations in the model structure are of the form $(\mathbb{C}A) \wedge (D^n, \partial D^n)_+$ where

$$\mathbb{C}A = \bigvee_{k \ge 0} \Phi_{p^k} A$$

for $A \in \mathcal{A}_p^{\text{cyc}}$, or in the pc case

$$\mathbb{C}A = \bigvee_{n \ge 1} \Phi_n A$$

for $A \in \mathcal{A}^{\text{cyc}}$. We see in both cases that these generating cofibrations are cofibrations in the corresponding model structure of Theorem 1.2, which implies the following proposition.

PROPOSITION 2.2. The forgetful functor from ppc spectra (resp., pc spectra) to \mathbb{T} -spectra preserves cofibrations and acyclic cofibrations for the relevant model structures in Theorem 2.1.

Proposition 1.6 implies that cofibrant objects in the model structure above are in the class \mathcal{E}_{Φ} of 1.5. By definition, if $X \in \mathcal{E}_{\Phi}$ and Y is any T-spectrum, then the canonical map


$$\Phi_n X \wedge \Phi_n Y \longrightarrow \Phi_n (X \wedge Y)$$

is an isomorphism. For X and Y ppc spectra whose underlying T-spectra are in \mathcal{E}_{Φ} , $X \wedge Y$ gets a ppc structure map

$$\Phi_p(X \wedge Y) \cong \Phi_p X \wedge \Phi_p Y \longrightarrow X \wedge Y$$

from the smash product of the ppc structure maps on X and Y. This constructs a symmetric monoidal product on the full subcategory of ppc spectra whose underlying objects are in \mathcal{E}_{Φ} . In particular, this establishes part (i) of Theorem D. The analogue for pc spectra holds by the same observation.

For part (ii) of Theorem D, we recall the definition of associative ring and commutative ring ppc spectra. An associative ring ppc (respectively, pc) spectrum consists of a ppc (resp., pc) spectrum A together with a *unit* map $\mathbb{S} \to A$ that is a map of ppc (resp., pc) spectra and a *multiplication* map of \mathbb{T} -spectra $A \wedge A \to A$ such that the unit and multiplication give A the structure of an associative ring \mathbb{T} -spectrum and the diagram



constructed from the multiplication and structure maps commutes (for all primes p in the pc case). An associative ring ppc (resp., pc) spectrum is a commutative ring when the multiplication is commutative.

Proposition 1.8 is the key to the construction of the model structure for commutative ring ppc spectra. The generating cofibrations in this model structure are obtained by applying the free commutative ring functor

$$\mathbb{P}X = \bigvee_{k \ge 0} X^{(k)} / \Sigma_k$$

to the generating cofibrations for the model structure on ppc spectra of Theorem 2.1. In the resulting map of commutative ring ppc spectra, both the source and target have underlying \mathbb{T} -spectra formed as the wedge of \mathbb{S} with a wedge of objects in \mathcal{A}_p^{cyc} (by Lemma 1.4), and in particular are in \mathcal{E}_{Φ} . If we write such a cell as $A \to B$, then for any commutative ring ppc spectrum R, and map $A \to R$, the cell attachment $R \wedge_A B$ satisfies

$$\Phi_p(R \wedge_A B) \cong \Phi_p R \wedge_{\Phi_p A} \Phi_p B$$

and we can give it the ppc structure map induced by the ppc structure maps on R, A, and B (which by hypothesis are maps of commutative ring T-spectra). Thus, we can form the required cell attachments in commutative ring ppc spectra to reproduce the usual argument constructing a model structure on the commutative ring objects. This gives the model structure for part (ii) of Theorem D. The case for commutative ring pc spectra is entirely similar.

THEOREM 2.3. The category of commutative ring ppc spectra (resp., pc spectra) admits a model structure with fibrations and weak equivalences created in the model structure of ppc spectra (resp., pc spectra) of Theorem 2.1.

The cells that we used to construct cofibrant commutative ring $p\mathbf{p}\mathbf{c}$ spectra are of the form

$$\mathbb{PC}(A \wedge \partial D^n_+) \longrightarrow \mathbb{PC}(A \wedge D^n_+)$$

for $A \in \mathcal{A}_p^{\text{cyc}}$. On underlying commutative ring T-spectra, this is a cofibration of cofibrant objects in the $\mathcal{A}_p^{\text{cyc}}$ -variant $\mathcal{V}_{\pi^{\dagger}}$ -constrained \mathcal{F}_p -model structure (by Proposition 2.2). We see that the forgetful functor from commutative ring ppc spectra to commutative ring T-spectra preserves cofibrations. Analogous observations hold in the case of pc spectra.

PROPOSITION 2.4. The forgetful functor from commutative ring ppc spectra (resp., pc spectra) to commutative ring \mathbb{T} -spectra preserves cofibrations and acyclic cofibrations for the model structure in Theorem 2.3 on the source and the $\mathcal{A}_p^{\text{cyc}}$ -variant $\mathcal{V}_{\pi^{\text{l}}}$ -constrained \mathcal{F}_p -model structure (resp., \mathcal{A}^{cyc} -variant $\mathcal{V}_{\pi^{\text{l}}}$ -constrained \mathcal{F}_{fin} -model structure) on the target.

To finish the proof of Theorem D, we need to see that for a cofibrant commutative ring ppc spectrum, the underlying ppc spectrum is cofibrant in the undercategory of S. This easily reduces to showing that for $A_1, \ldots, A_r \in \mathcal{A}_p^{cyc}$,

$$\mathbb{C}A_1 \wedge \cdots \wedge \mathbb{C}A_r$$

is a wedge of ppc spectra of the form $\mathbb{C}A$ with $A \in \mathcal{A}_p^{cyc}$. This is then an easy combinatorial argument using the fact that \mathcal{A}_p^{cyc} is closed (up to isomorphism) under Φ_{p^k} .

For Addendum D, recall that for R an associative ring ppc (resp., pc) spectrum an R-module ppc (resp., pc) spectrum consists of an R-module M (in \mathbb{T} -spectra) together with a ppc (resp., pc) structure on M such that the diagram

$$\begin{split} \Phi_p R \wedge \Phi_p M & \longrightarrow \Phi_p (R \wedge M) & \longrightarrow \Phi_p M \\ & \downarrow & \qquad \qquad \downarrow \\ R \wedge M & \longrightarrow M \end{split}$$

constructed from the structure maps and the action map commutes (for all primes p in the pc case). When R is commutative, we also have notions of associative R-algebras and commutative R-algebras in ppc spectra (resp., pc spectra): an associative R-algebra consists of a map of associative ring ppc spectra (resp., pc spectra) $\eta: R \to A$ with the property that η is also a map of associative ring ppc spectrum (resp., pc spectra (resp., pc spectra)) is just a map of commutative ring ppc spectra (resp., pc spectra) $R \to A^{\text{op}}$. A commutative ring ppc spectra (resp., pc spectra) $R \to A$. The statements in the Addendum and the corresponding results for pc spectra are clear from the work above. We also have the following generalization to the case when R is just in \mathcal{E}_{Φ} .

THEOREM 2.5. Let R be a commutative ring ppc (resp., pc) spectrum whose underlying \mathbb{T} -spectrum is in \mathcal{E}_{Φ} (e.g., if it cofibrant in the model structure of Theorem 2.3).

- (i) The category of R-module ppc (resp., pc) spectra admits a topological model structure with fibrations and weak equivalences created by the forgetful functor to the model structure on ppc (resp., pc) spectra of Theorem 2.1. This model category is enriched over the standard model category of non-equivariant spectra.
- (ii) The category of R-algebra ppc (resp., pc) spectra admits a topological model structure with fibrations and weak equivalences created by the forgetful functor to the model structure on ppc (resp., pc) spectra of Theorem 2.1.
- (iii) The category of commutative R-algebra ppc (resp., pc) spectra admits a topological model structure with fibrations and weak equivalences created by the forgetful functor to the model structure on ppc (resp., pc) spectra of Theorem 2.1.

Moreover, in each of these model structures, the underlying \mathbb{T} -spectra of the cofibrant objects in this model category lie in \mathcal{E}_{Φ} .

When R is a cofibrant commutative ring ppc spectrum, or more generally any commutative ring ppc spectrum whose underlying T-spectrum is in \mathcal{E}_{Φ} , the smash product \wedge_R can be partially defined just like the smash product \wedge on ppc spectra. When M is a cofibrant R-module in ppc spectra and X is any R-module in ppcspectra, we get a canonical ppc structure on the smash product $M \wedge_R X$ of their underlying T-spectra with the structure map

$$\Phi_p(M \wedge_R X) \cong \Phi_p M \wedge_{\Phi_n R} \Phi_p X \longrightarrow M \wedge_R X$$

induced by the structure maps on M, R, and X, using the isomorphism of Proposition 1.8. This in particular constructs a symmetric monoidal smash product on the full subcategory of cofibrant R-modules in ppc spectra.

Before turning to Theorem B, we make a brief remark about monadicity. While equivariant commutative ring spectra may be defined in terms of the monad \mathbb{P} on equivariant orthogonal spectra, there is no corresponding definition for commutative ring *p*pc spectra, because the smash product does not extend to the whole category of *p*pc spectra. However, by the work above, if we restrict to the subcategory of *p*pc spectra under S that are cofibrant, we do have a smash product, and on this category we can define the monad \tilde{P} by

$$\tilde{P}X = \prod_{n=1}^{\infty} X^{(n)} / \Sigma_n$$

(where the coproduct is taken in the undercategory of S, i.e., it is wedge with all copies of S identified). Analogous observations hold for pc spectra. Formally, we get the following observation.

PROPOSITION 2.6. The category of algebras over the monad P in the category of cofibrant objects in the undercategory of S in ppc spectra (respectively, pc spectra) with the model structure of Theorem 2.1 is the full subcategory of the category of commutative ring ppc spectra (respectively, commutative ring pc spectra) whose underlying ppc spectra (respectively, pc spectra) under S are cofibrant.

For Theorem B, the main ingredient beyond Proposition 2.4 we need is that the model structure above is topological, which implies that maps into a fibrant object convert cofibrations into fibrations. The proof of the corresponding result [4, 5.13] for mapping spectra of ppc spectra in the standard model structure works in this context to prove the mapping space result for commutative ring ppc spectra in this model structure. The following is a restatement of Theorem B naming the specific model structure.

THEOREM 2.7. If A and B are commutative ring ppc spectra where the underlying commutative ring T-spectra of A and B are cofibrant and fibrant, respectively, in the $\mathcal{A}_p^{\text{cyc}}$ -variant $\mathcal{V}_{\pi!}$ -constrained \mathcal{F}_p -model structure, then the derived space of maps of commutative ring ppc spectra from A to B, $\mathbb{RC}om^{p \operatorname{cyc}}(A, B)$, is given by the homotopy equalizer

 $\mathbf{RCom}^{p\operatorname{cyc}}(A,B) \simeq \operatorname{hoEq}\left(\operatorname{Com}^{\mathbb{T}}(A,B) \rightrightarrows \operatorname{Com}^{\mathbb{T}}(\Phi A,B)\right)$

where $Com^{\mathbb{T}}$ denotes the point-set mapping space in commutative ring \mathbb{T} -spectra. Here one map is induced by the cyclotomic structure map $\Phi A \to A$ and the other map is induced by applying Φ and post-composing with the cyclotomic structure map $\Phi B \rightarrow B$.

The proof works just like the proof of [4, 5.13]: A check of definitions shows that the point-set space of maps of ppc spectra from A to B, $Com^{p \operatorname{cyc}}(A, B)$ is the equalizer

$$\operatorname{Eq}\left(\operatorname{\mathcal{C}om}^{\mathbb{T}}(A,B)\rightrightarrows\operatorname{\mathcal{C}om}^{\mathbb{T}}(\Phi A,B)\right)$$

and the point-set mapping space represents the derived mapping space when A is cofibrant and B is fibrant in the model structure on commutative ring ppc spectra of Theorem 2.3. The proof of [4, 5.12] generalizes to show that the map from the equalizer to the homotopy equalizer is a weak equivalence under these hypotheses. If A and B are just cofibrant and fibrant, respectively, as commutative ring T-spectra, their cofibrant and fibrant approximation $A_c \to A$ and $B \to B_f$, respectively, in commutative ring ppc spectra calculate the derived mapping space, and the map from the homotopy equalizer for A, B to the homotopy equalizer for A_c, B_f is a weak equivalence.

In the case when R is cofibrant as a commutative ring ppc spectrum (or has underlying \mathbb{T} -spectrum in \mathcal{E}_{Φ}), we also get the corresponding result in the categories of R-modules and R-algebras. For convenience we write $\mathcal{A}lg_R^{\mathbb{T}}$ for the category of associative R-algebras in \mathbb{T} -spectra and $\mathcal{A}lg_R^{pcyc}$ for the category of associative R-algebras in ppc spectra. We also write $\mathcal{A}lg_R^{\mathbb{T}}(-,-)$ and $\mathcal{A}lg^{pcyc}(-,-)$ for the corresponding (point-set) mapping spaces and $\mathbf{R}\mathcal{A}lg_R^{\mathbb{T}}(-,-)$ and $\mathbf{R}\mathcal{A}lg_R^{pcyc}(-,-)$ for the derived mapping spaces. Checking definitions, we have that for R-algebras A and B, $\mathcal{A}lg_R^{pcyc}(A, B)$ is the (point-set) equalizer

$$\operatorname{Eq}\left(\mathcal{A}lg_{R}^{\mathbb{T}}(A,B) \rightrightarrows \mathcal{A}lg_{\Phi_{n}R}^{\mathbb{T}}(\Phi_{p}A,B)\right)$$

where on the right B becomes a $\Phi_p R$ -algebra via the structure map $\Phi_p R \to R$. In the equalizer one map is induced by the reduction of scalars map $\mathcal{A}lg_R^{\mathbb{T}} \to \mathcal{A}lg_{\Phi_p R}^{\mathbb{T}}$ and the structure map $\Phi_p A \to A$ and the other by applying the continuous functor $\Phi_p: \mathcal{A}lg_R^{\mathbb{T}} \to \mathcal{A}lg_{\Phi_p R}^{\mathbb{T}}$ and using the structure map $\Phi_p B \to B$. From here, the following result follows from the same argument outline as [4, 5.13].

THEOREM 2.8. Let R be a commutative ring ppc spectrum, whose underlying \mathbb{T} -spectrum is in \mathcal{E}_{Φ} (which holds for example if R is cofibrant). Let A and B be associative R-algebras in ppc spectra. Assume that on underlying \mathbb{T} -equivariant associative R-algebras, A is cofibrant as a \mathbb{T} -equivariant associative R-algebra and B is fibrant as a \mathbb{T} -equivariant associative R-algebra in the \mathcal{A}_p^{cyc} -variant $\mathcal{V}_{\pi!}$ -constrained \mathcal{F}_p -model structure. Then the derived mapping space $\mathbb{R}\mathcal{A}lg_R^{pcyc}(A, B)$ is given by the homotopy equalizer

$$\mathbf{R}\mathcal{A}lg_{R}^{p\,\mathrm{cyc}}(A,B) \simeq \operatorname{hoEq}\left(\mathcal{A}lg_{R}^{\mathbb{T}}(A,B) \rightrightarrows \mathcal{A}lg_{\Phi_{n}R}^{\mathbb{T}}(\Phi_{p}A,B)\right).$$

For R any associative ring ppc spectrum and X, Y any R-modules in ppc spectra (without hypotheses), we have the mapping spectrum $F_R^{p \operatorname{cyc}}(X, Y)$, which is defined to be the orthogonal spectrum obtained as the equalizer

$$\operatorname{Eq}\left(F_{R}^{\mathbb{T}}(X,Y) \rightrightarrows F_{\Phi_{n}R}^{\mathbb{T}}(\Phi_{p}X,Y)\right)$$

where $F_R^{\mathbb{T}}$ denotes the (non-equivariant) spectrum of *R*-module maps of the underlying \mathbb{T} -equivariant *R*-modules. The maps are analogous to those above: one map

is induced by the reduction of scalars from R-modules to $\Phi_p R$ -modules followed by the structure map $\Phi_p X \to X$ for X and the other is induced by the spectrally enriched functor Φ_p from R-modules to $\Phi_p R$ -modules and the structure map $\Phi_p Y \to Y$ for Y. The space of R-module ppc spectrum maps from X to Y is the zeroth space of the orthogonal spectrum $F_R^{p\,cyc}$ above (because $F_R^{\mathbb{T}}$ has this property for the category of R-modules and zeroth space commutes with equalizers). This construction satisfies the spectral version of Quillen's Axiom SM7. (This is precisely the assertion of Theorem 2.5.(i) of enrichment over non-equivariant spectra.)

This implies that the derived functor $\mathbf{R}F_R^{p\,\text{cyc}}(X,Y)$ exists and is calculated by cofibrant approximation of X and fibrant approximation of Y. Again the argument of [4, 5.13] generalizes to prove the following result.

THEOREM 2.9. Let R be an associative ring ppc spectrum, whose underlying \mathbb{T} -spectrum is in \mathcal{E}_{Φ} (which holds for example if R is cofibrant or cofibrant commutative). Let X and Y be R-modules in ppc spectra. Assume that on underlying \mathbb{T} -equivariant R-modules, A is cofibrant as a \mathbb{T} -equivariant R-module and B is fibrant as a \mathbb{T} -equivariant R-module in either the standard \mathcal{F}_p -model structure or the model structure from Theorem 2.5. Then the derived mapping space $\mathbb{R}\mathcal{M}od_R^{p\,cyc}(A, B)$ is given by the homotopy equalizer

$$\mathbf{R}\mathcal{M}od_{R}^{p\operatorname{cyc}}(A,B) \simeq \left(\mathcal{M}od_{R}^{\mathbb{T}}(A,B) \rightrightarrows \mathcal{M}od_{\Phi_{p}R}^{\mathbb{T}}(\Phi_{p}A,B)\right)$$

and the derived mapping spectrum $\mathbf{R}F_R^{p\operatorname{cyc}}(X,Y)$ is given by the homotopy equalizer of

$$\mathbf{R}F_R^{p\operatorname{cyc}}(X,Y) \simeq \operatorname{hoEq}\left(F_R^{\mathbb{T}}(A,B) \rightrightarrows F_{\Phi_p R}^{\mathbb{T}}(\Phi_p A,B)\right).$$

The pc case is more complicated than the ppc case in precisely the same way as in the unstructured case of [4, §5]. For commutative ring pc spectra, the proof of [4, 5.18] works in the current context to prove the following theorem, the pc analogue of Theorem B. See [4, 5.15] for notation.

THEOREM 2.10. If A and B are commutative ring pc spectra where the underlying commutative ring \mathbb{T} -spectra of A and B are cofibrant and fibrant, respectively, in the \mathcal{A}^{cyc} -variant $\mathcal{V}_{\pi!}$ -constrained \mathcal{F}_{fin} -model structure, then the derived space of maps of commutative ring pc spectra from A to B, $\mathbb{RC}om^{cyc}(A, B)$ is given by the homotopy limit

$$\mathbf{RCom}^{\mathrm{cyc}}(A, B) \simeq \operatorname{holim}_{\Theta} \mathcal{Com}^{\mathbb{T}}(\Phi_n A, B)$$

with maps as in [4, (5.14)].

The argument works the same for R-algebras and R-modules to give the following results.

THEOREM 2.11. Let R be a commutative ring pc spectrum, whose underlying \mathbb{T} -spectrum is in \mathcal{E}_{Φ} (which holds for example if R is cofibrant). Let A and B be associative R-algebras in pc spectra. Assume that on underlying \mathbb{T} -equivariant associative R-algebras, A is cofibrant as a \mathbb{T} -equivariant associative R-algebra and B is fibrant as a \mathbb{T} -equivariant associative R-algebra in the A^{cyc}-variant $\mathcal{V}_{\pi!}$ -constrained \mathcal{F}_{fin} -model structure. Then the derived mapping space $\mathbf{R}\mathcal{A}lg_R^{cyc}(A, B)$ is given by the homotopy limit

$$\mathbf{R}\mathcal{A}lg_{R}^{\mathrm{cyc}}(A,B) \simeq \operatorname{holim}_{\Theta}\mathcal{A}lg_{\Phi_{n}R}^{\mathbb{T}}(\Phi_{n}A,B)$$

with maps as in [4, (5.14)].

THEOREM 2.12. Let R be an associative ring pc spectrum, whose underlying \mathbb{T} -spectrum is in \mathcal{E}_{Φ} (which holds for example if R is cofibrant or cofibrant commutative). Let X and Y be R-modules in pc spectra. Assume that on underlying \mathbb{T} -equivariant R-modules, A is cofibrant and B is fibrant in either the standard \mathcal{F}_{fin} -model structure or the model structure of Theorem 2.5. Then the derived mapping space $\mathbb{R}\mathcal{M}od_{R}^{cyc}(A, B)$ is given by the homotopy limit

$$\mathbf{R}\mathcal{M}od_{R}^{\mathrm{cyc}}(A,B) \simeq \operatorname{holim}_{\Theta}\mathcal{M}od_{\Phi_{n}R}^{\mathbb{T}}(\Phi_{n}X,Y)$$

and the derived mapping spectrum $\mathbf{R}F_{R}^{\mathrm{cyc}}(X,Y)$ is given by the homotopy limit

$$\mathbf{R}F_R^{\mathrm{cyc}}(X,Y) \simeq \operatorname{holim}_{\Theta} F_{\Phi_n R}^{\mathbb{T}}(\Phi_n X,Y)$$

with maps as in [4, (5.14)].

3. Construction of $\mathcal{A}^{\mathrm{cyc}}$ and proof of Lemma 1.4

We construct \mathcal{A}^{cyc} and $\mathcal{A}_p^{\text{cyc}}$ by specifying explicit generalized orbit desuspension spectra that we can show have the symmetric power property required for the hypotheses of Theorem II.1.13 while at the same time form sets that are closed (up to isomorphism) under Φ_n (or Φ_{p^n} in the case of $\mathcal{A}_p^{\text{cyc}}$), smash products, and symmetric powers. This requires leaving the context of the Lie group T and working with finite extensions. The most convenient way to set this up is as follows. For $n \geq 1$, let

$$\mathbb{T}(n) := \mathbb{R}/n\mathbb{Z}, \qquad C_m(n) = \langle \frac{n}{m} \rangle < \mathbb{T}(n), \qquad \Lambda(n) = \langle 1 \rangle = C_n(n) < \mathbb{T}(n).$$

 $(C_m(n) < \mathbb{T}(n)$ is the subgroup with m elements.) We regard $\mathbb{T}(1)$ as canonically isomorphic to \mathbb{T} (via the map $t \mapsto e^{2\pi i t}$) and use this isomorphism to convert $\mathbb{T}(1)$ -spectra to \mathbb{T} -spectra without comment or notation. We also use the canonical isomorphism $\mathbb{T}(1) \cong \mathbb{T}(n)/\Lambda(n)$ without additional notation. Combining these conventions, we write \mathbb{T} -compatible and $J^{\mathbb{T}}$ for $\mathbb{T}(n)/\Lambda(n)$ -compatible and $J^{\mathbb{T}(n)/\Lambda(n)}$ in the context of Terminology V.1.7 and Notation V.1.8.

The following gives the first basic building block for the definition of the objects in \mathcal{A}^{cyc} and $\mathcal{A}_{p}^{\text{cyc}}$.

NOTATION 3.1. Let $n \ge 1$, $m \ge 1$, and let V be an orthogonal $C_m(n)$ representation. Let $\eta[n, m, V]$ be the $(\mathbb{T}(n), 1)$ vector bundle

$$\mathbb{T}(n) \times_{C_m(n)} V \longrightarrow \mathbb{T}(n) / C_m(n).$$

The definition of the objects in \mathcal{A}^{cyc} and $\mathcal{A}^{\text{cyc}}_{p}$ also needs the ideas from Definition V.2.13. Specifically, we need the following variant of the constructions there.

DEFINITION 3.2. For the purposes of this section, data consists of

- Integers $n \ge 1$, $r \ge 1$, $q_0 \ge 0$, $q_1, \ldots, q_r \ge 1$, $m_1, \ldots, m_r \ge 1$, $\ell_1, \ldots, \ell_r \ge 1$ with ℓ_i dividing both n and m_i .
- For i = 1, ..., r, a $C_{\ell_i}(n)$ -trivial orthogonal $C_{m_i}(n)$ -representation V_i .
- A subgroup $Q < \Sigma_{q_0} \times \cdots \times \Sigma_{q_r}$.

We abbreviate this data as

$$Q = (n, r, (q_0, \dots, q_r), (m_1, \dots, m_r), (\ell_1, \dots, \ell_r), (V_1, \dots, V_r), Q),$$

and write $\Sigma = \Sigma_{q_0} \times \cdots \times \Sigma_{q_r}$. Given data \mathcal{Q} write $n^{\mathcal{Q}}, r^{\mathcal{Q}}, q_i^{\mathcal{Q}}, m_i^{\mathcal{Q}}, \ell_i^{\mathcal{Q}}, V_i^{\mathcal{Q}}$, and $Q^{\mathcal{Q}}$ for its constituent elements. Let $\eta_{\mathcal{Q}}$ be the $(\mathbb{T}(n), Q)$ vector bundle obtained from

$$\operatorname{Sym}^{q_0}(\mathbb{R} \longrightarrow *) \times \operatorname{Sym}^{q_1} \eta[n, m_1, V_1] \times \cdots \times \operatorname{Sym}^{q_r} \eta[n, m_r, V_r]$$

by reducing the group for quotients from Σ to Q. Let $\operatorname{Hom}(Q)$ denote the set of homomorphisms $\sigma: \Lambda(n) \to Q$ such that for each $i = 1, \ldots, r$, the composite

$$\Lambda(n) \longrightarrow Q \longrightarrow \Sigma_{q_1} \times \cdots \times \Sigma_{q_r} \longrightarrow \Sigma_{q_i}$$

has kernel containing $C_{\ell_i}(n)$; we give $\operatorname{Hom}(\mathcal{Q})$ the right Q action by conjugation (and the trivial $\mathbb{T}(n)$ action). Writing $(-)^{\Lambda(n),\sigma}$ for σ -twisted $\Lambda(n)$ fixed points (as in Definition V.2.10), let θ_Q be the $(\mathbb{T}(n), Q)$ vector bundle

$$\prod_{\sigma \in \operatorname{Hom}(\mathcal{Q})} \eta_{\mathcal{Q}}^{\Lambda(n),}$$

with Q action coming from the Q action on both $\operatorname{Hom}(Q)$ and η_Q . Finally, when X is a subset of the base of θ_Q that is a $(\mathbb{T}(n) \times Q^{\operatorname{op}})$ -equivariant subcomplex for a $(\mathbb{T}(n) \times Q^{\operatorname{op}})$ -equivariant cell complex structure, then we let $\theta_{Q,X}$ be the $(\mathbb{T}(n), Q)$ vector bundle obtained by base change to X.

In other words, $\theta_{\mathcal{Q}}$ is the subset of

$$\eta_{\mathcal{Q}}(\Lambda(n)) = \prod_{\sigma \, : \, \Lambda(n) \to Q} \eta_{\mathcal{Q}}^{\Lambda(n),\sigma}$$

consisting of the summands indexed by $\sigma \in \text{Hom}(\mathcal{Q})$. In particular $\theta_{\mathcal{Q}}$ is T-compatible as is $\theta_{\mathcal{Q},X}$ for any X.

DEFINITION 3.3. Let $D(\mathcal{A}^{cyc})$ be the set of ordered pairs (\mathcal{Q}, X) where \mathcal{Q} is data as above, X is an equivariant subcomplex as above, Q satisfies the property

For any non-identity s ∈ Q, there exists j > 0 such that s is not in the kernel of the map Q → Σ → Σ_{q_j},

and for each i, the $C_{m_i^{\mathcal{Q}}}(n^{\mathcal{Q}})/C_{\ell_i^{\mathcal{Q}}}(n^{\mathcal{Q}})$ -representation $V_i^{\mathcal{Q}}$ satisfies the following properties:

- $V_i^{\mathcal{Q}}$ is a semiregular permutation representation (a permutation representation containing at least one copy of the regular representation), and
- $V_i^{\mathcal{Q}}$ has underlying non-equivariant inner product space given by \mathbb{R}^k for some k.

Let $D(\mathcal{A}_p^{\text{cyc}})$ be the subset of $D(\mathcal{A}^{\text{cyc}})$ consisting of those (\mathcal{Q}, X) where $n^{\mathcal{Q}}$ and $m_i^{\mathcal{Q}}$ are powers of p. Let \mathcal{A}^{cyc} be the set of \mathbb{T} -spectra $J^{\mathbb{T}}(\theta_{\mathcal{Q},X})$ for $(\mathcal{Q}, X) \in D(\mathcal{A}^{\text{cyc}})$ and let $\mathcal{A}_p^{\text{cyc}} \subset \mathcal{A}^{\text{cyc}}$ be the set of \mathbb{T} -spectra $J^{\mathbb{T}}(\theta_{\mathcal{Q},X})$ for $(\mathcal{Q}, X) \in D(\mathcal{A}_p^{\text{cyc}})$.

We note that $\mathcal{A}^{\text{cyc}} \supset \mathcal{A}^{\circ}(\mathcal{F}_{\text{fin}}, \mathcal{V}_{\pi !})$ and $\mathcal{A}_{p}^{\text{cyc}} \supset \mathcal{A}^{\circ}(\mathcal{F}_{p}, \mathcal{V}_{\pi !})$ with the elements of $\mathcal{A}^{\circ}(\mathcal{F}_{\text{fin}}, \mathcal{V}_{\pi !})$ and $\mathcal{A}^{\circ}(\mathcal{F}_{p}, \mathcal{V}_{\pi !})$ given by the $J^{\mathbb{T}}(\theta_{\mathcal{Q}})$ where $n^{\mathcal{Q}} = 1$, $r^{\mathcal{Q}} = 1$, $q_{0}^{\mathcal{Q}} = 0$, and $q_{1}^{\mathcal{Q}} = 1$. In general, for $A \in \mathcal{A}^{\text{cyc}}$, the $(\mathcal{Q}, X) \in D(\mathcal{A}^{\text{cyc}})$ for which $A = J^{\mathbb{T}}(\theta_{\mathcal{Q},X})$ may not be unique, but choosing one, we write $\mathcal{Q}_{A} = \mathcal{Q}, X_{A} = X$, $\eta_{A} = \eta_{\mathcal{Q}_{A}}, \ \theta_{A} = \theta_{\mathcal{Q}_{A}}, \ \theta_{A,X} = \theta_{\mathcal{Q},X}$, and in the notation for the data in \mathcal{Q}_{A} , abbreviate the superscript \mathcal{Q}_{A} to just A. (In the case when $A \in \mathcal{A}_{p}^{\text{cyc}}$, we choose $(\mathcal{Q}, X) \in D(\mathcal{A}_{p}^{\text{cyc}})$.)

To explain the idea behind the definition, we introduce the following construction and the following observation about it. CONSTRUCTION 3.4. Let \mathcal{Q} be data as in Definition 3.2 and let $k \geq 1$. define $k\mathcal{Q}$ by

$$n^{k\mathcal{Q}} := kn^{\mathcal{Q}}, r^{k\mathcal{Q}} := r^{\mathcal{Q}}, m_i^{k\mathcal{Q}} := km_i^{\mathcal{Q}}, \ell_i^{k\mathcal{Q}} := k\ell_i^{\mathcal{Q}}, q_i^{k\mathcal{Q}} := q_i^{\mathcal{Q}}, Q^{k\mathcal{Q}} := Q^{\mathcal{Q}},$$

and take $V_i^{k\mathcal{Q}}$ to be $V_i^{\mathcal{Q}}$ converted to a $C_{k\ell_i^{\mathcal{Q}}}(kn^{\mathcal{Q}})$ -acyclic orthogonal $C_{km_i^{\mathcal{Q}}}(kn^{\mathcal{Q}})$ -representation using the canonical isomorphism

$$C_{km_i^{\mathcal{Q}}}(kn^{\mathcal{Q}})/C_{k\ell_i^{\mathcal{Q}}}(kn^{\mathcal{Q}}) \cong C_{m_i^{\mathcal{Q}}}(n^{\mathcal{Q}})/C_{\ell_i^{\mathcal{Q}}}(n^{\mathcal{Q}}).$$

Let $\operatorname{Hom}_k(k\mathcal{Q}) \subset \operatorname{Hom}(k\mathcal{Q})$ consist of those homomorphisms $\sigma \colon \Lambda(n^{k\mathcal{Q}}) \to Q^{k\mathcal{Q}}$ where the composite

$$\Lambda(n^{k\mathcal{Q}}) \longrightarrow Q^{k\mathcal{Q}} \longrightarrow \Sigma_{q_0^{k\mathcal{Q}}} \times \dots \times \Sigma_{q_{r^{k\mathcal{Q}}}^{k\mathcal{Q}}} \longrightarrow \Sigma_{q_0^{k\mathcal{Q}}}$$

has kernel containing $C_k(n^{kQ})$. Let $B_k(kQ)$ denote the subset of the base of θ_{kQ} consisting of those components indexed by homomorphisms in $\operatorname{Hom}_k(kQ)$.

PROPOSITION 3.5. Let Q, X be as in Definition 3.2, let $k \ge 1$, and write $Q = Q^{Q} = Q^{kQ}$. Then:

(i) The action of $C_k(n^{kQ})$ on $B_k(kQ)$ is trivial and under the canonical isomorphism

$$\mathbb{T}(n^{k\mathcal{Q}})/C_k(n^{k\mathcal{Q}}) \times (Q^{k\mathcal{Q}})^{\mathrm{op}} \cong \mathbb{T}(n^{\mathcal{Q}}) \times (Q^{\mathcal{Q}})^{\mathrm{op}},$$

 $B_k(k\mathcal{Q})$ is equivariantly isomorphic to the base of $\theta_{\mathcal{Q}}$.

- (ii) There is an isomorphism of \mathbb{T} -spectra $J^{\mathbb{T}}\theta_{k\mathcal{Q},kX} \cong J^{\mathbb{T}}\theta_{\mathcal{Q},X}$, where kX is subcomplex of $B_k(k\mathcal{Q})$ corresponding to X under the isomorphism of (ii).
- (iii) When Q is in D(A^{cyc}) then so is kQ; when k is a power of p, if Q is in D(A^{cyc}_p), then so is kQ.

PROOF. Since

$$C_k(n^{k\mathcal{Q}}) < C_{\ell_i^{k\mathcal{Q}}}(n^{k\mathcal{Q}}) < C_{m_i^{k\mathcal{Q}}}(n^{k\mathcal{Q}}), \qquad C_k(n^{k\mathcal{Q}}) < \Lambda(n^{k\mathcal{Q}})$$

(and $C_k(n^{kQ})$ is abelian), we have that $C_k(n^{kQ})$ acts trivially on the vector bundle θ_{kQ} before restricting the base to $B_k(kQ)$. Since $C_k(n^{kQ}) < C_{\ell_k^{kQ}}(n^{kQ})$ for all i, any $\sigma \in \operatorname{Hom}_k(kQ)$ has $C_k(n^{kQ})$ in the kernel. The canonical isomorphism

$$\Lambda(n^{\mathcal{Q}}) \cong \Lambda(kn^{\mathcal{Q}})/C_k(kn^{\mathcal{Q}})$$

therefore induces a bijection between $\operatorname{Hom}(\mathcal{Q})$ and $\operatorname{Hom}_k(k\mathcal{Q})$. This proves statement (i). For the proof of statement (ii), we use the notation of Construction V.1.2 for the specifics of the \mathbb{T} -spectra J. Let W be an orthogonal \mathbb{T} -representation. The observations above about the relations of the subgroups then give isomorphisms of $(\mathbb{T}(n^{\mathcal{Q}}) \times Q^{\operatorname{op}})$ -equivariant fiber bundles

$$\mathcal{I}(\theta_{k\mathcal{Q},B_k(k\mathcal{Q})},W) \cong \mathcal{I}(\theta_{\mathcal{Q}},W)$$

and $(\mathbb{T}(n^{\mathcal{Q}}) \times Q^{\mathrm{op}})$ -equivariant vector bundles

$$\mathfrak{S}^{\perp}\mathcal{I}(\theta_{k\mathcal{Q},B_{k}}(k\mathcal{Q}),W) \cong \mathfrak{S}^{\perp}\mathcal{I}(\theta_{\mathcal{Q}},W).$$

Since J is constructed as the spectrum formed by the Thom spaces of these vector bundles, this proves (ii). For (iii), we note that if $V_i^{\mathcal{Q}}$ is a semiregular permutation representation of $C_{m_i^{\mathcal{Q}}}(n^{\mathcal{Q}})/C_{\ell_i^{\mathcal{Q}}}(n^{\mathcal{Q}})$, then $V_i^{k\mathcal{Q}}$ is a semiregular permutation representation of $C_{m_i^{k\mathcal{Q}}}(n^{k\mathcal{Q}})/C_{\ell_i^{k\mathcal{Q}}}(n^{k\mathcal{Q}})$.

42 III. THE CIRCLE GROUP AND CYCLOTOMIC SPECTRA (THEOREMS B, C, D)

NOTATION 3.6. For \mathcal{Q}, X as in Definition 3.2 and $k \ge 1$, let $k(\mathcal{Q}, X) = (k\mathcal{Q}, kX)$ where kX is as in (iii) above.

The idea behind the definition of \mathcal{A}^{cyc} is as follows. Looking at data \mathcal{Q} (and dropping the \mathcal{Q} superscripts from the notation), when $q_0 = 0$ and $Q = Q_1 \times \cdots \times Q_r$ for $Q_i < \Sigma_{q_i}$, the previous proposition combined with Proposition V.1.4 and Theorem V.2.4 implies that under faithfulness hypotheses (that we will show to hold when $\mathcal{Q} \in D(\mathcal{A}^{\text{cyc}})$)

$$J^{\mathbb{T}}\theta_{\mathcal{Q}} \cong \Phi_{n/\ell_1}((J\eta[\frac{n}{\ell_1},\frac{m_1}{\ell_1},V_1'])^{(q_1)}/Q_1) \wedge \dots \wedge \Phi_{n/\ell_r}((J\eta[\frac{n}{\ell_r},\frac{m_r}{\ell_r},V_r'])^{(q_r)}/Q_r)$$

where V' denotes V converted to the appropriate type of representation by the canonical isomorphism of quotient groups (the reverse of the conversion in Construction 3.4). The purpose of the ℓ_i is thus to put together a smash product of geometric fixed points for differing subgroups. At the cost of passing to a retract (which we handle by using a subspace X of the base), the functors Φ_n pull outside a symmetric power, and allowing Q to be a more general subgroup of Σ than a product treats the case of a smash product inside of a symmetric power. (The restriction on the group Q when $q_0 > 0$ is purely technical and arises from the restriction imposed by the requirement that the Q action be faithful; see the proof of part (i) of Lemma 1.4 below.)

It remains to prove Lemma 1.4. We do this one part at a time.

PROOF OF LEMMA 1.4.(I). Most of the statement follows by construction: we just need to check that θ_{Q_A,X_A} is Q^A -faithful and also that for every $k \geq 1$, $\theta_{Q_A,X_A}(C_{kn}(n)|\Lambda(n))$ is Q^A -faithful. For this, it is enough to show that θ_A and $\theta_A(C_{kn}(n)|\Lambda(n))$ are Q^A -faithful. Let $\mathcal{Q} = \mathcal{Q}_A$. Since only one object is involved, we drop the superscripts \mathcal{Q} or A on the data of \mathcal{Q} and the subscripts \mathcal{Q} or A on the bundles. Fix k, and let \mathcal{H}_k be the set of homomorphisms $\sigma: C_{kn}(n) \to Q$ such that the composite to Σ_{q_i} contains $C_{\ell_i}(n)$ for all $i = 1, \ldots, r$, and let

$$\theta\langle k\rangle = \prod_{\sigma\in\mathcal{H}_k} \eta^{C_{kn}(n),\sigma}$$

Then when k = 1, $\mathcal{H}_1 = \text{Hom}(\mathcal{Q})$ and $\theta \langle 1 \rangle = \theta$, and so it suffices to show that $\theta \langle k \rangle$ is *Q*-faithful: the case k = 1 shows that θ is *Q*-faithful, and then the proof of Theorem V.2.5 in Section VI.1 shows that $\theta \langle k \rangle = \theta(C_{kn}(n)|\Lambda(n))$.

We use the notation $E(\eta)$ and $B(\eta)$ for the base and total spaces of η and Eand B for the base and total spaces of $\theta\langle k \rangle$: $\eta \colon E(\eta) \to B(\eta)$ and $\theta\langle k \rangle \colon E \to B$. Let

$$s \in Q < \Sigma_{q_1} \times \cdots \times \Sigma_{q_r}$$

not the identity, and let $b \in B$ with bs = b; we need to show there exists an element $e \in E_b$ with $es \neq e$. Let $\sigma \in \mathcal{H}_k$ be the homomorphism indexing the summand containing b. Let b' be the corresponding element of $B(\eta)^{C_{kn}(n),\sigma}$. Since s is not the identity, there exists an i > 0 so that the projection s_i of s to Σ_{q_i} is not the identity; let $\sigma_i \colon C_{kn}(n) \to \Sigma_{q_i}$ be the composite of σ with the projection. By definition of \mathcal{H}_k , σ_i has kernel containing $C_{\ell_i}(n)$.

While we have defined $\eta[n, m_i, V_i]$ as a $(\mathbb{T}(n), 1)$ vector bundle, because V_i is $C_{\ell_i}(n)$ -trivial, the group of equivariance restricts to $\mathbb{T}(n)/C_{\ell_i}(n) \cong \mathbb{T}(n/\ell_i)$. Viewing $\eta[n, m_i, V_i]$ as a $(\mathbb{T}(n/\ell_i), 1)$ vector bundle, it is canonically isomorphic to

$$\mathbb{T}(n/\ell_i) \times_{C_{m_i}(n)/C_{\ell_i}(n)} V_i \longrightarrow \mathbb{T}(n/\ell_i)/C_{m_i/\ell_i}(n/\ell_i),$$

and the $C_{m_i/\ell_i}(n/\ell_i) \cong C_{m_i}(n)/C_{\ell_i}(n)$ -representation V_i contains a copy of the regular representation of $C_{m/\ell_i}(n/\ell_i)$. Then

$$\xi := (\operatorname{Sym}^{q_i} \eta[n, m_i, V_i])(C_{kn/\ell_i}(n/\ell_i)) = \prod_{\tau : \ C_{kn/\ell_i}(n/\ell_i) \to \Sigma_{q_i}} (\operatorname{Sym}^{q_i} \eta[n, m_i, V_i])^{C_{kn/\ell_i}(n/\ell_i), \tau}$$

is Σ_{q_i} -faithful by Proposition V.3.6.(a) and Theorem V.3.5. Write

$$\bar{\sigma}_i \colon C_{kn/\ell_i}(n/\ell_i) \longrightarrow \Sigma_{q_i}$$

for the composite of the canonical isomorphism $C_{kn/\ell_i}(n/\ell_i) \cong C_{kn}(n)/C_{\ell_i}(n)$ and the homomorphism $C_{kn}(n)/C_{\ell_i}(n) \to \Sigma_{q_i}$ induced by σ_i . Let b_i be the element in the base of ξ in the $\bar{\sigma}_i$ summand corresponding to the projection of b'. Because σ is fixed by the action of s on \mathcal{H}_k , $\bar{\sigma}$ is fixed by the action of s_i on $\operatorname{Hom}(C_{kn/\ell_i}(n/\ell_i), \Sigma_{q_i})$, and because $bs = b, \ b_i s_i = b_i$. We can therefore find e_i in the fiber of ξ at b_i such that $e_i s_i \neq e_i$.

Let e' be the element of $E(\eta)_{b'}$ that is zero in the coordinates for $\eta[n, m_j, V_j]^{q_j}$ with $j \neq i$ and is given by e_i in the coordinate for $\eta[n, m_i, V_i]^{q_i}$. Because e_i is a $\bar{\sigma}_i$ -twisted $C_{kn/\ell_i}(n/\ell_i)$ fixed point of the total space of $\operatorname{Sym}^{q_i} \eta[n, m_i, V_i]$ viewed as a $(\mathbb{T}(n/\ell_i), \Sigma_{q_i})$ vector bundle, it is a σ_i -twisted $C_{kn}(n)$ fixed point of the total space of $\operatorname{Sym}^{q_i} \eta[n, m_i, V_i]$ viewed as a $(\mathbb{T}(n), \Sigma_{q_i})$ vector bundle. It follows that e'is a σ -twisted $C_{kn}(n)$ fixed point of $E(\eta)_{b'}$. We then have a corresponding point ein E_b , and because $e_i s_i \neq e_i$, $es \neq e$.

PROOF OF LEMMA 1.4.(II). Let $A, B \in \mathcal{A}^{\text{cyc}}$. Define \mathcal{Q} by taking $n = \text{lcm}(n^A, n^B)$ and concatenating the data for $(n/n^A)\mathcal{Q}_A$ and $(n/n^B)\mathcal{Q}_B$:

$$r = r^{A} + r^{B} \qquad \qquad Q = Q^{A} \times Q^{B}$$

$$m_{i} = \begin{cases} (n/n^{A})m_{i}^{A} & i \leq r^{A} \\ (n/n^{B})m_{i-r_{A}}^{B} & i > r^{A} \end{cases} \qquad \qquad q_{i} = \begin{cases} q_{0}^{A} + q_{0}^{B} & i = 0 \\ q_{i}^{A} & 0 < i \leq r^{A} \\ q_{i-r_{A}}^{B} & i > r^{A} \end{cases}$$

$$\ell_{i} = \begin{cases} (n/n^{A})\ell_{i}^{A} & i \leq r^{A} \\ (n/n^{B})\ell_{i-r_{A}}^{B} & i > r^{A} \end{cases} \qquad \qquad V_{i} = \begin{cases} V_{i}^{(n/n^{A})}\mathcal{Q}_{A} & i \leq r^{A} \\ V_{i-r_{A}}^{(n/n^{B})}\mathcal{Q}_{B} & i > r^{A} \end{cases}$$

(We use block sum to identify $\Sigma_{q_0^A} \times \Sigma_{q_0^B}$ as a subgroup of Σ_{q_0} , and define Q as the subgroup of $\Sigma_{q_0} \times \cdots \times \Sigma_{q_r}$ corresponding to $Q^A \times Q^B$.) Since $Q_A, Q_B \in D(\mathcal{A}^{cyc})$, so are $(n/n^A)Q_A, (n/n^B)Q_B$ and it is easy to see that $Q \in D(\mathcal{A}^{cyc})$. We then have $n^{(n/n^A)Q_A} = n = n^{(n/n^B)Q_B}$, and an isomorphism of $(\mathbb{T}(n), Q)$ vector bundles

$$\theta_{\mathcal{Q}} \cong \theta_{(n/n^A)\mathcal{Q}_A} \times \theta_{(n/n^B)\mathcal{Q}_B}.$$

Let X be the subspace of the base of θ_Q corresponding to $X_A \times X_B$ under this isomorphism (and the isomorphism of Proposition 3.5.(ii)). Combining this isomorphism with the isomorphisms from Proposition V.1.6 and 3.5.(iii), we get an isomorphism

$$J^{\mathbb{T}}(\theta_{\mathcal{Q},X}) \cong J^{\mathbb{T}}(\theta_{(n/n^A)(\mathcal{Q}_A,X_A)}) \wedge J^{\mathbb{T}}(\theta_{(n/n^B)(\mathcal{Q}_B,X_B)}) \cong A \wedge B.$$

When A and B are in $\mathcal{A}_n^{\text{cyc}}$, n and m_i are powers of p and so $\mathcal{Q} \in D(\mathcal{A}_n^{\text{cyc}})$.

For the proof of Lemma 1.4.(iii), write $\rho_k(kn)$ for the kth root isomorphism $\mathbb{T}(kn) \cong \mathbb{T}(n)$ and $\rho_k(kn)^*$ (universally) for the resulting isomorphisms from categories of $\mathbb{T}(n)$ -equivariant objects to the corresponding categories of $\mathbb{T}(kn)$ -equivariant

objects. These change of group functors $\rho_k(kn)^*$ and the change of group functors ρ_k^* (for $\mathbb{T} \to \mathbb{T}/C_k$) typically commute in the expected way with reasonable functors between $\mathbb{T}(n)$ -spectra and $\mathbb{T}(n')$ -spectra. For example, we have a natural isomorphism

$$\Phi_k = \rho_k^* \circ \Phi^{C_k} \cong \Phi^{\Lambda(k)} \circ \rho_k(k)^*$$

We have $(\rho_k(kn))^{-1}(\Lambda(n)) = C_n(kn)$, so for \mathcal{Q}, X as in Definition 3.2, we have that $\rho_k(kn)^*(\theta_{\mathcal{Q},X})$ is a $\mathbb{T}(k) \cong \mathbb{T}(kn)/C_n(kn)$ -compatible $(\mathbb{T}(kn), Q^{\mathcal{Q}})$ vector bundle and we get an isomorphism

$$\rho_k(k)^*(J^{\mathbb{T}}\theta_{\mathcal{Q},X}) \cong J^{\mathbb{T}(k)}(\rho_k(kn)^*(\theta_{\mathcal{Q},X}))$$

Combining the previous two isomorphisms, we get an isomorphism

$$\Phi_k(J^{\mathbb{T}}\theta_{\mathcal{Q},X}) \cong \Phi^{\Lambda(k)}(J^{\mathbb{T}(k)}(\rho_k(kn)^*(\theta_{\mathcal{Q},X})))$$

PROOF OF LEMMA 1.4.(III). Given $(\mathcal{Q}, X) \in D(\mathcal{A}^{cyc})$, it suffices to construct a pair $(\mathcal{Q}(k), X(k))$ in $D(\mathcal{A}^{cyc})$ and an isomorphism

$$\Phi^{\Lambda(k)}(J^{\mathbb{T}(k)}(\rho_k(kn)^*(\theta_{\mathcal{Q},X}))) \cong J^{\mathbb{T}}\theta_{\mathcal{Q}(k),X(k)}$$

where $n := n^{\mathcal{Q}}$. Since $\theta_{\mathcal{Q},X}$ is $Q := Q^{\mathcal{Q}}$ -faithful, so is $\rho_k(kn)^*(\theta_{\mathcal{Q},X})$, and we will take $Q^{\mathcal{Q}(k)} = Q$. Then by Theorem V.2.4, it suffices to construct an isomorphism of $(\mathbb{T}(kn), Q)$ vector bundles

$$(\rho_k(kn)^*(\theta_{\mathcal{Q},X}))(\Lambda(kn)|C_n(kn)) \cong \theta_{\mathcal{Q}(k),X(k)}.$$

We begin with the definition of $\mathcal{Q}(k)$. Let

$$n^{\mathcal{Q}(k)} := kn, \qquad r^{\mathcal{Q}(k)} := r := r^{\mathcal{Q}}, \\ m_i^{\mathcal{Q}(k)} := m_i := m_i^{\mathcal{Q}}, \qquad \ell_i^{\mathcal{Q}(k)} := \ell_i := \ell_i^{\mathcal{Q}}, \\ q_i^{\mathcal{Q}(k)} := q_i := q_i^{\mathcal{Q}}, \qquad Q^{\mathcal{Q}(k)} := Q := Q^{\mathcal{Q}}, \\ V_i^{\mathcal{Q}(k)} := \rho_k (kn)^* (V_i^{\mathcal{Q}}).$$

We note that since $V_i^{\mathcal{Q}}$ is a semiregular permutation representation of $C_{m_i}(n)/C_{\ell_i}(n)$, $V_i^{\mathcal{Q}(k)}$ is a semiregular permutation representation of $C_{m_i}(kn)/C_{\ell_i}(kn)$. We see that since $(\mathcal{Q}, X) \in D(\mathcal{A}^{cyc})$, so is $(\mathcal{Q}(k), X(k))$ for any choice of X(k), and when k is a power of p, if $(\mathcal{Q}, X) \in D(\mathcal{A}^{cyc}_p)$, then so is $(\mathcal{Q}(k), X(k))$.

Next we analyze $\rho_k(kn)^*(\theta_Q)$. First we note that

$$\rho_k(kn)^*(\eta[n,m,V]) = \eta[kn,m,\rho_k(kn)^*(V)]$$

and so we get

$$\rho_k(kn)^*(\eta_{\mathcal{Q}}) = \eta_{\mathcal{Q}(k)}.$$

Let $\rho_k(kn)^* \operatorname{Hom}(\mathcal{Q})$ denote the set of homomorphisms $C_n(kn) \to Q$ such that for each $i = 1, \ldots, r$ the composite

$$C_n(kn) \longrightarrow Q \longrightarrow \Sigma_{q_0} \times \cdots \times \Sigma_{q_r} \longrightarrow \Sigma_{q_i}$$

has kernel containing $C_{\ell_i}(kn)$. Then we have

$$\rho_k(kn)^*(\theta_{\mathcal{Q}}) \cong \prod_{\tau \in \rho_k(kn)^* \operatorname{Hom}(\mathcal{Q})} \eta_{\mathcal{Q}(k)}^{C_n(kn),\tau}$$

Write $B = \coprod B_{\tau}$ and $E = \coprod E_{\tau}$ for the base and total space of $\rho_k(kn)^*(\theta_Q)$ where we write B_{τ} and E_{τ} for the base and total space of $\eta_{Q(k)}^{C_n(kn),\tau}$. By Definition V.2.13, $(\rho_k(kn)^*(\theta_Q))(\Lambda(kn)|C_n(kn))$ is the restriction of

$$(\rho_k(kn)^*(\theta_{\mathcal{Q}}))(\Lambda(kn)) = \prod_{\sigma: \Lambda(kn) \to Q} (\rho_k(kn)^*\theta(\mathcal{Q}))^{\Lambda(kn),\sigma}$$

along the inclusion of the subset

$$\prod_{\sigma: \Lambda(kn) \to Q} B^{\Lambda(kn),\sigma} \{ C_n(kn) \} \subset \prod_{\sigma: \Lambda(kn) \to Q} B^{\Lambda(kn),\sigma}$$

of the base of $(\rho_k(kn)^*(\theta_Q))(\Lambda(n))$. By definition, an element $b \in B^{\Lambda(kn),\sigma}$ in the σ summand is in the subset $B^{\Lambda(kn),\sigma}\{C_n(kn)\}$ when the $\sigma|_{C_n(kn)}$ -twisted $C_n(kn)$ action on the fiber E_b is trivial. Since $\rho_k(kn)^*(\theta_Q)$ is Q-faithful, this happens exactly when b is in the $\tau = \sigma|_{C_n(kn)}$ summand of B. This subset is either empty (when $\sigma|_{C_n(kn)} \notin \rho_k(kn)^* \operatorname{Hom}(Q))$ or consists of the summand $B_{\sigma|_{C_n(kn)}} \subset B$ (when $\sigma|_{C_n(kn)} \in \rho_k(kn)^* \operatorname{Hom}(Q)$). The condition that a homomorphism $\sigma \colon \Lambda(kn) \to Q$ satisfy $\sigma|_{C_n(kn)} \in \rho_k(kn)^* \operatorname{Hom}(Q)$ is a hypothesis on kernels that coincides with the condition $\sigma \in \operatorname{Hom}(Q(k))$. This gives an isomorphism

$$(\rho_k(kn)^*(\theta_{\mathcal{Q}}))(\Lambda(kn)|C_n(kn)) \cong \prod_{\sigma \in \operatorname{Hom}(\mathcal{Q}(k))} \eta_{\mathcal{Q}(k)}^{\Lambda(kn),\sigma} = \theta_{\mathcal{Q}(k)}$$

Taking X(k) to be the subspace of the base of $\theta_{\mathcal{Q}(k)}$ corresponding under this isomorphism to the subspace of the base of $(\rho_k(kn)^*(\theta_{\mathcal{Q}}))(\Lambda(kn)|C_n(kn))$ given by the base of $(\rho_k(kn)^*(\theta_{\mathcal{Q},X}))(\Lambda(kn)|C_n(kn))$, the isomorphism above restricts to give the isomorphism

$$(\rho_k(kn)^*(\theta_{\mathcal{Q},X}))(\Lambda(kn)|C_n(kn)) \cong \theta_{\mathcal{Q}(k),X(k)}.$$

PROOF OF LEMMA 1.4.(IV). Given \mathcal{Q}, X as in Definition 3.2 and $k \geq 1$, we construct $S^k(\mathcal{Q}, X) := (S^k \mathcal{Q}, S^k X)$ as follows. Let $S^k \mathcal{Q}$ be the data with the same values for n, r, m_i, ℓ_i , and V_i as \mathcal{Q} , but with $q_i^{S^k \mathcal{Q}} = kq_i$ and

$$Q^{S^k\mathcal{Q}} < \Sigma_k \wr \Sigma_{q_0} \times \cdots \times \Sigma_k \wr \Sigma_{q_r} < \Sigma_{kq_0} \times \cdots \times \Sigma_{kq_r}$$

the subset where the projection to $(\Sigma_k)^r$ lands in the diagonal and for each $j = 1, \ldots, k$ the *j*th factor projection

$$\Sigma_k \wr \Sigma_{q_0} \times \cdots \times \Sigma_k \wr \Sigma_{q_r} = \Sigma_k \ltimes \Sigma_{q_0}^k \times \cdots \times \Sigma_k \ltimes \Sigma_{q_r}^k \longrightarrow \Sigma_{q_0} \times \cdots \times \Sigma_{q_r}$$

lands in Q. In other words, the image of Q^{S^kQ} in $\Sigma_{kq_1+\cdots+kq_r}$ is conjugate to the image of $\Sigma_k \wr Q$ via the permutation associated to the canonical bijection

$$(\{1,\ldots,q_0\}\amalg\cdots\amalg\{1,\ldots,q_r\})^{\amalg k}\cong\{1,\ldots,q_0\}^{\amalg k}\amalg\cdots\amalg\{1,\ldots,q_r\}^{\amalg k}.$$

Writing ψ for the isomorphism $Q^{S^k \mathcal{Q}} \cong \Sigma_k \wr Q$, we get an isomorphism of $(\mathbb{T}(n), Q^{S^k \mathcal{Q}})$ vector bundles

$$\psi^*(\operatorname{Sym}^k \eta_{\mathcal{Q}}) \cong \eta_{S^k \mathcal{Q}}.$$

Let $S^k \operatorname{Hom}(\mathcal{Q})$ be the subset of $\operatorname{Hom}(S^k \mathcal{Q})$ of those homomorphisms $\sigma \colon \Lambda(n) \to Q^{S^k \mathcal{Q}}$ whose composite with ψ

$$\Lambda(n) \longrightarrow Q^{S^k \mathcal{Q}} \xrightarrow{\cong} \Sigma_k \wr Q$$

lands in Q^k . Then we get an isomorphism of $(\mathbb{T}(n), \mathcal{Q}^{S^k \mathcal{Q}})$ vector bundles

$$\psi^*(\operatorname{Sym}^k \theta_{\mathcal{Q}}) \cong \coprod_{\sigma \in S^k \operatorname{Hom}(\mathcal{Q})} \eta_{S^k \mathcal{Q}}^{\Lambda(n), \sigma}$$

This is the restriction of $\theta_{S^k\mathcal{Q}}$ to the summands indexed by $\sigma \in S^k \operatorname{Hom}(\mathcal{Q}) \subset \operatorname{Hom}(S^k\mathcal{Q})$. Let S^kX be the subspace of the base of $\theta_{S^k\mathcal{Q}}$ corresponding to the subspace ψ^*X^k in the base of $\psi^*\operatorname{Sym}^k\theta_{\mathcal{Q}}$. The isomorphism above then induces an isomorphism

$$J^{\mathbb{T}}_{(\mathbb{T}(n),\Sigma_k\wr Q)}(\operatorname{Sym}^k \theta_{\mathcal{Q},X}) \cong J^{\mathbb{T}}_{(\mathbb{T}(n),Q^{S^k}\mathcal{Q})}(\psi^*(\operatorname{Sym}^k \theta_{\mathcal{Q},X})) \longrightarrow J^{\mathbb{T}}\theta_{S^k(\mathcal{Q},X)}.$$

We note that when (\mathcal{Q}, X) is in $D(\mathcal{A}^{cyc})$ or $D(\mathcal{A}^{cyc}_p)$, then so is $S^k(\mathcal{Q}, X)$.

PROOF OF LEMMA 1.4. (V). This follows from part (i) and Theorem V.2.8. \Box

PROOF OF LEMMA 1.4.(VI). Given \mathcal{Q}, X as in Definition 3.2, define \mathcal{Q}' to have the same values for n, r, m_i, ℓ_i, V_i , and q_i for $i \geq 1$ as \mathcal{Q} , but with $q_0^{\mathcal{Q}'} = q_0^{\mathcal{Q}} + 1$. Regarding Σ_q as a subgroup of Σ_{q+1} in the usual way, we can view $Q^{\mathcal{Q}}$ as a subgroup of

$$\Sigma_{q_0^{\mathcal{Q}}+1} \times \Sigma_{q_1^{\mathcal{Q}}} \times \cdots \times \Sigma_{q_{r\mathcal{Q}}^{\mathcal{Q}}} = \Sigma_{q_0^{\mathcal{Q}'}} \times \Sigma_{q_{r\mathcal{Q}'}^{\mathcal{Q}'}}$$

and we take $Q^{\mathcal{Q}'} = Q^{\mathcal{Q}}$. Then $\theta_{\mathcal{Q}'}$ and $\theta_{\mathcal{Q}}$ have the same base space, so $\theta_{\mathcal{Q}',X}$ makes sense, and it is easy to see that $J^{\mathbb{T}}\theta_{\mathcal{Q}',X}$ is isomorphic to $J^{\mathbb{T}}\theta_{\mathcal{Q},X} \wedge F_{\mathbb{R}}S^0$.

4. The positive convenient Σ -model structure and Theorem 1.3

This section defines the positive convenient Σ -model structure on non-equivariant spectra and proves Theorem 1.3 which compares cofibrations in this structure with cofibrations in the model structures on \mathbb{T} -spectra from Theorem 1.2.

We begin with the definition of the model structure. The model category results from Section II.II apply in particular to non-equivariant spectra by taking the compact Lie group G to be the trivial group 1. Using this theory, we specify the positive convenient Σ -model structure by specifying a set of spectra containing the set $\mathcal{A}(\mathcal{F}, \mathcal{V}_+)$ (where \mathcal{F} is the unique family of subgroups of the trivial group and \mathcal{V}_+ is the representation constraint of Example II.1.2). In the definition, we use the Σ_q action on $F_{\mathbb{R}^q}S^0$ induced by the action of O(q) on \mathbb{R}^q .

DEFINITION 4.1. Let \mathcal{A}_{Σ} denote the set of spectra $(F_{\mathbb{R}^q}S^0)/Q$ where $Q < \Sigma_q$. The positive complete Σ model structure on spectra is the \mathcal{A}_{Σ} -variant \mathcal{V}_+ -constrained model structure of Theorem II.1.11.

One direction of Theorem 1.3 is easy. Applying the free functor to a spectrum $(F_{\mathbb{R}^n}S^0)/Q$ in \mathcal{A}_{Σ} , we get the T-spectrum $\mathbb{T}_+ \wedge (F_{\mathbb{R}^n}S^0)/Q$. This is isomorphic to the T-spectrum $J\theta_{\mathcal{Q},\Delta}$ in $\mathcal{A}_p^{\text{cyc}}$ where Δ is the diagonal of $\mathbb{T}^q = \mathbb{T}(1)^q$ and \mathcal{Q} is the data where $n = 1, r = 1, q_0 = 0, q_1 = q, m_1 = 1, \ell_1 = 1, Q = Q$, and $V = \mathbb{R}^1$. In other words, it is the restriction to the diagonal T in the base of the Q^{op} -equivariant vector bundle

$$\operatorname{Sym}^q(\mathbb{T} \times \mathbb{R} \longrightarrow \mathbb{T})$$

with the group for quotients reduced to $Q < \Sigma_q$. Since the free functor sends spectra in \mathcal{A}_{Σ} to spectra in $\mathcal{A}_p^{\text{cyc}}$ (up to isomorphism), it follows that it sends cofibrations in the positive convenient Σ -model structure on spectra to cofibrations in either of the model structures on \mathbb{T} -spectra in Theorem 1.2.

The proof of Theorem 1.3 is then completed by showing that the forgetful functor sends objects in \mathcal{A}^{cyc} to cofibrant objects in the positive convenient Σ -model structure. The main observation we need for that is the following lemma.

LEMMA 4.2. Let $Q \in D(\mathcal{A}^{cyc})$, let $Q = Q^{\mathcal{Q}}$, and let f be a Q^{op} -equivariant map from an orbit Q^{op}/P^{op} to the base of θ_Q . Then the spectrum $J_{(1,Q)}f^*\theta_Q$ constructed from the $(1, Q^{op})$ vector bundle $f^*\theta_Q$ is isomorphic to an object in \mathcal{A}_{Σ} .

PROOF OF THEOREM 1.3 FROM LEMMA 4.2. As observed above, it suffices to show that the forgetful functor sends objects in \mathcal{A}^{cyc} to cofibrant objects in the positive convenient Σ -model structure. Let $(\mathcal{Q}, X) \in D(\mathcal{A}^{cyc})$. The underlying nonequivariant spectrum of the corresponding element $J^{\mathbb{T}}\theta_{\mathcal{Q},X}$ of \mathcal{A}^{cyc} is the generalized orbit desuspension spectrum $J_{(1,Q)}\theta_{\mathcal{Q},X}$ (for the trivial group). Ignoring the $\mathbb{T}(n)$ action, X is a Q^{op} -cell complex and for any Q^{op} -cell α : $Q^{op}/P^{op} \times D^s \to X$, base change gives a (1,Q) vector bundle $\alpha^*\theta_{\mathcal{Q},X}$ and a spectrum $J_{(1,Q)}(\alpha^*\theta_{\mathcal{Q},X})$. We write $J_{(1,Q)}(\partial \alpha^* \theta_{\mathcal{Q},X})$ for the spectrum obtained the same way on the boundary $\partial \alpha: Q/P \times \partial D^s \to X$. The spectrum $J_{(1,Q)}\theta_{\mathcal{Q},X}$ is built as a sequential colimit of pushouts over cofibrations of these restrictions. It is therefore enough to show that for every Q^{op} -cell α , the pair

$$(J_{(1,Q)}(\alpha^*\theta_{\mathcal{Q},X}), J_{(1,Q)}(\partial\alpha^*\theta_{\mathcal{Q},X}))$$

is isomorphic to a cell $A \wedge (D^s, \partial D^s)$ for A in \mathcal{A}_{Σ} . Using the contraction to the center of the disk, we see that

$$(\alpha^* \theta_{\mathcal{O},X}, \partial \alpha^* \theta_{\mathcal{O},X}) \cong \eta \times (D^s, \partial D^s)$$

for the (1, Q) vector bundle η over $Q^{\text{op}}/P^{\text{op}}$ obtained by base change over the restriction of α to the center point. The result now follows from Lemma 4.2.

The rest of the section proves Lemma 4.2. We fix

$$\mathcal{Q} = (n, r, (q_0, \dots, q_r), (m_1, \dots, m_r), (\ell_1, \dots, \ell_r), (V_1, \dots, V_r)) \in D(\mathcal{A}^{cyc})$$

and a point $x = (x_{1,1}, \ldots, x_{r,q_r})$ in the base of θ_Q . Let $\sigma \colon \Lambda(n) \to Q$ in Hom(Q)denote the homomorphism indexing the component of x. We further fix a subgroup P < Q which is contained in the isotropy subgroup of x for the right Q action on θ_Q , which then specifies the Q^{op} -equivariant map f from $Q^{\text{op}}/P^{\text{op}}$ to the base of θ_Q (sending the base point of the orbit to the point x). Let $\theta = f^* \theta_Q$; we need to show that $J_{(1,Q)}\theta$ is isomorphic to an object of \mathcal{A}_{Σ} .

The argument proceeds by studying the fiber of θ over the basepoint and the inherent right P action on it. Writing E_x for this fiber and $V_{i,j}$ for the fiber of $\eta[n, m_i, V_i]$ over the point $x_{i,j} \in \mathbb{T}(n)/C_{m_i}(n)$, the construction of θ_Q identifies E_x as

$$(\mathbb{R}^{q_0} \times V_{1,1} \times \cdots \times V_{r,q_r})^{\Lambda(n),\sigma}.$$

Writing $\sigma_i \colon \Lambda(n) \to \Sigma_{q_i}$ for the composite map

$$\sigma_i \colon \Lambda(n) \longrightarrow Q \longrightarrow \Sigma_{q_0} \times \cdots \times \Sigma_{q_r} \longrightarrow \Sigma_{q_i},$$

we can further identify E_x as

$$E_x = (\mathbb{R}^{q_0})^{\Lambda(n),\sigma_0} \times (V_{1,1} \times \cdots \times V_{1,q_1})^{\Lambda(n),\sigma_1} \times \cdots \times (V_{r,1} \times \cdots \times V_{r,q_r})^{\Lambda(n),\sigma_r}.$$

The zeroth factor $(\mathbb{R}^{q_0})^{\Lambda(n),\sigma_0}$ is the fixed point set of the action of the subgroup $\sigma_0(\Lambda(n)) < \Sigma_{q_0}$ given by the image of σ_0 . Since $\mathbb{R}^{q_0} = \mathbb{R}\langle \{1, \ldots, q_0\} \rangle$ is a permutation representation of Σ_{q_0} , the fixed point set is also a permutation representation

$$(\mathbb{R}^{q_0})^{\Lambda(n),\sigma_0} \cong \mathbb{R}\langle \{1,\ldots,q_0\} / \sigma_0(\Lambda(n)) \rangle.$$

A choice of enumeration of $\{1, \ldots, q_0\}/\sigma_0(\Lambda(n))$ as $\{1, \ldots, k_0\}$ defines a homomorphism $\Sigma_{q_0} \to \Sigma_{k_0}$ and an isomorphism of Σ_{q_0} -representations

$$(\mathbb{R}^{q_0})^{\Lambda(n),\sigma_0} \cong \mathbb{R}^{k_0}$$

with the action of Σ_{q_0} on \mathbb{R}^{k_0} via the homomorphism, and in particular, by permutation of coordinates.

To complete the proof of Lemma 4.2, it suffices to show that for each i there exists an orthonormal basis of

$$(V_{i,1} \times \cdots \times V_{i,q_i})^{\Lambda(n),\sigma_i}$$

such that the P action on the vector space restricts to an action on the basis: the choice of these bases specify a homomorphism

$$P \longrightarrow \Sigma_{k_1} \times \cdots \times \Sigma_{k_r} \longrightarrow \Sigma_k$$

(for $k = \sum k_r$) and an isomorphism of (1, Q) vector bundles

$$\theta \cong \mathbb{R}^k \times_P Q \longrightarrow P \backslash Q,$$

which then induces an isomorphism of spectra

$$J_{(1,Q)}\theta \cong (F_{\mathbb{R}^k}S^0)/P \cong (F_{\mathbb{R}^k}S^0)/H$$

where H is the image of P in Σ_k .

We can now concentrate on an individual i, which amounts to studying the σ_i twisted $\Lambda(n)$ fixed points of the vector bundle $\operatorname{Sym}^{q_i} \eta[n, m_i, V_i]$. Fixing i, we can tighten up notation significantly. As a first step, since the action of $C_{\ell_i}(n)$ on V_i is trivial and $C_{\ell_i}(n)$ is in the kernel of σ_i , we get an equivalent problem if we replace $\mathbb{T}(n)$ with $\mathbb{T}(n)/C_{\ell_i}(n)$, $\Lambda(n)$ with $\Lambda(n)/C_{\ell_i}(n)$ and $C_{m_i}(n)$ with $C_{m_i}(n)/C_{m_i}(\ell_i)$: let

$$\Gamma = \mathbb{T}(n)/C_{\ell_i}(n), \quad \Lambda = \Lambda(n)/C_{\ell_i}(n), \quad C = C_{m_i}(n)/C_{\ell_i}(n).$$

We also let $q = q_i$, $V = V_i$, $W_j = V_{i,j}$, and $\sigma' = \sigma_i$. In this notation, the hypothesis that $Q \in D(\mathcal{A}^{\text{cyc}})$ implies that V and each W_j is a non-zero permutation representation of C, and we are studying the pullback (1, Q) vector bundle

$$\begin{array}{c} Q^{\mathrm{op}} \times_{P^{\mathrm{op}}} (V^q)^{\Lambda,\sigma'} & & \cdots & ((\Gamma \times_C V)^q)^{\Lambda,\sigma} \\ & & \theta' \downarrow & & \downarrow \\ & Q^{\mathrm{op}}/P^{\mathrm{op}} & & \cdots & ((\Gamma/C)^q)^{\Lambda,\sigma'}. \end{array}$$

Let $x' = (x'_1, \ldots, x'_q)$ for $x'_j = x_{i,j}$ be the point in θ' obtained by restricting coordinates. We now recycle the symbols i, ℓ, m, n , and r for reuse.

Section VI.6 studies twisted fixed points on cartesian powers of bundles. (To apply this section, understand the group Q in the work there to be the trivial group.) The homomorphism $\sigma' \colon \Lambda \to \Sigma_q$ endows the set $\{1, \ldots, q\}$ with a Λ action. Let O_1, \ldots, O_k be the orbits for this action, and choose an element $m_j \in O_j$ for each $j = 1, \ldots, k$. Let $H_j < \Lambda$ be the isotropy subgroup of m_j . Since Λ is abelian, H_j is the isotropy subgroup for every element in O_j . Moreover, we note that $H_j < \Lambda < \Gamma$ is also a subgroup of C since every element in it fixes $x'_{m_j} \in \Gamma/C$. For each $i \in \{1, \ldots, q\}$, define j_i by $i \in O_{j_i}$ and choose $\ell_i \in \Lambda$ such that $\ell_i \cdot m_{j_i} = i$, with $\ell_{m_j} = id$. Then for the inherent action of Γ on the vector bundle $\Gamma \times_C V \to \Gamma/C$, ℓ_i sends $W_{m_{j_i}}$ (the fiber over $x_{m_{j_i}}$) to W_i (the fiber over x_i). Define

$$\phi \colon W_{m_i}^{H_{m_1}} \times \cdots \times W_{m_k}^{H_{m_k}} \longrightarrow W_1 \times \cdots \times W_q$$

by

$$\phi(w_1, \dots, w_k) := \left(\frac{1}{\sqrt{\#O_{j_1}}} \ell_1 w_{j_1}, \dots, \frac{1}{\sqrt{\#O_{j_q}}} \ell_q w_{j_q}\right).$$

The scaling ensures that this is an isometry, and Lemma VI.6.1 shows that ϕ is an isomorphism onto the σ' -twisted Λ fixed points $(W_1 \times \cdots \times W_q)^{\Lambda,\sigma'}$.

For each $i = 1, \ldots, q$, choose an element \bar{x}_i in Γ lifting the element $x'_i \in \Gamma/C$. The choice of \bar{x}_i induces a *C*-equivariant isomorphism $\psi_i : V \to W_i$. If $x'_i = x'_{i'}$ for some i', then $\bar{x}_{i'}\bar{x}_i^{-1}$ is an element of *C* and the composite endomorphism

$$V \xrightarrow{\psi_i} W_i = W_{i'} \xrightarrow{\psi_{i'}^{-1}} V$$

is induced by $\bar{x}_i^{-1}\bar{x}_{i'} \in C$ for the inherent C action on V. More generally if under the σ' -twisted Λ action, the element $\lambda \in \Lambda$ takes x'_i to $x'_{i'}$, then $\bar{x}_{i'}\lambda^{-1}\bar{x}_i^{-1}$ with the composite endomorphism of V given by multiplication by $\bar{x}_i^{-1}\lambda^{-1}\bar{x}_{i'}$. (These formulas and the C-equivariance of the ψ_i strongly use the fact that Γ is abelian.)

The isomorphisms ψ_i together with the isomorphism ϕ above identify the fiber of x' in θ' as

$$V^{H_{m_1}} \times \cdots \times V^{H_{m_k}}$$

Because V is a permutation representation of C, each V^{H_j} is a permutation representation of C/H_j . We can therefore choose an orthonormal basis of V^{H_j} on which the C action restricts. If $H_j = H_{j'}$, we choose the same basis for V^{H_j} and $V^{H_{j'}}$. We assemble the bases of V^{H_j} to an orthonormal basis of the fiber of x' in θ' .

The action of P on θ' is induced by the permutation action. Since P is contained in the isotropy subgroup of x' for the Q action on the base of θ' , every element of P < Q fixes the homomorphism σ' , and that is equivalent to saying that the right action on P on $\{1, \ldots, q\}$ commutes with the Λ action induced by σ' . In particular, if some element of P sends i to i', then it induces an isomorphism of Λ -sets $O_{j_i} \to O_{j_{i'}}$ and we must have $H_{j_i} = H_{j_{i'}}$. Because for each j, the elements ℓ_i for $i \in O_j$ form a complete list of coset representatives for Λ/H_j , under the identification above, the action of each element $\rho \in P$ on $V^{H_{m_1}} \times \cdots \times V^{H_{m_k}}$ does the permutation on the orbits followed by the action of an element of C on each factor (namely, on the jth factor, it acts by

$$\ell_i \bar{x}_{m_{j_i}} \bar{x}_i^{-1} \bar{x}_i \bar{x}_{m_j}^{-1} = \ell_i \bar{x}_{m_{j_i}} \bar{x}_{m_j}^{-1},$$

where $\rho \in P < Q < \Sigma_q$ sends m_j to *i*). It follows that the action of *P* on the fiber of x' restricts to an action the orthonormal basis constructed above.

CHAPTER IV

The Multiplicative tom Dieck Splitting (Theorem A)

A fundamental structural property of the equivariant stable category is the tom Dieck splitting, which in its original form gave a formula for the categorical fixed points of a suspension spectrum. It arises from the existence of transfers on categorical fixed points. In the multiplicative context, equivariant commutative ring orthogonal spectra come with certain transfers on their geometric fixed points, which we review below. When the equivariant commutative ring spectrum comes from a non-equivariant commutative ring spectrum, these transfers are independent of each other and assemble to a multiplicative splitting that is a multiplicative analogue of the tom Dieck splitting. We gave a concrete statement in Theorem A, which we restate as Theorem 1.4 in Section 1. This chapter just scratches the surface of the properties of the multiplicative transfers, which we believe to be foundational in the equivariant multiplicative theory and intend to return to in future work. Section 2 provides some additional information on the pieces in the splitting in special cases; little beyond this seems to currently be known.

1. Construction of the multiplicative splitting

We begin with the construction of the multiplicative transfer in the context of commutative ring G-spectra (with G a finite group). For H < G, the norm N_H^G restricts to a functor from commutative ring H-spectra to commutative ring G-spectra and it is left adjoint to the reduction of structure functor i_H^* from commutative ring G-spectra to commutative ring H-spectra

$$\mathcal{C}om_H \xleftarrow{N_H^G}{i_H^*} \mathcal{C}om_G$$

In particular, for A a commutative ring G-spectrum, the counit of the adjunction gives a natural map of commutative ring G-spectra

$$N_H^G i_H^* A \longrightarrow A.$$

Applying Φ^G to this map and composing with the Hill-Hopkins-Ravenel diagonal map

$$\Phi^H i_H^* A \longrightarrow \Phi^G N_H^G i_H^* A$$

(which is an isomorphism when A is nice enough) we get a natural map

$$\tau_H \colon \Phi^H i_H^* A \longrightarrow \Phi^G N_H^G i_H^* A \longrightarrow \Phi^G A.$$

of non-equivariant commutative ring orthogonal spectra on the point-set level, or working with derived functors, on the homotopy category level. DEFINITION 1.1. The multiplicative transfer for H < G is the natural map τ_H displayed above.

For the multiplicative tom Dieck splitting, the first observation we need is that τ_H is actually equivariant, where we regard $\Phi^H A$ with its natural WH action and $\Phi^G A$ with its trivial action. The following gives a precise statement.

THEOREM 1.2. The multiplicative transfer $t_H: \Phi^H A \to \Phi^G A$ is a map of left WH-objects in the category of non-equivariant commutative ring orthogonal spectra, where $\Phi^H A$ is a WH-object by neglect of structure (from its commutative ring WH-spectrum structure) and $\Phi^G A$ is a WH-object with trivial action.

PROOF. The key observations are:

- (i) $N_{H}^{G}i_{H}^{*}A$ has a natural action of WH in the category of commutative ring G-spectra,
- (ii) The counit map $N_{H}^{G}i_{H}^{*}A \rightarrow A$ is a map of WH-objects in the category of commutative ring G-spectra where we give the target the trivial action, and
- (iii) The diagonal map $\Phi^H A \to \Phi^G N_H^G i_H^* A$ is a map of WH-objects in non-equivariant commutative ring orthogonal spectra.

For point (i), we use the fact [15, 2.28] that $N_H^G i_H^* A$ is naturally isomorphic to the tensor $G/H \otimes A$ of A with the left G-space G/H, and under this isomorphism, the map $N_H^G i_H^* A \to A$ is induced by the G-map $G/H \to *$. The commutative ring G-spectrum $N_H^G i_H^* A$ then obtains a WH action (in that category) from the right action of WH on G/H. For (ii), formulated this way, the map is obviously a map of WH-objects where the target has the trivial action. For (iii), [15, B.190] implies that the wedge of maps from $\Phi^H F_V X(V)$ to $\Phi^H X$ is a categorical epimorphism, so it suffices to check (iii) in the case when A is free on $F_V G/H_+$, where it is easy from the description of the diagonal in [15, B.209].

The previous theorem was a point set statement. While it is true that the pointset category of left WH-objects in non-equivariant orthogonal spectra is equivalent to the point-set category of WH-equivariant orthogonal spectra, for our purposes, the corresponding homotopy theory statement we need is in terms of the WH-Borel equivariant category of commutative ring spectra. The Borel category is the category of WH-objects in non-equivariant commutative ring orthogonal spectra where we understand the weak equivalences to be the maps that are weak equivalences of the underlying non-equivariant spectra. Homotopically, then we should consider the map

$$\Phi^H A \otimes EWH \longrightarrow \Phi^G A$$

obtained by tensoring with the universal free WH-space. Since the target is trivial, this factors through the quotient (in the category of commutative ring equivariant spectra)

(1.3)
$$\Phi^H A \otimes_{WH} EWH \longrightarrow \Phi^G A.$$

We can now begin to study the case when the commutative ring G-spectrum comes from a non-equivariant commutative ring spectrum R. We have a point-set functor ϵ^* that takes non-equivariant commutative ring spectra to commutative ring G-spectra by giving it the trivial action: precisely, it is the restriction of action along the final homomorphism $G \to 1$. A check of basic definitions shows that ϵ^*

52

is left adjoint to the categorical G fixed point functor, and since categorical fixed point functors preserve fibrations and acyclic fibrations, ϵ^* is a Quillen left adjoint. In particular, ϵ^* admits a left derived functor that is calculated by applying it to a cofibrant object. In what follows, we use ϵ^* to denote this left derived functor, and for convenience, we assume without loss of generality that R is a cofibrant commutative ring spectrum, so that $\epsilon^* R$ is represented by the point-set functor applied to R.

We want to study the left derived geometric fixed points of $\epsilon^* R$, and for this we take the commutative ring *G*-spectrum *A* in the above to be a cofibrant approximation in a model structure satisfying the hypotheses of Theorem II.2.5. We get a map in the homotopy category of non-equivariant commutative ring spectra

$$R \longrightarrow L\Phi^H \epsilon^* R \simeq \Phi^H A$$

as the composite with the canonical map from R to the derived categorical H fixed points of $\epsilon^* R$ with the canonical map from the categorical fixed points to the geometric fixed points. Formally, these are maps in the WH-Borel category of commutative ring orthogonal spectra where R has the trivial action. Composing with the map (1.3) above, we get a well defined map in the homotopy category of non-equivariant commutative ring orthogonal spectra

$$R \otimes^{L} BWH \cong R \otimes^{L}_{WH} EWH \longrightarrow \Phi^{H}A \otimes^{L}_{WH} EWH \longrightarrow L\Phi^{G}\epsilon^{*}R.$$

(This is not a point-set map.) Taking the coproduct over conjugacy classes of subgroups H, we get a map of non-equivariant commutative ring orthogonal spectra

$$\bigwedge_{[H]} R \otimes^L BWH \longrightarrow L\Phi^G \epsilon^* R$$

We prove the following theorem.

THEOREM 1.4 (Multiplicative tom Dieck splitting for ϵ^*). Let R be a non-equivariant commutative ring orthogonal spectrum, and let ϵ^* denote the derived functor from non-equivariant to commutative ring G-spectra. The map

$$\bigwedge_{[H]} R \otimes^L BWH \longrightarrow L\Phi^G \epsilon^* R.$$

induced by the multiplicative transfers is a weak equivalence.

PROOF. We take R to be a cofibrant non-equivariant commutative ring orthogonal spectrum. Both the domain and codomain of the natural map in the statement preserve homotopy colimits in commutative ring spectra, and so it suffices to check the case when R is the domain and codomain of a cell, that is, when $R = \mathbb{P}(F_m X_+), q \ge 1$, the free commutative ring spectrum on $F_m X_+$ for $X = D^n$ or ∂D^n . We can then take A to be the free commutative ring G-spectrum $\mathbb{P}Y$ on $Y = F_{\mathbb{R}^m \oplus \mathbb{R}\langle G \rangle} \Sigma^{\mathbb{R}\langle G \rangle} X_+$, where we have written $\mathbb{R}\langle G \rangle$ for the regular representation of G. While the symmetric powers $Y^{(q)}/\Sigma_q$ of Y are not themselves generalized orbit desuspension spectra, they are each the quotient of a Hurewicz cofibration of generalized orbit desuspension spectra, for the (globally trivial) (G, Σ_q) vector bundles

$$\begin{array}{c} (X^q \times \partial D(\mathbb{R}\langle G \rangle^q)) \times (\mathbb{R}^m \oplus \mathbb{R}\langle G \rangle)^q & \longrightarrow (X^q \times D(\mathbb{R}\langle G \rangle^q)) \times (\mathbb{R}^m \oplus \mathbb{R}\langle G \rangle)^q \\ \downarrow & \downarrow \\ X^q \times \partial D(\mathbb{R}\langle G \rangle^q) & \longrightarrow X^q \times D(\mathbb{R}\langle G \rangle^q) \end{array}$$

for $q \ge 1$. Since the geometric fixed points functor Φ^G commutes with quotients by spacewise closed inclusions, Theorem V.2.4 then gives the isomorphism

$$\Phi^{G}(Y^{(q)}/\Sigma_{q}) \cong \left(\bigvee_{\sigma: \ G \to \Sigma_{q}} F_{((\mathbb{R}^{m} \oplus \mathbb{R}\langle G \rangle)^{q})^{(G,\sigma)}} (\Sigma^{\mathbb{R}\langle G \rangle^{q}} X^{q}_{+})^{(G,\sigma)}\right)/\Sigma_{q}$$

for $q \geq 1$.

The remainder of the argument is a version of the proof of Theorem V.3.1 where (in the notation there) Q is the trivial group. The argument becomes significantly more concrete in this case and we include it to make this section more self-contained.

If we think of a homomorphism $G \to \Sigma_q$ as a G action on $\{1, \ldots, q\}$, much of the quotient by Σ_q above is relating isomorphic G-sets, and we can rewrite the above in terms of isomorphism classes of G-sets of size q:

$$\Phi^{G}(Y^{(q)}/\Sigma_{q}) \cong \bigvee_{[Z], \#Z=q} (F_{((\mathbb{R}^{m} \oplus \mathbb{R}\langle G \rangle) \otimes \mathbb{R}\langle Z \rangle)^{G}} (\Sigma^{\mathbb{R}\langle G \times Z \rangle} X_{+}^{Z})^{G} / \operatorname{Aut}_{G}(Z))$$

where the action is the combined action of G on Z and the other terms. The previous formula was only for $q \ge 1$, but when Z is the empty set the formula reduces to a copy of \mathbb{S} , which is the zero summand of A. Adding up over $q \ge 0$, we get a wedge over isomorphism classes of finite G-sets Z

$$\Phi^G A \cong \bigvee_{[Z]} (F_{((\mathbb{R}^m \oplus \mathbb{R}\langle G \rangle) \otimes \mathbb{R}\langle Z \rangle)^G} (\Sigma^{\mathbb{R}\langle G \times Z \rangle} X^Z_+)^G / \operatorname{Aut}_G(Z)).$$

The construction

Ì

$$F_{((\mathbb{R}^m \oplus \mathbb{R}\langle G \rangle) \otimes \mathbb{R}\langle Z \rangle)^G} (\Sigma^{\mathbb{R}\langle G \times Z \rangle} X_+^Z)^G$$

takes coproducts in Z to smash products of spectra. Reorganizing the wedge above in terms of isomorphism classes of orbits then gives

$$\Phi^G A \cong \bigwedge_{[H]} \left(\bigvee_{q \ge 0} (F_{(\mathbb{R}^m \oplus \mathbb{R}\langle G/H \rangle)^q} (\Sigma^{\mathbb{R}\langle G/H \rangle} X_+)^{(q)}) / \Sigma_q \wr WH \right)$$

with the outer smash product indexed over conjugacy classes of subgroups and where we have identified

$$\operatorname{Aut}_G(G/H_1^{q_1}\amalg\cdots\amalg G/H_r^{q_r})\cong (\Sigma_{q_1}\wr WH_1)\times\cdots\times (\Sigma_{q_r}\wr WH_r)$$

for pairwise non-conjugate subgroups H_1, \ldots, H_r . Finally, moving the quotients by WH inside the quotients by Σ_q , we get

$$\Phi^G A \cong \bigwedge_{[H]} \left(\bigvee_{q \ge 0} ((F_{\mathbb{R}^m \oplus \mathbb{R} \langle G/H \rangle} (\Sigma^{\mathbb{R} \langle G/H \rangle} X_+)) / WH)^{(q)} / \Sigma_q \right)$$

The map in the homotopy category of non-equivariant commutative ring orthogonal spectra $R \to \Phi^G A$ induced by τ_H is determined by a map in the derived category of non-equivariant spectra

$$F_{\mathbb{R}^m}X_+ \longrightarrow \Phi^G A.$$

54

Using the description of the diagonal map from [15, B.209], we can identify this as the map

$$F_{\mathbb{R}^m}X_+ \xleftarrow{\simeq} F_{\mathbb{R}^m \oplus \mathbb{R}\langle G/H \rangle} \Sigma^{\mathbb{R}\langle G/H \rangle}X_+ \longrightarrow (F_{\mathbb{R}^m \oplus \mathbb{R}\langle G/H \rangle}(\Sigma^{\mathbb{R}\langle G/H \rangle}X_+))/WH$$

followed by the inclusion of the latter as the q = 1 summand in the smash factor for G/H. Because the action of WH on G/H is faithful, the constituent spaces of $F_{\mathbb{R}^m \oplus \mathbb{R}\langle G/H \rangle}(\Sigma^{\mathbb{R}\langle G/H \rangle}X)$ are free based WH-CW complexes and the maps

$$F_{\mathbb{R}^m} X_+ \wedge BWH = (F_{\mathbb{R}^m} X_+) \wedge_{WH} EWH$$
$$\stackrel{\simeq}{\longleftarrow} (F_{\mathbb{R}^m \oplus \mathbb{R}\langle G/H \rangle} (\Sigma^{\mathbb{R}\langle G/H \rangle} X_+)) \wedge_{WH} EWH$$
$$\longrightarrow (F_{\mathbb{R}^m \oplus \mathbb{R}\langle G/H \rangle} (\Sigma^{\mathbb{R}\langle G/H \rangle} X_+))/WH$$

are weak equivalences. It follows that the map $R \otimes BWH \to \Phi^G A$ induces a weak equivalence into the G/H smash factor, and the smash of these maps is a weak equivalence.

2. The factors in the multiplicative splitting

For R a (non-equivariant) commutative ring orthogonal spectrum, the commutative ring orthogonal spectra $R \otimes^L BWH$ in the tom Dieck splitting are examples of *Loday constructions* on R. Most work to date on Loday constructions $R \otimes^L X$ study examples when X is a finite complex, most often a sphere or product of spheres. The multiplicative tom Dieck splitting of the previous section provides new motivation to study Loday constructions when X is the classifying space of a finite group. While little seems to be known calculationally about this case, we do offer the following result.

THEOREM 2.1. Let R be a connective commutative ring orthogonal spectrum and let G be a finite group. If the cardinality of G is invertible in $\pi_0 R$, then the map $R \to R \otimes^L BG$ induced by the inclusion of the base point of BG is a weak equivalence.

The theorem is an immediate consequence of the following theorem, which is well-known to experts.

THEOREM 2.2. Let R be a commutative ring orthogonal spectrum and $X \to Y$ a map of CW spaces that induces an isomorphism on R-homology. Then $R \otimes^L X \to R \otimes Y$ is a weak equivalence.

For the proof, we follow the outline of [10, 2.7] (and [9, 2.2.1.3]).

PROOF. We can assume without loss of generality that R is a cofibrant commutative ring spectrum, and then the derived tensor $R \otimes^L X$ is calculated by the point-set tensor $R \otimes X$ for any CW complex X. Because $R \otimes X$ is continuous in X, it preserves homotopies in X, and so it suffices to consider the case when X, Y are geometric realizations of simplicial sets X_{\bullet}, Y_{\bullet} , and the map $X \to Y$ is the geometric realization of a simplicial map. We also assume that X_{\bullet} and Y_{\bullet} are non-empty.

Let **F** denote the skeleton of the category of non-empty finite sets, with objects $\mathbf{n} = \{1, \ldots, n\}$ for n > 0. Since R is a commutative monoid object for \wedge , this structure gives a covariant functor $\mathbf{n} \mapsto R^{(n)}$, while the structure on a non-empty space Z of being a commutative comonoid object for \times (with diagonal map $Z \to$

 $Z \times Z$ and unique map $Z \to *$) gives a contravariant functor $\mathbf{n} \mapsto Z^n$. Regarding \mathbf{n} as a discrete space, Z^n is the space of maps from \mathbf{n} to Z. The coend

$$CE(Z) := \int^{\mathbf{n} \in \mathbf{F}} R^{(n)} \wedge Z^n_+$$

is then easily identified as the left Kan extension of the functor $R^{(-)}$ along the inclusion of **F** in the category of non-empty spaces. Since the tensor functor $Z \mapsto R \otimes Z$ is a functor from non-empty spaces to orthogonal spectra that restricts to $\mathbf{n} \mapsto R^{(n)}$ on **F**, we get a natural transformation $CE(Z) \to R \otimes Z$, which is an isomorphism when $Z = \mathbf{n}$. Commuting colimits then show that it is an isomorphism when the non-empty space Z is discrete, and the properties of geometric realization (commuting with colimits and finite products) show that it is an isomorphism when Z is the geometric realization of a non-empty simplicial set. (In fact, it is an isomorphism for all non-empty spaces, but we do not need that here.)

Next consider the homotopy coend

$$HCE(Z) := B(R^{(-)}, \mathbf{F}, Z^{-})$$

constructed using the categorical bar construction: it is the geometric realization of the simplicial orthogonal spectrum $HCE_{\bullet}(Z)$ that in degree q is

$$HCE_q(Z) := \bigvee_{n_0,\dots,n_q} R^{(n_q)} \wedge \mathbf{F}(\mathbf{n}_{q-1},\mathbf{n}_q)_+ \wedge \dots \wedge \mathbf{F}(\mathbf{n}_0,\mathbf{n}_1)_+ \wedge Z^{n_0}_+$$

(with degeneracies induced by inserting identity maps in \mathbf{F} and face maps induced by the composition in \mathbf{F} and the action of \mathbf{F} on $R^{(-)}$ and Z^{-}). We have a canonical map from the homotopy coend to the coend

$$HCE(Z) \longrightarrow CE(Z)$$

and the usual extra degeneracy argument shows that it is a homotopy equivalence when $Z = \mathbf{n}$. Because R is cofibrant as a commutative ring orthogonal spectrum, the map is a weak equivalence when Z is discrete: the homotopy groups of both sides can be calculated as the colimit over the non-empty finite subsets. When Z is discrete, $CE(Z) = R \otimes Z$ is the coproduct in commutative ring orthogonal spectra of Z copies of R. Since R is cofibrant, it follows that for any inclusion of discrete spaces $Z \to Z'$, the map $CE(Z) \to CE(Z')$ is a cofibration of cofibrant commutative ring orthogonal spectra. In particular, for any simplicial set Z_{\bullet} , the degeneracies in the simplicial object $CE(Z_{\bullet})$ are cofibrations of cofibrant commutative ring orthogonal spectra and in particular spacewise Hurewicz cofibrations of non-degenerately based spaces. The degeneracies of $HCE(\mathbb{Z}_{\bullet})$ are likewise spacewise Hurewicz cofibrations of non-degenerately based spaces. These technical checks suffice to imply that geometric realization preserves level weak equivalences. Because geometric realization commutes with both HCE(-) and CE(-), we see that the map is a weak equivalence whenever Z is the geometric realization of a simplicial set.

Returning to the proof of the theorem, by the work above it now suffices to show that the map $HCE(X) \to HCE(Y)$ is a weak equivalence, and for this, it suffices to show that the map $HCE_q(X) \to HCE_q(Y)$ is a weak equivalence for all q. To show this, it is enough to see that

$$R^{(m)} \wedge X^n \longrightarrow R^{(m)} \wedge Y^n$$

is a weak equivalence for all m, n > 0. This is an easy consequence of the hypothesis that $X \to Y$ is an *R*-homology isomorphism.

CHAPTER V

Generalized Orbit Desuspension Spectra: Main Properties

The purpose of this chapter is to introduce a class of equivariant orthogonal spectra generalizing the orbit desuspension spectra with similarly nice point-set and homotopical properties, particularly with respect to the multiplicative geometric fixed point functor and smash product. In a departure from our standard notation, in this chapter and the next we write Γ rather than G for a general compact Lie group with the idea that $G \cong \Gamma/K$ for some $K \triangleleft \Gamma$, as this is the case needed for arguments in Section III.3. The first section provides the basic definitions. The next two sections explore how these spectra behave under geometric fixed point functors, smash products, symmetric power functors, and the interactions of these functors.

1. Generalized orbit desuspension spectra

Classically, an orbit desuspension spectra is a Γ -spectrum of the form $\Gamma/H_+ \wedge S^{-V}$ for some orthogonal Γ -representation V and closed subgroup $H < \Gamma$. The object $F_V S^0$ models S^{-V} in the Γ -equivariant stable category, where F_V is the functor of [**22**, II.4.6], left adjoint to evaluation at V. (Indeed, [**15**] denotes $F_V S^0$ as S^{-V} ; see, for example, p. 146.) The description of Γ -equivariant orthogonal spaces as equivariant diagram spaces in [**22**, II§4] allows a concise description of $F_V S^0$ as the representable functor

$$F_V S^0(W) = \mathcal{J}_{\Gamma}(V, W)$$

where \mathcal{J}_{Γ} is the (based Γ -equivariantly enriched) category defined in [22, II.4.1]. Precisely, $\mathcal{J}_{\Gamma}(V, W)$ is the Thom space of the Γ -equivariant vector bundle $\mathfrak{S}^{\perp}\mathcal{I}(V, W)$ whose base space is the Γ -space $\mathcal{I}(V, W)$ of isometric embeddings of V in W and whose fiber over $f: V \to W$ is the orthogonal complement of f(V). The based Γ -spaces $\mathcal{J}_{\Gamma}(V, W)$ give the morphisms in the category \mathcal{J}_{Γ} (with objects the orthogonal Γ -representations), where composition is induced by composition of embeddings and (internal) direct sum of orthogonal complement subspaces. The gist of [22, II§4] is that the category of Γ -spectra is isomorphic to the category of based Γ -space enriched functors from \mathcal{J}_{Γ} to based Γ -spaces.

To generalize orbit desuspension spectra, we replace the vector space V with a vector bundle η and we also take the quotient by a finite group action; to be concise about the input data, we use the following terminology.

TERMINOLOGY 1.1. Let Γ be a compact Lie group and Q a finite group. For the purposes of this paper, we use the terminology (Γ, Q) vector bundle to mean a $(\Gamma \times Q^{\text{op}})$ -equivariant real vector bundle $E \to B$ with $(\Gamma \times Q^{\text{op}})$ -invariant inner product, where B is a Hausdorff (and not just weak Hausdorff) space. We require $E \to B$ to be locally finite dimensional but do not require the dimension to be constant over the different components of the base or the dimension function to be bounded. When needed, we refer to Γ as the group of equivariance and Q as the group for the quotient.

We then have the following construction.

CONSTRUCTION 1.2. Let η be a (Γ, Q) vector bundle $E \to B$. For a vector space W with inner product, let $\mathcal{I}(\eta, W) \to B$ be the locally trivial fiber bundle with fiber over b the space $\mathcal{I}(E_b, W)$ of linear isometries $E_b \to W$. (The homeomorphism class of the fiber may vary with the component of B.) We then have a vector bundle $\mathfrak{S}^{\perp}\mathcal{I}(\eta, W) \to \mathcal{I}(\eta, W)$ where the fiber over an isometry $f: E_b \to W$ is the orthogonal complement of the image. Let $\mathcal{J}(\eta, W)$ be the Thom space of this vector bundle. We have a left Q action on $\mathcal{J}(\eta, W)$ via the right Q action on η and we convert it back to a right Q action using the inverse:

$$\alpha \cdot \sigma := \alpha \circ \sigma_*^{-1}$$

where $\sigma_* : E_b \to E_{b\sigma}$ is the action of σ on E. We can then form the quotient $\mathcal{J}(\eta, W)/Q$. When W has an orthogonal (left) Γ action, $\mathcal{J}(\eta, W)/Q$ obtains a left Γ action via action on W and (inverse) action on η . Letting W vary, $\mathcal{J}(\eta, -)/Q$ obtains the structure of a based Γ -space enriched functor from \mathcal{J}_{Γ} to based Γ -spaces using composition of isometries and (internal) direct sum of vector spaces as in the definition of \mathcal{J}_{Γ} . Let $J\eta$ denote the corresponding Γ -spectrum. We write $J_{(\Gamma,Q)}\eta$ when Γ or Q need to be specified.

EXAMPLE 1.3. Taking Q to be the trivial group, an orthogonal Γ -representation V gives a (Γ, Q) vector bundle $V \to *$, and $J(V \to *)$ is canonically isomorphic to $F_V S^0$. More generally, for $H < \Gamma$ a closed subgroup, $J(\Gamma/H \times V \to \Gamma/H)$ is canonically isomorphic to the orbit desuspension spectrum $F_V(\Gamma/H_+) \cong \Gamma/H_+ \land F_V S^0$. When V is an orthogonal H-representation $\eta \colon \Gamma \times_H V \to \Gamma/H$ is a (Γ, Q) vector bundle, and $J\eta$ is canonically isomorphic to the induced orbit desuspension spectrum $\Gamma_+ \land_H F_V S^0$ (denoted in [15] as $\Gamma_+ \land_H S^{-V}$).

The following proposition gives natural examples with Q non-trivial that include the basic building blocks for the underlying equivariant orthogonal spectra of cells for commutative ring equivariant orthogonal spectra.

PROPOSITION 1.4. For $\eta: E \to B$ a (Γ, Q) vector bundle and $q \ge 1$, let Sym^q η denote the $(\Gamma, \Sigma_q \wr Q)$ vector bundle obtained by taking the qth cartesian power

$$\operatorname{Sym}^q \eta \colon E^q \longrightarrow B^q.$$

Then

$$J_{(\Gamma,\Sigma_q \wr Q)}(\operatorname{Sym}^q \eta) \cong (J_{(\Gamma,Q)}\eta)^{(q)} / \Sigma_q$$

More generally, if Σ is any subgroup of Σ_q , let $\Sigma_{\Sigma}^q \eta$ denote the $(\Gamma, \Sigma \wr Q)$ vector bundle obtained by restring the group for the quotient; then

$$J_{(\Gamma,\Sigma)Q}(\operatorname{Sym}_{\Sigma}^{q}\eta) \cong (J_{(\Gamma,Q)}\eta)^{(q)}/\Sigma$$

PROOF. Consider the map

$$\mathcal{J}(\eta, W_1) \wedge \cdots \wedge \mathcal{J}(\eta, W_1) \longrightarrow \mathcal{J}(\eta^q, W_1 \oplus \cdots \oplus W_q)$$

induced by cartesian product of isometries and complementary subspaces. Applying the universal property of the left Kan extension defining the smash product, we get a $\Sigma_q \wr Q$ equivariant map of Γ -spectra

(1.5)
$$(J_{(\Gamma,1)}\eta)^{(q)} \longrightarrow J_{(\Gamma,1)}(\eta^q)$$

(where 1 denotes the trivial group). Since smash product commutes with colimits in each variable, taking the quotient by $\Sigma \wr Q$ action, we get a map of Γ -spectra

$$(J_{(\Gamma,Q)}\eta)^{(q)}/\Sigma \longrightarrow J_{(\Gamma,\Sigma \wr Q)}(\operatorname{Sym}_{\Sigma}^{q} \eta).$$

To show that it is an isomorphism, it suffices to work non-equivariantly, and indeed to show that the map (1.5) is an isomorphism of non-equivariant spectra. Choosing a cover $\mathcal{U} = \{U_{\alpha}\}$ of B by trivial neighborhoods for η , \mathcal{U}^q gives a cover of B^q by trivial neighborhoods of η^q . Choosing trivializations $\eta_{U_{\alpha}} \cong U_{\alpha} \times \mathbb{R}^{k_{\alpha}}$, for each such neighborhood $U_{\alpha_1} \times \cdots \times U_{\alpha_q}$, we get isomorphisms

$$\mathcal{J}(\eta|_{\alpha_i}) \cong F_{\mathbb{R}^{k\alpha_i}}(U_{\alpha_i})_+$$
$$\mathcal{J}(\eta|_{U_{\alpha_1}} \times \dots \times \eta|_{U_{\alpha_q}}) \cong F_{\mathbb{R}^{k\alpha_1} \times \dots \times \mathbb{R}^{k\alpha_q}}(U_{\alpha_1} \times \dots \times U_{\alpha_q})_+$$

and the composite of these isomorphisms with the map in (1.5)

$$F_{\mathbb{R}^{k_{\alpha_1}}}(U_{\alpha_1})_+ \wedge \dots \wedge F_{\mathbb{R}^{k_{\alpha_q}}}(U_{\alpha_1})_+ \longrightarrow F_{\mathbb{R}^{k_{\alpha_1}} \times \dots \times \mathbb{R}^{k_{\alpha_q}}}(U_{\alpha_1} \times \dots \times U_{\alpha_q})_+$$

is the standard isomorphism. Without loss of generality, we may assume that \mathcal{U} is closed under intersection, and make it a partially ordered set under inclusion. The map in (1.5) is then the colimit over \mathcal{U} of the local maps, and so it is also an isomorphism.

An argument similar to the proof of the previous proposition also proves the following result.

PROPOSITION 1.6. Let η and η' be (Γ, Q) and (Γ, Q') vector bundles, respectively. Then

$$J_{(\Gamma,Q\times Q')}(\eta_1\times\eta_2)\cong J\eta_1\wedge J\eta_2.$$

We have one more basic piece of notation and terminology for generalized orbit desuspension spectra, relevant to the case when we have a fixed isomorphism $G \cong \Gamma/K$ for some normal subgroup $K \triangleleft \Gamma$. In the next section, we will construct an important class of examples of (Γ, Q) vector bundles η with the property that when W is a K-trivial orthogonal Γ -representation, the based Γ -spaces $\mathcal{J}(\eta, W)/Q$ have trivial K action. We use the following terminology.

TERMINOLOGY 1.7. Let $K < \Gamma$ be a closed subgroup. We say that a (Γ, Q) vector bundle η is Γ/K -compatible if the based Γ -space $\mathcal{J}(\eta, W)/Q$ has trivial K action whenever the orthogonal Γ -representation W has trivial K action.

Orthogonal Γ/K -representations are essentially the same thing as K-trivial orthogonal Γ -representations. By slight abuse of notation, we therefore use $\mathcal{J}_{\Gamma/K}$ for the full subcategory of \mathcal{J}_{Γ} consisting of the K-trivial orthogonal Γ -representations. When V and W are K-trivial orthogonal Γ -representations, the Γ -space $\mathcal{J}_{\Gamma}(V, W)$ is a Γ/K -spaces and coincides with $\mathcal{J}_{\Gamma/K}(V, W)$. In particular, when η is Γ/K -compatible, $\mathcal{J}(\eta, W)/Q$ is a based Γ/K space for all K fixed orthogonal Γ -representations W, and the restriction of $\mathcal{J}(\eta, -)/Q$ to these representations has the structure of a based Γ/K -space enriched functor from $\mathcal{J}_{\Gamma/K}$ to based Γ/K -spaces, that is to say, it defines a Γ/K -spectrum.

NOTATION 1.8. For $\eta: E \to B$ a Γ/K -compatible (Γ, Q) vector bundle, we write $J^{\Gamma/K}\eta$ for the corresponding Γ/K -spectrum. We write $J_{(\Gamma,Q)}^{\Gamma/K}\eta$ for $J^{\Gamma/K}\eta$ when Γ or Q need to be indicated.

2. Geometric fixed points and smash products

This sections studies the point-set and homotopical properties of the multiplicative geometric fixed point functor on the generalized orbit desuspension spectra introduced in the previous section. The multiplicative geometric fixed point functor is well-behaved on orbit desuspension spectra, and therefore on those equivariant orthogonal spectra nicely built out of them, that is, the cofibrant equivariant orthogonal spectra in the standard model structure. But it is not known to have well-behaved point-set or homotopical properties on arbitrary spectra. The purpose of this section is to expand the results that hold for orbit desuspension spectra to a wide class of generalized orbit desuspension spectra. This section contains definitions and statements only; proofs appear in Chapter VI.

In this section and the next, we use the following notational convention. It is meant to capture the situation when the Lie group of equivariance G is isomorphic to the quotient Γ/K of a Lie group Γ by a closed normal subgroup; a normal subgroup $\Lambda \lhd \Gamma$ containing K then corresponds to a normal subgroup $L \lhd G$ with the isomorphism $G \cong \Gamma/K$ taking L to Λ/K . In many cases of interest, K is the trivial group and then the Γ/K -compatibility hypotheses (where it appears) is vacuous.

NOTATION 2.1. In this section Γ denotes a compact Lie group, and K, Λ denote closed normal subgroups of Γ with $K < \Lambda$.

We are particularly interested in the generalized orbit desuspension spectra that satisfy the following faithfulness property.

DEFINITION 2.2. We say that a (Γ, Q) vector bundle $\eta: E \to B$ is Q-faithful when for every $b \in B$, the isotropy subgroup $Q_b < Q$ acts faithfully on the fiber E_b .

Because the condition is defined fiberwise and equivariant maps do not decrease stabilizer subgroups, the faithfulness property above is preserved by all the usual kinds of restriction:

PROPOSITION 2.3. Let $\eta: E \to B$ be a Q-faithful (Γ, Q) vector bundle.

- (i) For any Q' < Q, and any homomorphism Γ' → Γ, the restriction of η to a (Γ', Q') vector bundle is Q'-faithful.
- (ii) For any Hausdorff $(\Gamma \times Q^{\text{op}})$ -space B' and any $(\Gamma \times Q^{\text{op}})$ -equivariant map $f: B' \to B$, the pullback (Γ, Q) vector bundle $f^*\eta$ is Q-faithful.

We use the Q-faithfulness hypothesis for most results in this section. In particular, it appears in the following theorem underlying all work on the multiplicative geometric fixed points.

THEOREM 2.4. Let η be a Γ/K -compatible Q-faithful (Γ, Q) vector bundle. Then there exists an isomorphism of Γ/Λ -spectra

$$\Phi^{\Lambda/K} J^{\Gamma/K} \eta \cong J^{\Gamma/\Lambda}(\eta(\Lambda|K))$$

where $\eta(\Lambda|K)$ is the Γ/Λ -compatible (Γ, Q) vector bundle of Definition 2.13 below.

In the case of a non-normal closed subgroup $H < \Gamma$ containing K, K is a normal subgroup of the normalizer $N_{\Gamma}H$ of H in Γ . Then restricting the group of equivariance, η becomes a Q-faithful $(N_{\Gamma}H, Q)$ vector bundle and the previous theorem then asserts an isomorphism of $N_{\Gamma}H/H$ -spectra.

The previous theorem now gives some control over the multiplicative geometric fixed points on generalized orbit desuspension spectra with enough faithfulness hypotheses. For example, the following theorem asserts that iterating multiplicative geometric fixed point functors behaves properly when its hypotheses hold. (It is not known to hold for arbitrary Γ -spectra, cf. [4, 2.6].)

THEOREM 2.5. Let M be a closed normal subgroup of Γ with $\Lambda < M$, and let η be a Γ/K -compatible (Γ, Q) vector bundle. If η and $\eta(\Lambda|K)$ are both Q-faithful, then the canonical map of Γ/M -spectra

$$\Phi^{M/K}(J^{\Gamma/K}\eta) \longrightarrow \Phi^{M/\Lambda}(\Phi^{\Lambda/K}(J^{\Gamma/K}\eta))$$

is an isomorphism.

A technical issue throughout [4] is that although the multiplicative geometric fixed point functor is lax symmetric monoidal, it is not known to be strong symmetric monoidal, and a difficult technical result in *ibid*. is that it is strong symmetric monoidal when one variable is an orbit desuspension spectrum. The following theorem generalizes this to Q-faithful generalized orbit desuspension spectra.

THEOREM 2.6. Suppose η is a Q-faithful Γ/K -compatible (Γ, Q) vector bundle where the base B admits the structure of a $(\Gamma \times Q^{\text{op}})$ -equivariant cell complex. Then for any Γ/K -spectrum X, the canonical map of orthogonal spectra

$$\Phi^{\Lambda/K}(J^{\Gamma/K}\eta) \wedge \Phi^{\Lambda/K}X \longrightarrow \Phi^{\Lambda/K}(J^{\Gamma/K}\eta \wedge X)$$

is an isomorphism.

The previous theorems study point-set properties. The next three theorems study the homotopical properties. The first of these gives hypotheses that ensure that the multiplicative geometric fixed point functor represents the derived geometric fixed point functor. We state it for the normal subgroup Λ , but as in the remarks following Theorem 2.4, restricting from Γ to the normalizer gives the desired theorem for a general closed subgroup.

THEOREM 2.7. Let η be a Γ/K -compatible (Γ, Q) vector bundle. If η and $\eta(\Lambda|K)$ are both Q-faithful, then the canonical map in the genuine Γ/Λ -equivariant stable category $L\Phi^{\Lambda/K}(J^{\Gamma/K}\eta) \to \Phi^{\Lambda/K}(J^{\Gamma/K}\eta)$ is an isomorphism.

The final two theorems do not involve the multiplicative geometric fixed point functor but share much of the same work as the previous results. The first is about derived functors of symmetric powers.

THEOREM 2.8. Suppose η is a Q-faithful Γ/K -compatible (Γ, Q) vector bundle with each fiber positive dimensional and $X \to J^{\Gamma/K} \eta$ is a cofibrant approximation in the positive complete model structure on G/K-spectra. Then for any $k \ge 1$, and any subgroup $\Sigma < \Sigma_k$, the map $X^{(k)}/\Sigma \to (J^{\Gamma/K}\eta)^{(k)}/\Sigma$ is a weak equivalence.

We also have the following flatness result.

THEOREM 2.9. Suppose η is a Q-faithful Γ/K -compatible (Γ, Q) vector bundle and assume the base B admits the structure of a $(\Gamma \times Q^{\text{op}})$ -equivariant cell complex. Then $J^{\Gamma/K}\eta$ is flat for the smash product in Γ/K -spectra.

The rest of the section is devoted to the construction and basic properties of $\eta(\Lambda|K)$. We start with the easier construction of $\eta(\Lambda)$ which corresponds to the case when K is the trivial subgroup.

DEFINITION 2.10. Let Λ be a closed normal subgroup of Γ and let η be a Q-faithful (Γ, Q) vector bundle $E \to B$. For a continuous homomorphism $\sigma \colon \Lambda \to Q$, we get a σ -twisted Λ action on $E \to B$ from the isomorphism of Λ to the graph subgroup of $\Gamma \times Q^{\text{op}}$ associated to the homomorphism $\sigma((-)^{-1})$. Concretely,

$$\lambda \cdot_{\sigma} e := \lambda e \sigma(\lambda^{-1})$$

We write $\eta^{\Lambda,\sigma} \colon E^{\Lambda,\sigma} \to B^{\Lambda,\sigma}$ for the fixed points under this action. Define the map

$$\eta(\Lambda) \colon E(\Lambda) \longrightarrow B(\Lambda)$$

as

$$\coprod_{\sigma \colon \Lambda \to Q} (\eta^{\Lambda, \sigma} \colon E^{\Lambda, \sigma} \longrightarrow B^{\Lambda, \sigma}).$$

We assert that the map $\eta(\Lambda)$ in the previous definition has a canonical Γ/Λ compatible (Γ, Q) vector bundle structure, with the $\Gamma \times Q^{\text{op}}$ action inherited from η and the conjugation action on $\text{Hom}(\Lambda, Q)$.

For the vector bundle structure, we use the (trivial) fact that for any $\sigma \colon \Lambda \to Q$, the σ -twisted Λ action on $\eta|_{B^{\Lambda,\sigma}}$ is fiberwise. Then for any $b \in B^{\Lambda,\sigma}$, there is an open set $b \in U \subset B^{\Lambda,\sigma}$ on which the restriction $E|_U \to U$ is Λ -equivariantly isomorphic (with the σ -twisted Λ action) to the product vector bundle $U \times V \to U$ for the orthogonal Λ -representation $V = E_b$. (This follows for example by character theory.) Taking Λ fixed points, we get the local trivialization for $\eta(\Lambda)$.

We construct the $\Gamma \times Q^{\text{op}}$ action as follows. An element x of $E(\Lambda)$ consists of a homomorphism $\sigma \colon \Lambda \to Q$ together with an element $e \in E^{\Lambda,\sigma}$; we write $x = (\sigma, e)$. For $\gamma \in \Gamma$,

(2.11)
$$\gamma(\sigma, e) = (\sigma^{\gamma}, \gamma e)$$

where $\sigma^{\gamma} \colon \Lambda \to Q$ is the homomorphism

$$\sigma^{\gamma} \colon \lambda \mapsto \sigma(\gamma^{-1}\lambda\gamma).$$

To see that this specifies a left Γ action, we need to check that $\gamma e \in E^{\Lambda,\sigma^{\gamma}}$. For $\lambda \in \Lambda$,

$$\lambda \cdot_{\sigma^{\gamma}} \gamma e := \lambda \gamma e \sigma^{\gamma}(\lambda^{-1}) := \lambda \gamma e \sigma(\gamma^{-1}\lambda^{-1}\gamma) = \gamma \gamma^{-1}\lambda \gamma e \sigma(\gamma^{-1}\lambda^{-1}\gamma) = \gamma e$$

with the last equality holding because $e \in E^{\Lambda,\sigma}$. For the right Q action, given $s \in Q$,

(2.12)
$$(\sigma, e)s = (\sigma_s, es)$$

where

$$\sigma_s(\lambda) = s^{-1}\sigma(\lambda)s.$$

To see that this specifies a right Q action, we need to check that $es \in E^{\Lambda,\sigma_s}$. For $\lambda \in \Lambda$,

$$\lambda \cdot_{\sigma_s} es := \lambda es \sigma_s(\lambda^{-1}) := \lambda es s^{-1} \sigma(\lambda^{-1}) s = \lambda e \sigma(\lambda^{-1}) s = es$$

with the last equality holding because $e \in E^{\Lambda,\sigma}$. It is clear from the formulas that the left Γ action and right Q action commute.

Finally, we need to check that $\eta(\Lambda)$ is Γ/Λ -compatible. It suffices to show that if W is a Λ -trivial orthogonal Γ -representation, the Γ space $\mathfrak{S}^{\perp}\mathcal{J}(\eta(\Lambda), W)/Q$ has trivial Λ action. An element of $\mathfrak{S}^{\perp}\mathcal{J}(\eta(\Lambda), W)$ is specified by a homomorphism $\sigma \colon \Lambda \to Q$, a point $b \in B^{\Lambda,\sigma}$, an isometry $f \colon E^{\Lambda,\sigma}|_b \to W$, and a point w in the complement of its image. We write this as a tuple (σ, b, f, w) , and we write $[\sigma, b, f, w]$ for the image in the quotient by Q. For $\lambda \in \Lambda$,

$$\lambda(\sigma, b, f, w) = (\sigma^{\lambda}, \lambda b, f^{\lambda}, \lambda w) = (\sigma^{\lambda}, \lambda b, f \circ \lambda_{*}^{-1}, w)$$
$$= (\sigma^{\lambda}, b\sigma(\lambda), f \circ \sigma(\lambda)_{*}^{-1}, w)$$

where the first equality follows from the fact that W is Λ -trivial and the second equality follows from the fact that b and the domain of f lie in the σ -twisted Λ fixed points. Since

$$\sigma^{\lambda}(-) = \sigma(\lambda^{-1}(-)\lambda) = \sigma(\lambda)^{-1}\sigma(-)\sigma(\lambda) = \sigma_{\sigma(\lambda)}(-),$$

we see that

$$\lambda(\sigma, b, f, w) = (\sigma, b, f, w)\sigma(\lambda)$$

and so $[\sigma, b, f, w]$ is fixed by λ .

We now construct $\eta(\Lambda|K)$ by restricting $\eta(\Lambda)$ to a subset of its base space.

DEFINITION 2.13. Let $K, \Lambda \lhd \Gamma$, with $K < \Lambda$ and let η be a Γ/K -compatible (Γ, Q) vector bundle. Given $\sigma : \Lambda \to Q$, the σ -twisted Λ action on the restricted bundle $\eta|_{B^{\Lambda,\sigma}}$ is fiberwise, and for each component of $B^{\Lambda,\sigma}$, the isomorphism class of the representation of the fiber is constant. Let $B^{\Lambda,\sigma}\{K\}$ denote the subspace of $B^{\Lambda,\sigma}$ consisting of those components where the fiberwise $\sigma|_K$ -twisted K action is trivial. Write

$$\eta^{\Lambda,\sigma}\{K\} := (\eta|_{B^{\Lambda,\sigma}\{K\}})^{\Lambda,\sigma}$$

for the σ -twisted Λ fixed points of the restriction of η to $B^{\Lambda,\sigma}\{K\}$, and note that $\eta^{\Lambda,\sigma}\{K\}$ is also the restriction of $\eta^{\Lambda,\sigma}$ to the components $B^{\Lambda,\sigma}\{K\} \subset B^{\Lambda,\sigma}$ of its base. Write

$$\eta(\Lambda|K) := \coprod_{\sigma \colon \Lambda \to Q} \eta^{\Lambda,\sigma}\{K\}.$$

By construction, $\eta(\Lambda|K)$ is the restriction of $\eta(\Lambda)$ to a certain $(\Gamma \times Q^{\text{op}})$ -stable subset of components in its base. We give $\eta(\Lambda)\{K\}$ the Γ/Λ -compatible (Γ, Q) vector bundle structure inherited from $\eta(\Lambda)$.

As a special case of Proposition 2.3, we point out the following observation.

PROPOSITION 2.14. Let η be a Γ/K -compatible (Γ, Q) vector bundle. If $\eta(\Lambda)$ is Q-faithful, then so is $\eta(\Lambda|K)$

3. Geometric fixed points of symmetric powers

This section again follows the notational conventions of Notation 2.1. Its purpose is to provide results on and tools for working with the point-set geometric fixed points of symmetric powers of generalized orbit suspension spectra $J^{\Gamma/K}\eta$ under certain faithfulness hypotheses we state below. The main theorem, Theorem 3.1, and its proof give a complete description of $\Phi^{\Lambda/K}((J^{\Gamma/K}\eta)^{(q)}/\Sigma_q)$ when $\operatorname{Sym}^q \eta$ is $(\Sigma_q \wr Q)$ -faithful. As discussed in Proposition 1.4 and Theorem 2.4, under

this faithfulness hypotheses, $\Phi^{\Lambda/K}((J^{\Gamma/K}\eta)^{(q)}/\Sigma_q)$ is itself the generalized orbit desuspension spectrum $J^{\Gamma/\Lambda}((\operatorname{Sym}^q \eta)(\Lambda|K))$ and we describe a criterion (Proposition 3.6, or more generally, Theorem 3.5) for $(\operatorname{Sym}^q \eta)(\Lambda)$ to be $(\Sigma_q \wr Q)$ -faithful. We need these results for the bundles considered in the arguments in Section III.3. Both Theorem 3.1 and 3.5 are proved in later sections.

Let \mathbb{P} denote the free functor from equivariant orthogonal spectra to equivariant commutative ring orthogonal spectra; it is the wedge sum of symmetric powers

$$\mathbb{P}X = \bigvee_{n \ge 0} X^{(n)} / \Sigma_n$$

where we understand $X^{(0)}/\Sigma_0$ as the sphere spectrum S. We prove the following theorem in Section VI.5. Along the way to the proof, we give more details on the individual symmetric powers including an intrinsic description of the spectra X_q in terms of η and Λ .

THEOREM 3.1. Let η be a Γ/K -compatible (Γ, Q) vector bundle and assume that Sym^q η is $(\Sigma_q \wr Q)$ -faithful for every $q \ge 1$. Then there exist Γ/Λ -spectra $X_q(\eta)$, $1 \le q \le \#(\pi_0\Lambda)$, such that $X_q(\eta)$ is a wedge summand of $\Phi^{\Lambda/K}((J^{\Gamma/K}\eta)^{(q)}/\Sigma_q)$ and the inclusions induce an isomorphism

$$\bigwedge_{q=1}^{\#\pi_0\Lambda} \mathbb{P}(X_q(\eta)) \cong \mathbb{P}\left(\bigvee_q X_q(\eta)\right) \longrightarrow \Phi^{\Lambda/K}(\mathbb{P}(J^{\Gamma/K}\eta))$$

of commutative ring Γ/Λ -spectra. Moreover, if D is a (non-equivariant) Hausdorff space, then $X_q(\eta \times D) \cong X_q(\eta) \wedge D^q_+$, naturally in D.

The hypothesis that each $\operatorname{Sym}^q \eta$ is $(\Sigma_q \wr Q)$ -faithful is not usually much more stringent than the hypothesis that η is Q-faithful. An easy argument gives the following observation.

PROPOSITION 3.2. If $\eta: E \to B$ is Q-faithful and either E_b is positive dimensional for all $b \in B$ or Q is not the trivial group, then $\operatorname{Sym}^q \eta$ is $(\Sigma_q \wr Q)$ -faithful for all $q \ge 1$.

Theorem 2.7 allows us to deduce homotopical conclusions from Theorem 3.1 when the $(\Gamma, \Sigma_q \wr Q)$ vector bundle $(\operatorname{Sym}^q \eta)(\Lambda)$ (or its base restriction $(\operatorname{Sym}^q \eta)(\Lambda|K))$ is $(\Sigma_q \wr Q)$ -faithful for all $q \ge 1$. We use the following terminology.

DEFINITION 3.3. A (Γ, Q) vector bundle is inheritably faithful for Φ^{Λ} means that for every $q \geq 1$, $(\text{Sym}^q \eta)(\Lambda)$ is $(\Sigma_q \wr Q)$ -faithful. (See Proposition 1.4 and Definitions 2.2, 2.10.)

The previous definition asks for a minimal hypothesis to check but leads to the following wider conclusion as an application of Proposition 2.3.

PROPOSITION 3.4. Let η be a (Γ, Q) vector bundle and assume that $(\operatorname{Sym}^q \eta)(\Lambda)$ is $(\Sigma_q \wr Q)$ -faithful. Then for any $\Sigma < \Sigma_q$, $(\operatorname{Sym}_{\Sigma}^q \eta)(\Lambda)$ is $(\Sigma \wr Q)$ -faithful. Moreover if η is Γ/K -compatible then for any $\Sigma < \Sigma_q$, $(\operatorname{Sym}_{\Sigma}^q \eta)(\Lambda|K)$ is $(\Sigma \wr Q)$ -faithful.

To employ this theory, we offer the following result for identifying inheritably faithful vector bundles. In the statement Λ_0 denotes the identity component of Λ . We prove the following theorem in Section VI.6.

THEOREM 3.5. Let $\eta: E \to B$ be a (Γ, Q) vector bundle such that for every closed subgroup $\Lambda_0 < H < \Lambda$, every $\sigma: H \to Q$, and every $b \in B^{H,\sigma}$:

- (i) $(E_b)^{H,\sigma}$ is positive dimensional,
- (ii) Given $\lambda \in \Lambda H$, $s \in Q$ such that $\lambda b = bs$, there exists $v \in (E_b)^{H,\sigma}$ for which $\lambda v \neq vs$, and
- (iii) Given $s \in Q$, not the identity element, such that bs = b and s commutes with $\sigma(h)$ for all $h \in H$, there exists $v \in (E_b)^{H,\sigma}$ for which $v \neq vs$.

Then η is inheritably faithful for Φ^{Λ} .

We note that condition (iii) is equivalent to $\eta(H)$ being Q-faithful for all $\Lambda_0 < H < \Lambda$. The conditions given in the previous theorem are technical; it is helpful to have concrete special cases that provide examples for when the conditions hold. We provide these in the following proposition. We note that in the first set of examples, the hypothesis does not involve Λ , and so it gives examples of (Γ, Q) vector bundles that are inheritably faithful for Φ^H for all subgroups H (even non-normal subgroups after restricting Γ to the normalizer of H).

PROPOSITION 3.6. If $\eta: E \to B$ is a (Γ, Q) vector bundle that satisfies either of the following hypotheses, then η satisfies the conditions of Theorem 3.5 and is therefore inheritably faithful for Φ^{Λ} .

- (a) For every $b \in B$, the isotropy subgroup $(\Gamma \times Q^{\text{op}})_b$ is finite and the orthogonal $(\Gamma \times Q^{\text{op}})_b$ -representation E_b contains a copy of the regular representation.
- (b) Q is the trivial group and for every b ∈ B whose isotropy subgroup Γ_b < Γ contains Λ₀, the (restricted) orthogonal (Γ_b ∩ Λ)-representation E_b contains a copy of the regular representation of π₀(Γ_b ∩ Λ).

PROOF. Let $\Lambda_0 < H < \Lambda$, $\sigma: H \to Q$, and $b \in B^{H,\sigma}$; we check conditions (i), (ii), and (iii) of Theorem 3.5.

For hypothesis (a), since $b \in B^{H,\sigma}$, the subgroup $H_{\sigma} = \{(h, \sigma(h^{-1}))\}$ of $\Gamma \times Q^{\text{op}}$ is contained in the isotropy subgroup $L := (\Gamma \times Q^{\text{op}})_b$. The H_{σ} fixed points of the regular *L*-representation are positive dimensional, so (i) holds. Now suppose that $\lambda b = b\tau$ for $\tau \in Q$ and either $\lambda \in \Lambda - H$ or $\lambda = 1$ with $\tau \neq 1$. In either case, $(\lambda, \tau^{-1}) \in L$ but not in H_{σ} so (λ, τ^{-1}) acts non-trivially on the H_{σ} fixed points of the regular *L*-representation.

For hypothesis (b), since H fixes b, we have $H < \Gamma_b$ and in particular $\Lambda_0 < \Gamma_b$. Condition (iii) holds vacuously and condition (i) holds because the fixed points of the regular representation form a positive dimensional subspace. For condition (ii), let $\lambda \in \Lambda - H$ with $\lambda b = b$; then $\lambda \in \Gamma_b \cap (\Lambda - H)$ and the image of λ in $\pi_0(\Gamma_b \cap \Lambda)$ is disjoint from $\pi_0 H$. It follows that λ acts non-trivially on the H fixed points of the regular $\pi_0(\Gamma_b \cap \Lambda)$ -representation, establishing condition (ii).

CHAPTER VI

Generalized Orbit Desuspension Spectra: Proofs

This chapter provides the proofs for the main theorems of Chapter V.

1. The quotient fixed point lemma and Theorems V.2.4, V.2.5, and V.2.7

This section proves Theorems V.2.4, V.2.5, and V.2.7. We follow the notational convention of Notation V.2.1: Γ is a compact Lie group with closed normal subgroups K, Λ satisfying $K < \Lambda$. The proofs have in common a lemma about fixed points of the quotients of the isometry bundles intrinsic to the construction of generalized orbit desuspension spectra for Q-faithful (Γ, Q) vector bundles. This lemma in turn is based on the following (easy) observation.

PROPOSITION 1.1. If η is a Q-faithful (Γ, Q) vector bundle, then for any orthogonal Γ -representation W, the action of Q on $\mathcal{I}(\eta, W)$ and $\mathfrak{S}^{\perp}\mathcal{I}(\eta, W)$ is free.

The following is the main technical result we need in the proofs of the theorems. As in the definition of the bundle $\eta(\Lambda)$ (Definition V.2.10), we write $(-)^{\Lambda,\sigma}$ for the fixed points of the σ -twisted Λ action for a continuous homomorphism $\sigma \colon \Lambda \to Q$.

LEMMA 1.2. Let $\eta: E \to B$ be a Q-faithful (Γ, Q) vector bundle, and let W be an orthogonal Γ -representation. Then we have a canonical isomorphism

$$(\mathcal{J}(\eta, W)/Q)^{\Lambda} \cong \left(\bigvee_{\sigma \colon \Lambda \to Q} \mathcal{J}(\eta, W)^{\Lambda, \sigma}\right)/Q$$

where Q acts by conjugation on the set of homomorphisms, acts as usual on η , and acts trivially on W. Letting Γ act on the right hand side by conjugation on the set of homomorphisms and as usual on η and W, the isomorphism is Γ - and therefore Γ/Λ -equivariant. Furthermore, for each summand on the right hand side (that is, for each continuous homomorphism $\sigma \colon \Lambda \to Q$):

- (i) $(\mathfrak{S}^{\perp}\mathcal{I}(\eta, W))^{\Lambda,\sigma} \to \mathcal{I}(\eta, W)^{\Lambda,\sigma}$ is a vector bundle with Thom space $\begin{array}{l} \mathcal{J}(\eta, W)^{\Lambda, \sigma}. \\ \text{(ii)} \quad \mathcal{I}(\eta, W)^{\Lambda, \sigma} \to B^{\Lambda, \sigma} \text{ is a locally trivial fiber bundle with fiber } \mathcal{I}(E_b, W)^{\Lambda, \sigma}. \end{array}$
- (The fiber may vary with the component of $B^{\Lambda,\sigma}$.)

PROOF. The numbered statements follow from the fact that the σ -twisted Λ action on

$$\mathcal{I}(\eta|_{B^{\Lambda,\sigma}}, W) \longrightarrow B^{\Lambda,\sigma}$$

is fiberwise, and locally in $b \in B^{\Lambda,\sigma}$, $\eta|_{B^{\Lambda,\sigma}}$ is a product bundle $U \times V \to U$ for an orthogonal Λ -representation V (that may vary in the component of $B^{\Lambda,\sigma}$).

To study $(\mathcal{J}(\eta, W)/Q)^{\Lambda}$, we use its identification as the Thom space of the vector bundle

$$(\Im^{\perp}\mathcal{I}(\eta, W)/Q)^{\Lambda} \longrightarrow (\mathcal{I}(\eta, W)/Q)^{\Lambda}$$

Let $q: \mathcal{I}(\eta, W) \to \mathcal{I}(\eta, W)/Q$ and consider

$$q^{-1}((\mathcal{I}(\eta, W)/Q)^{\Lambda}) \subset \mathcal{I}(\eta, W)$$

An element on the left consists of a point $b \in B$ and an isometry $f: E_b \to W$ such that for all $\lambda \in \Lambda$, there exists $\sigma(\lambda) \in Q$ such that $\lambda b = b\sigma(\lambda)$ and $\lambda \cdot f = f \circ \sigma(\lambda)^{-1}_*$ as isometries $E_{\lambda b} = E_{b\sigma(\lambda)} \to W$. Since Q acts freely on $\mathcal{I}(\eta, W)$ (Proposition 1.1), $\sigma(\lambda)$ is unique and $\lambda \mapsto \sigma(\lambda)$ defines a continuous homomorphism $\sigma: \Lambda \to Q$. In other words, f is a σ -twisted Λ fixed point for a unique $\sigma: \Lambda \to Q$ (where we understand the Q action on W to be trivial). Likewise, for any $\sigma: \Lambda \to Q$, the σ -twisted Λ fixed points of $\mathcal{I}(\eta, W)$ lie in $q^{-1}((\mathcal{I}(\eta, W)/Q)^{\Lambda})$. This shows

$$(\mathcal{I}(\eta, W)/Q)^{\Lambda} = \left(\coprod_{\sigma \colon \Lambda \to Q} \mathcal{I}(\eta, W)^{\Lambda, \sigma}\right)/Q$$

and we can similarly identify $(\mathfrak{T}(\eta, W)/Q)^{\Lambda}$ as

$$\left(\coprod_{\sigma \colon \Lambda \to Q} (\mathfrak{S}^{\perp} \mathcal{I}(\eta, W))^{\Lambda, \sigma}\right) / Q.$$

The identification of $\mathcal{J}(\eta, W)/Q$ in the statement is now clear. The proof that the isomorphism is Γ -equivariant is a straightforward check of the action on the righthand side.

Each of the theorems we are proving in this section concerns a multiplicative geometric fixed point functor of the form Φ^N from the category of *G*-spectra to the category of *G*/*N*-spectra, where *N* is a normal subgroup of a compact Lie group *G*; in the theorems, we have in various cases (in the respective notation there), $G = \Gamma/K$ and $H = \Lambda/K$, $G = \Gamma/K$ and H = M/K, or $G = \Gamma/\Lambda$ and $H = M/\Lambda$. The arguments require some background in the construction of these functors. We recall that Φ^N is a composite of a functor Fix^N and a left Kan extension as follows.

Let G be a compact Lie group with $N \triangleleft G$ normal subgroup. As in previous sections, we take the perspective that G-spectra are G-equivariant \mathcal{J}_G -spaces: based G-space enriched functors from the category \mathcal{J}_G (reviewed in Section V.1) to based G-spaces. Following [22, V§4] (with slightly altered notation), let \mathcal{J}_G^N be the category enriched in based G/N-spaces obtained as the N fixed points of \mathcal{J}_G . It has the same objects (namely, the orthogonal G-representations) and for objects V, W, the based G/N-space of morphisms $\mathcal{J}_G^N(V, W) = (\mathcal{J}_G(V, W))^N$. We define the category of G/N-equivariant \mathcal{J}_G^N -spaces to be the category of based G/N-space enriched functors from \mathcal{J}_G^N to based G/N-spaces. The functor Fix^N from G-spectra to G/N-equivariant \mathcal{J}_G^N -spaces is defined by

$$(\operatorname{Fix}^N X)(W) := (X(W))^N$$

for W an orthogonal G-representation.

For the left Kan extension, we use the based G/N-space enriched functor $\phi: \mathcal{J}_G^N \to \mathcal{J}_{G/N}$ that on objects takes V to its N fixed points V^N and on morphisms restricts to fixed points. Precisely, the morphism space $\mathcal{J}_G^N(V,W)$ is the Thom space of the vector bundle

$$(\mathfrak{S}^{\perp}\mathcal{I}(V,W))^N \longrightarrow (\mathcal{I}(V,W))^N.$$

where an element in the total space consists of a N-equivariant isometry $V \to W$ together with an element of W^N in the orthogonal complement of the image. On morphisms ϕ is the map on Thom spaces induced by the map of vector bundles

$$\left((\Im^{\perp}\mathcal{I}(V,W))^{N} \to (\mathcal{I}(V,W))^{N}\right) \longrightarrow \left(\Im^{\perp}\mathcal{I}(V^{N},W^{N}) \longrightarrow \mathcal{I}(V^{N},W^{N})\right)$$

that takes the N-equivariant isometry $f: V \to W$ to the restriction of the map on N fixed points $f^N: V^N \to W^N$ and maps $f^{\perp} \cap W^N$ to $(f|_{V^N})^{\perp} \subset W^N$ by the identity. The based G/N-space enriched left Kan extension along ϕ is then a based G/N-space enriched functor from G/N-equivariant \mathcal{J}_G^N -spaces to G/N-spectra.

G/N-space enriched functor from G/N-equivariant \mathcal{J}_G^N -spaces to G/N-spectra. The point-set multiplicative geometric fixed point functor Φ^N from G-spectra to G/N-spectra is the composite of Fix^N from G-spectra to G/N-equivariant \mathcal{J}_G^N -spaces and based G/N-space enriched left Kan extension along ϕ .

Returning to the context of Theorems V.2.4, V.2.5, and V.2.7, the functor $\Phi^{\Lambda/K}$ from Γ -spectra to Γ/Λ -spectra is the instance of this functor for $\Lambda/K \lhd \Gamma/\Lambda$ combined with the canonical isomorphism of compact Lie groups $\Gamma/\Lambda \cong (\Gamma/K)/(\Lambda/K)$. As in Section V.1, we switch between orthogonal Γ/K -representations and K-trivial orthogonal Γ -representations without notation and comment. We are now ready for the proof of Theorem V.2.4.

PROOF OF THEOREM V.2.4. The left Kan extension along ϕ is left adjoint to the functor u_{ϕ} given by restriction along the functor ϕ ,

$$u_{\phi}X = X \circ \phi$$

In this context, u_{ϕ} is a based Γ/Λ -space enriched functor from Γ/Λ -spectra to Γ/Λ -equivariant $\mathcal{J}_{\Gamma/K}^{\Lambda/K}$ -spaces. Thus, to construct a map of Γ/Λ -spectra

$$\Phi^{\Lambda/K} J^{\Gamma/K} \eta \longrightarrow J^{\Gamma/\Lambda}(\eta(\Lambda|K))$$

it suffices to construct a map of Γ/Λ -equivariant $\mathcal{J}_{\Gamma/K}^{\Lambda/K}$ -spaces

$$\operatorname{Fix}^{\Lambda/K} J^{\Gamma/K} \eta \longrightarrow (J^{\Gamma/\Lambda} \eta(\Lambda|K)) \circ \phi.$$

For W a $K\text{-trivial orthogonal }\Gamma\text{-representation}, we need to construct a <math display="inline">\Gamma/\Lambda\text{-equivariant}$ map

(1.3)
$$(\mathcal{J}(\eta, W)/Q)^{\Lambda} \longrightarrow \mathcal{J}(\eta(\Lambda|K), W^{\Lambda})/Q$$

natural in W in $\mathcal{J}_{\Gamma/K}^{\Lambda/K}$.

Before constructing this map, we study $(\mathcal{J}(\eta, W)/Q)^{\Lambda}$ further. From the lemma, $(\mathcal{J}(\eta, W)/Q)^{\Lambda}$ is the Thom space of the vector bundle

$$\left(\coprod_{\sigma: \Lambda \to Q} (\mathfrak{S}^{\perp} \mathcal{I}(\eta, W))^{\Lambda, \sigma} \longrightarrow \mathcal{I}(\eta, W)^{\Lambda, \sigma}\right) / Q.$$

We note that on base spaces

$$\mathcal{I}(\eta, W)^{\Lambda, \sigma} = \mathcal{I}(\eta|_{B^{\Lambda, \sigma}}, W)^{\Lambda, \sigma} = \mathcal{I}(\eta|_{B^{\Lambda, \sigma}\{K\}}, W)^{\Lambda, \sigma}$$

where $\eta: E \to B$ and $B^{\Lambda,\sigma}\{K\}$ is as in Definition V.2.13. This is because for $b \in B$, a σ -twisted Λ fixed point, the σ -twisted Λ action on E_b does not admit any equivariant isometries into a K-trivial representation unless $b \in B^{\Lambda,\sigma}\{K\}$. Thus, $(\mathcal{J}(\eta, W)/Q)^{\Lambda}$ is the Thom space of the vector bundle

$$\prod_{\Lambda \to Q} (\mathfrak{S}^{\perp} \mathcal{I}(\eta|_{B^{\Lambda,\sigma}\{K\}}, W))^{\Lambda,\sigma} \longrightarrow \mathcal{I}(\eta|_{B^{\Lambda,\sigma}\{K\}}, W))^{\Lambda,\sigma} \bigg)/Q.$$

We construct (1.3) as a variant of ϕ as follows. We note that in the vector bundle displayed above, on each fiber, the vector space is a subspace of W^{Λ} ; specifically, over a point $b, f: E_b \to W$, it is the orthogonal complement of $f(E_b^{\Lambda,\sigma})$ in W^{Λ} . We define the map (1.3) to be the induced map on Thom spaces that takes $f: E_b \to W$ to $f|_{E_b^{\Lambda,\sigma}}$ and is the identity on the fiber vector spaces (viewed as subspaces of W^{Λ}). The resulting map is clearly natural in W in $\mathcal{J}_{\Gamma/K}^{\Lambda/K}$. To see that it is Γ/Λ equivariant, it suffices to check that it is Γ -equivariant, and this follows from the formula in (V.2.11) for the Γ action on $\eta(\Lambda)$: acting by $\gamma \in \Gamma$ on a σ -twisted fixed point $f: E_b \to W$ of $\mathcal{I}(\eta, W)$ takes it to the σ^{γ} -twisted fixed point $\gamma \cdot f = \gamma \circ f \circ \gamma^{-1}$.

Having constructed the comparison map of $\Gamma/\Lambda\text{-spectra}$

$$\Phi^{\Lambda/K} J^{\Gamma/K} \eta \longrightarrow J^{\Gamma/\Lambda}(\eta(\Lambda|K)),$$

it suffices to show that it is an isomorphism non-equivariantly. Consider the categories of non-equivariant $\mathcal{J}_{\Gamma/K}^{\Lambda/K}$ -spaces and $\mathcal{J}_{\Gamma/\Lambda}$ -spaces defined to be categories of based space enriched functors from $\mathcal{J}_{\Gamma/K}^{\Lambda/K}$ or $\mathcal{J}_{\Gamma/\Lambda}$ to based spaces, respectively. (The category of non-equivariant $\mathcal{J}_{\Gamma/\Lambda}$ -spaces is equivalent to the category of nonequivariant spectra by the functor that restricts to the full subcategory of trivial representations.) The functor ϕ induces a restriction functor from non-equivariant $\mathcal{J}_{\Gamma/\Lambda}$ -spaces to non-equivariant $\mathcal{J}_{\Gamma/K}^{\Lambda/K}$ -spaces which has a left adjoint given by left Kan extension. For formal reasons, forgetting equivariance commutes with left Kan extension. Thus, forgetting Γ/Λ -equivariance, we can study $\Phi^{\Lambda/K} J^{\Gamma/K} \eta$ by applying the non-equivariant left Kan extension along ϕ to the underlying nonequivariant $\mathcal{J}_{\Gamma/K}^{\Lambda/K}$ -space of Fix^{$\Lambda/K} J^{\Gamma/K} \eta$. We denote the left Kan extension along ϕ as P_{ϕ} .</sup>

Non-equivariantly, Fix^{Λ/K} $J^{\Gamma/K}\eta$ and $J^{\Gamma/\Lambda}(\eta(\Lambda|K))$ decompose as the coproduct over $\sigma \colon \Lambda \to Q$ of sub- $\mathcal{J}_{\Gamma/K}^{\Lambda/K}$ -spaces and sub- $\mathcal{J}_{\Gamma/\Lambda}$ -spaces (respectively), and the comparison map preserves this decomposition. As a left adjoint, P_{ϕ} preserves coproducts, so it suffices to study each summand separately, and we now fix $\sigma \colon \Lambda \to Q$. Let $B' = B^{\Lambda,\sigma}\{K\}$, and let $\eta' \colon E' \to B'$ be the vector bundle obtained from η by restriction. When we regard η' as a Λ -equivariant vector bundle with the σ -twisted Λ action, the Λ action is K-trivial, making $\eta' \neq \Lambda/K$ -equivariant vector bundle; moreover, the vector bundle $\eta^{\Lambda,\sigma}$ is precisely its Λ/K fixed points. For every $b \in B'$, there exists an open set U around b where the restriction $\eta'|_U$ is Λ/K -equivariant vector bundle. Let \mathcal{U} be the partially ordered set of open subsets U of B' on which η' admits a Λ/K -equivariant trivialization. Both Fix^{$\Lambda/K} J^{<math>\Gamma/K}\eta$ and $J^{<math>\Gamma/\Lambda}(\eta(\Lambda|K))$ decompose as a colimit over \mathcal{U} , compatibly with the comparison map, and so it suffices to show that for $U \in \mathcal{U}$, the restriction of the comparison map</sup></sup>

$$P_{\phi}(\mathcal{J}(U \times V \to U, -)^{\Lambda/K}) \cong P_{\phi}(\mathcal{J}(\eta'|_{U}, -)^{\Lambda/K})$$
$$\longrightarrow \mathcal{J}(\eta^{\Lambda, \sigma}\{K\}|_{U}, -) \cong \mathcal{J}(U \times V^{\Lambda/K} \to U, -)$$

is an isomorphism of non-equivariant $\mathcal{J}_{\Gamma/\Lambda}$ -spaces, where $V = E_b$ denotes the fiber Λ/K -representation of η' at an arbitrarily chosen point $b \in U$. Since P_{ϕ} commutes with smash product with spaces, the map in the previous display is easily identified as the canonical map

$$U_+ \wedge P_{\phi}(\mathcal{J}(V, -)^{\Lambda/K}) \longrightarrow U_+ \wedge \mathcal{J}(V^{\Lambda/K}, -),$$

which is an isomorphism by [22, V.4.5] (as extended by [19, 3.8.12]).

Theorem V.2.5 is now an easy consequence.

PROOF OF THEOREM V.2.5. By Theorem V.2.4

$$\Phi^{M/\Lambda} \Phi^{\Lambda/K} J^{\Gamma/K} \eta \cong \Phi^{M/\Lambda} (J^{\Gamma/\Lambda}(\eta(\Lambda|K))) \cong J^{\Gamma/M}((\eta(\Lambda|K))(M|\Lambda))$$

$$\Phi^{M/K} J^{\Gamma/K} \eta \cong J^{\Gamma/M}(\eta(M|K)).$$

In the notation of Definition V.2.13, we have

$$(\eta(\Lambda|K))(M|\Lambda) = \prod_{\sigma \colon M \to Q} (\eta(\Lambda|K))^{M,\sigma} \{\Lambda\}$$
$$= \prod_{\sigma \colon M \to Q} \left(\left(\prod_{\tau \colon \Lambda \to Q} \eta^{\Lambda,\tau} \{K\} \right)^{M,\sigma} \{\Lambda\} \right).$$

Consider $\sigma \colon M \to Q, \tau \colon \Lambda \to Q$, and an element $b \in B^{\Lambda,\tau}\{K\}$, which when viewed as an element of the base of $\eta(\Lambda|K)$ (in the τ summand) is fixed by the σ -twisted M action. In particular, b is then also in $B^{M,\sigma}$ and the $\sigma|_{\Lambda}$ -twisted Λ action on the total space of $\eta(\Lambda|K)$ restricts to an action on the fiber $E_b^{\Lambda,\tau}$. Since $\eta(\Lambda|K)$ is assumed to be Q-faithful, the $\sigma|_{\Lambda}$ -twisted Λ action on $E_b^{\Lambda,\tau}$ is trivial if and only if $\sigma|_{\Lambda} = \tau$. Moreover, $\sigma|_{\Lambda}$ is a fixed point for the σ -twisted M action on the set of homomorphisms $\Lambda \to Q$, and

$$(\eta^{\Lambda,\sigma|_{\Lambda}}\{K\})^{M,\sigma} = \eta^{M,\sigma}\{K\}.$$

It follows that the inclusion of $\sigma|_{\Lambda}$ in the set of all homomorphisms $\tau \colon \Lambda \to Q$ induces an isomorphism of (Γ, Q) vector bundles

$$\eta(M) = \prod_{\sigma \colon M \to Q} \eta^{M,\sigma} \{K\} = \prod_{\sigma \colon M \to Q} (\eta^{\Lambda,\sigma|_{\Lambda}} \{K\})^{M,\sigma}$$
$$\longrightarrow \prod_{\sigma \colon M \to Q} \left(\prod_{\tau \colon \Lambda \to Q} \eta^{\Lambda,\tau} \{K\} \right)^{M,\sigma} \{\Lambda\} = (\eta(\Lambda|K))(M|\Lambda).$$

This induces an isomorphism $\Phi^{M/K} J^{\Gamma/K} \eta \to \Phi^{M/\Lambda} \Phi^{\Lambda/K} J^{\Gamma/K} \eta$ and it is straightforward to identify this map as the canonical one.

For the proof of Theorem V.2.7, recall the geometric homotopy groups: for a compact Lie group G, closed subgroup H < G, and G-spectrum X, let

$$\pi_q^{\Phi H}(X) = \begin{cases} \operatorname{colim}_V \pi_q(\Omega^{V^H}(X(V)^H)) & q \ge 0\\ \operatorname{colim}_V \pi_0(\Omega^{V^H}(X(\mathbb{R}^{-q} \oplus V)^H)) & q < 0 \end{cases}$$

where the colimit ranges over the finite dimensional subspaces of an H-universe, ordered by inclusion; this depends on the universe but any other choice of universe with the same isomorphism classes of representations gives canonically isomorphic groups. By convention, except when we explicitly mention a universe, we always understand this colimit to be over a complete universe.

The functor $\pi_*^{\Phi H}$ is denoted ρ_*^K in [22, V.4.8.(iii)] for K the trivial group in this case; then [22, V.4.12–16] gives a natural isomorphism from $\pi_q^{\Phi H} X$ to the homotopy groups of the underlying non-equivariant spectrum of the derived geometric fixed point spectrum,

(1.4)
$$\pi_q^{\Phi H} X \cong \pi_q(L\Phi^H X).$$

According to [24, XVI.6.4], these homotopy groups detect weak equivalences and family equivalences of G-spectra; the following proposition holds for any G-universe provided we use the same universe to define weak equivalences and (its restriction to subgroups) to construct $\pi_*^{\Phi H}$:

PROPOSITION 1.5. Let G be a compact Lie group and let \mathcal{F} be a family of subgroups. A map of G-spectra $X \to Y$ is an \mathcal{F} -equivalence if and only if it induces an isomorphism $\pi_*^{\Phi H} X \to \pi_*^{\Phi H} Y$ for all $H \in \mathcal{F}$.

For N a normal subgroup of G contained in H, we have a natural map

$$\pi_q^{\Phi H}(X) \longrightarrow \pi_q^{\Phi(H/N)}(\Phi^N X).$$

induced by the map $\operatorname{Fix}^N X \to \Phi^N X \circ \phi$ as follows. For $q \ge 0$, and V an orthogonal G-representation, we use the map

$$\pi_q(\Omega^{V^H}(X(V)^H)) \cong \pi_q(\Omega^{(V^N)^{H/N}}((X(V)^N)^{H/N}))$$
$$\longrightarrow \pi_q(\Omega^{(V^N)^{H/N}}((\Phi^N X(V^N))^{H/N})) \longrightarrow \pi_q^{\Phi(H/N)}(\Phi^N X)$$

(where we take the colimit implicit in $\pi_q^{\Phi(H/N)}$ over the inclusions in the complete H/N-universe obtained as the N fixed points of the chosen complete H-universe). Similar formulas apply when q < 0. Considering the case when X is cofibrant in the standard model structure, the point-set geometric fixed point functors model the derived geometric fixed point functors, and (using [**22**, V.4.13]) we see that the diagram of natural transformations

$$\begin{array}{ccc} \pi_q^{\Phi H}(X) & \longrightarrow \pi_q^{\Phi(H/N)}(\Phi^N X) \\ \cong & & \downarrow & & \downarrow \\ \pi_q(L\Phi^H(X)) & \longrightarrow \pi_q(L\Phi^{H/N}(L\Phi^N X)) & \longrightarrow \pi_q(L\Phi^{H/N}(\Phi^N X)) \end{array}$$

commutes. In the context of Notation V.2.1, taking $G = \Gamma/K$, the previous proposition now gives the following conclusion.

PROPOSITION 1.6. Let X be a Γ/K -spectrum. The universal map $L\Phi^{\Lambda/K}X \to \Phi^{\Lambda/K}X$ in the Γ/Λ -equivariant stable category is an isomorphism if and only if the natural maps $\pi^{\Phi(M/K)}_*X \to \pi^{\Phi(M/\Lambda)}_*(\Phi^{\Lambda/K}X)$ are isomorphisms for all closed subgroups $M < \Gamma$ containing Λ .

Our strategy for the proof of Theorem V.2.7 is to apply the previous proposition and calculate the maps.

PROOF OF THEOREM V.2.7. By Theorem V.2.4, it suffices to show that the map

$$\pi^{\Phi(M/K)}_* J^{\Gamma/K} \eta \longrightarrow \pi^{\Phi(M/\Lambda)}_* (J^{\Gamma/\Lambda} \eta(\Lambda|K))$$

is an isomorphism for all $M < \Gamma$ containing Λ . We now fix one such M. For V an orthogonal Γ/K -representation, by Lemma 1.2, we have

$$(J^{\Gamma/K}\eta(V))^M \cong \left(\bigvee_{\sigma \colon M \to Q} \mathcal{J}(\eta, V)^{M,\sigma}\right)/Q = \left(\bigvee_{\sigma \colon M \to Q} \mathcal{J}(\eta|_{B^{M,\sigma}\{K\}}, V)^{M,\sigma}\right)/Q$$

1. THE QUOTIENT FIXED POINT LEMMA AND THEOREMS V.2.4, V.2.5, AND V.2.7 75

since V is K-trivial and η is Q-faithful. Likewise, we have

$$((J^{\Gamma/\Lambda}\eta(\Lambda|K))(V^{\Lambda}))^{M} \cong \left(\bigvee_{\sigma \colon M \to Q} \mathcal{J}(\eta^{\Lambda,\sigma|_{\Lambda}}|_{B^{M,\sigma}\{K\}}, V^{\Lambda})^{M,\sigma}\right)/Q;$$

as in the proof of Theorem V.2.5, the summands $\tau \colon \Lambda \to Q$ for $\tau \neq \sigma|_{\Lambda}$ disappear since V^{Λ} is Λ -trivial and $\eta(\Lambda|K)$ is Q-faithful. The map we are calculating is induced by the map

$$(\eta|_{B^{M,\sigma}\{K\}}, V)^{M,\sigma} \longrightarrow \mathcal{J}(\eta^{\Lambda,\sigma|_{\Lambda}}|_{B^{M,\sigma}\{K\}}, V^{\Lambda})^{M,\sigma}$$

which is the induced map on Thom spaces from

J

$$\mathfrak{F}^{\perp}\mathcal{I}(\eta|_{B^{M,\sigma}\{K\}}, V)^{M,\sigma} \longrightarrow \mathfrak{F}^{\perp}\mathcal{I}(\eta^{\Lambda,\sigma|_{\Lambda}}|_{B^{M,\sigma}\{K\}}, V^{\Lambda})^{M,\sigma}$$

which takes a σ -twisted M-equivariant isometry $E_b \to V$ to its $\sigma|_{\Lambda}$ -twisted Λ fixed points $E_b^{\Lambda,\sigma|_{\Lambda}} \to V^{\Lambda}$ and is the identity on the complementary (σ -twisted M fixed) vector space.

To calculate the map on homotopy groups of colimits, consider a complete M/K-universe $U = U_0 \oplus U_1$, where $U_0 = \mathbb{R}^{\infty}$ and $U_1^M = 0$. Then for X a Γ/K -spectrum we can calculate $\pi_q^{\Phi M/K} X$ as

$$\operatorname{colim}_{n \gg -q} \operatorname{colim}_{W < U_1} \pi_{n+q} (X(\mathbb{R}^n \oplus W)^{M/K})$$

for any $q \in \mathbb{Z}$. If we extend the definition of $\mathcal{J}(\eta, -)$ to infinite dimensional inner product spaces by taking the colimit over finite dimensional subspaces, then we get

$$\pi^{\Phi M/K}(J^{\Gamma/K}\eta) \cong \operatorname{colim}_{n} \pi_{n+q} \left(\left(\bigvee_{\sigma \colon M \to Q} \mathcal{J}(\eta|_{B^{M,\sigma}\{K\}}, \mathbb{R}^{n} \oplus U_{1})^{M,\sigma} \right) / Q \right)$$
$$\pi^{\Phi M/\Lambda}(J^{\Gamma/K}\eta(\Lambda|K)) \cong \operatorname{colim}_{n} \pi_{n+q} \left(\left(\bigvee_{\sigma \colon M \to Q} \mathcal{J}(\eta^{\Lambda,\sigma|_{\Lambda}}|_{B^{M,\sigma}\{K\}}, \mathbb{R}^{n} \oplus U_{1}^{\Lambda})^{M,\sigma} \right) / Q \right)$$

Thus, it suffices to show that the map

$$\bigvee_{\sigma \colon M \to Q} \mathcal{J}(\eta|_{B^{M,\sigma}\{K\}}, \mathbb{R}^n \oplus U_1)^{M,\sigma} \longrightarrow \bigvee_{\sigma \colon M \to Q} \mathcal{J}(\eta^{\Lambda,\sigma|_\Lambda}|_{B^{M,\sigma}\{K\}}, \mathbb{R}^n \oplus U_1^{\Lambda})^{M,\sigma}$$

is a Q-equivariant weak equivalence. Both sides are Q-free because the bundles are Q-faithful, so it suffices to show for each $\sigma: M \to Q$, the map

$$\mathcal{J}(\eta|_{B^{M,\sigma}\{K\}},\mathbb{R}^n\oplus U_1)^{M,\sigma}\longrightarrow \mathcal{J}(\eta^{\Lambda,\sigma|_{\Lambda}}|_{B^{M,\sigma}\{K\}},\mathbb{R}^n\oplus U_1^{\Lambda})^{M,\sigma}$$

is a non-equivariant homotopy equivalence.

We understand the M action on the vector bundle $\eta|_{B^{M,\sigma}\{K\}}$ to be the σ twisted M action. This is a fiberwise K fixed M action and the usual integration argument decomposes $\eta|_{B^{M,\sigma}\{K\}}$ into an internal Whitney sum

$$\eta|_{B^{M,\sigma}\{K\}} = \theta_0 \oplus \theta_1$$

where $\theta_0 = (\eta|_{B^{M,\sigma}\{K\}})^{M/K}$ and $\theta_1 = \theta_0^{\perp}$ satisfies $\theta_1^{M/K}$ is zero dimensional. Because Λ is normal in M, θ_1 decomposes further as

$$\theta = \theta_1^{\Lambda/K} \oplus \theta_1'.$$

We then have

$$\mathcal{J}(\eta|_{B^{M,\sigma}\{K\}}, \mathbb{R}^n \oplus U_1)^{M,\sigma} \cong \mathcal{J}(\theta_0, \mathbb{R}^n) \times \mathcal{I}(\theta_1, U_1)^{M/K}$$
$$\mathcal{J}(\eta^{\Lambda,\sigma|_\Lambda}|_{B^{M,\sigma}\{K\}}, \mathbb{R}^n \oplus U_1^{\Lambda})^{M,\sigma} \cong \mathcal{J}(\theta_0, \mathbb{R}^n) \times \mathcal{I}(\theta_1^{\Lambda/K}, U_1^{\Lambda/K})^{M/K};$$

the map preserves this product decomposition, is the identity on the first factor, and is a homotopy equivalence on the second factors because both are contractible. \Box

2. Reduction of Theorems V.2.6 and V.2.9

This section reduces Theorems V.2.6 and V.2.9 to a simpler but more technical result stated as Theorem 2.2 below and proved in the next section. As in the statements, we follow the notational convention of Notation V.2.1: Γ is a compact Lie group with a closed normal subgroups K, Λ satisfying $K < \Lambda$.

Theorem 2.2 is stated in terms of a variant of the functor $\operatorname{Fix}^{\Gamma}$ that we write as $\operatorname{Fix}_{K}^{\Gamma}$. The domain of $\operatorname{Fix}_{K}^{\Gamma}$ is the category of Γ -spectra indexed on K-trivial orthogonal Γ -representations, or equivalently, orthogonal Γ/K -representations. It is equivalent to the category of Γ -equivariant $\mathcal{J}_{\Gamma/K}$ -spaces, and we mostly refer to it this way, except in some headline statements, where we write Γ -spectra indexed on K-trivial orthogonal Γ -representations for historical and philosophical reasons. Starting with a Γ -equivariant $\mathcal{J}_{\Gamma/K}$ -space and taking Γ fixed points levelwise gives a (non-equivariant) $\mathcal{J}_{\Gamma/K}^{\Gamma/K}$ -space. (We emphasize that $\mathcal{J}_{\Gamma/K}^{\Gamma/K}$ -spaces here are always understood to be non-equivariant.)

DEFINITION 2.1. Let $\operatorname{Fix}_{K}^{\Gamma}$ be the functor from Γ -equivariant $\mathcal{J}_{\Gamma/K}$ -spaces to $\mathcal{J}_{\Gamma/K}^{\Gamma/K}$ -spaces defined by

$$(\operatorname{Fix}_{K}^{\Gamma} X)(W) = (X(W))^{\Gamma}.$$

We often want to apply $\operatorname{Fi}_{K}^{\Gamma}$ to a Γ -spectrum Y; to avoid confusion, we write iY for the underlying Γ -equivariant $\mathcal{J}_{\Gamma/K}$ -space obtained by restricting Y to the K-trivial orthogonal Γ -representations. In parallel, for a Γ/K -spectrum X, we get a Γ -equivariant $\mathcal{J}_{\Gamma/K}$ -space jX obtained by regarding Γ/K -spaces as Γ -spaces with trivial K action.

The category of $\mathcal{J}_{\Gamma/K}^{\Gamma/K}$ -spaces has a smash product defined by Day convolution for the smash product in $\mathcal{J}_{\Gamma/K}^{\Gamma/K}$ induced by direct sum; see [23, §21]. The functor Fix_K^{Γ} is lax symmetric monoidal for this smash product, with structure map

$$\operatorname{Fix}_{K}^{\Gamma} X \wedge \operatorname{Fix}_{K}^{\Gamma} Y \longrightarrow \operatorname{Fix}_{K}^{\Gamma} (X \wedge Y)$$

induced by the natural transformation

$$(\operatorname{Fix}_{K}^{\Gamma} X)(V) \wedge (\operatorname{Fix}_{K}^{\Gamma} Y)(W) = (X(V))^{\Gamma} \wedge (Y(W))^{\Gamma}$$
$$\longrightarrow ((X \wedge Y)(V \oplus W))^{\Gamma} = (\operatorname{Fix}_{K}^{\Gamma} (X \wedge Y))(V \oplus W)$$

where the middle map is the induced map on Γ fixed points of the universal map $X(V) \wedge Y(W) \rightarrow (X \wedge Y)(V \oplus W)$ for the smash product of Γ -spectra indexed on K-trivial orthogonal Γ -representations.

For some of the work, we will need the homotopy theory of $\mathcal{J}_{\Gamma/K}^{\Gamma/K}$ -spaces. We understand a *weak equivalence* as a π_* -isomorphism, that is an isomorphism on homotopy groups: for a $\mathcal{J}_{\Gamma/K}^{\Gamma/K}$ -space X, we define homotopy groups by

$$\pi_q X := \begin{cases} \operatorname{colim}_V \pi_q \Omega^{V^{\Gamma/K}} X(V) & q \ge 0\\ \operatorname{colim}_V \pi_0 \Omega^{V^{\Gamma/K}} X(\mathbb{R}^{-q} \oplus V) & q < 0 \end{cases}$$

where the colimit ranges over the finite dimensional subspaces of a complete Γ/K universe (cf. [22, V.4.8.i]). These are the weak equivalences in a model structure whose cofibrant objects are the retracts of cell complexes built by cells of the form $\mathcal{J}_{\Gamma/K}^{\Gamma/K}(V,-) \wedge (D^n, \partial D^n)_+$.

In the following theorem, $F = \mathcal{F}(1 \times Q^{\text{op}})$ denotes the family of graph subgroups, and EF denotes the universal space for this family (sometimes elsewhere written $E_{\Gamma}Q^{op}$). The graph subgroups are precisely those subgroups whose intersection with $1 \times Q^{\text{op}}$ is the trivial subgroup. The universal space EF is a $(\Gamma \times Q^{\text{op}})$ -CW complex characterized up to equivariant homotopy equivalence by the condition that EF^{H} is contractible when H is a graph subgroup of $\Gamma \times Q^{\text{op}}$ and is empty when H is not.

THEOREM 2.2. Let η be the (Γ, Q) -vector bundle

$$(\Gamma \times Q^{\mathrm{op}}) \times_M V \times Y \longrightarrow (\Gamma \times Q^{\mathrm{op}})/M \times Y$$

for M a closed subgroup of $\Gamma \times Q^{\text{op}}$, V an orthogonal M-representation, and Y a $(\Gamma \times Q^{\text{op}})$ -equivariant cell complex. Assume that η is Q-faithful. Then for any Γ -spectrum X indexed on K-trivial orthogonal Γ -representations:

(i) The canonical map

$$\operatorname{Fix}_{K}^{\Gamma}(\imath J\eta) \wedge \operatorname{Fix}_{K}^{\Gamma}(X) \longrightarrow \operatorname{Fix}_{K}^{\Gamma}(\imath J\eta \wedge X)$$

is an isomorphism.

(ii) The map $EF \rightarrow *$ induces a weak equivalence

$$\operatorname{Fix}_{K}^{\Gamma}(iJ(EF \times \eta) \wedge X) \longrightarrow \operatorname{Fix}_{K}^{\Gamma}(iJ\eta \wedge X).$$

We now deduce Theorems V.2.6 and V.2.9 from Theorem 2.2.

PROOF OF THEOREM V.2.6 FROM THEOREM 2.2. Let η and X be as in the statement of Theorem V.2.6. Since $\Phi^{\Lambda/K}$ is the composite of Fix^{Λ/K} with left Kan extension, it suffices to show that the map

$$\operatorname{Fix}^{\Lambda/K}(J^{\Gamma/K}\eta)\wedge\operatorname{Fix}^{\Lambda/K}X\longrightarrow\operatorname{Fix}^{\Lambda/K}(J^{\Gamma/K}\eta\wedge X)$$

is an isomorphism of non-equivariant $\mathcal{J}_{\Gamma/K}^{\Lambda/K}$ -spaces. Non-equivariantly, the restriction of equivariance functor induces an equivalence of (non-equivariant) topological categories from $\mathcal{J}_{\Gamma/K}^{\Lambda/K}$ to a full subcategory of $\mathcal{J}_{\Lambda/K}^{\Lambda/K}$, and so it suffices to show that the map above is an isomorphism of non-equivariant $\mathcal{J}_{\Lambda/K}^{\Lambda/K}$ -spaces for any Λ equivariant $\mathcal{J}_{\Lambda/K}$ -space X. Thus, by a change of notation, we can assume without loss of generality that $\Lambda = \Gamma$. Then

$$\begin{split} \operatorname{Fix}^{\Lambda/K}(J^{\Gamma/K}\eta) &\cong \operatorname{Fix}_{K}^{\Gamma}(\imath J\eta), \quad \operatorname{Fix}^{\Lambda/K}(X) &\cong \operatorname{Fix}_{K}^{\Gamma}(\jmath X), \quad \text{and} \\ \operatorname{Fix}^{\Lambda/K}(J^{\Gamma/K}\eta \wedge X) &\cong \operatorname{Fix}_{K}^{\Gamma}(\imath J\eta \wedge \jmath X), \end{split}$$

compatibly with the universal map, so it suffices to show that the map

$$\operatorname{Fix}_{K}^{\Gamma}(\imath J\eta) \wedge \operatorname{Fix}_{K}^{\Gamma}(\jmath X) \longrightarrow \operatorname{Fix}_{K}^{\Gamma}(\imath J\eta \wedge \jmath X).$$

is an isomorphism. Since both sides commute with cofiber sequences and sequential colimits over Hurewicz cofibrations, working a cell at a time, it suffices to consider the case when the base of η consists of a single cell or its boundary, i.e., when η is as in the statement of Theorem 2.2 for $Y = D^n$ or ∂D^n . Part (i) of Theorem 2.2 establishes this case.

For Theorem V.2.9, we need the following additional observation.

PROPOSITION 2.3. Let $\eta: E \to B$ be a (Γ, Q) vector bundle such that B is a $(\Gamma \times Q^{\text{op}})$ -equivariant CW complex with isotropy groups in $\mathcal{F}(1 \times Q^{\text{op}})$. Then $\operatorname{Fix}_{K}^{\Gamma}(iJ\eta)$ is a cellular $\mathcal{J}_{\Gamma/K}^{\Gamma/K}$ -space, i.e., is built from cells of the form $\mathcal{J}(V, -) \wedge (D^{n}, \partial D^{n})_{+}$ for V an orthogonal Γ/K -representation.

PROOF. Working over the cells of B, it suffices to check the case when $B = (\Gamma \times Q^{\text{op}})/\tilde{H}$ for $\tilde{H} \in \mathcal{F}(1 \times Q^{\text{op}})$. Let $H < \Gamma$ be the image of \tilde{H} under the projection; then the projection $\tilde{H} \to H$ is an isomorphism and there exists a unique homomorphism $\sigma: H \to Q$ such that

$$\tilde{H} = H_{\sigma} = \{ (h, \sigma(h^{-1})) \mid h \in H \}.$$

The description of B as an orbit implicitly chooses a point with isotropy subgroup \tilde{H} . Let V denote the fiber over that point; then V has the structure of an orthogonal \tilde{H} -representation. Because $\tilde{H} \cap (1 \times Q^{\text{op}})$ is trivial, for any orthogonal Γ/K -representation W, we have

$$(\operatorname{Fix}_{K}^{\Gamma}(iJ\eta))(W) = (\mathcal{J}(\eta, W)/Q)^{\Gamma} \cong (((\Gamma \times Q^{\operatorname{op}})_{+} \wedge_{\tilde{H}} \mathcal{J}(V, W))/Q)^{\Gamma} \\ \cong (\Gamma_{+} \wedge_{H} \mathcal{J}(V, W))^{\Gamma}$$

where in the last isomorphism H acts on V via the isomorphism $\tilde{H} \to H$. We see that this based space is trivial unless $H = \Gamma$, in which case we get $\mathcal{J}(V, W)^{\Gamma}$. Since the K action on W is trivial, there are no Γ -equivariant isometries $V \to W$ unless the K action on V is also trivial, in which case

$$\mathcal{J}(V,W)^{\Gamma} = \mathcal{J}_{\Gamma/K}^{\Gamma/K}(V,W).$$

Thus, $\operatorname{Fix}_{K}^{\Gamma}(iJ\eta)$ is either * or $\mathcal{J}_{\Gamma/K}^{\Gamma/K}(V,-)$ for an orthogonal Γ/K -representation V.

The argument of [23, 12.3] shows that cellular $\mathcal{J}_{\Gamma/K}^{\Gamma/K}$ -spaces are flat for the smash product. To make this argument work, it suffices to see that for any K-trivial Γ -representation V, the map

$$S^{(V\oplus V)^{\Gamma}} \longrightarrow \mathcal{J}_{\Gamma/K}^{\Gamma/K}(V, V \oplus V) \wedge S^{V^{\Gamma}}$$

(induced by the inclusion of V as the first summand) is a $2(\dim V^{\Gamma})$ -equivalence. This is easy to see by decomposing V into irreducible Γ -representations: each non-trivial irreducible representation contributes a smash factor of $O(2n)/O(n)_+$, $U(2n)/U(n)_+$, or $Sp(2n)/Sp(n)_+$ (where n is the number of summands of that type), and the trivial representation together with $S^{V^{\Gamma}}$ contribute a smash factor of

$$(O(2n)_+ \wedge_{O(n)} S^n) \wedge S^n \cong O(2n)/O(n)_+ \wedge S^{2n}$$

(where $n = \dim(V^{\Gamma})$). We can now deduce Theorem V.2.9. In fact, we prove the following more general statement.

THEOREM 2.4. Suppose η is a Q-faithful (Γ, Q) vector bundle and assume the base B admits the structure of a $(\Gamma \times Q^{\text{op}})$ -equivariant cell complex. Then $iJ\eta$ is flat for the smash product in Γ -spectra indexed on K-trivial orthogonal Γ representations. Theorem V.2.9 is a consequence of the previous theorem because of the following easy facts:

- (i) When η is Γ/K -compatible, $iJ\eta = jJ^{\Gamma/K}\eta$.
- (ii) When X and Y are Γ/K -spectra $jX \wedge jY \cong j(X \wedge Y)$
- (iii) When X and Y are Γ/K -spectra a map $X \to Y$ is a weak equivalence of Γ/K -spectra if and only if applying j takes it to a weak equivalence of Γ -spectra indexed on K-trivial orthogonal Γ -representations.

PROOF OF THEOREM 2.4 FROM THEOREM 2.2. Let η be as in the statement of Theorem 2.4. To show that $iJ\eta$ is flat, by definition, we need to show that for any weak equivalence $X \to X'$ of Γ -equivariant $\mathcal{J}_{\Gamma/K}$ -spaces, the induced map

$$iJ\eta \wedge X \longrightarrow iJ\eta \wedge X$$

is a weak equivalence. For this, by [24, XVI.6.4] (see also [22, V.4.12–16] or Proposition 1.5), it suffices to show that for every closed subgroup H of Γ , the map

$$\operatorname{Fix}_{H\cap K}^{H}(j_{H}^{*}iJ\eta \wedge j_{H}^{*}X) \longrightarrow \operatorname{Fix}_{H\cap K}^{H}(j_{H}^{*}iJ\eta \wedge j_{H}^{*}X')$$

is a weak equivalence, where j_H denotes the inclusion of H in Γ , and j_H^* denotes the reduction of structure functor from Γ -equivariant $\mathcal{J}_{\Gamma/K}$ -spaces to H-equivariant $\mathcal{J}_{H/(H\cap K)}$ -spaces. Working one H at a time, we can change notation, replacing Γ with H, and it suffices to show that the induced map

$$\operatorname{Fix}_{K}^{\Gamma}(iJ\eta \wedge X) \longrightarrow \operatorname{Fix}_{K}^{\Gamma}(iJ\eta \wedge X')$$

is a weak equivalence (for any $X \to X'$ and η as in the statement of Theorem 2.4). Working a cell at a time over the base of η , it suffices to prove the case when η is as in the statement of Theorem 2.2 for $Y = D^n$ or ∂D^n . If EF is a universal F-space for $F = \mathcal{F}(1 \times Q^{\text{op}})$ (as above), then $\operatorname{Fix}_K^{\Gamma}(iJ(EF \times \eta))$ is a cellular $\mathcal{J}_{\Gamma/K}^{\Gamma/K}$ -space by the Proposition 2.3, and hence the top map in the commuting diagram

1

is a weak equivalence. Applying part (ii) of Theorem 2.2, we see that the vertical maps are weak equivalences, and we conclude that the bottom horizontal map is a weak equivalence. Part (i) of Theorem 2.2 then finishes the proof. \Box

3. Proof of Theorem 2.2

Let η , M, V, X, and Y be as in the statement of Theorem 2.2. For the proof of the theorem, it suffices to treat the case when the projection map $M \to \Gamma$ is onto, as otherwise all $\mathcal{J}_{\Gamma/K}^{\Gamma/K}$ -spaces mentioned in (i) and (ii) are trivial.

Let $H \triangleleft M$ be the kernel of the projection $M \rightarrow \Gamma$. Let $L \triangleleft M$ be the inverse image of K. Then

$$M/L \xrightarrow{\cong} \Gamma/K$$

and we use this isomorphism to pass back and forth between L-trivial M actions, K-trivial Γ actions, and Γ/K actions without comment or notation.

As a subgroup of $\Gamma \times Q^{\text{op}}$, $H = M \cap (1 \times Q^{\text{op}})$ and so in particular its projection to Q^{op} is an injection; write H' for the subgroup of Q opposite to H. Whenever we have a left action of H, we have a corresponding right action by H' that we will use without further comment. For any orthogonal Γ/K -representation W, we then have an Γ -equivariant isomorphism

$$\mathcal{J}(\eta, W)/Q \cong (\mathcal{J}(V, W) \wedge Y_+)/H'$$

which is a natural isomorphism of based Γ -space enriched functors from $\mathcal{J}_{\Gamma/K}$ to Γ -equivariant based spaces, i.e., a map of Γ -equivariant $\mathcal{J}_{\Gamma/K}$ -spaces. Using this isomorphism and the fact that smash product commutes with colimits, as in [23, 4.4] (or see the proofs of [4, A.1] or [22, III.8.4]) we get an isomorphism

(3.1)
$$(iJ\eta \wedge X)(W) \cong (\mathcal{I}(V \oplus \mathbb{R}^n, W) \times Y)_+ \wedge_{H' \times O(n)} (S^0 \wedge X(\mathbb{R}^n))_+$$

where dim $W = n + \dim V$. The right action on $\alpha \in \mathcal{I}(V \oplus \mathbb{R}^n, W)$ is defined by

$$\alpha \cdot (\sigma, r) := \alpha \circ (\sigma_*^{-1} \oplus r)$$

for $\sigma \in H'$ and $r \in O(n)$. The right action of O(n) on Y is trivial. We note that since the action of H' < Q on η is faithful by hypothesis, the action of $H' \times O(n)$ on

$$\mathcal{I}(\eta, W) = \mathcal{I}(V \oplus \mathbb{R}^n, W) \times Y$$

is free.

We use the isomorphism in (3.1) to analyze $\operatorname{Fix}_{K}^{\Gamma}(iJ\eta \wedge X)$ as follows. To compress notation, we write

$$\Xi(W) := (\mathcal{I}(V \oplus \mathbb{R}^n, W) \times Y)_+ \wedge (S^0 \wedge X(\mathbb{R}^n)),$$

so that

$$\bar{\Xi}(W)/(H' \times O(n)) \cong (\mathcal{I}(V \oplus \mathbb{R}^n, W) \times Y)_+ \wedge_{H' \times O(n)} (S^0 \wedge X(\mathbb{R}^n))$$

and per (3.1)

(3.2)
$$(\bar{\Xi}(W)/(H' \times O(n)))^{\Gamma} \cong \operatorname{Fix}_{K}^{\Gamma}(iJ\eta \wedge X)(W).$$

We also work with the space

$$\Xi(W) := \mathcal{I}(V \oplus \mathbb{R}^n, W) \times Y \times (S^0 \wedge X(\mathbb{R}^n)).$$

Let

$$\Psi(W) \subset \Xi(W), \qquad \overline{\Psi}(W) \subset \overline{\Xi}(W)$$

denote the inverse image of the Γ fixed point subspaces of $\Xi(W)/(H' \times O(n))$ and $\overline{\Xi}(W)/(H' \times O(n))$, respectively.

We next observe that $\Psi(W)$ and $\Psi(W)$ have the obvious quotient relationship, but this takes a bit of work because of the point-set topology of quotients of (compactly generated) weak Hausdorff spaces. Since being Γ fixed is a closed condition, these are closed subspaces. Let $\Psi(W)_* \subset \Psi(W)$ denote the inverse image of the basepoint of $S^0 \wedge X(\mathbb{R}^n)$ under the projection. Then $\Psi(W)_* \subset \Psi(W)$ is closed and the map from the pushout to the union of subspaces

$$\Psi(W) \cup_{\Psi(W)_*} (\mathcal{I}(V \oplus \mathbb{R}^n, W) \times Y \times *) \longrightarrow \Psi(W) \cup (\mathcal{I}(V \oplus \mathbb{R}^n, W) \times Y \times *)$$

is an isomorphism. In particular, the map $\Psi(W)/\Psi(W)_* \to \overline{\Psi}(W)$ is an isomorphism. Because the group $H' \times O(n)$ is compact, the equivalence relation associated to an $(H' \times O(n))$ action is closed. The underlying set of a quotient (in the category

of compactly generated weak Hausdorff spaces) is then the quotient of the underlying set; moreover, the quotient of a closed $(H' \times O(n))$ -stable subspace includes as a closed subspace of the quotient. Specifically, we have that the maps

(3.3)
$$(\Psi(W)/\Psi(W)_*)/(H' \times O(n)) \xrightarrow{\cong} \bar{\Psi}(W)/(H' \times O(n))$$
$$\xrightarrow{\cong} (\bar{\Xi}(W)/(H' \times O(n)))^{\Gamma}$$

are isomorphisms. This lets us work with $\Psi(W)$ to study $\operatorname{Fix}_{K}^{\Gamma}(iJ\eta \wedge X)(W)$. Our strategy is to decompose $\Psi(W)$ into a disjoint union of pieces corresponding to graph subgroups as in the study of fixed points in Section 1 and then to analyze the individual pieces to identify $\Psi(W)$ and $\overline{\Psi}(W)$. (The precise statement is given as Theorem 3.7 below.)

We write (α, y, x) for an element of $\mathcal{I}(V \oplus \mathbb{R}^n, W) \times Y \times (S^0 \wedge X(\mathbb{R}^n))$ with $\alpha \in \mathcal{I}(V \oplus \mathbb{R}^n, W), y \in Y$, and $x \in X(\mathbb{R}^n)$. By definition, (α, y, x) is in Ψ if and only if its image $[\alpha, y, x]$ in the quotient $\Xi/(H' \times O(n))$ is fixed by Γ if and only if that image is fixed by the restriction action of M if and only if for each $g \in M$, there exists a $\tilde{\sigma}(g) \in H'$ and $r(g) \in O(n)$ such that

(3.4)
$$g \cdot \alpha = \alpha \cdot (\tilde{\sigma}(g), r(g)), \quad gy = y\tilde{\sigma}(g), \quad gx = r(g)^{-1}x$$

Because the action of $H' \times O(n)$ on $\mathcal{I}(V \oplus \mathbb{R}^n, W) \times Y$ is free, $\tilde{\sigma}(g)$ and r(g) are uniquely determined, and the existence of local sections shows that

$$\tilde{\sigma} \times r \colon M \longrightarrow H' \times O(n)$$

is continuous.

Looking at what happens for the product of elements in M, we see that r is a homomorphism. Indeed, because M acts trivially on \mathbb{R}^n and L acts trivially on W, r descends to a homomorphism $\Gamma/K \to O(n)$. On the other hand, $\tilde{\sigma}$ is not a homomorphism but rather satisfies the following formulas:

- (i) Restricted to $H \lhd M$, $\tilde{\sigma}$ is the projection anti-isomorphism $H^{\mathrm{op}} \cong H'$, and
- (ii) For $g, h \in M$, $\tilde{\sigma}(gh) = \tilde{\sigma}(g)({}^{g}\tilde{\sigma}(h))$, where ${}^{g}(-)$ denotes the left action of g on H' corresponding to the conjugation action $h \mapsto ghg^{-1}$ on H.

Equivalently, the graph $h \mapsto (h, \tilde{\sigma}(h))$ defines a continuous graph homomorphism from M into the semidirect product $H' \rtimes M$ extending the diagonal on H; for the purposes of this argument we call a continuous map $M \to H'$ satisfying (i) and (ii) above an *sdde-graph homomorphism* and write $\tilde{\sigma} \colon M \rightsquigarrow H'$. We were led to this formulation by the fixed point formulas, but we can rewrite the structure in terms of graph subgroups of $M < \Gamma \times Q^{\text{op}}$ and σ -twisted Γ actions as follows.

PROPOSITION 3.5. There is a one-to-one correspondence between the set of sdde-graph homomorphisms $\tilde{\sigma} \colon M \rightsquigarrow H'$ and the set of homomorphisms $\sigma \colon \Gamma \to Q$ for which the corresponding graph subgroup

$$\Gamma_{\sigma} = \{(\gamma, \sigma(\gamma^{-1})) \mid \gamma \in \Gamma\} < \Gamma \times Q^{\mathrm{op}}$$

lies in M. Given $\tilde{\sigma}$ and an M-space Z, the corresponding σ -twisted Γ action on $z \in Z$ is given by

$$\gamma \cdot_{\sigma} z = (gz)\tilde{\sigma}(g)^{-1}$$

where the projection $M \to \Gamma$ sends g to γ .

PROOF. The formulas are easier when we use the anti-isomorphism to write $\tilde{\sigma}$ in terms of H: let $\tilde{\sigma}': M \to H$ denote the composite of $\tilde{\sigma}$ with the anti-isomorphism $H' \cong H$. The defining rules for a sdde-graph homomorphism then become

- (i) Restricted to $H \triangleleft M$, $\tilde{\sigma}'$ is the identity, and
- (ii) For $g_1, g_2 \in M$, $\tilde{\sigma}'(g_1g_2) = g_1\tilde{\sigma}'(g_2)g_1^{-1}\tilde{\sigma}'(g_1)$.

Given $\tilde{\sigma}' \colon M \to H$, define $\sigma \colon \Gamma \to Q$ by

$$(\gamma, \sigma(\gamma)^{-1}) = \tilde{\sigma}'(g)^{-1}g \in \Gamma \times Q^{\mathrm{op}}$$

where $g \mapsto \gamma$; this is well-defined because if $h \in H$, then

$$\tilde{\sigma}'(hg)^{-1}hg = (h\tilde{\sigma}'(g)h^{-1}\tilde{\sigma}'(h))^{-1}hg = (h\tilde{\sigma}'(g))^{-1}hg = \tilde{\sigma}(g)^{-1}g$$

and σ is a homomorphism because

$$\begin{aligned} (\gamma_1, \sigma(\gamma_1)^{-1})(\gamma_2, \sigma(\gamma_2)^{-1}) &= \tilde{\sigma}'(g_1)^{-1}g_1\tilde{\sigma}'(g_2)^{-1}g_2 \\ &= \tilde{\sigma}'(g_1)^{-1}g_1\tilde{\sigma}'(g_2)^{-1}g_1^{-1}g_1g_2 \\ &= (g_1\tilde{\sigma}'(g_2)g_1^{-1}\tilde{\sigma}'(g_1))^{-1}g_1g_2 \\ &= \tilde{\sigma}'(g_1g_2)^{-1}g_1g_2 = (\gamma_1\gamma_2, \sigma(\gamma_1\gamma_2)^{-1}), \end{aligned}$$

noting that the multiplication rule in $\Gamma \times Q^{\text{op}}$ satisfies

$$(\gamma_1, a_1^{-1})(\gamma_2, a_2^{-1}) = (\gamma_1 \gamma_2, (a_1 a_2)^{-1}).$$

On the other hand, given $\sigma \colon \Gamma \to Q$ such that $\Gamma_{\sigma} \subset M$, define

$$\tilde{\sigma}'(g) = g(\gamma^{-1}, \sigma(\gamma))$$

where $g \mapsto \gamma$. Since the right side is in M and the projection $M \to \Gamma$ sends $\tilde{\sigma}'(g)$ to the identity, $\tilde{\sigma}'(g) \in H$. If $g \in H$, then γ is the identity and $\tilde{\sigma}'(g) = g$. Given $g_1, g_2 \in M$,

$$\begin{split} \tilde{\sigma}'(g_1g_2) &= g_1g_2((\gamma_1\gamma_2)^{-1}, \sigma(\gamma_1\gamma_2)) \\ &= g_1g_2(\gamma_2^{-1}, \sigma(\gamma_2))(\gamma_1^{-1}, \sigma(\gamma_1)) \\ &= g_1g_2(\gamma_2^{-1}, \sigma(\gamma_2))g_1^{-1}g_1(\gamma_1^{-1}, \sigma(\gamma_1)) \\ &= g_1\tilde{\sigma}'(g_2)g_1^{-1}\tilde{\sigma}'(g_1). \end{split}$$

These rules are obviously inverse, establishing the one-to-one correspondence. For $z \in Z$, by definition, the σ -twisted Γ action on z is given by

$$\gamma \cdot_{\sigma} z := (\gamma, \sigma(\gamma)^{-1}) z$$

which by the above is $\tilde{\sigma}'(g)^{-1}gz$, or written in terms of the right H' action $\gamma \cdot_{\sigma} z = (gz)\tilde{\sigma}(g)^{-1}$.

We call a continuous homomorphism with the property that the corresponding graph subgroup $\Gamma_{\sigma} < \Gamma \times Q^{\text{op}}$ lies M an M graph subgroup homomorphism and write $\sigma \colon \Gamma \xrightarrow{M} Q$. Given such a homomorphism σ , we write $\tilde{\sigma} \colon M \rightsquigarrow H'$ for the corresponding sdde-graph homomorphism. We note that care must be taken with the action formula $\gamma \cdot_{\sigma} z = (gz)\tilde{\sigma}(g)^{-1}$ as the left action of M and the right action of H' do not commute, rather $g(z\sigma) = (gz)({}^{g}\sigma)$. Returning to the study of Ψ , given $\sigma \colon \Gamma \xrightarrow{M} Q$ and $r \colon \Gamma/K \to O(n)$, define the (σ, r) -twisted Γ actions on $\mathcal{I}(V \oplus \mathbb{R}^n, W)$, Y and X(W) by

$$\begin{split} \gamma \cdot_{(\sigma,r)} \alpha &:= (\gamma \cdot_{\sigma} \alpha) \cdot (\mathrm{id}, r(\gamma)) = (g \cdot \alpha) \cdot (\tilde{\sigma}(g), r(\gamma))^{-1} \\ \gamma \cdot_{(\sigma,r)} y &:= \gamma \cdot_{\sigma} y = \gamma y \sigma(\gamma^{-1}) = (gy) \tilde{\sigma}(g)^{-1} \\ \gamma \cdot_{(\sigma,r)} x &:= r(\gamma) \cdot \gamma x \end{split}$$

where $g \mapsto \gamma$; these formulas give well-defined actions because the σ -twisted Γ action on $\mathcal{I}(V \oplus \mathbb{R}^n, W)$ and the natural Γ action on X(W) commute with the O(n) action. We then put these together to define the (σ, r) -twisted Γ action on Ξ ,

$$\gamma \cdot_{(\sigma,r)} (\alpha, y, x) := (\gamma \cdot_{(\sigma,r)} \alpha, \gamma \cdot_{(\sigma,r)} y, \gamma \cdot_{(\sigma,r)} x).$$

The following proposition now summarizes the formulas in (3.4).

PROPOSITION 3.6. An element ξ of Ξ is in Ψ if and only if there exists a continuous homomorphisms $r: \Gamma/K \to O(n)$ and an M graph subgroup homomorphism $\sigma: \Gamma \xrightarrow{M} Q$ such that ξ is fixed by the (σ, r) -twisted Γ action on Ξ . When r and σ exist, they are unique.

We can rewrite the (σ, r) -twisted Γ action of the previous paragraph more intrinsically to develop an intrinsic identification of $\operatorname{Fix}_{K}^{\Gamma}(iJ\eta \wedge X)$. Given a homomorphism $r \colon \Gamma/K \to O(n)$, we get an orthogonal Γ/K -representation R_r with underlying inner product space \mathbb{R}^n and Γ/K action given by r. Given $\sigma \colon \Gamma \xrightarrow{M} Q$, we get σ -twisted Γ actions on V and Y; we write V_{σ} for the orthogonal Γ -representation with underlying inner product space V and Γ action given by the σ -twisted Γ action on V, and we write Y_{σ} for the space Y with Γ action given by the σ twisted Γ action on Y. Then the (σ, r) -twisted Γ action on $\mathcal{I}(V \oplus \mathbb{R}^n, W)$ is precisely the canonical Γ action on $\mathcal{I}(V_{\sigma} \oplus R_r, W)$ and the (σ, r) -twisted Γ action on $(\mathcal{I}(V \oplus \mathbb{R}^n, W) \times Y)_+ \wedge (S^0 \wedge X(\mathbb{R}^n))$ is precisely the canonical Γ action on $(\mathcal{I}(V_{\sigma} \oplus R_r, W) \times Y_{\sigma})_+ \wedge (S^0 \wedge X(\mathbb{R}^n))$. We see from this reformulation that if there exists an (σ, r) -twisted fixed point of $\mathcal{I}(V \oplus \mathbb{R}^n, W)$, then $V_{\sigma} \oplus R_r$ is Γ -isomorphic to W, and this implies that K acts trivially V_{σ} as well as R_r . In particular, R_r and V_{σ} are both orthogonal Γ/K -representations.

We say that an M graph subgroup homomorphism is W-compatible when V_{σ} is Γ -isomorphic to a subrepresentation of W, and write $\sigma \colon \Gamma \xrightarrow{W} Q$. By the previous proposition, only the W-compatible M graph subgroup homomorphisms are relevant to $\Psi(W) \subset \Xi(W)$ and $\xi \in \Xi(W)$ cannot be a (σ, r) -twisted Γ fixed point unless σ is W-compatible.

The (conjugation) action of H' on the set of M graph subgroup homomorphisms preserves the subset of W-compatible ones. Indeed, if $\alpha: V_{\sigma} \oplus R_r \to W$ is an isomorphism of orthogonal Γ/K -representations, then for all $\tau \in H'$, $\alpha \cdot (\tau, \mathrm{id})$ is an isomorphism of orthogonal Γ/K -representations $V_{\sigma_{\tau}} \oplus R_r \to W$. For each H'orbit of W-compatible M graph subgroup homomorphisms $[\sigma]$, choose and fix a homomorphism $r[\sigma]$ so that $V_{\sigma} \oplus R_{r[\sigma]} \cong W$. We then get a map

$$(\mathcal{I}(V_{\sigma} \oplus R_{r[\sigma]}, W) \times Y_{\sigma} \times X(R_{r[\sigma]}))^{\Gamma} \longrightarrow \Psi(W)^{\Gamma, \sigma} \subset \Xi(W)$$

which is $O(R_{r[\sigma]})^{\Gamma/K}$ -equivariant, where

$$O(R_{r[\sigma]})^{\Gamma/K} \subset O(R_{r[\sigma]}) = O(n)$$

denotes the Lie subgroup of Γ/K -equivariant isometries (the final equality holds because the underlying inner product space of $R_{r[\sigma]}$ is \mathbb{R}^n). Inducing up to O(n), we get an O(n)-equivariant map; let T(W) be the disjoint union of these spaces over the W-compatible M graph subgroup homomorphisms

$$T(W) := \coprod_{\sigma \colon \Gamma \xrightarrow{W} Q} (\mathcal{I}(V_{\sigma} \oplus R_{r[\sigma]}, W) \times Y_{\sigma} \times X(R_{r[\sigma]}))^{\Gamma} \times_{O(R_{r[\sigma]})^{\Gamma/K}} O(n)$$

We give this the right H' action where $\tau \in H'$ sends $v \in V_{\sigma}$ to $v\tau \in V_{\sigma_{\tau}}$ and $y \in Y_{\sigma}$ to $y\tau \in Y_{\sigma_{\tau}}$ (acting trivially on $R_{r[\sigma]}$). Likewise, let

$$\bar{T}(W) = \bigvee_{\sigma \colon \Gamma \xrightarrow{W} Q} \left(\left(\mathcal{I}(V_{\sigma} \oplus R_{r[\sigma]}, W) \times Y_{\sigma} \right)_{+} \wedge X(R_{r[\sigma]}) \right)^{\Gamma} \wedge_{O(R_{r[\sigma]})^{\Gamma/K}} O(n)_{+} \right)$$

with the analogous H' action. The induced maps

$$T(W) \longrightarrow \Psi(W), \qquad \overline{T}(W) \longrightarrow \overline{\Psi}(W)$$

are then $(H' \times O(n))$ -equivariant. We prove the following theorem.

THEOREM 3.7. The $(H' \times O(n))$ -equivariant maps

$$T(W) \longrightarrow \Psi(W), \qquad \overline{T}(W) \longrightarrow \overline{\Psi}(W)$$

are isomorphisms.

PROOF. We construct an inverse isomorphism $\Psi(W) \to T(W)$ as follows. For fixed $\sigma \colon \Gamma \xrightarrow{W} Q$, let

$$T(\sigma) = (\mathcal{I}(V_{\sigma} \oplus R_{r[\sigma]}, W) \times Y_{\sigma} \times X(R_{r[\sigma]}))^{\Gamma} \times_{O(R_{r[\sigma]})^{\Gamma/K}} O(n) \subset T(W)$$

(a summand in the disjoint union), let

$$\mathcal{I}(\sigma) \subset \mathcal{I}(V \oplus \mathbb{R}^n, W) \times Y$$

denote the union of the (σ, r) -twisted M fixed points as r varies, and let

 $\Psi(\sigma) \subset \Psi(W)$

denote the subspace that projects to $\mathcal{I}(\sigma)$. The subsets $\mathcal{I}(\sigma)$ are then disjoint and they are closed since the inclusion of $\mathcal{I}(\sigma)$ in $\mathcal{I}(V \oplus \mathbb{R}^n, W) \times Y$ is the inverse image of the inclusion of the σ -twisted Γ fixed points in $\mathcal{I}(V \oplus \mathbb{R}^n, W)/O(n) \times Y$. The action of O(n) on $\mathcal{I}(V \oplus \mathbb{R}^n, W)$ restricts to each $\mathcal{I}(\sigma)$. These observations for $\mathcal{I}(\sigma)$ imply the corresponding observations for $\Psi(\sigma)$. In particular, $\Psi(W)$ is the disjoint union of $\Psi(\sigma)$, and since by construction the map

$$T(\sigma) \subset T(W) \longrightarrow \Psi(W)$$

factors through $\Psi(\sigma)$, it suffices to construct an inverse isomorphism $\Psi(\sigma) \to T(\sigma)$.

For each $(\alpha, y) \in \mathcal{I}(\sigma)$, there is a unique $r_{\alpha} \colon \Gamma/K \to O(n)$ for which α is a (σ, r_{α}) -twisted Γ fixed point (independently of $y \in Y$), and the function $\alpha \mapsto r_{\alpha}$ is a continuous map r from $\mathcal{I}(\alpha)$ to the space $\operatorname{Hom}(\Gamma/K, O(n))$ of continuous homomorphisms $\Gamma/K \to O(n)$ (with the compact-open topology). It is convenient to write $\operatorname{Hom}(\sigma)$ for the subspace of $\operatorname{Hom}(\Gamma/K, O(n))$ of elements $r \colon \Gamma/K \to O(n)$ such that $R_r \cong R_{r[\sigma]}$.

Because $V_{\sigma} \oplus R_{r_{\alpha}}$ is Γ/K -isomorphic to W, $R_{r_{\alpha}}$ is Γ/K -isomorphic to $R_{r[\sigma]}$. Choose an arbitrary Γ/K -isomorphism $f_{\alpha} \colon R_{r_{\alpha}} \to R_{r[\sigma]}$. Since $R_{r_{\alpha}}$ and $R_{r[\sigma]}$ both have underlying inner product space \mathbb{R}^n , $f_{\alpha} \in O(n)$. In general there is no way to choose the f_{α} so that the resulting function $f: \mathcal{I}(\sigma) \to O(n)$ is continuous; however, we can choose f so that it factors as

$$\mathcal{I}(\sigma) \xrightarrow{r} \operatorname{Hom}(\sigma) \xrightarrow{f} O(n)$$

and the usual compact Lie group integration argument shows that for any fixed r_0 and fixed choice of $\tilde{f}_{r_0}: R_{r_0} \cong R_{r[\sigma]}$, there exists a neighborhood U of r_0 in $\operatorname{Hom}(\sigma)$ where we can choose \tilde{f}_r to assemble to a continuous function $\tilde{f}: U \to O(n)$ extending the chosen value at r_0 .

Now consider the function $\theta: \Psi(\sigma) \to T(\sigma)$ defined by

$$\begin{aligned} \theta \colon (\alpha, y, x) &\mapsto \left((\alpha \circ (\mathrm{id}_V \oplus f_\alpha^{-1}), y, f_\alpha \cdot x), f_\alpha \right) \\ &\in \left(\mathcal{I}(V_\sigma \oplus R_{r[\sigma]}, W) \times Y_\sigma \times X(R_{r[\sigma]}) \right)^{\Gamma} \times_{O(R_{r[\sigma]})^{\Gamma/K}} O(n). \end{aligned}$$

It is not obvious that θ is continuous, but it is well defined:

- By hypothesis α is an (σ, r_{α}) -twisted Γ fixed point of $\mathcal{I}(V \oplus \mathbb{R}^{n}, W)$ and hence a Γ fixed element of $\mathcal{I}(V_{\sigma} \oplus R_{r_{\alpha}}, W)$; then $a \circ (\mathrm{id}_{V} \oplus f_{\alpha}^{-1})$ is a Γ fixed element of $\mathcal{I}(V_{\sigma} \oplus R_{r[\sigma]}, W)$ since f_{α} is Γ -equivariant.
- By hypothesis y is a (σ, r_{α}) -twisted Γ fixed point of Y and hence is a Γ fixed element of Y_{σ} .
- By hypothesis x is a (σ, r_{α}) -twisted Γ fixed point of $X(\mathbb{R}^n)$ and hence is a Γ fixed element of $X(R_{r_{\alpha}})$; then $f_{\alpha} \cdot x$ is a Γ fixed element of $X(R_{r[\sigma]})$ since f_{α} is Γ -equivariant.

Next we check that θ is continuous. If f'_{α} is another choice of Γ/K -equivariant isomorphism $R_{r_{\alpha}} \to R_{r[\sigma]}$, then $\psi := f'_{\alpha} \circ f^{-1}_{\alpha}$ is an element of $O(R_{r[\sigma]})^{\Gamma/K}$ and

$$\begin{aligned} ((\alpha \circ (\mathrm{id}_V \oplus f_\alpha^{-1}), y, f_\alpha \cdot x), f_\alpha) &= ((\alpha \circ (\mathrm{id}_V \oplus f_\alpha^{-1}), y, f_\alpha \cdot x) \cdot \psi^{-1}, \psi \circ f_\alpha) \\ &= ((\alpha \circ (\mathrm{id}_V \oplus f_\alpha^{-1} \circ \psi^{-1}), y, ((\psi \circ f_\alpha) \cdot x)), \psi \circ f_\alpha) \\ &= ((\alpha \circ (\mathrm{id}_V \oplus (f')_\alpha^{-1}), y, (f'_\alpha) \cdot x), f'_\alpha). \end{aligned}$$

Thus, the function θ is independent of the choice of f. Since we can choose f to be continuous in a neighborhood of α for any fixed α , θ is continuous in a neighborhood of every point (α, y, x) , and so θ is continuous on all of $\Psi(\sigma)$.

By inspection, θ is inverse to the original map $T(\sigma) \to \Psi(\sigma)$, and this proves the statement for $\Psi(W)$. The statement for $\bar{\Psi}(W)$ follows from the statement for $\Psi(W)$.

We can now prove Theorem 2.2.

PROOF OF THEOREM 2.2. For part (i), the previous theorem applied in the case when X = S gives

$$\operatorname{Fix}_{K}^{\Gamma}(iJ\eta)(W) = \left(\bigvee_{\sigma \colon \Gamma \stackrel{W}{\to} Q} (\mathcal{I}(V_{\sigma} \oplus R_{r[\sigma]}, W)^{\Gamma} \times Y_{\sigma}^{\Gamma})_{+} \wedge_{O(R_{r[\sigma]})^{\Gamma/K}} S^{R_{r[\sigma]}^{\Gamma/K}} \right) / H'$$
$$= \left(\bigvee_{\sigma \colon \Gamma \stackrel{W}{\to} Q} \mathcal{J}_{\Gamma/K}^{\Gamma/K}(V_{\sigma}, W) \wedge (Y_{\sigma}^{\Gamma})_{+} \right) / H'$$

and it is straightforward to check that the latter formula specifies an isomorphism functorial for $W \in \mathcal{J}_{\Gamma/K}^{\Gamma/K}$, i.e., describes $\operatorname{Fix}_{K}^{\Gamma}(iJ\eta)$ as a $\mathcal{J}_{\Gamma/K}^{\Gamma/K}$ -space. (This also

follows from the proof of Theorem V.2.4.) Since smash product commutes with colimits in each variable, this isomorphism induces an isomorphism

$$\operatorname{Fix}_{K}^{\Gamma}(iJ\eta) \wedge \operatorname{Fix}_{K}^{\Gamma}(X) \cong \left(\bigvee_{\sigma \colon \Gamma \xrightarrow{W} Q} (\mathcal{J}_{\Gamma/K}^{\Gamma/K}(V_{\sigma}, -) \wedge (Y_{\sigma}^{\Gamma})_{+}) \wedge \operatorname{Fix}_{K}^{\Gamma}(X) \right) / H'.$$

As in [23, 4.4] (or as in the proof of [4, A.1]), we get for the W space

$$(\operatorname{Fix}_{K}^{\Gamma}(iJ\eta) \wedge \operatorname{Fix}_{K}^{\Gamma}(X))(W) \\ \cong \left(\bigvee_{\sigma \colon \Gamma \xrightarrow{W} Q} (\mathcal{J}_{\Gamma/K}^{\Gamma/K}(V_{\sigma}, W) \wedge (Y_{\sigma}^{\Gamma})_{+}) \wedge_{O(R_{r[\sigma]})^{\Gamma/K}} X(R_{r[\sigma]})^{\Gamma} \right) / H' \\ \cong \bar{T}(W) / (H' \times O(n))$$

with the universal map

$$(\operatorname{Fix}_{K}^{\Gamma}(\iota J\eta))(W') \wedge (\operatorname{Fix}_{K}^{\Gamma}(X))(W'') \longrightarrow \overline{T}(W)/(H' \times O(n))$$
 for $W = W' \oplus W''$ induced by the inclusion of W' in W

$$\mathcal{J}_{\Gamma/K}^{\Gamma/K}(V_{\sigma}, W') \wedge (Y_{\sigma}^{\Gamma})_{+} \cong (\mathcal{I}(V_{\sigma} \oplus R, W') \times Y_{\sigma})_{+}^{\Gamma} \wedge_{O(R)^{\Gamma/K}} S^{R^{\Gamma/K}} \longrightarrow (\mathcal{I}(V_{\sigma} \oplus R \oplus W'', W) \times Y_{\sigma})_{+}^{\Gamma} \wedge_{O(R)^{\Gamma/K}} S^{R^{\Gamma/K}}$$

in the $J\eta$ factor and the left suspension map

$$S^{R^{\Gamma/K}} \wedge X(W'')^{\Gamma} \longrightarrow X(R \oplus W'')^{\mathrm{I}}$$

in the X factor. (To compare notation, note that $R_{r[\sigma]} \cong R \oplus W''$ since they both equivariantly complement V_{σ} in W.) The previous theorem (combined with (3.2) and (3.3)) gives an isomorphism

$$\overline{T}(W)/(O(n) \times H') \longrightarrow (\operatorname{Fix}_{K}^{\Gamma}(\imath J\eta \wedge X))(W)$$

and by inspection, the composite

$$(\operatorname{Fix}_{K}^{\Gamma}(\imath J\eta))(W') \wedge (\operatorname{Fix}_{K}^{\Gamma}X)(W'') \longrightarrow \overline{T}(W)/(H' \times O(n)) \longrightarrow (\operatorname{Fix}_{K}^{\Gamma}(\imath J\eta \wedge X))(W)$$

is the universal map for the natural transformation

$$\operatorname{Fix}_{K}^{\Gamma}(\imath J\eta) \wedge \operatorname{Fix}_{K}^{\Gamma}(X) \longrightarrow \operatorname{Fix}_{K}^{\Gamma}(\imath J\eta \wedge X).$$

This proves part (i).

For part (ii), the previous theorem identifies the map

$$(\operatorname{Fix}_{K}^{\Gamma}(\imath J(EF \times \eta) \wedge X))(W) \longrightarrow (\operatorname{Fix}_{K}^{\Gamma}(\imath J\eta \wedge X))(W)$$

as the map

$$\left(\bigvee_{\sigma: \ \Gamma \stackrel{W}{\to} Q} ((\mathcal{I}(V_{\sigma} \oplus R_{r[\sigma]}, W)^{\Gamma} \times (EF \times Y)^{\Gamma}_{\sigma})_{+} \wedge_{O(R_{r[\sigma]})^{\Gamma/K}} X(R_{r[\sigma]})^{\Gamma})\right) / H' \longrightarrow \left(\bigvee_{\sigma: \ \Gamma \stackrel{W}{\to} Q} ((\mathcal{I}(V_{\sigma} \oplus R_{r[\sigma]}, W)^{\Gamma} \times Y^{\Gamma}_{\sigma})_{+} \wedge_{O(R_{r[\sigma]})^{\Gamma/K}} X(R_{r[\sigma]})^{\Gamma})\right) / H'$$

For any homomorphism $\sigma: \Gamma \to Q$, the graph subgroup $\Gamma_{\sigma} < \Gamma \times Q^{\text{op}}$ is in the family $\mathcal{F} = \mathcal{F}(1 \times Q^{\text{op}})$ and so the map $(EF \times Y)^{\Gamma}_{\sigma} \to Y^{\Gamma}_{\sigma}$ is a homotopy equivalence. Let O_1, \ldots, O_q be the set of orbits for the action of H' on the set of W-compatible M graph subgroup homomorphisms $\Gamma \xrightarrow{W} Q$, and for each i, choose a representative $\sigma_i \in O_i$. Because $H' \times O(n)$ acts freely on $\mathcal{I}(V \oplus \mathbb{R}^n, W) \times Y$, for each $i, H' \times (O(R_{r[\sigma_i]}))^{\Gamma/K}$ acts freely on

$$\prod_{\sigma \in O_i} \mathcal{I}(V_\sigma \oplus R_{r[\sigma]}, W)^{\Gamma} \times Y_{\sigma}^{\Gamma} \subset \mathcal{I}(V \oplus \mathbb{R}^n, W) \times Y$$

so for each i, the map

$$\coprod_{\sigma \in O_i} \mathcal{I}(V_{\sigma} \oplus R_{r[\sigma]}, W)^{\Gamma} \times (EF \times Y)^{\Gamma}_{\sigma} \longrightarrow \coprod_{\sigma \in O_i} \mathcal{I}(V_{\sigma} \oplus R_{r[\sigma]}, W)^{\Gamma} \times Y^{\Gamma}_{\sigma}$$

is an $H' \times (O(R_{r[\sigma]})^{\Gamma/K})$ -equivariant homotopy equivalence. The maps

$$\left(\bigvee_{\sigma\in O_{i}} \left(\mathcal{I}(V_{\sigma}\oplus R_{r[\sigma]}, W)^{\Gamma} \times (EF \times Y)_{\sigma}^{\Gamma}\right)_{+} \wedge_{O(R_{r[\sigma]})^{\Gamma/K}} X(R_{r[\sigma]})^{\Gamma}\right) / H' \\ \longrightarrow \left(\bigvee_{\sigma\in O_{i}} \left(\mathcal{I}(V_{\sigma}\oplus R_{r[\sigma]}, W)^{\Gamma} \times Y_{\sigma}^{\Gamma}\right)_{+} \wedge_{O(R_{r[\sigma]})^{\Gamma/K}} X(R_{r[\sigma]})^{\Gamma}\right) / H'$$

are therefore based homotopy equivalences. Taking a wedge over i, we see that the map

$$\operatorname{Fix}_{K}^{\Gamma}(iJ(EF \times \eta) \wedge X) \longrightarrow \operatorname{Fix}_{K}^{\Gamma}(iJ\eta \wedge X)$$

induces a based homotopy equivalence on W spaces for all W. This proves part (ii). $\hfill \Box$

4. Proof of Theorem V.2.8

The strategy for the proof of Theorem V.2.8 is an indirect approach using Proposition 4.1 below, which gives criteria to identify when symmetric powers preserve a weak equivalence out of a cofibrant object.

The criterion involves indexed smash powers and HHR diagonal maps, which are reviewed and slightly generalized in Section 7. In the current context, for G a compact Lie group, Y a G-spectrum, and S a finite G-set, we can form the indexed smash power $Y^{(S)}$ which gets its G-spectrum structure both from Y and the Gaction on S. When we decompose S as a disjoint union of orbits

$$S \cong G/H_1 \amalg \cdots \amalg G/H_r,$$

we then have an *HHR diagonal map* [15, B.209] (or Proposition 7.6) of non-equivariant spectra

$$\Phi^{H_1}Y\wedge\cdots\wedge\Phi^{H_r}Y\longrightarrow\Phi^G(Y^{(S)}).$$

We remark that the map depends on the choice of isomorphism, that is, the choice of the elements of S that correspond to the identity cosets eH_1, \ldots, eH_r : the map is analogous to and induced by the diagonal map for based G-spaces

$$A^{H_1} \wedge \dots \wedge A^{H_r} \longrightarrow (A^{(S)})^G$$

that sends $a_1 \wedge \cdots \wedge a_r$ to the element of the smash product $A^{(S)}$ that is given by $g_{\zeta}a_i$ in the $\zeta \in S$ coordinate, where ζ corresponds to the element $g_{\zeta}H_i$ in the identification of S as a disjoint union of orbits above.

In the statement of the following proposition, for a finite group Σ , we write $E_G \Sigma$ for the universal space for the family of subgroups of $G \times \Sigma$ whose intersection with $1 \times \Sigma$ is trivial. It is a $(G \times \Sigma)$ -CW complex whose fixed points are contractible for subgroups in the family and empty for subgroups not in the family.

PROPOSITION 4.1. Let G be a compact Lie group, let k > 1, and let $\Sigma < \Sigma_k$. Let Y be a G-spectrum that satisfies the following conditions:

- (i) The quotient map $E_G \Sigma_+ \wedge_{\Sigma} Y^{(k)} \to Y^{(k)} / \Sigma$ is a weak equivalence.
- (ii) For every M < G and every M-set $S = M/H_1 \amalg \cdots \amalg M/H_r$ of cardinality k, the map

$$L\Phi^{H_1}Y\wedge^L\cdots\wedge^L L\Phi^{H_r}Y\longrightarrow\Phi^{H_1}Y\wedge\cdots\wedge\Phi^{H_r}Y\longrightarrow\Phi^M(Y^{(S)})$$

in the non-equivariant stable category is an isomorphism.

Then for any cofibrant approximation $X \to Y$ in the positive complete model structure on G-spectra, the induced map $X^{(k)}/\Sigma \to Y^{(k)}/\Sigma$ is a weak equivalence.

PROOF. Let $X \to Y$ be a cofibrant approximation in the positive complete model structure on G-spectra. In the diagram

the top horizontal map is a weak equivalence by [15, B.116] and the bottom horizontal map is a weak equivalence by hypothesis (i). We want to show that the right horizontal map is a weak equivalence, so it suffices to show that the left horizontal map is a weak equivalence. The $(G \times \Sigma)$ -CW complex $E_G \Sigma$ is built by cells of the form $(G \times \Sigma)/\tilde{H} \times (D^n, \partial D^n)$ where $\tilde{H} \cap (1 \times \Sigma)$ is trivial, and working a cell at a time, it suffices to show that the map

$$(((G \times \Sigma)/\tilde{H}) \wedge X^{(k)})/\Sigma \longrightarrow (((G \times \Sigma)/\tilde{H}) \wedge Y^{(k)})/\Sigma$$

is a weak equivalence for every such subgroup \tilde{H} . Let H be the image of \tilde{H} in Gunder the projection. Because $\tilde{H} \cap (1 \times \Sigma)$ is trivial, the projection is an isomorphism $\tilde{H} \to H$ and there exists a homomorphism $\sigma \colon H \to \Sigma$ with \tilde{H} the graph of this homomorphism:

$$\tilde{H} = H_{\sigma} = \{(h, \sigma(h)) \mid h \in H\} < G \times \Sigma.$$

Let $S = \{1, ..., k\}$ be the *H*-set obtained using σ to define the action. We then have a commutative diagram

$$\begin{array}{ccc} (((G \times \Sigma)/\tilde{H}) \wedge X^{(k)})/\Sigma & \xrightarrow{\cong} & G_{+} \wedge_{H} X^{(S)} \\ & \downarrow & & \downarrow \\ (((G \times \Sigma)/\tilde{H}) \wedge Y^{(k)})/\Sigma & \xrightarrow{\cong} & G_{+} \wedge_{H} Y^{(S)} \end{array}$$

with the horizontal maps isomorphisms (see, for example, Proposition 7.1 with H playing the role of G there). We are now reduced to showing that the map of H-spectra

$$X^{(S)} \longrightarrow Y^{(S)}$$

is a weak equivalence. By [24, XVI.6.4] (or Proposition 1.5), it is now enough to show that for every M < H, the map $L\Phi^M(X^{(S)}) \to L\Phi^M(Y^{(S)})$ is a nonequivariant weak equivalence, and this follows from hypothesis (ii). We now start the argument for Theorem V.2.8, using the notational convention of Notation V.2.1: Γ is a compact Lie group with a closed normal subgroups K, Λ satisfying $K < \Lambda$. We treat the case k > 1, noting that in the statement of Theorem V.2.8, there is nothing to show in the case k = 1. The following proposition gives the first step.

PROPOSITION 4.2. Let η be a Γ/K -compatible Q-faithful (Γ, Q) vector bundle where each fiber is positive dimensional. Then for every $\Sigma < \Sigma_k$, the map

$$E_{\Gamma/K}\Sigma_+ \wedge_{\Sigma} (J^{\Gamma/K}\eta)^{(k)} \longrightarrow (J^{\Gamma/K}\eta)^{(k)}/\Sigma$$

is a level equivalence.

PROOF. By Proposition V.3.2, $\operatorname{Sym}^k \eta$ is $(\Sigma_k \wr Q)$ -faithful. In particular, the Σ_k action on

$$(J^{\Gamma/K}\eta)^{(k)}(V) \cong (J^{\Gamma/K}\eta^k)(V) = \mathcal{J}(\eta^k, V)/Q^k$$

is free for every K-trivial orthogonal Γ -representation V. We get that

$$E_{\Gamma/K}\Sigma_+ \wedge \mathcal{J}(\eta^k, V)/Q^k \longrightarrow \mathcal{J}(\eta^k, V)/Q^k$$

is a $(\Gamma/K \times \Sigma)$ -equivalence; taking quotients by Σ , the map

$$E_{\Gamma/K}\Sigma_+ \wedge_{\Sigma} \mathcal{J}(\eta^k, V)/Q^k \longrightarrow (\mathcal{J}(\eta^k, V)/Q^k)/\Sigma$$

is a Γ/K -equivalence.

Next we study the geometric fixed points $\Phi^{\Gamma/K}$ of $(J^{\Gamma/K}\eta)^{(S)}$ for finite Γ/K sets S. As in Section 2, we need the extra flexibility of considering Γ actions which are not K fixed and the appropriate context for this is the category of Γ -spectra indexed on K-trivial spectra, or in other words, Γ -equivariant $\mathcal{J}_{\Gamma/K}$ -spaces. As in that section, we use i for the functor from Γ -spectra to Γ -equivariant $\mathcal{J}_{\Gamma/K}$ -spaces that restricts to the K-trivial orthogonal Γ -representations, and j for the functor from Γ/K -spectra to Γ -equivariant $\mathcal{J}_{\Gamma/K}$ -spaces that regards a Γ/K action as a Γ action. For $\eta \in \Gamma/K$ -compatible (Γ, Q) vector bundle, we then have

$$iJ\eta = jJ^{\Gamma/K}\eta.$$

Section 7 reviews the construction of the indexed smash product and HHR diagonal in this context. To interface with the terminology and notation there, we let U be a complete Γ/K -universe, regarded as a Γ -universe (a K-trivial Γ -universe containing a copy of every K-trivial orthogonal G-representation). While we can define the indexed smash product for any finite Γ -set, we will restrict to considering finite Γ/K -sets. Then when X is a Γ/K -spectrum, we have a natural isomorphism

$$(jX)^{(S)} \cong j(X^{(S)}).$$

For generalized orbit desuspension spectra $J\eta$, using the construction of indexed smash powers in Proposition 7.1, we can identify $(\imath J\eta)^{(S)}$ as $\imath J(\eta^{(S)})$ with $\eta^{(S)}$ defined as follows.

DEFINITION 4.3. Let S be a finite Γ/K -set and let η be a (Γ, Q) vector bundle. Write |S| for the underlying set of S. Let $\Sigma_{|S|}$ denote the group of automorphisms of the set |S|, let $\sigma_S \colon \Gamma \to \Sigma_{|S|}$ be the tautological homomorphism for the Γ action on S, and let $\Sigma_{|S|}(\sigma_S)$ be the $(\Gamma \times \Sigma_{|S|}^{op})$ -set of Proposition 7.1, given by $\Sigma_{|S|}$ with

its natural right action and left Γ action induced by σ_S . Define $\eta^{(S)}$ to be the $(\Gamma, \Sigma_{|S|} \wr Q)$ vector bundle

$$\eta^{(S)} = \Sigma_{|S|}(\sigma_S) \times \eta^{|S|}$$

where the left Γ action is the diagonal of the left Γ action on $\Sigma_{|S|}(\sigma_S)$ and the action of Γ on each copy of η coordinatewise; the right action of $\Sigma_{|S|} \wr Q$ uses the right action of $\Sigma_{|S|}$ on $\Sigma_{|S|}(\sigma_S)$ (with trivial $Q^{|S|}$ action), the action of $\Sigma_{|S|}$ on $\eta^{|S|}$ that permutes the coordinates, and the product action of $Q^{|S|}$ on $\eta^{|S|}$.

As a consequence of Proposition V.1.6 and Proposition 7.1, we get the following observation.

PROPOSITION 4.4. Let S be a finite Γ/K -set and let η be a (Γ, Q) vector bundle. Then $(iJ\eta)^{(S)} \cong iJ(\eta^{(S)})$.

We use this in the following proposition, which allows us to study the indexed smash power of a generalized orbit desuspension spectrum $J^{\Gamma/K}\eta$ in terms of the indexed smash power of $iJ\eta'$ where we have taken an equivariant CW approximation of the base space. This kind of CW approximation does not preserve Γ/K -compatibility and is the reason we consider generalizations to Γ -equivariant $\mathcal{J}_{\Gamma/K}$ -spaces. In the statement, $\mathcal{F}(1 \times Q^{\mathrm{op}})$ denotes the family of subgroups of $\Gamma \times Q^{\mathrm{op}}$ whose intersection with $1 \times Q^{\mathrm{op}}$ is trivial.

PROPOSITION 4.5. Let η be a Q-faithful (Γ, Q) vector bundle, let $f: B' \to B$ be an $\mathcal{F}(1 \times Q^{\mathrm{op}})$ -equivalence of $(\Gamma \times Q^{\mathrm{op}})$ -spaces. Assume that B' is Hausdorff and let $f^*\eta$ denote the pullback (Γ, Q) -vector bundle with base space B'. Then:

(i) $f^*\eta$ is Q-faithful.

 σ

- (ii) The induced map $J(f^*\eta) \to J\eta$ is a level equivalence.
- (iii) For any finite Γ/K -set S, the induced map $(iJ(f^*\eta))^{(S)} \to (iJ\eta)^{(S)}$ is a level equivalence.

PROOF. Conclusion (i) is clear from the fiberwise definition of Q-faithful. For conclusion (ii), we need to show that for an arbitrary orthogonal Γ -representation W and an arbitrary closed subgroup $H < \Gamma$, the map

$$(\mathcal{J}(f^*\eta, W)/Q)^H \longrightarrow (\mathcal{J}(\eta, W)/Q)^H$$

is a (non-equivariant) weak equivalence. By Lemma 1.2, this is the Q-quotient of the induced map on Thom spaces of the map of vector bundles (of the same local dimension) which on total spaces is

$$\prod_{H \to Q} \Im^{\perp} \mathcal{I}(f^*\eta, W)^{H, \sigma} \longrightarrow \prod_{\sigma \colon H \to Q} \Im^{\perp} \mathcal{I}(\eta, W)^{H, \sigma}$$

Since each side is a Q-stable subspace of a free Q-space (Proposition 1.1), the Q actions are free. Since these are Hausdorff spaces, to see that the map induces a weak equivalence on Q-quotients, it is enough to see that the map itself is a weak equivalence, and for this it is enough to see that for each $\sigma: H \to Q$, the map

$$\mathcal{I}(f^*\eta, W)^{H,\sigma} \longrightarrow \mathcal{I}(\eta, W)^{H,\sigma}$$

is a weak equivalence. Applying Lemma 1.2 again, the previous map is the map on total spaces of a map of locally trivial fiber bundles which is the restriction $f^{H,\sigma}$ of f on the base spaces

$$f^{H,\sigma}\colon (B')^{H,\sigma}\longrightarrow B^{H,\sigma}$$

(with the identity map on the fibers). The σ -twisted H fixed points are the fixed points of the subgroup

$$(H,\sigma) = \{(h,\sigma(h^{-1})) \mid h \in H\} < \Gamma \times Q^{\mathrm{op}}$$

and hence by hypothesis, $f^{H,\sigma}$ is a weak equivalence. This establishes (ii). To prove (iii), we apply Proposition 4.4 and observe that $(f^*\eta)^{(S)}$ is obtained from $\eta^{(S)}$ as the pullback bundle of the $(\Sigma_{|S|} \wr Q)$ -equivariant map of base spaces

$$\Sigma_{|S|}(\sigma_S) \times (B')^{|S|} \longrightarrow \Sigma_{|S|}(\sigma_S) \times B^{|S|}.$$

Because $B' \to B$ is assumed to be an $\mathcal{F}(1 \times Q^{\mathrm{op}})$ -equivalence, the displayed map is a $\mathcal{F}(1 \times (\Sigma_{|S|} \wr Q)^{\mathrm{op}})$ -equivalence: for any space X, the fixed points of X^k for graph subgroups of $\Sigma_k \wr Q$ can be written in terms of products of fixed points of X for subgroups of Q (see, for example, Lemma 6.1). Next we observe that $\eta^{(S)}$ is $(\Sigma_{|S|} \wr Q)$ -faithful: this is clear when Q is the trivial group since $\Sigma_{|S|}$ acts freely on $\Sigma_{|S|}(\sigma_S)$ and is covered by Proposition V.3.2 when Q is non-trivial. Conclusion (ii) now implies conclusion (iii).

We defined the functor $\operatorname{Fix}_{K}^{\Gamma}$ from Γ -equivariant $\mathcal{J}_{\Gamma/K}$ -spaces to non-equivariant $\mathcal{J}_{\Gamma/K}^{\Gamma}$ -spaces in Definition 2.1. Although not needed in that section, left Kan extension along the functor $\phi: \mathcal{J}_{\Gamma/K}^{\Gamma} \to \mathcal{J}$ produces a functor Φ_{K}^{Γ} from Γ -equivariant $\mathcal{J}_{\Gamma/K}$ -spaces to non-equivariant spectra; the functor Φ_{K}^{Γ} ; it is canonically naturally isomorphic to the functor denoted Φ_{U}^{Γ} in Section 7 for the universe U described above. More generally, for any closed subgroup H of Γ containing K, we likewise have a functor $\Phi_{K}^{H} = \Phi_{U}^{H}$ defined using the functor $\operatorname{Fix}_{K}^{H}$ from $N_{\Gamma}H$ -equivariant $\mathcal{J}_{N_{\Gamma}H/K}$ -spaces to non-equivariant $\mathcal{J}_{N_{\Gamma}H/K}^{H}$ -spaces by the formula

$$(\operatorname{Fix}_{K}^{H} Y)(W) = Y(W)^{H}$$

and left Kan extension along the functor $\phi: \mathcal{J}_{N_{\Gamma}H/K}^{H} \to \mathcal{J}$ defined as usual (on objects it takes V to V^{H} regarded as a non-equivariant inner product space, and on maps it takes the restriction of the equivariant isometry to its fixed points and sends the embedding complement element by the identity). The functors Φ_{K}^{H} are instances of the geometric fixed point functors of [22, V§4] when we identify Γ -equivariant $\mathcal{J}_{\Gamma/K}$ -spaces as Γ -equivariant orthogonal spectra indexed on the Ktrivial orthogonal Γ -representations. The work in *ibid*. does not restrict to the complete universe and the work there establishes various point-set and homotopical properties of these functors, including the existence of left derived functors, computed by applying the point-set functors to cofibrant approximations (in the standard stable model structure).

For Γ/K -spectra, we have natural isomorphisms relating $\operatorname{Fix}_{K}^{H}$ and Φ_{K}^{H} to the underlying non-equivariant spectra of $\operatorname{Fix}^{H/K}$ and $\Phi^{H/K}$ which are essentially the identity map. As reviewed in Section 7, the construction of the HHR diagonal extends to the case of Γ -equivariant $\mathcal{J}_{\Gamma/K}$ -spaces to produce a natural diagonal map

$$\Phi_K^H(Y) \longrightarrow \Phi_K^{\Gamma}(Y^{(\Gamma/H)}),$$

and for X a Γ/K -spectrum, the diagram of comparison isomorphisms

$$\begin{array}{ccc}
\Phi_{K}^{H}(jX) \longrightarrow \Phi_{K}^{\Gamma}((jX)^{(\Gamma/H)}) \\
\cong & \downarrow & \downarrow \cong \\
\Phi^{H/K}(X) \longrightarrow \Phi^{\Gamma/K}(X^{(\Gamma/H)})
\end{array}$$

commutes. The following proposition generalizes [15, B.209] in the current context.

THEOREM 4.6. Let $\eta: E \to B$ be a (Γ, Q) vector bundle such that B is a $(\Gamma \times Q^{\text{op}})$ -equivariant CW complex with isotropy groups in $F = \mathcal{F}(1 \times Q^{\text{op}})$. Then for any closed subgroup H of Γ containing K, the diagonal map

$$\Phi_K^H i J \eta \longrightarrow \Phi_K^\Gamma((i J \eta)^{(\Gamma/H)})$$

is an isomorphism.

PROOF. By hypothesis, the base B is built out of cells of the form

$$(\Gamma \times Q^{\mathrm{op}})/M_{\sigma} \times (D^n, \partial D^n)$$

for a M_{σ} a subgroup of $\Gamma \times Q^{\text{op}}$ in the family F. Restricted to such a cell, the fiber $V = E_b$ at a point has an action of M_{σ} and a trivialization giving an isomorphism

$$\eta|_{D^n} \cong ((\Gamma \times Q^{\operatorname{op}}) \times_{M_{\sigma}} V \longrightarrow (\Gamma \times Q^{\operatorname{op}})/M_{\sigma}) \times D^n.$$

We then see that η is built out of cells of the form

$$((\Gamma \times Q^{\mathrm{op}})_+ \wedge_{M_{\sigma}} \mathcal{J}(V, -))/Q^{\mathrm{op}} \wedge (D^n, \partial D^n)_+$$

where M_{σ} is in F, and V is an orthogonal M_{σ} -representation. Since M_{σ} is in the family F, letting M denote the image of M_{σ} under the projection to Γ , we have

$$M_{\sigma} = \{(h, \sigma(h)^{-1})) \mid h \in M\}$$

for a unique homomorphism $\sigma: M \to Q$. Using the projection isomorphism $M_{\sigma} \to M$ to regard V as an orthogonal M-representation, we get an isomorphism of Γ -spectra

$$((\Gamma \times Q^{\mathrm{op}})_+ \wedge_{M_{\sigma}} \mathcal{J}(V, -))/Q^{\mathrm{op}} \cong \Gamma_+ \wedge_M \mathcal{J}(V, -).$$

This shows that $J\eta$ is cofibrant in the complete model structure on Γ -spectra; the result here is now a special case of Corollary 7.8.

Finally, we prove Theorem V.2.8.

PROOF OF THEOREM V.2.8. Let $\eta: E \to B$ be a Q-faithful Γ/K -compatible (Γ, Q) vector bundle where each fiber is positive dimensional. To prove the statement of Theorem V.2.8, we just need to check that $J^{\Gamma/K}\eta$ satisfies hypotheses (i) and (ii) of Proposition 4.1 for every k > 1 and every $\Sigma < \Sigma_k$. Proposition 4.2 establishes hypothesis (i).

For hypothesis (ii), by restricting the group of equivariance and changing notation if necessary, it suffices to consider finite Γ/K -sets S. Let $f: B' \to B$ be an $\mathcal{F}(1 \times Q^{\mathrm{op}})$ -colocal CW-approximation, i.e., an $\mathcal{F}(1 \times Q^{\mathrm{op}})$ -equivalence from a CW complex with cells whose isotropy groups are in $\mathcal{F}(1 \times Q^{\mathrm{op}})$. By Proposition 4.5, $J(f^*\eta) \to J\eta$ induces a weak equivalence of Γ -equivariant $\mathcal{J}_{\Gamma/K}$ -spaces

$$(iJ(f^*\eta))^{(S)} \longrightarrow (iJ\eta)^{(S)}$$

for all finite Γ/K -sets S. Proposition 4.5 also asserts that $J(f^*\eta) \to J\eta$ is a level equivalence, and so induces non-equivariant weak equivalences

$$\operatorname{Fix}_{K}^{H} iJ(f^{*}\eta) \longrightarrow \operatorname{Fix}_{K}^{H} iJ\eta$$

for all closed subgroups $H < \Gamma$ containing K. This gives a commuting diagram in the stable category of Γ -spectra indexed on K-trivial representations

for $S = \Gamma/H_1 \amalg \cdots \amalg \Gamma/H_r$, with the indicated arrows isomorphisms. Noting that the functor j is a left Quillen adjoint (for the standard stable model structures of [**22**, III§4]) and that it preserves and reflects weak equivalences, the isomorphism $\Phi^{\Gamma/K} \cong \Phi_K^{\Gamma} j$ induces an isomorphism of derived functors

$$L\Phi^{\Gamma/K} \cong (L\Phi^{\Gamma}_{K}) \circ j$$

In particular, the bottom map in the square above is an isomorphism in the stable category of Γ -spectra indexed on K-trivial representations if and only if the map

$$L\Phi^{H_1/K}(J^{\Gamma/K}\eta) \wedge^L \dots \wedge^L L\Phi^{H_r/K}(J^{\Gamma/K}\eta) \longrightarrow (J^{\Gamma/K}\eta)^{(S)}$$

is an isomorphism is the Γ/K -equivariant stable category. Thus, we are reduced to showing that the map

$$L\Phi_K^{H_1}(\imath J(f^*\eta)) \wedge^L \dots \wedge^L L\Phi_K^{H_r}(\imath J(f^*\eta)) \longrightarrow (\imath J(f^*\eta))^{(S)}$$

is an isomorphism in the stable category of $\Gamma\mbox{-spectra}$ indexed on $K\mbox{-trivial}$ representations.

By Theorem 2.4, each $iJ(f^*\eta)$ is flat for the smash product of Γ -equivariant $\mathcal{J}_{\Gamma/K}$ -spaces, which implies that its smash product with any Γ -equivariant $\mathcal{J}_{\Gamma/K}$ -space represents the derived smash product. Since

$$(iJ(f^*\eta))^{(S\amalg S')} \cong (iJ(f^*\eta))^{(S)} \wedge (iJ(f^*\eta))^{(S')}$$

it suffices to consider just the case when $S = \Gamma/H$. Although $iJ(f^*\eta)$ is generally not a cofibrant Γ -equivariant $\mathcal{J}_{\Gamma/K}$ -space, by Proposition 2.3, $\operatorname{Fix}_{K}^{H}(iJ(f^*\eta))$ is a cofibrant non-equivariant $\mathcal{J}_{\Gamma/K}^{H}$ -space and this is enough to ensure that the point-set functor Φ_{K}^{H} represents the derived functor $L\Phi_{K}^{H}$. The argument is now completed by Theorem 4.6.

5. Proof of Theorem V.3.1

This section proves Theorem V.3.1 and along the way provides more details about the spectra in the decomposition given there. It is independent of the quotient fixed point lemma, and could have been placed just after Section V.3: it uses only the statement of Theorem V.2.4 and not the techniques of its argument.

To compress notation, we write $\Sigma'_q := \Sigma_q \wr Q$. The key idea in the construction is that homomorphisms $\Lambda \to \Sigma'_q$ generalize the theory of finite Λ -sets. When Q is the

trivial group, a homomorphism $\sigma \colon \Lambda \to \Sigma_q$ specifies an action of Λ on $\{1, \ldots, q\}$; if we denote the corresponding Λ -set as $S(\sigma)$, then the (left) action by an element $s^{-1} \in \Sigma_q$ specifies an isomorphism of Λ -sets $S(\sigma) \to S(\sigma_s)$, where (as in (V.2.12)) σ_s denotes the homomorphism obtain by right conjugation by s,

$$\sigma_s \colon \lambda \mapsto s^{-1} \sigma(\lambda) s.$$

When conjugation by s fixes σ , we get an automorphism of $S(\sigma)$. The corresponding theory when Q is non-trivial goes as follows.

Given a homomorphism $\sigma: \Lambda \to \Sigma'_q$, we call q the *cardinality* of σ . We say that homomorphisms σ and σ' of the same cardinality q are *conjugate* when they are conjugate under the Σ'_q action on the set of all homomorphisms $\Lambda \to \Sigma'_q$; homomorphisms of different cardinality are never conjugate.

Composing with the projection homomorphism $\Sigma'_q \to \Sigma_q$, a homomorphism $\sigma \colon \Lambda \to \Sigma'_q$ gives a homomorphism $\Lambda \to \Sigma_q$, which defines an action on $\{1, \ldots, q\}$. We say that σ is *irreducible* when the action on $\{1, \ldots, q\}$ is transitive. Given any homomorphisms $\sigma_i \colon \Lambda \to \Sigma'_{q_i}$, $i = 1, \ldots, r$, we get a homomorphism $\sigma \colon \Lambda \to \Sigma'_q$, $q = q_1 + \cdots + q_r$ by taking the block sum; we then write $\sigma = \sigma_1 \oplus \cdots \oplus \sigma_r$ and call $\sigma_1, \ldots, \sigma_r$ a block sum decomposition of σ .

An arbitrary homomorphism $\sigma \colon \Lambda \to \Sigma'_q$ is conjugate to a homomorphism where the orbits O_1, \ldots, O_k consists of intervals in $\{1, \ldots, q\}$. When the orbits are intervals, then σ has a block sum decomposition by irreducible homomorphisms $\sigma_i \to \Sigma'_{q_i}, i = 1, \ldots, k$, with $q_i = \#O_i$.

For $\sigma: \Lambda \to \Sigma'_q$, we write $C(\sigma)$ for the subgroup of Σ'_q whose conjugation action on σ is trivial. The number of distinct conjugates of σ is then $\#(\Sigma'_q/C(\sigma))$. If $\sigma: \Lambda \to \Sigma'_q$ is irreducible then any element in $C(\sigma^{\oplus n}) < \Sigma'_{nq}$ must have image in Σ_{nq} that permutes the blocks

$$\{1, \ldots, q\}, \{q+1, \ldots, 2q\}, \ldots, \{(n-1)q+1, \ldots, nq\}$$

Numbering these blocks in the obvious way and using the induced isomorphism between Σ_n and the subgroup of Σ_{nq} that permutes these blocks, we get an isomorphism

$$\Sigma_n \wr C(\sigma) \longrightarrow C(\sigma^{\oplus n}) < \Sigma'_{nq}.$$

If $\sigma_i \colon \Lambda \to \Sigma'_{q_i}$, $i = 1, \ldots, r$ are indecomposable and pairwise non-conjugate, then the isomorphism above and block sum induces an isomorphism

(5.1)
$$(\Sigma_{n_1} \wr C(\sigma_1)) \times \cdots \times (\Sigma_{n_r} \wr C(\sigma_r)) \longrightarrow C(\sigma_1^{\oplus n_1} \oplus \cdots \oplus \sigma_r^{\oplus n_r}).$$

When $q > \#(\pi_0 \Lambda)$, no homomorphism of cardinality q can be irreducible and so there are only finitely many total irreducible homomorphisms of any cardinality. Let T be the number of conjugacy classes of irreducible homomorphisms; we choose one in each conjugacy class and enumerate them τ_1, \ldots, τ_T . Let t_i denote the cardinality of τ_i , and let $\vec{t} = (t_1, \ldots, t_T) \in \mathbb{N}^T$. For $\vec{n} = (n_1, \ldots, n_T)$, define

$$\tau(\vec{n}) := \tau_1^{\oplus n_1} \oplus \dots \oplus \tau_T^{\oplus n_T} \colon \Lambda \longrightarrow \Sigma'_{\vec{n} \cdot \vec{t}}$$

when at least one of the n_i is non-zero. The cardinality of $\tau(\vec{n})$ is then given by the vector dot product $\vec{n} \cdot \vec{t}$. Let

$$\eta[\vec{n}] \colon (E^{\vec{n}\cdot\vec{t}})^{\Lambda,\tau(\vec{n})}\{K\} \longrightarrow (B^{\vec{n}\cdot\vec{t}})^{\Lambda,\tau(\vec{n})}\{K\}$$

be the $(\Gamma, C(\tau(\vec{n})))$ vector bundle obtained by raising η to the power of the cardinality of $\tau(\vec{n})$, taking the $\tau(\vec{n})$ -twisted Λ fixed points, and restricting the base to those points for which the $\tau(\vec{n})|_{K}$ -twisted K action on the fiber is trivial. (The $C(\tau(\vec{n}))$ action is the restriction to the fixed points of the $C(\tau(\vec{n})) < \Sigma'_{\vec{n}\cdot\vec{t}}$ action on $\eta^{\vec{n}\cdot\vec{t}}$.) When $q = \vec{n}\cdot\vec{t}$ and we write $(\text{Sym}^{q}\eta)(\Lambda|K)|_{C(\tau(\vec{n}))}$ for the restriction of $(\text{Sym}^{q}\eta)(\Lambda|K)$ to a $(\Gamma, C(\tau(\vec{n})))$ vector bundle, $\eta[\vec{n}]$ includes as a $(\Gamma, C(\tau(\vec{n})))$ vector bundle in $(\text{Sym}^{q}\eta)(\Lambda|K)|_{C(\tau(\vec{n}))}$ as the summand indexed by $\tau(\vec{n})$. Let $I\eta[\vec{n}]$ be the (Γ, Σ'_{q}) vector bundle

$$I\eta[\vec{n}] := (\eta[\vec{n}]) \times_{C(\tau(\vec{n}))} \Sigma'_q.$$

Then formally, we get a map of (Γ, Σ'_q) vector bundles $I\eta[\vec{n}] \to (\text{Sym}^q \eta)(\Lambda|K)$. The summand of $I\eta[\vec{n}]$ indexed by the coset $C(\tau(\vec{n}))s$ maps to the summand of $(\text{Sym}^q \eta)(\Lambda|K)$ indexed by $\tau(\vec{n})_s$. Since the cosets are in bijective correspondence with the conjugates of $\tau(\vec{n})$, the map $I\eta[\vec{n}] \to (\text{Sym}^q \eta)(\Lambda|K)$ is an inclusion. Because every $\sigma \colon \Lambda \to \Sigma'_q$ is conjugate to $\tau(\vec{n})$ for some (unique) \vec{n} with $\vec{n} \cdot \vec{t} = q$, we get the following result.

PROPOSITION 5.2. For $q \geq 1$, the map of (Γ, Σ'_q) vector bundles

$$\coprod_{\vec{n}\cdot\vec{t}=q} I\eta[\vec{n}] \longrightarrow (\operatorname{Sym}^q \eta)(\Lambda|K)$$

is an isomorphism.

Because $J_{(\Gamma,C(\tau(\vec{n})))}(\eta[\vec{n}]) \cong J_{(\Gamma,\Sigma'_{\vec{n}\cdot\vec{t}})}I\eta[\vec{n}]$, we then get an isomorphism of Γ/Λ -spectra

$$\bigvee_{\vec{n}\cdot\vec{t}=q} J^{\Gamma/\Lambda}(\eta[\vec{n}]) \xrightarrow{\cong} J^{\Gamma/\Lambda}((\operatorname{Sym}^{q}\eta)(\Lambda|K)).$$

For $i = 1, \ldots, T$, let \vec{e}_i be the vector with 1 in the *i* position and zero elsewhere so that $\tau(\vec{e}_i) = \tau_i$. The decomposition of $\tau(\vec{n})$ into indecomposable pieces decomposes the $\tau(\vec{n})$ -twisted Λ action on $\eta^{\vec{n}\cdot\vec{t}}$ into a product, and we get the following further identification of $\eta[\vec{n}]$. In the formula, when $n_i = 0$, we should understand $\operatorname{Sym}^{n_i} \eta[\vec{e}_i]$ as the trivial bundle $* \to *$.

PROPOSITION 5.3. Let $f_{\vec{n}}$ denote the group isomorphism (5.1) and $f_{\vec{n}}^*$ the induced equivalence of $(\Sigma_{n_1} \wr C(\tau_1)) \times \cdots \times \Sigma_{n_r} \wr C(\tau_r))$ -spaces with $C(\tau(\vec{n}))$ -spaces. We then have an isomorphism of $(\Gamma, C(\tau(\vec{n})))$ vector bundles

$$\eta[\vec{n}] \cong f_{\vec{n}}^*((\operatorname{Sym}^{n_1}\eta[\vec{e}_1]) \times \dots \times (\operatorname{Sym}^{n_T}\eta[\vec{e}_T]))$$

for any non-zero \vec{n} .

We can now prove Theorem V.3.1

PROOF OF THEOREM V.3.1. We use the notation established above. We assume that $\operatorname{Sym}^q \eta$ is Σ'_q -faithful for all q so that by Theorem V.2.4 (and Proposition V.1.4), we have an isomorphism of Γ/Λ -spectra

$$\Phi^{\Lambda}((J^{\Gamma/K}\eta)^{(q)}/\Sigma_q) \cong J^{\Gamma/\Lambda}((\operatorname{Sym}^q \eta)(\Lambda|K))$$

for all $q \ge 1$. For each $q \ge 1$, if there is no irreducible homomorphism $\sigma \colon \Lambda \to \Sigma'_q$ then set $X_q = *$; otherwise let

$$X_q = \bigvee_{t_i = q} J^{\Gamma/\Lambda}(\eta[\vec{e_i}]).$$

In particular, $X_q = *$ for $q > \#(\pi_0 \Lambda)$. By Proposition 5.2 and the observation below it, each X_q is then a wedge summand of $\Phi^{\Lambda}((J^{\Gamma/K}\eta)^{(q)}/\Sigma_q)$. Propositions 5.2 and 5.3 (together with Proposition V.1.4) then imply that the inclusion of the X_q in $\Phi^{\Lambda}((J^{\Gamma/K}\eta)^{(q)}/\Sigma_q)$ induces an isomorphism

$$\mathbb{P}\left(\bigvee_{q} X_{q}\right) \longrightarrow \Phi^{\Lambda}(\mathbb{P}(J^{\Gamma/K}\eta))$$

of commutative ring Γ/Λ -spectra.

6. The diagonal fixed point lemma and Theorem V.3.5

In this section, we prove a general lemma about fixed points of cartesian powers of equivariant spaces with respect to subgroups of wreath product groups. The main application in this section is the proof of Theorem V.3.5, which like the arguments in the previous section uses the statement of Theorem V.2.4 but not the ideas in its proof. We also used a version of the lemma in Section III.4.

The setup for the diagonal fixed point lemma is as follows. Let Λ and Q be groups, and $q \geq 1$. Let X be a left $(\Lambda \times Q^{\text{op}})$ -space, and let $\sigma \colon \Lambda \to \Sigma_q \wr Q$ be a homomorphism. By definition, the σ -twisted Λ action on X^q is the diagonal of the given left Λ action and the right action via σ .

$$\lambda \cdot_{\sigma} (x_1, \dots, x_q) := (\lambda x_1, \dots, \lambda x_q) \sigma(\lambda^{-1})$$

We denote the fixed points of this action as $(X^q)^{\Lambda,\sigma}$ and the lemma below gives a diagonal description of this action. Since Λ and Q are involved only in taking fixed points, any topology on these groups is irrelevant; we only need to assume their actions on X are continuous.

To fix notation with simplest formulas, we write the wreath product $\Sigma_q \wr Q$ as the semidirect product $Q^q \rtimes \Sigma_q$ and write an element as $(a_1, \ldots, a_q; s)$ with $a_i \in Q$ and $s \in \Sigma_q$. Then

$$(a_1, \dots, a_q; s)(a'_1, \dots, a'_q; s') = (a_1 a'_{s^{-1}(1)}, \dots, a_q a_{s^{-1}(q)}; ss').$$

We also use the a_i and s notation for writing the coordinates of σ :

$$(a_1(\lambda),\ldots,a_q(\lambda);s(h)):=\sigma(\lambda),$$

noting that $s: \Lambda \to \Sigma_q$ is a homomorphism, but the multiplication intertwines the other coordinates and s. To avoid confusion in notation with multiple arguments, we write the action of $s \in \Sigma_q \wr Q$ on $\{1, \ldots, q\}$ using brackets: $s(\lambda)$ sends i to $s(\lambda)[i]$. The formula for multiplication in the a_i coordinates is then

$$a_i(\lambda\mu) = a_i(\lambda)a_{s(\lambda^{-1})[i]}(\mu)$$

The action of an element $(a_1, \ldots, a_q; s)$ on an element (x_1, \ldots, x_q) in X^q in this notation is

$$(x_1, \ldots, x_q)(a_1, \ldots, a_q; s) = (x_{s[1]}a_{s[1]}, \ldots, x_{s[q]}a_{s[q]}).$$

For $\lambda \in \Lambda$, we then have

$$\lambda \cdot_{\sigma} (x_1, \dots, x_q) = (\lambda x_{s(\lambda^{-1})[1]} a_{s(\lambda^{-1})[1]} (\lambda^{-1}), \dots, \lambda x_{s(\lambda^{-1})[q]} a_{s[q]} (\lambda^{-1})).$$

In general, given an element of, or formula in X^q , we write a subscript *i* for its *i*th component; e.g., if $x \in X^q$, x_i denotes its *i*th component, and for $\lambda \in \Lambda$, $(\lambda \cdot_{\sigma} x)_i$ denotes the *i*th component of $\lambda \cdot_{\sigma} x$.

For $i = 1, \ldots, q$, let H_i denote the subgroup of Λ that is the inverse image under s of the subgroup of Σ_q that fixes i; restricted to H_i , a_i becomes a homomorphism, which we denote as $\alpha_i \colon H_i \to Q$. (We note that in the typical case, when Λ is a compact Lie group, Q is a discrete group, and the homomorphism σ is continuous, we have that the identity component Λ_0 of Λ is contained in H_i for every i.) We then have an α_i -twisted H_i action on X defined by

$$h \cdot_{\alpha_i} x := h x \alpha_i (h^{-1})$$

(for $x \in X$, $h \in H_i$) using the original left Λ action and right Q action on X.

The homomorphism s gives an action of Λ on the set $\{1, \ldots, q\}$; let O_1, \ldots, O_k denote its orbits. For $i = 1, \ldots, q$, define $j_i \in \{0, \ldots, k\}$ by $i \in O_{j_i}$. For each $j = 1, \ldots, k$, choose an element $m_j \in O_j$. For each $i = 1, \ldots, q$, choose an element $\ell_i \in \Lambda$ such that $s(\ell_i)[m_{j_i}] = i$. We can now state the lemma.

LEMMA 6.1. In the notation above, the map

$$f: X^{H_{m_1},\alpha_{m_1}} \times \cdots \times X^{H_{m_k},\alpha_{m_k}} \longrightarrow X^q$$

defined by

$$f(x_1, \dots, x_k)_i = \ell_i x_{j_i} a_{m_{j_i}}(\ell_i^{-1})$$

induces a homeomorphism onto the σ -twisted fixed points $(X^q)^{\Lambda,\sigma}$. Moreover, this map is independent of the choice of the elements ℓ_i above.

The map does depend on the choice of elements $m_j \in O_j$.

PROOF. To show that every σ -twisted Λ fixed point is in the image, assume $x \in (X^q)^{\Lambda,\sigma}$ and observe that

$$(\ell_i \cdot_{\sigma} x)_i = \ell_i x_{s(\ell_i^{-1})[i]} a_{s(\ell_i^{-1})[i]} (\ell_i^{-1})$$
$$= \ell_i x_{m_{j_i}} a_{m_{j_i}} (\ell_i^{-1}).$$

In the case when $i = m_j$ for some j and $h \in H_{m_j}$, we have

$$(h \cdot_{\sigma} x)_{m_j} = h x_{s(h^{-1})[m_j]} a_{s(h^{-1})[m_j]} (h^{-1})$$
$$= h x_{m_j} a_{m_j} (h^{-1}).$$

Since $h \in H_{m_j}$, we have $\alpha_{m_j}(h^{-1}) := a_{m_j}(h^{-1})$, and since $x \in (X^q)^{\Lambda,\sigma}$, we conclude that

$$x_{m_j} = (h \cdot_\sigma x)_{m_j} = h x_{m_j} \alpha_{m_j} (h^{-1}) = h \cdot_{\alpha_{m_j}} x_{m_j},$$

i.e., $x_{m_j} \in X^{H_{m_j},\alpha_{m_j}}$. So $x \in (X^q)^{\Lambda,\sigma}$ implies $x = f(x_{m_1},\ldots,x_{m_k})$ with $x_{m_j} \in X^{H_{m_j},\alpha_{m_j}}$. Next to show that f lands in the σ -twisted Λ fixed points, for $\lambda \in \Lambda$, writing $s = s(\lambda^{-1})$, we have

$$\begin{aligned} (\lambda \cdot_{\sigma} f(x_1, \dots, x_k))_i &= (\lambda \ell_{j_{s[i]}}) x_{j_{s[i]}} (a_{m_{j_{s[i]}}} (\ell_{j_{s[i]}}^{-1}) a_{s[i]} (\lambda^{-1})) \\ &= (\lambda \ell_{j_{s[i]}}) x_{j_{s[i]}} (a_{m_{j_{s[i]}}} (\ell_{j_{s[i]}}^{-1} \lambda^{-1})) \end{aligned}$$

since $s(\ell_{s[i]})[m_{j_{s[i]}}] = s[i]$. Since $s(\lambda \ell_{s[i]})[m_{j_i}] = s^{-1}s[i] = i, \lambda \ell_{j_{s[i]}}$ is an alternative choice for the element $\ell_i \in \Lambda$. Thus, to see that $f(x_1, \ldots, x_k)$ is fixed by λ (under the σ -twisted action), we just need to verify the last statement: f is independent of the choice of ℓ_i . Let ℓ'_i be another choice and $\mu = \ell_i^{-1}\ell'_i$ so that $\ell_i\mu = \ell'_i$. To reduce subscripts, let $j = j_i$. Then $s(\mu)[m_j] = m_j$ and $\mu \in H_{m_j}$, so

$$\mu x_j a_{m_j}(\mu^{-1}) = x_j$$

since $x_j \in X^{H_{m_j}, \alpha_{m_j}}$. Then

$$\ell_i x_j a_{m_j}(\ell_i^{-1}) = \ell_i \mu x_j a_{m_j}(\mu^{-1}) a_{m_j}(\ell_i^{-1})$$
$$= (\ell_i \mu) x_j a_{m_j}(\mu^{-1}\ell_i^{-1})$$
$$= \ell'_i x_j a_{m_j}((\ell'_i)^{-1})$$

with the second equality holding because μ fixes m_i .

We now move on to Theorem V.3.5, and we use the notations and conventions from there, so Q is a finite group, and Λ is a closed normal subgroup of a compact Lie group Γ . As in the previous section, we write Σ'_q for $\Sigma_q \wr Q$. We look at a homomorphism $\sigma \colon \Lambda \to \Sigma_{q'}$ and the σ -twisted Λ action on $\operatorname{Sym}^q \eta$ for a (Γ, Q) vector bundle η . The previous lemma gives us a diagonal description of the σ twisted Λ fixed points of both the total space E^q and the base space B^q . If we restrict to an element $b = (b_1, \ldots, b_q) \in (B^q)^{\Lambda, \sigma}$, the σ -twisted Λ action is fiberwise on E_b , and the lemma identifies the space $E_b^{\Lambda, \sigma}$ as

$$E_{b_{m_1}}^{H_{m_1},\alpha_{m_1}} \times \dots \times E_{b_{m_k}}^{H_{m_k},\alpha_{m_k}}$$

By inspection of the formula, this homeomorphism is an isomorphism of vector spaces. It may not be an isometry, but this is easy to correct by scaling. In the following proposition, for an element v of E_{b_i} , let $\delta_i v$ denote the element of E_b that is v in the *i*th coordinate and zero elsewhere; notation aside, the homomorphism in it is just the composite of the one in the lemma with the rescaling $v_j \mapsto v_j/\sqrt{\#O_j}$.

PROPOSITION 6.2. Define the isometry

$$f \colon E_{b_{m_1}}^{H_{m_1},\alpha_{m_1}} \times \cdots \times E_{b_{m_k}}^{H_{m_k},\alpha_{m_k}} \longrightarrow E_b^q$$

by the formula

$$f(v_{m_1}, \dots, v_{m_k}) = \sum_{j=1}^k \sum_{i \in O_j} \frac{\ell_i \cdot_\sigma (\delta_{m_j} v_{m_j})}{\sqrt{\#O_j}} = \sum_{j=1}^k \sum_{i \in O_j} \frac{\delta_i f_i(v_{m_j})}{\sqrt{\#O_j}}.$$

Then f is an isomorphism onto $(E_h^q)^{\Lambda,\sigma}$.

We can now prove theorem V.3.5.

PROOF OF THEOREM V.3.5. We fix $q \ge 1$ and show that $\eta' := (\text{Sym}^q \eta)(\Lambda)$ is Σ'_q -faithful. We use the notation of the discussion above: we study the coproduct factor corresponding to a fixed $\sigma \colon \Lambda \to \Sigma'_q$ at a point $b = (b_1, \ldots, b_q) \in (B^q)^{\Lambda, \sigma}$. For this, it suffices to show that for every $\tau = (c_1, \ldots, c_q; t)$, not the identity, with $b\tau = b$, there exists $w \in (E_b^q)^{\Lambda, \sigma}$ such that $w\tau \neq w$. We treat this in three cases, one corresponding to each of the conditions in Theorem V.3.5.

The first case is when t sends an element i in O_j to an element i' in $O_{j'}$ for some $j \neq j'$. Then by Proposition 6.2 and condition (i), we can choose an element w where the components in O_j are non-zero and the components in $O_{j'}$ are zero. For such a $w, w\tau \neq w$.

The second case is when t preserves each of the orbits O_1, \ldots, O_k but is not the identity, say $t(i) = i' \neq i$ for $i \in O_j$ for some j. Since the numbering of the orbits was arbitrary and the choice of orbit representatives above was arbitrary, we can

assume without loss of generality that j = 1 and $m_1 = i'$; we note that $\ell_i \notin H_{m_1}$. Since $b\tau = b$ and $\ell_i \cdot_{\sigma} b = b$, looking at *i*th coordinates, we see that

$$b_i = b_{m_1} c_{m_1} = \ell_i b_{m_1} a(\ell_i^{-1})$$

and so, writing $a := c_{m_1} a(\ell_i^{-1})^{-1}$, we have $\ell_i b_{m_1} = b_{m_1} a$. By condition (ii), there exists $v \in E_{b_{m_1}}^{H_{m_1},\alpha_{m_1}}$ such that $\ell_{m_1} v \neq v a$. Let $w = f(\sqrt{\#O_1}v, 0, \dots, 0) \in (E_b^q)^{\Lambda,\sigma}$. Then

$$w_i = \ell_{m_1} va(\ell_{m_1}^{-1}), \qquad (w\tau)_i = vc_{m_1}$$

and we have chosen v so that these two elements are unequal.

The final case is when t is the identity. Then there exists some i for which c_i is not the identity. Without loss of generality $m_1 = i$. By condition (iii), we can choose $v \in E_{b_m}^{H_{m_1},\alpha_{m_1}}$ so that $vc_{m_1} \neq v$. Then

$$f(v, 0, \dots, 0)\tau = f(vc_{m_1}, 0, \dots, 0) \neq f(v, 0, \dots, 0).$$

7. Indexed smash powers, HHR norms, geometric fixed points, and the HHR diagonal

The purpose of this section is to generalize the HHR diagonal map [15, B.209]

$$\Phi^H X \longrightarrow \Phi^G(X^{(G/H)})$$

that computes the geometric fixed points of an indexed smash power. We generalize in two directions: we consider universes other than the complete universe and we consider compact Lie groups that are not finite. This map does not exist for arbitrary universes and arbitrary orbit spaces G/H; we specify below when results hold. The results in this section are known to experts, but statements have not appeared in the generality we consider here, though the proofs in the existing literature generalize to cover them. Since we expect these results to be of interest beyond the scope of the rest of the paper, we have made this section as self-contained as possible at the cost of a bit of redundancy.

In this section G denotes a compact Lie group. We let U be a G-universe. We write V < U to mean that V is a finite dimensional G-stable subspace of U; for finite dimensional G-inner product spaces, we also write V < W to indicate that V is a subspace of W (with inherited G action and inner product) and we emphasize that in this case the notation does not imply proper subspace: W < W. For H < G a closed subgroup, we write U_H for U viewed as an H-universe (and write $V < U_H$ for V a finute dimensional H-stable subspace). We use the notation of Section V.1 (the unadorned form of the notation in [22]) for the categories \mathcal{I} and \mathcal{J} : for vector spaces with inner products $V, W, \mathcal{I}(V, W)$ denotes the space of isometric embeddings $V \to W$, $\Im^{\perp} \mathcal{I}(V, W) \to \mathcal{I}(V, W)$ denotes the vector bundle over $\mathcal{I}(V,W)$ whose fiber is the orthogonal complement of the image of an embedding, and $\mathcal{J}(V,W)$ denotes the Thom space of this vector bundle. When V and W are orthogonal G-representations, $\mathcal{I}(V, W)$ gets the usual G action on a mapping space, $\Im^{\perp} \mathcal{I}(V, W) \to \mathcal{I}(V, W)$ becomes an G-equivariant vector bundle with its natural action, and $\mathcal{J}(V,W)$ becomes a based G-space. By [22, II§4], the functor $\mathcal{J}(V,-)$ specifies a G-equivariant orthogonal spectrum that represents the Vth space functor $X \mapsto X(V)$. In [22], $\mathcal{J}(V, -)$ is more usually denoted $F_V S^0$, and in [15], it is denoted $S^{-\dot{V}}$.

The results in this section are of a point-set rather than homotopical nature, and therefore, when convenient, we can treat *G*-equivariant orthogonal spectra as indexed on \mathbb{R}^{∞} instead of *U*: the forgetful functor that only remembers the indexing on \mathbb{R}^{∞} is an equivalence of categories [**22**, V.1.5]. The convenience of this is that the category of *G*-equivariant orthogonal spectra indexed on \mathbb{R}^{∞} is just the category of left *G*-objects in (non-equivariant) orthogonal spectra and this makes constructions like indexed smash powers and HHR norms significantly easier to describe. In so doing, however, we need to be careful about which concepts or constructions depend on *U* and which do not. The concepts/constructions that are independent of the universe *U* include:

- The smash product \wedge of G-equivariant orthogonal spectra and the smash product of a G-equivariant orthogonal spectrum with a based G-space.
- Indexed smash powers / HHR norms (when defined).
- Limits and colimits.
- Restriction / (additive) induction along inclusions of subgroups.

The concepts/constructions that depend in the universe U include:

- Weak equivalences / homotopy groups.
- Cofibrancy / fibrancy in standard or complete model structures.
- The tautological presentation (coequalizer).
- The geometric fixed point functor (both point-set and derived).

It is in the last two points that the universe plays a role in this section. When indexing G-equivariant orthogonal spectra on \mathbb{R}^{∞} and working with the universe U, we understand the Vth space X(V) of a G-equivariant orthogonal spectrum to be

$$X(V) := \mathcal{I}(\mathbb{R}^n, V)_+ \wedge_{O(n)} X(\mathbb{R}^n) = \mathcal{J}(\mathbb{R}^n, V) \wedge_{O(n)} X(\mathbb{R}^n)$$

when V is *n*-dimensional; see $[22, V\S1]$.

Indexing on \mathbb{R}^{∞} , finite *G*-set indexed smash powers become easy to describe. For a finite *G*-space *S* and a *G*-equivariant orthogonal spectrum *X*, we get an orthogonal spectrum $X^{(S)}$ with *n*th space

$$X^{(S)}(\mathbb{R}^n) := (X(\mathbb{R}^n))^{(S)}$$

the smash power indexed on S, with G action both on $X(\mathbb{R}^n)$ and on S. In some contexts, it is useful to have a description of the indexed smash power that is universe U intrinsic in the sense that it does not require reindexing. We gives such a construction in the following proposition; its proof is clear when indexing on the universe \mathbb{R}^{∞} and the construction is preserved under the reindexing to U.

PROPOSITION 7.1. Let G be a compact Lie group. Let $k \ge 1$, $\Sigma < \Sigma_k$, and $\sigma: G \to \Sigma$ a continuous homomorphism. Let S be the finite G-set obtained from $\{1, \ldots, k\}$ using σ for the action, and let $\Sigma(\sigma)$ be the $(G \times \Sigma^{\text{op}})$ -set whose underlying right Σ -set is Σ and which has left Γ action induced by the homomorphism σ . Then for G-equivariant symmetric spectra X, there is a natural isomorphism

$$X^{(S)} \cong (\Sigma(\sigma)_+ \wedge X^{(k)}) / \Sigma_-$$

When S = G/H for H < G a codimension zero subgroup, the indexed smash power $X^{(G/H)}$ is a functor of the underlying *H*-equivariant orthogonal spectrum, namely the HHR norm [15, A§4]. In the case of finite groups, the HHR norm can be made without choices; when *G* is a compact Lie group, we seem to have to choose

100

a set of coset representatives $\gamma_{\zeta} \in G$ for $\zeta \in G/H$ (with $\zeta = \gamma_{\zeta} H$). We then get a continuous homomorphism

$$\sigma_{G/H} \colon G \longrightarrow \Sigma_{G/H} \wr H$$

defined by $\sigma_{G/H}(g) = (s, (h_{\zeta}))$ where s is the permutation on G/H induced by right multiplication by g, and h_{ζ} is defined by $\gamma\gamma_{\zeta} = \gamma_{s\zeta}h_{\zeta}$. We note that for an *H*-equivariant orthogonal spectrum X, the spaces of the kth smash power $X^{(k)}(\mathbb{R}^n)$ come with an action of the group $\Sigma_k \wr H$, making it naturally a $(\Sigma_k \wr H)$ -equivariant orthogonal spectrum; then, regarding G/H as just a set, the smash power $X^{(G/H)}$ has the structure of a $(\Sigma_{G/H} \wr H)$ -equivariant orthogonal spectrum, and we use $\sigma_{G/H}$ to make it a *G*-equivariant orthogonal spectrum. This defines the HHR norm $N_H^G X$.

DEFINITION 7.2. Let G be a compact Lie group and H < G a codimension zero subgroup. Define the norm N_H^G from H-equivariant orthogonal spectra to G-equivariant orthogonal spectra by

$$N_H^G X(\mathbb{R}^n) = X^{(G/H)}(\mathbb{R}^n)$$

with the G action defined by $\sigma_{G/H}$ above.

We have a completely analogous construction of N_H^G as a functor from based H-spaces to based G-spaces. When $X = \mathcal{J}(V, -) \wedge A$ where V is a finite dimensional H-inner product space and A is a based H-spaces, then by inspection

$$N_H^G(\mathcal{J}(V,-)\wedge A)\cong \mathcal{J}(\operatorname{Ind}_H^G V,-)\wedge N_H^G A$$

where $\operatorname{Ind}_{H}^{G} V$ is the (finite dimensional) *G*-inner product space defined by *H* to *G* induction; it is characterized by the universal property $\mathcal{I}(\operatorname{Ind}_{H}^{G} V, -)^{G} \cong \mathcal{I}(V, -)^{H}$. (It is constructed analogously to N_{H}^{G} but using direct sum in place of smash product.)

While the construction in Definition 7.2 depends on the choice of coset representatives, a different choice gives a naturally isomorphic functor via multiplication by the element of $H^{G/H}$ that transforms the representatives.

In the case when X is a G-equivariant orthogonal spectrum, the norm N_H^G is isomorphic to the indexed smash power $X^{(G/H)}$, where the isomorphism multiplies by the element of $G^{G/H}$ given by the coset representatives (for the indexing on the universe \mathbb{R}^{∞}).

The geometric fixed point functor does depend on the universe U, and for H < G a closed subgroup, we write Φ_U^H to keep track of the universe when it is not complete. The functor is constructed in [22, V§4] as the composite of the spacewise H fixed point functor and a left Kan extension. Rather than write out the Kan extension, we follow [15, B§10] and describe it in terms of the resulting coequalizer. This coequalizer is well-motivated in terms of the the *tautological presentation* of a G-equivariant orthogonal spectrum X, which is the coequalizer

$$\bigvee_{V,W < U} \mathcal{J}(W, -) \land \mathcal{J}(V, W) \land X(V) \Longrightarrow \bigvee_{V < U} \mathcal{J}(V, -) \land X(V) \longrightarrow X.$$

In the case of H fixed points, we restrict the G action to H and consider the tautological presentation for that group

$$\bigvee_{V,W < U_H} \mathcal{J}(W, -) \land \mathcal{J}(V, W) \land X(V) \Longrightarrow \bigvee_{V < U_H} \mathcal{J}(V, -) \land X(V) \longrightarrow X.$$

The idea is that $\Phi_U^H \mathcal{J}(V, -) \cong \mathcal{J}(V^H, -)$ for every $V < U_H$ and although Φ_U^H does not preserve all reflexive coequalizers, it does preserve reflexive coequalizers that are spacewise split on the underlying spaces indexed on U_H (see [1, 2.11]). The geometric fixed point spectrum $\Phi_U^H X$ is then the coequalizer

$$\bigvee_{V,W < U_H} \mathcal{J}(W^H, -) \wedge \mathcal{J}(V, W)^H \wedge X(V)^H \Longrightarrow \bigvee_{V < U_H} \mathcal{J}(V^H, -) \wedge X(V)^H$$

where one map is induced by the *H* fixed points of the action map $\mathcal{J}(V, W) \land X(V) \to X(W)$ and the other is induced by the map

$$\mathcal{J}(W^H, -) \land \mathcal{J}(V, W)^H \longrightarrow \mathcal{J}(W^H, -) \land \mathcal{J}(V^H, W^H) \longrightarrow \mathcal{J}(V^H, -)$$

that takes an H fixed element of $\mathcal{J}(V, W)$, restricts to the fixed points in V and W to get an element of $\mathcal{J}(V^H, W^H)$, and composes. This defines $\Phi_U^H X$ as a functor to non-equivariant orthogonal spectra, but in the case of interest to us when H < G is codimension zero, we can also recover the action of the Weyl group $W_G H$ via the evident (continuous) action of the normalizer $N_G H$ on the wedge sums (which involves permutation of the indexing $V, W < U_H$ and $V < U_H$).

If we want the equivariance for more general H < G, we restrict the G action to the normalizer $N = N_G H$ and use the coequalizer

$$\bigvee_{V,W < U_N} \mathcal{J}(W^H, -) \wedge \mathcal{J}(V, W)^H \wedge X(V)^H \Longrightarrow \bigvee_{V < U_N} \mathcal{J}(V^H, -) \wedge X(V)^H$$

deriving from the tautological presentation for the group N. This coequalizer is isomorphic to the one above using the group H by [19, 3.8.12] (the statement is phrased in terms of Kan extensions, but the argument is exactly a comparison of these coequalizers). We restate this using the notation of [22].

PROPOSITION 7.3 ([19, 3.8.12]). When $N \triangleleft G$ is a normal subgroup, the functor Φ_U^N from G-equivariant orthogonal spectra (indexed on U) to non-equivariant orthogonal spectra is naturally isomorphic to the composite of the geometric fixed point functor Φ^N of [22, V.4.3] from G-equivariant orthogonal spectra (indexed on U) to G/N-equivariant orthogonal spectra (indexed on U^N) and the forgetful functor to non-equivariant orthogonal spectra.

REMARK 7.4. The statement of [19, 3.8.12] includes a hypothesis on N fixed points of N-representations versus N fixed points of G-representations that amounts to the following: If N is a closed normal subgroup of a compact Lie group G, and Vis a non-trivial irreducible (finite dimensional) N-representation, then there exits an irreducible (finite dimensional) G-representation W that contains V as a sub N-representation and satisfies $W^N = 0$. This hypothesis always holds. The Peter-Weyl theorem implies that V is contained as a sub N-representation of some finite dimensional G-representation W, which without loss of generality can be taken to be irreducible. Choosing any non-zero v in V, the underlying vector space of W is then generated by elements of the form gv for $g \in G$. The sub N-representation generated by gv is isomorphic to $\operatorname{Ad}_g^* V$, where $\operatorname{Ad}_g: N \to N$ is automorphism of N given by conjugation by g. Since by hypothesis, V is a non-trivial irreducible N-representation, so is $\operatorname{Ad}_g^* V$, and this writes the underlying N-representation of W as a sum of non-trivial irreducible N-representations, showing that $W^N = 0$.

REMARK 7.5. If H is not normal in G, the coequalizer using the tautological presentation for the group G does not in general give an isomorphic orthogonal

spectrum. To give a specific example, consider the case when $G = A_5$ and $H = \langle (12)(34) \rangle$, with U a complete G-universe, and $\mathbb{R}(\sigma) < U_H$ the sign representation of H; in this example, the coequalizers

$$\bigvee_{V,W
$$\bigvee_{V,W$$$$

are non-isomorphic: the bottom is a split coequalizer, giving $\mathcal{J}(\mathbb{R}(\sigma)^H, 0) = S^0$ as the resulting space, but in the top, every A_5 representation V containing the sign representation also contains copies of the trivial representation, so we get * for the coequalizer. (The *G*-equivariant orthogonal spectrum $X = G_+ \wedge_H \mathcal{J}(\mathbb{R}(\sigma), -)$ likewise gives different answers for the two coequalizers.) The point is that when H < G is normal, V is a non-trivial irreducible *H*-representation, and W is an irreducible *G*-representation containing V as a sub *H*-representation, we have $W^H = 0$, but this can fail when *H* is not normal.

To see that Φ_U^H depends on the U (and that it does not preserve arbitrary reflexive coequalizers), consider the case $X = \mathcal{J}(V, -)$ for various finite dimensional G-inner product spaces V. As already noted, we have

$$\Phi_U^H \mathcal{J}(V, -) \cong \mathcal{J}(V^H, -) \qquad \text{for } V < U_H$$

but if V is not H-isomorphic to a subspace of U, then $\mathcal{I}(V, W)^H$ is empty for all $W < U_H$ and so $X(W)^H = \mathcal{J}(V, W)^H = *$ for all $W < U_H$. This gives

$$\Phi_U^H \mathcal{J}(V, -) \cong * \quad \text{for } \mathcal{I}(V, U)^H = \{\}.$$

Now we restrict to the case when H < G is zero codimensional so that G/H is a finite G-set. Under the hypothesis that the G-universe U is isomorphic to the G-universe $U \otimes G/H$, we obtain a diagonal natural transformation

$$\Phi_U^H \longrightarrow \Phi_U^G((-)^{(G/H)})$$

for G-equivariant orthogonal spectra as a special case of the natural transformation

$$\Phi^H_{U_H} \longrightarrow \Phi^G_U(N^G_H(-))$$

defined as follows, using the outline given in [15, B.209]. First, as already observed, for a based *H*-space *A*, we have a norm construction N_H^G defined using $\sigma_{G/H}$ in the same manner as the spectral construction in Definition 7.2. The diagonal map

$$A^H \xrightarrow{\cong} (N^G_H A)^G$$

that sends an H fixed point a to the element of the smash product which is a in each factor is an isomorphism. Likewise, on a finite dimensional H-inner product space V, we get an isomorphism of vector spaces

$$V^H \xrightarrow{\cong} (\operatorname{Ind}_H^G V)^G$$

which with the appropriate scaling becomes an isometry. Then for a based G-space A and $V < U_H$, we get isomorphisms

$$\begin{split} \Phi^G_U(N^G_H(\mathcal{J}(V,-)\wedge A)) &\cong \Phi^G_U(\mathcal{J}(\mathrm{Ind}_H^G V,-)\wedge N^G_H A) \\ &\cong \mathcal{J}((\mathrm{Ind}_H^G V)^G,-)\wedge (N^G_H A)^G \cong \mathcal{J}(V^H,-)\wedge A^H. \end{split}$$

Applying the composite functor $\Phi_U^G N_H^G$ to the tautological presentation (for the group H) and applying the isomorphism above, we precisely get the coequalizer diagram for $\Phi_{U_H}^H$, and that induces the diagonal map from $\Phi_{U_H}^H \to \Phi_U^G N_H^G$. We state the observation here in the following proposition for easy reference.

PROPOSITION 7.6. Let G be a compact Lie group and let H < G be a codimension zero subgroup. Assume the G-universe U is isomorphic to the G-universe $U \otimes G/H$. Then for H-equivariant orthogonal spectra indexed on U_H , there is a natural diagonal map of non-equivariant orthogonal spectra

$$\Phi^H_{U_H} \longrightarrow \Phi^G_U N^G_H$$

and for G-equivariant orthogonal spectra indexed on U, a natural diagonal map of non-equivariant orthogonal spectra

$$\Phi_U^H \longrightarrow \Phi_U^G((-)^{(G/H)}).$$

As in the case covered in [15, B.209], for nice X, the diagonal map is an isomorphism. It is asserted and proved there that (in the setting of a finite group and the complete universe), whenever X is cofibrant for the positive complete model structure, the diagonal is an isomorphism; moreover, [15, 2.23] remarks that positivity is not needed, and a careful check of the proof shows that it applies to objects that are cofibrant in the complete model structure, that is, retracts of cell complexes built out of cells of the form

$$G_+ \wedge_M \mathcal{J}(V, -) \wedge (D^n, \partial D^n)_+$$

where M < G and V is an M-inner product space. (Positivity would require V to have a non-zero M fixed vector.) The same result holds with essentially the same proof for the diagonal map in any of the universes above.

THEOREM 7.7. Let G be a compact Lie group, H < G a codimension zero subgroup, and U a G-universe, not necessarily complete, with $U \otimes G/H$ isomorphic to U. If X is an H-equivariant orthogonal spectrum that is cofibrant in the complete model structure on the complete universe, then the U universe diagonal

$$\Phi^H_{U_H} X \longrightarrow \Phi^G_U N^G_H X$$

is an isomorphism.

PROOF. The outline of the argument for the finite group version of the theorem in [15, B.209] goes through without difficulty to reduce to the case when X is a single cell of the form $H_+ \wedge_M \mathcal{J}(V, -)$. In this case, when M is a proper subgroup of H, both sides are clearly the trivial orthogonal spectrum *, and the result holds. In the case when M = H, and V is not isomorphic to a subspace of the universe U_H , $\operatorname{Ind}_H^G V$ is also not isomorphic to a subspace of the universe $U_H \cong (U \otimes G/H)_H$, and again both sides are the trivial orthogonal spectrum *. Finally, in the last case $M = H, V < U_H$, and we have that the diagonal map

$$\mathcal{J}(V^H, -) \cong \Phi^H_{U_H} \mathcal{J}(V, -) \longrightarrow \Phi^G_U N^G_H \mathcal{J}(V, -) \cong \mathcal{J}((\operatorname{Ind}_H^G V)^G, -)$$

is an isomorphism by construction.

Working with G-equivariant orthogonal spectra, we get the following immediate corollary.

COROLLARY 7.8. Let G be a compact Lie group, H < G a codimension zero subgroup, and U a G-universe, not necessarily complete, with $U \otimes G/H$ isomorphic to U. If X is an G-equivariant orthogonal spectrum that is cofibrant in the complete model structure on the complete universe, then the U universe diagonal

$$\Phi_U^H X \longrightarrow \Phi_U^G (X^{(G/H)})$$

is an isomorphism.

PROOF. Under the hypothesis, the underlying H-equivariant orthogonal spectrum is cofibrant in the complete model structure on the complete universe. \Box

We state one more result on the diagonal in the case when H is the trivial group, a version of Theorem 7.7 for more general cofibrant objects. A non-equivariant orthogonal spectrum is *convenient O-model cofibrant* if it is a retract of an orthogonal spectrum built as cell complex using cells of the form

$$A \wedge (D^n, \partial D^n)_+$$

where

$$A = (F_{\mathbb{R}^n} S^0) / M$$

for M a closed subgroup of O(n). (These are the cofibrant objects in a model structure, the further details of which are not relevant.) As a special case we note that the cofibrant objects in the positive convenient Σ -model structure of Definition III.4.1 are convenient O-model cofibrant as are the underlying orthogonal spectra of cofibrant objects in the standard model structure on commutative ring orthogonal spectra.

THEOREM 7.9. Let G be a finite group and U a complete G-universe. If X is a non-equivariant orthogonal spectrum that is convenient O-model cofibrant, then the diagonal

$$X \longrightarrow \Phi_U^G N_1^G X$$

is an isomorphism.

PROOF. As in the proof of Theorem 7.7, it suffices to treat the case when $X = (F_{\mathbb{R}^n} S^0)/M$ for M a closed subgroup of O(n). This case reduces to showing that for every orthogonal G-representation W in U, the map

$$\mathcal{J}(\mathrm{Ind}_1^G \mathbb{R}^n, W)^G / M \longrightarrow (\mathcal{J}(\mathrm{Ind}_1^G \mathbb{R}^n, W) / N_1^G M)^G$$

is an isomorphism, where $N_1^G M$ is the group object in *G*-spaces given by the cartesian product of copies of M indexed by G. Since both sides are compact Hausdorff spaces and the map is obviously injective, it suffices to show that it is surjective. Let $(\phi, w) \in \mathcal{J}(\operatorname{Ind}_1^G \mathbb{R}^n, W)$ be a non-base point whose image in $\mathcal{J}(\operatorname{Ind}_1^G \mathbb{R}^n, W)/N_1^G M$ is G fixed. Here ϕ denotes an isometry $\operatorname{Ind}_1^G \mathbb{R}^n \to W$ and w a point in its complement. Since the image of (ϕ, w) is G fixed, w must be G fixed and for each $g \in G$, there exists an element $\sigma(g)$ in $N_1^G M$ such that

(7.10)
$$g \circ \phi \circ g^{-1} = \phi \circ \sigma(g)^{-1}.$$

In the identification of $\operatorname{Ind}_1^G \mathbb{R}^n$ and $N_1^G M$ as cartesian products indexed on G, let ϕ_1 denote the restriction of ϕ to the coordinate indexed by the identity and let $\sigma_1(g)$ denote the coordinate of $\sigma(g)$ indexed by the identity. Define $\psi \colon \operatorname{Ind}_1^G \mathbb{R}^n \to W$ by sending the coordinate indexed by g to $g \circ \phi_1$. We see from (7.10), the image of $g \circ \phi_1$ is the image of ϕ on gth coordinate of $\operatorname{Ind}_1^G \mathbb{R}^n$, and so ψ is a linear isometry

and (ψ, w) specifies an element of $\mathcal{J}(\operatorname{Ind}_1^G \mathbb{R}^n, W)$, which is in the same $N_1^G M$ coset as (ϕ, w) . But by inspection, $(\psi, w) \in \mathcal{J}(\operatorname{Ind}_1^G \mathbb{R}^n, W)^G$.

Bibliography

- Vigleik Angeltveit, Andrew J. Blumberg, Teena Gerhardt, Michael A. Hill, Tyler Lawson, and Michael A. Mandell. Topological cyclic homology via the norm. *Doc. Math.*, 23:2101–2163, 2018.
- [2] Bhargav Bhatt, Matthew Morrow, and Peter Scholze. Integral p-adic Hodge theory. Publ. Math. Inst. Hautes Études Sci., 128:219–397, 2018.
- [3] Andrew J. Blumberg and Michael A. Hill. Operadic multiplications in equivariant spectra, norms, and transfers. Adv. Math., 285:658–708, 2015.
- [4] Andrew J. Blumberg and Michael A. Mandell. The homotopy theory of cyclotomic spectra. Geom. Topol., 19(6):3105–3147, 2015.
- [5] Andrew J. Blumberg, Michael A. Mandell, and Allen Yuan. Relative cyclotomic structures and equivariant complex cobordism. Preprint, arXiv:2310.02348, 2023.
- [6] M. Bökstedt, W. C. Hsiang, and I. Madsen. The cyclotomic trace and algebraic K-theory of spaces. Invent. Math., 111(3):465–539, 1993.
- [7] Morten Brun, Bjørn Ian Dundas, and Martin Stolz. Equivariant structure on smash powers. Preprint, arXiv:1604.05939, 2022.
- [8] Gunnar Carlsson. Equivariant stable homotopy and Segal's Burnside ring conjecture. Ann. of Math. (2), 120(2):189–224, 1984.
- [9] Bjørn Ian Dundas, Thomas G. Goodwillie, and Randy McCarthy. The local structure of algebraic K-theory, volume 18 of Algebra and Applications. Springer-Verlag London, Ltd., London, 2013.
- [10] Bjørn Ian Dundas and Andrea Tenti. Higher Hochschild homology is not a stable invariant. Math. Z., 290(1-2):145–154, 2018.
- [11] Daniel S. Freed, Michael J. Hopkins, and Constantin Teleman. Loop groups and twisted K-theory III. Ann. of Math. (2), 174(2):947–1007, 2011.
- [12] Daniel S. Freed, Michael J. Hopkins, and Constantin Teleman. Loop groups and twisted K-theory II. J. Amer. Math. Soc., 26(3):595–644, 2013.
- [13] J. P. C. Greenlees and J. P. May. Localization and completion theorems for MU-module spectra. Ann. of Math. (2), 146(3):509–544, 1997.
- [14] Lars Hesselholt and Ib Madsen. On the K-theory of local fields. Ann. of Math. (2), 158(1):1– 113, 2003.
- [15] M. A. Hill, M. J. Hopkins, and D. C. Ravenel. On the nonexistence of elements of Kervaire invariant one. Ann. of Math. (2), 184(1):1–262, 2016.
- [16] Michael A. Hill, XiaoLin Danny Shi, Guozhen Wang, and Zhouli Xu. The slice spectral sequence of a C_4 -equivariant height-4 Lubin-Tate theory. Mem. Amer. Math. Soc., 288(1429):v+119, 2023.
- [17] Michael J. Hopkins, Nicholas J. Kuhn, and Douglas C. Ravenel. Generalized group characters and complex oriented cohomology theories. J. Amer. Math. Soc., 13(3):553–594, 2000.
- [18] D. Kaledin. Motivic structures in non-commutative geometry. In Proceedings of the International Congress of Mathematicians. Volume II, pages 461–496, New Delhi, 2010. Hindustan Book Agency.
- [19] Tore August Kro. Involutions on $s[\omega m]$. Preprint, arXiv:math/0510221, 2005.
- [20] L. G. Lewis, Jr., J. P. May, M. Steinberger, and J. E. McClure. Equivariant stable homotopy theory, volume 1213 of Lecture Notes in Mathematics. Springer-Verlag, Berlin, 1986. With contributions by J. E. McClure.
- [21] L. Gaunce Lewis, Jr. and Michael A. Mandell. Modules in monoidal model categories. J. Pure Appl. Algebra, 210(2):395–421, 2007.

BIBLIOGRAPHY

- [22] M. A. Mandell and J. P. May. Equivariant orthogonal spectra and S-modules. Mem. Amer. Math. Soc., 159(755):x+108, 2002.
- [23] M. A. Mandell, J. P. May, S. Schwede, and B. Shipley. Model categories of diagram spectra. Proc. London Math. Soc. (3), 82(2):441–512, 2001.
- [24] J. P. May. Equivariant homotopy and cohomology theory, volume 91 of CBMS Regional Conference Series in Mathematics. Published for the Conference Board of the Mathematical Sciences, Washington, DC, 1996. With contributions by M. Cole, G. Comezaña, S. Costenoble, A. D. Elmendorf, J. P. C. Greenlees, L. G. Lewis, Jr., R. J. Piacenza, G. Triantafillou, and S. Waner.
- [25] Stefan Schwede and Brooke E. Shipley. Algebras and modules in monoidal model categories. Proc. London Math. Soc. (3), 80(2):491–511, 2000.
- [26] Brooke Shipley. A convenient model category for commutative ring spectra. In *Homotopy theory: relations with algebraic geometry, group cohomology, and algebraic K-theory*, volume 346 of *Contemp. Math.*, pages 473–483. Amer. Math. Soc., Providence, RI, 2004.
- [27] Martin Stolz. Equivariant structure on smash powers of commutative ring spectra. University of Begen thesis, 2011.

108