

# $L^2$ -BETTI NUMBERS OF DEHN FILLINGS

NANSEN PETROSYAN AND BIN SUN

ABSTRACT. We initiate the study of the  $L^2$ -Betti numbers of group-theoretic Dehn fillings. For a broad class of virtually special groups  $G$ , we prove that the  $L^2$ -Betti numbers of sufficiently deep Dehn fillings  $\overline{G}$  are equal to those of  $G$ . As applications, we verify the Singer Conjecture for certain Einstein manifolds, establish a virtual fibering criterion for  $\overline{G}$ , obtain bounds on deficiency of  $\overline{G}$ , and provide new examples of hyperbolic groups with exotic subgroups that arise as Dehn fillings of any cusped arithmetic hyperbolic manifold of dimension at least four.

## 1. INTRODUCTION

Dehn filling is a process that produces a closed manifold from one with boundary. Let  $M$  be an  $n$ -manifold with boundary an  $(n - 1)$ -torus. A *Dehn filling* of  $M$  is a manifold obtained by gluing an  $n$ -dimensional solid torus  $D^2 \times S^{n-1}$  to  $M$  along their boundaries via a homeomorphism. It is particularly central to the study of 3-manifolds. Notably, Thurston's Hyperbolic Dehn Filling Theorem [Thu82] states that if  $M$  is a 3-dimensional hyperbolic knot complement, then most Dehn fillings of  $M$  admit a hyperbolic Riemannian metric, producing examples of non-Haken hyperbolic 3-manifolds and giving evidence to his Geometrization Conjecture.

There is an analogous construction in group theory, called (*group-theoretic*) *Dehn filling*, which also finds important applications in group theory and topology. Remarkably, one of the crucial steps in Agol's proof [Ago13] of Thurston's Virtual Haken Conjecture is to show that a virtually compact cubical hyperbolic group is in fact virtually compact special, for which Dehn filling is an indispensable tool. Other applications of group-theoretic Dehn filling can be found in [DGO17, ACG18, DG18, CIO23].

In its most general form, group-theoretic Dehn filling involves a triple  $(G, H, N)$ , where  $G$  is a group,  $H \leq G$  is a subgroup, and  $N \triangleleft H$  is a normal subgroup. The Dehn filling process produces the quotient group  $G/\langle\langle N \rangle\rangle$  as its output, where  $\langle\langle N \rangle\rangle$  is the normal closure of  $N$  in  $G$ . This algebraic framework mirrors the topological notion of Dehn filling, as seen through the Seifert–van Kampen theorem. Specifically, let  $G = \pi_1(M)$ ,  $H$  be the image of  $\pi_1(\partial M)$  in  $\pi_1(M)$ , and  $N$  the image of  $\pi_1(\partial D^2)$ . The gluing process in topology then corresponds to the group-theoretic construction.

The general setting of Dehn filling is overly broad to allow for meaningful conclusions. In practice, it is often assumed that  $H$  is a hyperbolically embedded subgroup of  $G$ , which is a generalization of  $G$  being hyperbolic relative to  $H$ , and proved that for many choices of  $N \triangleleft H$ , the quotient  $G/\langle\langle N \rangle\rangle$  will have certain properties. For example, a critical ingredient of Agol's work [Ago13] is Wise's Malnormal Special Quotient Theorem [Wis21, AGM16], which states that if  $G$  is a virtually compact special hyperbolic group, and  $G$  is hyperbolic

relative to its subgroup  $H$ , then for many choices of  $N \triangleleft H$ , the quotient  $G/\langle\langle N \rangle\rangle$  will be hyperbolic virtually compact special.

In this paper, we will consider the case where  $H$  is a hyperbolically embedded subgroup of  $G$  and compute the  $L^2$ -Betti numbers of  $G/\langle\langle N \rangle\rangle$  for many choices of  $N \triangleleft H$ .

Most of our results in the introduction are stated for a single peripheral for simplicity of exposition; however, they hold more generally for multiple peripherals.

**$L^2$ -Betti numbers of Dehn filling.** For any group  $G$ , the  $L^2$ -Betti number  $b_*^{(2)}(G)$  can be thought of as a normalized version of the usual Betti number. For example, for residually finite groups,  $L^2$ -Betti numbers can be computed using a nested sequence of finite-index normal subgroups, as described by Lück's Approximation Theorem [L94].

We will use  $H \hookrightarrow_h G$  to indicate that  $H$  is a hyperbolically embedded subgroup of  $G$ . To simplify the notation, for  $N \triangleleft H$ , we will often denote  $\overline{H} := H/N$  and  $\overline{G} := G/\langle\langle N \rangle\rangle$ . The reader is referred to Section 4.1 for a proof of Theorem 1.1 below.

**Theorem 1.1** (see also Corollary 5.5). *Let  $G$  be a virtually locally indicable group of type  $F_\infty$  and has a hyperbolically embedded amenable subgroup  $H \hookrightarrow_h G$ . Then for every  $\delta > 0$  and  $n \geq 0$ , there exists a finite subset  $\mathcal{F}_{n,\delta} \subset H \setminus \{1\}$  such that if a normal subgroup  $N \triangleleft H$  satisfies  $N \cap \mathcal{F}_{n,\delta} = \emptyset$  and  $\overline{H}$  is of type  $F_\infty$ , then*

$$|b_n^{(2)}(\overline{G}) - b_n^{(2)}(G)| < \delta.$$

Recall that a group is *locally indicable* if every non-trivial finitely generated subgroup of it has a homomorphism onto  $\mathbb{Z}$ , and a group *virtually* has property  $\mathcal{P}$  if it has a finite-index subgroup that has property  $\mathcal{P}$ .

The above theorem provides an approximation result for  $L^2$ -Betti numbers. Under the stronger assumption that  $G$  is virtually compact cubical, however, it becomes possible to compute the  $L^2$ -Betti numbers explicitly. Here, we say that a group is *compact cubical* if it is the fundamental group of a compact locally CAT(0) cube complex.

To simplify the statements of our next results, we will say that a property  $\mathcal{P}$  holds for all *sufficiently deep* Dehn fillings or for all *sufficiently deep* normal subgroups  $N \triangleleft H$  if there exists a finite subset  $\mathcal{F} \subset H \setminus \{1\}$  such that  $\mathcal{P}$  holds whenever  $N \cap \mathcal{F} = \emptyset$ .

**Theorem 1.2** (Theorem 4.13 (see also Theorem 5.7)). *Let  $G$  be a virtually compact cubical hyperbolic group, and suppose that  $G$  is hyperbolic relative to a subgroup  $H$ . Then there exists a finite-index torsion-free normal subgroup  $K \triangleleft H$  such that for every sufficiently deep normal subgroup  $N \triangleleft H$  that satisfies*

- $N \leq K$ , and
- $K/N$  is torsion-free virtually compact cubical hyperbolic,

we have

$$b_1^{(2)}(\overline{G}) \leq b_1^{(2)}(G).$$

By replacing hyperbolicity with the assumption that  $G$  hyperbolic relative to a virtually abelian subgroup  $H$ , we obtain equality of  $L^2$ -Betti numbers in all degrees.

**Theorem 1.3** (Theorem 6.9). *Let  $G$  be a virtually compact cubical group that is hyperbolic relative to a virtually abelian subgroup  $H \leq G$ . Then there exists a finite-index torsion-free normal subgroup  $K \triangleleft H$  such that for every sufficiently deep normal subgroup  $N \triangleleft H$  that satisfies  $N \leq K$  and  $K/N$  is torsion-free, we have*

$$b_*^{(2)}(\overline{G}) = b_*^{(2)}(G).$$

*Remark 1.4.* The torsion-free assumption on  $K/N$  in Theorem 1.2 and Theorem 1.3 cannot be removed. We refer to Remark 4.20 for an example. The passage from  $H$  to its finite-index subgroup  $K$  is more subtle – it can be omitted if the Atiyah Conjecture is preserved by sufficiently deep Dehn fillings (see Remark 4.15).

In several applications, we compute the  $L^2$ -Betti numbers of Dehn fillings associated with cusped arithmetic hyperbolic manifolds. While Theorem 1.3 is a key input, it is itself insufficient for this purpose, as it is still open whether non-uniform arithmetic hyperbolic lattices are virtually compact cubical in general. To address this, we extend Theorem 1.3 to include manifolds with virtually special sparse fundamental groups (see Theorem 6.9). Importantly, Bergeron and Wise [BW12] showed that all cusped arithmetic hyperbolic manifolds have virtually sparse special fundamental groups (see Theorem 7.4). Furthermore, we note that any virtually compact cubical group that is hyperbolic relative to virtually abelian subgroups is itself virtually sparse special (see Proposition 4.12 and Remark 6.4).

**Applications.** Our first application concerns the algebraic fibering of groups, which is intimately related to  $L^2$ -Betti numbers [Kie20, Fis21]. Recall that a group  $G$  is said to  $FP_k(\mathbb{Q})$ -fiber if there is an epimorphism  $G \rightarrow \mathbb{Z}$  whose kernel is of type  $FP_k(\mathbb{Q})$ . In particular,  $FP_1(\mathbb{Q})$ -fibering means that the kernel is finitely generated.

This notion of algebraic fibering originally emerged in the study of 3-manifolds that fiber over the circle. More recently, Kielak [Kie20] and Fisher [Fis21] characterized the virtual algebraic fibering of residually finite rationally solvable (RFRS) groups in terms of the vanishing of the first  $L^2$ -Betti number. Their results offer an algebraic counterpart to Agol's theorem, which was instrumental in proving Thurston's Virtual Fibering Conjecture.

**Corollary 1.5** (Corollary 7.2). *Let  $G$  be a virtually compact cubical hyperbolic group that virtually  $FP_1(\mathbb{Q})$ -fibers, and suppose that  $G$  is hyperbolic relative to its subgroup  $H$ . Then there exists a finite-index torsion-free normal subgroup  $K \triangleleft H$  such that for every sufficiently deep normal subgroup  $N \triangleleft H$  that satisfies  $N \leq K$  and  $K/N$  is torsion-free virtually compact cubical hyperbolic, we have that the group  $G/\langle\langle N \rangle\rangle$  virtually  $FP_1(\mathbb{Q})$ -fibers.*

**Corollary 1.6** (Corollary 7.3). *Let  $G$  be a virtually compact cubical group that is hyperbolic relative to a virtually abelian subgroup  $H \leq G$ , and let  $k \in \mathbb{N}^+$ . Then there exists a finite-index torsion-free normal subgroup  $K \triangleleft H$  such that for every sufficiently deep normal subgroup  $N \triangleleft H$  such that*

- $N \leq K$ , and
- $K/N \cong \mathbb{Z}$ ,

*we have that the group  $G/\langle\langle N \rangle\rangle$  virtually  $FP_k(\mathbb{Q})$ -fibers if and only if so does  $G$ .*

Our next application concerns the Singer Conjecture for Einstein manifolds. Note that Thurston's Hyperbolic Dehn Filling Theorem only holds in dimension 3. This is because

if  $M$  is a manifold with torus boundary and  $\dim M \geq 4$ , then for any Dehn filling  $\overline{M}$  of  $M$ , the fundamental group  $\pi_1(\overline{M})$  contains a  $\mathbb{Z}^2$  subgroup, and thus  $\overline{M}$  cannot admit any hyperbolic Riemannian metric. Nonetheless, Anderson [And06] proved that most Dehn fillings of  $M$  still admit Einstein metrics.

**Corollary 1.7** (Corollary 7.11). *For every cusped arithmetic hyperbolic manifold  $M$  of dimension at least four, there exists a sequence of pairwise non-homeomorphic Anderson-type closed Einstein manifolds  $\{W_n\}_{n \geq 1}$  such that each  $W_n$  satisfies the Singer Conjecture and  $\{W_n\}_{n \geq 1}$  converges to a finite-sheeted cover  $W$  of  $M$  under the pointed Gromov–Hausdorff metric with based point any  $y \in W$ .*

Recall that the Singer Conjecture predicts that the  $L^2$ -Betti numbers of the fundamental group of any closed aspherical manifold are concentrated in the middle dimension. To the best of our knowledge, the above corollary gives the first examples of Anderson-type Einstein manifolds that satisfy the Singer Conjecture.

In recent years, there has been significant progress in understanding the finiteness properties of subgroups of hyperbolic groups. Italiano–Martelli–Migliorini, Isenrich–Martelli–Py, and Isenrich–Py in [IMM23, IMP24, LIP24] have provided new examples of subgroups of hyperbolic groups with unusual finiteness properties, offering counterexamples to long-standing conjectures. By applying Corollary 7.3, we present a new construction of hyperbolic groups with exotic subgroups. An advantage of this approach is that the hyperbolic groups constructed in this way arise as Dehn fillings of any cusped arithmetic hyperbolic manifold of dimension four or higher.

**Theorem 1.8** (Theorem 7.12). *Let  $G$  be the fundamental group of a cusped arithmetic hyperbolic manifold of dimension  $n$ . Then there is a hyperbolic Dehn filling quotient  $\overline{G}$  of  $G$  such that*

- (i) *If  $n = 2q \geq 4$ , then there exist a finite-index subgroup  $\overline{G}_1 \leq \overline{G}$  and an epimorphism  $\psi : \overline{G}_1 \rightarrow \mathbb{Z}$  such that  $\ker \psi$  is of type  $FP_{q-1}(\mathbb{Q})$  but not of type  $FP_q(\mathbb{Q})$ .*
- (ii) *If  $n = 2q + 1 \geq 5$ , then there exist a finite-index subgroup  $\overline{G}_1 \leq \overline{G}$  and an epimorphism  $\psi : \overline{G}_1 \rightarrow \mathbb{Z}$  such that  $\ker \psi$  is of type  $FP(\mathbb{Q})$  but not hyperbolic.*

In [LIMP24], Isenrich, Martelli and Py constructed the first examples of hyperbolic groups which contain subgroups that are of type  $F_3$  but are not of type  $F_4$ . These groups are obtained by Dehn filling a certain non-uniform lattice in  $PO(8, 1)$ . They state that an alternative approach would be to compute the  $L^2$ -Betti numbers of Dehn fillings and pose the following:

**Question 1.9** (Isenrich–Martelli–Py, [LIMP24]). *Let  $\Gamma < PO(2k, 1)$  be a torsion-free non-uniform lattice with purely unipotent parabolic subgroups. Is  $b_k^{(2)}(\overline{\Lambda}) \neq 0$  for a Dehn filling  $\overline{\Lambda}$  of a deep enough finite-index subgroup  $\Lambda < \Gamma$ ?*

We answer this question in the affirmative for arithmetic lattices and sufficiently deep fillings by establishing:

**Corollary 1.10** (Theorem 7.6 (i)). *Let  $\Gamma < PO(2k, 1)$  be a non-uniform arithmetic lattice. Then for every  $\delta > 0$ , all sufficiently deep fillings  $\bar{\Gamma}$  of  $\Gamma$  satisfy*

$$\left| b_k^{(2)}(\bar{\Gamma}) - |\chi(\Gamma)| \right| < \delta.$$

*In particular, if  $\delta$  is sufficiently small, then  $b_k^{(2)}(\bar{\Gamma}) \neq 0$ .*

Finally, we use  $L^2$ -Betti numbers to give an upper bound for deficiency of groups. For a finitely presented group  $G$ , given by a finite presentation, the *deficiency* of the presentation is the number of generators minus the number of relators, and the *deficiency* of  $G$ , denoted as  $\text{defi}(G)$ , is the maximum of the deficiency of all finite presentations of  $G$ .

**Corollary 1.11** (Corollary 7.14). *Let  $G$  be a non-uniform arithmetic lattice in  $SO(n, 1)$ ,  $n \geq 3$ . Denote by  $\{H_i\}_{i=1}^k$  the collection of peripheral subgroups. Then for every family of sufficiently deep normal subgroups  $\{N_i \triangleleft H_i\}_{i=1}^k$ , we have*

$$\text{defi}(\bar{G}) \leq 1.$$

*In addition, if  $G < SO(4, 1)$ , then*

$$\text{defi}(\bar{G}) \leq 1 - \frac{3}{4\pi^2} \text{vol}(\mathbb{H}^4/G).$$

We note that the bounds obtained in Corollary 1.11 do not depend on the chosen Dehn filling and coincide with the known bounds for the deficiency of the lattice  $G$ . Lubotzky in [Lub83, Proposition 6.2] showed that any lattice in  $SO(n, 1)$ ,  $n \geq 3$  has deficiency at most one. Lott in [Lot99] improved Lubotzky's result and in particular showed that any lattice in  $SO(4, 1)$  has a deficiency bound as above.

**Classifying space and finiteness properties.** A key step in our approach of computations of  $L^2$ -Betti numbers is the construction of a tailored classifying space.

**Theorem 1.12** (Corollary 4.3 (see also Corollary 5.2)). *Let  $G$  be a group with a hyperbolically embedded subgroup  $H$ . Let  $\phi: BH \rightarrow BG$  be a cellular map induced by the inclusion  $H \hookrightarrow G$ . For every normal subgroup  $N \triangleleft H$ , let  $\psi_N: BH \rightarrow B\bar{H}$  be a cellular map induced by  $H \rightarrow \bar{H}$ . Let  $X_N$  be the CW-complex obtained by gluing the mapping cylinders  $M_\phi$  and  $M_{\psi_N}$  along their common subcomplex  $BH$ . Then for sufficiently deep  $N \triangleleft H$ , the complex  $X_N$  is a  $K(\bar{G}, 1)$ -space.*

In addition to the computation of  $L^2$ -Betti numbers, the above theorem allows us to deduce geometric finiteness properties of Dehn fillings, which are analogs of the algebraic finiteness properties obtained by the authors in [PS24, Corollary 2.2 (iii)].

**Corollary 1.13** (Corollary 4.4 (see also Corollary 5.3)). *Let  $G$  be a group with a hyperbolically embedded subgroup  $H \hookrightarrow_h G$ .*

- (i) *If  $G$  is of type  $F_n$  for some  $n \in \mathbb{N}^+ \cup \{\infty\}$ , then for all sufficiently deep normal  $N \triangleleft H$ , we have that  $\bar{G}$  is of type  $F_n$  if and only if  $\bar{H}$  is of type  $F_n$ .*
- (ii) *If both  $G$  and  $H$  are of type  $F$ , then for all sufficiently deep normal  $N \triangleleft H$  such that  $\bar{H}$  is of type  $F$ , we have that  $\bar{G}$  is of type  $F$ .*

*Remark 1.14.* Analogs of other results of [PS24] with geometric finiteness properties in place of algebraic finiteness properties can be deduced using the above corollary. Since the proofs of those results are almost identical with their counterparts in [PS24], we will not include them in this paper.

**Our approach on computing  $L^2$ -Betti numbers.** For any group  $G$  and a  $G$ -CW-complex  $X$ , we denote the  $L^2$ -Betti numbers associated with the action of  $G$  on  $X$  by  $b_*^{(2)}(X; G)$ . Let  $EG$  be the universal cover of a  $K(G, 1)$ -CW-complex. A special case of Lück's Approximation Conjecture for  $L^2$ -Betti numbers, proved by Jaikin-Zapirain and López-Álvarez in [JZLA20], asserts that, if  $\{M_i\}_{i \geq 1}$  is a nested sequence of normal subgroups of  $G$  with trivial intersection, then under certain conditions, one obtains

$$b_*^{(2)}(EG; G) = \lim_{i \rightarrow \infty} b_*^{(2)}(M_i \backslash EG; G/M_i).$$

Note that the above result deals with a sequence of normal subgroups rather than a single one. To apply this in the Dehn filling setting, we first slightly reformulate their approximation result by showing that, for every  $\delta > 0$ , there exists a finite subset  $\mathcal{S}_\delta \subset G \setminus \{1\}$  such that if a normal subgroup  $M \triangleleft G$  satisfies  $M \cap \mathcal{S}_\delta = \emptyset$ , then

$$|b_*^{(2)}(M \backslash EG; G/M) - b_*^{(2)}(EG; G)| < \delta.$$

Two problems still remain. First, the above only gives an estimate rather than a computation of  $L^2$ -Betti numbers. Secondly,  $b_*^{(2)}(M \backslash EG; G/M)$  may not equal  $b_*^{(2)}(G/M)$  as  $M \backslash EG$  is not contractible unless  $M = \{1\}$ . We use the properties of Dehn fillings to resolve these as follows:

- 1) Assuming  $G$  is virtually compact cubical, and is either hyperbolic or relatively hyperbolic with respect to abelian subgroups, we use different versions of the Malnormal Special Quotient Theorem, proved by Agol–Groves–Manning [AGM16] and Wise [Wis21], to ensure that  $G/M$  satisfies the Atiyah Conjecture, which asserts that  $b_*^{(2)}(M \backslash EG; G/M)$  takes value in a discrete subgroup of  $\mathbb{Q}$ , and thus must be equal to  $b_*^{(2)}(EG; G)$  provided  $\delta$  is sufficiently small.
- 2) By using the Cohen–Lyndon property for sufficiently deep Dehn fillings of hyperbolically embedded subgroups of  $G$ , established by the second authored in [Sun20], we construct a  $K(G/M, 1)$ -space suitable for our applications. We refer to this construction as a *Dehn filling space*, which is summarized in Theorem 1.12. An excision argument on this space relates  $b_*^{(2)}(M \backslash EG; G/M)$  with  $b_*^{(2)}(G/M)$ .

Finally, it is worth highlighting that the choice of finite subsets that the sufficiently deep normal subgroups must avoid in the main results depends not only on the Cohen–Lyndon property but also on the quantitative version of Lück's Approximation Conjecture.

### Ingredients in the proof of Theorem 1.3 and its generalisation Theorem 6.9:

- Lück's Approximation Conjecture for virtually locally indicable groups [JZLA20] and its quantitative version Theorem 3.2.



- Cohen–Lyndon property for acylindrically [Sun20] and relatively [GMS19] hyperbolic groups.
- Construction of the Dehn filling space for Dehn filling quotients, Theorem 4.2.
- Geometric finiteness properties of Dehn filling quotients, Corollary 4.4.
- Inequalities for  $L^2$ -Betti numbers of Dehn filling quotients of acylindrically hyperbolic virtually locally indicable groups, Theorem 4.7.
- Embeddability of finitely generated special groups in finitely generated right-angled Artin groups, [HW08, Theorem 1.1].
- The Malnormal Special Quotient Theorem [Wis21, AGM16] and its relative version [Wis21].
- A weaker version of the Atiyah Conjecture for certain virtually locally indicable groups, Proposition 3.5.
- The Atiyah Conjecture for virtually compact special groups, [Sch15].
- Separability of abelian subgroups in virtually special groups, [Wis21, Corollary 6.8].
- Structure of sparse virtually special cube complexes whose fundamental group is hyperbolic relative to virtually abelian groups, [Wis21, Lemma 7.51, Theorem 7.54].

**Structure of the paper.** We recall the necessary background and notation in Section 2, and then, in Section 3, we derive a quantitative version of the result in [JZLA20] on Lück’s Approximation Conjecture. In Section 4, we first prove Theorem 1.12, and then use it to deduce Theorem 1.1 and Theorem 1.2. The proof of Theorem 1.3 proceeds in two steps. First, we prove Theorem 4.18, which is a special case of Theorem 1.3 where the peripheral subgroup is abelian. Then, in Section 5, we state the general versions of Theorem 1.1, Theorem 1.2 and Theorem 4.18 for a finite family of peripheral subgroups. In Section 6, we consider virtually sparse special groups and use the general results of Section 5 to prove a general version of Theorem 1.3. Finally, we deduce our applications in Section 7.

**Acknowledgments.** First, the authors thank Kevin Li, who asked us whether the algebraic excision formula obtained in [PS24] could be derived using a topological excision. The pursuit of an answer to this question led us to the construction of the Dehn filling space, which became the catalyst for the remaining results of this paper. We are indebted to Mark Hagen for explaining to us how virtually sparse special groups may be used to understand cubulations of hyperbolic arithmetic lattices. The authors are also grateful to Robert Bell, Luca Di Cerbo, Chris Hruska, Wolfgang Lück, Jason Manning, Ashot Minasyan and Alan Reid for their helpful conversations.

## 2. PRELIMINARIES

**2.1. Finiteness properties.** Throughout this paper, all actions will be left actions. Let  $G$  be a group. We use the terminology  $G$ -CW-complex for a  $G$ -space  $X$  as defined by Lück [Lüc02, Definition 1.25]. This is equivalent to the notion of an *admissible  $G$ -complex* in the sense of [Bro94, Chapter IX, §10], characterised by the property that for any cell  $\sigma$  of  $X$ , if an element of  $G$  stabilises  $\sigma$ , then it must fix  $\sigma$  pointwise.

The subspace  $X_n$  is called the  $n$ -skeleton of  $X$ . We say that  $X$  is a *free  $G$ -CW complex* if in addition the action  $G \curvearrowright X$  is free. The  $G$ -CW complex  $X$  is *of type  $F_n$*  for some

$n \in \mathbb{N}$  (resp. *type*  $F_\infty$ ) if  $G \setminus X$  has finite  $n$ -skeleton (resp. has finitely many cells in each dimension).

Next, we define the necessary finiteness notions of groups and mention their relevant properties. For more details, we refer the reader to Chapter VIII of [Bro94].

Let  $R$  denote a commutative ring with unit and  $G$  denote a discrete group. The *cohomological dimension of  $G$  over  $R$*  can be defined by

$$\text{cd}_R G = \sup\{n \in \mathbb{N} \mid H^n(G, M) \neq 0 \text{ for some } RG\text{-module } M\}.$$

When  $R = \mathbb{Z}$ , we omit it from the notation and denote the cohomological dimension by  $\text{cd } G$ . A group  $G$  is said to have finite *virtual cohomological dimension* equal to  $n$ , denoted  $\text{vcd } G$ , if  $G$  has a subgroup  $H$  of finite index with  $\text{cd } H = n$ . This is a well-defined notion by [Bro94, VIII.2.4]. A group  $G$  is said to be of *type*  $FP_n(R)$  for some  $n \in \mathbb{Z}_{\geq 0} \cup \{\infty\}$  if there is a projective  $RG$ -resolution

$$\cdots \rightarrow P_2 \rightarrow P_1 \rightarrow P_0 \twoheadrightarrow R$$

such that  $P_i$  is finitely generated and projective as an  $RG$ -module for all  $i \leq n$ . A projective  $RG$ -resolution  $P_* \twoheadrightarrow R$  is said to be of *length*  $n$ , if  $P_n \neq 0$  and  $P_i = 0$  for all  $i > n$ . A group  $G$  is said to be of *type*  $FP(R)$ , if there is a projective  $RG$ -resolution of  $R$  by finitely generated projective  $RG$ -modules of finite length. It is immediate from the definitions that if  $G$  is of type  $FP(R)$ , then it is of type  $FP_\infty(R)$ .

Denote  $\text{cd}_R G = n$ . By [Bro94, VIII.2.1], it follows that  $G$  is of type  $FP(R)$  if and only if  $n < \infty$  and  $G$  is of type  $FP_n(R)$ .

There are analogous topological finiteness properties which we explain next.

An Eilenberg-MacLane space or  $K(G, 1)$ -complex is an aspherical CW complex with fundamental group equal to  $G$ . A group  $G$  is of *type*  $F_n$  if there is a  $K(G, 1)$ -complex  $X$  with finite  $n$ -skeleton and it is of *type*  $F$  if  $X$  is finite. If  $G$  is of type  $F_n$  or  $F$ , then it is of type  $FP_n(\mathbb{Z})$  or  $FP$ , respectively. A group  $G$  is finitely generated if and only if it is of type  $F_1$  or  $FP_1(R)$  [Bie81, 1.2.1]. A group  $G$  is finitely presented if and only if it is of type  $F_2$ .

For the rest of this paper, by  $EG$  we mean a contractible free  $G$ -CW complex, and by  $BG$  we mean the quotient  $G \setminus EG$ . If we do not specify the choice of  $EG$ , then we simply choose an arbitrary contractible free  $G$ -CW complex as  $EG$ . Such a complex always exists for any group  $G$  and is a terminal object in the homotopy category of free  $G$ -CW complexes. In particular, any two choices for  $EG$  are  $G$ -homotopy equivalent [LÖ5]. In certain situations we may also specify the choice so that  $EG$  (resp.  $BG$ ) has a certain property, say  $\mathcal{P}$ , in which case we will say that let  $EG$  (resp.  $BG$ ) be a contractible free  $G$ -CW complex (resp.  $K(G, 1)$ -space) satisfying  $\mathcal{P}$ .

**2.2. Von Neumann rank of a matrix.** Let  $G$  be a discrete group. Recall that  $\ell^2(G)$  denotes the Hilbert space of square-summable complex-valued functions on  $G$  and there is a left  $G$ -action on  $\ell^2(G)$ , called the *left regular representation*. The *group von Neumann algebra* of  $G$ , denoted  $\mathcal{N}(G)$ , is the set of all bounded  $G$ -equivariant linear operators on  $\ell^2(G)$ . The *von Neumann dimension*  $\dim_G$  of  $G$  is a function from the set of  $\mathcal{N}(G)$ -modules to  $[0, \infty]$ . We refer to [Lüc02, Chapter 6] for the precise definition of  $\dim_G$ . For the purpose of this paper, we only collect some properties of  $\dim_G$ .



**Proposition 2.1.** *Let  $G$  be a group. The following properties hold.*

- (i) *If  $M$  is a finitely generated  $\mathcal{N}(G)$ -module, then  $\dim_G M < \infty$ .*
- (ii) *If  $M$  and  $N$  are isomorphic  $\mathcal{N}(G)$ -modules, then  $\dim_G M = \dim_G N$ .*
- (iii) *If  $0 \rightarrow M \rightarrow N \rightarrow P \rightarrow 0$  is an exact sequence of  $\mathcal{N}(G)$ -modules, then*

$$\dim_G N = \dim_G M + \dim_G P,$$

*where the laws of the summation of  $[0, \infty)$  extends naturally to  $[0, \infty]$ .*

- (iv) *Let  $\{M_i \mid i \in I\}$  be a cofinal system of  $\mathcal{N}(G)$ -submodules of an  $\mathcal{N}(G)$ -module  $M$ , i.e.,  $M = \bigcup_{i \in I} M_i$  and for every two indices  $i, j \in I$  there exists an index  $k \in I$  such that  $M_i, M_j \subset M_k$ . Then  $\dim_G M = \sup\{\dim_G M_i \mid i \in I\}$ .*

Let  $I, J$  be indexing sets. A *tamed matrix* over  $\mathbb{C}G$  of dimension  $I \times J$  is a (possibly infinite) matrix  $A = (a_{ij})_{i \in I, j \in J}$  such that for all  $i, j$ , we have  $a_{ij} \in \mathbb{C}G$ , and for all  $i \in I$ , there are only finitely many  $j \in J$  such that  $a_{ij} \neq 0$ . If  $|I|, |J| < \infty$ , then we call  $A$  a *finite matrix*.

Consider the  $\mathcal{N}(G)$ -modules  $(\mathcal{N}(G))^I$  and  $(\mathcal{N}(G))^J$ , whose elements are of the form  $(x_i)_{i \in I}$  and  $(y_j)_{j \in J}$ , respectively, where  $x_i, y_j \in \mathcal{N}(G)$  for all  $i, j$  and  $x_i = y_j = 0$  for all but finitely many  $i \in I, j \in J$ .

The right-multiplication of a tamed matrix  $A = (a_{ij})_{i \in I, j \in J}$  over  $\mathbb{C}G$  induces a  $\mathcal{N}(G)$ -module homomorphism  $r_A: (\mathcal{N}(G))^I \rightarrow (\mathcal{N}(G))^J$ , i.e.,

$$r_A(x) := x \cdot A = \left( \sum_{i \in I} x_i \cdot a_{ij} \right)_{j \in J}.$$

The *von Neumann rank* of  $A$  is

$$\text{rk}_G(A) := \dim_G \text{im}(r_A) \in [0, \infty].$$

If  $N \triangleleft G$  is a normal subgroup, then the quotient map  $G \rightarrow G/N$  extends to a ring homomorphism  $\mathbb{C}G \rightarrow \mathbb{C}[G/N]$ , which, when applied to each entry of  $A$ , gives us a matrix  $\epsilon_N(A)$ . For simplicity, we will denote

$$\text{rk}_{G/N}(A) := \text{rk}_{G/N}(\epsilon_N(A)).$$

**2.3. L<sup>2</sup>-Betti numbers and (Strong) Atiyah Conjecture.** Let  $G$  be a group, let  $X$  be a  $G$ -CW complex, and let  $C_*(X)$  be the cellular  $\mathbb{Z}G$ -chain complex of  $X$ , with boundary map  $\partial_*$ . Suppose  $M$  is a  $\mathbb{Z}G$ -module. We can thus form the tensor product chain complex  $M \otimes_{\mathbb{Z}G} C_*(X)$ , whose boundary map is  $\tilde{\partial}_* = \text{id}_M \otimes \partial_*$ . The homology of this chain complex is denoted by  $H_*^G(X; M)$ . Define

$$H_*(G; M) := H_*^G(EG; M).$$

If  $Y \subset X$  is a  $G$ -CW subcomplex, then we can form a relative chain complex  $M \otimes_{\mathbb{Z}G} C_*(X, Y)$ , where  $C_*(X, Y)$  is the relative cellular chain complex of the pair  $(X, Y)$ . The homology of this chain complex will be denoted by  $H_*^G(X, Y; M)$ . For any subgroup  $H \leq G$ , the inclusion  $H \hookrightarrow G$  induces a classifying map  $\phi: G \times_H EH \rightarrow EG$ . Let  $M_\phi$  be the mapping cylinder of  $\phi$ . Define

$$H_*(G, H; M) := H_n^G(M_\phi, G \times_H EH; M).$$

This definition is equivalent to the original definition of relative group homology given by Bieri–Eckmann [BE78]. The cohomology groups  $H_G^*(X; M)$  and  $H_G^*(X, Y; M)$  are defined similarly where instead of tensoring with  $M \otimes_{\mathbb{Z}G} -$ , one applies  $\text{Hom}_{\mathbb{Z}G}(-, M)$  to the corresponding chain complexes.

There is a natural embedding  $\mathbb{Z}G \hookrightarrow \mathcal{N}(G)$ , and thus the group von Neumann algebra  $\mathcal{N}(G)$  becomes a right  $\mathbb{Z}G$ -module. If we let  $M = \mathcal{N}(G)$ , then the homology  $H_*^G(X; \mathcal{N}(G))$ , is naturally a left  $\mathcal{N}(G)$ -module. In this case, to emphasize the  $\mathcal{N}(G)$ -module structure, we denote

$$\begin{aligned}\mathcal{H}_*(X; \mathcal{N}(G)) &:= H_*^G(X; \mathcal{N}(G)), \\ \mathcal{H}_*(G; \mathcal{N}(G)) &:= H_*(G; \mathcal{N}(G)).\end{aligned}$$

We denote

$$b_*^{(2)}(X; G) := \dim_G \mathcal{H}_*(X; \mathcal{N}(G)), \quad b_*^{(2)}(G) := \dim_G \mathcal{H}_*(G; \mathcal{N}(G))$$

and call them the  $L^2$ -Betti numbers of  $X$  and  $G$ , respectively. Similarly, denote

$$b_*^{(2)}(X, Y; G) := \dim_G \mathcal{H}_*(X, Y; \mathcal{N}(G)), \quad b_*^{(2)}(G, H) := \dim_G \mathcal{H}_*(G, H; \mathcal{N}(G)).$$

*Remark 2.2.* For any two choices of  $EG$ , there exists a natural  $\mathcal{N}(G)$ -module isomorphism between the corresponding  $\mathcal{H}_*(G; \mathcal{N}(G))$ , and for any two choices of  $EG$  and any two choices of  $EH$ , there exists a natural  $\mathcal{N}(G)$ -module isomorphism between the corresponding  $\mathcal{H}_n(G, H; \mathcal{N}(G))$ .

The next lemma follows easily from [Lüc02, Theorem 6.54 (6)].

**Lemma 2.3.** *Let  $G$  be a group. Then for all  $n \geq \text{vcd}(G) + 1$ , we have  $b_n^{(2)}(G) = 0$ .*

For any group  $G$ , we will denote

$$\mathcal{FLN}(G) := \{n \in \mathbb{N}^+ \mid \exists \text{ subgroup } H \leq G \text{ with } |H| = n\}.$$

Let

$$\frac{1}{\mathcal{FLN}(G)} \cdot \mathbb{Z}$$

be the subgroup of the additive group of  $\mathbb{Q}$  generated by elements in the set

$$\left\{ \frac{1}{n} \mid n \in \mathcal{FLN}(G) \right\}.$$

**Conjecture 2.4** ((Strong) Atiyah Conjecture). *For every free  $G$ -CW complex  $X$  of type  $F_{k+1}$  for some  $k \in \mathbb{N}$  and every  $n \leq k$ ,*

$$b_n^{(2)}(X; G) \in \frac{1}{\mathcal{FLN}(G)} \cdot \mathbb{Z}.$$

The Atiyah conjecture offers a powerful framework for understanding topological and homological properties of spaces. We will exploit the specific cases for which it is already known to support some of our main arguments.

**2.4. Hyperbolically embedded subgroups.** Let  $G$  be a group,  $\{H_\lambda\}_{\lambda \in \Lambda}$  a family of subgroups of  $G$ ,  $X$  a subset of  $G$  such that  $G$  is generated by  $X$  together with the union of all  $H_\lambda$  (in which case  $X$  is called a *relative generating set* of  $G$  with respect to  $\{H_\lambda\}_{\lambda \in \Lambda}$ ), and  $\mathcal{H} = \bigsqcup_{\lambda \in \Lambda} H_\lambda$ . Consider the Cayley graph  $\Gamma(G, X \sqcup \mathcal{H})$ . Note that, for  $\lambda \in \Lambda$  there is a natural embedding  $\Gamma(H_\lambda, H_\lambda) \hookrightarrow \Gamma(G, X \sqcup \mathcal{H})$  whose image is the subgraph of  $\Gamma(G, X \sqcup \mathcal{H})$  with vertices and edges labeled by elements of  $H_\lambda$ .

*Remark 2.5.* For distinct  $\lambda, \mu \in \Lambda$ , we allow  $X \cap H_\lambda \neq \emptyset$  and  $H_\lambda \cap H_\mu \neq \{1\}$ , in which case there will be multiple edges between some pairs of vertices of  $\Gamma(G, X \sqcup \mathcal{H})$ .

For  $\lambda \in \Lambda$ , an edge path in  $\Gamma(G, X \sqcup \mathcal{H})$  between vertices of  $\Gamma(H_\lambda, H_\lambda)$  is called  $H_\lambda$ -*admissible* if it does not contain any edge of  $\Gamma(H_\lambda, H_\lambda)$ . Note that an  $H_\lambda$ -admissible path is allowed to pass through vertices of  $\Gamma(H_\lambda, H_\lambda)$ .

**Definition 2.6.** For every pair of elements  $h, k \in H_\lambda$ , let  $\hat{d}_\lambda(h, k) \in [0, \infty]$  be the length of a shortest  $H_\lambda$ -admissible path connecting the vertices labeled by  $h$  and  $k$ . If no such path exists, set  $\hat{d}_\lambda(h, k) = \infty$ . The laws of summation on  $[0, \infty)$  extend naturally to  $[0, \infty]$  and it is easy to verify that  $\hat{d}_\lambda : H_\lambda \times H_\lambda \rightarrow [0, +\infty]$  defines a metric on  $H_\lambda$ , which is called the *relative metric on  $H_\lambda$  with respect to  $X$* .

**Definition 2.7.** We say that the family  $\{H_\lambda\}_{\lambda \in \Lambda}$  *hyperbolically embeds into  $(G, X)$*  (denoted by  $\{H_\lambda\}_{\lambda \in \Lambda} \hookrightarrow_h (G, X)$ ) if the following hold:

- (i)  $G$  is generated by the set  $X$  together with the union of all  $H_\lambda$ ,  $\lambda \in \Lambda$ .
- (ii) The Cayley graph  $\Gamma(G, X \sqcup \mathcal{H})$  is a Gromov hyperbolic space.
- (iii) For each  $\lambda \in \Lambda$ , the metric space  $(H_\lambda, \hat{d}_\lambda)$  is proper, i.e., every ball of finite radius contains only finitely many elements

Further, we say that the family  $\{H_\lambda\}_{\lambda \in \Lambda}$  *hyperbolically embeds into  $G$* , denoted by  $\{H_\lambda\}_{\lambda \in \Lambda} \hookrightarrow_h G$ , if there exists some subset  $X \subset G$  such that  $\{H_\lambda\}_{\lambda \in \Lambda} \hookrightarrow_h (G, X)$ . And we say that  $G$  is *hyperbolic relative to the family  $\{H_\lambda\}_{\lambda \in \Lambda}$*  or the family  $\{H_\lambda\}_{\lambda \in \Lambda}$  is a *peripheral structure* on  $G$  if  $|\Lambda| < \infty$  and there exists a finite set  $X \subset G$  such that  $\{H_\lambda\}_{\lambda \in \Lambda} \hookrightarrow_h (G, X)$ . If  $\{H_\lambda\}_{\lambda \in \Lambda}$  consists of a single element  $H$  we will drop the braces and write that  $H \hookrightarrow_h (G, X)$  (resp.  $H \hookrightarrow_h G$ ,  $G$  is hyperbolic relative to  $H$ ) instead of  $\{H\} \hookrightarrow_h (G, X)$  (resp.  $\{H\} \hookrightarrow_h G$ ,  $G$  is hyperbolic relative to  $\{H\}$ ).

*Remark 2.8.* Our definition of relative hyperbolicity is not one of the standard definitions, but is equivalent to those by [DGO17, Proposition 4.28].

**2.5. Dehn filling.** Dehn filling is a process which takes input a group  $G$ , a family of subgroups  $\{H_\lambda\}_{\lambda \in \Lambda}$  of  $G$ , and a family of normal subgroups  $\{N_\lambda \triangleleft H_\lambda\}_{\lambda \in \Lambda}$ , and outputs the quotient group  $G/\langle\langle \bigcup_{\lambda \in \Lambda} N_\lambda \rangle\rangle_G$ . Here and in the sequel, for any subset  $S$  of a group  $G$ , we denote by  $\langle\langle S \rangle\rangle_G$  the normal closure of  $S$  in  $G$ . If  $G$  is understood, we will simply write  $\langle\langle S \rangle\rangle$ .

In practice, it is often assumed that  $G$  and  $\{H_\lambda\}_{\lambda \in \Lambda}$  satisfy certain conditions, so that for most choices of  $\{N_\lambda \triangleleft H_\lambda\}_{\lambda \in \Lambda}$ , the quotient  $G/\langle\langle \bigcup_{\lambda \in \Lambda} N_\lambda \rangle\rangle_G$  satisfy certain properties. In this paper, we always assume that  $\{H_\lambda\}_{\lambda \in \Lambda}$  hyperbolically embeds into  $G$ , in which case the Cohen–Lyndon property arises naturally. To simplify the statements of our results, we use the following terminology, which aligns with the terminology of 3-manifold theory.

**Definition 2.9.** Let  $G \geq H$  be groups, and let  $\mathcal{P} = \mathcal{P}(G, H, N)$ ,  $\mathcal{Q} = \mathcal{Q}(G, H, N)$  be properties formulated for all normal subgroups  $N \triangleleft H$ . We say that  $\mathcal{P}$  holds for sufficiently deep normal subgroups  $N \triangleleft H$  if there exists a finite subset  $\mathcal{F} \subset H \setminus \{1\}$  such that  $\mathcal{P}$  holds whenever  $N \cap \mathcal{F} = \emptyset$ . We say that  $\mathcal{P}$  holds for sufficiently deep normal subgroups  $N \triangleleft H$  such that  $\mathcal{Q}$  holds if there exists a finite subset  $\mathcal{F} \subset H \setminus \{1\}$  such that, if  $N \cap \mathcal{F} = \emptyset$  and  $\mathcal{Q}$  holds, then  $\mathcal{P}$  holds.

*Remark 2.10.* Let  $G \geq H$  be groups. Note that, if  $|H| < \infty$  then a property  $\mathcal{P}$  holds for sufficiently deep normal subgroup  $N \triangleleft H$  if and only if it holds for  $N = \{1\}$ .

There are two key ingredients in our study of Dehn filling. The first one is the following result which roughly says that hyperbolicity is preserved by sufficiently deep Dehn filling:

**Theorem 2.11** (Dahmani–Guirardel–Osin [DGO17]). *Let  $G$  be a group with a subgroup  $H \hookrightarrow_h G$ . Then for sufficiently deep  $N \triangleleft H$ , the natural homomorphism  $H/N \rightarrow G/\langle\langle N \rangle\rangle$  maps  $H/N$  injectively onto a hyperbolically embedded subgroup of  $G/\langle\langle N \rangle\rangle$ .*

The second is a structural result about the kernel associated with a sufficiently deep Dehn filling, termed as a *Cohen–Lyndon triple*, which was introduced by [Sun20], but the idea dates back to a paper by Cohen–Lyndon [CL63], hence the name. This property is crucial for our construction of a classifying space of the quotient group arising from a Dehn filling process.

**Definition 2.12.** Let  $N \triangleleft H \leq G$  be a triple of groups. We say that the triple  $(G, H, N)$  is a *Cohen–Lyndon triple* if there exists a left transversal  $T$  of  $H\langle\langle N \rangle\rangle$  in  $G$  such that  $\langle\langle N \rangle\rangle$  decomposes as a free product of the following form  $\langle\langle N \rangle\rangle = \ast_{t \in T} t N t^{-1}$ .

**Theorem 2.13** ([Sun20, Theorem 2.5]). *Let  $G$  be a group with a hyperbolically embedded subgroup  $H \hookrightarrow_h G$ . Then for all sufficiently deep normal subgroups  $N \triangleleft H$ , the triple  $(G, H, N)$  is a Cohen–Lyndon triple.*

In the subsequent sections, we will often need to pass to a finite-index normal subgroup of a (sufficiently deep) Dehn filling quotient of a group  $G$ . Lemma 2.16 below roughly states that such a finite-index normal subgroup arises as a (sufficiently deep)  $G$ -equivariant Dehn filling of a finite-index normal subgroup of  $G$ . Let us start with the related definitions.

**Definition 2.14.** Let  $G$  be a group that is hyperbolic relative to a family of subgroups  $\{H_i\}_{i=1}^k$ , let  $G_0 \triangleleft G$  be a finite-index normal subgroup. For all  $i$ , let  $H_{0i} = G_0 \cap H_i$ , and let  $\mathcal{H}_i$  be the family of subgroups of  $G_0$  obtained by choosing one element from each  $G_0$ -conjugacy class of the set  $\{gH_{0i}g^{-1} \mid g \in G\}$ . Let  $\mathcal{H} = \bigcup_{i=1}^k \mathcal{H}_i$ . Then  $\mathcal{H}$  is a finite family of subgroups of  $G_0$  and  $G_0$  is hyperbolic relative to  $\mathcal{H}$  by [Hru10, Theorem 9.1]. We call  $\mathcal{H}$  the *peripheral structure induced by  $\{H_i\}_{i=1}^k$* .

**Definition 2.15.** Let  $G$  be a group hyperbolic relative to a family of subgroups, and let  $G_0 \triangleleft G$  be a finite-index normal subgroup with the induced peripheral structure  $\mathcal{H}$ . We say that a family of normal subgroups  $\{N_H \triangleleft H \mid H \in \mathcal{H}\}$  is  $G$ -equivariant or the Dehn filling of  $G_0$  induced by this family is  $G$ -equivariant if, for all  $H \in \mathcal{H}$  and  $g \in G$  such that  $gHg^{-1} \in \mathcal{H}$ , we have  $N_{gHg^{-1}} = gN_Hg^{-1}$ .

In the following lemma, as we will discuss normal closures in different groups, we will use  $\langle\langle \cdot \rangle\rangle_G$  instead of  $\langle\langle \cdot \rangle\rangle$ .

**Lemma 2.16.** *Let  $G$  be a group that is hyperbolic relative to a family of infinite subgroups  $\{H_i\}_{i=1}^k$ , and let  $G_0 \triangleleft G$  be a finite-index normal subgroup, endowed with the induced peripheral structure  $\{K_j\}_{j=1}^\ell$  from  $G$ . Then the following hold:*

- (i) *There is a bijection  $\varphi: \mathcal{T} \rightarrow \mathcal{S}$ , where  $\mathcal{S}$  is the set of all families of normal subgroups  $\{N_i \triangleleft H_i\}_{i=1}^k$  such that  $N_i \leq G_0 \cap H_i$  for all  $i$ , and  $\mathcal{T}$  is the set of all  $G$ -equivariant families of normal subgroups  $\{M_j \triangleleft K_j\}_{j=1}^\ell$ .*
- (ii) *For  $k = 1, 2$ , let  $S_k = \{N_{k,i} \triangleleft H_i\}_{i=1}^k \in \mathcal{S}$  such that  $N_{1,i} \leq N_{2,i}$  for all  $i$ , and let  $\varphi^{-1}(S) = \{M_{k,j} \triangleleft K_j\}_{j=1}^\ell$ . Then for each  $j \in \{1, \dots, \ell\}$ , we have  $M_{1,j} \leq M_{2,j}$  and there exists  $i \in \{1, \dots, k\}$  and  $g \in G$  such that*

$$gN_{k,i}g^{-1} = M_{k,j}, \quad K_j/M_{k,j} \cong (G_0 \cap H_i)/N_{k,i}, \quad N_{2,i}/N_{1,i} \cong M_{2,j}/M_{1,j}.$$

- (iii) *For each  $T \in \mathcal{T}$ , the following diagram commutes*

$$\begin{array}{ccc} G_0 & \longrightarrow & G_0 / \langle\langle \bigcup_{M \in T} M \rangle\rangle_{G_0} \\ \downarrow & & \downarrow \\ G & \longrightarrow & G / \langle\langle \bigcup_{N \in \varphi(T)} N \rangle\rangle_G \end{array}$$

and we have

$$(1) \quad [G / \langle\langle \bigcup_{N \in \varphi(T)} N \rangle\rangle_G : G_0 / \langle\langle \bigcup_{M \in T} M \rangle\rangle_{G_0}] = [G : G_0].$$

- (iv) *For each family of finite subsets  $\{\mathcal{F}_i \subset H_i \setminus \{1\}\}_{i=1}^k$ , there exists a family of finite subsets  $\{\mathcal{F}'_j \subset K_j \setminus \{1\}\}_{j=1}^\ell$  such that the following hold: let  $S = \{N_i \triangleleft H_i\}_{i=1}^k \in \mathcal{S}$ , and let  $\varphi^{-1}(S) = \{M_j \triangleleft K_j\}_{j=1}^\ell$ . Then  $M_j \cap \mathcal{F}'_j = \emptyset$  for all  $j$  implies  $N_i \cap \mathcal{F}_i = \emptyset$  for all  $i$ .*

*Conversely, for each family of finite subsets  $\{\mathcal{F}'_j \subset K_j \setminus \{1\}\}_{j=1}^\ell$ , there exists a family of finite subsets  $\{\mathcal{F}_i \subset H_i \setminus \{1\}\}_{i=1}^k$  such that the following hold: let  $T = \{M_j \triangleleft K_j\}_{j=1}^\ell \in \mathcal{T}$  and let  $\varphi(T) = \{N_i \triangleleft H_i\}_{i=1}^k$ . Then  $N_i \cap \mathcal{F}_i = \emptyset$  for all  $i$  implies  $M_j \cap \mathcal{F}'_j = \emptyset$  for all  $j$ .*

- (v) *For each family of finite-index subgroups  $\{P_i \leq H_i\}_{i=1}^k$ , there exists a family of finite-index subgroups  $\{Q_j \leq K_j\}_{j=1}^\ell$  such that the following hold: let  $S = \{N_i \triangleleft H_i\}_{i=1}^k \in \mathcal{S}$ , and let  $\varphi^{-1}(S) = \{M_j \triangleleft K_j\}_{j=1}^\ell$ . Then  $M_j \leq Q_j$  for all  $j$  implies  $N_i \leq P_i$  for all  $i$ .*

*Conversely, for each family of finite-index subgroups  $\{Q'_j \leq K_j\}$ , there exists a family of finite-index subgroups  $\{P'_i \leq H_i\}_{i=1}^k$  such that the following hold: let  $T = \{M_j \triangleleft K_j\}_{j=1}^\ell \in \mathcal{T}$  and let  $\varphi(T) = \{N_i \triangleleft H_i\}_{i=1}^k$ . Then  $N_i \leq P'_i$  for all  $i$  implies  $M_j \leq Q'_j$  for all  $j$ .*

*Proof.* For each  $j \in \{1, \dots, \ell\}$ , choose  $\alpha(j) \in \{1, \dots, k\}$  and  $g_j \in G$  such that

$$K_j = g_j(G_0 \cap H_{\alpha(j)})g_j^{-1}.$$

For each  $i \in \{1, \dots, k\}$ , choose  $\beta(i) \in \{1, \dots, \ell\}$  such that  $\alpha \circ \beta(i) = i$ . Therefore,

$$K_{\beta(i)} = g_{\beta(i)}(G_0 \cap H_i)g_{\beta(i)}^{-1}.$$

(i). Let  $\{M_j \triangleleft K_j\}_{j=1}^\ell$  be a  $G$ -equivariant family of normal subgroups. For each  $i$ , let

$$N_i = g_{\beta(i)}^{-1} M_{\beta(i)} g_{\beta(i)}.$$

Then  $N_i \leq G_0 \cap H_i$ . We need to show that  $N_i$  is normal in  $H_i$ . Let  $h \in H_i$ . As  $G_0 \triangleleft G$  we have  $g_{\beta(i)}^{-1} K_{\beta(i)} g_{\beta(i)} = G_0 \cap H_i \triangleleft H_i$  and thus  $g_{\beta(i)} h g_{\beta(i)}^{-1} K_{\beta(i)} g_{\beta(i)} h^{-1} g_{\beta(i)}^{-1} = K_{\beta(i)}$ . As the family  $\{M_j \triangleleft K_j\}_{j=1}^\ell$  is  $G$ -equivariant, we have  $g_{\beta(i)} h g_{\beta(i)}^{-1} M_{\beta(i)} g_{\beta(i)} h^{-1} g_{\beta(i)}^{-1} = M_{\beta(i)}$ , which is equivalent to  $h N_i h^{-1} = N_i$  as desired. Next, define

$$\varphi(\{M_j \triangleleft K_j\}_{j=1}^\ell) := \{N_i\}_{i=1}^k = \{g_{\beta(i)}^{-1} M_{\beta(i)} g_{\beta(i)}\}_{i=1}^k.$$

We need to construct the inverse of  $\varphi$ . For each  $i \in \{1, \dots, k\}$ , let  $N'_i \triangleleft H_i$  such that  $N'_i \leq G_0 \cap H_i$ . For  $j = 1, \dots, \ell$ , let

$$M'_j = g_j N'_{\alpha(j)} g_j^{-1}.$$

We verify that  $\{M'_j\}_{j=1}^\ell$  is a  $G$ -equivariant family. Let  $j_1, j_2 \in \{1, \dots, \ell\}$  such that there exists  $g \in G$  with  $g K_{j_1} g^{-1} = K_{j_2}$ . We have  $K_{j_1} \leq g_{j_1} H_{\alpha(j_1)} g_{j_1}^{-1}$ , and thus  $K_{j_2} = g K_{j_1} g^{-1} \leq g g_{j_1} H_{\alpha(j_1)} g_{j_1}^{-1} g^{-1}$ . Therefore,  $G_0 \cap H_{\alpha(j_2)} = g_{j_2}^{-1} K_{j_2} g_{j_2} \leq g_{j_2}^{-1} g g_{j_1} H_{\alpha(j_1)} g_{j_1}^{-1} g^{-1} g_{j_2}$ . Being a finite-index subgroup of the infinite group  $H_{\alpha(j_2)}$ , the group  $G_0 \cap H_{\alpha(j_2)}$  is infinite, and thus [DGO17, Proposition 4.33] implies that  $\alpha(j_1) = \alpha(j_2)$  and  $g_{j_2}^{-1} g g_{j_1} \in H_{\alpha(j_1)}$ . Therefore,

$$\begin{aligned} M'_{j_2} &= g_{j_2} N'_{\alpha(j_2)} g_{j_2}^{-1} = g_{j_2} N'_{\alpha(j_1)} g_{j_2}^{-1} = g_{j_2} (g_{j_2}^{-1} g g_{j_1}) N'_{\alpha(j_1)} (g_{j_2}^{-1} g g_{j_1})^{-1} g_{j_2}^{-1} \\ &= g g_{j_1} N'_{\alpha(j_1)} g_{j_1}^{-1} g^{-1} = g M'_{j_1} g^{-1}, \end{aligned}$$

as desired.

(ii). For every  $j \in \{1, \dots, \ell\}$ , we have

$$\begin{aligned} g_j N_{k, \alpha(j)} g_j^{-1} &= M_{k, j}, \\ M_{1, j} &= g_j N_{1, \alpha(j)} g_j^{-1} \leq g_j N_{2, \alpha(j)} g_j^{-1} = M_{2, j}, \\ M_{2, j} / M_{1, j} &= (g_j N_{2, \alpha(j)} g_j^{-1}) / (g_j N_{1, \alpha(j)} g_j^{-1}) \cong N_{2, \alpha(j)} / N_{1, \alpha(j)}, \\ K_j / M_{k, j} &= (g_j (G_0 \cap H_{\alpha(j)}) g_j^{-1}) / (g_j N_{k, \alpha(j)} g_j^{-1}) \cong (G_0 \cap H_{\alpha(j)}) / N_{k, \alpha(j)}. \end{aligned}$$

(iii). Let  $T = \{M_j \triangleleft K_j\}_{j=1}^\ell \in \mathcal{T}$  and let  $\varphi(T) = \{N_i \triangleleft H_i\}_{i=1}^k$ . For  $j = 1, \dots, \ell$ , we have  $M_j = g_j N_{\alpha(j)} g_j^{-1}$ . In particular, we have  $\langle\langle \bigcup_{j=1}^\ell M_j \rangle\rangle_{G_0} \leq \langle\langle \bigcup_{i=1}^k N_i \rangle\rangle_G$  and thus there is a natural homomorphism  $\psi: G_0 / \langle\langle \bigcup_{j=1}^\ell M_j \rangle\rangle_{G_0} \rightarrow G / \langle\langle \bigcup_{i=1}^k N_i \rangle\rangle_G$  that fits into a commutative diagram

$$\begin{array}{ccc} G_0 & \longrightarrow & G_0 / \langle\langle \bigcup_{j=1}^\ell M_j \rangle\rangle_{G_0} \\ \downarrow & & \downarrow \psi \\ G & \longrightarrow & G / \langle\langle \bigcup_{i=1}^k N_i \rangle\rangle_G \end{array}$$



To prove that  $\psi$  is injective, it suffices to prove  $\langle\langle \bigcup_{i=1}^k N_i \rangle\rangle_G \leq \langle\langle \bigcup_{j=1}^\ell M_j \rangle\rangle_{G_0}$ . A general element  $n \in \langle\langle \bigcup_{i=1}^k N_i \rangle\rangle_G$  has the form  $n = \prod_{p=1}^q g'_p n_p (g'_p)^{-1}$ . For each  $p$ , there exists  $i(p) \in \{1, \dots, k\}$  such that  $n_p \in N_{i(p)}$ . We have

$$N_{i(p)} = g_{\beta(i(p))}^{-1} M_{\beta(i(p))} g_{\beta(i(p))}.$$

So,

$$n \in \prod_{p=1}^q g'_p g_{\beta(i(p))}^{-1} M_{\beta(i(p))} g_{\beta(i(p))} (g'_p)^{-1}.$$

By the definition of the induced peripheral structure, there exists  $j(p) \in \{1, \dots, \ell\}$  and  $g''_p \in G_0$  such that

$$g'_p g_{\beta(i(p))}^{-1} K_{\beta(i(p))} g_{\beta(i(p))} (g'_p)^{-1} = g''_p K_{j(p)} (g''_p)^{-1}.$$

As  $\{M_j\}_{j=1}^\ell$  is a  $G$ -equivariant family, we have

$$g'_p g_{\beta(i(p))}^{-1} M_{\beta(i(p))} g_{\beta(i(p))} (g'_p)^{-1} = g''_p M_{j(p)} (g''_p)^{-1}.$$

Therefore,

$$n \in \prod_{p=1}^q g''_p M_{j(p)} (g''_p)^{-1} \subset \langle\langle \bigcup_{j=1}^\ell M_j \rangle\rangle_{G_0},$$

which finishes the proof that  $\psi$  is injective and also proves (1).

(iv). Let  $\{\mathcal{F}_i \subset H_i \setminus \{1\}\}_{i=1}^k$  be a family of finite subsets. For  $j \in \{1, \dots, \ell\}$ , let

$$\mathcal{F}'_j = g_j (\mathcal{F}_{\alpha(j)} \cap G_0) g_j^{-1}.$$

Then the family  $\{\mathcal{F}'_j\}_{j=1}^\ell$  satisfies the requirement.

Conversely, let  $\{\mathcal{F}'_j \subset K_j \setminus \{1\}\}_{j=1}^\ell$  be a family of finite subsets. For  $i \in \{1, \dots, k\}$ , let

$$\mathcal{F}_i = \bigcup_{j \in \alpha^{-1}(\{i\})} g_j^{-1} \mathcal{F}'_j g_j.$$

Then the family  $\{\mathcal{F}_i\}_{i=1}^k$  satisfies the requirement.

(v). Let  $\{P_i \leq H_i\}_{i=1}^k$  be a family of finite-index subgroups. For  $j \in \{1, \dots, \ell\}$ , let

$$Q_j = g_j (P_{\alpha(j)} \cap G_0) g_j^{-1}.$$

Then the family  $\{Q_j\}_{j=1}^\ell$  satisfies the requirement.

Conversely, let  $\{Q'_j \leq K_j\}_{j=1}^\ell$  be a family of finite-index subgroups. For  $i \in \{1, \dots, k\}$ , let

$$P'_i = \bigcap_{j \in \alpha^{-1}(\{i\})} g_j^{-1} Q'_j g_j.$$

Then the family  $\{P'_i\}_{i=1}^k$  satisfies the requirement.  $\square$

## 3. QUANTIFYING LÜCK'S APPROXIMATION

We will need an approximation result for  $L^2$ -Betti numbers for the space of marked groups, proved by Jaikin-Zapirain–López-Álvarez [JZLA20].

Let  $F$  be a group freely generated by a finite set  $S$ . The space of marked groups  $\text{MG}(F)$  can be identified with the set of normal subgroups of  $F$  with the metric  $d(M_1, M_2) = e^{-n}$ , where  $M_1 \neq M_2 \triangleleft F$  are two normal subgroups and  $n$  is the largest number such that the balls of radius  $n$  in the Cayley graphs  $\Gamma(F/M_1, S)$  and  $\Gamma(F/M_2, S)$  are isomorphic as labeled graphs. The metric  $d$  induces a topology on  $\text{MG}(F)$ . Below, the notion of convergence is discussed with respect to this topology.

**Theorem 3.1** (Jaikin-Zapirain–López-Álvarez, 2020). *Let  $F$  be a finitely generated free group and assume that a sequence  $\{M_i\}_{i \geq 1}$  of normal subgroups of  $F$  converges to a normal subgroup  $M \triangleleft F$  in  $\text{MG}(F)$ . Suppose that  $F/M$  is virtually locally indicable. Then for every finite matrix  $A$  over  $\mathbb{C}F$ , we have*

$$\lim_{i \rightarrow \infty} \text{rk}_{F/M_i}(A) = \text{rk}_{F/M}(A).$$

For applications to Dehn fillings, we need to quantify the above theorem.

**Theorem 3.2.** *Let  $G$  be a finitely generated, virtually locally indicable group. Then for every finite matrix  $A$  over  $\mathbb{C}G$  and every  $\delta > 0$ , there exists a finite subset  $\mathcal{F}_\delta \subset G \setminus \{1\}$  such that if a normal subgroup  $N \triangleleft G$  satisfies  $N \cap \mathcal{F}_\delta = \emptyset$ , then*

$$|\text{rk}_{G/N}(A) - \text{rk}_G(A)| < \delta.$$

*Proof.* Suppose, to the contrary, that there exists a matrix  $A$  over  $\mathbb{C}G$  and  $\delta_0 > 0$  such that for each finite subset  $\mathcal{F} \subset G \setminus \{1\}$ , there exists a normal subgroup  $N \triangleleft G$  such that  $N \cap \mathcal{F} = \emptyset$  but

$$|\text{rk}_{G/N}(A) - \text{rk}_G(A)| > \delta_0.$$

Fix a finite generating set  $S$  of  $G$ . Let  $F$  be the free group on basis  $S$ , let  $\epsilon: F \rightarrow G$  be the natural surjection, and let  $M = \ker \epsilon$ . Then  $\epsilon$  can be naturally seen to be a map from the set of square matrices over  $\mathbb{C}F$  to the set of square matrices over  $\mathbb{C}G$ . Let  $B$  be a matrix over  $\mathbb{C}F$  such that  $A = \epsilon(B)$ .

For every  $n \in \mathbb{N}^+$ , let  $B_n \subset \Gamma(G, S)$  be the subgraph consisting of all points within distance  $n$  from the vertex 1, and let  $\mathcal{F}_n = G \cap B_n \setminus \{1\}$ . By assumption, there exists a normal subgroup  $N_n \triangleleft G$  such that  $N_n \cap \mathcal{F}_n = \emptyset$  and

$$(2) \quad |\text{rk}_{G/N_n}(A) - \text{rk}_G(A)| > \delta_0.$$

Let  $M_n \triangleleft F$  be the preimage of  $N_n$  under  $\epsilon$ . We claim that the sequence of normal subgroups  $\{M_n\}_{n \geq 1}$  converges to  $M$  in  $\text{MG}(F)$ . Indeed, for every  $R \in \mathbb{N}^+$ , let  $n > 2R$ . Consider the subgraph  $\overline{B}_R \subset \Gamma(F/M_n, S) = \Gamma(G/N_n, S)$  consisting of all points within distance  $R$  from 1, and the subgraph  $B_R \subset \Gamma(F/M, S) = \Gamma(G, S)$ . Let  $g \neq h$  be two vertices of  $B_R$ , where we think of  $g, h$  as elements of  $G$ , and let  $\bar{g}, \bar{h}$  be their natural images in  $G/N_n$ . Then  $\bar{g}, \bar{h}$  are vertices in  $\overline{B}_R$ . Note that  $g$  and  $h$  are a distance less than  $2R$  apart from each other. As  $n > 2R$ , we have  $g^{-1}h \notin N_n$ , and thus  $\bar{g} \neq \bar{h}$ . Therefore, the natural map from  $B_R$  to  $\overline{B}_R$  is injective. As this natural map is obviously surjective, the labeled graphs  $B_R$  and  $\overline{B}_R$  are isomorphic, whenever  $n > 2R$ .

By Theorem 3.1, we have

$$(3) \quad \lim_{n \rightarrow \infty} \text{rk}_{F/M_n}(B) = \text{rk}_{F/M}(B).$$

On the other hand, we have  $\text{rk}_{F/M_n}(B) = \text{rk}_{G/N_n}(A)$  and  $\text{rk}_{F/M}(B) = \text{rk}_G(A)$ . Thus, (2) and (3) contradict each other.  $\square$

In the above theorem, if  $A$  is not finite but just a tamed matrix, one can still obtain a lower bound on  $\text{rk}_{G/N}(A)$ .

**Corollary 3.3.** *Let  $G$  be a finitely generated, virtually locally indicable group. Then for each tamed matrix  $A$  over  $\mathbb{C}G$  such that  $\text{rk}_G(A) < \infty$  and each  $\delta > 0$ , there exists a finite subset  $\mathcal{F}_\delta \subset G \setminus \{1\}$  such that if a normal subgroup  $N \triangleleft G$  satisfies  $N \cap \mathcal{F}_\delta = \emptyset$ , then*

$$\text{rk}_{G/N}(A) > \text{rk}_G(A) - \delta.$$

*Proof.* Write  $A$  as  $A = (a_{ij})_{i \in I, j \in J}$  and let  $r_A: (\mathcal{N}(G))^I \rightarrow (\mathcal{N}(G))^J$  be the  $\mathcal{N}(G)$ -module homomorphism given by the right-multiplication of  $A$ . By definition,

$$(4) \quad \text{rk}_G(A) = \dim_G \text{im}(r_A).$$

For each  $i \in I$ , let  $e_i = (x_{i'})_{i' \in I} \in (\mathcal{N}(G))^I$  be the element such that

$$x_{i'} = \begin{cases} 1, & \text{if } i' = i \\ 0, & \text{if } i' \neq i. \end{cases}$$

Let  $\delta > 0$ , and let  $\mathcal{S}$  be the family of non-empty finite subsets of  $I$ . For every  $S \in \mathcal{S}$ , let  $M_S$  be the  $\mathcal{N}(G)$ -submodule of  $(\mathcal{N}(G))^I$  generated by  $e_i, i \in S$ . Then  $\{r_A(M_S) \mid S \in \mathcal{S}\}$  is a cofinal family of  $\text{im}(r_A)$ , and thus

$$\dim_G \text{im}(r_A) = \sup\{\dim_G r_A(M_S) \mid S \in \mathcal{S}\}$$

by Proposition 2.1 (iv). So there exists  $S \in \mathcal{S}$  such that

$$(5) \quad \dim_G \text{im}(r_A) < \dim_G r_A(M_S) + \frac{\delta}{2}.$$

For each  $i \in S$ , there are only finitely many  $j \in J$  such that  $a_{ij} \neq 0$ . Let  $T \subset J$  be the subset consisting of  $j \in J$  such that there exists  $i \in S$  with  $a_{ij} \neq 0$ . As  $|S| < \infty$ , we also have  $|T| < \infty$ . Let  $B = (a_{ij})_{i \in S, j \in T}$ , and let  $r_B: (\mathcal{N}(G))^S \rightarrow (\mathcal{N}(G))^T$  be the  $\mathcal{N}(G)$ -module homomorphism induced by the right-multiplication of  $B$ . Then  $\text{im}(r_B) \cong r_A(M_S)$  as  $\mathcal{N}(G)$ -modules, and thus

$$(6) \quad \dim_G r_A(M_S) = \dim_G \text{im}(r_B) = \text{rk}_G(B).$$

By Theorem 3.2, there exists a finite subset  $\mathcal{F}_\delta \subset G \setminus \{1\}$  such that if a normal subgroup  $N \triangleleft G$  satisfies  $N \cap \mathcal{F}_\delta = \emptyset$ , then

$$(7) \quad \text{rk}_{G/N}(B) > \text{rk}_G(B) - \frac{\delta}{2}.$$

For every  $N \triangleleft G$ , let  $\epsilon_N: G \rightarrow G/N$  be the natural homomorphism, and let  $r_{\epsilon_N(A)}: (\mathcal{N}(G/N))^I \rightarrow (\mathcal{N}(G/N))^J$  (resp.  $r_{\epsilon_N(B)}: (\mathcal{N}(G/N))^S \rightarrow (\mathcal{N}(G/N))^T$ ) be the  $\mathcal{N}(G/N)$ -module homomorphism induced by the right-multiplication of  $\epsilon_N(A)$  (resp.  $\epsilon_N(B)$ ).

For each  $i \in I$ , let  $\bar{e}_i = (\bar{x}_{i'})_{i' \in I} \in (\mathcal{N}(G/N))^I$  be the element such that

$$\bar{x}_{i'} = \begin{cases} 1, & \text{if } i' = i \\ 0, & \text{if } i' \neq i. \end{cases}$$

And let  $\overline{M}_S$  be the submodule of  $(\mathcal{N}(G/N))^I$  generated by  $\bar{e}_i, i \in S$ . Then  $\text{im}(r_{\epsilon_N(B)}) \cong r_{\epsilon_N(A)}(\overline{M}_S)$  as  $\mathcal{N}(G)$ -modules, and thus

$$(8) \quad \begin{aligned} \text{rk}_{G/N}(B) &= \dim_{G/N} \text{im}(r_{\epsilon_N(B)}) = \dim_{G/N} \bar{r}_{\epsilon_N(A)}(\overline{M}_S) \\ &\leq \dim_{G/N} \text{im}(r_{\epsilon_N(A)}) = \text{rk}_{G/N}(A). \end{aligned}$$

The desired result follows from (4), (5), (6), (7) and (8).  $\square$

**Corollary 3.4.** *Let  $G$  be a finitely generated virtually locally indicable group and let  $X$  be a free  $G$ -CW complex of type  $F_{k+1}$  for some  $k \in \mathbb{N}$ . Then for every  $\delta > 0$ , there exists a finite subset  $\mathcal{F}_\delta \subset G \setminus \{1\}$  such that if a normal subgroup  $N \triangleleft G$  satisfies  $N \cap \mathcal{F}_\delta = \emptyset$ , then*

$$b_{k+1}^{(2)}(X; G) + \delta > b_{k+1}^{(2)}(N \setminus X; G/N),$$

and for all  $n \leq k$ , we have

$$|b_n^{(2)}(X; G) - b_n^{(2)}(N \setminus X; G/N)| < \delta.$$

*Proof.* Let  $\partial_*$  be the boundary map of  $C_*(X)$ . For each  $n \in \mathbb{Z}$  and each  $G$ -orbit of the  $n$ -cells of  $X$ , pick one representative. The chosen representatives form a  $\mathbb{Z}G$ -basis  $B_n$  of the module  $C_n(X)$  of  $X$ , yielding an isomorphism  $C_n(X) \cong (\mathbb{Z}G)^{B_n}$ .

Let  $A_n$  be the tamed matrix over  $\mathbb{Z}G$  such that

$$\partial_n: C_n(X) \cong (\mathbb{Z}G)^{B_n} \rightarrow C_{n-1}(X) \cong (\mathbb{Z}G)^{B_{n-1}}$$

is given by the right-multiplication of  $A_n$ . Then for  $n \leq k+1$ , we have

$$b_n^{(2)}(X; G) = |B_n| - \text{rk}_G(A_n) - \text{rk}_G(A_{n+1}),$$

$$b_n^{(2)}(N \setminus X; G/N) = |B_n| - \text{rk}_{G/N}(A_n) - \text{rk}_{G/N}(A_{n+1}).$$

For  $n \leq k$ , let  $\mathcal{F}_{n1} \subset G \setminus \{1\}$  (resp.  $\mathcal{F}_{n2} \subset G \setminus \{1\}$ ) be the finite subset given by Theorem 3.2 with respect to  $A_n$  (resp.  $A_{n+1}$ ) and  $\delta/2$ . Let  $\mathcal{F}_{k+1} \subset G \setminus \{1\}$  be the finite subset given by Corollary 3.3 with respect to  $A_{k+2}$  and  $\delta/2$ . Let

$$\mathcal{F}_\delta = \mathcal{F}_{k+1} \cup \left( \bigcup_{n=0}^{k+1} (\mathcal{F}_{n1} \cup \mathcal{F}_{n2}) \right),$$

and let  $N \triangleleft G$  be a normal subgroup such that  $N \cap \mathcal{F}_\delta = \emptyset$ . By Theorem 3.2 and Corollary 3.3, we have

$$\text{rk}_{G/N}(A_{k+2}) > \text{rk}_G(A_{k+2}) - \frac{\delta}{2},$$

and

$$|\text{rk}_G(A_n) - \text{rk}_{G/N}(A_n)| < \delta/2$$

for  $n \leq k+1$ . Therefore,

$$b_{k+1}^{(2)}(N \setminus X; G/N) = |B_{k+1}| - \text{rk}_{G/N}(A_{k+1}) - \text{rk}_{G/N}(A_{k+2})$$

$$\begin{aligned} &< |B_{k+1}| - \text{rk}_G(A_{k+1}) - \text{rk}_G(A_{k+2}) + \delta \\ &= b_{k+1}^{(2)}(X; G) + \delta, \end{aligned}$$

and

$$|b_n^{(2)}(X; G) - b_n^{(2)}(N \setminus X; G/N)| \leq |\text{rk}_G A_n - \text{rk}_{G/N} A_n| + |\text{rk}_G A_{n+1} - \text{rk}_{G/N} A_{n+1}| < \delta$$

for  $n \leq k$ . □

The above results provide estimates of  $L^2$ -Betti numbers. To obtain actual computation rather than just estimation, we will combine them with the Atiyah Conjecture. We will need the following result:

**Proposition 3.5.** *Let  $G$  be a finitely generated virtually locally indicable group. Suppose that there exists  $C \in \mathbb{N}$  such that for all finite subset  $\mathcal{S} \subset G \setminus \{1\}$ , there exists a normal subgroup  $N_{\mathcal{S}} \triangleleft G$  such that*

- (i)  $N_{\mathcal{S}} \cap \mathcal{S} = \emptyset$ ;
- (ii)  $G/N_{\mathcal{S}}$  satisfies the Atiyah Conjecture; and
- (iii)  $\max \mathcal{FLN}(G/N_{\mathcal{S}}) \leq C$ , where  $\max$  means taking the maximum of a set.

Let  $X$  be a free  $G$ -CW complex of type  $F_{k+1}$  for some  $k \in \mathbb{N}$ . Then for all  $n \leq k$ , we have

$$b_n^{(2)}(X; G) \in \frac{1}{C!} \cdot \mathbb{Z}.$$

*Proof.* For each finite subset  $\mathcal{S} \subset G \setminus \{1\}$ , let  $N_{\mathcal{S}} \triangleleft G$  be a normal subgroup such that the above (i) through (iii) hold. The family

$$\mathcal{N} = \{N_{\mathcal{S}} \mid \mathcal{S} \text{ is a finite subset of } G\}$$

is a directed set under the partial order  $N_{\mathcal{S}} < N_{\mathcal{S}'}$  if and only if  $\mathcal{S} \subset \mathcal{S}'$ . For each finite matrix  $A$  over  $\mathbb{Z}G$ , we have

$$\text{rk}_G(A) = \varinjlim \text{rk}_{G/N_{\mathcal{S}}}(A) \in \mathbb{N} \in \frac{1}{(\max_{N_{\mathcal{S}} \in \mathcal{N}} \mathcal{FLN}(G/N_{\mathcal{S}}))!} \cdot \mathbb{Z} \leq \frac{1}{C!} \cdot \mathbb{Z}$$

by Theorem 3.2.

Now fix  $n \leq k$ . Let  $B_n$  (resp.  $B_{n+1}, B_{n-1}$ ) be the set obtained by choosing one representative of each  $G$ -orbit of the  $n$ -cells (resp.  $(n+1)$ -cells,  $(n-1)$ -cells) of  $X$ . Then  $B_n$  (resp.  $B_{n+1}, B_{n-1}$ ) is a  $\mathbb{Z}G$ -basis of  $C_n(X)$  (resp.  $C_{n+1}(X), C_{n-1}(X)$ ). Let  $\partial_*$  be the boundary map of  $C_*(X)$ , and let  $A_n$  (resp.  $A_{n+1}$ ) be the matrix representative of  $\partial_n$  (resp.  $\partial_{n+1}$ ) under the bases  $B_n$  and  $B_{n-1}$  (resp.  $B_{n+1}$  and  $B_n$ ). Then

$$b_n^{(2)}(X; G) = |B_n| - \text{rk}_G A_n - \text{rk}_G A_{n+1} \in \frac{1}{C!} \cdot \mathbb{Z}.$$

□

**3.1. The first  $\ell^2$ -Betti number.** Our next results yield an upper bound for the first  $L^2$ -Betti number of Dehn filling (see Corollary 3.8). Estimation of the other  $L^2$ -Betti numbers needs an additional ingredient called a Dehn filling space, which will be studied in Section 4.

**Lemma 3.6.** *Let  $G$  be any group and let  $N \triangleleft G$  be a normal subgroup. Then*

$$b_1^{(2)}(N \setminus EG; G/N) \geq b_1^{(2)}(G/N).$$

*Proof.* Let  $EG$  (resp.  $E(G/N)$ ) be a free contractible  $G$ -CW complex (resp.  $(G/N)$ -CW complex). The quotient map  $G \rightarrow G/N$  induces a  $G$ -equivariant cellular map  $f: EG \rightarrow E(G/N)$ , which factors through a  $G/N$ -equivariant cellular map  $\tilde{f}: N \backslash EG \rightarrow E(G/N)$ .

We will use [Lüc02, Theorem 6.54 (1)]. To this end, let us consider a subgroup  $H \leq G/N$  and let  $(N \backslash EG)^H$  (resp.  $(E(G/N))^H$ ) be the subcomplex of  $N \backslash EG$  (resp.  $E(G/N)$ ) consisting of the global fixed points of  $H$ . If  $H \neq \{1\}$ , then  $(N \backslash EG)^H = (E(G/N))^H = \emptyset$  and the assumptions of [Lüc02, Theorem 6.54 (1)] are trivially satisfied. If  $H = \{1\}$ , then  $(N \backslash EG)^H = N \backslash EG$  and  $(E(G/N))^H = E(G/N)$ . Clearly,  $\tilde{f}$  induces an isomorphism  $H_0(N \backslash EG; \mathbb{C}) \rightarrow H_0(E(G/N); \mathbb{C})$ . As  $H_1(E(G/N); \mathbb{C}) = 0$ ,  $\tilde{f}$  induces an epimorphism  $H_1(N \backslash EG; \mathbb{C}) \rightarrow H_1(E(G/N); \mathbb{C})$ . So the assumptions of [Lüc02, Theorem 6.54 (1)] are met, which yields

$$b_1^{(2)}(N \backslash EG; G/N) \geq b_1^{(2)}(E(G/N); G/N) = b_1^{(2)}(G/N).$$

□

**Corollary 3.7.** *Let  $G$  be a finitely generated virtually locally indicable group. Then for every  $\delta > 0$ , there exists a finite subset  $\mathcal{F}_\delta \subset G \setminus \{1\}$  such that if a normal subgroup  $N \triangleleft G$  satisfies  $N \cap \mathcal{F}_\delta = \emptyset$ , then*

$$b_1^{(2)}(G/N) < b_1^{(2)}(G) + \delta$$

*Proof.* Let  $X$  be the Cayley 2-complex of  $G$  with respect to any presentation of  $G$  with finitely many generators, and let  $\mathcal{F}_\delta \subset G \setminus \{1\}$  be the finite subset provided by Corollary 3.4 with respect to  $k = 0$  and  $X$ . Then if a normal subgroup  $N \triangleleft G$  satisfies  $N \cap \mathcal{F}_\delta = \emptyset$ , we have

$$b_1^{(2)}(N \backslash X; G/N) < b_1^{(2)}(X; G) + \delta.$$

By Lemma 3.6, we have

$$b_1^{(2)}(N \backslash EG; G/N) \geq b_1^{(2)}(G/N).$$

Note that

$$\begin{aligned} b_1^{(2)}(N \backslash EG; G/N) &= b_1^{(2)}(N \backslash X; G/N), \\ b_1^{(2)}(G) &= b_1^{(2)}(X; G), \end{aligned}$$

from which the desired result follows. □

**Corollary 3.8.** *Let  $G$  be a finitely generated virtually locally indicable group, and let  $H \hookrightarrow_h G$  be a hyperbolically embedded subgroup. Then for every  $\delta > 0$ , there exists a finite subset  $\mathcal{F}_\delta \subset H \setminus \{1\}$  such that if a normal subgroup  $N \triangleleft H$  satisfies  $N \cap \mathcal{F}_\delta = \emptyset$ , then*

$$(9) \quad b_1^{(2)}(G/\langle\langle N \rangle\rangle) < b_1^{(2)}(G) + \delta.$$

*Proof.* By Corollary 3.7, there exists a finite subset  $\mathcal{S}_\delta \subset G \setminus \{1\}$  such that if a normal subgroup  $M \triangleleft G$  satisfies  $M \cap \mathcal{S}_\delta = \emptyset$ , then

$$(10) \quad b_1^{(2)}(G/M) < b_1^{(2)}(G) + \delta.$$

By [DGO17, Theorem 7.19 (c)], there exists a finite subset  $\mathcal{F}_\delta \subset H \setminus \{1\}$  such that if a normal subgroup  $N \triangleleft H$  satisfies  $N \cap \mathcal{F}_\delta = \emptyset$ , then  $\langle\langle N \rangle\rangle \cap \mathcal{S}_\delta = \emptyset$ , which implies the desired result. □



4. DEHN FILLING SPACE AND  $L^2$ -BETTI NUMBERS

4.1. **Dehn filling space.** Let  $G$  be a group,  $H \leq G$  a subgroup, and  $N \triangleleft H$  a normal subgroup such that  $(G, H, N)$  is a Cohen–Lyndon triple, i.e., there exists a left transversal  $T$  of  $H\langle\langle N \rangle\rangle$  in  $G$  such that

$$\langle\langle N \rangle\rangle = \ast_{t \in T} t N t^{-1}.$$

As in the introduction, we denote  $\overline{G} = G/\langle\langle N \rangle\rangle$  and  $\overline{H} = H/N$ . The goal of this subsection is to build a specific  $K(\overline{G}, 1)$ -space.

Note that the natural map  $\overline{H} \rightarrow \overline{G}$  is injective (see e.g., [Sun20, Lemma 6.4]), and below we will view  $\overline{H}$  as a subgroup of  $\overline{G}$ . Let  $BG$  (resp.  $BH, B\overline{H}$ ) be a  $K(G, 1)$  (resp.  $K(H, 1), K(\overline{H}, 1)$ ) CW-complex.




FIGURE 1. Dehn filling space associated to the triple  $(G, H, N)$ . The mappings cylinders  $M_\psi$  and  $M_\phi$  are shaded blue and grey, respectively. Their bases  $B\overline{H}$  and  $BG$  are represented by larger discs. The common subspace  $BH$  is shaded red.

**Definition 4.1.** There is a classifying map  $\phi: BH \rightarrow BG$  (resp.  $\psi: BH \rightarrow B\overline{H}$ ) induced by the inclusion  $H \hookrightarrow G$  (resp. the quotient map  $q: H \rightarrow \overline{H}$ ). Let  $X$  be the complex obtained by gluing the mapping cylinders  $M_\phi, M_\psi$  along their common subcomplex  $BH$  (see Figure 1). We call the complex  $X$  a *Dehn filling space* associated to the triple  $(G, H, N)$ .

For simplicity, we will use the following notation. Let

$$\tilde{\phi}: \overline{G} \times_{\overline{H}} (N \setminus EH) \cong \langle\langle N \rangle\rangle \backslash (G \times_H EH) \rightarrow \langle\langle N \rangle\rangle \backslash EG$$

and

$$\tilde{\psi}: \overline{G} \times_{\overline{H}} (N \setminus EH) \rightarrow \overline{G} \times_{\overline{H}} E\overline{H}$$

be the  $\overline{G}$ -equivariant maps induced by  $\phi$  and  $\psi$ , respectively. Also let  $M_{\tilde{\phi}}$  and  $M_{\tilde{\psi}}$  be the mapping cylinder of  $\tilde{\phi}$  and  $\tilde{\psi}$ , respectively.

**Theorem 4.2.** *Let  $(G, H, N)$  be a Cohen–Lyndon triple. Then every Dehn filling space associated to the triple  $(G, H, N)$  is a  $K(\overline{G}, 1)$ -space.*

By combining the above theorem with Theorem 2.13 we get the following:

**Corollary 4.3.** *Let  $G$  be a group with a hyperbolicly embedded subgroup  $H \hookrightarrow_h G$ . Then for sufficiently deep  $N \triangleleft H$ , every Dehn filling space associated to the triple  $(G, H, N)$  is a  $K(\overline{G}, 1)$ -space.*

*Proof of Theorem 4.2.* We will use the notation from Definition 4.1. We first prove that  $\pi_1(X) = \overline{G}$ . By the Seifert–van Kampen theorem,  $\pi_1(X)$  has the presentation

$$(11) \quad \pi_1(X) = \langle G, \overline{H} \mid h = q(h), \forall h \in H \leq G \rangle.$$

In particular, there is an epimorphism  $\tau: G \twoheadrightarrow \pi_1(X)$  that maps every  $n \in N$  to 1. Thus,  $\tau$  descends to an epimorphism (still denoted by)  $\tau: \overline{G} \twoheadrightarrow \pi_1(X)$ .

Let  $p: G \rightarrow \overline{G}$  be the natural quotient map. Then from the presentation (11) we see that there is a homomorphism  $\eta: \pi_1(X) \rightarrow \overline{G}$  given by  $\eta(g) = p(g)$  for all  $g \in G$  and  $\eta(\bar{h}) = \bar{h}$  for all  $\bar{h} \in \overline{H}$ . It is easy to see that  $\tau$  and  $\eta$  are mutual inverses. Thus,  $\pi_1(X) = \overline{G}$ .

It remains to prove that the universal cover  $\tilde{X}$  of  $X$  is contractible. Since  $\tilde{X}$  is a simply connected CW-complex, it suffices to prove that  $H_n(\tilde{X}) = 0$  for all  $n \geq 2$ . For simplicity, all homology groups have integral coefficients (where the group action on  $\mathbb{Z}$  is trivial) for the rest of this proof.

The space  $\tilde{X}$  can be obtained by gluing the mapping cylinders  $M_{\tilde{\phi}}, M_{\tilde{\psi}}$  along their common subcomplex  $\overline{G} \times_{\overline{H}} (N \setminus EH)$ . We thus have a Mayer–Vietoris sequence:

$$\cdots \rightarrow H_{n+1}(\tilde{X}) \rightarrow H_n(\overline{G} \times_{\overline{H}} (N \setminus EH)) \xrightarrow{\lambda} H_n(M_{\tilde{\phi}}) \oplus H_n(M_{\tilde{\psi}}) \rightarrow H_n(\tilde{X}) \rightarrow \cdots$$

It is enough to prove that for all  $n \geq 2$ ,  $\lambda: H_n(\overline{G} \times_{\overline{H}} (N \setminus EH)) \rightarrow H_n(M_{\tilde{\phi}}) \oplus H_n(M_{\tilde{\psi}})$  is an isomorphism. Note that  $M_{\tilde{\psi}}$  deformation retracts onto  $\overline{G} \times_{\overline{H}} E\overline{H}$  and  $H_n(\overline{G} \times_{\overline{H}} E\overline{H}) = 0$ . Thus,  $H_n(M_{\tilde{\psi}}) = 0$ . So it suffices to prove that the homomorphism

$$\theta: H_n(\overline{G} \times_{\overline{H}} (N \setminus EH)) \rightarrow H_n(M_{\tilde{\phi}})$$

induced by  $\lambda$  is an isomorphism. Note that  $M_{\tilde{\phi}}$  deformation retracts onto  $\langle\langle N \rangle\rangle \setminus EG$ , and thus is a  $K(\langle\langle N \rangle\rangle, 1)$ -space. By the Cohen–Lyndon property,

$$H_n(\langle\langle N \rangle\rangle \setminus EG) = \bigoplus_{t \in T} H_n(tNt^{-1}),$$

where  $T$  is a left transversal of  $H\langle\langle N \rangle\rangle$  in  $G$ . The space  $\overline{G} \times_{\overline{H}} (N \setminus EH)$  is a disjoint union of copies of  $N \setminus EH$ . Each of these copies is labeled by a unique element  $p(t)$  for some  $t \in T$ . The inclusion  $\overline{G} \times_{\overline{H}} (N \setminus EH) \hookrightarrow M_{\tilde{\phi}}$ , which induces the homology map  $\theta$ , can be seen as a disjoint union of inclusions of each copy of  $N \setminus EH$ . Consider one such copy  $(N \setminus EH)_{p(t)}$  labeled by  $p(t)$ . The copy  $(N \setminus EH)_{p(t)}$  is a  $K(tNt^{-1}, 1)$ -space and the inclusion  $i_t: (N \setminus EH)_{p(t)} \hookrightarrow M_{\tilde{\phi}}$  is the classifying map. So the induced map on homology maps  $H_n((N \setminus EH)_{p(t)}) = H_n(tNt^{-1})$  isomorphically onto the summand  $H_n(tNt^{-1})$  of  $H_n(\langle\langle N \rangle\rangle \setminus EG)$ . Since this holds for all  $t \in T$  we conclude that  $\theta$  is an isomorphism, as desired.  $\square$

**Corollary 4.4.** *Let  $G$  be a group with a hyperbolically embedded subgroup  $H \hookrightarrow_h G$ .*

- (i) *If  $G$  is of type  $F_n$  for some  $n \in \mathbb{N}^+ \cup \{\infty\}$ , then for all sufficiently deep normal  $N \triangleleft H$ , we have that  $\overline{G}$  is of type  $F_n$  if and only if  $\overline{H}$  is of type  $F_n$ .*
- (ii) *If both  $G$  and  $H$  are of type  $F$ , then for all sufficiently deep normal  $N \triangleleft H$  such that  $\overline{H}$  is of type  $F$ , we have that  $\overline{G}$  is of type  $F$ .*

*Proof.*

(i). Suppose  $\overline{G}$  is of type  $F_n$ . By Theorem 2.11,  $\overline{H}$  is hyperbolically embedded in  $\overline{G}$ . Applying, Corollary 4.32 of [DGO17], we obtain that  $\overline{H}$  is of type  $F_n$ .

Now, suppose  $\overline{H}$  is of type  $F_n$ . We will show that the Dehn filling space has finite  $n$ -skeleton. It follows that there is a model for  $B\overline{H}$  with finite  $n$ -skeleton. Since  $G$  is of type  $F_n$ , by Corollary 4.32 of [DGO17],  $H$  is of type  $F_n$ . Thus, there are models for  $BG$  and  $BH$  with finite  $n$ -skeleta. Using the notation of Definition 4.1, it follows that the mapping cylinders  $M_\phi$  and  $M_\psi$  have finite  $n$ -skeleta. Since they are glued along  $BH$ , the resulting Dehn filling space has the desired finiteness properties. By Corollary 4.3, this Dehn filling space is a  $K(\overline{G}, 1)$  space provided  $N$  is sufficiently deep.

(ii). The proof is analogous to part (i). There are models for  $BG, BH$  and  $B\overline{H}$  that are finite complexes. Using the notation of Definition 4.1, it follows that the mapping cylinders  $M_\phi$  and  $M_\psi$  are finite complexes. Since they are glued along  $BH$ , the resulting Dehn filling space is a finite complex. By Corollary 4.3, this Dehn filling space is a  $K(\overline{G}, 1)$  space provided  $N$  is sufficiently deep.  $\square$

In [PS24], we gave a rather elaborate argument based on the spectral sequence derived in [PS24, Theorem 4.3] to obtain the next result, which now follows easily from excision on Dehn filling space.

**Corollary 4.5** ([PS24, Theorem A (ii)]). *Let  $(G, H, N)$  be a Cohen–Lyndon triple and  $A$  be a  $\overline{G}$ -module. Then, there is natural isomorphism induced by the quotient maps  $G \rightarrow \overline{G}$  and  $H \rightarrow \overline{H}$ ,*

$$H^n(G, H; A) \cong H^n(\overline{G}, \overline{H}; A), \quad \forall n \geq 0.$$

*Proof.* Recall that the space  $\tilde{X}$  of Theorem 4.2 is obtained by gluing the mapping cylinders  $M_{\tilde{\phi}}, M_{\tilde{\psi}}$  along  $\overline{G} \times_{\overline{H}}(N \setminus EH)$ . By excision [Hat02, Corollary 2.24], for all  $\overline{G}$ -modules  $A$ , we have a natural isomorphism

$$H_G^n(M_{\tilde{\phi}}, \overline{G} \times_{\overline{H}}(N \setminus EH); A) \cong H_G^n(\tilde{X}, M_{\tilde{\psi}}; A), \quad \forall n \geq 0.$$

Since  $\tilde{X}$  is a model for  $E\overline{G}$ , by definition,

$$H^n(\overline{G}, \overline{H}; A) = H_G^n(\tilde{X}, M_{\tilde{\psi}}; A) = H_G^n(\tilde{X}, M_{\tilde{\phi}}; A).$$

Since  $\langle\langle N \rangle\rangle$  acts trivially on  $A$ , the pair  $(M_{\tilde{\phi}}, \overline{G} \times_{\overline{H}}(N \setminus EH))$  and its  $\langle\langle N \rangle\rangle$ -cover  $(\tilde{M}_{\tilde{\phi}}, G \times_H EH)$  have the same cohomology, i.e.

$$H_G^n(\tilde{M}_{\tilde{\phi}}, G \times_H EH; A) = H_G^n(M_{\tilde{\phi}}, \overline{G} \times_{\overline{H}}(N \setminus EH); A).$$

By definition,  $H^n(G, H; A) = H_G^n(\tilde{M}_{\tilde{\phi}}, G \times_H EH; A)$ .  $\square$

**Lemma 4.6.** *Let  $G$  be a group with a hyperbolically embedded subgroup  $H \hookrightarrow_h G$ . Then for all sufficiently deep normal  $N \triangleleft H$ , we have the following commutative diagram with exact rows*

$$(12) \quad \begin{array}{ccccc} \dots & \mathcal{N}(\overline{G}) \otimes_{\mathcal{N}(\overline{H})} \mathcal{H}_n(N \setminus EH; \mathcal{N}(\overline{H})) & \longrightarrow & \mathcal{H}_n(\langle\langle N \rangle\rangle \setminus EG; \mathcal{N}(\overline{G})) & \longrightarrow & \mathcal{H}_n(\overline{G}, \overline{H}; \mathcal{N}(\overline{G})) & \dots \\ & \downarrow & & \downarrow & & \downarrow = & \\ \dots & \mathcal{N}(\overline{G}) \otimes_{\mathcal{N}(\overline{H})} \mathcal{H}_n(\overline{H}; \mathcal{N}(\overline{H})) & \longrightarrow & \mathcal{H}_n(\overline{G}; \mathcal{N}(\overline{G})) & \longrightarrow & \mathcal{H}_n(\overline{G}, \overline{H}; \mathcal{N}(\overline{G})) & \dots \end{array}$$

Moreover, if both  $G$  and  $\overline{H}$  are of type  $F_k$  for some  $k \in \mathbb{N}$ , then for  $n \leq k$  we have the following inequalities

$$(13) \quad \begin{aligned} b_n^{(2)}(\langle\langle N \rangle\rangle \backslash EG; \overline{G}) - b_n^{(2)}(N \backslash EH; \overline{H}) &\leq b_n^{(2)}(\overline{G}, \overline{H}) \\ &\leq b_n^{(2)}(\langle\langle N \rangle\rangle \backslash EG; \overline{G}) + b_{n-1}^{(2)}(N \backslash EH; \overline{H}), \end{aligned}$$

$$(14) \quad b_n^{(2)}(\overline{G}) - b_n^{(2)}(\overline{H}) \leq b_n^{(2)}(\overline{G}, \overline{H}) \leq b_n^{(2)}(\overline{G}) + b_{n-1}^{(2)}(\overline{H}).$$

*Proof.* We use the notation from Definition 4.1. The map of CW pairs

$$(M_{\tilde{\phi}}, \overline{G} \times_{\overline{H}} N \backslash (EH)) \rightarrow (\tilde{X}, M_{\tilde{\psi}})$$

gives rise to a commutative diagram with exact rows

$$(15) \quad \begin{array}{ccccc} \dots & \mathcal{H}_n(\overline{G} \times_{\overline{H}}(N \backslash EH); \mathcal{N}(\overline{G})) & \longrightarrow & \mathcal{H}_n(M_{\tilde{\phi}}; \mathcal{N}(\overline{G})) & \longrightarrow & \mathcal{H}_n(M_{\tilde{\phi}}, \overline{G} \times_{\overline{H}}(N \backslash EH); \mathcal{N}(\overline{G})) & \dots \\ & \downarrow & & \downarrow & & \downarrow & \\ \dots & \mathcal{H}_n(M_{\tilde{\psi}}; \mathcal{N}(\overline{G})) & \longrightarrow & \mathcal{H}_n(\tilde{X}; \mathcal{N}(\overline{G})) & \longrightarrow & \mathcal{H}_n(\tilde{X}, M_{\tilde{\psi}}; \mathcal{N}(\overline{G})) & \dots \end{array}$$

By [Lüc02, Theorem 6.54 (7)], we have natural isomorphisms

$$\begin{aligned} \mathcal{H}_n(\overline{G} \times_{\overline{H}}(N \backslash EH); \mathcal{N}(\overline{G})) &\cong \mathcal{N}(\overline{G}) \otimes_{\mathcal{N}(\overline{H})} \mathcal{H}_n(N \backslash EH; \mathcal{N}(\overline{H})), \\ \mathcal{H}_n(M_{\tilde{\psi}}; \mathcal{N}(\overline{G})) &\cong \mathcal{N}(\overline{G}) \otimes_{\mathcal{N}(\overline{H})} \mathcal{H}_n(\overline{H}; \mathcal{N}(\overline{H})). \end{aligned}$$

By Definition 4.1 and Corollary 4.3, we also have

$$\begin{aligned} \mathcal{H}_n(M_{\tilde{\phi}}; \mathcal{N}(\overline{G})) &\cong \mathcal{H}_n(\langle\langle N \rangle\rangle \backslash EG; \mathcal{N}(\overline{G})), \\ \mathcal{H}_n(\tilde{X}; \mathcal{N}(\overline{G})) &\cong \mathcal{H}_n(\overline{G}; \mathcal{N}(\overline{G})). \end{aligned}$$

By excision, we have the natural isomorphism

$$\mathcal{H}_n(M_{\tilde{\phi}}, \overline{G} \times_{\overline{H}}(N \backslash EH); \mathcal{N}(\overline{G})) \cong \mathcal{H}_n(\tilde{X}, M_{\tilde{\psi}}; \mathcal{N}(\overline{G})) \cong \mathcal{H}_n(\overline{G}, \overline{H}; \mathcal{N}(\overline{G})).$$

The above isomorphisms transform (15) to (12).

If both  $G$  and  $\overline{H}$  are of type  $F_k$  for some  $k \in \mathbb{N}$ , then from (15) and Proposition 2.1 (iii) we deduce the inequalities for all  $n \leq k$ ,

$$(16) \quad \begin{aligned} b_n^{(2)}(M_{\tilde{\phi}}; \overline{G}) - b_n^{(2)}(\overline{G} \times_{\overline{H}}(N \backslash EH); \overline{G}) &\leq b_n^{(2)}(M_{\tilde{\phi}}, \overline{G} \times_{\overline{H}}(N \backslash EH); \overline{G}) \\ &\leq b_n^{(2)}(M_{\tilde{\phi}}; \overline{G}) + b_{n-1}^{(2)}(\overline{G} \times_{\overline{H}}(N \backslash EH); \overline{G}), \end{aligned}$$

$$(17) \quad \begin{aligned} b_n^{(2)}(\tilde{X}; \overline{G}) - b_n^{(2)}(M_{\tilde{\psi}}; \overline{G}) &\leq b_n^{(2)}(\tilde{X}, M_{\tilde{\psi}}; \overline{G}) \\ &\leq b_n^{(2)}(\tilde{X}; \overline{G}) + b_{n-1}^{(2)}(M_{\tilde{\psi}}; \overline{G}), \end{aligned}$$

as these numbers are all finite by Proposition 2.1 (i).

By [Lüc02, Theorem 6.54 (7)], we have

$$\begin{aligned} b_n^{(2)}(\overline{G} \times_{\overline{H}}(N \backslash EH); \overline{G}) &= b_n^{(2)}(N \backslash EH; \overline{H}), \\ b_{n-1}^{(2)}(\overline{G} \times_{\overline{H}}(N \backslash EH); \overline{G}) &= b_{n-1}^{(2)}(N \backslash EH; \overline{H}), \\ b_n^{(2)}(M_{\tilde{\psi}}; \overline{G}) &= b_n^{(2)}(\overline{H}), \end{aligned}$$

$$b_{n-1}^{(2)}(M_{\bar{\psi}}; \overline{G}) = b_{n-1}^{(2)}(\overline{H}).$$

The above equalities turn (16) and (17) into (13) and (14), respectively.  $\square$

We now ready to prove our first general result on  $L^2$ -Betti numbers of groups, providing bounds for the  $L^2$ -Betti numbers of Dehn fillings of virtually locally indicable groups.

**Theorem 4.7.** *Let  $G$  be a virtually locally indicable group of type  $F_{k+1}$  for some  $k \geq 0$ , and let  $H \hookrightarrow_h G$  be a hyperbolically embedded subgroup. Then for every  $\delta > 0$ , there exists a finite subset  $\mathcal{F}_\delta \subset H \setminus \{1\}$  such that if a normal subgroup  $N \triangleleft H$  satisfies  $N \cap \mathcal{F}_\delta = \emptyset$  and  $H/N$  is of type  $F_{k+1}$ , then*

$$(18) \quad b_{k+1}^{(2)}(G/\langle\langle N \rangle\rangle) < b_{k+1}^{(2)}(G) + b_k^{(2)}(H) + b_{k+1}^{(2)}(H/N) + \delta,$$

and for  $n \leq k$ , we have

$$(19) \quad -b_n^{(2)}(H) - b_{n-1}^{(2)}(H/N) - \delta < b_n^{(2)}(G/\langle\langle N \rangle\rangle) - b_n^{(2)}(G) < b_{n-1}^{(2)}(H) + b_n^{(2)}(H/N) + \delta.$$

*Proof.* By Corollary 3.4, there exists a finite subset  $\mathcal{S}_\delta \subset G \setminus \{1\}$  such that if a normal subgroup  $M \triangleleft G$  satisfies  $M \cap \mathcal{S}_\delta = \emptyset$ , then we have

$$(20) \quad b_{k+1}^{(2)}(M \setminus EG; G/M) < b_{k+1}^{(2)}(G) + \delta/2,$$

and

$$(21) \quad |b_n^{(2)}(G) - b_n^{(2)}(M \setminus EG; G/M)| < \delta/2 \text{ for } n \leq k.$$

By [DGO17, Theorem 7.19 (c)], there exists a finite subset  $\mathcal{F}_1 \subset H \setminus \{1\}$  such that if a normal subgroup  $N \triangleleft H$  satisfies  $N \cap \mathcal{F}_1 = \emptyset$ , then  $\langle\langle N \rangle\rangle \cap \mathcal{S}_\delta = \emptyset$ .

By [DGO17, Corollary 4.32 (b)],  $H$  is of type  $F_{k+1}$ . Therefore, by Corollary 3.4, there exists a finite subset  $\mathcal{F}_2 \subset G \setminus \{1\}$  such that if a normal subgroup  $N \triangleleft H$  satisfies that  $N \cap \mathcal{F}_2 = \emptyset$ , then we have

$$(22) \quad b_{k+1}^{(2)}(N \setminus EH; H/N) < b_{k+1}^{(2)}(H) + \delta/2,$$

and

$$(23) \quad |b_n^{(2)}(H) - b_n^{(2)}(N \setminus EH; H/N)| < \delta/2 \text{ for } n \leq k.$$

By Lemma 4.6, there exists a finite subset  $\mathcal{F}_3 \subset H \setminus \{1\}$  such that if a normal subgroup  $N \triangleleft H$  satisfies  $N \cap \mathcal{F}_3 = \emptyset$ , then (13) and (14) hold for all  $n \leq k+1$ .

Let  $\mathcal{F}_\delta = \mathcal{F}_1 \cup \mathcal{F}_2 \cup \mathcal{F}_3$  and fix a normal subgroup  $N \triangleleft H$  such that  $N \cap \mathcal{F}_\delta = \emptyset$  and  $H/N$  is of type  $F_{k+1}$ .

By (20), (21) and the choice of  $N$ , we have

$$(24) \quad b_{k+1}^{(2)}(\langle\langle N \rangle\rangle \setminus EG; G/\langle\langle N \rangle\rangle) < b_{k+1}^{(2)}(G) + \delta$$

and

$$(25) \quad |b_n^{(2)}(G) - b_n^{(2)}(\langle\langle N \rangle\rangle \setminus EG; G/\langle\langle N \rangle\rangle)| < \delta/2 \text{ for } n \leq k.$$

Inequalities (18) and (19) follow by combining (13), (14), (22) (23), (24), and (25).  $\square$

We will prove Theorem 1.1 using Theorem 4.7. In preparation, we first prove the following lemma:

**Lemma 4.8.** *Let  $G$  be a finitely generated infinite group. Then for every  $n > 0$ , there exists a finite subset  $\mathcal{F} \subset G \setminus \{1\}$  such that if a normal subgroup  $N \triangleleft G$  satisfies  $N \cap \mathcal{F} = \emptyset$ , then  $|G/N| > n$ , and thus  $b_0^{(2)}(G/N) < 1/n$ .*

*Proof.* Let  $S$  be a finite generating set of  $G$ . For  $r \in \mathbb{N}^+$  we denote by  $B_r$  the subgraph of the Cayley graph  $\Gamma(G, S)$  consisting of all points within distance  $r$  from the vertex 1. Choose  $R \in \mathbb{N}^+$  such that  $B_R$  has more than  $n$  vertices. Let  $\mathcal{F} = G \cap B_{2R} \setminus \{1\}$ . Then for every two vertices  $u \neq v \in B_R$ , we have  $uv^{-1} \in \mathcal{F}$ . So if  $N \triangleleft G$  is a normal subgroup such that  $N \cap \mathcal{F} = \emptyset$ , then  $u$  and  $v$  have different images in  $G/N$ , whence  $|G/N| > |G \cap B_R| > n$ . The last inequality of the lemma follows from [Lüc02, Theorem 6.54 (8b)].  $\square$

*Proof of Theorem 1.1.* First consider the case where  $|H| < \infty$ . Fix  $n \geq 0$  and let  $\mathcal{F}_{n,\delta} = H \setminus \{1\}$ . Then the only normal subgroup  $N \triangleleft H$  that satisfies  $N \cap \mathcal{F}_{n,\delta} = \emptyset$  is  $\{1\}$ . So  $\overline{G} = G$  and the result follows.

Now consider the case where  $|H| = \infty$ . Since  $H$  is amenable, then so is  $H/N$ , being a quotient of an amenable group. By [Lüc02, Theorem 6.37], we have

$$(26) \quad b_n^{(2)}(H) = b_n^{(2)}(H/N) = 0 \text{ for all } n \geq 1.$$

and by [Lüc02, Theorem 6.54 (8b)], we have

$$(27) \quad b_0^{(2)}(H) = 0.$$

By Lemma 4.8, there exists a finite subset  $\mathcal{F}'_\delta \subset H \setminus \{1\}$  such that

$$(28) \quad b_0^{(2)}(\overline{H}) < \delta/2$$

whenever  $N \cap \mathcal{F}'_\delta = \emptyset$ . Now fix  $n \geq 0$ . Theorem 4.7 provides us a finite subset  $\mathcal{F}'_{n,\delta} \subset H \setminus \{1\}$  such that if  $N \cap \mathcal{F}'_{n,\delta} = \emptyset$ , then

$$(29) \quad -b_n^{(2)}(H) - b_{n-1}^{(2)}(H/N) - \delta/2 < b_n^{(2)}(G/\langle\langle N \rangle\rangle) - b_n^{(2)}(G) < b_{n-1}^{(2)}(H) + b_n^{(2)}(H/N) + \delta/2.$$

Let  $\mathcal{F}_{n,\delta} = \mathcal{F}'_\delta \cup \mathcal{F}'_{n,\delta}$ . So, if  $N \cap \mathcal{F}_{n,\delta} = \emptyset$ , then all of (26), (27), (28) and (29) hold, which yields the desired result.  $\square$

**4.2. Bounding finite subgroups.** Next, we refine the above approximation result to an equality by applying the Atiyah Conjecture. To achieve this, we proceed as follows:

- (i) bound the order of finite subgroups arising in Dehn fillings (this will be addressed in the current subsection), and
- (ii) apply the theory of virtually compact special groups to show that certain Dehn fillings satisfy the Atiyah Conjecture (this will be discussed in the following subsection).

**Lemma 4.9.** *Let  $G$  be a group and let  $N \triangleleft G$  be a normal subgroup. Then*

$$\max \mathcal{FLN}(G) \leq \max \mathcal{FLN}(N) \cdot \max \mathcal{FLN}(G/N).$$

*Proof.* Let  $K \leq G$  be a finite subgroup and let  $\phi: G \rightarrow G/N$  be the natural quotient map. Then  $|K| = |K \cap \ker \phi| \cdot |\phi(K)| \leq \max \mathcal{FLN}(N) \cdot \max \mathcal{FLN}(G/N)$ .  $\square$



**Lemma 4.10.** *Let  $G$  be a group that is hyperbolic relative to a subgroup  $H$ . Then there exists a finite subset  $\mathcal{F} \subset H \setminus \{1\}$  such that if a normal subgroup  $N \triangleleft H$  satisfies  $N \cap \mathcal{F} = \emptyset$ , then*

$$\max \mathcal{FLN}(G/\langle\langle N \rangle\rangle) \leq \max(\mathcal{FLN}(G) \cup \mathcal{FLN}(H/N)).$$

*Proof.* For every  $N \triangleleft H$ , let  $\phi_N: G \rightarrow G/\langle\langle N \rangle\rangle$  be the natural quotient map. By [DG18, Lemma 4.3], there exists a finite subset  $\mathcal{S} \subset G \setminus \{1\}$  and a collection  $\mathcal{C}$  of finite subgroups of  $G$  such that if a normal subgroup  $N \triangleleft H$  satisfies  $\langle\langle N \rangle\rangle \cap \mathcal{S} = \emptyset$ , then the following holds:

- (\*) for every finite subgroup  $K \leq G/\langle\langle N \rangle\rangle$ , there exists a subgroup  $L \leq G$  such that  $K \leq \phi_N(L)$  and  $L$  either conjugates to  $H$  or belongs to  $\mathcal{C}$ .

By [Osi07, Theorem 1.1], there exists a finite subset  $\mathcal{F} \subset H \setminus \{1\}$  such that if a normal subgroup  $N \triangleleft H$  satisfies  $N \cap \mathcal{F} = \emptyset$ , then  $\langle\langle N \rangle\rangle \cap \mathcal{S} = \emptyset$  and  $H \cap \langle\langle N \rangle\rangle = N$ .

Now let  $N \triangleleft H$  be a normal subgroup such that  $N \cap \mathcal{F} = \emptyset$ , and let  $K \leq G/\langle\langle N \rangle\rangle$  be a finite subgroup. Let  $L$  and  $\mathcal{C}$  be given by (\*). If  $L$  conjugates to  $H$ , then  $K$  conjugates to a finite subgroup of  $\phi_N(H) \cong H/N$ , and thus  $|K| \leq \max \mathcal{FLN}(H/N)$ . If  $L \in \mathcal{C}$ , then  $|K| \leq |L| \leq \max \mathcal{FLN}(G)$ .  $\square$

### 4.3. Compact special Dehn filling.

**Definition 4.11.** A group  $G$  is *virtually compact cubical* if there is a finite-index subgroup  $H \leq G$  such that  $H$  is the fundamental group of a compact locally CAT(0) cube complex  $X$ .

The group  $G$  is *virtually compact special* if in addition  $X$  is special. Note that [AGM16, GM22] call such groups *virtually special*, while [Sch15] calls such groups *virtually cocompact special*.

A group  $G$  is *virtually special* if there is a finite-index subgroup  $H \leq G$  such that  $H$  is the fundamental group of a special locally CAT(0) cube complex.

Below, we collect some known facts about virtually (compact) special groups that will be used later.

**Proposition 4.12.** *The following hold.*

- (i) *Any cover of a special cube complex is special. In particular, any subgroup of a virtually special group is virtually special.*
- (ii) *Every virtually compact cubical group is of type  $F_\infty$  and has finite virtual cohomological dimension.*
- (iii) *Every finitely generated virtually special group is virtually locally indicable.*
- (iv) *Every finitely generated virtually special group is virtually residually finite rationally solvable (RFRS).*
- (v) ([GM22]) *Let  $G$  be a virtually compact cubical group that is hyperbolic relative to a family of virtually abelian groups. Then  $G$  is virtually compact special. In particular, virtually compact cubical hyperbolic groups are virtually compact special, which was first proved by [Ago13, Theorem 1.1].*

*Proof.*

- (i) Let  $X$  be a special cube complex. By [Wis21, Proposition 6.2],  $X$  admits a local-isometry to a Salvetti complex  $S$ . Let  $Y$  be a cover of  $X$ . Then, the covering  $Y \rightarrow X$ , followed by the local-isometry  $X \rightarrow S$ , induces a local-isometry  $Y \rightarrow S$ . Thus, by [Wis21, Proposition 6.2],  $Y$  is special.

Let  $G$  be a special group and  $H$  a subgroup. Consider a finite-index subgroup  $G_0 \leq G$  that is a fundamental group of special cube complex  $X$ . Let  $Y$  be the cover of  $X$  whose fundamental group is  $H_0 := H \cap G_0$ , which is a finite-index subgroup of  $H$ . Then, by the above,  $Y$  is special.

- (ii) Let  $G$  be a virtually compact special group. Then  $G$  has a finite-index subgroup  $G_1$  that is the fundamental group of some compact locally CAT(0) cube complex. In particular,  $G_1$  is of type  $F$ .
- (iii) Let  $G$  be a finitely generated virtually special group. By [HW08, Theorem 1.1],  $G$  has a finite-index subgroup  $G_1$  that embeds into a finitely generated right-angled Artin group. By [DK92, p. 387 and p.389], every finitely generated right-angled Artin group is residually torsion-free nilpotent, and thus is residually  $p$  for any prime  $p$  by [Gru57, Theorem 2.1], and thus is bi-orderable by [Rhe73], and thus is locally indicable by [RR02, Corollary 1.4]. So  $G_1$  is locally indicable.
- (iv) Let  $G$  be a finitely generated virtually special group. By (iii),  $G$  has a finite-index subgroup  $G_1$  that embeds into a finitely generated right-angled Artin group. By [Ago08, Corollary 2.3], finitely generated right-angled Artin groups are virtually RFRS, and thus so is the subgroup  $G_1$ .
- (v) Let  $\{H_i\}_{i=1}^n$  be a family of virtually abelian subgroups of  $G$  such that  $G$  is hyperbolic relative to  $\{H_i\}_{i=1}^n$ , let  $X$  be a CAT(0) cube complex with a free geometric action of a finite-index normal subgroup  $G_0 \leq G$ , and let  $\{K_j\}_{j=1}^m$  be the peripheral structure of  $G_0$  induced by  $\{H_i\}_{i=1}^n$ . As each  $H_i$  is abelian, so is each  $K_j$ , and thus by [GM22, Remark 1.4] the relatively hyperbolic pair  $(G_0, \{K_j\}_{j=1}^m)$  and the action  $G_0 \curvearrowright X$  satisfy Assumptions 1.1 and 1.2 of [GM22]. Hence, by [GM22, Theorem A], the group  $G_0$ , and therefore the group  $G$ , is virtually compact special.

The last statement follows by noting that every hyperbolic group is hyperbolic relative to the trivial subgroup  $\{1\}$ .

□

**Theorem 4.13.** *Let  $G$  be a virtually compact cubical hyperbolic group, and let  $H \leq G$  be an almost malnormal quasi-convex subgroup. Then there exists a finite-index torsion-free normal subgroup  $K \triangleleft H$  such that for every sufficiently deep normal subgroup  $N \triangleleft H$  that satisfies*

- $N \leq K$ , and
- $K/N$  is torsion-free virtually compact cubical hyperbolic,

we have

$$b_1^{(2)}(G/\langle\langle N \rangle\rangle) \leq b_1^{(2)}(G).$$

*Remark 4.14.* Note that in Theorem 1.2 we assume that  $G$  is hyperbolic relative to  $H$ , whereas here we assume that  $H$  is an almost malnormal quasi-convex subgroup. These assumptions are equivalent, by [Bow12, Theorem 7.11].

*Proof.* If  $H$  is finite we can simply take  $K = \{1\}$ , and the theorem holds trivially. Below, we assume  $|H| = \infty$ . Then, by [Bow12, Theorem 7.11],  $G$  is hyperbolic relative to  $H$ . For every  $N \triangleleft H$ , let  $\phi_N: G \rightarrow G/\langle\langle N \rangle\rangle$  be the natural quotient map.

**Claim 4.14.1.** *There is a finite-index normal subgroup  $K \triangleleft H$  such that if a normal subgroup  $N \triangleleft H$  satisfies that  $N \leq K$  and  $\phi_N(H)$  is virtually compact cubical hyperbolic, then  $G/\langle\langle N \rangle\rangle$  is virtually compact special hyperbolic.*

*Proof of claim.* The claim follows from [AGM16, Theorem 2.7]. Note that [AGM16, Theorem 2.7] requires both  $G$  and  $\phi_N(H)$  to be virtually compact special hyperbolic, but by Proposition 4.12, this is equivalent to being virtually compact cubical and hyperbolic.  $\square$

Fix a  $K \triangleleft H$  that satisfies Claim 4.14.1. As  $G$  is virtually compact cubical, it is virtually torsion-free. Let  $G_1 \triangleleft G$  be a torsion-free finite-index normal subgroup. By Proposition 4.12,  $G$  is virtually compact special, and thus is virtually locally indicable, again by Proposition 4.12. By Corollary 3.8, there exists a finite subset  $\mathcal{F}_1 \subset H \setminus \{1\}$  such that if a normal subgroup  $N \triangleleft H$  satisfies  $N \cap \mathcal{F}_1 = \emptyset$ , then

$$(30) \quad b_1^{(2)}(G/\langle\langle N \rangle\rangle) < b_1^{(2)}(G) + \frac{1}{\max\{|G/G_1|, |H/K|\}}.$$

By [Osi07, Theorem 1.1], there exists a finite subset  $\mathcal{F}_2 \subset H \setminus \{1\}$  such that if a normal subgroup  $N \triangleleft H$  satisfies  $N \cap \mathcal{F}_2 = \emptyset$ , then  $\phi_N(H) \cong H/N$ . Let  $\mathcal{F}_3 \subset H \setminus \{1\}$  be the finite subset given by Lemma 4.10. Let

$$\mathcal{F}_\delta = \mathcal{F}_1 \cup \mathcal{F}_2 \cup \mathcal{F}_3$$

and consider a normal subgroup  $N \triangleleft H$  such that  $N \leq K$ ,  $N \cap \mathcal{F}_\delta = \emptyset$  and  $K/N$  is torsion-free virtually compact cubical hyperbolic. As  $\phi_N(K) \cong K/N$  is a finite-index subgroup of  $\phi_N(H) \cong H/N$ , we have that  $\phi_N(H)$  is virtually compact cubical hyperbolic. Claim 4.14.1 implies that  $G/\langle\langle N \rangle\rangle$  is virtually compact special, and thus satisfies the Atiyah Conjecture [Sch15]. Therefore, by Lemma 4.10

$$(31) \quad b_1^{(2)}(G/\langle\langle N \rangle\rangle) \in \frac{1}{\mathcal{FLN}(G/\langle\langle N \rangle\rangle)} \cdot \mathbb{Z} \leq \frac{1}{(\max(\mathcal{FLN}(G) \cup \mathcal{FLN}(H/N)))!} \cdot \mathbb{Z}.$$

As  $G_1$  is a torsion-free normal subgroup of  $G$ , Lemma 4.9 implies

$$(32) \quad \max \mathcal{FLN}(G) \leq \max \mathcal{FLN}(G_1) \cdot \max \mathcal{FLN}(G/G_1) = |G/G_1|.$$

Similarly, by considering  $K/N$  in  $H/N$  we have

$$\mathcal{FLN}(H/N) \leq \mathcal{FLN}(K/N) \cdot \mathcal{FLN}(H/K) = |H/K|.$$

Combining these with (31), we get

$$(33) \quad b_1^{(2)}(G/\langle\langle N \rangle\rangle) \in \frac{1}{\max\{|G/G_1|, |H/K|\}} \cdot \mathbb{Z}.$$

As  $G$  is virtually compact special, it satisfies the Atiyah Conjecture [Sch15], and thus

$$(34) \quad b_1^{(2)}(G) \in \frac{1}{\mathcal{FLN}(G)} \cdot \mathbb{Z} \leq \frac{1}{\max\{|G/G_1|, |H/K|\}} \cdot \mathbb{Z}$$

by (32).

The theorem follows from (30), (33), and (34).  $\square$

*Remark 4.15.* In the proof of Theorem 4.13, the passage from  $H$  to the finite-index subgroup  $K$  is a consequence of the Malnormal Special Quotient Theorem, which guarantees that the Dehn filling quotient is virtually compact special, which in turn guarantees that the quotient satisfies the Atiyah Conjecture. If the Atiyah Conjecture is preserved by sufficiently deep Dehn fillings (i.e., whenever  $G$  is a group that satisfies the Atiyah Conjecture and has a hyperbolically embedded subgroup  $H \hookrightarrow_h G$ , the quotient  $G/\langle\langle N \rangle\rangle$  satisfies the Atiyah Conjecture for sufficiently deep  $N \triangleleft H$ ), one can assume  $K = H$ .

Our next result concerns groups hyperbolic relative to abelian subgroups. The lemma below follows by the work of Groves–Manning [GM22] and [Wis21].

**Lemma 4.16.** *Let  $G$  be a virtually compact cubical group that is hyperbolic relative to a finite family of abelian subgroups  $\{H_i\}_{i=1}^k$ . Then there exists a family of finite-index torsion-free subgroups  $\{P_i \triangleleft H_i\}_{i=1}^k$  such that for every family of subgroups  $\{N_i \leq P_i\}_{i=1}^k$  that satisfies that  $P_i/N_i$  is virtually  $\mathbb{Z}$  for all  $i$ , the group  $G/\langle\langle \bigcup_{i=1}^k N_i \rangle\rangle$  is virtually compact special.*

*Proof.* By Proposition 4.12, the group  $G$  is virtually compact special. The desired result then follows from [Wis21, Theorem 15.6]. Note that [Wis21, Theorem 15.6] is stated for virtually sparse special groups, but virtually compact special groups are virtually sparse special (see, e.g., Remark 6.4).  $\square$

**Lemma 4.17.** *Let  $n \geq 2$ . Then for any finite subset  $\mathcal{F} \subset \mathbb{Z}^n \setminus \{0\}$ , there exists a subgroup  $N \leq \mathbb{Z}^n$  such that  $N \cap \mathcal{F} = \emptyset$  and  $\mathbb{Z}^n/N \cong \mathbb{Z}$ .*

*Proof.* Let  $e_1, \dots, e_n$  be a basis of the free abelian group  $\mathbb{Z}^n$ . Let  $p_1, \dots, p_n$  be mutually distinct primes such that the subgroup  $G$  of  $\mathbb{Z}^n$  generated by  $\{p_i e_i \mid i = 1, \dots, n\}$  satisfies  $G \cap \mathcal{F} = \emptyset$ . Consider the natural quotient homomorphisms

$$\alpha: \mathbb{Z}^n \rightarrow \mathbb{Z}^n/G \cong \prod_{i=1}^n (\mathbb{Z}/p_i \mathbb{Z}) \cong \mathbb{Z}/(p_1 \cdots p_n \cdot \mathbb{Z})$$

and

$$\beta: \mathbb{Z} \rightarrow \mathbb{Z}/(p_1 \cdots p_n \cdot \mathbb{Z}).$$

As  $\mathbb{Z}^n$  is free abelian, there exists a homomorphism  $\gamma: \mathbb{Z}^n \rightarrow \mathbb{Z}$  such that  $\beta \circ \gamma = \alpha$ . Let  $N = \ker \gamma$ . Then  $N \cap \mathcal{F} \subset \ker \alpha \cap \mathcal{F} = \emptyset$ .  $\square$

We now state our next main result.

**Theorem 4.18.** *Let  $G$  be a virtually compact cubical group that is hyperbolic relative to an abelian subgroup  $H \leq G$ . Then there exists a finite-index torsion-free subgroup  $K \leq H$  such that for every sufficiently deep subgroup  $N \leq H$  that satisfies  $N \leq K$  and  $K/N$  is torsion-free, we have for all  $n$ ,*

$$(35) \quad b_n^{(2)}(G/\langle\langle N \rangle\rangle) = b_n^{(2)}(G).$$

*Remark 4.19.* In Theorem 4.18, if  $H$  is finite, we can choose  $K$  to be  $\{1\}$ . If  $H$  is virtually  $\mathbb{Z}$ , we can choose the group  $K$  to be an infinite cyclic subgroup of  $H$  and the finite set  $\mathcal{F}$  to consist of just a generator of  $K$ . In this case, the only subgroup  $N \leq K$ , that satisfies that both  $N \cap \mathcal{F} = \emptyset$  and  $K/N$  is torsion-free, is  $\{1\}$ . The desired conclusion clearly follows.

*Proof.* Let  $G_1 \triangleleft G$  be a finite-index normal subgroup that is the fundamental group of some compact locally CAT(0) cube complex, and let  $\{H_i\}_{i \in I}$  be the induced peripheral structure on  $G_1$ . The main idea of the proof is to apply Theorem 4.7, which, by itself, only gives (35) for finitely many  $n$ . The claim below takes care of all other values of  $n$ .

**Claim 4.19.1.** *There exists a finite subset  $\mathcal{F}_1 \subset H \setminus \{1\}$  such that if a subgroup  $N \leq G_1 \cap H$  satisfies that  $N \cap \mathcal{F}_1 = \emptyset$  and  $|H/N| = \infty$ , then*

$$b_n^{(2)}(G/\langle\langle N \rangle\rangle_G) = 0,$$

for all  $n > \text{vcd}(G) + 1$ .

*Proof of claim.* By [Osi07, Theorem 1 (1)] and [PS24, Corollary 4.6], there exists a family of finite subsets  $\{\mathcal{F}_{1i} \subset H_i \setminus \{1\}\}_{i \in I}$  such that whenever a family of groups  $\{N_i \leq H_i\}_{i \in I}$  satisfies that  $N_i \cap \mathcal{F}_{1i} = \emptyset$  for all  $i \in I$ , then the natural maps  $H_i/N_i \rightarrow \overline{G}_1$  are all injective, where

$$\overline{G}_1 := G_1/\langle\langle \bigcup_{i \in I} N_i \rangle\rangle_{G_1},$$

and for all  $n > \text{cd}(G_1) + 1 = \text{vcd}(G) + 1$ , by [PS24, Corollary 4.6 (a)], we have

$$\begin{aligned} \mathcal{H}_n(\overline{G}_1; \mathcal{N}(\overline{G}_1)) &\cong \bigoplus_{i \in I} H_n(H_i/N_i; \mathcal{N}(\overline{G}_1)) \\ &\cong \bigoplus_{i \in I} \mathcal{H}_n(\overline{G}_1 \times_{H_i/N_i} E(H_i/N_i); \mathcal{N}(\overline{G}_1)). \end{aligned}$$

as  $\mathcal{N}(\overline{G}_1)$ -modules. By [Lüc02, Theorem 6.54 (7)], we get

$$(36) \quad b_n^{(2)}(\overline{G}_1) = \sum_{i \in I} b_n^{(2)}(\overline{G}_1 \times_{H_i/N_i} E(H_i/N_i); \mathcal{N}(\overline{G}_1)) = \sum_{i \in I} b_n^{(2)}(H_i/N_i)$$

for  $n > \text{vcd}(G) + 1$ .

We now choose the family  $\{N_i \leq H_i\}_{i \in I}$ . Let  $\mathcal{F}_1 \subset H \setminus \{1\}$  be the subset given by Lemma 2.16 (iv) with respect to the family  $\{\mathcal{F}_{1i}\}_{i \in I}$ , let  $N \leq G_1 \cap H$  be a subgroup that satisfies that  $N \cap \mathcal{F}_1 = \emptyset$  and  $|H/N| = \infty$ , and let  $\{N_i \leq H_i\}_{i \in I} = \varphi^{-1}(\{N\})$ , where  $\varphi$  is the map given by Lemma 2.16. Then

$$b_n^{(2)}(G_1/\langle\langle \bigcup_{i \in I} N_i \rangle\rangle_{G_1}) = \sum_{i \in I} b_n^{(2)}(H_i/N_i) = \sum_{i \in I} b_n^{(2)}((G_1 \cap H)/N) = 0$$

for all  $n > \text{vcd}(G) + 1$ , by [Lüc02, Theorem 6.37] and the fact that  $(G_1 \cap H)/N$  is infinite abelian, being a finite-index subgroup of  $H/N$ .

By Lemma 2.16 (iii),  $G_1/\langle\langle \bigcup_{i \in I} N_i \rangle\rangle_{G_1}$  is a finite-index subgroup of  $G/\langle\langle N \rangle\rangle_G$ , and thus by [Lüc02, Theorem 6.54 (6)] and Lemma 2.16,

$$b_n^{(2)}(G/\langle\langle N \rangle\rangle_G) = b_n^{(2)}(E(G/\langle\langle N \rangle\rangle_G); G/\langle\langle N \rangle\rangle_G)$$

$$\begin{aligned}
&= \frac{1}{[G/\langle\langle N \rangle\rangle_G : G_1/\langle\langle \bigcup_{i \in I} N_i \rangle\rangle_{G_1}]} \cdot b_n^{(2)}\left(E(G/\langle\langle N \rangle\rangle_G) ; G_1/\langle\langle \bigcup_{i \in I} N_i \rangle\rangle_{G_1}\right) \\
&= \frac{1}{[G/\langle\langle N \rangle\rangle_G : G_1/\langle\langle \bigcup_{i \in I} N_i \rangle\rangle_{G_1}]} \cdot b_n^{(2)}(G_1/\langle\langle \bigcup_{i \in I} N_i \rangle\rangle_{G_1}) = 0.
\end{aligned}$$

□

We now apply Theorem 4.7 along with the Atiyah Conjecture. By Proposition 4.12, the group  $G$  is virtually compact special, and thus is virtually locally indicable, again by Proposition 4.12. By Theorem 4.7, we have the following:

**Claim 4.19.2.** *There exists a finite subset  $\mathcal{F}_2 \subset H \setminus \{1\}$  such that if a subgroup  $N \triangleleft H$  satisfies  $N \cap \mathcal{F}_2 = \emptyset$ , then for all  $n \leq \text{vcd}(G) + 1$ ,*

$$\begin{aligned}
&b_n^{(2)}(G) - b_n^{(2)}(H) - b_{n-1}^{(2)}(H/N) - \frac{1}{\max\{|G/G_1|, |H/K|\}} \\
&< b_n^2(G/\langle\langle N \rangle\rangle_G) \\
&< b_n^{(2)}(G) + b_{n-1}^{(2)}(H) + b_n^{(2)}(H/N) + \frac{1}{\max\{|G/G_1|, |H/K|\}}.
\end{aligned}$$

Now, let  $K \leq H$  be the finite-index torsion-free subgroup given by Lemma 4.16. If  $|K/N| = \infty$ , then  $H/N$  is infinite abelian. So by [Lüc02, Theorem 1.44], we have the following:

**Claim 4.19.3.** *If  $|K/N| = \infty$ , then for all  $n \in \mathbb{N}$ ,*

$$b_n^{(2)}(H) = b_{n-1}^{(2)}(H) = b_n^{(2)}(H/N) = b_{n-1}^{(2)}(H/N) = 0.$$

Therefore,

$$|b_n^2(G/\langle\langle N \rangle\rangle_G) - b_n^{(2)}(G)| < \frac{1}{\max\{|G/G_1|, |H/K|\}}!$$

whenever  $N \cap \mathcal{F}_2 = \emptyset$  and  $n \leq \text{vcd}(G) + 1$ .

By Lemma 4.9, we have the following:

**Claim 4.19.4.** *If a subgroup  $N \leq K$  satisfies that  $K/N$  is torsion-free, then*

$$\max \mathcal{FLN}(H/N) \leq \max \mathcal{FLN}(K/N) \cdot \max \mathcal{FLN}(H/K) = |H/K|.$$

We will also need the following:

**Claim 4.19.5.** *There exists a finite subset  $\mathcal{F}_3 \subset H \setminus \{1\}$  such that if a subgroup  $N \leq K$  satisfies that  $N \cap \mathcal{F}_3 = \emptyset$  and  $K/N$  is torsion-free, then*

$$\max \mathcal{FLN}(G/\langle\langle N \rangle\rangle_G) \leq \max\{|G/G_1|, |H/K|\}.$$

*Proof of claim.* By Lemma 4.9, we have

$$(37) \quad \max \mathcal{FLN}(G) \leq \max \mathcal{FLN}(G_1) \cdot \max \mathcal{FLN}(G/G_1) = |G/G_1|.$$

By Lemma 4.10, there exists a finite subset  $\mathcal{F}_3 \subset H \setminus \{1\}$  such that if a subgroup  $N \leq K$  satisfies that  $N \cap \mathcal{F}_3 = \emptyset$  and  $K/N$  is torsion-free, then

$$(38) \quad \max \mathcal{FLN}(G/\langle\langle N \rangle\rangle_G) \leq \max(\mathcal{FLN}(G) \cup \mathcal{FLN}(H/N)).$$

The claim follows from Claim 4.19.4, (37) and (38).  $\square$

For any subgroup  $N \leq H$ , let  $\phi_N: G \rightarrow G/\langle\langle N \rangle\rangle_G$  be the natural quotient map. The following claim follows from [Osi07, Theorem 1.1].

**Claim 4.19.6.** *There exists a finite subset  $\mathcal{F}_4 \subset H \setminus \{1\}$  such that if a subgroup  $N \leq H$  satisfies  $N \cap \mathcal{F}_4 = \emptyset$ , then  $\phi_N(H) \cong H/N$  and  $G/\langle\langle N \rangle\rangle_G$  is hyperbolic relative to  $\phi_N(H)$ .*

Next, we let

$$\mathcal{F} = \mathcal{F}_1 \cup \mathcal{F}_2 \cup \mathcal{F}_3 \cup \mathcal{F}_4.$$

By Remark 4.19, we may assume that  $|H| = \infty$ . So, by enlarging  $\mathcal{F}_1$ , we may assume that  $\mathcal{F} \cap K \neq \emptyset$ . We have the following:

(\*) If a subgroup  $N \leq K$  satisfies  $N \cap \mathcal{F} = \emptyset$ , then  $K/N \neq \{1\}$ .

Now consider a subgroup  $N \leq K$  such that  $N \cap \mathcal{F} = \emptyset$  and  $K/N$  is torsion-free.

**Claim 4.19.7.**

$$b_i^{(2)}(G/\langle\langle N \rangle\rangle_G) \in \frac{1}{\max\{|G/G_1|, |H/K|\}} \cdot \mathbb{Z}.$$

*Proof of claim.* By (\*), the group  $K/N$  is infinite finitely generated free abelian. If  $K/N \cong \mathbb{Z}$ , then Lemma 4.16 implies that  $G/\langle\langle N \rangle\rangle_G$  is virtually compact special, and thus satisfies the Atiyah Conjecture [Sch15]. In particular,

$$b_i^{(2)}(G/\langle\langle N \rangle\rangle_G) \in \frac{1}{\mathcal{FLN}(G/\langle\langle N \rangle\rangle_G)} \cdot \mathbb{Z} \subset \frac{1}{\max\{|G/G_1|, |H/K|\}} \cdot \mathbb{Z},$$

by Claim 4.19.5.

Below, we assume that  $K/N$  is free abelian of finite rank at least 2. We will use Proposition 3.5. Let  $\mathcal{S} \subset G/\langle\langle N \rangle\rangle_G \setminus \{1\}$  be any finite subset. We need to find a normal subgroup of  $G/\langle\langle N \rangle\rangle_G$  satisfying the assumptions in Proposition 3.5. By Claim 4.19.6, Theorem 2.13 and [Osi07, Theorem 1.1], there is a finite subset  $\mathcal{F}' \subset \phi_N(H) \setminus \{1\}$  such that if a subgroup  $\overline{M} \leq \phi_N(H)$  satisfies  $\overline{M} \cap \mathcal{F}' = \emptyset$ , then  $\langle\langle \overline{M} \rangle\rangle_{G/\langle\langle N \rangle\rangle_G} \cap \mathcal{S} = \emptyset$  and  $(G/\langle\langle N \rangle\rangle_G, \phi_N(H), \overline{M})$  is a Cohen–Lyndon triple. By enlarging  $\mathcal{F}'$ , we may assume

$$(39) \quad \phi_N(\mathcal{F}) \subset \mathcal{F}'$$

By assumption,  $\phi_N(K) \cong K/N$  is free abelian of rank at least 2. Use Lemma 4.17 to choose a subgroup  $\overline{M}_{\mathcal{S}} \leq \phi_N(K)$  such that  $\overline{M}_{\mathcal{S}} \cap \mathcal{F}' = \emptyset$  and  $\phi_N(K)/\overline{M}_{\mathcal{S}} \cong \mathbb{Z}$ . Then  $\langle\langle \overline{M}_{\mathcal{S}} \rangle\rangle_{G/\langle\langle N \rangle\rangle_G} \triangleleft G/\langle\langle N \rangle\rangle_G$  and  $\langle\langle \overline{M}_{\mathcal{S}} \rangle\rangle_{G/\langle\langle N \rangle\rangle_G} \cap \mathcal{S} = \emptyset$ .

Let  $M_{\mathcal{S}} \leq K$  be the preimage of  $\overline{M}_{\mathcal{S}} \leq \phi_N(K)$  under the restriction  $\phi_N|_K$ . Then  $M_{\mathcal{S}} \cap \mathcal{F} = \emptyset$  and thus all other claims in the current proof can be applied to  $M_{\mathcal{S}}$ . Note that  $K/M_{\mathcal{S}} \cong \phi_N(K)/\overline{M}_{\mathcal{S}} \cong \mathbb{Z}$ . By Lemma 4.16, we have that

$$(G/\langle\langle N \rangle\rangle_G)/\langle\langle \overline{M}_{\mathcal{S}} \rangle\rangle_{G/\langle\langle N \rangle\rangle_G} \cong G/\langle\langle M_{\mathcal{S}} \rangle\rangle_G$$

is virtually compact special. In particular,  $(G/\langle\langle N \rangle\rangle_G)/\langle\langle \overline{M}_{\mathcal{S}} \rangle\rangle_{G/\langle\langle N \rangle\rangle_G}$  satisfies the Atiyah Conjecture [Sch15]. Another consequence of the virtual compact specialness of  $(G/\langle\langle N \rangle\rangle_G)/\langle\langle \overline{M}_{\mathcal{S}} \rangle\rangle_{G/\langle\langle N \rangle\rangle_G}$  is that  $(G/\langle\langle N \rangle\rangle_G)/\langle\langle \overline{M}_{\mathcal{S}} \rangle\rangle_{G/\langle\langle N \rangle\rangle_G}$  is virtually locally indicable by Proposition 4.12. By Theorem 2.13 and our choice of  $\mathcal{F}'$ , the triple  $(G/\langle\langle N \rangle\rangle_G, \phi_N(H), \overline{M}_{\mathcal{S}})$



has the Cohen–Lyndon property, which implies that  $\langle\langle \overline{M}_{\mathcal{S}} \rangle\rangle_{G/\langle\langle N \rangle\rangle_G}$  is a free product of groups isomorphic to  $\overline{M}_{\mathcal{S}}$ . In particular,  $\langle\langle \overline{M}_{\mathcal{S}} \rangle\rangle_{G/\langle\langle N \rangle\rangle_G}$  is locally indicable. From the short exact sequence

$$1 \rightarrow \langle\langle \overline{M}_{\mathcal{S}} \rangle\rangle_{G/\langle\langle N \rangle\rangle_G} \rightarrow G/\langle\langle N \rangle\rangle_G \rightarrow (G/\langle\langle N \rangle\rangle_G)/\langle\langle \overline{M}_{\mathcal{S}} \rangle\rangle_{G/\langle\langle N \rangle\rangle_G} \rightarrow 1$$

we conclude that  $G/\langle\langle N \rangle\rangle_G$  is virtually locally indicable.

By Claim 4.19.5, we have

$$\max \mathcal{FLN}(G/\langle\langle M_{\mathcal{S}} \rangle\rangle_G) \leq \max\{|G/G_1|, |H/K|\}.$$

The above shows that for every finite subset  $\mathcal{S} \subset G/\langle\langle N \rangle\rangle_G \setminus \{1\}$ , there is a normal subgroup  $\langle\langle \overline{M}_{\mathcal{S}} \rangle\rangle_{G/\langle\langle N \rangle\rangle_G} \triangleleft G/\langle\langle N \rangle\rangle_G$  satisfying the conditions in Proposition 3.5 with  $C = \max\{|G/G_1|, |H/K|\}$ , which then yields the desired result.  $\square$

The theorem then follows from Claim 4.19.3 and Claim 4.19.7.  $\square$

*Remark 4.20.* The torsion-free condition in Theorem 1.2 and Theorem 1.3 cannot be dropped, as shown by the following example. Let  $G'$  be the fundamental group of a hyperbolic 3-manifold and let  $H$  be a maximal cyclic subgroup of  $G'$ . Let  $G = G' *_H G'$  be the amalgam of two copies of  $G'$  glued via the identity map on  $H$ . Then by [Bow12, Theorem 7.11],  $G'$  is hyperbolic relative to  $H$ . So by [Dah03, Theorem 0.1 (3)],  $G$  is hyperbolic relative to  $H$ . As  $H$  is hyperbolic, so is  $G$  by [Osi06, Corollary 2.41]. Note that  $G'$  is virtually compact cubical by [AFW15, Theorem 5.4]. Thus, by [Wis21, Theorem 13.1],  $G$  is virtually compact cubical. So the pair of groups  $G$  and  $H$  satisfy the assumptions of both Theorem 1.2 and Theorem 1.3.

Using a Mayer–Vietoris series argument, we get that  $b_1^{(2)}(G) = 0$ . Let  $h$  be a generator of  $H$  and let  $n \in \mathbb{N}$  be a large integer. By [Osi07, Theorem 1.1], the group  $G'/\langle\langle t^n \rangle\rangle_{G'}$  is infinite and contains  $H/\langle t^n \rangle$  as a subgroup. Then  $G/\langle\langle t^n \rangle\rangle_G$  is the amalgam of two copies of  $G'/\langle\langle t^n \rangle\rangle_{G'}$  along their common subgroup  $H/\langle t^n \rangle$ . By [PT11, Lemma 3.1], we have  $b_1^{(2)}(G/\langle\langle t^n \rangle\rangle_G) \geq 1/n$ , and so  $G/\langle\langle t^n \rangle\rangle_G$  does not satisfy the conclusions of Theorem 1.2 and Theorem 1.3.

## 5. MAIN RESULTS FOR MULTIPLE PERIPHERALS

None of the results involved in the proof of Theorem 4.13 and Theorem 4.18 are confined to the case of a single peripheral subgroup. Below, we state their general versions. Their proofs are completely analogous and therefore left to the reader.

**5.1. Dehn filling space.** Let  $G$  be a group, let  $\{H_\lambda\}_{\lambda \in \Lambda}$  be a family of subgroups of  $G$ , and let  $\{N_\lambda \triangleleft H_\lambda\}_{\lambda \in \Lambda}$  be a family of normal subgroups. For the rest of this subsection, we assume that the triple  $(G, \{H_\lambda\}_{\lambda \in \Lambda}, \{N_\lambda\}_{\lambda \in \Lambda})$  is a *Cohen–Lyndon triple*, i.e., for each  $\lambda$  there exists a family of left transversal of the subgroup  $T_\lambda$  of  $H_\lambda \langle\langle \bigcup_{\lambda \in \Lambda} N_\lambda \rangle\rangle$  in  $G$  such that

$$\langle\langle \bigcup_{\lambda \in \Lambda} N_\lambda \rangle\rangle = *_{\lambda \in \Lambda, t \in T_\lambda} t N_\lambda t^{-1}.$$

Let  $\overline{G} := G/\langle\langle \bigcup_{\lambda \in \Lambda} N_\lambda \rangle\rangle$ , and let  $\overline{H}_\lambda := H_\lambda/N_\lambda$  for each  $\lambda$ . Note that the natural maps  $\overline{H}_\lambda \rightarrow \overline{G}$  are injective (see e.g., [Sun20, Lemma 6.4]), and below we will view each  $\overline{H}_\lambda$  as a

subgroup of  $\overline{G}$ . Let  $BG$  (resp.  $BH_\lambda, B\overline{H}_\lambda$ ) be a  $K(G, 1)$  (resp.  $K(H_\lambda, 1), K(\overline{H}_\lambda, 1)$ ) space that has the homotopy type of a CW complex.

For each  $\lambda$ , there is a classifying map  $\phi_\lambda: BH_\lambda \rightarrow BG$  (resp.  $\psi_\lambda: BH_\lambda \rightarrow B\overline{H}_\lambda$ ) induced by the inclusion  $H_\lambda \hookrightarrow G$  (resp. the quotient map  $q: H_\lambda \rightarrow \overline{H}_\lambda$ ). Let

$$\phi: \bigsqcup_{\lambda \in \Lambda} BH_\lambda \rightarrow BG$$

be the map induced by  $\phi_\lambda, \lambda \in \Lambda$ , and let

$$\psi: \bigsqcup_{\lambda \in \Lambda} BH_\lambda \rightarrow \bigsqcup_{\lambda \in \Lambda} B\overline{H}_\lambda$$

be the disjoint union of  $\psi_\lambda, \lambda \in \Lambda$ . Let  $X$  be the complex obtained by gluing all the mapping cylinders  $M_\phi$  and  $M_\psi$  along the common subcomplex  $\bigsqcup_{\lambda \in \Lambda} BH_\lambda$ . We call  $X$  a *Dehn filling space* associated to the triple  $(G, \{H_\lambda\}_{\lambda \in \Lambda}, \{N_\lambda\}_{\lambda \in \Lambda})$ .

**Theorem 5.1.** *Let  $(G, \{H_\lambda\}_{\lambda \in \Lambda}, \{N_\lambda\}_{\lambda \in \Lambda})$  be a Cohen–Lyndon triple. Then every Dehn filling space associated to the triple  $(G, \{H_\lambda\}_{\lambda \in \Lambda}, \{N_\lambda\}_{\lambda \in \Lambda})$  is a  $K(G/\langle\langle \bigcup_{\lambda \in \Lambda} N_\lambda \rangle\rangle, 1)$ -space.*

**Corollary 5.2.** *Let  $G$  be a group with a family of hyperbolically embedded subgroups  $\{H_\lambda\}_{\lambda \in \Lambda} \hookrightarrow_h G$ . Then for sufficiently deep  $\{N_\lambda \triangleleft H_\lambda\}_{\lambda \in \Lambda}$ , every Dehn filling space associated to the triple  $(G, \{H_\lambda\}_{\lambda \in \Lambda}, \{N_\lambda\}_{\lambda \in \Lambda})$  is a  $K(G/\langle\langle \bigcup_{\lambda \in \Lambda} N_\lambda \rangle\rangle, 1)$ -space.*

**Corollary 5.3.** *Let  $G$  be a group with a finite family hyperbolically embedded subgroups  $\{H_i\}_{i=1}^k \hookrightarrow_h G$ .*

- (i) *If  $G$  is of type  $F_n$  for some  $n \in \mathbb{N}^+ \cup \{\infty\}$ , then there exists a family of finite subsets  $\{\mathcal{F}_i \subset H_i \setminus \{1\}\}_{i=1}^k$  such that if, for every  $i$ , we have  $N_i \cap \mathcal{F}_i = \emptyset$ , then  $G/\langle\langle \bigcup_{i=1}^k N_i \rangle\rangle$  is of type  $F_n$  if and only if all  $H_i/N_i$  are of type  $F_n$ .*
- (ii) *If  $G$  and every  $H_i$  are of type  $F$ , then there exists a family of finite subsets  $\{\mathcal{F}_i \subset H_i \setminus \{1\}\}_{i=1}^k$  such that if for every  $i$ , we have  $N_i \cap \mathcal{F}_i = \emptyset$  and  $H_i/N_i$  is of type  $F$ , then  $G/\langle\langle \bigcup_{i=1}^k N_i \rangle\rangle$  is of type  $F$ .*

**5.2.  $L^2$ -Betti number.** The following results are generalizations of our main results on  $L^2$ -Betti numbers in the presence of multiple peripheral subgroups.

**Theorem 5.4.** *Let  $k \geq 0$  and let  $G$  be a type  $F_{k+1}$  virtually locally indicable group with a finite family of hyperbolically embedded subgroups  $\{H_i\}_{i=1}^\ell \hookrightarrow_h G$ . Then for every  $\delta > 0$ , there exists a family of finite subsets  $\{\mathcal{F}_{i,\delta} \subset H_i \setminus \{1\}\}_{i=1}^\ell$  such that if a family of normal subgroups  $\{N_i \triangleleft H_i\}_{i=1}^\ell$  satisfies  $N_i \cap \mathcal{F}_{i,\delta} = \emptyset$  and  $H_i/N_i$  is of type  $F_{k+1}$  for all  $i$ , then  $\overline{G} := G/\langle\langle \bigcup_{i=1}^\ell N_i \rangle\rangle$  satisfies*

$$b_1^{(2)}(\overline{G}) < b_1^{(2)}(G) + \delta,$$

$$b_{k+1}^{(2)}(\overline{G}) < b_{k+1}^{(2)}(G) + \sum_{i=1}^\ell b_k^{(2)}(H_i) + \sum_{i=1}^\ell b_{k+1}^{(2)}(H_i/N_i) + \delta,$$

and

$$-\sum_{i=1}^\ell b_n^{(2)}(H_i) - \sum_{i=1}^\ell b_{n-1}^{(2)}(H_i/N_i) - \delta < b_n^{(2)}(\overline{G}) - b_n^{(2)}(G) < \sum_{i=1}^\ell b_{n-1}^{(2)}(H_i) + \sum_{i=1}^\ell b_n^{(2)}(H_i/N_i) + \delta$$

for all  $n \leq k$ .

**Corollary 5.5.** *Let  $G$  be a virtually locally indicable group of type  $F_\infty$  and has a finite family of hyperbolically embedded amenable subgroups  $\{H_i\}_{i=1}^m \hookrightarrow_h G$ . Then for every  $\delta > 0$  and  $n \geq 0$ , there exists a family of finite subsets  $\{\mathcal{F}_{n,\delta,i} \subset H \setminus \{1\}\}_{i=1}^m$  such that if a family of normal subgroups  $\{N_i \triangleleft H_i\}_{i=1}^m$  satisfies that, for all  $i$ ,  $N_i \cap \mathcal{F}_{n,\delta,i} = \emptyset$  and  $H_i/N_i$  is of type  $F_\infty$ , then*

$$\left| b_n^{(2)}(G/\langle\langle \bigcup_{i=1}^m N_i \rangle\rangle) - b_n^{(2)}(G) \right| < \delta.$$

*Remark 5.6.* One can prove Corollary 5.5 by proceeding as follows: First, prove Corollary 5.5 under the additional assumption that each  $H_i$  is infinite. The proof will be a generalization of and completely analogous to the proof of Theorem 1.1. Now consider the general case. Suppose that there exists  $\ell < k$  such that  $|H_i| < \infty$  for  $i = \ell + 1, \dots, k$  and  $|H_i| = \infty$  for  $i = 1, \dots, \ell$ . We can then let  $\mathcal{F}_i = H \setminus \{1\}$  for  $i = \ell + 1, \dots, k$ , from which it follows that  $N_i = \{1\}$  for  $i = \ell + 1, \dots, k$ . Moreover, it follows from the definition that, if  $X \subset G$  is a subset such that  $\{H_i\}_{i=1}^k \hookrightarrow_h (G, X)$ , then  $\{H_i\}_{i=1}^\ell \hookrightarrow_h (G, X \cup \bigcup_{i=\ell+1}^k H_i)$ . In particular,  $\{H_i\}_{i=1}^\ell \hookrightarrow_h G$ . So we are reduced to the previously solved special case where the hyperbolically embedded family consists of only infinite amenable groups.

**Theorem 5.7.** *Let  $G$  be a virtually compact cubical hyperbolic group that is hyperbolic relative to a finite family of subgroups  $\{H_i\}_{i=1}^k < G$ . Then there exists a family of finite-index torsion-free normal subgroups  $\{K_i \triangleleft H_i\}_{i=1}^k$  such that the following  $(\star)$  holds.*

$(\star)$  *For each  $i$ , let  $L_i \leq K_i$  be any finite-index subgroup. Then there exists a family of finite subsets  $\{\mathcal{F}_i \subset L_i \setminus \{1\}\}_{i=1}^k$  such that if a family of normal subgroups  $\{N_i \triangleleft H_i\}_{i=1}^k$  satisfies for all  $i$ ,*

- $N_i \leq L_i$ ,
- $N_i \cap \mathcal{F}_i = \emptyset$ , and
- $L_i/N_i$  is torsion-free virtually compact cubical hyperbolic,

then

$$b_1^{(2)}(G/\langle\langle \bigcup_{i=1}^k N_i \rangle\rangle) \leq b_1^{(2)}(G).$$

**Theorem 5.8.** *Let  $G$  be a virtually compact cubical group that is hyperbolic relative to a finite family of abelian subgroups  $\{H_i\}_{i=1}^k$ . Then there exists a family of finite-index torsion-free subgroups  $\{K_i \triangleleft H_i\}_{i=1}^k$  such that the following  $(\star)$  holds.*

$(\star)$  *For each  $i$ , let  $L_i \leq K_i$  be any finite-index subgroup. Then there exist finite subsets  $\{\mathcal{F}_i \subset L_i \setminus \{1\}\}_{i=1}^k$  such that if a family of normal subgroups  $\{N_i \triangleleft H_i\}_{i=1}^k$  satisfies for all  $i$ ,*

- (i)  $N_i \leq L_i$ ,
- (ii)  $N_i \cap \mathcal{F}_i = \emptyset$ , and
- (iii)  $L_i/N_i$  is torsion-free,

then, for all  $n \in \mathbb{N}$ ,

$$b_n^{(2)}(G/\langle\langle \bigcup_{i=1}^k N_i \rangle\rangle) = b_n^{(2)}(G).$$

*Remark 5.9.* Theorem 5.8 can be proved by proceeding along the the following line: First prove the theorem under the additional assumption that each  $H_i$  is virtually  $\mathbb{Z}^m$  for  $m \geq 2$ . The proof will be a generalization of and completely analogous to the proof of Theorem 4.18. Now consider the general case. Suppose that there exists  $\ell < k$  such that, for  $i = \ell + 1, \dots, k$ , the group  $H_i$  is virtually cyclic; and for  $i = 1, \dots, \ell$ , the group  $H_i$  is virtually  $\mathbb{Z}^m$  for some  $m \geq 2$ . Then by [Osi06, Theorem 2.40],  $G$  is hyperbolic relative to the family  $\{H_i\}_{i=1}^\ell$ . Moreover, if for some  $i$ , the group  $H_i$  is finite, then we can take  $K_i = \{1\}$ , from which it follows that  $N_i = \{1\}$ . And if for some  $i$ , the group  $H_i$  is virtually  $\mathbb{Z}$ , then we can choose  $K_i$  to be isomorphic to  $\mathbb{Z}$ . In this case any finite-index subgroup  $L_i \leq K_i$  will be isomorphic to  $\mathbb{Z}$  and we can let  $\mathcal{F}_i$  to consists of the generator of  $L_i$ . The only  $N_i$  that satisfies conditions (i), (ii) and (iii) in Theorem 5.8 is  $\{1\}$ . Therefore,  $N_i = \{1\}$  for  $i = \ell + 1, \dots, k$  and we are reduced to the previously solved special case where  $G$  is hyperbolic relative to the family  $\{H_i\}_{i=1}^\ell$  with each  $H_i$  virtually  $\mathbb{Z}^m$  for some  $m \geq 2$ .

## 6. VIRTUALLY SPARSE SPECIAL GROUPS

We will study  $L^2$ -Betti numbers of Dehn fillings of cusped hyperbolic manifolds. For this purpose, Theorem 4.18 is not satisfactory because it only deals with virtually compact cubical groups, and, to the best of the authors' knowledge, only a finite number of cusped hyperbolic manifolds of dimension at least 4 with virtually compact cubical fundamental group have been constructed so far. So, in this section, we will generalize Theorem 4.18 to deal with multiple cusps and virtually sparse special groups. Notably, Bergeron–Wise [BW12] proved that all cusped arithmetic hyperbolic manifolds have virtually sparse special fundamental group (see Theorem 7.4). We start with the definition of a sparse group. For more details on this topic, we refer to [Wis21].

**Definition 6.1.** A *quasiflat* is a pair  $(X, A)$ , where  $A$  is a finitely generated virtually abelian group and  $X$  is a locally finite CAT(0) cube complex with a proper action by  $A$  such that there are finitely many  $A$ -orbits of hyperplanes.

**Definition 6.2.** A quasiflat  $(X, A)$  is *closable* if there exists a quasiflat  $(Y, B)$  such that  $B$  acts properly and cocompactly on  $Y$ , and there is an inclusion  $A \hookrightarrow B$  and an equivariant convex embedding  $X \rightarrow Y$ .

The quasiflat  $(X, A)$  is called *abelian closable* if additionally  $(Y, B)$  can be chosen such that  $B$  is free abelian. Note that if  $(X, A)$  is a closable quasiflat, then there is a finite-index normal abelian subgroup  $A_0 \triangleleft A$  such that the quasiflat  $(X, A_0)$  is abelian closable. Moreover, for every finite-index subgroup  $A_1 \leq A_0$ , the quasiflat  $(X, A_1)$  is abelian closable.

**Definition 6.3.** Let  $G$  be a group that is hyperbolic relative to a finite family of virtually abelian groups  $\{H_i\}_{i=1}^n$ . An action  $G \curvearrowright X$  on a CAT(0) cube complex  $X$  is called *cosparse* if there exists a compact subcomplex  $K \subset X$  and a family of quasiflats  $\{(F_i, H_i)\}_{i=1}^n$  with  $H_i = \text{Stab}_G(F_i)$  for each  $i$  such that

- (i)  $X = GK \cup (\bigcup_{i=1}^n GF_i)$ ;
- (ii) for each  $i$  we have  $(F_i \cap GK) \subset H_i K_i$  for some compact subcomplex  $K_i \subset X$ ; and
- (iii) for  $i, j \in \{1, \dots, n\}$  and  $g \in G$ , either  $(F_i \cap gF_j) \subset GK$  or  $i = j$  and  $F_i = gF_j$ .

The group  $G$  is called *sparse special* if additionally the action  $G \curvearrowright X$  is free and the quotient  $G \backslash X$  is special.

*Remark 6.4.* We note that any compact special group  $G$  that is hyperbolic relative to a family of virtually abelian groups  $\{H_i\}_{i=1}^n$  is sparse special. To see this, consider the special CAT(0) cube complex  $X$  on which  $G$  acts properly and cocompactly. Let  $K$  be a compact subcomplex so that  $X = GK$ . Then, the assumptions (i) and (iii) of Definition 6.3 are immediately satisfied. By Theorem 3.6 of [WW17], for each  $i$ , there is an  $H_i$ -invariant convex subcomplex  $F_i$  on which  $H_i$  acts cocompactly, which gives (ii).

**Lemma 6.5.** *Let  $G$  be a group that is hyperbolic relative to a finite family of virtually abelian groups  $\{H_i\}_{i=1}^n$ . Suppose  $G$  acts cosparsely on a CAT(0) cube complex  $X$ . Then, the restriction of the action to any finite index subgroup  $G' \leq G$  is cosparse.*

*In particular, any finite index subgroup of a sparse special group is sparse special.*

*Proof.* Let  $T$  be a right transversal of  $G'$  in  $G$ . For  $t \in T$  and  $i \in \{1, \dots, n\}$ , let  $H'_{i,t} = tH_it^{-1} \cap G'$ . Then  $G'$  is hyperbolic relative to the family of virtually abelian groups  $\{H'_{i,t}\}_{\{1 \leq i \leq n, t \in T\}}$  by [Hru10, Theorem 9.1]. Following the notation of Definition 6.3, we define  $K' = TK$  and the family of quasiflats  $\{(tF_i, H'_{i,t})\}_{\{1 \leq i \leq n, t \in T\}}$ . Note that  $H'_{i,t} = \text{Stab}_{G'}(tF_i)$  for each  $i$  and  $t$ . Next, we verify the conditions of Definition 6.3.

(i).  $X = GK \cup (\bigcup_{i=1}^n GF_i) = G'K' \cup (\bigcup_{\{1 \leq i \leq n, t \in T\}} G'tF_i)$ .

(ii). For each  $i$  and  $t$ , we have  $(tF_i \cap G'K') \subset tH_iK_i \subset H'_{i,t}K'_i$ , for some compact subcomplex  $K'_i \subset X$ .

(iii). For  $i, j \in \{1, \dots, n\}$ ,  $t_1, t_2 \in T$  and  $g \in G$ , either  $(t_1F_i \cap gt_2F_j) = t_1(F_i \cap t_1^{-1}gt_2F_j) \subset G'K'$  or  $i = j$  and  $F_i = t_1^{-1}gt_2F_j$  implying  $t_1F_i = gt_2F_j$ .

For the last claim of the lemma suppose  $\Gamma'$  is finite index subgroup of a sparse special group  $\Gamma$ . Then  $\Gamma$  acts cosparsely on a CAT(0) cube complex  $X$  such that the quotient is special. By the above, the group  $\Gamma'$  acts cosparsely on  $X$  and the quotient is a finite cover of the special complex  $X/\Gamma$  and hence, it is special by Proposition 4.12 (i).  $\square$

**Lemma 6.6.** *Let  $G$  be a finitely generated virtually special group that is hyperbolic relative to a finite family of virtually abelian subgroups  $\{H_i\}_{i=1}^k$ . For each  $i$ , let  $K_i \leq H_i$  be any finite-index abelian subgroup. Then there exists a finite-index special normal subgroup  $G_0 \leq G$  such that  $G_0 \cap H_i \leq K_i$  for all  $i$ . If  $G$  is virtually sparse (resp. compact) special, then we can take  $G_0$  to be sparse (resp. compact) special.*

*Proof.* Consider one of the  $K_i$ . By [Wis21, Corollary 6.8], there exists a nested sequence of finite-index subgroups of  $G$

$$G_{i0} > G_{i1} > \dots$$

such that  $\bigcap_{j \geq 0} G_{ij} = K_i$ . The sequence of numbers  $\{[H_i : H_i \cap G_{ij}]\}_{j \geq 0}$  is a monotone increasing sequence of integers, with upper bound  $[H_i : K_i]$ . Since

$$\bigcap_{j \geq 0} (H_i \cap G_{ij}) = H_i \cap \left( \bigcap_{j \geq 0} G_{ij} \right) = K_i,$$

this sequence of numbers will eventually stabilize, so there exists  $j_i$  such that  $[H_i : H_i \cap G_{ij_i}] = [H_i : K_i]$ , i.e.,  $H_i \cap G_{ij_i} = K_i$ .

Let  $G'_0 \leq G$  be a finite-index special subgroup, and let  $G_0$  be the normal core of  $G'_0 \cap (\bigcap_{i=1}^k G_{ij_i})$ . Then by Proposition 4.12 (i),  $G_0$  satisfies all requirements. If  $G$  is virtually

sparse (resp. compact) special, then we can take  $G'_0$  to be sparse (resp. compact) special, in which case  $G_0$  will also be sparse (resp. compact) special by Lemma 6.5.  $\square$

**Lemma 6.7.** *Let  $G$  be a finitely generated virtually sparse special group that is hyperbolic relative to a finite family of infinite virtually abelian groups  $\{H_i\}_{i=1}^n$ . Then there exist a finite-index normal subgroup  $G_0 \triangleleft G$  and a compact cubical group  $G'_0$  such that the following hold:*

- (i)  $G'_0$  splits as a graph of groups  $\mathcal{T}$  whose underlying graph is a star, i.e., obtained by gluing a finite number of edges to a central vertex.
- (ii) The central vertex group of  $\mathcal{T}$  is  $G_0$ .
- (iii) Every non-central vertex group is infinite finitely generated free abelian.
- (iv) Let  $\{K_j\}_{j=1}^m$  be the induced peripheral structure on  $G_0$ . Then each edge group of  $\mathcal{T}$  is isomorphic to some  $K_j$ , and different edge groups correspond to different  $K_j$ .
- (v) For all  $i$ , the group  $G_0 \cap H_i$  is infinite finitely generated free abelian.
- (vi)  $G_0$  is hyperbolic relative to the family of edge groups of  $\mathcal{T}$ .

*Proof.* Let  $G_1 \triangleleft G$  be a finite-index normal sparse special subgroup. So there exists a CAT(0) cube complex  $X$  with a cosparsely cospecial action of  $G_1$ . By [Wis21, Lemma 7.51], each quasiflat of  $X$  is closable.

Let  $\{L_k\}_{k=1}^p$  be the induced peripheral structure of  $G_1$ . Each  $L_k$  corresponds to a quasiflat  $Y_k$  of  $X$ . There exists a finite-index free abelian normal subgroup  $N_k \triangleleft L_k$  such that the quasiflat  $(Y_k, N_k)$  is abelian closable. By Lemma 6.6, there exists a finite-index normal subgroup  $G_0 \triangleleft G$  such that  $G_0 \cap L_k \leq N_k$  for all  $k$ . Then, by Lemma 6.5, the action  $G_0 \curvearrowright X$  is cosparsely, the quotient  $G_0 \backslash X$  is special, and every quasiflat of the action  $G_0 \backslash X$  is abelian closable. The desired result follows from [Wis21, Theorem 7.54]. Note that [Wis21, Theorem 7.54] only asserts that the non-central vertex groups of  $\mathcal{T}$  are virtually abelian, but a close inspection of its proof shows that, if each quasiflat is abelian closable, then  $G'_0$  can be constructed in such a way that the non-central vertex groups of  $\mathcal{T}$  are abelian: There is a CAT(0) cube complex  $X$  with a free cosparsely action of  $G_0$  such that the quotient  $G_0 \backslash X$  is special and every quasiflat of  $X$  is abelian closable. Each quasiflat of  $X$  has the form  $(F_j, K_j)$  for some  $j \in \{1, \dots, m\}$ , and there exists a quasiflat  $(F'_j, K'_j)$  such that  $K'_j$  is finitely generated free abelian and there is an inclusion  $K_j \hookrightarrow K'_j$  and an equivariant embedding  $F_j \hookrightarrow F'_j$ . The group  $G'_0$  is the quotient of  $G_0 * (\ast_{j=1}^m K'_j)$  by identifying, for each  $j$ , the two embeddings  $K'_j \hookrightarrow K_j \hookrightarrow G_0$  of  $K_j$ .  $\square$

**Lemma 6.8.** *Let  $A \leq B$  be finitely generated free abelian groups. Then there exists a finite-index subgroup  $C \leq B$  such that  $A \leq C$  and for every subgroup  $D \leq A$  such that  $A/D$  is torsion-free, we have that  $C/D$  is also torsion-free.*

*Proof.* Considering  $A'$  as a  $\mathbb{Z}$ -module,  $A' \otimes_{\mathbb{Z}} \mathbb{Q}$  is a  $\mathbb{Q}$ -vector space. Let  $v_1, \dots, v_n$  be elements of  $B \subset B \otimes_{\mathbb{Z}} \mathbb{Q}$  that together with a basis of  $A \otimes_{\mathbb{Z}} \mathbb{Q}$  give a basis of  $B \otimes_{\mathbb{Z}} \mathbb{Q}$ . Let  $C \leq B$  be the subgroup generated by  $A$  and  $v_1, \dots, v_n$ . Then  $C$  has finite-index in  $B$ . Moreover,  $A$  is a direct factor of  $C$  and thus if  $D \leq A$  is a subgroup then  $A/D$  is a direct factor of  $C/D$ . Thus, if  $A/D$  is torsion-free then so is  $C/D$ .  $\square$

**Theorem 6.9.** *Let  $G$  be a finitely generated virtually sparse special group that is hyperbolic relative to a finite family of virtually abelian subgroups  $\{H_i\}_{i=1}^n$ . Then there exists a family of finite-index normal subgroups  $\{K_i \leq H_i\}_{i=1}^n$  such that the following holds:*

(★) *For each  $i$ , let  $L_i \triangleleft H_i$  be any finite-index normal subgroup such that  $L_i \leq K_i$ . Then there exists a family of finite subsets  $\{\mathcal{F}_i \subset L_i \setminus \{1\}\}_{i=1}^n$  such that if a family of normal subgroups  $\{N_i \triangleleft H_i\}$  satisfies that, for all  $i$ ,*

- $N_i \leq L_i$ ,
- $N_i \cap \mathcal{F}_i = \emptyset$ , and
- $L_i/N_i$  is torsion-free,

then, for all  $n \in \mathbb{N}$ ,

$$b_n^{(2)}(G/\langle\langle \bigcup_{i=1}^n N_i \rangle\rangle) = b_n^{(2)}(G).$$

*Proof.* As we will take normal closures in different groups, we will use  $\langle\langle \cdot \rangle\rangle_G$  instead of  $\langle\langle \cdot \rangle\rangle$ .

If some  $H_i$  is finite, for each choice of finite-index normal subgroup  $L_i \triangleleft H_i$ , we can let  $\mathcal{F}_i = L_i \setminus \{1\}$ . Moreover, by [Osi06, Theorem 2.40], we can exclude  $H_i$  from the peripheral structure of  $G$ . Therefore, we may assume that every  $H_i$  is infinite.

By Lemma 6.7, there exists a finite-index normal subgroup  $G_0 \triangleleft G$  and a compact special group  $G'_0$  which splits as a graph of groups  $\mathcal{T}$  such that  $G_0, G'_0$  and  $\mathcal{T}$  satisfy the conclusions of Lemma 6.7.

Let  $\{H_{0j}\}_{j=1}^m$  be the peripheral structure of  $G_0$  induced by  $\{H_i\}_{i=1}^n$ . Then each  $H_{0j}$  is an edge group of  $\mathcal{T}$ . The edge of  $\mathcal{T}$  corresponding to  $H_{0j}$  is incident to a unique non-central vertex, whose vertex group will be denoted by  $H'_{0j}$ . By [Dah03, Theorem 0.1 (2)],  $G'_0$  is hyperbolic relative to the family of abelian subgroups  $\{H'_{0j}\}_{j=1}^m$ .

**Claim 6.9.1.**  $b_*^{(2)}(G'_0) = b_*^{(2)}(G_0)$ .

*Proof of the claim.* Consider the Mayer–Vietoris series:

$$\cdots \rightarrow \bigoplus_{j=1}^m H_r(H_{0j}; \mathcal{N}(G'_0)) \rightarrow \left( \bigoplus_{j=1}^m H_r(H'_{0j}; \mathcal{N}(G'_0)) \right) \oplus H_r(G_0; \mathcal{N}(G'_0)) \rightarrow H_r(G'_0; \mathcal{N}(G'_0)) \rightarrow \cdots$$

Since  $H_{0j}$  and  $H'_{0j}$  are infinite abelian, the desired result follows by taking  $\dim_{G'_0}$  and using Theorem 6.37 and Theorem 6.54 (7) of [Lüc02].  $\square$

Let  $\{K'_{0j} \triangleleft H'_{0j}\}_{j=1}^m$  be the family of finite-index normal subgroups given by Theorem 5.8 such that the (★) property of Theorem 5.8 holds.

For the rest of this proof, we will apply Lemma 2.16 multiple times. In each of these applications, we are always applying Lemma 2.16 with respect to the pair of groups  $G_0 \triangleleft G$  and the peripheral structures  $\{H_i\}_{i=1}^n$  and  $\{H_{0j}\}_{j=1}^m$ .

Let  $\{\overline{K}_i \leq H_i\}_{i=1}^n$  be the family of finite-index subgroups given by Lemma 2.16 (v) with respect to the family of finite-index subgroups  $\{K'_{0j} \cap H_{0j}\}_{j=1}^m$ . For each  $i$ , let  $K_i$  be the normal core of  $\overline{K}_i$  in  $H_i$ . Then  $K_i$  has finite-index in  $H_i$ .



Now consider a family of finite-index subgroups  $\{L_i \leq K_i\}_{i=1}^n$  such that each  $L_i$  is normal in  $H_i$ . Let  $\{L_{0j} \triangleleft H_{0j}\}_{j=1}^m$  be the family of normal subgroups given by Lemma 2.16 (i) with respect to the family  $\{L_i\}_{i=1}^n$ . Then by Lemma 2.16 (ii), for each  $j$ , the group  $H_{0j}/L_{0j}$  is isomorphic to some  $(G_0 \cap H_i)/L_i$ , and thus  $L_{0j}$  has finite-index in  $H_{0j}$ . Let  $\{K_{0j} \triangleleft H_{0j}\}_{j=1}^m$  be the family of normal subgroups given by Lemma 2.16 (i) with respect to the family  $\{K_i \triangleleft H_i\}_{i=1}^n$ . Then Lemma 2.16 (v) implies that  $K_{0j} \leq K'_{0j} \cap H_{0j}$  for all  $j$ , and thus  $L_{0j} \leq K'_{0j}$ .

For each  $j$ , Lemma 6.8, applied to the pair of finitely generated free abelian groups  $L_{0j} \leq K'_{0j}$ , gives rise to a finite-index subgroup  $L'_{0j} \leq K'_{0j}$  such that for every subgroup  $N_{0j} \leq L_{0j}$  with  $L_{0j}/N_{0j}$  torsion-free, the group  $L'_{0j}/N_{0j}$  is torsion-free as well.

By [Osi07, Theorem 1.1], Theorem 5.8 and our choice of  $\{K'_{0j}\}_{j=1}^m$ , there exists a family of finite subsets  $\{F'_{0j} \subset L'_{0j} \setminus \{1\}\}_{j=1}^m$  such that if a family of subgroups  $\{N_{0j} \leq L'_{0j}\}_{j=1}^m$  satisfies that for all  $j$ ,

- $N_{0j} \cap F'_{0j} = \emptyset$ , and
- $L'_{0j}/N_{0j}$  is torsion-free,

then

$$(40) \quad b_*^{(2)}(G'_0 / \langle\langle \bigcup_{j=1}^m N_{0j} \rangle\rangle_{G'_0}) = b_*^{(2)}(G'_0),$$

and

$$(41) \quad \langle\langle \bigcup_{j=1}^m N_{0j} \rangle\rangle_{G'_0} \cap K'_{0j} = N_{0j} \text{ for all } j.$$

We will specify our choice of  $\{N_{0j} \leq L'_{0j}\}_{j=1}^m$  later.

Let  $\{\mathcal{F}_i \subset H_i \setminus \{1\}\}_{i=1}^n$  be the family of finite subsets given by Lemma 2.16 (iv) with respect to the family  $\{F'_{0j} \cap H_{0j}\}_{j=1}^m$ . Now fix a family of normal subgroups  $\{N_i \triangleleft H_i\}_{i=1}^n$  such that for all  $i$ ,

- $N_i \leq L_i$ ,
- $N_i \cap \mathcal{F}_i = \emptyset$ , and
- $L_i/N_i$  is torsion-free.

Let  $\{N_{0j} \triangleleft H_{0j}\}_{j=1}^m$  be the family of normal subgroups given by Lemma 2.16 (i) with respect to the family  $\{N_i\}_{i=1}^n$ . Then  $N_{0j} \cap F'_{0j} = \emptyset$  for all  $j$ . Moreover, by Lemma 2.16 (ii), we have that, for every  $j$ , the group  $L_{0j}/N_{0j}$  is isomorphic to some  $L_i/N_i$ , and thus is torsion-free. Therefore, so is  $L'_{0j}/N_{0j}$ , by Lemma 6.8 and our choice of  $L'_{0j}$ . Hence, (40) and (41) hold.

Note that, as  $N_{0j} \leq K_{0j} \leq K'_{0j}$ , (41) implies

$$N_{0j} \leq \langle\langle \bigcup_{j=1}^m N_{0j} \rangle\rangle_{G'_0} \cap K_{0j} \leq \langle\langle \bigcup_{j=1}^m N_{0j} \rangle\rangle_{G'_0} \cap K'_{0j} = N_{0j}.$$

In particular,

$$(42) \quad \langle\langle \bigcup_{j=1}^m N_{0j} \rangle\rangle_{G'_0} \cap K_{0j} = N_{0j} \text{ for all } j.$$

By enlarging  $\mathcal{F}_i$  if necessary, we may assume that

$$(43) \quad |K_i/N_i| = \infty \text{ for all } i,$$

and thus

$$(44) \quad |K'_{0j}/N_{0j}| = |K_{0j}/N_{0j}| = \infty \text{ for all } j.$$

**Claim 6.9.2.**  $G'_0/\langle\langle\bigcup_{j=1}^m N_{0j}\rangle\rangle_{G'_0}$  splits as a graph of groups  $\overline{\mathcal{T}}$  with the following properties:

- The underlying graph of  $\overline{\mathcal{T}}$  is the same as that of  $\mathcal{T}$ .
- The central vertex group is  $G_0/\langle\langle\bigcup_{j=1}^m N_{0j}\rangle\rangle_{G_0}$ .
- The non-central vertex groups, as well as the edge groups, are infinite abelian.

*Proof of the claim.* Construct a graph of groups  $\overline{\mathcal{T}}$  from  $\mathcal{T}$  by the following procedure:

- Let  $v$  be a non-central vertex of  $\mathcal{T}$ . Then  $G_v = K'_{0j}$  for some  $j$ . Replace  $G_v$  by  $K'_{0j}/N_{0j}$ .
- Let  $v$  be the central vertex of  $\mathcal{T}$ . Replace  $G_v$  by  $G_0/\langle\langle\bigcup_{j=1}^m N_{0j}\rangle\rangle_{G_0}$ .
- Let  $e$  be an edge of  $\mathcal{T}$ . Then  $G_e = K_{0j}$  for some  $j$ . Replace  $G_e$  by  $K_{0j}/N_{0j}$ .

By (42), for each  $j$ , the group  $K_{0j}/N_{0j}$  is a subgroup of both  $G_0/\langle\langle\bigcup_{j=1}^m N_{0j}\rangle\rangle_{G_0}$  and  $K'_{0j}/N_{0j}$ . Therefore, the above procedure does produce a graph of groups  $\overline{\mathcal{T}}$ . By (43) and (44), the non-central vertex groups and the edge groups of  $\mathcal{T}$  are infinite abelian. It is now easy to check that the fundamental group of  $\overline{\mathcal{T}}$  is  $G'_0/\langle\langle\bigcup_{j=1}^m N_{0j}\rangle\rangle_{G'_0}$ .  $\square$

Similar to Claim 6.9.1, we have

$$(45) \quad b_*^{(2)}(G'_0/\langle\langle\bigcup_{j=1}^m N_{0j}\rangle\rangle_{G'_0}) = b_*^{(2)}(G_0/\langle\langle\bigcup_{j=1}^m N_{0j}\rangle\rangle_{G_0}).$$

By Lemma 2.16 (iii), we have

$$(46) \quad [G/\langle\langle\bigcup_{i=1}^n N_i\rangle\rangle_G : G_0/\langle\langle\bigcup_{j=1}^m N_{0j}\rangle\rangle] = [G : G_0].$$

The desired result follows from Claim 6.9.1, (40), (45), (46) and [Lüc02, Theorem 6.54 (6)].  $\square$

*Remark 6.10.* In Theorem 6.9, we may assume that the family  $\{K_i\}_{i=1}^n$  satisfies the following additional properties:

- (i) There exists a finite-index sparse special normal subgroup  $G_0 \triangleleft G$  such that  $K_i = G_0 \cap H_i$  for all  $i$ .
- (ii) If  $L_i/N_i \cong \mathbb{Z}$  for all  $i$ , then the group  $G/\langle\langle\bigcup_{i=1}^n N_i\rangle\rangle$  is virtually compact special.

Indeed, by [Wis21, Theorem 15.6], there exists a family of finite-index normal subgroups  $\{K'_i \triangleleft H_i\}_{i=1}^n$  such that if a family of normal subgroups  $\{N_i \triangleleft H_i\}_{i=1}^n$  satisfies that, for all  $i$ ,

- $N_i \leq K'_i$ , and
- $K'_i/N_i$  is virtually cyclic,

then the quotient  $G/\langle\langle \bigcup_{i=1}^n N_i \rangle\rangle$  is virtually compact special.

By Lemma 6.6, there exists a finite-index sparse special normal subgroup  $G_0 \triangleleft G$  such that  $G_0 \cap H_i \leq K_i \cap K'_i$  for all  $i$ . Now simply note that the  $(\star)$  condition continues to hold with  $G_0 \cap H_i$  in place of  $K_i$ .

## 7. APPLICATIONS

Next, we present some applications of our main results.

**7.1. Algebraic fibering.** For  $n \in \mathbb{N}^+ \cup \{\infty\}$ , we say that a group  $G$  *virtually  $FP_n(\mathbb{Q})$ -fibers* if  $G$  has a finite-index subgroup  $G_0$  that has a homomorphism onto  $\mathbb{Z}$  with kernel of type  $FP_k(\mathbb{Q})$ . Algebraic fibrations of groups are intimately related to  $L^2$ -Betti numbers. The following theorem is proved for  $n = 1$  by Kielak, and then extended to all  $n$  by Fisher.

**Theorem 7.1** ([Kie20, Fis21]). *For a virtually RFRS group  $G$  of type  $FP_n(\mathbb{Q})$ , the following are equivalent:*

- (i)  $G$  *virtually  $FP_n(\mathbb{Q})$ -fibers.*
- (ii)  $b_i^{(2)}(G) = 0$  for  $i \leq n$ .

By combining the above theorem with our computations on  $L^2$ -Betti numbers, we obtain:

**Corollary 7.2.** *Let  $G$  be a virtually compact cubical hyperbolic group. Suppose that  $G$  *virtually  $FP_1(\mathbb{Q})$ -fibers* and is hyperbolic relative to a family of subgroups  $\{H_i\}_{i=1}^n$ . Then there exists a family of finite-index torsion-free normal subgroups  $\{K_i \triangleleft H_i\}_{i=1}^n$  such that for every family of sufficiently deep normal subgroups  $\{N_i \triangleleft H_i\}_{i=1}^n$  that satisfies that, for all  $i$ ,*

- $N_i \leq K_i$ , and
- $K_i/N_i$  is torsion-free virtually compact cubical hyperbolic,

*then the group  $G/\langle\langle \bigcup_{i=1}^n N_i \rangle\rangle$  *virtually  $FP_1(\mathbb{Q})$ -fibers.**

The assumption of Corollary 7.2 seemingly differs from that of Corollary 1.5, but these assumptions are in fact equivalent (see Remark 4.14).

*Proof.* By Proposition 4.12,  $G$  is of type  $F_\infty$ , and thus is of type  $FP_\infty(\mathbb{Q})$ . First consider the case where  $G$  is virtually  $\mathbb{Z}$ . For each  $i$ , let  $\mathcal{F}_i = \emptyset$ . It remains to choose a finite-index normal subgroup  $K_i \triangleleft H_i$  for each  $i$ . If  $|H_i| < \infty$ , then we let  $K_i = \{1\}$  and  $\mathcal{F}_i = \emptyset$ . If  $|H_i| = \infty$ , then  $H_i = G$  and we will let  $K_i = G$ .

Now suppose that a family of normal subgroups  $\{N_i \triangleleft H_i\}_{i=1}^n$  is given such that, for all  $i$ ,

- $N_i \leq K_i$ , and
- $K_i/N_i$  is torsion-free.

If  $|H_i| < \infty$  for all  $i$ , then by construction  $N_i = \{1\}$  for all  $i$  and thus the group  $G/\langle\langle \bigcup_{i=1}^n N_i \rangle\rangle = G$  *virtually  $FP_1(\mathbb{Q})$ -fibers*. If some  $H_{i_0}$  is infinite, then for  $i \neq i_0$  we will have  $|H_i| < \infty$ . Therefore,  $G/\langle\langle \bigcup_{i=1}^n N_i \rangle\rangle = G/\langle\langle N_{i_0} \rangle\rangle = K_{i_0}/N_{i_0}$ . As the group  $K_{i_0}/N_{i_0}$  is torsion-free virtually cyclic, it is either  $\{1\}$  or  $\mathbb{Z}$ . In either case, the group  $K_{i_0}/N_{i_0}$ , and thus the group  $G/\langle\langle \bigcup_{i=1}^n N_i \rangle\rangle$  *virtually  $FP_1(\mathbb{Q})$ -fibers.*

It remains to consider the case where  $G$  is not virtually  $\mathbb{Z}$ . As  $G$  virtually  $FP_1(\mathbb{Q})$ -fibers, [Lüc02, Theorem 6.63] implies that  $b_1^{(2)}(G) = 0$ .

By Theorem 4.13, [Osi07, Theorem 1.1] and [AGM16, Theorem 2.7], there exists a family of finite-index torsion-free normal subgroup  $\{K_i \triangleleft H_i\}_{i=1}^n$  such that for every family of sufficiently deep normal subgroup  $\{N_i \triangleleft H_i\}_{i=1}^n$  that satisfies that, for all  $i$ ,  $N_i \leq K_i$  and  $K_i/N_i$  is torsion-free virtually compact cubical hyperbolic, we have that

- (i)  $G/\langle\langle \bigcup_{i=1}^n N_i \rangle\rangle$  is virtually compact special (and thus is of type  $FP_\infty(\mathbb{Q})$  and virtually RFRS by Proposition 4.12);
- (ii)  $G/\langle\langle \bigcup_{i=1}^n N_i \rangle\rangle$  is non-elementary hyperbolic, in particular,  $|G/\langle\langle \bigcup_{i=1}^n N_i \rangle\rangle| = \infty$ , and thus  $b_0^{(2)}(G/\langle\langle \bigcup_{i=1}^n N_i \rangle\rangle) = 0$  by [Lüc02, Theorem 6.54 (8b)]; and
- (iii)  $b_1^{(2)}(G/\langle\langle \bigcup_{i=1}^n N_i \rangle\rangle) = 0$ .

The desired result then follows from Theorem 7.1.  $\square$

**Corollary 7.3.** *Let  $G$  be a finitely generated virtually sparse special group that is hyperbolic relative to a family of virtually abelian subgroups  $\{H_i\}_{i=1}^n$ , and let  $k \in \mathbb{N}^+$ . Then there exists a family of finite-index torsion-free normal subgroups  $\{K_i \triangleleft H_i\}_{i=1}^n$  such that for every family of sufficiently deep normal subgroups  $\{N_i \triangleleft H_i\}_{i=1}^n$  such that for all  $i$ ,*

- $N_i \leq K_i$ , and
- $K_i/N_i \cong \mathbb{Z}$ ,

*we have that the group  $G/\langle\langle \bigcup_{i=1}^n N_i \rangle\rangle$  virtually  $FP_k(\mathbb{Q})$ -fibers if and only if so does  $G$ .*

*Proof.* By Theorem 2.11 and Remark 2.13 of [Pat23], each  $H_i$  is finitely generated virtually abelian, and thus is of type  $F_\infty$ . By Theorem 2.11 and Remark 2.13 of [Pat23] again, the group  $G$  is of type  $F_\infty$  (in particular,  $FP_\infty(\mathbb{Q})$ ). By Proposition 4.12, the group  $G$  is virtually RFRS. Therefore,  $G$  virtually  $FP_n(\mathbb{Q})$ -fibers if and only if  $b_j^{(2)}(G) = 0$  for  $j \leq k$  by Theorem 7.1.

By Corollary 4.4, Lemma 4.16, and Theorem 6.9, there exists a family of finite-index normal subgroups  $\{K_i \triangleleft H_i\}_{i=1}^n$  such that for every family of sufficiently deep normal subgroups  $\{N_i \triangleleft H_i\}_{i=1}^n$  such that for all  $i$ ,  $N_i \leq K_i$  and  $K_i/N_i \cong \mathbb{Z}$ , we have the following:

- The group  $G/\langle\langle \bigcup_{i=1}^n N_i \rangle\rangle$  is virtually compact special, and thus is virtually RFRS and of type  $F_\infty$  by Proposition 4.12.
- For all  $j$ , we have  $b_j^{(2)}(G/\langle\langle N \rangle\rangle) = 0$  if and only if  $b_j^{(2)}(G) = 0$ .

Note that by Remark 6.10, we may assume that the subgroups  $K_i$  are contained in a finite-index sparse special subgroup of  $G$ . In particular, we may assume that the  $K_i$  are torsion-free.

Now, Theorem 7.1 implies that  $G/\langle\langle N \rangle\rangle$  virtually  $FP_k(\mathbb{Q})$ -fibers if and only if  $b_j^{(2)}(G/\langle\langle N \rangle\rangle_G) = 0$  for  $j \leq k$ , which is equivalent to  $b_j^{(2)}(G) = 0$  for  $j \leq k$ , which in turn is equivalent to that  $G$  virtually  $FP_k(\mathbb{Q})$ -fibers.  $\square$

**7.2. Arithmetic lattices in  $SO(n, 1)$ .** We apply the results of Section 6 to hyperbolic arithmetic lattices.

**Theorem 7.4** (Bergeron–Wise [BW12]). *Let  $G < SO(n, 1)$  be a non-uniform arithmetic lattice. Then  $G$  is virtually sparse special.*

*Proof.* In [BW12, Theorem 6.2], Bergeron–Wise proved that  $G$  is virtually special by the following argument: The group  $G$  has a quasi-isometrically embedded codimension-1 subgroup that intersects every peripheral subgroup of  $G$  along an infinite subgroup. The authors applied [BW12, Proposition 5.2] to obtain an action  $G \curvearrowright X$  of  $G$  on a CAT(0) cube complex  $X$  and argue that the action is proper and that the quotient  $G_0 \backslash X$  is special, where  $G_0$  is a finite-index torsion-free subgroup given by the Selberg’s Lemma. Note that [BW12, Proposition 5.2] is strengthened by [Wis21, Theorem 7.9] which asserts that the action  $G \curvearrowright X$  is cosparsely as well.  $\square$

The following theorem is a direct consequence of [JX00, Theorem 3.3] and [Lüc02, Theorem 6.54 (6b)]. We provide the proof for the convenience of the reader.

**Theorem 7.5.** *Let  $G < SO(n, 1)$  be a lattice. Then*

$$b_i^{(2)}(G) = \begin{cases} 0, & \text{if } n \neq n/2 \\ |\chi(G)| = \frac{2 \cdot \text{vol}(\mathbb{H}^n/G)}{\Omega_n}, & \text{if } i = n/2, \end{cases}$$

where  $\Omega_n$  is the volume of the unit  $n$ -sphere and  $\chi(G)$  is the rational-valued Euler characteristic of  $G$ .

*Proof.* By the Selberg Lemma,  $G$  has a finite-index torsion-free subgroup  $H$ . Thus,  $\mathbb{H}^n/H$  is a hyperbolic  $n$ -manifold. By [JX00, Theorem 2.3],

$$b_i^{(2)}(H) = \begin{cases} 0, & \text{if } n \neq n/2 \\ |\chi(\mathbb{H}^n/H)|, & \text{if } i = n/2. \end{cases}$$

By [KZ01, Theorem 3.3],  $|\chi(\mathbb{H}^n/H)| = 2 \cdot \text{vol}(\mathbb{H}^n/H)/\Omega_n$ . Note that

$$\begin{aligned} \text{vol}(\mathbb{H}^n/H) &= [G : H] \cdot \text{vol}(\mathbb{H}^n/G) \\ \chi(H) &= [G : H] \cdot \chi(G) && \text{by [Bro94, p. 249],} \\ b_*^{(2)}(H) &= [G : H] \cdot b_*^{(2)}(G) && \text{by [Lüc02, Theorem 6.54 (6b)],} \end{aligned}$$

from which the desired result follows.  $\square$

The theorem below summarizes our results in the setting of hyperbolic lattices.

**Theorem 7.6.** *Let  $G < SO(n, 1)$  be any non-uniform lattice with  $n \geq 2$ , and let  $\{H_i\}_{i=1}^k$  be the peripheral subgroups of  $G$ .*

(i) *Assume  $G$  is virtually locally indicable (e.g.  $G$  is arithmetic). For every  $\delta > 0$ , there exists a family of finite subsets  $\{\mathcal{F}_{\delta,i} \subset H_i \setminus \{1\}\}_{i=1}^k$  such that if a family of normal subgroups  $\{N_i \triangleleft H_i\}_{i=1}^k$  satisfies that  $N_i \cap \mathcal{F}_{\delta,i} = \emptyset$  for all  $i$ , then for  $\ell \leq n$  we have*

$$(47) \quad \left| b_\ell^{(2)}(G/\langle\langle \bigcup_{i=1}^k N_i \rangle\rangle_G) - b_\ell^{(2)}(G) \right| < \delta.$$

In particular, if additionally  $n$  is even and  $\delta$  is sufficiently small, then  $b_{n/2}^{(2)}(G/\langle\langle \bigcup_{i=1}^k N_i \rangle\rangle_G) > 0$ .

(ii) Assume  $G$  is arithmetic. There exists a finite-index sparse special normal subgroup  $G_0 \triangleleft G$  such that the following holds: For each  $i$ , let  $H'_i = G_0 \cap H_i$ . Then there exists a family of finite subsets  $\{\mathcal{F}'_i \subset H'_i \setminus \{1\}\}_{i=1}^k$  such that if a family of normal subgroups  $\{N_i \triangleleft H_i\}_{i=1}^k$  satisfies that, for all  $i$ ,

- $N_i \cap \mathcal{F}'_i = \emptyset$ ,
- $N_i \leq H'_i$ , and
- $H'_i/N_i$  is torsion-free,

then

$$b_*^{(2)}(G/\langle\langle \bigcup_{i=1}^k N_i \rangle\rangle_G) = b_*^{(2)}(G).$$

Moreover, if additionally  $H'_i/N_i \cong \mathbb{Z}$  for all  $i$ , then the quotient  $G/\langle\langle \bigcup_{i=1}^k N_i \rangle\rangle_G$  is virtually compact special. In particular, when  $n$  is odd, the group  $G/\langle\langle \bigcup_{i=1}^k N_i \rangle\rangle_G$  is virtually  $FP_\infty(\mathbb{Q})$ -fibers; and when  $n$  is even, the group  $G/\langle\langle \bigcup_{i=1}^k N_i \rangle\rangle_G$  is virtually  $FP_{n/2-1}(\mathbb{Q})$ -fibers but does not virtually  $FP_{n/2}(\mathbb{Q})$ -fiber.

*Proof.*

(i). For  $\delta > 0$  and each  $\ell \in \{0, 1, \dots, n\}$ , Corollary 5.5 provides a family of finite subsets  $\{\mathcal{F}_{\delta,i,\ell} \subset H_i \setminus \{1\}\}_{i=1}^k$  such that if a family of normal subgroups  $\{N_i \triangleleft H_i\}_{i=1}^k$  satisfies that  $N_i \cap \mathcal{F}_{\delta,i,\ell} = \emptyset$  for all  $i$ , then

$$\left| b_\ell^{(2)}(G/\langle\langle \bigcup_{i=1}^k N_i \rangle\rangle_G) - b_\ell^{(2)}(G) \right| < \delta.$$

Inequality (47) follows by letting  $\mathcal{F}_{\delta,i} = \bigcup_{\ell=0}^n \mathcal{F}_{\delta,i,\ell}$ . The last statement of the theorem follows from (47) and Theorem 7.5.

In the case when  $G$  is arithmetic, by Theorem 7.4, it is finitely generated virtually special, and thus it is virtually locally indicable by Proposition 4.12.

(ii). By Theorem 7.4, the group  $G$  is finitely generated virtually sparse special. So the desired conclusion follows from Theorem 6.9, Remark 6.10, Proposition 4.12, and Theorem 7.1.  $\square$

**7.3. Einstein manifolds.** In this and the next subsections, we will apply our result to cusped arithmetic hyperbolic manifolds. There are two notions of Dehn filling, one for manifolds and the other for groups. To distinguish them, we will call the former *manifold filling*, as defined below.

**Definition 7.7.** Let  $M$  be a compact topological  $n$ -manifold such that  $\partial M$  can be decomposed as a disjoint union  $\partial M = \bigsqcup_{i=1}^k C_i$ , with each  $C_i$  homeomorphic to the  $(n-1)$ -torus  $(S^1)^{n-1}$ .

Let  $T_1, \dots, T_k$  be  $k$ -copies of the solid  $n$ -torus  $D^2 \times (S^1)^{n-2}$ . A *manifold filling*  $\overline{M}$  of  $M$  is a closed  $n$ -manifold obtained by gluing  $T_1, \dots, T_k$  to  $M$  by a homeomorphism

identifying  $\partial T_i$  with  $C_i$  for each  $i$ . For each  $i$ ,  $\partial D^2$  represents an element  $x_i \in \pi_1(\partial T_i)$ , via the homeomorphism  $T_i \cong D^2 \times (S^1)^{n-2}$ . Let  $H_i$  be the image of  $\pi_1(C_i)$  in  $\pi_1(M)$  under the inclusion  $C_i \hookrightarrow M$ . The gluing sends  $x_i$  to an element  $y_i \in H_i$ .

We say that a property  $\mathcal{P}$  holds for all sufficiently deep manifold fillings  $\overline{M}$  of  $M$  if there exists a family of finite subsets  $\{\mathcal{F}_i \subset H_i \setminus \{1\}\}_{i=1}^k$  such that  $\mathcal{P}$  holds whenever  $\langle y_i \rangle \cap \mathcal{F}_i = \emptyset$  for all  $i$ .

**Definition 7.8.** We say that a Dehn filling  $\overline{M}$  of  $M$  is a  $G$ -equivariant manifold filling if  $\overline{M}$  is a manifold filling of  $M$  and  $\pi_1(\overline{M})$  is a  $G$ -equivariant filling of  $G_0$  (see Definition 2.15).

The following Theorem 7.9 is a consequence of Theorem 7.6 (i).

**Theorem 7.9.** Let  $M$  be a cusped hyperbolic manifold with torus cusps and virtually locally indicable fundamental group, then for every  $\delta > 0$ , every sufficiently deep manifold filling  $\overline{M}$  of  $M$  satisfies

- (a) if  $n \neq \dim(M)/2$ , then  $b_n^{(2)}(\pi_1(\overline{M})) < \delta$ ;
- (b) if  $n = \dim M/2$ , then  $|b_n^{(2)}(\pi_1(\overline{M})) - |\chi(M)|| < \delta$ .

Suppose  $M = \mathbb{H}^n/\Gamma$  is a finite volume cusped hyperbolic manifold. Then  $M$  is called *arithmetic* if  $\Gamma$  is commensurable with a non-uniform arithmetic lattice in  $\mathrm{SO}(n, 1)$ .

**Theorem 7.10.** Let  $M$  be a cusped arithmetic hyperbolic manifold, then there is a finite-sheeted cover  $W$  of  $M$  such that  $W$  has torus cusps and sufficiently deep  $\pi_1(M)$ -equivariant manifold fillings  $\overline{W}$  of  $W$  satisfy

$$b_*^{(2)}(\pi_1(\overline{W})) = [\pi_1(M) : \pi_1(W)] \cdot b_*^{(2)}(\pi_1(M)).$$

*Proof.* Let  $G_0 \triangleleft \pi_1(M)$  be the finite-index normal subgroup provided by Theorem 7.6 (ii), and let  $W$  be the cover of  $M$  with  $\pi_1(W) = G_0$ . Then  $W$  satisfies the requirements.  $\square$

**Corollary 7.11.** For every cusped arithmetic hyperbolic manifold  $M$  of dimension at least four, there exists a sequence of pairwise non-homeomorphic Anderson-type closed Einstein manifolds  $\{W_n\}_{n \geq 1}$  such that each  $W_n$  satisfies the Singer Conjecture and  $\{W_n\}_{n \geq 1}$  converges to a finite-sheeted cover  $W$  of  $M$  under the pointed Gromov–Hausdorff metric with based point any  $y \in W$ .

*Proof.* Let  $W$  be the finite-sheeted cover of  $M$  given by Theorem 7.10, and let  $H_1, \dots, H_m$  be the fundamental groups of the cusps of  $M$ , viewed as subgroups of  $\pi_1(M)$ . For  $i = 1, \dots, m$ , let  $a_i, b_i \in \pi_1(W) \cap H_i$  be two elements that can be extended to a basis of the free abelian group  $\pi_1(W) \cap H_i$ . For  $n \in \mathbb{N}^+$ , let  $x_{i,n} = a_i + n \cdot b_i$ . Then the sequence of elements  $\{x_{i,n}\}_{n \in \mathbb{N}^+}$  satisfies the following:

- (i) Each  $x_{i,n}$  is not a proper power of another element of  $\pi_1(W) \cap H_i$ .
- (ii) For each finite subset  $\mathcal{S} \subset H_i$ , there exists  $m > 0$  such that for all  $n > m$  we have  $\langle x_{i,n} \rangle \cap \mathcal{S} = \emptyset$ .

Let  $\{K_j\}_{j=1}^\ell$  be the peripheral structure on  $\pi_1(W)$  induced by  $\{H_i\}_{i=1}^m$ . For each  $n$ , let  $\{Q_{j,n} \leq K_j\}_{j=1}^\ell$  be the family given by Lemma 2.16 (i) with respect to the family  $\{\langle x_{i,n} \rangle \leq H_i\}_{i=1}^m$ . Then for all  $j$  and  $n$ , we have that  $Q_{j,n} \cong \mathbb{Z}$  and  $K_j/Q_{j,n}$  is torsion-free,



by Lemma 2.16 (ii). Therefore, each of the quotient groups  $\pi_1(W)/\langle\langle \bigcup_{j=1}^{\ell} Q_{j,n} \rangle\rangle_{\pi_1(W)}$  is the fundamental group of a  $\pi_1(M)$ -equivariant manifold filling of  $W$ , which will be denoted by  $W_n$ .

By the main result of [And06] and [FM10, Theorem 2.7], there exists a family of finite subsets  $\{\mathcal{F}_j \subset K_j \setminus \{1\}\}_{j=1}^{\ell}$  such that if a family of subgroups  $\{Q_j \leq K_j\}_{j=1}^{\ell}$  induces a manifold filling  $\overline{W}$  of  $W$  and satisfies that for each  $j$ , we have  $Q_j \cap \mathcal{F}_j = \emptyset$ , then  $\overline{W}$  admits both an Einstein metric and a locally CAT(0) metric. The latter implies that  $\overline{W}$  is aspherical.

By Lemma 2.16 (iv), for  $n$  large enough we have that  $Q_{j,n} \cap \mathcal{F}_j = \emptyset$ . By Theorem 7.10 and Theorem 7.5, for large  $n$ , we have that each  $W_n$  satisfies the Singer Conjecture. By [And06, Proposition 3.9], we may assume that the manifolds  $W_n$  are pairwise non-homeomorphic. By [And06], the sequence  $\{W_n\}_{n \geq 1}$  converges to  $W$  under the pointed Gromov–Hausdorff metric with based point any  $y \in W$ .  $\square$

**7.4. Exotic subgroups of hyperbolic groups.** We present a new construction of exotic subgroups of hyperbolic groups that arise from hyperbolic Dehn filling.

**Theorem 7.12.** *Let  $G$  be the fundamental group of a cusped arithmetic hyperbolic manifold  $M$  of dimension  $n$ . Then there is a hyperbolic Dehn filling quotient  $\overline{G}$  of  $G$  such that*

- (i) *If  $n = 2q \geq 4$ , then there exist a finite-index subgroup  $\overline{G}_1 \leq \overline{G}$  and an epimorphism  $\psi: \overline{G}_1 \rightarrow \mathbb{Z}$  such that  $\ker \psi$  is of type  $FP_{q-1}(\mathbb{Q})$  but not of type  $FP_q(\mathbb{Q})$ .*
- (ii) *If  $n = 2q + 1 \geq 5$ , then there exist a finite-index subgroup  $\overline{G}_1 \leq \overline{G}$  and an epimorphism  $\psi: \overline{G}_1 \rightarrow \mathbb{Z}$  such that  $\ker \psi$  is of type  $FP(\mathbb{Q})$  but not hyperbolic.*

*Proof.* Let  $H_i, i = 1, \dots, \ell$ , be the the fundamental groups of the cusps of  $M$ , viewed as subgroups of  $G$ . Theorem 7.6 (ii) provides us with a finite-index sparse special normal subgroup  $G_0 \triangleleft G$  and a family of finite subsets  $\{\mathcal{F}_i \subset G_0 \cap H_i \setminus \{1\}\}_{i=1}^{\ell}$ .

(i). In this case, for each  $i$ , the group  $G_0 \cap H_i$  is free abelian of rank at least 3. For each  $i$ , use Lemma 4.17 to choose a subgroup  $N_i \leq (G_0 \cap H_i)$  such that  $(G_0 \cap H_i)/N_i \cong \mathbb{Z}$  and  $N_i \cap \mathcal{F}_i = \emptyset$ . Then the quotient  $\overline{G} := G/\langle\langle \bigcup_{i=1}^{\ell} N_i \rangle\rangle_G$  virtually  $FP_{q-1}(\mathbb{Q})$ -fibers, i.e., it has a finite-index subgroup  $\overline{G}_1 \leq \overline{G}$  with an epimorphism  $\psi: \overline{G}_1 \rightarrow \mathbb{Z}$  such that  $\ker \psi$  is of type  $FP_{q-1}(\mathbb{Q})$ . By Theorem 7.6 (ii), the group  $\overline{G}_1$  does not virtually  $FP_q(\mathbb{Q})$ -fiber. So  $\ker \psi$  is not of type  $FP_q(\mathbb{Q})$ .

(ii). By Theorem 7.6 (ii), there exists a finite-index sparse special normal subgroup  $G_0 \triangleleft G$ , with an induced peripheral structure  $\{H_{0j}\}_{j=1}^m$ , and a family of finite subsets  $\{\mathcal{F}_{0j} \subset H_{0j} \setminus \{1\}\}_{j=1}^m$  such that if a  $G$ -equivariant family of subgroups  $\{N_{0j} \leq H_{0j}\}_{j=1}^m$  satisfies, for all  $j$ ,

- $N_{0j} \cap \mathcal{F}_{0j} = \emptyset$ , and
- $H_{0j}/N_{0j}$  is torsion-free,

then

- (a) the group  $\overline{G}_0 := G_0/\langle\langle \bigcup_{j=1}^m N_{0j} \rangle\rangle_{G_0}$  virtually  $FP_{\infty}(\mathbb{Q})$ -fibers.

By Corollary 4.7 and Theorem 4.10 (ii) of [PS24] and Corollary 5.3, there exists a family of finite subsets  $\{\mathcal{F}'_{0j} \subset H_{0j} \setminus \{1\}\}_{j=1}^m$  such that if a family of subgroups  $\{N_{0j}\}_{j=1}^m$  satisfies that  $N_{0j} \cap \mathcal{F}'_{0j} = \emptyset$  for all  $j$ , then the following hold:

- (b)  $\text{cd}(\overline{G}_0) < \infty$ , in particular,  $\overline{G}_0$  is torsion-free

And

$$(48) \quad H^n(\overline{G}_0, \{\overline{H}_{0j}\}_{j=1}^m; \mathbb{Z}/2\mathbb{Z}) \cong H^n(G_0, \{H_{0j}\}_{j=1}^m; \mathbb{Z}/2\mathbb{Z}) \cong \mathbb{Z}/2\mathbb{Z},$$

where, for  $j = 1, \dots, m$ , the group  $\overline{H}_{0j}$  is the natural image of  $H_{0j}$  in  $\overline{G}_0$ , and the last isomorphism follows from Poincaré–Lefschetz duality: as  $G_0$  is sparse special, it is torsion-free, and thus is the fundamental group of a cusped hyperbolic manifold. We will specify our choice of  $\{N_{0j}\}_{j=1}^m$  later.

By applying Lemma 2.16 (iv) to the pair of groups  $G_0 \triangleleft G$ , we obtain a family of finite subsets  $\{\mathcal{F}_{1i} \subset H_i \setminus \{1\}\}_{i=1}^\ell$  with respect to the family  $\{\mathcal{F}_{0j} \cup \mathcal{F}'_{0j}\}_{j=1}^m$ .

By [Osi07, Theorem 1.1], there exists a family of finite subsets  $\{\mathcal{F}_{2i} \subset H_i \setminus \{1\}\}_{i=1}^\ell$  such that if a family of subgroups  $\{N_i \leq H_i\}_{i=1}^\ell$  satisfies that, for all  $i$ ,

- $N_i \cap \mathcal{F}_{2i} = \emptyset$ , and
- $H_i/N_i$  is virtually  $\mathbb{Z}$ ,

then

- (d) the group  $G/\langle\langle \bigcup_{i=1}^\ell N_i \rangle\rangle$  is hyperbolic.

We now specify our choice of  $\{N_i\}_{i=1}^\ell$  and  $\{N_{0j}\}_{j=1}^m$ . Without loss of generality, we may assume that  $n \geq 4$ . Therefore, for each  $i$ , the group  $G_0 \cap H_i$  is free abelian of rank at least 3. For each  $i$ , use Lemma 4.17 to choose a subgroup  $N_i \leq G_0 \cap H_i$  such that  $N_i \cap (\mathcal{F}_{1i} \cup \mathcal{F}_{2i}) = \emptyset$  and  $(G_0 \cap H_i)/N_i \cong \mathbb{Z}$ . Let  $\{N_{0j} \leq H_{0j}\}_{j=1}^m$  be the  $G$ -equivariant family of subgroups given by Lemma 2.16 (i) with respect to the family  $\{N_i\}_{i=1}^\ell$ . Then  $N_{0j} \cap (\mathcal{F}_{0j} \cup \mathcal{F}'_{0j}) = \emptyset$  for all  $j$ . Therefore, (a), (b), and (d) hold. Let

$$\overline{G} := G/\langle\langle \bigcup_{i=1}^k N_i \rangle\rangle_G, \quad \overline{G}_0 := G_0/\langle\langle \bigcup_{j=1}^\ell N_{0j} \rangle\rangle_{G_0}.$$

As  $\overline{G}$  virtually  $FP_\infty(\mathbb{Q})$ -fibers, so does  $\overline{G}_0$ . So there is a finite-index subgroup  $\overline{G}_1 \leq \overline{G}_0$  and an epimorphism  $\psi: \overline{G}_1 \rightarrow \mathbb{Z}$  such that  $\ker \psi$  is of type  $FP_\infty(\mathbb{Q})$ . By (b) and  $\ker \psi < \overline{G}_0$ , we have  $\text{cd}(\ker \psi) < \infty$  and thus  $\ker \psi$  is of type  $FP(\mathbb{Q})$ . Assume, for the sake of contradiction, that  $\ker \psi$  is hyperbolic. Since  $\overline{G}_0$  is torsion-free by (b), so is its subgroup  $\overline{G}_1$ . It follows from [Kud23, Proposition 3.3] that  $\text{cd}(\ker \psi) \leq 2$ , and thus  $\text{cd}(\overline{G}_1) \leq 3$ . On the other hand, the fact that  $\overline{H}_{0j} \cong \mathbb{Z}$  for all  $j$  and (48) imply that  $\text{cd}(\overline{G}_0) \geq 5$  and thus  $\text{cd}(\overline{G}_1) \geq 5$ , which is a contradiction. So  $\ker \psi$  is not hyperbolic.  $\square$

**7.5. Deficiency.** Deficiency is an important invariant of a finitely presented group. A positive deficiency suggests that the group may be large (i.e. a finite index subgroup admits a homomorphism onto a non-cyclic free group) or possess a certain degree of freedom [BP78], which often manifests in its homological or geometric properties. In contrast, groups with

non-positive deficiency tend to have a more rigid and constrained structure, such as many lattices in simple Lie groups [Lot99].

**Corollary 7.13.** *Let  $G$  be a virtually locally indicable group of type  $F_3$  that has a finite family of hyperbolically embedded subgroups  $\{H_i\}_{i=1}^n$ . Then for sufficiently deep families of normal subgroups  $\{N_i \triangleleft H_i\}_{i=1}^n$  such that  $H_i/N_i$  is of type  $F_3$  for all  $i$ , the group  $G/\langle\langle \bigcup_{i=1}^n N_i \rangle\rangle$  is finitely presented and we have*

$$\begin{aligned} & \text{defi}(G/\langle\langle \bigcup_{i=1}^n N_i \rangle\rangle) \\ & \leq b_1^{(2)}(G) - b_2^{(2)}(G) + \sum_{i=1}^n b_1^{(2)}(H_i/N_i) + \sum_{i=1}^n b_2^{(2)}(H) + 1 \\ & \leq b_1^{(2)}(G) - b_2^{(2)}(G) + \sum_{i=1}^n b_1^{(2)}(H_i) + \sum_{i=1}^n b_2^{(2)}(H_i) + 1. \end{aligned}$$

*Proof.* For simplicity, denote

$$\begin{aligned} s & := b_1^{(2)}(G) - b_2^{(2)}(G) + \sum_{i=1}^n b_1^{(2)}(H_i/N_i) + \sum_{i=1}^n b_2^{(2)}(H) + 1 \\ t & := b_1^{(2)}(G) - b_2^{(2)}(G) + \sum_{i=1}^n b_1^{(2)}(H_i) + \sum_{i=1}^n b_2^{(2)}(H_i) + 1. \end{aligned}$$

Choose  $\delta > 0$  such that each of the open intervals  $(s, s + 2\delta)$  and  $(t, t + (n + 2)\delta)$  does not contain any integer.

By Corollary 5.3 and Theorem 5.4, there exists a family of finite subsets  $\{\mathcal{F}_{1,i} \subset H_i \setminus \{1\}\}_{i=1}^n$  such that if a family of normal subgroups  $N_i \triangleleft H_i$  satisfies that, for all  $i$ ,  $N_i \cap \mathcal{F}_{1,i} = \emptyset$  and  $H_i/N_i$  is of type  $F_3$ , then  $G/\langle\langle \bigcup_{i=1}^n N_i \rangle\rangle$  is of type  $F_3$  and

$$(49) \quad b_2^{(2)}(G/\langle\langle \bigcup_{i=1}^n N_i \rangle\rangle) > b_2^{(2)}(G) - \sum_{i=1}^n b_1^{(2)}(H_i/N_i) - \sum_{i=1}^n b_2^{(2)}(H) - \delta.$$

For each  $i$ , by Corollary 3.7, there exists a finite subset  $\mathcal{F}_{2,i} \subset H_i \setminus \{1\}$  such that if a normal subgroup  $N_i \triangleleft H_i$  satisfies  $N_i \cap \mathcal{F}_{2,i} = \emptyset$ , then

$$(50) \quad b_1^{(2)}(H_i/N_i) < b_1^{(2)}(H_i) + \delta.$$

By Corollary 3.8, there exists a family of finite subsets  $\{\mathcal{F}_{3,i} \subset H_i \setminus \{1\}\}_{i=1}^n$  such that if a family of normal subgroups  $\{N_i \triangleleft H_i\}_{i=1}^n$  satisfies that, for all  $i$ ,  $N_i \cap \mathcal{F}_{3,i} = \emptyset$ , then

$$(51) \quad b_1^{(2)}(G/\langle\langle \bigcup_{i=1}^n N_i \rangle\rangle) < b_1^{(2)}(G) + \delta.$$

Let  $\{N_i \triangleleft H_i\}$  be a family of normal subgroups such that, for all  $i$ ,

$$N_i \cap (\mathcal{F}_{1,i} \cup \mathcal{F}_{2,i} \cup \mathcal{F}_{3,i}) = \emptyset$$

and  $H_i/N_i$  is of type  $F_3$ . Then from [Lüc02, Lemma 7.16], we conclude that

$$\begin{aligned} \text{defi}(G/\langle\langle \bigcup_{i=1}^n N_i \rangle\rangle) &\leq b_1^{(2)}(G/\langle\langle \bigcup_{i=1}^n N_i \rangle\rangle) - b_2^{(2)}(G/\langle\langle \bigcup_{i=1}^n N_i \rangle\rangle) + 1 \\ &< s + 2\delta && \text{by (49) and (51)} \\ &< t + (n + 2)\delta && \text{by (50)} \end{aligned}$$

The desired result follows as  $\text{defi}(G/\langle\langle \bigcup_{i=1}^n N_i \rangle\rangle)$  is an integer. □

**Corollary 7.14.** *Let  $G$  be a non-uniform arithmetic lattice in  $SO(n, 1)$ ,  $n \geq 3$ . Denote by  $\{H_i\}_{i=1}^k$  the collection of peripheral subgroups. Then for every family of sufficiently deep normal subgroups  $\{N_i \triangleleft H_i\}_{i=1}^k$ , we have*

$$\text{defi}(\overline{G}) \leq 1.$$

In addition, if  $n = 4$ , then

$$\text{defi}(\overline{G}) \leq 1 - \frac{3}{4\pi^2} \text{vol}(\mathbb{H}^4/G).$$

*Proof.* By combining Theorem 7.4 and Proposition 4.12, we see that  $G$  is virtually locally indicable. Also, Theorem 7.5 implies that  $b_1^{(2)}(G) = 0$ . Since the peripheral subgroups of  $G$  are virtually abelian, they have vanishing  $L^2$ -Betti numbers in positive degrees. The first claim now follows from applying Corollary 7.13.

When  $n = 4$ , one gets a sharper bound as follows:

$$\begin{aligned} \text{defi}(\overline{G}) &\leq 1 - b_2^{(2)}(G) && \text{by Corollary 7.13,} \\ &\leq 1 - \chi(G) && \text{by Theorem 7.5,} \\ &\leq 1 - \frac{3}{4\pi^2} \text{vol}(\mathbb{H}^4/G) && \text{by (2.13) and (2.14) of [Lot99].} \end{aligned}$$

□

## REFERENCES

- [ACG18] Y. Antolín, R. Coulon, and G. Gandini. Farrell-Jones via Dehn fillings. *J. Topol. Anal.*, 10(4):873–895, 2018.
- [AFW15] M. Aschenbrenner, S. Friedl, and H. Wilton. *3-manifold groups*. EMS Series of Lectures in Mathematics. European Mathematical Society (EMS), Zürich, 2015.
- [AGM16] I. Agol, D. Groves, and J. Manning. An alternate proof of Wise’s malnormal special quotient theorem. *Forum Math. Pi*, 4:e1, 54, 2016.
- [Ago08] I. Agol. Criteria for virtual fibering. *J. Topol.*, 1(2):269–284, 2008.
- [Ago13] I. Agol. The virtual Haken conjecture. *Doc. Math.*, 18:1045–1087, 2013. With an appendix by I. Agol, D. Groves, and J. Manning.
- [And06] M. Anderson. Dehn filling and Einstein metrics in higher dimensions. *J. Differential Geom.*, 73(2):219–261, 2006.
- [BE78] R. Bieri and B. Eckmann. Relative homology and Poincaré duality for group pairs. *J. Pure Appl. Algebra*, 13(3):277–319, 1978.
- [Bie81] Robert Bieri. *Homological dimension of discrete groups*. Queen Mary College Mathematics Notes. Queen Mary College, Department of Pure Mathematics, London, second edition, 1981.
- [Bow12] B. Bowditch. Relatively hyperbolic groups. *Internat. J. Algebra Comput.*, 22(3):1250016, 66, 2012.

- [BP78] Benjamin Baumslag and Stephen J. Pride. Groups with two more generators than relators. *J. London Math. Soc. (2)*, 17(3):425–426, 1978.
- [Bro94] K. Brown. *Cohomology of groups*, volume 87 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1994. Corrected reprint of the 1982 original.
- [BW12] N. Bergeron and D. Wise. A boundary criterion for cubulation. *Amer. J. Math.*, 134(3):843–859, 2012.
- [CIOS23] I. Chifan, A. Ioana, D. Osin, and B. Sun. Wreath-like products of groups and their von Neumann algebras I:  $W^*$ -superrigidity. *Ann. of Math. (2)*, 198(3):1261–1303, 2023.
- [CL63] D. Cohen and R. Lyndon. Free bases for normal subgroups of free groups. *Trans. Amer. Math. Soc.*, 108:526–537, 1963.
- [Dah03] F. Dahmani. Combination of convergence groups. *Geom. Topol.*, 7:933–963, 2003.
- [DG18] F. Dahmani and V. Guirardel. Recognizing a relatively hyperbolic group by its Dehn fillings. *Duke Math. J.*, 167(12):2189–2241, 2018.
- [DGO17] F. Dahmani, V. Guirardel, and D. Osin. Hyperbolically embedded subgroups and rotating families in groups acting on hyperbolic spaces. *Mem. Amer. Math. Soc.*, 245(1156):v+152, 2017.
- [DK92] G. Duchamp and D. Krob. The lower central series of the free partially commutative group. In *Semigroup Forum*, volume 45, pages 385–394. Springer, 1992.
- [Fis21] S. Fisher. Improved algebraic fibrings. *arXiv preprint arXiv:2112.00397*, 2021.
- [FM10] K. Fujiwara and J. F. Manning.  $CAT(0)$  and  $CAT(-1)$  fillings of hyperbolic manifolds. *J. Differential Geom.*, 85(2):229–269, 2010.
- [GM22] D. Groves and J. F. Manning. Specializing cubulated relatively hyperbolic groups. *J. Topol.*, 15(2):398–442, 2022.
- [GMS19] D. Groves, J. F. Manning, and A. Sisto. Boundaries of Dehn fillings. *Geom. Topol.*, 23(6):2929–3002, 2019.
- [Gru57] K. Gruenberg. Residual properties of infinite soluble groups. *Proc. London Math. Soc. (3)*, 7:29–62, 1957.
- [Hat02] A. Hatcher. *Algebraic Topology*. Cambridge University Press, Cambridge, 2002.
- [Hru10] C. Hruska. Relative hyperbolicity and relative quasiconvexity for countable groups. *Algebr. Geom. Topol.*, 10(3):1807–1856, 2010.
- [HW08] F. Haglund and D. Wise. Special cube complexes. *Geom. Funct. Anal.*, 17(5):1551–1620, 2008.
- [IMM23] Giovanni Italiano, Bruno Martelli, and Matteo Migliorini. Hyperbolic 5-manifolds that fiber over  $S^1$ . *Invent. Math.*, 231(1):1–38, 2023.
- [IMP24] Claudio Llosa Isenrich, Bruno Martelli, and Pierre Py. Hyperbolic groups containing subgroups of type  $\mathcal{F}_3$  not  $\mathcal{F}_4$ . *J. Differential Geom.*, 127(3):1121–1147, 2024.
- [JX00] J. Jost and Y. Xin. Vanishing theorems for  $L^2$ -cohomology groups. *J. Reine Angew. Math.*, 525:95–112, 2000.
- [JZLA20] A. Jaikin-Zapirain and D. López-Álvarez. The strong Atiyah and Lück approximation conjectures for one-relator groups. *Math. Ann.*, 376(3-4):1741–1793, 2020.
- [Kie20] D. Kielak. Residually finite rationally solvable groups and virtual fibring. *J. Amer. Math. Soc.*, 33(2):451–486, 2020.
- [Kud23] M. Kudlinska. An infinite family of exotic subgroups of hyperbolic groups. *arXiv:2303.11218*, 2023.
- [KZ01] Ruth Kellerhals and Thomas Zehrt. The Gauss-Bonnet formula for hyperbolic manifolds of finite volume. *Geom. Dedicata*, 84(1-3):49–62, 2001.
- [L94] W. Lück. Approximating  $L^2$ -invariants by their finite-dimensional analogues. *Geom. Funct. Anal.*, 4(4):455–481, 1994.
- [L05] Wolfgang Lück. Survey on classifying spaces for families of subgroups. In *Infinite groups: geometric, combinatorial and dynamical aspects*, volume 248 of *Progr. Math.*, pages 269–322. Birkhäuser, Basel, 2005.
- [LIMP24] Claudio Llosa Isenrich, Bruno Martelli, and Pierre Py. Hyperbolic groups containing subgroups of type  $\mathcal{F}_3$  not  $\mathcal{F}_4$ . *J. Differential Geom.*, 127(3):1121–1147, 2024.
- [LIP24] Claudio Llosa Isenrich and Pierre Py. Subgroups of hyperbolic groups, finiteness properties and complex hyperbolic lattices. *Invent. Math.*, 235(1):233–254, 2024.

- [Lot99] John Lott. Deficiencies of lattice subgroups of Lie groups. *Bull. London Math. Soc.*, 31(2):191–195, 1999.
- [Lub83] Alexander Lubotzky. Group presentation,  $p$ -adic analytic groups and lattices in  $\mathrm{SL}_2(\mathbf{C})$ . *Ann. of Math. (2)*, 118(1):115–130, 1983.
- [Lüc02] W. Lück. *L<sup>2</sup>-invariants: theory and applications to geometry and K-theory*, volume 44 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 2002.
- [Osi06] D. Osin. Relatively hyperbolic groups: intrinsic geometry, algebraic properties, and algorithmic problems. *Mem. Amer. Math. Soc.*, 179(843):vi+100, 2006.
- [Osi07] D. Osin. Peripheral fillings of relatively hyperbolic groups. *Invent. Math.*, 167(2):295–326, 2007.
- [Pat23] H. Patil. Finiteness properties and relatively hyperbolic groups. *arXiv preprint arXiv:2308.08353*, 2023.
- [PS24] N. Petrosyan and B. Sun. Cohomology of group theoretic Dehn fillings II. *Adv. Math.*, 437:Paper No. 109412, 56, 2024.
- [PT11] J. Peterson and A. Thom. Group cocycles and the ring of affiliated operators. *Invent. Math.*, 185(3):561–592, 2011.
- [Rhe73] A. Rhemtulla. Residually  $F_p$ -groups, for many primes  $p$ , are orderable. *Proc. Amer. Math. Soc.*, 41:31–33, 1973.
- [RR02] A. Rhemtulla and D. Rolfsen. Local indicability in ordered groups: braids and elementary amenable groups. *Proc. Amer. Math. Soc.*, 130(9):2569–2577, 2002.
- [Sch15] K. Schreve. *The L<sup>2</sup>-cohomology of Discrete Groups*. ProQuest LLC, Ann Arbor, MI, 2015. Thesis (Ph.D.)—The University of Wisconsin - Milwaukee.
- [Sun20] B. Sun. Cohomology of group theoretic Dehn fillings I: Cohen-Lyndon type theorems. *J. Algebra*, 542:277–307, 2020.
- [Thu82] W. Thurston. Three-dimensional manifolds, Kleinian groups and hyperbolic geometry. *Bull. Amer. Math. Soc. (N.S.)*, 6(3):357–381, 1982.
- [Wis21] D. T. Wise. *The structure of groups with a quasiconvex hierarchy*, volume 209 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, [2021] ©2021.
- [WW17] Daniel T. Wise and Daniel J. Woodhouse. A cubical flat torus theorem and the bounded packing property. *Israel J. Math.*, 217(1):263–281, 2017.

SCHOOL OF MATHEMATICS, UNIVERSITY OF SOUTHAMPTON, SOUTHAMPTON SO17 1BJ, UK

*Email address:* `n.petrosyan@soton.ac.uk`

DEPARTMENT OF MATHEMATICS, MICHIGAN STATE UNIVERSITY, EAST LANSING, MI, 48824, USA

*Email address:* `sunbin1@msu.edu`