arXiv:2412.16111v1 [q-bio.NC] 20 Dec 2024

How connection probability shapes fluctuations of neural population dynamics

Nils E. Greven,^{1,2} Jonas Ranft,^{3, *} and Tilo Schwalger^{1,2,†}

¹Institute of Mathematics, Technical University Berlin, 10623 Berlin, Germany.

²Bernstein Center for Computational Neuroscience Berlin, 10115 Berlin, Germany.

³Institut de Biologie de l'ENS, Ecole normale supérieure, PSL University, CNRS, Paris.

(Dated: December 23, 2024)

Mean-field models of neuronal populations in the brain have proven extremely useful to understand network dynamics and response to stimuli, but these models generally lack a faithful description of the fluctuations in the biologically relevant case of finite network size and connection probabilities p < 1 (non-full connectivity). To gain insight into the different fluctuation mechanisms underlying the neural variability of populations of spiking neurons, we derive and analyze a stochastic mean-field model for finite-size networks of Poisson neurons with random, non-full connectivity, external noise and disordered mean inputs. We treat the quenched disorder of the connectivity by an annealed approximation that enables a reduction to a low-dimensional closed system of coupled Langevin equations for the mean and variance of the neuronal membrane potentials as well as a variable capturing finite-size fluctuations arising specifically in the case p < 1. Comparing to microscopic simulations, we find that the mesoscopic model describes the fluctuations and nonlinearities well and outperforms previous mesoscopic models that neglected the recurrent noise effect caused by the non-full connectivity. This effect can be analytically understood by a softening of the effective nonlinearity and the multiplicative character of finitesize spiking noise. The mesoscopic theory shows that quenched disorder can stabilize the asynchronous state, and it correctly predicts large quantitative and non-trivial qualitative effects of connection probability on the variance of the population firing rate and its dependence on stimulus strength. Our theory thus elucidates how disordered connectivity shapes nonlinear dynamics and fluctuations of neural populations at the mesoscopic scale and showcases a useful mean-field method to treat non-full connectivity in finite-size, spiking neural networks.

I. INTRODUCTION

Biological systems often exhibit significant fluctuations and variability in their dynamics, in contexts ranging from gene expression to neural activity in the brain. The possible roles of fluctuations for the dynamics and functions of such systems have thus received increasing attention [1–7]. An important tool to understand the collective dynamics of biological systems consisting of many interacting units are low-dimensional descriptions of the effective dynamics at the population level. Taking fluctuations properly into account can be necessary to establish a correct understanding of the system and can provide crucial insight into the origins of observed biological variability. At the population level, fluctuations may then be accounted for by stochastic low-dimensional models. In the context of living systems, examples range from Langevin equations for gene regulation [8] to coupled molecular motors [9–11] to ecology [12].

In neuroscience, neural-mass models (also called firingrate or neural-population models) such as the Wilson-Cowan model [13] have been highly successful in describing various phenomena in cortical circuits such as oscillations [14], multistability [15], and nonlinear reponse properties [16]. Stochastic versions of neural-mass models have been critical for understanding fluctuating neural population dynamics such as metastability [15, 17, 18], stochastic oscillations [19], and response properties of cortical variability [20, 21]. While neural-mass models and their stochastic variants stand out due to their mathematical tractability, they are often phenomenological, lacking a clear link to an underlying network of spiking neurons at the microscopic level.

Neural population models that maintain a link to biophysical properties can be obtained through mean-field modeling [22–28], where the macroscopic or mesoscopic dynamics is analytically derived from a microscopic model. The resulting mean-field models are amenable to mathematical analysis [29-32] or may enable efficient and accurate simulations of large spiking neural networks [33-36]. In current meanfield models, however, the treatment of fluctuations is typically based on the (implicit) assumptions of either infinitely large network size or homogeneous, full (all-to-all) connectivity. These assumptions are often not applicable to biologically realistic recurrent neural networks. For example, in rat barrel cortex, the number of neurons per cell type in a given layer of a column has been counted to be on the order of $N \sim 100$ to $N \sim 1000$ neurons per cell type [37] – a population size for which finite-size fluctuations cannot be neglected. Furthermore, the local connection probabilities p in mouse visual cortex vary on a wide range ($0 \leq p \leq 0.7$) with low average $(p \sim 0.1)$ and larger connection probabilities $(p \sim 0.5)$ for similarly tuned neurons [38].

Both finite network sizes and connection probabilities < 1 cause intrinsically generated (endogenous) fluctuations of the recurrent, synaptic input currents, and may thus strongly affect the nonlinear population dynamics. However, their distinct and combined effects are still poorly understood. Moreover, these intrinsic fluctuations act as multiplicative noise whose strength increases with population firing rates. On the other hand, neural variability in the cortex often decreases upon stimulation [39] – a phenomenon that could be explained, among others, with a model, where variability is gen-

^{*} Contact author: jonas.ranft@ens.psl.eu

[†] Contact author: tilo.schwalger@bccn-berlin.de

erated by an external, additive noise [21]. To clarify the role of intrinsic fluctuations for such cortical phenomena under biologically realistic neuron numbers and connection probabilities requires a mean-field model that accounts for these different origins of fluctuations.

A mesoscopic theory for finite-size spiking neural networks providing accurate population models with fluctuations has recently been developed for generalized integrate-and-fire neurons with escape noise [32, 35]. This theory suggests a multiscale modeling framework for efficient forward [35] and inverse [40, 41] modeling of cortical circuits based on realistic neuron dynamics [42, 43]. However, this and other meanfield theories [32, 35, 44-48] for finite-size neural networks are based on homogeneous, fully connected networks (p = 1, "global coupling"). Even though the full connectivity may be regarded as an effective, mean-connectivity approximation of a non-fully connected network [35, 49, 50], the validity of such approximation remains unclear. In particular, in a homogeneous fully connected network, the fluctuations caused by a finite network size $N < \infty$ are common to all neurons because neurons receive identical recurrent inputs when neurons are connected all-to-all. On the population level, these "coherent" fluctuations lead to stochastic population equations with finite-size noise of order $1/\sqrt{N}$. Clearly, in random networks with connection probability p < 1, different neurons no longer receive identical synaptic inputs, and therefore the implicit assumption of perfectly coherent noise made in the effective, mean-connectivity approximation cannot hold fully true.

A mean-field description of non-fully connected, random networks (p < 1) is difficult because it requires the treatment of fluctuations resulting from a "quenched" (i.e. temporally constant) random synaptic connectivity. A classical approach to deal with random networks is to consider the macroscopic limit $N \to \infty$. Depending on the concrete limiting procedure and network architecture, the recurrent fluctuations seen by different neurons either vanish [22, 28, 30, 31] or become perfectly independent ("incoherent") in this limit [51-55] unless correlations are induced externally. A famous example is the mean-field theory for sparsely connected networks by Brunel and co-workers [29, 51]. In this theory, the neural population dynamics is deterministic and given by a nonlinear Fokker-Planck equation. The effect of incoherent fluctuations of the recurrent synaptic input appears as an additional contribution to the diffusion coefficient in this equation. Although this approach neglects temporal correlations of the spike trains caused by the quenched disorder, the nonlinear Fokker-Planck equation has been successful to capture the influence of intrinsic fluctuations on the nonlinear population dynamics (for a treatment of temporal correlations, see e.g. [56]). Using this theory, intrinsic fluctuations have been shown to crucially shape nonlinear response properties [57], network oscillations [29, 58] and multistability [59].

Again, even though mean-field theories for $N \to \infty$ have been used as simple mean-field descriptions for real neural networks of finite size (see e.g. [60]), they cannot describe finite-size fluctuations at the population level. Furthermore, finite-size networks are necessarily outside the sparse limit (i.e. p > 0) which entails a non-vanishing probability of shared recurrent inputs among neurons. Therefore the assumption of perfectly incoherent noise in the mean-field theory cannot hold fully true for $N < \infty$. While finite-size extensions of the mean-field theory for sparsely connected networks have been proposed in [29, 51, 61], low-dimensional neural population models that reveal and take into account the distinct effects of coherent and incoherent fluctuations are currently lacking.

In this paper, we derive and analyze a simple bottomup mean-field model that describes incoherent and coherent recurrent fluctuations for arbitrary connection probabilities $p \in [0,1]$ in a finite-size network. To this end, we study a network of Poisson neurons with "annealed" random connectivity [29, 62] as an approximation to a corresponding network with "quenched" random connectivity. The annealed approximation neglects temporal correlations of incoherent fluctuations but captures the effect of non-full connectivity and finite network size surprisingly well. Importantly, the annealed approximation enables a simple mean-field description of the fluctuations in form of a three-dimensional Langevin equation. We compare this model with a naive mean-field model in which the incoherent fluctuations due to random connectivity are neglected and which precisely corresponds to the ad hoc mean-field approximation used in previous models [35, 49, 50]. While the focus of the paper is the Langevin dynamics for a homogeneous population with quenched or annealed random connectivity and common external noise, the theoretical framework can be extended to heterogeneous populations, where the resting potentials or external currents of the neurons exhibit quenched Gaussian disorder. In summary, we present a simple theoretical framework for the effects of various sources of noise and quenched disorder on the neural population dynamics at the mesoscopic scale $(1 \ll N < \infty)$.

The paper is organized as follows. We introduce the microscopic network model with quenched random connectivity and its annealed approximation in Sec. II. By reducing the annealed network to a mean-connectivity network with incoherent dynamical noise, we derive in Sec. III a mesoscopic population model as a system of three coupled stochastic differential equations. In Sec. IV, we then analyze the mesoscopic model with respect to fixed-point solutions and their stability, the linear response to dynamic stimuli and second-order statistics. In particular, we compare the obtained analytical results to the corresponding statistics of the quenched and annealed microscopic models using extensive simulations. We conclude and discuss our results in Sec. V. The extension to quenched Gaussian disorder as well as longer, detailed calculations are provided in the Appendix.

II. MICROSCOPIC MODELS

A. Quenched network model

At the microscopic level, we consider a random network of N interacting Poisson neurons (also called nonlinear Hawkes processes). In this model, neurons fire spikes stochastically

with conditional intensity $\lambda_i(t) = \phi(h_i(t^-))$, i = 1, ..., N, where $h_i(t)$ is the input potential of neuron i and ϕ is a nonnegative hazard function. The dynamics of the input potentials is given by a system of coupled first-order equations with delay,

$$\tau \frac{\mathrm{d}h_i}{\mathrm{d}t} = -h_i + \mu(t) + \frac{w}{C} \sum_{j=1}^N a_{ij} \dot{Z}_j(t-d), \qquad (1)$$

 $i = 1, \ldots, N$. Here, τ is the time constant of the lowpass filter dynamics, $\mu(t)$ is an external drive and $\dot{Z}_i(t) = \sum_l \delta(t - t_{i,l})$ is the spike train of neuron i with $\{t_{i,l}\}$ being the individual spike times. The presence or absence of synaptic connections between neurons are described by the random adjacency matrix $[a_{ij}]$ with fixed in-degree $C = \sum_{j=1}^{N} a_{ij}$ and elements $a_{ij} \in \{0, 1\}$ that are marginally Bernoulli distributed with mean $\langle a_{ij} \rangle = p = C/N$ representing the connection probability. This random connectivity can be constructed by choosing for each neuron C presynaptic neurons randomly, and the resulting connectivity is fixed ("quenched") in time. Furthermore, d is the synaptic delay and w = CJ is the total coupling strength with J being the efficacy of a single synapse (i.e. J/τ is the jump size in millivolt of the postsynaptic potential caused by a single presynaptic spike). Since we focus on inhibitory networks in this paper, we assume that J < 0.

As a concrete hazard function, we choose here a sigmoidal function in the form of an error function [22]:

$$\phi(h) = r_{\rm m} \Phi(\beta(h - \vartheta)), \qquad (2)$$

where $\Phi(x) = \left[1 + \operatorname{erf}\left(x/\sqrt{2}\right)\right]/2$. Although not essential for the general theory, this choice will allow us to analytically calculate the first and second moments of $\phi(h_i)$. The parameter β determines the steepness around the inflection point. Without loss of generality, we choose the position of the inflection point ϑ to be zero because we can always measure voltages with respect to the potential ϑ . The sigmoidal shape with an upper limit $r_{\rm m}$ prevents the Poisson neurons to fire with arbitrarily high rates. While the concave behavior and saturation of the sigmoidal hazard function for h > 0 is technically useful, we are in the following mainly interested in dynamical regimes operating in the convex part of the hazard function $\phi(h)$, i.e. corresponding to mean inputs below the inflection point, h < 0. This is because biologically realistic hazard functions of cortical neurons are typically convex [63, 64]. As default parameters of the model, we choose $\tau = 20$ ms, $r_{\rm m} = 100$ Hz and d = 0 ms unless otherwise noted.

The aim of this paper is to derive a mesoscopic population model that generates population activities that statistically match the population activities obtained from a microscopic network simulation. We define the population activity $A_N(t, \Delta t)$ with respect to a time discretization with time step Δt as the total number of spikes per neuron and time step:

$$A_N(t;\Delta t) = \frac{\Delta Z(t)}{N\Delta t}.$$
(3)



FIG. 1. Response of the population rate variance to a change of the mean stimulus strength in the original, quenched random network (a) and in a corresponding fully-connected, mean-connectivity network that only accounts for the mean synaptic weight (b). Top: Noisy external stimulus $\mu(t)$ whose mean increases at t = 0.08 s in a steplike fashion from $\mu_0 = 28$ mV to $\mu_0 = 50$ mV; the noise strength $\sigma_{\rm ext}^2 = 1 \,\,{\rm mV}^2$ is kept fixed. (a) Schematic of a quenched network with fixed adjacency matrix a_{ij} (left) and the trial-averaged variance of the population firing rate as a function of time (right) for different network sizes (green line: $N = 5 \cdot 10^4$, blue line: $N = 10^3$, in both cases C = 100). (b) Same for a mean-connectivity network, Eq. (5), as an ad hoc approximation of the quenched network, where the quenched connectivity a_{ij} is replaced by the average connectivity $\langle a_{ij} \rangle$. Note the different scaling of the y-axis in a) and b). Thus, while the mean stationary population rates after the step (as indicated in the right panels) are equal in both models, the variances of the population rate strongly differ.

Here, $\Delta Z(t)$ denotes the total number of spikes in the time interval $(t, t + \Delta t]$.

B. Mean-connectivity network

A major obstacle in deriving population models analytically is the quenched randomness of the connectivity $J_{ij} := Ja_{ij}$. A standard mean-field approach to tackle this problem is to neglect fluctuations in the recurrent synaptic input by scaling the synaptic weights inversely proportional to the network size, $J_{ij} = w_{ij}/N$, where the coupling strength w_{ij} is assumed to be of order 1. Then, in the limit $N \to \infty$, the synaptic input $N^{-1} \sum_j w_{ij} \dot{Z}_j$ converges to the deterministic input $\langle w_{ij} \rangle \langle \dot{Z}_j \rangle$ [27, 28], where $\langle \cdot \rangle$ denotes the average over both quenched disorder and Poisson noise. In our case, this corresponds to a network where the quenched connectivity w_{ij} (Fig. 1a, left) is replaced by the mean connectivity $\langle w_{ij} \rangle = NJ \langle a_{ij} \rangle = w$ (Fig. 1b, left). Importantly, the mean connectivity effectively yields a fully connected network with homogeneous coupling w. In the limit $N \to \infty$, the synaptic input is thus not affected by the variance

$$\sigma_w^2 = w^2 (1 - p)/p$$
 (4)

of the disordered couplings w_{ij} but only by the mean $\langle w_{ij} \rangle$.

The replacement of all elements of the adjacency matrix a_{ij} with their mean p has been used ad hoc also for finite network size $N < \infty$ [35, 49, 50]. However, this ad hoc approach is no longer exact for finite N but must be regarded as an approximation that retains the mean synaptic connectivity $\langle J_{ij} \rangle$ but ignores all higher-order cumulants of the random variables J_{ij} . In the following, we will refer to this effective, fully connected network as the *mean-connectivity network*, whereas the original model, Eq. (1), will be referred to as the *quenched network*. Replacing the connection strengths w_{ij} with their means w, we obtain from Eq. (1) the dynamics of the mean-connectivity network:

$$\tau dh_i(t) = \left[-h_i + \mu(t)\right] dt + \frac{w}{N} \sum_{j=1}^N dZ_j(t-d), \quad (5)$$

where $dZ_j(t) \sim \text{Pois}(\phi(h_j(t))dt)$ are the conditionally independent Poisson increments of the spike count of neuron j. The mean-connectivity network of Poisson neurons, Eq. (5), is a microscopic model that admits an exact reduction to a mesoscopic model for the total spike count $dZ(t) = \sum_{i=1}^{N} dZ_i(t)$. To see this, we note that the sum of independent Poisson numbers in Eq. (5) is a Poisson number with mean Nr(t-d)dt, where

$$r(t) = \frac{1}{N} \sum_{i=1}^{N} \phi(h_i(t)).$$
 (6)

is the stochastic population rate. Furthermore, because all neurons are driven by the same synaptic input, their input potentials $h_i(t)$ coincide (or converge after an initial transient). The common input potential $\bar{h}(t)$ thus obeys the mesoscopic dynamics

$$\tau d\bar{h}(t) = \left[-\bar{h} + \mu(t)\right] dt + \frac{w}{N} dZ(t-d), \tag{7}$$

where now $dZ(t) \sim \text{Pois}(N\phi(\bar{h}(t))dt)$ describes the increment of the total spike count of the population. Equation (7) describes the mesoscopic dynamics as it only involves the population activity dZ(t)/(Ndt).

Simulations of the quenched and the mean-connectivity network yield similar results for the stationary mean firing rates and mean input potentials. However, the meanconnectivity network may differ drastically compared to the quenched network when it comes to the non-stationary response (linear response function) and the fluctuations (second-order statistics). To illustrate the failure of the meanconnectivity approximation, let us consider the variance of the stochastic population rate r(t). This variance represents the part of the mesoscopic neural variability (variance of the population activity $A_N(t, \Delta t)$) that is caused by the rate variability rather than the Poisson spiking noise. It can be measured from population activity data (see Appendix, Sec. B), or directly computed from a simulated time series of r(t) in our model. A non-vanishing variance of the population rate can occur for two reasons in our model: first, because of intrinsic finite-size fluctuations, and second, because of externally injected common noise. Here, we model the common external noise as a Gaussian white noise (more precisely, $\mu(t) = \bar{\mu}(t) + \sqrt{\tau \sigma_{\text{ext}}^2} \hat{\zeta}(t)$, where $\bar{\mu}(t)$ is the mean stimulus and $\hat{\zeta}(t)$ is a standard Gaussian white noise process).

Interestingly, simulations of the mean-connectivity network (Eq. (5) or (7)) strongly overestimate the population-rate variance of the quenched network by more than one order of magnitude (Fig. 1, note the different scale of the y axes). In addition to this large quantitative difference, we also observe a qualitatively different response to stimulus changes. When the mean stimulus strength is increased from 28 mV to 50 mV, the variance of the population rate exhibits a small but significant decrease in the quenched network (Fig. 1a, right). This suppression of variability is in marked contrast to the prediction by the mean-connectivity network, which exhibits instead an increase in the variance (Fig. 1b, right). We stress that the marked differences concern the second-order statistics but not the first-order statistics: The stationary mean population rates $\langle r \rangle$ are the same in both networks as indicated in Fig. 1a,b.

How can we understand non-stationary responses and neural variability of mesoscopic variables theoretically? In this paper, we present a "second-order" mesoscopic mean-field theory that also captures second-order statistics and nonstationary responses quantitatively, and explains non-trivial, qualitative phenomena such as the suppression of variability observed in Fig. 1a. In particular, the explanation of the failure of the mean-connectivity approximation with respect to the stationary variances in Fig. 1b and its resolution by the "second-order" mean-field theory will be presented towards the end of the paper in Sec. IV E.

C. Annealed network model

The quenched connectivity causes heterogeneity of neural firing with spatial and temporal correlations which often make a direct mean-field treatment infeasible. Here, we follow a different approach based on a dynamical resampling of the connectivity ("annealing") at each spike [29, 62] (Fig. 2a). This procedure largely ignores the part of the spatial and temporal correlations that is caused by the quenched connectivity, but leads to a tractable model of a fully connected neural network with additional dynamical noise that, as we will see, retains and captures the main statistical features of the recurrent synaptic input to a neuron (Fig. 2b). We expect that the annealed approximation is better in dense networks of strongly correlated neurons. In this case, the temporal correlations in the spike input of a given neuron are partially retained upon randomization of the presynaptic neurons. In contrast, in sparse networks, the membrane potentials are independent among neurons, and therefore, the resampling of presynap-



FIG. 2. Schematic illustration of the annealed network and its approximation by a mean-connectivity network with additional noise. (a) In the annealed network model, Eq. (8), the random connectivity is re-sampled at each spike time resulting in an adjacency matrix a_{ijk} that varies at each spike k. (b) The annealed model can be mapped to a mean-connectivity network with additional noise, Eq. (12). The network is all-to-all connected with an average connectivity $\langle a_{ij} \rangle$. Additionally, each neuron is subject to an independent noise that captures the random connectivity. (c) Example trajectories of the input potentials in the mean-connectivity network with additional noise (grey). The distribution P of the input potentials follows a Gaussian distribution with a time-dependent mean $\bar{h}(t)$ and a time-dependent variance $\sigma^2(t)$. The second-order mean-field model (20) aims to describe the stochastic dynamics of these two mean-field variables.

tic neurons in time destroys the temporal correlations in the presynaptic spike train of a fixed presynaptic neuron.

In the annealed approximation, the connections a_{ij} are resampled at each spike occurring in the network. Thus, the time-varying adjacency matrix can be written as $[\hat{a}_{ijk}]$, where the additional index k labels the k-th spike occurring in the network. Then, for a fixed k, the $N \times N$ -adjacency matrix $[\hat{a}_{ij}]_k$ is still a random matrix with fixed in-degree, and therefore has independent rows but dependent elements \hat{a}_{ijk} within a given row *i*. However, the dependence caused by the fixedin-degree constraint is immaterial for the recurrent inputs in the annealed approximation. This is because only the j_k -th column of the matrix $[\hat{a}_{ij}]_k$, which has independent elements, matters for the transmission of the k-th spike, where j_k is the index of the neuron that fired the k-th spike, and because of the re-sampling of the adjacency matrix. The latter implies that for different spike times, the columns are independent. From the perspective of the recurrent inputs, we can thus equally well re-sample the elements of the adjacency matrix as independent Bernoulli variables $a_{ijk} \sim \text{Ber}(p)$ with success probability p = C/N. Thus, our model, Eq. (1), changes in the

annealed approximation to

$$\tau dh_i(t) = [-h_i + \mu(t)]dt + \frac{w}{C}dY_i(t-d),$$
 (8)

$$\frac{dY_i}{dt}(t) = \sum_{j=1}^N \sum_l a_{ijk_{j,l}} \delta(t - t_{j,l}) \tag{9}$$

where the index $k_{j,l}$ delivers the location of the *l*-th spike time $t_{j,l}$ of neuron *j* within the sequence of all spikes in the network. The quantity $Y_i(t)$ is the number of spikes received by neuron *i* until time *t* and, dY_i/dt is the corresponding spike train. We refer to this model as the *annealed network*. We note that in simulations of the annealed network, it is not necessary to re-sample the entire adjacency matrix at each spike but only the column that corresponds to the neuron that fired the spike. In other words, whenever a neuron fires a spike, this spike is transmitted independently to postsynaptic neurons with probability p = C/N.

Apart from the random connectivity, another source of quenched disorder that can be included in our theory is heterogeneity in the external current μ_0 . For the sake of clarity, we will begin our analysis without heterogeneity in μ_0 and will defer the straightforward extension to heterogeneity to a brief section in the Appendix (Sec. A).

III. DERIVATION OF THE MESOSCOPIC MODEL

A. A mean-connectivity network with additional noise that accounts for second-order connectivity statistics

The first step of the derivation is a temporal coarse-graining of the annealed model leading to an effective, fully connected network with homogeneous weights (mean-connectivity network) and additional noise. The additional noise will be the crucial difference to the mean-connectivity network discussed in Fig. 1. To coarse-grain, we integrate Eq. (8) over a small time step of length Δt :

$$\tau \Delta h_i(t) = \int_t^{t+\Delta t} \left[-h_i(s) + \mu(s)\right] ds + \frac{w}{C} \Delta Y_i(t-d).$$
(10)

Here, $\Delta h_i(t) = h_i(t + \Delta t) - h_i(t)$ is the increment of the input potential and $\Delta Y_i(t) = Y_i(t + \Delta t) - Y_i(t)$ is the number of spikes received by neuron *i* during the time step Δt . We now want to relate $\Delta Y_i(t)$ to the total number of spikes $\Delta Z(t)$ in the population during the same time step because the latter corresponds to the desired mesoscopic population activity via Eq. (3). The increment $\Delta Y_i(t)$ can be regarded as the result of a doubly stochastic process: On the one hand, given the total spike count $\Delta Z(t)$, each $\Delta Y_i(t)$ is the sum of $\Delta Z(t)$ independent Bernoulli variables according to Eq. (9) integrated over one time step. Hence, the increments $\Delta Y_i(t)$ are independent and binomially distributed with conditional mean and covariance

$$\begin{aligned} \langle \Delta Y_i(t) | \Delta Z(t) \rangle &= p \Delta Z(t), \\ \operatorname{cov}(\Delta Y_i(t), \Delta Y_i(t) | \Delta Z(t)) &= p \left(1 - p\right) \Delta Z(t) \delta_{ij}. \end{aligned}$$

If N is large such that $p\Delta Z(t) \gg 1$, the number of synaptic input spikes $\Delta Y_i(t)$ for a given $\Delta Z(t)$ is approximately Gaussian distributed. Under this condition, we can therefore write

$$\Delta Y_i(t) \approx p \Delta Z(t) + \sqrt{p (1-p) \Delta Z(t)} n_i(t), \qquad (11)$$

where $n_i(t) \sim \mathcal{N}(0, 1)$ are independent standard normal random variables. The first term in Eq. (11) is proportional to the total number of spikes or the mesoscopic population activity and thus corresponds to the input in an effective fully connected network. This term is common to all neurons. In contrast, the second term is different among neurons and only arises when p < 1. It thus captures the individual differences of the synaptic inputs in a non-fully connected ("diluted") network.

On the other hand, $\Delta Z(t)$ is itself a Poisson random variable with mean $Nr(t)\Delta t$. From Eq. (11), or, alternatively, from the law of total expectation and total covariance, we find that the synaptic inputs $\Delta Y_i(t)$ exhibit the following mean and covariance for a given stochastic population rate r(t):

$$\langle \Delta Y_i(t) \mid r(t) \rangle = Cr(t)\Delta t \cos(\Delta Y_i(t), \Delta Y_j(t) \mid r(t)) = [p + (1-p)\delta_{ij}]Cr(t)\Delta t,$$

where δ_{ij} denotes the Kronecker delta. The last equation shows that pairwise correlations of the synaptic input to different neurons are non-zero for p > 0. These correlations arise from the first term in Eq. (11) representing the part of the recurrent input that is proportional to the population activity. The fluctuations of the population activity are common to all neurons, and thus yield a source of coherent noise.

The covariance structure of the synaptic inputs $\Delta Y_i(t)$ is preserved if their representation by Eq. (11) is replaced by

$$\Delta \hat{Y}_i(t) = p\Delta Z(t) + \sqrt{C(1-p)r(t)\Delta t}n_i(t).$$

This representation allows to take the temporal continuum limit. Substituting $\Delta \hat{Y}_i(t-d)$ for $\Delta Y_i(t-d)$ in Eq. (10), dividing by Δt and taking the limit $\Delta t \rightarrow 0$, we obtain

$$\tau \frac{dh_i}{dt} = -h_i + \mu(t) + \frac{w}{N} \sum_{j=1}^N \dot{Z}_j(t-d) + w \sqrt{\frac{1-p}{C}r(t-d)} \zeta_i(t).$$
(12)

Here, we have used that the population activity, Eq. (3), can be rewritten as

$$A_N(t,\Delta t) = \frac{1}{N\Delta t} \int_t^{t+\Delta t} \sum_{i=1}^N dZ_i(t'), \qquad (13)$$

where the spike trains $\dot{Z}_i(t)$ have stochastic intensities $\phi(h_i(t^-))$. Furthermore, $\zeta_i(t)$ are independent Gaussian white noises obeying $\langle \zeta_i(t)\zeta_j(t')\rangle = \delta_{ij}\delta(t-t')$.

Equation (12) represents an effective, fully connected network with rescaled synaptic efficacy w/N = pJ and additional dynamic noise whose intensity is proportional to the

variance σ_w^2 of the coupling strengths w_{ij} (cf. Eq. (4)). Therefore, the dynamics is the same as the mean-connectivity network, Eq. (5), but with an additional noise term that captures the fluctuations caused by the random connectivity. This representation also reveals the distinct effects of finite network size $(N < \infty)$ and random dilution (p < 1) on the recurrent synaptic fluctuations. First, the stochastic input spikes from all neurons in the network contribute a common synaptic input current proportional to the population activity (third term on the right-hand side of Eq. (12)), which is stochastic for finite N and thus exposes the coherent part of the synaptic fluctuations. More precisely, the population activity has conditional mean r(t) (given all input potentials $h_i(t^-)$, $i = 1, \ldots, N$) and finite-size fluctuations of order $N^{-1/2}$ because the spike trains $Z_i(t)$ are conditionally Poisson (given $h_i(t^-)$) with conditional mean r(t).

In contrast, the last term of Eq. (12) contains a Gaussian white noise $\zeta_i(t)$ that is independent for each neuron, and thus represents the incoherent part of the synaptic fluctuations. Because the noise intensity is proportional to $\sigma_w^2 = w^2(1-p)/p$, the noise term captures the effect of the quenched random connectivity ($\sigma_w^2 > 0$), and, specifically, the random dilution (p < 1). It therefore vanishes in the limit of fully connected, homogeneous networks ($p \to 1, \sigma_w^2 \to 0$), in which case the mean-connectivity network is recovered.

In the following, we assume a large network size N such that a Gaussian approximation can be made: noting that $\Delta Z(t)$ is conditionally Poisson with mean and variance equal to $Nr(t)\Delta t$, the population activity can be approximated for large N as

$$A_N(t;\Delta t) \approx \frac{1}{\Delta t} \int_t^{t+\Delta t} A(t') dt',$$
$$A(t) = r(t) + \sqrt{\frac{r(t)}{N}} \eta(t), \quad (14)$$

where $\eta(t)$ is a common Gaussian white noise with $\langle \eta(t)\eta(t')\rangle = \delta(t-t')$. In this form, the finite-size fluctuations of order $1/\sqrt{N}$ become explicit. We note that the Gaussian approximation of the population activity, Eq. (14), can in principle become negative. However, if N is sufficiently large such that $Nr(t)\Delta t \gg 1$, the probability that such event happens is negligibly small.

B. Second-order mesoscopic mean-field dynamics

The mean-connectivity network with additional noise derived in the last section is still an N-dimensional microscopic model and thus too complicated to analyze its dynamics. We now derive a mesoscopic mean-field model that is analytically tractable. To this end, we proceed in two steps: first, we consider the dynamics of the input potentials

$$\tau \frac{\mathrm{d}h_i}{\mathrm{d}t} = -h_i + f(t) + g(t)\zeta_i(t),\tag{15}$$

driven by fixed, time-dependent input signals f(t) and g(t), and derive the dynamics of the mean and variance conditioned on these inputs. We then derive the dynamics of the stochastic rate r(t) for fixed input signals f and g such that it depends on these input signals only through the conditional mean and variance and an additional colored noise. In the second step, in turn, the input signals will be expressed in terms of the stochastic rate r(t) according to Eqs. (12), (13) and (14):

$$f(t) = \mu(t) + w \left[r(t-d) + \sqrt{\frac{r(t-d)}{N}} \eta(t-d) \right],$$

$$g(t) = w \sqrt{\frac{1-p}{C}} r(t-d)$$
(16)

This step will close the system for the conditional mean and variance and the colored noise because the stochastic rate depends itself only on these variables. In this way, we will obtain a self-consistent mean-field description for the annealed model, Eq. (12), as detailed in the following. In view of Eq. (16), we also note that the conditioning on the input signals f and g will allow for the interpretation of the mean-field variables at time t as conditional averages given the history of the noise $\eta(t'-d)$, the stochastic rate r(t'-d) and the external input $\mu(t')$ for t' < t.

In the first step, when f and g are fixed, the input potentials h_i represent N independent, time-inhomogeneous Ornstein-Uhlenbeck processes for which the ensemble mean \bar{h} and ensemble variance σ^2 obey the dynamics

$$\tau \frac{\mathrm{d}\bar{h}}{\mathrm{d}t} = -\bar{h} + f(t), \quad \tau \frac{\mathrm{d}\sigma^2}{\mathrm{d}t} = -2\sigma^2 + \frac{g^2(t)}{\tau}. \tag{17}$$

These variables are the conditional mean and variance described above. To relate the stochastic rate to these variables, we first consider the limit $N \rightarrow \infty$. In this limit, the stochastic rate, Eq. (6), converges to its conditional mean

$$F\left(\bar{h}(t),\sigma^{2}(t)\right) := \langle \phi(h_{i}(t)) \mid \bar{h}(t),\sigma^{2}(t) \rangle$$
$$= \int_{-\infty}^{\infty} \phi(h)g_{\bar{h},\sigma^{2}}(h) \, dh \qquad (18)$$

where $g_{\bar{h},\sigma^2}(h) = \exp[-(h-\bar{h})^2/(2\sigma^2)]/\sqrt{2\pi\sigma^2}$. Note that the conditional mean depends on the functions f and g only through the conditional mean $\bar{h}(t)$ and conditional variance $\sigma^2(t)$. Next, we consider the case of finite N. In this case, the stochastic rate r(t) has finite-size fluctuations of order $1/\sqrt{N}$, hence we rewrite Eq. (6) as

$$r(t) = \left[F\left(\bar{h}(t), \sigma^2(t)\right) + \frac{1}{\sqrt{N}}\xi(t)\right]_+$$
(19)

with $\xi(t) = \sqrt{N} \left[\frac{1}{N} \sum_{i=1}^{N} \phi(h_i(t)) - F\left(\bar{h}(t), \sigma^2(t)\right) \right]$. The rectification bracket $[\cdot]_+ = \max(0, \cdot)$ could be inserted because $N^{-1} \sum_i \phi(h_i)$ is always a non-negative quantity. Including the rectification is necessary for a Gaussian approximation of ξ to enforce a non-negative population rate r(t), albeit negative values are extremely rare events for biologically relevant network sizes N. For fixed input functions f and g, the second term $F\left(\bar{h}(t), \sigma^2(t)\right)$ is deterministic, and hence,

for large N, the variable $\xi(t)$ is an approximately Gaussian, colored noise with mean 0 and auto-covariance function $c_{\epsilon}^{f,g}(t,t') = \operatorname{cov}(\phi(h_i(t)), \phi(h_i(t')))$ (again taken with fixed functions f and g). Although there exist explicit expressions for the auto-covariance function of a nonlinear transformation of a Gaussian process with given auto-covariance function (see e.g. [65]), these expressions are in the form of infinite series or double integrals, and are not directly usable for the derivation of mesoscopic dynamics. Therefore, we follow a simpler ad hoc approach here: we assume that $\phi(h_i(t))$ has approximately the same temporal correlation structure as $h_i(t)$. Therefore, we make the heuristic approximation $c_{\xi}^{f,g}(t,t') \approx \frac{c_{\xi}^{f,g}(t,t)}{\sigma^2} c_h^{f,g}(t,t')$, where $c_h^{f,g}$ denotes the auto-correlation function of the process Eq. (15). Because $\xi(t)$ is Gaussian and $h_i(t)$ has an exponential auto-correlation function with correlation time τ , we hence model the colored noise as an Ornstein-Uhlenbeck process of the form

$$\tau \frac{d\xi}{dt} = -\xi + \sqrt{2\tau \sigma_{\lambda}^2(t)} \zeta(t).$$

Here, $\zeta(t)$ is Gaussian white noise with $\langle \zeta(t) \rangle = 0$ and $\langle \zeta(t)\zeta(t') \rangle = \delta(t-t')$, and

$$\begin{aligned} \sigma_{\lambda}^{2}(t) &= \left\langle \phi(h_{i}(t))^{2} \mid \bar{h}(t), \sigma^{2}(t) \right\rangle - \left\langle \phi(h_{i}(t)) \mid \bar{h}(t), \sigma^{2}(t) \right\rangle^{2} \\ &= G\left(\bar{h}(t), \sigma^{2}(t)\right) \end{aligned}$$

is the conditional variance of the stochastic intensity $\lambda(t)=\phi(h_i(t))$ given by

$$G\left(\bar{h},\sigma^{2}\right) := \int_{-\infty}^{\infty} \phi^{2}(h)g_{\bar{h},\sigma^{2}}(h) \, dh - F^{2}\left(\bar{h},\sigma^{2}\right).$$

We note again that the conditional variance depends on the functions f and g only through the conditional mean $\bar{h}(t)$ and conditional variance $\sigma^2(t)$.

We now proceed with the second step of the mean-field derivation, where we close the system in the mean-field variables \bar{h} , σ^2 and ξ self-consistently. To this end, the functions f and g are related back to the stochastic mean-field rate r(t) using Eq. (16). Inserting this equation into Eq. (17) we finally arrive at the mesoscopic mean-field dynamics

$$\tau \frac{\mathrm{d}\bar{h}}{\mathrm{d}t} = -\bar{h} + \mu(t) + w \left[r(t-d) + \sqrt{\frac{r(t-d)}{N}} \eta(t) \right]$$

$$\tau \frac{\mathrm{d}\sigma^2}{\mathrm{d}t} = -2\sigma^2 + \frac{w^2 \left(1-p\right)}{\tau p N} r(t-d)$$

$$\tau \frac{\mathrm{d}\xi}{\mathrm{d}t} = -\xi + \sqrt{2\tau G \left(\bar{h}(t), \sigma^2(t)\right)} \zeta(t),$$

(20)

where $r(t) = [F(\bar{h}(t), \sigma^2(t)) + \xi(t)/\sqrt{N}]_+$. The population activity $A_N(t, \Delta t)$ associated with this mesoscopic dynamics is given by Eq. (14). Because we consider the population mean $\bar{h}(t)$ and population variance $\sigma^2(t)$, we call Eq. (20) the 2nd-order mean-field (MF) model.

The 2nd-order MF model reveals the effects of coherent finite-size fluctuations and incoherent fluctuations caused by

the random disorder of the connectivity on the population dynamics. Firstly, finite-size noise appears in the equation for \bar{h} as an explicit, multiplicative noise term, whose magnititude is proportional to $\sqrt{r/N}$. It thus increases with larger firing rates reflecting the Poisson noise in the underlying microscopic model, and it vanishes in the large-N limit. Secondly, the incoherent fluctuations arising from the randomness of the connectivity are reflected by the equation for σ^2 , which is essentially driven by the variance of the synaptic weights $\sigma_w^2 = w^2(1-p)/p$. In turn, the variance σ^2 impacts the population rate r through the averaged nonlinearity F, Eq. (18). However, the population average, Eq. (6), is not perfectly equal to F for finite N and p < 1. The imperfect averaging leads to finite-size fluctuations $\xi(t)$ caused by both the finite network size and the random connectivity. This combined effect becomes manifest in our mean-field model by the noise strength proportional to $\sqrt{G/N}$ since G vanishes for $\sigma^2 = 0.$

If we had neglected the dispersion of the input potentials due to the random connectivity, i.e. assuming $\sigma^2 \equiv 0$, we would have arrived at a first-order MF model with only one equation for the mean $\bar{h}(t)$:

$$\tau \frac{\mathrm{d}}{\mathrm{d}t}\bar{h} = -\bar{h} + \mu(t) + w \left[r(t-d) + \sqrt{\frac{r(t-d)}{N}} \eta(t) \right]$$
$$r(t) = \phi(\bar{h}(t)). \tag{21}$$

Here, we have used that, for $\sigma = 0$, the population transfer function $F(\bar{h}, \sigma^2)$ simplifies to the transfer function of a single Poisson neuron $\phi(\bar{h}) = F(\bar{h}(t), 0)$ and that $G(\bar{h}, 0) = 0$. The 1st-order MF model (21) precisely corresponds to the limit $p \to 1$ while keeping the network size N = C/p and the coupling strength w = JpN constant. In the following, the 1st-order MF theory can thus always be obtained as a limit case of the 2nd-order theory by first re-expressing any occurrence of the parameters C and J by pN and w/(pN), respectively, and then taking the limit $p \to 1$.

Apart from a Gaussian approximation for large N, the 1st-order MF model is essentially equivalent to the meanconnectivity network, Eq. (5). Indeed, the 1st-order MF model is recovered from the exact mesoscopic mean-field dynamics, Eq. (7), of the mean-connectivity network, if the Gaussian approximation Eq. (14) is applied to the Poisson noise in Eq. (7). By comparing simulations of the microscopic models (quenched and annealed networks) with the 1st-order and 2nd-order MF models, we will thus be able to judge the mean-connectivity approximation used in previous studies [35, 49, 50] and the correction for the effect of random dilution (p < 1) given by the 2nd-order MF theory.

In the limit $N \to \infty$, the 2nd-order mean-field model, Eq. (20), converges to either a one- or a two-dimensional, deterministic model depending on whether the connectivity is dense or sparse. In the dense limit, where p is kept constant and, hence, $C = pN \to \infty$, the variance σ^2 as well as the fluctuations $N^{-1/2}\eta$ and $N^{-1/2}\xi$ vanish leaving a onedimensional, deterministic firing rate model. This case is equivalent to the large-N limit of the 1st-order MF model. In contrast, in the sparse limit, where C is kept constant and, hence, $p = C/N \rightarrow 0$, the noise terms $N^{-1/2}\eta$ and $N^{-1/2}\xi$ vanish but the variance σ^2 does not. The result is a twodimensional, deterministic mean-field model for the mean and variance of the input potentials similar to the classical model of Amari [22].

1. Functions F and G for an error-function nonlinearity

For the error function nonlinearity, Eq. (2), the functions F and G can be calculated explicitly as (see appendix, Sec. H)

$$F(h,\sigma^2) = r_{\rm m}\Phi\left(\beta_{\rm eff}(\sigma^2)h\right),\tag{22}$$

$$G(h,\sigma^2) = r_{\rm m}^2 \Phi \left(\beta_{\rm eff}(\sigma^2)h\right) - 2r_{\rm m}^2 T \left(\beta_{\rm eff}(\sigma^2)h, \beta_{\rm eff}(2\sigma^2)h\right)$$
(23)

with T being the Owen T-function and

$$\beta_{\rm eff}(\sigma^2) = \frac{\beta}{\sqrt{1+\beta^2\sigma^2}}$$

being the effective steepness of the sigmoidal nonlinearity at the population level. Because $\beta_{\text{eff}}(\sigma^2) \leq \beta_{\text{eff}}(0) = \beta$, the effective nonlinearity that governs the population dynamics has a reduced steepness compared to the the single-neuron nonlinearity (Fig. 3) and this reduction is caused by the strictly positive spread $\sigma > 0$ of the membrane potentials in a nonfully connected random network.

IV. ANALYSIS OF THE MESOSCOPIC MODEL

The reduction of the microscopic network model to a mesoscopic MF model in form of the system Eq. (20) allows us to calculate and analyze the first- and second-order statistics of mesoscopic quantities such as the population activity $A_N(t, \Delta t)$, the stochastic population rate r(t) and the population-averaged input potential $\bar{h}(t)$.

A. Stationary mean population activity

We first analyze the fixed-point solutions $\bar{h}(t) = \bar{h}_0$, $\sigma^2(t) = \sigma_0^2$ and $\xi(t) = 0$ of the deterministic system when the noise terms in Eq. (20) are set to zero and the external drive is constant, $\mu(t) = \mu_0$. In the following, we assume an inhibitory network (w < 0) balanced with a positive external drive $\mu_0 > 0$. In this case, the system exhibits a unique fixed point as shown in the Appendix, Sec I. If the fluctuations are sufficiently small and the fixed point is stable, the quantity $r_0 := F(h_0, \sigma_0^2)$ then provides an approximation of the mean stationary population activity (or equivalently, the mean stationary firing rate of neurons). At equilibrium, the noiseless version of Eq. (20) gives

$$\bar{h}_0 = \mu_0 + wr_0, \quad \sigma_0^2 = \frac{w^2(1-p)}{2\tau pN}r_0, \quad r_0 = F(\bar{h}_0, \sigma_0^2)$$
(24)



FIG. 3. Mean stationary population rate of the quenched and annealed networks and fixed-point solutions of the noiseless meanfield dynamics. (a) Graphical solution of the fixed point equation (25). Solid lines: function $F(\bar{h}, \sigma^2(\bar{h}))$ for the 1st-order (orange, $\sigma^2(\bar{h}) \equiv 0$ and 2nd-order mean-field model (blue, $\sigma^2(\bar{h})$ as in Eq. (25)). Black dashed line: left-hand side of Eq. (25). Cyan dashdotted line: F with the second argument fixed at the fixed-point value σ_0^2 . Mean population rates as a function of (b) the external input μ_0 , (c) the connection probability p, and (d) the coupling strength w. Orange (blue) line: 1st-order (2nd-order) MF theory; red squares (green circles): annealed (quenched) network simulation; dotted line: analytical approximation (26). The inset in (d) shows the same data, but for a larger range of w in a double logarithmic scaling. Grey dashed line: 2nd-order theory r_{∞} for the firing rate in the limit $w \to -\infty$ (Eq. (29)). Parameters: $N = 1000, \beta = 5 \text{ mV}^{-1}$ (a): p = 0.1,w = -0.7 mVs, and $\mu_0 = 5$ mV. (b): p = 0.1 and w = -1 mVs, (c): $\mu_0 = 10 \text{ mV}$, w = -1 mVs, (d): p = 0.1, $\mu_0 = 10 \text{ mVs}$.

which yields a fixed-point equation for h_0 :

$$\frac{\bar{h}_0 - \mu_0}{w} = F\left(\bar{h}_0, \frac{w(1-p)}{2\tau pN} \left[\bar{h}_0 - \mu_0\right]\right).$$
(25)

Note that both sides of the equation correspond to the stationary firing rate r_0 . Furthermore, the fixed points of the 1storder MF model are obtained by setting p = 1 in Eq. (25). The solutions for both 1st- and 2nd order MF models can be determined graphically (Fig. 3a): While the left-hand side is identical for the two MF models and is described by a line, the nonlinear function F on the right-hand side of Eq. (25) differs in its second argument σ_0^2 . For the 1st-order mean-field theory, we have $\sigma_0^2 = 0$ and therefore the sigmoidal function is steeper compared to the 2nd-order theory, for which σ_0^2 is strictly positive. Because we assume an inhibitory network (w < 0), the linear function on the left-hand side of Eq. (25) has a negative slope. If we choose w such that the intersection occurs for values h_0 lower than the inflection point $\vartheta = 0$ of the hazard function $\phi(h)$ (corresponding to the biologically relevant convex part of $\phi(h)$), then from the graphical solution it is clear that the 2nd-order model exhibits a fixed-point with lower mean input potential \bar{h}_0 and higher population firing rate r_0 compared to the 1st-order MF model. This effect becomes stronger for smaller connection probability p.

The mean stationary firing rates of the quenched and annealed microscopic models are well captured by the numerical solution of the fixed-point equation (25) (Fig. 3b-d). As a function of the external drive μ_0 , we observe a strikingly linear dependence between the saturation regimes at low and high values of μ_0 (Fig 3b). Furthermore, we find that the firing rate is roughly constant when the connection probability p is varied while N is fixed (Fig 3c). The weak dependence on p is slightly better predicted by the 2nd-order MF theory than by the 1st-order MF theory, which, by definition, has no dependence on p. We also observe a power-law behavior of the mean firing rate in both microscopic simulations and fixed-point solutions if the coupling strength is not too strong (Fig. 3d). We note that, in general, the relative error of the 1storder MF model (and hence the mean-connectivity network) with respect to the stationary mean population rate is rather small compared to the much larger error with respect to the second-order statistics reported in Fig. 1. The second-order statistics will be analyzed later in Sec. IV E.

1. Analytical approximation of fixed points

The linear dependence on μ_0 , the weak dependence on pand the power-law dependence on w can be explained with our MF theory as follows. Firstly, we notice that for a sufficiently large steepness $\beta \gg 1/\sigma_0^2$ of the single-neuron transfer function ϕ , the 2nd-order MF dynamics remains practically unaltered if we take the limit $\beta \to \infty$. In this limit, the singleneuron transfer function tends to $r_m \theta(\bar{h})$, where θ denotes the Heaviside step function. Indeed, for the values of β considered in this paper, taking the limit $\beta \to \infty$ does not noticeably alter the effective nonlinearity F and hence the 2nd-order MF dynamics. The limiting value of the fixed point can be calculated analytically from the graphical representation (Fig. 3) which yields an approximate analytical expression for the firing rate in the 1st-order MF model,

$$r_{0} \approx \begin{cases} 0, & \mu_{0} \leq 0 \\ -\frac{\mu_{0}}{w}, & 0 < \mu_{0} < -wr_{m} \\ r_{m}, & \mu_{0} \geq -wr_{m}. \end{cases}$$
(26)

This analytical expression predicts a power-law behaviour as a function of coupling strength in the balanced regime $0 < \mu_0 < -wr_m$. Indeed, the numerical fixed-point solution for the 1st-order MF model exhibits an excellent quantitative agreement with this prediction (Fig. 3d).

Secondly, we note from the geometry of the graphical solution of the fixed points that the intersection point of the 1storder MF model serves as a rough approximation for the firing rate r_0 of the 2nd-order MF model as well (Fig. 3a) unless wis extremely negative as discussed below. Indeed, the horizontal stretching of the transfer function in the 2nd-order theory mainly affects the value of the fixed point \bar{h}_0 rather than the firing rate r_0 . More precisely, a sufficient condition for the validity of our approximation is that $-\mu_0/w \gg (1-p)/(\tau C)$ and $-\mu_0/w > 0.023 r_{\rm m}$ (see Appendix C). We expect that the approximation (26) slightly underestimates the firing rates if the intersection is below the inflection point ($\bar{h}_0 < 0$) and slightly overestimates the firing rates if the intersection is above the inflection point ($\bar{h}_0 > 0$). Our approximation Eq. (26) well explains the piecewise linear behavior of the firing rate as a function of the external input as observed in Fig. 3b. Furthermore, Eq. (26) also explains why the firing rates are roughly independent of the connection probability p(Fig. 3c), and why we see the power-law $-\mu_0/w$ as a function of coupling strength w even in non-fully connected networks (Fig. 3d) unless w is extremely negative (this case is treated below). In general we note that, compared to the 1st-order model, the 2nd-order MF model shows a better agreement with the firing rate in microscopic simulations, in particular, it fits the annealed network almost perfectly.

Because closed-form analytical approximation of the fixed points will be crucial for a qualitative—if not quantitative discussion of the linear analysis around fixed points in the following sections, we provide here also approximations for the fixed point values \bar{h}_0 and σ_0^2 . Using our approximation for the firing rate, Eq. (26), we obtain from Eq. (24) an approximation for the dispersion:

$$\sigma_0^2 \approx -\frac{w\mu_0(1-p)}{2\tau Np} \tag{27}$$

if $0 < \mu_0 < -wr_m$. To obtain a corresponding approximation for \bar{h}_0 in the 2nd-order MF theory, we solve the fixed point equation $r_0 = F(\bar{h}_0, \sigma_0^2)$ (cf. Eq. (24)) for \bar{h}_0 using our approximations for r_0 and σ_0^2 resulting in

$$\bar{h}_0 \approx \Phi^{-1} \left(-\frac{\mu_0}{w r_{\rm m}} \right) \sqrt{\beta^{-2} + \sigma_0^2}.$$
 (28)

2. Limit of strongly inhibitory networks

Finally, the regime of strong inhibitory coupling reveals a qualitative difference between the the 1st- and 2nd-order mean-field prediction. In the limit $w \to -\infty$, the right-hand side of Eq. (25) is asymptotic to

$$r_{\rm m}\Phi\left(-\sqrt{\frac{2\tau pN(\bar{h}_0-\mu_0)}{(1-p)w}}
ight),$$

and thus the mean stationary firing rate in a strongly inhibitory network, is given by the solution r_{∞} of the limiting equation

$$r_{\infty} = r_{\rm m} \Phi\left(-\sqrt{\frac{2\tau p N r_{\infty}}{1-p}}\right). \tag{29}$$

The graphical solution of this equation clearly shows a vanishing firing rate in the 1st-order MF theory $(p \rightarrow 1)$, whereas the limiting equation has a non-vanishing solution for p < 1 (2ndorder MF theory, Fig. 3d, inset). The non-vanishing, limiting firing rate in the 2nd-order MF model is in good agreement with simulations of the annealed and quenched microscopic



FIG. 4. Second-order MF theory explains stable asynchronous activity and oscillatory instability in the presence of a transmission delay d > 0. (a) Sample trajectories for quenched (green) and annealed network (red), 2nd-order MF model (blue) and 1st-order MF model (orange) with d = 2 ms. (b) (d, w)-phase diagram with Hopf boundaries (onset of oscillations) for N = 20000 neurons. The heat map shows the standard deviation of the population rate r(t) in the annealed network to indicate the presence of oscillations. Green line: contour line for the quenched network at which the standard deviation of r(t) equals a threshold of 2 Hz, which was visually inferred from panel (d). A non-zero threshold was chosen because of finitesize fluctuations. Blue (orange) line: Hopf boundary for 2nd- (1st-) order MF model, calculated from Eq. (31) (c) Standard deviation of the stochastic population rate for different connection strengths w at a delay d = 2 ms calculated from simulations (circles: quenched network, squares: annealed network, diamonds: 2nd-order MF, triangles: 1st-order MF). (d) Standard deviation of the stochastic population rate for different connection strengths d and fixed coupling strength w = -1 mVs. Parameters: C = 100, $\mu = 10$ mV, $\beta = 5 \ \mathrm{mV}^{-1}.$

model. In contrast, the 1st-order MF model generally underestimates the firing rates in strongly inhibitory networks: for $w \to -\infty$, the 1st-order MF model has a rate that converges to zero like $-\mu_0/w$ as predicted by Eq. (26).

B. Stability of the fixed-point solutions

In the presence of a non-zero transmission delay d > 0, the fixed-point solution of the mean-field model may become unstable, leading to an oscillatory regime, see e.g. [66]. For such an instability to correctly predict oscillatory behavior of our microscopic network model, our mean-field theory must accurately describe the recurrent fluctuations in the network. Comparing simulations of the microscopic network models with simulations of the first- and second-order mean-field models (Fig. 4a), we observe that for the chosen parameters the 1storder MF model wrongly predicts strong oscillations when the actual population activity of the network is nearly constant corresponding to an asynchronous firing regime. In contrast, the 2nd-order theory correctly reproduces the constant population activity.

The onset of oscillations can be understood by a linear stability analysis. To this end, we linearize the noiseless system around the fixed point. Small deviations of the mesoscopic variables, $\delta \bar{h} = \bar{h}(t) - \bar{h}_0$, $\delta \sigma^2 = \sigma^2(t) - \sigma_0^2$, obey the linearized dynamics

$$\frac{\mathrm{d}}{\mathrm{d}t}\boldsymbol{X}(t) = \mathrm{T}\,\boldsymbol{X}(t) + \mathrm{W}\,\boldsymbol{X}(t-d), \tag{30}$$

where we introduced the deviation vector $\boldsymbol{X}(t) = (\delta \bar{h}(t), \delta \sigma^2(t), \xi(t))^T$ and the matrices

$$\mathbf{T} = -\frac{\operatorname{diag}(1,2,1)}{\tau}, \quad \mathbf{W} = \left(\frac{w}{\tau}, \frac{w^2 (1-p)}{\tau^2 p N}, 0\right)^T \boldsymbol{L}^T$$

with $\boldsymbol{L}^T = (F_h, F_\sigma, N^{-1/2})$. Here, F_h and F_σ denote the partial derivatives of $F(h, \sigma^2)$ with respect to h and σ^2 at the fixed point (exact and approximate expressions for F_h and F_σ are given in the Appendix D). We note that any perturbation in ξ exponentially decays to zero with a time constant τ . It is therefore sufficient to look for solutions of the form $\boldsymbol{X}(t) = e^{\lambda t} \left(\hat{h}(\lambda), \hat{\sigma}(\lambda), 0 \right)^T$ with some constant parameter λ . The sign of the real part of λ then determines the stability of the fixed point; in particular, if $\operatorname{Re}(\lambda) > 0$ the fixed-point is unstable. As shown in Appendix E, the parameter λ must satisfy the characteristic equation

$$\lambda \tau = -1 + wF_h e^{-\lambda d} + \frac{wF_h \hat{F}_\sigma e^{-2\lambda d}}{2 + \lambda \tau - \hat{F}_\sigma e^{-\lambda d}}, \quad (31)$$

where $\hat{F}_{\sigma} := w^2 \frac{1-p}{\tau C} F_{\sigma}$.

To find the onset of oscillations, we look for a Hopf bifurcation, at which $\lambda = i\omega$ with a frequency $\omega \in \mathbb{R} \setminus \{0\}$. Imposing such purely imaginary values of λ in Eq. (31) allows us to find the "Hopf boundary" of the oscillatory phase in the parameter space. In the following, we investigate the stability depending on the transmission delay d and the coupling strength w (Fig. 4b). In the (d, w)-space, we can see that the 2nd-order MF model significantly differs from the first-order one. For sufficiently strong coupling strength, the 1st-order MF model exhibits a Hopf boundary at small values of d, predicting oscillations already at small delays. In contrast, the 2nd-order model predicts oscillation onset for much larger delays. Interestingly, for sparse connectivity (sufficiently small p), a stronger coupling strength w requires a larger delay to enable oscillations, whereas the first-order model requires a smaller delay. To assess whether the microscopic models are in an oscillatory state in our simulations, we calculate the standard deviation of a time series of the stochastic population rate r(t) as given in Eq. (6). We find that the oscillatory regime of the annealed network (Fig. 4b, red-colored area) coincides

with the area enclosed by the Hopf boundary obtained from the 2nd-order MF theory. A qualitatively similar oscillation boundary is observed in the quenched network model.

For a more detailed comparison of the oscillation amplitude beyond the Hopf bifurcation, we also computed the standard deviation of r(t) in the mean-field models. We find that the 2nd-order MF model matches well the annealed network model (Fig. 4c,d), whereas the 1st-order mean-field model fails to describe the oscillations already at small delays. The quenched microscopic model has an oscillation onset at even larger delays, however, with a qualitatively similar dependence on the delay and coupling strength as the annealed microscopic model.

In conclusion, our stability analysis of the 2nd-order MF model shows that a randomly diluted connectivity leads to a stabilisation of the non-oscillatory state. In particular, the random dilution leads to incoherent fluctuations of the recurrent input, which flattens the effective nonlinearity of the macroscopic dynamics and thus changes the stability properties of the fixed point.

C. Linear response of the mean activity

We now analyse how the network responds to a small, timedependent perturbation of the external current, $\mu(t) = \mu_0 + \mu_1(t)$ by calculating the linear response function [67]. To this end, we linearize our mesoscopic mean-field model, Eq. (20), around the fixed point and obtain a generalization of Eq. (30):

$$\frac{\mathrm{d}}{\mathrm{d}t}\boldsymbol{X}(t) = \mathrm{T}\,\boldsymbol{X}(t) + \mathrm{W}\,\boldsymbol{X}(t-d) + \boldsymbol{M}(t) + \mathrm{B}\,\boldsymbol{\zeta}(t).$$
 (32)

Here, the vector X(t) represents again the deviation from the fixed point as in the previous section and the threedimensional vector M(t) represents a general small perturbation to the system. Furthermore, the last term represents the finite-size noise, where

$$\mathbf{B} = \begin{pmatrix} \frac{w}{\tau} \sqrt{\frac{\tau_0}{N}} & 0\\ 0 & 0\\ 0 & \sqrt{\frac{2G_0}{\tau}} \end{pmatrix}, \quad \boldsymbol{\zeta}(t) = \begin{pmatrix} \eta(t)\\ \boldsymbol{\zeta}(t) \end{pmatrix}$$

and $G_0 = G(\bar{h}_0, \sigma_0^2)$. In the linearized dynamics we neglected terms of order O(1/N) and higher. In the Fourier domain, the solution reads

$$\tilde{\boldsymbol{X}}(\omega) = \tilde{\chi}(\omega) \left(\tilde{\boldsymbol{M}}(\omega) + \mathbf{B} \, \tilde{\boldsymbol{\zeta}}(\omega) \right), \tag{33}$$

where we introduced the susceptibility matrix

$$\tilde{\chi}(\omega) = \left[i\omega \,\mathbf{I}_3 - \mathbf{T} - \mathbf{W} \,e^{-i\omega d}\right]^{-1}.\tag{34}$$

Here, I₃ denotes the 3 × 3 identity matrix and the tilde notation $f(\omega)$ for a function f(t) denotes its Fourier transform, i.e. $\tilde{f}(\omega) = \int_{-\infty}^{\infty} f(t)e^{-i\omega t} dt$. The explicit expressions for the elements of $\tilde{\chi}$ are given in the appendix, Eq. (F1).

The susceptibility matrix provides the response of the ensemble mean $\langle \mathbf{X}(t) \rangle$ of the deviation from the fixed point



FIG. 5. Linear response of the time-dependent mean population rate ("rate susceptibility") to a weak modulation of the external current. (a) $\tilde{\chi}_r$ for the microscopic model with quenched (green circle) and annealed (red squares) networks for p = 0.1, d = 0 ms, and the corresponding 1st- (orange) and 2nd-order (blue) mean-field theories, Eq. (35) (where p is set to 1 for the 1st-order MF theory). Top and bottom row: amplitude $|\tilde{\chi}_r|$ and phase ϕ_r of the susceptibility, respectively. Dashed horizontal line: $\phi_r = -\pi/2$. (b) Same for p = 0.4, d = 0.001 s. Other parameters in (a) and (b): N = 1000, w = -1 mVs.

of the full nonlinear system to an infinitesimally small perturbation $\boldsymbol{M}(t)$ in the Fourier domain via the relation $\langle \tilde{\boldsymbol{X}}(\omega) \rangle = \tilde{\chi}(\omega) \tilde{\boldsymbol{M}}(\omega)$. Similarly, the response of the ensemble-averaged firing rate $\langle r(t) \rangle = F(h_0, \sigma_0^2) + \delta r(t)$ is given in the Fourier domain by $\delta \tilde{r}(\omega) = \boldsymbol{L}^T \tilde{\chi}(\omega) \tilde{\boldsymbol{M}}(\omega)$. For an external stimulus $\mu(t) = \mu_0 + \mu_1(t)$, the perturbation is $\tilde{\boldsymbol{M}}(\omega) = (\tilde{\mu}_1(\omega)/\tau, 0, 0)^T$. In this case, we then have $\delta \tilde{r}(\omega) = \tilde{\chi}_r(\omega) \tilde{\mu}_1(\omega)$, where we introduced the rate susceptibility

$$\tilde{\chi}_r(\omega) = \frac{1}{\tau} \left[F_h \tilde{\chi}_{11}(\omega) + F_\sigma \tilde{\chi}_{21}(\omega) \right].$$
(35)

Here, we used that $\tilde{\chi}_{31} \equiv 0$.

The rate susceptibility is in general complex-valued with absolute value $|\tilde{\chi}_r(\omega)|$ and phase $\phi_r(\omega)$. A straightforward interpretation is that the system driven by a sinusoidal external input modulation $\mu_1(t) = \epsilon \sin(\omega t)$ with small amplitude ϵ will respond with a sinusoidal rate modulation with amplitude $\epsilon |\tilde{\chi}_r(\omega)|$ and a phase shift $\phi_r(\omega)$. For all models studied here, the amplitude decreases for high frequencies with a power law with exponent -1 (Fig. 5). The annealed network and the 2nd-order MF model agree very well, in contrast to the 1st-order MF model which massively overestimates the amplitude at high frequencies. The quenched network shows a slightly smaller amplitude than predicted by the 2nd-order MF model, but is still reasonably well approximated by the latter. While the phase shift vanishes at low frequencies as expected, the phase shift approaches $-\frac{\pi}{2}$ in the limit of large frequencies (Fig. 5). We can understand this behaviour analytically by considering the limits of Eq. (35). For $\omega = 0$ we obtain

$$\tilde{\chi}_r(0) = \frac{F_h}{1 - wF_h - \frac{w^2}{2\tau}F_\sigma \left(\frac{1}{C} - \frac{1}{N}\right)},$$
(36)

which is real-valued and has therefore a phase zero. The amplitude in the low-frequency limit is approximately the negative inverse of the coupling strength $-w^{-1}$, under the condition that $1 \ll wF_h$ and C large. Similarly, for large frequencies we find asymptotically

$$\tilde{\chi}_r(\omega) \sim -i \frac{F_h}{\tau} \cdot \frac{1}{\omega}, \qquad \omega \to \infty.$$
(37)

In this limit, the phase converges to $-\frac{\pi}{2}$ and the amplitude follows a power law ω^{-1} scaled by F_h/τ . The different values for the derivative F_h explain the difference between the 1storder and the 2nd-order MF model. To understand this difference analytically, we use the following approximation for the slope of the transfer function at the fixed point (see Appendix D):

$$F_h \approx \frac{r_{\rm m} Q\left(-\frac{\mu_0}{w r_{\rm m}}\right)}{\sqrt{2\pi \left(\beta^{-2} + \sigma_0^2\right)}},\tag{38}$$

where σ_0^2 is given by Eq. (27). Furthermore, we introduced the non-dimensional function

$$Q(x) := \exp\left[-\frac{1}{2}\Phi^{-1}(x)^2\right]$$
 (39)

defined on the interval [0, 1]. This function has an inverted-U shape and is symmetric with respect to $x = \frac{1}{2}$ (a graph of Q is shown in Fig. 8c, dotted lines). Equation (38) clearly shows that the factor F_h increases with the connection probability p because σ_0^2 decreases with p, cf. Eq. (27). The first-order MF theory corresponding to the mean-connectivity network (without additional noise) is obtained by setting p = 1 in Eq. (27). In this case, the dispersion σ_0^2 vanishes, and hence the slope F_h is maximized. Therefore, the 1st- and 2nd-order models predict the same power-law behaviour, but with a higher



FIG. 6. The first-order mean-field theory strongly underestimates the power spectral density of the population activity. Simulations of the power spectral densities of the population activity $A_N(t; \Delta t)$ for the microscopic (green circles: quenched network, red squares: annealed network) and mesoscopic models (orange triangle: 1storder MF model, blue diamonds: 2nd-order MF model). Solid lines depict the 1st- and 2nd-order MF theory, Eqs. (G1) and (42), respectively. Parameters: w = -1 mVs a) N = 500, p = 0.07, $\mu(t) = \mu_0 = 10$ mV.

magnitude for the 1st-order model (Fig. 5a). With increasing connection probability p the susceptibility approaches the 1st-order MF theory. The low-frequency limit is in both mesoscopic models almost independent of p as the low-frequency limit is roughly given by $-w^{-1}$. Simulations of the annealed network model match the predictions of the 2nd-order MF model while the quenched network model has slightly lower amplitude and slightly lower phase for intermediate frequencies. Overall, the 2nd-order MF theory yields a strongly improved prediction compared to the 1st-order MF model.

The synaptic delay d appears in $\tilde{\chi}_r$ only via the complex exponential in Eq. (34). On the other hand, the delay does not influence the low-frequency limit and the high-frequency asymptotics, Eqs. (36) and (37) of the response, respectively. For intermediate frequencies the complex exponential causes a modulation and a peak in the amplitude as a function of ω (Fig. 5b). Simulations of the annealed network model perfectly confirm the prediction of the 2nd-order MF model, whereas the quenched network model has a less pronounced peak at a lower frequency.

D. Power spectral density

Now we turn to the second-order statistics of the stationary dynamics. We assume that the external stimulus $\mu(t)$ is a stationary process with given mean μ_0 and power spectral density $S_{\mu\mu}(\omega)$. The second-order statistics can be characterized, in the frequency domain, by the power spectral density matrix of the mean-field state vector X(t) (cf. Eq. (30)). This matrix involves, e.g., the power spectrum of the average input potential $\bar{h}(t) = \bar{h}_0 + X_1(t)$ but also cross spectra such as between $\bar{h}(t)$ and $\sigma^2(t) = \sigma_0^2 + X_2(t)$. Furthermore, we are interested in the power spectra of the average intensity r(t) and the population activity A(t), as well as in the variance of r(t). The latter is studied in the subsequent section.

The power spectral density matrix $\mathbf{S}(\omega)$ is defined via the relation

$$\left\langle \tilde{\boldsymbol{X}}(\omega)\tilde{\boldsymbol{X}}^{*}(\omega')\right\rangle = 2\pi\,\mathrm{S}(\omega)\delta(\omega-\omega')$$
 (40)

where * denotes the conjugate transpose of a matrix. Using Eq. (33) for the linearized system, we find the following approximation for $S(\omega)$ describing weak fluctuations around the fixed points (see Appendix, Sec. G):

$$S_{ij}(\omega) = \frac{1}{\tau^2} \left[S_{\mu\mu}(\omega) + \frac{w^2 r_0}{N} \right] \tilde{\chi}_{i1}(\omega) \tilde{\chi}_{j1}^*(\omega) + \frac{2G\left(\bar{h}_0, \sigma_{h,0}^2\right)}{\tau} \tilde{\chi}_{i3}(\omega) \tilde{\chi}_{j3}^*(\omega).$$

The matrix S gives us the (cross-) spectral densities between \bar{h} , σ^2 and ξ . For the power spectral densities of the stochastic firing rate r(t) and the activity A(t), we use the linear approximation

$$\tilde{r} \approx F_h \tilde{h} + F_\sigma \widetilde{\sigma^2} + \frac{1}{\sqrt{N}} \tilde{\xi}, \quad \tilde{A} \approx \tilde{r} + \sqrt{\frac{r_0}{N}} \tilde{\eta}.$$
 (41)

These linear equations allow us to compute the power spectral density $S_{rr}(\omega)$ of the stochastic population rate r(t) via the relation $\langle \tilde{r}(\omega) \tilde{r}^*(\omega') \rangle = 2\pi S_{rr}(\omega) \delta(\omega - \omega')$, which yields

$$\begin{split} S_{rr}(\omega) &\approx F_h^2 S_{11}(\omega) + F_{\sigma}^2 S_{22}(\omega) + \frac{1}{N} S_{33}(\omega) + \\ 2F_h F_{\sigma} \Re \mathfrak{e}(S_{12}(\omega)) + 2 \frac{F_h \Re \mathfrak{e}(S_{13}(\omega)) + F_{\sigma} \Re \mathfrak{e}(S_{23}(\omega))}{\sqrt{N}} \end{split}$$

Further, we find, using Eq. (41), the power spectrum of A(t) within the linear approximation as

$$S_{AA}(\omega) \approx S_{rr}(\omega) + \frac{r_0}{N} + 2\sqrt{\frac{r_0}{N}} \mathfrak{Re}(S_{r\eta}(\omega)).$$
(42)

The cross spectral density $S_{r\eta}$ that appears here can be calculated using Eq. (41) and (33):

$$S_{r\eta}(\omega) \approx \frac{w}{\tau} \sqrt{\frac{r_0}{N}} \left[F_h \tilde{\chi}_{11}(\omega) + F_\sigma \tilde{\chi}_{21}(\omega) \right].$$

In Fig. 6, we compare the theory for the power spectrum of A(t), Eq. (42), to simulations of $A_N(t, \Delta t)$ in the microscopic and mesoscopic models. For all models, the power spectra exhibit a trough at low frequencies. The 1st-order MF model massively underestimates the observed low-frequency power, whereas the 2nd-order MF model matches well the microscopic model with annealed connectivity. Compared to the quenched network, the low-frequency power of the annealed network is still significantly reduced. Thus, despite the considerable improvement compared to the 1st-order MF theory, the underestimation of low-frequency power reveals a short-coming of the annealed approximation.



FIG. 7. Stationary variance of the population rate and mean input potential for varying connection probability p. (a) The number of neurons is fixed to N = 1000 (hence C = 1000p). (b) The indegree is fixed to C = 100 (hence N = 100/p). Left panels of (a) and (b): Variance of the population rate r(t) (Symbols: simulations of microscopic networks and the mesoscopic MF models as indicated in the legend; orange and blue solid lines: full theory Eq. (43) and Eqs. (G4); black dashed line: analytical approximation for the variances based on the 2nd-order MF theory, Eq. (45), (46) and Eqs. (44), (48) for var(r) and var(\bar{h}), respectively). Right panels of (a) and (b): Same as left panels but for variance of \bar{h} . Parameters: $\mu_0 = 10$ mV, w = -1 mVs, $\sigma_{\text{ext}} = 0$ mV, $\beta = 5$ mV⁻¹

E. Variances

The stationary variances of r(t) and h(t) can be calculated by standard methods for the linearized system in the absence of synaptic delay, d = 0 (see Appendix, Sec. G). Furthermore, we assume that the external stimulus $\mu(t)$ has mean μ_0 and white Gaussian fluctuations of strength σ_{ext}^2 as in Sec. II B. Under these assumptions, one eventually obtains

$$\operatorname{var}(r) \approx F_{h}^{2} \sigma_{11} + F_{\sigma}^{2} \sigma_{22} + 2F_{h} F_{\sigma} \sigma_{12} + \frac{2F_{h}}{\sqrt{N}} \sigma_{13} + \frac{2F_{\sigma}}{\sqrt{N}} \sigma_{23} + \frac{1}{N} G(\bar{h}_{0}, \sigma_{0}^{2}). \quad (43)$$

where $\sigma_{ij} = \langle X_i(t)X_j(t) \rangle$ are the elements of the covariance matrix of the state vector $\mathbf{X}(t)$. In particular, σ_{11} yields the variance of the population-averaged input potential $\bar{h}(t)$. Approximate analytical expressions for σ_{ij} can be calculated using the linearized system, Eq. (32), and are given in the appendix, Sec. G. Our theory, Eq. (43), agrees well with microscopic simulations (Fig. 7 and 8), discussed in more detail below. In general, we find a good quantitative match between the theory and simulations of the annealed network. While, in the case of quenched random connectivity, the theory again exhibits quantitative deviations, our 2nd-order MF theory still captures the parameter dependence of the variance qualitatively and yields a much better quantitative prediction than the 1st-order MF theory.

1. Dependence on the connection probability

Which connection probability causes the largest variability in our network model? To address this question, we analyze the variances of the population rate r and the average input potential \bar{h} as a function of p (Fig. 7). To avoid a trivial dependence through the mean firing rate, we vary p such that the mean firing rates remain roughly unchanged. According to Eq. (26), a constant rate can be achieved by keeping the coupling strength w = NpJ constant. In an experimental setting, where the synaptic efficacy J is the basic physiological parameter, this constraint could be realized in two different scenarios, either by fixing the network size N and changing the synaptic efficacy J such that Jp = const., or by fixing both the number of input connections C = Np and the synaptic efficacy J.

In the first scenario with constant N, we find that the variance of the rate r(t) increases with connection probability in microscopic network simulations (Fig. 7a, left), although the variance of the input potential \bar{h} decreases (Fig. 7a, right). The 1st-order MF model cannot explain this behavior because the model equation (21) does not depend on the parameter p when w is fixed and N is constant. Thus, for all $p \in [0, 1]$, the variance predicted by the 1st-order MF model corresponds to the fully connected case p = 1, i.e. to the mean-connectivity network. In contrast, the dependence on pof the variances measured from simulations of the annealed network are quantitatively well reproduced by the 2nd-order MF theory, Eq. (43). The 2nd-order theory also captures the variances of the quenched network qualitatively. The quantitative agreement is also reasonably good, however, we find that the rate variance is overestimated and the variance of the population-averaged input potentials is underestimated by the annealed network. Nevertheless, the prediction of variances by the 2nd-order theory yields a significant improvement over the 1st-order MF theory.

In order to gain an analytical understanding, we use a simplified theory based on an approximation of the variance of \bar{h} when $-wF_h \gg 1$ (see Appendix, Sec. G) :

$$\operatorname{var}(\bar{h}) \equiv \sigma_{11} \approx \left(\frac{\mu_0}{2\tau N} + \frac{\sigma_{\text{ext}}^2}{-2w}\right) \frac{1}{F_h}.$$
 (44)

Furthermore, for the parameters used in this study, we find that the variance of r is vastly dominated by the first term in Eq. (43), while the other terms are small enough to be ignored. With Eq.(44), we thus obtain the following approximation

$$\operatorname{var}(r) \approx \left(\frac{\mu_0}{2\tau N} + \frac{\sigma_{\text{ext}}^2}{-2w}\right) F_h. \tag{45}$$

The variance formulas (44) and (45) share a common factor that represents the sum of two contributions to the variability. The first term of the sum corresponds to the intrinsically generated variability caused by finite-size fluctuations of order $\sqrt{r_0/N}$ in the mesoscopic MF model, Eq. (20). We recall that the dependence of these fluctuations on the firing rate $r_0 = -\mu_0/w$, arises from the Poisson spiking noise at the microscopic level. Thus, the Poisson property that the variance scales proportionally to the mean causes finite-size fluctuations that increase with the mean stimulus strength μ_0 (see below). We emphasize that this contribution to the variability is a clear finite-size effect that vanishes in the limit $N \to \infty$. In contrast, the second term of the sum in Eqs. (44) and (45) represents externally generated variability proportional to the variance of the common external stimulus.

For constant N, the *p*-dependence in both formulas, Eq. (44) and (45), is solely contained in the slope F_h of the transfer function. As discussed in Eq. (38) above, the slope F_h is a monotonously increasing function of *p*. The slope dependence thus explains the increase of the variance of the population rate (Fig. 7a left) and the decrease of the variance of the mean input potential (Fig. 7a right), respectively.

In the second scenario with constant C and varying population size N = C/p, we observe different dependencies of the variances on p compared to the first scenario with constant N (Fig. 7b). In simulations of microscopic networks and 2nd-order MF model, the rate variance increases supralinearly with p and the variance of the mean input potential \bar{h} exhibits a non-monotonic behavior with a maximum inside the range 0 . Again, the 1st-order MF theory does not capture these behaviors. Note that there is a good qualitative agreement between 2nd-order MF theory and microscopic network simulations. As in the first scenario, however, the variance of the average input potential in the quenched network is underestimated by the annealed network, which, in turn, is quantitatively well matched by the 2nd-order MF theory.

To gain a theoretical understanding of these observations, we use our analytical approximation Eq. (45). For constant C, the approximation can be rewritten as

$$\operatorname{var}(r) \approx \frac{r_{\rm m}^2 \hat{\beta}}{\sqrt{8\pi}} \left(\frac{\hat{\mu}_0 p}{\hat{\tau} C} + \hat{\sigma}_{\rm ext}^2 \right) \hat{F}_h. \tag{46}$$

where we have introduced the dimensionless parameters $\hat{\mu}_0 = -\mu_0/(wr_{\rm m})$, $\hat{\sigma}_{\rm ext}^2 = \sigma_{\rm ext}^2/(wr_{\rm m})^2$, $\hat{\beta} = -\beta wr_{\rm m}$, $\hat{\tau} = \tau r_{\rm m}$, as well as the dimensionless slope of the transfer function

$$\hat{F}_{h} := \sqrt{2\pi} \frac{F_{h}}{\beta r_{m}} \approx \frac{Q(\hat{\mu}_{0})}{\sqrt{1 + \frac{\hat{\beta}^{2}(1-p)}{2\hat{\tau}C}\hat{\mu}_{0}}}$$
(47)

(cf. Eq. (38)). The expression for the rate variance in Eq. (46) is a product of two monotonously increasing functions of p – a linear function (the prefactor of \hat{F}_h) and a strictly convex function (\hat{F}_h). Hence, the variance of the population rate is a monotonosly increasing, strictly convex function of p explaining our observation in Fig. 7b (left). In contrast, the variance of the mean input potential, Eq. (44), can be rewritten for constant C and dimensionless parameters as

$$\operatorname{var}(\bar{h}) \approx \frac{\sqrt{2\pi}w^2 r_{\rm m}^2}{2\hat{\beta}} \left(\frac{\hat{\mu}_0 p}{\hat{\tau} C} + \hat{\sigma}_{\rm ext}^2\right) \frac{1}{\hat{F}_h}.$$
 (48)

This expression is a product of a monotonously increasing and a monotonously decreasing function of p (the prefactor of $1/\hat{F}_h$ and $1/\hat{F}_h$, respectively). The product may therefore exhibit a non-monotonic behavior. A closer inspection





FIG. 8. Variance of the stochastic population rate r(t) for varying mean external stimulus μ_0 . (a) Moderate-size network of $N = 10^3$ neurons whose inputs exhibit internally generated, finite-size noise but no external noise, $\sigma_{\text{ext}} = 0$. Left: Response to a step stimulus, where μ_0 jumps from 28 mV to 50 mV at time t = 0.04 s. Blue and orange line: simulations of the 2nd-order MF model for C = 100and C = 1000, respectively (the latter case (p = 1) corresponds to the 1st-order MF model or mean-connectivity approximation of the former case (p = 0.1); black dotted lines: theoretical prediction of stationary variances, Eq. (43); gray dashed line: stimulus onset. The inset magnifies the same data for C = 100. Right: Stationary variance of r(t) vs. mean stimulus strength μ_0 . (Symbols: simulations of quenched network (green circles occluded by red squares), annealed network (red squares), both for C = 100, 1st- (yellow triangles) and 2nd-order MF model (blue diamonds). Solid lines: full 1st- (vellow) and 2nd-order MF theory (blue); black dashed lines: corresponding analytical approximations, Eq. (46).) The gray dashed line marks the border of the region of interest $\mu_0 \leq -wr_m/2$, where the hazard function at the fixed point, $\phi(\bar{h}_0)$, is convex. (b) Analogous to (a) but for a much larger network ($N = 5 \cdot 10^4$) with negligible, internally generated, finite-size noise but additional external noise, $\sigma_{\text{ext}}^2 = 1 \text{ mV}^2$. The connection probability is varied by different values of C as indicated. The yellow line corresponds to the common 1st-order MF model or mean-connectivity network. (c) Slope F_h vs. mean stimulus $\hat{\mu}_0$. Dotted line: p = 1 corresponding to the function Q, solid line: p = 0.7. Gray dashed line as in (a),(b) right. (d) Same for the product $\hat{\mu}_0 F_h$.

of Eq. (48) reveals a maximum at an intermediate connection probability

$$p_{\max} = \frac{2}{3} \left(1 + \frac{2\tau C}{-w\mu_0} \left(\beta^{-2} - \frac{\sigma_{\exp}^2}{4} \right) \right)$$

in the interval (0,1) if $-2/3 < \frac{2\tau C}{-w\mu_0} \left(\beta^{-2} - \sigma_{\text{ext}}^2/4\right) < 1/2$. In the limit of a steep single-neuron transfer function $(\beta \rightarrow \infty)$ and in the absence of external noise, the maximum is attained at $p_{\text{max}} = 2/3$. Such maximum is indeed verified by simulations (Fig. 7b, right). Furthermore, the variance $\operatorname{var}(\bar{h})$ approaches zero for $p \rightarrow 0$ and approaches a non-vanishing value for $p \rightarrow 1$.

In contrast to the 2nd-order theory, the 1st-order MF model cannot explain the strictly convex and non-monotonic behavior of the variances of the population rates and input potentials, respectively (Fig. 7b). In the second scenario with constant C, the 1st-order MF theory is obtained from the above formulas by fixing the ratio p/C and letting $p \rightarrow 1$. Thus, the common factor $\left(\frac{\tilde{\mu}op}{\tilde{\tau}C} + \hat{\sigma}_{ext}^2\right)$ in Eqs. (46) and (48) remains unchanged, whereas the second factor becomes independent of p. Because the common factor is an increasing linear function of p, the variances are thus also increasing linear functions, in line with simulations of the 1st-order mean-field model (but in contradiction to microscopic simulations and 2nd-order MF theory).

2. Dependence on the mean stimulus strength

We started our paper with network simulations pointing out marked quantitative and qualitative discrepancies of the meanconnectivity network with respect to the stimulus dependence of the rate variability (Fig. 1). In particular, we observed a strong overestimation of the rate variance by the meanconnectivity network compared to the original quenched network. Furthermore, for a noisy external stimulus and large network size (N = 50000), we observed a suppression of the rate variability by an increase in the mean stimulus strength in the quenched network. In contrast, the mean-connectivity network incorrectly predicted an increase in rate variability rather than a suppression. As we will see now, these observations can be well explained by our 2nd-order MF theory.

To this end, we study two opposing scenarios: first, we consider a small ("mesoscopic") network without noise in the external stimulus ($\sigma_{ext} = 0$). In this case, the rate variability solely originates intrinsically from the finite-size fluctuations. Second, we consider a large ("macroscopic") network with a noisy external stimulus ($\sigma_{ext} > 0$). In this case, the rate variability is purely externally generated. In both cases, we find that the first-order MF theory largely overestimates the variance of the population rate in the quenched and annealed network, whereas the 2nd-order MF theory, Eq. (43), correctly predicts the magnitude of the rate variance (Fig. 8a,b). Because the 1st-order MF model is the exact mean-field model for the mean-connectivity network, this large deviation explains our observation in Fig. 1. Furthermore, in the two different scenarios, we find a different monotonicity of the rate variance if we restrict ourselves to the biologically interesting range where the fixed point is in the convex part of the sigmoidal hazard function, $\bar{h}_0 < 0$ and $r_0 < r_m/2$, i.e. below the inflection point of $\phi(h)$ (cf. Fig. 3a). The upper boundary of this range given by the inflection point corresponds to a single-neuron firing rate at half maximum, $r_m/2$, and hence an external stimulus $\mu_0^* = -wr_m/2$ (Fig. 8a,b right, dashed vertical line). In the first scenario without external noise, the finite-size induced rate variability increases monotonically for $\mu_0 < \mu_0^*$ for both 1st- and 2nd-order MF theory. In contrast, in the second scenario with external noise and without finite-size fluctuations, the variance of the rate shows a non-monotonic behavior with a maximum below μ_0^* if p < 1.

These observations can be understood analytically using the approximation Eq. (46), (47). Let us first discuss how the factor F_h in Eq. (46) depends on $\hat{\mu}_0$. For p = 1, i.e. in the 1storder MF theory, \hat{F}_h is exactly given by the function $Q(\hat{\mu}_0)$ displayed by the dotted line in Fig. 8c. This function, and thus \hat{F}_h , grows monotonotically for $\hat{\mu}_0 > 0$ until it reaches a maximum at $\hat{\mu}_0^* = -\mu_0^*/(wr_m) = 0.5$, which corresponds to the upper bound $r_{\rm m}/2$ of the biologically relevant dynamical range. For p < 1, the factor F_h is the quotient of the symmetric function Q and an increasing function, which shifts the maximum of \hat{F}_h to a value < 1/2 (Fig. 8c, solid line). Hence, \hat{F}_h exhibits a non-monotonic behavior in the biologically relevant range $0 < \mu_0 < \mu_0^*$. Besides the shift of the maximum to smaller values of the mean stimulus strength $\hat{\mu}_0$, the maximum also decreases when the connection probability is lowered from p = 1 to smaller values. This decrease is a consequence of dividing the function $Q(\hat{\mu}_0)$ by the square-root term that is strictly larger than one for p < 1. Importantly, \hat{F}_h can decrease by a large factor if the single-neuron transfer function is steep, i.e. $\hat{\beta} \gg 1$, which explains the drastic decrease of the rate variance from 1st- to 2nd-order MF theory (Fig. 8a,b, where $\hat{\beta} = 500$).

The behavior of \hat{F}_h just described explains the variance of the population rate for an infinitely large network with external noise (second scenario, Fig. 8b). In fact, the prefactor of \hat{F}_h in Eq. (46) becomes independent of $\hat{\mu}_0$ in the limit when $N = C/p \rightarrow \infty$, and hence the dependence on $\hat{\mu}_0$ is fully captured by \hat{F}_h . In particular, the stark difference in magnitude of var(r) between the 1st- and 2nd-order MF prediction, as well as the qualitative response of the rate variability (suppression or amplification) to increasing stimuli, is explained by how the monotonicity and magnitude of the slope of the population transfer function \hat{F}_h varies with p.

In contrast, in the first scenario when $\sigma_{\text{ext}} = 0$ and N = C/p is finite, Eq. (46) tells us that the variance of the population rate is proportional to

$$\hat{\mu}_0 \hat{F}_h = \frac{\hat{\mu}_0}{\sqrt{1 + \frac{\hat{\beta}^2 (1-p)}{2\hat{\tau}C} \hat{\mu}_0}} Q(\hat{\mu}_0).$$

The right-hand side is a product of a monotonically increasing function of $\hat{\mu}_0$ and the function $Q(\hat{\mu}_0)$, which is maximized at

 $\hat{\mu}_0^* = 1/2$. Therefore, the maximum of $\hat{\mu}_0 \hat{F}_h$ is shifted to values $\hat{\mu}_0 > 1/2$ for all $p \in [0, 1]$ (Fig. 8d). As a consequence, the variance of the population rate var(r) is a monotonically increasing function of the mean stimulus strength in the biologically relevant range $0 \le \hat{\mu}_0 \le 1/2$ (Fig. 8a right, $\mu_0 < 50$ mV), and thus an increase of the mean stimulus strength always increases the rate variability in the absence of external common noise (Fig. 8a left).

To keep the arguments simple, we studied here two clearcut cases – one with and another without external noise. According to our theory, Eq. (45), the rate variance is a sum of two terms, where each term on its own corresponds to one of these cases. In general, the theory also applies to the mixed case, where the rate variance is the sum of externally and intrinsically induced variability. For example, in the simulations of Fig. 1, we also compared two networks of size N = 1000and N = 50000 but the external common noise was present in both cases. Nonetheless, the mesoscopic case of N = 1000with external noise can still be qualitatively understood by the limit case $\sigma_{\text{ext}} = 0$ because the finite-size fluctuations dominate over the external noise.

V. CONCLUSIONS

In this work, we set out to derive a low-dimensional, stochastic population dynamics for random recurrent networks of spiking neurons that correctly accounts for fluctuations when the network is neither fully connected (p = 1) nor infinitely large $(N \rightarrow \infty)$. While such regime is biologically highly relevant, to our knowledge it has not yet been systematically investigated within the frameworks of previously developed mean-field theories [29, 51]. By deriving a low-dimensional, stochastic mean-field model that allows one to study specifically the combined population effects of quenched disorder and stochastic neuronal firing in spiking networks, we thus hope to close an important gap in the use cases of simplified descriptions of networks and to provide the foundations for a mean-field modeling framework of neural variability at the mesoscopic population level.

Based on a Poisson assumption for neural firing and a "dynamically annealed" description of the connectivity, we approximated the recurrent input to each neuron by its (population) mean, a coherent fluctuating part shared among all neurons, and an individually fluctuating contribution reflecting the connectivity disorder. Eventually, we obtained three coupled stochastic differential equations (SDEs) that describe the evolution of the mean input potential \bar{h} within the population, the input potential's variance σ^2 , as well as a colored noise ξ in the population rate that vanishes in both limits $N \to \infty$ and $p \to 1$. Our theory thus goes beyond previous, deterministic mean-field dynamics for first- and second-order cumulants [22] or pseudo-cumulants [55] that similarly captured the quenched variability of synaptic weights but assumed the macroscopic limit $N \to \infty$.

The analysis of the new "second-order" mean-field model showed that the disorder of the connectivity for p < 1 has a drastic effect on the neural variability, the response to timedependent stimuli, and the stability of the network in biologically relevant regimes. Specifically, our main findings with regard to the effects of a finite connectivity 0 andfinite network size $N < \infty$ are the following: (i) A finite variance in the input currents ($\sigma^2 > 0$) caused by the disorder of a non-full, random connectivity with finite C leads to an effective broadening of the neuron transfer function, but the stationary mean firing rate remains rather unaffected (Fig. 3). (ii) In the presence of synaptic delays, the stability properties of fixed points of the dynamics and locations of bifurcations can change considerably. In particular, we showed that the connectivity disorder can stabilize networks in regimes where a "first-order" mean-field theory corresponding to a mean-connectivity approximation would predict oscillatory dynamics (Fig. 4). (iii) The population-rate response to highfrequency stimuli and (iv) the variance of the population rate are significantly lower than predicted by a "first-order" meanfield theory that does not correctly account for the incoherent fluctuations caused by the connectivity disorder (Fig. 5 and 7). Moreover, (v) in the presence of shared noise due to common external input, an increase in the mean external input can actually lead to a *decrease* of the variance of the population rate (Figs. 1 and 8) – an effect which again is not captured in a "first-order" mean-field theory. Overall, we provided simple analytical explanations for the effects of finite connectivity and finite network size in terms of the slope of the population transfer function and the multiplicative character of the finite-size noise. In particular, we found that a decreasing connection probability p lowers the slope F_h through an increase of the membrane potential variance σ^2 and that, at finite network size N, the neural variability has a rate-dependent contribution that is proportional to the product of mean stimulus strength (or firing rate) and the slope of the transfer function (Fig. 8d, Eq. (45)).

To derive a simple mean-field model with fluctuations that highlights the effects of finite connectivity and finite size, we made several simplifications. In particular, we used (i) an annealed approximation of the quenched random connectivity, (ii) only one cell type (single population), and (iii) a first-order Poisson model of the spiking-neuron dynamics. In the following, we will briefly discuss these simplifications and possible extensions.

In our "dynamically annealed" approximation, we assumed that each presynaptic spike would be randomly distributed among all neurons according to the connection probability p, independent of the actual, fixed (quenched) synaptic connectivity matrix of a given network. A systematic comparison of our theory with simulations of quenched networks showed that our second-order mean-field theory captures reliably the behavior of quenched networks with respect to the main findings described above. Deviations of the annealed network (from the quenched network) and, correspondingly, the 2ndorder MF model, are noticeable for several statistics, though. However, these deviations were considerably smaller than the respective deviations of the mean-connectivity network and the corresponding 1st-order MF model used in previous studies [35, 49, 50]. The remaining deviations of the annealed network can most likely be attributed to the neglect of temporal

correlations in the recurrent input caused by the annealing, i.e. the incessant resampling of the connections in time. This explanation would be consistent with the underestimation of the low-frequency power by the annealed network which we observed in Fig. 6. For infinitely large, sparsely connected networks, a proper account of temporal correlations can, in principle, be obtained through a self-consistent treatment of the auto-correlations of the recurrent fluctuations [56, 68], which typically adds low-frequency power [56]. Such self-consistent theory would lead to colored noise rather than white noise for the incoherent fluctuations in Eq. (12). However, how to build a stochastic, dynamical population model for finite-size, random networks that accounts for temporal correlations in a selfconsistent manner is an open theoretical problem which goes beyond the scope of the present study.

Interestingly, perhaps not surprisingly, the comparison of our theory with simulations of "dynamically annealed" networks shows that it can be considered almost exact for the annealed case. One may ask whether a biological realization of such a "dynamically annealed" network, where our theory becomes exact, might actually exist. A possible origin of such randomly distributed spikes might be the probabilistic vesicle release: A fully connected network where synapses would transmit a spike with probability p would be an exact realization of the system we describe by our approximation. Thus, the annealed network model may also be regarded as a simple caricature of a spiking neural network with probabilistic synaptic transmission. Another type of annealing occurs in networks with full connectivity but dynamic synapses (shortterm synaptic plasticity). Recently, such dynamically changing networks have also been reduced to low-dimenensional Langevin dynamics based on a similar Poisson neuron model [69, 70].

For the sake of a systematic and thorough development of the theory, we here began with the case of one population corresponding to a single neuron type; especially, we focused on the case of an inhibitory network. However, our theory suggests a straightforward extension to multiple populations. Cortical networks typically consist of several cell types and are generally described as excitatory-inhibitory networks that are, in the simplest case, organized as two coupled populations. Rate models of such two-population E-I networks remain a popular tool to describe network dynamics up to this day, whether to characterize inhibition stabilization in cortical networks [71] or spatial patterns of oscillatory activity observed in motor cortex [50, 72]. It will certainly be worthwhile to investigate how the effects of finite connectivity and network size we began to explore here may shape the more complicated dynamics of networks that comprise more than one population. Furthermore, a multi-population extension would offer the possibility to model specific cortical microcircuits, such as canonical circuit models [73] or cortical column models with multiple interneuron types [43], on the mesoscopic scale, which accounts for fluctuations.

To develop the theory, we used Poisson neurons based on a one-dimensional dynamics for the membrane potential of each neuron. While a Poisson model neglects spike-history effects such as neuronal refractoriness, it allowed us to derive a low-dimensional, closed set of mean-field equations for the first- and second-order cumulants. We expect that in regimes where the neuron's firing is Poissonian, similar observations can be made in more biophysically grounded models of spiking networks. While this is beyond the scope of this work, it would certainly be worthwhile to systematically investigate the effects described here in simulations of such networks.

The main difference of our Poisson model compared to more realistic integrate-and-fire (IF) models is the absence of a reset mechanism (in fact, with reset, our model would be equivalent to a leaky IF model with escape noise [28, 30, 32, 74, 75]). While the resetting in IF neurons captures refractoriness, it significantly complicates the population dynamics. Specifically, the population dynamics of IF neurons is usually infinite dimensional, represented by partial differential equations [29, 75], integral equations [23], infinite systems of SDEs [61, 76], or, in the case of finite N, stochastic versions of these [32, 35, 61, 77]. In principle, it seems possible to carry over our theory to these population equations by again using the same annealed approximation. It would be interesting to study how well the annealed approximation works for neurons with refractoriness. One may speculate that similar second-order mean-field theories may provide a quantitatively correct account for networks e.g. of generalized integrate-and-fire neurons in the presence of finite connectivity where effective first-order mean-field theories have already proven to be quantitatively precise for fully connected networks [32, 35, 50].

For simulations and concreteness, we made some further specific modeling choices: For example, we studied random networks with fixed in-degree, i.e. an identical number of presynaptic connections, and chose a specific transfer function (or f - I curve) $\phi(h)$ that relates the instantaneous firing rate to the intensity or input current h. While our specific choice of the transfer function also enabled us to analytically evaluate the effect of a finite variance in the input currents on the population rate, the reduction of the spiking network dynamics to a low-dimensional system of coupled SDEs should remain valid for arbitrary choices of the transfer function, albeit probably more difficult to compute. The assumption of fixed in-degree allowed us to assume that the mean input current as well as the strengths of the fluctuating parts are identical among neurons, but in practice we observed that a purely random Erdös-Réniy network shows almost identical statistics as those described in the previous sections for networks with fixed in-degree.

Here, we showed how to derive low-dimensional stochastic models that describe fluctuations in a complex, biological model system, namely a finite-size spiking neural networks with non-full connectivity. Already in the simple case of one population and Poissonian spiking, we found highly non-trivial effects such as the non-monotonic dependence of the variance of the recurrent input on connection probability (Fig. 7b) and the weak suppression of variability by external stimuli (Fig. 8b). Thus, even though a single population and Poisson spiking dynamics may be too simple to quantitatively model certain phenomena in cortex (such as a stronger suppression of variability [39]), the fact that our 2nd-order MF theory explains non-trivial fluctuation effects demonstrates the power of the approach as a proof of principle. Importantly, with the biological extensions discussed above, we believe that our theory may pave the way for a useful modeling framework for variability in cortical circuits at the mesoscopic population level.

VI. ACKNOWLEDGMENT

We are grateful to Jakob Stubenrauch for useful comments on the manuscript.

Appendix A: Mesoscopic dynamics in the presence of heterogeneous external drive

An additional source of disorder in the microscopic models is the heterogeneity in the parameters. For Eq. (1), we include heterogeneity in the external drive as follows:

$$\mu_{\text{ext},i}(t) = \mu(t) + \hat{\mu}_i. \tag{A1}$$

The random variables $\hat{\mu}_i$ for each neuron *i* have mean zero and are independently and identically distributed with probability density $\rho(\hat{\mu})$. They can be interpreted as heterogeneity of the external drives, the resting potentials or the thresholds ϑ of the different neurons. Importantly, the variables $\hat{\mu}_i$ are constant in time, and thus represent quenched disorder. The derivation of the microscopic annealed network (12) does not explicitly depend on the value of $\mu(t)$ and we therefore obtain the same results with $\mu(t)$ replaced with Eq. (A1). Specifically, Eq. (15) is modified as follows

$$\tau \frac{\mathrm{d}h_i}{\mathrm{d}t} = -h_i + \hat{\mu}_i + f(t) + g(t)\zeta_i(t).$$

Splitting off the heterogeneous part from $h_i(t)$ yields a new variable $x_i(t) = h_i(t) - \hat{\mu}_i$ that obeys

$$\tau \dot{x}_i = -x_i + f(t) + g(t)\zeta_i(t).$$

The last equation is of the same form as Eq. (15) and can be dealt with accordingly. For the self-consistent closure, the probability density of the input potentials h_i is needed for the calculation of $\langle \phi(h_i(t)) \rangle$. At each time point we assume $h_i(t) = x_i(t) + \hat{\mu}_i$ is the sum of two independent random variables and thus its density is the convolution $g_{\bar{x},\sigma_x^2} * \rho$. For simplicity, we assume that the heterogeneity is Gaussian distributed $\hat{\mu}_i \sim \mathcal{N}(0, \sigma_{\mu}^2)$ with a given fixed variance σ_{μ}^2 . In this case, we have a Gaussian probability for the distribution h_i with mean $\bar{x}(t)$ and variance $\sigma_x^2 + \sigma_{\mu}^2$. We therefore have

$$\begin{aligned} \tau \frac{\mathrm{d}\bar{x}}{\mathrm{d}t} &= -\bar{x} + \mu(t) + w \left[r(t-d) + \sqrt{\frac{r(t-d)}{N}} \eta(t) \right] \\ \tau \frac{\mathrm{d}\sigma_x^2}{\mathrm{d}t} &= -2\sigma_x^2 + \frac{w^2 \left(1-p\right)}{\tau p N} r(t-d) \\ \tau \frac{\mathrm{d}\xi}{\mathrm{d}t} &= -\xi + \sqrt{2\tau G\left(\bar{x}(t), \sigma^2(t)\right)} \zeta(t) \end{aligned}$$

where

$$r(t) = F\left(\bar{x}(t), \sigma^2(t)\right) + \frac{1}{\sqrt{N}}\xi(t)$$
$$\sigma^2(t) = \sigma_x^2(t) + \sigma_\mu^2.$$

In other words, we obtain the very same mean-field dynamics as in the case without heterogeneity, Eq. (20), but with an increased variance σ^2 . See also the classical mean-field theory of Amari [22] for a similar treatment of heterogeneity.

Appendix B: Estimation of the rate variance from population activities

The variance of the population firing rate is not directly accessible from measurements in real biological networks. However, we can esimate the rate variance by the variance of the emphirical population activity. We assume that in small time bins $(t, t + \Delta t]$, neurons fire spikes independently with conditional intensities $\lambda_i(t)$ given the past. Then, from the law of total variance and the conditionally Poisson statistics, we have for the variance of the total number of spikes in a time bin

$$\operatorname{var}(\Delta Z) = \langle \operatorname{var}(\Delta Z | \{\lambda_i\}) \rangle + \operatorname{var}(\langle \Delta Z | \{\lambda_i\} \rangle)$$
$$= N \langle r \rangle \Delta t + N^2 \operatorname{var}(r) \Delta t^2,$$

where $r(t) = N^{-1} \sum_{i=1}^{N} \lambda_i(t)$ is the population firing rate and $\langle \cdot \rangle$ denotes the trial average. Hence,

$$\begin{aligned} \operatorname{var}(r) &= \frac{\operatorname{var}(\Delta Z)}{N^2 \Delta t^2} - \frac{\langle r \rangle}{N \Delta t} \\ &= \operatorname{var}(A_N(t, \Delta t)) - \frac{\langle A_N(t, \Delta t) \rangle}{N \Delta t} \end{aligned}$$

where we used the definition of the empirical population activity, Eq. (3). For stationary data, the trial average can be replaced by the time average.

Appendix C: Condition for the validity of the firing-rate approximation (26)

In Eq. (26), a simple approximation for the fixed-point firing rate is given, namely $r_0 \approx -\mu_0/w$, which we used for an inhibitory network (w < 0) in the non-saturated regime ($0 < \mu_0 \le -wr_m$). This approximation corresponds to the solution of the 1st-order MF model in the limit $\beta \to \infty$. We argued that under certain sufficient conditions, the approximation is good for the 2nd-order MF model as well. Here we provide a justification for these conditions. For biological reasons, we make a slightly stronger assumption on the working point of our system: we assume that the fixed point is in the convex part of the transfer function, i.e. $h_0 < 0$ and $r_0 < r_m/2$, as drawn in Fig. 3.

For the firing-rate approximation to be applicable also to the 2nd-order MF model, we require that the relative difference

between the firing rates of the 1st- and 2nd-order MF theory is small:

$$r_0^{2nd} - r_0^{1st} \ll r_0^{1st}.$$
 (C1)

Here, $r_0^{1\text{st}} = -\mu_0/w$ and $r_0^{2\text{nd}} = F(h_0, \sigma_0^2)$. Furthermore, we have used that $r_0^{2\text{nd}} > r_0^{1\text{st}}$ because of our assumption, $h_0 < 0$ (Fig. 3a). To estimate the left hand side of (C1) from above, we want to obtain an upper bound approximation of the rate $r_0^{2\text{nd}} = (h_0 - \mu_0)/w$. To this end, we lower bound the input potential h_0 (note that w < 0). Because h_0 is expected to be on the order of the width σ_0 of the transfer function, we write the lower bound as $-a\sigma_0 < h_0$, where a is a positive number that is still unknown. This lower bound of h_0 yields an upper bound estimate for the mean firing rate: $r_0^{2\text{nd}} < (-a\sigma_0 - \mu_0)/w$. Hence, the condition (C1) is surely met if

$$\frac{-a\sigma_0 - \mu_0}{w} - \frac{-\mu_0}{w} = -a\frac{\sigma_0}{w} \ll -\frac{\mu_0}{w},$$

hence $a\sigma_0 \ll \mu_0$. Unfortunately, we do not know the exact value of σ_0 for the 2nd-order model. However, if our approximation holds true, we can self-consistently use our approximation for σ_0^2 , Eq. (27), and the condition becomes

$$a\sqrt{\frac{-w(1-p)}{2\tau\mu_0 C}} \ll 1.$$
 (C2)

The larger the value of a, the more conservative this sufficient condition becomes. However, to obtain a less conservative condition, we could lower the value of a, as long as $h_0 > -a\sigma_0$. We check for this latter condition in the same manner as above, by self-consistently using the approximation of h_0 in Eq. (28):

$$h_0 \approx \Phi^{-1} \left(\frac{-\mu_0}{w r_{\rm m}} \right) \sigma_0 > -a \sigma_0 ,$$

where we have used the limit $\beta \to \infty$. Eventually, this leads to

$$\frac{-\mu_0}{wr_{\rm m}} > \Phi(-a). \tag{C3}$$

The last step was possible, because Φ is strictly monotonously increasing. If there exists a > 0 such that both Eq. (C2) and Eq. (C3) are fulfilled, we consider the approximations for the fixed points to be valid. For the parameter choice used in our simulations, a = 2 was sufficient. The numerical evaluation of the error function yields $\Phi(-2) \approx 0.023$. Fixing a = 2yields the sufficient condition reported in Sec. IV A.

Appendix D: Partial derivatives of the function F and approximations thereof

Using the definition of the function F defined in Eq. (22), we find for the partial derivative with respect to h at the fixed point (\bar{h}_0, σ_0^2)

$$F_{h} := \frac{\partial F}{\partial h} (\bar{h}_{0}, \sigma_{0}^{2} + \sigma_{\mu}^{2}) = \frac{r_{\mathrm{m}} \beta \exp\left[-\frac{\beta^{2} h_{0}^{2}}{2\left(1 + \beta^{2}(\sigma_{0}^{2} + \sigma_{\mu}^{2})\right)}\right]}{\sqrt{2\pi \left(1 + \beta^{2}(\sigma_{0}^{2} + \sigma_{\mu}^{2})\right)}}.$$

Note that $\sigma_{\mu}^2 = 0$ in the absence of heterogeneity of μ_i . Similarly, for the partial derivative with respect to the second argument, we find

$$F_{\sigma} := \frac{\partial F}{\partial \sigma^2} (\bar{h}_0, \sigma_0^2 + \sigma_{\mu}^2) \\ = \frac{-r_{\rm m} \beta^3 h_0 \exp\left[-\frac{\beta^2 h_0^2}{2(1+\beta^2(\sigma_0^2 + \sigma_{\mu}^2))}\right]}{2\sqrt{2\pi}\sqrt{1+\beta^2(\sigma_0^2 + \sigma_{\mu}^2)^3}}$$

The approximate expressions Eq. (28) and Eq. (27) for h and σ^2 at the fixed point can be used for further simplifications of the partial derivatives:

$$F_h \approx \frac{r_{\rm m}\beta Q\left(-\frac{\mu_0}{r_{\rm m}w}\right)}{\sqrt{2\pi}\sqrt{1+\beta^2\left(\sigma_0^2+\sigma_\mu^2\right)}}$$

where σ_0^2 and the function Q are given by Eq. (27) and Eq. (39), respectively. Analogously, we can approximate the partial derivative with respect to σ^2 at the fixed point:

$$F_{\sigma} \approx \frac{-r_{\rm m}\beta^2 \Phi^{-1} \left(\frac{-\mu_0}{r_{\rm m}w}\right) Q\left(\frac{-\mu_0}{r_{\rm m}w}\right)}{2\sqrt{2\pi} \left(1 + \beta^2 (\sigma_0^2 + \sigma_\mu^2)\right)}$$

Appendix E: Linear stability analysis

For the linear stability analysis, we consider the linearized system with no noise and constant $\mu(t) = \mu$ which reads

$$\tau \frac{\mathrm{d}}{\mathrm{d}t} \delta h(t) = -\delta h(t) + w \delta r(t-d)$$
(E1a)

$$\tau \frac{\mathrm{d}}{\mathrm{d}t} \delta \sigma^2(t) = -2\delta \sigma^2(t) + \frac{w^2(1-p)}{\tau C} \delta r(t-d) \quad \text{(E1b)}$$

$$\delta r(t) = F_h \delta h(t) + F_\sigma \delta \sigma^2(t), \qquad (E1c)$$

where we neglected the variable ξ as it decays exponentially. To solve the linearized system, we insert the exponential ansatz

$$\delta h(t) = \hat{h}(\lambda)e^{\lambda t}$$
$$\delta \sigma^2(t) = \hat{\sigma}(\lambda)e^{\lambda t},$$

with a constant parameter λ . Solving Eq. (E1b) for $\hat{\sigma}$ yields

$$\hat{\sigma} = \frac{w^2 B F_h e^{-\lambda d}}{\lambda \tau + 2 - w^2 B F_\sigma e^{-\lambda d}} \hat{h}, \quad B := \frac{1 - p}{\tau C}.$$
 (E2)

We insert this expression for $\hat{\sigma}$ into the first equation (E1a) and obtain an equation for the eigenvalue

$$\lambda \tau = -1 + wF_h e^{-\lambda d} + \frac{\hat{F}_\sigma wF_h e^{-2\lambda d}}{2 + \lambda \tau - \hat{F}_\sigma e^{-\lambda d}}$$
(E3)

where we used the abbreviation $\hat{F}_{\sigma} := w^2 B F_{\sigma}$. This equation depends on the fixed points through the functions F_h and

 F_{σ} . To find an oscillatory instability, we look for a Hopf bifurcation, at which $\lambda = i\omega$ for some real-valued frequency ω . Applying the condition on the complex-valued equation (E3) we obtain two conditions, one for the amplitude

$$1 = \frac{4 + \omega^2 \tau^2}{1 + \omega^2 \tau^2} \frac{w^2 F_h^2}{(2 - \cos(\omega d)\hat{F}_{\sigma})^2 + (\omega \tau + \sin(\omega d)\hat{F}_{\sigma})^2}$$
(E4)

and one for the phase

$$\arctan(\omega\tau) = \arctan\left(\frac{\omega\tau}{2}\right) + \arg(wF_h) - \omega d + 2\pi k$$
$$-\operatorname{atan2}\left(2 - \hat{F}_{\sigma}\cos(\omega d), \omega\tau + \hat{F}_{\sigma}\sin(\omega d)\right)$$

Both must be satisfied simultaneously. Here, atan2 is the twoargument arctangent and $k \in \mathbb{N}_0$. In the (d, w)-parameter space, these conditions correspond to a family of curves, one curve for each k. At the k-th curve an oscillatory perturbation with frequency ω_k becomes unstable. For the overall instability boundary it is sufficient that at least one mode becomes unstable. We found empirically that the overall instability boundary is given by the mode k = 0, whereas higher modes yield regions of instability that are contained already in the instability region given by k = 0. For the solution of both conditions a numerical solution of the fixed point equations is necessary. In Fig. 4b) we scan through each parameter point and solve the phase condition for ω . In the investigated parameter range we were always able to find a unique solution for the phase condition for each mode. We then determined the boundary in the (d, w)-space that separated the regions where the right-hand side of the amplitude condition, Eq. (E4), is larger and smaller than 1, respectively. The equations are more simple for the 1st-oder MF model where $F_{\sigma} = 0$. The simplified amplitude condition reads in this case

$$1 = \frac{w^2 F_h^2}{1 + \omega^2 \tau^2} \quad \Leftrightarrow \quad \omega = \pm \frac{\sqrt{w^2 F_h^2 - 1}}{\tau}.$$

This explicit solution for ω can be used in the phase condition to determine the value of the delay at instability boundary for a given coupling strength w:

$$d = \frac{1}{\omega} \left(-\arctan(\omega\tau) + \arg(wF_h) + 2\pi k \right).$$

Appendix F: Susceptibility matrix

Here, we present explicit expressions for the susceptibility matrix $\tilde{\chi}$ defined in Eq. (34). Using Cramer's rule we obtain

$$\tilde{\chi}_{11} = \frac{1}{\mathcal{D}} \left(i\omega + \frac{2}{\tau} - F_{\sigma}\hat{\beta}e^{-i\omega d} \right)$$
(F1a)

$$\tilde{\chi}_{12} = \frac{1}{\mathcal{D}} F_{\sigma} \hat{\alpha} e^{-i\omega d} \tag{F1b}$$

$$\tilde{\chi}_{13} = \frac{1}{\mathcal{D}} \hat{\gamma} \hat{\alpha} e^{-i\omega d} \frac{i\omega + \frac{2}{\tau}}{i\omega + \frac{1}{\tau}}$$
(F1c)

$$\tilde{\chi}_{21} = \frac{1}{\mathcal{D}} F_h \hat{\beta} e^{-i\omega d} \tag{F1d}$$

$$\tilde{\chi}_{22} = \frac{1}{\mathcal{D}} \left(i\omega + \frac{1}{\tau} - F_h \hat{\alpha} e^{-i\omega d} \right)$$
(F1e)

$$\tilde{\chi}_{23} = \frac{1}{\mathcal{D}} \hat{\gamma} \hat{\beta} e^{-i\omega d} \tag{F1f}$$

$$\tilde{\chi}_{31} = \tilde{\chi}_{32} = 0 \tag{F1g}$$

$$\tilde{\chi}_{33} = \frac{1}{\mathcal{D}} \left(i\omega + \frac{2}{\tau} \right) - \frac{1}{\mathcal{D}} F_{\sigma} \hat{\beta} e^{-i\omega d}$$
(F1h)

$$-\frac{1}{\mathcal{D}}\frac{i\omega+\frac{2}{\tau}}{i\omega+\frac{1}{\tau}}F_h\hat{\alpha}e^{-i\omega d}$$
(F1i)

with the abbreviations

$$\mathcal{D}(\omega) = \left(i\omega + \frac{1}{\tau}\right) \left(i\omega + \frac{2}{\tau}\right) - \left(i\omega + \frac{2}{\tau}\right) F_h \hat{\alpha} e^{-i\omega d} - \left(i\omega + \frac{1}{\tau}\right) F_\sigma \hat{\beta} e^{-i\omega d} \hat{\alpha} = \frac{w}{\tau}, \quad \hat{\beta} = \frac{w^2}{\tau^2} \left(\frac{1}{C} - \frac{1}{N}\right), \quad \hat{\gamma} = \frac{1}{\sqrt{N}}.$$

Appendix G: Variance and power spectral density in the stationary state

With the susceptibility we can express the power spectral density of the system. From Eq. (33), we obtain

$$\left\langle \tilde{\boldsymbol{X}}(\omega)\tilde{\boldsymbol{X}}^{*}(\omega)\right\rangle = \tilde{\chi}\left\langle \left(\mathrm{B}\,\tilde{\boldsymbol{\zeta}}+\tilde{\boldsymbol{M}}\right)\left(\tilde{\boldsymbol{\zeta}}^{*}\,\mathrm{B}^{*}+\tilde{\boldsymbol{M}}^{*}\right)\right\rangle \tilde{\chi}^{*}$$

and with the definition of the power spectral density matrix $S(\omega)$, Eq. (40),

$$S(\omega) = \tilde{\chi}(\omega) [S_M(\omega) + BB^*] \tilde{\chi}^*(\omega).$$

Here, S_M is the 3 × 3 spectral density matrix of the external stimulus. In our case, the external stimulus only acts on the \bar{h} -variable and thus $(S_M)_{ij} = S_{\mu\mu}\delta_{1i}\delta_{1j}$ has only one non-zero entry with the power spectral density $S_{\mu\mu}(\omega)$ of the Gaussian stimulus $\mu(t)$. Explicitly,

$$\begin{split} \mathbf{S}_{ij}(\omega) &= \frac{1}{\tau^2} \left[S_{\mu\mu}(\omega) + \frac{w^2 r_0}{N} \right] \tilde{\chi}_{i1}(\omega) \tilde{\chi}^*_{j1}(\omega) \\ &+ \frac{2G \left(\bar{h}_0, \sigma^2_{h,0} \right)}{\tau N} \tilde{\chi}_{i3}(\omega) \tilde{\chi}^*_{j3}(\omega). \end{split}$$

In the main text, we used this expression to compute the power spectrum of the population activity A(t), given by Eq. (42). Taking the limit $p \rightarrow 1$ yields an explicit expression for the corresponding power spectrum of the 1st-order MF theory

$$S_{AA} \approx \frac{r_0}{N} + \frac{S_{\mu\mu} + w^2 \frac{r_0}{N} \phi_h^2 + 2w \frac{r_0}{N} \phi_h (1 - \cos(\omega d) w \phi_h)}{(\omega \tau - \sin(\omega d) w \phi_h)^2 + (1 - \cos(\omega d) w \phi_h)^2}.$$
(G1)

Here, ϕ_h is the partial derivative $\partial \phi / \partial h$ at the fixed point.

In principle, the stationary variance of h(t) can be computed from the integral over the power spectral density $S_{11}(\omega)$. However, the integral is difficult to solve for nonzero transmission delay and general spectral statistics $S_M(\omega)$ of the stimulus $\mu(t)$. If we restrict ourselves to the case of zero delay, d = 0, and a Gaussian white noise stimulus, $\mu(t) = \mu_0 + \sqrt{\tau \sigma_{ext}^2} \hat{\zeta}(t)$, the linearized sytem can be rewritten as

$$\frac{\mathrm{d}}{\mathrm{d}t}\boldsymbol{X} = \Gamma \,\boldsymbol{X} + \begin{pmatrix} \sqrt{\frac{\sigma_{\mathrm{ext}}^2}{\tau}}\hat{\zeta}(t) \\ 0 \\ 0 \end{pmatrix} + \begin{pmatrix} \frac{w}{\tau}\sqrt{\frac{r_0}{N}}\eta(t) \\ 0 \\ \sqrt{\frac{2G_0}{\tau}}\zeta(t) \end{pmatrix}, \quad (G2)$$

where $\Gamma = T + W$, or explicitly

$$\Gamma = \begin{pmatrix} \frac{w}{\tau} F_h - \frac{1}{\tau} & \frac{w}{\tau} F_\sigma & \frac{w}{\tau} \frac{1}{\sqrt{N}} \\ \frac{w^2(1-p)}{\tau^2 N p} F_h & \frac{w^2(1-p)}{\tau^2 N p} F_\sigma - \frac{2}{\tau} & \frac{w^2(1-p)}{\tau^2 N p} \frac{1}{\sqrt{N}} \\ 0 & 0 & -\frac{1}{\tau} \end{pmatrix}.$$

The two Gaussian white noise processes in the first component of Eq. (G2) can be lumped together into a single Gaussian white noise process:

$$\frac{\mathrm{d}}{\mathrm{d}t}\boldsymbol{X} = \Gamma \,\boldsymbol{X} + \hat{\mathrm{B}} \begin{pmatrix} \eta(t) \\ \zeta(t) \end{pmatrix}, \quad \hat{\mathrm{B}} = \begin{pmatrix} \sqrt{\frac{w^2}{\tau^2}} \frac{r_0}{N} + \frac{\sigma_{\mathrm{ext}}^2}{\tau} & 0 \\ 0 & 0 \\ 0 & \sqrt{\frac{2G_0}{\tau}} \end{pmatrix}$$

This is a three-dimensional Ornstein-Uhlenbeck process for which the stationary variance is known to be the solution of the linear system of equations [78] (Lyapunov equation)

$$\Gamma \,\sigma + \sigma \,\Gamma^T = -\hat{B}\hat{B}^T. \tag{G3}$$

Because the covariance matrix σ is symmetric, $\sigma_{ij} = \sigma_{ji}$, we have a system of six linear equations for the variables σ_{11}, σ_{12} , $\sigma_{13}, \sigma_{22}, \sigma_{23}$ and σ_{33} . For a systematic solution, we notice that the matrix equation (G3) reads in the position (3, 3)

$$2(\Gamma_{31}\sigma_{13} + \Gamma_{32}\sigma_{23} + \Gamma_{33}\sigma_{33}) = -\frac{2G_0}{\tau},$$

which reduces to $\sigma_{33} = G_0$ because $\Gamma_{31} = \Gamma_{32} = 0$ and $\Gamma_{33} = -1/\tau$. Therefore, we only need to solve a system of 5 linear equations

$$\begin{pmatrix} 2\Gamma_{11} & 2\Gamma_{12} & 0 & 2\Gamma_{13} & 0 \\ 0 & 2\Gamma_{21} & 2\Gamma_{22} & 0 & 2\Gamma_{23} \\ \Gamma_{21} & \Gamma_{11} + \Gamma_{22} & \Gamma_{12} & \Gamma_{23} & \Gamma_{13} \\ 0 & 0 & 0 & \Gamma_{11} + \Gamma_{33} & \Gamma_{12} \\ 0 & 0 & 0 & \Gamma_{21} & \Gamma_{22} + \Gamma_{33} \end{pmatrix} \begin{pmatrix} \sigma_{11} \\ \sigma_{12} \\ \sigma_{22} \\ \sigma_{13} \\ \sigma_{23} \end{pmatrix}$$
$$= \left(-\left(\frac{w^2}{\tau^2} \frac{r_0}{N} + \frac{\sigma_{\text{ext}}^2}{\tau}\right), \ 0, \ 0, \ -\Gamma_{13}G_0, \ -\Gamma_{23}G_0 \right)^T.$$

We can solve the last two equations separately as they only contain two variables:

$$\begin{pmatrix} \sigma_{13} \\ \sigma_{23} \end{pmatrix} = -\frac{G_0}{D} \begin{pmatrix} \Gamma_{13}(\Gamma_{22} + \Gamma_{33}) - \Gamma_{23}\Gamma_{12} \\ -\Gamma_{13}\Gamma_{21} + \Gamma_{23}(\Gamma_{11} + \Gamma_{33}) \end{pmatrix}, \quad (G4)$$

where $D = (\Gamma_{11} + \Gamma_{33})(\Gamma_{22} + \Gamma_{33}) - \Gamma_{21}\Gamma_{12}$. This equation gives us the values of σ_{13} and σ_{23} . The remaining three equations of the form

$$\begin{pmatrix} 2\Gamma_{11} & 2\Gamma_{12} & 0\\ 0 & 2\Gamma_{21} & 2\Gamma_{22}\\ \Gamma_{21} & \Gamma_{11} + \Gamma_{22} & \Gamma_{12} \end{pmatrix} \begin{pmatrix} \sigma_{11}\\ \sigma_{12}\\ \sigma_{22} \end{pmatrix} \\ = \begin{pmatrix} -\frac{w^2}{\tau^2} \frac{r_0}{N} - \frac{\sigma_{\text{ext}}^2}{\tau}\\ 0\\ 0 \end{pmatrix} - \begin{pmatrix} 2\Gamma_{13}\sigma_{13}\\ 2\Gamma_{23}\sigma_{23}\\ \Gamma_{23}\sigma_{13} + \Gamma_{13}\sigma_{23} \end{pmatrix}$$

have the solution

$$\begin{pmatrix} \sigma_{11}, \ \sigma_{12}, \ \sigma_{22} \end{pmatrix}^{T} = \frac{-2}{\tau E} \begin{pmatrix} \frac{w^{2} r_{0}}{\tau N} + \sigma_{\text{ext}}^{2} \end{pmatrix} \begin{pmatrix} \Gamma_{12} \Gamma_{21} - \Gamma_{22} (\Gamma_{11} + \Gamma_{22}) \\ \Gamma_{22} \Gamma_{21} \\ -\Gamma_{21}^{2} \end{pmatrix} \\ - \frac{4 \sigma_{13}}{E} \begin{pmatrix} \Gamma_{13} [\Gamma_{21} \Gamma_{12} - \Gamma_{22} (\Gamma_{11} + \Gamma_{22})] + \Gamma_{23} \Gamma_{12} \Gamma_{22} \\ \Gamma_{13} \Gamma_{22} \Gamma_{21} - \Gamma_{23} \Gamma_{11} \Gamma_{22} \\ -\Gamma_{13} \Gamma_{21}^{2} + \Gamma_{23} \Gamma_{11} \Gamma_{21} \end{pmatrix} \\ - \frac{4 \sigma_{23}}{E} \begin{pmatrix} -\Gamma_{23} \Gamma_{12}^{2} + \Gamma_{13} \Gamma_{12} \Gamma_{22} \\ \Gamma_{23} \Gamma_{11} \Gamma_{12} - \Gamma_{13} \Gamma_{11} \Gamma_{22} \\ -\Gamma_{23} [\Gamma_{11} (\Gamma_{11} + \Gamma_{22}) - \Gamma_{21} \Gamma_{12}] + \Gamma_{13} \Gamma_{11} \Gamma_{21} \end{pmatrix}$$
(G5)

with the determinant of the remaining 3×3 subsystem

$$E = -4(\Gamma_{11}\Gamma_{22} - \Gamma_{12}\Gamma_{21})(\Gamma_{11} + \Gamma_{22})$$

The formula for the variances of the mean input potential hand the firing rate r can be more easily interpreted if some further, heuristic simplifications are made. For the parameters used in our study, we observed that Γ_{11} is typically much larger than Γ_{22} , so that we could safely replace the terms $\Gamma_{11} + \Gamma_{22}$ in Eq. (G5) by Γ_{11} . Likewise, we found that the terms proportional to σ_{13} and σ_{23} in Eq. (G5) are small enough to be ignored. These simplifications yield

$$\sigma_{11} \approx \frac{1}{2} \frac{1}{(1 - wF_h)} \left(\frac{w^2 r_0}{\tau N} + \sigma_{\text{ext}}^2 \right)$$

To simplify even further, we note that in our simulations we have that $|w|F_h \gg 1$, and hence

$$\sigma_{11} \approx \left(\frac{-wr_0}{2\tau N} + \frac{\sigma_{\text{ext}}^2}{-2w}\right) \frac{1}{F_h} \tag{G6}$$

As mentioned in the main text, the variance of the population firing rate is dominated by the first term in Eq. (43), so that we obtain from Eq. (G6)

$$\operatorname{var}(r) \approx \left(\frac{-wr_0}{2\tau N} + \frac{\sigma_{\operatorname{ext}}^2}{-2w}\right) F_h. \tag{G7}$$

The variance in the 1st-order MF theory can be either calculated directly from Eq. (20) or can be understood as a special case of the formulas Eqs. (G6), (G7) for $p \rightarrow 1$ while keeping N constant. The variances for the mean input and the population firing rate read

$$\begin{split} \mathrm{var}(\bar{h}) &= \frac{1}{2(1 - w\phi_h)} \left(\frac{w^2 r_0}{\tau N} + \sigma_{\mathrm{ext}}^2 \right), \\ \mathrm{var}(r) &= \frac{1}{2(1 - w\phi_h)} \left(\frac{w^2 r_0}{\tau N} + \sigma_{\mathrm{ext}}^2 \right) \phi_h^2, \end{split}$$

where ϕ_h is the derivative of the single neuron transfer function at the fixed point. For the the approximation $1 - w\phi_h \approx -w\phi_h$ we arrive at the same form as Eqs. (G6), (G7).

Appendix H: Calculation of the functions F and G

For the evaluation of Eq (22) and Eq (23) for the given nonlinearity (2), we use two known formulas [79] for Gaussian integrals

$$\int_{-\infty}^{\infty} \Phi(a+bx)g_{0,1}(x)dx = \Phi\left(\frac{a}{\sqrt{1+b^2}}\right)$$
$$\int_{-\infty}^{\infty} \Phi^2(a+bx)g_{0,1}(x)dx = \Phi\left(\frac{a}{\sqrt{1+b^2}}\right)$$
$$-2T\left(\frac{a}{\sqrt{1+b^2}}, \frac{1}{\sqrt{1+2b^2}}\right)$$

with $a, b \in \mathbb{R}$ and $g_{0,1}$ being the standard normal distribution. Here,

$$T(h,a) = g_{0,1}(h) \int_0^a \frac{g_{0,1}(hx)}{1+x^2} \, dx$$

is the Owen's T-function. For our integral (22) we need to substitute $x:=(h-\bar{h})/\sigma$

$$\begin{split} F(\bar{h},\sigma^2) &= r_{\rm m} \int_{-\infty}^{\infty} \frac{\Phi\left(\beta(h-\vartheta)\right)}{\sqrt{2\pi\sigma^2}} \exp\left[-\frac{(h-\bar{h})^2}{2\sigma^2}\right] \,\mathrm{d}h \\ &= \frac{r_{\rm m}}{2\pi} \int_{-\infty}^{\infty} \Phi\left(\beta(\bar{h}-\vartheta) + \beta\sqrt{\sigma^2}x\right) e^{-x^2/2} \,\mathrm{d}x \\ &= r_{\rm m} \Phi\left(\frac{\beta(\bar{h}-\vartheta)}{\sqrt{1+\beta^2\sigma^2}}\right). \end{split}$$

The second integral (23) is computed analogously using the same substitution.

Appendix I: There exists exactly one fixed point for inhibitory networks with positive external drive

For the calculation of the fixed points of the 2nd-order model (20), we set the left-hand side to zero and solve the following equations for h_0 :

$$r_0 = F(h_0, \sigma^2(h_0)) = r_{\rm m} \Phi\left(\frac{\beta h_0}{\sqrt{1 + \alpha(\mu_0 - h_0)}}\right) =: F(h_0)$$
(11)

$$r_0 = \frac{h_0 - \mu_0}{w} \tag{12}$$

where $\alpha = -\beta^2 w(1-p)/(2\tau C) > 0$. The strictly positive non-linear function F in Eq. (I1) can only intersect the linear function in Eq. (I2) at a value for the fixed point $h_0 < \mu_0$. Equation (I2) represents a strictly monotonically decreasing function of h_0 for w < 0. We will show that for $\mu_0 > 0$, the function F(h) is strictly monotonically increasing for $h_0 < \mu_0$. The consequence is that both graphs will only intersect at exactly one point providing a unique fixed point. The proof for the monotonicity of $F(h_0)$ is straightforward. The derivative is given by the derivative of the error-function:

$$\frac{\mathrm{d}F}{\mathrm{d}h} = \frac{r_{\mathrm{m}}\beta \exp\left[-\frac{\beta^{2}h^{2}}{1+\alpha(\mu_{0}-h)}\right]}{\sqrt{2\pi}\sqrt{1+\alpha(\mu_{0}-h)}} \left(1+\frac{1}{2}\cdot\frac{\alpha h}{1+\alpha(\mu_{0}-h)}\right).$$

The first factor is always positive, the remaining factor provides the condition for the monotonic increase:

$$\frac{2}{\alpha} + 2\mu_0 > h$$

For $\mu_0 > 0$, this condition is fulfilled because

$$\frac{2}{\alpha} + 2\mu_0 > \mu_0 > h.$$

Here, we used $\alpha > 0$, $\mu_0 > 0$ and $\mu_0 > h_0$. With F(h) > 0 strictly monotonically increasing and the linear function in Eq. (I1) monotonically decreasing there must be exactly one fixed point solution $h_0 \in]-\infty, \mu_0[$ with a positive firing rate $r_0 > 0$.

- M. B. Elowitz, A. J. Levine, E. D. Siggia, and P. S. Swain, Stochastic Gene Expression in a Single Cell, Science 297, 1183 (2002).
- [2] A. Raj and A. Van Oudenaarden, Nature, Nurture, or Chance: Stochastic Gene Expression and Its Consequences, Cell 135, 216 (2008).
- [3] S. Taheri-Araghi, S. Bradde, J. T. Sauls, N. S. Hill, P. A. Levin, J. Paulsson, M. Vergassola, and S. Jun, Cell-Size Control and Homeostasis in Bacteria, Current Biology 25, 385 (2015).
- [4] S. Arzash, I. Tah, A. J. Liu, and M. L. Manning, Rigidity of epithelial tissues as a double optimization problem (2024), arXiv:2312.11683 [cond-mat.soft].
- [5] A. A. Faisal, L. P. J. Selen, and D. M. Wolpert, Noise in the nervous system, Nat Rev Neurosci 9, 292 (2008).
- [6] B. Doiron, A. Litwin-Kumar, R. Rosenbaum, G. K. Ocker, and K. Josić, The mechanics of state dependent neural correlations, Nat. Neurosci. 19, 383 (2016).
- [7] L. Waschke, N. A. Kloosterman, J. Obleser, and D. D. Garrett, Behavior needs neural variability, Neuron 109, 751 (2021).
- [8] T. B. Kepler and T. C. Elston, Stochasticity in transcriptional regulation: Origins, consequences, and mathematical representation, Biophys. J. 81, 3116 (2001).
- [9] T. Guérin, J. Prost, and J.-F. Joanny, Bidirectional motion of motor assemblies and the weak-noise escape problem, Phys. Rev. E 84, 041901 (2011).
- [10] C. Touya, T. Schwalger, and B. Lindner, Relation between cooperative molecular motors and active brownian particles, Phys. Rev. E 83, 051913 (2011).
- [11] R. Ma, G. S. Klindt, I. H. Riedel-Kruse, F. Jülicher, and B. M. Friedrich, Active phase and amplitude fluctuations of flagellar beating, Phys. Rev. Lett. **113**, 048101 (2014).
- [12] A. Dobrinevski and E. Frey, Extinction in neutrally stable stochastic lotka-volterra models, Phys. Rev. E 85, 051903 (2012).
- [13] H. R. Wilson and J. D. Cowan, Excitatory and inhibitory interactions in localized populations of model neurons, Biophys. J. 12, 1 (1972).
- [14] Y. Lu and J. Rinzel, Firing rate models for gamma oscillations in ii and ei networks, J. Comput. Neurosci., 1 (2024).
- [15] D. Levenstein, G. Buzsáki, and J. Rinzel, Nrem sleep in the rodent neocortex and hippocampus reflects excitable dynamics, Nature Comm. 10, 1 (2019).
- [16] Y. Ahmadian, D. B. Rubin, and K. D. Miller, Analysis of the stabilized supralinear network, Neural Comput. 25, 1994 (2013).
- [17] K.-F. Wong and X.-J. Wang, A recurrent network mechanism of time integration in perceptual decisions, J. Neurosci. 26, 1314 (2006).
- [18] A. Shpiro, R. Moreno-Bote, N. Rubin, and J. Rinzel, Balance between noise and adaptation in competition models of perceptual bistability, J. Comput. Neurosci. 27, 37 (2009).
- [19] G. Dumont, G. Northoff, and A. Longtin, Linear noise approximation for oscillations in a stochastic inhibitory network with delay, Phys. Rev. E 90, 012702 (2014).
- [20] T. Kanashiro, G. K. Ocker, M. R. Cohen, and B. Doiron, Attentional modulation of neuronal variability in circuit models of cortex, Elife 6, e23978 (2017).
- [21] G. Hennequin, Y. Ahmadian, D. B. Rubin, M. Lengyel, and K. D. Miller, The dynamical regime of sensory cortex: Stable dynamics around a single stimulus-tuned attractor account for patterns of noise variability, Neuron 98, 846 (2018).

- [22] S.-I. Amari, Characteristics of random nets of analog neuronlike elements, IEEE Trans. Syst. Man Cybern. SMC-2, 643 (1972).
- [23] W. Gerstner, W. M. Kistler, R. Naud, and L. Paninski, Neuronal Dynamics: From Single Neurons to Networks and Models of Cognition (Cambridge University Press, Cambridge, 2014).
- [24] G. Deco, V. K. Jirsa, P. A. Robinson, M. Breakspear, and K. Friston, The dynamic brain: from spiking neurons to neural masses and cortical fields, PLoS Comput. Biol. 4, e1000092 (2008).
- [25] O. D. Faugeras, J. D. Touboul, and B. Cessac, A constructive mean-field analysis of multi population neural networks with random synaptic weights and stochastic inputs, Front. Comput. Neurosci. 3, 323 (2009).
- [26] S. El Boustani and A. Destexhe, A master equation formalism for macroscopic modeling of asynchronous irregular activity states, Neural Comput. 21, 46 (2009).
- [27] G. La Camera, The mean field approach for populations of spiking neurons, in *Computational Modelling of the Brain: Modelling Approaches to Cells, Circuits and Networks*, edited by M. Giugliano, M. Negrello, and D. Linaro (Springer International Publishing, Cham, 2022) pp. 125–157.
- [28] G. K. Ocker, Republished: Dynamics of stochastic integrateand-fire networks, Phys. Rev. X 13, 041047 (2023).
- [29] N. Brunel and V. Hakim, Fast global oscillations in networks of integrate-and-fire neurons with low firing rates, Neural Comput. 11, 1621 (1999).
- [30] W. Gerstner, Population dynamics of spiking neurons: Fast transients, asynchronous states, and locking, Neural Comput. 12, 43 (2000).
- [31] E. Montbrió, D. Pazó, and A. Roxin, Macroscopic description for networks of spiking neurons, Phys. Rev. X 5, 021028 (2015).
- [32] V. Schmutz, E. Löcherbach, and T. Schwalger, On a finite-size neuronal population equation, SIAM J. Appl. Dyn. Syst. 22, 996 (2023).
- [33] D. Q. Nykamp and D. Tranchina, A population density approach that facilitates large-scale modeling of neural networks: Analysis and an application to orientation tuning, J. Comput. Neurosci. 8, 19 (2000).
- [34] N. Cain, R. Iyer, C. Koch, and S. Mihalas, The computational properties of a simplified cortical column model, PLoS Comput. Biol. 12, e1005045 (2016).
- [35] T. Schwalger, M. Deger, and W. Gerstner, Towards a theory of cortical columns: From spiking neurons to interacting neural populations of finite size, PLoS Comput. Biol. 13, e1005507 (2017).
- [36] M. Augustin, J. Ladenbauer, F. Baumann, and K. Obermayer, Low-dimensional spike rate models derived from networks of adaptive integrate-and-fire neurons: Comparison and implementation, PLoS Comput. Biol. 13, e1005545 (2017).
- [37] S. Lefort, C. Tomm, J. C. F. Sarria, and C. C. H. Petersen, The excitatory neuronal network of the c2 barrel column in mouse primary somatosensory cortex, Neuron 61, 301 (2009).
- [38] H. Ko, S. B. Hofer, B. Pichler, K. A. Buchanan, P. J. Sjöström, and T. D. Mrsic-Flogel, Functional specificity of local synaptic connections in neocortical networks, Nature 473, 87 (2011).
- [39] M. M. Churchland, M. Y. Byron, J. P. Cunningham, L. P. Sugrue, M. R. Cohen, G. S. Corrado, W. T. Newsome, A. M. Clark, P. Hosseini, B. B. Scott, *et al.*, Stimulus onset quenches neural variability: a widespread cortical phenomenon, Nat. Neurosci.

13, 369 (2010).

- [40] A. René, A. Longtin, and J. H. Macke, Inference of a mesoscopic population model from population spike trains, Neural Comput. 32, 1448 (2020).
- [41] S. Wang, V. Schmutz, G. Bellec, and W. Gerstner, Mesoscopic modeling of hidden spiking neurons, Adv. Neural Inf. Process. Syst. 35, 23566 (2022).
- [42] C. Pozzorini, S. Mensi, O. Hagens, R. Naud, C. Koch, and W. Gerstner, Automated high-throughput characterization of single neurons by means of simplified spiking models, PLoS Comput Biol 11, e1004275 (2015).
- [43] Y. N. Billeh, B. Cai, S. L. Gratiy, K. Dai, R. Iyer, N. W. Gouwens, R. Abbasi-Asl, X. Jia, J. H. Siegle, S. R. Olsen, C. Koch, S. Mihalas, and A. Arkhipov", Systematic integration of structural and functional data into multi-scale models of mouse primary visual cortex, Neuron https://doi.org/10.1016/j.neuron.2020.01.040 (2020).
- [44] M. A. Buice, J. D. Cowan, and C. C. Chow, Systematic fluctuation expansion for neural network activity equations, Neural Comput. 22, 377 (2010).
- [45] M. A. Buice and C. C. Chow, Dynamic finite size effects in spiking neural networks, PLoS Comput. Biol. 9, e1002872 (2013).
- [46] G. Dumont, A. Payeur, and A. Longtin, A stochastic-field description of finite-size spiking neural networks, PLoS Comput. Biol. 13, e1005691 (2017).
- [47] S. Heesen and W. Stannat, Fluctuation limits for mean-field interacting nonlinear hawkes processes, Stochastic Processes and their Applications (2021).
- [48] V. V. Klinshov and S. Y. Kirillov, Shot noise in next-generation neural mass models for finite-size networks, Phys. Rev. E 106, L062302 (2022).
- [49] M. Deger, T. Schwalger, R. Naud, and W. Gerstner, Fluctuations and information filtering in coupled populations of spiking neurons with adaptation, Phys. Rev. E 90, 062704 (2014).
- [50] A. Kulkarni, J. Ranft, and V. Hakim, Synchronization, stochasticity, and phase waves in neuronal networks with spatiallystructured connectivity, Frontiers in Computational Neuroscience 14, 10.3389/fncom.2020.569644 (2020).
- [51] N. Brunel, Sparsely connected networks of spiking neurons, J. Comput. Neurosci. 8, 183 (2000).
- [52] C. van Vreeswijk and H. Sompolinsky, Chaos in neuronal networks with balanced excitatory and inhibitory activity, Science 274, 1724 (1996).
- [53] A. Renart, R. Moreno-Bote, X.-J. Wang, and N. Parga, Meandriven and fluctuation-driven persistent activity in recurrent networks, Neural Computat. 19, 1 (2007).
- [54] M. Helias, T. Tetzlaff, and M. Diesmann, The correlation structure of local neuronal networks intrinsically results from recurrent dynamics, PLoS Comput Biol 10, e1003428 (2014).
- [55] D. S. Goldobin, M. di Volo, and A. Torcini, Reduction methodology for fluctuation driven population dynamics, Phys. Rev. Lett. **127**, 038301 (2021).
- [56] S. Wieland, D. Bernardi, T. Schwalger, and B. Lindner, Slow fluctuations in recurrent networks of spiking neurons., Phys. Rev. E 92, 040901 (2015).
- [57] A. Sanzeni, M. H. Histed, and N. Brunel, Response nonlinearities in networks of spiking neurons, PLoS Comput. Biol. 16, e1008165 (2020).
- [58] N. Schieferstein, T. Schwalger, B. Lindner, and R. Kempter, Intra-ripple frequency accommodation in an inhibitory network model for hippocampal ripple oscillations, PLOS Comput. Biol. 20, e1011886 (2024).

- [59] L. Mazzucato, A. Fontanini, and G. La Camera, Dynamics of multistable states during ongoing and evoked cortical activity, J. Neurosci. 35, 8214 (2015).
- [60] P. Ekelmans, N. Kraynyukova, and T. Tchumatchenko, Targeting operational regimes of interest in recurrent neural networks, PLOS Computational Biology 19, e1011097 (2023).
- [61] M. Mattia and P. Del Giudice, Population dynamics of interacting spiking neurons, Phys. Rev. E 66, 051917 (2002).
- [62] B. Derrida and Y. Pomeau, Random networks of automata: a simple annealed approximation, Europhys. Lett. 1, 45 (1986).
- [63] R. Jolivet, A. Rauch, H. R. Lüscher, and W. Gerstner, Predicting spike timing of neocortical pyramidal neurons by simple threshold models, J. Comput. Neurosci. 21, 35 (2006).
- [64] N. J. Priebe and D. Ferster, Inhibition, spike threshold, and stimulus selectivity in primary visual cortex, Neuron 57, 482 (2008).
- [65] S. P. Muscinelli, W. Gerstner, and T. Schwalger, How single neuron properties shape chaotic dynamics and signal transmission in random neural networks, PLOS Comput. Biol. 15, e1007122 (2019).
- [66] N. Brunel and V. Hakim, Sparsely synchronized neuronal oscillations, Chaos 18 (2008).
- [67] E. Ledoux and N. Brunel, Dynamics of networks of excitatory and inhibitory neurons in response to time-dependent inputs, Front. Comput. Neurosci. 5, 25 (2011).
- [68] J. Kadmon and H. Sompolinsky, Transition to chaos in random neuronal networks, Phys. Rev. X 5, 041030 (2015).
- [69] V. Schmutz, W. Gerstner, and T. Schwalger, Mesoscopic population equations for spiking neural networks with synaptic short-term plasticity, J. Math. Neurosc. 10, 1 (2020).
- [70] B. Pietras, V. Schmutz, and T. Schwalger, Mesoscopic description of hippocampal replay and metastability in spiking neural networks with short-term plasticity, PLOS Comput. Biol. 18, 1 (2022).
- [71] A. Sanzeni, B. Akitake, H. C. Goldbach, C. E. Leedy, N. Brunel, and M. H. Histed, Inhibition stabilization is a widespread property of cortical networks, eLife 9, e54875 (2020).
- [72] L. Kang, J. Ranft, and V. Hakim, Beta oscillations and waves in motor cortex can be accounted for by the interplay of spatially structured connectivity and fluctuating inputs, Elife 12, e81446 (2023).
- [73] G. Hahn, A. Kumar, H. Schmidt, T. R. Knösche, and G. Deco, Rate and oscillatory switching dynamics of a multilayer visual microcircuit model, Elife 11, e77594 (2022).
- [74] L. Yu and T. O. Taillefumier, Metastable spiking networks in the replica-mean-field limit, PLoS Computational Biology 18, e1010215 (2022).
- [75] G. Dumont, J. Henry, and C. O. Tarniceriu, Oscillations in a Fully Connected Network of Leaky Integrate-and-Fire Neurons with a Poisson Spiking Mechanism, J. NonLin. Sci. 34, 18 (2024).
- [76] B. W. Knight, D. Manin, and L. Sirovich, Dynamical models of interacting neuron populations in visual cortex, in *Sympo*sium on Robotics and Cybernetics: Computational Engineering in Systems Applications, Vol. 54, edited by E. C. Geil (Lille, France: Cite Scientifique, 1996) pp. 4–8.
- [77] G. V. Vinci, R. Benzi, and M. Mattia, Self-consistent stochastic dynamics for finite-size networks of spiking neurons, Phys. Rev. Lett. 130, 097402 (2023).
- [78] H. Risken, *The Fokker-Planck Equation* (Springer, Berlin, 1984).
- [79] D. B. Owen, A table of normal integrals, Commun. Stat. -Simul. Comput. 9, 389 (1980).