FILTERED COLIMITS OF COMPLETE INTERSECTION ALGEBRAS.

DORIN POPESCU

In the memory of Lucian Bădescu.

ABSTRACT. This is mainly a small exposition on extensions of valuation rings.

Key words : immediate extensions, smooth morphisms, Henselian rings, complete intersection algebras, pseudo convergent sequences, pseudo limits. 2020 Mathematics Subject Classification: Primary 13F30, Secondary 13A18, 13F20, 13B40.

1. INTRODUCTION

After Zariski [25] a valuation ring V containing a field K of characteristic zero is a filtered union of its smooth K-subalgebras. When V contains a field K of positive characteristic this result may fail, as we believe. A possible support for this idea is given by Example 8.

Then we ask if V is a filtered union of its complete intersection K-subalgebras essentially of finite type. This could hold as our Theorem 19 hints. We remind that for a given complete intersection local ring (R, \mathfrak{m}, k) a minimal free resolution over R of the residue field k is described in [24, Theorem 4] and others as [3, Theorem 2.7], [4, §6], [7, Proposition 1.5.4], [8, §3], [5, Theorem 2.5], [11, Theorem 4.1]. As Oana Veliche asked, we wonder if a minimal free resolution of the residue field of V could be described using somehow our Theorem 19.

We owe thanks to Oana Veliche and the referee who hinted some misprints in our paper.

2. Henselian rings and smooth algebras

A Henselian local ring is a local ring (A, \mathfrak{m}) for which the Implicit Function Theorem hods, that is for every system of polynomials $f = (f_1, \ldots, f_r)$ of A[Y], $Y = (Y_1, \ldots, Y_N), N \ge r$ over A and every solution $y = (y_1, \ldots, y_N) \in A^N$ modulo \mathfrak{m} of f in A such that an $r \times r$ -minor M of Jacobian matrix $(\partial f_i / \partial Y_j)_{1 \le i \le r, 1 \le j \le N}$ satisfies $M(y) \notin \mathfrak{m}$ there exists a solution \tilde{y} of f = 0 in A with $y \equiv \tilde{y}$ modulo \mathfrak{m} .

An A-algebra B is smooth if B is a localization of an A-algebra of type $(A[Y]/(f))_M$, where $f = (f_1, \ldots, f_r)$ is a system of polynomials in $Y = (Y_1, \ldots, Y_N)$, $N \ge r$ over A and M is an $r \times r$ -minor of $(\partial f/\partial Y)$. Thus (A, \mathfrak{m}) is Henselian if for any smooth A-algebra B any A-morphism $B \to A/\mathfrak{m}$ can be lifted to an A-morphism $w: B \to A$. Thus w is a retraction of the canonical morphism $u: A \to B$, that is $wu = 1_A$. A retraction of $A \to B$ maps the solutions of a system of polynomials $g = (g_1, \ldots, g_s) \in A[Y]^s$ in B in solutions of g in A.

We remind that a filtered colimit is a limit indexed by a small category that is filtered (see [22, 002V] or [22, 04AX]). A filtered union is a filtered direct limit in which all objects are subobjects of the final colimit, so that in particular all the transition arrows are monomorphisms.

Proposition 1. Let $(A, \mathfrak{m}) \to (A', \mathfrak{m}')$ be a morhism of local rings with $A/\mathfrak{m} \cong A'/\mathfrak{m}'$, which is a filtered colimit of smooth A-algebras and $g \in A[Y]^s$ a system of polynomials which has solutions in A'. If (A, \mathfrak{m}) is Henselian, then g has also solutions in A.

Proof. Note that a solution y in A' of a system of polynomials g over A comes from a solution y' of g in a smooth A-algebra B, which is mapped in a solution of g in A by a retraction of $A \to B$.

Examples of morphisms $u : (A, \mathfrak{m}) \to (A', \mathfrak{m}')$, which are filtered colimits of smooth A-algebras are given by the following theorems.

The following result is the General Neron Desingularization [13], [14], [23].

Theorem 2. Let $u : (A, \mathfrak{m}) \to (A', \mathfrak{m}')$ be a flat morphism of Noetherian local rings. The following statements are equivalent:

- (1) *u* is a filtered colimit of smooth A-algebras.
- (2) *u* is regular, that is for every prime ideal $\mathfrak{p} \in \operatorname{Spec} A$ and every finite field extension K of the fraction field of A/\mathfrak{p} the ring $K \otimes_{A/\mathfrak{p}} A'$ is regular.

Let (A, \mathfrak{m}) be a Noetherian local ring and A its completion in the \mathfrak{m} -adic topology. A is *excellent* if the completion map $A \to \hat{A}$ is regular. The following corollary holds by Proposition 1.

Corollary 3. ([13]) Let (A, \mathfrak{m}) be an excellent Henselian local ring and \hat{A} its completion. Then every system of polynomials over A which has a solution in \hat{A} has also one in A. In particular A has the property of Artin approximation; this was conjectured by M. Artin in [2].

Theorem 4. (Zariski [25]) A valuation ring containing a field K of characteristic zero is a a filtered union of its smooth K-subalgebras.

In the same frame, the following result holds.

Theorem 5. ([18, Theorem 2]) Let $V \subset V'$ be an extension of valuation rings containing $\mathbf{Q}, K \subset K'$ its fraction field extension, $\Gamma \subset \Gamma'$ the value group extension of $V \subset V'$ and val : $K'^* \to \Gamma'$ the valuation of V'. Then V' is a filtered colimit of smooth V-algebras if and only if the following statements hold:

- (1) For each $q \in \operatorname{Spec} V$ the ideal qV' is prime,
- (2) For any prime ideals $q_1, q_2 \in \operatorname{Spec} V$ such that $q_1 \subset q_2$ and $\operatorname{height}(q_2/q_1) = 1$ and any $x' \in q_2 V' \setminus q'_1$ there exists $x \in V$ such that $\operatorname{val}(x') = \operatorname{val}(x)$, where $q'_1 \in \operatorname{Spec} V'$ is the prime ideal corresponding to the maximal ideal of $V_{q_1} \otimes_V V'$, that is the maximal prime ideal of V' lying on q_1 .

An *immediate extension* of valuation rings is an extension inducing trivial extensions on residue fields and group value extensions. As a consequence of Theorem 5 we can prove the following:

Corollary 6. ([16, Theorem 21]) If $V \subset V'$ is an immediate extension of valuation rings containing \mathbf{Q} then V' is a filtered colimit of smooth V-algebras.

The following corollary holds by Proposition 1.

Corollary 7. ([18, Proposition 18]) Let $V \subset V'$ be an extension of valuation rings containing \mathbf{Q} with the same residue field, $K \subset K'$ its fraction field extension, $\Gamma \subset \Gamma'$ the value group extension of $V \subset V'$ and val : $K'^* \to \Gamma'$ the valuation of V'. Assume that V is Henselian and the statements (1), (2) of Theorem 5 hold (for example when $V \subset V'$ is immediate). Then every system of polynomials over V which has a solution in V' has also one in V.

Let V be a valuation ring, λ be a fixed limit ordinal and $v = \{v_i\}_{i < \lambda}$ a sequence of elements in V indexed by the ordinals *i* less than λ . Then v is called *pseudo convergent* if

 $\operatorname{val}(v_i - v_{i''}) < \operatorname{val}(v_{i'} - v_{i''})$ for $i < i' < i'' < \lambda$ (see [9], [21]). A pseudo limit of v is an element $w \in V$ with

 $\operatorname{val}(w - v_i) < \operatorname{val}(w - v_{i'})$ (that is, $\operatorname{val}(w - v_i) = \operatorname{val}(v_i - v_{i'})$ for $i < i' < \lambda$. The following example shows that Corollary 6 fails in positive characteristic.

Example 8. ([15, Example 3.1.3],[12], [17]) . Let k be a field of characteristic p > 0, X a variable, $\Gamma = \mathbf{Q}$ and K the fraction field of the group algebra $k[\Gamma]$, that is the rational functions in $(X_q)_{q \in \mathbf{Q}}$. Let P be the field of all formal sums $z = \sum_{n \in \mathbf{N}} c_n X^{\alpha_n}$, where $(\alpha_n)_{n \in \mathbf{N}}$ is a monotonically increasing sequence from Γ and $c_n \in k$. Set $\operatorname{val}(z) = \alpha_s$, where $s = \min\{n \in \mathbf{N} : c_n \neq 0\}$ if $z \neq 0$ and let V be the valuation ring defined by $\operatorname{val}: P^* \to \Gamma, z \to \operatorname{val}(z)$.

Let $\rho_n = (p^{n+1}-1)/(p-1)p^{n+1}$, $y = -1 + \sum_{n\geq 0} (-1)^n X^{\rho_n}$ and $a_i = -1 + \sum_{0\leq n\leq i} (-1)^n X^{\rho_n}$, We have $1+\rho_n = p(\rho_{n+1})$ for $n\geq 0$ and $p\rho_0 = 1$ and y is a pseudo limit of the pseudo convergent sequence $a = (a_i)_{i\in\mathbb{N}}$, which has no pseudolimit in K. Then y is a root of the separable polynomial $g = Y^p + XY + 1 \in K[Y]$ and the algebraic separable extension $V_0 = V \cap K \subset V_1 = V \cap K(y)$ is not dense, in particular V_1 is not a filtered colimit of smooth V_0 -algebras (apply [15, Theorem 6.9], or [17, Theorem 2]).

However, there exist an important result when char k = p > 0.

Theorem 9. (B. Antieau, R. Datta [1, Theorem 4.1.1]) Every perfect valuation ring of characteristic p > 0 is a filtered union of its smooth \mathbf{F}_p -subalgebras.

We mention also the following result.

Theorem 10. ([19]) Let V be a one dimensional valuation ring with value group Γ containing its residue field and U a set with $\operatorname{card}(U) > \operatorname{card}(\Gamma)$. Then there exists an ultrafilter \mathcal{U} on U such that taking the ultrapower \tilde{V} of V with respect to \mathcal{U} and $\bar{V} = \tilde{V} / \bigcap_{z \in V, z \neq 0} z \tilde{V}$ the following assertions hold:

- (1) \overline{V} contains the completion of V.
- (2) If V' is an immediate extension of V contained in \overline{V} with V'/V separable then V' is a filtered colimit of smooth V-algebras.

3. Complete intersection algebras

For a general immediate extension of valuatin rings $V \subset V'$ we should expect that V' is a filtered union of its complete intersection V-subalgebras. In the Noetherian case a morphism of rings is a filtered direct limit of smooth algebras iff it is a regular morphism (see Theorem 2).

A complete intersection V-algebra essentially of finite type is a local V-algebra of type C/(P), where C is a localization of a polynomial V-algebra of finite type and P is a regular sequence of elements of C. Theorem 18 stated below says that V' is a filtered union of its V-subalgebras of type C/(P). Since V' is local it is enough to say that V' is a filtered union of its V-subalgebras of type $T_h/(P)$, T being a polynomial V-algebra of finite type, $0 \neq h \in T$ and P is a regular sequence of elements of T. Clearly, T_h is a smooth V-algebra and in fact it is enough to say that V' is a filtered union of its V-subalgebras of type G/(P), where G is a smooth V-algebra of finite type and P is a regular sequence of elements of G. Conversely, a V-algebra of such type G/(P) has the form $T_h/(P)$ for some T, h, P using [23, Theorem 2.5]. By abuse we understand by a complete intersection V-algebra of finite type a V-algebra of such type G/(P), or $T_h/(P)$ which are not assumed to be flat over V.

The following lemma is a variant of Ostrowski ([12, page 371, IV and III], see also [21, (II,4), Lemma 8]) and [20, Lemma 2].

Lemma 11. (Ostrowski) Let β_1, \ldots, β_m be any elements of an ordered abelian group G, λ a limit ordinal and let $\{\gamma_s\}_{s<\lambda}$ be an increasing sequence of elements of G. Let t_1, \ldots, t_m , be distinct integers. Then there exists an ordinal $\nu < \lambda$ such that $\beta_i + t_i \gamma_s$ are different for all $s > \nu$. Also there exists an integer $1 \le r \le m$ such that

$$\beta_i + t_i \gamma_s > \beta_r + t_r \gamma_s$$

for all $i \neq r$ and $s > \nu$.

The following two results use the above lemma.

Lemma 12. ([20, Lemma 3]) Let $V \subset V'$ be an immediate extension of valuation rings, $K \subset K'$ its fraction field extension and $(v_i)_{i < \lambda}$ an algebraic pseudo convergent sequence in V, which has a pseudo limit x in V', but no pseudo limit in K. Set $x_i = (x - v_i)/(v_{i+1} - v_i)$. Let $s \in \mathbb{N}$ be the minimal degree of the polynomials $f \in V[Y]$ such that $val(f(v_i)) < val(f(v_j))$ for large $i < j < \lambda$ and $g \in V[Y]$ a polynomial with deg g < s. Then there exist $d \in V \setminus \{0\}$ and $u \in V[x_i]$ for some $i < \lambda$ with g(x) = du and val(u) = 0.

Lemma 13. ([20, Lemma 4]) Let $V \subset V'$ be an immediate extension of valuation rings, $K \subset K'$ its fraction field extension and $(v_i)_{i < \lambda}$ an algebraic pseudo convergent sequence in K, which has a pseudo limit x in V', algebraic over V, but no pseudo limit in K. Assume that h = Irr(x, K) is from V[X]. Then

(1) $\operatorname{val}(h(v_i)) < \operatorname{val}(h(v_j))$ for large $i < j < \lambda$,

(2) if h has minimal degree among the polynomials $f \in V[X]$ such that $val(f(v_i)) < val(f(v_j))$ for large $i < j < \lambda$, then $V'' = V' \cap K(x)$ is a filtered union of its complete intersection V-subalgebras.

Remark 14. The extension $V \subset V''$ from (2) of the above lemma is isomorphic with the one constructed in [9, Theorem 3].

Lemma 15. ([10, Lemma 5.2]) Assume that $(v_j)_j$ is a pseudo-convergent sequence in V with a pseudo limit x in an extension V' of V but with no pseudo limit in V. Let $f \in V[X]$ be a polynomial. Then $(f(v_j))_j$ is ultimately pseudo-convergent and f(x) is a pseudo limit of it.

An interesting situation is given below.

Example 16. (Anonymous Referee of a previous paper) Let $R = \mathbf{F}_2[[t]]$ be the formal power series ring in t over $\mathbf{F}_2 = \mathbf{Z}/2\mathbf{Z}$. Then the fraction field $\mathbf{F}_2((t))$ of R has a t-adic valuation which has a canonical extension denoted by val to the perfect hull $K = \mathbf{F}_2((t))^{1/2^{\infty}}$ of $\mathbf{F}_2((t))$. Let V be the valuation ring defined by val on K. Let a be a root of the polynomial $f = X^2 - X - (1/t) \in K[X]$. Then the extension of val to K(a) gives an immediate extension $K \subset K(a)$ with a value group Γ . This extension is closed to an example of [12] (see Example 8). We can assume that the root a is a pseudo limit of the pseudo convergent sequence $(a_n)_{n \in \mathbf{N}}$ given by

$$a_n = \sum_{i=1}^n t^{-2^{-i}},$$

which has no pseudo limit in K. Note that $val(a_n) < 0$ and so $a_n \notin V$ for all $n \in \mathbb{N}$.

Let $0 < \alpha \in \Gamma$ and $b = (b_j)_{j < \lambda}$ be a transcendental pseudo convergent sequence over K(a) such that $\operatorname{val}(b_j) = \alpha$ for high enough j. By [9, Theorem 2] there exists an unique immediate transcendental extension $K(a) \subset L = K(a, z)$ such that z is a pseudo limit of b (the valuation of L is still denoted by val). Then $\operatorname{val}(z) = \alpha > 0$. Set $x = z + a \in L$. We have

$$\operatorname{val}(f(x)) = \operatorname{val}((x-a)^2 - (x-a) + a^2 - a - (1/t)) = \operatorname{val}(x-a) + \operatorname{val}(x-a-1) = \alpha + 0 > 0.$$

Note that

$$\operatorname{val}(x - a_n) = \min\{\operatorname{val}(x - a), \operatorname{val}(a - a_n)\} = \operatorname{val}(a_{n+1} - a_n)$$

because $\operatorname{val}(x-a) = \alpha > 0$ and $\operatorname{val}(a-a_n) = \operatorname{val}(a_{n+1}-a_n) < 0$, that is x is also a pseudo limit of $(a_n)_n$. Also we see that $\operatorname{val}(a_n^2 - a_m - (1/t)) < 0$ for all $n, m \in \mathbb{N}$ with $m \neq n-1$ and $a_n^2 - a_m - (1/t) = 0$ when m = n-1. Consequently, $\operatorname{val}(a_n^2 - a_m - (1/t))$ is either < 0, or it is ∞ for all $n, m \in \mathbb{N}$ and it follows that $\operatorname{val}(a_n^2 - a_m - (1/t))$ is never $\operatorname{val}(f(x))$.

We see that the ultimately pseudo convergent sequence $(a_n^2)_{n \in \mathbb{N}}$ (using Lemma 15) has no pseudo limit in K because $K^2 = K$ and a pseudo limit of (a_n^2) in K would give a pseudo limit of (a_n) in K, which is false. Set $y_0 = x$, $y_1 = y_0^2$, $g = tY_1 - tY_0 - 1 \in K[Y_0, Y_1]$. We have $g(y_0, y_1) = tf(x)$. Then $\operatorname{val}(g(a_n^2, a_m))$, $n, m \in \mathbb{N}$ is never $\operatorname{val}(g(y_0, y_1))$ because $\operatorname{val}(a_n^2 - a_m - (1/t))$ is never $\operatorname{val}(f(x))$ as we have seen. So Lemma 12 does not work for polynomials in two variables.

Lemma 17. ([20, Lemma 6]) Let B be a complete intersection algebra over a ring A and C a complete intersection algebra over B. Then C is a complete intersection algebra over A.

Using Lemma 13 and the above lemma we proved the following result.

Theorem 18. ([20, Theorem 1]) Let V' be an immediate extension of a valuation ring V and $K \subset K'$ the fraction field extension. If K'/K is algebraic then V' is a filtered union of its complete intersection V-subalgebras of finite type.

As a consequence the next theorem follows.

Theorem 19. Let V be a valuation ring containing its residue field k with a free value group (for example when it is finitely generated) Γ . Assume that y is a system of elements of V such that val(y) is a basis of Γ and the fraction field of V is an algebraic extension of k(y). Then V is a filtered union of its complete intersection k-subalgebras of finite type.

Proof. By [16, Lemma 27 (1)] we see that V is an immediate extension of the valuation ring $V_0 = V \cap k(y)$, which is a filtered union of localizations of its polynomial k-subalgebras (see [6, Theorem 1, in VI (10.3)], or [16, Lemma 26 (1)]). In fact, important here is that there exists a cross-section $s : \Gamma \to K^*$ (a cross-section of V is a section in the category of abelian groups of the valuation map val : $K^* \to \Gamma$). Since Γ is free we define s by val $(y) \to y$. Now it is enough to apply Theorem 18 for $V_0 \subset V$.

The above theorem connects the theory of valuation rings with the theory of complete intersection local rings. We remind that for a given complete intersection local ring (R, \mathfrak{m}, k) a minimal free resolution over R of the residue field k is described in [24, Theorem 4] and others as [3, Theorem 2.7], [4, §6], [7, Proposition 1.5.4], [8, §3], [5, Theorem 2.5], [11, Theorem 4.1].

References

- B. Antieau, R. Datta, Valuation rings are derived splinters, Math. Zeitschrift, 299, (2021, 827-851.
- [2] M. Artin, Construction techniques for algebraic spaces, Actes Congres Intern. Math. 1, (1970), 419-423.
- [3] E. Assmus, On the homology of local rings, Illinois J. Math., 3(2) (1959), 187–199.
- [4] L. Avramov, Infinite free resolutions, Six Lectures on Commutative Algebra (Bellaterra, 1996), Progr. Math., 166 (1998), Birkhauser, Basel, 1–118.
- [5] L. Avramov, I. Henriques, L. Sega, Quasi-complete intersection homomorphisms, Pure Appl. Math. 9 (2013), no. 4, 579–612
- [6] N. Bourbaki, Éléments de mathématique. Algèbre commutative, chap. I-VII, Hermann (1961, 1964, 1965); chap. VIII-X, Springer (2006, 2007) (French).
- [7] T. Gulliksen, G. Levin, Homology of local rings, Queen's Papers in Pure Appl. Math. 20 (1969),

- [8] J. Herzog, A. Martsinkovsky, Gluing Cohen-Macaulay modules with applications to quasihomogeneous complete intersections with isolated singularities, *Comment. Math. Helv.* 68 (1993), no. 3, 365–384.
- [9] I. Kaplansky, Maximal fields with valuations, Duke Math. J. 9 (1942), 303-321.
- [10] F. V. Kuhlmann, I. Vlahu The relative approximation degree in valued function fields, Math. Z., 276, (2014), 203–235.
- [11] V. Nguyen, O. Veliche, Iterated mapping cones on the Koszul complex and their application to complete intersection rings, Journal of Algebra and Its Applications, https://www.worldscientific.com/doi/10.1142/S0219498825502342.
- [12] A. Ostrowski, Untersuchungen zur arithmetischen Theorie der Körper, Math.
 Z. 39 (1935), no. 1, 321-404.
- [13] D. Popescu, General Neron Desingularization and approximation, Nagoya Math. J., 104, (1986), 85-115.
- [14] D. Popescu, Letter to the Editor, General Neron Desingularization and approximation, Nagoya Math. J., 118, (1990), 45-53.
- [15] D. Popescu, Algebraic extensions of valued fields, J. Algebra 108, (1987), no. 2, 513-533.
- [16] D. Popescu, Néron desingularization of extensions of valuation rings with an Appendix by Kęstutis Česnavičius, in "Transcendence in Algebra, Combinatorics, Geometry and Number Theory", Eds. Alin Bostan, Kilian Raschel, Springer Proceedings in Mathematics and Statistics 373, (2021), 275-307.
- [17] D. Popescu, Valuation rings of dimension one as limits of smooth algebras, Bull. Math. Soc. Sci. Math. Roumanie, 64(112), (2021), 65-74.
- [18] D. Popescu, Extensions of valuation rings containing Q as limits of smooth algebras, Bull. Math. Soc. Sci. Math. Roumanie, 65(113), No 2, (2022), 253-262.
- [19] D. Popescu, Immediate extensions of valuation rings and ultrapowers, Math. Reports 25(75), (2023), 505–512.
- [20] D. Popescu, Algebraic valuation ring extensions as limits of complete intersection algebras, Revista Matematica Complutense, **37**, No 2, (2024), 467-472.
- [21] O. F. G. Schilling, The theory of valuations, Mathematical Surveys, Number IV, American Math. Soc., (1950).
- [22] A. J. de Jong et al., The Stacks Project. Available at http://stacks.math.columbia.edu.
- [23] R. Swan, Neron-Popescu desingularization, in "Algebra and Geometry", Ed. M. Kang, International Press, Cambridge, (1998), 135-192.
- [24] J. Tate, Homology of Noetherian rings and local rings, Illinois J. Math., 1 (1957), 14–27.
- [25] O. Zariski, Local uniformization on algebraic varieties, Ann. of Math. 41, (1940), 852-896.

SIMION STOILOW INSTITUTE OF MATHEMATICS OF THE ROMANIAN ACADEMY, RESEARCH UNIT 5, P.O. Box 1-764, BUCHAREST 014700, ROMANIA,

University of Bucharest, Faculty of Mathematics and Computer Science Str. Academiei 14, Bucharest 1, RO-010014, Romania,

 ${\rm EMAIL:} \ dorin.m.popescu@gmail.com$