A RENORMALIZATION GROUP APPROACH TO HIGHER ORDER CORRECTIONS TO THE DECAY OF SOLUTIONS TO NONLINEAR INTEGRAL EQUATIONS

Gastão A. Braga¹, Jussara M. Moreira¹, Antônio Marcos da Silva³, and Camila F. Souza⁴

¹Departamento de Matemática, Universidade Federal de Minas Gerais ³Departamento de Matemática, Universidade Federal de Ouro Preto ⁴Departamento de Matemática, Centro Federal de Educação Tecnológica de Minas Gerais

Abstract

In this paper we employ the Renormalization Group (RG) method to study higher order corrections to the long-time asymptotics of a class of nonlinear integral equations with a generalized heat kernel and with time-dependent coefficients. This is a follow up of the papers [1, 2, 3].

1 Introduction

This paper is a follow up of [1, 2], where the long-time asymptotics of solutions to the following class of nonlinear integral equations

$$u(x,t) = \int G(x-y,s(t))f(y)dy +$$
$$\lambda \int_{1}^{t} \int G(x-y,s(t)-s(\tau))F(u,u_{x})(y,\tau)\,dyd\tau$$
(1)

was investigated using the Renormalization Group (RG) method developed by Bricmont et al. in [4]. In the above equation, G(x,t) is required to satisfy some properties (see conditions (G) listed below), s(t) is given by

$$s(t) = \int_{1}^{t} c(\tau) d\tau, \qquad (2)$$

where $c(t) = t^p + o(t^p)$, with p > 0 and $o(t^p)$ is a little order of t^p as $t \to \infty$, with fin an appropriate Banach space. In [1, 2], the nonlinear term $F(u, u_x)$ depends only on u and is given by $F(u) = -\mu u^{\alpha_c} + \sum_{j>\alpha_c} a_j u^j$, where $\alpha_c = (p+1+d)/(p+1)$, with p and d being positive exponents related to s(t) and G(x, t), respectively. The asymptotic results in [1], for $\mu = 0$ in the above definition of F(u), and in [2] for $\mu > 0$, are described by

$$u(x,t) \sim \frac{A}{[t\ r(t)]^{(p+1)/d}} G\left(\frac{x}{t^{(p+1)/d}}, \frac{1}{p+1}\right) \text{ when } t \to \infty.$$
 (3)

where r(t) = 1 if $\mu = 0$ and $r(t) = \ln t$ if $\mu > 0$. The behavior (3) is an outcome of the RG operator flow analysis around its linearization's fixed point G(x, 1/(p+1)).

The aim of this paper is to keep exploiting the RG method to study the asymptotic behavior of solutions to (1), with the nonlinearity

$$F(u, u_x) = \sum_{j > \alpha} a_j u^j u_x, \quad \alpha > \alpha_c, \tag{4}$$

where

$$\alpha_c = \frac{d - (p+1)}{2(p+1)} \tag{5}$$

and F being analytic at the origin (0,0), but considering the initial data f under the zero mass condition $\int_{\mathbb{R}} f(x) dx = 0$.

Under the zero mass condition, A = 0 in (3) and, in order to get the right asymptotic behavior, it is necessary to find the next leading order contribution to the long time behavior and this will be done by using the RG multiscale analysis. Following [3] we show that, also in this case, there is a well defined RG operator so that the long time behavior of u(x, t) is dictated by a fixed point of its linearization and is given by

$$u(x,t) \sim \frac{A}{t^{2(p+1)/d}} G'\left(\frac{x}{t^{(p+1)/d}}, \frac{1}{p+1}\right) \text{ when } t \to \infty,$$
 (6)

where $G'(x,t) \equiv \partial_x G(x,t)$ and G'(x,1/(p+1)) is a fixed point of the linear RG operator.

The RG method is a multiscale method that has shown to be quite appropriate for getting fine information on the asymptotics of solutions to time evolution problems Concerning the asymptotics, via the RG method, under the zero mass condition, we point out the work [5] of Bona et al. Using the standard RG operator, see [4], Bona et al. analyzed the asymptotic behavior of solutions to a generalized Kortweg-de Vries equation but the long time profile function is not a fixed point of their RG operator. Subsequently, based on the numerical results presented in [6], Braga et al. [3] redefined the RG operator to study a generalized Burgers equation considering the zero mass.

The kernel G = G(x, t) satisfies the following conditions which we denote by (G):

 (G_1) There are integers q > 1 and M > 0 such that $G(\cdot, 1) \in C^{q+2}(\mathbb{R})$ and

$$\sup_{x \in \mathbb{R}} \{ (1+|x|+x^2)(1+|x|)^{M+1} | G^{(j)}(x,1) | \} < \infty, \quad j = 0, 1, ..., q+2,$$

where $G^{(j)}(x,1)$ denotes the j-th derivative $(\partial_x^j G)(x,1)$.

 (G_2) There is a positive constant d such that

$$G(x,t) = t^{-\frac{1}{d}} G\left(t^{-\frac{1}{d}}x,1\right), \quad x \in \mathbb{R}, \ t > 0;$$

$$(G_3) \ G(x,t) = \int_{\mathbb{R}} G(x-y,t-\tau)G(y,\tau)dy \quad \text{ for } x \in \mathbb{R} \text{ and } t > \tau > 0.$$

From these properties, we can now present the heuristics for obtaining the asymptotics in the linear case. Consider the solution to (1) with $\lambda = 0$ and $s(t) \approx \frac{t^{p+1}}{p+1}$ for large t. Using condition (G₂) in Fourier space $\widehat{G}\left(\omega, \frac{t^{p+1}}{p+1}\right) = \widehat{G}\left(t^{\frac{(p+1)}{d}}\omega, \frac{1}{p+1}\right)$, we can write

$$\begin{split} u(x,t) &= \frac{1}{2\pi} \int_{|\omega| < \epsilon} \widehat{G}\left(t^{\frac{(p+1)}{d}}\omega, \frac{1}{p+1}\right) e^{i\omega x} \widehat{f}(\omega) d\omega \\ &+ \frac{1}{2\pi} \int_{|\omega| > \epsilon} \widehat{G}\left(t^{\frac{(p+1)}{d}}\omega, \frac{1}{p+1}\right) e^{i\omega x} \widehat{f}(\omega) d\omega. \end{split}$$

The smoothness of G(x,t) implies a certain polynomial decay rate of $\widehat{G}(\omega,1)$ so that, after expanding \widehat{f} around $\omega = 0$, we get

$$u(x,t) \approx \frac{1}{2\pi} \int_{|\omega| < \epsilon} \widehat{G}\left(t^{\frac{(p+1)}{d}}\omega, \frac{1}{p+1}\right) e^{i\omega x} \left[\widehat{f}(0) + \omega\widehat{f}'(0) + O\left(\omega^2\right)\right] d\omega.$$

Assuming the zero mass hypothesis and dropping off $O(\omega^2)$ terms from the above integral, after a change of variables,

$$u(x,t) \approx \frac{1}{2\pi} \left[\frac{1}{t^{(p+1)/d}} \right]^2 \int_{\left| \frac{\omega}{t^{(p+1)/d}} \right| < \epsilon} \omega \ \widehat{G}\left(\omega, \frac{1}{p+1}\right) \ e^{i\left(\frac{\omega}{t^{(p+1)/d}}\right)x} \ \widehat{f}'(0) \ d\omega,$$

which gives us the right asymptotic behavior of u

$$u(x,t) \approx \frac{A}{t^{2(p+1)/d}} G'\left(\frac{x}{t^{(p+1)/d}}\right), \ t \gg 1,$$
(7)

where $A = -i\hat{f}'(0)$ and G'(x) stands for the x-derivative of $G\left(x, \frac{1}{p+1}\right)$. The conclusion, from (7), is that as $t \to \infty$, the solution to the linear problem ((1) with $\lambda = 0$) decays with rate $t^{2(p+1)/d}$, diffuses with rate $t^{(p+1)/d}$ and, up to a constant $A = -i\int_R xf(x)dx$, it has G'(x) as profile function.

Motivated by equation (7), we define the Renormalization Group operator as an operator acting in the space of initial conditions as follows

$$(R_L f)(x) = L^{2(p+1)/d} u\left(L^{(p+1)/d} x, L\right),$$
(8)

where L > 1 and u is the solution to (1). Accordingly to the above definition, nonlinearities of the form $u^{j}u_{x}$, $j \geq 1$, are irrelevant in the RG sense. In particular, the nonlinearity (4), which will be treated in this paper, is irrelevant, implying that the asymptotics of solutions to (1) is, basically, the one given by (7) and that is what we will prove in this paper. Also, as it will be shown in Lemma 2.6 the profile function G'(x) in (7) is a RG fixed point.

In order to state our results, we now introduce the space for the initial data. Since we are considering f with zero mass, it is necessary to incorporate the second derivative term in the definition below. Therefore, given q > 1, we consider

$$\mathcal{B}_q = \left\{ f \in L^1(\mathbb{R}) : \widehat{f}(\omega) \in C^2(\mathbb{R}) \text{ and } \|f\|_q < +\infty \right\},\tag{9}$$

where $||f|| = \sup_{\omega \in \mathbb{R}} (1 + |\omega|^q) \left(|\widehat{f}(\omega)| + |\widehat{f'}(\omega)| + |\widehat{f''}(\omega)| \right).$

Theorem 1.1. Let u(x,t) be the solution to (1) with $\lambda \in [-1,1]$, where $f \in B_q$ satisfies the zero mass hypothesis, G(x,t) satisfies the conditions (**G**) with q > 2and F is given by (4). There exists positive constants A and ϵ such that, if $||f|| < \epsilon$, then

$$\lim_{n \to \infty} \|t^{2n(p+1)/d} u(t^{n(p+1)/d}, t^n) - AG_p\| = 0,$$
(10)

where $A = A(d, f_0, F, p, q)$ and

$$G_p = G'\left(\cdot, \frac{1}{p+1}\right). \tag{11}$$

Remark: The above prefactor A, which we will show to be the limit of a sequence of prefactors A_n , can be found explicitly.

2 The Linear Case

In order to be able to fully discuss equation (1), we must present some properties of the kernel G and consider first the employment of the RG method to u(x,t) given by the linear component of (1), that is, u satisfies (1) with $\lambda = 0$, for t > 1, $x \in \mathbb{R}$, G = G(x, t) and s(t) satisfying, respectively, hypotheses (G) and equation (2) and with $f \in \mathcal{B}_q$.

Our goal in this section is to prove the following:

Theorem 2.1. Let G be a kernel satisfying conditions (G), $f \in \mathcal{B}_q$ satisfying the zero mass condition, $A = -i\hat{f}'(0)$ and u solution to (1) with $\lambda = 0$. Then,

$$\lim_{t \to \infty} \| t^{\frac{2(p+1)}{d}} u(t^{\frac{p+1}{d}}, t) - AG_p(\cdot) \| = 0,$$
(12)

where G_p is given by (11).

To prove Theorem 2.1 we will establish some properties of the kernel G which follow from conditions (G). We denote $\widehat{G}'(\omega, t) \equiv \partial \widehat{G}/\partial \omega(\omega, t)$.

Lemma 2.1. If $G(x,t) : \mathbb{R} \times (0,\infty) \to \mathbb{R}$ satisfies property (G_1) , then, for j = 0, 1, 2, $\widehat{G}^{(j)}(\omega, 1) \in L^{\infty}(\mathbb{R})$ and

$$\sup_{\omega \in \mathbb{R}} (1 + |\omega|^q) |\omega|^2 |\widehat{G}^{(j)}(\omega, 1)| < \infty.$$

Proof: It follows from property (G_1) of (\mathbf{G}) and from Fourier Transform results.

We then denote throughout the paper:

$$K_j \equiv \sup_{\omega \in \mathbb{R}} |\widehat{G}^{(j)}(\omega, 1)|, \text{ for } j = 0, 1, 2.$$
(13)

It follows from Lemma 2.1 and from property (G_2) of **(G)**, that $\widehat{G}(\omega, t)$ is well defined for t > 0 and

$$\widehat{G}^{(j)}(\omega,t) = t^{j/d} \partial_{\omega}^{j} \widehat{G}(t^{1/d}\omega,1), \quad \text{for } j = 0, 1, 2, \ \forall t > 0, \ \omega \in \mathbb{R}.$$
(14)

Also, condition (G_3) of (\mathbf{G}) in the Fourier space is

$$\widehat{G}(\omega, t) = \widehat{G}(\omega, t - s)\widehat{G}(\omega, s) \quad t > s > 0 \text{ and } \omega \in \mathbb{R}.$$
(15)

These results together lead to the following lemmas:

Lemma 2.2. Suppose $G(x,t) : \mathbb{R} \times (0,\infty) \to \mathbb{R}$ satisfies properties (G_1) , (G_2) and (G_3) . Given $t_1, t_2 \in (0,\infty)$, with $t_1 < t_2$, then, for all $\omega \in \mathbb{R}$,

$$|\widehat{G}(\omega, t_2)| \le K_0 |\widehat{G}(\omega, t_1)|,$$

$$|\widehat{G}'(\omega, t_2)| \le K_1(t_2 - t_1)^{\frac{1}{d}} |\widehat{G}(\omega, t_1)| + K_0 |\widehat{G}'(\omega, t_1)|$$

and

$$|\widehat{G}''(\omega, t_2)| \le K_2(t_2 - t_1)^{\frac{2}{d}} |\widehat{G}(\omega, t_1)| + 2K_1(t_2 - t_1)^{\frac{1}{d}} |\widehat{G}'(\omega, t_1)| + K_0 |\widehat{G}''(\omega, t_1)|$$

with K_j , for j = 0, 1, 2, is given by (13).

Lemma 2.3. If $G(x,t) : \mathbb{R} \times (0,\infty) \to \mathbb{R}$ satisfies conditions (G_1) , (G_2) and (G_3) , then, for all t > 0,

$$\int_{\mathbb{R}} G(x,t) dx = 1$$

2.1 The RG Operator for the Integral Linear Equation

In Braga et al. [1], we have established the RG method to integral equations. Unlike the case where we look for the long-time behavior of solutions to PDEs, for integral equations, the definition of the RG operator is slightly more refined. In fact, for equation equation (1), with $\lambda = 0$, not even the kernel is scale invariant. In order to define our RG transformation, let us consider u given by equation (1), with $\lambda = 0$, and define, for $t \in (1, L]$, $f_0 \equiv f$, $u_0(x, t) = \int G(x - y, s(t)) f_0(y) dy$ and, for $n = 1, 2, \dots$,

$$u_n(x,t) \equiv \int G(x-y,s(t)-s(L^n))u_{n-1}(y,L^n)dy, \ t \in (L^n,L^{n+1}],$$

and, for $n = 0, 1, \cdots$,

$$u_{f_n}(x,t) \equiv \int G\left(x-y, s_n(t)\right) f_n(y) dy, \ t \in (1,L],$$

where

$$s_n(t) = \frac{s(L^n t) - s(L^n)}{L^{n(p+1)}}$$
(16)

and

$$f_{n+1} \equiv R_{L,n}^0 f_n \equiv L^{2(p+1)/d} u_{f_n}(L^{(p+1)/d}, L).$$
(17)

With the above definitions, it is not hard to see that the RG operator satisfies the semigroup property, that is:

$$R^{0}_{L^{n},0}f_{0} = (R^{0}_{L,n-1} \circ \dots \circ R^{0}_{L,1} \circ R^{0}_{L,0})f_{0}$$

$$(18)$$

and so, the limit in (12) will be obtained by studying the dynamics of operators $R_{L,n}^0 f_n$, $n = 0, 1, 2, \cdots$ in the initial data space \mathcal{B}_q .

It follows from definition (2) that we can write the function s(t) as

$$s(t) = \frac{t^{p+1} - 1}{p+1} + r(t), \tag{19}$$

where $r(t) = o(t^{p+1})$. Therefore, there is $L_0 > 1$ such that, for $L > L_0$, $|r(L)|/L^{p+1} < 1/[4(p+1)]$. Furthermore, from (16), we can write

$$s_n(t) \equiv \frac{t^{p+1} - 1}{p+1} + r_n(t)$$
, where $r_n(t) = \frac{r(L^n t) - r(L^n)}{L^{n(p+1)}}$.

Then, for $L > L_0 > 1$, $|r_n(L)| < 1/[2(p+1)]$. Defining

$$L_1 \equiv \max\{L_0, 3^{1/(p+1)}\}\tag{20}$$

we get, for $L > L_1$,

$$\frac{1}{6(p+1)} < \frac{s_n(L)}{L^{p+1}} < \frac{3}{2(p+1)},\tag{21}$$

for all $n \ge 0$. Furthermore, from the properties of G and the Fourier transform, $R_{L,n}^0 g \in \mathcal{B}_q$, for all $g \in \mathcal{B}_q$ and $n = 0, 1, \cdots$. In the next lemma, we will show that the derivative with respect to the variable x of the function G(x, t) at t = 1/(p+1)is also in \mathcal{B}_q for q > 1 given by (G_1) .

Lemma 2.4. For q > 1 given by (G_1) , the function G_p in (11) is in \mathcal{B}_q and

$$\|G_p\| < C_{d,p,q},\tag{22}$$

where

$$C_{d,p,q} = (p+1)^{\frac{q}{d}} \sup_{k \in \mathbb{R}} \left((1+|k|^q) \sum_{j=0}^2 |\widehat{G}^{(j)}(k,1)| \right),$$
(23)

Proof: It follows from (14), with $t = \frac{1}{p+1}$ and $j \in \{0, 1, 2\}$, that

$$||G_p|| = \sup_{\omega \in \mathbb{R}} (1+|\omega|^q) \left[\sum_{j=0}^2 \left(\frac{1}{p+1} \right)^{j/d} \left| \widehat{G_p}^{(j)} \left(\left(\frac{1}{p+1} \right)^{\frac{1}{d}} \omega, 1 \right) \right| \right].$$

Taking $k = [1/(p+1)]^{\frac{1}{d}}\omega$ and the supreme in k, we obtain (22).

From now on, L_1 will always denotes the constant given by (20). In the next lemma we prove that $R_{L^n}^0 G_p \in \mathcal{B}_q$:

Lemma 2.5. There is a constant \tilde{K} such that $||R_{L^n}^0G_p|| \leq \tilde{K}$ for all $L > L_1$.

Proof: It follows from (11), (15) and the properties of the Fourier transform that

$$\mathcal{F}[R_{L^n}^0 G_p](\omega) = i\omega \widehat{G}\left(\frac{\omega}{L^{n(p+1)/d}}, \frac{1}{p+1} + s_0(L^n)\right).$$
(24)

Taking the derivatives of the above equation and using (14) for j = 0, 1 and 2 and (21), it follows from Lemma 2.2 that

$$\begin{aligned} \|R_{L^{n}}^{0}G_{p}\| &\leq \sup_{\omega \in \mathbb{R}} (1+|\omega|^{q})(|\omega|+1) \left(2K_{0}+2K_{1}\left(\frac{7}{3(p+1)}\right)^{\frac{1}{d}}+K_{2}\left(\frac{7}{3(p+1)}\right)^{\frac{2}{d}}\right) \\ &\left[\left|\widehat{G}\left(\omega,\frac{1}{6(p+1)}\right)\right|+\left|\widehat{G}'\left(\omega,\frac{1}{6(p+1)}\right)\right|+\left|\widehat{G}''\left(\omega,\frac{1}{6(p+1)}\right)\right|\right].\end{aligned}$$

Applying the change of variable $k = 1/[6(p+1)]^{\frac{1}{d}}\omega$, we get, as an upper bound for $||R_{L^n}^0G_p||$:

$$\sup_{k \in \mathbb{R}} \qquad [6(p+1)]^{\frac{q+1}{d}} (1+|k|^q) (1+|k|) \left[2K_0 + 2K_1 \left(\frac{7}{3(p+1)} \right)^{\frac{1}{d}} + K_2 \left(\frac{7}{3(p+1)} \right)^{\frac{2}{d}} \right]$$
$$\left\{ |\widehat{G}(k,1)| + |\widehat{G}'(k,1)| + |\widehat{G}''(k,1)| \right\}.$$

From Lemma 2.1, the right hand side of the above inequality is finite, which finishes the proof.

In the next lemma we show that G_p is an asymptotic fixed point for the RG operator $R_{L^n}^0 G_p$, that is, $R_{L^n}^0 G_p \to G_p$ when $n \to \infty$.

Lemma 2.6. There are positive constants M = M(p,q,d) and N = N(p,q,d) such that

$$\|R_{L^{n}}^{0}G_{p} - G_{p}\| \le M \left| \frac{r(L^{n})}{L^{n(p+1)}} \right|^{\frac{1}{d}} + N \left| \frac{r(L^{n})}{L^{n(p+1)}} \right|^{\frac{2}{d}},$$
(25)

where $G_p(x) = G'(x, 1/(p+1)).$

Proof: It follows from (24), (15) and (14) with j = 0, that

$$\mathcal{F}[R_{L^n}^0 G_p - G_p](\omega) = i\omega \widehat{G}\left(\omega, \frac{1}{p+1}\right) \left[\widehat{G}\left(\left(\frac{r(L^n)}{L^{n(p+1)}}\right)^{\frac{1}{d}}\omega, 1\right) - 1\right].$$
 (26)

From Lemmas 2.1 and 2.3 and from the Mean Value Theorem, we conclude that

$$\left|\widehat{G}\left(\omega\left(\frac{r(L^n)}{L^{n(p+1)}}\right)^{\frac{1}{d}}, 1\right) - 1\right| \le K_1 \left|\frac{r(L^n)}{L^{n(p+1)}}\right|^{\frac{1}{d}} |\omega|.$$
(27)

Using the above results, we bound $|\mathcal{F}[R^0_{L^n}G_p-G_p]^{(j)}(\omega)|$, for j=0,1,2, respectively by

$$K_{1}\omega^{2} \left| \frac{r(L^{n})}{L^{n(p+1)}} \right|^{\frac{1}{d}} \left| \widehat{G} \left(\omega, \frac{1}{p+1} \right) \right|,$$
$$\left| \frac{r(L^{n})}{L^{n(p+1)}} \right|^{\frac{1}{d}} \left[2K_{1} \left| \omega \right| \left| \widehat{G} \left(\omega, \frac{1}{p+1} \right) \right| + K_{1}\omega^{2} \left| \widehat{G}' \left(\omega, \frac{1}{p+1} \right) \right| \right]$$

and

$$\begin{split} K_1 \left| \frac{r(L^n)}{L^{n(p+1)}} \right|^{\frac{1}{d}} \left[4|\omega| \left| \widehat{G}'\left(\omega, \frac{1}{p+1}\right) \right| + \omega^2 \left| \widehat{G}''\left(\omega, \frac{1}{p+1}\right) \right| + 2 \left| \widehat{G}\left(\omega, \frac{1}{p+1}\right) \right| \right] \\ + K_2 |\omega| \left| \frac{r(L^n)}{L^{n(p+1)}} \right|^{\frac{2}{d}} \left| \widehat{G}\left(\omega, \frac{1}{p+1}\right) \right|. \end{split}$$

Finally, the result follows from the change of variable $y = 1/(p+1)^{\frac{1}{d}}\omega$ and from the definitions: $M = K_1(M_1 + M_2 + M_3)$, where $M_1 = (p+1)^{\frac{q+2}{d}} \sup_{y \in \mathbb{R}} (1+|y|^q) y^2 \left(|\hat{G}(y,1)| + |\hat{G}'(y,1)| + |\hat{G}''(y,1)| \right)$, $M_2 = 2(p+1)^{\frac{q+1}{d}} \sup_{y \in \mathbb{R}} (1+|y|^q) |y| \left(|\hat{G}(y,1)| + 2|\hat{G}'(y,1)| \right)$, $M_3 = 2(p+1)^{\frac{q}{d}} \sup_{y \in \mathbb{R}} (1+|y|^q) |\hat{G}(y,1)|$ and $N = K_2(p+1)^{\frac{q+1}{d}} \sup_{y \in \mathbb{R}} (1+|y|^q) |y| |\hat{G}(y,1)|$.

The next lemma states that, for L sufficiently large, $R_{L,n}^0$ is a contraction when acting on the space of functions $g \in \mathcal{B}_q$ with zero mass and null first momentum $(\hat{g}(0) = \hat{g}'(0) = 0).$

Lemma 2.7. (Contraction Lemma) Given q > 1, if $g \in \mathcal{B}_q$ is such that $\widehat{g}(0) = \widehat{g}'(0) = 0$, then, there is a constant C = C(d, p, q) > 0 such that

$$\|R_{L,n}^{0}g\| \le \frac{C}{L^{(p+1)/d}} \|g\|, \quad \forall \ L > L_{1}.$$
(28)

The proof of the above theorem follows basically from the Fundamental Theorem of Calculus, the definition of the \mathcal{B}_q space and the properties of the Fourier Transform.

The method in order to obtain Theorem 2.1 consists in decomposing the initial data f into two components, the first a multiple of $R_{L,n}G_p$. We will use the next lemma to conclude that, as the second component of this decomposition has zero mass and null first momentum, its norm goes to zero when $n \to \infty$. It will follow from this fact and from Lemma 2.6 that the function u(x,t), defined by (1), with $\lambda = 0$, and properly rescaled, will behave, when $t \to \infty$, as a multiple of G_p .

Lemma 2.8. Let $f \in \mathcal{B}_q$, $f_0 \equiv f$, $A \equiv -i\hat{f}'_0(0)$ and $f_n = R^0_{L,n-1}f_{n-1}$, $n = 1, 2, \cdots$. Given $L > L_1$, there are functions $g_n \in \mathcal{B}_q$, $n = 0, 1, 2, \ldots$, such that $\hat{g}_n(0) = \hat{g}'_n(0) = 0, \forall n, and$

$$f_0 = AG_p + g_0, \qquad f_n = AR_{L^n}^0 G_p + g_n.$$
 (29)

Furthermore,

$$||g_n|| \le \left(\frac{C}{L^{(p+1)/d}}\right)^n ||g_0||, \ n = 0, 1, 2, \cdots$$
 (30)

where C is the constant from Lemma 2.7.

The proof of Lemma 2.8 follows directly from induction on n and we are finally ready to prove Theorem 2.1 and then analyze the nonlinear case.

Proof of Theorem 2.1: Let C be the constant from Lemma 2.7 and define $L_2 \equiv \max\{L_1, C^{d/(p+1)}\}$. If $L > L_2$, it follows from (25), (29) and (30) that

$$\|L^{2n(p+1)/d}u(L^{n(p+1)/d},L^n) - AG_p\| \le \left(\frac{C}{L^{(p+1)/d}}\right)^n \|g_0\| + M|A| \left|\frac{r(L^n)}{L^{n(p+1)}}\right|^{\frac{1}{d}}.$$

Given $\delta \in (0,1)$, we take $L_3 > L_2$ such that $L_3^{\delta(p+1)/d} > C$. Then, for $L > L_3$, $(CL^{-(p+1)/d})^n \leq L^{-n(p+1)(1-\delta)/d}$ and, for $t = L^n$ with $L > L_3$,

$$\|t^{2(p+1)/d}u(t^{(p+1)/d},t) - AG_p\| \le \frac{\|g_0\|}{t^{(p+1)(1-\delta)/d}} + M|A| \left|\frac{r(t)}{t^{p+1}}\right|^{1/d}.$$
 (31)

To extend this result, we just replace L with $\tau^{\frac{1}{n}}L$ in the estimates, where $\tau \in [1, L]$

and $L > L_3$. Since the constants are independent from L, the above inequality holds for $t = \tau L^n$.

3 The Nonlinear Case

Our goal here is to consider equation (1), with F given by (4) and therefore to prove Theorem 1.1. First we prove that there is a unique local solution to the integral equation (1).

3.1 Local existence and uniqueness of the solution

Our goal in this section is to prove that there is a unique local solution to the integral equation (1). In order to do that, let u_f be the solution to the linear integral equation (1), with $\lambda = 0$, and, given L, q > 1, define

$$B^{(L)} = \{ u : \mathbb{R} \longrightarrow \mathbb{R} \times [1, L]; \ u(\cdot, t) \in B_q, \ \forall t \in [1, L] \}$$
(32)

equipped with the norm $||u||_L = \sup_{t \in [1,L]} ||u(\cdot,t)||$, and

$$B_f \equiv \{ u \in B^{(L)} : \| u - u_f \|_L \le \| f \| \}.$$
(33)

Furthermore, consider the operator $T(u) \equiv u_f + N(u)$, where

$$N(u)(x,t) = \lambda \int_1^t \int G(x-y,s(t)-s(\tau))F(u,u_x)(y,\tau)dyd\tau.$$
 (34)

We will prove the following theorem:

Theorem 3.1. Suppose $F(u, u_x)$ is given by (4), with positive convergence radius r. Given L > 1, d > 1, q > 2 and $\lambda \in [-1, 1]$, there exists $\epsilon = \epsilon(L, q, r, F, d) > 0$ such that, if $f \in \mathcal{B}_q$ and $||f|| < \epsilon$, then there is a unique solution to the integral equation (1), for $t \in [1, L]$ in B_f .

Theorem 3.1 follows immediately from Lemmas 3.2 and 3.3, since the former establishes that T maps B_f into itself and the latter that T is a contraction. Before proving both lemmas, we need the following result: **Lemma 3.1.** Let G(x,t) be a generalized integral kernel G = G(x,t) satisfying conditions (G) and $\phi(\tau) = s(t) - s(\tau)$, where s(t) is given by (2). Then,

$$\left|\int_0^{t-1} \omega \widehat{G}(\omega, \phi(\tau)) d\tau\right| \le B_t$$

for all $\omega \in \mathbb{R}$ and t > 0, where

$$B_t \equiv B(t, d, p, \delta) = K_0(t-1) + C \left[\frac{d}{(d-1)t^{\frac{p}{d}}} + \frac{t-1}{\phi(\delta)^{\frac{1}{d}}} \right],$$
(35)

with K_0 given by (13), δ and C positive constants and d > 1 is the constant given in condition (G₂).

Proof: From equation (14), with j = 0, we get $\left| \int_{0}^{t-1} \omega \widehat{G}(\omega, \phi(\tau)) d\tau \right| \leq K_{0}(t-1)$, when $|\omega| \leq 1$. For $|\omega| > 1$, notice that, from Lemma 2.1, there is C > 0 such that $\widehat{G}(\omega, t) \leq C/(1+|\omega|^{q})|\omega|$. Also, taking $0 < \delta \ll 1$, we have

$$\left|\int_{0}^{t-1} \omega \widehat{G}(\omega, \phi(\tau)) d\tau\right| \leq \left(\int_{0}^{\delta} + \int_{\delta}^{t-1}\right) \frac{C}{[\phi(\tau)]^{\frac{1}{d}}} d\tau = I_{1} + I_{2}, \ t > 1.$$

Then, for $\tau \approx 0$, there is k > 0 such that $\phi(\tau) \geq kt^p \tau$, which leads to $I_1 \leq Ct^{-p/d}d/(d-1)$. Since $\phi(\tau)$ is an increasing function and d > 1, $I_2 \leq C(t-1)[\phi(\delta)]^{-1/d}$, concluding the proof.

Lemma 3.2. Suppose $F(u, u_x)$ is given by (4), with positive convergence radius r. Given L, d > 1, q > 2 and $\lambda \in [-1, 1]$, there exists $\epsilon_1 > 0$ such that, if $f \in \mathcal{B}_q$ and $\|f\| < \epsilon_1$, then, $\|N(u)\| < \|f\|$, for all $u \in B_f$.

Proof: Given $q > 2, f \in B_q, u \in B_f$ and $\alpha > 1$, we define

$$H_{\alpha}(u)(x,t) = c_{\alpha} \int_{1}^{t} \int G(x-y,\phi(\tau)) \left(u^{\alpha}\right)_{x} (y,\tau) dy d\tau, \ x \in \mathbb{R}, \ t \in [1,L],$$
(36)

where $\phi(\tau) = s(t) - s(t - \tau)$. Since $N(u) = \sum_{\alpha>1} H_{\alpha}(u)$, from Lemma 3.1, we get

$$\left|\widehat{N(u)}(\omega,t)\right| < \frac{B_t}{1+|\omega|^q} \sum_{\alpha>1} |c_\alpha| \left(\frac{G_q}{2\pi}\right)^{\alpha-1} \|u\|_L^{\alpha},\tag{37}$$

where B_t is given by (35) and

$$G_q = \left(2^{q+1} + 3\right) \int_{\mathbb{R}} \frac{1}{1 + |x|^q} dx.$$
 (38)

We notice that, given q > 2, if $f \in \mathcal{B}_q$ and $u \in B_f$, then $||u||_L \leq \overline{C}_L ||f||$, where

$$\bar{C}_L = 1 + K_0 + 2K_1 \left(s(L) \right)^{\frac{1}{d}} + K_2 \left(s(L) \right)^{\frac{2}{d}},$$
(39)

with K_i , for $i \in \{0, 1, 2\}$, given by (13). Also, there is K > 0 such that $|u(x,t)|, |u_x(x,t)| \leq K\bar{C}_L ||f||$ (see remark after Proposition 3.2 in [3]). Similarly, defining

$$\overline{B}_{t} \equiv (t-1) \left[K_{0} + K_{1} \left(\phi(t-1) \right)^{1/d} + C \right]$$
(40)

and

$$\overline{\overline{B}}_t \equiv 2K_0(t-1)K_1\left[(t-1) + 2\left(\phi(t-1)\right)^{1/d}(t-1) + 2(1+C)\right],\tag{41}$$

$$+K_2 \left(\phi(t-1)\right)^{1/d} \left(t-1\right) \left[\left(\phi(t-1)\right)^{1/d} + C \right] + 2C \left[\frac{d}{(d-1)t^{\frac{p}{d}}} + \frac{t-1}{\phi(\delta)^{\frac{1}{d}}} \right].$$

we get

$$\left|\partial_{\omega}\widehat{N(u)}(\omega,t)\right| < \frac{(2B_t + \overline{B}_t)}{(1+|\omega|^q)} \sum_{\alpha>1} |c_{\alpha}| \left(\frac{G_q}{2\pi}\right)^{\alpha-1} \|u\|_L^{\alpha}$$
(42)

and

$$\left|\partial_{\omega}^{2}\widehat{N(u)}(\omega,t)\right| < \frac{(3B_{t} + \overline{B}_{t} + \overline{\overline{B}}_{t})}{(1+|\omega|^{q})} \sum_{\alpha>1} |c_{\alpha}| \left(\frac{G_{q}}{2\pi}\right)^{\alpha-1} \|u\|_{L}^{\alpha}$$
(43)

Now define

$$r_0 = \min\left\{\frac{r}{K}, \frac{2\pi r}{G_q}\right\},$$

where r is the convergence radius of the sum (4). Since B_t, \overline{B}_t and \overline{B}_t are increasing functions for $t \in [1, L]$ and $\alpha \geq 2$, taking $||f|| < \overline{C_L}^{-1}r_0$, it follows that $||u||_L < r_0$ and, from (37), (42) and (43), we get

$$||N(u)|| < K_{L,q,d} \bar{C}_L^2 ||f||^2,$$
(44)

where

$$K_{L,q,d} = \left(\overline{\overline{B}}_L + 2\overline{B}_L + 6B_L\right) \sum_{\alpha>1} |c_\alpha| \left(\frac{G_q}{2\pi}\right)^{\alpha-1} r_0^{\alpha-2}.$$
 (45)

The result follows by defining $\epsilon_1 \equiv \min \left\{ (K_{L,q,d} \bar{C}_L^2)^{-1}, \bar{C}_L^{-1} r_0 \right\}.$

Lemma 3.3. Under the hypothesis of Lemma 3.2, there exists $\epsilon_2 > 0$ such that, if $f \in \mathcal{B}_q$ and $||f|| < \epsilon_2$, then $||N(u) - N(v)|| < \frac{1}{2}||u - v||$, for all $u, v \in B_f$.

Proof: Given q > 2 and $f \in B_q$, let $u, v \in B_f$ and consider $H_{\alpha}(u)(x,t)$ given by (36), for all $x \in \mathbb{R}$ and $t \in [1, L]$. Then, $[\widehat{H}_{\alpha}(u) - \widehat{H}_{\alpha}(v)](\omega, t)$ is given by

$$\frac{c_{\alpha} \cdot i}{(2\pi)^{\alpha-1}} \int_{1}^{t} \omega \widehat{G}(x-y,\phi(\tau)) \left(\widehat{u} \ast \cdots \ast \widehat{u} \ast \widehat{u} - \widehat{v} \ast \cdots \ast \widehat{v} \ast \widehat{v}\right) (y,\tau) d\tau, \qquad (46)$$

where in the integrand there are $\alpha - 1$ convolutions of \hat{u} and $\alpha - 1$ convolutions of \hat{v} . We obtain the following upper bound for $[\widehat{H_{\alpha}}(u) - \widehat{H_{\alpha}}(v)](\omega, t)$:

$$\left(\frac{G_t}{2\pi}\right)^{\alpha-1} \frac{|c_{\alpha}| B_t ||u-v||_L}{1+|\omega|^q} \left\{ ||u||_L^{\alpha-1} + ||u||_L^{\alpha-2} ||v||_L + \dots + ||u||_L ||v||_L^{\alpha-2} + ||v||_L^{\alpha-1} \right\},$$

where G_q and B_t are given, respectively, by (38) and (35). Therefore, we get, for $||f||_q < \bar{C_L}^{-1} r_0$,

$$\left| \left[\widehat{N(u)} - \widehat{N(v)} \right] (\omega, t) \right| < \frac{B_t}{1 + |\omega|^q} \|u - v\|_L \|u\|_L \sum_{\alpha > 1} |c_\alpha| \left(\frac{G_q}{2\pi} \right)^{\alpha - 2} r_0^{\alpha - 2}.$$

and, similarly,

$$\left|\partial_{\omega}\left[\widehat{N(u)}-\widehat{N(v)}\right](\omega,t)\right| < \frac{(2B_t+\overline{B}_t)}{(1+|\omega|^q)} \|u-v\|_L \|u\|_L \sum_{\alpha>1} |c_{\alpha}| \left(\frac{G_q}{2\pi}\right)^{\alpha-2} r_0^{\alpha-2}$$

and

$$\left|\partial_{\omega}^{2}\left[\widehat{N(u)}-\widehat{N(v)}\right](\omega,t)\right| < \frac{(3B_{t}+\overline{B}_{t}+\overline{\overline{B}}_{t})}{(1+|\omega|^{q})} \|u-v\|_{L} \|u\|_{L} \sum_{\alpha>1} |c_{\alpha}| \left(\frac{G_{q}}{2\pi}\right)^{\alpha-2} r_{0}^{\alpha-2},$$

which leads to

$$||N(u) - N(v)||_L < \overline{K}_{L,q,d} ||f||_q ||u - v||_L.$$

Defining

$$\overline{K}_{L,q,d} = \left(\overline{\overline{B}}_L + 2\overline{B}_L + 6B_L\right) \overline{C}_L \sum_{\alpha > 1} |c_\alpha| \left(\frac{G_q}{2\pi}\right)^{\alpha - 1} r_0^{\alpha - 2} \tag{47}$$

and $\epsilon_2 \equiv \min\{\left(2\overline{K}_{L,q,d}\right)^{-1}, \bar{C}_L^{-1}r_0\}$, we get the desired result.

3.2 Renormalization

In order to obtain the asymptotic behavior of solutions using the renormalization group approach, one must analyze the existence and stability of fixed points of an appropriate RG operator. Once such operator has been found for a particular problem, the method is iterative and the application of the RG transformation progressively evolves the solution in time and at the same time renormalizes the terms of the equation.

For each defined RG operator, we classify the nonlinearities of the problem according to their importance in the asymptotic behavior of the solution. This classification in the case of initial value problems is quite straightforward, with just a substitution of the renormalized solution to the equation. However, in the case of the integral equation, it is necessary to explore the properties of the kernel of the equation to obtain the classification, as we will show below. Let us consider equation (1) with $F(u) = u^a u^b_x$ and, given L > 1, define

$$u_n(x,t) \equiv L^{2n(p+1)/d} u(L^{n(p+1)/d}x, L^n t), \quad t \in [1,L], \quad n = 0, 1, 2, \cdots.$$
(48)

We will prove that $u_n(x,t)$ satisfies the renormalized equation

$$u_n(x,t) = \int \qquad G(x-y,s_n(t))f_n(y)dy + \\ \lambda_n \int_1^t \int G(x-y,s_n(t)-s_n(q))F(u_n(y,q))dydq,$$

where λ_n is the generalized coupling constant

$$\lambda_n = L^{n((-2a-3b+2)(p+1)+d)/d}\lambda,$$
(49)

$$F(u_n)(y,q) = L^{n(2a+3b)(p+1)/d} u^a (L^{n/d}y, L^n q) u^b_x (L^{n/d}y, L^n q)$$
(50)

and f_n is the renormalized initial data

$$f_n(x) \equiv L^{2n(p+1)/d} u(L^{n(p+1)/d}x, L^n).$$
(51)

In fact, if we write $u_n(x,t) = a(x,t) + b(x,t)$ where

$$a(x,t) \equiv L^{2n(p+1)/d} \int G(L^{n(p+1)/d}x - y, s(L^n t))f(y)dy + \lambda L^{2n(p+1)/d} \int_1^{L^n} \int G(L^{n(p+1)/d}x - y, s(L^n t) - s(\tau))u^a(y,\tau)u^b_x(y,\tau)dyd\tau$$

and

$$b(x,t) \equiv \lambda L^{2n(p+1)/d} \int_{L^n}^{L^n t} \int G(L^{n(p+1)/d} x - y, s(L^n t) - s(\tau)) u^a(y,\tau) u^b_x(y,\tau) dy d\tau,$$

it follows from conditions (G_2) and (G_3) that

$$a(x,t) = L^{2n(p+1)/d} \left[\int \int G(x-\omega,s_n(t))G(L^{n(p+1)/d}\omega-y,s(L^n))d\omega f(y)dy + \lambda \int_1^{L^n} \int \int G(x-\omega,s_n(t))G(L^{n(p+1)/d}\omega-y,s(L^n)-s(\tau))d\omega u^a(y,\tau)u^b_x(y,\tau)dyd\tau \right].$$

Using Fubini's theorem and definition (51) we get that $a(x,t) = \int G(x-\omega,s_n(t))f_n(\omega)d\omega$. Now if we consider the change of variables $y = L^{n(p+1)/d}\omega$, $\tau = L^n q$ and we divide and multiply the definition of b(x,t) by $L^{n\alpha(p+1)/d}$, where $\alpha = 2a + 3b$, then, it follows from condition (G₂) that

$$L^{n(p+1)/d}G(L^{n(p+1)/d}(x-\omega), s(L^n t) - s(L^n q)) = G(x-\omega, s_n(t) - s_n(q)).$$

Therefore, b(x,t) can be written as the second term of the equation (49), where $F(u_n)$ is given by (50), which concludes the proof.

We are then able to classify the nonlinearity $F(u, u_x)$ into its universality class, according to its asymptotic behavior: subcritical (or relevant) and critical (or marginal) nonlinearities are obtained when $2a + 3b < \frac{2(p+1)+d}{p+1}$ and $2a + 3b = \frac{2(p+1)+d}{p+1}$, respectively. We are interested in considering irrelevant perturbations, namely, those where $F(u) \sim u^a u_x^b$, with $2a + 3b > \frac{2(p+1)+d}{p+1}$.

After classifying the perturbations, we are able to obtain the renormalization step, which will guarantee the possibility of iterating the procedure. Given L > 1, consider $B^{(L)}$ and f_n given, respectively, by (32) and (51), and define $B_{f_n} \equiv \{u_n \in$ $B^{(L)} : ||u_n - u_{f_n}|| \le ||f_n||\}$ and $T_n(u_n) \equiv u_{f_n} + N_n(u_n)$, where u_{f_n} is the solution to the linear integral equation, that is, (49) with $\lambda_n = 0$, and

$$N_n(u_n)(x,t) = \lambda_n \int_1^t \int G(x-y, s_n(t) - s_n(\tau)) F_{L,n}(u_n(y,\tau)) dy d\tau$$
(52)

where $s_n(t)$, λ_n and $F_{L,n}(u_n)$ are respectively given by (16), (49) and (50). To simplify the notation, we define $\nu_n(x) \equiv N_n(u_n)(x, L)$. Note that, taking $||f_n|| < \epsilon$, Theorem 3.1 guarantees that the renormalized integral equation (49) has a unique solution that can be written in time t = L as $u_n(x, L) = u_{f_n}(x, L) + \nu_n(x)$. It follows that, for every positive integer n, the renormalization group operator for the integral equation (49) is well defined:

$$(R_{L,n}f_n)(x) \equiv L^{2(p+1)/d} u_n(L^{(p+1)/d}x, L),$$
(53)

which leads to the definition

$$f_0 = f$$
 and $f_{n+1} = R_{L,n} f_n$, for $n = 0, 1, 2, \cdots$ (54)

Using definitions (53) and (54) we get that

$$f_{n+1}(x) = R_{L,n}^0 f_n(x) + L^{(p+1)/d} \nu_n(L^{(p+1)/d}x),$$
(55)

where $R_{L,n}^0$ is the linear operator RG given by (17).

From now on, we will consider the space \mathcal{B}_q with q given by property (G_1) . We will also consider $\epsilon > 0$ given by Theorem 3.1, L_1 such that (21) holds and α_c given by (5).

Lemma 3.4. (Renormalization Lemma) Let q > 2, d > 1, $\alpha > \alpha_c$ and $L > L_1$ be given and consider the integral renormalized equation (49), where the initial data f_n , given by (54) has zero mass. If f_n satisfies

$$f_0 = A_0 G_p + g_0$$
 and $f_n = A_n R_{L^n}^0 G_p + g_n$, for all $n = 1, 2, \cdots$, (56)

where A_n is a constant, G_p is given by (12) and $g_n \in B_q$, with $\widehat{g_n}(0) = \widehat{g_n}'(0) = 0$, and $||f_n|| < \epsilon$, then

(a) f_{n+1} given by (54), can be decomposed as $f_{n+1} = A_{n+1}R_{L^{n+1}}^0G_p + g_{n+1}$, where A_{n+1} is a constant and

$$g_{n+1} = R_{L,n}^0 g_n + L^{2(p+1)/d} \nu_n (L^{(p+1)/d} \cdot) + i \hat{\nu}'_n(0) R_{L^n}^0 G_p.$$

Therefore, $g_{n+1} \in \mathcal{B}_q$ and $\widehat{g_{n+1}}(0) = \widehat{g_{n+1}}'(0) = 0$, In particular, f_{n+1} has zero mass.

(b)
$$|A_{n+1} - A_n| \leq L^{-n((2\alpha+1)(p+1)-d)/d} K_{L,q,d} \bar{C}_L^2 ||f_n||^2.$$

(c) $||g_{n+1}|| \leq \frac{C}{L^{(p+1)/d}} ||g_n|| + EL^{-n((2\alpha+1)(p+1)-d)/d} ||f_n||^2, where
 $E = E_{L,d,q,p,\alpha} \equiv (L^{(q+1)(p+1)/d} + C_{L,q,p}) K_{L,q,d} \bar{C}_L^2.$ (57)$

Proof: Consider the initial data f_n of (49) given by (54). By adding and subtracting the term $i\widehat{\nu_n}'(0)R_{L^n}^0G_p(x)$ on the right hand side of (55), we obtain the decomposition (56), where $A_{n+1} = A_n - i\widehat{\nu_n}'(0)$ and $g_{n+1}(x) = R_{L,n}^0g_n(x) + L^{2(p+1)/d}\nu_n(L^{(p+1)/d}x) + i\widehat{\nu_n}'(0)R_{L^n}^0G_p(x)$. Notice that $g_{n+1} \in B_q$ and $\widehat{g_{n+1}}(0) = \widehat{g_{n+1}}'(0) = 0$, since $\widehat{g_n}(0) = \widehat{g_n}'(0) = \widehat{G_p}(0) = 0$ and $\widehat{G_p}'(0) = i$. In particular, f_{n+1} has zero mass. Item (b) is straightforward from (44) and (49). Item (c) follows from the decomposition of g_{n+1} , the Contraction Lemma and (44).

Given $\alpha > \alpha_c$, consider $\delta \in (0, 1)$ such that

$$(1-\delta)(p+1) < (2\alpha+1)(p+1) - d,$$
(58)

and define

$$L_{\delta} = \max\left\{L_{1}, 2C\left(1 + C_{d,q,p}\right)^{\frac{d}{\delta(p+1)}}\right\},$$
(59)

where C is the constant given in the Contraction Lemma 2.7 and $C_{d,q,p}$ is given by (23). Throughout the rest of this paper, L_{δ} will be referring to the constant above.

Lemma 3.5. Given $\delta \in (0,1)$, α satisfying (58) and $L > L_{\delta}$, if f_0 satisfies

$$\|f_0\| < \frac{1}{2L^{(1-\delta)(p+1)/d}ED^2},\tag{60}$$

with E given by (57) and

$$D \equiv 1 + \tilde{K} \sum_{j=0}^{+\infty} \frac{1}{L^{j(p+1)(1-\delta)/d}},$$
(61)

then

$$\frac{ED^2 \|f_0\|}{L^{n[(2\alpha+1)(p+1)-d]/d}} < \frac{1}{2L^{(n+1)(1-\delta)(p+1)/d}},$$

for all $n \in \mathbb{Z}^+$.

Proof: Since δ and α satisfy (58) and f_0 satisfies (60):

$$\frac{ED^2 \|f_0\|}{L^{n[(2\alpha+1)(p+1)-d]/d}} < \frac{1}{2L^{(n+1)(1-\delta)(p+1)/d}}.$$

Lemma 3.6. Suppose (60) holds and define

$$D_1 \equiv \frac{1}{L^{(1-\delta)(p+1)/d}} + \tilde{K} \left(1 + K_{L,q,d} \bar{C_L}^2 \| f_0 \| \right)$$

and

$$D_{n+1} = \frac{1}{L^{(1-\delta)(p+1)(n+1)/d}} + \tilde{K}\left(1 + K_{L,q,d}\bar{C_L}^2 \|f_0\| + K_{L,q,d}\bar{C_L}^2 \|f_0\| \sum_{j=1}^n \frac{D_j^2}{L^{j[(2\alpha+1)(p+1)-d]/d}}\right)$$

where constants \bar{C}_L , $K_{L,q,d}$ and \tilde{K} are defined in (39), (45) and Lemma 2.5. Then,

$$D_{n+1} < D, \forall n \in \mathbb{Z}^+$$

Proof: Since $E > K_{L,q,d} \bar{C_L}^2$ and f_0 satisfies inequality (60), then

$$D_1 < \frac{1}{L^{(1-\delta)(p+1)/d}} + \tilde{K}\left(1 + E \|f_0\|\right) < 1 + \tilde{K}\left(1 + \frac{1}{2L^{(1-\delta)(p+1)/d}D^2}\right) < D.$$

Now suppose that, for some $n_0 \in \mathbb{Z}^+_*$, $D_n < D$ for $n \in \{1, \dots, n_0 - 1\}$. Then, using the induction hypothesis and the previous lemma, we can bound D_{n_0} by

$$\frac{1}{L^{(1-\delta)(p+1)(n+1)/d}} + \tilde{K}\left(1 + \frac{1}{2L^{(1-\delta)(p+1)/d}} + \sum_{j=1}^{n_0-1} \frac{1}{2L^{j(n+1)(1-\delta)(p+1)/d}}\right),$$

which finishes the proof

The results of the next theorem will guarantee that all renormalized equations will have a unique solution so that the algorithm iterations will be well defined.

Theorem 3.2. Given $\delta \in (0, 1)$ satisfying inequality (58) and $L > L_{\delta}$, if f_0 is a zero mass function, there exists $\bar{\epsilon} > 0$ such that, if $||f_0|| < \bar{\epsilon}$, then, for all $n \in \{1, 2, \dots\}$, f_n given by (54), is well defined, has zero mass and admits the representation (56), where g_n has zero mass, zero first moment and satisfies

$$\|g_n\| \le \frac{\|f_0\|}{L^{n(p+1)(1-\delta)/d}}.$$
(62)

Furthermore, f_n satisfies

$$||f_n|| \le D||f_0||,\tag{63}$$

where D is the constant given by (61). In particular, $||f_n|| < \epsilon$, with $\epsilon > 0$ given by Theorem 3.1.

Proof: Define

$$\bar{\epsilon} \equiv \min\left\{\frac{1}{2L^{(p+1)(1-\delta)/d}ED^2}, \frac{\epsilon}{D}\right\}$$
(64)

where $\epsilon > 0$ is given by Theorem 3.1 and the constants E and D are given by (57) and (61), respectively. Suppose that, given $n_0 \in \mathbb{Z}^+_*$, the hypotheses are valid for all $n \in \{0, 1, \dots, n_0\}$. Then, since $||f_0|| < \bar{\epsilon}$, it follows that $||f_n|| < \epsilon$, for all $n \in \{0, 1, \dots, n_0\}$. From Lemma 3.4, for all $n \in \{0, 1, \dots, n_0\}$, f_{n+1} can be decomposed as $f_{n+1} = A_{n+1}R_{L^{n+1}}^0G_p + g_{n+1}$, where

$$g_{n+1} = R^0_{L,n}g_n + L^{2(p+1)/d}\nu_n(L^{(p+1)/d}\cdot) + i\hat{\nu}'_n(0)R^0_{L^n}G_p.$$

Moreover, $g_{n+1} \in \mathcal{B}_q$ and $\widehat{g_{n+1}}(0) = \widehat{g_{n+1}}'(0) = 0$. We also have

$$|A_{n+1} - A_n| \le \frac{K_{L,q,d}\bar{C}_L^2}{L^{n((2\alpha+1)(p+1)-d)/d}} \|f_n\|^2 \text{ and } \|g_{n+1}\| \le \frac{\|f_0\|}{L^{(n+1)(p+1)(1-\delta)/d}},$$

since $L > L_{\delta}$ and $||f_0|| < \bar{\epsilon}$. Therefore, since

$$|f_{n+1}|| \le ||g_n|| + \tilde{K} \left[\sum_{i=0}^n (|A_{i+1} - A_i|) + |A_0| \right],$$

it follows inequality (63).

Proof of Theorem 1.1: In order to prove Theorem 1.1, we first prove that, under the hypotheses of Theorem 3.2, there exists $A = A(d, f_0, F, p, q)$ such that, if $||f_0|| < \bar{\epsilon}$, then,

$$\lim_{n \to \infty} \|L^{2n(p+1)/d} u(L^{n(p+1)/d}, L^n) - AG_p\| = 0.$$

In fact, as $||f_0|| < \bar{\epsilon}$, it follows from the Renormalization Lemma, from the semigroup property of the RG operator and from (63) that $f_n = A_n R_{L^n}^0 G_p + g_n = L^{2n(p+1)/d} u(L^{n(p+1)/d}.,L^n)$ and $|A_{n+1} - A_n| \leq L^{2n(p+1)/d} u(L^{n(p+1)/d}.,L^n)$

 $L^{-n[(2\alpha+1)(p+1)-d]/d}K_{L,q,d}\bar{C}_L^2D^2||f_0||^2$. Since $n((2\alpha+1)(p+1)-d)/d > 0$, there is A such that $A_n \to A$ and, using that $||f_0|| < \bar{\epsilon}, E > K_{L,q,d}\bar{C}_L^2$ and $L > L_{\delta}$, we get

$$|A_n - A| < \frac{1}{2L^{(p+1)(1-\delta)}} \cdot \frac{L^{-n[(2\alpha+1)(p+1)-d]/d}}{1 - L^{-[(2\alpha+1)(p+1)-d]/d}} ||f_0||.$$
(65)

Using the triangle inequality, we bound $||L^{2n(p+1)/d}u(L^{n(p+1)/d}.,L^n) - AG_p||$ by

$$|A| ||R_{L^n}^0 G_p - G_p|| + ||L^{2n(p+1)/d} u(L^{n(p+1)/d}, L^n) - A_n R_{L^n}^0 G_p|| + |A_n - A| ||R_{L^n}^0 G_p||.$$

Inequality (10) will follow by taking the limit $n \longrightarrow +\infty$. Now, using Lemmas 2.5, 2.6 and inequalities (62), (65), we get

$$\begin{split} \|t^{2(p+1)/d}u(t^{(p+1)/d},t) - AG_p(.)\| &\leq |A| \left(M \left| \frac{r(t)}{t^{(p+1)}} \right|^{\frac{1}{d}} \right. \\ &+ N \left| \frac{r(t)}{t^{(p+1)}} \right|^{\frac{2}{d}} \right) + \frac{\|f_0\|}{t^{(p+1)(1-\delta)/d}} + \frac{1}{2L_{\delta}^{(p+1)(1-\delta)}} \cdot \frac{t^{-[(2\alpha+1)(p+1)-d]/d}}{1 - L_{\delta}^{-[(2\alpha+1)(p+1)-d]/d}} \tilde{K} \|f_0\|. \end{split}$$

References

- G. A. Braga, J. M. Moreira and Camila F. Souza. Asymptotics for Nonlinear Integral Equations with Generalized Heat Kernel and Time Dependent Coefficients Using Renormalization Group Technique. Journal of Mathematical Physics, 60, Issue 1, 013507, 2019.
- [2] G. A. Braga, J. M. Moreira and Camila F. Souza. Asymptotics for Nonlinear Integral Equations with a Generalized Heat Kernel using Renormalization Group Technique II: Marginal Perturbations and Logarithmic Corrections to the Time Decay of Solutions. Journal of Mathematical Physics, 62, 083507, 2021, https://doi.org/10.1063/5.0043505.
- [3] Braga, G.A., Furtado, F., Moreira, J.M. and Da Silva, A. M. The Asymptotic Leading Term of Decaying Solutions of Some Generalized Burgers' Equations. Mediterr. J. Math., 22, 10, 2025, https://doi.org/10.1007/s00009-024-02772-4
- [4] J. Bricmont, A. Kupiainen and G. Lin. Renormalization group and asymptotics of solutions of nonlinear parabolic equations. *Communications in Pure and Applied Mathematics*, 47, 893–922, 1994.
- [5] J. L. Bona, K. S. Promislow and C. E. Wayne. Higher-order asymptotics of decaying solutions of some nonlinear, dispersive, dissipative wave equations. *Nonlinearity*, 8, 1179-1206, 1995.
- [6] G. Braga, F. Furtado, V. Isaia, and L. Lee. Numerical Renormalization Group Algorithms for Self-Similar Asymptotics of Partial Differential Equations. *Multiscale Modeling and Simulation*, SIAM, 18, p. 131-162, 2020.