

Information Limits of Joint Community Detection and Finite Group Synchronization

Yifeng Fan*

Zhizhen Zhao*

Abstract

The emerging problem of joint community detection and group synchronization, with applications in signal processing and machine learning, has been extensively studied in recent years. Previous research on this topic has predominantly focused on a statistical model that extends the well-known stochastic block model (SBM) by incorporating additional group transformations. In its simplest form, the model randomly generates a network of size n that consists of two equal-sized communities, where each node i is associated with an unknown group element $g_i^* \in \mathcal{G}_M$ for some finite group \mathcal{G}_M of order M . The connectivity between nodes follows a probability p if they belong to the same community, and a probability q otherwise. Moreover, a group transformation $g_{ij} \in \mathcal{G}_M$ is observed on each edge (i, j) , where $g_{ij} = g_i^* - g_j^*$ if nodes i and j are within the same community, and $g_{ij} \sim \text{Uniform}(\mathcal{G}_M)$ otherwise. The goal of the joint problem is to recover both the underlying communities and group elements. Under this setting, when $p = a \log n/n$ and $q = b \log n/n$ with $a, b > 0$, we establish the following sharp information-theoretic threshold for exact recovery by maximum likelihood estimation (MLE):

$$(i): \frac{a+b}{2} - \sqrt{\frac{ab}{M}} > 1 \quad \text{and} \quad (ii): a > 2$$

where the exact recovery of communities is possible with high probability only if (i) is satisfied, and the recovery of group elements is achieved with high probability only if both (i) and (ii) are satisfied. Our theory indicates the recovery of communities greatly benefits from the extra group transformations. Also, it demonstrates a significant performance gap exists between the MLE and all the existing approaches, including algorithms based on semidefinite programming and spectral methods.

1 Introduction

Community detection and group synchronization are both fundamental problems in signal processing, machine learning, and computer vision. Recently, there has been increasing interest in the joint problem of these two areas [24, 26, 23, 12, 6, 66]. That is, in the presence of heterogeneous data where each sample is associated with an unknown group element (e.g. the orthogonal groups $O(d)$) and falls into multiple underlying communities, a network is observed that represents the interactions between data samples including the group transformations. Then, the joint problem aims to simultaneously recover the underlying community structures and the unknown group elements. A motivating example is the 2D class averaging process in *cryo-electron microscopy single-particle reconstruction* [29, 62, 69, 22], whose goal is to group and rotationally align projections images of

*Y. Fan is with the Department of Electrical and Computer Engineering, University of Illinois at Urbana-Champaign, Champaign, IL, 94404, USA (email: yifengf2@illinois.edu). Z. Zhao is with the Department of Electrical and Computer Engineering, University of Illinois at Urbana-Champaign, Champaign, IL, 94404, USA (email: zhizhenz@illinois.edu)




Figure 1: An illustration of our statistical model. We present a network that consists of two communities of equal sizes, shown in circles and squares respectively. Each node i is associated with an unknown group element $g_i^* \in \mathcal{G}_M$ for some finite group \mathcal{G}_M of size M . Each pair of nodes within the same community (resp. across different communities) are independently connected with probability p (resp. q) as shown in solid (resp. dash) lines. A group transformation g_{ij} is observed on each edge which satisfies $g_{ij} = g_i^*(g_j^*)^{-1}$ when nodes i and j are within the same community, otherwise $g_{ij} \sim \text{Unif}(\mathcal{G}_M)$ that is uniformly drawn from \mathcal{G}_M . Given the network, our goal is to recover the underlying communities and the associated group elements $\{g_i^*\}_{i=1}^n$.

a single particle with similar viewing angles, in order to improve their signal-to-noise ratio (SNR). Another application in computer vision is *simultaneous permutation group synchronization and clustering* on heterogeneous object collections consisting of 2D images or 3D shapes [6, 41, 40].

This paper centers on a hybrid statistical model initially proposed in [24], which combines the celebrated stochastic block model (SBM) (e.g. [17, 18, 19, 28, 39, 43, 50, 51, 53, 54]) for community detection and the random rewiring model (e.g. [62, 26, 21, 48, 65, 14, 13, 40]) for group synchronization. Notably, most of the existing methods [24, 23, 12, 66] for solving the joint problem are designed and evaluated based on this model. Formally, the model generates a random network (graph) of size n with K underlying communities, each node i is associated with its community assignment $\kappa_i^* \in \{1, \dots, K\}$, along with an unknown group element $g_i^* \in \mathcal{G}_M$ for some finite group \mathcal{G}_M of order M (e.g. the symmetric group S_M and cyclic group Z_M). Then, edges are placed randomly and independently between nodes (i, j) with probability p if i and j belong to the same community, and with probability q otherwise. In addition, a group transformation $g_{ij} \in \mathcal{G}_M$ is observed on each edge connection (i, j) in a similar way that $g_{ij} = g_i^*(g_j^*)^{-1}$ if i and j are within the same community, which represents the clean measurement, otherwise $g_{ij} \in \text{Unif}(\mathcal{G}_M)$ that is uniformly drawn from \mathcal{G}_M implying the measurement is completely noisy. In this paper, we focus on the case when $K = 2$ and the two communities are of equal sizes $n/2$, see Figure 1 for an illustration.

Given the network observed, our goal is to recover the underlying communities as well as the unknown group elements. Notably, a naive two-stage approach, which first applies classical graph-clustering algorithms (e.g. [37, 30, 27, 2]) followed by synchronization within each identified community, leads to sub-optimal results, since it does not leverage the consistency (resp. inconsistency) of group transformations within each community (resp. across different communities). To address this limitation, several improved optimization criteria are proposed [6, 24, 66] incorporating these consistencies. Due to the non-convex and computationally intractable nature of directly solving such optimization programs, convex relaxation techniques such as semidefinite programming (SDP) [24] and spectral method [23, 6, 12, 66] have emerged, which provide approximated solutions with

polynomial time complexity. Importantly, such efficient methods have significantly enhanced the recovery accuracy compared to the aforementioned two-stage approach.

Despite the progress made before, various fundamental questions remain unsettled such as:

1. *What is the fundamental limit for achieving exact recovery of the clusters or group elements?*
2. *Is the fundamental limit achieved by the existing algorithms?*

Finding this limit enables us to benchmark the performance of existing algorithms and assess their potential for improvement. Notably, the optimal algorithm that maximizes the probability of exact recovery points to the *maximum a posteriori* (MAP) estimation. Meanwhile, since there is no preference regarding the underlying community assignment and group elements, the MAP estimation is equivalent to the *maximum likelihood* estimation (MLE) under a uniform prior. Therefore, the performance of MLE serves as the *information-theoretic limits* such that if MLE fails in exact recovery with high probability as the network size n grows, no algorithm, regardless of its complexity, can succeed with high probability.

Our contribution In this work, we answer the two questions above by establishing the information-theoretic limits for the exact recovery of (i) communities, (ii) group elements, and both of them, under the statistical model aforementioned. For ease of presentation, we restrict our attention to the simplest scenario of two communities ($K = 2$) of equal size, which allows us to convey the most informative findings without introducing unnecessary complexity. In a nutshell, on the sparse regime such that $p = \frac{a \log n}{n}$ and $q = \frac{b \log n}{n}$ for some constants $a, b > 0$, and as n is sufficiently large, we identify sharp information-theoretic thresholds for the exact recovery of (i) and (ii) above as follows:

- (i) Recovering the communities is possible if and only if

$$\frac{a+b}{2} - \sqrt{\frac{ab}{M}} > 1$$

- (ii) Recovering the group elements is possible if and only if

$$\frac{a+b}{2} - \sqrt{\frac{ab}{M}} > 1 \quad \text{and} \quad \frac{a}{2} > 1$$

Here, several remarks are worthy to be highlighted:

1. The recovery of communities is the *prerequisite* for recovering the group elements. In other words, one cannot hope to identify the group elements without determining the communities at first.
2. When $M = 1$, our model degenerates to the SBM and the condition in (i) becomes

$$(\sqrt{a} - \sqrt{b})^2 > 2$$

which agrees with the existing result of information limit on the SBM [1, 3, 50, 54].

3. By examining the performance of existing efficient algorithms [23, 6, 12, 66, 24], our theory demonstrates a significant performance gap between the information limit and those approaches (see Figure 2), suggesting a considerable room of improvement exists.
4. Our proof strategy is substantially distinct from (and more difficult than) previous works (e.g. [14, 1, 54, 61]) which focuses on community detection or group synchronization individually, and the difficulty comes from dealing with the coupling of these two problems.

Organization The rest of this paper is organized as follows: In the remaining of Section 1, we introduce some additional related works and the notations for our analysis. Then, in Section 2 we formulate the MLE program for recovery on our statistical model and define the evaluation measurement. Next, in Section 3 we present our theory on the information-theoretic limits, and Section 4 is devoted to discussions on the result. We end up with the proofs in Section 5, and leave the technical details to the Appendix.

1.1 Related works

Given its practical importance to various scientific applications, either community detection or group synchronization has been extensively studied over the past decades. Therefore, this section by no means provides a comprehensive review of all the previous works but only highlights the most related ones.

Community detection is commonly studied on the SBM (e.g. [17, 18, 19, 28, 39, 43, 50, 51, 53, 54]) where maximum likelihood is equivalent to minimizing the number of edges between different communities. Then, a major line of research (e.g. [9, 19, 8, 63, 42, 16, 10, 51, 58, 15]) attempts to find the information-theoretic limits for exact recovery, especially on the case of two communities, until the two seminal papers [1, 52] first characterize a sharp threshold for the case of two equal-sized communities, which greatly inspires the present work and is further extended to more general scenarios (e.g. [3, 38]). In terms of algorithms for recovery, as exactly solving the MLE is often NP-hard, different approaches such as semidefinite programming (SDP) [1, 37, 38, 57, 36, 4, 7], spectral method [2, 64, 67, 50, 44, 55, 32], and belief propagation [17, 3] are proposed for obtaining an approximated solution to the MLE. However, many of them have been proven optimal such that they can achieve the information-theoretic limits at least on two equal-sized communities [37, 38, 2, 67, 46, 57, 32].

On the side of group synchronization, the goal is to recover the underlying group elements from a set of noisy pairwise measurements. A large body of literature focuses on the aforementioned random rewiring model (e.g. [62, 26, 21, 48, 65, 14, 13, 40]) which independently replaces each pairwise clean measurement with some random element. The fundamental information limits for exact recovery are generally studied in [14]. Regarding the algorithms, similar to the development of community detection, various approaches are developed for recovery depending on the specific group of interest, including semidefinite relaxation [61, 40], spectral methods [61, 5, 11, 56, 59, 34], and message passing [60], along with many theoretical investigations on their performances [61, 70, 48, 40, 49, 33, 68].

The joint problem of community detection and group synchronization is an emerging research topic (e.g. [26, 21, 6, 45]) motivated by recent scientific applications such as the cryo-electron microscopy single-particle reconstruction [29, 62, 69, 22] aforementioned. In particular, the statistical model adopted in this paper was initially proposed in [24], which also studied exact recovery by SDP with the performance guarantee provided. Recently, several spectral method-based approaches [23, 12, 66] are designed for efficient recovery. As there is a lack of indicators on the

Table 1: A summary of notations for the model. The notations with * indicate the ground truth parameters.

Definition	Description
$\boldsymbol{\kappa}^* := \{\kappa_1^*, \dots, \kappa_n^*\}, \quad \boldsymbol{\kappa} := \{\kappa_1, \dots, \kappa_n\}$	The set of true (estimated) community assignment
$\boldsymbol{g}^* := \{g_1^*, \dots, g_n^*\}, \quad \boldsymbol{g} := \{g_1, \dots, g_n\}$	The set of true (estimated) group elements
$S_k^* := \{i \kappa_i^* = k\}, \quad S_k := \{i \kappa_i = k\}$	The set of nodes in the k -th true (estimated) community
$n_k^* := S_k^* , \quad n_k := S_k $	The size of the k -th true (estimated) community
$\boldsymbol{x}^* := \{\boldsymbol{\kappa}^*, \boldsymbol{g}^*\}, \quad \boldsymbol{x} := \{\boldsymbol{\kappa}, \boldsymbol{g}\}$	The true (estimated) parameters
\mathcal{E}	The set of edge connections
$\mathfrak{G} := \{g_{ij} (i, j) \in \mathcal{E}\}$	The set of group transformations
$\boldsymbol{y} := \{\mathcal{E}, \mathfrak{G}\}$	The observation of the network
$\mathcal{E}(S)$	The set of edges within the node set S
$\mathcal{E}(S_1, S_2)$	The set of edges across the node sets S_1 and S_2

performance of these algorithms, the information-theoretic limit obtained in this work fills in the blanks.

1.2 Notations

Here, we define a series of notations for our statistical model that will be frequently used. Given the network of size n with K underlying communities, we denote $\boldsymbol{\kappa}^* := \{\kappa_1^*, \dots, \kappa_n^*\}$ as the set of true community assignment, and $\boldsymbol{g}^* := \{g_1^*, \dots, g_n^*\}$ as the set of true group elements. Similarly, the estimated community assignments and group elements are denoted as $\boldsymbol{\kappa} := \{\kappa_1, \dots, \kappa_n\}$ and $\boldsymbol{g} := \{g_1, \dots, g_n\}$, respectively. We use n_k^* and S_k^* to denote the size of the k -th community and the set of nodes belonging to it, respectively, i.e. $S_k^* := \{i | \kappa_i^* = k\}$ and $n_k^* := |S_k^*|$ for $k = 1, \dots, K$, and we define $S_k := \{i | \kappa_i = k\}$ and $n_k := |S_k|$ analogous to S_k^* and n_k^* . Then, we denote $\boldsymbol{x}^* := \{\boldsymbol{\kappa}^*, \boldsymbol{g}^*\}$ and $\boldsymbol{x} := \{\boldsymbol{\kappa}, \boldsymbol{g}\}$ as the true and estimated parameters. By denoting \mathcal{E} as the set of edge connections and $\mathfrak{G} := \{g_{ij} | (i, j) \in \mathcal{E}\}$ as the set of group transformations observed on each edge, we define $\boldsymbol{y} := \{\mathcal{E}, \mathfrak{G}\}$ as the whole observation of the network. Lastly, given any set of nodes S , we define $\mathcal{E}(S) = \{(i, j) | i, j \in S, (i, j) \in \mathcal{E}\}$ as the set of edge connections with S . For any two sets of nodes S_1 and S_2 , we denote $\mathcal{E}(S_1, S_2) = \{(i, j) | i \in S_1, j \in S_2, (i, j) \in \mathcal{E}\}$ as the set of edges connecting S_1 and S_2 . We summarize these notations in Table 1.

Besides, we use the following standard notations for our analysis: for two non-negative functions $f(n)$ and $g(n)$, $f(n) = O(g(n))$ means there exists an absolute positive constant C such that $f(n) \leq Cg(n)$ for all sufficiently large n ; $f(n) = o(g(n))$ indicates for every positive constant C , the inequality $f(n) \leq Cg(n)$ holds for all sufficiently large n ; $f(n) = \omega(g(n))$ denotes for every positive constant C , the inequality $f(n) \geq Cg(n)$ for all sufficiently large n ; and $f(n) = \Theta(g(n))$ represents there exists two absolute positive constants C_1, C_2 such that $C_1g(n) \leq f(n) \leq C_2g(n)$.

2 Preliminary

2.1 Maximum likelihood estimation

Given an observation $\boldsymbol{y} = \{\mathcal{E}, \mathfrak{G}\}$ of the network, our goal is to recover the true parameters $\boldsymbol{x}^* = \{\boldsymbol{\kappa}^*, \boldsymbol{g}^*\}$ including both the community assignment $\boldsymbol{\kappa}^*$ and the group elements \boldsymbol{g}^* . Recall that in

our statistical model, each pair of nodes are independently connected by following

$$\mathcal{P}((i, j) \in \mathcal{E}) = \begin{cases} p, & \kappa_i^* = \kappa_j^* \\ q, & \text{otherwise} \end{cases}$$

Also, the group transformation g_{ij} observed on each edges $(i, j) \in \mathcal{E}$ follows

$$g_{ij} = \begin{cases} g_i^* (g_j^*)^{-1}, & \kappa_i^* = \kappa_j^* \\ \text{Uniform}(\mathcal{G}_M), & \text{otherwise} \end{cases}$$

Given the above, this work focuses on the maximum likelihood estimator (MLE) that maximizes $\mathcal{P}(\mathbf{y}|\mathbf{x})$. Formally, we have

Claim 2.1. *The maximum likelihood estimator (MLE) for estimating $\mathbf{x}^* = \{\boldsymbol{\kappa}^*, \mathbf{g}^*\}$ from $\mathbf{y} = \{\mathcal{E}, \mathfrak{G}\}$ is*

$$\psi_{\text{MLE}}(\mathbf{y}) = \begin{cases} \arg \max_{\boldsymbol{\kappa}, \mathbf{g}} |\mathcal{E}_{\text{inner}}|, & \frac{Mp(1-q)}{q(1-p)} > 1 \\ \arg \min_{\boldsymbol{\kappa}, \mathbf{g}} |\mathcal{E}_{\text{inner}}|, & \text{otherwise} \end{cases} \quad (1)$$

$$\text{s.t. } n_k = n_k^*, \quad k = 1, \dots, K$$

$$g_{ij} = g_i g_j^{-1}, \quad \forall (i, j) \in \mathcal{E}_{\text{inner}}$$

where

$$\mathcal{E}_{\text{inner}} := \{(i, j) \mid (i, j) \in \mathcal{E}, i < j, \kappa_i = \kappa_j\} \quad (2)$$

denotes the set of edges within the same community.

The derivation of (1) is deferred to Section 5. Here, we assume the cluster sizes $\{n_k^*\}_{k=1}^K$ are given and therefore the constraint $n_k = n_k^*$ ensures our recovery is valid.

According to (1), finding the MLE is equivalent to performing the following two tasks simultaneously:

1. (*For clustering*) Partitioning the data network into K disjoint clusters such that the number of in-cluster edges is *maximized* or *minimized*.
2. (*For synchronization*) On each in-cluster edge, the observed group transformation g_{ij} should satisfy the *consistency* $g_{ij} = g_i g_j^{-1}$.

Here, as a special case when $M = 1$ such that no group synchronization is needed, (1) essentially becomes the MLE on the SBM:

$$\psi_{\text{MLE}}(\mathbf{y}) = \begin{cases} \arg \max_{\boldsymbol{\kappa}} |\mathcal{E}_{\text{inner}}|, & p > q \\ \arg \min_{\boldsymbol{\kappa}} |\mathcal{E}_{\text{inner}}|, & \text{otherwise} \end{cases} \quad (3)$$

$$\text{s.t. } n_k = n_k^*, \quad k = 1, \dots, K$$

which simply maximizes or minimizes the number of in-cluster edges. As a result, (1) can be treated as an extension of (3) with additional constraints on the group elements included.

In this work, although we focus on the scenario when \mathcal{G}_M is a finite group with order M , notice that by letting $M \rightarrow \infty$ that goes to infinity, (1) becomes

$$\begin{aligned} \psi_{\text{MLE}}(\mathbf{y}) &= \arg \max_{\boldsymbol{\kappa}, \mathbf{g}} |\mathcal{E}_{\text{inner}}| \\ \text{s.t. } \quad n_k &= n_k^*, \quad k = 1, \dots, K \\ g_{ij} &= g_i g_j^{-1}, \quad \forall (i, j) \in \mathcal{E}_{\text{inner}} \end{aligned} \tag{4}$$

which is the MLE for the case of an infinite group, e.g. the orthogonal group $O(d)$ and the rotation group $SO(d)$. In this scenario, we always aim to maximize the number of in-cluster edges regardless of the model parameters.

From an algorithm perspective, the program (1) is obviously non-convex and is thus computationally intractable to be exactly solved. Moreover, even solving (3) is well-known to be NP-hard, including the famous *minimum bisection* problem (see e.g. [35]) as a special case when the network consists of two equal-sized communities. This challenge gives rise to several approximate but efficient algorithms proposed in [23, 12, 66, 25].

2.2 Evaluation measurement

This paper centers on exact information recovery that the true parameter $\boldsymbol{\kappa}^*$ and \mathbf{g}^* are precisely recovered by (27). For evaluating the recovery of clusters, given any cluster assignments $\boldsymbol{\kappa}$, we introduce the following *zero-one distance* between $\boldsymbol{\kappa}^*$ and $\boldsymbol{\kappa}$:

$$\text{dist}_c(\boldsymbol{\kappa}, \boldsymbol{\kappa}^*) := 1 - \max_{\pi \in \Pi_K} \mathbb{1}(\kappa_i^* = \pi(\kappa_i); i = 1, \dots, n) \tag{5}$$

where Π_K denotes the permutation group on $\{1, \dots, K\}$. Notably, Π_K is introduced to eliminate the ambiguity on the permutation of cluster labels. That is, any permutation on the labels does not affect the underlying partition, and thus the evaluation metric should be invariant to permutations.

The same idea applies to the metric for group elements: one can only identify them within each community up to some global offset $\bar{g} \in \mathcal{G}_M$, as there is no basis to distinguish g_i with $g_i \bar{g}$ given the pairwise transformations. Therefore, similar to (5), we define

$$\text{dist}_g(\mathbf{g}, \mathbf{g}^*) := \sum_{k=1}^K \left[1 - \max_{\bar{g}_k \in \mathcal{G}_M} \mathbb{1}(g_i^* = g_i \bar{g}_k; \forall i \in S_k^*) \right] \tag{6}$$

as the distance between \mathbf{g} and \mathbf{g}^* . Here, we compute the distance on each cluster by maximizing over its offset \bar{g}_k , and then sum over all clusters. As a result, (6) is zero only if all the group elements are exactly recovered.

With all the metrics in place, we define, for any recovery procedure $\psi(\mathbf{y})$ which takes some observation $\mathbf{y} = \{\mathcal{E}, \mathcal{G}\}$ as input and output the estimation $\mathbf{x} = \{\boldsymbol{\kappa}, \mathbf{g}\}$, the error probability of $\boldsymbol{\kappa}$ and \mathbf{g} as

$$\begin{aligned} \mathcal{P}_{e,c}(\psi) &= \max_{\mathbf{x}^* = \{\boldsymbol{\kappa}^*, \mathbf{g}^*\}} \mathcal{P}(\text{dist}_c(\boldsymbol{\kappa}, \boldsymbol{\kappa}^*) \neq 0 \mid \mathbf{x}^*), \\ \mathcal{P}_{e,g}(\psi) &= \max_{\mathbf{x}^* = \{\boldsymbol{\kappa}^*, \mathbf{g}^*\}} \mathcal{P}(\text{dist}_g(\mathbf{g}, \mathbf{g}^*) \neq 0 \mid \mathbf{x}^*). \end{aligned} \tag{7}$$

In other words, $\mathcal{P}_{e,c}(\psi)$ (resp. $\mathcal{P}_{e,g}(\psi)$) stands for the maximum probability of giving a wrong estimation $\boldsymbol{\kappa}$ (resp. \mathbf{g}) by ψ over all possible ground truth \mathbf{x}^* . We remark that in our model setting, the error probability does not depend on the specific choice of \mathbf{x}^* as there is no preference. Therefore, one should expect both $\mathcal{P}(\text{dist}_c(\boldsymbol{\kappa}, \boldsymbol{\kappa}^*) \neq 0 \mid \mathbf{x}^*)$ and $\mathcal{P}(\text{dist}_g(\mathbf{g}, \mathbf{g}^*) \neq 0 \mid \mathbf{x}^*)$ are invariant to \mathbf{x}^* .

3 Main results

This section presents the information-theoretic limits for exact recovery on the statistical model defined in Section 1. Notably, even though we aim to recover both the (i) communities and (ii) group elements simultaneously, there might exist scenarios where the recovery of one is possible but impossible for the another. Because of this, we study the conditions for recovering (i) and (ii) separately. Specifically, for each of them, we provide

- Information-theoretic *upper* bound, above which the MLE (1) achieves exact recovery with high probability (w.h.p.)
- Information-theoretic *lower* bound, below which the MLE (1) fails to recover with a probability bounded away from zero

In the rest of our analysis, we focus on the scenario of having two communities of equal sizes, and the order of group M is assumed to be a fixed integer as n grows.

3.1 Exact recovery of communities

Theorem 3.1 (Upper bound). *Under the setting of two equal-sized clusters. Let $p = \frac{a \log n}{n}$ and $q = \frac{b \log n}{n}$ for $a, b \geq 0$. For any $c > 0$, if*

$$\frac{a+b}{2} - (1+o(1))\sqrt{\frac{ab}{M}} > 1+c \quad (8)$$

then the MLE (1) exactly recovers the communities w.p. at least $1 - n^{-c}$, i.e. $\mathcal{P}_{e,c}(\psi_{\text{MLE}}) \leq n^{-c}$.

Theorem 3.2 (Lower bound). *Under the same setting of Theorem 3.1, if*

$$\frac{a+b}{2} - \sqrt{\frac{ab}{M}} < 1 \quad (9)$$

then the MLE (1) fails to recover the communities w.p. at least $\frac{3}{5}$, i.e. $\mathcal{P}_{e,c}(\psi_{\text{MLE}}) \geq \frac{3}{5}$.

Proof strategy The proofs of Theorems 3.1 and 3.2 are deferred to Section 5.2 and Section 5.3, respectively. Notably, our proof technique for the upper bound significantly differs from those traditional approaches employed in community detection or group synchronization (e.g. [1, Theorem 2] for SBM and [14, Theorem 1] for synchronization) in the way of handling different hypotheses. To be concrete, since the MLE fails when there exists a *wrong* hypothesis $\mathbf{x} = \{\boldsymbol{\kappa}, \mathbf{g}\}$ with a higher likelihood than the ground truth $\mathbf{x}^* = \{\boldsymbol{\kappa}^*, \mathbf{g}^*\}$, i.e.

$$\mathcal{P}(\mathbf{y}|\mathbf{x}) > \mathcal{P}(\mathbf{y}|\mathbf{x}^*) \quad \text{and} \quad \text{dist}_c(\boldsymbol{\kappa}, \boldsymbol{\kappa}^*) > 0,$$

a common path towards an upper bound of $\mathcal{P}_{e,c}(\psi_{\text{MLE}})$ is by first applying the union bound over all wrong hypothesis then conducting a sharp analysis on each one individually. That is

$$\mathcal{P}_{e,c}(\psi_{\text{MLE}}) = \mathcal{P}\left(\exists \mathbf{x} : \frac{\mathcal{P}(\mathbf{y}|\mathbf{x})}{\mathcal{P}(\mathbf{y}|\mathbf{x}^*)} \geq 1, \text{dist}_c(\boldsymbol{\kappa}, \boldsymbol{\kappa}^*) > 0\right) \quad (10)$$

$$\leq \sum_{\mathbf{x}:\text{dist}_c(\boldsymbol{\kappa}, \boldsymbol{\kappa}^*) > 0} \mathcal{P}\left(\frac{\mathcal{P}(\mathbf{y}|\mathbf{x})}{\mathcal{P}(\mathbf{y}|\mathbf{x}^*)} \geq 1\right). \quad (11)$$




Figure 2: An illustration of the information-theoretic limits for exact recovery of the communities. We plot the phase transition threshold (12) obtained from Theorems 3.1 and 3.2 with different choices of M in red dash curves. The gray area represents the region where the MLE (1) fails to recover the communities with a certain probability bounded away from zero, indicating the exact recovery is impossible, and the remaining area represents the region where (1) achieves exact cluster recovery with high probability.

This proof strategy offers a simple but sharp analysis in the context of community detection or synchronization e.g. [1, 14], where cluster memberships or group elements are considered separately. However, in our situation when these two components are coupled, (11) fails to provide a tight result due to the exponentially large space of the wrong hypothesis. To illustrate this, consider that for any hypothesis of the cluster assignment κ , there are M^n combinations of the associated group elements \mathbf{g} . Therefore, (11) explodes exponentially as $n \rightarrow \infty$. To address this issue, we instead analyze the original probability (10) that involves different \mathbf{g} as a whole, leading to a substantially more complicated proof scheme.

In addition, our proof relies on several sharp analyses related to random graph theory (see e.g. Theorem A.4 in Appendix A), which serves as one of the main technical challenges. We leave some discussion in Section 4.

Phase transition Theorems 3.1 and 3.2 reveal the presence of a phase transition phenomenon for the exact recovery of communities. That is, as the network size n grows, both the upper bound (8) and the lower bound (9) converge to the same threshold given by

$$\frac{a+b}{2} - \sqrt{\frac{ab}{M}} = 1 \quad (12)$$

where exact recovery is possible above and impossible below, indicating the optimality of our analysis.

To provide further insights, we plot the threshold (12) in Figure 2 for different choice of M . As a special case when $M = 1$, our model simplifies to the SBM without synchronization needed, and the threshold becomes

$$(\sqrt{a} - \sqrt{b})^2 = 2 \quad (13)$$

which agrees with the existing information-theoretic limit on the SBM proved in [1, 52]. Moreover, when $M > 1$, the threshold (12) exhibits a distinct behavior compared to (13), where exact recovery is possible as long as either a or b , that indicates the density of in-cluster or across clusters connections, is sufficiently large. To understand this, recall the MLE (1) explicitly checks the consistency $g_{ij} = g_i g_j^{-1}$ on each in-cluster edge and then rule out unsatisfied hypotheses. As a result, even edges that span different communities and possess “noisy” group transformations are helpful to recovery by serving as negative labels. On the contrary, such edges becomes irrelevant and provide no information in the absence of synchronization when $M = 1$.

In addition, one can observe that as M increases, the gray area in Figure 2 that represents the impossible region for recovery shrinks rapidly. This implies that the consistency check $g_{ij} = g_i g_j^{-1}$ is progressively effective for recovering the communities as M grows. To interpret this fact, recall the noisy group transformation across communities is uniformly drawn from \mathcal{G}_M . As a result, the probability for a noisy observation to satisfy the consistency is exactly M^{-1} , indicating the increasing difficulty for a wrong hypothesis to pass the check as M is large. Moreover, the threshold (12) would eventually converge to

$$\frac{a+b}{2} = 1 \quad (14)$$

when M is large¹. Then, as a basic result derived from [20], (14) corresponds to the threshold for a random graph being connected with high probability, where the graph is generated from the SBM with parameters $p = \frac{a \log n}{n}$ and $q = \frac{b \log n}{n}$. As a result, we conjecture that in the extreme case when $M \rightarrow \infty$, Theorem 3.1 still holds and thus the exact recovery of clusters becomes possible as long as the whole network is connected.

3.2 Exact recovery of group elements

Theorem 3.3 (Upper bound). *Under the same setting of Theorem 3.1. For any $c > 0$, if*

$$\frac{a+b}{2} - (1+o(1))\sqrt{\frac{ab}{M}} > 1+c \quad \text{and} \quad \frac{a}{2} > 1+c \quad (15)$$

then the MLE (1) exactly recovers the group elements w.p. at least $1 - n^{-c+o(1)}$, i.e. $\mathcal{P}_{e,g}(\psi_{\text{MLE}}) \leq n^{-c+o(1)}$.

¹Notice that in our analysis, we only consider the finite groups and thus M must be a constant as opposed to n that grows to infinity. Therefore, it is somewhat not rigorous to let $M \rightarrow \infty$ and achieve the threshold (14). Here, we present this merely for the purpose of a better interpretation of our theory.

Theorem 3.4 (Lower bound). *Under the same setting of Theorem 3.1. If*

$$\frac{a+b}{2} - \sqrt{\frac{ab}{M}} < 1 \quad \text{or} \quad \frac{a}{2} < 1 \quad (16)$$

then the MLE (1) fails to recover the group elements w.p. at least 0.45, i.e. $\mathcal{P}_{e,g}(\psi_{\text{MLE}}) \geq 0.45$.

The proofs of Theorems 3.3 and 3.4 are deferred to Section 5.4 and Section 5.5 respectively, which are simply extended from the ones of Theorems 3.1 and 3.2. Similar to the threshold (12), Theorems 3.3 and 3.4 suggest a phase transition exists for the exact recovery of group elements, given as

$$\frac{a+b}{2} - \sqrt{\frac{ab}{M}} = 1 \quad \text{and} \quad a = 2. \quad (17)$$

Here, one can see that (17) is an extension of (12) with the additional condition $a = 2$, implying that the exact recovery of communities is a *prerequisite* for recovering the group elements. In other words, one cannot hope to recover the group elements before recovering the communities at first. As a result, (17) can be also viewed as the threshold for recovering both communities and group elements.

Besides, the condition $a = 2$ can be interpreted from a random graph perspective: recall the nodes within each community of size $\frac{n}{2}$ are connected with the same probability $p = \frac{a \log n}{n}$, essentially following the Erdős–Rényi model. Then, it is well known that $a = 2$ represents the sharp threshold for the community being connected. Therefore, this implies that the exact recovery of group elements is possible only if each community is connected, which makes sense as one cannot hope the synchronize successfully over disconnected components due to the lack of information.

In summary, the recovery of group elements is possible only if (1) the two communities are recovered correctly and (2) each community is connected.

4 Discussions

Algorithm design From an algorithm perspective, since directly solving the MLE (1) is non-convex and computationally intractable, finding an efficient algorithm with a competitive performance becomes necessary in practice. In the context of community detection, efficient approaches based on semidefinite programming (SDP) [1, 37, 38, 4, 57, 47] and spectral method [2, 67, 46] have been proposed. In particular, they are known to achieve the information-theoretic limit (13) for exact recovery on the SBM [1, 2, 37, 57]. In light of this, similar algorithms [24, 23, 12, 66] have been applied to the joint problem, and it is natural to expect that these methods also work well all the way down to the information limit derived in this paper.

Unfortunately, our result suggests that none of these methods are likely to reach the information limit, indicating a *performance gap* exists compared to the MLE (1). For instance, the SDP developed in [24] for solving the joint problem exhibits a sharp threshold for exact recovering both communities and group elements, which is given as

$$a - \sqrt{2b} \log \left(\frac{ea}{\sqrt{2b}} \right) = 2. \quad (18)$$

as shown in Figure 3. Notably, the statistical model studied in [24] is slightly different to ours, where [24] focuses on $\mathcal{G}_M = \text{SO}(2)$ that is infinite, as apposed to the finite group considered in this work. Therefore, approximately, one may compare the threshold (18) by the SDP to the information limit (17) with a large M . As a result, the performance gap is clearly demonstrated.




Figure 3: The sharp threshold in blue dash curve for exactly recovering both communities and group elements, by the SDP proposed in [24]. The gray area represents the region where the SDP fails in the exact recovery with high probability.

To further understand the performance gap, our conjecture is that all existing approaches essentially differ from the MLE (1) and instead aim to solve an alternative optimization problem that allows (convex) relaxations to be smoothly applied. As an example, the authors in [24, 23] consider the orthogonal group $O(d)$ and design the following program for recovery:

$$\max_{\{\kappa_i\}_{i=1}^n, \{\mathbf{O}_i\}_{i=1}^n} \sum_{(i,j) \in \mathcal{E}, \kappa_i = \kappa_j} \langle \mathbf{O}_{ij}, \mathbf{O}_i \mathbf{O}_j^\top \rangle. \quad (19)$$

Here, \mathbf{O}_i and $\mathbf{O}_{ij} \in \mathbb{R}^{d \times d}$ are orthogonal matrices that represent g_i and g_{ij} , respectively. As a result, (19) attempts to recover by maximizing the total consistency between \mathbf{O}_{ij} and $\mathbf{O}_i \mathbf{O}_j^\top$ for all in-cluster edges. Obviously, a major difference between (19) and the corresponding MLE (4) is that the strict constraint $g_{ij} = g_i g_j^{-1}$ or equivalently $\mathbf{O}_{ij} = \mathbf{O}_i \mathbf{O}_j^\top$ given in (4) has been weakened² to maximizing the inner product $\langle \mathbf{O}_{ij}, \mathbf{O}_i \mathbf{O}_j^\top \rangle$, which instead tolerates a certain amount of error between \mathbf{O}_{ij} and $\mathbf{O}_i \mathbf{O}_j^\top$. As a result, the relaxed program (19) may not fully leverage the consistency requirement, resulting in the performance gap.

Giant component of Erdős–Rényi graphs An important technical ingredient of our theory is analyzing the giant component of Erdős–Rényi graphs [20]. As aforementioned, it is well-known that an Erdős–Rényi graph $G(n, p)$ of size n and connection probability $p = \frac{a \log n}{n}$ for some constant a is connected with high probability if and only if $a > 1$. Furthermore, when $a < 1$ such as the graph is disconnected, a giant component of size close to n is expected to exist with high probability. To be specific, let Z_n denote the size of the largest connected component in the graph, a common result [31, Theorem 2.14] is

$$\mathcal{P} \left(Z_n > \left(1 - n^{-a+\epsilon+o(1)} \right) n \right) > 1 - n^{-\epsilon} \quad (20)$$

²Equivalently, the constraint $g_{ij} = g_i g_j^{-1}$ can be transferred to the objective function in (4) and becomes a binary-valued penalty $f(g_i, g_j)$ such that $f(g_i, g_j) = \begin{cases} -\infty & g_{ij} \neq g_i g_j^{-1} \\ 0, & \text{otherwise} \end{cases}$.

which holds for any $\epsilon < a$. However, (20) is too loose to be applied for our analysis in Section 5.2. Therefore, we improve (20) with the following sharp result obtained

$$\mathcal{P}\left(Z_n \geq n - \Theta\left(\frac{n}{\log \log n}\right)\right) \geq 1 - n^{-\Theta(\log n)}, \quad (21)$$

which is necessary for acquiring the sharp threshold (12), see Theorem A.4 in Appendix A for the details. Remarkably, (21) cannot be derived by the conventional moment methods such as Markov's or Chebyshev's inequality, which are generally applied in random graph theory (e.g. see [7]). Instead, we prove (21) by conducting a delicate combinatorial analysis, which serves as one of the main technical challenges in our proof.

Future works Our analysis presented in this work can be generalized in several aspects, listed as follows:

1. Our theory given in Section 3 is based on the assumption that \mathcal{G}_M is a finite group and M stays as a constant as n grows, extending the result to infinite groups e.g. the orthogonal group $O(d)$ is left as a future work.
2. Although we restrict our attention on the simplest scenario of two equal-sized communities, our proof framework can be naturally extended to more generalized cases of multiple clusters with different sizes
3. Depending on the applications, one may consider a more involved statistical model such as the one studied in [23] where the group transformation g_{ij} observed on an edge (i, j) within the same community is perturbed by noise rather than being exact i.e., $g_{ij} = g_i^* \left(g_j^*\right)^{-1}$.
4. Based on our discussion in Section 4, finding an efficient algorithm that achieves the information limit is still an open problem. Here, our result suggests that the key might be focusing on the recovery of communities which is the prerequisite for recovering group elements.

5 Proof of the theorems

In this section, we describe the proofs of our theorems presented in Section 3. For ease of presentation, some technical details are deferred to the Appendices.

5.1 Preliminaries

We first introduce several important ingredients that support our proof.

Four subsets. Under the setting of two equal-sized communities, given the ground truth of the community assignment κ^* and any alternative hypothesis κ , we define the following four subsets of nodes, which will be frequently used in our proof:

$$\begin{aligned} S_{11} &:= S_1 \cap S_1^* = \{i \mid \kappa_i = 1, \kappa_i^* = 1\}, \\ S_{12} &:= S_1 \cap S_2^* = \{i \mid \kappa_i = 1, \kappa_i^* = 2\}, \\ S_{21} &:= S_2 \cap S_1^* = \{i \mid \kappa_i = 2, \kappa_i^* = 1\}, \\ S_{22} &:= S_2 \cap S_2^* = \{i \mid \kappa_i = 2, \kappa_i^* = 2\}. \end{aligned} \quad (22)$$




Figure 4: An illustration of the four subsets S_{11} , S_{12} , S_{21} , and S_{22} defined in (22). We use green and orange dash lines to indicate the set of edges corresponding to r and r^* defined in (30), respectively.

For instance, S_{12} stands for the set of nodes that is assigned to S_1 in κ but actually belongs to S_2^* in κ^* . See Figure 4 for an illustration. Notably, all the four subsets are mutually disjoint and their union covers all nodes in the network. Also, one can immediately observe the following equations:

$$|S_{11}| = |S_{22}|, \quad \text{and} \quad |S_{21}| = |S_{12}| = \frac{n}{2} - |S_{11}|, \quad (23)$$

which hold for any valid hypothesis κ .

Derivation of the MLE (1) We start by expanding the likelihood $\mathcal{P}(\mathcal{E}, \mathfrak{G}|\kappa, \mathbf{g})$ as

$$\begin{aligned} \mathcal{P}(\mathcal{E}, \mathfrak{G}|\kappa, \mathbf{g}) &= \mathcal{P}(\mathcal{E}|\kappa, \mathbf{g})\mathcal{P}(\mathfrak{G}|\kappa, \mathbf{g}, \mathcal{E}) \\ &= \mathcal{P}(\mathcal{E}|\kappa)\mathcal{P}(\mathfrak{G}|\kappa, \mathbf{g}, \mathcal{E}) \end{aligned} \quad (24)$$

which is separated into the likelihood of edge connections $\mathcal{P}(\mathcal{E}|\kappa)$ and the one of group transformations given the edges, i.e. $\mathcal{P}(\mathfrak{G}|\kappa, \mathbf{g}, \mathcal{E})$. According to the statistical model, $\mathcal{P}(\mathcal{E}|\kappa)$ is given as

$$\begin{aligned} \mathcal{P}(\mathcal{E}|\kappa) &= \prod_{\substack{(i,j) \in \mathcal{E} \\ \kappa_i = \kappa_j}} p \prod_{\substack{(i,j) \notin \mathcal{E} \\ \kappa_i = \kappa_j}} (1-p) \prod_{\substack{(i,j) \in \mathcal{E} \\ \kappa_i \neq \kappa_j}} q \prod_{\substack{(i,j) \notin \mathcal{E} \\ \kappa_i \neq \kappa_j}} (1-q) \\ &\propto \left(\frac{p}{1-p}\right)^{|\mathcal{E}_{\text{inner}}|} \left(\frac{q}{1-q}\right)^{|\mathcal{E}_{\text{inter}}|} \\ &= \left(\frac{p}{1-p}\right)^{|\mathcal{E}_{\text{inner}}|} \left(\frac{q}{1-q}\right)^{|\mathcal{E}| - |\mathcal{E}_{\text{inner}}|} \\ &\propto \left(\frac{p(1-q)}{q(1-p)}\right)^{|\mathcal{E}_{\text{inner}}|} \end{aligned} \quad (25)$$

where $\mathcal{E}_{\text{inter}}$ denotes the set of edges across different communities, as opposed to $\mathcal{E}_{\text{inner}}$ defined in (2), and we use the fact that $|\mathcal{E}| = |\mathcal{E}_{\text{inner}}| + |\mathcal{E}_{\text{inter}}|$. Also, for $\mathcal{P}(\mathfrak{G}|\mathbf{g}, \boldsymbol{\kappa}, \mathcal{E})$, we have

$$\begin{aligned} \mathcal{P}(\mathfrak{G}|\mathbf{g}, \boldsymbol{\kappa}, \mathcal{E}) &= \prod_{\substack{(i,j) \in \mathcal{E} \\ \kappa_i = \kappa_j}} \mathbb{1}\left(g_{ij} = g_i g_j^{-1}\right) \prod_{\substack{(i,j) \in \mathcal{E} \\ \kappa_i \neq \kappa_j}} M^{-1} \\ &\propto \mathbb{1}\left(g_{ij} = g_i g_j^{-1}; \forall (i,j) \in \mathcal{E}_{\text{inner}}\right) \cdot M^{-|\mathcal{E}_{\text{inter}}|} \\ &\propto \mathbb{1}\left(g_{ij} = g_i g_j^{-1}; \forall (i,j) \in \mathcal{E}_{\text{inner}}\right) \cdot M^{|\mathcal{E}_{\text{inner}}|} \end{aligned} \quad (26)$$

where $\mathbb{1}(\cdot)$ denotes the indicator function.

Given the above, by plugging (25) and (26) into (24), the MLE can be written as

$$\begin{aligned} \psi_{\text{MLE}}(\mathcal{E}, \mathfrak{G}) &= \arg \max_{\boldsymbol{\kappa}, \mathbf{g}} \mathcal{P}(\mathcal{E}, \mathfrak{G}|\boldsymbol{\kappa}, \mathbf{g}) \\ &= \arg \max_{\boldsymbol{\kappa}, \mathbf{g}} \left(\frac{Mp(1-q)}{q(1-p)} \right)^{|\mathcal{E}_{\text{inner}}|} \cdot \mathbb{1}\left(g_{ij} = g_i g_j^{-1}; \forall (i,j) \in \mathcal{E}_{\text{inner}}\right) \\ \text{s.t. } & n_k = n_k^*, \quad k = 1, \dots, K \end{aligned} \quad (27)$$

where we apply the assumption that all cluster sizes are given.

At first glance, solving (27) requires the prior knowledge of the model parameters p, q , and M . However, by assuming $Mp(1-q) > q(1-p)$ or $Mp(1-q) \leq q(1-p)$, and making the indicator function as constraints, (27) becomes (1) which does not involve any model parameter specifically.

Likelihood ratio Given the true parameter $\mathbf{x}^* = \{\boldsymbol{\kappa}^*, \mathbf{g}^*\}$ and a hypothesis $\mathbf{x} = \{\boldsymbol{\kappa}, \mathbf{g}\}$, our proof is based on the likelihood ratio defined as

$$\frac{\mathcal{P}(\mathbf{y}|\mathbf{x})}{\mathcal{P}(\mathbf{y}|\mathbf{x}^*)} = \frac{\mathcal{P}(\mathcal{E}|\boldsymbol{\kappa})}{\mathcal{P}(\mathcal{E}|\boldsymbol{\kappa}^*)} \cdot \frac{\mathcal{P}(\mathfrak{G}|\boldsymbol{\kappa}, \mathbf{g}, \mathcal{E})}{\mathcal{P}(\mathfrak{G}|\boldsymbol{\kappa}^*, \mathbf{g}^*, \mathcal{E})}.$$

To this end, by using the result in (25), we get

$$\frac{\mathcal{P}(\mathcal{E}|\boldsymbol{\kappa})}{\mathcal{P}(\mathcal{E}|\boldsymbol{\kappa}^*)} = \left(\frac{Mp(1-q)}{q(1-p)} \right)^{|\mathcal{E}_{\text{inner}}| - |\mathcal{E}_{\text{inner}}^*|} \quad (28)$$

where $\mathcal{E}_{\text{inner}}^*$ denotes the set of in-cluster edges according to $\boldsymbol{\kappa}^*$. Furthermore, recall the four subsets defined in (22), then notice that

$$\begin{aligned} |\mathcal{E}_{\text{inner}}| - |\mathcal{E}_{\text{inner}}^*| &= |\mathcal{E}(S_{11}, S_{12})| + |\mathcal{E}(S_{21}, S_{22})| - |\mathcal{E}(S_{11}, S_{21})| - |\mathcal{E}(S_{12}, S_{22})| \\ &= r - r^* \end{aligned} \quad (29)$$

where r and r^* are defined as

$$r := |\mathcal{E}(S_{11}, S_{12})| + |\mathcal{E}(S_{21}, S_{22})|, \quad r^* := |\mathcal{E}(S_{11}, S_{21})| + |\mathcal{E}(S_{12}, S_{22})|. \quad (30)$$

One can interpret r (resp. r^*) as the number of connections that are within the same community in $\boldsymbol{\kappa}$ (resp. $\boldsymbol{\kappa}^*$) but across different ones in $\boldsymbol{\kappa}^*$ (resp. $\boldsymbol{\kappa}$). See Figure 4 for an illustration. Also, by using (26), we obtain

$$\frac{\mathcal{P}(\mathfrak{G}|\boldsymbol{\kappa}, \mathbf{g}, \mathcal{E})}{\mathcal{P}(\mathfrak{G}|\boldsymbol{\kappa}^*, \mathbf{g}^*, \mathcal{E})} = \frac{\mathbb{1}\left(g_{ij} = g_i g_j^{-1}; \forall (i,j) \in \mathcal{E}_{\text{inner}}\right)}{\mathbb{1}\left(g_{ij} = g_i^* (g_j^*)^{-1}; \forall (i,j) \in \mathcal{E}_{\text{inner}}^*\right)}.$$

Combining the results above yields the likelihood ratio

$$\frac{\mathcal{P}(\mathbf{y}|\mathbf{x})}{\mathcal{P}(\mathbf{y}|\mathbf{x}^*)} = \left(\frac{Mp(1-q)}{q(1-p)} \right)^{r-r^*} \cdot \frac{\mathbb{1}(g_{ij} = g_i g_j^{-1}; \forall (i, j) \in \mathcal{E}_{\text{inner}})}{\mathbb{1}(g_{ij} = g_i^* (g_j^*)^{-1}; \forall (i, j) \in \mathcal{E}_{\text{inner}}^*)} = r - r^* \quad (31)$$

which will be served as the starting point of our analysis.

5.2 Proof of Theorem 3.1

For ease of presentation, without loss of generality, we assume all the ground truth group elements $\mathbf{g}^* = \{g_i^*\}_{i=1}^n$ satisfies

$$g_i^* = g_0, \quad \forall i \in \{1, \dots, n\} \quad (32)$$

where g_0 stands for the neutral (zero) element of the group \mathcal{G}_M . Also, we denote

$$\mathcal{P}_{\mathbf{x}^*}(\cdot) := \mathcal{P}(\cdot|\mathbf{x}^*), \quad \mathcal{P}_{\mathbf{x}}(\cdot) := \mathcal{P}(\cdot|\mathbf{x})$$

as the probability measure conditional on the ground truth \mathbf{x}^* and the hypothesis \mathbf{x} , respectively.

Step 1: A union bound To begin with, given an observation \mathbf{y} , the MLE (1) fails to exactly recover $\boldsymbol{\kappa}^*$ if and only if there exists an alternative hypothesis $\mathbf{x} = \{\boldsymbol{\kappa}, \mathbf{g}\}$ such that

$$\text{dist}_c(\boldsymbol{\kappa}, \boldsymbol{\kappa}^*) > 0 \quad \text{and} \quad \mathcal{P}_{\mathbf{x}}(\mathbf{y}) \geq \mathcal{P}_{\mathbf{x}^*}(\mathbf{y}).$$

In other words, \mathbf{x} is a wrong hypothesis with a likelihood higher than or equal to the ground truth \mathbf{x}^* . As a result, the error probability defined in (7) can be written as

$$\mathcal{P}_{e,c}(\psi_{\text{MLE}}) = \mathcal{P}_{\mathbf{x}^*}(\exists \mathbf{x} = \{\boldsymbol{\kappa}, \mathbf{g}\}: \text{dist}_c(\boldsymbol{\kappa}, \boldsymbol{\kappa}^*) > 0, \mathcal{P}_{\mathbf{x}}(\mathbf{y}) \geq \mathcal{P}_{\mathbf{x}^*}(\mathbf{y})). \quad (33)$$

To proceed, recall the definition of the two subsets sets S_{12} and S_{21} in Section 5.1 which includes the nodes with different labels between $\boldsymbol{\kappa}$ and $\boldsymbol{\kappa}^*$. Let us denote $\Delta := |S_{12}| = |S_{21}|$ as their sizes, then for any wrong hypothesis $\boldsymbol{\kappa}$ with $\text{dist}_c(\boldsymbol{\kappa}, \boldsymbol{\kappa}^*) > 0$, it satisfies $1 \leq \Delta \leq \frac{n}{2} - 1$ (when $\Delta = \frac{n}{2}$, exact recovery can be achieved by swapping the labels of S_1 and S_2). In this way, for a fixed Δ , we define

$$\mathcal{H}_{\boldsymbol{\kappa}, \Delta} := \{\boldsymbol{\kappa}: |S_{12}| = |S_{21}| = \Delta\},$$

as the set of all such hypotheses, and there are $\binom{n/2}{\Delta}^2$ number of unique $\boldsymbol{\kappa}$, i.e., $|\mathcal{H}_{\boldsymbol{\kappa}, \Delta}| = \binom{n/2}{\Delta}^2$. Furthermore, let $\boldsymbol{\kappa}_{\Delta} \in \mathcal{H}_{\boldsymbol{\kappa}, \Delta}$ be any hypothesis in $\mathcal{H}_{\boldsymbol{\kappa}, \Delta}$, we denote

$$\mathcal{H}_{\mathbf{x}, \Delta} := \{\mathbf{x} = \{\boldsymbol{\kappa}, \mathbf{g}\}: \boldsymbol{\kappa} = \boldsymbol{\kappa}_{\Delta}\},$$

as the set of hypotheses \mathbf{x} with a fixed $\boldsymbol{\kappa}$ and different group elements \mathbf{g} . Given the above, the error probability in (33) can be bounded as

$$\begin{aligned}
\mathcal{P}_{e,c}(\psi_{\text{MLE}}) &\stackrel{(a)}{\leq} \sum_{\Delta=1}^{n/2-1} \mathcal{P}_{\mathbf{x}^*}(\exists \mathbf{x} = \{\boldsymbol{\kappa}, \mathbf{g}\} : \boldsymbol{\kappa} \in \mathcal{H}_{\boldsymbol{\kappa},\Delta}, \mathcal{P}_{\mathbf{x}}(\mathbf{y}) \geq \mathcal{P}_{\mathbf{x}^*}(\mathbf{y})) \\
&\stackrel{(b)}{\leq} \sum_{\Delta=1}^{n/2-1} \sum_{\boldsymbol{\kappa} \in \mathcal{H}_{\boldsymbol{\kappa},\Delta}} \mathcal{P}_{\mathbf{x}^*}(\exists \mathbf{x} = \{\boldsymbol{\kappa}, \mathbf{g}\} : \mathcal{P}_{\mathbf{x}}(\mathbf{y}) \geq \mathcal{P}_{\mathbf{x}^*}(\mathbf{y})) \\
&\stackrel{(c)}{=} \sum_{\Delta=1}^{n/2-1} \underbrace{\binom{n/2}{\Delta}^2}_{=: \mathcal{P}_{\Delta}} \mathcal{P}_{\mathbf{x}^*}(\exists \mathbf{x} \in \mathcal{H}_{\boldsymbol{\kappa},\Delta} : \mathcal{P}_{\mathbf{x}}(\mathbf{y}) \geq \mathcal{P}_{\mathbf{x}^*}(\mathbf{y})) \\
&\stackrel{(d)}{\leq} 2 \sum_{\Delta=1}^{\lceil n/4 \rceil} \binom{n/2}{\Delta}^2 \mathcal{P}_{\Delta}. \tag{34}
\end{aligned}$$

Here, (a) holds by applying the union bound over different Δ ; (b) comes by further applying this over the set $\mathcal{H}_{\boldsymbol{\kappa},\Delta}$; (c) stems from the fact that any hypothesis $\boldsymbol{\kappa} \in \mathcal{H}_{\boldsymbol{\kappa},\Delta}$ should contribute the same amount of error probability under a fixed Δ ; (d) uses the symmetry between Δ and $\frac{n}{2} - \Delta$ by swapping the labels in $\boldsymbol{\kappa}$.

Step 2: Bound on each \mathcal{P}_{Δ} Our next job is to tightly bound \mathcal{P}_{Δ} for each $1 \leq \Delta \leq \lceil \frac{n}{4} \rceil$. To this end, by using the likelihood ratio obtained in (31) and plugging in the scaling $p = \frac{a \log n}{n}, q = \frac{b \log n}{n}$, we have

$$\begin{aligned}
\mathcal{P}_{\Delta} &= \mathcal{P}_{\mathbf{x}^*} \left(\exists \mathbf{x} \in \mathcal{H}_{\boldsymbol{\kappa},\Delta} : \frac{\mathcal{P}_{\mathbf{x}}(\mathbf{y})}{\mathcal{P}_{\mathbf{x}^*}(\mathbf{y})} \geq 1 \right) \\
&\leq \mathcal{P}_{\mathbf{x}^*} \left(\exists \mathbf{x} \in \mathcal{H}_{\boldsymbol{\kappa},\Delta} : \left(\frac{Ma}{b} \right)^{r-r^*} \cdot \frac{\mathbb{1}(g_{ij} = g_i g_j^{-1}; \forall (i,j) \in \mathcal{E}_{\text{inner}})}{\mathbb{1}(g_{ij} = g_i^* (g_j^*)^{-1}; \forall (i,j) \in \mathcal{E}_{\text{inner}}^*)} > 1 \right) \\
&= \mathcal{P}_{\mathbf{x}^*} \left(\exists \mathbf{x} \in \mathcal{H}_{\boldsymbol{\kappa},\Delta} : \left(\frac{Ma}{b} \right)^{r-r^*} \cdot \mathbb{1}(g_{ij} = g_i g_j^{-1}; \forall (i,j) \in \mathcal{E}_{\text{inner}}) > 1 \right) \tag{35}
\end{aligned}$$

where the last step uses the fact that

$$\mathcal{P}_{\mathbf{x}^*}(g_{ij} = g_i^* (g_j^*)^{-1}; \forall (i,j) \in \mathcal{E}_{\text{inner}}^*) = 1.$$

To move forward, for any wrong hypothesis \mathbf{x} , let us define the following two events in the probability space:

$$\begin{aligned}
E_1 &: \left(\frac{Ma}{b} \right)^{r-r^*} > 1, \\
E_{\mathbf{x},2} &: g_{ij} = g_i g_j^{-1}, \forall (i,j) \in \mathcal{E}_{\text{inner}}.
\end{aligned} \tag{36}$$

which are the necessary and sufficient conditions for \mathbf{x} with a likelihood higher than \mathbf{x}^* . Notably, for different hypothesis $\mathbf{x} \in \mathcal{H}_{\boldsymbol{\kappa},\Delta}$ with the fixed community assignment $\boldsymbol{\kappa} = \boldsymbol{\kappa}_{\Delta}$, the event E_1 does not depend on \mathbf{x} but is invariant in the sample space, therefore we omit the subscript \mathbf{x} . In this

way, (35) can be written as

$$\begin{aligned} \mathcal{P}_\Delta &\leq \mathcal{P}_{\mathbf{x}^*} \left(\bigcup_{\mathbf{x} \in \mathcal{H}_{\kappa, \Delta}} (E_1 \cap E_{\mathbf{x}, 2}) \right) = \mathcal{P}_{\mathbf{x}^*} \left(E_1 \cap \underbrace{\left(\bigcup_{\mathbf{x} \in \mathcal{H}_{\kappa, \Delta}} E_{\mathbf{x}, 2} \right)}_{=: E_2} \right) \\ &= \mathcal{P}_{\mathbf{x}^*} (E_1 \cap E_2) \end{aligned} \quad (37)$$

where we denote E_2 as the union over $\mathbf{x} \in \mathcal{H}_{\kappa, \Delta}$. To proceed, recall the definitions of r and r^* in (30), we further define $r = r_1 + r_2$ where

$$r_1 := |\mathcal{E}(S_{11}, S_{12})|, \quad r_2 := |\mathcal{E}(S_{21}, S_{22})|.$$

Under the ground truth \mathbf{x}^* , the three r_1 , r_2 , and r^* are independent binomial random variables since they involve mutually disjoint sets of edges. Therefore, by conditioning on r_1 , r_2 , and r^* , (37) satisfies

$$\begin{aligned} \mathcal{P}_\Delta &\leq \sum_{r_1} \sum_{r_2} \sum_{r^*} \mathcal{P}_{\mathbf{x}^*}(r_1) \mathcal{P}_{\mathbf{x}^*}(r_2) \mathcal{P}_{\mathbf{x}^*}(r^*) \mathcal{P}_{\mathbf{x}^*} (E_1 \cap E_2 \mid r_1, r_2, r^*) \\ &= \sum_{r_1} \sum_{r_2} \sum_{r^*} \mathcal{P}_{\mathbf{x}^*}(r_1) \mathcal{P}_{\mathbf{x}^*}(r_2) \mathcal{P}_{\mathbf{x}^*}(r^*) \mathcal{P}_{\mathbf{x}^*} \left(\left(\left(\frac{Ma}{b} \right)^{r_1+r_2-r^*} > 1 \right) \cap E_2 \mid r_1, r_2, r^* \right) \\ &= \sum_{r_1} \sum_{r_2} \sum_{r^*} \mathcal{P}_{\mathbf{x}^*}(r_1) \mathcal{P}_{\mathbf{x}^*}(r_2) \mathcal{P}_{\mathbf{x}^*}(r^*) \cdot \mathbb{1} \left(\left(\frac{Ma}{b} \right)^{r_1+r_2-r^*} > 1 \right) \cdot \mathcal{P}_{\mathbf{x}^*} (E_2 \mid r_1, r_2, r^*) \end{aligned} \quad (38)$$

where the last step comes from the fact that given r_1 , r_2 , and r^* , the probability $\mathcal{P}_{\mathbf{x}^*}(E_1 \mid r_1, r_2, r^*)$ is either 0 or 1. Therefore we can represent it as an indicator function. Furthermore, by introducing an extra factor $\alpha > 0$, one can see that it satisfies

$$\begin{aligned} \mathbb{1} \left(\left(\frac{Ma}{b} \right)^{r_1+r_2-r^*} > 1 \right) &= \mathbb{1} \left(\left(\frac{Ma}{b} \right)^{\alpha(r_1+r_2-r^*)} > 1 \right) \\ &\leq \left(\frac{Ma}{b} \right)^{\alpha(r_1+r_2-r^*)}. \end{aligned}$$

Plugging this into (38) yields

$$\mathcal{P}_\Delta \leq \min_{\alpha > 0} \left(\sum_{r_1} \sum_{r_2} \sum_{r^*} \mathcal{P}_{\mathbf{x}^*}(r_1) \mathcal{P}_{\mathbf{x}^*}(r_2) \mathcal{P}_{\mathbf{x}^*}(r^*) \cdot \left(\frac{Ma}{b} \right)^{\alpha(r_1+r_2-r^*)} \cdot \mathcal{P}_{\mathbf{x}^*} (E_2 \mid r_1, r_2, r^*) \right). \quad (39)$$

Notably, α is significant for obtaining the sharp result since we can minimize over all $\alpha > 0$.

Step 3: Cycle analysis Now, we need to upper bound $\mathcal{P}_{\mathbf{x}^*} (E_2 \mid r_1, r_2, r^*)$. Importantly, a naive way by applying the union bound over all possible hypotheses in $\mathcal{H}_{\mathbf{x}, \Delta}$ such that

$$\mathcal{P}_{\mathbf{x}^*} (E_2 \mid r_1, r_2, r^*) \leq \sum_{\mathbf{x} \in \mathcal{H}_{\kappa, \Delta}} \mathcal{P}_{\mathbf{x}^*} (E_{\mathbf{x}, 2} \mid r_1, r_2, r^*)$$

fails to give a sharp result, since $|\mathcal{H}_{\kappa,\Delta}| = M^n$ where each node has M choices of its group elements, then the summation over $\mathcal{H}_{\kappa,\Delta}$ explodes as n is large. Instead, we have to bound $\mathcal{P}_{\mathbf{x}^*}(E_2 | r_1, r_2, r^*)$ directly as a whole. To this end, let us restate the event E_2 defined in (37) as

$$E_2 : \quad \exists \mathbf{x} \in \mathcal{H}_{\kappa,\Delta} : g_{ij} = g_i g_j^{-1}, \quad \forall (i, j) \in \mathcal{E}_{\text{inner}}. \quad (40)$$

Here, recall $\mathcal{E}_{\text{inner}}$ denotes the set of edges within the two communities S_1 and S_2 , i.e. $\mathcal{E}_{\text{inner}} = \mathcal{E}(S_1) \cup \mathcal{E}(S_2)$, then we can consider S_1 and S_2 separately by further defining the following two events:

$$\begin{aligned} E_{2,S_1} : \quad & \exists \mathbf{x} \in \mathcal{H}_{\kappa,\Delta} : g_{ij} = g_i g_j^{-1}, \quad \forall (i, j) \in \mathcal{E}(S_1), \\ E_{2,S_2} : \quad & \exists \mathbf{x} \in \mathcal{H}_{\kappa,\Delta} : g_{ij} = g_i g_j^{-1}, \quad \forall (i, j) \in \mathcal{E}(S_2), \end{aligned} \quad (41)$$

and one can see that $E_2 = E_{2,S_1} \cap E_{2,S_2}$. In this way, we have

$$\begin{aligned} \mathcal{P}_{\mathbf{x}^*}(E_2 | r_1, r_2, r^*) &\stackrel{(a)}{=} \mathcal{P}_{\mathbf{x}^*}(E_2 | r_1, r_2) = \mathcal{P}_{\mathbf{x}^*}(E_{2,S_1} \cap E_{2,S_2} | r_1, r_2) \\ &\stackrel{(b)}{=} \mathcal{P}_{\mathbf{x}^*}(E_{2,S_1} | r_1) \cdot \mathcal{P}_{\mathbf{x}^*}(E_{2,S_2} | r_2) \end{aligned} \quad (42)$$

where (a) holds since E_2 does not involve r^* ; (b) comes from the fact that E_{2,S_1} and E_{2,S_2} are independent given r_1 and r_2 . As a result, because of the symmetry, we can focus on bounding $\mathcal{P}_{\mathbf{x}^*}(E_{2,S_1} | r_1)$ and the similar result applies to $\mathcal{P}_{\mathbf{x}^*}(E_{2,S_2} | r_2)$.

To this end, by definition E_{2,S_1} occurs as long as there exists a hypothesis of the group elements \mathbf{g} that satisfies the consistency $g_{ij} = g_i g_j^{-1}$ on each edge in S_1 . However, determining the conditional probability $\mathcal{P}_{\mathbf{x}^*}(E_{2,S_1} | r_1)$ appears to be challenging as it involves two sources of randomness:

1. For any pair of node (i, j) within S_{11} or S_{12} , they are connected with probability $p = \frac{a \log n}{n}$, and the group transformation observed is $g_{ij} = g_i^* (g_j^*)^{-1} = 0$ with probability 1
2. For any pair of nodes (i, j) between S_{11} and S_{12} , the edges are randomly assigned under the constraint that the total number of edges is r_1 , and g_{ij} is uniformly drawn from \mathcal{G}_M

Here, our analysis on $\mathcal{P}_{\mathbf{x}^*}(E_{2,S_1} | r_1)$ starts from several simple examples shown in Figure 5, which is illustrated as follows:

- (a) We start from Figure 5 *a* with $r_1 = 0$ such that no edges connect between S_{11} and S_{12} . In this case, recall the assumption (32) that $g_i^* = 0$, $i = 1, \dots, n$, then g_{ij} on all edges within S_{11} and S_{12} satisfy

$$g_{ij} = g_i^* (g_j^*)^{-1} = 0.$$

As a result, no matter how the nodes within S_{11} or S_{12} are connected, we have

$$\mathcal{P}_{\mathbf{x}^*}(E_{2,S_1} | r_1 = 0) = 1$$

as a hypothesis $g_i = 0$, $i = 1, \dots, n$ always satisfies the consistency.

- (b) Now, suppose $r_1 = 1$ where there is an edge (l_1, l_2) connecting $l_1 \in S_{12}$ and $l_2 \in S_{11}$. In this case, the group transformation $g_{l_1 l_2}$ is uniformly drawn from \mathcal{G}_M . However, for any realization of $g_{l_1 l_2}$, there always exists a hypothesis \mathbf{g} satisfy the consistency by taking

$$g_i = 0, \quad \forall i \in S_{11} \quad \text{and} \quad g_i = g_{l_1 l_2}, \quad \forall i \in S_{12}$$




Figure 5: Simple examples on E_{2,S_1} . We consider five different scenarios: (a) and (b): no edge or only one edge connected between S_{11} and S_{12} , then E_{2,S_1} occurs with probability 1; (c): assuming the subgraph formed by S_{11} is connected, and two edges connected to the same node in S_{12} form a cycle, then E_{2,S_1} occurs when the cycle consistency satisfies with probability M^{-1} ; (d) and (e): more edges between S_{11} and S_{12} leads to more cycles and further lower probability of E_{2,S_1} . See text for details.

as shown in Figure 5 b. Therefore, we still have

$$\mathcal{P}_{\mathbf{x}^*}(E_{2,S_1} | r_1 = 1) = 1.$$

(c) Next, we consider when $r_1 = 2$ and further assume

1. The subgraph formed by S_{11} , denoted as $G(S_{11})$, is connected
2. The $r_1 = 2$ edges between S_{11} and S_{12} connect the same node $l_1 \in S_{12}$

Let us denote this event as E_c . As a result, let l_2 and $l_3 \in S_{11}$ denote the two nodes that connect with l_1 . Then, l_2 and l_3 are connected by some path in $G(S_{11})$ due to the assumption that $G(S_{11})$ is connected. This path, together with the two edges (l_1, l_2) and (l_1, l_3) , form a cycle across S_{11} and S_{12} :

$$l_1 \xrightarrow{g_{l_1 l_2}} l_2 \longrightarrow \cdots \longrightarrow l_3 \xrightarrow{g_{l_3 l_1}} l_1 \quad (43)$$

which is shown in Figure 5 c. In this case, one can notice that E_2 occurs only if the aggregation of group transformations on the cycle is 0, i.e.

$$g_{l_1 l_2} \cdots g_{l_3 l_1} = 0 \quad (44)$$

which is so-called the *cycle consistency*. However, since $g_{l_1 l_2}$ and $g_{l_3 l_1}$ are both randomly drawn from \mathcal{G}_M , no matter what the hypothesis \mathbf{g} is, the probability that (44) holds is only M^{-1} . Hence, we have

$$\mathcal{P}_{\mathbf{x}^*}(E_{2,S_1} | E_c, r_1 = 2) = M^{-1}$$

given that E_c occurs.

- (d) Now, we consider $r_1 = 3$ and assume an event E_d similar to E_c above occurs, i.e. $G(S_{11})$ is connected and all the three edges connect the same node $l_1 \in S_{22}$. We denote the new node connect to l_1 as $l_4 \in S_{11}$. As a result, this gives rise another cycle

$$l_1 \xrightarrow{g_{l_1 l_4}} l_4 \longrightarrow \cdots \longrightarrow l_3 \xrightarrow{g_{l_3 l_1}} l_1$$

as shown in Figure 5 *d*. Again, by applying the same argument as in part (c), one can see that the new cycle satisfies the cycle consistency with probability M^{-1} and is independent with the old one. Therefore, we have

$$\mathcal{P}_{\mathbf{x}^*}(E_{2,S_1} | E_d, r_1 = 3) = M^{-2}$$

given that E_d occurs.

- (e) Lastly, we consider a more involved example when two nodes in S_{12} connect with S_{11} , and the subgraph $G(S_{11})$ is still connected, as shown in Figure 5 *e*. We denote such an event as E_e . Here, each node has three edge connections and therefore bring us two cycles. In this case, one can see that with probability M^{-4} , all the cycles satisfy the consistency, and thus

$$\mathcal{P}_{\mathbf{x}^*}(E_{2,S_1} | E_e, r_1 = 6) \leq M^{-4}.$$

Notably, here we use ‘ \leq ’ instead of ‘ $=$ ’ since we have ignored the edge connections in S_{12} , and there might be more cycles exist by considering those edges. However, such an upper bound is sufficient to our analysis.

Now, our analysis in Figure 5 can be generalized to the following result

Lemma 5.1. *Let E_{con} denotes the event that $G(S_{11})$ is connected. Then,*

$$\mathcal{P}_{\mathbf{x}^*}(E_{2,S_1} | r_1, E_{\text{con}}) \leq \min \{1, M^{\Delta - r_1}\} \quad (45)$$

where $\Delta = |S_{12}|$.

Proof. For each node $i \in S_{12}$, let $r_{1,i}$ denotes the number of edges it connects to S_{11} . Importantly, since $G(S_{11})$ is assumed to be connected, there are always at least $\max\{0, r_{1,i} - 1\}$ number of cycles formed by i like (43). Therefore, let $E_{\text{cycle},i}$ denotes the event that all of them satisfy the cycle consistency such as (44). Then,

$$\begin{aligned} \mathcal{P}_{\mathbf{x}^*}(E_{2,S_1} | r_1, E_{\text{con}}) &\leq \mathcal{P} \left(\bigcap_{i \in S_{12}} E_{\text{cycle},i} \right) \stackrel{(a)}{=} \prod_{i \in S_{12}} \mathcal{P}(E_{\text{cycle},i}) \\ &= \prod_{i \in S_{12}} M^{-\max\{0, r_{1,i} - 1\}} = \prod_{i \in S_{12}} \min \{1, M^{1 - r_{1,i}}\} \\ &\leq \min \left\{ 1, \prod_{i \in S_{12}} M^{1 - r_{1,i}} \right\} = \min \left\{ 1, M^{\sum_{i \in S_{12}} (r_{1,i} - 1)} \right\} \\ &\stackrel{(b)}{=} \min \{1, M^{\Delta - r_1}\} \end{aligned}$$

where (a) uses the fact that the set of events $\{E_{\text{cycle},i}\}_{i \in S_{12}}$ are mutually independent as they involve disjoint sets of edges; (b) holds since $\sum_{i \in S_{12}} r_{1,i} = r_1$. \square

However, Lemma 5.1 is still one step away from the desired bound as it requires $G(S_{11})$ to be connected. Fortunately, as $G(S_{11})$ follows an Erdős–Rényi model and when $p = \Theta\left(\frac{\log n}{n}\right)$, we can show that at least a giant component of size

$$(1 - o(1)) \cdot |S_{11}|$$

exists in $G(S_{11})$ w.h.p. (see Theorem A.4). As a result, an upper bound similar to (45) can be obtained as follows:

Lemma 5.2. *Given $p = \Theta\left(\frac{\log n}{n}\right)$, it satisfies*

$$\mathcal{P}_{\mathbf{x}^*}(E_{2,S_1} \mid r_1) \leq n^{-\Theta(\log n)} + \min\left\{1, M^{\Delta-(1-o(1))r_1}\right\} \quad (46)$$

which holds for $\Delta \leq \lceil n/4 \rceil$.

Lemma 5.2 does not require any assumption on the graph and can be applied to (42) directly. The proof is deferred to Appendix B.1. Similarly,

$$\mathcal{P}_{\mathbf{x}^*}(E_{2,S_2} \mid r_2) \leq n^{-\Theta(\log n)} + \min\left\{1, M^{\Delta-(1-o(1))r_2}\right\}.$$

Plugging this into (42) yields

$$\begin{aligned} \mathcal{P}_{\mathbf{x}^*}(E_2 \mid r_1, r_2, r^*) &\leq \left(n^{-\Theta(\log n)} + \min\left\{1, M^{\Delta-(1-o(1))r_1}\right\}\right) \cdot \left(n^{-\Theta(\log n)} + \min\left\{1, M^{\Delta-(1-o(1))r_2}\right\}\right) \\ &\leq n^{-\Theta(\log n)} + M^{2\Delta-(1-o(1))(r_1+r_2)} \\ &= n^{-\Theta(\log n)} + M^{2\Delta-(1-o(1))r}. \end{aligned} \quad (47)$$

As a result, the upper bound (47) does not depend on r_1 and r_2 but only their sum r .

Now, by plugging (47) into (39) and taking the summation over r and r^* , we obtain the following bound on \mathcal{P}_Δ . The details are left in Appendix B.2.

Lemma 5.3. \mathcal{P}_Δ in (39) satisfies

$$\mathcal{P}_\Delta \leq M^{2\Delta} \cdot \exp\left(-\frac{N_\Delta \log n}{n} \left(a + b - (2 + o(1))\sqrt{\frac{ab}{M}}\right)\right)$$

where $N_\Delta := 2\Delta\left(\frac{n}{2} - \Delta\right)$.

Step 4: Sum over Δ Now, we are ready to bound the error probability $\mathcal{P}_{e,c}(\psi_{\text{MLE}})$ in (34), which is given as

$$\begin{aligned} \mathcal{P}_{e,c}(\psi_{\text{MLE}}) &\leq 2 \sum_{\Delta=1}^{\lceil n/4 \rceil} \binom{n/2}{\Delta}^2 \cdot M^{2\Delta} \cdot \exp\left(-\frac{N_\Delta \log n}{n} \left(a + b - (2 + o(1))\sqrt{\frac{ab}{M}}\right)\right) \\ &\stackrel{(a)}{\leq} 2 \underbrace{\sum_{\Delta=1}^{\lceil n/4 \rceil} \left(\frac{en}{2\Delta}\right)^{2\Delta} \cdot M^{2\Delta} \cdot \exp\left(-\frac{N_\Delta \log n}{n} \left(a + b - (2 + o(1))\sqrt{\frac{ab}{M}}\right)\right)}_{=:g(\Delta)} \\ &\leq 2 \sum_{\Delta=1}^{\lceil n/4 \rceil} g(\Delta) \end{aligned} \quad (48)$$

where (a) uses the property of binomial coefficient that $\binom{n}{k} \leq \left(\frac{n}{k}\right)^k$ and we define each term in the summation as $g(\Delta)$. To proceed, we take the logarithm of $g(\Delta)$ as

$$\begin{aligned}\log g(\Delta) &= 2\Delta \log\left(\frac{en}{2\Delta}\right) + 2\Delta \log M - \frac{N_\Delta \log n}{n} \cdot \left(a + b - (2 + o(1))\sqrt{\frac{ab}{M}}\right) \\ &= 2\Delta \left[\log\left(\frac{en}{2\Delta}\right) + \log M - \left(\frac{1}{2} - \frac{\Delta}{n}\right) \cdot \left(a + b - (2 + o(1))\sqrt{\frac{ab}{M}}\right) \log n \right].\end{aligned}$$

Then, we study $g(\Delta)$ on the following two cases depending Δ :

- **Case 1:** When $1 \leq \Delta \leq \delta_1 n$, where $\delta_1 = \frac{c-c_1}{2(1+c)}$ for any $c_1 \in [\frac{2c}{3}, c]$ (recall that c is given in the condition (8)). Then,

$$\begin{aligned}\log g(\Delta) &\stackrel{(a)}{\leq} 2\Delta \left(\log n - \left(\frac{1}{2} - \delta_1\right) \cdot \left(a + b - (2 + o(1))\sqrt{\frac{ab}{M}}\right) \log n + O(1) \right) \\ &\stackrel{(b)}{\leq} 2\Delta (\log n - (1 - 2\delta_1)(1 + c) \log n + O(1)) \\ &= -2(1 - o(1)) \cdot c_1 \Delta \log n\end{aligned}\tag{49}$$

where (a) comes from the range $1 \leq \Delta \leq \delta_1 n$ and (b) holds by plugging in the condition (8). As a result, $g(\Delta)$ converges to zero exponentially fast in this case.

- **Case 2:** When $\delta_1 n < \Delta \leq \lceil \frac{n}{4} \rceil$, similarly

$$\begin{aligned}\log g(\Delta) &\leq 2\Delta \left(\log\left(\frac{2e}{\delta_1}\right) + \log M - \frac{1}{4} \cdot \left(a + b - (2 + o(1))\sqrt{\frac{ab}{M}}\right) \log n \right) \\ &\leq 2\Delta \left(-\left(\frac{1+c}{2}\right) \log n + O(1) \right) \\ &= -(1+c)\Delta \log n\end{aligned}\tag{50}$$

which implies $g(\Delta)$ converges to zero super-exponentially fast in this case.

Finally, assembling (49) and (50) and plugging them into (48) yields the upper bound on $\mathcal{P}_{e,c}(\psi_{\text{MLE}})$ as

$$\begin{aligned}\mathcal{P}_{e,c}(\psi_{\text{MLE}}) &\leq 2 \sum_{\Delta=1}^{\lfloor \delta_1 n \rfloor} n^{-2(1-o(1))c_1 \Delta} + 2 \sum_{\Delta=\lfloor \delta_1 n \rfloor+1}^{\lceil n/4 \rceil} n^{-(1+c)\Delta} \\ &\leq n^{-2(1-o(1))c_1} \leq n^{-c},\end{aligned}$$

where the last step uses $c \in [\frac{2c}{3}, c]$. This completes the proof.

5.3 Proof of Theorem 3.2

Our proof technique is mainly inspired by the one of [1, Theorem 1]. Similar to the proof of Theorem 5.2, we still follow the assumption (32). To begin with, recall the MLE (27) fails to exactly recover the communities $\boldsymbol{\kappa}^*$ if and only if there exists an alternative hypothesis $\boldsymbol{x} = \{\boldsymbol{\kappa}, \boldsymbol{g}\}$ that

$$\text{dist}_c(\boldsymbol{\kappa}, \boldsymbol{\kappa}^*) > 0 \quad \text{and} \quad \mathcal{P}(\boldsymbol{y}|\boldsymbol{x}) \geq \mathcal{P}(\boldsymbol{y}|\boldsymbol{x}^*).$$

Also, recall the likelihood ratio given in (31):

$$\frac{\mathcal{P}(\mathbf{y}|\mathbf{x})}{\mathcal{P}(\mathbf{y}|\mathbf{x}^*)} = \left(\frac{Mp(1-q)}{q(1-p)} \right)^{r-r^*} \cdot \frac{\mathbb{1}(g_{ij} = g_i g_j^{-1}; \forall (i, j) \in \mathcal{E}_{\text{inner}})}{\mathbb{1}(g_{ij} = g_i^* (g_j^*)^{-1}; \forall (i, j) \in \mathcal{E}_{\text{inner}}^*)}.$$

Then, depending on the value of $\frac{Mp(1-q)}{q(1-p)}$, we consider the following two cases:

Case 1 We first consider when

$$\frac{Mp(1-q)}{q(1-p)} \geq 1.$$

By plugging in the scaling $p = \frac{a \log n}{n}$ and $q = \frac{b \log n}{n}$ this is equivalent to $\frac{Ma}{b} \geq 1$, and the MLE fails only if $r \geq r^*$.

For ease of presentation, without specification, we denote $\mathcal{P}(\cdot)$ as the probability conditioning on the ground truth \mathbf{x}^* and omit the subscript. Then, let us define the following events:

F : the MLE (27) fails

$E_A^{(i)}$: node $i \in S_1^*$ (resp. S_2^*) is connecting to more nodes in S_2^* (resp. S_1^*) than in S_1^* (resp. S_2^*)
i.e.

$$|\mathcal{E}(i, S_1^* \setminus i)| < |\mathcal{E}(i, S_2^*)|$$

(resp. $|\mathcal{E}(i, S_2^* \setminus i)| < |\mathcal{E}(i, S_1^*)|$)

$E_B^{(i, g_i)}$: node $i \in S_1^*$ (resp. S_2^*) satisfies $g_{ij} = g_i$, for any $(i, j) \in \mathcal{E}(i, S_2^*)$ (resp. $\mathcal{E}(i, S_1^*)$)

We further define

$$\begin{aligned} E^{(i)} &:= E_A^{(i)} \cap \left(\bigcup_{g_i \in \mathcal{G}_M} E_B^{(i, g_i)} \right), \\ E_1 &:= \bigcup_{i \in S_1^*} E^{(i)}, \\ E_2 &:= \bigcup_{i \in S_2^*} E^{(i)}. \end{aligned} \tag{51}$$

Given the above, we have the following relation between E_1 and F :

Lemma 5.4. *If $\mathcal{P}(E_1) \geq \frac{4}{5}$, then $\mathcal{P}(F) \geq \frac{3}{5}$.*

Proof. By symmetry, we have $\mathcal{P}(E_1) = \mathcal{P}(E_2) \geq \frac{4}{5}$. Then, consider the event $E = E_1 \cap E_2$ occurs, which indicates there exists a pair of nodes $i \in S_1^*$ and $j \in S_2^*$ with the two events $E^{(i)}$ and $E^{(j)}$ occur with some g_i and g_j . As a result, one can construct an alternative hypothesis \mathbf{x} from \mathbf{x}^* by swapping the community assignment of i and j i.e. $i \in S_2$ and $j \in S_1$, and let the associated group elements be g_i and g_j , then $\mathcal{P}(\mathbf{y}|\mathbf{x}) > \mathcal{P}(\mathbf{y}|\mathbf{x}^*)$, indicating F occurs. Therefore,

$$\mathcal{P}(F) \geq \mathcal{P}(E_1 \cap E_2) \geq \mathcal{P}(E_1) + \mathcal{P}(E_2) - 1 \geq \frac{3}{5}$$

which completes the proof. \square

To proceed, let us denote the following two quantities:

$$\gamma := \log^3 n, \quad \delta := \frac{\log n}{\log \log n}$$

and we assume them to be integers (otherwise round them to the nearest ones). Then, let H be a fixed subset of S_1^* of size $\frac{n}{\gamma}$, we further define the following two events

F_H : no node in H connects to at least δ number of other nodes in H

$E_{H,A}^{(i)}$: node $i \in H$ satisfies $|\mathcal{E}(i, S_1^* \setminus H)| + \delta \leq |\mathcal{E}(i, S_2^*)|$

and we denote

$$E_H^{(i)} := E_{H,A}^{(i)} \cap \left(\bigcup_{g_i \in \mathcal{G}_M} E_B^{(i,g_i)} \right),$$

$$E_H := \bigcup_{i \in H} E_H^{(i)}.$$

Then we have

Lemma 5.5. *If $\mathcal{P}(E_H) \geq \frac{9}{10}$, then $\mathcal{P}(F) \geq \frac{3}{5}$.*

Proof. By definition, E_H occurs when there exists a node $i \in H$ with some g_i that such that $E_{H,A}^{(i)}$ and $E_B^{(i,g_i)}$ occur. Then one can see that $F_H \cap E_H \subseteq E_1$, as $F_H \cap E_H^{(i)}$ indicates $E^{(i)}$ occurs. Therefore, by applying $\mathcal{P}(F_H) \geq \frac{9}{10}$ from [1, Lemma 10], we have

$$\mathcal{P}(E_1) \geq \mathcal{P}(F_H \cap E_H) \geq \mathcal{P}(F_H) + \mathcal{P}(E_H) - 1 \geq \frac{4}{5}.$$

By further applying Lemma 5.4 we complete the proof. \square

Now, we focus on bounding $\mathcal{P}(E_H)$. Importantly, it satisfies

$$\begin{aligned} \mathcal{P}(E_H) &= \mathcal{P} \left(\bigcup_{i \in H} E_H^{(i)} \right) = 1 - \mathcal{P} \left(\bigcap_{i \in H} (E_H^{(i)})^c \right) \stackrel{(a)}{=} 1 - \prod_{i \in H} \mathcal{P} \left((E_H^{(i)})^c \right) \\ &= 1 - \prod_{i \in H} \left(1 - \mathcal{P} \left(E_H^{(i)} \right) \right) = 1 - \prod_{i \in H} \left(1 - \mathcal{P} \left(E_{H,A}^{(i)} \cap \left(\bigcup_{g_i \in \mathcal{G}_M} E_B^{(i,g_i)} \right) \right) \right) \\ &\stackrel{(b)}{\geq} 1 - \prod_{i \in H} \left(1 - \mathcal{P} \left(E_{H,A}^{(i)} \cap E_B^{(i,0)} \right) \right) \end{aligned} \quad (52)$$

where (a) holds since the set of events $\{E_H^{(i)}\}_{i \in H}$ are independent, as they involve mutually disjoint sets of edge connections, and (b) comes by focusing on the case of $g_i = 0$. Then, our next job is to lower bound $\mathcal{P} \left(E_{H,A}^{(i)} \cap E_B^{(i,0)} \right)$. To this end, let us denote

$$r_1 = |\mathcal{E}(i, S_1^* \setminus H)|, \quad r_2 = |\mathcal{E}(i, S_2^*)|. \quad (53)$$

Notice that r_1 and r_2 are two independent binomial random variables such that

$$r_1 \sim \text{Binom} \left(\frac{n}{2} - \frac{n}{\gamma}, p \right), \quad r_2 \sim \text{Binom} \left(\frac{n}{2}, q \right). \quad (54)$$

Then, it satisfies

$$\begin{aligned}
\mathcal{P}\left(E_{H,A}^{(i)} \cap E_B^{(i,0)}\right) &= \sum_{r_1} \sum_{r_2} \mathcal{P}(r_1) \mathcal{P}(r_2) \mathcal{P}\left(E_{H,A}^{(i)} \cap E_B^{(i,0)} \mid r_1, r_2\right) \\
&\stackrel{(a)}{=} \sum_{r_1} \sum_{r_2} \mathcal{P}(r_1) \mathcal{P}(r_2) \mathcal{P}\left(E_{H,A}^{(i)} \mid r_1, r_2\right) \cdot \mathcal{P}\left(E_B^{(i,0)} \mid r_1, r_2\right) \\
&\stackrel{(b)}{=} \sum_{r_1} \sum_{r_2} \mathcal{P}(r_1) \mathcal{P}(r_2) \cdot \mathbb{1}(r_2 \geq r_1 + \delta) \cdot M^{-r_2}.
\end{aligned} \tag{55}$$

Here, (a) comes from the fact that the two events $E_{H,A}^{(i)}$ and $E_B^{(i,0)}$ occurs independently given r_1 and r_2 ; (b) holds as given r_1 and r_2 , $E_{H,A}^{(i)}$ is deterministic, and by definition $E_B^{(i,0)}$ occurs only if $g_{ij} = 0$ for all $(i, j) \in \mathcal{E}(i, S_2^*)$, and each $g_{ij} = 0$ occurs independent with probability M^{-1} . Following this way with some heavy calculation, we obtain

Lemma 5.6. *Given $p = \frac{a \log n}{n}$ and $q = \frac{b \log n}{n}$, (55) satisfies*

$$\log \mathcal{P}\left(E_{H,A}^{(i)} \cap E_B^{(i,0)}\right) \geq \left(-\frac{a+b}{2} + \sqrt{\frac{ab}{M}}\right) \log n - o(\log n). \tag{56}$$

The proof of Lemma 5.6 is deferred to Appendix B.3. As a result, plugging (56) into (52) yields:

$$\begin{aligned}
\mathcal{P}(E_H) &\geq 1 - \prod_{i \in H} \left(1 - \mathcal{P}\left(E_{H,A}^{(i)} \cap E_B^{(i,0)}\right)\right) \\
&\stackrel{(a)}{\geq} 1 - \prod_{i \in H} \exp\left(-\mathcal{P}\left(E_{H,A}^{(i)} \cap E_B^{(i,0)}\right)\right) \\
&\stackrel{(b)}{=} 1 - \exp\left(-|H| \cdot \mathcal{P}\left(E_{H,A}^{(i)} \cap E_B^{(i,0)}\right)\right) \\
&\geq 1 - \exp\left(-n^{-\frac{a+b}{2} + \sqrt{\frac{ab}{M}} + 1 - o(1)}\right) \\
&\stackrel{(c)}{\geq} \frac{9}{10}
\end{aligned} \tag{57}$$

which holds as n is sufficiently large. Here, (a) holds by using $1 - x \leq e^{-x}$, $\forall x \in \mathbb{R}$; (b) comes from the fact that $\mathcal{P}\left(E_{H,A}^{(i)} \cap E_B^{(i,0)}\right)$ should be identical to all $i \in H$; (c) applies the condition in (9). Therefore, by further applying Lemma 5.5 we complete the proof for this case.

Case 2. Now we consider

$$\frac{Mp(1-q)}{q(1-p)} < 1$$

or equivalently $\frac{Ma}{b} < 1$, which indicates MLE fails only if $r \leq r^*$. The proof is very similar to case 1 where most of the result still applies at here. Therefore, we will skip those repeated analyses. To begin with, we reuse all the notations of events in case 1 but redefine the following two:

$E_A^{(i)}$: node $i \in S_1^*$ (resp. S_2^*) is connecting to more nodes in S_1^* (resp. S_2^*) than in S_2^* (resp. S_1^*)
i.e.

$$|\mathcal{E}(i, S_1^* \setminus i)| > |\mathcal{E}(i, S_2^*)|$$

(resp. $|\mathcal{E}(i, S_2^* \setminus i)| < |\mathcal{E}(i, S_1^*)|$)

$E_{H,A}^{(i)}$: node $i \in H$ satisfies $|\mathcal{E}(i, S_1^* \setminus H)| > |\mathcal{E}(i, S_2^*)|$

where H is still the fixed subset of S_1^* of size $\frac{n}{\gamma}$ and $\gamma = \log^3 n$. In this way, Lemmas 5.4 and 5.5 still hold in this case.³ Furthermore, we follow the steps in (55) and provide a lower bound on $\mathcal{P}\left(E_{H,A}^{(i)} \cap E_B^{(i,0)}\right)$ similar to Lemma 5.6. To this end, by using r_1 and r_2 defined in (53), we have

$$\begin{aligned}
\mathcal{P}\left(E_{H,A}^{(i)} \cap E_B^{(i,0)}\right) &= \sum_{r_1} \sum_{r_2} \mathcal{P}(r_1) \mathcal{P}(r_2) \mathcal{P}\left(E_{H,A}^{(i)} \cap E_B^{(i,0)} \mid r_1, r_2\right) \\
&= \sum_{r_1} \sum_{r_2} \mathcal{P}(r_1) \mathcal{P}(r_2) \mathcal{P}\left(E_{H,A}^{(i)} \mid r_1, r_2\right) \cdot \mathcal{P}\left(E_B^{(i,0)} \mid r_1, r_2\right) \\
&= \sum_{r_1} \sum_{r_2} \mathcal{P}(r_1) \mathcal{P}(r_2) \cdot \mathbb{1}(r_1 > r_2) \cdot M^{-r_2} \\
&= \sum_{r_2} \mathcal{P}(r_2) \mathcal{P}(r_1 > r_2) \cdot M^{-r_2}.
\end{aligned} \tag{58}$$

Then, similar to Lemma 5.6, we obtain

Lemma 5.7. *Given $p = \frac{a \log n}{n}$ and $q = \frac{b \log n}{n}$, (58) satisfies*

$$\begin{aligned}
\log \mathcal{P}\left(E_{H,A}^{(i)} \cap E_B^{(i,0)}\right) &\geq \left(-\frac{a+b}{2} + \sqrt{\frac{ab}{M}}\right) \log n \\
&\quad - o(\log n).
\end{aligned} \tag{59}$$

The proof of Lemma 5.7 is deferred to Appendix B.4, which is highly similar to the one of Lemma 5.6. As a result, by following the same steps as in (57) we complete the proof.

5.4 Proof of Theorem 3.3

To begin with, we follow the same path of proving Theorem 3.1 in Section 5.2 by writing out the probability that the MLE (27) fails to exactly recover the group elements as

$$\begin{aligned}
\mathcal{P}_{e,g}(\psi_{\text{MLE}}) &= \mathcal{P}\left(\exists \mathbf{x} = \{\boldsymbol{\kappa}, \mathbf{g}\} : \text{dist}_g(\mathbf{g}, \mathbf{g}^*) > 0, \frac{\mathcal{P}(\mathbf{y}|\mathbf{x})}{\mathcal{P}(\mathbf{y}|\mathbf{x}^*)} \geq 1\right) \\
&\stackrel{(a)}{\leq} \mathcal{P}\left(\exists \mathbf{x} = \{\boldsymbol{\kappa}, \mathbf{g}\} : \text{dist}_c(\boldsymbol{\kappa}, \boldsymbol{\kappa}^*) > 0, \text{dist}_g(\mathbf{g}, \mathbf{g}^*) > 0, \frac{\mathcal{P}(\mathbf{y}|\mathbf{x})}{\mathcal{P}(\mathbf{y}|\mathbf{x}^*)} \geq 1\right) \\
&\quad + \mathcal{P}\left(\exists \mathbf{x} = \{\boldsymbol{\kappa}, \mathbf{g}\} : \text{dist}_c(\boldsymbol{\kappa}, \boldsymbol{\kappa}^*) = 0, \text{dist}_g(\mathbf{g}, \mathbf{g}^*) > 0, \frac{\mathcal{P}(\mathbf{y}|\mathbf{x})}{\mathcal{P}(\mathbf{y}|\mathbf{x}^*)} \geq 1\right) \\
&\stackrel{(b)}{\leq} n^{-c} + \mathcal{P}\left(\exists \mathbf{x} = \{\boldsymbol{\kappa}^*, \mathbf{g}\} : \text{dist}_g(\mathbf{g}, \mathbf{g}^*) > 0, \frac{\mathcal{P}(\mathbf{y}|\mathbf{x})}{\mathcal{P}(\mathbf{y}|\mathbf{x}^*)} \geq 1\right).
\end{aligned} \tag{60}$$

Here, in step (a) we split it into two cases whether the community assignment $\boldsymbol{\kappa}$ is correct or not, then apply the union bound; (b) uses the fact that the condition

$$\frac{a+b}{2} - (1+o(1))\sqrt{\frac{ab}{M}} > 1+c$$

³A slight difference is that we no longer need the event F_H , and therefore E_H implies E_1 directly.

satisfies and therefore from Theorem 3.1, $\boldsymbol{\kappa}$ is incorrect w.p. at most n^{-c} . It remains to consider when $\boldsymbol{\kappa} = \boldsymbol{\kappa}^*$. In this case, the likelihood ratio in (31) satisfies

$$\frac{\mathcal{P}(\mathbf{y}|\mathbf{x})}{\mathcal{P}(\mathbf{y}|\mathbf{x}^*)} = \frac{\mathbb{1}(g_{ij} = g_i g_j^{-1}; \forall (i, j) \in \mathcal{E}_{\text{inner}}^*)}{\mathbb{1}(g_{ij} = g_i^* (g_j^*)^{-1}; \forall (i, j) \in \mathcal{E}_{\text{inner}}^*)}$$

where we use the fact that $r = r^*$ and $\mathcal{E}_{\text{inner}} = \mathcal{E}_{\text{inner}}^*$. Therefore, recall the definition $\mathcal{E}_{\text{inner}}^* = \mathcal{E}(S_1^*) \cup \mathcal{E}(S_2^*)$, for any $\mathbf{x} = \{\boldsymbol{\kappa}^*, \mathbf{g}\}$ with a wrong hypothesis \mathbf{g} , one can see that $\mathcal{P}(\mathbf{y}|\mathbf{x}) \geq \mathcal{P}(\mathbf{y}|\mathbf{x}^*)$ occurs only if both subgraphs formed by S_1^* and S_2^* are disconnected (otherwise there exists an edge $(i, j) \in \mathcal{E}_{\text{inner}}^*$ with $g_{ij} \neq g_i g_j^{-1}$). Therefore, by denoting E_1 and E_2 as the events that the subgraph of S_1^* and S_2^* are connected respectively, we have

$$\begin{aligned} \mathcal{P}\left(\exists \mathbf{x} = \{\boldsymbol{\kappa}^*, \mathbf{g}\} : \text{dist}_{\mathbf{g}}(\mathbf{g}, \mathbf{g}^*) > 0, \frac{\mathcal{P}(\mathbf{y}|\mathbf{x})}{\mathcal{P}(\mathbf{y}|\mathbf{x}^*)} \geq 1\right) &\leq \mathcal{P}(E_1^c \cap E_2^c) \\ &= 1 - \mathcal{P}(E_1 \cap E_2) \\ &= 1 - \mathcal{P}(E_1) \cdot \mathcal{P}(E_2) \\ &\leq 1 - (1 - n^{-c+o(1)}) \cdot (1 - n^{-c+o(1)}) \\ &= n^{-c+o(1)}. \end{aligned} \tag{61}$$

Here, we apply Lemma A.1 which suggests

$$\mathcal{P}(E_1) = \mathcal{P}(E_2) \geq 1 - n^{-c+o(1)}$$

given the condition $\frac{a}{2} > 1 + c$ satisfies. Plugging (61) into (60) completes the proof.

5.5 Proof of Theorem 3.4

We will show the necessity of the two conditions for the exact recovery of group elements separately. First, when $\frac{a}{2} < 1$, Lemma A.1 suggests that the two subgraphs formed by S_1^* and S_2^* are disconnected with prob. $1 - o(1)$. As a result, when either one is disconnected, there exist at least two components in the subgraph, and furthermore, there is no way to synchronize the group elements between them but only guess and randomly draw from \mathcal{G}_M . Therefore, in this case, the probability of having a wrong \mathbf{g} estimated is at least $1 - M^{-1} \geq 1/2$, where the equality holds when $M = 2$.

Secondly, when the condition

$$\frac{a+b}{2} - \sqrt{\frac{ab}{M}} < 1$$

satisfies, Theorem 3.2 suggests that the estimated community assignment $\boldsymbol{\kappa}$ is inaccurate with probability at least $3/5$. When this occurs, there exists at least one pair of wrong classified vertices (i, j) such that $i \in S_1^*$ and $j \in S_2^*$ but $i \in S_2$ and $j \in S_1$. As a result, the estimated group elements on i and j , denoted by g_i and g_j , are completely determined by those edges across S_1^* and S_2^* , whose group transformations are randomly drawn from \mathcal{G}_M . Therefore, the probability of having a wrong estimated g_i or g_j is at least $1 - M^{-2}$. Combining this with the event of having a wrong community assignment $\boldsymbol{\kappa}$ yields the total probability of having the wrong group elements \mathbf{g} at least $\frac{3}{5} \cdot (1 - M^{-2}) \geq 0.45$, where the equality holds when $M = 2$. This completes the proof.

Acknowledgments

ZZ and YF acknowledge the support from NSF grant DMS-1854791 and Alfred P. Sloan foundation. We thank Yuehaw Khoo for valuable discussions.

References

- [1] Emmanuel Abbe, Afonso S Bandeira, and Georgina Hall. Exact recovery in the stochastic block model. *IEEE Transactions on Information Theory*, 62(1):471–487, 2015.
- [2] Emmanuel Abbe, Jianqing Fan, Kaizheng Wang, Yiqiao Zhong, et al. Entrywise eigenvector analysis of random matrices with low expected rank. *Annals of Statistics*, 48(3):1452–1474, 2020.
- [3] Emmanuel Abbe and Colin Sandon. Community detection in general stochastic block models: Fundamental limits and efficient algorithms for recovery. In *2015 IEEE 56th Annual Symposium on Foundations of Computer Science*, pages 670–688. IEEE, 2015.
- [4] Arash A Amini and Elizaveta Levina. On semidefinite relaxations for the block model. *The Annals of Statistics*, 46(1):149–179, 2018.
- [5] Federica Arrigoni, Beatrice Rossi, and Andrea Fusiello. Spectral synchronization of multiple views in se (3). *SIAM Journal on Imaging Sciences*, 9(4):1963–1990, 2016.
- [6] Chandrajit Bajaj, Tingran Gao, Zihang He, Qixing Huang, and Zhenxiao Liang. SMAC: simultaneous mapping and clustering using spectral decompositions. In *International Conference on Machine Learning*, pages 324–333, 2018.
- [7] Afonso S Bandeira. Random laplacian matrices and convex relaxations. *Foundations of Computational Mathematics*, 18(2):345–379, 2018.
- [8] Ravi B Boppana. Eigenvalues and graph bisection: An average-case analysis. In *28th Annual Symposium on Foundations of Computer Science (sfcs 1987)*, pages 280–285. IEEE, 1987.
- [9] Thang Bui, Soma Chaudhuri, Tom Leighton, and Michael Sipser. Graph bisection algorithms with good average case behavior. In *25th Annual Symposium on Foundations of Computer Science, 1984.*, pages 181–192. IEEE, 1984.
- [10] Ted Carson and Russell Impagliazzo. Hill-climbing nds random planted bisections. In *Proceedings of the 12th Annual SIAM Symposium on Discrete Algorithms*. Citeseer, 2001.
- [11] Kunal N Chaudhury, Yuehaw Khoo, and Amit Singer. Global registration of multiple point clouds using semidefinite programming. *SIAM Journal on Optimization*, 25(1):468–501, 2015.
- [12] Sijin Chen, Xiwei Cheng, and Anthony Man-Cho So. Non-convex joint community detection and group synchronization via generalized power method. *arXiv preprint arXiv:2112.14204*, 2021.
- [13] Yuxin Chen, Leonidas Guibas, and Qixing Huang. Near-optimal joint object matching via convex relaxation. In *International Conference on Machine Learning*, pages 100–108. PMLR, 2014.
- [14] Yuxin Chen, Changho Suh, and Andrea J Goldsmith. Information recovery from pairwise measurements. *IEEE Transactions on Information Theory*, 62(10):5881–5905, 2016.
- [15] David S Choi, Patrick J Wolfe, and Edoardo M Airoldi. Stochastic blockmodels with a growing number of classes. *Biometrika*, 99(2):273–284, 2012.

- [16] Anne Condon and Richard M Karp. Algorithms for graph partitioning on the planted partition model. *Random Structures & Algorithms*, 18(2):116–140, 2001.
- [17] Aurelien Decelle, Florent Krzakala, Christopher Moore, and Lenka Zdeborová. Asymptotic analysis of the stochastic block model for modular networks and its algorithmic applications. *Physical Review E*, 84(6):066106, 2011.
- [18] Patrick Doreian, Vladimir Batagelj, and Anuska Ferligoj. *Generalized blockmodeling*, volume 25. Cambridge university press, 2005.
- [19] Martin E. Dyer and Alan M. Frieze. The solution of some random np-hard problems in polynomial expected time. *Journal of Algorithms*, 10(4):451–489, 1989.
- [20] Paul Erdős, Alfréd Rényi, et al. On the evolution of random graphs. *Publ. Math. Inst. Hung. Acad. Sci*, 5(1):17–60, 1960.
- [21] Yifeng Fan, Tingran Gao, and Zhizhen Zhao. Unsupervised co-learning on G -manifolds across irreducible representations. In *Advances in Neural Information Processing Systems*, pages 9041–9053, 2019.
- [22] Yifeng Fan, Tingran Gao, and Zhizhen Zhao. Representation theoretic patterns in multi-frequency class averaging for three-dimensional cryo-electron microscopy. *Information and Inference: A journal of IMA*, page accepted, 2021.
- [23] Yifeng Fan, Yuehaw Khoo, and Zhizhen Zhao. A spectral method for joint community detection and orthogonal group synchronization. *arXiv preprint arXiv:2112.13199*, 2021.
- [24] Yifeng Fan, Yuehaw Khoo, and Zhizhen Zhao. Joint community detection and rotational synchronization via semidefinite programming. *SIAM Journal on Mathematics of Data Science*, 4(3):1052–1081, 2022.
- [25] Yifeng Fan, Yuehaw Khoo, and Zhizhen Zhao. Joint community detection and rotational synchronization via semidefinite programming. *SIAM Journal on Mathematics of Data Science*, 4(3):1052–1081, 2022.
- [26] Yifeng Fan and Zhizhen Zhao. Multi-frequency vector diffusion maps. In *International Conference on Machine Learning*, pages 1843–1852. PMLR, 2019.
- [27] Uriel Feige and Eran Ofek. Spectral techniques applied to sparse random graphs. *Random Structures & Algorithms*, 27(2):251–275, 2005.
- [28] Stephen E Fienberg, Michael M Meyer, and Stanley S Wasserman. Statistical analysis of multiple sociometric relations. *Journal of the american Statistical association*, 80(389):51–67, 1985.
- [29] Joachim Frank. *Three-Dimensional Electron Microscopy of Macromolecular Assemblies: Visualization of Biological Molecules in Their Native State*. Oxford University Press, New York, 2nd edition, 2006.
- [30] Brendan J Frey and Delbert Dueck. Clustering by passing messages between data points. *science*, 315(5814):972–976, 2007.
- [31] Alan Frieze and Michał Karoński. *Introduction to random graphs*. Cambridge University Press, 2016.

- [32] Chao Gao, Zongming Ma, Anderson Y Zhang, and Harrison H Zhou. Achieving optimal misclassification proportion in stochastic block models. *The Journal of Machine Learning Research*, 18(1):1980–2024, 2017.
- [33] Chao Gao and Anderson Y Zhang. Optimal orthogonal group synchronization and rotation group synchronization. *arXiv preprint arXiv:2109.13491*, 2021.
- [34] Tingran Gao and Zhizhen Zhao. Multi-frequency phase synchronization. In *International Conference on Machine Learning*, pages 2132–2141. PMLR, 2019.
- [35] Michael R Garey, David S Johnson, and Larry Stockmeyer. Some simplified np-complete problems. In *Proceedings of the sixth annual ACM symposium on Theory of computing*, pages 47–63, 1974.
- [36] Olivier Guédon and Roman Vershynin. Community detection in sparse networks via grothendieck’s inequality. *Probability Theory and Related Fields*, 165(3):1025–1049, 2016.
- [37] Bruce Hajek, Yihong Wu, and Jiaming Xu. Achieving exact cluster recovery threshold via semidefinite programming. *IEEE Transactions on Information Theory*, 62(5):2788–2797, 2016.
- [38] Bruce Hajek, Yihong Wu, and Jiaming Xu. Achieving exact cluster recovery threshold via semidefinite programming: Extensions. *IEEE Transactions on Information Theory*, 62(10):5918–5937, 2016.
- [39] Paul W Holland, Kathryn Blackmond Laskey, and Samuel Leinhardt. Stochastic blockmodels: First steps. *Social networks*, 5(2):109–137, 1983.
- [40] Qi-Xing Huang and Leonidas Guibas. Consistent shape maps via semidefinite programming. In *Computer Graphics Forum*, volume 32, pages 177–186. Wiley Online Library, 2013.
- [41] Qi-Xing Huang, Guo-Xin Zhang, Lin Gao, Shi-Min Hu, Adrian Butscher, and Leonidas Guibas. An optimization approach for extracting and encoding consistent maps in a shape collection. *ACM Transactions on Graphics (TOG)*, 31(6):1–11, 2012.
- [42] Mark Jerrum and Gregory B Sorkin. The metropolis algorithm for graph bisection. *Discrete Applied Mathematics*, 82(1-3):155–175, 1998.
- [43] Brian Karrer and Mark EJ Newman. Stochastic blockmodels and community structure in networks. *Physical review E*, 83(1):016107, 2011.
- [44] Florent Krzakala, Cristopher Moore, Elchanan Mossel, Joe Neeman, Allan Sly, Lenka Zdeborová, and Pan Zhang. Spectral redemption in clustering sparse networks. *Proceedings of the National Academy of Sciences*, 110(52):20935–20940, 2013.
- [45] Roy R Lederman and Amit Singer. A representation theory perspective on simultaneous alignment and classification. *Applied and Computational Harmonic Analysis*, 2019.
- [46] Jing Lei and Alessandro Rinaldo. Consistency of spectral clustering in stochastic block models. *The Annals of Statistics*, 43(1):215–237, 2015.
- [47] Xiaodong Li, Yudong Chen, Jiaming Xu, et al. Convex relaxation methods for community detection. *Statistical Science*, 36(1):2–15, 2021.

- [48] Shuyang Ling. Near-optimal performance bounds for orthogonal and permutation group synchronization via spectral methods. *arXiv preprint arXiv:2008.05341*, 2020.
- [49] Shuyang Ling. Solving orthogonal group synchronization via convex and low-rank optimization: Tightness and landscape analysis. *arXiv preprint arXiv:2006.00902*, 2020.
- [50] Laurent Massoulié. Community detection thresholds and the weak ramanujan property. In *Proceedings of the forty-sixth annual ACM symposium on Theory of computing*, pages 694–703, 2014.
- [51] Frank McSherry. Spectral partitioning of random graphs. In *Proceedings 42nd IEEE Symposium on Foundations of Computer Science*, pages 529–537. IEEE, 2001.
- [52] Elchanan Mossel, Joe Neeman, and Allan Sly. Consistency thresholds for binary symmetric block models. *arXiv preprint arXiv:1407.1591*, 3(5), 2014.
- [53] Elchanan Mossel, Joe Neeman, and Allan Sly. Reconstruction and estimation in the planted partition model. *Probability Theory and Related Fields*, 162(3):431–461, 2015.
- [54] Elchanan Mossel, Joe Neeman, and Allan Sly. A proof of the block model threshold conjecture. *Combinatorica*, 38(3):665–708, 2018.
- [55] Andrew Y Ng, Michael I Jordan, and Yair Weiss. On spectral clustering: Analysis and an algorithm. In *Advances in neural information processing systems*, pages 849–856, 2002.
- [56] Deepti Pachauri, Risi Kondor, and Vikas Singh. Solving the multi-way matching problem by permutation synchronization. In *Advances in neural information processing systems*, pages 1860–1868. Citeseer, 2013.
- [57] Amelia Perry and Alexander S Wein. A semidefinite program for unbalanced multisection in the stochastic block model. In *2017 International Conference on Sampling Theory and Applications (SampTA)*, pages 64–67. IEEE, 2017.
- [58] Karl Rohe, Sourav Chatterjee, and Bin Yu. Spectral clustering and the high-dimensional stochastic blockmodel. 2011.
- [59] Yanyao Shen, Qixing Huang, Nati Srebro, and Sujay Sanghavi. Normalized spectral map synchronization. *Advances in Neural Information Processing Systems*, 29:4925–4933, 2016.
- [60] Yunpeng Shi and Gilad Lerman. Message passing least squares framework and its application to rotation synchronization. *arXiv preprint arXiv:2007.13638*, 2020.
- [61] Amit Singer. Angular synchronization by eigenvectors and semidefinite programming. *Applied and computational harmonic analysis*, 30(1):20–36, 2011.
- [62] Amit Singer, Zhizhen Zhao, Yoel Shkolnisky, and Ronny Hadani. Viewing angle classification of cryo-electron microscopy images using eigenvectors. *SIAM Journal on Imaging Sciences*, 4(2):723–759, 2011.
- [63] Tom AB Snijders and Krzysztof Nowicki. Estimation and prediction for stochastic blockmodels for graphs with latent block structure. *Journal of classification*, 14(1):75–100, 1997.
- [64] Van Vu. A simple svd algorithm for finding hidden partitions. *arXiv preprint arXiv:1404.3918*, 2014.

- [65] Lanhui Wang and Amit Singer. Exact and stable recovery of rotations for robust synchronization. *Information and Inference: A Journal of the IMA*, 2(2):145–193, 2013.
- [66] Lingda Wang and Zhizhen Zhao. Multi-frequency joint community detection and phase synchronization. *IEEE Transactions on Signal and Information Processing over Networks*, 2023.
- [67] Se-Young Yun and Alexandre Proutiere. Accurate community detection in the stochastic block model via spectral algorithms. *arXiv preprint arXiv:1412.7335*, 2014.
- [68] Anderson Ye Zhang. Exact minimax optimality of spectral methods in phase synchronization and orthogonal group synchronization. *arXiv preprint arXiv:2209.04962*, 2022.
- [69] Zhizhen Zhao and Amit Singer. Rotationally invariant image representation for viewing direction classification in cryo-EM. *Journal of structural biology*, 186(1):153–166, 2014.
- [70] Yiqiao Zhong and Nicolas Boumal. Near-optimal bounds for phase synchronization. *SIAM Journal on Optimization*, 28(2):989–1016, 2018.

A The giant component of Erdős–Rényi graphs

In this section, we briefly review several classical properties of Erdős–Rényi graph [20] that support our theory. Remarkably, most of the results in this section are by no means original, but we still provide our proofs in detail for completeness.

To begin with, given a network of size n , the Erdős–Rényi model denoted by $G(n, p)$ ⁴ generates a random graph (denoted by \mathcal{G}_n) by connecting each pair of vertices independently with some probability p . In this section, our study focuses on the size of the largest connected component in \mathcal{G}_n , denoted by Z_n .

First, when $p = \frac{a \log n}{n}$ follows the scaling $\Theta\left(\frac{\log n}{n}\right)$ for some constant $a > 0$, it is well-known that $a = 1$ serves as the sharp threshold for \mathcal{G}_n being connected. Formally, we have:

Lemma A.1 (Connectivity). *For any $c > 0$,*

$$\begin{aligned} \mathcal{P}(\mathcal{G}_n \text{ is connected}) &\geq 1 - n^{-c+o(1)}, \quad \text{when } a > 1 + c \\ \mathcal{P}(\mathcal{G}_n \text{ is disconnected}) &\geq 1 - n^{-c+o(1)}, \quad \text{when } a < 1 - c \end{aligned}$$

where the second one also assumes $c < 1$.

For completeness, a proof of Lemma A.1 is given in Appendix A.1, which is basically built from the analysis provided in [31]. Lemma A.1 indicates that the largest connected component will be the whole network w.h.p. when $a > 1$. Furthermore, when $a < 1$, it can be shown that the largest component will be giant w.h.p. such that

$$\mathcal{P}(Z_n > (1 - o(1))n) = 1 - o(1).$$

To be concrete, [31, Theorem 2.14] states

$$\mathcal{P}\left(Z_n > \left(1 - n^{-a+\epsilon+o(1)}\right)n\right) > 1 - n^{-\epsilon} \tag{62}$$

⁴The model originally proposed by Erdős and Rényi is slightly different: instead of connecting each pair of vertices independently, it generates the graph by randomly assigning exactly m edges to the network, and the resulting model is denoted by $G(n, m)$. However, the two models $G(n, m)$ and $G(n, p)$ are closely related, see [31, Chapter 1].

which holds for any $\epsilon < a$. (62) suggests that a giant component exists with a probability at least $n^{-\Theta(1)}$.

Unfortunately, (62) is too loose to be applied for our analysis in Section 5.2 and a tighter result is necessary. For example, we hope

$$\mathcal{P}(Z_n > (1 - o(1))n) > 1 - n^{-\omega(n)} \quad (63)$$

where $\omega(n)$ stands for some slowly growing function such that $\omega(n) \rightarrow \infty$ as $n \rightarrow \infty$. To this end, we derive the following results that meet the requirement in (63):

Lemma A.2 (No medium component when disconnected). *Given the Erdős–Rényi model $G(n, p)$ with $p = \frac{a \log n}{n}$ for some $a \in (0, 1)$, the largest connected component Z_n satisfies*

$$\mathcal{P}\left(\beta_1 \log n \leq Z_n \leq n - \frac{\beta_2 n}{\log \log n}\right) \leq n^{-\Theta(\log n)} \quad (64)$$

for any constants $\beta_1, \beta_2 > 0$.

Lemma A.3 (No small component when disconnected). *Under the same setting of Lemma A.2, it satisfies*

$$\mathcal{P}(Z_n \leq \beta \log n) \leq n^{-\frac{an(1-o(1))}{2}} \quad (65)$$

for any constant $\beta > 0$.

Due to the complexity, the proofs of Lemmas A.2 and A.3 are deferred to Appendices A.2 and A.3, respectively. Remarkably, the proof of Lemma A.3 is original and involves a significant amount of calculation and optimization, because traditional approaches based on the moment method do not work in this case.

Given the above, combining Lemmas A.2 and A.3 yields a desired result like (63) as follows:

Theorem A.4 (Giant component only when disconnected). *Under the same setting of Lemma A.2,*

$$\mathcal{P}\left(Z_n > n - \frac{\beta n}{\log \log n}\right) > 1 - n^{-\Theta(\log n)} \quad (66)$$

for any constant $\beta > 0$.

Proof of Theorem A.4. Plugging in (64) and (65) as

$$\begin{aligned} \mathcal{P}\left(Z_n > n - \frac{\beta n}{\log \log n}\right) &= 1 - \mathcal{P}\left(Z_n \leq n - \frac{\beta n}{\log \log n}\right) \\ &= 1 - \mathcal{P}\left(\beta \log n \leq Z_n \leq n - \frac{\beta n}{\log \log n}\right) - \mathcal{P}(Z_n \leq \beta \log n) \\ &> 1 - n^{-\Theta(\log n)} \end{aligned}$$

completes the proof. □

A.1 Proof of Lemma A.1

We first consider the case when $a > 1 + c$ for any $c > 0$. Given the graph \mathcal{G}_n , let r_k denote the number of connected components of size k . The proof strategy is to sharply bound the expectation $\mathbb{E}[r_k]$ for $1 \leq k \leq n/2$ then apply Markov's inequality. That is

$$\begin{aligned}
\mathcal{P}(\mathcal{G}_n \text{ is connected}) &= \mathcal{P}(r_k = 0, k = 1, \dots, \lfloor n/2 \rfloor) \\
&= \mathcal{P}\left(\sum_{k=1}^{\lfloor n/2 \rfloor} r_k = 0\right) \\
&= 1 - \mathcal{P}\left(\sum_{k=1}^{\lfloor n/2 \rfloor} r_k \geq 1\right) \\
&\geq 1 - \sum_{k=1}^{\lfloor n/2 \rfloor} \mathbb{E}[r_k], \tag{67}
\end{aligned}$$

where the last step applies Markov's inequality. To determine $\mathbb{E}[r_k]$, let \mathcal{G}_k denotes any subgraph of \mathcal{G}_n with size k , and let \mathcal{G}_k^c be the subgraph that contains the remaining vertices, then we have

$$\begin{aligned}
\mathbb{E}[r_k] &\stackrel{(a)}{\leq} \binom{n}{k} \cdot \mathcal{P}(\mathcal{G}_k \text{ is connected and isolated to } \mathcal{G}_k^c) \\
&\stackrel{(b)}{\leq} \binom{n}{k} \cdot \mathcal{P}(\text{A spanning tree of } \mathcal{G}_k \text{ exists and is isolated to } \mathcal{G}_k^c) \\
&\stackrel{(c)}{\leq} \binom{n}{k} \cdot k^{k-2} \cdot \mathcal{P}(\text{The spanning tree exists and is isolated to } \mathcal{G}_k^c) \\
&\leq \binom{n}{k} \cdot k^{k-2} \cdot p^{k-1} \cdot (1-p)^{n-k}. \tag{68}
\end{aligned}$$

Here, step (a) holds by applying the union bound over all subgraphs of size k ; (b) comes from the fact that \mathcal{G}_k is connected only if a spanning tree of \mathcal{G}_k exists; (c) uses Cayley's formula that there are k^{k-2} ways of choosing a tree among \mathcal{G}_k , and then applies the union bound over them. Now, by plugging $p = \frac{a \log n}{n}$ into (68) we obtain

$$\begin{aligned}
\mathbb{E}[r_k] &\leq \left(\frac{en}{k}\right)^k \cdot k^{k-2} \cdot \left(\frac{a \log n}{n}\right)^{k-1} \cdot e^{-\frac{a \log n}{n} \cdot k(n-k)} \\
&= \left(\frac{e^k}{k^2}\right) \cdot (a \log n)^{k-1} \cdot n^{-ak(1-\frac{k}{n})+1} \tag{69}
\end{aligned}$$

where the first step uses $\binom{n}{k} \leq \left(\frac{en}{k}\right)^k$ and $(1-p) \leq e^{-p}$. To proceed, notice that when $a > 1 + c$, (69) satisfies

$$\mathbb{E}[r_k] \leq n^{-(1+c)k(1-\frac{k}{n})(1-o(1))+1}$$

which holds for any $1 \leq k \leq n/2$. Furthermore, when $k < k_0$ for some large constant k_0 , say $k_0 = 10$, it satisfies

$$\mathbb{E}[r_k] \leq n^{-(1+c)(k-1)(1-o(1))-c+o(1)} \leq n^{-c+o(1)},$$

and when $k \geq 10$ and $k \leq n/2$, we have

$$\mathbb{E}[r_k] \leq n^{-(1+c) \cdot \frac{k(1-o(1))}{2} + 1} \leq n^{-5(1+c)(1-o(1))+1} \leq n^{-c+1}.$$

Given the above, (67) satisfies

$$\begin{aligned}
\mathcal{P}(\mathcal{G}_n \text{ is connected}) &\geq 1 - \sum_{k=1}^9 \mathbb{E}[r_k] - \sum_{k=10}^{\lfloor n/2 \rfloor} \mathbb{E}[r_k] \\
&\geq 1 - 9 \cdot n^{-c+o(1)} - \frac{n}{2} \cdot n^{-c+1} \\
&\geq 1 - n^{-c+o(1)}
\end{aligned}$$

which completes the proof for $a > 1 + c$.

Now, we consider when $a < 1 - c$ for any $c \in (0, 1)$. In this case, we have

$$\begin{aligned}
\mathcal{P}(\mathcal{G}_n \text{ is disconnected}) &\geq \mathcal{P}(r_1 \geq 0) \\
&= 1 - \mathcal{P}(r_1 = 0).
\end{aligned} \tag{70}$$

Furthermore, $\mathcal{P}(r_1 = 0)$ can be upper bounded by

$$\begin{aligned}
\mathcal{P}(r_1 = 0) &= \mathcal{P}(r_1 - \mathbb{E}[r_1] = -\mathbb{E}[r_1]) \leq \mathcal{P}((r_1 - \mathbb{E}[r_1])^2 = \mathbb{E}^2[r_1]) \\
&\leq \mathcal{P}((r_1 - \mathbb{E}[r_1])^2 \geq \mathbb{E}^2[r_1]) \leq \frac{\text{Var}(r_1)}{\mathbb{E}^2[r_1]}
\end{aligned} \tag{71}$$

where the last step comes from Markov's inequality. Our next job is to determine or bound $\mathbb{E}^2[r_1]$ and $\text{Var}(r_1)$. To this end, for each vertex i in \mathcal{G}_n , let us define the binary random variable x_i as

$$x_i = \begin{cases} 1, & i \text{ is isolated} \\ 0, & \text{otherwise} \end{cases}$$

then immediately we have $r_1 = \sum_{i=1}^n x_i$ and

$$\mathbb{E}[r_1] = \sum_{i=1}^n \mathbb{E}[x_i] = n(1-p)^{n-1} = n \cdot e^{-p(n-1)(1-o(1))} = n \cdot n^{a(1-o(1))} = n^{c-o(1)}$$

where we plugged in $p = a \log n/n$ and $a = 1 + c$. Also,

$$\mathbb{E}[r_1^2] = \sum_{i=1}^n \mathbb{E}[x_i^2] + \sum_i \sum_{j \neq i} \mathbb{E}[x_i x_j] = \mathbb{E}[r_1] + n(n-1) \cdot (1-p)^{2(n-2)+1}.$$

Given the above, (71) satisfies

$$\begin{aligned}
\mathbb{P}(r_1 = 0) &\leq \frac{\mathbb{E}[r_1^2] - \mathbb{E}^2[r_1]}{\mathbb{E}^2[r_1]} \leq \frac{\mathbb{E}[r_1^2]}{\mathbb{E}^2[r_1]} - 1 \leq \frac{1}{\mathbb{E}[r_1]} + \frac{n(n-1) \cdot (1-p)^{2(n-2)+1}}{n^2 \cdot (1-p)^{2(n-1)}} - 1 \\
&= n^{-c+o(1)} + (1 - n^{-1}) \cdot (1-p) - 1 = n^{-c+o(1)}.
\end{aligned}$$

Plugging this back into (70) completes the proof.

A.2 Proof of Lemma A.2

Again, given the graph \mathcal{G}_n , we denote r_k as the number of connected component with size k , then we have

$$\begin{aligned}
\mathcal{P}\left(\beta_1 \log n \leq Z_n \leq n - \frac{\beta_2 n}{\log \log n}\right) &= \sum_{\beta_1 \log n \leq k \leq n - \frac{\beta_2 n}{\log \log n}} \mathcal{P}(Z_n = k) \\
&\leq \sum_{\beta_1 \log n \leq k \leq n - \frac{\beta_2 n}{\log \log n}} \mathcal{P}(r_k \geq 1) \\
&\leq \sum_{\beta_1 \log n \leq k \leq n - \frac{\beta_2 n}{\log \log n}} \mathbb{E}[r_k] \tag{72}
\end{aligned}$$

where the last step holds by Markov's inequality. Now, recall the result obtained in (69) that

$$\mathbb{E}[r_k] = \left(\frac{e^k}{k^2}\right) \cdot (a \log n)^{k-1} \cdot n^{-ak(1-\frac{k}{n})+1},$$

by taking the logarithm we obtain

$$\log \mathbb{E}[r_k] \leq -\left[ak\left(1 - \frac{k}{n}\right) - 1\right] \log n + (k-1) \log \log n + O(k). \tag{73}$$

To proceed, on the one hand, consider any k that satisfies $\beta_1 \log n \leq k \leq \beta n$ for some constant $\beta > 0$, then (73) satisfies

$$\begin{aligned}
\log \mathbb{E}[r_k] &\leq -ak(1-\beta) \log n + (k-1) \log \log n + O(k) \\
&\leq -(1-o(1)) \cdot a\beta_1(1-\beta) \log^2 n \\
&\leq -\Theta(\log^2 n)
\end{aligned}$$

which further implies $\mathbb{E}[r_k] \leq n^{-\Theta(\log n)}$. On the other hand, let $\beta n \leq k \leq n - \frac{\beta_2 n}{\log \log n}$, similarly we have

$$\begin{aligned}
\log \mathbb{E}[r_k] &\leq -a \cdot \beta n \cdot \frac{\beta_2}{\log \log n} \cdot \log n + n \log \log n + O(n) \\
&\leq -\Theta\left(\frac{n \log n}{\log \log n}\right)
\end{aligned}$$

which implies $\mathbb{E}[r_k] \leq n^{-\Theta\left(\frac{n}{\log \log n}\right)}$. Finally, plugging the bounds on $\mathbb{E}[r_k]$ into (72) yields

$$\mathcal{P}\left(\beta_1 \log n \leq Z_n \leq n - \frac{\beta_2 n}{\log \log n}\right) \leq n \cdot n^{-\Theta(\log n)} = n^{-\Theta(\log n)}$$

which completes the proof.

A.3 Proof of Lemma A.3

First of all, the technique used for proving Theorem A.2 does not apply in this case, since it can be shown that $\mathbb{E}[r_k] \geq 1$ when k is small and then the upper bound by Markov's inequality (68) becomes trivial. Therefore, we have to upper bound $\mathcal{P}(Z_n \leq \beta \log n)$ more delicately. To this end,

given the graph \mathcal{G}_n , let r_k denotes the number of connected components of size k , and $k_m := \beta \log n$ as the largest size⁵. Then, one can observe the following three constraints on r_k that always hold:

$$\begin{aligned}
(a) : \quad & \sum_{l=1}^{k_m} r_k \geq \left\lceil \frac{n}{k_m} \right\rceil, \\
(b) : \quad & \sum_{k=1}^{k_m} k \cdot r_k = n, \\
(c) : \quad & r_k \geq 0, \quad k = 1, \dots, k_m.
\end{aligned} \tag{74}$$

Next, in order to determine $\mathcal{P}(k \leq \beta \log n)$ by using r_k , for each feasible combination $\mathbf{r} := \{r_k\}_{k=1}^{k_m}$, the following two questions need to be answered:

- Q1: what is the number of arrangements on $\{r_k\}_{k=1}^{k_m}$?
- Q2: what is the probability of each arrangement occurs?

Answer for Q1. Let $m_{1,\mathbf{r}}$ denotes the number of arrangement, then we claim

$$\begin{aligned}
m_{1,\mathbf{r}} &= \binom{n}{r_1, 2r_2, \dots, k_m r_{\max}} \cdot \prod_{k=1}^{k_m} \left[\binom{kr_k}{k, k, \dots, k} \cdot \frac{1}{(r_k)!} \right] \\
&= \frac{n!}{\prod_{k=1}^{k_m} (k!)^{r_k} \cdot (r_k)!}.
\end{aligned} \tag{75}$$

To interpret (75), first notice that

$$\binom{n}{r_1, 2r_2, \dots, k_m r_{\max}}$$

assigns kr_k nodes for those r_k components of size k . Then, for each kr_k nodes, $\binom{kr_k}{k, k, \dots, k}$ further splits them into r_k pieces. However, this overcounts the actual number by introducing the permutation over the r_k pieces, therefore we divide it by $(r_k)!$.

Answer for Q2. Notice that a specific arrangement of $\{r_k\}_{k=1}^{k_m}$ occurs if and only if the following two events happen:

- E1: each component is connected,
- E2: each one is isolated to others.

For E1, let \mathcal{P}_k denotes the probability that a subgraph of size k is connected, then we have

$$\begin{aligned}
\mathcal{P}(\text{E1}) &= \prod_{k=1}^{k_m} (\mathcal{P}_k)^{r_k} \leq \prod_{k=1}^{k_m} (\mathcal{P}(\text{A spanning tree of size } k \text{ exists}))^{r_k} \\
&\leq \prod_{k=1}^{k_m} \left(k^{k-2} \cdot p^{k-1} \right)^{r_k}
\end{aligned}$$

⁵Here, we assume $\beta \log n$ is an integer, otherwise round it to the nearest one.

where the last step comes from the union bound over the total k^{k-2} number of trees as Cayley's formula suggests. For E2, we have

$$\mathcal{P}(\text{E2}) = (1 - p)^{m_{2,\mathbf{r}}}$$

where $m_{2,\mathbf{r}}$ stands for the number of node pairs across different components that satisfies

$$\begin{aligned} m_{2,\mathbf{r}} &\stackrel{(a)}{=} \sum_{k=1}^{k_m} k^2 \cdot \left(\frac{r_k(r_k - 1)}{2} \right) + \frac{1}{2} \sum_{k_1=1}^{k_m} \sum_{k_2=k_1+1}^{k_m} k_1 r_{k_1} \cdot k_2 r_{k_2} \\ &= \frac{1}{2} \sum_{k_1=1}^{k_m} \sum_{k_2=1}^{k_m} k_1 r_{k_1} \cdot k_2 r_{k_2} - \frac{1}{2} \sum_{k=1}^{k_m} r_k \cdot k^2 \\ &\geq \frac{1}{2} \left(\sum_{k=1}^{k_m} k r_k \right)^2 - \frac{k_m}{2} \sum_{k=1}^{k_m} k r_k \\ &\stackrel{(b)}{=} \frac{n(n - k_m)}{2} \end{aligned}$$

where (a) holds by first counting number of pairs that across the r_k components of size k , then adding the pairs that across different k ; (b) applies the constraint $\sum_{k=1}^{k_m} k \cdot r_k = n$ in (74). Given the above, the probability of a specific arrangement \mathbf{r} occurs, denoted by $\mathcal{P}_{\mathbf{r}}$, is given as

$$\begin{aligned} \mathcal{P}_{\mathbf{r}} &= \mathcal{P}(\text{E1} \cap \text{E2}) = \mathcal{P}(\text{E1})\mathcal{P}(\text{E2}) \\ &\leq (1 - p)^{m_{2,\mathbf{r}}} \cdot \prod_{k=1}^{k_m} \left(k^{k-2} \cdot p^{k-1} \right)^{r_k} \end{aligned}$$

where we use the fact that E1 and E2 are independent as they involve two disjoint sets of edges.

Now, we are able to write out $\mathcal{P}(Z_n \leq \beta \log n)$ as

$$\begin{aligned} \mathcal{P}(Z_n \leq \beta \log n) &= \sum_{\mathbf{r}} m_{1,\mathbf{r}} \cdot \mathcal{P}_{\mathbf{r}} \\ &\leq \sum_{\mathbf{r}} \left(\frac{n!}{\prod_{k=1}^{k_m} (k!)^{r_k} \cdot (r_k)!} \right) \cdot (1 - p)^{m_{2,\mathbf{r}}} \cdot \prod_{k=1}^{k_m} \left(k^{k-2} \cdot p^{k-1} \right)^{r_k} \\ &= \sum_{\mathbf{r}} n! (1 - p)^{m_{2,\mathbf{r}}} \prod_{k=1}^{k_m} \left(\frac{k^{k-2} p^{k-1}}{k!} \right)^{r_k} \cdot \frac{1}{(r_k)!} \\ &\leq \sum_{\mathbf{r}} n! (1 - p)^{m_{2,\mathbf{r}}} \prod_{k=1}^{k_m} \frac{(pk)^{r_k (k-1)}}{(r_k)!} \\ &\leq \sum_{\mathbf{r}} n! \cdot e^{-p \cdot m_{2,\mathbf{r}}} \cdot \left(\frac{(pk)^{n - \sum_{k=1}^{k_m} r_k}}{\prod_{k=1}^{k_m} (r_k)!} \right) \end{aligned} \tag{76}$$

where in the last step we applies $(1 - p) \leq e^{-p}$ and the constraint $\sum_{k=1}^{k_m} k \cdot r_k = n$ in (74), and we remark that summation is over all combinations \mathbf{r} that satisfy the three constraints in (74). To proceed, by applying $p = \frac{a \log n}{n}$ and Stirling's approximation that $n! = \sqrt{2\pi n} \left(\frac{n}{e}\right)^n (1 + O(\frac{1}{n}))$,

(76) can be rewritten as

$$\begin{aligned}
\mathcal{P}(k \leq \beta \log n) &\leq \sum_{\mathbf{r}} \sqrt{2\pi n} \left(\frac{n}{e}\right)^n \left(1 + O\left(\frac{1}{n}\right)\right) \cdot n^{-\frac{a(n-k)}{2}} \cdot \left(\frac{ak \log n}{n}\right)^{n - \sum_{k=1}^{k_m} r_k} \cdot \frac{1}{\prod_{k=1}^{k_m} (r_k)!} \\
&\stackrel{(a)}{=} \sum_{\mathbf{r}} \left(\frac{\sqrt{2\pi n}}{e^n}\right) \cdot n^{-\frac{an(1-o(1))}{2}} \cdot (a\beta \log^2 n)^{n - \sum_{k=1}^{k_m} r_k} \cdot \left(\frac{n^{\sum_{k=1}^{k_m} r_k}}{\prod_{k=1}^{k_m} (r_k)!}\right) \\
&\stackrel{(b)}{=} \sum_{\mathbf{r}} n^{-\frac{an(1-o(1))}{2}} \cdot \underbrace{\left(\frac{n^{\sum_{k=1}^{k_m} r_k}}{\prod_{k=1}^{k_m} (r_k)!}\right)}_{=: f(\mathbf{r})} \\
&= \sum_{\mathbf{r}} n^{-\frac{an(1-o(1))}{2}} \cdot f(\mathbf{r})
\end{aligned} \tag{77}$$

where in step (a) we plug in $k_m = \beta \log n$ and (b) holds as $n^{-\frac{an(1-o(1))}{2}}$ is dominant and thus can absorb all other terms but only one denoted as

$$f(\mathbf{r}) := \frac{n^{\sum_{k=1}^{k_m} r_k}}{\prod_{k=1}^{k_m} (r_k)!}. \tag{78}$$

At first glance, it is not clear if $f(\mathbf{r})$ can be also absorbed. To confirm this, we derive the following lemma that sharply bounds $f(\mathbf{r})$:

Lemma A.5. *Given $f(\mathbf{r})$ defined in (78) with $k_m = \Theta(\log n)$, it satisfies*

$$f(\mathbf{r}) \leq n^{o(n)} \tag{79}$$

which holds for any $\mathbf{r} = \{r_k\}_{k=1}^{k_m}$ that satisfies the three constraints in (74).

For ease of presentation, the proof of Lemma A.5 is deferred to Appendix A.4. By plugging (79) into (77) we get

$$\begin{aligned}
\mathcal{P}(k \leq \beta \log n) &\leq \sum_{\mathbf{r}} n^{-\frac{an(1-o(1))}{2}} \cdot n^{o(n)} = \sum_{\mathbf{r}} n^{-\frac{an(1-o(1))}{2}} \\
&\stackrel{(a)}{\leq} (n+1)^{k_m} \cdot n^{-\frac{an(1-o(1))}{2}} \\
&= (n+1)^{\beta \log n} \cdot n^{-\frac{an(1-o(1))}{2}} \\
&= n^{-\frac{an(1-o(1))}{2}}.
\end{aligned}$$

Here, (a) applies the union bound over the set of all possible \mathbf{r} and the size of the set is upper bounded by $(n+1)^{k_m}$, since each r_k takes values from 0 to n . This completes the proof.

A.4 Proof of Lemma A.5

To begin with, we denote

$$m := \sum_{k=1}^{k_m} r_k$$

that represents the number of isolated components. Our proof strategy is to study $f(\mathbf{r})$ under different choices of m . In this way, we can immediately prove Lemma A.5 on two special cases:

1. when $m = n$, this indicates that only $r_1 = n$ is nonzero and all other $r_k = 0$ for $k = 2, \dots, k_m$. Then we have $f(\mathbf{r}) \leq e^n = n^{o(n)}$ which satisfies (79),
2. when $m = o(n)$, then $f(\mathbf{r}) \leq n^m = n^{o(n)}$, which is the desired (79).

Now, we consider the remaining cases of m that satisfy $m < n$ and $m = \Theta(n)$, which excludes the two special cases mentioned above. First, $f(\mathbf{r})$ can be upper bounded as

$$f(\mathbf{r}) = \frac{n^m}{\prod_{k=1}^{k_m} (r_k)!} \stackrel{(a)}{\leq} \frac{n^m}{\prod_{k=1}^{k_m} \frac{r_k^{r_k}}{e^{r_k-1}}} \leq n^m / \underbrace{\prod_{k=1}^{k_m} \left(\frac{r_k}{e}\right)^{r_k}}_{=:g(\mathbf{r})} = \frac{n^m}{g(\mathbf{r})}$$

where step (a) uses the fact that $n! \geq \frac{n^n}{e^{n-1}}$. Then, under a fixed m , we focus on finding the minimum of the defined $g(\mathbf{r})$ under the constraints of \mathbf{r} in (74) and $\sum_{k=1}^{k_m} r_k = m$. To this end, by noticing that

$$\log g(\mathbf{r}) = \sum_{k=1}^{k_m} r_k (\log r_k - 1)$$

we can formulate the following convex optimization program:

$$\begin{aligned} \min_{r_k \in \mathbb{R}_+, k=1, \dots, k_m} \quad & \sum_{k=1}^{k_m} r_k (\log r_k - 1) \\ \text{s.t.} \quad & \sum_{k=1}^{k_m} r_k = m \\ & \sum_{k=1}^{k_m} k r_k = n \\ & r_k \geq 0, \quad k = 1, \dots, k_m \end{aligned} \tag{80}$$

Notably, (80) ignores the constraint that $r_k \in \mathbb{Z}_+$ is an integer and instead extends to real number i.e. $r_k \in \mathbb{R}_+$, which significantly facilitates our analysis. The resulting optima is still a lower bound of $g(\mathbf{r})$ under $r_k \in \mathbb{Z}_+$.

We can solve (80) from its KKT conditions, where the Lagrangian is given as

$$L(\mathbf{r}, \lambda, \nu, \{\mu_k\}_{k=1}^{k_m}) = \sum_{k=1}^{k_m} r_k (\log r_k - 1) + \lambda \left(\sum_{k=1}^{k_m} r_k - m \right) + \nu \left(\sum_{k=1}^{k_m} k r_k - n \right) - \sum_{k=1}^{k_m} \mu_k r_k.$$

Then, by stationarity, the optimum $\mathbf{r}^* = \{r_k^*\}_{k=1}^{k_m}$ must satisfy

$$0 = \left. \frac{\partial L}{\partial r_k} \right|_{r_k=r_k^*} = \log r_k^* + \lambda + \nu k - \mu_k, \quad \Rightarrow \quad r_k^* = e^{\nu k + \lambda - \mu_k}, \quad k = 1, \dots, k_m. \tag{81}$$

Meanwhile, from the complementary slackness we have $\mu_k r_k = 0$ and $\mu_k \geq 0$ for all $k = 1, \dots, k_m$. Combining this fact with (81) yields $\mu_k = 0$ and

$$r_k^* = c \cdot s^k, \quad k = 1, \dots, k \tag{82}$$

for some $c, s > 0$. As a result, the optima is given as

$$\begin{aligned}\log g(\mathbf{r}^*) &= \sum_{k=1}^{k_m} r_k (\log cs^k - 1) = \sum_{k=1}^{k_m} r_k (\log c + k \log s - 1) \\ &= m \log c + n \log s - m\end{aligned}\tag{83}$$

where in the last step we apply the constraints in (80). In order to determine c and s , plugging (82) into the constraints of (80) yields

$$\begin{aligned}m &= \sum_{k=1}^{k_m} r_k^* = \sum_{k=1}^{k_m} c \cdot s^k, \quad \Rightarrow \quad c = \frac{m(1-s)}{s(1-s^{k_m})}, \\ n &= \sum_{k=1}^{k_m} kr_k^* = c \cdot \sum_{k=1}^{k_m} ks^k, \quad \Rightarrow \quad \frac{n}{m} = \frac{1}{1-s} - \frac{k_m s^{k_m}}{1-s^{k_m}}.\end{aligned}\tag{84}$$

However, finding a solution of c and s from (84) is nontrivial, and thus some approximation is necessary. To this end, let us define the following function

$$h(x) := \frac{1}{1-x} - \frac{k_m x^{k_m}}{1-x^{k_m}}\tag{85}$$

and $x = s$ is the solution to $h(x) = \frac{n}{m}$. Before we determine s , several importantly properties of $h(x)$ are listed below:

- P1: $h(0) = 1$, $h(1) = \frac{k_m+1}{2}$, and $\lim_{x \rightarrow \infty} h(x) = k_m$.
- P2: $h(x)$ is monotonically increasing as $x \geq 0$.
- P3: a unique solution $x = s$ to $h(x) = \frac{n}{m}$ exists for any $\frac{n}{k_m} < m < n$.

Here, we provide some justifications for each property one by one. For P1, $h(0)$ and $\lim_{x \rightarrow \infty} h(x)$ can be observed immediately. The remaining $h(1) = \frac{k_m+1}{2}$ is obtained from L'Hôpital's rule. For P2, the derivative of $h(x)$ is given as

$$\begin{aligned}h'(x) &= \frac{1}{(1-x)^2} - \frac{k_m^2 x^{k_m-1}}{(1-x^{k_m})^2} = \frac{(1+x+\dots+x^{k_m-1})^2 - k_m^2 x^{k_m-1}}{(1-x^{k_m})^2} \\ &= \frac{(1+x+\dots+x^{k_m-1} - k_m x^{k_m-1})(1+x+\dots+x^{k_m-1} + k_m x^{k_m-1})}{(1-x^{k_m})^2} \\ &\geq 0\end{aligned}$$

where the last step stems from Jensen's inequality⁶ that $1+x+\dots+x^{k_m-1} \geq k_m x^{k_m-1}$, and the equality only holds when $x = 1$, which implies $h'(x) = 0$ only when $x = 1$. For P3, one can see that $1 < \frac{n}{m} < k_{\max}$ when $\frac{n}{k_m} < m < n$, which agrees with the range of $h(x)$ when $x \geq 0$, therefore the solution to $h(x) = \frac{n}{m}$ always exists and P2 suggests its uniqueness.

Now, we can (approximately) determine the solution $x = s$ and the corresponding optima $g(\mathbf{r}^*)$ in (83) for different choices of m as follows:

⁶By defining the convex function $f_0(k) = x^k$ and a random variable y that uniformly takes values on $0, 1, \dots, k_m - 1$, then Jensen's inequality suggests $\mathbb{E}[f_0(y)] \geq f_0(\mathbb{E}[y])$, which is the desired inequality.

- **Case 1:** When $m = n - n_0$ for any $n_0 = o(n)$. In this case,

$$\frac{n}{m} = 1 + \frac{n_0}{n} + o\left(\frac{n_0}{n}\right).$$

Notice that when $x = o(1)$, it satisfies $h(x) = 1 + x - o(x)$, therefore the solution to $h(x) = \frac{n}{m}$ is given as

$$s = \frac{n_0}{n}(1 + o(1)),$$

plugging this back into (84) gives

$$c = \frac{m(1-s)}{s(1-s^{k_m})} \geq \frac{nm}{n_0} \cdot (1 - o(1)).$$

Therefore, the optima given in (83) satisfies

$$\begin{aligned} \log g(\mathbf{r}^*) &= m \log c + n \log s - m \\ &\geq m \log \left(\frac{nm}{n_0} \cdot (1 - o(1)) \right) + n \log \left(\frac{n_0}{n} \cdot (1 + o(1)) \right) \\ &= n(1 - o(1)) \cdot \log \left(\frac{n^2}{n_0} \cdot (1 - o(1)) \right) - (1 + o(1)) \cdot n \log n \\ &\geq (1 - o(1))n \log n. \end{aligned}$$

This further leads to $g(\mathbf{r}^*) \geq n^{(1-o(1))n}$ and $f(\mathbf{r}^*) \leq n^{o(n)}$, which is the desired result.

- **Case 2:** When $m = n - n_0$ for $n_0 = \Theta(n)$ and $n_0 \leq n - \frac{2n}{k_m+1}$, this indicates that

$$1 < \frac{n}{m} \leq \frac{k_m + 1}{2}.$$

To find the solution s , recall that $h(1) = \frac{k_m+1}{2}$ and $h(x)$ is monotonically increasing, then we get $s \leq 1$. Also, notice that when $s \leq 1$, it satisfies

$$h(s) = \frac{1}{1-s} - \frac{k_m s^{k_m}}{1-s^{k_m}} \leq \frac{1}{1-s},$$

and $h(s) = \frac{n}{m} = \frac{1}{1-\frac{n_0}{n}}$, this further leads to $s \geq \frac{n_0}{n}$. Given the above, we obtain the range of s as

$$s \in \left[\frac{n_0}{n}, 1 \right].$$

Next, to determine c in (84), notice that $c(s) = \frac{m(1-s)}{s(1-s^{k_m})}$ as a function of s is monotonically decreasing when $s > 0$. To see this, again by taking the derivative we have

$$c'(s) = \frac{(k_m + 1)s^{k_m} - k_m s^{k_m+1} - 1}{s^2(1-s^{k_m})^2} \leq 0$$

where the last steps holds by Jensen's inequality that $(k_m + 1)s^{k_m} - k_m s^{k_m+1} - 1 \leq 0$. Therefore, we have

$$c \geq c(s)|_{s=1} = \frac{m(1-s)}{s(1-s^{k_m})} \Big|_{s=1} = \frac{m}{k_m}.$$

Now, we can determine the optima in (83) as

$$\begin{aligned}\log g(\mathbf{r}^*) &= m \log c + n \log s - m \stackrel{(a)}{\geq} m \log \left(\frac{m}{k_m} \right) + n \log \left(\frac{n_0}{n} \right) - m \\ &\stackrel{(b)}{\geq} m \log \left(\frac{2n}{k_m(k_m + 1)} \right) - O(n) \\ &= (1 - o(1)) \cdot m \log n\end{aligned}$$

where (a) comes by plugging $c \geq \frac{m}{k_m}$ and $a \geq \frac{n_0}{n}$; (b) holds by using $m \geq \frac{2}{k_m+1}$ and $\frac{n_0}{n} = \Theta(1)$. In this way, we further obtain $f(\mathbf{r}^*) = \frac{n^m}{g(\mathbf{r}^*)} \leq n^{o(n)}$.

Notably, cases 1 and 2 together cover all the scenarios when $m = \Theta(n)$, and we have shown that $f(\mathbf{r}) = n^{o(n)}$, which completes the proof.

B Proof of lemmas

B.1 Proof of Lemma 5.2

To begin with, recall the subgraph $G(S_{11})$ with size $|S_{11}| = \frac{n}{2} - \Delta = \Theta(n)$ follows Erdős–Rényi model. Then, let z denote the size of the largest connected component of $G_{S_{11}}$, Theorem A.4 suggests a giant component in $G(S_{11})$ exists with high probability:

$$\mathcal{P} \left(z \geq |S_{11}| - \frac{\beta n}{\log \log n} \right) \geq 1 - n^{-\Theta(\log n)} \quad (86)$$

for some constant $\beta > 0$. In this case, let us denote

$$E_{\text{giant}} : z \geq |S_{11}| - \frac{\beta n}{\log \log n}$$

as the event of a giant component and E_{giant}^c as its complement. Then, $\mathcal{P}_{\mathbf{x}^*}(E_{2,S_1} | r_1)$ can be written as

$$\begin{aligned}\mathcal{P}_{\mathbf{x}^*}(E_{2,S_1} | r_1) &= \mathcal{P}_{\mathbf{x}^*}(E_{\text{giant}}^c) \mathcal{P}_{\mathbf{x}^*}(E_{2,S_1} | r_1, E_{\text{giant}}^c) + \mathcal{P}_{\mathbf{x}^*}(E_{\text{giant}}) \mathcal{P}_{\mathbf{x}^*}(E_{2,S_1} | r_1, E_{\text{giant}}) \\ &\leq n^{-\Theta(\log n)} + \min\{1, \mathcal{P}_{\mathbf{x}^*}(E_{2,S_1} | r_1, E_{\text{giant}})\}\end{aligned} \quad (87)$$

where in the last step we apply $\mathcal{P}_{\mathbf{x}^*}(E_{\text{giant}}^c) = 1 - \mathcal{P}_{\mathbf{x}^*}(E_{\text{giant}}) \leq n^{-\Theta(\log n)}$ from (86).

Now, we focus on bounding $\mathcal{P}_{\mathbf{x}^*}(E_{2,S_1} | r_1, E_{\text{giant}})$ with a giant component exists. For each node $i \in S_{12}$, let us denote $r_{1,i}$ as the number of edges that it connects to S_{11} , and further, let $r_{1,\text{giant},i} \leq r_{1,i}$ denote the number of edges that connects to the giant component. Then, under a fixed set of $\{r_{1,i}\}_{i=1}^{\Delta}$ and $\{r_{1,\text{giant},i}\}_{i=1}^{\Delta}$, the event E_{2,S_1} occurs with probability

$$\mathcal{P} \left(E_{2,S_1} | r_1, \{r_{1,i}\}_{i=1}^{\Delta}, \{r_{1,\text{giant},i}\}_{i=1}^{\Delta}, E_{\text{giant}} \right) \leq \prod_{i=1}^{\Delta} M^{1-r_{1,\text{giant},i}}. \quad (88)$$

(88) can be obtained by using the same argument as in the proof of Lemma 5.1. That is, for each number of $r_{1,\text{giant},i}$ edges that connects to the giant component, it creates at least $r_{1,\text{giant},i} - 1$

number of cycles, thus the corresponding probability that the cycle consistency satisfies is upper bounded by $M^{1-r_{1,\text{giant},i}}$. To proceed, we have

$$\begin{aligned} \mathcal{P}_{\mathbf{x}^*} \left(E_{2,S_1} \mid r_1, \{r_{1,i}\}_{i=1}^\Delta, E_{\text{giant}} \right) &\leq \prod_{i=1}^\Delta \mathbb{E} \left[M^{1-r_{1,\text{giant},i}} \mid r_1, \{r_{1,i}\}_{i=1}^\Delta, E_{\text{giant}} \right] \\ &= \prod_{i=1}^\Delta \mathbb{E} \left[M^{1-r_{1,\text{giant},i}} \mid r_{1,i}, E_{\text{giant}} \right] \end{aligned} \quad (89)$$

where the last step holds as $r_{1,\text{giant},i}$ only depends on $r_{1,i}$.

Our next job is to upper bound $\mathbb{E} \left[M^{1-r_{1,\text{giant},i}} \mid r_{1,i}, E_{\text{giant}} \right]$. First, recall that z is denoted as the giant component size. We also define $z' := |S_{11}| - z$ as the size of the residual. Then, under a fixed $r_{1,i}$, the distribution of $r_{1,\text{giant},i}$ is given as

$$\mathcal{P} (r_{1,\text{giant},i} = k \mid r_{1,i}, E_{\text{giant}}) = \frac{\binom{z}{k} \cdot \binom{z'}{r_{1,i}-k}}{\binom{|S_{11}|}{r_{1,i}}}$$

where we use the fact that given $r_{1,i}$, each pair of nodes (i, j) for $j \in S_{11}$ is assigned with an edge with the same probability. Then, we have

$$\mathbb{E} \left[M^{1-r_{1,\text{giant},i}} \mid r_{1,i}, E_{\text{giant}} \right] = \sum_{k=\max\{0, r_{1,i}-z'\}}^{\min\{r_{1,i}, z\}} \mathcal{P} (r_{1,\text{giant},i} = k \mid r_{1,i}, E_{\text{giant}}) \cdot M^{1-k}. \quad (90)$$

It is important to note that the summation over k ranges from $\max\{0, r_{1,i} - z'\}$ to $\min\{r_{1,i}, z\}$ since there is at least $r_{1,i} - z'$ edges connecting to the giant component when $r_{1,i} > z'$, and also $r_{1,i}$ cannot exceed z . Then, in order to calculate (90), we consider two cases depending on the relation between $r_{1,i}$ and z' :

- **Case 1:** When $z'/r_{1,i} = o(1)$, (90) satisfies

$$\begin{aligned} \mathbb{E} \left[M^{1-r_{1,\text{giant},i}} \mid r_{1,i}, E_{\text{giant}} \right] &= \sum_{k=r_{1,i}-z'}^{\min\{r_{1,i}, z\}} \mathcal{P} (r_{1,\text{giant},i} = k \mid r_{1,i}, E_{\text{giant}}) \cdot M^{1-k} \\ &\leq M^{1-(r_{1,i}-z')} = M^{1-(1-o(1))r_{1,i}} \end{aligned}$$

which is a desired upper bound.

- **Case 2:** When $z'/r_{1,i} = \Omega(1)$, let us denote

$$f(k) := \mathcal{P} (r_{1,\text{giant},i} = k \mid r_{1,i}, E_{\text{giant}}) \cdot M^{1-k}.$$

To proceed, we study the ratio $f(k-1)/f(k)$, which is given as

$$\frac{f(k-1)}{f(k)} = \frac{\binom{z}{k-1} \cdot \binom{z'}{r_{1,i}-k+1}}{\binom{z}{k} \cdot \binom{z'}{r_{1,i}-k}} \cdot M = \frac{Mk(z' - r_{1,i} + k)}{(z - k + 1)(r_{1,i} - k + 1)}. \quad (91)$$

Then, we consider $k \leq (1 - \epsilon)r_{1,i}$ for some $\epsilon = o(1)$, plugging this into (91) yields

$$\begin{aligned}
\frac{f(k-1)}{f(k)} &\leq \frac{Mk(z' - \epsilon r_{1,i})}{(z - k + 1)(\epsilon r_{1,i} + 1)} \stackrel{(a)}{=} \frac{Mk \cdot z'(1 - o(1))}{z(1 - o(1)) \cdot (\epsilon r_{1,i} + 1)} \\
&\leq \frac{Mkz'}{z\epsilon r_{1,i}} \cdot (1 + o(1)) = \frac{Mz'(1 - \epsilon)}{z\epsilon} (1 + o(1)) \\
&\leq \frac{Mz'}{z\epsilon} (1 + o(1))
\end{aligned} \tag{92}$$

where (a) comes from the assumption that $z'/r_{1,i} = \Omega(1)$ and therefore $\epsilon r_{1,i} = o(z')$. Now, given that the event E_{giant} occurs, from (86) we have $z \geq |S_{11}| - \frac{\beta n}{\log \log n}$ and $z' \leq \frac{\beta n}{\log \log n}$ for some $\beta > 0$. Then, by taking

$$\epsilon = \Theta\left(\frac{1}{\log \log \log n}\right),$$

(92) satisfies

$$\frac{f(k-1)}{f(k)} = O\left(\frac{\log \log \log n}{\log \log n}\right) = o(1)$$

which holds for any $k \leq (1 - \epsilon)r_{1,i}$. As a result, let $k_0 := (1 - \epsilon)r_{1,i} = (1 - o(1))r_{1,i}$ and assume it to be an integer, the summation in (90) satisfies

$$\begin{aligned}
\mathbb{E} [M^{1-r_{1,\text{giant},i}} \mid r_{1,i}, E_{\text{giant}}] &= \sum_{k=\min\{0, r_{1,i}-1\}}^{r_{1,i}} f(k) = \sum_{k=\min\{0, r_{1,i}-1\}}^{k_0-1} f(k) + \sum_{k=k_0}^{r_{1,i}} f(k) \\
&= o(1) \cdot f(k_0) + \sum_{k=k_0}^{r_{1,i}} \mathcal{P}(r_{1,\text{giant},i} = k \mid r_{1,i}, E_{\text{giant}}) \cdot M^{1-k} \\
&\leq o(1) \cdot \mathcal{P}(r_{1,\text{giant},i} = k_0 \mid r_{1,i}, E_{\text{giant}}) \cdot M^{1-k_0} + M^{1-k_0} \\
&\leq M^{1-(1-o(1))r_{1,i}}
\end{aligned}$$

which is the desired upper bound.

Given the above, (89) can be written as

$$\begin{aligned}
\mathcal{P}_{\mathbf{x}^*} \left(E_{2,S_1} \mid r_1, \{r_{1,i}\}_{i=1}^{\Delta}, E_{\text{giant}} \right) &\leq \prod_{i=1}^{\Delta} M^{1-(1-o(1))r_{1,i}} = M^{\sum_{i=1}^{\Delta} [1-(1-o(1))r_{1,i}]} \\
&= M^{\Delta-(1-o(1))r_1}
\end{aligned} \tag{93}$$

where the last step applies $\sum_{i=1}^{\Delta} r_{1,i} = r_1$. Therefore, since the upper bound in (93) does not depend on the specific set $\{r_{1,i}\}_{i=1}^{\Delta}$, we have

$$\begin{aligned}
\mathcal{P}_{\mathbf{x}^*} (E_{2,S_1} \mid r_1, E_{\text{giant}}) &\leq \mathbb{E} \left[M^{\Delta-(1-o(1))r_1} \mid r_1, E_{\text{giant}} \right] \\
&= M^{\Delta-(1-o(1))r_1}.
\end{aligned}$$

Plugging this back into (87) completes the proof.

B.2 Proof of Lemma 5.3

To begin with, plugging (47) into (39) yields

$$\begin{aligned}
P_\Delta &\leq \min_{\alpha>0} \left(\sum_{r_1} \sum_{r_2} \sum_{r^*} \mathcal{P}_{\mathbf{x}^*}(r_1) \mathcal{P}_{\mathbf{x}^*}(r_2) \mathcal{P}_{\mathbf{x}^*}(r^*) \cdot \left(\frac{Ma}{b}\right)^{\alpha(r_1+r_2-r^*)} \cdot \left(n^{-\Theta(\log n)} + M^{2\Delta-(1-o(1))(r_1+r_2)}\right) \right) \\
&= \min_{\alpha>0} \left(\sum_r \sum_{r^*} \mathcal{P}_{\mathbf{x}^*}(r) \mathcal{P}_{\mathbf{x}^*}(r^*) \cdot \left(\frac{Ma}{b}\right)^{\alpha(r-r^*)} \cdot \left(n^{-\Theta(\log n)} + M^{2\Delta-(1-o(1))r}\right) \right) \\
&= \min_{\alpha>0} \left(n^{-\Theta(\log n)} \cdot \mathbb{E} \left[\left(\frac{Ma}{b}\right)^{\alpha(r-r^*)} \right] + M^{2\Delta} \cdot \mathbb{E} \left[\left(\frac{Ma}{b}\right)^{\alpha(r-r^*)} \cdot M^{-(1-o(1))r} \right] \right). \tag{94}
\end{aligned}$$

Recall that r and r^* defined in (30) are independent and follow binomial distributions as

$$r \sim \text{Binom} \left(2\Delta \left(\frac{n}{2} - \Delta \right), q \right), \quad r^* \sim \text{Binom} \left(2\Delta \left(\frac{n}{2} - \Delta \right), p \right).$$

Then we apply the following standard result:

Lemma B.1. *Given a binomial random variable $r \sim \text{Binom}(N, p)$, it satisfies*

$$\mathbb{E}[t^r] \leq e^{-pN(1-t)}$$

for any constant $t > 0$.

Proof. By definition,

$$\begin{aligned}
\mathbb{E}[t^r] &= \sum_{k=0}^N \binom{N}{k} p^k (1-p)^{N-k} \cdot t^k = \sum_{k=0}^N \binom{N}{k} (pt)^k (1-p)^{N-k} \\
&= (1-p+pt)^N = (1-(1-t)p)^N \\
&\leq e^{-(1-t)pN}
\end{aligned}$$

where the last step uses the $1-x \leq e^{-x}$ for any $x \in \mathbb{R}$. □

As a result, by denoting $N_\Delta := 2\Delta \left(\frac{n}{2} - \Delta \right)$, the expectations in (94) satisfy

$$\begin{aligned}
\mathbb{E} \left[\left(\frac{Ma}{b}\right)^{\alpha(r-r^*)} \right] &= \mathbb{E} \left[\left(\frac{Ma}{b}\right)^{\alpha r} \right] \cdot \mathbb{E} \left[\left(\frac{Ma}{b}\right)^{-\alpha r^*} \right] \\
&\leq \exp \left(-qN_\Delta \left(1 - \left(\frac{Ma}{b}\right)^\alpha \right) \right) \cdot \exp \left(-pN_\Delta \left(1 - \left(\frac{Ma}{b}\right)^{-\alpha} \right) \right) \\
&= \exp \left(-\frac{N_\Delta \log n}{n} \left(a + b - a \left(\frac{Ma}{b}\right)^{-\alpha} - b \left(\frac{Ma}{b}\right)^\alpha \right) \right) \\
&= \exp \left(-\left(\frac{N_\Delta \log n}{n} \right) f_1(\alpha) \right) \tag{95}
\end{aligned}$$

where we use the fact $p = \frac{a \log n}{n}$, $q = \frac{b \log n}{n}$ and define

$$f_1(\alpha) := a + b - a \left(\frac{Ma}{b}\right)^{-\alpha} - b \left(\frac{Ma}{b}\right)^\alpha.$$

Similarly,

$$\begin{aligned}
\mathbb{E} \left[\left(\frac{Ma}{b} \right)^{\alpha(r-r^*)} \cdot M^{-(1-o(1))r} \right] &= \mathbb{E} \left[\left(\frac{Ma}{b} \right)^{\alpha r} \cdot M^{-(1-o(1))r} \right] \cdot \mathbb{E} \left[\left(\frac{Ma}{b} \right)^{-\alpha r^*} \right] \\
&\leq \exp \left(-qN_\Delta \left(1 - \left(\frac{Ma}{b} \right)^\alpha \cdot M^{-(1-o(1))} \right) \right) \cdot \exp \left(-pN_\Delta \left(1 - \left(\frac{Ma}{b} \right)^{-\alpha} \right) \right) \\
&= \exp \left(-\frac{N_\Delta \log n}{n} \left(a + b - a \left(\frac{Ma}{b} \right)^{-\alpha} - b \left(\frac{Ma}{b} \right)^\alpha \cdot M^{-(1-o(1))} \right) \right) \\
&= \exp \left(-\left(\frac{N_\Delta \log n}{n} \right) f_2(\alpha) \right) \tag{96}
\end{aligned}$$

where we define

$$f_2(\alpha) := a + b - a \left(\frac{Ma}{b} \right)^{-\alpha} - b \left(\frac{Ma}{b} \right)^\alpha \cdot M^{-(1-o(1))}.$$

Notably, since α can be chosen arbitrarily, $f_2(\alpha)$ satisfies

$$f_2(\alpha) \leq a + b - 2\sqrt{\frac{ab}{M^{1-o(1)}}} = a + b - (2 + o(1))\sqrt{\frac{ab}{M}} = O(1)$$

where the equality holds when $\alpha = \alpha_0$ such that

$$a \left(\frac{Ma}{b} \right)^{-\alpha_0} = b \left(\frac{Ma}{b} \right)^{\alpha_0} \cdot M^{-(1-o(1))} \quad \Rightarrow \quad \left(\frac{Ma}{b} \right)^{\alpha_0} = \sqrt{\frac{aM^{1-o(1)}}{b}}.$$

Then, plugging $\alpha = \alpha_0$ into $f_1(\alpha)$ yields

$$f_1(\alpha_0) = -a - b + \sqrt{\frac{ab}{M^{1-o(1)}}} + \sqrt{abM^{1-o(1)}} = O(1).$$

Now, plugging (95) and (96) into (94) and letting $\alpha = \alpha_0$ yields

$$\begin{aligned}
\mathcal{P}_\Delta &\leq n^{-\Theta(\log n)} \cdot \exp \left(-\left(\frac{N_\Delta \log n}{n} \right) f_1(\alpha_0) \right) + M^{2\Delta} \cdot \exp \left(-\left(\frac{N_\Delta \log n}{n} \right) f_2(\alpha_0) \right) \\
&= n^{-\Theta(\log n)} \cdot \exp \left(-\left(\frac{N_\Delta \log n}{n} \right) \cdot O(1) \right) + M^{2\Delta} \cdot \exp \left(-\left(\frac{N_\Delta \log n}{n} \right) \cdot f_2(\alpha_0) \right) \\
&= (1 + o(1)) \cdot M^{2\Delta} \cdot \exp \left(-\left(\frac{N_\Delta \log n}{n} \right) f_2(\alpha_0) \right) \\
&= M^{2\Delta} \cdot \exp \left(-\frac{N_\Delta \log n}{n} \left(a + b - (2 + o(1))\sqrt{\frac{ab}{M}} \right) \right)
\end{aligned}$$

which completes the proof.

B.3 Proof of Lemma 5.6

We start from (55), and recall that r_1 and r_2 are two binomial distributions that follow (54). Here, we further denote another binomial random variable r'_1 that facilitates our analysis

$$r'_1 \sim \text{Binom} \left(\frac{n}{2}, p \right)$$

which slightly differs from $r_1 \sim \text{Binom}\left(\frac{n}{2} - \frac{n}{\gamma}, p\right)$. Then, (55) satisfies

$$\begin{aligned}
\mathcal{P}\left(E_{H,A}^{(i)} \cap E_B^{(i,0)}\right) &= \sum_{r_1} \sum_{r_2} \mathcal{P}(r_1) \mathcal{P}(r_2) \cdot \mathbb{1}(r_2 \geq r_1 + \delta) \cdot M^{-r_2} \\
&= \sum_{r_2} \mathcal{P}(r_2) \cdot \mathcal{P}(r_1 \leq r_2 - \delta) \cdot M^{-r_2} \\
&\stackrel{(a)}{\geq} \sum_{r_2} \mathcal{P}(r_2) \cdot \mathcal{P}(r'_1 \leq r_2 - \delta) \cdot M^{-r_2} \\
&\geq \sum_{r_2=\delta}^{n/2} \mathcal{P}(r_2) \cdot \mathcal{P}(r'_1 = r_2 - \delta) \cdot M^{-r_2}
\end{aligned} \tag{97}$$

where (a) comes from the fact that $\mathcal{P}(r_1 \leq x) \geq \mathcal{P}(r'_1 \leq x)$ for any x as r' has more trials than r . To proceed, by plugging in the binomial distribution of r'_1 and r_2 , (97) is given as

$$\begin{aligned}
\mathcal{P}\left(E_{H,A}^{(i)} \cap E_B^{(i,0)}\right) &\geq \sum_{k=\delta}^{n/2} \mathcal{P}(r_2 = k) \cdot \mathcal{P}(r'_1 = k - \delta) \cdot M^{-k} \\
&= \sum_{k=\delta}^{n/2} \binom{n/2}{k} \left(\frac{q}{M}\right)^k (1-q)^{\frac{n}{2}-k} \cdot \binom{n/2}{k-\delta} p^{k-\delta} (1-p)^{\frac{n}{2}-k+\delta} \\
&= \sum_{k=0}^{n/2-\delta} \binom{n/2}{k} \binom{n/2}{k+\delta} p^k (1-p)^{\frac{n}{2}-k} \left(\frac{q}{M}\right)^{k+\delta} (1-q)^{\frac{n}{2}-k-\delta}.
\end{aligned} \tag{98}$$

Now, in order to calculate (98), let us define

$$\begin{aligned}
f(k) &:= \binom{n/2}{k} \binom{n/2}{k+\delta} \left(\frac{pq}{M(1-p)(1-q)}\right)^k \cdot \underbrace{[(1-p)(1-q)]^{\frac{n}{2}} \cdot \left[\frac{q}{M(1-q)}\right]^\delta}_{=:C} \\
&= C \binom{n/2}{k} \binom{n/2}{k+\delta} \left(\frac{pq}{M(1-p)(1-q)}\right)^k.
\end{aligned} \tag{99}$$

Then, (98) satisfies

$$\mathcal{P}\left(E_{H,A}^{(i)} \cap E_B^{(i,0)}\right) = \sum_{k=0}^{n/2-\delta} f(k) \geq \max_{0 \leq k \leq \frac{n}{2}-\delta} f(k). \tag{100}$$

Our next job is to determine $\max_k f(k)$. To this end, we first show the following result:

Lemma B.2. *Given $f(k)$ defined in (99), it satisfies*

$$k^* = \arg \max_{0 \leq k \leq \frac{n}{2}-\delta} f(k) = \Theta(\log n)$$

Proof. We consider the ratio $f(k+1)/f(k)$, which is given as

$$\begin{aligned} \frac{f(k+1)}{f(k)} &= \frac{\binom{n/2}{k+1}}{\binom{n/2}{k}} \cdot \frac{\binom{n/2}{k+\delta+1}}{\binom{n/2}{k+\delta}} \cdot \frac{pq}{M(1-p)(1-q)} \\ &= \frac{\left(\frac{n}{2}-k\right)\left(\frac{n}{2}-k-\delta\right)}{(k+1)(k+\delta+1)} \cdot \left(\frac{ab \cdot \log^2 n}{Mn^2}\right) (1+o(1)) \\ &= \frac{\left(\frac{1}{2}-\frac{k}{n}\right)\left(\frac{1}{2}-\frac{k-\delta}{n}\right)}{(k+1)(k+\delta+1)} \cdot \left(\frac{ab \cdot \log^2 n}{M}\right) (1+o(1)). \end{aligned} \quad (101)$$

where we plugged in $p = \frac{a \log n}{n}$ and $q = \frac{b \log n}{n}$. Then, when $k = \omega(\log n)$, (101) satisfies

$$\frac{f(k+1)}{f(k)} \leq \left(\frac{\log^2 n}{k^2}\right) \cdot \left(\frac{ab}{4M}\right) \cdot (1+o(1)) = o(1)$$

which indicates $k^* = O(\log n)$. Also, when $k = o(\log n)$, (101) satisfies

$$\frac{f(k+1)}{f(k)} \geq \frac{\log^2 n}{(k+\delta+1)^2} \cdot \left(\frac{ab}{4M}\right) \cdot (1+o(1)) = \omega(1)$$

which indicates $k^* = \Omega(\log n)$ and completes the proof. \square

Lemma B.2 suggests us to focus on $f(k)$ with $k = \Theta(\log n)$. To proceed, by taking the logarithm of $f(k)$ we get

$$\log f(k) = \log \binom{n/2}{k} + \log \binom{n/2}{k+\delta} + k \log \left(\frac{pq}{M(1-p)(1-q)}\right) + \log C. \quad (102)$$

Next, we consider each term separately. First, by using the binomial coefficient bound [31, Lemma 23.1] that $\binom{n}{k} \geq \frac{n^k}{k!} \left(1 - \frac{k(k-1)}{2n}\right)$, we have

$$\begin{aligned} \log \binom{n/2}{k} &\geq k \log \left(\frac{n}{2}\right) - \log k! + \log \left(1 - \frac{k(k-1)}{2n}\right) \\ &\stackrel{(a)}{\geq} k \log \left(\frac{n}{2}\right) - (k+1) \log k + (k-1) - \frac{k(k-1)}{2n} \\ &= k \left[\log \left(\frac{n}{2}\right) - \log \left(\frac{k}{e}\right) - o(1) \right] \end{aligned} \quad (103)$$

where (a) applies the factorial bound that $\log k! \leq (k+1) \log k - (k-1)$ and $\log(1+x) \leq x, \forall x > -1$. In a similar manner,

$$\log \binom{n/2}{k+\delta} \geq (k+\delta) \left[\log \left(\frac{n}{2}\right) - \log \left(\frac{k+\delta}{e}\right) - o(1) \right]. \quad (104)$$

Also, we have

$$\begin{aligned} k \log \left(\frac{pq}{M(1-p)(1-q)}\right) &= k \log \left(\frac{ab \cdot \log^2 n}{Mn^2}\right) + o(k) \\ &= k \left(\log \left(\frac{ab}{M}\right) + 2 \log \log n - 2 \log n + o(1) \right), \end{aligned} \quad (105)$$

and

$$\begin{aligned}
\log C &= \binom{n}{2} \cdot \log[(1-p)(1-q)] + \delta \cdot \log\left(\frac{q}{M(1-q)}\right) \\
&\geq -\binom{n}{2} \cdot \frac{(a+b)\log n}{n} + \delta(\log \log n - \log n) - o(\log n) \\
&= \left(-\frac{a+b}{2} + 1\right) \log n - \delta \log n - o(\log n).
\end{aligned} \tag{106}$$

Now, assembling (103) - (106) and plugging them into (102) yields

$$\begin{aligned}
\log f(k) &\geq \underbrace{-k \log\left(\frac{2k}{e}\right) - (k+\delta) \log\left(\frac{2(k+\delta)}{e}\right) + k \left[\log\left(\frac{ab}{M}\right) + 2 \log \log n\right]}_{=:h(k)} \\
&\quad + \left(-\frac{a+b}{2} + 1\right) \log n - o(\log n) \\
&= h(k) + \left(-\frac{a+b}{2} + 1\right) \log n - o(\log n)
\end{aligned} \tag{107}$$

Then, we can estimate k^* by finding the maximum of $h(k)$. Also, since $h(k)$ is concave, this can be done by solving $h'(k) = 0$, which gives us

$$k_0 = \arg \max_k h(k) = (1 + o(1)) \left(\sqrt{\frac{ab}{4M}}\right) \log n.$$

Now, by plugging $k = k_0$ into (107) we obtain

$$\begin{aligned}
\max_k \log f(k) &\geq h(k_0) + \left(-\frac{a+b}{2} + 1\right) \log n - o(\log n) \\
&= \left(-\frac{a+b}{2} + \sqrt{\frac{ab}{M}}\right) \log n - o(\log n).
\end{aligned}$$

Plugging this back into (100) completes the proof.

B.4 Proof of Lemma 5.7

The proof is very similar to the one of Lemma 5.6 given in Appendix B.3, therefore we will skip those repeated analyses. To begin with, (58) satisfies

$$\begin{aligned}
\mathcal{P}\left(E_{H,A}^{(i)} \cap E_B^{(i,0)}\right) &= \sum_{r_2} \mathcal{P}(r_2) \mathcal{P}(r_1 \geq r_2 + 1) M^{-r_2} \geq \sum_{r_2} \mathcal{P}(r_2) \mathcal{P}(r_1 = r_2 + 1) M^{-r_2} \\
&= \sum_{k=0}^{\frac{n}{2} - \frac{n}{r} - 1} \binom{n/2}{k} q^k (1-p)^{\frac{n}{2}-k} \cdot \binom{\frac{n}{2} - \frac{n}{r}}{k+1} p^{k+1} (1-p)^{\frac{n}{2} - \frac{n}{r} - k - 1} \cdot M^{-k} \\
&= \sum_{k=0}^{\frac{n}{2} - \frac{n}{r} - 1} \binom{n/2}{k} \cdot \binom{\frac{n}{2} - \frac{n}{r}}{k+1} \cdot \left(\frac{pq}{M(1-p)(1_q)}\right)^k \cdot \underbrace{[(1-p)(1-q)]^{\frac{n}{2}} \cdot \frac{p}{(1-p)^{\frac{n}{r}+1}}}_{=:C} \\
&=: \sum_{k=0}^{\frac{n}{2} - \frac{n}{r} - 1} f(k).
\end{aligned}$$

To proceed, we take the logarithm over $f(k)$,

$$\log f(k) = \log \binom{n/2}{k} + \log \binom{\frac{n}{2} - \frac{n}{r}}{k+1} + k \log \left(\frac{pq}{M(1-p)(1-q)} \right) + \log C.$$

By considering $k = \Theta(\log n)$ and following the similar analysis in Appendix B.3, we obtain

$$\log \binom{\frac{n}{2} - \frac{n}{r}}{k+1} \geq (k+1) \left[\log \left(\frac{n}{2} \right) - \log \left(\frac{k}{e} \right) \right] - o(\log n)$$

and

$$\log C = \left(-\frac{a+b}{2} + 1 \right) \log n - o(\log n).$$

The other two terms are identical to (103) and (105). Given the above, we have

$$\begin{aligned} \log f(k) &\geq k \underbrace{\left[2 \log \log n + \log \left(\frac{ab}{M} \right) - 2k \log \left(\frac{k}{e} \right) - 2 \log 2 \right]}_{=:h(k)} - \left(\frac{a+b}{2} \right) \log n - o(\log n) \\ &\geq \left(-\frac{a+b}{2} + \sqrt{\frac{ab}{M}} \right) \log n - o(\log n) \end{aligned}$$

where the last step comes by finding the minimum of $h(k)$. This completes the proof.