Systematic discrepancies between reference methods for non-covalent interactions within the S66 dataset

Benjamin X. Shi,^{1, a)} Flaviano Della Pia,^{1, a)} Yasmine S. Al-Hamdani,^{2, 3, 4} Angelos Michaelides,¹ Dario Alfè,^{2, 5, 3, 4} and Andrea Zen^{2, 4, b)}

¹⁾Yusuf Hamied Department of Chemistry, University of Cambridge, Lensfield Road,

Cambridge CB2 1EW, United Kingdom

²⁾Dipartimento di Fisica Ettore Pancini, Universita di Napoli Federico II,

Monte Sant'Angelo, I-80126 Napoli, Italy

³⁾Thomas Young Centre, University College London, London WC1E 6BT,

United Kingdom

⁴⁾Department of Earth Sciences, University College London, London WC1E 6BT,

United Kingdom

⁵⁾London Centre for Nanotechnology, University College London, London WC1E 6BT, United Kingdom

(Dated: 24 December 2024)

The accurate treatment of non-covalent interactions is necessary to model a wide range of applications, from molecular crystals to surface catalysts to aqueous solutions and many more. Quantum diffusion Monte Carlo (DMC) and coupled cluster theory with single, double and perturbative triple excitations [CCSD(T)] are considered two widely-trusted methods for treating non-covalent interactions. However, while they have been well-validated for small molecules, recent work has indicated that these two methods can disagree by more than 7.5 kcal/mol for larger systems. The origin of this discrepancy remains unknown. Moreover, the lack of systematic comparisons, particularly for medium-sized complexes, has made it difficult to identify which systems may be prone to such disagreements and the potential scale of these differences. In this work, we leverage the latest developments in DMC to compute interaction energies for the entire S66 dataset, containing 66 medium-sized complexes with a balanced representation of dispersion and electrostatic interactions. Comparison to previous CCSD(T) references reveals systematic trends, with DMC predicting stronger binding than CCSD(T) for electrostatic-dominated systems, while the binding becomes weaker for dispersion-dominated systems. We show that the relative strength of this discrepancy is correlated to the ratio of electrostatic and dispersion interactions, as obtained from energy decomposition analysis methods. Finally, we pinpoint systems in the S66 dataset where these discrepancies are particularly prominent, offering cost-effective benchmarks to guide future developments in DMC, CCSD(T) as well as the wider electronic structure theory community.

^{a)}These authors contributed equally to this work.

^{b)}Electronic mail: andrea.zen@unina.it

I. INTRODUCTION

Non-covalent interactions play a crucial role in many areas of science. These interactions govern the structure of molecular crystals¹ (e.g., in pharmaceutical drugs), biomolecules² like DNA and proteins and are relevant to supramolecular³ science and nanotechnology.^{4,5} They also underlie important processes across chemistry and biology, from protein-ligand binding⁶, to catalytic reactions, both on the surface⁷ and in solution.³ Understanding and unlocking new processes for these applications will increasingly rely on accurate computational modeling tools that can treat non-covalent interactions.⁸

Two methods of choice for modeling non-covalent interactions are quantum diffusion Monte Carlo⁹ (DMC) and coupled cluster theory¹⁰ with single, double and perturbative triple excitations [CCSD(T)]. While these methods may not be as affordable as density functional theory¹¹ (DFT), the reference data they provide are pivotal for benchmarking and parametrizing the density functional approximations (DFAs) necessary for practical routine simulations. For example, the local density approximation (LDA) and many extensions build upon a DMC-based parametrization of the correlation energy,¹² while CCSD(T) interaction energy datasets have helped aid in the development of many modern dispersion corrections.^{13–16} In particular, the applicability of these methods to larger systems have rapidly expanded in recent years, arising from computer hardware improvements and, more importantly, algorithmic/methodological developments to both DMC^{17–26} and CCSD(T).^{27–34}

DMC and CCSD(T) solve the Schrödinger equation to model the systems with distinct approaches and corresponding approximations. Despite these differences, there are many examples where DMC and CCSD(T) have come into alignment. For example, besides small molecules,^{35–37} agreement has been obtained for graphene bilayer binding energies³⁸, molecular crystal lattice energies,^{39–41} molecule-surface interactions^{42–47} and vacancy formation energies.⁴⁸ Recently, this agreement has been shown to start to falter^{49–52} for large dispersion-bound molecules, with differences as large as 7.5 kcal/mol for a buckyball-ring (C₆₀@[6]CPPA) complex.

The origin of the discrepancy between DMC and CCSD(T) for large dispersion-bound molecules is a topic of current debate,^{50,53–55} particularly on the validity of the perturbative triples (T) contribution in CCSD(T). Schäfer *et al.*⁵⁰ have suggested that part of this discrepancy arises from missing contributions in (T) that can be accounted by the (cT) approach. In addition, Semidalas *et al.*⁵⁶ have reported non-trivial discrepancies between CCSD(T) and post-CCSD(T) methods such as CCSDT(Q). Conversely, Lambie *et al.*⁵⁴ have found that CCSD(T) does not differ significantly against CCSDT(Q) using the Pariser-Parr-Pople (PPP) model^{57–59} for large conjugated systems. Similarly, Fishman *et al.*⁵³ and Lao⁵⁵ report only a slight overbinding of CCSD(T) against CCSDT(Q) that cannot explain the discrepancy against DMC.

Understanding these discrepancies between DMC and CCSD(T) for large molecules requires cross-validating these methods across systematic datasets, particularly those involving medium to large sized molecules^{51,60,61} which sample a range of non-covalent interactions. While DMC and CCSD(T) have both been compared (to great agreement) for the A24⁶² and S22⁶³ datasets of small molecular complexes, DMC has not been frequently applied to study medium-sized datasets. In particular, it has not been used to study the S66 dataset,⁶⁴ a compilation of 66 dimers that probes the two major types of non-covalent interactions: dispersion and hydrogen-bonding together with those of mixed character. As well as covering a range of interactions, many of the molecules considered form the building blocks for larger biomolecules along different binding configurations. Furthermore, the parallel-displaced benzene dimer⁶⁵ is included in this set of complexes, making it an interesting modeling challenge. Such a dataset has been pivotal towards benchmarking^{66–74} DFAs in DFT as well as lower-level approximations to wave-function methods^{56,75–80} and even machine-learning models.^{81,82}

In this work, we leverage the latest developments in DMC to compute interaction energies for the entire S66 dataset. When compared to CCSD(T) estimates (taken from the literature), we reveal a consistent weaker binding of dispersion interactions and consistent stronger binding of electrostatic interactions in DMC. In particular, we show that their differences are correlated to the ratio of electrostatic and dispersive interactions within the system. The discrepancies in dispersion-dominated systems are shown to be reduced when utilizing an (empirically fitted) CCSD(cT) formulation,⁵⁰ although notable differences remain. We identify specific systems with well-defined differences between DMC and CCSD(T) that can serve as model systems for testing future developments in both methods, setting the stage towards resolving their discrepancies.

II. METHODS

A. Diffusion Monte Carlo

The DMC interaction energies of the $S66^{64}$ dataset are computed as:

$$\Delta E_{\rm int.} = E_{\rm dimer} - E_{\rm mon.\ 1} - E_{\rm mon.\ 2},\tag{1}$$

where E_{dimer} is the total energy of the dimer, and $E_{\text{mon. 1}}$, $E_{\text{mon. 2}}$ are the total energies of the constituent monomers. In the S66 dataset, these monomers are kept fixed to their geometry in the dimer, which is in general different from their equilibrium geometry. In this work, we first computed the energies of the monomers with DMC at a chosen reference geometry. Subsequently, we added the deformation energy, i.e. the energy difference between the geometry of the monomer in the dimer and against this reference geometry using CCSD(T). We provide further details on these calculations in Sec. S2.1 of the supplementary material, and show for a subset of the S66 complexes that differences between DMC and CCSD(T) predictions of the deformation energies are within ~ 0.12 kcal/mol.

A detailed description of the DMC method can be found in Ref. 9. In this work, we compute fixed-node DMC interaction energies by using the CASINO code²³. We use energy-consistent correlated electron pseudopotentials⁸³ (eCEPP) with the determinant locality approximation (DLA)²². The trial wave-functions were of the Slater–Jastrow type with single Slater determinants, and the single-particle orbitals obtained from DFT local-density approximation (LDA) plane-wave calculations performed with PWscf^{84,85} using an energy cut-off of 600 Ry and reexpanded in terms of B-splines⁸⁶. The Jastrow factor included a two-body electron–electron (e–e) term, two-body electron-nucleus (e-n) terms, and three-body electron-electron-nucleus (e-e-n) terms. The variational parameters of the Jastrow have been optimized by minimizing the variance of each system. The final DMC estimates of $\Delta E_{int.}$ were extrapolated towards the zero time step limit $(\tau \rightarrow 0)$ by making a cubic fit to a series of time step estimates from 0.11 au down to 0.003 au. We have estimated errors on this fit with two approaches: either by taking the 1σ standard deviation of the fit at zero time step or the difference w.r.t. the zero time step limit given by a linear fit from the two smallest (0.01 au and 0.003 au) time steps; the latter accounts for systems where the time step behavior may change close to the zero time step limit. The final error estimate is taken to be the larger of the two, as discussed in Sec. S2.4 of the supplementary material.

The parameters chosen within the present work follow from previous DMC calculations for

large molecules in Ref. 49 as well as molecular crystals in Refs. 40 and 41. Within these studies of non-covalent interactions, the LDA trial wave-function was shown to be valid, either by comparison to experiments or when using trial wave-functions with other DFAs. For the case of the AcOH dimer system (ID 20), we have performed our own validation tests on the choice of trial wave-function as well as localization approximation in Sec. S7 of the supplementary material.

B. Coupled Cluster Theory

Several CCSD(T) estimates^{56,64,75,77,79,87–91} of the S66 interaction energies are available in the literature. Here, we compare DMC to the average of three recent CCSD(T) calculations^{75,88,91}: the revised calculations from Řezáč *et al.*⁷⁵; the "SILVER" estimates from Kesharwani *et al.*⁸⁸; and the "14k-GOLD" estimates from Nagy *et al.*⁹¹. A brief description of the three different CCSD(T) calculations is reported in Sec. S3 of the supplementary material.⁹²

Schäfer *et al.*⁵⁰ have recently demonstrated that there exists an empirical relationship between the (cT) and the (T) correlation contributions to the total energy using the CCSD and MP2 correlation energies. The resulting (cT)-fit is of the form:

$$\frac{(\mathrm{T})}{(\mathrm{cT})-\mathrm{fit}} = a + b \cdot \frac{\mathrm{MP2 \ corr.}}{\mathrm{CCSD \ corr.}},$$
(2)

where *a* and *b* were parameters fitted from comparing CCSD(cT) to CCSD(T) calculations for a set of dispersion-bound complexes. We have recomputed the CCSD(cT)-fit values from the original CCSD(T) "SILVER" estimates from Kesharwani *et al.*, adding the difference between (cT) and (T) to the final (averaged) CCSD(T) estimates, as given in Sec. S4 of the supplementary material.

III. RESULTS AND DISCUSSION

This dataset comprises of a diverse range of interactions and we have separated the systems according to the original S66 categories of hydrogen-bonded, dispersion-bonded and "mixed"-character systems in Figs. 1, 2 and 3, respectively. The corresponding dimer complexes are visualized in Fig. S1 of the supplementary material. We report the DMC estimate of the interaction energy $\Delta E_{int.}$ above the label of each S66 complex. In all cases, the rrors on $\Delta E_{int.}$ estimates are below 0.26 kcal/mol, with the majority below 0.10 kcal/mol, facilitating reliable comparisons to other CCSD(T).



FIG. 1. Comparison between DMC interaction energies $\Delta E_{int.}$ calculated in the present work against CCSD(T) for a subset of systems in the S66 dataset with hydrogen-bonds. The CCSD(T) estimate is taken as an average from three previous calculations^{75,88,91}, with corresponding standard deviation as error. The deviation of CCSD(T) from the DMC is plotted with grey crosses, with the statistical errors (corresponding to one standard deviation σ). The complex ID and label are provided below the x-axis, while the number above each x-axis tick represents the DMC $\Delta E_{int.}$ estimate, with the error on the last reported digit given in parentheses. The uracil dimer (ID 17) is in its base-pair (BP) configuration.

The strength of $\Delta E_{\text{int.}}$ varies significantly across the systems, from as large as $-20.17 \pm 0.11 \text{ kcal/mol}$ for complex 20 (acetic acid dimer) to as weak as $-1.11 \pm 0.06 \text{ kcal/mol}$ for complex 30 (benzene-ethene dimer), being stronger in the H-bonded systems. With gray crosses, we plot the difference between DMC and CCSD(T) estimates (as described in the Methods) for the three classes of interactions. We use DMC as the reference (i.e., zero), and plot blue vertical bars along the horizontal zero-axis representing the errors on the DMC estimates. There is overall excellent agreement, with a mean absolute deviation (MAD) of 0.21 kcal/mol across the entire S66 dataset. We find systematic trends in the differences between CCSD(T) and DMC, with CCSD(T) predicting weaker binding compared to DMC for hydrogen-bonded systems in Fig. 1, with an MAD of $\sim 0.24 \text{ kcal/mol}$, while predicting a stronger binding for dispersion dominated systems in Fig. 2, with an MAD of $\sim 0.24 \text{ kcal/mol}$. For the "mixed" character systems in Fig. 3, the MAD is lower

at 0.14 kcal/mol.

	1.2 - 0.6 -	××	CCS CCS	5D(T) 5D(c1	[MA [)-fit	D: 0. [MAI	24] D: 0.()9]		1															
cal/mol]	0.0 -] ×	×	×	- 1 - ×	-] - ×	×	×	-¥-	- X -	Ť	ž	-¥-	¥	¥	×	÷	X	÷	-Ŧ-	- 1	×	¥	- <u>+</u> - ×	
Difference against DMC [kc	-0.6 -	Benzene…Benzene (π-π) - 242.33(8)	PyridinePyridine (π-π) - 253.53(7)	Uracil…Uracil (π - π) - 269.33(8)	BenzenePyridine (π - π) - 273.04(8)	Benzene…Uracil (π-π) - 285.15(11)	Pyridine…Uracil (π-π) - 296.40(17)	Benzene…Ethene - 30 -1.11(6)	UracilEthene - 313.18(9)	UracilEthyne - 323.59(8)	Pyridine…Ethene - 33 -1.69(7)	Pentane…Pentane - 343.53(7)	Neopentane…Pentane - 352.46(8)	Neopentane…Neopentane - 361.67(8)	Cyclopentane…Neopentane - 372.17(7)	Cyclopentane…Cyclopentane - 382.75(7)	Benzene…Cyclopentane - 39 – -3.14(8)	Benzene…Neopentane - 40 -2.68(8)	UracilPentane - 414.46(9)	Uracil…Cyclopentane - 423.59(9) 🗙	Uracil…Neopentane - 43 – -3.51(16)	EthenePentane - 441.79(6)	Ethyne…Pentane - 451.57(6)	PeptidePentane - 463.82(11)	
												S	yste	m											

Dispersion-dominated systems

FIG. 2. Comparison between DMC interaction energies $\Delta E_{int.}$ calculated in the present work against CCSD(T) for a subset of systems in the S66 dataset dominated by dispersion interactions. The first 6 dimers are $\pi - \pi$ stacked. Refer to the caption of Fig. 1 for the plot details. The CCSD(T) estimate is taken as an average from three previous calculations^{75,88,91}, with corresponding standard deviation as error. Additional CCSD(cT)-fit estimates are are reported with golden crosses. These are calculated by scaling the CCSD(T) estimates based on their MP2 and CCSD contributions with the approach described in Ref. 50.

The stronger binding of DMC over CCSD(T) has not been (systematically) reported before, with the acetic acid dimer (ID 20) giving the maximum deviation of 0.8 kcal/mol across all S66 systems. Within Sec. S7 of the supplementary material, we have confirmed that the computed $\Delta E_{int.}$ estimate (-20.17 ± 0.11 kcal/mol) does not depend on the chosen pseudopotential (eCEPP), localization scheme (DLA) or trial wave-function (LDA). For example, we have also performed allelectron calculations, giving an estimate of -20.19 ± 0.11 kcal/mol (at τ =0.0015 au that is within the statistical uncertainties of our original estimate. We have also performed tests using PBE and PBE0 trial wave-functions, showing that the DMC $\Delta E_{int.}$ has a negligible (<0.15 kcal/mol) dependence on the nodal surface for the DFAs considered. Furthermore, we also computed estimates



FIG. 3. Comparison between DMC interaction energies $\Delta E_{int.}$ calculated in the present work against CCSD(T) for a subset of systems in the S66 dataset with mixed bonding character. The systems consist of T-shaped (TS) aromatic ring complexes as well as X-H··· π (X = C,O,N) interactions. Refer to the caption of Fig. 1 for the plot details. The CCSD(T) estimate is taken as an average from three previous calculations^{75,88,91}, with corresponding standard deviation as error.

for two-other localization schemes: T-move and determinant localization T-move, both of which were within the statistical uncertainties of our original estimate. It should be noted that while the absolute value of the difference can be significant for some hydrogen-bonded systems, the relative difference (normalized against the DMC $\Delta E_{int.}$) is significantly smaller, with a mean relative difference of 2.45% compared to 8.17% for the dispersion-dominated systems (see Sec. S6 of the supplementary material).

The weaker binding of DMC over CCSD(T) for dispersion-dominated systems is now relatively well-documented^{49,50}, and there is evidence that it can be improved by replacing the perturbative triples (T) contribution with the recent $(cT)^{32}$ contribution. We plot the difference between an empirically CCSD(cT)-fit formulation as yellow crosses for the dispersion-dominated systems. In all cases, CCSD(cT) has a weaker binding than CCSD(T), leaning closer towards DMC, leading to an MAD of 0.09kcal/mol and mean relative difference of 3.43% However, this does not



FIG. 4. Error decomposition analysis. We report the absolute difference between DMC and CCSD(T) relative to the DMC values, i.e. $(E_{DMC} - E_{CCSD(T)}) / |E_{DMC}|$, as a function of the natural logarithm of the electrostatic (ELST) to dispersion (DISP) ratio contribution to the binding energy. The ELST to DISP ratio is determined from the SAPT analysis from Ref. 93. The color code is red for H-bonded systems (ID from 1 to 23), blue for dispersion dominated systems (ID from 24 to 46), and green for mixed systems (ID from 47 to 66).

fully resolve the discrepancies across all of the dispersion systems, with significant discrepancy of ~0.4 kcal/mol remaining for the uracil-cyclopentane dimer (ID 42). Such a significant discrepancy makes this a worthwhile system to investigate further and could give clues on remaining discrepancies between DMC and CCSD(T) observed in Ref. 50. The analysis reported above highlights an important outcome of this work: the identification of smaller, simpler systems that show notable discrepancies between DMC and coupled cluster methods. Specifically, we found a discrepancy of approximately ~0.9 kcal/mol for the hydrogen-bonded acetic acid dimer (ID 20) and ~0.5 kcal/mol for the dispersion-dominated uracil–cyclopentane dimer (ID 43), which contain 64 and 98 (total) electrons, respectively. These medium-sized systems represent an almost tenfold reduction in electron count compared to the larger C₆₀@[6]CPPA buckyball-ring system (672 electrons) studied in Ref. 49. Thus, they might offer practical, cost-effective models for further exploring the discrepancy between DMC and coupled-cluster.

Finally, we now focus on the difference between DMC and CCSD(T) as a function of the dispersion and electrostatic contribution to the interaction energy. In particular, we find that the relative differences between DMC and CCSD(T) for each system within the S66 dataset can be correlated to the relative strength of the dispersion and electrostatic interactions that make up its $\Delta E_{int.}$. We used the Symmetry Adapted Perturbation Theory (SAPT) calculations from Burns *et al.*⁹³, which decomposes $\Delta E_{int.}$ into contributions from electrostatics (ELST), exchange, induction, and dispersion (DISP). Notably, we show in Fig. 2 that there is a strong linear trend (R^2 =0.78) between the natural logarithm of the ELST and DISP contributions, log ($\frac{ELST}{DISP}$), and the relative difference (in %) between CCSD(T) and DMC. This analysis confirms our prior observations on the trends between DMC and CCSD(T). For example, the more dominant the DISP contribution to $\Delta E_{int.}$ (i.e., a more negative log ($\frac{ELST}{DISP}$)), the more CCSD(T) is found to underbind with respect to DMC. Similarly, the stronger the ELST contribution to $\Delta E_{int.}$ (i.e., a more positive log ($\frac{ELST}{DISP}$)), the more CCSD(T) is found to overbind with respect to DMC. We expect that this cheap descriptor can be used in the future to identify more challenging systems with larger discrepancies between DMC and CCSD(T).

IV. CONCLUSIONS

To summarize, we have computed highly accurate estimates for the S66 dateset – one of the most widely used databases for non-covalent interactions in biological and organic molecules – with fixed-node diffusion quantum Monte Carlo. These estimates have provided new insights into recent discussions on its discrepancies with another widely-trusted method: coupled cluster theory with single, double and perturbative triple excitations [CCSD(T)]. Our data shows systematic trends, with DMC predicting stronger binding in hydrogen-bonded systems than CCSD(T), and weaker binding in dispersion dominated systems. We show that there is a correlation between the relative strength of these discrepancies with the nature of the interaction, specifically the relative ratio of the electrostatic and dispersion contributions to the interaction energy as provided by previous Symmetry Adapted Perturbation Theory (SAPT) calculations.⁹³ In addition, we show that the discrepancy between DMC and CCSD(T) on dispersion-dominated systems can be reduced using a recently proposed CCSD(cT) formulation, albeit with still significant remaining

differences. While this work does not identify the origin of the disagreement between DMC and CCSD(T), it has identified the type of interactions where it is particularly prevalent and importantly, we have identified model systems within the S66 dataset where these errors are prominent. These results have strong implications for the electronic structure theory community, addressing the knowledge gap on the trends of DMC interaction energies for non-covalent molecular complexes. Furthermore, the accurate reference data produced within this work is expected to benefit the wider materials modeling community, being instrumental for benchmarking applications ranging from the development of machine learned interatomic potentials to crystal structure prediction, drug design, and renewable energy.

SUPPLEMENTARY MATERIAL

See the supplementary material for details on the DMC calculations, comprising the convergence of the calculations with respect to the time step, the influence of the choice of the monomer geometry on the dimer interaction energy, as well as tests on the pseudopotential localization error and the Jastrow optimization.

DATA AVAILABILITY

The data that support the findings of this study are available within the article and its supplementary material. All analysis can be found on GitHub (made public upon acceptance of the manuscript) at zenandrea/FNDMC-S66 and can be viewed interactively online through associated Jupyter notebooks (via Google Colab), with links provided in the corresponding GitHub repository.

ACKNOWLEDGMENTS

We acknowledge the computational resources from Cambridge Service for Data Driven Discovery (CSD3) operated by the University of Cambridge Research Computing Service, provided by Dell EMC and Intel using Tier-2 funding from the Engineering and Physical Sciences Research Council (capital grant EP/T022159/1 and EP/P020259/1), and DiRAC funding from the Science and Technology Facilities Council (www.dirac.ac.uk). We are further grateful for computational support from the UK national high performance computing service, ARCHER2, for which access was obtained via the UKCP consortium and funded by EPSRC grant ref EP/X035891/1. This research also used resources of the Oak Ridge Leadership Computing Facility at the Oak Ridge National Laboratory, which is supported by the Office of Science of the U.S. Department of Energy under Contract No. DE-AC05-00OR22725. A.M. and B.X.S acknowledge support from the European Union under the "n-AQUA" European Research Council project (Grant No. 101071937). D.A. and A.Z. acknowledges support from Leverhulme grant no. RPG-2020-038, and from the European Union under the Next generation EU (projects 20222FXZ33 and P2022MC742).

REFERENCES

- ¹G. J. O. Beran, "Modeling Polymorphic Molecular Crystals with Electronic Structure Theory," Chem. Rev. **116**, 5567–5613 (2016).
- ²K. E. Riley and P. Hobza, "Noncovalent interactions in biochemistry," WIREs Comput. Mol. Sci. **1**, 3–17 (2011).
- ³M. Raynal, P. Ballester, A. Vidal-Ferran, and P. W. N. M. van Leeuwen, "Supramolecular catalysis. Part 1: Non-covalent interactions as a tool for building and modifying homogeneous catalysts," Chem. Soc. Rev. **43**, 1660–1733 (2014).
- ⁴N. Planas, A. L. Dzubak, R. Poloni, L.-C. Lin, A. McManus, T. M. McDonald, J. B. Neaton, J. R. Long, B. Smit, and L. Gagliardi, "The Mechanism of Carbon Dioxide Adsorption in an Alkylamine-Functionalized Metal–Organic Framework," J. Am. Chem. Soc. **135**, 7402–7405 (2013).
- ⁵R. Poloni, B. Smit, and J. B. Neaton, "Ligand-Assisted Enhancement of CO2 Capture in Metal– Organic Frameworks," J. Am. Chem. Soc. **134**, 6714–6719 (2012).
- ⁶X. Du, Y. Li, Y.-L. Xia, S.-M. Ai, J. Liang, P. Sang, X.-L. Ji, and S.-Q. Liu, "Insights into Protein–Ligand Interactions: Mechanisms, Models, and Methods," Int. J. Mol. Sci. **17**, 144 (2016).
- ⁷F. R. Rehak, G. Piccini, M. Alessio, and J. Sauer, "Including dispersion in density functional theory for adsorption on flat oxide surfaces, in metal–organic frameworks and in acidic zeolites," Phys. Chem. Chem. Phys. 22, 7577–7585 (2020).
- ⁸K. Müller-Dethlefs and P. Hobza, "Noncovalent Interactions: A Challenge for Experiment and Theory," Chem. Rev. **100**, 143–168 (2000).
- ⁹W. M. C. Foulkes, L. Mitas, R. J. Needs, and G. Rajagopal, "Quantum Monte Carlo simulations of solids," Rev. Mod. Phys. **73**, 33–83 (2001).

- ¹⁰R. J. Bartlett and M. Musiał, "Coupled-cluster theory in quantum chemistry," Rev. Mod. Phys. **79**, 291–352 (2007).
- ¹¹A. E. Mattsson, P. A. Schultz, M. P. Desjarlais, T. R. Mattsson, and K. Leung, "Designing meaningful density functional theory calculations in materials science—a primer," Modelling Simul. Mater. Sci. Eng. 13, R1–R31 (2004).
- ¹²D. M. Ceperley and B. J. Alder, "Ground State of the Electron Gas by a Stochastic Method," Phys. Rev. Lett. 45, 566–569 (1980).
- ¹³S. Grimme, "Semiempirical GGA-type density functional constructed with a long-range dispersion correction," J. Comp. Chem. 27, 1787–1799 (2006).
- ¹⁴S. Grimme, J. Antony, S. Ehrlich, and H. Krieg, "A consistent and accurate ab initio parametrization of density functional dispersion correction (DFT-D) for the 94 elements H-Pu," J. Chem. Phys. **132**, 154104 (2010).
- ¹⁵S. Grimme, A. Hansen, J. G. Brandenburg, and C. Bannwarth, "Dispersion-Corrected Mean-Field Electronic Structure Methods," Chem. Rev. **116**, 5105–5154 (2016).
- ¹⁶A. J. A. Price, A. Otero-de-la-Roza, and E. R. Johnson, "XDM-corrected hybrid DFT with numerical atomic orbitals predicts molecular crystal lattice energies with unprecedented accuracy," Chem. Sci. 14, 1252–1262 (2023).
- ¹⁷A. Zen, S. Sorella, M. J. Gillan, A. Michaelides, and D. Alfè, "Boosting the accuracy and speed of quantum Monte Carlo: Size consistency and time step," Phys. Rev. B **93**, 241118 (2016).
- ¹⁸J. T. Krogel, "Nexus: A modular workflow management system for quantum simulation codes," Comput. Phys. Commun. **198**, 154–168 (2016).
- ¹⁹M. C. Bennett, C. A. Melton, A. Annaberdiyev, G. Wang, L. Shulenburger, and L. Mitas, "A new generation of effective core potentials for correlated calculations," J. Chem. Phys. **147**, 224106 (2017).
- ²⁰M. C. Bennett, G. Wang, A. Annaberdiyev, C. A. Melton, L. Shulenburger, and L. Mitas, "A new generation of effective core potentials from correlated calculations: 2nd row elements," J. Chem. Phys. **149**, 104108 (2018).
- ²¹A. Annaberdiyev, G. Wang, C. A. Melton, M. C. Bennett, L. Shulenburger, and L. Mitas, "A new generation of effective core potentials from correlated calculations: 3d transition metal series," J. Chem. Phys. 149, 134108 (2018).
- ²²A. Zen, J. G. Brandenburg, A. Michaelides, and D. Alfè, "A new scheme for fixed node diffusion quantum Monte Carlo with pseudopotentials: Improving reproducibility and reducing the trial-

wave-function bias," J. Chem. Phys. 151, 134105 (2019).

- ²³R. J. Needs, M. D. Towler, N. D. Drummond, P. López Ríos, and J. R. Trail, "Variational and diffusion quantum Monte Carlo calculations with the CASINO code," J. Chem. Phys. 152, 154106 (2020).
- ²⁴K. Nakano, C. Attaccalite, M. Barborini, L. Capriotti, M. Casula, E. Coccia, M. Dagrada, C. Genovese, Y. Luo, G. Mazzola, A. Zen, and S. Sorella, "TurboRVB: A many-body toolkit for ab initio electronic simulations by quantum Monte Carlo," J. Chem. Phys. **152**, 204121 (2020).
- ²⁵P. R. C. Kent, A. Annaberdiyev, A. Benali, M. C. Bennett, E. J. Landinez Borda, P. Doak, H. Hao, K. D. Jordan, J. T. Krogel, I. Kylänpää, J. Lee, Y. Luo, F. D. Malone, C. A. Melton, L. Mitas, M. A. Morales, E. Neuscamman, F. A. Reboredo, B. Rubenstein, K. Saritas, S. Upadhyay, G. Wang, S. Zhang, and L. Zhao, "QMCPACK: Advances in the development, efficiency, and application of auxiliary field and real-space variational and diffusion quantum Monte Carlo," J. Chem. Phys. **152**, 174105 (2020).
- ²⁶K. Nakano, O. Kohulák, A. Raghav, M. Casula, and S. Sorella, "TurboGenius: Python suite for high-throughput calculations of ab initio quantum Monte Carlo methods," J. Chem. Phys. **159**, 224801 (2023).
- ²⁷C. Riplinger and F. Neese, "An efficient and near linear scaling pair natural orbital based local coupled cluster method," J. Chem. Phys. **138**, 034106 (2013).
- ²⁸C. Riplinger, P. Pinski, U. Becker, E. F. Valeev, and F. Neese, "Sparse maps—A systematic infrastructure for reduced-scaling electronic structure methods. II. Linear scaling domain based pair natural orbital coupled cluster theory," J. Chem. Phys. **144**, 024109 (2016).
- ²⁹Q. Ma and H.-J. Werner, "Explicitly correlated local coupled-cluster methods using pair natural orbitals," Wiley Interdiscip. Rev.: Comput. Mol. Sci. 8, e1371 (2018).
- ³⁰P. R. Nagy, G. Samu, and M. Kállay, "Optimization of the Linear-Scaling Local Natural Orbital CCSD(T) Method: Improved Algorithm and Benchmark Applications," J. Chem. Theory Comput. 14, 4193–4215 (2018).
- ³¹P. R. Nagy and M. Kállay, "Approaching the Basis Set Limit of CCSD(T) Energies for Large Molecules with Local Natural Orbital Coupled-Cluster Methods," J. Chem. Theory Comput. 15, 5275–5298 (2019).
- ³²N. Masios, A. Irmler, T. Schäfer, and A. Grüneis, "Averting the Infrared Catastrophe in the Gold Standard of Quantum Chemistry," Phys. Rev. Lett. **131**, 186401 (2023).

- ³³A. Jiang, Z. L. Glick, D. Poole, J. M. Turney, C. D. Sherrill, and H. F. Schaefer, III, "Accurate and efficient open-source implementation of domain-based local pair natural orbital (DLPNO) coupled-cluster theory using a t1-transformed Hamiltonian," J. Chem. Phys. 161, 082502 (2024).
- ³⁴H.-Z. Ye and T. C. Berkelbach, "Periodic Local Coupled-Cluster Theory for Insulators and Metals," J. Chem. Theory Comput. **20**, 8948–8959 (2024).
- ³⁵M. Dubecký, P. Jurečka, R. Derian, P. Hobza, M. Otyepka, and L. Mitas, "Quantum Monte Carlo Methods Describe Noncovalent Interactions with Subchemical Accuracy," J. Chem. Theory Comput. 9, 4287–4292 (2013).
- ³⁶J. Řezáč, M. Dubecký, P. Jurečka, and P. Hobza, "Extensions and applications of the A24 data set of accurate interaction energies," Phys. Chem. Chem. Phys. **17**, 19268–19277 (2015).
- ³⁷A. Raghav, R. Maezono, K. Hongo, S. Sorella, and K. Nakano, "Toward Chemical Accuracy Using the Jastrow Correlated Antisymmetrized Geminal Power Ansatz," J. Chem. Theory Comput. 19, 2222–2229 (2023).
- ³⁸E. Mostaani, N. D. Drummond, and V. I. Fal'ko, "Quantum Monte Carlo Calculation of the Binding Energy of Bilayer Graphene," Phys. Rev. Lett. **115**, 115501 (2015).
- ³⁹A. Zen, J. G. Brandenburg, J. Klimeš, A. Tkatchenko, D. Alfè, and A. Michaelides, "Fast and accurate quantum Monte Carlo for molecular crystals," Proc. Natl. Acad. Sci. U. S. A. 115, 1724–1729 (2018).
- ⁴⁰F. Della Pia, A. Zen, D. Alfè, and A. Michaelides, "DMC-ICE13: Ambient and high pressure polymorphs of ice from diffusion Monte Carlo and density functional theory," J. Chem. Phys. **157**, 134701 (2022).
- ⁴¹F. Della Pia, A. Zen, D. Alfè, and A. Michaelides, "How Accurate Are Simulations and Experiments for the Lattice Energies of Molecular Crystals?" Phys. Rev. Lett. **133**, 046401 (2024).
- ⁴²O. Karalti, D. Alfè, M. J. Gillan, and K. D. Jordan, "Adsorption of a water molecule on the MgO(100) surface as described by cluster and slab models," Phys. Chem. Chem. Phys. 14, 7846– 7853 (2012).
- ⁴³Y. S. Al-Hamdani, D. Alfè, and A. Michaelides, "How strongly do hydrogen and water molecules stick to carbon nanomaterials?" J. Chem. Phys. **146**, 094701 (2017).
- ⁴⁴T. Tsatsoulis, F. Hummel, D. Usvyat, M. Schütz, G. H. Booth, S. S. Binnie, M. J. Gillan, D. Alfè, A. Michaelides, and A. Grüneis, "A comparison between quantum chemistry and quantum Monte Carlo techniques for the adsorption of water on the (001) LiH surface," J. Chem. Phys. 146, 204108 (2017).

- ⁴⁵Y. S. Al-Hamdani, M. Rossi, D. Alfè, T. Tsatsoulis, B. Ramberger, J. G. Brandenburg, A. Zen, G. Kresse, A. Grüneis, A. Tkatchenko, and A. Michaelides, "Properties of the water to boron nitride interaction: From zero to two dimensions with benchmark accuracy," J. Chem. Phys. 147, 044710 (2017).
- ⁴⁶J. G. Brandenburg, A. Zen, M. Fitzner, B. Ramberger, G. Kresse, T. Tsatsoulis, A. Grüneis, A. Michaelides, and D. Alfè, "Physisorption of Water on Graphene: Subchemical Accuracy from Many-Body Electronic Structure Methods," J. Phys. Chem. Lett. 10, 358–368 (2019).
- ⁴⁷B. X. Shi, A. Zen, V. Kapil, P. R. Nagy, A. Grüneis, and A. Michaelides, "Many-Body Methods for Surface Chemistry Come of Age: Achieving Consensus with Experiments," J. Am. Chem. Soc. 145, 25372–25381 (2023).
- ⁴⁸B. X. Shi, V. Kapil, A. Zen, J. Chen, A. Alavi, and A. Michaelides, "General embedded cluster protocol for accurate modeling of oxygen vacancies in metal-oxides," J. Chem. Phys. **156**, 124704 (2022).
- ⁴⁹Y. S. Al-Hamdani, P. R. Nagy, A. Zen, D. Barton, M. Kállay, J. G. Brandenburg, and A. Tkatchenko, "Interactions between large molecules pose a puzzle for reference quantum mechanical methods," Nat. Commun. **12**, 3927 (2021).
- ⁵⁰T. Schäfer, A. Irmler, A. Gallo, and A. Grüneis, "Understanding Discrepancies of Wavefunction Theories for Large Molecules," (2024), arXiv:2407.01442.
- ⁵¹F. Ballesteros, S. Dunivan, and K. U. Lao, "Coupled cluster benchmarks of large noncovalent complexes: The L7 dataset as well as DNA–ellipticine and buckycatcher–fullerene," J. Chem. Phys. **154**, 154104 (2021).
- ⁵²C. Villot, F. Ballesteros, D. Wang, and K. U. Lao, "Coupled Cluster Benchmarking of Large Noncovalent Complexes in L7 and S12L as Well as the C60 Dimer, DNA–Ellipticine, and HIV– Indinavir," J. Phys. Chem. A **126**, 4326–4341 (2022).
- ⁵³V. Fishman, M. Lesiuk, J. M. L. Martin, and A. D. Boese, "A New Angle on Benchmarking Noncovalent Interactions," (2024), arXiv:2410.12603.
- ⁵⁴S. Lambie, D. Kats, D. Usyvat, and A. Alavi, "On the applicability of CCSD(T) for dispersion interactions in large conjugated systems," (2024), arXiv:2411.13986 [physics].
- ⁵⁵K. U. Lao, "Canonical coupled cluster binding benchmark for nanoscale noncovalent complexes at the hundred-atom scale," J. Chem. Phys. **161**, 234103 (2024).
- ⁵⁶E. Semidalas, G. Santra, N. Mehta, and J. M. L. Martin, "S66 noncovalent interactions benchmark re-examined: Composite localized coupled cluster approaches," AIP Conf. Proc. **2611**,

020016 (2022).

- ⁵⁷R. Pariser and R. G. Parr, "A Semi-Empirical Theory of the Electronic Spectra and Electronic Structure of Complex Unsaturated Molecules. I." J. Chem. Phys. **21**, 466–471 (1953).
- ⁵⁸R. Pariser and R. G. Parr, "A Semi-Empirical Theory of the Electronic Spectra and Electronic Structure of Complex Unsaturated Molecules. II," J. Chem. Phys. 21, 767–776 (1953).
- ⁵⁹J. A. Pople, "Electron interaction in unsaturated hydrocarbons," Trans. Faraday Soc. 49, 1375– 1385 (1953).
- ⁶⁰R. Sure and S. Grimme, "Comprehensive Benchmark of Association (Free) Energies of Realistic Host–Guest Complexes," J. Chem. Theory Comput. **11**, 3785–3801 (2015).
- ⁶¹S. Grimme, "Supramolecular Binding Thermodynamics by Dispersion-Corrected Density Functional Theory," Chem. - Eur. J. **18**, 9955–9964 (2012).
- ⁶²J. Řezáč and P. Hobza, "Describing Noncovalent Interactions beyond the Common Approximations: How Accurate Is the "Gold Standard," CCSD(T) at the Complete Basis Set Limit?" J. Chem. Theory Comput. 9, 2151–2155 (2013).
- ⁶³P. Jurečka, J. Šponer, J. Černý, and P. Hobza, "Benchmark database of accurate (MP2 and CCSD(T) complete basis set limit) interaction energies of small model complexes, DNA base pairs, and amino acid pairs," Phys. Chem. Chem. Phys. 8, 1985–1993 (2006).
- ⁶⁴J. Řezáč, K. E. Riley, and P. Hobza, "S66: A Well-balanced Database of Benchmark Interaction Energies Relevant to Biomolecular Structures," J. Chem. Theory Comput. 7, 2427–2438 (2011).
- ⁶⁵M. O. Sinnokrot and C. D. Sherrill, "Highly Accurate Coupled Cluster Potential Energy Curves for the Benzene Dimer: Sandwich, T-Shaped, and Parallel-Displaced Configurations," J. Phys. Chem. A **108**, 10200–10207 (2004).
- ⁶⁶L. Goerigk, H. Kruse, and S. Grimme, "Benchmarking Density Functional Methods against the S66 and S66x8 Datasets for Non-Covalent Interactions," ChemPhysChem 12, 3421–3433 (2011).
- ⁶⁷T. Gao, H. Li, W. Li, L. Li, C. Fang, H. Li, L. Hu, Y. Lu, and Z.-M. Su, "A machine learning correction for DFT non-covalent interactions based on the S22, S66 and X40 benchmark databases," J. Cheminform. **8**, 24 (2016).
- ⁶⁸H. Peng, Z.-H. Yang, J. P. Perdew, and J. Sun, "Versatile van der Waals Density Functional Based on a Meta-Generalized Gradient Approximation," Phys. Rev. X **6**, 041005 (2016).
- ⁶⁹F. Yu and Y. Wang, "Dual-hybrid direct random phase approximation and second-order screened exchange with nonlocal van der Waals correlations for noncovalent interactions," J. Comp.

Chem. 41, 1018–1025 (2020).

- ⁷⁰S. Grimme, A. Hansen, S. Ehlert, and J.-M. Mewes, "r2SCAN-3c: A "Swiss army knife" composite electronic-structure method," J. Chem. Phys. **154**, 064103 (2021).
- ⁷¹S. Ehlert, U. Huniar, J. Ning, J. W. Furness, J. Sun, A. D. Kaplan, J. P. Perdew, and J. G. Brandenburg, "r2SCAN-D4: Dispersion corrected meta-generalized gradient approximation for general chemical applications," J. Chem. Phys. **154**, 061101 (2021).
- ⁷²M. Müller, A. Hansen, and S. Grimme, "ωB97X-3c: A composite range-separated hybrid DFT method with a molecule-optimized polarized valence double-ζ basis set," J. Chem. Phys. **158**, 014103 (2023).
- ⁷³T. Lu and Q. Chen, "Simple, Efficient, and Universal Energy Decomposition Analysis Method Based on Dispersion-Corrected Density Functional Theory," J. Phys. Chem. A 127, 7023–7035 (2023).
- ⁷⁴M. Lee, B. Kim, M. Sim, M. Sogal, Y. Kim, H. Yu, K. Burke, and E. Sim, "Correcting Dispersion Corrections with Density-Corrected DFT," J. Chem. Theory Comput. **20**, 7155–7167 (2024).
- ⁷⁵J. Řezáč, K. E. Riley, and P. Hobza, "Extensions of the S66 Data Set: More Accurate Interaction Energies and Angular-Displaced Nonequilibrium Geometries," J. Chem. Theory Comput. 7, 3466–3470 (2011).
- ⁷⁶K. E. Riley, J. Řezáč, and P. Hobza, "The performance of MP2.5 and MP2.X methods for nonequilibrium geometries of molecular complexes," Phys. Chem. Chem. Phys. 14, 13187– 13193 (2012).
- ⁷⁷A. Altun, F. Neese, and G. Bistoni, "Extrapolation to the Limit of a Complete Pair Natural Orbital Space in Local Coupled-Cluster Calculations," J. Chem. Theory Comput. 16, 6142–6149 (2020).
- ⁷⁸J. Shee, M. Loipersberger, A. Rettig, J. Lee, and M. Head-Gordon, "Regularized Second-Order Møller–Plesset Theory: A More Accurate Alternative to Conventional MP2 for Noncovalent Interactions and Transition Metal Thermochemistry for the Same Computational Cost," J. Phys. Chem. Lett. 12, 12084–12097 (2021).
- ⁷⁹J. Lupi, S. Alessandrini, C. Puzzarini, and V. Barone, "junChS and junChS-F12 Models: Parameter-free Efficient yet Accurate Composite Schemes for Energies and Structures of Noncovalent Complexes," J. Chem. Theory Comput. **17**, 6974–6992 (2021).

- ⁸⁰G. J. O. Beran, C. Greenwell, C. Cook, and J. Řezáč, "Improved Description of Intra- and Intermolecular Interactions through Dispersion-Corrected Second-Order Møller–Plesset Perturbation Theory," Acc. Chem. Res. 56, 3525–3534 (2023).
- ⁸¹A. S. Christensen, S. K. Sirumalla, Z. Qiao, M. B. O'Connor, D. G. A. Smith, F. Ding, P. J. Bygrave, A. Anandkumar, M. Welborn, F. R. Manby, and T. F. Miller, III, "OrbNet Denali: A machine learning potential for biological and organic chemistry with semi-empirical cost and DFT accuracy," J. Chem. Phys. 155, 204103 (2021).
- ⁸²C. Villot and K. U. Lao, "Ab initio dispersion potentials based on physics-based functional forms with machine learning," J. Chem. Phys. **160**, 184103 (2024).
- ⁸³J. R. Trail and R. J. Needs, "Shape and energy consistent pseudopotentials for correlated electron systems," J. Chem. Phys. **146**, 204107 (2017).
- ⁸⁴P. Giannozzi, S. Baroni, N. Bonini, M. Calandra, R. Car, C. Cavazzoni, D. Ceresoli, G. L. Chiarotti, M. Cococcioni, I. Dabo, A. D. Corso, S. de Gironcoli, S. Fabris, G. Fratesi, R. Gebauer, U. Gerstmann, C. Gougoussis, A. Kokalj, M. Lazzeri, L. Martin-Samos, N. Marzari, F. Mauri, R. Mazzarello, S. Paolini, A. Pasquarello, L. Paulatto, C. Sbraccia, S. Scandolo, G. Sclauzero, A. P. Seitsonen, A. Smogunov, P. Umari, and R. M. Wentzcovitch, "QUAN-TUM ESPRESSO: A modular and open-source software project for quantum simulations of materials," J. Phys.: Condens. Matter 21, 395502 (2009).
- ⁸⁵P. Giannozzi, O. Baseggio, P. Bonfà, D. Brunato, R. Car, I. Carnimeo, C. Cavazzoni, S. de Gironcoli, P. Delugas, F. Ferrari Ruffino, A. Ferretti, N. Marzari, I. Timrov, A. Urru, and S. Baroni, "Quantum ESPRESSO toward the exascale," J. Chem. Phys. **152**, 154105 (2020).
- ⁸⁶D. Alfè and M. J. Gillan, "Efficient localized basis set for quantum Monte Carlo calculations on condensed matter," Phys. Rev. B 70, 161101 (2004).
- ⁸⁷B. Brauer, M. K. Kesharwani, S. Kozuch, and J. M. L. Martin, "The S66x8 benchmark for noncovalent interactions revisited: Explicitly correlated ab initio methods and density functional theory," Phys. Chem. Chem. Phys. 18, 20905–20925 (2016).
- ⁸⁸M. K. Kesharwani, A. Karton, N. Sylvetsky, and J. M. L. Martin, "The S66 Non-Covalent Interactions Benchmark Reconsidered Using Explicitly Correlated Methods Near the Basis Set Limit*," Aust. J. Chem. 71, 238–248 (2018).
- ⁸⁹Q. Ma and H.-J. Werner, "Accurate Intermolecular Interaction Energies Using Explicitly Correlated Local Coupled Cluster Methods [PNO-LCCSD(T)-F12]," J. Chem. Theory Comput. 15, 1044–1052 (2019).

- ⁹⁰G. Santra, E. Semidalas, N. Mehta, A. Karton, and J. M. L. Martin, "S66x8 noncovalent interactions revisited: New benchmark and performance of composite localized coupled-cluster methods," Phys. Chem. Chem. Phys. 24, 25555–25570 (2022).
- ⁹¹P. R. Nagy, L. Gyevi-Nagy, B. D. Lőrincz, and M. Kállay, "Pursuing the basis set limit of CCSD(T) non-covalent interaction energies for medium-sized complexes: Case study on the S66 compilation," Mol. Phys. **121**, e2109526 (2023).
- ⁹²Note that the difficulty of reaching the complete basis set (CBS) limit is the principle reason for several CCSD(T) evaluations of S66. As reaching the CBS limit remains a challenge, particularly in dispersion bound complexes.
- ⁹³L. A. Burns, J. C. Faver, Z. Zheng, M. S. Marshall, D. G. A. Smith, K. Vanommeslaeghe, A. D. MacKerell, Jr., K. M. Merz, Jr., and C. D. Sherrill, "The BioFragment Database (BFDb): An open-data platform for computational chemistry analysis of noncovalent interactions," J. Chem. Phys. 147, 161727 (2017).

Supporting Information: Systematic discrepancies between reference methods for non-covalent interactions within the S66 dataset

Benjamin X. Shi,^{†,⊥} Flaviano Della Pia,^{†,⊥} Yasmine S. Al-Hamdani,^{‡,¶,§} Angelos

Michaelides,[†] Dario Alfè,^{\ddagger , \parallel , \P ,\$ and Andrea Zen^{*, \ddagger ,\$}}

[†]Yusuf Hamied Department of Chemistry, University of Cambridge, Lensfield Road, Cambridge CB2 1EW, United Kingdom

‡Dipartimento di Fisica Ettore Pancini, Universita di Napoli Federico II, Monte Sant'Angelo, I-80126 Napoli, Italy

¶ Thomas Young Centre, University College London, London WC1E 6BT, United Kingdom

§Department of Earth Sciences, University College London, London WC1E 6BT, United

Kingdom

||London Centre for Nanotechnology, University College London, London WC1E 6BT, United Kingdom

 \perp These authors contributed equally to this work.

E-mail: andrea.zen@unina.it

Contents

S1	The S66 dataset	S- 4
S2	Validating diffusion Monte Carlo	S-6
	S2.1 Computing the interaction energy	S-6
	S2.2 Setup for the $CCSD(T)$ calculations used to evaluate the deformation energy	S-12
	S2.3 Brief summary	S-18
	S2.4 Reaching the time step limit and estimating errors	S-19
S 3	CCSD(T) estimates from the literature	S-20
$\mathbf{S4}$	Final CCSD(T) and CCSD(cT)-fit estimates	S-25
$\mathbf{S5}$	Final DMC estimates	S-26
S6	Mean relative difference of DMC and CCSD(T)	S-32
S 7	Validation tests for the AcOH dimer	S-3 4
S 8	Convergence of the total and binding energy with respect to the DM	С
	simulation time step	S-36
S 9	Interaction energy decomposition analysis	S-59
Re	eferences	S-63

We provide here additional supporting data as well as contextual information for the manuscript "On the systematic discrepancies between reference methods on non-covalent interaction energies within the S66 dataset". All output files are provided on GitHub, which contains a Jupyter Notebook file that analyzes the data. This data can also be viewed and analyzed on the browser with Colab.

In particular, in this supplemental material we provide:

- a brief description of the three previous CCSD(T) estimates of the binding energy of the S66 dataset in Sec. S3, with the final estimates given in S4.
- the total energy of each dimer and the corresponding monomers used to compute the binding energies reported in the main manuscript in Sec. S5;
- an analysis on the mean relative differences between DMC and CCSD(T) on the S66 dataset in Sec. S6;
- a quantitative analysis of the localization error on the binding energy for the case of acetic acid in Sec. S7;
- the convergence of the DMC estimates with respect to the simulation time step for all the dimers in Sec. S8.
- an energy decomposition analysis (into electrostatic, dispersive, induction and exchange contributions) of the S66 dataset with the SAPT method in Sec. S9.

S1 The S66 dataset

The entire S66 dataset is visualised in Fig. S1. It consists of 66 dimer complexes, composed from combinations of 14 monomer molecules. These monomers consist of only carbon, oxygen, nitrogen, and hydrogen – the most commonly encountered elements in biochemistry. Within the dimers, the monomers are combined and placed at different geometries, for example through parallel π - π stacking or in a T-shape (TS) or with NH, CH or OH groups pointing perpendicular to the plane of an aromatic π ring, among others. The geometries of these dimers were obtained from second-order Møller-Plesset perturbation theory (MP2) performed with the Dunning cc-pVTZ basis set.

The dimers of the S66 dataset were chosen to sample a balanced range of non-covalent interactions, consisting of 23 electrostatic-dominated systems [IDs 1-23], 23 dispersion-dominated systems [IDs 24-46], and 20 systems [IDs 47-66] with mixed (electrostatic/dispersion) interactions. It should be noted all these classifications are rather arbitrary and can differ based on the choice of energy decomposition analysis schemes. Regardless, they have been chosen to sample some important types of interactions within each category. For examples, the electrostatic-dominated systems covers all possible combinations of hydrogen bonding donors and acceptors of the water molecule, hydroxyl group, amine group, and carbonyl group, alongside the type of hydrogen bonding expected in nucleic acid base pairs. The dispersion-dominated systems consists of combinations between planar aromatic molecules and aliphatic hydrocarbons, leading to three types of interactions: $\pi-\pi$ stacking (10 systems), aliphatic–aliphatic (5 systems), and π –aliphatic (8 systems) interactions.



Figure S1: A visualization of the 66 dimer complexes within the S66 dataset. The IDs for each system is provided on the top right with additional description of their geometry given in the bottom right.

S2 Validating diffusion Monte Carlo

S2.1 Computing the interaction energy

The DMC interaction energies of the $S66^{S1}$ dataset are computed as:

$$\Delta E_{\text{int.}}^{\text{DMC}} = E_{\text{dimer}}^{\text{DMC}} - E_{\text{mon. 1}}^{\text{DMC}} - E_{\text{mon. 2}}^{\text{DMC}},\tag{1}$$

where E_{dimer} is the total energy of the dimer, and $E_{\text{mon. 1}}$, $E_{\text{mon. 2}}$ are the total energies of the constituent monomers. Here, the constituent monomers take on the geometry they adopt in the dimer – away from their equilibrium position.

Table S1: Total energy of the 14 monomers which make up the S66 dataset. These geometries are taken from specific dimer complexes within the S66 dataset that are identified in the table and the order in which the monomer appears (important for dimers consisting of the same molecule) is reported. The method (see Sec. S2.4) used to estimate the error on the DMC estimate is also provided.

Monomer	Dimer Geometry	Order	Total Energy [kcal/mol]	Error Type
$AcNH_2$	$AcNH_2 \cdots AcNH_2 (ID 21)$	1	-25290.30 ± 0.03	$\sigma_{ m cubic\ fit}$
AcOH	$AcOH \cdots AcOH (ID 20)$	1	-28725.26 ± 0.05	$\Delta_{\rm cubic\ fit}^{\rm linear\ fit}$
Benzene	Benzene \cdots Benzene $(\pi - \pi)$ (ID 24)	1	-23624.42 ± 0.04	$\sigma_{ m cubic\ fit}$
Cyclopentane	Cyclopentane \cdots Neopentane (ID 37)	1	-21586.06 ± 0.03	$\sigma_{ m cubic\ fit}$
Ethene	Benzene···Ethene (ID 30)	2	-8610.42 ± 0.02	$\sigma_{ m cubic\ fit}$
Ethyne	$\text{Uracil} \cdots \text{Ethyne} (\text{ID } 32)$	2	-7823.07 ± 0.02	$\sigma_{ m cubic\ fit}$
$MeNH_2$	Benzene···MeNH ₂ (NH- π) (ID 56)	2	-11671.48 ± 0.04	$\Delta_{\text{cubic fit}}^{\text{linear fit}}$
MeOH	Benzene···MeOH (OH- π) (ID 55)	2	-15103.82 ± 0.02	$\sigma_{ m cubic\ fit}$
Neopentane	Neopentane \cdots Neopentane (ID 36)	1	-22350.00 ± 0.04	$\sigma_{ m cubic\ fit}$
Pentane	Pentane \cdots Pentane (ID 34)	1	-22346.69 ± 0.03	$\sigma_{ m cubic\ fit}$
Peptide	Benzene···Peptide (NH- π) (ID 57)	2	-29604.33 ± 0.04	$\sigma_{ m cubic\ fit}$
Pyridine	Pyridine \cdots Pyridine $(\pi - \pi)$ (ID 25)	1	-25905.64 ± 0.05	$\Delta_{\rm cubic\ fit}^{\rm linear\ fit}$
Uracil	Uracil··· Uracil $(\pi - \pi)$ (ID 26)	1	-48309.04 ± 0.12	$\Delta_{\rm cubic\ fit}^{\rm linear\ fit}$
Water	Water \cdots Water (ID 1)	2	-10799.44 ± 0.01	$\sigma_{ m cubic fit}$

There are a total of only 14 different monomer species (listed in Table S1 that are combined to make up the 66 dimers. Importantly, the above definition requires the calculation of the total energy of 132 monomers, which can add significant manual expense and cost to compute with DMC. We reach an estimate of the energy of each monomer by computing the DMC total energy at a reference geometry (chosen from a dimer in the S66 dataset) $E_{\text{monomer 1/2, ref.}}^{\text{DMC}}$, combined with a deformation energy (to reach its geometry in the dimer) computed at the CCSD(T) level (details given in section S2.2):

$$E_{\text{mon. 1}}^{\text{DMC}} = E_{\text{mon. 1, ref.}}^{\text{DMC}} + \Delta E_{\text{mon. 1, def.}}^{\text{CCSD(T)}}$$
(2)

Thus, this requires DMC estimates on the total energy of only 14 monomers. We give the final estimate to the DMC total energy for each of the 14 monomers in Table S1, with the corresponding dimer geometry where this monomer was taken from identified and the type of extrapolation used to reach the zero time step limit.

Figs. S2– S15 illustrate the time step dependence of the total energy for each individual monomer. Table S2 illustrates the CCSD(T) deformation energy calculated for each of the two monomers of the dimers of the S66 dataset with respect to the corresponding geometries used with DMC. We show in Table S3 that the CCSD(T) deformation energy matches DMC estimates to within 0.12 kcal/mol for a subset of systems. The DMC estimates were reported for the 0.01 au time step.

ID	Dimer Name	$\Delta E_{\rm mon.~1,~def.}^{\rm CCSD(T)}~[\rm kcal/mol]$	$\Delta E_{\rm mon.~2,~def.}^{\rm CCSD(T)}~[\rm kcal/mol]$
1	Water···Water	0.031	0.000
2	$Water \cdots MeOH$	0.042	-0.016
3	$\mathrm{Water} \cdots \mathrm{MeNH}_2$	0.109	-0.026
4	$Water \cdots Peptide$	0.087	0.067
5	$\mathrm{MeOH} \cdots \mathrm{MeOH}$	0.056	-0.022
6	$\mathrm{MeOH} \cdots \mathrm{MeNH}_2$	0.222	-0.026

Table S2: Deformation energy for the two monomers within each of the dimers of the S66 dataset. This energy is with respect to the geometry used in Table S1.

Continued on next page

Table S2: (continued)

7	$MeOH \cdots Peptide$	0.147	-0.006
8	$MeOH \cdots Water$	0.038	-0.001
9	$MeNH_2\cdots MeOH$	-0.003	-0.033
10	$\mathrm{MeNH}_2\!\cdots\mathrm{MeNH}_2$	0.005	-0.015
11	$MeNH_2 \cdots Peptide$	0.018	-0.102
12	$MeNH_2 \cdots Water$	-0.018	0.116
13	$\operatorname{Peptide}\cdots\operatorname{MeOH}$	-0.048	-0.034
14	$\operatorname{Peptide}\cdots\operatorname{MeNH}_2$	0.076	-0.016
15	$\operatorname{Peptide}\cdots\operatorname{Peptide}$	0.160	0.078
16	$\operatorname{Peptide}\cdots\operatorname{Water}$	0.050	-0.003
17	$\text{Uracil}\cdots\text{Uracil}(\text{BP})$	0.348	0.230
18	$Water \cdots Pyridine$	0.101	0.004
19	$MeOH \cdots Pyridine$	0.208	0.008
20	$AcOH \cdots AcOH$	0.000	-0.002
21	$AcNH_2 \cdots AcNH_2$	0.000	-0.002
22	$AcOH\cdots Uracil$	0.070	0.390
23	$AcNH_2\cdots Uracil$	0.056	0.500
24	Benzene···Benzene $(\pi - \pi)$	0.000	0.000
25	Pyridine \cdots Pyridine $(\pi - \pi)$	0.000	-0.003
26	Uracil··· Uracil $(\pi - \pi)$	0.000	0.000
27	Benzene···Pyridine $(\pi-\pi)$	-0.002	-0.005
28	Benzene····Uracil $(\pi$ - π)	0.010	-0.305
29	Pyridine · · · Uracil $(\pi$ - π)	0.014	-0.239
30	$Benzene \cdots Ethene$	-0.006	0.000
31	$Uracil \cdots Ethene$	-0.318	-0.000

Continued on next page

Table S2: (continued)

32	$\text{Uracil}\cdots\text{Ethyne}$	-0.246	0.000
33	$Pyridine \cdots Ethene$	-0.016	-0.000
34	$Pentane \cdots Pentane$	0.000	-0.000
35	$Neopentane \cdots Pentane$	0.000	-0.002
36	$Neopentane \cdots Neopentane$	0.000	0.000
37	${\rm Cyclopentane} \cdots {\rm Neopentane}$	0.000	0.000
38	$Cyclopentane \cdots Cyclopentane$	0.007	0.007
39	$Benzene \cdots Cyclopentane$	-0.006	0.011
40	$Benzene \cdots Neopentane$	-0.004	0.006
41	$\text{Uracil}\cdots\text{Pentane}$	-0.338	0.051
42	$Uracil \cdots Cyclopentane$	-0.349	0.024
43	$\mathrm{Uracil}\cdots\mathrm{Neopentane}$	-0.311	0.012
44	$Ethene \cdots Pentane$	-0.003	0.003
45	$Ethyne \cdots Pentane$	-0.027	0.035
46	$\operatorname{Peptide} \cdots \operatorname{Pentane}$	-0.013	0.029
47	Benzene \cdots Benzene (TS)	-0.003	0.004
48	$Pyridine \cdots Pyridine (TS)$	-0.003	0.002
49	Benzene \cdots Pyridine (TS)	-0.001	0.004
50	Benzene···Ethyne (CH- π)	0.001	-0.017
51	Ethyne \cdots Ethyne (TS)	-0.030	-0.026
52	Benzene···AcOH (OH- π)	0.018	-1.276
53	Benzene···AcNH ₂ (NH- π)	0.042	-0.646
54	Benzene···Water (OH- π)	0.000	0.044
55	Benzene···MeOH (OH- π)	0.004	0.000
56	Benzene····MeNH ₂ (NH- π)	-0.001	0.000

Continued on next page

Table S2: (continued)

Benzene···Peptide (NH- π)	0.003	0.000
$Pyridine \cdots Pyridine (CH-N)$	0.018	0.018
Ethyne··· Water (CH-O)	-0.010	-0.001
Ethyne··· AcOH (OH- π)	0.024	-1.253
$Pentane \cdots AcOH$	0.033	-1.330
$Pentane \cdots AcNH_2$	0.031	-0.705
$Benzene \cdots AcOH$	0.002	-1.295
$Peptide \cdots Ethene$	-0.027	0.018
$Pyridine \cdots Ethyne$	-0.006	0.029
$MeNH_2 \cdots Pyridine$	0.002	-0.001
	Benzene \cdots Peptide (NH- π) Pyridine \cdots Pyridine (CH-N) Ethyne \cdots Water (CH-O) Ethyne \cdots AcOH (OH- π) Pentane \cdots AcOH Pentane \cdots AcOH Benzene \cdots AcOH Peptide \cdots Ethene Pyridine \cdots Ethyne MeNH ₂ \cdots Pyridine	Benzene ··· Peptide (NH- π) 0.003 Pyridine ··· Pyridine (CH-N) 0.018 Ethyne ··· Water (CH-O) -0.010 Ethyne ··· AcOH (OH- π) 0.024 Pentane ··· AcOH 0.033 Pentane ··· AcOH 0.031 Benzene ··· AcOH 0.031 Peptide ··· Ethene -0.027 Pyridine ··· Ethyne -0.006 MeNH ₂ ··· Pyridine 0.002

Table S3: Comparison between DMC (0.01 au time step) and CCSD(T) for the deformation energy $E_{def.}$ of a subset of AcNH₂, AcOH, cyclopentane, peptide and urcail monomers found in the S66 dataset. The order in which the monomer appears in the dimer (in the provided .xyz geometry) is given. The reference monomer configuration to calculate $E_{def.}$ is given in Table S2.

Monomer	Dimer Geometry	Order	$\Delta E_{\rm def.}^{\rm DMC}$	$\Delta E_{\rm def.}^{\rm CCSD(T)}$	Deviation
$AcNH_2$	$Pentane \cdots AcNH_2$	2	-0.64 ± 0.05	-0.70	-0.07 ± 0.05
$AcNH_2$	$AcNH_2 \cdots AcNH_2$	1	0.00 ± 0.00	0.00	0.00 ± 0.00
$AcNH_2$	$AcNH_2 \cdots Uracil$	1	0.10 ± 0.05	0.06	-0.04 ± 0.05
$AcNH_2$	Benzene···AcNH ₂ (NH- π)	2	-0.68 ± 0.04	-0.65	0.03 ± 0.04
$AcNH_2$	$AcNH_2 \cdots AcNH_2$	2	-0.07 ± 0.05	-0.00	0.07 ± 0.05
AcOH	$AcOH \cdots AcOH$	2	0.07 ± 0.05	-0.00	-0.07 ± 0.05
AcOH	$Benzene \cdots AcOH$	2	-1.18 ± 0.04	-1.29	-0.12 ± 0.04
AcOH	Ethyne···AcOH (OH- π)	2	-1.18 ± 0.06	-1.25	-0.07 ± 0.06
AcOH	AcOH···Uracil	1	0.15 ± 0.06	0.07	-0.08 ± 0.06
AcOH	Benzene···AcOH (OH- π)	2	-1.18 ± 0.05	-1.28	-0.10 ± 0.05
AcOH	AcOH···AcOH	1	0.00 ± 0.00	0.00	0.00 ± 0.00
AcOH	$Pentane \cdots AcOH$	2	-1.27 ± 0.05	-1.33	-0.06 ± 0.05
Cyclopentane	$Cyclopentane \cdots Cyclopentane$	1	0.06 ± 0.05	0.01	-0.05 ± 0.05
Cyclopentane	$Cyclopentane \cdots Cyclopentane$	2	0.07 ± 0.05	0.01	-0.06 ± 0.05
Cyclopentane	$Cyclopentane \cdots Neopentane$	1	0.00 ± 0.00	0.00	0.00 ± 0.00
Cyclopentane	Uracil···Cyclopentane	2	0.08 ± 0.05	0.02	-0.06 ± 0.05
Cyclopentane	$Benzene \cdots Cyclopentane$	2	0.08 ± 0.05	0.01	-0.07 ± 0.05
Peptide	$Peptide \cdots Peptide$	1	0.18 ± 0.05	0.16	-0.02 ± 0.05
Peptide	$Peptide \cdots Ethene$	1	0.00 ± 0.05	-0.03	-0.03 ± 0.05
Peptide	$\tilde{Water} \cdots Peptide$	2	0.08 ± 0.05	0.07	-0.02 ± 0.05
Peptide	$Peptide \cdots Peptide$	2	0.11 ± 0.05	0.08	-0.03 ± 0.05
Peptide	Peptide · · · MeOH	1	-0.05 ± 0.06	-0.05	-0.00 ± 0.06
Peptide	$MeOH \cdots Peptide$	2	0.11 ± 0.05	-0.01	-0.11 ± 0.05
Peptide	$\operatorname{Peptide} \cdots \operatorname{Water}$	1	0.07 ± 0.05	0.05	-0.02 ± 0.05
Peptide	$\operatorname{Peptide} \cdots \operatorname{MeNH}_2$	1	0.14 ± 0.05	0.08	-0.06 ± 0.05
Peptide	Peptide · · · Pentane	1	0.06 ± 0.05	-0.01	-0.07 ± 0.05
Peptide	$MeNH_2 \cdots Peptide$	2	-0.03 ± 0.05	-0.10	-0.07 ± 0.05
Peptide	Benzene · · · Peptide (NH- π)	2	0.00 ± 0.00	0.00	0.00 ± 0.00
Uracil	Benzene \cdots Uracil $(\pi - \pi)$	2	-0.41 ± 0.05	-0.31	0.10 ± 0.05
Uracil	Uracil···Ethyne	1	-0.30 ± 0.06	-0.25	0.06 ± 0.06
Uracil	$AcOH \cdots Uracil$	2	0.33 ± 0.06	0.39	0.06 ± 0.06
Uracil	$\text{Uracil} \cdots \text{Uracil} (BP)$	2	0.18 ± 0.06	0.23	0.05 ± 0.06
Uracil	Uracil · · · Uracil $(\pi - \pi)$	2	-0.02 ± 0.06	0.00	0.02 ± 0.06
Uracil	$AcNH_2\cdots Uracil$	2	0.49 ± 0.06	0.50	0.02 ± 0.06
Uracil	Uracil··· Uracil $(\pi$ - π)	1	0.00 ± 0.00	0.00	0.00 ± 0.00
Uracil	$\text{Uracil} \cdots \text{Uracil} (BP)$	1	0.25 ± 0.06	0.35	0.10 ± 0.06
Uracil	$\mathrm{Uracil}\cdots\mathrm{Ethene}$	1	-0.43 ± 0.05	-0.32	0.12 ± 0.05
Uracil	Pyridine \cdots Uracil $(\pi - \pi)$	2	-0.34 ± 0.06	-0.24	0.10 ± 0.06
Uracil	$Uracil \cdots Pentane$	1	-0.34 ± 0.05	-0.34	0.00 ± 0.05
Uracil	$\text{Uracil}\cdots\text{Cyclopentane}^{S-1}$	11 1	-0.42 ± 0.06	-0.35	0.07 ± 0.06
Uracil	$Uracil \cdots Neopentane$	1	-0.36 ± 0.05	-0.31	0.05 ± 0.05

S2.2 Setup for the CCSD(T) calculations used to evaluate the deformation energy

The deformation energy, appearing in Eq. 2, had been estimated using the Orca program system^{S2} version 4.2.1. In particular, we performed Domain-Based Local Pair Natural Orbital Coupled Cluster with Single, Double, and Perturbative Triple excitations, ^{S3,S4} or DLPNO-CCSD(T), calculations. We used Dunning's correlation consistent polarized valence triple-zeta (cc-pVTZ) and quadruple-zeta (cc-pVQZ) basis sets, and we extrapolated the complete basis set limit independently for the self-consistent field energy, with the scheme defined in Ref. S5, and for the correlation energy, with the scheme defined in Ref. S7.



Figure S2: The time step dependence of the $AcNH_2$ monomer in the $AcNH_2 \cdots AcNH_2$ dimer (ID 21) geometry.



Figure S3: The time step dependence of the AcOH monomer in the AcOH $\cdot \cdot \cdot$ AcOH dimer (ID 20) geometry.



Figure S4: The time step dependence of the Benzene monomer in the Benzene \cdots Benzene $(\pi - \pi)$ dimer (ID 24) geometry.



Figure S5: The time step dependence of the Cyclopentane monomer in the Cyclopentane \cdots Neopentane dimer (ID 37) geometry.



Figure S6: The time step dependence of the Ethene monomer in the Benzene \cdots Ethene dimer (ID 30) geometry.



Figure S7: The time step dependence of the Ethyne monomer in the Uracil \cdots Ethyne dimer (ID 32) geometry.



Figure S8: The time step dependence of the MeNH₂ monomer in the Benzene···MeNH₂ (NH- π) dimer (ID 56) geometry.



Figure S9: The time step dependence of the MeOH monomer in the Benzene··· MeOH $(OH-\pi)$ dimer (ID 55) geometry.



Figure S10: The time step dependence of the Neopentane monomer in the Neopentane \cdots Neopentane dimer (ID 36) geometry.



Figure S11: The time step dependence of the Pentane monomer in the Pentane \cdots Pentane dimer (ID 34) geometry.


Figure S12: The time step dependence of the Peptide monomer in the Benzene···Peptide (NH- π) dimer (ID 57) geometry.



Figure S13: The time step dependence of the Pyridine monomer in the Pyridine \cdots Pyridine $(\pi - \pi)$ dimer (ID 25) geometry.



Figure S14: The time step dependence of the Uracil monomer in the Uracil \cdots Uracil $(\pi - \pi)$ dimer (ID 26) geometry.



Figure S15: The time step dependence of the Water monomer in the Water \cdots Water dimer (ID 1) geometry.

S2.3 Brief summary

In this section, we briefly describe the fixed node DMC algorithm and the main factors affecting its accuracy within practical calculations.

Fixed-node DMC, is a stochastic projector method for solving the imaginary-time manybody Schrodinger equation, where a trial many-electron wave-function $\Psi_{\rm T}(\mathbf{R})$, where \mathbf{R} is the electronic configuration, is chosen and used to define a trial many-electron nodal surface (the hyper-surface where $\Psi_{\rm T}(\mathbf{R}) = 0$). With the given nodal surface, fixed-node DMC will project out the lowest-energy many-electron state. The trial wave-function has a critical role in determining the accuracy of fixed-node DMC. The trial wave-function is the product $\Psi_{\rm T}(\mathbf{R}) = \mathcal{D}(\mathbf{R}) * \exp{\{\mathcal{J}(\mathbf{R})\}}$ of an antisymmetric function $\mathcal{D}(\mathbf{R})$ and a symmetric (bosonic) function $\exp{\{\mathcal{J}(\mathbf{R})\}}$, called the Jastrow factor, describing the dynamical correlation between the electrons by including explicit functions of the electron-electron distances and electronnucleus distances. The common practice for the Jastrow factor is to decide a functional form for \mathcal{J} and optimize its parameters by minimizing either the energy or the variance, using the variational Monte Carlo (VMC)^{S8} scheme. The stochastic optimization of the Jastrow factor implies an optimization uncertainty on its parameters.

When dealing with large systems (> 100 atoms), fixed-node DMC employs pseudopotentials to substantially improve its efficiency. To deal with nonlocal terms of pseudopotentials, the fixed-node DMC algorithm must use an additional approximation, leading to the socalled localization error. The first approximation to solve the localization error consisted of "localizing" nonlocal pseudopotential operators using the trial wave function, ^{S9} or part of the wave function. ^{S10} Currently there are four schemes based on this approach: the locality approximation (LA), ^{S9} the T-move (TM), ^{S11,S12} the determinant locality approximation (DLA), ^{S13} and the determinant locality T-move (DTM). ^{S13} Here, we use the DLA scheme, and report additional tests against TM and DTM in Table S4.

		$\Delta E_{\rm int.}$ [kcal/mol]
$H2O \cdots H2O$	TM	-5.06 ± 0.04
	DLA	-5.17 ± 0.03
AcOH···AcOH	TM	-19.98 ± 0.06
	DLA	-20.16 ± 0.11
	DTM	-20.30 ± 0.09

Table S4: Comparison of the extrapolated interaction energy $\Delta E_{\text{int.}}$ for the TM and DLA localization schemes for the H₂O···H₂O (ID 1) and AcOH···AcOH dimers (ID 20).

S2.4 Reaching the time step limit and estimating errors

A key aspect affecting the accuracy of DMC is the simulation time step. In fact, as mentioned above, in DMC a propagation according to the imaginary time Schrödinger equation is performed to project out the exact ground state from a trial wave-function. A time step τ must be chosen, but the projection is exact only in the continuous limit $\tau \to 0$. The bias due to the finite time step is usually called the time step error. We note that the time step dependence can be affected by the chosen Jastrow, the trial wave-function as well as the algorithm used to perform the walker propagation. Many such algorithms exist and we use the ZSGMA^{S14,S15} DMC algorithm.

We extrapolate to the zero time step limit $(\tau \rightarrow 0)$ using a set of time steps (0.1, 0.08, 0.06, 0.05, 0.04, 0.03, 0.02, 0.01 and 0.003 au). In calculating any of the energy terms, for all the time steps up until and including 0.1 au, we fit a curve to a cubic polynomial of the form:

$$E(\tau) = A + B\tau + C\tau^2 + D\tau^3, \tag{3}$$

where A, B, C, and D are fit parameters, with A being the value in the limit of $\tau \to 0$. Here E can be either a total energy (i.e., $E_{\text{dimer}}^{\text{DMC}}$, $E_{\text{mon. 1}}^{\text{DMC}}$ and $E_{\text{mon. 2}}^{\text{DMC}}$) or an interaction energy ΔE_{int} . For the two smaller time steps (0.01 and 0.003 au), we also fit a (linear) line.

We use the zero time step estimate from the cubic fit for all of our estimates. For the majority of systems, we set the error on this estimate to the predicted standard deviation of the cubic fit at zero time step: $\sigma_{\text{cubic fit}}$. However, for some systems, the predicted zero

time step prediction with the linear fit can be outside the standard deviation of the cubic fit, indicating that there is a significant change in behavior at smaller time steps. Here, we instead estimate the error as the difference in the zero time step prediction between the linear and cubic fits: $\Delta_{\text{cubic fit}}^{\text{linear fit}}$

S3 CCSD(T) estimates from the literature

Reference values at the CCSD(T) level for the S66 dataset has been computed within several studies. The largest dimer of S66 dataset – the uracil dimer – can reach up to 34 atoms and 116 electrons (of which 84 are valence). This can pose considerable difficulty with performing CCSD(T) on the calculations, particularly when aiming to reach the complete basis set (CBS) limit. For example, the uracil dimer with a saug-ano-pVQZ basis set calculation 'took eight days wall clock time running in parallel on 96 CPUs with a total of 1.5 TB RAM and 18 TB of solid state scratch disk'. In fact, reaching accurate estimates at the CBS limit requires going beyond the above quadruple- ζ (QZ) basis set. As such, the previous studies take on various composite schemes to approximate the CBS limit, performing larger basis set or extrapolated calculations for the MP2 or CCSD contribution to the binding energy, with the remaining contributions to CCSD(T) performed at a smaller basis set. In particular, they exploit the decomposition of the CCSD(T) binding energy into a Hartree-Fock (HF) component $\Delta E_{\rm HF}$, a MP2 correlation component $\Delta E_{\rm MP2}$, a CCSD correction to MP2 $\Delta E_{\rm CCSD-MP2}$ alongside a final (T) contribution $\Delta E_{\rm (T)}$:

$$\Delta E_{\rm int}^{\rm CCSD(T)} = \Delta E_{\rm int}^{\rm HF} + \Delta E_{\rm int}^{\rm MP2} + \Delta E_{\rm int}^{\rm CCSD-MP2} + \Delta E_{\rm int}^{\rm (T)}.$$
(4)

Each of these components has a differing dependence on the basis set, with higher order correlations [i.e., (T)] typically requiring smaller basis sets.

The most common class of basis sets used to treat non-covalent interactions are the Dunning cc-pVXZ and aug-cc-pVXZ (augmented with diffuse functions) basis, which we

will refer to as XZ and aXZ respectively. In many studies, The XZ basis sets are used on the H atoms, with aXZ on the remaining elements, leading to the heavy-aug-cc-pVXZ basis set, shortened to haXZ. The cardinal number X can be either a double- ζ (DZ), triple- ζ (TZ), quadruple- ζ (QZ) or quintuple- ζ (5Z) basis set, in order of increasing basis set size. Adjacent pairs of basis sets can be combined within a two-point extrapolation scheme to approximate the CBS limit, which we shall indicate with CBS(DZ/TZ) for the DZ and TZ pair. Additionally, there is the choice of employing a counterpoise correction, where within the calculation of Δ_{int} , the energies of the individual monomers are computed together with 'ghost' basis functions from the other monomer, removing some errors arising from basis-set superposition error. Calculations can either employ no counterpoise correction (no-CP), counterpoise correction (CP) or the average of the two (half-CP). Regardless all these estimates should reach the same value in the CBS limit.

In this work, we reach a final CCSD(T) estimate which takes the average of CCSD(T) estimates from three separate studies, all of which approximate the CCSD(T) CBS limit with differing treatments. These differences are summarized below:

- Řezáč et al.^{S16} The S66 dataset and its first CCSD(T) estimates were introduced by Řezáč et al. in Ref. S1, and these CCSD(T) estimates were subsequently revised and improved in Ref. S16. Here, both ΔE_{int}^{HF} and ΔE_{int}^{MP2} were computed with a two-point extrapolation using the aTZ and aQZ basis sets [i.e., CBS(aTZ/aQZ)]. The remaining $\Delta E_{int}^{CCSD-MP2}$ and $\Delta E_{int}^{(T)}$ contributions were computed with a CBS(haDZ/haTZ) treatment. All contributions utilised CP corrections.
- Kesharwani *et al.*^{S17} Kesharwani *et al.* re-evaluated the S66 dataset using explicitly correlated F12-based methods. They came up with four different tiers which trade accuracy for cost, namely the 'GOLD', 'SILVER', 'BRONZE' and 'STERLING' levels. As GOLD was only feasible for a subset of 18 systems, we focus here on the SILVER estimates, which were computed for the entire dataset. The ΔE_{int}^{HF} and ΔE_{int}^{MP2} components were computed using MP2-F12 with the cc-pV5Z-F12 basis set, with further

(minor) corrections to the HF treatment using a complementary auxiliary basis set (CABS) treatment. The $\Delta E_{\text{int}}^{\text{CCSD-MP2}}$ contribution was computed with CCSD(F12*) employing the aug-cc-pVTZ-F12 basis set. The final $\Delta E_{\text{int}}^{(\text{T})}$ contribution did not employ any F12 treatment and was reached using a CBS(haDZ/haTZ) extrapolation. All contributions utilised a half-CP correction.

• Nagy et al.^{S18} — Nagy et al. reported an improvement upon the previous S66 CCSD(T) references – termed '14k-GOLD'. Here, $\Delta E_{\rm int}^{\rm HF}$ was treated with using the aQZ-F12 basis set together with a CABS treatment, while $\Delta E_{\rm int}^{\rm MP2}$ was treated using a CBS(aTZ-F12/aQZ-F12) two-point extrapolation. The $\Delta E_{\rm int}^{\rm CCSD-MP2}$ contribution was computed with CCSD(F12*) and MP2-F12 using a CBS(haDZ-F12/haTZ-F12) extrapolation. The final $\Delta E_{\rm int}^{\rm (T)}$ contribution did not employ any F12 treatment and was reached using a CBS(haTZ/haQZ) extrapolation. The $\Delta E_{\rm int}^{\rm HF}$ was performed with half-CP while all other contributions were performed with (full) CP correction.

Table S5: CCSD(T) references for the S66 dataset. The final CCSD(T) and CCSD(cT)-fit values are computed as the average of the values from the three references. The error is computed as twice the standard deviation of the values from the three references.

	System	Řezáč et al. (2006)	Kesharwani <i>et al.</i> (2018)	Nagy et al. (2023)	Final CCSD(T)	Final CCSD(cT)-fit
1	$Water \cdots Water$	-5.01	-4.98	-4.99	-4.99 ± 0.01	-4.96 ± 0.01
2	$Water \cdots MeOH$	-5.70	-5.67	-5.67	-5.68 ± 0.01	-5.63 ± 0.01
3	$Water \cdots MeNH_2$	-7.04	-6.99	-7.00	-7.01 ± 0.02	-6.94 ± 0.02
4	$Water \cdots Peptide$	-8.22	-8.18	-8.19	-8.20 ± 0.02	-8.15 ± 0.02

Table S5: (continued)

5	${\rm MeOH} \cdots {\rm MeOH}$	-5.85	-5.82	-5.83	-5.83 ± 0.01	-5.78 ± 0.01
6	$\mathrm{MeOH} \cdots \mathrm{MeNH}_2$	-7.67	-7.62	-7.62	-7.64 ± 0.02	-7.55 ± 0.02
7	$MeOH \cdots Peptide$	-8.34	-8.31	-8.31	-8.32 ± 0.01	-8.25 ± 0.01
8	$MeOH \cdots Water$	-5.09	-5.06	-5.07	-5.08 ± 0.01	-5.03 ± 0.01
9	$\mathrm{MeNH}_2\!\cdots\!\mathrm{MeOH}$	-3.11	-3.09	-3.09	-3.10 ± 0.01	-3.05 ± 0.01
10	$\mathrm{MeNH}_2\!\cdots\mathrm{MeNH}_2$	-4.22	-4.18	-4.19	-4.20 ± 0.01	-4.13 ± 0.01
11	$MeNH_2 \cdots Peptide$	-5.48	-5.44	-5.44	-5.45 ± 0.02	-5.37 ± 0.02
12	$\mathrm{MeNH}_2\!\cdots\!\mathrm{Water}$	-7.40	-7.35	-7.36	-7.37 ± 0.02	-7.29 ± 0.02
13	$\operatorname{Peptide}\cdots\operatorname{MeOH}$	-6.28	-6.25	-6.25	-6.26 ± 0.01	-6.20 ± 0.01
14	$\operatorname{Peptide}\cdots\operatorname{MeNH}_2$	-7.56	-7.52	-7.52	-7.53 ± 0.02	-7.44 ± 0.02
15	$\operatorname{Peptide}\cdots\operatorname{Peptide}$	-8.72	-8.69	-8.69	-8.70 ± 0.01	-8.61 ± 0.01
16	$\operatorname{Peptide}\cdots\operatorname{Water}$	-5.20	-5.18	-5.18	-5.19 ± 0.01	-5.15 ± 0.01
17	$\text{Uracil}\cdots\text{Uracil}(\text{BP})$	-17.45	-17.41	-17.40	-17.42 ± 0.02	-17.29 ± 0.02
18	$Water \cdots Pyridine$	-6.97	-6.93	-6.93	-6.94 ± 0.02	-6.87 ± 0.02
19	$MeOH \cdots Pyridine$	-7.51	-7.47	-7.46	-7.48 ± 0.02	-7.39 ± 0.02
20	$AcOH \cdots AcOH$	-19.41	-19.36	-19.38	-19.39 ± 0.02	-19.27 ± 0.02
21	$AcNH_2 \cdots AcNH_2$	-16.52	-16.47	-16.48	-16.49 ± 0.02	-16.40 ± 0.02
22	$AcOH \cdots Uracil$	-19.78	-19.74	-19.75	-19.75 ± 0.02	-19.64 ± 0.02
23	$AcNH_2 \cdots Uracil$	-19.47	-19.42	-19.42	-19.44 ± 0.02	-19.33 ± 0.02
24	Benzene···Benzene $(\pi$ - π)	-2.72	-2.68	-2.69	-2.70 ± 0.02	-2.46 ± 0.02
25	Pyridine \cdots Pyridine $(\pi - \pi)$	-3.80	-3.75	-3.76	-3.77 ± 0.02	-3.51 ± 0.02
26	Uracil···Uracil $(\pi$ - π)	-9.75	-9.67	-9.72	-9.71 ± 0.03	-9.39 ± 0.03
27	Benzene···Pyridine $(\pi - \pi)$	-3.34	-3.30	-3.30	-3.31 ± 0.02	-3.07 ± 0.02
28	Benzene··· Uracil $(\pi - \pi)$	-5.59	-5.52	-5.54	-5.55 ± 0.03	-5.25 ± 0.03
29	Pyridine····Uracil $(\pi$ - π)	-6.70	-6.63	-6.66	-6.66 ± 0.03	-6.37±0.03

Table S5: (continued)

30	$Benzene \cdots Ethene$	-1.36	-1.36	-1.34	-1.35 ± 0.01	-1.24 ± 0.01
31	$\mathrm{Uracil}\cdots\mathrm{Ethene}$	-3.33	-3.29	-3.31	-3.31 ± 0.02	-3.17 ± 0.02
32	$\mathrm{Uracil}\cdots\mathrm{Ethyne}$	-3.69	-3.65	-3.68	-3.67 ± 0.02	-3.54 ± 0.02
33	$Pyridine \cdots Ethene$	-1.80	-1.78	-1.78	-1.79 ± 0.01	-1.66 ± 0.01
34	$Pentane \cdots Pentane$	-3.76	-3.74	-3.73	-3.74 ± 0.01	-3.63 ± 0.01
35	$Neopentane \cdots Pentane$	-2.60	-2.58	-2.58	-2.59 ± 0.01	-2.50 ± 0.01
36	$Neopentane \cdots Neopentane$	-1.76	-1.74	-1.75	-1.75 ± 0.01	-1.69 ± 0.01
37	$Cyclopentane \cdots Neopentane$	-2.40	-2.38	-2.38	-2.38 ± 0.01	-2.30 ± 0.01
38	$Cyclopentane \cdots Cyclopentane$	-2.99	-2.97	-2.96	-2.97 ± 0.01	-2.87 ± 0.01
39	$Benzene \cdots Cyclopentane$	-3.51	-3.49	-3.48	-3.49 ± 0.01	-3.33 ± 0.01
40	$Benzene \cdots Neopentane$	-2.85	-2.82	-2.82	-2.83 ± 0.01	-2.71 ± 0.01
41	$\mathrm{Uracil} \cdots \mathrm{Pentane}$	-4.81	-4.76	-4.77	-4.78 ± 0.02	-4.59 ± 0.02
42	${\rm Uracil} \cdots {\rm Cyclopentane}$	-4.09	-4.05	-4.06	-4.07 ± 0.02	-3.90 ± 0.02
43	$\mathrm{Uracil}\cdots\mathrm{Neopentane}$	-3.69	-3.65	-3.66	-3.67 ± 0.02	-3.53 ± 0.02
44	$Ethene \cdots Pentane$	-1.99	-1.97	-1.98	-1.98 ± 0.01	-1.91 ± 0.01
45	$Ethyne \cdots Pentane$	-1.72	-1.70	-1.70	-1.71 ± 0.01	-1.63 ± 0.01
46	$\operatorname{Peptide}\cdots\operatorname{Pentane}$	-4.26	-4.22	-4.22	-4.23 ± 0.02	-4.09 ± 0.02
47	Benzene \cdots Benzene (TS)	-2.83	-2.80	-2.81	-2.81 ± 0.01	-2.68 ± 0.01
48	$Pyridine \cdots Pyridine (TS)$	-3.51	-3.47	-3.48	-3.49 ± 0.02	-3.35 ± 0.02
49	Benzene \cdots Pyridine (TS)	-3.29	-3.26	-3.27	-3.27 ± 0.01	-3.15 ± 0.01
50	Benzene···Ethyne (CH- π)	-2.86	-2.83	-2.84	-2.84 ± 0.01	-2.76 ± 0.01
51	Ethyne \cdots Ethyne (TS)	-1.54	-1.52	-1.53	-1.53 ± 0.01	-1.49 ± 0.01
52	Benzene···AcOH (OH- π)	-4.73	-4.69	-4.69	-4.70 ± 0.02	-4.59 ± 0.02
53	Benzene···AcNH ₂ (NH- π)	-4.40	-4.38	-4.38	-4.38 ± 0.01	-4.29 ± 0.01
54	Benzene··· Water (OH- π)	-3.29	-3.27	-3.26	-3.27 ± 0.01	-3.21 ± 0.01

Table S5: (continued)

55	Benzene··· MeOH (OH- π)	-4.17	-4.14	-4.14	-4.15 ± 0.02	-4.04 ± 0.02
56	Benzene···MeNH ₂ (NH- π)	-3.20	-3.17	-3.17	-3.18 ± 0.01	-3.07 ± 0.01
57	Benzene···Peptide (NH- π)	-5.26	-5.22	-5.22	-5.23 ± 0.02	-5.08 ± 0.02
58	$Pyridine \cdots Pyridine (CH-N)$	-4.24	-4.19	-4.19	-4.21 ± 0.02	-4.12 ± 0.02
59	Ethyne··· Water (CH-O)	-2.93	-2.90	-2.91	-2.92 ± 0.01	-2.89 ± 0.01
60	Ethyne····AcOH (OH- π)	-4.97	-4.92	-4.93	$-4.94{\pm}0.02$	-4.86 ± 0.02
61	$Pentane \cdots AcOH$	-2.91	-2.88	-2.88	-2.89 ± 0.02	-2.79 ± 0.02
62	$\mathrm{Pentane} \cdots \mathrm{AcNH}_2$	-3.53	-3.49	-3.50	-3.51 ± 0.02	-3.39 ± 0.02
63	$Benzene \cdots AcOH$	-3.75	-3.71	-3.72	-3.72 ± 0.02	-3.59 ± 0.02
64	$\operatorname{Peptide}\cdots\operatorname{Ethene}$	-3.00	-2.97	-2.98	-2.98 ± 0.01	-2.90 ± 0.01
65	$Pyridine \cdots Ethyne$	-4.10	-4.06	-4.07	-4.08 ± 0.02	-4.02 ± 0.02
66	$MeNH_2 \cdots Pyridine$	-3.97	-3.93	-3.93	$-3.94{\pm}0.02$	-3.83 ± 0.02

S4 Final CCSD(T) and CCSD(cT)-fit estimates

As discussed above, in this work, we arrive at a final estimate of CCSD (T) taking the average of the CCSD(T) estimates from Kesharwani *et al.*, Řezáč *et al.* and Nagy *et al.*. The resulting error bars are the standard deviation of the three references. The final CCSD(T) estimates are reported in Table S5. In addition, we also give an estimate at the CCSD(cT) for the dispersion-dominated systems. CCSD(cT) incorporates additional higher-order terms to the triples excitation amplitudes compared CCSD(T), crucial for studying systems with large polarizability. In particular, it has been shown that CCSD(cT) can be approximated from the Δ_{int}^{MP2} , $\Delta_{int}^{CCSD-MP2}$ and $\Delta_{int}^{(T)}$ values. Specifically, it is given by the following expression:

$$\frac{\Delta_{\rm int}^{\rm (T)}}{\Delta_{\rm int}^{\rm (cT)}} = a + b \cdot \frac{\Delta_{\rm int}^{\rm MP2}}{\Delta_{\rm int}^{\rm CCSD}},\tag{5}$$

where a = 0.7764 and b = 0.2780 were fitted to CCSD(cT) data.

S5 Final DMC estimates

We report the final DMC estimates in Table S6. This calculates the interaction energy as given in Eq. 1 with deformation energies computed as in Eq. 2 and Table S2. We note that the extrapolation towards the zero time step limit is performed directly on ΔE_{int} rather than the individual total energy components. In Table S7, we have computed the deviation of these final estimates to the CCSD(T) estimates in Table S5.

Table S6: Final DMC ΔE_{int} estimates for the S66 dataset. The method (see Sec. S2.4) used to estimate the error on the DMC estimate is also provided.

	System	$\Delta E_{\rm int} \; [\rm kcal/mol]$	Fit type
1	Water···Water	-5.17 ± 0.03	$\sigma_{ m cubic\ fit}$
2	$Water \cdots MeOH$	-5.82 ± 0.04	$\sigma_{ m cubic\ fit}$
3	$Water \cdots MeNH_2$	-7.18 ± 0.04	$\sigma_{ m cubic\ fit}$
4	$Water \cdots Peptide$	-8.59 ± 0.06	$\sigma_{ m cubic\ fit}$
5	$MeOH \cdots MeOH$	-5.93 ± 0.10	$\Delta_{\rm cubic\ fit}^{\rm linear\ fit}$
6	$\mathrm{MeOH} \cdots \mathrm{MeNH}_2$	-7.83 ± 0.07	$\Delta_{\rm cubic\ fit}^{\rm linear\ fit}$
7	$MeOH \cdots Peptide$	-8.57 ± 0.07	$\sigma_{ m cubic\ fit}$
8	$MeOH \cdots Water$	-5.24 ± 0.08	$\Delta_{\rm cubic\ fit}^{\rm linear\ fit}$
9	$MeNH_2\cdots MeOH$	-3.12 ± 0.08	$\Delta_{\rm cubic\ fit}^{\rm linear\ fit}$
10	$MeNH_2\cdots MeNH_2$	-4.20 ± 0.10	$\Delta_{\rm cubic\ fit}^{\rm linear\ fit}$
11	$MeNH_2\cdots Peptide$	-5.42 ± 0.12	$\Delta_{\rm cubic\ fit}^{\rm linear\ fit}$
12	$MeNH_2 \cdots Water$	-7.53 ± 0.05	$\sigma_{ m cubic\ fit}$
13	$\operatorname{Peptide}\cdots\operatorname{MeOH}$	-6.32 ± 0.07	$\Delta_{\rm cubic\ fit}^{\rm linear\ fit}$
14	$\operatorname{Peptide}\cdots\operatorname{MeNH}_2$	-7.50 ± 0.10	$\Delta_{\rm cubic\ fit}^{\rm linear\ fit}$
		Continued on a	next page

Table S6: (continued)

15	$\operatorname{Peptide}\cdots\operatorname{Peptide}$	-8.88 ± 0.08	$\sigma_{ m cubic\ fit}$
16	$Peptide \cdots Water$	-5.37 ± 0.06	$\sigma_{ m cubic\ fit}$
17	$\text{Uracil}\cdots\text{Uracil}$ (BP)	-17.79 ± 0.26	$\Delta^{\rm linear fit}_{\rm cubic fit}$
18	$Water \cdots Pyridine$	-7.30 ± 0.09	$\Delta_{\rm cubic\ fit}^{\rm linear\ fit}$
19	$MeOH \cdots Pyridine$	-7.88 ± 0.07	$\sigma_{ m cubic\ fit}$
20	$AcOH \cdots AcOH$	-20.17 ± 0.11	$\Delta^{\rm linear \ fit}_{\rm cubic \ fit}$
21	$AcNH_2 \cdots AcNH_2$	-16.83 ± 0.07	$\sigma_{ m cubic\ fit}$
22	$AcOH \cdots Uracil$	-20.40 ± 0.09	$\Delta^{\rm linear \ fit}_{\rm cubic \ fit}$
23	$AcNH_2\cdots Uracil$	-19.82 ± 0.09	$\sigma_{ m cubic\ fit}$
24	Benzene···Benzene $(\pi$ - π)	-2.33 ± 0.08	$\sigma_{ m cubic\ fit}$
25	Pyridine \cdots Pyridine $(\pi - \pi)$	-3.53 ± 0.07	$\sigma_{ m cubic\ fit}$
26	Uracil···Uracil $(\pi$ - π)	-9.33 ± 0.08	$\sigma_{ m cubic\ fit}$
27	Benzene···Pyridine $(\pi - \pi)$	-3.04 ± 0.08	$\sigma_{ m cubic\ fit}$
28	Benzene··· Uracil $(\pi$ - π)	-5.15 ± 0.11	$\Delta^{\rm linear \ fit}_{\rm cubic \ fit}$
29	Pyridine · · · Uracil $(\pi$ - π)	-6.40 ± 0.17	$\Delta^{\rm linear \ fit}_{\rm cubic \ fit}$
30	$Benzene \cdots Ethene$	-1.11 ± 0.06	$\sigma_{ m cubic\ fit}$
31	$\mathrm{Uracil}\cdots\mathrm{Ethene}$	-3.18 ± 0.09	$\Delta^{\rm linear fit}_{\rm cubic fit}$
32	$\mathrm{Uracil}\cdots\mathrm{Ethyne}$	-3.59 ± 0.08	$\sigma_{ m cubic\ fit}$
33	$Pyridine \cdots Ethene$	-1.69 ± 0.07	$\sigma_{ m cubic\ fit}$
34	$Pentane \cdots Pentane$	-3.53 ± 0.07	$\sigma_{ m cubic\ fit}$
35	$Neopentane \cdots Pentane$	-2.46 ± 0.08	$\sigma_{ m cubic\ fit}$
36	$Neopentane \cdots Neopentane$	-1.67 ± 0.08	$\sigma_{ m cubic\ fit}$
37	$Cyclopentane \cdots Neopentane$	-2.17 ± 0.07	$\sigma_{ m cubic\ fit}$
38	$Cyclopentane \cdots Cyclopentane$	-2.75 ± 0.07	$\sigma_{ m cubic\ fit}$
39	$Benzene \cdots Cyclopentane$	-3.14 ± 0.08	$\sigma_{ m cubic\ fit}$
		Continued on a	next page

Table S6: (continued)

40	$Benzene \cdots Neopentane$	-2.68 ± 0.08	$\Delta_{\text{cubic fit}}^{\text{linear fit}}$
41	$\mathrm{Uracil}\cdots\mathrm{Pentane}$	-4.46 ± 0.09	$\sigma_{ m cubic\ fit}$
42	$\mathrm{Uracil}\cdots\mathrm{Cyclopentane}$	-3.59 ± 0.09	$\sigma_{ m cubic\ fit}$
43	$\mathrm{Uracil}\cdots\mathrm{Neopentane}$	-3.51 ± 0.16	$\Delta^{\rm linear \ fit}_{\rm cubic \ fit}$
44	$Ethene \cdots Pentane$	-1.79 ± 0.06	$\sigma_{ m cubic\ fit}$
45	$Ethyne \cdots Pentane$	-1.57 ± 0.06	$\Delta^{\rm linear \ fit}_{\rm cubic \ fit}$
46	$\operatorname{Peptide} \cdots \operatorname{Pentane}$	-3.82 ± 0.11	$\Delta^{\rm linear \ fit}_{\rm cubic \ fit}$
47	Benzene \cdots Benzene (TS)	-2.61 ± 0.08	$\sigma_{ m cubic\ fit}$
48	$Pyridine \cdots Pyridine (TS)$	-3.44 ± 0.08	$\sigma_{ m cubic\ fit}$
49	Benzene \cdots Pyridine (TS)	-3.11 ± 0.08	$\sigma_{ m cubic\ fit}$
50	Benzene···Ethyne (CH- π)	-2.90 ± 0.06	$\sigma_{ m cubic\ fit}$
51	Ethyne \cdots Ethyne (TS)	-1.54 ± 0.08	$\Delta^{\rm linear \ fit}_{\rm cubic \ fit}$
52	Benzene···AcOH (OH- π)	-4.64 ± 0.16	$\Delta^{\rm linear \ fit}_{\rm cubic \ fit}$
53	Benzene···AcNH ₂ (NH- π)	-4.25 ± 0.08	$\sigma_{ m cubic\ fit}$
54	Benzene···Water (OH- π)	-3.20 ± 0.06	$\sigma_{ m cubic\ fit}$
55	Benzene····MeOH (OH- π)	-3.93 ± 0.07	$\sigma_{ m cubic\ fit}$
56	Benzene···MeNH ₂ (NH- π)	-3.03 ± 0.07	$\sigma_{ m cubic\ fit}$
57	Benzene···Peptide (NH- π)	-5.10 ± 0.10	$\Delta^{\rm linear \ fit}_{\rm cubic \ fit}$
58	$Pyridine \cdots Pyridine (CH-N)$	-4.22 ± 0.22	$\Delta^{\rm linear \ fit}_{\rm cubic \ fit}$
59	Ethyne··· Water (CH-O)	-3.04 ± 0.04	$\sigma_{ m cubic\ fit}$
60	Ethyne··· AcOH (OH- π)	-4.98 ± 0.07	$\Delta^{\rm linear \ fit}_{\rm cubic \ fit}$
61	$Pentane \cdots AcOH$	-2.63 ± 0.18	$\Delta^{\rm linear \ fit}_{\rm cubic \ fit}$
62	$Pentane \cdots AcNH_2$	-3.08 ± 0.07	$\sigma_{ m cubic\ fit}$
63	$Benzene \cdots AcOH$	-3.52 ± 0.08	$\sigma_{ m cubic\ fit}$
64	$\operatorname{Peptide} \cdots \operatorname{Ethene}$	-2.77 ± 0.07	$\sigma_{ m cubic\ fit}$
		Continued on a	next page

Table S6: (continued)

65	$Pyridine \cdots Ethyne$	-4.26 ± 0.06	$\sigma_{ m cubic\ fit}$
66	$MeNH_2 \cdots Pyridine$	-3.81 ± 0.09	$\Delta^{\rm linear \ fit}_{\rm cubic \ fit}$

Table S7: Final DMC and CCSD(T) $\Delta E_{\text{int.}}$ estimates for the S66 dataset in kcal/mol, with their deviation of CCSD(T) from DMC given.

	System	$\Delta E_{\rm int.}^{\rm DMC}$ [kcal/mol]	$\Delta E_{\rm int.}^{\rm CCSD(T)}$ [kcal/mol]	Deviation [kcal/mol]
1	Water···Water	-5.17 ± 0.03	-4.99 ± 0.01	$0.18 {\pm} 0.03$
2	$Water \cdots MeOH$	-5.82 ± 0.04	-5.68 ± 0.01	$0.14{\pm}0.04$
3	$Water \cdots MeNH_2$	-7.18 ± 0.04	-7.01 ± 0.02	$0.18 {\pm} 0.05$
4	$Water \cdots Peptide$	-8.59 ± 0.06	-8.20 ± 0.02	$0.39{\pm}0.06$
5	$MeOH \cdots MeOH$	-5.93 ± 0.10	-5.83 ± 0.01	$0.09 {\pm} 0.10$
6	$\mathrm{MeOH} \cdots \mathrm{MeNH}_2$	-7.83 ± 0.07	-7.64 ± 0.02	$0.19{\pm}0.07$
7	$MeOH \cdots Peptide$	-8.57 ± 0.07	-8.32 ± 0.01	$0.25{\pm}0.07$
8	$MeOH \cdots Water$	-5.24 ± 0.08	-5.08 ± 0.01	$0.16{\pm}0.08$
9	$\mathrm{MeNH}_2\!\cdots\!\mathrm{MeOH}$	-3.12 ± 0.08	-3.10 ± 0.01	$0.02{\pm}0.08$
10	$\mathrm{MeNH}_2\!\cdots\!\mathrm{MeNH}_2$	-4.20 ± 0.10	-4.20 ± 0.01	$0.00 {\pm} 0.11$
11	$MeNH_2 \cdots Peptide$	-5.42 ± 0.12	-5.45 ± 0.02	-0.04 ± 0.13
12	$MeNH_2 \cdots Water$	-7.53 ± 0.05	-7.37 ± 0.02	$0.16{\pm}0.05$
13	$\operatorname{Peptide} \cdots \operatorname{MeOH}$	-6.32 ± 0.07	-6.26 ± 0.01	$0.06{\pm}0.07$
14	$\mathrm{Peptide} \cdots \mathrm{MeNH}_2$	-7.50 ± 0.10	-7.53 ± 0.02	-0.03 ± 0.10
15	$\operatorname{Peptide}\cdots\operatorname{Peptide}$	-8.88 ± 0.08	-8.70 ± 0.01	$0.18 {\pm} 0.09$
16	$\operatorname{Peptide}\cdots\operatorname{Water}$	-5.37 ± 0.06	-5.19 ± 0.01	$0.19{\pm}0.06$
17	$\text{Uracil}\cdots\text{Uracil}\ (\text{BP})$	-17.79 ± 0.26	-17.42 ± 0.02	$0.37 {\pm} 0.27$
18	$Water \cdots Pyridine$	-7.30 ± 0.09	$-6.94{\pm}0.02$	$0.35{\pm}0.09$
19	$MeOH \cdots Pyridine$	$-7.88 {\pm} 0.07$	-7.48 ± 0.02	$0.40 {\pm} 0.08$
20	$AcOH \cdots AcOH$	-20.17 ± 0.11	-19.39 ± 0.02	$0.78 {\pm} 0.11$

Table S7: (continued)

21	$AcNH_2 \cdots AcNH_2$	$-16.83 {\pm} 0.07$	-16.49 ± 0.02	$0.34 {\pm} 0.07$
22	$AcOH \cdots Uracil$	-20.40 ± 0.09	-19.75 ± 0.02	$0.65 {\pm} 0.09$
23	$AcNH_2\cdots Uracil$	-19.82 ± 0.09	-19.44 ± 0.02	$0.39 {\pm} 0.09$
24	Benzene···Benzene $(\pi - \pi)$	-2.33 ± 0.08	-2.70 ± 0.02	-0.37 ± 0.08
25	Pyridine \cdots Pyridine $(\pi - \pi)$	-3.53 ± 0.07	-3.77 ± 0.02	-0.24 ± 0.08
26	Uracil··· Uracil $(\pi$ - π)	-9.33 ± 0.08	-9.71 ± 0.03	-0.39 ± 0.09
27	Benzene···Pyridine $(\pi$ - π)	$-3.04{\pm}0.08$	-3.31 ± 0.02	-0.28 ± 0.08
28	Benzene··· Uracil $(\pi$ - π)	-5.15 ± 0.11	-5.55 ± 0.03	-0.40 ± 0.12
29	Pyridine · · · Uracil $(\pi - \pi)$	-6.40 ± 0.17	-6.66 ± 0.03	-0.27 ± 0.17
30	$Benzene \cdots Ethene$	-1.11 ± 0.06	$-1.35 {\pm} 0.01$	-0.24 ± 0.06
31	$\mathrm{Uracil}\cdots\mathrm{Ethene}$	-3.18 ± 0.09	-3.31 ± 0.02	-0.13 ± 0.09
32	$\mathrm{Uracil}\cdots\mathrm{Ethyne}$	$-3.59 {\pm} 0.08$	-3.67 ± 0.02	-0.09 ± 0.08
33	$\operatorname{Pyridine} \cdots \operatorname{Ethene}$	$-1.69 {\pm} 0.07$	$-1.79 {\pm} 0.01$	-0.10 ± 0.07
34	$Pentane \cdots Pentane$	$-3.53 {\pm} 0.07$	-3.74 ± 0.01	-0.22 ± 0.07
35	$Neopentane \cdots Pentane$	-2.46 ± 0.08	$-2.59 {\pm} 0.01$	-0.13 ± 0.08
36	$Neopentane \cdots Neopentane$	-1.67 ± 0.08	$-1.75 {\pm} 0.01$	-0.08 ± 0.08
37	$Cyclopentane \cdots Neopentane$	-2.17 ± 0.07	-2.38 ± 0.01	-0.21 ± 0.07
38	$Cyclopentane \cdots Cyclopentane$	-2.75 ± 0.07	-2.97 ± 0.01	-0.23 ± 0.07
39	$Benzene \cdots Cyclopentane$	-3.14 ± 0.08	-3.49 ± 0.01	-0.36 ± 0.08
40	$\mathrm{Benzene} \cdots \mathrm{Neopentane}$	$-2.68 {\pm} 0.08$	-2.83 ± 0.01	-0.15 ± 0.08
41	$\mathrm{Uracil}\cdots\mathrm{Pentane}$	-4.46 ± 0.09	-4.78 ± 0.02	-0.32 ± 0.09
42	$\mathrm{Uracil}\cdots\mathrm{Cyclopentane}$	-3.59 ± 0.09	-4.07 ± 0.02	-0.47 ± 0.09
43	$\mathrm{Uracil}\cdots\mathrm{Neopentane}$	-3.51 ± 0.16	-3.67 ± 0.02	-0.15 ± 0.16
44	$Ethene \cdots Pentane$	$-1.79 {\pm} 0.06$	$-1.98 {\pm} 0.01$	-0.19 ± 0.06
45	$Ethyne\cdots Pentane$	-1.57 ± 0.06	$-1.71 {\pm} 0.01$	-0.14 ± 0.07
46	$\operatorname{Peptide} \cdots \operatorname{Pentane}$	-3.82 ± 0.11	-4.23 ± 0.02	-0.41±0.11
47	Benzene \cdots Benzene (TS)	-2.61 ± 0.08	-2.81 ± 0.01	-0.20 ± 0.08

Table S7: (continued)

48	$Pyridine \cdots Pyridine (TS)$	-3.44 ± 0.08	-3.49 ± 0.02	-0.05 ± 0.08
49	Benzene \cdots Pyridine (TS)	-3.11 ± 0.08	-3.27 ± 0.01	-0.16 ± 0.08
50	Benzene···Ethyne (CH- π)	-2.90 ± 0.06	$-2.84{\pm}0.01$	$0.06{\pm}0.06$
51	Ethyne \cdots Ethyne (TS)	-1.54 ± 0.08	-1.53 ± 0.01	$0.01{\pm}0.08$
52	Benzene···AcOH (OH- π)	-4.64 ± 0.16	-4.70 ± 0.02	-0.07 ± 0.16
53	Benzene···AcNH ₂ (NH- π)	-4.25 ± 0.08	-4.38 ± 0.01	-0.13 ± 0.08
54	Benzene··· Water (OH- π)	-3.20 ± 0.06	-3.27 ± 0.01	-0.08 ± 0.07
55	Benzene···MeOH (OH- π)	-3.93 ± 0.07	-4.15 ± 0.02	-0.22 ± 0.07
56	Benzene···MeNH ₂ (NH- π)	-3.03 ± 0.07	-3.18 ± 0.01	-0.16 ± 0.07
57	Benzene···Peptide (NH- π)	-5.10 ± 0.10	-5.23 ± 0.02	-0.13 ± 0.10
58	$Pyridine \cdots Pyridine (CH-N)$	-4.22 ± 0.22	-4.21 ± 0.02	$0.01{\pm}0.22$
59	Ethyne··· Water (CH-O)	-3.04 ± 0.04	-2.92 ± 0.01	$0.13 {\pm} 0.04$
60	Ethyne····AcOH (OH- π)	$-4.98 {\pm} 0.07$	$-4.94{\pm}0.02$	$0.04{\pm}0.07$
61	$Pentane \cdots AcOH$	-2.63 ± 0.18	-2.89 ± 0.02	-0.26 ± 0.18
62	$\mathrm{Pentane}\cdots\mathrm{AcNH}_2$	-3.08 ± 0.07	-3.51 ± 0.02	-0.43 ± 0.07
63	Benzene····AcOH	-3.52 ± 0.08	-3.72 ± 0.02	-0.21 ± 0.08
64	$\operatorname{Peptide}\cdots\operatorname{Ethene}$	-2.77 ± 0.07	-2.98 ± 0.01	-0.21 ± 0.07
65	$Pyridine \cdots Ethyne$	-4.26 ± 0.06	-4.08 ± 0.02	$0.18{\pm}0.06$
66	$MeNH_2 \cdots Pyridine$	-3.81 ± 0.09	$-3.94{\pm}0.02$	-0.13 ± 0.09

S6 Mean relative difference of DMC and CCSD(T)

In the main manuscript, we report the difference between the DMC binding energy computed in this work and three previous CCSD(T) estimates from Refs. S17,S19 and S18. Here, in Fig. S16 we show the relative difference between DMC and CCSD(T) for each system in S66. Overall, Fig. S16 confirms the conclusion of the main manuscripts, i.e. the overall good agreement between CCSD(T) and DMC. The Mean Relative Deviation (MRD) is in fact 2.4% for H-bonded systems, 8.2% for dispersion dominated systems, and 4.4% for mixed systems. As shown in the main manuscript, the CCSD(cT) approach^{S20} significantly improves the agreement between coupled cluster and DMC for the dispersion dominated systems, with an MRD of ~ 3.4%.



Figure S16: Relative difference between DMC and CCSD(T) binding energies of the S66 dataset. We report the difference between previously computed $CCSD(T)^{S17-S19}$ values and our DMC estimates of the interaction energies for each system in S66. The DMC statistical error bar is reported in blue. CCSD(T) estimates from Ref. are reported with grey stars. CCSD(cT) values estimated in this work according to the approach described in Ref. S20 are reported with golden stars. The interaction energies are split in three different panels according to the prevalent interaction in the molecular complex: hydrogen bond (top), dispersion (centre), and mixed (bottom).

S7 Validation tests for the AcOH dimer

The acetic acid (AcOH) dimer (entry 20 of the S66 dataset) is found to have the largest deviation (~ 1 kcal/mol) between CCSD(T) and DMC. We thus perform additional validation tests to validate the accuracy of our estimate, utilising the DLA localization scheme^{S13} with an LDA trial wave-function and the eCEPP pseudopotential.^{S21} In Fig. S17 we report the binding energy of acetic acid as a function of the simulation time step computed with following set-ups: (i) the DLA localization scheme and the eCEPP pseudopotentials with the CASINO code using an LDA nodal surface (i.e., our original setup); (ii) the TM localization scheme and the ccECP pseudopotentials with the QMCPACK code using the LDA, PBE and PBE0 nodal surface; (iii) the DTM localization scheme and the eCEPP pseudopotentials with the CASINO code for an LDA nodal surface; (iv) the all electron (AE) calculation, i.e. with no pseudopotentials, with the QMCPACK code using the LDA nodal surface. We summarize the smallest time step estimate in Table S8 for each of the above methods; we did not perform extrapolations as the AE LDA calculations had large statistical uncertainties which prevented such extrapolations, and for the rest, estimates with τ below 0.01 au appear well-converged to within 0.15 kcal/mol. It can be seen that there is only a small range of only 0.28 kcal/mol between all of these procedures, suggesting that the results and conclusions reported in the main manuscript are not influenced by the simulation set up (code, localization scheme and choice of pseudopotentials) when fully converged.

Table S8: Validation of the DLA localization scheme with an LDA trial wave-function for the AcOH \cdots AcOH dimer (ID 20). The smallest time step τ and the corresponding interaction energy ΔE_{int} are reported using various trial wave-functions, localization schemes as well as with all-electron LDA

	au	$\Delta E_{\rm int}$
LDA//DLA(eCEPP)//CASINO	0.003	-20.27 ± 0.08
LDA//TM(eCEPP)//CASINO	0.003	$-19.99 {\pm} 0.08$
LDA//DTM(eCEPP)//CASINO	0.003	-20.15 ± 0.10
LDA//TM(ccECP)//QMCPACK	0.002	-20.25 ± 0.14
PBE//TM(ccECP)//QMCPACK	0.010	-20.10 ± 0.06
PBE0//TM(ccECP)//QMCPACK	0.010	-20.17 ± 0.06
LDA//AE//QMCPACK	0.002	-20.19 ± 0.11
8.0		



Figure S17: Analysis of the localization error for acetic acid. We report the dimer binding energy as a function of the DMC simulation time step, computed with the CASINO code and three different localization schemes (TM, DLA, and DTM) using the eCEPP^{S21} pseudopotentials, respectively in purple, red, and orange. In addition, we show the results with the TM algorithm for both the ccECP pseudopotentials (gray) and the all electron calculation, i.e. with no pseudopotentials (black), computed with the QMCPACK code. The quadratic fits of the DMC data are plotted with dashed lines. The different CCSD(T) estimates used for comparison with DMC are plotted respectively with a blue, ^{S19} gray, ^{S17} and gold line^{S18} horizontal lines.

S8 Convergence of the total and binding energy with respect to the DMC simulation time step

In this section, we show one figure for each dimer in S66, reporting the binding energy (left panel) and the total energy (right panel) as a function of the simulation time step. In each binding energy plot we also report the final CCSD(T) estimate obtained as described in Sec. S3. In each plot of the binding and total energy, we also show a cubic fit (red) over the range $\tau \sim [0, 0.1]$ au and a linear fit (blue) over the range $\tau \sim [0, 0.01]$ au. The cubic fit is always used by the linear fit can be used to gauge the expected level of error for systems where the time step behavior changes at small time steps, as discussed in Sec. S2.4. We observe that the 0.003 au estimates are converged to within 0.15 kcal/mol w.r.t. the zero time step limit estimates across the entire S66 dataset, with an MAD of 0.03 kcal/mol.



Figure S18: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Water \cdots Water (ID 1) dimer.



Figure S19: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Water · · · MeOH (ID 2) dimer.



Figure S20: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Water \cdots MeNH₂ (ID 3) dimer.



Figure S21: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Water. Peptide (ID 4) dimer.



Figure S22: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the MeOH \cdots MeOH (ID 5) dimer.



Figure S23: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the MeOH···MeNH₂ (ID 6) dimer.



Figure S24: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the MeOH···Peptide (ID 7) dimer.



Figure S25: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the MeOH···Water (ID 8) dimer.



Figure S26: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the MeNH₂··· MeOH (ID 9) dimer.



Figure S27: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the MeNH₂··· MeNH₂ (ID 10) dimer.



Figure S28: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the MeNH₂··· Peptide (ID 11) dimer.



Figure S29: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the MeNH₂··· Water (ID 12) dimer.



Figure S30: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Peptide · · · MeOH (ID 13) dimer.



Figure S31: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Peptide · · · MeNH₂ (ID 14) dimer.



Figure S32: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Peptide \cdots Peptide (ID 15) dimer.



Figure S33: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Peptide · · · Water (ID 16) dimer.



Figure S34: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Uracil... Uracil (BP) (ID 17) dimer.



Figure S35: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Water. Pyridine (ID 18) dimer.



Figure S36: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the MeOH···Pyridine (ID 19) dimer.



Figure S37: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the AcOH \cdots AcOH (ID 20) dimer.



Figure S38: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the AcNH₂··· AcNH₂ (ID 21) dimer.



Figure S39: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the AcOH··· Uracil (ID 22) dimer.



Figure S40: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the AcNH₂··· Uracil (ID 23) dimer.



Figure S41: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Benzene \cdots Benzene $(\pi - \pi)$ (ID 24) dimer.



Figure S42: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Pyridine \cdots Pyridine $(\pi - \pi)$ (ID 25) dimer.



Figure S43: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Uracil \cdots Uracil $(\pi - \pi)$ (ID 26) dimer.



Figure S44: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Benzene... Pyridine $(\pi - \pi)$ (ID 27) dimer.



Figure S45: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Benzene... Uracil $(\pi - \pi)$ (ID 28) dimer.



Figure S46: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Pyridine. Uracil $(\pi - \pi)$ (ID 29) dimer.



Figure S47: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Benzene. \cdots Ethene (ID 30) dimer.



Figure S48: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Uracil... Ethene (ID 31) dimer.



Figure S49: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Uracil... Ethyne (ID 32) dimer.



Figure S50: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Pyridine. $\cdot \cdot \cdot$ Ethene (ID 33) dimer.



Figure S51: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Pentane. Pentane (ID 34) dimer.



Figure S52: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Neopentane \cdots Pentane (ID 35) dimer.



Figure S53: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Neopentane (ID 36) dimer.



Figure S54: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Cyclopentane \cdots Neopentane (ID 37) dimer.



Figure S55: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Cyclopentane \cdots Cyclopentane (ID 38) dimer.



Figure S56: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Benzene. Cyclopentane (ID 39) dimer.



Figure S57: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Benzene... Neopentane (ID 40) dimer.



Figure S58: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Uracil... Pentane (ID 41) dimer.



Figure S59: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Uracil...Cyclopentane (ID 42) dimer.



Figure S60: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Uracil... Neopentane (ID 43) dimer.



Figure S61: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Ethene. Pentane (ID 44) dimer.



Figure S62: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Ethyne... Pentane (ID 45) dimer.



Figure S63: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Peptide \cdots Pentane (ID 46) dimer.


Figure S64: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Benzene \cdots Benzene (TS) (ID 47) dimer.



Figure S65: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Pyridine \cdots Pyridine (TS) (ID 48) dimer.



Figure S66: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Benzene... Pyridine (TS) (ID 49) dimer.



Figure S67: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Benzene... Ethyne (CH- π) (ID 50) dimer.



Figure S68: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Ethyne... Ethyne (TS) (ID 51) dimer.



Figure S69: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Benzene · · · AcOH (OH- π) (ID 52) dimer.



Figure S70: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Benzene · · · AcNH₂ (NH- π) (ID 53) dimer.



Figure S71: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Benzene... Water (OH- π) (ID 54) dimer.



Figure S72: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Benzene · · · MeOH (OH- π) (ID 55) dimer.



Figure S73: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Benzene \cdots MeNH₂ (NH- π) (ID 56) dimer.



Figure S74: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Benzene \cdots Peptide (NH- π) (ID 57) dimer.



Figure S75: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Pyridine \cdots Pyridine (CH-N) (ID 58) dimer.



Figure S76: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Ethyne... Water (CH-O) (ID 59) dimer.



Figure S77: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Ethyne...AcOH (OH- π) (ID 60) dimer.



Figure S78: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Pentane · · · AcOH (ID 61) dimer.



Figure S79: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Pentane · · · AcNH₂ (ID 62) dimer.



Figure S80: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Benzene···AcOH (ID 63) dimer.



Figure S81: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Peptide \cdots Ethene (ID 64) dimer.



Figure S82: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the Pyridine. $\cdot \cdot$ Ethyne (ID 65) dimer.



Figure S83: The time step dependence of ΔE_{int} and the total energy of the dimer complex for the MeNH₂···Pyridine (ID 66) dimer.

S9 Interaction energy decomposition analysis

Table S9: Symmetry-adapted perturbation theory (SAPT) energy decomposition for the S66 dataset taken from Ref. S22 using the SAPT0S-SA-jadz level of theory. The electrostatic (ELST), exchange (EXCH), induction (IND) and dispersion (DISP) energy components to the interaction energy are reported. The natural logarithm of the ratio between the electrostatic and dispersion energy is also reported.

System	ELST	EXCH	IND	DISP	LOG(ELST/DISP)
Water···Water	-8.569	6.651	-1.992	-1.222	1.947
$Water \cdots MeOH$	-9.517	8.040	-2.456	-1.747	1.695
$\mathrm{Water} \cdots \mathrm{MeNH}_2$	-12.719	11.830	-3.785	-2.120	1.792
$Water \cdots Peptide$	-13.376	11.329	-3.791	-2.639	1.623
$\mathrm{MeOH} \cdots \mathrm{MeOH}$	-9.547	8.413	-2.586	-2.059	1.534
$\mathrm{MeOH}\cdots\mathrm{MeNH}_2$	-13.210	13.167	-4.194	-2.930	1.506
$MeOH \cdots Peptide$	-13.224	12.114	-3.984	-3.199	1.419
$MeOH \cdots Water$	-8.454	6.853	-2.092	-1.447	1.765
$\mathrm{MeNH}_2\!\cdots\!\mathrm{MeOH}$	-4.350	4.261	-0.991	-1.622	0.986
$\mathrm{MeNH}_2\!\cdots\!\mathrm{MeNH}_2$	-5.970	6.435	-1.489	-2.472	0.882
$MeNH_2\cdots Peptide$	-7.334	7.561	-1.766	-3.416	0.764
$MeNH_2 \cdots Water$	-12.935	12.208	-3.900	-2.469	1.656
$\operatorname{Peptide}\cdots\operatorname{MeOH}$	-8.503	7.199	-2.122	-2.687	1.152
$\operatorname{Peptide}\cdots\operatorname{MeNH}_2$	-11.186	10.961	-3.310	-3.422	1.184
$\operatorname{Peptide}\cdots\operatorname{Peptide}$	-11.743	10.602	-3.379	-4.002	1.076
$Peptide \cdots Water$	-7.371	5.350	-1.655	-1.611	1.521
$\text{Uracil}\cdots\text{Uracil}$ (BP)	-27.486	26.002	-10.821	-6.076	1.509
$Water \cdots Pyridine$	-11.574	10.775	-3.610	-2.482	1.540
$MeOH \cdots Pyridine$	-12.117	11.904	-3.965	-3.196	1.333

(Continued on next page)

Table S9: (c	ontinued)
--------------	-----------

System	ELST	EXCH	IND	DISP	LOG(ELST/DISP)
AcOH····AcOH	-33.623	34.750	-15.334	-6.168	1.696
$AcNH_2 \cdots AcNH_2$	-26.449	24.247	-9.552	-5.031	1.660
$AcOH \cdots Uracil$	-32.158	30.809	-13.446	-6.119	1.659
$AcNH_2 \cdots Uracil$	-30.530	27.251	-11.514	-5.770	1.666
Benzene···Benzene $(\pi - \pi)$	-1.674	6.316	-0.686	-6.794	-1.401
Pyridine \cdots Pyridine $(\pi - \pi)$	-3.379	7.458	-0.809	-7.290	-0.769
Uracil···Uracil $(\pi$ - π)	-9.511	11.714	-1.645	-9.648	-0.014
Benzene···Pyridine $(\pi - \pi)$	-2.693	7.030	-0.762	-7.116	-0.972
Benzene··· Uracil $(\pi - \pi)$	-5.521	10.078	-1.126	-8.824	-0.469
Pyridine · · · Uracil $(\pi - \pi)$	-6.619	9.909	-1.215	-8.736	-0.278
$Benzene \cdots Ethene$	-0.892	4.271	-0.481	-3.707	-1.425
$\text{Uracil}\cdots\text{Ethene}$	-3.788	5.816	-0.545	-4.334	-0.135
$\text{Uracil}\cdots\text{Ethyne}$	-4.676	5.709	-0.584	-3.999	0.156
$\operatorname{Pyridine} \cdots \operatorname{Ethene}$	-1.712	4.865	-0.521	-3.954	-0.837
$Pentane \cdots Pentane$	-1.649	5.536	-0.500	-5.765	-1.252
$Neopentane \cdots Pentane$	-1.164	3.911	-0.378	-4.014	-1.238
$Neopentane \cdots Neopentane$	-0.659	2.637	-0.284	-2.850	-1.465
$Cyclopentane \cdots Neopentane$	-1.144	3.890	-0.393	-3.894	-1.225
$Cyclopentane \cdots Cyclopentane$	-1.324	4.371	-0.438	-4.554	-1.235
$\mathrm{Benzene} \cdots \mathrm{Cyclopentane}$	-2.394	5.864	-0.646	-5.972	-0.914
$\mathrm{Benzene} \cdots \mathrm{Neopentane}$	-1.814	4.464	-0.521	-4.635	-0.938
$\text{Uracil}\cdots\text{Pentane}$	-2.468	6.869	-0.813	-6.973	-1.039
$\mathrm{Uracil}\cdots\mathrm{Cyclopentane}$	-1.990	5.843	-0.608	-6.154	-1.129

(Continued on next page)

System	ELST	EXCH	IND	DISP	LOG(ELST/DISP)
Uracil··· Neopentane	-2.350	5.004	-0.490	-4.913	-0.738
$Ethene \cdots Pentane$	-1.018	3.218	-0.335	-2.956	-1.065
$Ethyne \cdots Pentane$	-1.241	3.069	-0.342	-2.542	-0.717
$\operatorname{Peptide} \cdots \operatorname{Pentane}$	-2.332	6.161	-0.931	-5.673	-0.889
Benzene \cdots Benzene (TS)	-2.025	4.016	-0.574	-4.277	-0.748
$Pyridine \cdots Pyridine (TS)$	-3.033	4.592	-0.723	-4.350	-0.361
Benzene \cdots Pyridine (TS)	-2.626	4.241	-0.707	-4.330	-0.500
Benzene···Ethyne (CH- π)	-2.490	3.106	-0.805	-2.798	-0.117
Ethyne \cdots Ethyne (TS)	-2.111	2.203	-0.521	-0.981	0.767
Benzene···AcOH (OH- π)	-4.395	5.735	-1.792	-3.786	0.149
Benzene···AcNH ₂ (NH- π)	-5.326	6.017	-1.515	-3.321	0.473
Benzene··· Water (OH- π)	-3.360	3.520	-0.976	-2.203	0.422
Benzene··· MeOH (OH- π)	-3.680	5.051	-1.168	-3.761	-0.022
Benzene··· MeNH ₂ (NH- π)	-2.649	4.407	-0.677	-3.780	-0.356
Benzene···Peptide (NH- π)	-4.265	6.178	-1.329	-5.458	-0.247
$Pyridine \cdots Pyridine (CH-N)$	-4.964	5.017	-1.336	-3.047	0.488
Ethyne···· Water (CH-O)	-4.482	3.051	-0.950	-0.878	1.630
Ethyne····AcOH (OH- π)	-7.787	7.906	-2.401	-2.516	1.130
$Pentane \cdots AcOH$	-1.574	4.290	-0.533	-3.843	-0.893
$Pentane \cdots AcNH_2$	-2.141	5.398	-0.992	-4.457	-0.733
Benzene··· AcOH	-3.688	5.718	-0.821	-4.734	-0.250
$\operatorname{Peptide}\cdots\operatorname{Ethene}$	-2.991	4.378	-0.804	-2.948	0.015
Pyridine···Ethyne	-6.249	5.446	-1.749	-1.926	1.177

Table S9: (continued)

(Continued on next page)

	Table S9: (continued)					
System	ELST	EXCH	IND	DISP	LOG(ELST/DISP)	
$MeNH_2 \cdots Pyridine$	-4.622	6.071	-1.035	-3.863	0.179	

References

- (S1) Rezáč, J.; Riley, K. E.; Hobza, P. S66: A well-balanced database of benchmark interaction energies relevant to biomolecular structures. <u>Journal of Chemical Theory and</u> Computation **2011**, 7, 2427–2438.
- (S2) Neese, F. Software update: the ORCA program system, version 4.0. <u>WIREs</u> Computational Molecular Science **2018**, 8, e1327.
- (S3) Riplinger, C.; Neese, F. An efficient and near linear scaling pair natural orbital based local coupled cluster method. The Journal of Chemical Physics 2013, 138, 034106.
- (S4) Riplinger, C.; Sandhoefer, B.; Hansen, A.; Neese, F. Natural triple excitations in local coupled cluster calculations with pair natural orbitals. <u>The Journal of Chemical</u> <u>Physics</u> 2013, 139, 134101.
- (S5) Zhong, S.; Barnes, E. C.; Petersson, G. A. Uniformly convergent n-tuple-zeta augmented polarized (nZaP) basis sets for complete basis set extrapolations. I. Selfconsistent field energies. The Journal of Chemical Physics 2008, 129, 184116.
- (S6) Helgaker, T.; Klopper, W.; Koch, H.; Noga, J. Basis-set convergence of correlated calculations on water. The Journal of Chemical Physics 1997, 106, 9639–9646.
- (S7) Neese, F.; Valeev, E. F. Revisiting the Atomic Natural Orbital Approach for Basis Sets: Robust Systematic Basis Sets for Explicitly Correlated and Conventional Correlated ab initio Methods? Journal of Chemical Theory and Computation 2011, 7, 33–43.
- (S8) Foulkes, W. M. C.; Mitas, L.; Needs, R. J.; Rajagopal, G. Quantum Monte Carlo simulations of solids. Reviews of Modern Physics 2001, 73, 33–83.
- (S9) Mitáš, L.; Shirley, E. L.; Ceperley, D. M. Nonlocal pseudopotentials and diffusion Monte Carlo. The Journal of Chemical Physics **1991**, 95, 3467–3475.

- (S10) Hammond, B.; Reynolds, P.; Lester, W. Valence quantum Monte Carlo with ab initio effective core potentials. The Journal of chemical physics **1987**, 87, 1130–1136.
- (S11) Casula, M. Beyond the locality approximation in the standard diffusion Monte Carlo method. Physical Review B 2006, 74, 161102.
- (S12) Casula, M.; Moroni, S.; Sorella, S.; Filippi, C. Size-consistent variational approaches to nonlocal pseudopotentials: Standard and lattice regularized diffusion Monte Carlo methods revisited. The Journal of Chemical Physics **2010**, 132, 154113.
- (S13) Zen, A.; Brandenburg, J. G.; Michaelides, A.; Alfè, D. A new scheme for fixed node diffusion quantum Monte Carlo with pseudopotentials: Improving reproducibility and reducing the trial-wave-function bias. <u>The Journal of Chemical Physics</u> 2019, <u>151</u>, 134105.
- (S14) Zen, A.; Sorella, S.; Gillan, M. J.; Michaelides, A.; Alfè, D. Boosting the accuracy and speed of quantum Monte Carlo: Size consistency and time step. <u>Physical Review B</u> 2016, <u>93</u>, 241118.
- (S15) Zen, A.; Brandenburg, J. G.; Klimeš, J.; Tkatchenko, A.; Alfè, D.; Michaelides, A. Fast and accurate quantum Monte Carlo for molecular crystals. <u>Proceedings of the</u> National Academy of Sciences **2018**, 115, 1724–1729.
- (S16) Rezáč, J.; Riley, K. E.; Hobza, P. Extensions of the S66 Data Set: More Accurate Interaction Energies and Angular-Displaced Nonequilibrium Geometries. <u>J. Chem.</u> Theory Comput. **2011**, 7, 3466–3470.
- (S17) Kesharwani, M. K.; Karton, A.; Sylvetsky, N.; Martin, J. M. L. The S66 Non-Covalent Interactions Benchmark Reconsidered Using Explicitly Correlated Methods Near the Basis Set Limit. Australian Journal of Chemistry 2018, 71, 238.

- (S18) Péter R. Nagy, B. D. L., László Gyevi-Nagy; Kállay, M. Pursuing the basis set limit of CCSD(T) non-covalent interaction energies for medium-sized complexes: case study on the S66 compilation. Molecular Physics **2023**, 121, e2109526.
- (S19) Rezáč, J.; Riley, K. E.; Hobza, P. Extensions of the S66 Data Set: More Accurate Interaction Energies and Angular-Displaced Nonequilibrium Geometries. <u>Journal of</u> <u>Chemical Theory and Computation</u> **2011**, <u>7</u>, 3466–3470.
- (S20) Schäfer, T.; Irmler, A.; Gallo, A.; Grüneis, A. Understanding Discrepancies of Wavefunction Theories for Large Molecules. 2024; https://arxiv.org/abs/2407.01442.
- (S21) Trail, J. R.; Needs, R. J. Shape and energy consistent pseudopotentials for correlated electron systems. The Journal of Chemical Physics 2017, 146, 204107.
- (S22) Burns, L. A.; Faver, J. C.; Zheng, Z.; Marshall, M. S.; Smith, D. G. A.; Vanommeslaeghe, K.; MacKerell, J., Alexander D.; Merz, J., Kenneth M.; Sherrill, C. D. The BioFragment Database (BFDb): An open-data platform for computational chemistry analysis of noncovalent interactions. <u>The Journal of Chemical Physics</u> 2017, <u>147</u>, 161727.