Wheel-like bricks and minimal matching covered graphs *

Xiaoling He 1, Fuliang Lu 2, Jinxin Xue 2

1. School of Mathematics and Statistics, Lanzhou University, Lanzhou, China

2. School of Mathematics and Statistics, Minnan Normal University, Zhangzhou, China

Abstract

A connected graph G with at least two vertices is matching covered if each of its edges lies in a perfect matching. We say that an edge e in a matching covered graph G is removable if G - e is matching covered. A pair $\{e, f\}$ of edges of a matching covered graph G is a removable doubleton if G - e - f is matching covered, but neither G - e nor G - f is. Removable edges and removable doubletons are called removable classes, introduced by Lovász and Plummer in connection with ear decompositions of matching covered graphs.

A 3-connected graph is a *brick* if the removal of any two distinct vertices, the left graph has a perfect matching. A brick G is *wheel-like* if G has a vertex h, such that every removable class of G has an edge incident with h. Lucchesi and Murty proposed a problem of characterizing wheel-like bricks. We show that every wheel-like brick may be obtained by splicing graphs whose underlying simple graphs are odd wheels in a certain manner.

A matching covered graph is *minimal* if the removal of any edge, the left graph is not matching covered. Lovász and Plummer proved that the minimum degree of a minimal matching covered bipartite graph different from K_2 is 2 by ear decompositions in 1977. By the properties of wheel-like bricks, we prove that the minimum degree of a minimal matching covered graph other than K_2 is 2 or 3.

Keywords: wheel-like bricks; minimal matching covered graphs; minimum degrees

1 Introduction

Graphs considered in this paper may have multiple edges, but no loops. We follow [1] for undefined notation and terminology. Let G be a graph with the vertex set V(G)

^{*}The research is partially supported by NSFC (No. 12271235).

E-mail addresses: xiaolinghe99@163.com (X. He), flianglu@163.com (F. Lu), xuejinxin00@163.com (J. Xue)

and the edge set E(G). A connected nontrivial graph G is matching covered if each of its edges lies in a perfect matching. A graph G with four or more vertices is bicritical if for any two distinct vertices u and v in $G, G - \{u, v\}$ has a perfect matching. Obviously, every bicritical graph is matching covered.

For $X, Y \subseteq V(G)$, by $E_G[X, Y]$ we mean the set of edges of G with one end vertex in X and the other end vertex in Y. Let $\partial_G(X) = E_G[X, \overline{X}]$ be an edge cut of G, where $\overline{X} = V(G) \setminus X$. (If G is understood, the subscript G is omitted.) If $X = \{u\}$, then we denote $\partial_G(\{u\})$, for brevity, by $\partial_G(u)$ or $\partial(u)$. The size of $|\partial_G(u)|$, denoted by $d_G(u)$, means the degree of u in G. Denote by $\delta(G)$ and $\Delta(G)$ the minimum degree and the maximum degree of G, respectively. An edge cut $\partial(X)$ is trivial if |X| = 1 or $|\overline{X}| = 1$. Let $\partial(X)$ be an edge cut of G. Denoted by $G/(X \to x)$, or simply G/X, the graph obtained from G by contracting X to a singleton x (and removing any resulting loops). The graphs G/X and G/\overline{X} are $\partial(X)$ -contractions of G.

An edge cut $\partial(X)$ is separating if both $\partial(X)$ -contractions of G are matching covered, and $\partial(X)$ is tight of G if $|\partial(X) \cap M| = 1$ for every perfect matching M of G. Obviously, a trivial edge cut is a tight cut and every tight cut is separating. A matching covered nonbipartite graph is a brick if every tight cut is trivial, and it is solid if every separating cut is a tight cut. Moreover, a graph G is a brick if and only if G is 3-connected and bicritical [6]. There is a procedure called a tight cut decomposition, due to Lovász [11], which can be applied to G to produce a list of unique bricks and braces (a matching covered bipartite graph in which every tight cut is trivial). In particular, any two decompositions of a matching covered graph G yield the same number of bricks; this number is denoted by b(G). A matching covered graph G is called a near-brick if b(G) = 1. Obviously, every brick is a near-brick.

We say that an edge e in a matching covered graph G is removable if G-e is matching covered. A pair $\{e, f\}$ of edges of a matching covered graph G is a removable doubleton if G-e-f is matching covered, but neither G-e nor G-f is. Removable edges and removable doubletons are called removable classes. Lovász [10] proved that every brick different from K_4 (the complete graph with 4 vertices) and $\overline{C_6}$ (the triangular prism) has a removable edge. Improving Lovász's result, Carvalho, Lucchesi and Murty obtained a lower bound of removable classes of a brick in terms of the maximum degree.

Theorem 1.1 ([2]). Every brick has at least $\Delta(G)$ removable classes. Moreover, every brick has at least $\Delta(G) - 2$ removable edges.

For an integer $k \ge 3$, the *wheel* W_k is the graph obtained from a cycle C of length k by adding a new vertex h and joining it to all vertices of C. The cycle C is the *rim* of

 W_k , the vertex h is its hub. Obviously, every wheel is planar. A wheel W_k is odd if k is odd. The graph K_4 is an odd wheel that every edge lies in a removable doubleton. For an odd wheel other than K_4 , it can be checked every edge on the rim is not removable, and every edge incident with the hub is removable (see Exercise 2.2.4 in [7]). We say that G is wheel-like if for every removable class R of G, there exists a vertex h of G, called its hub, such that $|R \cap \partial(h)| = 1$. Lucchesi and Murty proposed the following problem.

Problem 1.2. (see Page 216, and Unsolved Problems 10 in [7]) Characterize wheel-like bricks; characterize wheel-like bricks as a splicing of two bricks.

We obtain the following theorem in this paper, where the graph family \mathcal{G} is defined in Section 3.

Theorem 1.3. Let G be a wheel-like brick. Then $G \in \mathcal{G}$.

Let G be a matching covered graph. We say that G is minimal if G - e is not a matching covered graph for any edge e in G. Obviously, if G is minimal, then G has no removable edges. It is known that every matching covered graph on four or more vertices is 2-connected [9]. So, the minimum degree of a matching covered graph with more than two vertices is at least 2. Lovász and Plummer proved that $\delta(G) = 2$ if G is a minimal matching covered bipartite graph [8]. For example, a cycle with even number of vertices is a minimal matching covered bipartite graph [8]. For example, a cycle with even number of vertices is a minimal matching covered bipartite graph, the minimum degree may be 3; for example, K_4 and $\overline{C_6}$ are such graphs. Using the properties of wheel-like bricks, we prove the following result.

Theorem 1.4. Let G be a minimal matching covered graph with at least four vertices. Then $\delta(G) = 2$ or 3.

We will present some known results in Section 2. The propositions of wheel-like bricks will be presented in Section 3, and the proof of Theorem 1.4 will be given in Section 4.

2 Preliminaries

We begin with some notation. For a vertex $u \in V(G)$, denoted by $N_G(u)$ or simply N(u), the set of vertices in G adjacent to u. We denote by o(G) the number of components with odd number of vertices of a graph G. A nonempty vertex set B of a graph G that has a perfect matching is a *barrier* if o(G-B) = |B|. A component (or a barrier) is trivial if it contains exactly one vertex. Tutte proved the following theorem in 1947.

Theorem 2.1 ([13]). A graph G has a perfect matching if and only if $o(G - S) \leq |S|$, for every $S \subseteq V(G)$.

Using Tutte's Theorem, we have the following properties about matching covered graphs.

Corollary 2.2 ([11]). Let G be a matching covered graph and let S be a subset of V(G). Then, $o(G - S) \leq |S|$, with equality only if S is independent and G - S has no even components.

Proposition 2.3 ([9]). A matching covered graph different from K_2 is bicritical if and only if for every barrier B of G, |B| = 1.

A vertex set S of G is a 2-separation if |S| = 2, G - S is disconnected and each of the components of G - S is even. The following corollary can be gotten directly by Proposition 2.3.

Corollary 2.4. Let G be a bicritical graph different from K_2 and let $u, v \in V(G)$. If $G - \{u, v\}$ is disconnected, then $\{u, v\}$ is a 2-separation of G.

An edge cut C of G is a *barrier-cut* if there exists a barrier B of G and an odd component Q of G - B such that $C = \partial(V(Q))$. Let $\{u, v\}$ be a 2-separation of G, and let us divide the components of $G - \{u, v\}$ into two nonempty subgraphs G_1 and G_2 . The cuts $\partial(V(G_1) + u)$ and $\partial(V(G_1) + v)$ are both 2-separation cuts associated with $\{u, v\}$ of G. Barrier-cuts and 2-separation cuts, which are tight cuts, play an important role during tight cut decomposition. The following theorem, due to Edmonds, Lovász, and Pulleyblank [6], is a fundamental result of matching theory.

Theorem 2.5 ([6]). Every matching covered graph that has a nontrivial tight cut has a nontrivial barrier or a 2-separation.

2.1 The splicing of two graphs and Robust cuts

Let G and H be two vertex-disjoint graphs and let u and v be vertices of G and H, respectively, such that $d_G(u) = d_H(v)$. Moreover, let θ be a given bijection between $\partial_H(v)$ and $\partial_G(u)$. We denote by $(G(u) \odot H(v))_{\theta}$ the graph obtained from the union of G - uand H - v by joining, for each edge e in $\partial_H(v)$, the end of e in H belonging to V(H) - vto the end of $\theta(e)$ in G belonging to V(G) - u; and refer to $(G(u) \odot H(v))_{\theta}$ as the graph obtained by splicing G (at u), with H (at v), with respect to the bijection θ , for brevity, to $G(u) \odot H(v)$. We say u and v are the splicing vertices of G and H, respectively. In general, the graph resulted from splicing two graphs G and H depends on the choice of u, v and θ . The following proposition can be gotten by the definition of matching covered graphs directly (see Theorem 2.13 in [7] for example).

Proposition 2.6. The splicing of two matching covered graphs is also matching covered.

Let G be a matching covered graph. A separating cut C of G is a *robust cut* if C is not tight and both C-contractions of G are near-bricks.

Proposition 2.7 ([3]). Let G be a nonsolid brick. Then G has a robust cut.

Theorem 2.8 ([3]). Let G be a brick and $\partial(X)$ be a robust cut of G. Then, there exists a subset X' of X and a subset X'' of \overline{X} such that $G/\overline{X'}$ and $G/\overline{X''}$ are bricks and the graph H, obtained from G by contracting X' and X'' to single vertices x' and x'', respectively, is bipartite and matching covered, where x' and x'' lie in different color classes of H.

Lemma 2.9 ([12]). Let $\partial(X)$ and $\partial(Y)$ be two robust cuts of a brick G such that G/Xand G/Y are bricks, and $(G/(\overline{X} \to \overline{x}))/(\overline{Y} \to \overline{y})$ is a bipartite graph H. Then every edge incident with \overline{x} is removable in H.

2.2 Removable classes

We may assume that an edge e is removable in a matching covered graph G if $e \notin E(G)$. The following lemma is easy to verify by the definition (e.g., see Propositions 8.7 and 8.8 in [7]).

Lemma 2.10. Let G be a matching covered graph and C be a separating cut of G. If an edge e is removable in both C-contractions of G then e is removable in G. Moreover, if C is tight, then an edge e is removable in G if and only if e is removable in both C-contractions of G.

Lemma 2.11 (Lemma 3.1 in [3]). Let $C = \partial(X)$ be a separating cut but not a tight cut of a matching covered graph G and let $H = G/\overline{X}$. Suppose that H is a brick, and let R be a removable doubleton of H. If $R \cap C = \emptyset$ or if the edge of $R \cap C$ is removable in G/Xthen $R \setminus C$ contains an edge which is removable in G.

We shall denote a bipartite graph G with bipartition (A, B) by G[A, B]. The following proposition which can be derived from the definition of bipartite matching covered graphs will be used in the following text.

Proposition 2.12. Let G[A, B] be a matching covered graph. Assume that $X \subseteq V(G)$ such that $N(X \cap A) \subseteq X \cap B$ (X is not necessary nonempty). Then $|X \cap A| \leq |X \cap B|$. Moreover, $|X \cap A| = |X \cap B|$ if and only if $X = \emptyset$.

Proposition 2.13 ([11]). Let G[A, B] be a matching covered graph. An edge cut $\partial(X)$ of G is tight if and only if $||X \cap A| - |X \cap B|| = 1$ and every edge of $\partial(X)$ is incident with a vertex of the larger one between the two sets, $X \cap A$ and $X \cap B$.

Lemma 2.14 ([5]). Let G[A, B] be a matching covered graph, and $|E(G)| \ge 2$. An edge uv of G, with $u \in A$ and $v \in B$, is not removable in G if and only if there exist nonempty proper subsets A_1 and B_1 of A and B, respectively, such that:

1) the subgraph $G[A_1 \cup B_1]$ is matching covered, and

2) $u \in A_1$ and $v \in B \setminus B_1$, and $E[A_1, B \setminus B_1] = \{uv\}.$

Let G[A, B] be a matching covered graph with at least two edges and let X be a vertex set of G such that $|X \cap A| = |X \cap B|$. We say that X is a *P*-set of G if either $|E[X \cap A, \overline{X} \cap B]| = 1$ or $|E[\overline{X} \cap A, X \cap B]| = 1$. Obviously, \overline{X} is a *P*-set if X is a *P*-set. A *P*-set X of G is minimum if for each *P*-set Y in G different from X, $|X| \leq |Y|$. By Lemma 2.14, for every nonremoveble edge of a bipartite matching covered graph, there exist two *P*-sets associated with it.

Lemma 2.15. Let G[A, B] be a matching covered graph with at least 4 vertices and $\delta(G) \ge 3$. If X is a minimum P-set of G, then every edge of E(G[X]) is removable in G.

Proof. Without loss of generality, assume that $|E[\overline{X} \cap A, X \cap B]| = 1$. Let $\{ab\} = E[\overline{X} \cap A, X \cap B]$, where $a \in \overline{X} \cap A$ and $b \in X \cap B$. If |X| = 2, then E(G[X]) consists of some multiple edges, as $d(b) \geq 3$. As multiple edges are removable, the result holds. Now assume that $|X| \geq 4$. Suppose that there exists a nonremovable edge uv of G, such that $uv \in E(G[X])$. Let $Y = X \cup \{a\}$. It can be checked by Proposition 2.13 that $\partial(Y)$ is a tight cut. Noting that \overline{X} is also a P-set, $|\overline{X}| \geq |X| > 2$ by the minimality of X. Let $G' = G/(\overline{Y} \to \overline{y})$ (note that $G' \neq G$). Then uv is also a nonremovable edge of G' by Lemma 2.10. So there exists a P-set Z associated with uv in G'. Without loss of generality, assume that $E[Z \cap A, \overline{Z} \cap B] = \{uv\}$ and $u \in Z \cap A$. Note that $V(G') = \{a, \overline{y}\} \cup X$ and $\{u, v\} \subset X$. If $\{a, \overline{y}\} \subset Z$ (the case is the same if $\{a, \overline{y}\} \subset \overline{Z}$), then $\overline{Z} \subset X$ and $\overline{Z} \neq X$, that is, $|\overline{Z}| < |X|$. Note that \overline{Z} is also a P-set of G. It contradicts the assumption that X is a minimum P-set of G. So $|\{a, \overline{y}\} \cap Z| = 1$.

As $E[Z \cap A, \overline{Z} \cap B] = \{uv\}, a \neq u, \overline{y} \neq v \text{ and } a\overline{y} \in E(G'), \text{ we have } a \in \overline{Z} \cap A$ and $\overline{y} \in Z \cap B$. Suppose that $b \notin \overline{Z} \cap B$ or b = v. As $N_{G'}(a) = \{\overline{y}, b\}$ and $\overline{y} \notin \overline{Z}$, we have $N(\overline{Z} \cap B \setminus \{v\}) \subseteq \overline{Z} \cap A \setminus \{a\}$. Since $|\overline{Z} \cap B \setminus \{v\}| = |\overline{Z} \cap A \setminus \{a\}|$, we have $|\overline{Z} \cap B \setminus \{v\}| = |\overline{Z} \cap A \setminus \{a\}| = 0$ by Proposition 2.12. It means that $\overline{Z} = \{v, a\}$. If $b \notin \overline{Z} \cap B$, then $N_{G'}(v) = \{u\}$, contradicting the fact that G' is 2-connected. If b = v, then $N_{G'}(v) = \{u, a\}$. As uv and ab are nonremovable in G', we have $d_{G'}(v) = 2$, contradicting the assumption that $\delta(G') \ge 3$ (as $\delta(G) \ge 3$). Thus, $b \in \overline{Z} \cap B$ and $b \neq v$. Let $W = \overline{Z} \setminus \{a, b\}$. Then $|W \cap A| = |W \cap B|$ and $E[W \cap B, \overline{W} \cap A] = \{uv\}$ (as $N_{G'}(a) = \{\overline{y}, b\}$). It means that W is a P-set of G' associated with uv. Moreover, as $\overline{y} \notin W$ and $\overline{W} = Z \cup \{a, b\}$, W is also a P-set of G and $|W| \le |V(G')| - 4$. Note that |X| = |V(G')| - 2. Then |W| < |X|, contradicting the assumption that X is minimum. Therefore, the result follows.

Lemma 2.16. Let G[A, B] be a matching covered graph with at least 4 vertices. Assume that every vertex of A has degree at least 3. Then G has two nonadjacent removable edges or there exist two vertices u, v in V(G) such that $v \in B$, d(v) = 2, $d(u) \ge 4$ and every edge of $\partial(u)$ is removable.

Proof. We will prove the result by induction on |V(G)|. If |V(G)| = 4, then it can be checked that the result follows, as every vertex of A has degree at least 3 and at most two neighbors. Suppose that the result holds for $|V(G)| \le n$. Now we consider the case when |V(G)| = n + 2, where n is an even integer greater than 4.

Assume firstly that there exists a vertex $v \in B$ such that $d_G(v) = 2$. Let $N_G(v) = \{v_1, v_2\}$ and $Y = \{v, v_1, v_2\}$. It can be checked by Proposition 2.13 that $\partial(Y)$ is a nontrivial tight cut of G, as |V(G)| > 4. Let $G' = G/(\overline{Y} \to \overline{y})$ and $G'' = G/(Y \to y)$. Then G' and G'' are bipartite and matching covered. As $d_G(v) = 2$, $N_{G'}(v_i) \setminus \{v\} = \{\overline{y}\}$ (note that |V(G')| = 4 and G' is 2-connected) and every vertex of N(v) has degree at least 3, we have $|E[\{v_i\}, \{\overline{y}\}]| \ge 2$, for each $i \in \{1, 2\}$. It means that every edge of $\partial(\overline{y})$ is removable in G'. By induction, G'' has two nonadjacent removable edges or there exists a vertex w, such that $d_{G''}(w) \ge 4$ and every edge of $\partial_{G''}(w)$ is removable in G''. If G'' has two nonadjacent removable edges or the vertex $w \neq y$, then the result holds by Lemma 2.10, as every edge of $\partial(\overline{y})$ is removable in G'. Assume that w = y. Then every edge of $\partial(Y)$ is removable in G by Lemma 2.10 again. We can find two nonadjacent edges in $\partial(Y)$ as G is 2-connected. So the result follows.

Now assume that $\delta(G) \geq 3$. Let X be a minimum P-set of G. Without loss of generality, assume that $|E[\overline{X} \cap A, X \cap B]| = 1$. Let $\{ab\} = E[\overline{X} \cap A, X \cap B]$, where $a \in \overline{X} \cap A$ and $b \in X \cap B$. If $|X| \geq 4$, then by Lemma 2.15, E(G[X]) contains two nonadjacent removable edges of G, so the result holds. Assume that |X| = 2. Let

 $Z = X \cup \{a\}$. It can be checked by Proposition 2.13 that $\partial(Z)$ is a nontrivial tight cut as |V(G)| > 4 and $|E[Z \cap B, \overline{Z} \cap A]| = \emptyset$ (note that $E[\overline{X} \cap A, X \cap B] = \{ab\}$). Let $G_1 = G/(\overline{Z} \to \overline{z})$ and $G_2 = G/(Z \to z)$. As $d_G(a) \ge 3$ and $E_G[\overline{X} \cap A, X \cap B] = \{ab\}$, we have $|E_G[\{a\}, \overline{X}]| \ge 2$. Then $|E_{G_1}[\{a\}, \{\overline{z}\}]| \ge 2$, as $\overline{X} \setminus \{a\} \subseteq \overline{Z}$. It means that every edge of $E_{G_1}[\{a\}, \{\overline{z}\}]$ is removable in G_1 . Let $\{a'\} = X \setminus \{b\}$. As $\delta(G) \ge 3$, $N_G(b) = \{a', a\}$ and $|E_G[\{a\}, \{b\}]| = 1$, we have $|E_G[\{a'\}, \{b\}]| > 1$. Then a'b is also removable in G by Lemma 2.10.

By induction, G_2 has two nonadjacent removable edges, or there exists a vertex s such that $d_{G_2}(s) \ge 4$ and every edge of $\partial(s)$ is removable in G_2 . Recall that a'b is removable in G. If G_2 has two nonadjacent removable edges or the vertex $s \ne z$, that is, there exists a removable edge e of G_2 which lies in $E(G_2) \setminus \partial(z)$, then e is also removable in G by Lemma 2.10. Moreover, e and a'b is nonadjacent. So the result holds in this case. Assume that s = z. Recall that every edge of $E_{G_1}[\{a\}, \{\overline{z}\}]$ is removable in G_1 and every edge of $\partial_{G_2}(z)$ is removable in G_2 . Then there exists a removable edge of G which is incident with a by Lemma 2.10 again. Together with a'b, G has two nonadjacent removable edges. The result holds.

Lemma 2.17. Let $\partial(X)$ and $\partial(Y)$ be two robust cuts of a brick G such that G/X and G/Y are bricks, and $(G/(\overline{X} \to \overline{x}))/(\overline{Y} \to \overline{y})$ is a matching covered bipartite graph H. If $G/(X \to x)$ is wheel-like such that x is its hub and every edge of $\partial_{G/X}(x)$ belongs to a removable class of G/X, and $|N_H(\overline{x})| \ge 2$, then there exists a removable edge e of G such that both ends of e belong to $\overline{X} \cup N(\overline{X}) \setminus \overline{Y}$.

Proof. Let $G' = G/(X \to x)$. Note that $|N_H(\overline{x})| \ge 2$. Assume that $\overline{x}b$ is an edge in H such that $b \neq \overline{y}$. Then $\overline{x}b$ is removable in H by Lemma 2.9. Let $G/\overline{Y} = (G'(x) \odot H(\overline{x}))_{\theta}$. If $\theta(\overline{x}b)$ is removable in G', then $\overline{x}b$ is also removable in G/\overline{Y} by Lemma 2.10. Since $\overline{x}b \notin \partial(\overline{y}), \overline{x}b$ is removable in G by Lemma 2.10 again. Moreover, the ends of $\overline{x}b$ belong to $\overline{X} \cup N(\overline{X}) \setminus \overline{Y}$. The result follows by setting $e = \overline{x}b$ in this case. If $\theta(\overline{x}b)$ is an edge of a removable doubleton in G', assume that $\{e', \theta(\overline{x}b)\}$ is a removable doubleton in G'. Then e' is a removable edge in G/\overline{Y} by Lemma 2.11. So e' is removable in G by Lemma 2.10 once more. As the ends of e' belong to $\overline{X} \cup N(\overline{X}) \setminus \overline{Y}$, the result follows by setting e = e'.

3 Wheel-like bricks

In this section, we present some properties of wheel-like bricks and prove the Theorem 1.3. Lucchesi and Murty showed the following proposition.

Proposition 3.1 ([7]). Let G be a solid brick and let h be a vertex of G. Either G is a wheel having h as a hub, possibly with multiple edges incident with h, or G has two removable edges not incident with h.

Proposition 3.2 ([12]). Let G be a wheel-like brick and h is its hub. Then all the multiple edges are incident with h.

A matching covered graph G is *near-bipartite* if it has a pair of edges e and f such that the subgraph G - e - f obtained by the deletion of e and f is a matching covered bipartite graph. In fact, every brick with a removable doubleton is near-bipartite [11].

Theorem 3.3 (Theorem 9.17 in [7]). Every simple near-bipartite brick distinct from K_4 , $\overline{C_6}$ and R_8 (see Figure 1) has two nonadjacent removable edges.



Figure 1: R_8 .

Lemma 3.4 ([12]). 1) Let G be a simple near-bipartite brick. Then G is wheel-like if and only if G is isomorphic to K_4 .

2) Let G be a simple planar brick with six vertices. Then G is a wheel-like brick if and only if G is W_5 .

Proposition 3.5 ([4]). Let G be a simple brick on six vertices. Then G is either nonsolid or the 5-wheel W_5 .

Lemma 3.6. Let G be a wheel-like brick on 6 vertices and h be the hub of G. Then G is W_5 , possibly with multiple edges incident with h.

Proof. As G is wheel-like, the only possible multiple edges in G are incident with h by Proposition 3.2. If G is solid or planar, then the underlying simple graph of G is W_5 by Proposition 3.5 and Lemma 3.4. So the result is obviously.

Suppose that G is nonsolid and nonplanar. Then G has a nontrivial separating cut $\partial(X)$. As |V(G)| = 6, |X| = 3 and $|\overline{X}| = 3$. Note that G is 3-connected and both $\partial(X)$ contractions are matching covered. Then both $\partial(X)$ -contractions are isomorphic to K_4 (up to multiple edges). It can be checked that G is isomorphic to $\overline{C_6}$ or one of the graphs
in Figure 2. All the graphs in Figure 2 are not wheel-like (the bold edges are removable).
By Lemma 3.4, $\overline{C_6}$ is not wheel-like. Therefore, the result holds.



Figure 2: Nonplanar nonsolid bricks on six vertices, where the bold edges are removable.

Lemma 3.7 ([12]). Let G_1 and G_2 be two disjoint bricks and let $u \in V(G_1)$ and $v \in V(G_2)$. Assume that $G = G_1(u) \odot G_2(v)$ is a brick.

1) If G is wheel-like, then at least one of G_1 and G_2 is wheel-like such that u or v is its hub.

2) If G_1 is wheel-like, u is its hub, and every edge of $\partial_{G_1}(u)$ lies in some removable class of G_1 , then G_2 is also wheel-like.

Lu and Xue characterized wheel-like bricks that are obtained from the splicing of two odd wheels.

Lemma 3.8 ([12]). Let G and H be two odd wheels such that $V(G) = \{u_h, u_1, u_2, \ldots, u_s\}$ and $V(H) = \{v_h, v_1, v_2, \ldots, v_t\}$, where u_h and v_h are the hubs of G and H, respectively. Assume that $u \in V(G)$, $v \in V(H)$, $d_G(u) = d_H(v)$, and $G(u) \odot H(v)$ is a brick. The graph $G(u) \odot H(v)$ is wheel-like if and only if the following statements hold.

1). $|\{u,v\} \cap \{u_h,v_h\}| = 1$. Without loss of generality, assume that $u = u_h$, that is $v \neq v_h$. Then $|V(G)| \ge 6$.

2). All the multiple edges of G and H are incident with u_h and v_h , respectively.

3). Without loss of generality, assume that $v = v_t$ and $\{u_1v_1, u_rv_{t-1}\} \subset E(G(u) \odot H(v))$, where $1 \leq r \leq s$. Then $r \neq 1$ and $u_1u_r \notin E(G)$.

Let \mathcal{K}_4^+ be the family of graphs that the underling simple graph is K_4 , and all the multiple edges have the same end vertices. Then $K_4 \in \mathcal{K}_4^+$. And every graph in \mathcal{K}_4^+ is wheel-like; every graph in \mathcal{K}_4^+ , other than K_4 , has exactly two hubs. Let \mathcal{G}_1 be the family of wheel-like odd wheels. It should be noted that every graph in $\mathcal{G}_1 \setminus \mathcal{K}_4^+$ has exactly one hub, and all the multiple edges of it are incident with the hub. For an integer j (j > 0), let \mathcal{G}_{j+1} be the family of graphs with at least 8 vertices gotten by the splicing of one graph in \mathcal{G}_j and one graph in \mathcal{G}_1 , say \mathcal{G}_j and \mathcal{G}_{j+1} , such that 1) if $G_{j+1} = K_4$, then $\{v_j, v_{j+1}\} \cap (\bigcup_{i=1}^{j+1} H_i) = \emptyset$; if $G_{j+1} \in \mathcal{K}_4^+ \setminus \{K_4\}$, then $|\{v_j, v_{j+1}\} \cap (\bigcup_{i=1}^{j+1} H_i)| \ge 1$; if $G_{j+1} \notin \mathcal{K}_4^+$, then $|\{v_j, v_{j+1}\} \cap (\bigcup_{i=1}^{j+1} H_i)| = 1$; and

2) if $|V(G_{j+1})| = 4$ and $v_j \notin \{v_j, v_{j+1}\} \cap (\bigcup_{i=1}^{j+1} H_i)$, then for any nonremovable edge e of $\partial(v_{j+1})$ in G_{j+1} , the corresponding edge of e (incident with v_j) in G_j is not incident with any vertex of $(\bigcup_{i=1}^{j} H_i) \cap V(G_j)$.

Where if $G_i = K_4$ then $H_i = \emptyset$, otherwise H_i is the set of vertices with maximum degree in G_i $(i \in \{1, 2, ..., j+1\})$; v_j and v_{j+1} are the splicing vertices of G_j and G_{j+1} , respectively.

Let \mathcal{G}' be the family of graphs gotten by the splicing of two graphs in $\bigcup_{i=2}^{\infty} \mathcal{G}_i$, say G'_s and G'_t , such that $G'_s \in \mathcal{G}_s$, $G'_t \in \mathcal{G}_t$ and $|\{u'_s, u'_t\} \cap ((\cup_{i=1}^s H_i \cap V(G'_s)) \cup (\cup_{i=1}^t H_i \cap V(G'_t)))| =$ 1, where u'_s and u'_t are the splicing vertices of G'_s and G'_t , respectively $(G'_s$ and G'_t may be the same graph. In this case, $u'_s \neq u'_t$). Let $\mathcal{G} = \mathcal{G}' \cup (\cup_i \mathcal{G}_i)$.

It should be noted that the maximum degree of any graph different from K_4 in \mathcal{G} is at least 4. The vertex of a graph in \mathcal{G} with the maximum degree is called a *hub* of it. Moreover, we have the following lemma.

Lemma 3.9. Let $G \in \mathcal{G}$ and |V(G)| > 4. Then the following statements hold.

1). If $G \in \mathcal{G}_n$ $(n \in \{1, 2, 3, ...\})$, then $|(\bigcup_{i=1}^n H_i) \cap V(G)| = 1$; if $G = G'_j \odot G'_k \in \mathcal{G}'$, then $|((\bigcup_{i=1}^j H_i) \cup (\bigcup_{i=1}^k H_i)) \cap V(G)| = 1$ (where $G'_j \in \mathcal{G}_j$ and $G'_k \in \mathcal{G}_k$). Moreover, the vertex in $(\bigcup_i^n H_i) \cap V(G)$ or $((\bigcup_{i=1}^j H_i) \cup (\bigcup_{i=1}^k H_i)) \cap V(G)$ is the only vertex of maximum degree in G.

2). Every edge incident with the maximum degree of G is removable in G.

Proof. We will prove the result by considering the following two cases.

Case 1. $G \in \mathcal{G}_n$.

We will prove the result by induction on n. If n = 1, every graph in \mathcal{G}_1 is an odd wheel with at least 6 vertices. So, 1) and 2) hold for this case. Suppose that the result holds when n < s. Now we consider $G \in \mathcal{G}_s$, where $s \ge 2$. Then $G = G_{s-1}(v_{s-1}) \odot G_s(v_s)$, where $G_{s-1} \in \mathcal{G}_{s-1}$ and $G_s \in \mathcal{G}_1$. We may assume $|V(G_{s-1})| \ge 6$ (if $G_2 \in \mathcal{G}_1$, by interchanging G_1 with G_2 if necessary, such that $|V(G_1)| \ge 6$). Let $h_t \in (\bigcup_i^{s-1} H_i) \cap V(G_{s-1})$.

As $|V(G_{s-1})| \geq 6$, by induction hypothesis, $(\bigcup_{i=1}^{s-1}H_i) \cap V(G_{s-1}) = \{h_t\}$, and h_t is the only vertex of maximum degree in $V(G_{s-1})$. Obviously, $d_{G_{s-1}}(h_t) > 3$. If $G_s = K_4$, then $\{v_{s-1}, v_s\} \cap (\bigcup_{i=1}^{s}H_i) = \emptyset$, that is, $(\bigcup_{i=1}^{s}H_i) \cap V(G) = \{h_t\}$. As every vertex in K_4 is of degree 3, h_t is also the only vertex of maximum degree in G. If $G_s \in \mathcal{K}_4^+ \setminus \{K_4\}$, let h'_s and h_s be the hubs of G_s . If $v_{s-1} = h_t$, then $v_s \in \{h'_s, h_s\}$, as $d_{G_{s-1}}(v_{s-1}) \geq 4$ and every vertex of $V(G_s) \setminus \{h'_s, h_s\}$ is of degree 3 in $V(G_s)$. Without loss of generality, assume that $v_s = h_s$. Then $(\bigcup_{i=1}^{s}H_i) \cap V(G) = \{h'_s\}$. As $d_{G_s}(h'_s) = d_{G_s}(h_s) = d_{G_{s-1}}(h_t)$, h'_s and h_s are vertices of maximum degree in G_s and h_t is the only vertex of maximum degree in $V(G_{s-1})$, we have h'_s is the only vertex of maximum degree in G. If $v_{s-1} \neq h_t$, then $v_s \in \{h'_s, h_s\}$. As $d_G(h_t) = d_{G_{s-1}}(h_t) \geq d_{G_{s-1}}(v_{s-1}) = d_{G_s}(v_s)$, v_s is a vertex of maximum degree in $V(G_s)$ and $h_t \in V(G)$. So h_t is the only vertex of maximum degree in G. If $G_s \notin \mathcal{K}^+_4$, assume that h_s is the hub of G_s . Then $|\{v_{s-1}, v_s\} \cap \{h_t, h_s\}| = 1$. So $|(\cup_i^s H_i) \cap V(G)| = 1$ and the vertex of $(\cup_i^s H_i) \cap V(G)$ is the only vertex of maximum degree in G (recalling that $d_{G_{s-1}}(v_{s-1}) = d_G(v_s)$). Therefore, 1) holds.

As $|V(G_{s-1})| \ge 6$, by induction hypothesis, every edge of $\partial(h_t)$ is removable in G_{s-1} . If the underlying simple graph of G_s is not isomorphic to K_4 , then $|\{v_{s-1}, v_s\} \cap \{h_t, h_s\}| = 1$. Assume that $v_{s-1} = h_t$ and $v_s \neq h_s$ (the case when $v_s = h_s$ is similar). By 1) of this lemma again, h_s is the only hub of G. Then by induction hypothesis, every edge of $\partial_{G_s}(h_s)$ is removable in G_s . Therefore, every edge of $\partial(h_s)$ is removable in G by Lemma 2.10, and then 2) holds in this case. Now assume that the underlying simple graph of G_s is isomorphic to K_4 .

If $G_s \in \mathcal{K}_4^+$ and $v_{s-1} \neq h_t$, then h_t is the only hub of G by 1) of this lemma. As the corresponding edge (incident with v_{s-1}) of any nonremovable edge of G_s is not incident with h_t (h_t is also the only hub of G_{s-1}) and every edge of $\partial_{G_{s-1}}(h_t)$ is removable in G_{s-1} , by Lemma 2.10, every edge of $\partial_G(h_t)$ is also removable in G.

If $G_s \in \mathcal{K}_4^+$ and $v_{s-1} = h_t$, then $d_{G_s}(v_s) = d_{G_{s-1}}(v_{s-1}) \ge 4$ and so, $G_s \neq K_4$ and $\{v_{s-1}, v_s\} \subset \{h_t, h_s, h'_s\}$. Without loss of generality, assume that $v_s = h_s$. Then h'_s is the only hub of G. By Lemma 2.10, every removable edge of $\partial(h'_s)$ is removable in G. For any nonremovable edge e of $\partial(h'_s)$ in G_s , there exists an edge e' which is incident with v_s in G_s , such that $\{e, e'\}$ is a removable doubleton of G_s . By Lemma 2.11, e is removable in G. So every edge of $\partial_G(h'_s)$ is also removable in G.

Now assume that $G_s \notin \mathcal{K}_4^+$. Then h_s is the only hub of G_s . By induction hypothesis, $|\{v_{s-1}, v_s\} \cap \{h_t, h_s\}| = 1$. If $v_{s-1} = h_t$, then h_s is the only hub of G by 1) of this lemma. By induction and Lemma 2.10, every edge of $\partial(h_s)$ is removable in G. If $v_s = h_s$, then h_t is the only hub of G by 1) of this lemma again. As $G_s \notin \mathcal{K}_4^+$, v_s is incident with at most one nonremovable edge. Let f be the only possible nonremovable edge of $\partial(v_s)$ in G_s . Then the corresponding edge of f (incident with v_{s-1}) in G_{s-1} is not incident with h_t . So, every edge of $\partial(h_t)$ is removable in G. Therefore, 2) holds.

Case 2. $G = G'_i(u'_i) \odot G'_k(u'_k) \in \mathcal{G}'.$

As $G'_j, G'_k \in \bigcup_{i=2}^{\infty} \mathcal{G}_i, G'_j$ and G'_k have only one vertex of maximum degree in G'_j and G'_k , resp., by Case 1. Denote by h'_j and h'_k the vertices of maximum degree in G'_j and G'_k , resp.. Then $|\{u'_j, u'_k\} \cap \{h'_j, h'_k\}| = 1$, where $u'_j \neq u'_k$. As $d_{G'_j}(u'_j) = d_{G'_k}(u'_k)$, the vertex of

 $\{h'_j, h'_k\} \setminus (\{u'_j, u'_k\} \cap \{h'_j, h'_k\})$ is the only vertex of maximum degree in G. So 1) holds for this case.

Without loss of generality, assume that $u'_j = h'_j$. Then h'_k is the only hub of G by 1) of this Lemma. As $G'_j, G'_k \in \bigcup_{i=2}^{\infty} \mathcal{G}_i$, by Case 1 every edge of $\partial(u'_j)$ and $\partial(h'_k)$ is removable in G'_j and G'_k , resp.. By Lemma 2.10, every edge of $\partial(h'_k)$ is removable in G, 2 holds for this case.

Theorem 1.3. Let G be a wheel-like brick. Then $G \in \mathcal{G}$.

Proof. If G is solid, then by Proposition 3.1, the underlying simple graph of G is an odd wheel (all the multiple edges are incident with its hub). So the result holds. Now we assume that G is nonsolid. By Proposition 2.7, assume that $\partial(X)$ is a robust cut of G.

We will prove the result by induction on |V(G)|. By Lemma 3.6, the result holds when |V(G)| = 6. Now we assume that the result holds for every wheel-like brick with at most *n* vertices (*n* is even and n > 6). In the following we will consider the case when |V(G)| = n + 2.

As G is nonsolid, there exist a subset X' of \overline{X} and a subset X" of X such that $G/(\overline{X'} \to \overline{x'})$ and $G/(\overline{X''} \to \overline{x''})$ are bricks, and $(G/(X' \to x'))/(X'' \to x'')$ is a bipartite matching covered graph by Theorem 2.8. Let $G' = G/\overline{X'}$, $G'' = G/\overline{X''}$ and $H = (G/(X' \to x'))/(X'' \to x'')$. Then every edge of $\partial_H(x')$ and $\partial_H(x'')$ is removable in H by Lemma 2.9.

Claim 1. G' and G'' are wheel-likes.

Proof. By Theorem 1.1, G' and G'' contain at least three removable classes, respectively. Suppose that there exist removable classes R_1 and R_2 in G' and G'', respectively, such that $R_1 \cap \partial(\overline{x'}) = \emptyset$ and $R_2 \cap \partial(\overline{x''}) = \emptyset$. Then there exist two edges, e_1 and e_2 , such that $e_1 \in R_1$, $e_2 \in R_2$ and both of e_1 and e_2 are removable in G by Lemmas 2.10 and 2.11. Note that e_1 and e_2 are nonadjacent, contradicting the assumption that G is wheel-like. So at least one of G' and G'' is wheel-like such that $\overline{x'}$ or $\overline{x''}$ is its hub.

Without loss of generality, assume that G' is wheel-like with $\overline{x'}$ as its hub. By the induction hypothesis, every edge incident with $\overline{x'}$ belongs to a removable class (more exactly, every edge of $\partial(\overline{x'})$ is removable unless the underlying simple graph of G' is K_4 by Lemma 3.9). Note that every edge incident with x' is removable in H. If every removable class of G'' contains an edge incident with $\partial(\overline{x''})$, then G'' is wheel-like obviously. So we assume that there exists a removable class R_3 in G'' such that $R_3 \cap \partial(\overline{x''}) = \emptyset$. $|R_3|$ may be 1 or 2. In each case, by Lemmas 2.10 and 2.11, R_3 contains a removable edge e_0 in G. Suppose that $|N_H(x')| \geq 2$. Then there exists a removable edge e in G such that both ends of e belong to $X' \cup N(X') \setminus X''$ by Lemma 2.17. So e and e_0 are two nonadjacent

removable edges in G, contradicting the assumption that G is wheel-like. Therefore, we have $|N_H(x')| = 1$. As H is matching covered, |V(H)| = 2 and $V(H) = \{x', x''\}$. So G is isomorphic to $G'(\overline{x'}) \odot G''(\overline{x''})$. By Lemma 3.7, G'' is wheel-like since G is wheel-like. \Box

Let h' and h'' be hubs of G' and G'', resp.. By the proof of Claim 1, without loss of generality, assume that $\overline{x'} = h'$. Let $V(G'') \setminus h'' = \{v''_1, v''_2, \ldots, v''_n\}$. Then we have the following claim.

Claim 2. $V(H) = \{x', x''\}.$

Proof. Suppose to the contrary that $|N_H(x')| \ge 2$. By Lemma 2.17, there exists a removable edge e_1 in G such that both ends of e_1 belong to $X' \cup N(X') \setminus X''$. If $\overline{x''} = v_i''$ $(i \in \{1, 2, ..., n\})$, every edge of $\partial(h'') \setminus E[\{v_i''\}, \{h''\}]$ belongs to some removable class in G'' (in fact, if the underlying simple graph of G'' is not K_4 , every edge of $\partial(h'') \setminus E[\{v_i''\}, \{h''\}]$ is removable in G''). Let $e \in \partial(h'') \setminus E[\{v_i''\}, \{h''\}]$. Note that $e \notin \partial(\overline{x''})$. If e is a removable edge in G'', e is also a removable edge in G by Lemma 2.10; and if e lies in a removable doubleton of G'', one edge in this removable doubleton is removable edge in G by Lemma 2.11. So there exists an edge e_2 that is removable in G and both ends of e_2 belong to X''. Hence, e_1 and e_2 are two nonadjacent removable edges in G, contradicting the assumption that G is wheel-like. So $\overline{x''} = h''$. Suppose that $|N_H(x'')| \ge 2$. Then there exists an edge e_3 that is removable in G and both ends of e_3 belong to $X'' \cup N(X'') \setminus X'$. Hence, e_1 and e_3 are two nonadjacent removable edges in G, a contradiction. Therefore, we have $|N_H(x'')| = 1$. Since H is matching covered, it is connected. Then $|N_H(x'')| = |x'|$. So $V(H) = \{x', x''\}$.

By Claim 2, we have $G = G'(\overline{x'}) \odot G''(\overline{x''})$. As |V(G)| > 6, at least one of |V(G')| and |V(G'')| is at least 6. Moreover, we have the following claim.

Claim 3. Assume that G' has at least 6 vertices and the underlying simple graph of G'' is K_4 . Denote by H'' the set consisting of all hubs of G''. Then the following statements hold.

1) If $G'' = K_4$, then $\{\overline{x'}, \overline{x''}\} \cap (\{h'\} \cup H'') = \emptyset$; if $G'' \in \mathcal{K}_4^+ \setminus \{K_4\}$, then $|\{\overline{x'}, \overline{x''}\} \cap (\{h'\} \cup H'')| \ge 1$; if $G'' \notin \mathcal{K}_4^+$, then $|\{\overline{x'}, \overline{x''}\} \cap (\{h'\} \cup H'')| = 1$.

2) If $\overline{x'} \neq h'$, then for any nonremovable edge e of $\partial(\overline{x''})$ in G'', the corresponding edge of e (incident with $\overline{x'}$) in G' is not incident with h'.

Proof. By Claim 1, G' and G'' are wheel-like bricks. If $G'' = K_4$, then $\overline{x'} \neq h'$, as $d_{G'}(h') \geq 4$ and every vertex of V(G'') is of degree 3. Recall that $H'' = \emptyset$ (as $G'' = K_4$).

So $\{\overline{x'}, \overline{x''}\} \cap (\{h'\} \cup H'') = \emptyset$. Now assume that $G'' \neq K_4$. Then $H'' \neq \emptyset$. As G is a wheellike, by Lemma 3.7, $|\{\overline{x'}, \overline{x''}\} \cap (\{h'\} \cup H'')| \geq 1$. Assume that $|\{\overline{x'}, \overline{x''}\} \cap (\{h'\} \cup H'')| = 2$, that is, $\overline{x'} = h'$ and $\overline{x''} \in H''$. By induction hypothesis and Lemma 3.9, every edge of $\partial(h')$ is removable in G'. Then there exists a vertex s of G'', such that every removable edge of G'' belongs to $E[\{\overline{x''}\}, \{s\}]$, that is, $G'' \in \mathcal{K}_4^+$. Therefore, 1) holds.

Suppose, to the contrary, that there exists a nonremovable edge e of $\partial(\overline{x''})$ in G'', and the corresponding edge of e (incident with $\overline{x'}$) in G', say e', is incident with h'. As the underlying simple graph of G'' is K_4 , there exist two vertices s and t in $V(G'') \setminus \{\overline{x''}\}$, such that $\{e, st\}$ is a removable doubleton of G''. Note that every edge of $\partial(h')$ is removable in G'. By Lemma 2.11, st is removable in G and is not incident with h'. On the other hand, every removable edge of $\partial(h') \setminus \partial(\overline{x'})$ is also removable in G by Lemma 2.10. As $\partial(h') \setminus \partial(\overline{x'})$ contains at least two edges with exactly one common end h', and G is wheel-like, we have every removable edge of G is incident with h', a contradiction. So 2) holds.

By Claim 3, the result holds if the underlying simple graph of G' or G'' is K_4 . So assume that $|V(G')| \ge 6$ and $|V(G'')| \ge 6$. As $G', G'' \in \mathcal{G}$ (by induction hypothesis), each of G' and G'' has exactly one hub by 1) of Lemma 3.9. As G is wheel-like and both of G' and G'' are bricks, we have $|\{\overline{x'}, \overline{x''}\} \cap \{h', h''\}| \ge 1$ by Lemma 3.7. Suppose to the contrary that $|\{\overline{x'}, \overline{x''}\} \cap \{h', h''\}| = 2$. By 2) of Lemma 3.9, every edge of $\partial_{G'}(\overline{x'})$ and $\partial_{G''}(\overline{x''})$ is removable in G' and G'', resp.. Then by Lemma 2.10, every edge of $\partial_G(V(G') \setminus \{\overline{x'}\})$ is removable in G. As G is a brick, there exist at least two nonadjacent edges of $\partial_G(V(G') \setminus \{\overline{x'}\})$, contradicting the assumption that G is wheel-like. Therefore, $|\{\overline{x'}, \overline{x''}\} \cap \{h', h''\}| = 1$, that is, $G \in \mathcal{G}$.

It should be noted that not every graph in \mathcal{G} is wheel-like. Lemma 3.8 will help to determine when the splicing of two odd wheels is wheel-like. When n > 2, some edge not adjacent to the hub of \mathcal{G}_n will be removable, even if the splicing between any two odd wheels satisfying the condition in Lemma 3.8 (see Figure 3 for example).

Proposition 3.10. Let G be a brick such that every removable edge of it is incident with a vertex h. Then every edge of $\partial(h)$ is removable or there exists a vertex $u \in V(G) \setminus \{h\}$ such that $d_G(u) = 3$.

Proof. If G has a removable doubleton, then the underlying simple graph H of G is K_4 , $\overline{C_6}$ or R_8 by Theorem 3.3. If G = H, then the result holds as every vertex of K_4 , $\overline{C_6}$ and R_8 is of degree 3. Assume that G has a multiple edge. Then every multiple edge of G is incident with h and so, $d_G(h) \ge 4$. It can be checked that the result holds when $H \cong K_4$.



Figure 3: A brick in \mathcal{G}_3 (the bold edges are removable)

Assume that $H \cong \overline{C_6}$ or $H \cong R_8$. Note that there exists $v' \in V(G) \setminus \{h\}$, such that $v' \notin N_G(h)$. As every multiple edge of G belongs to $E[\{h\}, N_G(h)]$ and every vertex of H has degree 3, we have $d_G(v') = 3$.

If G has no removable doubletons, then G is wheel-like. Note that every edge incident with the hub in an odd wheel with at least 6 vertices is removable. Then the result holds by Lemma 3.6, Lemma 3.9 and Theorem 1.3.

Proposition 3.11 ([9]). Let G be a bicritical graph and let C be a 2-separation cut of G. Then the two C-contractions of G are both bicritical.

Proposition 3.12. Let G be a bicritical graph and let $\{u, v\}$ be a 2-separation of G. Then $d(u) \ge 4$ and $d(v) \ge 4$.

Proof. Suppose to the contrary that $d(u) \leq 3$. Let $\partial(X)$ be a 2-separation cut of G associated with $\{u, v\}$ such that $u \in X$. Then $|N(u) \cap X| = 1$ or $|N(u) \cap \overline{X}| = 1$. Without loss of generality, assume that $N(u) \cap X = \{s\}$. Then $G - \{s, v\}$ is disconnected. Moreover, the component of $G - \{s, v\}$ containing u is an odd component. So $G - \{s, v\}$ has no perfect matchings, contradicting the fact that G is bicritical.

Proposition 3.13 ([14]). Let G be a bicritical graph without removable edges. Then G has at least four vertices of degree three. As a consequence, every bicritical graph with minimum degree at least 4 has removable edges.

Lemma 3.14. Let G be a bicritical graph with a removable edge. Assume that every removable edge of G is incident with a vertex h. Then every edge of $\partial(h)$ is removable or there exists a vertex s in $V(G) \setminus \{h\}$ such that d(s) = 3.

Proof. If G is 3-connected, that is, G is a brick, then the result holds by Proposition 3.10. So we assume that G is not 3-connected. As G is bicritical, G has a 2-separation by

Corollary 2.4. Let a bicritical graph G be chosen with |V(G)| minimum such that some edge of $\partial(h)$ is not removable and every vertex in $V(G) \setminus \{h\}$ is of degree at least 4.

As K_4 is a brick, we have $|V(G)| \ge 6$. Let $\{u, v\}$ be a 2-separation of G. Assume that $\partial(X)$ is a 2-separation cut associated with $\{u, v\}$ such that $u \in X$. Let $H_1 = G/(\overline{X} \to \overline{x})$ and $H_2 = G/(X \to x)$. Then by Proposition 3.11, H_1 and H_2 are bicritical. Without loss of generality, assume that $h \in X$.

We first suppose that H_2 contains no removable edges. Then $V(H_2) \setminus \{x, v\}$ contains a vertex s_1 of degree 3 by Proposition 3.13. So $d_G(s_1) = d_{H_2}(s_1) = 3$. Now we suppose that H_2 contains a removable edge. By Lemma 2.10, the possible removable edges of H_2 are incident with the vertex x (in this case, h = u), or belong to $E[\{x\}, \{v\}]$. Then all the removable edges of H_2 are incident with x. As $|V(H_2)| < |V(G)|, V(H_2) \setminus \{x\}$ contains a vertex of degree 3 or every edge of $\partial(x)$ is removable in H_2 by the minimality of V(G). If $V(H_2) \setminus \{x\}$ contains a vertex of degree 3, then this vertex, different from h, is also of degree 3 in G. Now we consider the case when every edge of $\partial(x)$ is removable in H_2 . If every edge of $\partial_{H_1}(u)$ is removable in H_1 , then every edge of $\partial_G(u)$ is removable in G by Lemma 2.10. So we assume that some edge of $\partial_{H_1}(u)$ is not removable in H_1 . If every edge of $\partial_{H_1}(u)$ is not removable in H_1 , then H_1 contains no removable edges. Similar to the case when H_2 contains no removable edges, the result follows. So $\partial_{H_1}(u)$ contains removable edges and nonremovable edges in H_1 . Then every removable of H_1 is incident with u. As $|V(H_1)| < |V(G)|, V(H_1) \setminus \{u\}$ contains a vertex s_2 of degree 3 by the minimality of V(G). Then $d_G(s_2) = d_{H_2}(s_2) = 3$. The result follows.

4 Proof of Theorem 1.4

Let B be a maximal nontrivial barrier of a matching covered nonbipartite graph G. Denote by H(G, B) the graph obtained from G by contracting every nontrivial odd component of G - B to a singleton. By Corollary 2.2, B is an independent set and G - B has no even components. So H(G, B) is a bipartite graph with B as one of its color classes. Let $I = V(H(G, B)) \setminus B$. Note that $v \in V(G) \cap V(H)$ if and only if v is not gotten by contracting a nontrivial odd component of G - B. Let $U_{H(G,B)} = \{u \in I : u \text{ is incident with some removable edge of } H(G,B)\}$. As each odd component of G - B is a shore of a barrier cut of G, H(G,B) is a matching covered bipartite graph.

Theorem 1.4. Let G be a minimal matching covered graph with at least four vertices. Then $\delta(G) = 2$ or 3.

Proof. By Propositions 2.3 and 3.13, if a minimal matching covered graph has no nontriv-

ial barriers, then the result holds. So we consider the case when the minimal matching covered graph has a nontrivial barrier. Let a minimal matching covered G that has minimum degree at least 4 be chosen with |V(G)| minimum. If |V(G)| = 4, G is a cycle with 4 vertices or K_4 , both of which contain a vertex of degree less than 4. Then |V(G)| > 4. We have the following claim.

Claim A. For any maximal nontrivial barrier B of G, we have $|U_{H(G,B)}| \ge 2$. Moreover, for every vertex $u \in U_{H(G,B)}$, $u \notin V(G)$.

Proof. If $\delta(H(G, B)) \geq 3$, by Lemma 2.16, H(G, B) has two nonadjacent removable edges and then, $|U_{H(G,B)}| \geq 2$. So we will show that $\delta(H(G, B)) \geq 3$. Suppose, to the contrary, that there exists a vertex $k \in H(G, B)$ such that $d_{H(G,B)}(k) = 2$ (As H(G, B)is matching covered, H(G, B) is 2-connected). Then $k \in I$ and $k \notin V(G)$, as $B \subset V(G)$ and $\delta(G) \geq 4$. Assume that k is obtained by contracting the odd component K of G - B. Let $H_1 = G/(\overline{K} \to \overline{k})$. Then H_1 is matching covered as $\partial(V(K))$ is a barrier cut of G. Note that every edge of $E(H_1) \setminus \partial_{H_1}(\overline{k})$ is not removable in H_1 by Lemma 2.10, as G has no removable edges. As $d_{H_1}(\overline{k}) = 2$, it can be checked that every edge of $\partial_{H_1}(\overline{k})$ is not removable in H_1 . Then H_1 is minimal matching covered. Let $N_{H_1}(\overline{k}) = \{k_1, k_2\}$. Then $\partial_{H_1}(\{k, k_1, k_2\})$ is a tight cut of H_1 . Let $H_2 = H_1/(\{k, k_1, k_2\} \to t)$. As $\{k_1, k_2\} \subset V(G)$, the degrees of k_1 and k_2 are at least 4, respectively. So $d_{H_2}(t) \geq 4$. Since H_1 contains no removable edges, every edge incident with t is not removable in H_2 by Lemma 2.10. As $V(H_2) \setminus \{t\} \subset V(G), \, \delta(H_2) \geq 4$. Therefore, H_2 is a minimal matching covered graph with $|V(H_2)| < |V(G)|$, contradicting the choice that |V(G)| is minimum. So $\delta(H(G, B)) \geq 3$.

Suppose, to the contrary, that there exists a vertex $u \in V(G) \cap U_{H(G,B)}$. Since H(G, B)is bipartite, we may assume that ub is removable in H(G, B) where $b \in B$. As $u, b \in V(G)$, ub is removable in G by Lemma 2.10, contradicting the assumption that G has no removable edges. So $V(G) \cap U_{H(G,B)} = \emptyset$.

Let B_0 be a maximal nontrivial barrier of G. By Claim A, we may choose a nontrivial odd component of $G - B_0$, say Q_0 , such that $q_0 \in U_{H(G,B_0)}$, where the vertex q_0 is gotten by contracting the component Q_0 . Let $G_1 = G/(\overline{V(Q_0)} \to \overline{q_0})$. Then G_1 is not bipartite. Otherwise, suppose that A' and B' are two color classes of G_1 such that $\overline{q_0} \in B'$. As $|V(G_1)| \ge 4, B_0 \cup B' \setminus {\overline{q_0}}$ is a barrier of G satisfying $|B_0 \cup B' \setminus {\overline{q_0}}| > |B_0|$, which contradicts the maximality of B_0 . If G_1 is not bicritical, assume that B_1 is a maximal nontrivial barrier of G_1 . We have the following claim.

Claim B. There exists a nontrivial odd component Q_1 of $G_1 - B_1$ such that $\overline{q_0} \notin Q_1$ (*i.e.*, $V(Q_1) \subset V(G)$), and $q_1 \in U_{H(G_1,B_1)}$, where q_1 is obtained by contracting the odd component Q_1 of $G_1 - B_1$.

Proof. As B_0 is maximal and B_1 is not trivial, $\overline{q_0} \notin B_1$. Otherwise, $B_0 \cup B_1 \setminus \{\overline{q_0}\}$ is a barrier of G satisfying $|B_0 \cup B' \setminus \{\overline{q_0}\}| > |B_0|$, which contradicts the maximality of B_0 . So $\overline{q_0} \in V(H(G_1, B_1)) \setminus B_1$ or $\overline{q_0}$ lies in some component of $G_1 - B_1$. Let q be a vertex in $V(H(G_1, B_1)) \setminus B_1$ such that $q = \overline{q_0}$ if $\overline{q_0} \in V(H(G_1, B_1)) \setminus B_1$, otherwise q is the vertex gotten by contracting the nontrivial odd components of $G_1 - B_1$ that contains $\overline{q_0}$.

If $d_{H(G_1,B_1)}(q) = 2$, then by Lemma 2.16, there exists a vertex u in $H(G_1, B_1)$ such that d(u) > 3 and every edge of $\partial_{H(G_1,B_1)}(u)$ is removable in $H(G_1, B_1)$. Let e be an edge in $H(G_1, B_1)$ that is incident with u, but is not incident with q. Let u' be the end of e not in B_1 (u and u' may be the same vertex). Then the nontrivial odd components of $G_1 - B_1$ contracted to u' is that we need. If $d_{H(G_1,B_1)}(q) > 2$, similar to the proof of Claim A (with G_1 as G and B_1 as B), we have $|U_{H(G_1,B_1)}| \ge 2$. So there exists a vertex q_1 in $U_{H(G_1,B_1)}$ such that $q_1 \neq q$. Then the nontrivial odd components of $G_1 - B_1$ contracted to u' is that we need. If $d_{H(G_1,B_1)}| \ge 2$.

It is known that all the maximal barriers in G is a partition of V(G) (see Lemma 5.2.1 in [9]). We may contract several maximal barriers to get a bicritical graph with given property. Let $G_2 = G_1/(\overline{V(Q_1)} \to \overline{q_1})$. If G_2 has a maximal nontrivial barrier B_2 , similar to Claim B, we may assume that Q_2 is a nontrivial component of $G_2 - B_2$ such that $V(Q_2) \subset V(G)$, and $q_2 \in U_{H(G_2,B_2)}$, where q_2 is obtained by contracting the odd component Q_2 of $G_2 - B_2$. And then let $G_3 = G_2/(\overline{V(Q_2)} \to \overline{q_2})$ Continue above steps, we finally obtain a matching covered graph G_s satisfying the following:

1) G_s has a maximal nontrivial barrier B_s and $V(G_s) \setminus \{\overline{q_{s-1}}\} \subset V(G)$; and

2) there exists a nontrivial component Q_s of $G_s - B_s$, $G_s/(\overline{Q_s} \to \overline{q_s})$ has no nontrivial barriers, $V(Q_s) \subset V(G)$ and $q_s \in U_{H(G_s,B_s)}$, where q_s is obtained by contracting the odd component Q_s of $G_s - B_s$.

Let $G' = G_s/(\overline{Q_s} \to \overline{q_s})$. As G' is a bicritical graph, $V(G') \setminus \{\overline{q_s}\} \subset V(G)$ and G is minimal matching covered graph, if G' has a removable edge then this edge is incident with $\overline{q_s}$ by Lemma 2.10. Noting $q_s \in U_{H(G_s,B_s)}$, q_s is incident with a removable edge in $H(G_s, B_s)$, say e. Then the corresponding edge of e (incident with $\overline{q_s}$) in G' is not removable in G' by Lemma 2.10 again. By Lemma 3.14, there exists a vertex $v \in V(G') \setminus$ $\{\overline{q_s}\}$ such that $d_{G'}(v) = 3$. As $V(G') \setminus \{\overline{q_s}\} \subset V(G)$, we have $d_G(v) = 3$, contradicting the assumption that $\delta(G) \geq 4$. So the theorem holds.

References

- [1] J. A. Bondy and U. S. R. Murty, Graph Theory, Springer-Verlag, Berlin, 2008.
- [2] M. H. Carvalho, C. L. Lucchesi, and U. S. R. Murty, Ear decompositions of matching covered graphs, Combinatorica, 19: 151-174, 1999.
- [3] M. H. Carvalho, C. L. Lucchesi and U. S. R. Murty, On a conjecture of Lovász concerning bricks. II. Bricks of finite characteristic, J. Combin. Theory Ser. B, 85: 137-180, 2002.
- [4] M. H. Carvalho, C. L. Lucchesi and U. S. R. Murty, How to build a brick, Discrete Mathematics, 306: 2383-2410, 2006.
- [5] M. H. Carvalho, C. L. Lucchesi and U. S. R. Murty, Thin edges in braces, Electron. J. Combin., 22(4), #P4.14, 2015.
- [6] J. Edmonds, L. Lovász and W. R. Pulleyblank, Brick decompositions and the matching rank of graphs, Combinatorica, 2(3): 247-274, 1982.
- [7] C. L. Lucchesi and U. S. R. Murty, Perfect Matchings, Springer, 2024.
- [8] L. Lovász and M. D. Plummer, On minimal elementary bipartite graphs, J. Combin. Theory Ser. B, 23: 127-138, 1977.
- [9] L. Lovász and M. D. Plummer, Matching Theory, Annals of Discrete Mathematics, vol. 29, Elsevier Science, 1986.
- [10] L. Lovász. Ear decompositions of matching covered graphs, Combinatorica, 2: 105-117, 1983.
- [11] L. Lovász, Matching structure and the matching lattice, J. Combin. Theory Ser. B, 43: 187-222, 1987.
- [12] F. Lu and J. Xue, Planar wheel-like bricks, http://arxiv.org/abs/2410.20692.
- [13] W. T. Tutte, The factorization of linear graphs, J. Lond. Math. Soc. 22: 107-111, 1947.
- [14] Y. Zhang, X. Wang and J. Yuan, Bicritical graphs without removable edges, Discrete Applied Mathematics, 320: 1-10, 2022.