ON TORSION-FREE MODULES AND SEMI-HEREDITARY RINGS

RYOYA ANDO

Department of Mathematics, Faculty of Science and Technology, Tokyo University of Science, Chiba, Japan

SKILLUP NeXt Ltd., Chiyoda, Japan

Abstract. The class of semi-hereditary rings is an important class of rings in theories that do not assume the Noetherian condition, such as perfectoid ring theory. We prove several results concerning the structure theory of this class, focusing on the relationship between semi-hereditary rings and the flatness of torsion-free modules. We also consider Shimomoto's problem concerning the flatness of the Frobenius map.

1. INTRODUCTION

Throughout this paper, all rings are commutative with the identity element.

In recent years, rings without Noetherian conditions have been increasingly used in homological studies in commutative ring theory, exemplified by perfectoid ring theory. The class of coherent rings, which also appears in the context of perfectoid ring theory, is one of the important classes of rings that includes the class of Noetherian rings. For example, it is important in applications such as [\[AMM22\]](#page-8-0) and [\[Shi11\]](#page-9-0).

We discuss several observations on semi-hereditary rings, a subclass of coherent rings, which can be regarded as a generalization of Dedekind domains.

In §[2](#page-1-0) and §[3,](#page-1-1) we give a brief review of semi-hereditary rings, torsion-free modules, and some properties of Prüfer domains.

In $\S4$, we prove that the ring A is semi-hereditary if and only if every torsion-free A-module is flat [\(Theorem 4.10\)](#page-5-0). This theorem is an extension of Chase's theorem ([\[Cha60,](#page-9-1) Theorem 4.1]) , [Theorem 2.2](#page-1-2) of this paper).

In §[5,](#page-5-1) we will consider propositions concerning the decomposition of semihereditary rings into direct products in certain cases. While there are previous studies on the decomposition theorem for semi-hereditary rings, such as [\[Fai75\]](#page-9-2), [\[Dro80\]](#page-9-3), and [\[MO01\]](#page-9-4), we will attempt a different approach from those.

E-mail address: [6122701@ed.tus.ac.jp,](6122701@ed.tus.ac.jp) r_ando@skillupai.com. Date: December 24, 2024.

²⁰²⁰ Mathematics Subject Classification. 13C05 (Primary) 16E60 (Secondary).

Key words and phrases. Semi-hereditary rings, torsion-free modules, coherent rings.

Finally, in §[6,](#page-7-0) we show that the main theorem can be applied to several propositions in perfectoid ring theory. In perfectoid ring theory, torsionfree modules play an important role in several propositions originating from [\[Lur20,](#page-9-5) Theorem 3.2.], inspired by O. Gabber. The results of this work solve [\[Shi,](#page-9-6) Open Problem 7.13.].

2. Semi-hereditary rings and torsion-free modules

As for coherent rings, a well-known textbook is [\[Gla89\]](#page-9-7), which summarizes the properties of semi-hereditary rings. Additionally, the classes of rings such as semi-hereditary, Prüfer, and von Neumann regular that appear in this paper are also well summarized in [\[Lam99\]](#page-9-8). We will recall the definitions of these concepts.

Definition 2.1. Let A be a ring. A is called **hereditary** if every ideal of A is projective. Also, A is called **semi-hereditary** if every finitely generated ideal of A is projective.

As basic examples, valuation rings and Dedekind domains are semi-hereditary. We will introduce the characterization of semi-hereditary rings given by [\[Cha60\]](#page-9-1). Recall that an A-module M is called **torsion-less** if the following

map is injective;

 $M \to \text{Hom}(\text{Hom}(M, A), A); x \mapsto (f \mapsto f(x)).$

Theorem 2.2 ([\[Cha60,](#page-9-1) theorem 4.1]). *A ring* A *is semi-hereditary if and only if every torsion-less* A*-module is flat.*

Let us also consider torsion-free modules here. Let us recall the definition.

Definition 2.3. An A-module M is called **torsion-free** if for any non-zero divisor $a \in A$, the natural map $a \cdot : M \to M$ is injective.

Obviously, a torsion-less module is torsion-free, and a flat module is also torsion-free. When does the converse hold?

For example, the $\mathbb{Z}\text{-module} \mathbb{Q}$ is torsion-free but not torsionless (since $\text{Hom}_{\mathbb{Z}}(\mathbb{Q},\mathbb{Z})=0$. So, when does a torsion-free module become flat? Chase's theorem ([\[Cha60,](#page-9-1) Theorem 4.2], [Theorem 3.3](#page-2-0) of this paper) characterizes rings where flatness and torsion-free-ness are equivalent in the case of integral domains. Such rings are Prüfer domains.

Here, by [Theorem 2.2,](#page-1-2) we pay attention to the fact that a ring where flatness and torsion-free-ness are equivalent is semi-hereditary. In this paper, we extend this theorem and prove that the class of rings where flatness and torsion-free-ness are equivalent, even in the general case, is exactly the class of semi-hereditary rings [\(Theorem 4.10\)](#page-5-0). We also summarize the situation for the integral domain case, Noetherian case, and general case.

3. Integral domain cases

In the case of integral domains, the key concept is that of Prüfer domains, which are known as a generalization of Dedekind domains. Let us begin by reviewing the definition.

Definition 3.1. A domain A is called a Prüfer domain, if for any prime ideal P of A, A_P is a valuation ring.

This condition is equivalent to any finitely generated ideal being projective. Although this fact is well known, We will provide a proof.

Proposition 3.2. *Let* A *be a integral domain. The following conditions are equivalent:*

- (i) A *is a Prüfer domain*;
- (ii) *Every finitely generated ideal is projective;*
- (iii) *Every ideal is flat.*

Proof.

 $(i) \implies (ii)$

Let I be a finitely generated ideal of A. For any $P \in \text{Spec } A$, I_P is a finitely generated ideal of A_P , so it is principal. Since A_P is a domain, I_P is free, and hence projective. Since projectivity is a local property, I is projective. $(ii) \implies (iii)$

Let I be an ideal of A. It can be expressed as $I = \lim I_i$ by a family of finitely generated ideals $\{I_i\}$. Since each I_i is projective, for any A-module M, we have $\text{Tor}_1(I, M) = \text{Tor}_1(\varinjlim I_i, M) = \varinjlim \text{Tor}_1(I_i, M) = 0.$ Hence, I is flat.

 $(iii) \implies (i)$

For any $P \in \text{Spec } A$ and finitely generated ideal J of A_P , J is the localization of an ideal of A, so it is flat. Therefore, by Kaplansky's theorem [\[Kap58\]](#page-9-9), it must be free and principal. Hence, A_P is a valuation ring.

Let us give a simple proof of [\[Cha60,](#page-9-1) Theorem 4.2].

Theorem 3.3 ([\[Cha60,](#page-9-1) Theorem 4.2]). *Let* A *be a domain. The following conditions are equivalent:*

- (i) A *is a Prüfer domain*;
- (ii) *Every torsion-free* A*-module is flat.*

Proof. (i) \implies (ii): Let M be a torsion-free A-module. It suffices to show for any $P \in \text{Spec } A, M_P$ is flat. Now, M_P is a torsion-free A_P module. This is because if $a/s \neq 0 \in A_P$ and $x \in M$, and if $ax/s = 0$, then for some $h \notin P$, we have $ahx = 0$. Since A is a domain and $a/s \neq 0$ implies $ah \neq 0$, and because M is torsion-free, it follows that $x = 0$. Now, if J is a finitely generated ideal of A_P , since A_P is valuative, J is a principal ideal. Therefore, since M_P is torsion-free, the map $J \otimes M_P \to M_P$ is injective. This shows that M_P is flat.

(ii) \Longrightarrow (i): It suffices to show for any $P \in \text{Spec } A, A_P$ is a valuative ring. Let J be a finitely generated Ap ideal. Let I be the pullback of J along the natural map $A \to A_P$. Since A is a domain, I is torsion-free, i.e., flat. Hence, $J = I_P$ is also flat, and since A_P is local, J is free. Therefore, J is principal, and A_P is valuative.

The key point of this proof is that in a domain, the localization of a torsion-free module remains torsion-free. This perspective is important when extending this theorem. See [Proposition 4.5.](#page-4-0)

Let us organize the relationships between Prüfer domains and other classes. The following two facts are well known.

Noetherian Prüfer domain \iff Dedekind domain.

Prüfer domain \iff semi-hereditary domain.

4. General Cases

We will now discuss the case where we do not assume the integral domain condition, considering both the case where Noetherian condition is imposed and the more general case where it is not.

Proposition 4.1. *Let* A *be a ring. If every torsion-free* A*-module is flat, then* A *is reduced.*

Proof. Regarding $nil(A) = \{a \in A \mid a^n = 0 \text{ for some } n > 0\}, A/nil(A) \text{ is a }$ torsion-free A-module. By assumption, $A/\text{nil}(A)$ is flat, so for any $a \in$ nil(A), the natural map $aA \otimes_A (A/\text{nil}(A)) \to A/\text{nil}(A)$ is injective. However, this is a zero map, so $aA \otimes A/\text{nil}(A) = 0$. Therefore, we have $aA = \text{nil}(A)$. aA, and by NAK [\[Mat86,](#page-9-10) Theorem 2.2], it follows that $aA = 0$. Hence, $\mathrm{nil}(A)=0.$

This proposition shows that in the case where Krull dimension is 0, the class of rings for which torsion-free modules are flat is none other than the class of von Neumann regular rings. Let us recall the definition.

Definition 4.2. A ring A is called **von Neumann regular** (or absolutely flat) if for any element $a \in A$, there is a $b \in A$ satisfying $a^2b = a$.

Corollary 4.3. Let A be a ring with $\dim A = 0$. The following conditions *are equivalent:*

- (i) A *is von Neumann regular;*
- (ii) *Every torsion-free* A*-module is flat.*

Proof. (i) \implies (ii)

If A is von Neumann regular, then by [\[Gla89,](#page-9-7) Theorem 1.4.6], any Amodule is flat.

 $(ii) \implies (i)$

By [Proposition 4.1,](#page-3-1) A is reduced. A ring A being reduced and having $\dim A = 0$ is equivalent to being von Neumann regular [\[Lam99,](#page-9-8) Theorem 3.71], so A is von Neumann regular.

We introduce a restatement of the property of being semi-hereditary, as described by Glaz.

Theorem 4.4 ([\[Gla89,](#page-9-7) Corollary 4.2.19]). *Let* A *be a ring. The following conditions are equivalent:*

- (i) A *is semi-hereditary.*
- (ii) *The total quotient ring of* A*,* Q(A) *is von Neumann regular and for any maximal ideal* m *of* A*,* A^m *is a valuation ring.*

We note that if A is a reduced ring, then $Q(A)$ is also reduced. Therefore, if dim $Q(A) = 0$, $Q(A)$ is von Neumann regular. Furthermore, when A is a Noetherian reduced ring, we have dim $Q(A) = 0$ (see [\[Mat86,](#page-9-10) Exercise 6.5]). But note that this is not always the case in general, see [Example 4.8](#page-4-1) and [\[Que71\]](#page-9-11).

The condition that the dimension of total quotient ring is 0 plays an important role in ensuring that the localization of a torsion-free module remains torsion-free.

Proposition 4.5. Let A be a ring with $\dim Q(A) = 0$. For any torsion-free A-module M and $P \in \text{Spec } A$, M_P is a torsion-free A_P -module.

Proof. Let $\alpha = a/s$ be a non-zero divisor in Ap , and $\bar{x} = x/t \neq 0 \in Mp$ with $\alpha \bar{x} = 0$. Then, there exists some $h_1 \notin P$ such that $h_1ax = 0$.

We define $I \coloneqq \ker(A \to A_P) = \{a \in A \mid ha = 0 \text{ for some } h \notin P\}.$

Assume that $(a, I)Q(A) \neq Q(A)$. Then, there exists a maximal ideal m of $Q(A)$ containing (a, I) . Since $\dim Q(A) = 0$, m is minimal, so if we let $P' \coloneqq \mathfrak{m} \cap A \in \operatorname{Spec} A$, then P' is a minimal prime ideal containing (a, I) . Here, $P' A_{P'}$ is the unique prime ideal of $A_{P'}$, so all elements are nilpotent. Consequently, there exists some $n > 0$ and $b \notin P'$ such that $ba^n = 0$. Now, since a/s is a non-zero divisor in A_P , it follows that $ba^{n-1}/1 = 0 \in A_P$. Repeating this, we get $b/1 = 0 \in A_P$. In other words, $b \in I \subset P'$, which is a contradiction.

Therefore, $(a, I)Q(A) = Q(A)$. This implies that there exists some $b \in$ (a, I) which is a non-zero divisor of A. In this case, let $b = ra + c$ for some $r \in A$ and $c \in I$. Since there exists some $h_2 \notin P$ such that $h_2c = 0$, for $h \coloneqq h_1 h_2 \notin P$, we have $h b x = 0$. Now, since b is a non-zero divisor of A and M is a torsion-free A-module, it follows that $hx = 0$. Thus, $\bar{x} = 0 \in M_P$, and M_P is a torsion-free A_P -module.

In [Proposition 4.5,](#page-4-0) it remains a question whether the condition on the dimension of the total quotient ring can be weakened.

Open Problem 4.6. Characterize a ring A with the following property: If M is any torsion free A-module and $P \in \text{Spec } A$, then M_P is torsion-free A_P -module.

Several examples are presented where the localization of a torsion-free module is not torsion-free. The following example is given in a discussion we had with Ohashi and Matsumoto.

Example 4.7 (Noetherian and not reduced Example). Let k be a field. Let $\mathfrak{m} = (x, y, z) \subset k[x, y, z]$, and define $A = k[x, y, z]_{\mathfrak{m}}/(xy, yz, z^2)$. For $P = (x, y) \subset A$ and $M = A/xA$, the module M is a torsion-free A-module, but M_P is not a torsion-free A_P -module. Now, a regular element of A is a unit, so $Q(A) = A$ and dim $Q(A) = 2$.

[Example 4.7](#page-4-2) is an example in a Noetherian ring. As mentioned earlier, if a ring A is Noetherian, then if it is reduced, then $\dim Q(A) = 0$. Thus, to serve as a counterexample in a Noetherian ring, it must not be reduced. An example in a reduced ring is the following.

Example 4.8 (non Noetherian and reduced Example). Let k be a field. Define the ideals I of the infinite-variable polynomial ring $k[X_n | n = 1, 2, 3, \dots]$ as:

 $I = (X_i X_j | i \neq j \text{ and } \{i, j\} \neq \{1, 2\}).$

Minimal ideals of A are as follows:

 $P_{1,2} \coloneqq (X_n \mid n \neq 1, 2), P_i \coloneqq (X_n \mid n \neq i)$ for each $i \geq 3$.

A regular element of A is a polynomial with non-zero constant term. For $P := P_{1,2} + (X_2) = (X_n | n \neq 1)$ and $M = A/X_2A$, the module M is a torsion-free A-module, but M_P is not a torsion-free A_P -module. This ring A is reduced unlike [Example 4.7,](#page-4-2) and dim $Q(A) = 2$.

Note that the converse of [Proposition 4.5](#page-4-0) does not hold.

Example 4.9. Let k be a field. Let $\mathfrak{m} = (x, y) \subset k[x, y]$, and define $A =$ $k[x, y]_{m}/(x^{2}, xy)$. The prime ideals of A are $P = (x)$ and $\mathfrak{m} = (x, y)$ only. Since $A_m = A$ and $A_P = k(y)$, the regular elements in these rings are units. Therefore, the localization of any torsion-free module is also torsion-free.

On the other hand, as $A = Q(A)$ and $\dim Q(A) = 1$, this provides a counterexample to the converse of [Proposition 4.5.](#page-4-0)

We use [Proposition 4.5](#page-4-0) to characterize rings in which torsion-free modules become flat.

Theorem 4.10. *Let* A *be a ring . The following conditions are equivalent:*

- (i) A *is semi-hereditary ring.*
- (ii) *Every torsion-free* A*-module is flat.*
- *Proof.* (\implies)

Let M be a torsion-free A-module. Since a semi-hereditary ring is reduced and dim $Q(A) = 0$, by [Proposition 4.5,](#page-4-0) for any maximal ideal m, M_m is a torsion-free $A_{\mathfrak{m}}$ -module. Since $A_{\mathfrak{m}}$ is a valuation ring, by an argument similar to the proof of [Theorem 3.3,](#page-2-0) M_m is flat. Therefore, M is also flat. $(\Leftarrow$

If M is a torsion-less module, then since M is torsion-free, it follows that M is flat. Therefore, by [Theorem 2.2,](#page-1-2) A is semi-hereditary. \square

5. decomposition theorem for semi-hereditary rings

From the above, we have been able to reformulate a semi-hereditary ring in terms of torsion-free modules.

Using an idea similar to [Proposition 4.1,](#page-3-1) we can prove that a ring decomposes if there exists a special prime ideal. Since flatness is a local property, note that the following lemma holds.

Lemma 5.1. *Let* A *be a semi-hereditary ring, i.e. a ring with every torsionfree* A-module is flat. If there exist rings A_1 and A_2 such that $A \cong A_1 \times A_2$, *then they are semi-hereditary.*

Proposition 5.2. *Let* A *be a semi-hereditary ring. Suppose that there exists a prime ideal* $P \in \text{Spec } A$ *such that every element of* P *is a zero divisor. Then,* $P^2 = P$ *. Furthermore, if* P *is finitely generated, the following holds.*

- (i) *There exists an idempotent element* $e \in P$ *such that* $P = (e)$ *.*
- (ii) A *decomposes* as $A \cong A/eA \times A/(1-e)A$.
- (iii) A/eA *is a Prüfer domain, and* $A/(1-e)A$ *is semi-hereditary.*

Proof. Since all elements of P are zero divisors, A/P is a torsion-free Amodule. Thus, A/P is flat, just like [Proposition 4.1,](#page-3-1) the natural map $P \otimes_A$ $A/P \rightarrow A/P$ is injective and zero. Therefore, $P \otimes A/P = P/P^2 = 0$.

Now, suppose P is finitely generated.

- (i) By NAK [\[Mat86,](#page-9-10) Theorem 2.2], there exists some $a \in P$ such that $(a+1)P = 0$. Thus, $P = aA$. Since $P = P²$, there exists some $b \in A$ such that $a = a^2b$. Letting $e = ab$, we see that e is idempotent and $P = aA = eA$.
- (ii) Let $A \rightarrow A/eA \times A/(1-e)A; a \mapsto (a + eA, a + (1-e)A)$ and $A/eA \times A/(1-e)A \rightarrow A$; $(b+eA, c+(1-e)A) \rightarrow b(1-e) + ce$ be homomorphisms that are inverses of each other.
- (iii) From the lemma, A/eA and $A/(1-e)A$ are semi-hereditary. Here, since $A/eA = A/P$ is a domain, A/eA is a Prüfer domain by [Theorem 3.3.](#page-2-0) \Box

For a ring A , it is important to note that there always exists a minimal prime ideal, and all elements of the minimal prime ideal are zero divisors. Moreover, according to [\[Dut78,](#page-9-12) Proposition 4,5], if the minimal prime ideal P is finitely generated, then P becomes an associated prime, meaning that there exists some $a \neq 0 \in A$ such that $P = \text{Ann}(a)$.

Corollary 5.3. *Let* A *be a semi-hereditary ring. If every minimal prime ideal of* A *is finitely generated, then* A *is a finite product of Prüfer domains.*

Proof. Under this assumption, by [\[And94\]](#page-8-1), the minimal prime ideals are finite in number. Since each decomposition of the ring A as $A \cong A/P \otimes$ $A/(1-e)A$ using the minimal prime P decreases the number of minimal primes, this decomposition can occur at most finitely many times. \Box

From the above we can conclude that in the case of Noetherian rings, semi-hereditary rings are expressed as finite products of Dedekind domains. Before giving the proof, we will briefly summarise what is known about the relations between different classes.

Lemma 5.4. *Let* A *be a ring. The following conditions are equivalent:*

- (i) A *is a Noetherian Prüfer domain*:
- (ii) A *is a hereditary domain;*
- (iii) A *is a Dedekind domain.*

Proof. (i) \implies (ii)

Any ideal of A is finitely presented and flat, so it is projective. Therefore, A is hereditary.

 $(ii) \implies (iii)$

Any ideal of A is projective. Since A is a domain, (fractional) ideal of A is projective if and only if invertible (see [\[Mat86,](#page-9-10) Theorem 11.3]). So A is a Dedekind domain.

 $(iii) \implies (i)$

Since a Dedekind domain is Noetherian and every ideal of A is projective, so A is a Prüfer domain.

Corollary 5.5. *Let* A *be a Noetherian ring. The following conditions are equivalent:*

- (i) A *is a semi-hereditary ring.*
- (ii) A *is a finite direct product of Dedekind domains.*
- (iii) A *is a hereditary ring.*

Proof. (i) \implies (ii)

By [Corollary 5.3,](#page-6-0) \tilde{A} is a finite product of Prüfer domains. Now, from the construction, each Prüfer domain component of A is Noetherian. In a Noetherian Prüfer domain, every ideal is finitely presented and flat, hence projective. Therefore, A is a finite product of hereditary domains. Since hereditary domains are equivalent to Dedekind domains, A is a finite product of Dedekind domains.

 $(ii) \implies (iii)$

A direct product of hereditary rings is also hereditary. $(iii) \implies (i)$

It is obvious.

6. Some topics in perfectoid ring theory

It goes without saying that the absolute integral closure plays an extremely important role in perfectoid ring theory.

Definition 6.1. Let A be an integral domain with the field of fractions $Q(A)$. Let the algebraic closure of $Q(A)$ be denoted by $\overline{Q(A)}$. The integral closure of A in $\overline{Q(A)}$ is denoted by A^+ and is called the absolute integral closure of A.

It can be seen that the absolute integral closure of the valuation ring is an example of a Prüfer domain.

Proposition 6.2 ([\[Bou89,](#page-8-2) Chap.VI, §8.6, Prop.6]). *Let* A *be a valuation ring with a valuation* v *on a field* K*,* L *an algebraic extension of* K *and* A′ *the integral closure of* A *in* L*. There is a one-to-one correspondence between the valuation rings of valuations on* L *that extend* v *and the localizations* A′ m at each maximal ideal m of A'. In particular, A' is Prüfer.

Also, by virtue of a valuation ring A itself being a Prüfer domain, it can be shown that $A \to A^+$ is faithfully flat. Although this is well known, let us provide a proof.

Proposition 6.3. Let A be a valuation ring and let A^+ be the absolute *integral closure of A. The natural map* $A \rightarrow A^+$ *is faithflly flat.*

Proof. The map $A \to A^+$ is torsion-free. This is because, for any $a \neq 0$ in A and $\alpha \in A^+$, there exist elements $c_1, \ldots, c_r \in A$ such that

$$
\alpha^r + c_1 \alpha^{r-1} + \dots + c_r = 0
$$

Here, let r be the smallest degree of such a monic polynomial. By multiplying this equation by a, it follows that $c_r = 0$, which gives $\alpha(\alpha^{r-1} + c_1\alpha^{r-2} +$ $\cdots + c_{r-1}$ = 0. By the minimality of r, we conclude that $\alpha = 0$.

Since A is also a Prüfer domain, A^+ is flat over A.

Now, let v be a valuation of A on the field K. Then, for any maximal ideal \mathfrak{m}' of A^+ , by [Proposition 6.2,](#page-7-1) there exists an extension v' of v to an algebraic extension K' of K, along with a valuation ring A' such that $A' = A_{\mathfrak{m}'}^+$. Then, for any maximal ideal m of A, it must hold that for any $a \in \mathfrak{m}$ and $\alpha \in A^+$, we have $v'(a\alpha) = v(a) + v'(\alpha) > 0$. Consequently, $A^+ \neq \mathfrak{m}A^+$, and hence $A \rightarrow A^+$ is faithfully flat.

By Kunz's theorem [\[Kun76\]](#page-9-13), the importance of the flatness of the Frobenius map is widely recognized, and [\[Lur20\]](#page-9-5) provides an equivalent condition for the flatness of the Frobenius map $A/\varpi A \rightarrow A/\varpi^p A$ when A is Noetherian and ϖ is a regular element for which ϖ^p divides p. Inspired by this discussion, [\[Shi\]](#page-9-6) examines the flatness of the Frobenius map in the case of valuation rings, leveraging [Proposition 6.3.](#page-7-2)

Proposition 6.4 ([\[Shi,](#page-9-6) Proposition 7.12.]). Let A be a ring and let $p > 0$ *be a prime number.* Assume that $\varpi \in A$ *is an element such that* A/bA *is a ring of characteristic* $p > 0$ *, and assume that there is a unit* $u \in A^{\times}$ *suth that* $\varpi^p = bu$. Assume there is a faithfully flat A-algebra B such that the *Frobenius map* $B/bB \to B/bB$ *is surjective with kernel being equal to* (ϖ) *. Then the Frobenius map* $A/\varpi A \rightarrow A/pA$ *is flat. In paticular, if* A *is a valuation ring, then the Frobenius map is flat.*

In this context, the proof that the absolute integral closure of a valuation ring is a faithfully flat A-algebra [\(Proposition 6.3\)](#page-7-2) relies on the fact that torsion-free modules over a valuation ring are flat. This leads to the proposal of the following problem.

Open Problem 6.5 ([\[Shi,](#page-9-6) Open Problem 7.13.]). Characterize a ring A with the following property: If M is any torsion-free A -module, then M is flat.

This problem is solved by [Theorem 4.10](#page-5-0) in this paper. This property is equivalent to A being semi-hereditary.

The following problems are also related.

Open Problem 6.6 ([\[IS22,](#page-9-14) Question 1.]). Let A be a ring of mixed characteristic $p > 0$. Assume that there is an element $\varpi \in A$ together with an element $u \in A^{\times}$ such that $\varpi^{p} = pu$. Characterise a ring A such that the Frobenius map $A/\varpi A \rightarrow A/pA$ is faithfully flat, other than the case A is a regular ring, a valuation ring, or a ring whose p-adic completion is an integral perfectoid ring.

Open Problem 6.7 ([\[Shi,](#page-9-6) Open Problem 7.11.]). Let A be a ring and let $p > 0$ be a prime number. Assume that $\varpi \in A$ is an element such that A/bA is a ring of characteristic $p > 0$, and assume that there is a unit $u \in A^{\times}$ suth that $\varpi^p = bu$. Characterise a ring A such that the Frobenius map $A/\varpi A \rightarrow A/bA$ is flat.

Acknowledgments. I would like to thank Hisanori Ohashi and Yuya Matsumoto for attending the seminar and giving me various suggestions. I would also like to express my deepest gratitude to my company, SKILLUP NeXt Ltd, and its members.

REFERENCES

- [AMM22] B. Antieau, A. Mathew, and M. Morrow, The K-theory of perfectoid rings, Doc. Math. 27 (2022), 1923–1951.
- [And94] D. D. Anderson, A note on minimal prime ideals, Proc. Amer. Math. Soc. 122 (1994), 13–14, DOI 10.1090/S0002-9939-1994-1191864-2.
- [Bou89] N. Bourbaki, Commutative Algebra: Chapters 1-7, Springer Berlin, Heidelberg, 1989.
- [Cha60] S. U. Chase, *Direct products of modules*, Trans. Amer. Math. Soc. 97 (1960), no. 3, 457–473.
- [Dro80] Yu. A. Drozd, The structure of hereditary rings, Mat. Sb. 113(155) (1980). English translation: Math. Sbornik (USSR), 41, No. 1, 139–148 (1982).
- [Dut78] P. Dutton, Prime ideals attached to a module, The Quarterly Journal of Mathematics 29 (1978), no. 4, 403–413.
- [Fai75] C. Faith, On a theorem of Chatters, Communications in Algebra 3 (1975), no. 2, 169–184.
- [Gla89] S. Glaz, Commutative Coherent Rings, Lecture Notes in Mathematics, vol. 1371, Springer–Verlag, 1989.
- [IS22] S. Ishiro and K. Shimomoto, Another proof of the almost purity theorem for perfectoid valuation rings, Journal of Pure and Applied Algebra 226 (2022), no. 7, 106898.
- [Kap58] I. Kaplansky, Projective Modules, Ann. Math. 68 (1958), no. 2, 372–377.
- [Kun76] E. Kunz, On Noetherian Rings of Characteristic p, Amer. J. Math. 98 (1976), no. 4, 999–1013.
- [Lam99] T. Y. Lam, Lectures on Modules and Rings, Graduate Texts in Mathematics, vol. 189, Springer–Verlag, 1999.
- [Lur20] J. Lurie, Level Structures on Elliptic Curves, 2020. available at <https://www.math.ias.edu/~lurie/>.
- [MO01] K. Masaike and Y. Ozaki, Direct Product Decomposition of Semi-Hereditary Rings, Bulletin of Tokyo Gakugei University. Series IV, Mathematics and Natural Sciences 53 (2001), 19–23.
- [Mat86] H. Matsumura, Commutative Ring Theory, Cambridge Studies in Advanced Mathematics, vol. 8, Camb. Univ. Press, 1986.
- [Que71] Y. Quentel, Sur la compacité du spectre minimal d'un anneau, Bulletin de la Société Mathématique de France 99 (1971), 265–272.
- [Shi11] K. Shimomoto, *F-coherent rings with applications to tight closure theory*, Journal of Algebra 338 (2011), no. 1, 24-34.
	- [Shi] K. Shimomoto, Lectures on perfectoid geometry for commutative algebraists. in preparation.