

# Long-Time Behavior of Typical Pure States from Thermal Equilibrium Ensembles

Cornelia Vogel\*

December 24, 2024

## Abstract

We consider an isolated macroscopic quantum system in a pure state  $\psi_t \in \mathcal{H}$  evolving unitarily in a separable Hilbert space  $\mathcal{H}$ . Following von Neumann [46], we assume that different macro states  $\nu$  correspond to mutually orthogonal subspaces  $\mathcal{H}_\nu \subset \mathcal{H}$ . Let  $P_\nu$  be the projection to  $\mathcal{H}_\nu$ . “Normal typicality” is the statement (true for some Hamiltonians) that for all initial states  $\psi_0 \in \mathcal{H}$  and most  $t \geq 0$ ,  $\|P_\nu \psi_t\|^2$  is close to  $d_\nu/D$ , where  $d_\nu = \dim \mathcal{H}_\nu$  and  $D = \dim \mathcal{H} < \infty$ . It can be shown [42] that the statement becomes valid for all Hamiltonians if “all  $\psi_0 \in \mathcal{H}$ ” is replaced by “most  $\psi_0 \in \mathcal{H}_\mu$ ” (most w.r.t. the uniform distribution on the sphere of  $\mathcal{H}_\mu$ ) with an arbitrary macro state  $\mu$  and  $d_\nu/D$  by a  $t$ - and  $\psi_0$ -independent quantity  $M_{\mu\nu}$ . In the present work, we generalize this result from the uniform distribution to a much more general class of measures, so-called GAP measures. Let  $\rho$  be a density matrix on  $\mathcal{H}$ . Then  $\text{GAP}(\rho)$  is the most spread out distribution on the sphere of  $\mathcal{H}$  with density matrix  $\rho$ . If  $\rho$  is a canonical density matrix,  $\text{GAP}(\rho)$  arises as the thermal equilibrium distribution of wave functions and can be viewed as a quantum analog of the canonical ensemble. We show that also for  $\text{GAP}(\rho)$ -most  $\psi_0 \in \mathcal{H}$ , the superposition weight  $\|P_\nu \psi_t\|^2$  is close to a fixed value  $M_{\rho P_\nu}$  for most  $t \geq 0$ . Moreover, we prove a similar result for arbitrary bounded operators  $B$  instead of  $P_\nu$  and for finite times. The main ingredient of the proof is an improvement of bounds on the variance of  $\langle \psi | B | \psi \rangle$  w.r.t.  $\text{GAP}(\rho)$  which were first obtained by Reimann [29] and slightly generalized in [44].

**Key words:** von Neumann’s quantum ergodic theorem; equilibration; thermalization; Gaussian adjusted projected (GAP) measure; Scrooge measure; random wave function; quantum statistical mechanics; macroscopic quantum systems; long-time behavior

---

\*Mathematics Institute, Eberhard Karls University Tübingen, Auf der Morgenstelle 10, 72076 Tübingen, Germany. ORCID: 0000-0002-3905-4730, E-mail: cornelia.vogel@uni-tuebingen.de

# 1 Introduction

We consider an closed macroscopic quantum system in a pure state and investigate its long-time behavior. This approach in order to study equilibration and thermalization has been very fruitful over the last three decades; important discoveries include *dynamical typicality* [2, 26, 1, 32, 31, 33, 42], *canonical typicality* [24, 8, 9, 27, 17, 44], the *eigenstate thermalization hypothesis* (ETH) from a physicist’s as well as a mathematician’s point of view [6, 38, 5, 4, 7], and, only very recently, the rigorous study of thermalization of a system of free fermions [35, 41, 40, 34].

Another important result was the rediscovery and further elaboration of *normal typicality* [16, 18, 30] which goes back to von Neumann [46]. The goal of the present work is to generalize normal typicality from the uniform distribution on the sphere to so-called *GAP measures*. To any density matrix  $\rho$  on  $\mathcal{H}$  we can associate a measure  $\text{GAP}(\rho)$  which is, roughly speaking, the most spread-out distribution over  $\mathbb{S}(\mathcal{H})$  with density matrix  $\rho$ . For certain density matrices these measures arise as thermal distributions of quantum states and they can be regarded as a quantum analog of the canonical ensemble from classical statistical mechanics [21, 20].

We start with recalling the notion of normal typicality and first introduce the setting: Following von Neumann [46], we take for granted that the system’s Hilbert space  $\mathcal{H}$  can be decomposed into mutually orthogonal subspaces  $\mathcal{H}_\nu$  (“macro spaces”) associated with different macro states  $\nu$  such that

$$\mathcal{H} = \bigoplus_{\nu} \mathcal{H}_{\nu}. \quad (1)$$

Von Neumann thought of the macro spaces  $\mathcal{H}_\nu$  as being the joint eigenspaces of a set of mutually commuting self-adjoint and highly degenerate operators  $M_1, \dots, M_K$  (“macroscopic observables”). Each macro space is then characterized by a list of eigenvalues,  $\nu = (m_1, \dots, m_K)$  of the operators  $M_1, \dots, M_K$  and the elements from a macro space  $\mathcal{H}_\nu$  look “macroscopically the same” in the sense that every  $\psi \in \mathcal{H}_\nu$  is an eigenfunction of all  $M_j$  with corresponding eigenvalue  $m_j$ . Usually, a coarse-grained version of the system’s Hamiltonian  $H$  is among the  $M_j$  and therefore each  $\mathcal{H}_\nu$  is a subset of an *energy shell*  $\mathcal{H}_{\text{mc}} = \mathbb{1}_{[E-\Delta E, E]}(H)\mathcal{H}$ . Here,  $[E - \Delta E, E]$  is a micro-canonical energy interval with  $\Delta E$  (the resolution of macroscopic energy measurements) being small on the macroscopic scale but sufficiently large on the microscopic scale such that  $[E - \Delta E, E]$  contains a very large (but typically finite) number of eigenvalues of  $H$ . This implies in particular that even if we allow  $\mathcal{H}$  to be infinite-dimensional, the  $\mathcal{H}_\nu$  should be of finite dimension. Moreover, in each energy shell there usually is one macro space that contains most dimensions of the shell and which we associate with thermal equilibrium and denote by  $\mathcal{H}_{\text{eq}}$ .

Let  $P_\nu$  be the orthogonal projection to  $\mathcal{H}_\nu$  and let  $\mathbb{S}(\mathcal{H}) = \{\psi \in \mathcal{H} : \|\psi\| = 1\}$  denote the sphere in  $\mathcal{H}$ . If  $D := \dim \mathcal{H} < \infty$ , for most  $\phi \in \mathbb{S}(\mathcal{H})$  (where “most” refers

to the uniform distribution on  $\mathbb{S}(\mathcal{H})$  we have that

$$\|P_\nu\phi\|^2 \approx \frac{d_\nu}{D} \quad \forall \nu, \quad (2)$$

where  $d_\nu := \dim \mathcal{H}_\nu$  provided that  $d_\nu$  and  $D$  are sufficiently large [16]. If the eigenbasis of  $H$  is chosen purely randomly (i.e., according to the Haar measure) among all orthonormal bases (and under some further technical not very restrictive assumptions), every initial wave function  $\psi_0 \in \mathbb{S}(\mathcal{H})$  evolves such that for most times  $t \geq 0$ ,

$$\|P_\nu\psi_t\|^2 \approx \frac{d_\nu}{D}, \quad (3)$$

where  $\psi_t = \exp(-iHt)\psi_0$  (we set  $\hbar = 1$ ), provided that  $d_\nu$  and  $D$  are sufficiently large [16, 18, 30, 46]. This phenomenon is known as “normal typicality”. In particular, if the macro state  $\nu$  represents thermal equilibrium, we write  $d_\nu = d_{\text{eq}}$  and have that for every  $\psi_0 \in \mathbb{S}(\mathcal{H})$ ,

$$\|P_{\text{eq}}\psi_t\|^2 \approx \frac{d_{\text{eq}}}{D} \approx 1 \quad (4)$$

for most  $t \geq 0$ , i.e., every state spends most of the time in thermal equilibrium. Therefore we have thermalization in the sense that also non-equilibrium initial states sooner or later reach the thermal equilibrium macro space and stay there for most of the time. This sense of thermal equilibrium is often called “macroscopic thermal equilibrium” (MATE), see, e.g., [13, 14, 15, 39, 25, 19, 23]. Note that here it is important that the statement holds true for *every* initial wave function; if it was only true for “most”  $\psi_0$ , the statement would not tell us much about thermalization because “most” states in  $\mathbb{S}(\mathcal{H})$  are in thermal equilibrium anyway.

Von Neumann’s assumptions on the Hamiltonian are not physically realistic because the energy eigenbasis of  $H$  is unrelated to the decomposition of  $\mathcal{H}$  into the macro spaces and therefore the system goes very rapidly from any (possibly very non-equilibrium) macro space almost immediately to the thermal equilibrium macro space  $\mathcal{H}_{\text{eq}}$  [10, 11, 12]. However, if the system starts in a very non-equilibrium state, it should pass through larger and larger (and therefore less far away from thermal equilibrium) macro spaces until it finally reaches  $\mathcal{H}_{\text{eq}}$ . In [42], the notion of generalized normal typicality was therefore generalized in the following way: for a general Hamiltonian  $H$ , for most  $\psi_0 \in \mathbb{S}(\mathcal{H}_\mu)$  (where “most” again refers to the uniform distribution over  $\mathbb{S}(\mathcal{H}_\mu)$ ) with a possibly non-equilibrium macro state  $\mu$  and most times  $t \geq 0$ ,

$$\|P_\nu\psi_t\|^2 \approx M_{\mu\nu} \quad \forall \nu \quad (5)$$

for suitable values  $M_{\mu\nu}$  provided that the eigenvalues and eigenvalue gaps of  $H$  are not too highly degenerate. We remark that the  $M_{\mu\nu}$  depend also on  $\mu$  but we still expect that in relevant cases,  $M_{\mu\nu} \approx d_\nu/D$ . While (2) holds in the sense that the absolute as well as the relative errors are small, (5) was in [43] only proved in the sense that the

absolute error is small. In order to show that the relative errors are small as well, we need a lower bound on the quantities  $M_{\mu\nu}$ . In [43] such a lower bound was obtained in the case that the Hamiltonian is of the form  $H = H_0 + V$  where  $H_0$  is a deterministic Hermitian matrix and  $V$  is a Hermitian random Gaussian perturbation by making use of results from random matrix theory. We remark that (5) can also be shown for arbitrary bounded operators  $B$ , i.e., with  $\|P_\nu\psi_t\|^2$  replaced by  $\langle\psi_t|B|\psi_t\rangle$  and  $M_{\mu\nu}$  replaced by a suitable quantity  $M_{\mu B}$ . Moreover, a finite time result is available, however, the times required for a small error are extremely large, see, e.g., [42] for a detailed discussion.

In the present work we generalize (5) in the sense that we replace the notion of “most” which there refers to the uniform distribution over the sphere  $\mathbb{S}(\mathcal{H}_\mu)$  of some macro space  $\mathcal{H}_\mu$  with some other measure on  $\mathbb{S}(\mathcal{H})$ , namely with *GAP measures*. The acronym GAP stands for *Gaussian adjusted projected* measure [22, 21] and it refers to one possible way of how these measures can be constructed.<sup>1</sup> As mentioned above, to any density matrix  $\rho$  on  $\mathcal{H}$  we can associate a measure  $\text{GAP}(\rho)$  on  $\mathbb{S}(\mathcal{H})$ , see Section 2 for a mathematically precise definition and construction of  $\text{GAP}(\rho)$ . We remark already here that  $\text{GAP}(\rho)$  can, in contrast to the uniform distribution, also be defined on separable Hilbert spaces, i.e., the Hilbert space does not have to be of finite dimension. If  $\rho = \rho_{\text{can}}$ , i.e., if  $\rho$  is of the form

$$\rho = \rho_{\text{can}} = \frac{1}{Z} e^{-\beta H}, \quad (6)$$

where  $\beta$  is the inverse temperature and  $Z$  a normalization constant,  $\text{GAP}(\rho)$  arises as the thermal equilibrium distribution of the wave function, see [21, 20] for details.

Our main result in the present paper is that also for most  $\psi_0 \in \mathbb{S}(\mathcal{H})$ , where “most” now refers to  $\text{GAP}(\rho)$ , for most  $t \geq 0$ ,

$$\langle\psi_t|B|\psi_t\rangle \approx M_{\rho B} \quad (7)$$

for suitable quantities  $M_{\rho B}$  provided that  $\|\rho\|$  is small,  $\|B\|$  is not too large and the eigenvalues and eigenvalue gaps of  $H$  are not too highly degenerate. Here,  $B$  is any operator if  $\mathcal{H}$  is finite-dimensional. If  $\dim \mathcal{H} = \infty$ , we have to restrict the class of operators  $B$ , roughly speaking, to operators which act non-trivially only on finitely many eigenspaces of  $H$  and whose image is contained in the span of only finitely many eigenspaces. As discussed above, the projections  $P_\nu$  to the macro spaces should fulfill this assumption as the macro spaces are usually contained in an energy shell which typically is finite-dimensional. Note that here we restrict our considerations to the absolute errors; however, if  $B = P_\nu$  or  $\rho = P_\nu/d_\nu$  for some macro state  $\nu$ , the bounds from [43]

---

<sup>1</sup>GAP measures we first introduced by Jozsa, Robb and Wootters [22] in 1994 in an information theoretical context. More precisely, they showed that  $\text{GAP}(\rho)$  minimizes the so-called “accessible information” under the constraint that its density matrix is given by  $\rho$ . They named it *Scrooge measure*, referring to Ebenezer Scrooge, the protagonist of Charles Dickens’ novella *A Christmas Carol* (1843) who is a very stingy character. The authors chose this name as the GAP measure is “particularly stingy with its information”.

apply. Our result shows that for  $\text{GAP}(\rho)$ -most  $\psi_0 \in \mathbb{S}(\mathcal{H})$  the curve  $t \mapsto \langle \psi_t | B | \psi_t \rangle$  is nearly constant in the long run  $t \rightarrow \infty$ . We remark that for  $\text{GAP}(\rho)$ -most  $\psi_0 \in \mathbb{S}(\mathcal{H})$ , the curve  $t \mapsto \langle \psi_t | B | \psi_t \rangle$  is nearly deterministic (“dynamical typicality”); this has been shown in [44]. The generalization of (generalized) normal typicality to  $\text{GAP}(\rho)$  reveals that it is not a phenomenon specific for the uniform distribution but it is also valid for a broader class of natural and physically relevant distributions which can also be defined in infinite-dimensional Hilbert spaces. As some GAP measures are a quantum analog of the canonical ensemble whereas the uniform distribution on an energy shell  $\mathcal{H}_{\text{mc}}$  is an analog micro-canonical ensemble, our results can be regarded as expressing a version of equivalence of ensembles.

The remainder of this paper is organized as follows: In Section 2, we give some background and present the mathematical setup. In Section 3, we formulate and discuss our main result. In Section 4, we provide the proofs. Finally, in Section 5, we conclude.

## 2 Background and Mathematical Setup

In the following, we make precise some notions that appeared in the introduction, give a mathematically rigorous definition of the GAP measures and introduce the system’s Hamiltonian and its relevant quantities that we will need to state our main result in Section 3. Throughout this paper we assume that  $\mathcal{H}$  is a separable Hilbert space, i.e., that it has either a finite or countably infinite orthonormal basis.

**Measures of “most”.** Let  $\mathbb{P}$  be a probability measure on  $\mathbb{S}(\mathcal{H})$  and let  $\varepsilon > 0$ . We say that, w.r.t.  $\mathbb{P}$ , a statement  $s(\psi)$  is true for  $(1 - \varepsilon)$ -most  $\psi \in \mathbb{S}(\mathcal{H})$  if

$$\mathbb{P} \{ \psi \in \mathbb{S}(\mathcal{H}) : s(\psi) \text{ holds} \} \geq 1 - \varepsilon. \quad (8)$$

Let  $T, \delta > 0$ . Analogously we say that a statement  $S(t)$  is true for  $(1 - \delta)$ -most  $t \in [0, T]$  if

$$\frac{1}{T} \lambda \{ t \in [0, T] : S(t) \text{ holds} \} \geq 1 - \delta, \quad (9)$$

where  $\lambda$  denotes the Lebesgue measure on  $\mathbb{R}$ . Moreover, we say that a statement  $S(t)$  is true for  $(1 - \delta)$ -most  $t \in [0, \infty)$  (or also  $t \geq 0$ ) if

$$\liminf_{T \rightarrow \infty} \frac{1}{T} \lambda \{ t \in [0, T] : S(t) \text{ holds} \} \geq 1 - \delta. \quad (10)$$

**Time averages.** Let  $T > 0$  and let  $f : \mathbb{R}_+ \rightarrow \mathbb{C}$ . We define the finite time average  $\langle \cdot \rangle_T$  of  $f$  over the interval  $[0, T]$  by

$$\langle f(t) \rangle_T := \frac{1}{T} \int_0^T f(t) dt. \quad (11)$$

Moreover, the infinite time average of  $f$  is given by

$$\overline{f(t)} := \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T f(t) dt \quad (12)$$

whenever this limit exists.

**Norms.** Throughout this paper we need two norms, the operator norm and the trace norm. Let  $M$  be an operator on  $\mathcal{H}$ . Its *operator norm* is given by

$$\|M\| := \sup_{\|\psi\|=1} \|M\psi\|. \quad (13)$$

If  $M$  is self-adjoint,  $\|M\|$  is equal to the largest absolute eigenvalue of  $M$ . In general,  $\|M\|$  is equal to the square root of the largest eigenvalue of  $M^*M$  where  $M^*$  is the adjoint operator of  $M$ .

The *trace norm* of  $M$  is defined as

$$\|M\|_{\text{tr}} := \text{tr} |M| = \text{tr} \sqrt{M^*M}. \quad (14)$$

If  $M$  is self-adjoint,  $\|M\|_{\text{tr}}$  is equal to the sum of the absolute eigenvalues of  $M$ .

**Density matrix.** Let  $\mathbb{P}$  be a probability measure on  $\mathbb{S}(\mathcal{H})$ . From the measure  $\mathbb{P}$  we can obtain a density matrix  $\rho_{\mathbb{P}}$  on  $\mathcal{H}$  by

$$\rho_{\mathbb{P}} := \int_{\mathbb{S}(\mathcal{H})} \mathbb{P}(d\psi) |\psi\rangle\langle\psi| \quad (15)$$

and we say that the measure  $\mathbb{P}$  has density matrix  $\rho_{\mathbb{P}}$ . Note that this density matrix always exists, see, e.g., Lemma 1 in [45]. If the mean of  $\mathbb{P}$  is equal to zero,  $\rho_{\mathbb{P}}$  is the covariance matrix of  $\mathbb{P}$ .

**GAP measure.** In this paragraph, we give a mathematically rigorous definition of the measure  $\text{GAP}(\rho)$  on  $\mathbb{S}(\mathcal{H})$  for a density matrix  $\rho$  on  $\mathcal{H}$ . There are several equivalent definitions of  $\text{GAP}(\rho)$  in the case that  $\mathcal{H}$  is finite-dimensional, see [20]. In the following, we present the one which gave the GAP measures their name as this construction can be done in a similar way if  $\mathcal{H}$  is infinite-dimensional. For simplicity, we restrict ourselves to the finite-dimensional case and refer to [45] for the case that  $\dim \mathcal{H} = \infty$ .

Let  $\rho$  be a density matrix on a finite-dimensional Hilbert space  $\mathcal{H}$ . Then we can write it as

$$\rho = \sum_n p_n |n\rangle\langle n|, \quad (16)$$

where the  $|n\rangle$  form an orthonormal eigenbasis of  $\rho$  with corresponding eigenvalues  $p_n$ .

As the name *Gaussian adjusted projected measure* suggests, we start from a Gaussian measure  $G(\rho)$  on  $\mathcal{H}$ ; it is constructed as follows: Let  $(Z_n)$  be a sequence of independent complex-valued centered Gaussian random variables<sup>2</sup> with variances

$$\mathbb{E}|Z_n|^2 = p_n. \quad (17)$$

The Gaussian measure  $G(\rho)$  is then defined as the distribution of the random vector

$$\Psi^G := \sum_n Z_n |n\rangle. \quad (18)$$

While  $G(\rho)$  is centered and has density matrix  $\rho$ , it is not a distribution on the sphere  $\mathbb{S}(\mathcal{H})$ : It follows immediately from the definition of  $\Psi^G$  together with the fact that the  $p_n$  sum up to 1 that  $\mathbb{E}\|\Psi^G\|^2 = 1$  but in general  $\|\Psi^G\| \neq 1$ .

In the next step, we have to adjust the measure  $G(\rho)$ ; this adjustment is necessary because if we would project  $G(\rho)$  directly to the sphere  $\mathbb{S}(\mathcal{H})$ , the resulting measure would not have the property that it has density matrix  $\rho$ . We define the *adjusted Gaussian measure*  $\text{GA}(\rho)$  on  $\mathcal{H}$  by

$$\text{GA}(\rho)(d\psi) := \|\psi\|^2 G(\rho)(d\psi), \quad (19)$$

i.e., we multiply the density of  $G(\rho)$  by the factor  $\|\psi\|^2$ . It follows from  $\mathbb{E}\|\Psi^G\|^2 = 1$  that also  $\text{GA}(\rho)$  is a probability measure on  $\mathcal{H}$ .

In the last step of the construction, we project the measure  $\text{GA}(\rho)$  on  $\mathcal{H}$  to the sphere  $\mathbb{S}(\mathcal{H})$ . To this end, let  $\Psi^{\text{GA}}$  be a  $\text{GA}(\rho)$ -distributed random vector. Then we define  $\text{GAP}(\rho)$  to be the distribution of the random vector

$$\Psi^{\text{GAP}} := \frac{\Psi^{\text{GA}}}{\|\Psi^{\text{GA}}\|}. \quad (20)$$

Obviously, this measure is a distribution on  $\mathbb{S}(\mathcal{H})$  and a short computation shows that  $\text{GAP}(\rho)$  indeed has density matrix  $\rho$ :

$$\rho_{\text{GAP}(\rho)} = \int_{\mathbb{S}(\mathcal{H})} \text{GAP}(\rho)(d\psi) |\psi\rangle\langle\psi| = \int_{\mathcal{H}} \text{GA}(\rho)(d\psi) \frac{1}{\|\psi\|^2} |\psi\rangle\langle\psi| = \int_{\mathcal{H}} G(\rho)(d\psi) |\psi\rangle\langle\psi| = \rho. \quad (21)$$

**The Hamiltonian.** In this paper we consider Hamiltonians  $H$  with spectral decomposition

$$H = \sum_{e \in \mathcal{E}} e \Pi_e, \quad (22)$$

---

<sup>2</sup>Recall that a centered complex-valued Gaussian random variable  $Z$  with variance  $\sigma^2 > 0$  has independent real and imaginary part  $\text{Re } Z$  and  $\text{Im } Z$ ,  $\mathbb{E}\text{Re } Z = \mathbb{E}\text{Im } Z = 0$  and  $\mathbb{E}(\text{Re } Z)^2 = \mathbb{E}(\text{Im } Z)^2 = \sigma^2/2$ .

where  $\mathcal{E}$  denotes the set of distinct eigenvalues of  $H$  and  $\Pi_e$  is the projection onto the eigenspace of  $H$  with corresponding eigenvalue  $e$ . We define  $d_E := |\mathcal{E}|$ , where  $|\{\cdot\}|$  denotes the number of elements of the set  $\{\cdot\}$ , and  $D_E := \max_{e \in \mathcal{E}} \text{tr}(\Pi_e)$ , i.e.,  $d_E$  is the number of distinct eigenvalues of  $H$  and  $D_E$  is the maximum degeneracy of an eigenvalue of  $H$ . Moreover, we define the maximal gap degeneracy  $D_G$  by

$$D_G := \max_{E \in \mathbb{R}} |\{(e, e') \in \mathcal{E} \times \mathcal{E} : e \neq e' \text{ and } e - e' = E\}|. \quad (23)$$

Let  $\kappa > 0$ . The maximal number of gaps in an energy interval of length  $\kappa$  is given by

$$G(\kappa) := \max_{E \in \mathbb{R}} |\{(e, e') \in \mathcal{E} \times \mathcal{E} : e \neq e' \text{ and } e - e' \in [E, E + \kappa)\}| \quad (24)$$

and obviously  $D_G = \lim_{\kappa \rightarrow 0^+} G(\kappa)$ .

If  $\mathcal{H}$  is infinite-dimensional, these quantities are not necessarily finite. In particular,  $d_E$  and  $D_E$  cannot both be finite. Let  $B$  be an operator on  $\mathcal{H}$ . We will see in the proofs that only those eigenvalues  $e, e'$  with  $\Pi_e B \Pi_{e'} \neq 0$  contribute to the quantities we want to compute and we will require that there are only finitely many such  $e, e'$  which will imply that the sums over  $e, e'$  are effectively finite. We therefore define for an operator  $B$  the set of ‘‘contributing’’ eigenvalues by

$$\mathcal{E}_B := \{e \in \mathcal{E} : \exists e' \in \mathcal{E} \text{ such that } \Pi_e B \Pi_{e'} \neq 0 \text{ or } \Pi_{e'} B \Pi_e \neq 0\}. \quad (25)$$

Moreover, we also define the quantities  $d_E, D_E, D_G$  and  $G(\kappa)$  relative to  $B$ :

$$d_{E,B} := |\mathcal{E}_B|, \quad (26)$$

$$D_{E,B} := \max_{e \in \mathcal{E}_B} \text{tr}(\Pi_e), \quad (27)$$

$$D_{G,B} := \max_{E \in \mathbb{R}} |\{(e, e') \in \mathcal{E}_B \times \mathcal{E}_B : e \neq e' \text{ and } e - e' = E\}|, \quad (28)$$

$$G_B(\kappa) := \max_{E \in \mathbb{R}} |\{(e, e') \in \mathcal{E}_B \times \mathcal{E}_B : e \neq e' \text{ and } e - e' \in [E, E + \kappa)\}|. \quad (29)$$

### 3 Main Result

After having introduced the relevant quantities and notations in the previous section, we are now ready to state our main result.

**Theorem 1** (Normal Typicality for  $\text{GAP}(\rho)$ ). *Let  $B$  be an operator with  $d_{E,B} < \infty$  and let  $\rho$  be a density matrix with  $\|\rho\| < 1/4$  on a separable Hilbert space  $\mathcal{H}$ . Let  $\dim \mathcal{H} \geq 4$ , let  $\varepsilon, \delta, \kappa, T > 0$  and define*

$$M_{\rho B} := \sum_{e \in \mathcal{E}} \text{tr}(\rho \Pi_e B \Pi_e). \quad (30)$$

*Then w.r.t.  $\text{GAP}(\rho)$ ,  $(1 - \varepsilon)$ -most  $\psi_0 \in \mathbb{S}(\mathcal{H})$  are such that for  $(1 - \delta)$ -most  $t \in [0, T]$*

$$\left| \langle \psi_t | B | \psi_t \rangle - M_{\rho B} \right| \leq \left( \frac{188}{\varepsilon \delta} \|B\|^2 \|\rho\| D_{E,B} G_B(\kappa) \left( 1 + \frac{8 \log_2 d_{E,B}}{\kappa T} \right) \right)^{1/2} \quad (31)$$



Moreover, w.r.t.  $\text{GAP}(\rho)$ ,  $(1 - \varepsilon)$ -most  $\psi_0 \in \mathbb{S}(\mathcal{H})$  are such that for  $(1 - \delta)$ -most  $t \in [0, \infty)$ ,

$$\left| \langle \psi_t | B | \psi_t \rangle - M_{\rho B} \right| \leq \left( \frac{188}{\varepsilon \delta} \|B\|^2 \|\rho\| D_{E,B} D_{G,B} \right)^{1/2}. \quad (32)$$

Thus, as soon as  $\|\rho\| \ll D_{E,B} D_{G,B} \|B\|^2$ , i.e., as soon as the largest eigenvalue of  $\rho$  is small, no eigenvalue and no gap of  $H$  is hugely degenerate and  $\|B\|$  is not too large, for most initial states  $\psi_0 \in \mathbb{S}(\mathcal{H})$ , where “most” refers to  $\text{GAP}(\rho)$ , the expectation value  $\langle \psi_t | B | \psi_t \rangle$  of  $B$  in the state  $\psi_t$  is close to the fixed value  $M_{\rho B}$  for most times  $t \geq 0$ . Put another way, the curve  $t \mapsto \langle \psi_t | B | \psi_t \rangle$  is nearly constant in the long run  $t \rightarrow \infty$ . The finite-time result shows that as soon as  $\|\rho\| \ll D_{E,B} G_B(\kappa) \|B\|^2$  and  $T$  is large enough, for most (w.r.t.  $\text{GAP}(\rho)$ ) initial states  $\psi_0 \in \mathbb{S}(\mathcal{H})$  the expectation  $\langle \psi_t | B | \psi_t \rangle$  is close to  $M_{\rho B}$  for most times  $t \in [0, T]$ . Unfortunately, the times  $T$  required to make the upper bound in (31) small are usually extremely large: for example for a system of  $N$  particles we need that  $T \gg \exp(N)$ , see, e.g., the discussion in [42] after Theorem 4.

We remark that for  $\rho = P_\mu/d_\mu$  for some macro state  $\mu$ ,  $\text{GAP}(\rho)$  becomes the uniform distribution over the sphere  $\mathbb{S}(\mathcal{H}_\mu)$ , we have  $\|\rho\| = 1/d_\mu$  and in this case we basically recover the results from [42]. Note that in Theorem 1 for any  $B$  only the eigenvalues in  $\mathcal{E}_B$  and gaps in  $\mathcal{E}_B \times \mathcal{E}_B$  contribute to the quantities  $D_{E,B}$ ,  $d_{E,B}$ ,  $D_{G,B}$  and  $G_B(\kappa)$ . However, if  $\mathcal{H}$  is finite-dimensional, these quantities can obviously be replaced by  $D_E$ ,  $d_E$ ,  $D_G$  and  $G(\kappa)$ . The technical assumption on  $B$  that  $|\mathcal{E}_B| < \infty$  is needed to ensure that the sums over  $e$  appearing in the proofs are finite and it therefore is for example unproblematic to interchange the sums over  $e$  with the time average.

Motivated by the phenomenon of normal typicality [16, 18, 30, 46], in [42] the case that  $B = P_\nu$  for some macro state  $\nu$  was of particular interest. As discussed in the introduction, the  $P_\nu$  usually project to macro spaces  $\mathcal{H}_\nu$  which are contained in a finite-dimensional energy shell. Thus the  $P_\nu$  fulfill the assumption of Theorem 1 that  $|\mathcal{E}_{P_\nu}| < \infty$ . We therefore see that not only with respect to the uniform distribution over some subspace  $\mathcal{H}_\mu$  but also with respect to  $\text{GAP}(\rho)$ , for most initial wave functions  $\psi_0$  the superposition weights  $\|P_\nu \psi_t\|^2$  are, in the long run, close to some fixed value. In the case of normal typicality, this value was given by  $d_\nu/D$ , where  $D = \dim \mathcal{H}$ . Here, if  $D < \infty$ , we also expect that  $M_{\rho P_\nu}$  is usually close to  $d_\nu/D$ ; we can argue that this should be the case for example if  $H$  is non-degenerate and satisfies the eigenstate thermalization hypothesis in the sense that for every eigenstate  $|m\rangle$  of  $H$  we have that  $\langle m | P_\nu | m \rangle \approx \langle P_\nu \rangle_{\text{mic}} = \text{tr}(P_\nu)/D = d_\nu/D$  where  $\langle \cdot \rangle_{\text{mic}}$  denotes the micro-canonical expectation. Then we find

$$M_{\rho P_\nu} = \sum_m \langle m | \rho | m \rangle \langle m | P_\nu | m \rangle \approx \frac{d_\nu}{D} \sum_m \langle m | \rho | m \rangle = \frac{d_\nu}{D}. \quad (33)$$

If  $\dim \mathcal{H} = \infty$  (and there are infinitely many macro states  $\nu$ ), the  $\|P_\nu \psi_t\|^2$  form a null sequence for every  $\psi_0$  and  $t$ .

We now give a brief outline of the proof of Theorem 1. An important ingredient for the proof is an upper bound for the variance of  $\langle \psi | A | \psi \rangle$  where  $A$  is a bounded operator on  $\mathcal{H}$  and  $\psi \sim \text{GAP}(\rho)$ . Reimann [29] proved such an upper bound under the assumption that  $A$  is self-adjoint and  $\mathcal{H}$  is finite-dimensional. In [44] the proof was generalized to arbitrary bounded operators  $A$  on a separable Hilbert space  $\mathcal{H}$ . Unfortunately, the bounds obtained there are not sufficiently sharp for our purpose as it would lead in Theorem 1 to an upper bound of the order

$$\text{tr} \rho^2 \sum_{e \neq e'} \|\Pi_e B \Pi_{e'}\|^2 \leq \|\rho\| \sum_{e, e'} \text{tr}(\Pi_{e'} B^* \Pi_e B \Pi_{e'}) = \|\rho\| \text{tr}(B^* B) \quad (34)$$

and for example in the special case that  $\rho = P_\mu/d_\mu$  and  $B = P_\nu$  this would yield an upper bound of the order  $d_\nu/d_\mu$  which is not necessarily small (here we ignored the possible degeneracy of the eigenvalues and gaps of  $H$  to simplify the discussion). Therefore, using similar techniques as in [29, 44], we prove in Lemma 1 a slightly improved upper bound for the variance that allows us to obtain a bound of the order  $\|B\| \text{tr} \rho^2$  in Theorem 1. After having shown this improved upper bound for the variance of  $\langle \psi | A | \psi \rangle$ , we proceed to compute upper bounds for the expectation w.r.t.  $\text{GAP}(\rho)$  of the infinite time variance of  $\langle \psi_t | B | \psi_t \rangle$ , a related quantity where the outer infinite time average is replaced by a finite one and the variance w.r.t.  $\text{GAP}(\rho)$  of the infinite time average  $\langle \psi_t | B | \psi_t \rangle$  in Proposition 1 making use of the bound for the variance of  $\langle \psi | A | \psi \rangle$ . In these computations, many traces and products and sums over traces appear and we use similar methods as in [42] to bound them. Finally, with the help of these bounds and Markov's and Chebyshev's inequality we are able to prove Theorem 1.

**Remark 1** (Applying Lévy's Lemma). Another strategy of proof of Theorem 1 is to make use of Lévy's Lemma for GAP measures, a concentration-of-measure-type result for Lipschitz continuous functions on the sphere. We quote the result from [44]:

**Theorem 2** (Lévy's Lemma for GAP measures [44]). *Let  $\mathcal{H}$  be a separable Hilbert space, let  $f : \mathbb{S}(\mathcal{H}) \rightarrow \mathbb{C}$  be a Lipschitz continuous function with Lipschitz constant  $\eta$ , let  $\rho$  be a density matrix on  $\mathcal{H}$  and let  $\varepsilon \geq 0$ . Then,*

$$\text{GAP}(\rho) \left\{ \psi \in \mathbb{S}(\mathcal{H}) : |f(\psi) - \mathbb{E}_\rho(f)| > \varepsilon \right\} \leq 12 \exp \left( -\frac{C\varepsilon^2}{2\eta^2 \|\rho\|} \right), \quad (35)$$

where  $\frac{1}{288\pi^2}$ .

In fact, one can show that  $f(\psi_0) := \langle |\langle \psi_t | B | \psi_t \rangle - M_{\rho B}|^2 \rangle_T$  is Lipschitz continuous with Lipschitz constant bounded by  $8\|B\|^2$  (and the same holds true for the infinite time average as well). However, as it is also explained in [42], in such a situation Lévy's Lemma does in general not lead to better results than applying Markov's and Chebyshev's inequality. Markov's inequality gives

$$f(\psi_0) \leq \frac{\mathbb{E}_\rho f}{\varepsilon} \quad (36)$$

for  $(1 - \varepsilon)$ -most  $\psi_0 \in \mathbb{S}(\mathcal{H})$  while it follows from Theorem 2 that

$$f(\psi_0) \leq \mathbb{E}_\rho f + \sqrt{\frac{2\eta^2 \|\rho\| \log(12/\varepsilon)}{C}}. \quad (37)$$

The expectation  $\mathbb{E}_\rho f$  is of the order  $\|\rho\|$  while the square root in (37) is of the order  $\|\rho\|^{1/2}$ , so while the  $\varepsilon$ -dependence in the result from Lévy's Lemma is better, the dependence on  $\|\rho\|$  is better for Markov's inequality. Often,  $\|\rho\|$  is extremely small, for example for  $\rho = \mathbb{1}_{\mathcal{H}}/D$  where  $D$  is the dimension of the (finite-dimensional) Hilbert space and  $\mathbb{1}_{\mathcal{H}}$  the identity on  $\mathcal{H}$ , we have that  $\|\rho\| = 1/D$ . In such cases, the better  $\varepsilon$ -dependence does not compensate the worse  $\|\rho\|$ -dependence unless  $\varepsilon$  is extremely small and it is of little interest to consider, e.g.,  $\varepsilon$  smaller than  $10^{-200}$  [3].

Another possibility to obtain bounds on  $|\langle \psi_t | B | \psi_t \rangle - M_{\rho B}|$  with a better  $\varepsilon$ -dependence is to first apply Lévy's Lemma for GAP measures to the function  $f(\psi) = \langle \psi_t | B | \psi_t \rangle$  for fixed  $t \geq 0$ . Note that by Lemma 5 in [28],  $f$  is Lipschitz continuous with Lipschitz constant bounded by  $2\|B\|$ . Then it follows from Theorem 2 that for every  $t \geq 0$ , w.r.t.  $\text{GAP}(\rho)$ ,  $(1 - \varepsilon)$ -most  $\psi_0 \in \mathbb{S}(\mathcal{H})$  are such that

$$\left| \langle \psi_t | B | \psi_t \rangle - \text{tr}(e^{iHt} B e^{-iHt} \rho) \right| \leq \left( \frac{8\|B\|^2 \|\rho\| \log(12/\varepsilon)}{C} \right)^{1/2}. \quad (38)$$

Using similar estimates as in the proof of Proposition 1 to bound  $|\text{tr}(e^{iHt} B e^{-iHt}) - M_{\rho B}|$  for  $(1 - \delta)$ -most  $t \in [0, T]$  one can show that  $(1 - \delta)$ -most  $t \in [0, T]$  are such that for  $(1 - \varepsilon)$ -most  $\psi_0 \in \mathbb{S}(\mathcal{H})$ ,

$$\left| \langle \psi_t | B | \psi_t \rangle - M_{\rho B} \right| \leq 4 \left( \frac{\|B\|^2 \|\rho\| D_{E,B} G_B(\kappa) \log(12/\varepsilon)}{\delta C} \left( 1 + \frac{8 \log_2 d_{E,B}}{\kappa T} \right) \right)^{1/2}. \quad (39)$$

Note that similar estimates can be obtained for the infinite time interval.

As we are interested in a statement of the form that most  $\psi_0 \in \mathbb{S}(\mathcal{H})$  are such that for most times the error between  $\langle \psi_t | B | \psi_t \rangle$  and  $M_{\rho B}$  is small, we have to interchange “most  $\psi_0$ ” and “most  $t$ ”. When considering a finite time interval  $[0, T]$  (however, not for the infinite time interval  $[0, \infty)$ <sup>3</sup>) this is possible by Footnote 7 in [16] at the expense of

---

<sup>3</sup>While a statement that is true for most  $\psi_0 \in \mathbb{S}(\mathcal{H})$  for most  $t \in [0, \infty)$  is also true for most  $t \in [0, \infty)$  for most  $\psi_0 \in \mathbb{S}(\mathcal{H})$  (which can be proved similarly as Footnote 7 in [16] and by making use of Fatou's Lemma), the converse does not hold. In general, the quantifier “most  $t \in [0, \infty)$ ” and a “most”-quantifier referring to a probability measure on a probability space and cannot be interchanged. As a simple example consider the interval  $[0, 1]$  equipped with the Lebesgue measure and let  $N \in \mathbb{N}$  be large. We define  $I_1 := [0, 1/N)^c$ ,  $I_2 := [1/N, 2/N)^c, \dots, I_N := [(N-1)/N, 1]^c$ , where the complement is taken in  $[0, 1]$ . We define the set

$$S := \bigcup_{m=0}^{\infty} \bigcup_{n=1}^N \left( I_n \times [2^{mN+n-1}, 2^{mN+n}) \right) \subset [0, 1] \times [0, \infty).$$

Clearly, for every  $t > 1$  we have that  $(\psi, t) \in S$  for  $(1 - 1/N)$ -most  $\psi \in [0, 1]$ . Therefore, the statement

making the parameter  $\varepsilon$  and  $\delta$  worse. More precisely, if a statement is true for  $(1 - \delta)$ -most  $t \in [0, T]$  for  $(1 - \varepsilon)$ -most  $\psi_0 \in \mathbb{S}(\mathcal{H})$ , then it is also true for  $(1 - \varepsilon')$ -most  $\psi_0 \in \mathbb{S}(\mathcal{H})$  for  $(1 - \delta')$ -most  $t \in [0, T]$  where  $\varepsilon' \geq \frac{\varepsilon + \delta - \varepsilon\delta}{\delta'}$ .

We do not apply Footnote 7 in [16] directly to (39) but rather go back to (38) and apply it to this statement. We obtain that for every  $\varepsilon, \delta > 0$ , w.r.t.  $\text{GAP}(\rho)$ ,  $(1 - \varepsilon/\delta)$ -most  $\psi_0 \in \mathbb{S}(\mathcal{H})$  are such that for  $(1 - \delta/2)$ -most  $t \in [0, T]$ ,

$$\left| \langle \psi_t | B | \psi_t \rangle - \text{tr} (e^{iHt} B e^{-iHt} \rho) \right| \leq \left( \frac{8 \|B\|^2 \|\rho\| \log(12/\varepsilon)}{C} \right)^{1/2}. \quad (40)$$

Altogether we find that, w.r.t.  $\text{GAP}(\rho)$ ,  $(1 - \varepsilon/\delta)$ -most  $\psi_0 \in \mathbb{S}(\mathcal{H})$  are such that for  $(1 - \delta)$ -most  $t \in [0, T]$ ,

$$\left| \langle \psi_t | B | \psi_t \rangle - M_{\rho B} \right| \leq 5 \left( \frac{\|B\|^2 \|\rho\| D_{E,B} G_B(\kappa) \log(12/\varepsilon)}{\delta C} \left( 1 + \frac{8 \log_2 d_{E,B}}{\kappa T} \right) \right)^{1/2}. \quad (41)$$

Effectively, the  $1/\varepsilon$  in Theorem 1 is replaced by  $\log(12/(\varepsilon\delta))$  and in relevant cases (i.e.,  $\varepsilon \ll \delta$ ) this bound is slightly better than the one from Theorem 1 if  $\delta$  is sufficiently small.  $\diamond$

## 4 Proofs

This section is devoted to the proof of our main result. As already outlined in Section 3, we first prove an improved upper bound for the variance  $\text{Var}_\rho \langle \psi | A | \psi \rangle$  with respect to  $\text{GAP}(\rho)$  in Section 4.1. With the help of this bound, we prove in Section 4.2 an upper bound for the expected time variance of  $\langle \psi_t | B | \psi_t \rangle$ , i.e., for

$$\mathbb{E}_\rho \left( \left\langle \left| \langle \psi_t | B | \psi_t \rangle - \overline{\langle \psi_t | B | \psi_t \rangle} \right|^2 \right\rangle_T \right) \quad (42)$$

and also for the infinite time average instead of  $\langle \cdot \rangle_T$ . Moreover, an application of Lemma 1 also yields an upper bound for  $\text{Var}_\rho \langle \psi_t | B | \psi_t \rangle$ . After having proved these upper bounds in Section 4.2, the proof of Theorem 1 which we give in Section 4.3 basically reduces to a careful application of Markov's and Chebyshev's inequality.

---

“ $(\psi, t) \in S$ ” is true for most  $t \in [0, \infty)$  for most  $\psi \in [0, 1]$ .

Let  $\psi \in [0, 1]$ . Without loss of generality we assume that  $\psi \in [0, 1/N)$  which implies that  $\psi \notin I_1$  and  $\psi \in I_k$  for all  $k \geq 2$ . We define a sequence of times  $(T_l)_{l \geq 0}$  by  $T_l := 2^{lN+1}$ . Then, by construction,  $(\psi, t) \notin S$  for all  $t \in [2^{lN}, 2^{lN+1})$  and thus  $(\psi, t) \notin S$  for at least half of the times  $t \in [0, T_l]$ . This implies

$$\limsup_{T \rightarrow \infty} \frac{1}{T} \lambda \{t \in [0, T] : (\psi, t) \notin S\} \geq \lim_{l \rightarrow \infty} \frac{1}{T_l} \lambda \{t \in [0, T_l] : (\psi, t) \notin S\} \geq 1/2.$$

As  $\psi \in [0, 1]$  was arbitrary, we conclude that the statement “ $(\psi, t) \in S$ ” is false for every  $\psi \in [0, 1]$  for a substantial amount of times  $t \in [0, \infty)$ .

## 4.1 An Improved Upper Bound for $\text{Var}_\rho \langle \psi | A | \psi \rangle$

**Lemma 1.** *Let  $\rho$  be a density matrix on a separable Hilbert space  $\mathcal{H}$  with eigenvalues  $p_n > 0$  such that  $p_{\max} = \|\rho\| < 1/4$  and let  $\dim \mathcal{H} \geq 4$ . For  $\text{GAP}(\rho)$ -distributed  $\psi$  and any bounded operator  $A : \mathcal{H} \rightarrow \mathcal{H}$ ,*

$$\mathbb{E}_\rho \langle \psi | A | \psi \rangle = \text{tr}(A\rho), \quad (43)$$

$$\begin{aligned} \text{Var}_\rho \langle \psi | A | \psi \rangle &\leq \frac{1}{1 - p_{\max}} \left( \text{tr}(A\rho A^* \rho) + \frac{\text{tr}(A\rho^2 A^* \rho) + \text{tr}(A\rho A^* \rho^2)}{1 - 2p_{\max}} \right. \\ &+ \frac{2}{(1 - 2p_{\max})(1 - 3p_{\max})} \left[ \text{tr}(A\rho^3 A^* \rho) + \text{tr}(A\rho^2 A^* \rho^2) + \text{tr}(A\rho A^* \rho^3) \right. \\ &\left. \left. + \sum_{m,n} (|\text{tr}(A\rho^3 P_m) \text{tr}(A^* \rho P_n)| + |\text{tr}(A\rho^2 P_m) \text{tr}(A^* \rho^2 P_n)| + |\text{tr}(A\rho P_m) \text{tr}(A^* \rho^3 P_n)|) \right] \right), \end{aligned} \quad (44)$$

where  $\mathbb{E}_\rho$  and  $\text{Var}_\rho$  denote the expectation and variance with respect to  $\text{GAP}(\rho)$ ,  $P_n = |n\rangle\langle n|$  and  $\{|n\rangle\}$  is an orthonormal eigenbasis of  $\rho$ .

*Proof.* The formula for the expectation has already been proved in [44, Proposition 1]. For the proof of (44) we adapt the proof for the upper bound on the variance from [44] which in turn follows closely the proof of Reimann [29] who only considered the case that  $A$  is self-adjoint and  $\mathcal{H}$  is finite-dimensional. For convenience of the reader we give most of the details here.

We first assume that  $D := \dim \mathcal{H} < \infty$ . Recall that the variance of a complex-valued random variable  $Z$  is defined as

$$\text{Var}_\rho(Z) = \mathbb{E}_\rho(|Z - \mathbb{E}_\rho Z|^2) = \mathbb{E}_\rho |Z|^2 - |\mathbb{E}_\rho Z|^2. \quad (45)$$

Because of the invariance of  $\text{Var}_\rho$  under the addition of constants, i.e.,  $\text{Var}_\rho(Z + a) = \text{Var}_\rho(Z)$  for all constants  $a \in \mathbb{C}$ , we can assume without loss of generality that  $\mathbb{E}_\rho \langle \psi | A | \psi \rangle = \text{tr}(A\rho) = 0$ . As a consequence, we only have to compute  $\mathbb{E}_\rho(\langle \psi | A | \psi \rangle \langle \psi | A | \psi \rangle^*) = \mathbb{E}_\rho(\langle \psi | A | \psi \rangle \langle \psi | A^* | \psi \rangle)$ .

Let  $\{|n\rangle : n = 1, \dots, D\}$  be an orthonormal basis of  $\mathcal{H}$  consisting of eigenvectors of  $\rho$  and let  $\psi \in \mathbb{S}(\mathcal{H})$ . Then,

$$\langle \psi | A | \psi \rangle \langle \psi | A^* | \psi \rangle = \sum_{m,n,m',n'} \langle \psi | m \rangle \langle m | A | n \rangle \langle n | \psi \rangle \langle \psi | n' \rangle \langle n' | A^* | m' \rangle \langle m' | \psi \rangle \quad (46a)$$

$$= \sum_{m,n,m',n'} c_m^* c_n c_{n'}^* c_{m'} A_{mn} A_{m'n'}^*, \quad (46b)$$

where  $c_m = \langle m | \psi \rangle$  and  $A_{mn} = \langle m | A | n \rangle$ .

Reimann [29] computed the fourth moments  $\mathbb{E}_\rho(c_m^* c_n c_{n'}^* c_{m'})$  and obtained that they vanish except in the two cases (i)  $m = n$  and  $m' = n'$ , (ii)  $m = m'$  and  $n = n'$ , and that

$$\mathbb{E}_\rho(|c_m|^2 |c_n|^2) = p_m p_n (1 + \delta_{mn}) K_{mn}, \quad (47)$$

where

$$K_{mn} = \int_0^\infty (1 + xp_m)^{-1} (1 + xp_n)^{-1} \prod_{l=1}^D (1 + xp_l)^{-1} dx. \quad (48)$$

We compute

$$\begin{aligned} \mathbb{E}_\rho (\langle \psi | A | \psi \rangle \langle \psi | A^* | \psi \rangle) &= \sum_{m,n} p_n p_m (1 + \delta_{mn}) K_{mn} A_{mm} A_{nn}^* + \sum_{m,n} p_n p_m (1 + \delta_{mn}) K_{mn} |A_{mn}|^2 \\ &\quad - 2 \sum_n p_n^2 K_{nn} |A_{nn}|^2 \end{aligned} \quad (49a)$$

$$= \sum_{m,n} [A_{mm} A_{nn}^* + |A_{mn}|^2] p_n p_m K_{mn} \quad (49b)$$

and as in [29] we write  $K_{mn}$  as

$$K_{mn} = K^{(0)} - (p_m + p_n) K^{(1)} + 2\kappa_{mn} (p_m^2 + p_m p_n + p_n^2) K^{(2)}, \quad (50)$$

where

$$K^{(k)} = \frac{1}{k!} \int_0^\infty x^k \prod_{l=1}^D (1 + xp_l)^{-1} dx, \quad k = 0, 1, 2, \quad (51)$$

and  $\kappa_{mn} \in [0, 1]$ . This follows from a Taylor expansion of  $g_{mn}(x) := (1 + xp_m)^{-1} (1 + xp_n)^{-1}$  up to second order.

With (50) we obtain for the first term in (49b) that

$$\begin{aligned} &\sum_{m,n} A_{mm} A_{nn}^* p_n p_m (K^{(0)} - (p_m + p_n) K^{(1)} + 2\kappa_{mn} (p_m^2 + p_m p_n + p_n^2) K^{(2)}) \\ &= K^{(0)} \left( \sum_m A_{mm} p_m \right) \left( \sum_n A_{nn}^* p_n \right) - K^{(1)} \left[ \sum_m A_{mm} p_m^2 \sum_n A_{nn}^* p_n \right. \\ &\quad \left. + \sum_m A_{mm} p_m \sum_n A_{nn}^* p_n^2 \right] + 2K^{(2)} \sum_{m,n} A_{mm} A_{nn}^* \kappa_{mn} (p_m^3 p_n + p_m^2 p_n^2 + p_m p_n^3) \end{aligned} \quad (52a)$$

$$= 2K^{(2)} \sum_{m,n} A_{mm} A_{nn}^* \kappa_{mn} (p_m^3 p_n + p_m^2 p_n^2 + p_m p_n^3), \quad (52b)$$

where we used that

$$\begin{aligned} \sum_m A_{mm} p_m &= \sum_m \langle m | A | m \rangle \langle m | \rho | m \rangle = \sum_{m,m'} \langle m | A | m' \rangle \langle m' | \rho | m \rangle \\ &= \sum_m \langle m | A \rho | m \rangle = \text{tr}(A \rho) = 0, \end{aligned} \quad (53)$$

and similarly  $\sum_n A_{nn}^* p_n = 0$ . Therefore an upper bound for the absolute value of the first term in (49b) is given by

$$\begin{aligned} & 2K^{(2)} \sum_{m,n} |A_{mm} A_{nn}^*| (p_m^3 p_n + p_m^2 p_n^2 + p_m p_n^3) \\ &= 2K^{(2)} \sum_{m,n} (|\operatorname{tr}(A \rho^3 P_m) \operatorname{tr}(A^* \rho P_n)| + |\operatorname{tr}(A \rho^2 P_m) \operatorname{tr}(A^* \rho^2 P_n)| + |\operatorname{tr}(A \rho P_m) \operatorname{tr}(A^* \rho^3 P_n)|), \end{aligned} \quad (54)$$

where  $P_n = |n\rangle\langle n|$ . Here we used for the first term that

$$A_{mm} A_{nn}^* p_m^3 p_n = \langle m | A \rho^3 | m \rangle \langle n | A^* \rho | n \rangle = \operatorname{tr}(A \rho^3 P_m) \operatorname{tr}(A^* \rho P_n) \quad (55)$$

and similarly for the other terms.

For the second term in (49b) we find

$$\begin{aligned} & K^{(0)} \sum_{m,n} |A_{mn}|^2 p_n p_m - K^{(1)} \sum_{m,n} |A_{mn}|^2 (p_n^2 p_m + p_n p_m^2) \\ &+ 2K^{(2)} \sum_{m,n} |A_{mn}|^2 \kappa_{mn} (p_n p_m^3 + p_m^2 p_n^2 + p_n^3 p_m) \end{aligned} \quad (56a)$$

$$\begin{aligned} &= K^{(0)} \operatorname{tr}(A \rho A^* \rho) - K^{(1)} (\operatorname{tr}(A \rho^2 A^* \rho) + \operatorname{tr}(A \rho A^* \rho^2)) \\ &+ 2K^{(2)} \sum_{m,n} |A_{mn}|^2 \kappa_{mn} (p_n p_m^3 + p_m^2 p_n^2 + p_n^3 p_m), \end{aligned} \quad (56b)$$

where we used that

$$\sum_{m,n} |A_{nm}|^2 p_n p_m = \sum_{m,n} A_{mn} A_{mn}^* p_n p_m = \sum_{m,n} \langle m | A \rho | n \rangle \langle n | A^* \rho | m \rangle = \operatorname{tr}(A \rho A^* \rho) \quad (57)$$

and similarly  $\sum_{m,n} A_{mn} A_{mn}^* p_n^2 p_m = \operatorname{tr}(A \rho^2 A^* \rho)$  and  $\sum_{m,n} A_{mn} A_{mn}^* p_n p_m^2 = \operatorname{tr}(A \rho A^* \rho^2)$ .

An upper bound for (56b) is given by

$$K^{(0)} \operatorname{tr}(A \rho A^* \rho) + K^{(1)} (\operatorname{tr}(A \rho^2 A^* \rho) + \operatorname{tr}(A \rho A^* \rho^2)) + 2K^{(2)} \sum_{m,n} |A_{mn}|^2 (p_n p_m^3 + p_m^2 p_n^2 + p_n^3 p_m) \quad (58a)$$

$$\begin{aligned} &= K^{(0)} \operatorname{tr}(A \rho A^* \rho) + K^{(1)} (\operatorname{tr}(A \rho^2 A^* \rho) + \operatorname{tr}(A \rho A^* \rho^2)) \\ &+ 2K^{(2)} (\operatorname{tr}(A \rho^3 A^* \rho) + \operatorname{tr}(A \rho^2 A^* \rho^2) + \operatorname{tr}(A \rho A^* \rho^3)). \end{aligned} \quad (58b)$$

Thus we arrive at

$$\begin{aligned} & \operatorname{Var}_\rho \langle \psi | A | \psi \rangle \\ & \leq K^{(0)} \operatorname{tr}(A \rho A^* \rho) + K^{(1)} (\operatorname{tr}(A \rho^2 A^* \rho) + \operatorname{tr}(A \rho A^* \rho^2)) \\ & + 2K^{(2)} (\operatorname{tr}(A \rho^3 A^* \rho) + \operatorname{tr}(A \rho^2 A^* \rho^2) + \operatorname{tr}(A \rho A^* \rho^3)) \end{aligned}$$

$$+ \sum_{m,n} (|\operatorname{tr}(A\rho^3 P_m) \operatorname{tr}(A^* \rho P_n)| + |\operatorname{tr}(A\rho^2 P_m) \operatorname{tr}(A^* \rho^2 P_n)| + |\operatorname{tr}(A\rho P_m) \operatorname{tr}(A^* \rho^3 P_n)|) \Big). \quad (59)$$

In [29] it was shown that

$$K^{(k)} \leq \prod_{j=1}^{k+1} \frac{1}{1 - j p_{\max}}, \quad k = 0, 1, 2. \quad (60)$$

With the help of this inequality we finally obtain

$$\begin{aligned} \operatorname{Var}_\rho \langle \psi | A | \psi \rangle &\leq \frac{1}{1 - p_{\max}} \left( \operatorname{tr}(A\rho A^* \rho) + \frac{\operatorname{tr}(A\rho^2 A^* \rho) + \operatorname{tr}(A\rho A^* \rho^2)}{1 - 2p_{\max}} \right. \\ &\quad + \frac{2}{(1 - 2p_{\max})(1 - 3p_{\max})} \left[ \operatorname{tr}(A\rho^3 A^* \rho) + \operatorname{tr}(A\rho^2 A^* \rho^2) + \operatorname{tr}(A\rho A^* \rho^3) \right. \\ &\quad \left. \left. + \sum_{m,n} (|\operatorname{tr}(A\rho^3 P_m) \operatorname{tr}(A^* \rho P_n)| + |\operatorname{tr}(A\rho^2 P_m) \operatorname{tr}(A^* \rho^2 P_n)| + |\operatorname{tr}(A\rho P_m) \operatorname{tr}(A^* \rho^3 P_n)|) \right] \right). \end{aligned} \quad (61)$$

This concludes the proof in the finite-dimensional case.

Now suppose that  $\dim \mathcal{H} = \infty$ . We proceed similarly as in the proof of Proposition 1 in [44] and for convenience of the reader, we give in the following the full argument. Let  $\{|k\rangle : k \in \mathbb{N}\}$  be an orthonormal eigenbasis of  $\rho$  with corresponding positive eigenvalues  $p_{\max} = \|\rho\| = p_1 \geq p_2 \geq \dots$ . We approximate  $\rho$  in trace norm by finite-rank density matrices  $\rho_k$ ,  $k \in \mathbb{N}$ , defined by

$$\rho_k := \sum_{m=1}^{k-1} p_m |m\rangle \langle m| + \left( \sum_{m=k}^{\infty} p_m \right) |k\rangle \langle k|. \quad (62)$$

Because of  $\|\rho_k - \rho\|_{\operatorname{tr}} \rightarrow 0$  as  $k \rightarrow \infty$ , it follows from Theorem 3 in [45] that  $\operatorname{GAP}(\rho_k) \Rightarrow \operatorname{GAP}(\rho)$ , i.e., the measures  $\operatorname{GAP}(\rho_k)$  converge weakly to the measure  $\operatorname{GAP}(\rho)$ . Note that for  $k$  large enough,  $\sum_{m \geq k} p_m < p_1 = p_{\max}$  and therefore  $p_{\max,k} = p_{\max}$  where  $p_{\max,k}$  denotes the largest eigenvalue of  $\rho_k$ . We define the functions  $f_k, f : \mathbb{S}(\mathcal{H}) \rightarrow \mathbb{R}$  by

$$f_k(\psi) = |\langle \psi | A | \psi \rangle - \operatorname{tr}(A\rho_k)|^2, \quad (63)$$

$$f(\psi) = |\langle \psi | A | \psi \rangle - \operatorname{tr}(A\rho)|^2. \quad (64)$$

It follows from

$$|\operatorname{tr}(A\rho_k) - \operatorname{tr}(A\rho)| = |\operatorname{tr}(A(\rho_k - \rho))| \leq \|A\| \operatorname{tr}(|\rho_k - \rho|) = \|A\| \|\rho_k - \rho\|_{\operatorname{tr}} \rightarrow 0 \quad (65)$$



that  $f_k \rightarrow f$  uniformly in  $\psi$  and this implies  $\text{GAP}(\rho_k)(f_k) - \text{GAP}(\rho_k)(f) \rightarrow 0$  where we introduced the notation

$$\text{GAP}(\rho)(f) := \int_{\mathbb{S}(\mathcal{H})} f(\psi) d\text{GAP}(\rho)(\psi). \quad (66)$$

As the measures  $\text{GAP}(\rho_k)$  converge weakly to  $\text{GAP}(\rho)$  and  $f$  is continuous, we have that  $\text{GAP}(\rho_k)(f) \rightarrow \text{GAP}(\rho)(f)$  and therefore  $\text{Var}_{\rho_k} \langle \psi | A | \psi \rangle = \text{GAP}(\rho_k)(f_k) \rightarrow \text{GAP}(\rho)(f) = \text{Var}_{\rho} \langle \psi | A | \psi \rangle$ . By restricting to the subspace spanned by  $\{|n\rangle : n = 1, \dots, k\}$  we obtain from the finite-dimensional case that

$$\begin{aligned} \text{Var}_{\rho_k} \langle \psi | A | \psi \rangle &\leq \frac{1}{1 - p_{\max,k}} \left( \text{tr}(A\rho_k A^* \rho_k) + \frac{\text{tr}(A\rho_k^2 A^* \rho_k) + \text{tr}(A\rho_k A^* \rho_k^2)}{1 - 2p_{\max,k}} \right. \\ &+ \frac{2}{(1 - 2p_{\max,k})(1 - 3p_{\max,k})} \left[ \text{tr}(A\rho_k^3 A^* \rho_k) + \text{tr}(A\rho_k^2 A^* \rho_k^2) + \text{tr}(A\rho_k A^* \rho_k^3) \right. \\ &\left. \left. + \sum_{m,n} (|\text{tr}(A\rho_k^3 P_m) \text{tr}(A^* \rho_k P_n)| + |\text{tr}(A\rho_k^2 P_m) \text{tr}(A^* \rho_k^2 P_n)| + |\text{tr}(A\rho_k P_m) \text{tr}(A^* \rho_k^3 P_n)|) \right] \right), \end{aligned} \quad (67)$$

where we chose  $k$  large enough such that  $p_{\max,k} < 1/4$ . We have already discussed that  $p_{\max,k} = p_{\max}$  for  $k$  large enough. Moreover, we find

$$\begin{aligned} \|\rho_k^2 - \rho^2\|_{\text{tr}} &= \|(\rho_k + \rho)(\rho_k - \rho)\|_{\text{tr}} \leq \|\rho_k + \rho\| \|\rho_k - \rho\|_{\text{tr}} \leq 2\|\rho_k - \rho\|_{\text{tr}} \rightarrow 0, \quad (68) \\ \|\rho_k^3 - \rho^3\|_{\text{tr}} &= \|(\rho_k^2 - \rho^2)(\rho_k + \rho) + \rho_k(\rho - \rho_k)\rho\|_{\text{tr}} \leq 2\|\rho^2 - \rho_k^2\|_{\text{tr}} + \|\rho_k - \rho\|_{\text{tr}} \rightarrow 0 \end{aligned} \quad (69)$$

and therefore all traces in the third line in (67) converge to the corresponding trace with  $\rho$  instead of  $\rho_k$ ; this can be seen as in (65). The traces in the first and second line also converge to the same expression with  $\rho_k$  replaced by  $\rho$ ; this follows from

$$|\text{tr}(A\rho_k A^* \rho_k) - \text{tr}(A\rho A^* \rho)| \leq |\text{tr}(A\rho_k A^* (\rho_k - \rho))| + |\text{tr}(A(\rho - \rho_k) A^* \rho)| \quad (70a)$$

$$\leq 2\|A\|^2 \|\rho_k - \rho\|_{\text{tr}} \rightarrow 0 \quad (70b)$$

and similarly for the other terms. Thus taking the limit  $k \rightarrow \infty$  in (67) shows that the upper bound for the variance remains true in the infinite-dimensional case.  $\square$

## 4.2 Upper Bounds for Some Variances and Averaged Deviations

Before we state and prove these upper bounds, we first note that for any operator  $B$  on  $\mathcal{H}$  such that  $\Pi_{e'} B \Pi_e \neq 0$  only for finitely many  $e, e'$  and any  $\psi_0 \in \mathbb{S}(\mathcal{H})$  the limit

$$M_{\psi_0 B} = \overline{\langle \psi_t | B | \psi_t \rangle} := \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T \langle \psi_t | B | \psi_t \rangle dt \quad (71)$$

exists and can be computed as follows:

$$M_{\psi_0 B} = \overline{\left\langle \psi_0 \left| e^{iHt} \sum_{e \in \mathcal{E}} \Pi_e B \sum_{e' \in \mathcal{E}} \Pi_{e'} e^{-iHt} \right| \psi_0 \right\rangle} \quad (72a)$$

$$= \sum_{e, e' \in \mathcal{E}} \overline{e^{i(e-e')t} \langle \psi_0 | \Pi_e B \Pi_{e'} | \psi_0 \rangle} \quad (72b)$$

$$= \sum_{e \in \mathcal{E}} \langle \psi_0 | \Pi_e B \Pi_e | \psi_0 \rangle. \quad (72c)$$

An application of (43) immediately shows that

$$\mathbb{E}_\rho M_{\psi_0 B} = \sum_{e \in \mathcal{E}} \text{tr}(\rho \Pi_e B \Pi_e) = M_{\rho B}. \quad (73)$$

Note that interchanging the sums over  $\mathcal{E}$  with the time average and expectation with respect to  $\text{GAP}(\rho)$  are unproblematic as due to our assumption on  $B$  the sums are effectively sums over only finitely many  $e, e'$ .

**Proposition 1.** *Let  $\rho$  be a density matrix on a separable Hilbert space  $\mathcal{H}$  with eigenvalues  $p_n > 0$  and  $\|\rho\| < 1/4$ , let  $\dim \mathcal{H} \geq 4$  and let  $B$  be an operator on  $\mathcal{H}$  such that  $d_{E,B} < \infty$ . Then for every  $\kappa, T > 0$ ,*

$$\mathbb{E}_\rho \left( \left\langle \left| \langle \psi_t | B | \psi_t \rangle - \overline{\langle \psi_t | B | \psi_t \rangle} \right|_T^2 \right\rangle \right) \leq 24 \|B\|^2 \|\rho\| D_{E,B} G_B(\kappa) \left( 1 + \frac{8 \log_2 d_{E,B}}{\kappa T} \right), \quad (74)$$

$$\text{Var}_\rho \overline{\langle \psi_t | B | \psi_t \rangle} \leq 23 \|B\|^2 \|\rho\|. \quad (75)$$

Moreover,

$$\mathbb{E}_\rho \left( \overline{\left| \langle \psi_t | B | \psi_t \rangle - \overline{\langle \psi_t | B | \psi_t \rangle} \right|^2} \right) \leq 24 \|B\|^2 \|\rho\| D_{E,B} D_{G,B}. \quad (76)$$

*Proof.* We first assume that  $\mathcal{H}$  is finite-dimensional. The proof starts as the one of Proposition 1 in [42] but, for convenience of the reader, we repeat the steps here so that it is later, when we discuss the infinite-dimensional case, easier to see which steps remain true and where the subtleties are. We find that

$$\left\langle \left| \langle \psi_t | B | \psi_t \rangle - \overline{\langle \psi_t | B | \psi_t \rangle} \right|_T^2 \right\rangle = \left\langle \left| \sum_{e, e' \in \mathcal{E}} e^{i(e-e')t} \langle \psi_0 | \Pi_e B \Pi_{e'} | \psi_0 \rangle - \sum_{e \in \mathcal{E}} \langle \psi_0 | \Pi_e B \Pi_e | \psi_0 \rangle \right|_T^2 \right\rangle \quad (77a)$$

$$= \left\langle \left| \sum_{e \neq e'} e^{i(e-e')t} \langle \psi_0 | \Pi_e B \Pi_{e'} | \psi_0 \rangle \right|_T^2 \right\rangle \quad (77b)$$

$$= \sum_{\substack{e \neq e' \\ e'' \neq e'''}} \left\langle e^{i(e-e'-e''+e''')t} \right\rangle_T \langle \psi_0 | \Pi_e B \Pi_{e'} | \psi_0 \rangle \langle \psi_0 | \Pi_{e''} B^* \Pi_{e'''} | \psi_0 \rangle \quad (77c)$$

$$=: \sum_{\alpha, \beta} v_\alpha^* R_{\alpha\beta} v_\beta, \quad (77d)$$

where we defined for  $\alpha = (e, e'), \beta = (e'', e''') \in \mathcal{G} := \{(\bar{e}, \bar{e}') \in \mathcal{E} \times \mathcal{E}, \bar{e} \neq \bar{e}'\}$  the vector  $v_\alpha = \langle \psi_0 | \Pi_{e'} B^* \Pi_e | \psi_0 \rangle$  and the Hermitian matrix

$$R_{\alpha\beta} = \left\langle e^{i(G_\alpha - G_\beta)t} \right\rangle_T \quad (78)$$

with  $G_\alpha = e - e'$ . Written in this way, we immediately see that

$$\left\langle \left| \langle \psi_t | B | \psi_t \rangle - \overline{\langle \psi_t | B | \psi_t \rangle} \right|^2 \right\rangle_T \leq \|R\| \sum_{\alpha} |v_\alpha|^2 \leq \|R\| \sum_{e, e'} |\langle \psi_0 | \Pi_e B \Pi_{e'} | \psi_0 \rangle|^2. \quad (79)$$

It was proved by Short and Farrelly [36] that the operator norm of  $R$  can be bounded as

$$\|R\| \leq G(\kappa) \left( 1 + \frac{8 \log_2 d_E}{\kappa T} \right). \quad (80)$$

We further compute

$$\mathbb{E}_\rho \left( \sum_{e, e'} |\langle \psi_0 | \Pi_e B \Pi_{e'} | \psi_0 \rangle|^2 \right) = \sum_{e, e'} \text{Var}_\rho \langle \psi_0 | \Pi_e B \Pi_{e'} | \psi_0 \rangle + |\mathbb{E}_\rho \langle \psi_0 | \Pi_e B \Pi_{e'} | \psi_0 \rangle|^2 \quad (81a)$$

$$= \sum_{e, e'} \text{Var}_\rho \langle \psi_0 | \Pi_e B \Pi_{e'} | \psi_0 \rangle + |\text{tr}(\rho \Pi_e B \Pi_{e'})|^2, \quad (81b)$$

where we applied Lemma 1 in the second step. It follows again from Lemma 1 that

$$\begin{aligned} & \sum_{e, e'} \text{Var}_\rho \langle \psi_0 | \Pi_e B \Pi_{e'} | \psi_0 \rangle \\ & \leq \frac{1}{1 - p_{\max}} \sum_{e, e'} \left( \text{tr}(\Pi_e B \Pi_{e'} \rho \Pi_{e'} B^* \Pi_e \rho) + \frac{\text{tr}(\Pi_e B \Pi_{e'} \rho^2 \Pi_{e'} B^* \Pi_e \rho) + \text{tr}(\Pi_e B \Pi_{e'} \rho \Pi_{e'} B^* \Pi_e \rho^2)}{1 - 2p_{\max}} \right. \\ & \quad + \frac{2}{(1 - 2p_{\max})(1 - 3p_{\max})} [\text{tr}(\Pi_e B \Pi_{e'} \rho^3 \Pi_{e'} B^* \Pi_e \rho) + \text{tr}(\Pi_e B \Pi_{e'} \rho^2 \Pi_{e'} B^* \Pi_e \rho^2) \\ & \quad + \text{tr}(\Pi_e B \Pi_{e'} \rho \Pi_{e'} B^* \Pi_e \rho^3) + \sum_{m, n} (|\text{tr}(\Pi_e B \Pi_{e'} \rho^3 P_m) \text{tr}(\Pi_{e'} B^* \Pi_e \rho P_n)| \\ & \quad \left. + |\text{tr}(\Pi_e B \Pi_{e'} \rho^2 P_m) \text{tr}(\Pi_{e'} B^* \Pi_e \rho^2 P_n)| + |\text{tr}(\Pi_e B \Pi_{e'} \rho P_m) \text{tr}(\Pi_{e'} B^* \Pi_e \rho^3 P_n)|) \right]. \end{aligned} \quad (82)$$

The main tools for bounding the traces are the Cauchy-Schwarz inequality for the trace which states that  $|\operatorname{tr}(A^*B)| \leq \sqrt{\operatorname{tr}(A^*A)\operatorname{tr}(B^*B)}$  for any operators  $A$  and  $B$  and the inequality  $|\operatorname{tr}(AB)| \leq \|A\| \operatorname{tr}(|B|)$ ; for the latter see, e.g., Theorem 3.7.6 in [37]. We compute

$$\sum_{e,e'} \operatorname{tr}(\Pi_e B \Pi_{e'} \rho \Pi_{e'} B^* \Pi_e \rho) = \sum_e \operatorname{tr} \left( B \left( \sum_{e'} \Pi_{e'} \rho \Pi_{e'} \right) B^* \Pi_e \rho \Pi_e \right) \quad (83)$$

$$\leq \sum_e \|B\|^2 \left\| \sum_{e'} \Pi_{e'} \rho \Pi_{e'} \right\| \operatorname{tr}(\Pi_e \rho \Pi_e) \quad (84)$$

$$\leq \|B\|^2 \|\rho\|, \quad (85)$$

where we used that  $\|\sum_{e'} \Pi_{e'} \rho \Pi_{e'}\| \leq \|\rho\|$  which follows immediately from

$$\left\| \sum_{e'} \Pi_{e'} \rho \Pi_{e'} \psi \right\|^2 = \sum_{e'} \|\Pi_{e'} \rho \Pi_{e'} \psi\|^2 \leq \|\rho\|^2 \sum_{e'} \|\Pi_{e'} \psi\|^2 = \|\rho\|^2 \|\psi\|^2 \quad (86)$$

for all  $\psi \in \mathcal{H}$ . In the same way we get

$$\sum_{e,e'} \operatorname{tr}(\Pi_e B \Pi_{e'} \rho^2 \Pi_{e'} B^* \Pi_e \rho) \leq \|B\|^2 \|\rho\|^2, \quad (87)$$

$$\sum_{e,e'} \operatorname{tr}(\Pi_e B \Pi_{e'} \rho \Pi_{e'} B^* \Pi_e \rho^2) \leq \|B\|^2 \|\rho\|^2, \quad (88)$$

$$\sum_{e,e'} \operatorname{tr}(\Pi_e B \Pi_{e'} \rho^3 \Pi_{e'} B^* \Pi_e \rho) \leq \|B\|^2 \|\rho\|^3, \quad (89)$$

$$\sum_{e,e'} \operatorname{tr}(\Pi_e B \Pi_{e'} \rho^2 \Pi_{e'} B^* \Pi_e \rho^2) \leq \|B\|^2 \|\rho\|^2 \operatorname{tr} \rho^2 \leq \|B\|^2 \|\rho\|^3, \quad (90)$$

$$\sum_{e,e'} \operatorname{tr}(\Pi_e B \Pi_{e'} \rho \Pi_{e'} B^* \Pi_e \rho^3) \leq \|B\|^2 \|\rho\|^3. \quad (91)$$

Next we turn to the products of traces; we find that

$$\begin{aligned} & \sum_{e,e'} \sum_{m,n} |\operatorname{tr}(\Pi_e B \Pi_{e'} \rho^3 P_m) \operatorname{tr}(\Pi_{e'} B^* \Pi_e \rho P_n)| \\ &= \sum_{m,n} \sum_{e,e'} |\operatorname{tr}(\Pi_e B \Pi_{e'} \Pi_{e'} P_m \Pi_e) \operatorname{tr}(\Pi_{e'} B^* \Pi_e \Pi_e P_n \Pi_{e'})| p_m^3 p_n \end{aligned} \quad (92a)$$

$$\leq \sum_{m,n} p_m^3 p_n \sum_{e,e'} (\operatorname{tr}(\Pi_e B \Pi_{e'} B^*) \operatorname{tr}(\Pi_e P_m \Pi_{e'} P_m) \operatorname{tr}(\Pi_{e'} B^* \Pi_e B) \operatorname{tr}(\Pi_{e'} P_n \Pi_e P_n))^{1/2} \quad (92b)$$

$$= \sum_{m,n} p_m^3 p_n \sum_{e,e'} \operatorname{tr}(\Pi_e B \Pi_{e'} B^*) (\langle m | \Pi_e | m \rangle \langle m | \Pi_{e'} | m \rangle \langle n | \Pi_e | n \rangle \langle n | \Pi_{e'} | n \rangle)^{1/2} \quad (92c)$$

$$\leq \|B\|^2 \sum_{m,n} p_m^3 p_n \sum_{e,e'} \text{tr}(\Pi_e) (\langle m|\Pi_e|m\rangle \langle m|\Pi_{e'}|m\rangle \langle n|\Pi_e|n\rangle \langle n|\Pi_{e'}|n\rangle)^{1/2} \quad (92d)$$

$$\leq \|B\|^2 D_E \sum_{m,n} p_m^3 p_n \left( \sum_e (\langle m|\Pi_e|m\rangle \langle n|\Pi_e|n\rangle)^{1/2} \right)^2 \quad (92e)$$

$$\leq \|B\|^2 D_E \sum_{m,n} p_m^3 p_n \left( \sum_e \langle m|\Pi_e|m\rangle \right) \left( \sum_e \langle n|\Pi_e|n\rangle \right) \quad (92f)$$

$$= \|B\|^2 D_E \sum_{m,n} p_m^3 p_n \quad (92g)$$

$$\leq \|B\|^2 D_E \|\rho\|^2. \quad (92h)$$

In the same way we obtain

$$\sum_{e,e'} \sum_{m,n} |\text{tr}(\Pi_e B \Pi_{e'} \rho^2 P_m) \text{tr}(\Pi_{e'} B^* \Pi_e \rho^2 P_n)| \leq \|B\|^2 D_E \|\rho\|^2, \quad (93)$$

$$\sum_{e,e'} \sum_{m,n} |\text{tr}(\Pi_e B \Pi_{e'} \rho P_m) \text{tr}(\Pi_{e'} B^* \Pi_e \rho^3 P_n)| \leq \|B\|^2 D_E \|\rho\|^2. \quad (94)$$

Finally we estimate

$$\sum_{e,e'} |\text{tr}(\rho \Pi_e B \Pi_{e'})|^2 \leq \sum_{e,e'} \text{tr}(\Pi_e B \Pi_{e'} B^*) \text{tr}(\Pi_e \rho \Pi_{e'} \rho) \quad (95a)$$

$$\leq \|B\|^2 D_E \sum_{e,e'} \text{tr}(\Pi_e \rho \Pi_{e'} \rho) \quad (95b)$$

$$= \|B\|^2 D_E \text{tr} \rho^2 \quad (95c)$$

$$\leq \|B\|^2 D_E \|\rho\|. \quad (95d)$$

Putting everything together we arrive at

$$\begin{aligned} & \mathbb{E}_\rho \left( \left\langle \left| \langle \psi_t | B | \psi_t \rangle - \overline{\langle \psi_t | B | \psi_t \rangle} \right|^2 \right\rangle_T \right) \\ & \leq \|B\|^2 \|\rho\| D_E G(\kappa) \left( 1 + \frac{8 \log_2 d_E}{\kappa T} \right) \left( 1 + \frac{1}{1 - \|\rho\|} \left[ 1 + \frac{2\|\rho\|}{1 - 2\|\rho\|} + \frac{6(\|\rho\| + \|\rho\|^2)}{(1 - 2\|\rho\|)(1 - 3\|\rho\|)} \right] \right) \end{aligned} \quad (96a)$$

$$\leq 24 \|B\|^2 \|\rho\| D_E G(\kappa) \left( 1 + \frac{8 \log_2 d_E}{\kappa T} \right), \quad (96b)$$

where we used the assumption that  $\|\rho\| < 1/4$  to bound the bracket  $(1 + \frac{1}{1-\|\rho\|}[\dots])$  in (96a) by<sup>4</sup> 24.

---

<sup>4</sup>Note that the bracket is a monotone increasing function in  $\|\rho\|$  and therefore bounded by its value for  $\|\rho\| = 1/4$  which is  $71/3 < 24$ .

For the variance of  $\overline{\langle \psi_t | B | \psi_t \rangle}$  we see from (72c) that we can simply apply Lemma 1 with  $A = \sum_e \Pi_e B \Pi_e$  and estimate the occurring traces. We compute

$$\operatorname{tr} \left( \left( \sum_e \Pi_e B \Pi_e \right) \rho \left( \sum_{e'} \Pi_{e'} B^* \Pi_{e'} \right) \rho \right) \leq \left\| \left( \sum_e \Pi_e B \Pi_e \right) \rho \left( \sum_{e'} \Pi_{e'} B^* \Pi_{e'} \right) \right\| \operatorname{tr} \rho \quad (97a)$$

$$\leq \|B\|^2 \|\rho\|, \quad (97b)$$

where we used that  $\|\sum_e \Pi_e B \Pi_e\| \leq \|B\|$ . Similarly we find that

$$\operatorname{tr}(A \rho^2 A^* \rho), \operatorname{tr}(A \rho A^* \rho^2) \leq \|B\|^2 \|\rho\|^2, \quad (98)$$

$$\operatorname{tr}(A \rho^3 A^* \rho), \operatorname{tr}(A \rho^2 A^* \rho^2), \operatorname{tr}(A \rho A^* \rho^3) \leq \|B\|^2 \|\rho\|^3, \quad (99)$$

Moreover, we get

$$\sum_{m,n} \left| \operatorname{tr} \left( \sum_e \Pi_e B \Pi_e \rho^3 P_m \right) \operatorname{tr} \left( \sum_{e'} \Pi_{e'} B^* \Pi_{e'} \rho P_n \right) \right| \leq \sum_{m,n} p_m^3 p_n \|B\|^2 \operatorname{tr}(P_m) \operatorname{tr}(P_n) \quad (100a)$$

$$= \|B\|^2 \operatorname{tr} \rho^3 \quad (100b)$$

$$\leq \|B\|^2 \|\rho\|^2 \quad (100c)$$

and similarly

$$\sum_{m,n} |\operatorname{tr}(A \rho^2 P_m) \operatorname{tr}(A^* \rho^2 P_n)|, \sum_{m,n} |\operatorname{tr}(A \rho P_m) \operatorname{tr}(A^* \rho^3 P_n)| \leq \|B\|^2 \|\rho\|^2. \quad (101)$$

Thus we finally arrive at

$$\operatorname{Var}_\rho \overline{\langle \psi_t | B | \psi_t \rangle} \leq \frac{\|B\|^2 \|\rho\|}{1 - \|\rho\|} \left( 1 + \frac{2\|\rho\|}{1 - 2\|\rho\|} + \frac{6(\|\rho\| + \|\rho\|^2)}{(1 - 2\|\rho\|)(1 - 3\|\rho\|)} \right) \quad (102a)$$

$$\leq 23 \|B\|^2 \|\rho\|. \quad (102b)$$

Now suppose that  $\mathcal{H}$  is infinite-dimensional. Due to our assumption on  $B$ , all sums over  $e \in \mathcal{E}$  are effectively sums over  $e \in \mathcal{E}_B$  and therefore finite. In the finite-dimensional setting, the matrix  $R$  defined by (78) is a  $d_E(d_E - 1) \times d_E(d_E - 1)$  matrix and this is where the  $d_E$  in the upper bound for  $\|R\|$  comes from, see the proof in [36]. In the infinite-dimensional setting, the matrix  $R$  can be viewed as a  $d_{E,B}(d_{E,B} - 1) \times d_{E,B}(d_{E,B} - 1)$  matrix and therefore, in the bound for  $\|R\|$ ,  $d_E$  has to be replaced by  $d_{E,B}$ ; by the same argument,  $d_E$  can also be replaced by  $d_{E,B}$  in the finite-dimensional setting. Moreover, we can also change  $G(\kappa)$  to  $G_B(\kappa)$  and  $D_E$  to  $D_{E,B}$  as the only ‘‘contributing’’ eigenvalues are the ones in  $\mathcal{E}_B$ . Besides this change all other steps of the proof remain valid and this proves the bounds also in the infinite-dimensional case.

For the statements about the infinite time average, we want to take the limit  $T \rightarrow \infty$  in

$$\mathbb{E}_\rho \left\langle \left| \langle \psi_t | B | \psi_t \rangle - \overline{\langle \psi_t | B | \psi_t \rangle} \right|^2 \right\rangle_T \leq 24 \|B\|^2 \|\rho\| D_{E,B} G_B(\kappa) \left( 1 + \frac{8 \log_2 d_{E,B}}{\kappa T} \right). \quad (103)$$

To this end, we first choose  $\kappa$  so small that  $G_B(\kappa) = D_{G,B}$ . Then we take the limit  $T \rightarrow \infty$  and use dominated convergence to interchange the limit and  $\mathbb{E}_\rho$ . This gives (76) and finishes the proof.  $\square$

### 4.3 Proof of Theorem 1

Without loss of generality we assume that all eigenvalues of  $\rho$  are positive (otherwise we restrict our considerations to the subspace of  $\mathcal{H}$  spanned by the eigenvectors of  $\rho$  corresponding to its positive eigenvalues). It follows from Markov's inequality together with Proposition 1 that

$$\text{GAP}(\rho) \left\{ \left\langle \left| \langle \psi_t | B | \psi_t \rangle - \overline{\langle \psi_t | B | \psi_t \rangle} \right|^2 \right\rangle_T \geq \frac{48}{\varepsilon} \|B\|^2 \|\rho\| D_{E,B} G_B(\kappa) \left( 1 + \frac{8 \log_2 d_{E,B}}{\kappa T} \right) \right\} \leq \frac{\varepsilon}{2} \quad (104)$$

and

$$\text{GAP}(\rho) \left\{ \overline{\left| \langle \psi_t | B | \psi_t \rangle - \overline{\langle \psi_t | B | \psi_t \rangle} \right|^2} \geq \frac{48}{\varepsilon} \|B\|^2 \|\rho\| D_{E,B} D_{G,B} \right\} \leq \frac{\varepsilon}{2}. \quad (105)$$

Concerning the finite-time average, this shows that, w.r.t.  $\text{GAP}(\rho)$ ,  $(1 - \frac{\varepsilon}{2})$ -most  $\psi_0 \in \mathbb{S}(\mathcal{H})$  are such that

$$\left\langle \left| \langle \psi_t | B | \psi_t \rangle - \overline{\langle \psi_t | B | \psi_t \rangle} \right|^2 \right\rangle_T \leq \frac{48}{\varepsilon} \|B\|^2 \|\rho\| D_{E,B} G_B(\kappa) \left( 1 + \frac{8 \log_2 d_{E,B}}{\kappa T} \right). \quad (106)$$

Similarly, we have that, w.r.t.  $\text{GAP}(\rho)$ ,  $(1 - \frac{\varepsilon}{2})$ -most  $\psi_0 \in \mathbb{S}(\mathcal{H})$  are such that

$$\overline{\left| \langle \psi_t | B | \psi_t \rangle - \overline{\langle \psi_t | B | \psi_t \rangle} \right|^2} \leq \frac{48}{\varepsilon} \|B\|^2 \|\rho\| D_{E,B} D_{G,B}. \quad (107)$$

Let  $\lambda$  denote the Lebesgue measure on  $\mathbb{R}$ . An application of Markov's inequality and Proposition 1 shows that

$$\frac{1}{T} \lambda \left\{ t \in [0, T] : \left| \langle \psi_t | B | \psi_t \rangle - \overline{\langle \psi_t | B | \psi_t \rangle} \right|^2 \geq \frac{48}{\varepsilon \delta} \|B\|^2 \|\rho\| D_{E,B} G_B(\kappa) \left( 1 + \frac{8 \log_2 d_{E,B}}{\kappa T} \right) \right\} \leq \delta. \quad (108)$$

Thus, w.r.t.  $\text{GAP}(\rho)$ ,  $(1 - \frac{\varepsilon}{2})$ -most  $\psi_0 \in \mathbb{S}(\mathcal{H})$  are such that for  $(1 - \delta)$ -most  $t \in [0, T]$ ,

$$\left| \langle \psi_t | B | \psi_t \rangle - \overline{\langle \psi_t | B | \psi_t \rangle} \right| \leq \left( \frac{48}{\varepsilon \delta} \|B\|^2 \|\rho\| D_{E,B} G_B(\kappa) \left( 1 + \frac{8 \log_2 d_{E,B}}{\kappa T} \right) \right)^{1/2}. \quad (109)$$

In the same way, we obtain

$$\liminf_{T \rightarrow \infty} \frac{1}{T} \lambda \left\{ t \in [0, T] : \left| \langle \psi_t | B | \psi_t \rangle - \overline{\langle \psi_t | B | \psi_t \rangle} \right|^2 \geq \frac{48}{\varepsilon \delta} \|B\|^2 \|\rho\| D_{E,B} D_{G,B} \right\} \leq \delta \quad (110)$$

and conclude that, w.r.t.  $\text{GAP}(\rho)$ ,  $(1 - \frac{\varepsilon}{2})$ -most  $\psi_0 \in \mathbb{S}(\mathcal{H})$  are such that for  $(1 - \delta)$ -most  $t \in [0, \infty)$ ,

$$\left| \langle \psi_t | B | \psi_t \rangle - \overline{\langle \psi_t | B | \psi_t \rangle} \right| \leq \left( \frac{48}{\varepsilon \delta} \|B\|^2 \|\rho\| D_{E,B} D_{G,B} \right)^{1/2}. \quad (111)$$

It follows from Chebyshev's inequality together with Proposition 1 that

$$\text{GAP}(\rho) \left\{ \left| \overline{\langle \psi_t | B | \psi_t \rangle} - M_{\rho B} \right| \geq \left( \frac{46 \|B\|^2 \|\rho\|}{\varepsilon} \right)^{1/2} \right\} \leq \frac{\varepsilon}{2} \quad (112)$$

and thus, w.r.t.  $\text{GAP}(\rho)$ ,  $(1 - \frac{\varepsilon}{2})$ -most  $\psi_0 \in \mathbb{S}(\mathcal{H})$  are such that

$$\left| \overline{\langle \psi_t | B | \psi_t \rangle} - M_{\rho B} \right| \leq \left( \frac{46 \|B\|^2 \|\rho\|}{\varepsilon} \right)^{1/2}. \quad (113)$$

Now the claims follow from the triangle inequality.

## 5 Summary and Conclusions

Our result concerns the long-time behavior of  $\text{GAP}(\rho)$ -typical pure states  $\psi_0$  from the sphere of a separable Hilbert space where  $\rho$  is a density matrix on  $\mathcal{H}$  and  $\psi_0$  evolves unitarily according to  $\psi_t = \exp(-iHt)\psi_0$ . We have seen that for any operator  $B$  on  $\mathcal{H}$  that satisfies  $|\mathcal{E}_B| < \infty$ , for  $\text{GAP}(\rho)$ -most  $\psi_0 \in \mathbb{S}(\mathcal{H})$ , the curve  $t \mapsto \langle \psi_t | B | \psi_t \rangle$  is nearly constant in the long run  $t \rightarrow \infty$  provided that  $\|\rho\|$  is small,  $\|B\|$  is not too large and no eigenvalue or eigenvalue gap of the Hamiltonian  $H$  is too highly degenerate. In particular, we have argued that our result can be applied to the case that  $B = P_\nu$  for some macro state  $\nu$ . We have provided explicit error bounds that reveal the dependence of the error on these quantities. Our result shows that the concept of normal typicality can be generalized from the uniform distribution on the sphere of the Hilbert space to a much broader class of distributions which can also be defined on infinite-dimensional Hilbert spaces and which for certain density matrices arise naturally as the distribution of wave functions in thermal equilibrium.

**Acknowledgments.** It is a pleasure to thank Stefan Teufel and Roderich Tumulka for their helpful discussions. Financial support from the German Academic Scholarship Foundation is gratefully acknowledged.

**Data Availability Statement.** Data sharing is not applicable to this article as no datasets were generated or analyzed.

**Conflict of Interest Statement.** The author has no conflicts of interest.



## References

- [1] B. Balz, J. Richter, J. Gemmer, R. Steinigeweg, and P. Reimann. Dynamical typicality for initial states with a preset measurement statistics of several commuting observables. In F. Binger, L. A. Correa, C. Gogolin, J. Anders, and G. Adesso, editors, *Thermodynamics in the Quantum Regime*, chapter 17, pages 413–433. Springer, Cham, 2019. URL: <http://arxiv.org/abs/1904.03105>.
- [2] C. Bartsch and J. Gemmer. Dynamical typicality of quantum expectation values. *Physical Review Letters*, 102:110403, 2009. URL: <http://arxiv.org/abs/0902.0927>.
- [3] E. Borel. *Probabilities and Life*. Dover, 1962.
- [4] G. Cipolloni, L. Erdős, and J. Henheik. Eigenstate thermalisation at the edge for Wigner matrices, 2023. Preprint. URL: <https://arxiv.org/abs/2309.05488>.
- [5] G. Cipolloni, L. Erdős, and D. Schröder. Eigenstate Thermalization Hypothesis for Wigner Matrices. *Communications in Mathematical Physics*, 388:1005–1048, 2021. URL: <http://arxiv.org/abs/2012.13215>.
- [6] J. M. Deutsch. Quantum statistical mechanics in a closed system. *Physical Review A*, 43:2046–2049, 1991.
- [7] L. Erdős and V. Riabov. Eigenstate Thermalization Hypothesis for Wigner-Type Matrices. *Communications in Mathematical Physics*, 405(282), 2024. URL: <https://arxiv.org/abs/2403.10359>.
- [8] J. Gemmer and G. Mahler. Distribution of local entropy in the Hilbert space of bi-partite quantum systems: origin of Jaynes’ principle. *The European Physical Journal B*, 31:249–257, 2003. URL: <https://arxiv.org/abs/quant-ph/0201136>.
- [9] J. Gemmer, G. Mahler, and M. Michel. *Quantum Thermodynamics*. Springer, 2004.
- [10] S. Goldstein, T. Hara, and H. Tasaki. Time Scales in the Approach to Equilibrium of Macroscopic Quantum Systems. *Physical Review Letters*, 111:140401, 2013. URL: <http://arxiv.org/abs/1307.0572>.
- [11] S. Goldstein, T. Hara, and H. Tasaki. The approach to equilibrium in a macroscopic quantum system for a typical nonequilibrium subspace, 2014. Preprint, <http://arxiv.org/abs/1402.3380>.
- [12] S. Goldstein, T. Hara, and H. Tasaki. Extremely quick thermalization in a macroscopic quantum system for a typical nonequilibrium subspace. *New Journal of Physics*, 17:045002, 2015. URL: <http://arxiv.org/abs/1402.0324>.

- [13] S. Goldstein, D. A. Huse, J. L. Lebowitz, and R. Tumulka. Thermal equilibrium of a macroscopic quantum system in a pure state. *Physical Review Letters*, 115:100402, 2015. URL: <http://arxiv.org/abs/1506.07494>.
- [14] S. Goldstein, D. A. Huse, J. L. Lebowitz, and R. Tumulka. Macroscopic and microscopic thermal equilibrium. *Annalen der Physik*, 529:1600301, 2017. URL: <http://arxiv.org/abs/1610.02312>.
- [15] S. Goldstein, J. L. Lebowitz, C. Mastrodonato, R. Tumulka, and N. Zanghì. Approach to Thermal Equilibrium of Macroscopic Quantum Systems. *Physical Review E*, 81:011109, 2010. URL: <http://arxiv.org/abs/0911.1724>.
- [16] S. Goldstein, J. L. Lebowitz, C. Mastrodonato, R. Tumulka, and N. Zanghì. Normal Typicality and von Neumann’s Quantum Ergodic Theorem. *Proceedings of the Royal Society A*, 466(2123):3203–3224, 2010. URL: <http://arxiv.org/abs/0907.0108>.
- [17] S. Goldstein, J. L. Lebowitz, R. Tumulka, and N. Zanghì. Canonical Typicality. *Physical Review Letters*, 96:050403, 2006. URL: <http://arxiv.org/abs/cond-mat/0511091>.
- [18] S. Goldstein, J. L. Lebowitz, R. Tumulka, and N. Zanghì. Long-Time Behavior of Macroscopic Quantum Systems. *European Physical Journal H*, 35:173–200, 2010. URL: <http://arxiv.org/abs/1003.2129>.
- [19] S. Goldstein, J. L. Lebowitz, R. Tumulka, and N. Zanghì. Gibbs and Boltzmann entropy in classical and quantum mechanics. In *Statistical Mechanics and Scientific Explanation: Determinism, Indeterminism and Laws of Nature*, pages 519–581. World Scientific, 2020. URL: <http://arxiv.org/abs/1903.11870>.
- [20] S. Goldstein, J.L. Lebowitz, C. Mastrodonato, R. Tumulka, and N. Zanghì. Universal Probability Distribution for the Wave Function of a Quantum System Entangled with its Environment. *Communications in Mathematical Physics*, 342:965–988, 2016. URL: <https://arxiv.org/abs/1104.5482>.
- [21] S. Goldstein, J.L. Lebowitz, R. Tumulka, and N. Zanghì. On the Distribution of the Wave Function for Systems in Thermal Equilibrium. *Journal of Statistical Physics*, 125(5/6), 2006. URL: <https://arxiv.org/abs/quant-ph/0309021>.
- [22] R. Jozsa, D. Robb, and W.K. Wootters. Lower bound for accessible information in quantum mechanics. *Physical Review A*, 49(2):668, 1994.
- [23] N. Linden, S. Popescu, A.J. Short, and A. Winter. Quantum mechanical evolution towards thermal equilibrium. *Physical Review E*, 79:061103, 2009. URL: <http://arxiv.org/abs/0812.2385>.

- [24] S. Lloyd. *Black Holes, Demons, and the Loss of Coherence*. Ph.d. thesis, Department of Physics, Rockefeller University, 1988.
- [25] T. Mori, T.N. Ikeda, E. Kaminishi, and M. Ueda. Thermalization and prethermalization in isolated quantum systems: a theoretical overview. *Journal of Physics B*, 51:112001, 2018. URL: <http://arxiv.org/abs/1712.08790>.
- [26] M. P. Müller, D. Gross, and J. Eisert. Concentration of measure for quantum states with a fixed expectation value. *Communications in Mathematical Physics*, 303:785–824, 2011. URL: <http://arxiv.org/abs/1003.4982>.
- [27] S. Popescu, A. J. Short, and A. Winter. Entanglement and the foundation of statistical mechanics. *Nature Physics*, 21(11):754–758, 2006. URL: <http://arxiv.org/abs/quant-ph/0511225>.
- [28] S. Popescu, A.J. Short, and A. Winter. The foundations of statistical mechanics from entanglement: Individual states vs. averages, 2005. Preprint. URL: <http://arxiv.org/abs/quant-ph/0511225>.
- [29] P. Reimann. Typicality of Pure States Randomly Sampled According to the Gaussian Adjusted Projected Measure. *Journal of Statistical Physics*, 132:921–925, 2008. URL: <https://arxiv.org/abs/0805.3102>.
- [30] P. Reimann. Generalization of von Neumann’s approach to thermalization. *Physical Review Letters*, 115:010403, 2015. URL: <http://arxiv.org/abs/1507.00262>.
- [31] P. Reimann. Dynamical typicality approach to eigenstate thermalization. *Physical Review Letters*, 120:230601, 2018. URL: <http://arxiv.org/abs/1806.03193>.
- [32] P. Reimann. Dynamical typicality of isolated many-body quantum systems. *Physical Review E*, 97:062129, 2018. URL: <http://arxiv.org/abs/1805.07085>.
- [33] P. Reimann and J. Gemmer. Why are macroscopic experiments reproducible? Imitating the behavior of an ensemble by single pure states. *Physica A*, 552:121840, 2020. URL: <http://arxiv.org/abs/2005.14626>.
- [34] B. Roos, S. Teufel, R. Tumulka, and C. Vogel. Macroscopic Thermalization for Highly Degenerate Hamiltonians, 2024. Preprint. URL: <http://www.arxiv.org/abs/2408.15832>.
- [35] N. Shiraishi and H. Tasaki. Nature Abhors a Vacuum: A Simple Rigorous Example of Thermalization in an Isolated Macroscopic Quantum System. *Journal of Statistical Physics*, 191:82, 2024. URL: <http://arxiv.org/abs/2310.18880>.
- [36] A. J. Short and T. C. Farrelly. Quantum equilibration in finite time. *New Journal of Physics*, 14:013063, 2012.

- [37] B. Simon. *Operator Theory: A Comprehensive Course in Analysis, Part 4*. American Mathematical Society, 2015.
- [38] M. Srednicki. Chaos and quantum thermalization. *Physical Review E*, 50:888–901, 1994. URL: <http://arxiv.org/abs/cond-mat/9403051>.
- [39] H. Tasaki. Typicality of thermal equilibrium and thermalization in isolated macroscopic quantum systems. *Journal of Statistical Physics*, 163:937–997, 2016. URL: <http://arxiv.org/abs/1507.06479>.
- [40] H. Tasaki. Heat flows from hot to cold: A simple rigorous example of thermalization in an isolated macroscopic quantum system, 2024. Preprint. URL: <http://arxiv.org/abs/2404.04533>.
- [41] H. Tasaki. Macroscopic Irreversibility in Quantum Systems: ETH and Equilibration in a Free Fermion Chain, 2024. Preprint. URL: <http://arxiv.org/abs/2401.15263>.
- [42] S. Teufel, R. Tumulka, and C. Vogel. Time Evolution of Typical Pure States from a Macroscopic Hilbert Subspace. *Journal of Statistical Physics*, 190:69, 2023. URL: <http://arxiv.org/abs/2210.10018>.
- [43] S. Teufel, R. Tumulka, and C. Vogel. Typical Macroscopic Long-Time Behavior for Random Hamiltonians, 2023. Preprint. URL: <https://arxiv.org/abs/2303.13242>.
- [44] S. Teufel, R. Tumulka, and C. Vogel. Canonical Typicality for Other Ensembles than Micro-canonical. *Annales Henri Poincaré*, 2024. URL: <https://arxiv.org/abs/2307.15624>.
- [45] R. Tumulka. Thermal Equilibrium Distribution in Infinite-Dimensional Hilbert Spaces. *Reports on Mathematical Physics*, 86:303–313, 2020. URL: <http://arxiv.org/abs/2004.14226>.
- [46] J. von Neumann. Beweis des Ergodensatzes und des  $H$ -Theorems in der neuen Mechanik. *Zeitschrift für Physik*, 57:30–70, 1929. English translation: *European Physical Journal H*, 35: 201–237, 2010. URL: <http://arxiv.org/abs/1003.2133>.