THE TANNAKIAN RADICAL AND THE MANTLE OF A BRAIDED FUSION CATEGORY

JASON GREEN AND DMITRI NIKSHYCH

ABSTRACT. We define the Tannakian radical of a braided fusion category $\mathcal C$ as the intersection of its maximal Tannakian subcategories. The localization of $\mathcal C$ corresponding to the Tannakian radical, termed the mantle of \mathcal{C} , admits a canonical central extension that serves as a complete invariant of C. The mantle has a trivial Tannakian radical, and we refer to braided fusion categories with this property as reductive. We investigate the properties and structure of reductive categories and prove several classification results.

CONTENTS

7. Reductive p[-categories of small dimension](#page-26-0) 27 [References](#page-29-0) 30

1. INTRODUCTION

Let β be a braided fusion category. There is a well-known procedure of localizing β with respect to an étale algebra (i.e., a commutative separable algebra) A in \mathcal{B} , consisting of taking the category \mathcal{B}_{A}^{loc} of local A-modules in \mathcal{B} , see, e.g., [\[DMNO\]](#page-29-1). It produces a "smaller" braided fusion category in the Witt class of \mathcal{B} . In the special case where A is the regular algebra in a Tannakian subcategory $\mathcal{E} \subset \mathcal{B}$ this procedure coincides with the de-equivariantization [\[DGNO2\]](#page-29-2): $\mathcal{B}_{A}^{loc} \cong \mathcal{E}' \boxtimes_{\mathcal{E}} \mathcal{V}ect$, where \mathcal{E}' is the centralizer of \mathcal{E} . When \mathcal{E} is maximal among Tannakian subcategories of B, this localization is an invariant of B, called the *core*. It was explained in $[~~DGNO2~~ how the core can be used to extract the part of $\mathcal C$ that does not$ come from finite groups. This invariant turned out to be very useful in the classification of fusion categories. For example, non-degenerate braided fusion categories with a trivial core are precisely representation categories of twisted Drinfeld doubles [\[DGNO2\]](#page-29-2). In addition, Natale characterized weakly group-theoretical braided fusion categories in terms of their cores [\[Na\]](#page-30-0).

The category β can be recovered from its core by means of a gauging procedure [\[CGPW,](#page-29-3) [ENO2\]](#page-29-4) with respect to a group G such that $\mathcal{E} = \mathcal{R}ep(G) \subset \mathcal{B}$ is a maximal Tannakian subcategory. However, unlike the core, the isomorphism class of G is not an invariant of \mathcal{B} , see Remark [2.2.](#page-4-1) Thus, it is natural to replace $\mathcal E$ with an invariant Tannakian subcategory of β and investigate the corresponding localization. This is what we do in this paper, whose main results and organization are outlined below.

Sections [2](#page-2-0) and [3](#page-6-0) contain necessary results about gauging and localization of braided fusion categories and their maximal Tannakian subcategories.

We define the *Tannakian radical* $\mathcal{R}ad(\mathcal{B})$ of $\mathcal B$ as the intersection of all maximal Tannakian subcategories of \mathcal{B} (Section [4\)](#page-12-0). We say that \mathcal{B} is *reductive*, if $\mathcal{R}ad(\mathcal{B})$ is trivial, i.e., $\mathcal{R}ad(\mathcal{B})$ Vect. This is a new class of braided fusion categories with interesting properties. For example, $\mathcal{R}ep(D(G))$, the representation category of the Drinfeld double of a finite group G, is reductive if and only if G is generated by its normal Abelian subgroups, see Remark [4.8](#page-13-1) (so, in particular, G must be nilpotent in this case).

We introduce the *mantle* of $\mathcal B$ as its localization with respect to the radical: $\mathcal M$ antle $(\mathcal B)$ = $Rad(\mathcal{B})' \boxtimes_{Rad(\mathcal{B})} Vect.$ We show that the mantle is reductive (Corollary [4.15\)](#page-15-1). Clearly, the braided equivalence classes of $Rad(\mathcal{B})$ and $Mantle(\mathcal{B})$ are invariants of \mathcal{B} . Furthermore, there is a group $G_{\mathcal{B}}$, determined up to an isomorphism, such that $\mathcal{R}ad(\mathcal{B}) = \mathcal{R}ep(G_{\mathcal{B}})$ and

a canonical monoidal 2-functor $F_{\mathcal{B}} : G_{\mathcal{B}} \to Pic(\mathcal{M}antle(\mathcal{B}))$ such that \mathcal{B} is equivalent to the equivariantization of the corresponding central extension of $\mathcal{M}antle(\mathcal{B})$. In other words, **B** is a canonical G_B-gauging of its mantle. These data $(G_B, \text{Mantle}(\mathcal{B}), F_B)$ constitute a complete invariant of β (Section [4.3\)](#page-15-0). Since the center $\mathcal{Z}(\mathcal{A})$ of a fusion category \mathcal{A} is a complete invariant of its Morita equivalence class, by considering the radical and mantle of $\mathcal{Z}(\mathcal{A})$ we obtain new Morita invariants of \mathcal{A} (Corollary [4.19\)](#page-18-2).

We study the structure of reductive categories in detail in Section [5.](#page-18-0) We define a *coradical* of a braided fusion category as the subcategory generated by all its Tannakian subcategories. The coradical is a group-theoretical category (Proposition [5.2\)](#page-18-3). We classify reductive categories into three types based on the non-degeneracy properties of the categories themselves and their coradicals, see Definition [5.18.](#page-21-1) In Theorem [5.20](#page-21-2) we provide a decomposition of a reductive braided fusion category of each type in terms of its coradical and a certain anisotropic subcategory.

Pointed reductive braided fusion categories (i.e., pre-metric groups) are studied in Section [6.](#page-22-0) Their classification is given in Theorems [6.2](#page-23-1) and [6.9.](#page-25-0)

Finally, in Section [7](#page-26-0) we deal with reductive p-categories (i.e., categories whose Frobenius-Perron dimension is a power of a prime integer p). A motivation for this comes from Proposition [4.11](#page-14-0) which says that a reductive braided fusion category is weakly group-theoretical if and only if it is nilpotent, and so it is a direct product of p-categories by [\[DGNO1\]](#page-29-5). We show that all such categories of dimension $\leq p^5$ are pointed and classify non-pointed reductive *p*-categories of dimension p^6 in Proposition [7.6.](#page-28-0)

Acknowledgments. We are grateful to Alexei Davydov and Pavel Etingof for helpful discussions. The work of the second author was supported by the National Science Foundation under Grant No. DMS-2302267. This material is based upon work supported by the National Science Foundation under Grant No. DMS-1928930, while the second author was in residence at SLMath (the Mathematical Sciences Research Institute) in Berkeley, California, during the Summer of 2024.

2. Preliminaries

Throughout this paper we work over an algebraically closed field k of characteristic 0. Tensor and module categories are assumed to be finite semisimple and k-linear.

We refer the reader to [\[EGNO,](#page-29-6) [DGNO1\]](#page-29-5) for the theory of braided fusion categories and to [\[DN2,](#page-29-7) [ENO1\]](#page-29-8) for the theory of graded extensions.

2.1. Gauging and localization of braided fusion categories. Let β be a braided fusion category. Let $Pic(\mathcal{B})$ denote the 2-categorical Picard group of \mathcal{B} , consisting of invertible \mathcal{B} module categories [\[ENO2\]](#page-29-4).

Let G be a finite group. A *central G-extension* $\mathcal{C} = \bigoplus_{x \in G} \mathcal{B}_x$ of $\mathcal{B}_1 = \mathcal{B}$ (which is the same thing as a G-crossed extension [\[DN2,](#page-29-7) [JMPP\]](#page-29-9)) is a faithful G-graded extension along with an embedding $\mathcal{B} \hookrightarrow \mathcal{Z}(\mathcal{C})$ whose composition with the forgetful functor $\mathcal{Z}(\mathcal{C}) \to \mathcal{C}$ coincides with the embedding $\mathcal{B} \hookrightarrow \mathcal{C}$. It was shown in [\[ENO2,](#page-29-4) [DN2\]](#page-29-7) that the 2-groupoid of central extensions of B is 2-equivalent to the 2-groupoid of monoidal 2-functors $G \to \text{Pic}(\mathcal{B})$. A central extension has a canonical action of G such that $g(\mathcal{B}_x) = \mathcal{B}_{gxg^{-1}}$, $g, x \in G$. It restricts to an action of G on β by braided autoequivalences. The latter coincides with the composition of the corresponding functor $G \to Pic(B)$ with the canonical monoidal functor $Pic(\mathcal{B}) \rightarrow Aut^{br}(\mathcal{B}).$

The corresponding equivariantization

$$
\mathcal{G}(G,\mathcal{B},F):=\left(\bigoplus_{x\in G}\,\mathcal{B}_x\right)^G
$$

is called a G-*gauging* (or simply *gauging*) of B. It is a braided fusion category containing a Tannakian subcategory $\mathcal{R}ep(G)$. This category is non-degenerate if and only if $\mathcal B$ is nondegenerate.

Conversely, to a non-degenerate braided fusion category $\mathcal C$ containing a Tannakian subcategory $\mathcal{E} \cong \mathcal{R}ep(G)$, one associates a braided fusion category $\mathcal{B} = \mathcal{E}' \boxtimes_{\mathcal{E}} \mathcal{V}ect$, where \mathcal{E}' is the centralizer of $\mathcal E$ in $\mathcal C$. This category $\mathcal B$ is called a *localization* of $\mathcal C$ (with respect to $\mathcal E$). There is a canonical monoidal 2-functor $F: G \to Pic(B)$ such that C is canonically equivalent to $\mathcal{G}(G,\mathcal{B},F).$

2.2. Maximal Tannakian subcategories and the core of a braided fusion category. Let B be a braided fusion category. Let $T(\mathcal{B})$, $T_{max}(\mathcal{B})$ denote the sets of all Tannakian and maximal Tannakian subcategories of \mathcal{B} , respectively. If G is a group acting on \mathcal{B} by braided autoequivalenves, let $T^G(\mathcal{B})$, $T_{max}^G(\mathcal{B})$ denote the sets of all G-stable Tannakian and

A Tannakian subcategory E of a non-degenerate braided fusion category B is called *Lagrangian* if $\mathcal{E} = \mathcal{E}'$, where \mathcal{E}' is the centralizer of \mathcal{E} in \mathcal{B} . Lagrangian subcategories of \mathcal{B} are in bijection with isomorphism classes of braided equivalences between β and centers of pointed fusion categories [\[DGNO2\]](#page-29-2). These centers are precisely the representation categories of twisted Drinfeld doubles of finite groups, $\mathcal{R}ep(D^{\omega}(G)) \cong \mathcal{Z}(\mathcal{V}ect^{\omega}_G)$, where G is a finite group and ω is a 3-cocycle on G. Let $L(\mathcal{B})$ denote the set of Lagrangian subcategories of \mathcal{B} .

Definition 2.1. A non-degenerate category \mathcal{B} with $L(\mathcal{B}) \neq \emptyset$ is called *metabolic*.

maximal G-stable Tannakian subcategories of \mathcal{B} , respectively.

Since a Lagrangian subcategory is necessarily maximal, for metabolic categories one has $L(\mathcal{B})=T_{max}(\mathcal{B}).$

It was shown in [\[DGNO2\]](#page-29-2) that for any $\mathcal{E} = \mathcal{R}ep(G_{\mathcal{E}}) \in T_{max}(\mathcal{B})$ the braided equivalence class of the corresponding localization $\mathcal{E}' \boxtimes_{\mathcal{E}} \mathcal{V}$ ect does not depend on the choice of \mathcal{E} . This category is called the *core* of \mathcal{B} and is denoted $\mathcal{C}ore(\mathcal{B})$. Furthermore, the image of $G_{\mathcal{E}}$ in $Aut^{br}(Core(\mathcal{B}))$ is also independent of the choice of $\mathcal{E}.$

Remark 2.2. We saw that \mathcal{B} can be realized as a gauging of a canonical category $\text{Core}(\mathcal{B})$. But this gauging is not canonical, since the above group $G_{\mathcal{E}}$ is not an invariant of \mathcal{B} . For example, $\mathcal{R}ep(D(\mathbb{Z}/4\mathbb{Z}))$ contains maximal Tannakian subcategories $\mathcal{R}ep(\mathbb{Z}/4\mathbb{Z})$ and $\mathcal{R}ep(\mathbb{Z}/2\mathbb{Z}\times\mathbb{Z}/2\mathbb{Z})$ and so it is both a $\mathbb{Z}/4\mathbb{Z}$ - and $\mathbb{Z}/2\mathbb{Z}\times\mathbb{Z}/2\mathbb{Z}$ -gauging of the trivial category Vect. In other words, $\mathcal{R}ep(D(\mathbb{Z}/4\mathbb{Z})) \cong \mathcal{R}ep(D^{\omega}(\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}))$ for some 3-cocycle ω on $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$.

2.3. Metric groups. Recall that a *pre-metric group* (A, q) is a finite Abelian group A equipped with a quadratic form $q: A \to k^{\times}$. There is an equivalence between the groupoid of pre-metric groups and that of pointed braided fusion categories

$$
(A, q) \mapsto \mathcal{C}(A, q),
$$

where A is identified with the group of invertible objects of $\mathcal{C}(A, q)$ and the scalar $q(x)$ is the value of the self-braiding of $x \in A$.

A pre-metric group (A, q) is called a *metric group* if the quadratic form q is non-degenerate. This is equivalent to $\mathcal{C}(A, q)$ being a non-degenerate braided fusion category.

A pre-metric group (A, q) is *anisotropic* if $q(x) \neq 1$ for any $x \neq 0$.

For any prime p, let M_p be the monoid of metric p-groups. This monoid was described in [\[W\]](#page-30-1) and [\[KK\]](#page-29-10) by generators and relations as follows.

Suppose that p is odd. For any p^m -th root of unity $\zeta \in k^{\times}$ define a quadratic form

(1)
$$
q_{\zeta} : \mathbb{Z}/p^m \mathbb{Z} \to k^{\times}, \qquad q_{\zeta}(x) = \zeta^{x^2}.
$$

This form is non-degenerate if and only if ζ is a primitive p^m -th root of 1. When η is another such root, the metric groups $(\mathbb{Z}/p^m\mathbb{Z}, q_\zeta)$ and $(\mathbb{Z}/p^m\mathbb{Z}, q_\eta)$ are isomorphic if and only if $\eta = \zeta^{a^2}$ for some a relatively prime to p.

Fix a quadratic non-residue b modulo p^m and let $\bar{\zeta} = \zeta^b$. The isomorphism classes of metric groups $(\mathbb{Z}/p^m\mathbb{Z}, q_{\zeta})$ and $(\mathbb{Z}/p^m\mathbb{Z}, q_{\bar{\zeta}}), m \geq 1$, generate M_p . These generators obey the relations

(2)
$$
(\mathbb{Z}/p^m\mathbb{Z}, q_\zeta) \oplus (\mathbb{Z}/p^m\mathbb{Z}, q_\zeta) \cong (\mathbb{Z}/p^m\mathbb{Z}, q_{\bar{\zeta}}) \oplus (\mathbb{Z}/p^m\mathbb{Z}, q_{\bar{\zeta}}), \qquad m \ge 1.
$$

Consider the hyperbolic quadratic form

$$
h: \mathbb{Z}/p^m \mathbb{Z} \times \mathbb{Z}/p^m \mathbb{Z} \to k^{\times}, \qquad h(x, y) = \zeta^{xy}
$$

.

The corresponding metric group $(\mathbb{Z}/p^m\mathbb{Z}\times\mathbb{Z}/p^m\mathbb{Z}, h)$ is isomorphic to $(\mathbb{Z}/p^m\mathbb{Z}, q_\zeta)\oplus(\mathbb{Z}/p^m\mathbb{Z}, q_\zeta)$ if $p \equiv 1 \pmod{4}$ and to $(\mathbb{Z}/p^m \mathbb{Z}, q_\zeta) \oplus (\mathbb{Z}/p^m \mathbb{Z}, q_{\bar{\zeta}})$ if $p \equiv 3 \pmod{4}$.

The classification of anisotropic metric p-groups is well known. We recall it in Table [1.](#page-5-0)

	Odd prime p	Metric group	of number iso classes
⊥.	any	(\mathbb{Z}) $'p\mathbb{Z}$, q_{ζ}	
2.	$mod\;4$ $p \equiv 1($	$\setminus \oplus (\mathbb{Z}/p\mathbb{Z}, \overline{q_{\bar{e}}})$ $'p\mathbb{Z}$, 'Z, q_{ζ}	
3.	$p\equiv 30$ $mod\;4$	$(\mathbb{Z}/% \mathbb{Z})$ \mathbb{Z} , $'p\mathbb{Z}$, $/p\mathbb{Z}, q_c$ \oplus q_{ζ}	

TABLE 1. Anisotropic metric p-groups (for odd p). Here ζ is a primitive pth root of 1.

A description of M_2 is more involved. For any 2^{m+1} -th root of unity ξ and a 2^m -th root of unity ζ , $m \geq 1$, define quadratic forms

(3)
$$
q_{\xi}: \mathbb{Z}/2^m \mathbb{Z} \to k^{\times}, \qquad q_{\xi}(x) = \xi^{x^2},
$$

(4)
$$
h: \mathbb{Z}/2^m\mathbb{Z} \times \mathbb{Z}/2^m\mathbb{Z} \to k^{\times}, \qquad h(x, y) = \zeta^{xy},
$$

(5)
$$
f: \mathbb{Z}/2^m\mathbb{Z} \times \mathbb{Z}/2^m\mathbb{Z} \to k^{\times}, \qquad f(x, y) = \zeta^{x^2 + xy + y^2}.
$$

The quadratic form [\(3\)](#page-5-1) (respectively, [\(4\)](#page-5-1), [\(5\)](#page-5-1)) is non-degenerate if and only if ξ is a primitive 2^{m+1} -th root of unity (respectively, ζ is a primitive 2^m -th root of unity). There are 2 isomorphism classes of metric groups $(\mathbb{Z}/2\mathbb{Z}, q_i)$, $i^2 = -1$, and 4 isomorphism classes of metric groups $(\mathbb{Z}/2^m\mathbb{Z}, q_\xi)$ for each $m \geq 2$. Along with $(\mathbb{Z}/2^m\mathbb{Z} \times \mathbb{Z}/2^m\mathbb{Z}, h)$ and $(\mathbb{Z}/2^m\mathbb{Z} \times$ $\mathbb{Z}/2^m\mathbb{Z}, f$, $m \geq 1$, they generate M_2 . (Note that the isomorphism types of the last two metric groups are independent of the choice of ζ .) A complete list of relations between these generators is quite long and can be found in [\[KK,](#page-29-10) [Mir\]](#page-29-11). We only note the relation

(6)
$$
(\mathbb{Z}/2^m \mathbb{Z} \times \mathbb{Z}/2^m \mathbb{Z}, h)^{\oplus 2} \cong (\mathbb{Z}/2^m \mathbb{Z} \times \mathbb{Z}/2^m \mathbb{Z}, f)^{\oplus 2}, m \ge 1.
$$

Lemma 2.3. *A metric group* $((\mathbb{Z}/2^m\mathbb{Z})^k, q)$ *satisfies* $q^{2^m} = 1$ *if and only if it is a direct sum of some number of metric groups* [\(4\)](#page-5-1) *and* [\(5\)](#page-5-1)*.*

Proof. Such a metric group is isomorphic to an orthogonal sum of forms (3) , (4) , and (5) , but the condition $q^{2^m} = 1$ holds only for the last two.

Definition 2.4. A *fermion* in a pre-metric group (A, q) is an element $x \in A$ of order 2 such that $q(x) = -1$.

A proof of the classification of anisotropic pre-metric 2-groups given in Table [2](#page-6-2) can be found, e.g., in [\[DGNO2,](#page-29-2) Appendix A].

	Metric groups	Values of q	number of iso classes
1^*	$1, q_1$		
2^*	$\overline{\mathbb{Z}}/2\mathbb{Z}, q_i)$	1, i	$\overline{2}$
3^*	$(\mathbb{Z}/2\mathbb{Z}, q_i) \oplus (\mathbb{Z}/2\mathbb{Z}, q_i)$	$1, -1, i, i$	$\mathcal{D}_{\mathcal{L}}$
4^*	$\overline{\mathbb{Z}}/4\mathbb{Z}, q_{\xi})$	$-1, \xi, \xi$	4
5^*	$(\mathbb{Z}/4\mathbb{Z}, q_{\xi}) \oplus (\mathbb{Z}/2\mathbb{Z}, q_i)$	$-1, i, -i, \xi, -\xi, i\xi, i\xi$	$\overline{4}$
6	$(\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}, f)$	$-1, -1, -1$	
	$(\overline{\mathbb{Z}/2\mathbb{Z}} \times \mathbb{Z}/2\mathbb{Z}, f) \oplus (\mathbb{Z}/2\mathbb{Z}, q_i)$	$1, -1, -1, -1, i, -i, -i, -i$	$\overline{2}$
	Pre-metric groups		
$8*$	$(\overline{\mathbb{Z}}/2\mathbb{Z},\, q_{-1})$		
9^*	$\mathbb{Z}/2\mathbb{Z}, q_{-1}) \oplus (\mathbb{Z}/2\mathbb{Z}, q_i)$	$-1, i, -i$	

TABLE 2. Anisotropic pre-metric 2-groups. Here $i^2 = \xi^4 = -1$. Lines labelled by ∗ contain pre-metric groups with at most 1 fermion.

3. Maximal Tannakian subcategories of a gauged category

3.1. Subcategories of an equivariantization. Let $T: G \to \mathcal{A}ut^{\otimes}(\mathcal{A})$ be a monoidal functor, i.e., an action of a group G on a fusion category A. Let $N \subset G$ be a normal subgroup of G and let $\eta: T|_N \to I$ be a natural monoidal isomorphism between the restriction of T to N and the trivial monoidal functor $I: N \to \mathcal{A}ut^{\otimes}(\mathcal{A})$. We will call such η a *trivialization* of T on N. It provides a section $\mathcal{A} \to \mathcal{A}^N$ of the forgetful functor $\mathcal{A}^N \to \mathcal{A}$.

Following [\[GJ\]](#page-29-12), we say that η is *G-equivariant* if $\eta_{gng^{-1}} = T_g \eta_n T_{g^{-1}}$ for all $n \in N$, $g \in G$, where we identify $T_gT_nT_{g^{-1}}$ with $T_{gng^{-1}}$ using the monoidal functor structure of T.

One can check that G-equivariant trivializations of $T|_N$ are in bijection with isomorphism classes of factorizations of $T: G \to \mathcal{A}ut^{\otimes}(\mathcal{A})$ as a composition $G \to G/N \stackrel{\tilde{T}}{\to} \mathcal{A}ut^{\otimes}(\mathcal{A}),$ where \tilde{T} is a monoidal functor.

The following theorem was established by Galindo and Jones in [\[GJ\]](#page-29-12)

Theorem 3.1. Let A be a fusion category along with an action $T: G \to Aut^{\otimes}(\mathcal{A})$ of a *finite group* G. Fusion subcategories of A^G are in bijection with triples (D, N, η) , where D *is a G-stable fusion subcategory of* A, N *is a normal subgroup of G, and* $\eta: T|_N \to I$ *is a* G*-equinvariant trivialization.*

Remark 3.2. The subcategory of \mathcal{A}^G corresponding to the triple (\mathcal{D}, N, η) is $\mathcal{D}^{G/N}$, where the action of G/N on $\mathcal A$ is defined using η and the embedding of $\mathcal D^{G/N}$ into $\mathcal A^G$ is given by

$$
\mathcal{D}^{G/N} \hookrightarrow (\mathcal{D}^N)^{G/N} \cong \mathcal{D}^G \hookrightarrow \mathcal{A}^G.
$$

3.2. Subcategories of a central extension. Let \mathcal{B} be a braided fusion category, let G be a finite group, and let

(7)
$$
F: G \to \textbf{Pic}(\mathcal{B}): x \mapsto \mathcal{C}_x
$$

be a monoidal 2-functor. Let

(8)
$$
C := \bigoplus_{x \in G} C_x, \qquad C_1 = \mathcal{B},
$$

be the corresponding central extension of \mathcal{B} .

For a fusion subcategory $\tilde{\mathcal{C}} \subset \mathcal{C}$ let $\tilde{\mathcal{B}} = \mathcal{B} \cap \tilde{\mathcal{C}}$. There is a subgroup $H_{\tilde{\mathcal{C}}} \subset G$ such that $\tilde{\mathcal{C}}$ is a central $H_{\tilde{C}}$ -graded extension of \tilde{B} :

$$
\tilde{\mathcal{C}} = \bigoplus_{x \in H_{\tilde{\mathcal{C}}}} \tilde{\mathcal{C}}_x, \qquad \tilde{\mathcal{C}}_1 = \tilde{\mathcal{B}},
$$

where $\tilde{\mathcal{C}}_x \subset \mathcal{C}_x$ for all $x \in H_{\tilde{\mathcal{C}}}$. We will refer to $\tilde{\mathcal{C}}$ as a *subextension* of \mathcal{C} . We will say that a subextension $\tilde{\mathcal{C}} \subset \mathcal{C}$ is *faithful* if $H_{\tilde{\mathcal{C}}} = G$.

As an abstract central extension, $\tilde{\mathcal{C}}$ corresponds to a monoidal 2-functor

(9)
$$
\tilde{F}: H_{\tilde{\mathcal{C}}}\to \mathbf{Pic}(\tilde{\mathcal{B}}): x\mapsto \tilde{\mathcal{C}}_x.
$$

The tensor product $\mathcal{B} \boxtimes \tilde{\mathcal{C}} \to \mathcal{C}$ gives $\mathcal{B}\text{-module}$ equivalences $\mathcal{B} \boxtimes_{\tilde{\mathcal{B}}} \tilde{\mathcal{C}}_x \cong \mathcal{C}_x$ that combine into a pseudo-natural isomorphism of monoidal 2-functors:

$$
F|_{H_{\tilde{\mathcal{C}}}} \cong \text{Ind} \circ \tilde{F},
$$

where $\textsf{Ind}: \textbf{Pic}(\tilde{\mathcal{B}}) \to \textbf{Pic}(\tilde{\mathcal{C}})$ is the induction functor:

$$
\text{Ind}(\mathcal{M})=\mathcal{B}\boxtimes_{\tilde{\mathcal{B}}}\mathcal{M}.
$$

Conversely, an isomorphism (10) produces a subextension of \mathcal{C} :

$$
\tilde{\mathcal{C}}=\bigoplus_{x\in H_{\tilde{\mathcal{C}}}}\tilde{\mathcal{C}}_x\subset \bigoplus_{x\in H_{\tilde{\mathcal{C}}}}\mathcal{B}\boxtimes_{\tilde{\mathcal{B}}}\tilde{\mathcal{C}}_x\cong \bigoplus_{x\in H_{\tilde{\mathcal{C}}}}\mathcal{C}_x=\mathcal{C}.
$$

Definition 3.3. We will call a monoidal 2-functor $F : G \to Pic(B)$ *primitive* if it is not induced from any $F: G \to \text{Pic}(\tilde{\mathcal{B}})$ (as in [10\)](#page-7-1), where $\tilde{\mathcal{B}} \subset \mathcal{B}$ is a proper fusion subcategory. Similarly, we will call a central extension [\(8\)](#page-7-2) *primitive* if it has no proper faithful subextensions. A 2-functor (or a corresponding extension) is *imprimitive* if it is not primitive.

Let us describe subextensions of C in more explicit terms. Fix a fusion subcategory $\mathcal{B} \subset \mathcal{B}$. Let $\tilde{\mathcal{B}}^{co}$ be the *commutator* of $\tilde{\mathcal{B}}$ in C, i.e., the fusion subcategory of C generated by all simple objects $X \in \mathcal{C}$ such that $X \otimes X^* \in \tilde{\mathcal{B}}$ [\[GN\]](#page-29-13). This is the largest subcategory of \mathcal{C} that is a graded extension of $\ddot{\mathcal{B}}$:

(11)
$$
\tilde{\mathcal{B}}^{co} = \bigoplus_{y \in U(\tilde{\mathcal{B}})} (\tilde{\mathcal{B}}^{co})_y, \qquad (\tilde{\mathcal{B}}^{co})_1 = \tilde{\mathcal{B}},
$$

where $U(\tilde{\mathcal{B}})$ is the grading group of $\tilde{\mathcal{B}}^{co}$. Let $N(\tilde{\mathcal{B}}) \subset U(\tilde{\mathcal{B}})$ be the normal subgroup such that

(12)
$$
\tilde{\mathcal{B}}^{co} \cap \mathcal{B} = \bigoplus_{y \in N(\tilde{\mathcal{B}})} (\tilde{\mathcal{B}}^{co})_y
$$

and let $G(\tilde{\mathcal{B}}) \subset G$ be the projection of $U(\tilde{\mathcal{B}})$. We have a short exact sequence of groups:

(13)
$$
1 \to N(\tilde{\mathcal{B}}) \to U(\tilde{\mathcal{B}}) \stackrel{\pi}{\to} G(\tilde{\mathcal{B}}) \to 1.
$$

Let $O_{\tilde{\mathcal{B}}} \in H^2(G(\tilde{\mathcal{B}}), N(\tilde{\mathcal{B}}))$ be the cohomology class corresponding to the short exact sequence (13) .

Proposition 3.4. *Subextensions of* [\(8\)](#page-7-2) *are parameterized by the following data:*

- *a braided fusion subcategory* $\mathcal{B} \subset \mathcal{B}$,
- *a subgroup* $S \subset G(\tilde{\mathcal{B}})$ *, such that* $O_{\tilde{\mathcal{B}}}|_S$ *is cohomologically trivial, and*
- an element of an $H^1(S, N(\tilde{B}))$ -torsor.

Proof. A subextension $\tilde{\mathcal{C}} \subset \mathcal{C}$ such that $\tilde{\mathcal{C}} \cap \mathcal{B} = \tilde{\mathcal{B}}$ is contained in $\tilde{\mathcal{B}}^{co}$, which is a $U(\tilde{\mathcal{B}})$ graded extension of $\tilde{\mathcal{B}}$ [\(12\)](#page-8-2). If S is the support of $\tilde{\mathcal{C}}$ in G then $\pi^{-1}(S) \to S$ splits, which is equivalent to the restriction $O_{\tilde{B}}|_{S}$ being cohomologically trivial. Such splittings form a torsor over $H^1(S, N(\tilde{\mathcal{B}})).$

Conversely, if S is a subgroup of G and $\sigma : S \to \pi^{-1}(S)$ is a splitting, then

$$
\tilde{\mathcal{C}}_{\sigma} = \bigoplus_{z \in S} \, (\tilde{\mathcal{B}}^{co})_{\sigma(z)}
$$

is an S -graded subextension of C . The two above constructions are clearly inverses of each other. \Box

Remark 3.5. Any choice of a splitting $\sigma : S \to \pi^{-1}(S)$ determines a monoidal 2-functor

(14)
$$
F_{(\tilde{\mathcal{B}},S,\sigma)}:S\to \mathbf{Pic}(\tilde{\mathcal{B}}).
$$

Remark 3.6. Subextensions of C stable under the action of G correspond to a G -invariant version of the data from Proposition [3.4.](#page-8-3) Namely, the subcategory $\mathcal{B} \subset \mathcal{B}$ must be G-stable and the subgroup $S \subset G(\mathcal{B})$ must be normal in this case. Since the cohomology class of the obstruction $O_{\tilde{B}}$ is G-invariant, subextensions will be parameterized by a torsor over $H^1(S, N(\tilde{\mathcal{B}}))$ ^G.

3.3. Tannakian subcategories of a gauging. Let \mathcal{B} be a braided fusion category, let G be a finite group along with a monoidal 2-functor $F: G \to Pic(B)$, let

(15)
$$
\mathcal{C} = \bigoplus_{g \in G} \mathcal{C}_g, \qquad \mathcal{C}_1 = \mathcal{B},
$$

be the corresponding central extension, and let $\mathcal{G}(G,\mathcal{B},F) = \mathcal{C}^G$ be the corresponding gauging. The goal of this Section is to describe the set $T_{max}(\mathcal{G}(G,\mathcal{B}, F))$ of maximal Tannakian subcategories of $\mathcal{G}(G,\mathcal{B},F)$.

Definition 3.7. Let a group G act on a fusion category A. A braiding b on A is said to be G*-invariant* if G acts by braided (with respect to b) autoequivalences of A.

Proposition 3.8. $T_{max}(\mathcal{G}(G, \mathcal{B}, F))$ *is in bijection with the set of triples* (N, \mathcal{E}, b) *, where* N *is a normal Abelian subgroup of* G , $\mathcal{E} = \bigoplus_{n \in N} \mathcal{E}_n$ *is a G-stable subextension of* $\mathcal C$ *such that* $\mathcal{E}_1 \in T_{max}^G(\mathcal{B})$, and b is a G-invariant Tannakian braiding on $\mathcal{E}.$

Proof. Given a triple (N, \mathcal{E}, b) there is a trivialization of the action of N on $\mathcal E$ coming from the composition of natural isomorphisms

$$
X_n \otimes Y \xrightarrow{c_{X_n,Y}} T_n(Y) \otimes X_n \xrightarrow{b_{T_n(Y),X_n}} X_n \otimes T_n(Y), \qquad Y \in \mathcal{E}, X_n \in \mathcal{E}_n,
$$

where c denotes the G-crossed braiding of \mathcal{C} . This trivialization is G-invariant thanks to the G-invariance of the braiding b. It follows from Theorem [3.1](#page-6-3) and Remark [3.2](#page-6-4) that there is an action of G/N on $\mathcal E$ by braided autoequivalences and, consequently, a Tannakian subcategory

$$
\mathcal{T}(N,\,\mathcal{E},\,b):=\mathcal{E}^{G/N}\subset\bigoplus_{n\in N}\,\mathcal{C}_n.
$$

Note that \mathcal{E}_1^G is a maximal Tannakian subcategory of $\mathcal{G}(G,\mathcal{B},F)$. We compute:

 $FPdim(\mathcal{T}(N, \mathcal{E}, b)) = [G:N]FPdim(\mathcal{E}) = |G|FPdim(\mathcal{E}_1) = FPdim(\mathcal{E}_1^G).$

Since all maximal Tannakian subcategories of $\mathcal C$ have the same Frobenius-Perron dimension, it follows that $\mathcal{T}(N, \mathcal{E}, b) \in T_{max}(\mathcal{G}(G, \mathcal{B}, F)).$

Conversely, suppose that $\mathcal T$ is a maximal Tannakian subcategory of $\mathcal G(G,\mathcal B,F)$. Let $N_{\mathcal{T}} \subset G$ be a normal subgroup such that $\mathcal{T} \cap \mathcal{R}ep(G) = \mathcal{R}ep(G/N_{\mathcal{T}})$ and let

$$
\mathcal{E}_{\mathcal{T}} := \mathcal{T} \boxtimes_{\mathcal{R}ep(G/N_{\mathcal{T}})} \mathcal{V}ect \subset \mathcal{C}
$$

be the corresponding de-equivariantization. The Tannakian category $\mathcal{E}_{\mathcal{T}}$ is an $N_{\mathcal{T}}$ -graded extension of $\mathcal{T} \cap \mathcal{R}ep(G)$ and so the action of $N_{\mathcal{T}}$ on it is trivializable. Hence, the action of G on $\mathcal{E}_{\mathcal{T}}$ factors through the braided action of $G/N_{\mathcal{T}}$. In particular, $\mathcal{E}_{\mathcal{T}}$ is G-stable. Thus, we get a triple $(N_{\mathcal{T}}, \mathcal{E}_{\mathcal{T}}, b_{\mathcal{T}})$ with required properties, where $b_{\mathcal{T}}$ denotes the braiding of $\mathcal{E}_{\mathcal{T}}$.

Since equivariantization and de-equivariantization constructions are inverses of each other, the same is true for the assignments $(N, \mathcal{E}, b) \mapsto \mathcal{T}(N, \mathcal{E}, b)$ and $\mathcal{T} \mapsto (N_{\mathcal{T}}, \mathcal{E}_{\mathcal{T}}, b_{\mathcal{T}})$.

Definition 3.9. We will call a triple (N, \mathcal{E}, b) with $N \neq \{1\}$ from Proposition [3.8](#page-9-0) a *Tannakian imprimitivity datum* for the central extension [\(15\)](#page-8-4) (or for the corresponding monoidal 2-functor $F: G \to \text{Pic}(\mathcal{B})$ defining it). We say that an extension (or 2-functor) is *Tannakian primitive* if it does not admit a Tannakian imprimitivity datum.

- **Remark 3.10.** (1) If F admits a non-trivial Tannakian imprimitivity datum (N, \mathcal{E}, b) then $F|_N$ is imprimitive in the sense of Definition [3.3.](#page-7-3) In this case, the central extension corresponding to $F|_N$ is induced from a G-stable N-graded Tannakian subextension.
	- (2) We have $\mathcal{T}(N, \mathcal{E}, b) \cap \mathcal{R}ep(G) = \mathcal{R}ep(G/N)$. The support of the image of $\mathcal{T}(N, \mathcal{E}, b)$ under the de-equivariantization functor $\mathcal{C}^G \to \mathcal{C}$ is N.
	- (3) A parameterization of Tannakian imprimitivity data for a central extension [\(15\)](#page-8-4) can be given using the machinery of [\[DN2\]](#page-29-7), where the extension theory of symmetric categories was developed. We prefer not to do it here to avoid extra technical details.

3.4. Lagrangian subcategories of a twisted Drinfeld double. Here we apply results of the previous Section to describe Lagrangian subcategories of gaugings of the trivial fusion category Vect. By [\[DGNO2,](#page-29-2) Section 4.4.10], these are precisely representation categories of twisted Drinfeld doubles of finite groups.

Let N be a normal Abelian subgroup of a finite group G. Let $H^3(G, k^{\times})_1$ denote the kernel of the restriction homomorphism $H^3(G, k^{\times}) \to H^3(N, k^{\times})$.

Remark 3.11. There is a canonical homomorphism

(16)
$$
m: H^{3}(G, k^{\times})_{1} \to H^{1}(G, H^{2}(N, k^{\times})) : \omega \mapsto m_{\omega, N},
$$

where $H^2(N, k^{\times})$ is a right G-module via $(g \cdot \mu)(x, y) = \mu(gxg^{-1}, gyg^{-1}), g \in G, x, y \in N$. It is defined as follows. The cochain

(17)
$$
\mu_g(y,z) = \frac{\omega(gyg^{-1}, gzg^{-1}, g)\omega(g, y, z)}{\omega(gyg^{-1}, g, z)}
$$

defines a tensor structure on the conjugation autoequivalence $\text{ad}(g) : \oplus V_x \mapsto \oplus V_{gxg^{-1}}$, $g \in G$, of $\mathcal{V}ect_{G}^{\omega}$, that is, satisfies the identity

(18)
$$
\frac{\mu_g(xy, z)\mu_g(x, y)}{\mu_g(x, yz)\mu_g(y, z)} = \frac{\omega(gxg^{-1}, gyg^{-1}, gzg^{-1})}{\omega(x, y, z)}
$$

for all $g, x, y, z \in G$. Let d and ∂ denote differentials in $C^{\bullet}(G, k^{\times})$ and in $C^{\bullet}(G, H^2(N, k^{\times}))$, respectively. Let $\nu \in C^2(G, k^{\times})$ be such that $d\nu|_N = \omega|_N$. It follows from [\(18\)](#page-10-1) that

(19)
$$
m_{\omega,N}(g) := \mu_g|_N \times \frac{\nu}{\nu^g}, \qquad g \in G,
$$

is a 2-cocycle in $Z^2(N, k^{\times})$. The cochain $m_{\omega,N} \in C^1(G, H^2(N, k^{\times}))$ depends on the choice of ν , but its cohomology class does not. Indeed, if ν' is another choice, then $\nu'|_N = \nu|_N \times \eta$, where $\eta \in H^2(N, k^\times) = C^0(G, H^2(N, k^\times))$ and, hence, $m_{\omega,N}$ changes to $m_{\omega,N}\partial(\eta^{-1})$. To see that $m_{\omega,N}$ is a 1-cocycle, we compute

$$
\left(\partial(m_{\omega,N})(g,h)\right)(y,z) = \frac{\mu_{gh}(y,z)}{\mu_g(hyh^{-1},hzh^{-1})\mu_h(y,z)} = \frac{\gamma_{g,h}(yz)}{\gamma_{g,h}(y)\gamma_{g,h}(z)}, \qquad g,h,\in G, y,z \in N,
$$

where $\gamma_{g,h}(x) = \frac{\omega(g,hxh^{-1},h)}{\omega(g,h,x)\omega(ghxh^{-1}g^{-1},g,h)}, g,h,\in G, x \in N$. Thus, $\partial(m_{\omega,N})(g,h) = 0$ and the cohomology class of $m_{\omega,N}$ is a well defined element of $H^1(G, H^2(N, k^{\times}))$. This element depends on ω multiplicatively, hence the map [\(16\)](#page-10-2) is a group homomorphism.

Remark 3.12. Elements of $H^1(G, H^2(N, k^{\times}))$ correspond to isomorphism classes of monoidal functors $G \to \mathcal{A}ut^{\otimes}(\mathcal{V}ect_N)$ that restrict to the adjoint action of G on N.

Let

(20)
$$
\Omega(G;N) := \text{Ker}\left(H^3(G, k^{\times})_1 \xrightarrow{m} H^1(G, H^2(N, k^{\times}))\right).
$$

Remark 3.13. The group $\Omega(G; N)$ appeared in [\[U\]](#page-30-2) and is related to the problem of classification of categorical Morita equivalence classes of pointed fusion categories.

The following result was established in [\[NN\]](#page-29-14). We reformulate it in more transparent terms and derive it as a consequence of Proposition [3.8.](#page-9-0)

Proposition 3.14. *There is a bijection between* $T_{max}(\mathcal{Z}(\text{Vec}_G^{\omega}))$ *and the set of pairs* (N, b) *, where* N *is a normal Abelian subgroup of* G *such that* $\omega \in \Omega(G; N)$ *and* b *is an element of* an $H^2(N, k^{\times})^G$ -torsor.

Proof. In the notation of Proposition [3.8,](#page-9-0) we have $\mathcal{E}_1 = \mathcal{V}ect$ and $\mathcal{E} \cong \mathcal{V}ect_N$, where N is a normal Abelian subgroup of G. As in Remark [3.11,](#page-10-3) let $\nu \in C^2(G, k^{\times})$ be such that $d\nu|_N =$ $\omega|_N$ and let μ_g , $g \in G$, be defined by equation [\(17\)](#page-10-4). The Tannakian braiding on $\mathcal E$ is given by a function $b: N \times N \to k^{\times}$ such that $B := b$ Alt (ν) , where Alt $(\nu)(x, y) = \frac{\nu(x, y)}{\nu(y, x)}$, $x, y \in N$, is an alternating bilinear form on N . The G -invariance of this braiding is equivalent to the condition

(21)
$$
\frac{b^g}{b} \operatorname{Alt}(\mu_g) = 1,
$$

where $b^g(x, y) = b(gxg^{-1}, gyg^{-1}), g \in G, x, y \in N$. Equivalently,

(22)
$$
\frac{B^g}{B} \text{Alt}(m_{\omega,N}(g)) = 1, \qquad g \in G,
$$

where $m_{\omega,N}$ was defined in [\(19\)](#page-10-5). This translates to $\text{Alt}(m_{\omega,N}(g))$ being a coboundary in $C^1(G, H^2(N, k^{\times}))$, i.e., to $m_{\omega, N}(g)$ being cohomologically trivial. Different choices b, b' of the braiding on $\mathcal E$ give an alternating form B satisfying [\(22\)](#page-11-0) if and only if $\frac{b'}{b}$ $\frac{b'}{b}$ is a *G*-invariant alternating bilinear form on N. Since $H^2(N, k^{\times})^G \cong \text{Hom}(\bigwedge^2 N, k^{\times})^G$, we see that b is determined up to an element of an $H^2(N, k^*)^G$ -torsor.

Remark 3.15. Proposition [3.14](#page-11-1) can be interpreted as follows. Given a normal Abelian subgroup $N \subset G$ there is an obstruction $m_{\omega,N} \in H^1(G, H^2(N, k^{\times}))$ to the existence of Lagrangian subcategories of $\mathcal{Z}(\text{Vect}_{G}^{\omega})$ supported on N. When this obstruction vanishes,

the set of Lagrangian subcategories supported on N is parameterized by a torsor over $H^0(G, H^2(N, k^{\times})) = H^2(N, k^{\times})^G.$

Remark 3.16. When $\omega = 1$, the $H^2(N, k^{\times})^G$ -torsor is trivial and $L(\mathcal{Z}(\mathcal{V}ect_G))$ is in bijection with pairs (N, B) , where B is an alternating G-invariant bilinear form on N [\[NN\]](#page-29-14).

4. Tannakian radical, mantle, and reductive braided fusion categories

4.1. **Radical and mantle.** Let β be a braided fusion category.

Definition 4.1. The *Tannakian radical* of β is the intersection of all maximal Tannakian subcategories of B.

Let $Rad(\mathcal{B})$ denote the Tannakian radical of \mathcal{B} . This is a Tannakian category, so by Deligne's theorem there is canonical equivalence

(23)
$$
Rad(\mathcal{B}) \cong Rep(G_{\mathcal{B}})
$$

for a unique (up to an isomorphism) finite group $G_{\mathcal{B}}$.

Definition 4.2. The group $G_{\mathcal{B}}$ from [\(23\)](#page-12-2) will be called *the radical group* of \mathcal{B} .

Definition 4.3. The *mantle*^{[1](#page-12-3)} of \mathcal{B} is the localization of \mathcal{B} with respect to $\mathcal{R}ad(\mathcal{B})$, i.e.,

(24)
$$
\mathcal{M}\text{antle}(\mathcal{B}) = \mathcal{R}\text{ad}(\mathcal{B})' \boxtimes_{\mathcal{R}\text{ad}(\mathcal{B})} \mathcal{V}\text{ect.}
$$

Clearly, $G_{\mathcal{B}}$ and Mantle(\mathcal{B}) are invariants of \mathcal{B} . Furthermore, there is a canonical normal subgroup $H_{\mathcal{B}} \subset G_{\mathcal{B}}$ such that $\mathcal{B} \boxtimes_{\mathcal{R}ad(\mathcal{B})} \mathcal{V}ect$ is a faithfully $H_{\mathcal{B}}$ -graded extension of Mantle(\mathcal{B}) [\[DGNO2,](#page-29-2) Section 4.4.8]. Equivalently, there is a canonical monoidal 2-functor

(25)
$$
F_{\mathcal{B}}: H_{\mathcal{B}} \to \textbf{Pic}(\text{Mantle}(\mathcal{B})).
$$

Recall that a braided fusion category is *almost non-degenerate* if it is either non-degenerate or slightly degenerate. For an almost non-degenerate category \mathcal{B} we have $H_{\mathcal{B}} = G_{\mathcal{B}}$ and \mathcal{B} is equivalent to a $G_{\mathcal{B}}$ -gauging of Mantle(B). In this case, the triple $(G_{\mathcal{B}}, \text{Mantle}(\mathcal{B}), F_{\mathcal{B}})$ is a complete invariant of β .

Proposition 4.4. *Let* B *be an almost non-degenerate braided fusion category. The canonical monoidal* 2*-functor* $F_{\mathcal{B}}: G_{\mathcal{B}} \to \text{Pic}(\text{Mantle}(\mathcal{B}))$ *is Tannakian primitive.*

¹The term 'mantle' is used in geology to describe the layer surrounding the Earth's core. It extends outward and encompasses more material. This is similar to the relation between $Mantle(\mathcal{B})$ and $Core(\mathcal{B})$ in our setting since the latter is a localization of the former. We hope this analogy justifies our choice of the name.

Proof. Proposition [3.8](#page-9-0) gives a parametrization of $T_{max}(\mathcal{B})$ in terms of imprimitivity data (N, \mathcal{E}, b) associated to $F_{\mathcal{B}}$, see Definition [3.9.](#page-9-1) Let $\mathcal{T}(N, \mathcal{E}, b)$ denote the corresponding maximal Tannakian subcategory of \mathcal{B} . By Remark [3.10\(](#page-10-6)2),

$$
Rep(G_{\mathcal{B}}) \cap \mathcal{T}(N, \mathcal{E}, b) = Rep(G_{\mathcal{B}}/N).
$$

Since $\mathcal{R}ad(\mathcal{B}) = \mathcal{R}ep(G_{\mathcal{B}})$ is contained in every maximal Tannakian subcategory of \mathcal{B} , we conclude that $N = \{1\}$, so there is no non-trivial Tannakian imprimitivity data for $F_{\mathcal{B}}$. \Box

Let G be a finite group and let $N(G)$ be the subgroup of G generated by all normal Abelian subgroups of G. For $\omega \in H^3(G, k^{\times})$, let $N^{\omega}(G)$ be the subgroup of $N(G)$ generated by normal Abelian subgroups $N \subset G$ such that $\omega \in \Omega(G; N)$, see [\(20\)](#page-11-2). The groups $N(G)$ and $N^{\omega}(G)$ are nilpotent normal subgroups of G, as a product of nilpotent normal subgroups of a finite group is nilpotent.

Proposition 4.5. We have $\mathcal{R}ad(\mathcal{Z}(\mathcal{V}ect_G^{\omega})) \cong \mathcal{R}ep(G/N^{\omega}(G))$ and (26) Mantle($\mathcal{Z}(\mathcal{V}ect_G^{\omega}) \cong \mathcal{Z}(\mathcal{V}ect_{N^{\omega}(G)}^{\omega|_{N^{\omega}(G)}})$.

Proof. By Proposition [3.14,](#page-11-1) Lagrangian subcategories of $\mathcal{Z}(\mathcal{V}ect_G^{\omega})$ are parameterized by pairs (N, B) , where N is a normal Abelian subgroup of G and B is an element of a (possibly empty) $H^2(N, k^{\times})^G$ -torsor. By Remark [3.10\(](#page-10-6)2), the intersection of the Lagrangian category corresponding to (N, B) with $\mathcal{R}ep(G)$ is $\mathcal{R}ep(G/N)$. Therefore,

$$
Rad(\mathcal{Z}(\mathcal{V}ect_G^{\omega})) = \bigcap_{N \text{ is normal Abelian,} \atop \omega \in \Omega(G;N)} Rep(G/N) = Rep(G/N^{\omega}(G)).
$$

The centralizer $\mathcal{R}ep(G/N^{\omega}(G))'$ consists of objects of $\mathcal{Z}(\mathcal{V}ect^{\omega}_G)$ supported on $N^{\omega}(G)$ which is precisely $(\text{Vect}_{N^{\omega}(G)}^{\omega|_{N^{\omega}(G)})}$ ^G. The mantle of $\mathcal{Z}(\text{Vect}_{G}^{\omega})$ is the $G/N^{\omega}(G)$ -de-equivariantization of the latter, which is $\mathcal{Z}(\mathcal{V}ect^{ \omega|_{N^{\omega}(G)}}_{N^{\omega}(G)}$) and the contract of \Box

4.2. Reductive categories and their properties.

Definition 4.6. A braided fusion category \mathcal{B} is *reductive* if $\mathcal{R}ad(\mathcal{B}) = \mathcal{V}ect.$

Remark 4.7. Note that the maximal Tannakian subcategory of $\mathcal{Z}_{sym}(\mathcal{B})$ is contained in Rad(B). Therefore, if B is reductive, then $\mathcal{Z}_{sym}(\mathcal{B}) = \mathcal{V}ect$ or $\mathcal{S}\mathcal{V}ect$, i.e., B is almost nondegenerate.

Remark 4.8. It follows from Proposition [4.5](#page-13-2) that $\mathcal{Z}(\mathcal{V}ect_G^{\omega})$ is reductive if and only if $G = N^{\omega}(G)$. The latter condition implies that G is nilpotent. The converse is false in general. For example, if G is the dihedral group of order 2^n , $n \geq 4$, then $N(G) \subsetneq G$.

Example 4.9. Here are some examples of groups G such that $N(G) = G$.

- (1) Let G be a nilpotent group of nilpotency class \leq 2, i.e., such that $G/Z(G)$ is Abelian, where $Z(G)$ denotes the center of G. Any cyclic subgroup of $G/Z(G)$ gives rise to a normal Abelian subgroup of G. These subgroups generate G , so $N(G) = G$ in this case.
- (2) Let $G = UT(n, F)$ be the group of upper unitriangular $n \times n$ matrices, $n \geq 2$, with entries from a finite field F. This is a nilpotent group of class $n-1$ generated by matrices $e_{ij}(\lambda)$, $1 \leq i < j \leq n$, $\lambda \in F$, whose only nonzero entry off the main diagonal is λ in the (i, j) -position. The group G is generated by normal Abelian subgroups $N_k = \langle e_{ij}(\lambda) | 1 \leq k < j, \lambda \in F \rangle, k = 1, \ldots, n.$

Proposition 4.10. *A maximal Tannakian subcategory of a reductive braided fusion category is nilpotent.*

Proof. Let B be a reductive braided fusion category and let $\mathcal{E} = \mathcal{R}ep(G)$ be its maximal Tannakian subcategory. Then $\mathcal{B} \cong \mathcal{C}^G$, where C is a G-crossed braided fusion category. For any Tannakian subcategory $\mathcal{F} \subset \mathcal{B}$ let $N_{\mathcal{F}} \subset G$ be a normal subgroup such that $\mathcal{E} \cap$ $\mathcal{F} = \mathcal{R}ep(G/N_{\mathcal{F}}) \subset \mathcal{R}ep(G)$. By [\[DGNO2,](#page-29-2) Proposition 4.56], the support in G of the image of F under the de-equivariantization functor $\mathcal{C}^G \to \mathcal{C}$ is $N_{\mathcal{F}}$. Hence, $N_{\mathcal{F}}$ is Abelian. Since $\bigcap_{\mathcal{F} \in T_{max}(\mathcal{B})}$ $\mathcal{R}ep(G/N_{\mathcal{F}}) = \mathcal{R}ad(\mathcal{B}) = \mathcal{V}ect$, normal Abelian subgroups $\mathcal{N}_{\mathcal{F}}$ generate G , therefore, G is nilpotent. \Box

Proposition 4.11. *A reductive braided fusion category is weakly group-theoretical if and only if it is nilpotent.*

Proof. By definition, a nilpotent fusion category is weakly group-theoretical. Conversely, let **B** be a reductive weakly group-theoretical braided fusion category and let $\mathcal{E} = \mathcal{R}ep(G)$ be a maximal Tannakian subcategory of \mathcal{B} . By Proposition [4.11,](#page-14-0) G is nilpotent. Furthermore, the core of B is a nilpotent braided fusion category [\[Na\]](#page-30-0). Therefore, $C = \mathcal{B} \boxtimes_{\mathcal{E}} \mathcal{V}ect$ is nilpotent, since it is a graded extension of the core. Hence, $\mathcal{Z}(\mathcal{C})$ is nilpotent [\[GN\]](#page-29-13). Since $\mathcal{B} = \mathcal{C}^G$ is equivalent to a subcategory of $\mathcal{Z}(\mathcal{C})$ [\[DGNO2,](#page-29-2) Remark 4.43(i)], it follows from [\[GN\]](#page-29-13) that \mathcal{B} is nilpotent. \Box

Proposition 4.12. *Let* B *be a braided fusion category and let* G *be a group acting on* B *by braided autoequivalences. Then* $\mathcal{R}ad(\mathcal{B}) \subset \mathcal{R}ad(\mathcal{B}^G) \boxtimes_{\mathcal{R}ep(G)} \mathcal{V}ect.$

Proof. Any maximal Tannakian subcategory $\mathcal{T} \subset \mathcal{B}^G$ contains $\mathcal{R}ep(G)$ as a subcategory and so $\mathcal{T} = \mathcal{E}^G$, where $\mathcal E$ is a maximal G-stable Tannakian subcategory of $\mathcal B$. Hence, $\mathcal{R}ad(\mathcal{B})^G\subset \mathcal{T}$. Therefore,

$$
Rad(\mathcal{B})^G \subset \bigcap_{\mathcal{E} \in T_{max}^G(\mathcal{B})} \mathcal{E}^G = Rad(\mathcal{B}^G).
$$

The result follows by taking the de-equivariantization of both sides. \Box

Definition 4.13. Let \mathcal{B} be a braided fusion category and let G be a group acting on \mathcal{B} by braided autoequivalences. Define a G*-radical* of B by

$$
\mathcal{R}\hspace{-0.01cm}\mathrm{ad}^G(\mathcal{B}):=\bigcap_{\mathcal{E}\in T^G_{max}(\mathcal{B})}\mathcal{E}.
$$

If $\mathcal{R}ad^G(\mathcal{B}) = \mathcal{V}ect$ we will say that $\mathcal B$ is *G*-reductive.

Corollary 4.14. We have $\text{Rad}^G(\mathcal{B}) = \text{Rad}(\mathcal{B}^G) \boxtimes_{\text{Rep}(G)} \text{Vect}$ and $\text{Rad}(\mathcal{B}) \subset \text{Rad}^G(\mathcal{B})$. In *particular, if* B *is* G*-reductive, then it is reductive.*

Proof. This follows from the proof of Proposition [4.12](#page-14-1) since (de-)equivariantization commutes with taking the intersection of subcategories.

Corollary 4.15. Let β be a braided fusion category. Then Mantle(β) is G_{β} -reductive. In *particular,* Mantle(B) *is reductive.*

Proof. By definition of the radical, any maximal Tannakian subcategory of β is contained in $\mathcal{R}ad(\mathcal{B})'=\mathcal{M}antle(\mathcal{B})^{G_{\mathcal{B}}}$ and, hence,

$$
Rad(\mathcal{M}antle(\mathcal{B})^{G_{\mathcal{B}}})=\mathcal{R}ad(\mathcal{B})=\mathcal{R}ep(G_{\mathcal{B}}).
$$

Combining this with Corollary [4.15,](#page-15-1) we have

$$
Rad^{G_{\mathcal{B}}}(\mathcal{M}antle(\mathcal{B})) = Rad(\mathcal{M}antle(\mathcal{B})^{G_{\mathcal{B}}}) \boxtimes_{Rep(G_{\mathcal{B}})} Vect \cong Vect,
$$

as required. \square

4.3. A complete invariant of a braided fusion category.

Definition 4.16. A triple (G, \mathcal{C}, F) , where G is a finite group, C is a braided fusion category, and $F: G \to Pic(\mathcal{C})$ is a monoidal 2-functor such that

(1) F is Tannakian primitive in the sense of Definition [3.9](#page-9-1) and

(2) C is G-reductive with respect to the action of G on C coming from F.

will be called a *canonical gauging triple*.

A 1-isomorphism between canonical gauging triples (G, \mathcal{C}, F) and $(\tilde{G}, \tilde{\mathcal{C}}, \tilde{F})$ is a triple (i, I, ι) consisting of a group isomorphism $i: G \to \tilde{G}$, a braided equivalence $I: \tilde{\mathcal{C}} \to \mathcal{C}$, and a pseudo-natural isomorphism of monoidal 2-functors

(27)
$$
\begin{array}{ccc}\nG & \xrightarrow{F} & \text{Pic}(\mathcal{C}) \\
i & \xrightarrow{i} & \xrightarrow{t} \\
\tilde{G} & \xrightarrow{\tilde{F}} & \text{Pic}(\tilde{\mathcal{C}}),\n\end{array}
$$

where $I^* : Pic(\mathcal{C}) \to Pic(\tilde{\mathcal{C}})$ is a monoidal 2-equivalence induced from I.

If (i', I', ι') is another isomorphism between (G, \mathcal{C}, F) and $(\tilde{G}, \tilde{\mathcal{C}}, \tilde{F})$ then a 2-isomorphism between (i, I, ι) and (i', I', ι') is a pair (g, Γ) , where $g \in G$ is such that $i' = i \circ \text{ad}(g)$ and $\Gamma: I' \xrightarrow{\sim} g \circ I$ is a natural isomorphism of monoidal functors (here $g \in G$ is viewed as the autoequivalence of C obtained via the composition $G \to Pic(\mathcal{C}) \to Aut^{br}(\mathcal{C})$ such that the following prism of 2-cells commutes:

where Γ^* denotes the a pseudo-natural isomorphism of monoidal 2-functors induced by Γ and can is the canonical pseudo-natural isomorphism from [\[DN1,](#page-29-15) Theorem 3.10].

It follows that canonical gauging triples form a 2-groupoid which we will denote G.

Given a canonical gauging triple (G, \mathcal{C}, F) , the corresponding gauging $\mathcal{G}(G, \mathcal{C}, F)$ is a braided fusion category. The latter category is non-degenerate (respectively, slightly degenerate) if and only if $\mathcal C$ is non-degenerate (respectively, slightly degenerate).

Conversely, in Section [4.1](#page-12-1) to an almost non-degenerate braided fusion category β we associated a canonical gauging triple $\mathcal{T}(\mathcal{B}) := (G_{\mathcal{B}}, \text{Mantle}(\mathcal{B}), F_{\mathcal{B}}).$

Proposition 4.17. *Consider a triple* (G, \mathcal{C}, F) *, where* G *is a finite group,* C *is a reductive braided fusion category, and* $F : G \rightarrow Pic(C)$ *is a monoidal 2-functor. We have* $\mathcal{R}ad(\mathcal{G}(G,\mathcal{C}, F)) = \mathcal{R}ep(G)$ if and only if the following two conditions are satisfied:

- (1) *The functor* F *is Tannakian primitive in the sense of Definition [3.9,](#page-9-1)*
- (2) $\mathcal{R}ad^G(\mathcal{C}) = \mathcal{V}ect.$

Proof. Let $\mathcal{B} = \mathcal{G}(G, \mathcal{C}, F)$. The first condition is equivalent to $\mathcal{R}ep(G)$ being contained in every $\mathcal{E} \in T_{max}(\mathcal{B})$, while the second is equivalent to $\mathcal{R}ep(G)$ not being a proper subcategory of $Rad(\mathcal{B})$.

Let $\mathfrak B$ denote the 2-groupoid of almost non-degenerate braided fusion categories with braided equivalences as 1-morphisms and natural isomorphisms of tensor functors as 2 morphisms.

Theorem 4.18. *The assignments*

$$
(29)
$$

 $\mathcal{T} : \mathfrak{B} \to \mathfrak{G} : \mathcal{B} \mapsto (G_{\mathcal{B}}, \text{ Mantle}(\mathcal{B}), F_{\mathcal{B}})$ and $\mathcal{G} : \mathfrak{G} \to \mathfrak{B} : (G, \mathcal{C}, F) \mapsto \mathcal{G}(G, \mathcal{C}, F)$ *are mutually quasi-inverse* 2*-equivalences of* 2*-groupoids.*

Proof. It follows from Propositions [4.4](#page-12-4) and [4.17](#page-16-0) that $\mathcal{T}G(G,\mathcal{C},F) \cong (G,\mathcal{C},F)$. We have $\mathcal{GT}(\mathcal{B}) \cong \mathcal{B}$ since the equivariantization and de-equivariantization 2-functors are inverses of each other [\[DGNO2,](#page-29-2) Theorem 4.44].

An equivalence $\mathcal{B}_1 \xrightarrow{\sim} \mathcal{B}_2$ between braided fusion categories restricts to a braided equivalence $\mathcal{R}ad(\mathcal{B}_1) \xrightarrow{\sim} \mathcal{R}ad(\mathcal{B}_2)$ and, hence, gives an equivalence of crossed braided extensions

(30)
$$
\mathcal{B}_1 \boxtimes_{\mathcal{R}ad(\mathcal{B}_1)} \mathcal{V}ect \cong \mathcal{B}_2 \boxtimes_{\mathcal{R}ad(\mathcal{B}_2)} \mathcal{V}ect.
$$

In particular, there is a braided equivalence $I : \mathcal{M}antle(\mathcal{B}_1) \overset{\sim}{\to} \mathcal{M}antle(\mathcal{B}_2)$. The equivalence [\(30\)](#page-17-1) yields an isomorphism of the corresponding monoidal 2-functors [\[ENO2\]](#page-29-4). Thus, a 1 morphism in \mathfrak{B} yields a 1-morphism in \mathfrak{G} . Conversely, a 1-morphism between gauging triples gives rise to a braided equivalence between the corresponding gaugings. These assignments are inverse to each other, again thanks to the equivariantization - de-equivariantization correspondence.

Finally, a natural isomorphism η between braided autoequivalences $\alpha, \alpha': \mathcal{B}_1 \to \mathcal{B}_2$ yields an isomorphism $\eta|_{\text{Rad}(\mathcal{B}_1)}$ between their restrictions $\alpha|_{\text{Rad}(\mathcal{B}_1)}$, $\alpha'|_{\text{Rad}(\mathcal{B}_1)}$: $\text{Rad}(\mathcal{B}_1) \to \text{Rad}(\mathcal{B}_2)$. Let $\iota, \iota' : G_{\mathcal{B}_1} \to G_{\mathcal{B}_2}$ denote the corresponding isomorphisms between the radical groups. Then $\eta|_{\text{Rad}(\mathcal{B}_1)}$ is given by $\eta|_V = \iota(g)|_V$, $V \in \text{Rad}(\mathcal{B}_2) = \text{Rep}(G_{\mathcal{B}_2})$ for a unique $g \in G_{\mathcal{B}_1}$ such that $\iota' = \iota \circ \mathrm{ad}(g)$. Furthermore, a translation by g is an automorphism of the regular algebra $A_{\mathcal{B}_1} = \text{Fun}(G_{\mathcal{B}_1}, k) \in \mathcal{R}ep(G_{\mathcal{B}_1}).$ By definition, $\mathcal{M}antle(\mathcal{B}_1)$ is the category of $A_{\mathcal{B}_1}$ -modules in $\mathcal{R}ad(\mathcal{B}_1)'$, the centralizer of $\mathcal{R}ad(\mathcal{B}_1)$ in \mathcal{B}_1 , and the above action of g on $A_{\mathcal{B}_1}$ becomes a braided autoequivalence of **Mantle**(\mathcal{B}_1). Thus, $\eta|_{\mathcal{R}ad(\mathcal{B}_1)}$ yields a natural isomorphism $\Gamma: I' \to g \circ I$, where $I, I': \mathcal{M}antle(\mathcal{B}_2) \to \mathcal{M}antle(\mathcal{B}_1)$ are the braided equivalences coming from α, α' . Furthermore, there are induced equivalences of G-crossed braided categories

(31)
$$
\bar{\alpha}, \bar{\alpha}' : \mathcal{B}_1 \boxtimes_{\mathcal{R}ad(\mathcal{B}_1)} \mathcal{V}ect \xrightarrow{\sim} \mathcal{B}_2 \boxtimes_{\mathcal{R}ad(\mathcal{B}_2)} \mathcal{V}ect
$$

and η induces a natural isomorphism $\bar{\eta}$ between these. The fact that an $\bar{\eta}$ is an extension of Γ is equivalent to the commutativity of the polytope [\(28\)](#page-16-1).

Thus, $\mathcal T$ and $\mathcal G$ are mutually quasi-inverse 2-equivalences.

4.4. Towards a parametrization of Morita equivalence classes of fusion categories. It was shown in [\[ENO2\]](#page-29-4) that fusion categories A_1 , A_2 are categorically Morita equivalent if and only if their centers $\mathcal{Z}(\mathcal{A}_1)$ and $\mathcal{Z}(\mathcal{A}_2)$ are braided equivalent. In other words, the center is a complete invariant of the Morita equivalence class of a fusion category.

Corollary 4.19. *The canonical gauging triple* $(G_{\mathcal{Z}(A)}, \mathcal{M}$ *antle* $(\mathcal{Z}(A)), F_{\mathcal{Z}(A)})$ *is a complete invariant of the Morita equivalence class of a fusion category* A*. In particular, the radical* and mantle of $\mathcal{Z}(\mathcal{A})$ are Morita invariants of \mathcal{A} .

The mantle of $\mathcal{Z}(\mathcal{A})$ is a reductive metabolic braided fusion category, see Definition [2.1.](#page-3-1) In particular, it is braided equivalent to the representation category of a twisted Drinfeld double. This double is not uniquely defined. A classification of braided equivalence classes of Drinfeld doubles is equivalent to a classification of pointed fusion categories up to categorical Morita equivalence. The corresponding symmetric relation on the set of pairs (G, ω) , $\omega \in H^3(G, k^{\times})$, was described in [\[N,](#page-29-16) [U\]](#page-30-2). Our theory reduces the parametrization of these classes to the problem of classifying reductive metabolic braided fusion categories and computing their categorical Picard groups. As we saw in Remark [4.8](#page-13-1) and Proposition [4.11,](#page-14-0) such categories (and underlying groups G) are nilpotent. We discuss pointed reductive categories in Section [6](#page-22-0) and classify the smallest non-pointed metabolic reductive categories in Proposition [7.6.](#page-28-0)

5. The structure of reductive braided fusion categories

5.1. Tannakian-generated categories.

Definition 5.1. A braided fusion category B will be called a *Tannakian-generated category* or *TG-category* if it is generated by its Tannakian subcategories, i.e., $B = \bigvee_{\mathcal{E} \in T(\mathcal{B})} \mathcal{E}$.

Proposition 5.2. *A TG-category is group-theoretical.*

Proof. Let \mathcal{D}, \mathcal{E} be group-theoretical fusion subcategories of a TG-category \mathcal{B} . The tensor product of B determines a surjective tensor functor $\mathcal{D} \boxtimes \mathcal{E} \to \mathcal{D} \vee \mathcal{E}$ and the latter is grouptheoretical by [\[ENO1\]](#page-29-8). Hence, the induction on the number of Tannakian subcategories needed to generate $\mathcal B$ shows that $\mathcal B$ is group-theoretical.

Proposition 5.3. (i) *An almost non-degenerate TG-category is reductive.*

- (ii) *A metabolic braided fusion category is a TG-category if and only if it is reductive.*
- (iii) *A reductive TG-category is integral and nilpotent.*

Proof. (i) If a TG-category β is almost non-degenerate then

$$
\mathcal Rad(\mathcal B)=\bigcap_{\mathcal E\in T_{max}(\mathcal B)}\mathcal E\subset \left(\bigvee_{\mathcal E\in T_{max}(\mathcal B)}\mathcal E'\right)'\subset \left(\bigvee_{\mathcal E\in T_{max}(\mathcal B)}\mathcal E\right)'=\mathcal Z_{sym}(\mathcal B),
$$

so $Rad(B)$ must be trivial.

(ii) For a metabolic category B its subcategories $\bigcap_{\mathcal{E}\in T_{max}(\mathcal{B})} \mathcal{E}$ and $\bigvee_{\mathcal{E}\in T_{max}(\mathcal{B})} \mathcal{E}$ are centralizers of each other. So the former is trivial if and only if the latter coincides with \mathcal{B} .

(iii) This follows from Propositions [4.11](#page-14-0) and [5.2.](#page-18-3) \Box

Remark 5.4. It follows from Proposition [5.3\(](#page-18-4)iii) and [\[DGNO1,](#page-29-5) Theorem 6.10] that a reductive TG-category is a tensor product of TG-categories of prime power dimension.

Definition 5.5. The *coradical* of a braided fusion category β is the maximal TG-subcategory of \mathcal{B} :

(32)
$$
\mathcal{C} \text{orad}(\mathcal{B}) := \bigvee_{\mathcal{E} \in T(\mathcal{B})} \mathcal{E}.
$$

Remark 5.6. We have $\mathcal{R}ad(\mathcal{B}) = \mathcal{R}ad(\mathcal{C}orad(\mathcal{B}))$. In particular, \mathcal{B} is reductive if and only if \mathcal{C} *orad* (\mathcal{B}) is reductive.

Proposition 5.7. For any braided fusion category \mathcal{B} we have \mathcal{C} orad $(\mathcal{B}) \subset \mathcal{R}$ ad $(\mathcal{B})'$. When \mathcal{B} is metabolic, \mathcal{C} orad $(\mathcal{B}) = \mathcal{R}$ ad $(\mathcal{B})'$.

Proof. Since $\mathcal{E} \subset \mathcal{E}'$ for any Tannakian subcategory $\mathcal{E} \subset \mathcal{B}$, we have

$$
\mathcal{C}ord(\mathcal{B})=\bigvee_{\mathcal{E}\in T_{max}(\mathcal{B})}\mathcal{E}\subset \bigvee_{\mathcal{E}\in T_{max}(\mathcal{B})}\mathcal{E}'\subset \left(\bigcap_{\mathcal{E}\in T_{max}(\mathcal{B})}\mathcal{E}\right)'=\mathcal{R}ad(\mathcal{B}).'
$$

When B is metabolic, $\mathcal{E} = \mathcal{E}'$ for every $\mathcal{E} \in T_{max}(\mathcal{B})$ and the above inclusions become equalities. \Box

Corollary 5.8. Let β be a reductive braided fusion category. Then Corad(β)' is anisotropic, *i.e., has no non-trivial Tannakian subcategories. In particular, Corad* $(\mathcal{B})'$ *is almost nondegenerate.*

Proof. By Remark [5.6,](#page-19-1) $\text{Cora}(B)$ is reductive and, hence, is almost non-degenerate. Therefore, for any Tannakian subcategory $\mathcal{E} \subset \mathcal{C}$ *orad* $(\mathcal{B})'$ we have

$$
\mathcal{E} \cap \mathcal{C}orad(\mathcal{B}) = \mathcal{E} \cap \mathcal{Z}_{sym}(\mathcal{C}orad(\mathcal{B})) = \mathcal{V}ect.
$$

Since Corad(B) contains every Tannakian subcategory of \mathcal{B} , we conclude that \mathcal{E} must be trivial.

Thus, a reductive braided fusion category β has a canonical TG-subcategory Corad(β) and a canonical anisotropic subcategory \mathcal{C} *orad*(\mathcal{B})'.

5.2. Projectively isotropic objects. Let β be a braided fusion category.

Definition 5.9. We will say that an object $X \in \mathcal{B}$ is *symmetric* if $c_{X,X}^2 = id_{X \otimes X}$ and *projectively symmetric* if $c_{X,X}^2 = \lambda \, id_{X \otimes X}$ for some $\lambda \in k^{\times}$.

For any object X of B let $\langle X \rangle$ denote the fusion subcategory of B generated by X. Clearly, X is symmetric if and only if $\langle X \rangle$ is symmetric.

Definition 5.10. A symmetric object X is *isotropic* if $\langle X \rangle$ is Tannakian.

If X is projectively symmetric then $\langle X \otimes X^* \rangle$ is symmetric by [\[DGNO2,](#page-29-2) Proposition 3.22].

Definition 5.11. A projectively symmetric object X is *projectively isotropic* if $\langle X \otimes X^* \rangle$ is Tannakian.

Remark 5.12. Our terminology is consistent with the one used in [\[DGNO2,](#page-29-2) Section 3.3].

Lemma 5.13. Let X be a projectively isotropic object of \mathcal{B} . Then FPdim(X) $\in \mathbb{Z}$.

Proof. It follows from [\[DGNO2,](#page-29-2) Lemma 3.15] that X is a homogeneous object in a graded extension of some Tannakian subcategory $\mathcal{E} = \mathcal{R}ep(G) \subset \mathcal{B}$. But any homogeneous component of this extension is equivalent to the category of projective representations of G with a fixed Schur multiplier [\[DN2\]](#page-29-7). Therefore, $FPdim(X)$ is equal to the degree of a projective representation of G .

Let \mathcal{B}_{int} be the maximal integral fusion subcategory of \mathcal{B} , i.e., the subcategory generated by all objects $X \in \mathcal{B}$ such that $\text{FPdim}(X) \in \mathbb{Z}$. By [\[DGNO2,](#page-29-2) Corollary 2.24], there is a canonical ribbon structure θ on \mathcal{B}_{int} with respect to which categorical dimensions coincide with the Frobenius-Perron dimensions (this is true regardless of the existence of a ribbon structure on \mathcal{B}).

Lemma 5.14. *Let* X *be an object of* B*.*

- (i) *A symmetric object* $X \in \mathcal{B}$ *is isotropic if and only if* $\theta_X = id_X$.
- (ii) If X is a projectively isotropic simple object then $c_{X,X}^2 = \theta_X^2 id_{X \otimes X}$.

Proof. (i) This is a well-known fact of symmetric fusion categories.

(ii) We have $\theta_{X\otimes X^*} = (\theta_X \otimes \theta_{X^*})c_{X^*,X}c_{X,X^*}$. But $\theta_{X\otimes X^*} = 1$ by (i), so the statement follows by [\[DGNO2,](#page-29-2) Lemma 3.15(ii)]. \square

Lemma 5.15. Let X, Y be simple objects of B centralizing each other. Then $X \otimes Y$ is an *isotropic object of* \mathcal{B} *if and only if* X, Y *are projectively isotropic and* $\theta_X \theta_Y = 1$ *.*

Proof. The hexagon axioms of braiding imply that $c_{X \otimes Y, X \otimes Y}^2$ is a conjugate of $c_{X,X}^2 \otimes c_{Y,Y}^2$. So $X \otimes Y$ is symmetric if and only if X, Y are projectively isotropic and the scalars $c_{X,X}^2$ and $c_{Y,Y}^2$ are reciprocals of each other. But these scalars are θ_X^2 and θ_Y^2 , respectively, so the result follows from Lemma [5.14.](#page-20-0)

Definition 5.16. Let $\mathcal{B}_1, \mathcal{B}_2$ be braided fusion categories. We will say that two objects $X_1 \in \mathcal{B}_1, X_2 \in \mathcal{B}_2$ are *complementary* to each other if they are projectively isotropic and satisfy $\theta_{X_1}\theta_{X_2} = 1$. We will say that X_2 is *complementary* to \mathcal{B}_1 if it is complementary

to some $X_1 \in \mathcal{B}_1$ and *non-complementary* to \mathcal{B}_1 otherwise. We will say that \mathcal{B}_2 is *noncomplementary* to \mathcal{B}_1 if every simple object of \mathcal{B}_2 other than 1 is non-complementary to \mathcal{B}_1 . If \mathcal{B}_2 has a specified fermion object δ (e.g., if \mathcal{B}_2 is slightly degenerate) we will say that \mathcal{B}_2 is *super non-complementary* to \mathcal{B}_1 if every simple object of \mathcal{B}_2 other than 1 and δ is non-complementary to \mathcal{B}_1 .

Corollary 5.17. *Let* B *be a TG-category and* C *be a braided fusion category. Then*

(33)
$$
\mathcal{C} \text{orad}(\mathcal{B} \boxtimes \mathcal{C}) = \mathcal{B} \boxtimes \tilde{\mathcal{C}},
$$

where \tilde{C} *is the fusion subcategory of* C *generated by simple objects* Y *complementary to* \mathcal{B} *.*

5.3. A canonical decomposition of a reductive category.

Definition 5.18. Let \mathcal{B} be a reductive braided fusion category. We say that \mathcal{B} is

of type I if \mathcal{C} *orad* (\mathcal{B}) is non-degenerate,

of type II if both $\mathcal B$ and $\mathcal C$ *orad*($\mathcal B$) are slightly degenerate,

of type III if $\mathcal B$ is non-degenerate and \mathcal{C} or $\mathcal{A}(\mathcal B)$ is slightly degenerate.

By Remarks [4.7](#page-13-3) and [5.6,](#page-19-1) any reductive braided fusion category belongs to one of the above types.

Remark 5.19. A reductive braided fusion category β of type I is a tensor product of a TG-category Corad(\mathcal{B}) and an anisotropic category Corad(\mathcal{B}')', see Corollary [5.8.](#page-19-2) On the other hand, let \mathcal{B}_1 be a reductive TG-category, \mathcal{B}_2 be an anisotropic category, and consider $\mathcal{B} = \mathcal{B}_1 \boxtimes \mathcal{B}_2$. Then $\mathcal{B}_1 \subseteq \mathcal{C} \text{orad}(\mathcal{B})$, but, in general, $\mathcal{B}_1 \neq \mathcal{C} \text{orad}(\mathcal{B})$. For example, take $\mathcal{B}_1 =$ $\mathcal{Z}(\mathcal{R}ep(\mathbb{Z}/2\mathbb{Z}))$ and \mathcal{B}_2 an Ising category. Then $\mathcal{C}ora\mathcal{A}(\mathcal{B}_1 \boxtimes \mathcal{B}_2) = \mathcal{Z}(\mathcal{R}ep(\mathbb{Z}/2\mathbb{Z})) \boxtimes \mathcal{S}\mathcal{V}ect,$ cf. Corollary [5.17.](#page-21-3) In Theorem [5.20](#page-21-2) we will provide a decomposition of β in terms of its coradical and anisotropic parts.

Recall that a slightly degenerate braided fusion category β admits a minimal non-degenerate extension [\[JFR\]](#page-29-17). Such an extension is faithfully $\mathbb{Z}/2\mathbb{Z}$ -graded with β being the trivial component. There are 16 braided equivalence classes of such extensions, see, e.g., [\[DNO,](#page-29-18) Section 5.3]

A slightly degenerate braided fusion category B is *split* if $\mathcal{B} \cong \tilde{\mathcal{B}} \boxtimes \mathcal{S}\mathcal{V}$ where $\tilde{\mathcal{B}}$ is non-degenerate (the braided equivalence class of $\tilde{\mathcal{B}}$ is an invariant of \mathcal{B}). Otherwise, we say that B is *non-split*.

Theorem 5.20. *Let* B *be a reductive braided fusion category.*

(1) B *is of type I if and only if* $\mathcal{B} \cong \mathcal{B}_1 \boxtimes \mathcal{B}_2$ *, where* \mathcal{B}_1 *is a non-degenerate reductive TG-category and* \mathcal{B}_2 *is an anisotropic category non-complementary to* \mathcal{B}_1 *. In this case,* $\mathcal{B}_1 \cong \mathcal{C} \text{orad}(\mathcal{B})$ *and* $\mathcal{B}_2 \cong \mathcal{C} \text{orad}(\mathcal{B})'$ *.*

- (2) **B** *is of type II if and only if* $\mathcal{B} \cong \mathcal{B}_1 \boxtimes_{s\text{Vect}} \mathcal{B}_2$ *, where* \mathcal{B}_1 *is a reductive slightly degenerate TG-category and* \mathcal{B}_2 *is a slightly degenerate anisotropic category super non-complementary to* \mathcal{B}_1 *. In this case,* $\mathcal{B}_1 \cong \mathcal{C} \text{orad}(\mathcal{B})$ *and* $\mathcal{B}_2 \cong \mathcal{C} \text{orad}(\mathcal{B})'$ *.*
- (3) B *is of type III if and only if it is a minimal non-degenerate extension of a type II reductive category* \mathcal{B}_0 *such that either* \mathcal{B}_0 *is non-split, or* $\mathcal{B}_0 \cong \tilde{\mathcal{B}}_0 \boxtimes$ *sVect is split* and $\mathcal{B} \cong \tilde{\mathcal{B}}_0 \boxtimes \mathcal{C}$, where \mathcal{C} *is a minimal non-degenerate extension of s*Vect *super non-complementary to* \mathcal{B}_0 *. In this case,* $\mathcal{B}_0 = \text{Cora}(\mathcal{B}) \vee \text{Cora}(\mathcal{B})'$ *.*

Proof. (1) Since \mathcal{C} *orad*(\mathcal{B}) is non-degenerate, we have $\mathcal{B} \cong \mathcal{C}$ *orad*(\mathcal{B}) ⊠ \mathcal{C} *orad*(\mathcal{B})'. It follows from Lemma [5.15](#page-20-1) that \mathcal{C} *orad* $(\mathcal{B})'$ must be non-complementary to \mathcal{C} *orad* (\mathcal{B}) . Conversely, given categories \mathcal{B}_1 and \mathcal{B}_2 satisfying the hypothesis, $\text{Cora}(\mathcal{B}_1 \boxtimes \mathcal{B}_2) = \mathcal{B}_1$ by Corollary [5.17](#page-21-3) and $\mathcal{R}ad(\mathcal{B}_1 \boxtimes \mathcal{B}_2) = \mathcal{R}ad(\mathcal{B}_1) = \mathcal{V}ect.$

(2) We have $\mathcal{B} \cong \mathcal{C} \text{orad}(\mathcal{B}) \boxtimes_{sVect} \mathcal{C} \text{orad}(\mathcal{B})'$ by [\[DNO,](#page-29-18) Proposition 4.3]. Hence, every simple object of B is isomorphic to $X \boxtimes Y$, where $X \in \mathcal{C}$ and $Y \in \mathcal{C}$ and $(Y, \text{ so the})$ argument from part (1) applies here as well.

(3) When B is of type III, $\mathcal{B}_0 = \text{Cora}(B) \vee \text{Cora}(B)'$ is a type II reductive slightly degenerate subcategory of β of index 2, so β is a minimal extension of \mathcal{B}_0 . In the opposite direction, let us determine when a minimal extension β of a given type II reductive category \mathcal{B}_0 satisfies \mathcal{C} orad $(\mathcal{B}) = \mathcal{C}$ orad (\mathcal{B}_0) (in which case $\mathcal B$ is automatically reductive). Note that $Rad(\mathcal{B})$ is a (possibly trivial) $\mathbb{Z}/2\mathbb{Z}$ -graded extension of $Rad(\mathcal{B}) = Vect$, so either $Rad(\mathcal{B}) =$ Vect (and B is reductive) or $\mathcal{R}ad(\mathcal{B}) = \mathcal{R}ep(\mathbb{Z}/2\mathbb{Z})$. The latter situation is only possible when the non-trivial homogeneous component of $\mathcal B$ contains an invertible object, i.e., when $\mathcal{B}_0 \cong \tilde{\mathcal{B}}_0 \boxtimes$ sVect [\[JFR\]](#page-29-17). In this case $\mathcal{B} = \tilde{\mathcal{B}}_0 \boxtimes \mathcal{C}$, where \mathcal{C} is a minimal extension of sVect. By Corollary [5.17,](#page-21-3) \mathcal{C} orad(\mathcal{B}) = \mathcal{C} orad(\mathcal{B}_0) if and only if \mathcal{C} is super non-complementary to \mathcal{B} .

Remark 5.21. The direct factor C that appears in the split case of Theorem [5.20\(](#page-21-2)3) must be a non-degenerate anisotropic braided fusion category of the Frobenius-Perron dimension 4 with exactly one fermion. There are 14 such categories: 8 Ising categories and 6 pointed categories corresponding to metric groups in lines 3 and 4 of Table [2.](#page-6-2) Note that Ising categories are super non-complementary to any \mathcal{B}_0 , while for the pointed ones this property depends on the choice of \mathcal{B}_0 .

6. Reductive pointed braided fusion categories

Recall that pointed braided fusion categories correspond to pre-metric groups, i.e., pairs (A, q) , where $q : A \to k^{\times}$ is a quadratic form. These categories are denoted $\mathcal{C}(A, q)$. The goal of this Section is to classify pre-metric groups (A, q) such that $\mathcal{C}(A, q)$ is reductive.

6.1. Isotropically generated metric groups. We will say that a pre-metric group (A, q) is *isotropically generated* if A is generated by its *isotropic elements*, i.e., by those $x \in A$ for which $q(x) = 1$. The set of such elements is called the *light cone* in the literature.

Clearly, (A, q) is isotropically generated if and only if $\mathcal{C}(A, q)$ is a TG-category.

Example 6.1. For any finite Abelian group A let

(34) $h : A \oplus \widehat{A} \to k^{\times}, \qquad h(a, \phi) = \langle \phi, a \rangle, \quad a \in A, \phi \in \widehat{A},$

denote the canonical hyperbolic quadratic form on A. The *hyperbolic* metric group $(A \oplus \widehat{A}, h)$ is isotropically generated since A and \widehat{A} are isotropic subgroups of $A \oplus \widehat{A}$.

Theorem 6.2. Let (A, q) be a metric group and let n be the exponent of A. Then (A, q) is *isotropically generated if and only if the following two conditions are satisfied:*

- (i) $q^n = 1$ *, and*
- (ii) (A, q) *contains a hyperbolic metric subgroup* $(\mathbb{Z}/n\mathbb{Z} \times \mathbb{Z}/n\mathbb{Z}, h)$ *.*

Proof. We may assume that A is a *p*-group. Let

$$
B(x, y) = \frac{q(x + y)}{q(x)q(y)}, \qquad x, y \in A,
$$

be the non-degenerate symmetric bilinear form on A associated to q.

Suppose that (A, q) is isotropically generated. Every $x \in A$ can be written as $x =$ $x_1 + \cdots + x_k$ with $q(x_i) = 1$, $i = 1, \ldots k$. Therefore, $q(x) = \prod_{1 \leq i < j \leq k} B(x_i, x_j)$. But each factor $B(x_i, x_j)$ is an *n*th root of 1 and so $q(x)^n = 1$, proving (i).

Note that A must contain an isotropic element a of order n. Indeed, isotropic elements of A of order less than n generate a subgroup of exponent less than n, i.e., a proper subgroup of A. We claim that there is $c \in A$ such that

(35)
$$
q(c) = 1 \text{ and } B(a, c) \text{ is a primitive } n \text{th root of } 1.
$$

To see this, let $n_1 = n/p$ and suppose that $B(a, x)^{n_1} = 1$ for all $x \in A$. We have $B(n_1 a, x) =$ $B(a, x)^{n_1} = 1$, which contradicts the non-degeneracy of B. So there is $c' \in A$ such that $B(a, c')$ is a primitive nth root of 1. Hence, the order of c' in A is also n. Let us choose $k \in \mathbb{Z}$ such that $B(a, c')^k q(c') = 1$ and set $c = ka + c'$. Then c satisfies [\(35\)](#page-23-2) and the metric group $(\langle a, c \rangle, q|_{\langle a, c \rangle})$ is isomorphic to $(\mathbb{Z}_n \times \mathbb{Z}_n, h)$. This proves (ii).

Now suppose that (A, q) satisfies conditions (i) and (ii). Then

$$
(A, q) \cong (H, h) \oplus (A_1, q_1),
$$

where $(H, h) := (\mathbb{Z}_n \times \mathbb{Z}_n, h)$ and (A_1, q_1) is metric group such that $q_1^n = 1$. To prove that (A, q) is isotropically generated, note that the image of h consists of all nth roots of unity. Therefore, (A_1, q_1) is complementary to (H, h) in the sense of Definition [5.16](#page-20-2) (here we use the same terminology for metric groups and braided fusion categories). Hence, (A, q) is isotropically generated by Corollary [5.17.](#page-21-3)

Corollary 6.3. *Let* p *be an odd prime. Any isotropically generated metric* p*-group* (A, q) *of exponent* p e *can be written as*

$$
(A, q) = (\mathbb{Z}/p^e \mathbb{Z} \times \mathbb{Z}/p^e \mathbb{Z}, h) \bigoplus \{ sum \ of \ metric \ p\text{-groups of exponent} \leq p^e \}.
$$

Any isotropically generated metric 2*-group* (A, q) *of exponent* 2 e *can be written as*

 $(A, q) = (\mathbb{Z}/2^e \mathbb{Z} \times \mathbb{Z}/2^e \mathbb{Z}, h)^{\oplus n} \bigoplus (\mathbb{Z}/2^e \mathbb{Z} \times \mathbb{Z}/2^e \mathbb{Z}, f)^{\oplus e}$

 \bigoplus {sum of metric 2-groups of exponent $\langle 2^e \rangle$, $n \ge 1, \epsilon \in \{0, 1\},$

where the forms h *and* f *were defined in* [\(4\)](#page-5-1) *and* [\(5\)](#page-5-1)*.*

Proof. Condition (i) of Theorem [6.2](#page-23-1) is automatically satisfied for any odd p, which implies the first decomposition. The second one follows by Lemma [2.3.](#page-5-2)

Corollary 6.4. *Let* (A, q) *be an isotropically generated metric group and let* n *be the exponent of* A*.* For any nth root of unity $\zeta \in k^{\times}$ there is $x \in A$ such that $q(x) = \zeta$.

Proof. This is easily seen to be true for the hyperbolic group $(\mathbb{Z}/p^n\mathbb{Z} \times \mathbb{Z}/p^n\mathbb{Z}, h)$, so the result follows from Theorem [6.2.](#page-23-1)

Lemma 6.5. *Let* (A, q) *be an isotropically generated metric group and let* n *be the exponent of* A. Let (B, r) be a pre-metric group. Define $\tilde{B} = \langle y \in B | r(y)^n = 1 \rangle$. Then $A \oplus \tilde{B}$ is an *isotropically generated subgroup of* $(A \oplus B, q \oplus r)$ *.*

Proof. It follows from Corollary [6.4](#page-24-1) that for any $y \in \tilde{B}$ there is $x \in A$ such that $q(x)r(y) = 1$ and so (x, y) is an isotropic element of $A \oplus B$. Therefore, $A \oplus \tilde{B}$ is isotropic.

6.2. Reductive metric p -groups.

Definition 6.6. A pre-metric group is *reductive* if the intersection of its maximal isotropic subgroups is trivial.

Equivalently, a pre-metric group (A, q) is reductive if and only if the braided fusion category $\mathcal{C}(A, q)$ is reductive.

Proposition 6.7. *Let* p *be an odd prime. A metric* p*-group* (A, q) *is reductive if and only it is either isotropically generated or anisotropic.*

Proof. Since |A| is odd, the reductive pointed fusion category $C(A, q)$ is of type I, see Defini-tion [5.18.](#page-21-1) By Theorem [5.20\(](#page-21-2)1), $(A, q) = (A_1, r) \bigoplus (A_2, s)$, where (A_1, r) is an isotropically generated metric p-group and (A_2, s) is an anisotropic metric p-group non-complementary to (A_1, r) . If (A_1, r) is non-trivial it contains a hyperbolic summand by Theorem [6.2.](#page-23-1) So the values of r include all pth roots of 1. But the values of s are also pth roots of 1 and so if (A_2, s) is non-trivial, it must be complementary to (A_1, r) , a contradiction.

Lemma 6.8. *Let* (A, q) *be a slightly degenerate isotropically generated pre-metric* 2*-group. Then there is an isotropically generated metric* 2*-group* (\tilde{A}, \tilde{q}) *and an orthogonal isomorphism*

(36)
$$
(A, q) \cong (\tilde{A}, \tilde{q}) \oplus (\mathbb{Z}/2\mathbb{Z}, q_{-1}),
$$

where (Z/2Z, q−1) *is the unique non-isotropic degenerate metric group of order* 2*, cf.* [\(3\)](#page-5-1)*.*

Proof. By [\[DGNO1,](#page-29-5) Corollary A.19], (A, q) splits into an orthogonal direct sum as in [\(36\)](#page-25-1), so we only need to show that (A, \tilde{q}) can be chosen to be isotropically generated. If the exponent of A is ≥ 4 then (A, q) (and, hence, (A, \tilde{q})) contains a hyperbolic summand of the same exponent.

If the exponent of A is 2 then (\tilde{A}, \tilde{q}) is an orthogonal sum of metric groups [\(4\)](#page-5-1), [\(5\)](#page-5-1) by Lemma [2.3.](#page-5-2) Since

(37)
$$
(\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}, h) \oplus (\mathbb{Z}/2\mathbb{Z}, q_{-1}) \cong (\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}, f) \oplus (\mathbb{Z}/2\mathbb{Z}, q_{-1})
$$

we can assume that (\tilde{A}, \tilde{q}) has a hyperbolic summand.

By Theorem [6.2,](#page-23-1) in both cases the metric group (\tilde{A}, \tilde{q}) is isotropically generated. \Box

Theorem 6.9. *For a reductive pre-metric* 2*-group exactly one of the following is true:*

- (1) *it is isotropically generated,*
- (2) *it is non-trivial anisotropic (see Table [2\)](#page-6-2),*
- (3) *it is an orthogonal sum of an isotropically generated metric* 2*-group of exponent* 2 *and a non-trivial anisotropic pre-metric* 2*-group with at most one fermion (such groups are labelled by* [∗] *in Table [2\)](#page-6-2).*

Proof. All we need to do is to classify reductive pre-metric 2-groups (B, r) that are neither isotropically generated nor anisotropic. Let (A, q) be the maximal isotropically generated subgroup (i.e., the coradical) of (B, r) .

If (A, q) is non-degenerate then

$$
(B, r) = (A, q) \oplus (A', q'),
$$

where (A', q') is anisotropic and non-complementary to (A, q) , see Corollary [5.17.](#page-21-3) By Lemma [6.5,](#page-24-2) $q'(x)^n \neq 1$ for any non-zero $x \in A$, where *n* is the exponent of *A*. From Table [2](#page-6-2) we see that this is only possible when $n = 2$ and $(A', q') = (\mathbb{Z}/2\mathbb{Z}, q_i), i^2 = -1$.

If (A, q) is slightly degenerate then using Lemma [6.8](#page-25-2) we have

$$
(B, r) = (\tilde{A}, \tilde{q}) \oplus (A', q'),
$$

where (\tilde{A}, \tilde{q}) is an isotropically generated metric 2-group such that [\(36\)](#page-25-1) holds and (A', q') is an anisotropic pre-metric group containing a (necessarily unique) fermion. The same argument as above shows that the exponent of A is 2, so the statement follows. \Box

Remark 6.10. Equation [\(37\)](#page-25-3) implies that the choice of an isotropically generated summand in Theorem [6.9\(](#page-25-0)3) is non-unique. But the anisotropic summand is unique as it is the orthogonal complement of the coradical.

7. Reductive p-categories of small dimension

Let p be a prime. We say that a braided fusion category $\mathcal B$ is a p-category if $\text{FPdim}(\mathcal B) = p^n$ for some integer $n \geq 0$. Such categories are weakly group-theoretical by [\[ENO3\]](#page-29-19).

Lemma 7.1. Let \mathcal{B} be a reductive category with $FPdim(\mathcal{B}) = p^n$, $n \leq 5$. Then Corad(\mathcal{B}) is *pointed.*

Proof. Let $\mathcal{E} \subset \mathcal{C}$ *orad*(\mathcal{B}) be a maximal Tannakian subcategory.

If Corad(B) is non-degenerate, then $\mathrm{FPdim}(\mathcal{E})^2 \leq \mathrm{FPdim}(\mathcal{E})\mathrm{FPdim}(\mathcal{E}') = \mathrm{FPdim}(\mathcal{B}) \leq p^5$ and so FPdim $(\mathcal{E}) \leq p^2$. This implies that $\mathcal E$ is pointed. Since Corad(B) is generated by Tannakian subcategories of \mathcal{B} , it is pointed as well.

If $p = 2$ and \mathcal{C} *orad*(\mathcal{B}) is slightly degenerate, then any maximal *symmetric* subcategory $\mathcal{E} \subset \mathcal{B}$ satisfies $\mathrm{FPdim}(\mathcal{E})^2 \leq \mathrm{FPdim}(\mathcal{B})\mathrm{FPdim}(\mathcal{Z}_{sym}(\mathcal{B})) = 64$ and so $\mathrm{FPdim}(\mathcal{E}) \leq 8$. But $\mathcal E$ must contain $\mathcal Z_{sym}(\mathcal B) =$ svect, and so a maximal *Tannakian* subcategory of $\mathcal B$ has FP dimension at most 4 and, hence, is pointed. This implies that \mathcal{C} and (\mathcal{B}) is pointed.

Proposition 7.2. Let \mathcal{B} be a reductive category with $FPdim(\mathcal{B}) = p^n$, $n \leq 5$. Then either \mathcal{B} *is pointed or* $p = 2$ *and* \mathcal{B} *is a product of a pointed category and an Ising category.*

Proof. Recall three types of reductive categories introduced in Definition [5.18.](#page-21-1)

If B is of type I or II then by Theorem [5.20\(](#page-21-2)1,2), B is generated by $\text{Cond}(\mathcal{B})$ and $\text{Cond}(\mathcal{B})'$. Note that $\text{Cora}(B)$ is pointed by Lemma [7.1](#page-26-1) and $\text{Cora}(B)'$ is non-degenerate anisotropic. By [\[Na\]](#page-30-0), \mathcal{C} orad $(\mathcal{B})'$ is either pointed or is a product of a pointed braided category and an Ising category (for $p = 2$), which implies the statement in this case.

If B is of type III (so $p = 2$) then $\mathcal{C} \text{orad}(\mathcal{B})'$ must be pointed. Indeed, $\mathcal{C} \text{orad}(\mathcal{B})'$ is slightly degenerate anisotropic in this case. If it is not pointed, then it contains an Ising subcategory and a fermion centralizing it, which contradicts its anisotropy. Thus, the subcategory $\mathcal{B}_0 =$ Corad(B) \vee Corad(B)' \subset B is pointed. Therefore, \mathcal{B}_0 is split, i.e., $\mathcal{B}_0 = \tilde{\mathcal{B}}_0 \boxtimes$ sVect and by Theorem [5.20\(](#page-21-2)3), $\mathcal{B} = \tilde{\mathcal{B}}_0 \boxtimes \mathcal{C}$, where $\tilde{\mathcal{B}}_0$ is pointed and \mathcal{C} is a minimal extension of sVect (and so C is either pointed or Ising).

Corollary 7.3. An integral reductive category of dimension p^n , $n \leq 5$, is pointed.

It is straightforward to list equivalence classes of pointed reductive categories using results of Section [6.2.](#page-24-0) It is similarly not hard to list reductive categories that are products of Ising categories and pointed ones. Below we give a classification of *integral non-pointed* reductive categories of the Frobenius-Perron dimension p^6 . In view of Corollary [7.3](#page-26-2) this is the smallest dimension in which such categories exist.

For any prime p and $n \geq 1$ let E_{p^n} denote the elementary Abelian p-group of order p^n .

Lemma 7.4. Let $\mathcal B$ be an integral non-pointed reductive category with $FPdim(\mathcal B) = p^6$. Then $\mathcal{B} \cong \mathcal{Z}(\mathcal{V}ect_{E_{p^3}}^{\omega})$ for some $\omega \in H^3(E_{p^3}, k^{\times}).$

Proof. This statement is equivalent to β being non-degenerate and containing a Lagrangian subcategory equivalent to $\mathcal{R}ep(E_{p^3})$. Note that $\mathcal{C}ora\mathcal{A}(\mathcal{B})'$ is anisotropic integral and, hence, pointed. Let $\mathcal{E} \in T_{max}(\mathcal{B})$. We claim that $\text{FPdim}(\mathcal{E}) = p^3$. Indeed, if $\text{FPdim}(\mathcal{E}) \leq p^2$ then $\mathcal E$ is pointed and so both $\mathcal Conad}(\mathcal B)$ and $\mathcal B_0 = \mathcal Conad}(\mathcal B) \vee \mathcal Conad}(\mathcal B)'$ are pointed. We have either $\mathcal{B} = \mathcal{B}_0$ or \mathcal{B}_0 is slightly degenerate, and \mathcal{B} is its minimal extension. In the latter case β must contain an Ising subcategory, contradicting its integrality. On the other hand, there can be no symmetric subcategories $\mathcal{E} \subset \mathcal{B}$ with $FPdim(\mathcal{E}) \geq p^4$, since this would contradict β being almost non-degenerate. The last observation also implies that β is non-degenerate, since otherwise $\mathcal{E} \vee \mathcal{Z}_{sym}(\mathcal{B})$ is symmetric of dimension $\geq p^4$.

It follows from [\[DGNO2\]](#page-29-2), that $\mathcal{B} \cong \mathcal{Z}(\mathcal{V}ect_G^{\omega})$ for some group G of order p^3 . Since \mathcal{B} is reductive metabolic, it is a TG-category by Proposition [5.3\(](#page-18-4)ii). Since \mathcal{B} is non-pointed, it must contain non-pointed Lagrangian subcategories, so G can be chosen non-Abelian. For any such G its center is $Z(G) \cong E_p$ and the group of invertible objects in $\mathcal{R}ep(G)$ is $\widehat{G} \cong E_{p^2}$. Furthermore, the group $Inv(\mathcal{B})$ of invertible objects of \mathcal{B} fits into the exact sequence

(38)
$$
1 \to \widehat{G} \to \mathsf{Inv}(\mathcal{B}) \to Z(G),
$$

see, e.g., [\[GP,](#page-29-20) [MN\]](#page-29-21). In particular, $|\text{Inv}(\mathcal{B})| \leq p^3$. In fact, $|\text{Inv}(\mathcal{B})| = p^3$, since otherwise \mathcal{B}_{pt} is contained in every Lagrangian subcategory of \mathcal{B} , which is impossible since \mathcal{B} is reductive. For the same reason \mathcal{B}_{pt} must contain more than one Tannakian subcategory equivalent to $\mathcal{R}ep(E_{p^2})$. Therefore, $\mathcal{B}_{pt}=\mathcal{C}(E_{p^3}, q)$, where the pre-metric group (E_{p^3}, q) is isotropically generated and the set of its isotropic subgroups of order p^2 has a trivial intersection. This is only possible when $q = 1$, i.e., $\mathcal{B}_{pt} \cong \mathcal{R}ep(E_p)$ \Box

Lemma 7.5. Non-pointed braided fusion categories $\mathcal{Z}(\text{Vect}_{E_{p^3}}^{\omega_1})$ and $\mathcal{Z}(\text{Vect}_{E_{p^3}}^{\omega_2})$ are equiva*lent if and only if the cohomology classes of* ω_1, ω_2 *lie in the same orbit of* $GL_3(\mathbb{F}_p)$ *.*

Proof. Each of these categories has a unique pointed Lagrangian subcategory equivalent to $\mathcal{R}ep(E_{p^3})$ [\[MN\]](#page-29-21) and so a braided equivalence α between them restricts to a braided equiv-alence between these subcategories. By [\[MN,](#page-29-21) Theorem 3.3], this is equivalent to α being induced from a tensor equivalence between $\mathcal{V}ect_{E_{p^3}}^{\omega_1}$ and $\mathcal{V}ect_{E_{p^3}}^{\omega_2}$. Such an equivalence exists if an only if ω_2 is cohomologous to $f^*\omega_1$ for some automorphism f of E_p $3.$

Proposition 7.6. For any prime p there are precisely $p + 1$ braided equivalence classes of integral non-pointed reductive categories of the Frobenius-Perron dimension p^6 .

Proof. In view of Lemmas [7.4](#page-27-0) and [7.5,](#page-27-1) the number in question is equal to the number of $GL_3(\mathbb{F}_p)$ -orbits of $\omega \in H^3(E_{p^3}, k^{\times})$ such that $\mathcal{Z}(\mathcal{V}ect^{\omega}_{E_{p^3}})$ is reductive.

Suppose that p is odd. There is a $GL_3(\mathbb{F}_p)$ -equivariant isomorphism

$$
H^3(E_{p^3}, k^{\times}) \xrightarrow{\sim} \wedge^3(\mathbb{F}_p^3) \bigoplus \mathsf{S}^2(\mathbb{F}_p^3) : \omega \mapsto (\omega_{alt}, \omega_{sym}).
$$

The action of $A \in GL_3(\mathbb{F}_p)$ is given by

$$
A(\omega_{alt}, \omega_{sym}) = (\det(A)\omega_{alt}, A^t \omega_{sym} A),
$$

where we identify ω_{sym} with a symmetric matrix and A^t denotes the transpose of A. The category $\mathcal{Z}(\mathcal{V}ect_{E_{p3}}^{\omega})$ is non-pointed if and only if $\omega_{alt} \neq 0$ (i.e., $\omega_{alt} = \lambda \det, \lambda \in \mathbb{F}_p^{\times}$). In this case $\mathcal{Z}(\text{Vect}_{E_{p3}}^{\omega})_{pt} \cong \text{Rep}(E_{p^3})$, see [\[MN,](#page-29-21) Section 5.1]. Such a category $\mathcal{Z}(\text{Vect}_{E_{p^3}}^{\omega})$ is reductive if and only if E_{p^3} is generated by its subgroups N such that $\omega \in \Omega(E_{p^3}; N)$ [\(20\)](#page-11-2). This is equivalent to E_{p^3} being generated by its cyclic subgroups C such that $\omega_{sym}|_C = 1$. In other words, E_{p^3} is an isotropically generated metric group. There are precisely 4 congruence classes of symmetric 3×3 matrices ω_{sym} with this property

(39)
$$
A_0 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, A_1 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, A_2 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \text{ and } A_3 = \begin{pmatrix} \zeta & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix},
$$

where ζ is any quadratic non-residue modulo p. A description of $GL_3(\mathbb{F}_p)$ -orbits of the cohomology classes (λ det, ω_{sym}), $\lambda \neq 0$ is contained in [\[MCU\]](#page-29-22). These are (40)

 $\{(\lambda \det, A_0) \mid \lambda \in \mathbb{F}_p^{\times}\}, \{(\lambda \det, A_1) \mid \lambda \in \mathbb{F}_p^{\times}\}, \{(\pm \mu \det, A_2)\}, \{(\pm \mu \det, A_3)\}, \mu \in \mathbb{F}_p^{\times}.$ Thus, altogether there are $1 + 1 + \frac{p-1}{2} + \frac{p-1}{2} = p + 1$ orbits.

Now let $p = 2$. We can use the following description of the $GL_3(\mathbb{F}_2)$ -orbits on $H^3(E_8, k^{\times})$ from [\[GMN\]](#page-29-23). Namely, there is a bijection between $H^3(E_8, k^{\times})$ and subsets of the set S of order 2 subgroups of E_8 via

$$
\omega \mapsto \{C \in S \mid \text{ the class } \omega|_C \in H^3(C, k^{\times}) \text{ is non-trivial}\}.
$$

The corresponding subset of S is called the support of ω and denoted supp (ω) . The *weight* of ω is the cardinality $|\text{supp}(\omega)|$ of its support. The category $\mathcal{Z}(\text{Vect}_{E_8}^{\omega})$ is non-pointed if and only if the weight of ω is 1, 3, 5, or 7. The cohomology classes with the weight 1, 5, and 7 form three orbits, one for each value of the weight, while those of weight 3 form two orbits, see [\[GMN\]](#page-29-23). By Remark [4.8,](#page-13-1) the category $\mathcal{Z}(\mathcal{V}ect_{E_8}^{\omega})$ is reductive if and only if subgroups in S\supp(ω) generate E_8 , i.e., when the weight of ω is 1 or 3. Thus, there are 3 orbits in this \Box

Remark 7.7. Braided fusion categories from Proposition [7.6](#page-28-0) are metabolic in the sense of Definition [2.1.](#page-3-1)

REFERENCES

- [CGPW] S. Cui, C. Galindo, J. Plavnik, Z. Wang. On gauging symmetry of modular categories, Comm. Math. Phys., 348 (2016), 1043–1064.
- [DGNO1] V. Drinfeld, S. Gelaki, D. Nikshych, and V. Ostrik. Group-theoretical properties of nilpotent modular categories, [arXiv:0704.0195v](http://arxiv.org/abs/0704.0195)2 [math.QA] (2007).
- [DGNO2] V. Drinfeld, S. Gelaki, D. Nikshych, and V. Ostrik. On braided fusion categories I, Selecta Mathematica, 16 (2010), no. 1, 1 - 119.
- [DMNO] A. Davydov, M. Müger, D. Nikshych, and V. Ostrik, The Witt group of non-degenerate braided fusion categories, Journal für die reine und angewandte Mathematik, 677 (2013), 135–177.
- [DN1] A. Davydov, D. Nikshych, The Picard crossed module of a braided tensor category, Algebra and Number Theory, 7 (2013), no. 6, 1365–1403.
- [DN2] A. Davydov, D. Nikshych, Braided Picard groups and graded extensions of braided tensor categories, Sel. Math. New Ser.27, 65 (2021).
- [DNO] A. Davydov, D. Nikshych, V. Ostrik, On the structure of the Witt group of non-degenerate braided fusion categories, Selecta Mathematica 19 (2013), no. 1, 237–269.
- [EGNO] P. Etingof, S. Gelaki, D. Nikshych, and V. Ostrik, Tensor categories, Mathematical Surveys and Monographs, 205, American Mathematical Society (2015).
- [ENO1] P. Etingof, D. Nikshych, and V. Ostrik. On fusion categories, Annals of Mathematics 162 (2005), 581–642.
- [ENO2] P. Etingof, D. Nikshych, and V. Ostrik. Fusion categories and homotopy theory, Quantum Topology, 1 (2010), no. 3, 209-273.
- [ENO3] P. Etingof, D. Nikshych, and V. Ostrik, Weakly group-theoretical and solvable fusion categories, Adv. Math, 226 (2011), no. 1, 176–205
- [GP] C. Galindo, J. Plavnik, Tensor functor between Morita duals of fusion categories, Lett. Math. Phys. 107 (2017), 553–590. [arXiv:1407.2783](http://arxiv.org/abs/1407.2783) (2014).
- [GJ] C. Galindo and C. Jones, Equivariant fusion subcategories, [arXiv:2111.09116](http://arxiv.org/abs/2111.09116) [math.QA] (2021).
- [GN] S. Gelaki and D. Nikshych, Nilpotent fusion categories, Advances in Mathematics, 217 (2008), no. 3, 1053–1071.
- [GMN] C. Goff, G. Mason, S.-H. Ng, On the gauge equivalence of twisted quantum doubles of elementary abelian and extra-special 2-groups, Journal of Algebra 312 (2007), 849–875.
- [JFR] T. Johnson-Freyd, D. Reutter, Minimal nondegenerate extensions, Journal of the AMS, 37 (2024), no. 1, 81—150.
- [JMPP] C. Jones, S. Morrison, D. Penneys, J. Plavnik, Extension theory for braided-enriched fusion categories, International Mathematics Research Notices, 20 (2022), 15632—15683,
- [KK] A. Kawauchi and S. Kojima, Algebraic Classification of Linking Pairings on 3-Manifolds, Math. Ann. 253 (1980), 29-42.
- [MN] I. Marshall, D. Nikshych, On the Brauer-Picard groups of fusion categories, Mathematische Zeitschrift 288 (2018), 689-712.
- [MCU] K. Maya, A.M. Castaño, B. Uribe, Classification of Pointed Fusion Categories of dimension p3 up to weak Morita Equivalence, J. Algebra Appl. 20 (2021), no. 1, 2140008.
- [Mir] R. Miranda Nondegenerate Symmetric Bilinear Forms on Finite Abelian 2-Groups, Transactions of the American Mathematical Society, 284 (1984), no. 2, 535-542.
- [NN] D. Naidu, D. Nikshych, Lagrangian Subcategories and Braided Tensor Equivalences of Twisted Quantum Doubles of Finite Groups, Communications in Mathematical Physics, 279 (2008), 845-872.
- [N] D. Naidu, Categorical Morita equivalence for group-theoretical categories, Comm. Algebra 35 (2007), 3544–3565.
- [Na] S. Natale, On the core of a weakly group-theoretical braided fusion category, Internat. J. Math. 29 (2018), no. 2, 1850012.
- [U] B. Uribe, On the classification of pointed fusion categories up to weak Morita equivalence, Pacific Journal of Mathematics, 290 (2015), 437-466.
- [W] C.T.C. Wall, Quadratic forms on finite groups, and related topics, Topology, 2 (1963), 281–298.

Department of Mathematics and Statistics, University of New Hampshire, Durham, NH 03824, USA

Email address: jason.green@unh.edu

Department of Mathematics and Statistics, University of New Hampshire, Durham, NH 03824, USA

Email address: dmitri.nikshych@unh.edu