

THE GEOMETRY OF LOOP SPACES IV: CLOSED SASAKIAN MANIFOLDS

YOSHIAKI MAEDA AND STEVEN ROSENBERG

1. INTRODUCTION

It is well known that the dimension of the isometry group $\text{Isom}(M^n, g)$ of a closed Riemannian manifold (M^n, g) is at most $n(n+1)/2$, with the maximum achieved for the standard metric on S^n . Nevertheless, this large isometry group has small fundamental group: $\pi_1(\text{Isom}(S^{n+1}, g_{\text{std}})) = \mathbb{Z}_2$. (Throughout, all fundamental groups are based at the identity map.) In contrast, for a generic (M, g) , we expect that $\text{Isom}(M, g)$ is trivial. Thus it seems interesting to produce examples, other than flat tori, where $\pi_1(\text{Isom}(M, g))$ is infinite.

In this paper, we construct a one-parameter family of distinct metrics h_ρ on any closed Sasakian manifold M^{4k+1} such that $\pi_1(\text{Isom}(M, h_\rho))$ is infinite.

Theorem 5.1. *Let (M, g, ϕ, ξ, η) be a $(4k+1)$ -dimensional closed regular Sasakian manifold, where $k \geq 1$. The metric $h_\rho = g + \rho^2 \eta \otimes \eta$ has $|\pi_1(\text{Isom}(M, h_\rho))| = \infty$ for $\rho \in \mathbb{R}^+, \rho \gg 0$. For $\rho \neq \rho'$, h_ρ and $h_{\rho'}$ are not isometric.*

Theorem 5.1 will be proved in §5. We note that the proof does not work for $\dim(M) = 4k+3$.

For the case of S^{4k+1} , we refine Theorem 5.1:

Theorem 6.1. *Let $k \geq 1$. Let g_{std} be the standard metric on S^{4k+1} , with η the contact form on S^{4k+1} associated with the Hopf fibration. For the metric $h_\rho = g_{\text{std}} + \rho^2 \eta \otimes \eta$, $|\pi_1(\text{Isom}(S^{4k+1}, h_\rho))| = \infty$ for $\rho^2 \neq 0$.*

Thus the standard metric with small π_1 is the endpoint of a family with infinite π_1 .

Using these techniques, in §7 we obtain a similar result for an infinite dimensional group.

Theorem 7.1. *Let $\text{Diff}_{\eta, \text{str}}(S^{4k+1})$ be the group of strict contact diffeomorphisms of S^{4k+1} . Then $|\pi_1(\text{Diff}_{\eta, \text{str}}(S^{4k+1}))| = \infty$.*

In fact, a stronger result has been given in [3]; we reproduce this result in [6].

As an outline of the paper, in §2, we recall the setup of Sasakian manifolds. In §3, we outline the proof of Thm. 5.1, which is based on the Wodzicki-Chern-Simons classes on the loop space LM used in [7, 8]. In §4, we introduce h_ρ and compute its curvature. In §5 we prove that the pullback of the Wodzicki-Chern-Simons class on LM to a top class on M is nonvanishing for $\rho \gg 0$. This involves detailed calculations. By the results of §3, we obtain Thm. 5.1. In §6, we prove Thm. 6.1, using the special nature of the standard metric on the sphere. We obtain similar results for other Sasakian space forms. In §7, we prove Thm. 7.1.

2. SASAKIAN MANIFOLDS

We first recall the definition of a Sasakian manifold. We work in local coordinates in preparation for later calculations; coordinate-free versions of the definitions are in [2].

A $(2k + 1)$ -dimensional manifold M is called an *almost contact manifold* if there is an almost contact structure (ϕ, ξ, η) consisting of a $(1, 1)$ -tensor field ϕ , a *characteristic vector field* ξ , and a 1-form η satisfying

$$(1) \quad (\phi^2)_i^j = \phi_i^k \phi_k^j = -\delta_i^j + \eta_i \xi^j, \quad \eta_k \xi^k = 1, \quad \phi_j^k \xi^j = 0, \quad \phi_j^k \eta_k = 0.$$

((M, η) is a *contact manifold* if $\eta \wedge (d\eta)^k \neq 0$ pointwise.) Let g be a Riemannian metric on M associated with the almost contact structure (ϕ, ξ, η) , *i.e.*,

$$(2) \quad g_{ij} = g_{k\ell} \phi_i^k \phi_j^\ell + \eta_i \eta_j.$$

Such a metric always exists. Then (g, ϕ, ξ, η) is called an *almost contact metric structure* for the almost contact structure (ϕ, ξ, η) . We set

$$(3) \quad \phi_{ij} = \phi_i^k g_{kj}, \quad \phi^h_j = g^{h\ell} \phi_\ell^k g_{kj}.$$

We note that (2) implies

$$(4) \quad g_{ij} \xi^j = \eta_i,$$

so ξ and η are dual. Similarly, ϕ is skew-symmetric, since

$$\phi_{ji} = \phi_j^s g_{si} = \phi_j^s (g_{k\ell} \phi_s^k \phi_i^\ell + \eta_s \eta_i) = g_{k\ell} (-\delta_j^k + \eta_j \xi^k) \phi_i^\ell = -g_\ell \phi_i^\ell + \eta_j \eta_\ell \phi_i^\ell = -\phi_{ij}.$$

An almost contact metric structure is a *contact metric structure* (and (M, g, ϕ, ξ, η) is a *contact metric manifold*) if $\phi = -d\eta$, or equivalently,

$$(5) \quad \phi_{ij} = -\frac{1}{2}(\nabla_i \eta_j - \nabla_j \eta_i) := -\frac{1}{2}((\nabla_i \eta)_j - (\nabla_j \eta)_i),$$

where ∇ is the Levi-Civita connection for g . Finally, a contact metric manifold (M, g, ϕ, ξ, η) is *Sasakian* (or *normal*) if

$$(6) \quad \nabla_i \phi_j^k = g_{ij} \xi^k - \eta_j \delta_i^k.$$

As in [2, p. 87], writing (6) as $(\nabla_X \phi)(Y) = g(X, Y)\xi - \eta(Y)(X)$ and setting $Y = \xi$ gives $-\phi(\nabla_X \xi) = (\nabla_X \phi)(\xi) = \eta(X)\xi - X$. Apply ϕ to both sides, where $\phi^2(X) = -X + \eta(X)\xi$, and use $0 = g(\nabla_X \xi, \xi) = \eta(\nabla_X \xi)$ to obtain $\nabla_X \xi = -\phi(X)$. In coordinates, this becomes $\nabla_i \xi^j = -\phi_i^j$, or equivalently

$$(7) \quad -\phi_{ij} = \nabla_i \eta_j.$$

This refines (5) for Sasakian manifolds.

Since ϕ_{ij} is skew-symmetric, (7) implies

$$(8) \quad \nabla_i \eta_j + \nabla_j \eta_i = 0.$$

Therefore, $\xi^i = g^{ik} \eta_k$ is a unit Killing vector field on M which generates a one-parameter group of isometries.

The following result is standard.

Lemma 2.1. *If (M, g, ϕ, ξ, η) is a $(2k + 1)$ -dimensional contact metric manifold, then*

$$(9) \quad \eta \wedge (d\eta)^k = (-1)^k \eta \wedge \phi^k \neq 0.$$

$\eta \wedge (d\eta)^k$ is the volume form on M , so $\int_M \eta \wedge (d\eta)^k \neq 0$. Thus M is a contact manifold.

Proof. By (1) and (2), for $m \in M$ there is an orthonormal basis $\{e_1, \dots, e_{2k+1}\}$ on $T_m M$ such that $e_1 = \xi$, $\phi e_{1+\alpha} = e_{k+1+\alpha}$, $\phi e_{k+1+\alpha} = -e_{1+\alpha}$, for $\alpha = 1, \dots, k$. Thus,

$$\eta \wedge (d\eta)^k(e_1, e_2, \dots, e_{k+1}, e_{k+2}, \dots, e_{2k+1}) = \eta(e_1)g(\phi e_2, e_{k+1}) \cdots g(\phi e_{k+2}, e_{2k+1}) = 1 \neq 0.$$

□

A basic example of a Sasakian manifold is as follows.

Example 2.1. If $M = S^{2k+1}$, the $(2k+1)$ -dimensional standard sphere of radius 1, then M is a Sasakian manifold with respect to the Hopf fibration

$$(10) \quad \pi : S^{2k+1} \rightarrow \mathbb{C}\mathbb{P}^k,$$

where the characteristic vector field ξ is the unit vector along the fiber, and $\eta(\xi) = 1$ by (1).

In this paper, we assume that M is closed and *regular*, *i.e.*, the integral curves characteristic vector field ξ starting at any point $m \in M$ are periodic. By [2, p. 37], the period of the integral curves is independent of m . In this case, we have the circle fibration $S^1 \rightarrow M \rightarrow M/S^1$, and M/S^1 is a $2k$ -dimensional symplectic manifold. The example of the Hopf fibration is a special case. If ξ is regular, then there is a circle action

$$(11) \quad a : S^1 \times M \rightarrow M, \quad a(\theta, m) = R_\theta(m),$$

given by rotation by angle θ in the circle fiber. This action is via isometries for the metrics h_ρ below (see Cor. 4.1).

The Theorems in the Introduction entail proving that the loop of isometries $a^I : S^1 \rightarrow \text{Isom}(M, h_\rho)$, $a^I(\theta)(m) = a(\theta, m)$, has nontrivial class $[a^I] \in \pi_1(\text{Isom}(M, h_\rho))$ and that the powers $[a^I]^n$ are all distinct.

3. OUTLINE OF THE PROOFS

As explained in [8, §2.2], a Riemannian metric g on M determines the Wodzicki-Chern-Simons form $CS_g^W \in \Lambda^{2k-1}(LM)$ given by

$$(12) \quad CS_{2k-1}^W(g)(X_1, \dots, X_{2k-1}) \\ = \frac{k}{2^{k-2}} \sum_{\sigma} \text{sgn}(\sigma) \int_{S^1} \text{tr}[(R(X_{\sigma(1)}, \cdot)\dot{\gamma})(\Omega^M)^{k-1}(X_{\sigma(2)}, \dots, X_{\sigma(2k-1)})],$$

at the loop $\gamma \in LM$. Here $\sigma \in (1, \dots, 2k-1)$, the permutation group on $\{1, \dots, 2k-1\}$, R is the curvature tensor of g , Ω is the curvature two-form of g , and the $X_i \in T_\gamma LM$ are vector fields $X_{i,\theta} \in T_{\gamma(\theta)}M$ defined along γ . This form comes from replacing the usual trace in the definition of Chern-Simons classes with the Wodzicki residue, which is a trace on the Lie algebra of the structure group for natural metric connections on TLM .

Associated to the action a is the map $a^L : M \rightarrow LM$ from M to the free loop space $LM = C^\infty(S^1, M)$ given by $a^L(m)(\theta) = a(\theta, m)$. This produces a top form $a^{L,*}CS^W \in \Lambda^{2k-1}(M)$. In [4, Appendix A], we prove

Proposition 3.1.

$$(13) \quad \int_M a^{L,*}CS_{2k-1}^W(g) \neq 0 \Rightarrow 0 \neq [a^I] \in \pi_1(\text{Isom}(M, g)) \text{ and } [a^I]^n \neq [a^I]^m \text{ for } m \neq n.$$

for $g = h_\rho$.

The proof explicitly uses that the action is via isometries.

By the long calculation in §5, $a^{L,*}CS_{2k-1}^W(h_\rho)$ is a form-valued polynomial in ρ whose leading coefficient is a nonzero multiple of $\eta \wedge (d\eta)^k$ (see (19) and Prop. 5.1). Thus $\int_M a^{L,*}CS_{2k-1}^W(h_\rho)$ is a polynomial in ρ whose leading coefficient is a nonzero multiple of the h_ρ -volume of M . Therefore, $\int_M a^{L,*}CS_{2k-1}^W(h_\rho) \neq 0$ for $\rho \gg 0$, so Thm. 5.1 follows from (13). For the standard metric on S^{4k+1} , the form-valued polynomial consists of only one term, so $\int_M a^{L,*}CS_{2k-1}^W(h_\rho) \neq 0$ for all $\rho \neq 0$, which implies Thm. 5.1.

The tensorial calculation in §5 analyzes the integrand in (12), using the curvature calculations for h_ρ in §4. The integrand initially consists of an overwhelming number of terms, but the Bianchi identity and other symmetries are used repeatedly to eliminate most terms.

4. CURVATURE FORMULAS FOR THE METRIC h_ρ

Let (M, g, ϕ, η, ξ) be a regular Sasakian manifold. Define the new metric $h = h_\rho$ by

$$h_{ij} = g_{ij} + \rho^2 \eta_i \eta_j.$$

To motivate this metric, we recall from [8, §3.2] that a regular contact manifold M is the total space of an S^1 -bundle $L_k \xrightarrow{\pi} N$ over a Riemannian symplectic manifold (N, ω, g_N) with $[\omega] \in H^2(N, \mathbb{Z})$ and $c_1(L) = k\omega$ for our choice of $k \in \mathbb{Z}$. L_k has the contact metric $g_k = \pi^*g_N + \eta_k \otimes \eta_k$, where η_k is dual to ξ_k , the unit tangent vector field to the fibers. $L_1 \xrightarrow{\alpha} L_k$ is a k -fold cover, so g_k pulls back to a locally isometric metric on L_1 .

Noting that $\eta_1 = k\alpha^*\eta_k$, we have a \mathbb{Z} -family of metrics on L_1 of the form $\alpha^*g_k + k^2\eta_1 \otimes \eta_1$. This is a special case of h_ρ .

We will compute the Levi-Civita connection and the Riemannian curvature for h .

Lemma 4.1. *Let h^{ij} be the inverse of the metric h_{ij} . Then*

$$h^{ij} = g^{ij} + \alpha \xi^i \xi^j \text{ for } \alpha = -\frac{\rho^2}{1 + \rho^2}.$$

Proof.

$$h_{ik} h^{kj} = (g_{ik} + \rho^2 \eta_i \eta_k)(g^{kj} + \alpha \xi^k \xi^j) = \delta_i^j + (\rho^2 + \alpha + \rho^2 \alpha) \eta_i \xi^j = \delta_i^j,$$

for $\alpha = -\frac{\rho^2}{1 + \rho^2}$. □

Let Γ_{ij}^k and $\bar{\Gamma}_{ij}^k$ be the Christoffel symbols of g and h respectively:

$$\Gamma_{ij}^k = \frac{1}{2} g^{k\ell} (\partial_i g_{j\ell} + \partial_j g_{i\ell} - \partial_\ell g_{ij}), \quad \bar{\Gamma}_{ij}^k = \frac{1}{2} h^{k\ell} (\partial_i h_{j\ell} + \partial_j h_{i\ell} - \partial_\ell h_{ij}).$$

Lemma 4.2.

$$\bar{\Gamma}_{ij}^k = \Gamma_{ij}^k - \rho^2 (\phi_i^k \eta_j + \phi_j^k \eta_i).$$

Proof.

$$\begin{aligned} \bar{\Gamma}_{ij}^k &= \frac{1}{2} h^{k\ell} (\partial_i g_{j\ell} + \partial_j g_{i\ell} - \partial_\ell g_{ij}) + \frac{1}{2} \rho^2 h^{k\ell} (\partial_i (\eta_\ell \eta_j) + \partial_j (\eta_i \eta_\ell) - \partial_\ell (\eta_i \eta_j)) \\ &= \frac{1}{2} g^{k\ell} (\partial_i g_{j\ell} + \partial_j g_{i\ell} - \partial_\ell g_{ij}) + \frac{1}{2} \alpha \xi^k \xi^\ell (\partial_i g_{j\ell} + \partial_j g_{i\ell} - \partial_\ell g_{ij}) \\ &\quad + \frac{1}{2} g^{k\ell} \rho^2 (\partial_i (\eta_\ell \eta_j) + \partial_j (\eta_i \eta_\ell) - \partial_\ell (\eta_i \eta_j)) + \frac{1}{2} \alpha \rho^2 \xi^k \xi^\ell (\partial_i (\eta_\ell \eta_j) + \partial_j (\eta_i \eta_\ell) - \partial_\ell (\eta_i \eta_j)) \\ &:= \Gamma_{ij}^k + (b) + (c) + (d). \end{aligned}$$

Note that

$$\partial_i g_{\ell j} = \partial_i \langle \partial_\ell, \partial_j \rangle = \langle \nabla_i \partial_\ell, \partial_j \rangle + \langle \partial_\ell, \nabla_i \partial_j \rangle = \Gamma_{i\ell}^m g_{mj} + \Gamma_{ij}^m g_{\ell m}.$$

Using $\Gamma_{ij}^k = \Gamma_{ji}^k$, we have

$$\begin{aligned} (b) &= \frac{1}{2} \alpha \xi^k \xi^\ell (\partial_i g_{\ell j} + \partial_j g_{i\ell} - \partial_\ell g_{ij}) \\ &= \frac{1}{2} \alpha \xi^k \xi^\ell (\Gamma_{i\ell}^m g_{mj} + \Gamma_{ij}^m g_{\ell m} + \Gamma_{ji}^m g_{m\ell} + \Gamma_{j\ell}^m g_{im} - \Gamma_{\ell i}^m g_{mj} - \Gamma_{\ell j}^m g_{im}) \\ &= \alpha \xi^k \xi^\ell \Gamma_{ij}^m g_{\ell m} = \alpha \Gamma_{ij}^m \eta_m \xi^k, \end{aligned}$$

since $\xi^\ell g_{\ell m} = \eta_m$.

Using (5) and $\partial_i \eta_j = \nabla_i \eta_j + \Gamma_{ij}^k \eta_k$, we get

$$\begin{aligned} (c) &= \frac{\rho^2}{2} g^{k\ell} (\partial_i \eta_\ell \cdot \eta_j + \eta_\ell \cdot \partial_i \eta_j + \partial_j \eta_i \cdot \eta_\ell + \eta_i \cdot \partial_j \eta_\ell - \partial_\ell \eta_i \cdot \eta_j - \eta_i \cdot \partial_\ell \eta_j) \\ &= \frac{\rho^2}{2} g^{k\ell} (\nabla_i \eta_\ell \cdot \eta_j + \eta_\ell \cdot \nabla_i \eta_j + \nabla_j \eta_i \cdot \eta_\ell + \eta_i \cdot \nabla_j \eta_\ell - \nabla_\ell \eta_i \cdot \eta_j - \eta_i \cdot \nabla_\ell \eta_j \\ &\quad + \Gamma_{i\ell}^m \eta_m \eta_j + \eta_\ell \Gamma_{ij}^m \eta_m + \Gamma_{ji}^m \eta_m \eta_\ell + \eta_i \Gamma_{j\ell}^m \eta_m - \Gamma_{\ell i}^m \eta_m \eta_j - \eta_i \Gamma_{\ell j}^m \eta_m) \\ &= \frac{\rho^2}{2} g^{k\ell} (\nabla_i \eta_\ell \cdot \eta_j - \nabla_\ell \eta_i \cdot \eta_j + \eta_i \cdot \nabla_j \eta_\ell - \eta_i \cdot \nabla_\ell \eta_j + 2\Gamma_{ij}^m \eta_\ell \cdot \eta_m) \\ &= -\rho^2 (\phi_i^k \eta_j + \phi_j^k \eta_i) + \rho^2 \Gamma_{ij}^m \xi^k \eta_m, \end{aligned}$$

where (7) is used in the last line.

Similarly,

$$\begin{aligned} (d) &= \frac{1}{2} \alpha \rho^2 \xi^k \xi^\ell (\partial_i (\eta_\ell \eta_j) + \partial_j (\eta_i \eta_\ell) - \partial_\ell (\eta_i \eta_j)) = -\frac{\alpha}{2} \xi^k \xi^\ell \rho^2 (2\phi_{i\ell} \eta_j + 2\phi_{j\ell} \eta_i - 2\Gamma_{ij}^m \eta_\ell \eta_m) \\ &= \alpha \rho^2 \Gamma_{ij}^m \xi^k \eta_m, \end{aligned}$$

by (1).

Thus,

$$(b) + (c) + (d) = -\rho^2 (\phi_i^k \eta_j + \phi_j^k \eta_i) + (\alpha + \rho^2 + \alpha \rho^2) \Gamma_{ij}^m \xi^k \eta_m = -\rho^2 (\phi_i^k \eta_j + \phi_j^k \eta_i).$$

This gives the Lemma. \square

We want the action (11) to be an action by isometries for h_ρ . Since $\|\xi\|_{h_\rho}^2 = 1 + \rho^2$ by (1), (2) and the definition of h_ρ , $\bar{\xi} := (1 + \rho^2)^{-1/2} \xi$ has $\|\bar{\xi}\|_{h_\rho}^2 = 1$. We can consider (11) to be an action with $(d/d\theta)a(\theta, m) = \bar{\xi}$ by modifying the g -length of the orbits. If orbits have g -length λ , then these orbits now have h_ρ -length also equal to λ .

Corollary 4.1. *The S^1 -action $\{R_\theta\}$ in (11) is an action via isometries for h_ρ .*

Proof. We have $\bar{\eta} = \bar{\eta}_i dx^i$ is given by

$$\bar{\eta}_i = (h_\rho)_{ij} \bar{\xi}^j = (g_{ij} + \rho^2 \eta_i \eta_j) ((1 + \rho^2)^{-1/2} \xi^j) = (1 + \rho^2)^{1/2} \eta_i.$$

Let $\bar{\nabla}$ be the Levi-Civita connection for h_ρ . By Lem. 4.2,

$$\bar{\nabla}_i \bar{\eta}_j := (\bar{\nabla} \bar{\eta})_i = (1 + \rho^2)^{1/2} [\nabla_i \eta_j + \rho^2 (\phi_i^s \eta_j + \phi_j^s \eta_i) \eta_s] = (1 + \rho^2)^{1/2} (\nabla_i \eta_j),$$

so

$$\bar{\nabla}_i \bar{\eta}_j + \bar{\nabla}_j \bar{\eta}_i = (1 + \rho^2)^{1/2} (\nabla_i \eta_j + \nabla_j \eta_i) = 0.$$

Therefore, $\bar{\xi}$ is an h_ρ -unit Killing vector field and the rotation action is via isometry. \square

We now compute the curvature tensor for the metric h_ρ .

Lemma 4.3. *Let $\bar{R}_{kji}{}^h$ and $R_{kji}{}^h$ be the curvature tensor of the Riemannian metrics h_ρ and g respectively. Then*

$$(14) \quad \begin{aligned} \bar{R}_{kji}{}^h &= R_{kji}{}^h - \rho^2(\phi_{ki}\phi_j{}^h - \phi_k{}^h\phi_{ji} + 2\phi_{kj}\phi_i{}^h + 2\eta_k\eta_i\delta_j{}^h - 2\eta_j\eta_i\delta_k{}^h + g_{ki}\eta_j\xi^h - g_{ji}\eta_k\xi^h) \\ &\quad - \rho^4(\eta_k\eta_i\delta_j{}^h - \eta_j\eta_i\delta_k{}^h). \end{aligned}$$

Proof.

$$\begin{aligned} \bar{R}_{kji}{}^h &= \partial_k\bar{\Gamma}_{ji}{}^h - \partial_j\bar{\Gamma}_{ki}{}^h + \bar{\Gamma}_{kl}{}^h\bar{\Gamma}_{ji}{}^\ell - \bar{\Gamma}_{jl}{}^h\bar{\Gamma}_{ki}{}^\ell \\ &= \partial_k(\Gamma_{ji}{}^h - \rho^2(\phi_j{}^h\eta_i + \phi_i{}^h\eta_j)) - \partial_j(\Gamma_{ki}{}^h - \rho^2(\phi_k{}^h\eta_i + \phi_i{}^h\eta_k)) \\ &\quad + (\Gamma_{kl}{}^h - \rho^2(\phi_k{}^h\eta_\ell + \phi_\ell{}^h\eta_k))(\Gamma_{ji}{}^\ell - \rho^2(\phi_j{}^\ell\eta_i + \phi_i{}^\ell\eta_j)) \\ &\quad - (\Gamma_{jl}{}^h - \rho^2(\phi_j{}^h\eta_\ell + \phi_\ell{}^h\eta_j))(\Gamma_{ki}{}^\ell - \rho^2(\phi_k{}^\ell\eta_i + \phi_i{}^\ell\eta_k)) \end{aligned}$$

Therefore,

$$\begin{aligned} \bar{R}_{kji}{}^h &= R_{kji}{}^h - \rho^2(\partial_k(\phi_j{}^h\eta_i + \phi_i{}^h\eta_j) - \partial_j(\phi_k{}^h\eta_i + \phi_i{}^h\eta_k)) \\ &\quad + \rho^2(\Gamma_{kl}{}^h(\phi_j{}^\ell\eta_i + \phi_i{}^\ell\eta_j) + \Gamma_{ji}{}^\ell(\phi_k{}^h\eta_\ell + \phi_\ell{}^h\eta_k)) \\ &\quad - \rho^2(\Gamma_{jl}{}^h(\phi_k{}^\ell\eta_i + \phi_i{}^\ell\eta_k) + \Gamma_{ki}{}^\ell(\phi_j{}^h\eta_\ell + \phi_\ell{}^h\eta_j)) \\ &\quad + \rho^4(\phi_k{}^h\eta_\ell + \phi_\ell{}^h\eta_k)(\phi_j{}^\ell\eta_i + \phi_i{}^\ell\eta_j) - \rho^4(\phi_j{}^h\eta_\ell + \phi_\ell{}^h\eta_j)(\phi_k{}^\ell\eta_i + \phi_i{}^\ell\eta_k) \\ &= R_{kji}{}^h + (\rho_2) + (\rho_4) \end{aligned}$$

where (ρ_a) is the ρ^a -term of $\bar{R}_{kji}{}^h$. We have

$$\begin{aligned} (\rho_2) &= -[\rho^2(\partial_k(\phi_j{}^h\eta_i + \phi_i{}^h\eta_j) - \partial_j(\phi_k{}^h\eta_i + \phi_i{}^h\eta_k)) \\ &\quad + \rho^2(\Gamma_{kl}{}^h(\phi_j{}^\ell\eta_i + \phi_i{}^\ell\eta_j) + \Gamma_{ji}{}^\ell(\phi_k{}^h\eta_\ell + \phi_\ell{}^h\eta_k)) \\ &\quad - \rho^2(\Gamma_{jl}{}^h(\phi_k{}^\ell\eta_i + \phi_i{}^\ell\eta_k) + \Gamma_{ki}{}^\ell(\phi_j{}^h\eta_\ell + \phi_\ell{}^h\eta_j))] \\ &= -\rho^2[(\nabla_k\phi_j{}^h \cdot \eta_i + \phi_j{}^h \cdot \nabla_k\eta_i + \nabla_k\phi_i{}^h \cdot \eta_j + \phi_i{}^h \cdot \nabla_k\eta_j) \\ &\quad - \rho^2(\nabla_j\phi_k{}^h \cdot \eta_i + \phi_k{}^h \cdot \nabla_j\eta_i + \nabla_j\phi_i{}^h \cdot \eta_k + \phi_i{}^h \cdot \nabla_j\eta_k)], \end{aligned}$$

where we use

$$\nabla_k\eta_i := (\nabla_k\eta)_i = \partial_k\eta_i - \Gamma_{ki}{}^\ell\eta_\ell, \quad \nabla_k\phi_j{}^h := (\nabla_k\phi)_j{}^h = \partial_k\phi_j{}^h + \phi_j{}^\ell\Gamma_{k\ell}{}^h - \phi_\ell{}^h\Gamma_{kj}{}^\ell.$$

Using (6), (7), we have

$$(\rho_2) = -\rho^2(g_{ki}\eta_j\xi^h - g_{ji}\eta_k\xi^h - 2\delta_k{}^h\eta_j\eta_i + 2\delta_j{}^h\eta_k\eta_i + \phi_j{}^h\phi_{ki} + 2\phi_i{}^h\phi_{kj} - \phi_k{}^h\phi_{ji}).$$

For the ρ_4 -term, using (1), we easily get

$$\begin{aligned} (\rho_4) &= \rho^4(\phi_k{}^h\eta_\ell + \phi_\ell{}^h\eta_k)(\phi_j{}^\ell\eta_i + \phi_i{}^\ell\eta_j) - \rho^4(\phi_j{}^h\eta_\ell + \phi_\ell{}^h\eta_j)(\phi_k{}^\ell\eta_i + \phi_i{}^\ell\eta_k) \\ &= \rho^4(-\delta_j{}^h\eta_k\eta_i + \delta_k{}^h\eta_j\eta_i). \end{aligned}$$

Thus, we have the lemma. \square

5. COMPUTATION OF CS^W

In this section, we compute the Wodzicki-Chern-Simons form for (M, h_ρ) , or more precisely its pullback $(a^L)^*CS^W$ (19). As explained in Prop. 3.1, the nonvanishing of $(a^L)^*CS^W$ (Cor. 5.1) is the key step in the proofs of the main Thms. 5.1, 6.1.

Notation: We introduce some local coordinates notation. Let $\mathfrak{T}_P^Q(M)$ and $\Lambda^P(M)$ be the space of tensor fields of type (Q, P) and P -forms on M respectively. For an $(0, P)$ -tensor $S_{i_1 \dots i_P}$ and $(0, P')$ -tensor $S'_{i_1 \dots i_{P'}}$, its tensor product is

$$(S \otimes S')_{i_1 \dots i_{P+P'}} := S_{i_1 \dots i_P} \cdot S'_{i_{P+1} \dots i_{P+P'}} \in \mathfrak{T}_{P+P'}^0(M).$$

For an $(0, P)$ -tensor $S_{i_1 \dots i_P}$, its skew-symmetrization with respect to the indices i_1, \dots, i_P is denoted by

$$S_{[i_1 \dots i_P]} := \frac{1}{P!} \sum_{\sigma \in (1, 2, \dots, P)} \text{sgn}(\sigma) S_{i_{\sigma(1)} \dots i_{\sigma(P)}} \in \Lambda^P(M).$$

For $A \in \Lambda^P(M)$ and $A' \in \Lambda^{P'}(M)$, the exterior product is

$$(15) \quad \begin{aligned} (A \wedge A')_{i_1 \dots i_{P+P'}} &= A_{[i_1 \dots i_P} A'_{i_{P+1} \dots i_{P+P}]} \\ &:= \frac{1}{(P+P')!} \sum_{\sigma \in (1, \dots, P+P')} \text{sgn}(\sigma) A_{i_{\sigma(1)} \dots i_{\sigma(P)}} A'_{i_{\sigma(P+1)} \dots i_{\sigma(P+P')}}. \end{aligned}$$

tr means the contraction of the superscript index and the subscript index of the \mathfrak{T}_1^1 part of a tensor. For example, for $S = S_{i_1 \dots i_P k}^\ell \in \mathfrak{T}_P^0(M) \otimes \mathfrak{T}_1^1(M)$,

$$(16) \quad \text{tr}(S)_{i_1 \dots i_P} = S_{i_1 \dots i_P k}^k.$$

This notation extends to a “tensor and contract” operation on $\mathfrak{T}_P^0 \otimes \mathfrak{T}_1^1$. Namely, for $S_{i_1 \dots i_P \ell}^{\ell_\alpha} \in \mathfrak{T}_P^0 \otimes \mathfrak{T}_1^1$ and $S'_{i_1 \dots i_{P'} \ell_\beta}^{\ell'_\beta} \in \mathfrak{T}_{P'}^0 \otimes \mathfrak{T}_1^1$, set

$$(S \otimes S')_{i_1 \dots i_{P+P'} \ell_\beta}^{\ell_\alpha} = S_{i_1 \dots i_P \ell}^{\ell_\alpha} S'_{i_1 \dots i_{P'} \ell_\beta}^{\ell'_\beta} \in \mathfrak{T}_{P+P'}^0 \otimes \mathfrak{T}_1^1.$$

This similarly extends to a “wedge and contract” operation on $\Lambda^P(M) \otimes \mathfrak{T}_1^1$: For $A = A_{i_1 \dots i_P \ell}^{\ell_\alpha} \in \Lambda^P(M) \otimes \mathfrak{T}_1^1(M)$, and $B_{i_1 \dots i_{P'} \ell_\beta}^{\ell'_\beta} \in \Lambda^{P'}(M) \otimes \mathfrak{T}_1^1(M)$, set

$$\begin{aligned} (A \wedge B)_{i_1 \dots i_{P+P'} \ell_\beta}^{\ell_\alpha} &= (A \otimes B)_{[i_1 \dots i_{P+P'}] \ell_\beta}^{\ell_\alpha} \\ &= \frac{1}{(P+P')!} \sum_{\sigma \in (1, \dots, P+P')} \text{sgn}(\sigma) A_{i_{\sigma(1)} \dots i_{\sigma(P)} \ell}^{\ell_\alpha} B_{i_{\sigma(P+1)} \dots i_{\sigma(P+P')} \ell_\beta}^{\ell'_\beta} \\ &\in \Lambda^{P+P'}(M) \otimes \mathfrak{T}_1^1. \end{aligned}$$

For $S_{i_1 \dots i_P}, S'_{i_1 \dots i_P} \in \mathfrak{T}_P^0$, define an equivalence relation by

$$S_{i_1 \dots i_P} \sim S'_{i_1 \dots i_P} \text{ mod}(i_1 \dots i_P) \Leftrightarrow S_{[i_1 \dots i_P]} = S'_{[i_1 \dots i_P]},$$

and for $S_{i_1 \dots i_P \ell_0}^{\ell_1}, S'_{i_1 \dots i_P \ell_0}^{\ell_1} \in \mathfrak{T}_P^0 \otimes \mathfrak{T}_1^1$, similarly write

$$S_{i_1 \dots i_P \ell_0}^{\ell_1} \sim S'_{i_1 \dots i_P \ell_0}^{\ell_1} \text{ mod}(i_1 \dots i_P) \Leftrightarrow S_{[i_1 \dots i_P] \ell_0}^{\ell_1} = S'_{[i_1 \dots i_P] \ell_0}^{\ell_1}.$$

Let $LM = \{\gamma : S^1 \rightarrow M\}$ be the space of smooth loops on M . $T_\gamma LM = \{X_\gamma \in C^\infty(S^1, TM) : X_\gamma(\theta) \in T_{\gamma(\theta)}M\}$ is the set of smooth vector fields along γ .

We recall the definition of the Wodzicki-Chern-Simons class on LM [8]. For the metric h_ρ , set

$$(17) \quad \begin{aligned} K_{\nu\lambda_1 \dots \lambda_{2k+1} \kappa_0}^{\kappa_0'} &= \bar{R}_{\lambda_1 \kappa_1 \nu}^{\kappa_0'} \bar{R}_{\lambda_2 \lambda_3 \kappa_2}^{\kappa_1} \bar{R}_{\lambda_4 \lambda_5 \kappa_3}^{\kappa_2} \dots \bar{R}_{\lambda_{2k} \lambda_{2k+1} \kappa_0}^{\kappa_k} \in \Lambda^{2k+2}(M) \otimes \mathfrak{S}_1^1(M), \\ K_{\nu\lambda_1 \dots \lambda_{2k+1}} &= \text{tr}(K_{\nu\lambda_1 \dots \lambda_{2k+1} \kappa_0}^{\kappa_0'}) = K_{\nu\lambda_1 \dots \lambda_{2k+1} \kappa_0}^{\kappa_0}, \\ K_{\nu[\lambda_1 \dots \lambda_{2k+1}]} &\in \Lambda^1(M) \otimes \Lambda^{2k+2}(M). \end{aligned}$$

We define $CS^W \in \Omega^{2k+1}(LM)$ at the loop $\gamma \in LM$ by

$$(18) \quad CS^W(\gamma)(X_{\gamma,1}, \dots, X_{\gamma,2k+1}) = \int_0^{2\pi} K_{\nu[\lambda_1 \dots \lambda_{2k+1}]}(\gamma(\theta)) \dot{\gamma}^\nu(\theta) X_{\gamma,1}^{\lambda_1}(\theta) \dots X_{\gamma,2k+1}^{\lambda_{2k+1}}(\theta) d\theta.$$

Associated to a C^∞ map $a : S^1 \times M \rightarrow M$ are: (i) $a^D : S^1 \rightarrow \text{Maps}(M, M)$, $a^D(\theta)(m) := a(\theta, m)$; (ii) $a^L : M \rightarrow LM$, $a^L(m)(\theta) := a(\theta, m)$. If ξ is regular, the action (11) has $a^L(m)$ equal to the loop γ_m which is the circle fiber containing m and starting at m . From now on, we use this action.

We want to compute $(a^L)^*CS^W \in \Omega^{2k+1}(M)$. We have

$$((a^L)^*CS^W)_m = (a^L)^*CS^W(\partial_{x^{i_1}}, \dots, \partial_{x^{i_{2k+1}}})_m dx^{i_1} \wedge \dots \wedge dx^{i_{2k+1}}.$$

In our case, a is rotation in the S^1 fiber, and

$$\begin{aligned} &(a^L)^*CS^W(\partial_{x^{i_1}}, \dots, \partial_{x^{i_{2k+1}}})_m \\ &= \int_0^{2\pi} K_{\nu[\lambda_1 \dots \lambda_{2k+1}]}(a(\theta, m)) \frac{d}{d\theta} \Big|_{\theta} a^\nu(\theta, m) (a_*^L(\partial_{x^{i_1}}|_m))^{\lambda_1} \dots (a_*^L(\partial_{x^{i_{2k+1}}}|_m))^{\lambda_{2k+1}} d\theta \\ &= \int_0^{2\pi} K_{\nu[\lambda_1 \dots \lambda_{2k+1}]}(a(\theta, m)) \frac{\partial a^\nu}{\partial x^j} \Big|_{(\theta, m)} \xi^j|_m \frac{\partial a^{\lambda_1}}{\partial x^{i_1}} \Big|_{(\theta, m)} \dots \frac{\partial a^{\lambda_{2k+1}}}{\partial x^{i_{2k+1}}} \Big|_{(\theta, m)} d\theta, \end{aligned}$$

where $m = (x^1, \dots, x^{2k+1})$. In the last line, we use that a is an action, *i.e.*, $a(\theta + \psi, m) = a(\theta, a(\psi, m))$, to compute $(d/d\theta)a^\nu$.

For the action (11), $a^D : S^1 \rightarrow \text{Isom}(M, h)$, so by the tensorial nature of the curvature tensor, we have

$$K_{\nu\lambda_1 \dots \lambda_{2k+1}}(a(\theta, m)) \frac{\partial a^\nu}{\partial x^j} \Big|_{(\theta, m)} \xi^j|_m \frac{\partial a^{\lambda_1}}{\partial x^{i_1}} \Big|_{(\theta, m)} \dots \frac{\partial a^{\lambda_{2k+1}}}{\partial x^{i_{2k+1}}} \Big|_{(\theta, m)} = K_{j i_1 \dots i_{2k+1}} \xi^j(m),$$

for all θ . Thus, we have

$$(19) \quad \begin{aligned} ((a^L)^*CS^W)(m) &= \left(\int_0^{2\pi} K_{j[i_1 \dots i_{2k+1}]}(m) \xi^j(m) d\theta \right) dx^{i_1} \wedge \dots \wedge dx^{i_{2k+1}} \\ &= K_{j[i_1 \dots i_{2k+1}]}(m) \xi^j(m) dx^{i_1} \wedge \dots \wedge dx^{i_{2k+1}}. \end{aligned}$$

As explained at the beginning of this section, our goal is to show that $((a^L)^*CS^W)(m) \neq 0$ for h_ρ . This follows from the following Proposition, where it is more convenient to use indices i_1, \dots, i_{4k+1} .

Proposition 5.1. *Let $\dim(M) = 4k + 1$. For $K_{j[i_1 \dots i_{4k+1}]}$ given by (17), we have*

$$K_{j[i_1 \dots i_{4k+1}]} \xi^j = -(1 + \rho^2)^2 (a_{4k} \rho^{4k} + a_{4k-2} \rho^{4k-2} \dots + a_0),$$

for

$$\begin{aligned} a_{4k} &= (-1)^k 4^k (4k + 2) \cdot \eta_{[i_1 \phi_{i_2 i_3} \cdots \phi_{i_{4k} i_{4k+1}}]} \\ &= (-1)^k 4^k (4k + 2) (\eta \wedge (d\eta)^k)_{i_1 \cdots i_{4k+1}}, \end{aligned}$$

in the notation of (15).

We postpone the proof for the moment. From now on, we write $\eta \wedge (d\eta)^k$ just as $\eta \wedge d\eta^k$.

Corollary 5.1. *Let $\dim(M) = 4k + 1$. For $m \in M$ and $\rho \gg 0$, $((a^L)^*CS^W)(m) \neq 0$ for the metric h_ρ .*

Proof of Corollary 5.1. By Lem. 2.1, $a_{4k} \neq 0$. For $\rho \gg 0$, $K_{j[1 \cdots 4k+1]} \xi^j \neq 0$ by Prop 5.1. It follows from (19) that $((a^L)^*CS^W)(m) \neq 0$. \square

Using the Corollary, we can prove the main theorem:

Theorem 5.1. *Let (M, g, ϕ, ξ, η) be $(4k + 1)$ -dimensional closed regular Sasakian manifold, where $k \geq 1$. Then, there is a metric of the form $h_\rho = g + \rho^2 \eta \otimes \eta$ such that $|\pi_1(\text{Isom}(M, h_\rho))| = \infty$ for $\rho \in \mathbb{R}, \rho \gg 0$. For $\rho \neq \rho'$, h_ρ and $h_{\rho'}$ are not isometric.*

Proof of Theorem 5.1. By Cor. 5.1, $(a^L)^*CS^W$ for h_ρ is pointwise nonzero for $\rho \gg 0$. Since the rotation action (11) is an action by isometries for h_ρ (Cor. 4.1), then [8, Prop. 5.2] (as corrected in [7, Prop. 2.2]) implies that $\pi_1(\text{Isom}(M, h_\rho))$ is infinite for $\rho \gg 0$.

The curvature formula in Lem. 4.3 shows that h_ρ and $h_{\rho'}$ are not isometric for $\rho \gg 0$. \square

The rest of the section is devoted to the proof of Prop. 5.1. As an outline, by (24), $K_{j[i_1 \cdots i_{4k+1}]} \xi^j = -(1 + \rho^2) \eta_{[i_1} (\text{tr}(K1))_{i_2 \cdots i_{2k+1}]}$, where (K1) is defined in (22). By the notation defined in (25), (26), it suffices to prove (27). This is proven in Corollary 5.2(ii).

Proof of Proposition 5.1. We first work in an arbitrary odd dimension $2k + 1$. Recall that

$$(20) \quad K_{j i_1 \cdots i_{2k+1} \ell'_0} \xi^j = \bar{R}_{i_1 \ell_1 j}^{\ell_0} \bar{R}_{i_2 i_3 \ell_2}^{\ell_1} \bar{R}_{i_4 i_5 \ell_3}^{\ell_2} \cdots \bar{R}_{i_{2k} i_{2k+1} \ell'_0}^{\ell_k} \xi^j.$$

(6) implies $\nabla_j \xi^k = \phi_j^k [2, \text{p. 87}]$, so

$$\begin{aligned} (21) \quad R_{kji}^h \xi^i &= \nabla_k \nabla_j \xi^h - \nabla_j \nabla_k \xi^h = -(\nabla_k \phi_j^h - \nabla_j \phi_k^h) \\ &= -[(g_{kj} \xi^h - \eta_j \delta_k^h) - (g_{jk} \xi^h - \eta_k \delta_j^h)] \\ &= -(\eta_k \delta_j^h - \eta_j \delta_k^h). \end{aligned}$$

By Lem. 4.3, we get

$$\begin{aligned} \bar{R}_{kji}^h \xi^i &= R_{kji}^h \xi^i - \rho^2 (\phi_{ki} \phi_j^h - \phi_k^h \phi_{ji} + 2\phi_{kj} \phi_i^h + 2\eta_k \eta_i \delta_j^h - 2\eta_j \eta_i \delta_k^h + g_{ki} \eta_j \xi^h - g_{ji} \eta_k \xi^h) \xi^i \\ &\quad - \rho^4 (\eta_k \eta_i \delta_j^h - \eta_j \eta_i \delta_k^h) \xi^i \\ &= -(\eta_k \delta_j^h - \eta_j \delta_k^h) - 2\rho^2 (\eta_k \delta_j^h - \eta_j \delta_k^h) - \rho^4 (\eta_k \delta_j^h - \eta_j \delta_k^h) \\ &= -(1 + \rho^2)^2 (\eta_k \delta_j^h - \eta_j \delta_k^h), \end{aligned}$$

where we use (21) for the ρ^0 term, (1) and (2) for the ρ^2 term, and (1) for the ρ^4 term. Then

$$\begin{aligned} K_{j i_1 \cdots i_{2k+1} \ell'_0} \xi^j &= \xi^j \bar{R}_{i_1 \ell_1 j}^{\ell_0} \bar{R}_{i_2 i_3 \ell_2}^{\ell_1} \bar{R}_{i_4 i_5 \ell_3}^{\ell_2} \cdots \bar{R}_{i_{2k} i_{2k+1} \ell'_0}^{\ell_k} \xi^j \\ &= -(1 + \rho^2)^2 (\eta_{i_1} \delta_{\ell_1}^{\ell_0} - \eta_{\ell_1} \delta_{i_1}^{\ell_0}) \bar{R}_{i_2 i_3 \ell_2}^{\ell_1} \bar{R}_{i_4 i_5 \ell_3}^{\ell_2} \cdots \bar{R}_{i_{2k} i_{2k+1} \ell'_0}^{\ell_k} \\ &:= (\text{K1})_{i_1 \cdots i_{2k+1} \ell'_0}^{\ell_0} + (\text{K2})_{i_1 \cdots i_{2k+1} \ell'_0}^{\ell_0}, \end{aligned}$$

where

$$(22) \quad \begin{aligned} (\text{K1})_{i_1 \dots i_{2k+1} \ell_0^{\ell_0}} &= -(1 + \rho^2)^2 (\eta_{i_1} \delta_{\ell_1}^{\ell_0}) \bar{R}_{i_2 i_3 \ell_2}^{\ell_1} \bar{R}_{i_4 i_5 \ell_3}^{\ell_2} \dots \bar{R}_{i_{2k} i_{2k+1} \ell_0^{\ell_0}}^{\ell_k}, \\ (\text{K2})_{i_1 \dots i_{2k+1} \ell_0^{\ell_0}} &= (1 + \rho^2)^2 (\eta_{\ell_1} \delta_{i_1}^{\ell_0}) \bar{R}_{i_2 i_3 \ell_2}^{\ell_1} \bar{R}_{i_4 i_5 \ell_3}^{\ell_2} \dots \bar{R}_{i_{2k} i_{2k+1} \ell_0^{\ell_0}}^{\ell_k}. \end{aligned}$$

As in (16), we have

$$\begin{aligned} \text{tr}(\text{K1})_{i_1 \dots i_{2k+1}} &= -(1 + \rho^2)^2 \eta_{i_1} \bar{R}_{i_2 i_3 \ell_2}^{\ell_1} \bar{R}_{i_4 i_5 \ell_3}^{\ell_2} \dots \bar{R}_{i_{2k} i_{2k+1} \ell_1}^{\ell_k}, \\ \text{tr}(\text{K2})_{i_1 \dots i_{2k+1}} &= (1 + \rho^2)^2 (\eta_{\ell_1}) \bar{R}_{i_2 i_3 \ell_2}^{\ell_1} \bar{R}_{i_4 i_5 \ell_3}^{\ell_2} \dots \bar{R}_{i_{2k} i_{2k+1} i_1}^{\ell_k}. \end{aligned}$$

We continue the proof of Prop. 5.1 with a series of lemmas. Since the Proposition is concerned with the skew-symmetrization of $K \cdot \xi$, we consider the skew-symmetric parts of $\text{tr}(\text{K1})_{i_1 \dots i_{2k+1}}$ and $\text{tr}(\text{K2})_{i_1 \dots i_{2k+1}}$. The following proof is the first instance of our repeated use of the Bianchi identity.

Lemma 5.1.

$$(23) \quad \text{tr}(\text{K2})_{[i_1 \dots i_{2k+1}]} = 0.$$

Proof. Set

$$(\text{K2-1})_{i_2 \dots i_{2k-1} \ell_k} = \eta_{\ell_1} \bar{R}_{i_2 i_3 \ell_2}^{\ell_1} \bar{R}_{i_4 i_5 \ell_3}^{\ell_2} \dots \bar{R}_{i_{2k-2} i_{2k-1} \ell_k \ell_{k-1}},$$

and

$$\hat{R}_{i_{2k-2} i_{2k-1} i_1}^{\ell_k} = \bar{R}_{[i_{2k-2} i_{2k-1} i_1]}^{\ell_k}.$$

Then

$$\text{tr}(\text{K2})_{[i_1 \dots i_{2k+1}]} = \text{tr}((\text{K2-1}) \wedge \bar{R})_{[i_1 \dots i_{2k-1}]}.$$

Note that $\bar{R}_{[i_{2k-2} i_{2k-1} i_1]}^{\ell_k} = 0$ by the Bianchi identity. Thus, $\text{tr}(\text{K2})_{[i_1 \dots i_{2k+1}]} = 0$. \square

Therefore,

$$(24) \quad K_{j[i_1 \dots i_{2k+1}]} \xi^j = -(1 + \rho^2) \eta_{[i_1} (\text{tr}(\text{K1}))_{i_2 \dots i_{2k+1}]}$$

From now on, we focus on $\dim(M) = 4k + 1$. Let

$$(25) \quad N_{i_2 i_3 \dots i_{4k} i_{4k+1} \ell_0} = \bar{R}_{i_2 i_3 \ell_2}^{\ell_0} \bar{R}_{i_4 i_5 \ell_3}^{\ell_2} \dots \bar{R}_{i_{4k} i_{4k+1} \ell_1}^{\ell_{2k}}.$$

Then

$$(26) \quad \begin{aligned} (\text{K1})_{j[i_1 i_2 \dots i_{4k+1}]} \xi^j &= -(1 + \rho^2)^2 (\text{K1})_{[i_1 i_2 \dots i_{4k+1}]} \ell^\ell = -(1 + \rho^2)^2 \eta_{[i_1} N_{i_2 \dots i_{4k+1}]} \ell^\ell \\ &= -(1 + \rho^2)^2 \sum_{a=1}^{4k+1} (-1)^a \eta_{i_a} N_{[i_1 \dots \hat{i}_a \dots i_{4k+1}]} \ell^\ell, \end{aligned}$$

where \hat{i}_a means i_a is omitted. To get Proposition 5.1, it suffices to prove

$$(27) \quad \begin{aligned} \text{tr}(N)_{[i_2 \dots i_{4k+1}]} &= N_{[i_2 \dots i_{4k+1}]} \ell^\ell = b_{4k} \rho^{4k} + \dots + b_0, \\ \text{for } b_{4k} &= (-1)^k 4^k (4k + 2) (d\eta \wedge d\eta^k)_{i_2 \dots i_{4k+1}}. \end{aligned}$$

We regroup $N_{i_2 i_3 \dots i_{4k} i_{4k+1} \ell_0}^{\ell_1}$ as

$$(28) \quad \begin{aligned} &N_{i_2 i_3 \dots i_{4k} i_{4k+1} \ell_0}^{\ell_1} \\ &= A_{i_2 i_3 i_4 i_5 \ell_3}^{\ell_1} \cdot A_{i_6 i_7 i_8 i_9 \ell_5}^{\ell_3} \dots A_{i_{4k-6} i_{4k-5} i_{4k-4} i_{4k-3} \ell_{2k-1}}^{\ell_{2k-3}} \cdot A_{i_{4k-2} i_{4k-1} i_{4k} i_{4k+1} \ell_0}^{\ell_{2k-1}}, \end{aligned}$$

where

$$(29) \quad A_{i_{4a-2}i_{4a-1}i_{4a}i_{4a+1}\ell_{2a+1}}^{\ell_{2a-1}} := \bar{R}_{i_{4a-2}i_{4a-1}\ell_{2a}}^{\ell_{2a-1}} \cdot \bar{R}_{i_{4a}i_{4a+1}\ell_{2a+1}}^{\ell_{2a}}, \quad a = 1, \dots, k,$$

(with ℓ_{2k+1} replaced by ℓ_1 on the left hand side of (29) if $a = k$.)

We emphasize that this regrouping is not possible if $\dim(M) = 4k + 3$.

By Lem. 4.3, we have

$$\bar{R}_{kji}^h = (\text{R1})_{kji}^h + (\text{R2})_{kji}^h + (\text{R3})_{kji}^h,$$

where

$$(30) \quad \begin{aligned} (\text{R1})_{kji}^h &= R_{kji}^h, \\ (\text{R2})_{kji}^h &= -\rho^2(\phi_{ki}\phi_j^h - \phi_k^h\phi_{ji} + 2\phi_{kj}\phi_i^h + 2\eta_k\eta_i\delta_j^h - 2\eta_j\eta_i\delta_k^h + g_{ki}\eta_j\xi^h - g_{ji}\eta_k\xi^h), \\ (\text{R3})_{kji}^h &= \rho^4(\delta_k^h\eta_j\eta_i - \delta_j^h\eta_i\eta_k). \end{aligned}$$

In the notation of (30) (and relaxing the sub/superscripts on A), we have

$$(31) \quad A_{i_a i_{a+1} i_{a+2} i_{a+3} \ell_\beta}^{\ell_\alpha} = [(\text{R1})_{i_a i_{a+1} \ell_\gamma}^{\ell_\alpha} + (\text{R2})_{i_a i_{a+1} \ell_\gamma}^{\ell_\alpha} + (\text{R3})_{i_a i_{a+1} \ell_\gamma}^{\ell_\alpha}] \\ \cdot [(\text{R1})_{i_{a+2} i_{a+3} \ell_\beta}^{\ell_\gamma} + (\text{R2})_{i_{a+2} i_{a+3} \ell_\beta}^{\ell_\gamma} + (\text{R3})_{i_{a+2} i_{a+3} \ell_\beta}^{\ell_\gamma}].$$

Thus, we will compute $(\text{RE})_{i_a i_{a+1} \ell_\gamma}^{\ell_\alpha} \otimes (\text{RF})_{i_{a+2} i_{a+3} \ell_\beta}^{\ell_\gamma}$ for $E, F = 1, 2, 3$. For ease of notation, we will just compute $(\text{RE})_{i_2 i_3 \ell_2}^{\ell_1} \otimes (\text{RF})_{i_4 i_5 \ell_3}^{\ell_2}$, as this easily extends to the general case.

The Bianchi identity extends to each component of the curvature tensor.

Lemma 5.2.

$$(32) \quad (\text{RE})_{[kji]}^h = 0,$$

for $E = 1, 2, 3$.

Proof. The curvature tensor \bar{R}_{kji}^h of h_ρ has the Bianchi identity for each value of ρ . Letting ρ vary in (14), we must have the Bianchi identity for each RE. \square

We now eliminate curvature terms involving (R3).

Lemma 5.3.

$$(33) \quad ((\text{RE}) \otimes (\text{R3}))_{[i_2 i_3 i_4 i_5] \ell_3}^{\ell_1} = ((\text{R3}) \otimes (\text{RE}))_{[i_2 i_3 i_4 i_5] \ell_3}^{\ell_1} = 0,$$

for $E = 1, 2, 3$.

Proof. Ignoring powers of ρ , we have

$$\begin{aligned} (\text{RE})_{i_2 i_3 \ell_2}^{\ell_1} \cdot (\text{R3})_{i_4 i_5 \ell_3}^{\ell_2} &= (\text{RE})_{i_2 i_3 \ell_2}^{\ell_1} (\delta_{i_4}^{\ell_2} \eta_{i_5} \eta_{\ell_2} - \delta_{i_5}^{\ell_2} \eta_{i_4} \eta_{\ell_2}) \\ &= (\text{RE})_{i_2 i_3 i_4}^{\ell_1} \eta_{i_5} - (\text{RE})_{i_2 i_3 i_5}^{\ell_1} \eta_{i_4}. \end{aligned}$$

Using Lemma 5.2, we have

$$((\text{RE}) \otimes (\text{R3}))_{[i_2 i_3 i_4 i_5] \ell_3}^{\ell_1} = \eta_{i_5} (\text{RE})_{i_2 i_3 i_4}^{\ell_1} - \eta_{i_4} (\text{RE})_{i_2 i_3 i_5}^{\ell_1} = 0.$$

The second equation of 33 is obtained similarly. \square

We recall the regrouping of $N_{i_2 i_3 \dots i_{4k} i_{4k+1}}^{\ell_1 \ell_0}$ in (28) and that each $A_{i_{4a-2} i_{4a-1} i_{4a} i_{4a+1} \ell_{2a+1}}^{\ell_{2a-1}}$ is composed by (RE)'s.

For $a > 4k$, the terms in ρ^{2a} of $N_{i_2 i_3 \dots i_{4k} i_{4k+1}}^{\ell_1 \ell_0}$ should contain at least one (R3).

Thus, by Lemma 5.3, we have

$$(34) \quad \begin{aligned} N_{[i_2 i_3 \dots i_{4k} i_{4k+1}] \ell_0}^{\ell_1} &= \underbrace{((\text{R2}) \otimes \dots \otimes (\text{R2}))}_{2k} [i_2 i_3 \dots i_{4k} i_{4k+1}] \ell_0^{\ell_1} + l.o. \\ &= \underbrace{((\text{R2}) \wedge \dots \wedge (\text{R2}))}_{2k} i_2 i_3 \dots i_{4k} i_{4k+1} \ell_0^{\ell_1} + l.o. \end{aligned}$$

where $l.o.$ denotes the lower order terms in ρ^{2a} , $2a \leq 4k - 2$.

We have

$$(\text{R2})_{i_2 i_3 \ell_2}^{\ell_1} = -\rho^2((\text{R2-1})_{i_2 i_3 \ell_2}^{\ell_1} + (\text{R2-2})_{i_2 i_3 \ell_2}^{\ell_1}),$$

where

$$\begin{aligned} (\text{R2-1})_{i_2 i_3 \ell_2}^{\ell_1} &= \phi_{i_2 \ell_2} \phi_{i_3}^{\ell_1} - \phi_{i_2}^{\ell_1} \phi_{i_3 \ell_2} + 2\phi_{i_2 i_3} \phi_{\ell_2}^{\ell_1}, \\ (\text{R2-2})_{i_2 i_3 \ell_2}^{\ell_1} &= 2(\eta_{i_2} \delta_{i_3}^{\ell_1} - \eta_{i_3} \delta_{i_2}^{\ell_1}) \eta_{\ell_2} + (g_{i_2 \ell_2} \eta_{i_3} - g_{i_3 \ell_2} \eta_{i_2}) \xi^{\ell_1}. \end{aligned}$$

Lemma 5.4.

$$(35) \quad \begin{aligned} ((\text{R2-1}) \otimes (\text{R2-2}))_{[i_2 i_3 i_4 i_5] \ell_3}^{\ell_1} &= 0, \\ ((\text{R2-2}) \otimes (\text{R2-1}))_{[i_2 i_3 i_4 i_5] \ell_3}^{\ell_1} &= 0, \\ ((\text{R2-2}) \otimes (\text{R2-2}))_{[i_2 i_3 i_4 i_5] \ell_3}^{\ell_1} &= 0. \end{aligned}$$

Proof. A direct check gives

$$(36) \quad (\text{R2-1})_{[kji]}^h = 0, (\text{R2-2})_{[kji]}^h = 0.$$

For the first equation of (35), we get

$$\begin{aligned} (\text{R2-1})_{i_2 i_3 \ell_2}^{\ell_1} \times (\text{R2-2})_{i_4 i_5 \ell_3}^{\ell_2} &= (\text{R2-1})_{i_2 i_3 \ell_2}^{\ell_1} \times 2(\eta_{i_4} \delta_{i_5}^{\ell_2} - \eta_{i_5} \delta_{i_4}^{\ell_2}) \eta_{\ell_3} \\ &= 2(\eta_{i_4} (\text{R2-1})_{i_2 i_3 i_5}^{\ell_1} - \eta_{i_5} (\text{R2-1})_{i_2 i_3 i_4}^{\ell_1}) \eta_{\ell_3}, \end{aligned}$$

so

$$((\text{R2-1}) \otimes (\text{R2-2}))_{[i_2 i_3 i_4 i_5] \ell_3}^{\ell_1} = 2(\eta_{i_4} (\text{R2-1})_{i_2 i_3 i_5}^{\ell_1} - \eta_{i_5} (\text{R2-1})_{i_2 i_3 i_4}^{\ell_1}) = 0.$$

For the second equation of (35), we have

$$(\text{R2-2})_{i_2 i_3 \ell_2}^{\ell_1} \times (\text{R2-1})_{i_4 i_5 \ell_3}^{\ell_2} = ((\text{R2-1})_{i_4 i_5 \ell_3 i_2} \eta_{i_3} - (\text{R2-1})_{i_4 i_5 \ell_3 i_3} \eta_{i_2}) \xi^{\ell_1}.$$

so

$$((\text{R2-2}) \otimes (\text{R2-1}))_{[i_2 i_3 i_4 i_5] \ell_3}^{\ell_1} = -(\text{R2-1})_{[i_4 i_5 i_2] \ell_3}^{\ell_1} \eta_{[i_3]} + (\text{R2-1})_{[i_4 i_5 i_3] \ell_2}^{\ell_1} \eta_{[i_2]},$$

with the alternation on the right hand side over i_2, i_3, i_4, i_5 . By (36),

$$-(\text{R2-1})_{[i_4 i_5 i_2] \ell_3}^{\ell_1} \eta_{[i_3]} = (\text{R2-1})_{[i_4 i_5 i_3] \ell_2}^{\ell_1} \eta_{[i_2]} = 0.$$

For the third equation of (35), we get

$$\begin{aligned}
& (\text{R2-2})_{i_2 i_3 \ell_2}^{\ell_1} \times (\text{R2-2})_{i_4 i_5 \ell_3}^{\ell_2} \\
&= (2(\eta_{i_2} \delta_{i_3}^{\ell_1} - \eta_{i_3} \delta_{i_2}^{\ell_1}) \eta_{\ell_2} + (g_{i_2 \ell_2} \eta_{i_3} - g_{i_3 \ell_2} \eta_{i_2}) \xi^{\ell_1}) \\
&\quad \times (2(\eta_{i_4} \delta_{i_5}^{\ell_2} - \eta_{i_5} \delta_{i_4}^{\ell_2}) \eta_{\ell_3} + (g_{i_4 \ell_3} \eta_{i_5} - g_{i_5 \ell_3} \eta_{i_4}) \xi^{\ell_2}) \\
(37) \quad &= 2(\eta_{i_2} \delta_{i_3}^{\ell_1} - \eta_{i_3} \delta_{i_2}^{\ell_1})(g_{i_4 \ell_3} \eta_{i_5} - g_{i_5 \ell_3} \eta_{i_4}) \\
&\quad + 2(\eta_{i_4} \delta_{i_5}^{\ell_2} - \eta_{i_5} \delta_{i_4}^{\ell_2}) \eta_{\ell_3} (g_{i_2 \ell_2} \eta_{i_3} - g_{i_3 \ell_2} \eta_{i_2}) \xi^{\ell_1} \\
&= 2(\eta_{i_2} \eta_{i_5} \delta_{i_3}^{\ell_1} g_{i_4 \ell_3} - \eta_{i_3} \eta_{i_5} \delta_{i_2}^{\ell_1} g_{i_4 \ell_3} - \eta_{i_2} \eta_{i_4} \delta_{i_3}^{\ell_1} g_{i_5 \ell_3} + \eta_{i_3} \eta_{i_4} \delta_{i_2}^{\ell_1} g_{i_5 \ell_3}) \\
&\quad + 2(\eta_{i_4} \eta_{i_3} g_{i_5 i_2} - \eta_{i_5} \eta_{i_3} g_{i_2 i_4} - \eta_{i_4} \eta_{i_2} g_{i_3 i_5} + \eta_{i_5} \eta_{i_2} g_{i_3 i_4}) \xi^{\ell_1} \eta_{\ell_3}.
\end{aligned}$$

The term $\eta_{i_2} \eta_{i_5} \delta_{i_3}^{\ell_1} g_{i_4 \ell_3}$ in (37) is symmetric in i_2, i_5 , so

$$\eta_{[i_2} \eta_{i_5} \delta_{i_3}^{\ell_1} g_{i_4] \ell_3} = \eta_{i_2} \eta_{i_5} \delta_{[i_3}^{\ell_1} g_{i_4] \ell_3} - \eta_{i_5} \eta_{i_2} \delta_{[i_3}^{\ell_1} g_{i_4] \ell_3} = 0.$$

Every other term in the last two lines of (37) is symmetric in a pair of indices, so we conclude

$$((\text{R2-2}) \otimes (\text{R2-2}))_{[i_2 i_3 i_4 i_5] \ell_3}^{\ell_1} = 0.$$

□

By Lemmas 5.3 and 5.4, (34) becomes

$$(38) \quad N_{[i_2 i_3 \dots i_{4k} i_{4k+1}] \ell_0}^{\ell_1} = \rho^{4k} \underbrace{((\text{R2-1}) \wedge \dots \wedge (\text{R2-1}))}_{2k}{}_{i_2 i_3 \dots i_{4k} i_{4k+1} \ell_0}^{\ell_1} + l.o.$$

For

$$(\text{R2-1})'_{i_2 i_3 \ell_2}{}^{\ell_1} = 2(\phi_{i_2 \ell_2} \phi_{i_3}^{\ell_1} + \phi_{i_2 i_3} \phi_{\ell_2}^{\ell_1}),$$

we have

$$(\text{R2-1})_{[i_2 i_3] \ell_2}{}^{\ell_1} = (\text{R2-1})'_{[i_2 i_3] \ell_2}{}^{\ell_1}$$

Then (38) becomes

$$\begin{aligned}
(39) \quad N_{[i_2 i_3 \dots i_{4k} i_{4k+1}] \ell_0}^{\ell_1} &= \rho^{4k} ((\text{R2-1})' \wedge (\text{R2-1})' \dots \wedge (\text{R2-1})')_{i_2 i_3 \dots i_{4k} i_{4k+1} \ell_0}^{\ell_1} \\
&= \rho^{4k} \underbrace{(A' \wedge \dots \wedge A')}_k{}_{i_2 i_3 \dots i_{4k} i_{4k+1} \ell_0}^{\ell_1} + l.o.,
\end{aligned}$$

where

$$\begin{aligned}
A'_{i_2 i_3 i_4 i_5 \ell_3}{}^{\ell_1} &:= (\text{R2-1})'_{i_2 i_3 \ell_2}{}^{\ell_1} \times (\text{R2-1})'_{i_4 i_5 \ell_3}{}^{\ell_2} \\
&= 4(\phi_{i_2 \ell_2} \phi_{i_3}^{\ell_1} + \phi_{i_2 i_3} \phi_{\ell_2}^{\ell_1}) \times (\phi_{i_4 \ell_3} \phi_{i_5}^{\ell_2} + \phi_{i_4 i_5} \phi_{\ell_3}^{\ell_2}) \\
&= 4(\phi_{i_2 \ell_2} \phi_{i_3}^{\ell_1} \phi_{i_4 \ell_3} \phi_{i_5}^{\ell_2} + \phi_{i_2 \ell_2} \phi_{i_3}^{\ell_1} \phi_{i_4 i_5} \phi_{\ell_3}^{\ell_2} \\
&\quad + \phi_{i_2 i_3} \phi_{\ell_2}^{\ell_1} \phi_{i_4 \ell_3} \phi_{i_5}^{\ell_2} + \phi_{i_2 i_3} \phi_{\ell_2}^{\ell_1} \phi_{i_4 i_5} \phi_{\ell_3}^{\ell_2})
\end{aligned}$$

Since $\phi_{i_2 \ell_2} \phi_{i_3}^{\ell_1}$ is symmetric in i_2, i_5 ,

$$\phi_{i_2 \ell_2} \phi_{i_3}^{\ell_1} \phi_{i_4 \ell_3} \phi_{i_5}^{\ell_2} \sim 0 \pmod{(i_2 i_3 i_4 i_5)}.$$

Since exchanging the indices i_2, i_3 with i_4, i_5 does not change the sign of the permutation,

$$\phi_{i_2 \ell_2} \phi_{i_3}^{\ell_1} \phi_{i_4 i_5} \phi_{\ell_3}^{\ell_2} \sim \phi_{i_4 \ell_2} \phi_{i_5}^{\ell_1} \phi_{i_2 i_3} \phi_{\ell_3}^{\ell_2} \pmod{(i_2 i_3 i_4 i_5)}.$$

Therefore,

$$A'_{i_2 i_3 i_4 i_5 \ell_3}{}^{\ell_1} \sim 4(\phi_{i_4 \ell_2} \phi_{\ell_3}{}^{\ell_2} \phi_{i_5}{}^{\ell_1} + \phi_{i_4 \ell_3} \phi_{i_5}{}^{\ell_2} \phi_{\ell_2}{}^{\ell_1} + \phi_{i_4 i_5} \phi_{\ell_2}{}^{\ell_1} \phi_{\ell_3}{}^{\ell_2}) \phi_{i_2 i_3} \pmod{(i_2 i_3 i_4 i_5)},$$

which we will use to compute $(A' \wedge \dots \wedge A')_{i_2 i_3 \dots i_{4k} i_{4k+1} \ell}{}^{\ell}$ in (39). Since the wedge product involves skew-symmetrization, or modding out in our notation, we can relabel A' as

$$(40) \quad (A')_{i_2 i_3 i_4 i_5 \ell_3}{}^{\ell_1} := 4\phi_{i_2 i_3} (B^*)_{i_4 i_5 \ell_3}{}^{\ell_1} := 4\phi_{i_2 i_3} ((B1)_{i_4 i_5 \ell_3}{}^{\ell_1} + (B2)_{i_4 i_5 \ell_3}{}^{\ell_1} + (B3)_{i_4 i_5 \ell_3}{}^{\ell_1}),$$

where

$$(41) \quad (B^*)_{i_4 i_5 \ell_3}{}^{\ell_1} = (B1)_{i_4 i_5 \ell_3}{}^{\ell_1} + (B2)_{i_4 i_5 \ell_3}{}^{\ell_1} + (B3)_{i_4 i_5 \ell_3}{}^{\ell_1}, \\ (B1)_{i_4 i_5 \ell_3}{}^{\ell_1} = -(\phi^{(2)})_{i_4 \ell_3} \phi_{i_5}{}^{\ell_1}, \quad (B2)_{i_4 i_5 \ell_3}{}^{\ell_1} = \phi_{i_4 \ell_3} (\phi^{(2)})_{i_5}{}^{\ell_1}, \quad (B3)_{i_4 i_5 \ell_3}{}^{\ell_1} = \phi_{i_4 i_5} (\phi^{(2)})_{\ell_3}{}^{\ell_1}.$$

Here we introduce the notation

$$(\phi^{(2)})_k{}^h := \phi_k^{j_1} \phi_{j_1}{}^h = -\delta_k{}^h + \eta_k \xi^h, \quad (\phi^{(3)})_k{}^h := \phi_k^{j_1} \phi_{j_1}{}^{j_2} \phi_{j_2}{}^h = -\phi_k{}^h, \\ (\phi^{(4)})_k{}^h := \phi_k^{j_1} \phi_{j_1}{}^{j_2} \phi_{j_2}{}^{j_3} \phi_{j_3}{}^h = \delta_k{}^h - \eta_k \xi^h = -(\phi^{(2)})_k{}^h.$$

For

$$(42) \quad (A')_{i_2 \dots i_9 \ell_4}{}^{\ell_1} := (A')_{i_2 i_3 i_4 i_5 \ell_3}{}^{\ell_1} \times (A')_{i_6 i_7 i_8 i_9 \ell_4}{}^{\ell_3} = 4^2 \phi_{i_2 i_3} \phi_{i_6 i_7} (B^*)_{i_4 i_5 \ell_3}{}^{\ell_1} \times (B^*)_{i_8 i_9 \ell_4}{}^{\ell_3}$$

we have:

Lemma 5.5.

$$(A')_{i_2 \dots i_9 \ell_4}{}^{\ell_1} \sim -(4^2) \phi_{i_2 i_3} \phi_{i_4 i_5} \phi_{i_6 i_7} B_{i_8 i_9 \ell_4}^*{}^{\ell_1} \pmod{(i_2 \dots i_9)}.$$

Proof. We compute

$$(B1)_{i_4 i_5 \ell_3}{}^{\ell_1} \times (B1)_{i_8 i_9 \ell_4}{}^{\ell_3} = (\phi^{(2)})_{i_4 \ell_3} \phi_{i_9}{}^{\ell_3} (\phi^{(2)})_{i_8 \ell_4} \phi_{i_5}{}^{\ell_1} = -(\phi^{(3)})_{i_4 i_9} (\phi^{(2)})_{i_8 \ell_4} \phi_{i_5}{}^{\ell_1} \\ \sim \phi_{i_4 i_5}^{(3)} \phi_{i_8 \ell_4}^{(2)} \phi_{i_9}{}^{\ell_1} \sim -\phi_{i_4 i_5} \phi_{i_8 \ell_4}^{(2)} \phi_{i_9}{}^{\ell_1} \pmod{(i_4 i_5 i_8 i_9)}, \\ (B1)_{i_4 i_5 \ell_3}{}^{\ell_1} \times (B2)_{i_8 i_9 \ell_4}{}^{\ell_2} = -(\phi^{(2)})_{i_4 \ell_3} \phi_{i_5}{}^{\ell_1} \phi_{i_8 \ell_4} (\phi^{(2)})_{i_9}{}^{\ell_2} = -(\phi^{(4)})_{i_4 i_9} \phi_{i_5}{}^{\ell_1} \phi_{i_8 \ell_4} \\ \sim 0 \pmod{(i_4 i_5 i_8 i_9)}, \\ (B1)_{i_4 i_5 \ell_3}{}^{\ell_1} \times (B3)_{i_8 i_9 \ell_4}{}^{\ell_2} = -(\phi^{(2)})_{i_4 \ell_3} \phi_{i_5}{}^{\ell_1} (\phi^{(2)})_{\ell_4}{}^{\ell_2} \phi_{i_8 i_9} = -(\phi_{i_4 \ell_4}^{(4)}) \phi_{i_5}{}^{\ell_1} \phi_{i_8 i_9} \\ \sim -(\phi^{(4)})_{i_8 \ell_4} \phi_{i_9}{}^{\ell_1} \phi_{i_4 i_5} \sim (\phi^{(2)})_{i_8 \ell_4} \phi_{i_9}{}^{\ell_1} \phi_{i_4 i_5} \pmod{(i_4 i_5 i_8 i_9)}.$$

(For the second equation, note that $\phi_{i_4 i_9}^{(4)} = g_{i_4 i_9} - \eta_{i_4} \eta_{i_9}$ is symmetric in i_4, i_9 .)

Thus, we have

$$(43) \quad \sum_{E=1}^3 (B1)_{i_4 i_5 \ell_3}{}^{\ell_1} \times (BE)_{i_8 i_9 \ell_4}{}^{\ell_3} \sim 0 \pmod{(i_4 i_5 i_8 i_9)}.$$

Again using the symmetry of $(\phi^{(2)})_{i_a i_b}$, we have

$$\begin{aligned}
(\text{B2})_{i_4 i_5 \ell_3}^{\ell_1} \times (\text{B1})_{i_8 i_9 \ell_4}^{\ell_2} &= -\phi_{i_4 \ell_3}(\phi^{(2)})_{i_5}^{\ell_1}(\phi^{(2)})_{i_8 \ell_4} \phi_{i_9}^{\ell_2} = (\phi^{(2)})_{i_8 \ell_4}(\phi^{(2)})_{i_4 i_9}(\phi^{(2)})_{i_5}^{\ell_1} \\
&\sim 0 \pmod{(i_4 i_5 i_8 i_9)}, \\
(\text{B2})_{i_4 i_5 \ell_3}^{\ell_1} \times (\text{B2})_{i_8 i_9 \ell_4}^{\ell_2} \\
&= \phi_{i_4 \ell_3}(\phi^{(2)})_{i_5}^{\ell_1} \phi_{i_8 \ell_4}(\phi^{(2)})_{i_9}^{\ell_2} = -(\phi^{(3)})_{i_9 i_4}(\phi^{(2)})_{i_5}^{\ell_1} \phi_{i_8 \ell_4} = \phi_{i_9 i_4}(\phi^{(2)})_{i_5}^{\ell_1} \phi_{i_8 \ell_4} \\
&\sim -\phi_{i_5 i_4}(\phi^{(2)})_{i_9}^{\ell_1} \phi_{i_8 \ell_4} \sim \phi_{i_4 i_5}(\phi^{(2)})_{i_9}^{\ell_1} \phi_{i_8 \ell_4} \pmod{(i_4 i_5 i_8 i_9)}, \\
(\text{B2})_{i_4 i_5 \ell_3}^{\ell_1} \times (\text{B3})_{i_8 i_9 \ell_4}^{\ell_2} &= \phi_{i_4 \ell_3}(\phi^{(2)})_{i_5}^{\ell_1} \phi_{i_8 i_9}(\phi^{(2)})_{\ell_4}^{\ell_2} = \phi_{i_8 i_9} \phi_{i_4 \ell_3}(\phi^{(2)})_{\ell_4}^{\ell_2} (\phi^{(2)})_{i_5}^{\ell_1} \\
&= \phi_{i_8 i_9}(\phi^{(3)})_{i_4 \ell_4}(\phi^{(2)})_{i_5}^{\ell_1} \\
&\sim \phi_{i_4 i_5}(\phi^{(3)})_{i_8 \ell_4}(\phi^{(2)})_{i_9}^{\ell_1} \sim -\phi_{i_4 i_5} \phi_{i_8 \ell_4}(\phi^{(2)})_{i_9}^{\ell_1} \pmod{(i_4 i_5 i_8 i_9)}.
\end{aligned}$$

Therefore,

$$(44) \quad \sum_{E=1}^3 (\text{B2})_{i_4 i_5 \ell_3}^{\ell_1} \times (\text{BE})_{i_8 i_9 \ell_4}^{\ell_2} \sim 0 \pmod{(i_4 i_5 i_8 i_9)}.$$

Finally, we compute

$$\begin{aligned}
(\text{B3})_{i_4 i_5 \ell_3}^{\ell_1} \times (\text{B1})_{i_8 i_9 \ell_4}^{\ell_2} &= \phi_{i_4 i_5}(\phi^{(2)})_{\ell_3}^{\ell_1}(-(\phi^{(2)})_{i_8 \ell_4})\phi_{i_9}^{\ell_2} = \phi_{i_4 i_5}(\phi^{(3)})_{i_9}^{\ell_1}(\phi^{(2)})_{i_8 \ell_4} \\
&= \phi_{i_4 i_5}(-\phi_{i_9}^{\ell_1})(\phi^{(2)})_{i_8 \ell_4} = \phi_{i_4 i_5} \phi_{i_9}^{\ell_1}(\phi^{(2)})_{i_8 \ell_4} \\
&\sim -\phi_{i_4 i_5} \phi_{i_8}^{\ell_1}(\phi^{(2)})_{i_9 \ell_4} \pmod{(i_4 i_5 i_8 i_9)}, \\
(\text{B3})_{i_4 i_5 \ell_3}^{\ell_1} \times (\text{B2})_{i_8 i_9 \ell_4}^{\ell_2} &= \phi_{i_4 i_5}(\phi^{(2)})_{\ell_3}^{\ell_1} \phi_{i_8 \ell_4}(\phi^{(2)})_{i_9}^{\ell_2} = \phi_{i_4 i_5} \phi_{i_8 \ell_4}(\phi^{(4)})_{i_9}^{\ell_1} \\
&= -\phi_{i_4 i_5} \phi_{i_8 \ell_4}(\phi^{(2)})_{i_9}^{\ell_1}, \\
(\text{B3})_{i_4 i_5 \ell_3}^{\ell_1} \times (\text{B3})_{i_8 i_9 \ell_4}^{\ell_2} &= \phi_{i_4 i_5} \phi_{i_8 i_9}(\phi^{(2)})_{\ell_3}^{\ell_1}(\phi^{(2)})_{\ell_4}^{\ell_2} = \phi_{i_4 i_5} \phi_{i_8 i_9}(\phi^{(4)})_{\ell_4}^{\ell_1} \\
&= -\phi_{i_4 i_5} \phi_{i_8 i_9}(\phi^{(2)})_{\ell_4}^{\ell_1}.
\end{aligned}$$

By (43), (44), (45), we have

$$\begin{aligned}
(A')_{i_2 \dots i_9 \ell_4}^{(2) \ell_1} &\sim \phi_{i_2 i_3} \phi_{i_6 i_7} \sum_{E=1}^3 (\text{B3})_{i_4 i_5 \ell_3}^{\ell_1} \times (\text{BE})_{i_8 i_9 \ell_4}^{\ell_2} \pmod{(i_2 \dots i_9)} \\
&\sim -(4^2) \phi_{i_2 i_3} \phi_{i_4 i_5} \phi_{i_6 i_7} B_{i_8 i_9 \ell_4}^* \ell_1 \pmod{(i_2 \dots i_9)},
\end{aligned}$$

which proves Lemma 5.5. □

Corollary 5.2. For $(A')_{i_2 i_3 i_4 i_5 \ell_3}^{\ell_1}$ in (40) and $(A')_{i_2 \dots i_9 \ell_4}^{(2) \ell_1}$ in (42), set

$$(A')_{i_2 \dots i_{4r+1} \ell_{r+2}}^{(r) \ell_1} = (A')_{i_2 \dots i_{4r-3} \ell_{r+1}}^{(r-1) \ell_1} (A')_{i_{4r-2} i_{4r-1} i_{4r} i_{4r+1} \ell_{r+2}}^{\ell_{k+1}},$$

$r = 2, \dots, k$. Then (i)

$$(A')_{i_2 \dots i_{4r+1} \ell_{r+2}}^{(r) \ell_1} \sim (-1)^{r-1} 4^r \phi_{i_2 i_3} \phi_{i_4 i_5} \dots \phi_{i_{4r-2} i_{4r-1}} B_{i_{4r} i_{4r+1} \ell_{r+2}}^* \ell_1 \pmod{(i_2 \dots i_{4r+1})}$$

;

(ii) In the notation of (27),

$$\text{tr}(N)_{[i_2 \dots i_{4k+1}]} = N_{[i_2 \dots i_{4k+1}]} \ell^\ell = (-1)^{k-1} 4^k (4k+2) \rho^{4k} (\phi \wedge \dots \wedge \phi)_{i_2 \dots i_{4k+1}} + l.o.$$

Proof. (i) By Lemma 5.5, this follows by induction on r up to $r = k$, where $\dim(M) = 4k + 1$.

(ii) By (27), (28), (39), (40),

$$(46) \quad \begin{aligned} \operatorname{tr}(N)_{[i_2 \dots i_{4k+1}]} &= \rho^{4k} \operatorname{tr}(A')_{[i_2 \dots i_{4k+1}]}^{(k)} = \rho^{4k} (A')_{[i_2 \dots i_{4k+1}]}^{(k)\ell} \\ &\sim (-1)^{k-1} 4^k \rho^{4k} \phi_{[i_2 i_3} \phi_{i_4 i_5} \dots \phi_{i_{4k-2} i_{4k-1}} B_{i_{4k} i_{4k+1}}^* \ell^\ell \pmod{(i_2 \dots i_{4k+1})}. \end{aligned}$$

By (41),

$$\begin{aligned} B_{abl}^* \ell^\ell &= -(\phi^{(2)})_{al} \phi_b^\ell + \phi_{al} (\phi^{(2)})_b^\ell + \phi_{ab} (\phi^{(2)})_\ell^\ell = \phi_{ab} + \phi_{ab} + \phi_{ab} (\delta_\ell^\ell - \eta_\ell \xi^\ell) \\ &= 2\phi_{ab} + \phi_{ab} ((4k+1) - 1) = (4k+2)\phi_{ab}, \end{aligned}$$

and plugging this into (46) finishes the proof. \square

This proves (27), and so finishes the proof of Prop. 5.1 and of Thm. 5.1

6. SASAKIAN SPACE FORMS

For the Sasakian space forms defined below, the computations are more explicit. We start with the case of the sphere S^{4k+1} with the standard metric g_{std} , and then treat the other Sasakian space forms.

As in Ex. 2.1, S^{4k+1} has the Hopf fibration

$$(47) \quad S^1 \rightarrow S^{4k+1} \rightarrow \mathbb{C}\mathbb{P}^{2k}.$$

Let ξ be the unit vector field on S^{4k+1} tangent to the S^1 fiber, and let η be its dual 1-form. Then $(S^{4k+1}, g_{std}, \phi = \nabla\xi, \xi, \eta)$ is a Sasakian structure on S^{4k+1} .

We can refine Prop. 5.1 for S^{4k+1} .

Proposition 6.1. *Let $(S^{4k+1}, g_{std}, \phi, \xi, \eta)$ be as above. Then in the notation of Prop. 5.1*

$$(48) \quad K_{j[i_1 \dots i_{4k+1}]} \xi^j = (1 + \rho^2)^2 a_{4k} \rho^{4k},$$

with

$$a_{4k} = (-1)^k 4^k (4k+2) (\eta \wedge d\eta^{2k})_{i_1 \dots i_{4k+1}}.$$

Assuming the Proposition, we obtain:

Theorem 6.1. *Let S^{4k+1} be $(4k+1)$ -dimensional sphere with the standard metric g_{std} . Then, we have*

$$(49) \quad |\pi_1(\operatorname{Isom}(S^{4k+1}, h_\rho))| = \infty$$

for the metric $h_\rho = g_{std} + \rho^2 \eta \otimes \eta$, $\rho \neq 0$.

Note the contrast to the standard metric at $\rho = 0$, where $\pi_1(\operatorname{Isom}(S^{4k+1}, g_{std})) = \mathbb{Z}_2$.

The proof of the Theorem is almost the same as for Thm. 5.1: since $K_{j[i_1 \dots i_{4k+1}]} \xi^j \neq 0$ in (48) and the coefficients a_r of ρ^r in Prop. 5.1 vanish, $(a^L)^* CS^W \neq 0$ for any $\rho \neq 0$. Then we follow the proof of Thm. 5.1.

Prop. 6.1 will follow from the next Lemma.

Lemma 6.1. *For (S^{4k+1}, h_ρ) , in the notation of (30) we have*

$$\begin{aligned} (i) \quad & ((\text{R1}) \otimes (\text{RE}))_{[i_2 i_3 i_4 i_5] \ell_3}^{\ell_1} = 0; \\ (ii) \quad & ((\text{RE}) \otimes (\text{R1}))_{[i_2 i_3 i_4 i_5] \ell_3}^{\ell_1} = 0; \end{aligned}$$

for $E = 1, 2, 3$.

Proof. (i) (S^{4k+1}, g_{std}) has positive constant sectional curvature $g(R(X, Y)Y, X) = 1$, for $|X| = |Y| = 1, X \perp Y$, and curvature tensor

$$(50) \quad R_{kji}{}^h = \frac{c+3}{4}(g_{ki}\delta_j{}^h - g_{ji}\delta_k{}^h),$$

with $c = 1$ [5, I.V.2].

Noting that $(R1)_{kji}{}^h = R_{kji}{}^h$, we have

$$\begin{aligned} (R1)_{i_2i_3\ell_2}{}^{\ell_1} \times (RE)_{i_4i_5\ell_3}{}^{\ell_2} &= \frac{c+3}{4}(g_{i_2\ell_2}\delta_{i_3}{}^{\ell_1} - g_{i_3\ell_2}\delta_{i_2}{}^{\ell_1}) \times (RE)_{i_4i_5\ell_3}{}^{\ell_2} \\ &= \frac{c+3}{4}(\delta_{i_3}{}^{\ell_1}(RE)_{i_4i_5\ell_3i_2} - \delta_{i_2}{}^{\ell_1}(RE)_{i_4i_5\ell_3i_3}), \end{aligned}$$

where $(RE)_{kjih} = g_{lh}(RE)_{kji}{}^{\ell}$.

Note that $(RE)_{kjih} = -(RE)_{kjhi}$: for $\bar{R}_{kjih} = -\bar{R}_{kjhi}$ for any ρ , and (R1), (R2), (R3) come with different powers of ρ , so each (RE) is skew-symmetric in i, h . Thus,

$$((R1) \otimes (RE))_{[i_2i_3i_4i_5]\ell_3}{}^{\ell_1} = \frac{c+3}{4}(-\delta_{[i_3}{}^{\ell_1}(RE)_{i_4i_5i_2]\ell_3} - \delta_{[i_2}{}^{\ell_1}(RE)_{i_4i_5i_3]\ell_3})$$

where we skew-symmetrize over the i indices. Since $(RE)_{[kji]}{}^h = 0$ by the Bianchi identity,

$$((R1) \otimes (RE))_{[i_2i_3i_4i_5]\ell_3}{}^{\ell_1} = 0.$$

(ii) This is similar:

$$\begin{aligned} (RE)_{i_2i_3\ell_2}{}^{\ell_1} \times (R1)_{i_4i_5\ell_3}{}^{\ell_2} &= (RE)_{i_2i_3\ell_2}{}^{\ell_1} \times \frac{c+3}{4}(g_{i_4\ell_3}\delta_{i_5}{}^{\ell_2} - g_{i_5\ell_3}\delta_{i_4}{}^{\ell_2}) \\ &= \frac{c+3}{4}((RE)_{i_2i_3i_5}{}^{\ell_1}g_{i_4\ell_3} - (RE)_{i_2i_3i_4}{}^{\ell_1}g_{i_5\ell_3}), \end{aligned}$$

so again,

$$((RE) \otimes (R1))_{[i_2i_3i_4i_5]\ell_3}{}^{\ell_1} = 0. \quad \square$$

Proof of Prop. 6.1. By Lem. 5.3, (34), and Lem. 6.1, the lowest power of ρ in $K_{j[i_1 \dots i_{4k+1}]\xi^j}$ is ρ^{4k} . \square

We can generalize this result to Sasakian space forms. Recall that a Sasakian manifold (M, ϕ, ξ, η) is called a *Sasakian space form* if all the ϕ -sectional curvatures $K(X \wedge \phi X)$ are equal to a constant c , where X is a unit vector field orthogonal to ξ , $\{X, \phi X\}$ span a two-plane section of TM , and $K(X \wedge \phi X)$ denotes the sectional curvature. In this case, the Riemannian curvature tensor of M is given by

$$(51) \quad \begin{aligned} R_{kji}{}^h &= \frac{c+3}{4}(g_{ji}\delta_k{}^h - g_{ki}\delta_j{}^h) \\ &\quad + \frac{c-1}{4}(\eta_i\eta_k\delta_j{}^h - \eta_j\eta_i\delta_k{}^h + g_{ki}\eta_j\xi^h - g_{ji}\eta_k\xi^h + \phi_{ji}\phi_k{}^h - \phi_{ki}\phi_j{}^h - 2\phi_{kj}\phi_i{}^h) \end{aligned}$$

[2, Thm. 7.19].

As with ordinary space forms, Sasakian space forms fall into three classes [2, Thm. 7.20], [9] depending on whether $c > -3$, $c = -3$, or $c < -3$. For $\dim(M) = 4k + 1$, we have the following after scaling:

- $c+3 = 4$. M is locally isometric to $(4k+1)$ -dimensional sphere of constant sectional curvature 1.
- $c+3 = 0$. M is locally isometric to the Heisenberg nilpotent Lie group $N = \mathbb{R}^{4k} \times \mathbb{R}$, with $g = \frac{1}{4} \sum_{i=1}^{2k} ((dx^i)^2 + (dy^i)^2) + \eta \otimes \eta$, with $\eta = \frac{1}{2} \left(dx^{4k+1} - \sum_{i=1}^{2k} y^i dx^i \right)$.
- $c+3 = -1$. M is locally isometric to $\bar{B} = B \times \mathbb{R}$, where (B, J, G) is a simply connected bounded domain in \mathbb{C}^n with Kähler structure J and metric G , the curvature of G equals $\Omega = d\omega$, and $\eta = \pi^*\omega + dt$, $g = \pi^*G + \eta \otimes \eta$ on \bar{B} for the projection $\pi : \bar{B} \rightarrow B$.

Let (M, g, ϕ, ξ, η) be a $(4k+1)$ -dimensional Sasakian space form. Its curvature tensor (51) can be rewritten as

$$\begin{aligned} R_{kji}{}^h &= \frac{c+3}{4} (g_{ji}\delta_k{}^h - g_{ki}\delta_j{}^h) \\ &\quad - \frac{c-1}{4} (\phi_{ki}\phi_j{}^h - \phi_{ji}\phi_k{}^h + 2\phi_{kj}\phi_i{}^h + 2\eta_i\eta_k\delta_j{}^h - 2\eta_j\eta_i\delta_k{}^h + g_{ki}\eta_j\xi^h - g_{ji}\eta_k\xi^h) \\ &\quad + \frac{c-1}{4} (3\eta_i\eta_k\delta_j{}^h - 3\eta_j\eta_i\delta_k{}^h + 2g_{ki}\eta_j\xi^h - 2g_{ij}\eta_k\xi^h) \\ &= (\text{S1})_{kji}{}^h + (\text{S2})_{kji}{}^h + (\text{S3})_{kji}{}^h, \end{aligned}$$

where

$$\begin{aligned} (\text{S1})_{kji}{}^h &= \frac{c+3}{4} (g_{ji}\delta_k{}^h - g_{ki}\delta_j{}^h), \\ (\text{S2})_{kji}{}^h &= -\frac{c-1}{4} (\phi_{ki}\phi_j{}^h - \phi_{ji}\phi_k{}^h + 2\phi_{kj}\phi_i{}^h + 2\eta_i\eta_k\delta_j{}^h - 2\eta_j\eta_i\delta_k{}^h + g_{ki}\eta_j\xi^h - g_{ji}\eta_k\xi^h), \\ (\text{S3})_{kji}{}^h &= \frac{c-1}{4} (3\eta_i\eta_k\delta_j{}^h - 3\eta_j\eta_i\delta_k{}^h + 2g_{ki}\eta_j\xi^h - 2g_{ij}\eta_k\xi^h). \end{aligned}$$

Lemma 6.2.

$$(\text{S}\alpha)_{i_2i_3\ell_2}{}^{\ell_1} \times (\text{RE})_{i_4i_5\ell_3}{}^{\ell_2} \sim 0, \quad (\text{RE})_{i_2i_3\ell_2}{}^{\ell_1} \times (\text{S}\alpha)_{i_4i_5\ell_3}{}^{\ell_2} \sim 0,$$

for $\alpha = 1, \text{E} = 1, 2, 3$, and for $\text{E} = 3, \alpha = 1, 2, 3$.

Proof. For $\alpha = 1$, $(\text{S1}) = (\text{R1})$, so the proof of Lem. 6.1 carries over. For $\text{E} = 3$, each term in $(\text{S3})_{kji}{}^h$ contains either two η 's or an $\eta\xi$ pair, so these terms are similar to 0. \square

We note that

$$\bar{R}_{kji}{}^h (\text{R1})_{kji}{}^h + (\text{S2}')_{kji}{}^h + (\text{S3})_{kji}{}^h + (\text{R3})_{kji}{}^h,$$

where

$$(\text{S2}')_{kji}{}^h = -\left(\frac{c-1}{4} + \rho^2\right) (\phi_{ki}\phi_j{}^h - \phi_{ji}\phi_k{}^h + 2\phi_{kj}\phi_i{}^h + 2\eta_i\eta_k\delta_j{}^h - 2\eta_j\eta_i\delta_k{}^h + g_{ki}\eta_j\xi^h - g_{ji}\eta_k\xi^h).$$

Thus up to a scale factor $(\text{S2}')_{kji}{}^h = (\text{R2})_{kji}{}^h$ by (30). We can repeat the proof of Prop. 5.1 by replacing the power $\rho^{4k} = (\rho^2)^{2k}$ in Prop. 5.1 with $\frac{c-1}{4} + \rho^2$. We obtain:

Proposition 6.2. *Let $(M, g_0, \phi, \xi, \eta)$ be a Sasakian space form as above. Then in the notation of Prop. 5.1*

$$K_{j[i_1 \dots i_{4k+1}]} \xi^j = (1 + \rho^2)^2 a_{4k} \left(\frac{c-1}{4} + \rho^2\right)^{2k},$$

with

$$a_{4k} = (-1)^k 4k(4k+2) (\eta \wedge d\eta^{2k})_{i_1 \dots i_{4k+1}}.$$

As before, we conclude the following:

Theorem 6.2. *Let (M, g, ϕ, ξ, η) be $(4k + 1)$ -dimensional closed orientable Sasakian space form. Then $|\pi_1(\text{Isom}(M, g_\rho))| = \infty$ for (i) $c + 3 > 4, \rho \neq 0$, (ii) $c + 3 = 0, \rho \neq \pm 1$, or (iii) $c + 3 < 0, \rho^2 \neq -\frac{c-1}{4}$.*

Remark 6.1. It follows from [1, Note 7.5] that $|\pi(\text{Isom}(M, g_0))| = \infty$ for the Heisenberg infra-nilmanifold $M = N/\Gamma$ (i.e., a Sasakian space form with $c + 3 = 0$). This follows also from our result.

7. STRICT CONTACT DIFFEOMORPHISMS OF S^{4k+1}

We consider again the sphere S^{4k+1} with the standard metric g_{std} , and Sasakian structure $(g_{st}, \phi, \eta, \xi)$. The Reeb vector field ξ of the Hopf fibering

$$\pi : S^1 \rightarrow S^{4k+1} \rightarrow \mathbb{C}\mathbb{P}^{2k}$$

is the unit tangent vector field along the S^1 fiber. It generates not only an S^1 -action via isometries of g_{std} , but also an S^1 -action of strict contact diffeomorphisms of S^{4k+1} as a contact manifold. Here, we recall that a diffeomorphism ϕ of S^{4k+1} is strictly contact if $\phi^*\eta = \eta$. We denote the group of strict contact diffeomorphisms of (S^{4k+1}, η) by $\text{Diff}_{\eta, str}(S^{4k+1}, \eta)$. We note that this group is infinite dimensional.

Then we have the following:

Theorem 7.1.

$$|\pi_1(\text{Diff}_{\eta, str}(S^{4k+1}, \eta))| = \infty.$$

The proof is similar to those in other sections, and will be given below. As in §4, we take the family of metrics $h = g_\rho$ defined by $h_{ij} = g_{ij} + \rho^2 \eta_i \eta_j$ for $g = g_{std}$. We can explicitly compute the integrand $K_{\nu[\lambda_1 \lambda_2 \dots \lambda_{4k+1}]}$ in the definition of CS^W (18).

Proposition 7.1.

$$K_{\nu[\lambda_1 \lambda_2 \dots \lambda_{4k+1}]} = C \rho^{4k} g_{\nu[\lambda_1]} ((d\eta)^{2k})_{\lambda_1 \lambda_2 \dots \lambda_{4k+1}},$$

for some nonzero constant C .

The proof of Prop. 7.1 is given after the next two Lemmas.

Recall that

$$K_{\nu[\lambda_1 \lambda_2 \dots \lambda_{4k+1}]} = \bar{R}_{\lambda_1 \kappa_1 \nu}^{\kappa_0} \bar{R}_{\lambda_2 \lambda_3 \kappa_2}^{\kappa_1} \dots \bar{R}_{\lambda_{4k} \lambda_{4k+1} \kappa_0}^{\kappa_{2k}}.$$

Using (14) and (50), we have

$$\bar{R}_{kji}^h = (\text{CR1})_{kji}^h - \rho^2 (\text{CR2})_{kji}^h - \rho^2 (\text{CR3})_{kji}^h - (\rho^2 + \rho^4) (\text{CR4})_{kji}^h,$$

where

$$\begin{aligned} (\text{CR1})_{kji}^h &= g_{ji} \delta_k^h - g_{ki} \delta_j^h, & (\text{CR2})_{kji}^h &= \phi_{ki} \phi_j^h - \phi_k^h + 2\phi_{ki} \phi_i^h, \\ (\text{CR3})_{kji}^h &= g_{ki} \eta_j \xi^h - g_{ji} \eta_k \xi^h, & (\text{CR4})_{kji}^h &= \eta_k \eta_j \delta_j^h - \eta_j \eta_i \delta_k^h. \end{aligned}$$

Lemma 7.1.

$$(\text{CR1})_{i_1 \kappa_1 j}^{\kappa_0} \times \bar{R}_{i_2 i_3 \kappa_2}^{\kappa_1} \dots \bar{R}_{i_{4k} i_{4k+1} \kappa_0}^{\kappa_{2k}} \sim -g_{1j} \bar{R}_{i_2 i_3 \kappa_1}^{\kappa_0} \dots \bar{R}_{i_{4k} i_{4k+1} \kappa_0}^{\kappa_{2k}} \pmod{(i_1 i_2 i_3)}.$$

Proof.

$$\begin{aligned}
(\text{CR1})_{i_1 \kappa_1 j}^{\kappa_0} &\times \bar{R}_{i_2 i_3 \kappa_2}^{\kappa_1} \cdots \bar{R}_{i_{4k} i_{4k+1} \kappa_0}^{\kappa_{2k}} \\
&= (g_{\kappa_1 j} \delta_{i_1}^{\kappa_0} - g_{i_1 j} \delta_{\kappa_1}^{\kappa_0}) \bar{R}_{i_2 i_3 \kappa_2}^{\kappa_1} \cdots \bar{R}_{i_{4k} i_{4k+1} \kappa_0}^{\kappa_{2k}} \\
&= \bar{R}_{i_2 i_3 \kappa_2}^{\kappa_1} \cdots \bar{R}_{i_{4k} i_{4k+1} i_1}^{\kappa_{2k}} - g_{i_1 j} \bar{R}_{i_2 i_3 \kappa_2}^{\kappa_0} \cdots \bar{R}_{i_{4k} i_{4k+1} \kappa_0}^{\kappa_{2k}} \\
&\sim -g_{i_1 j} \bar{R}_{i_2 i_3 \kappa_2}^{\kappa_0} \cdots \bar{R}_{i_{4k} i_{4k+1} \kappa_0}^{\kappa_{2k}},
\end{aligned}$$

because $\bar{R}_{[i_{4k} i_{4k+1} i_1]}^{\kappa_{2k}} = 0$. □

Set

$$T_{j i_1 i_2 \cdots i_{4k+1}} = \bar{R}_{i_2 i_3 \kappa_2}^{\kappa_0} \bar{R}_{i_4 i_5 \kappa_3}^{\kappa_2} \cdots \bar{R}_{i_{4k} i_{4k+1} \kappa_0}^{\kappa_{2k}}.$$

By Lemma 7.1, we have

$$K_{j[i_1 i_2 \cdots i_{4k+1}]} = -g_{j[i_1} T_{i_2 i_3 \cdots i_{4k+1}]}.$$

Many terms in $T_{j i_1 i_2 \cdots i_{4k+1}}$ vanish by direct computations as in Lemmas 5.4, 6.2.

Lemma 7.2.

$$(\text{CRE})_{i_2 i_3 j_1}^{j'_0} \times (\text{CRF})_{i_4 i_5 j_2}^{j_1} \sim 0 \quad \text{mod}(i_1 i_2 i_3 i_4),$$

for $E, F \in \{1, 2, 3, 4\}$, $(E, F) \neq (2, 2)$.

Proof of Prop. 7.1. We have

$$T_{j i_1 i_2 \cdots i_{4k+1}} \sim \rho^{4k} (\text{CR2})_{i_2 i_3 j_2}^{j_0} (\text{CR2})_{i_2 i_3 j_2}^{j_0} \cdots (\text{CR2})_{i_{4k} i_{4k+1} j_0}^{j_{2k}} \quad \text{mod}(i_1 \cdots i_{4k+1}).$$

As in §5, there is a nonzero constant C such that

$$T_{j[i_1 i_2 \cdots i_{4k+1}]} = C \rho^{4k} ((d\eta)^{2k})_{i_2 i_3 \cdots i_{4k+1}}$$

□

Proof of Thm. 7.1. By Proposition 7.1, $K_{j[i_1 i_2 \cdots i_{4k+1}]} \xi^j$ is given explicitly in terms of η and $d\eta$ (see Prop. 5.1). Thus, $K_{j[i_1 i_2 \cdots i_{4k+1}]} \xi^j$ is invariant under strict contact diffeomorphisms. For the S^1 -action of the Reeb vector field ξ , the proof of Thm. 5.1 can be directly adapted to $\text{Diff}_{\eta, str}(S^{4k+1})$. □

Remark 7.1. (i) By a slight modification of these computations, Thm. 7.1 can be easily extended to the Sasakian space forms of §6.

(ii) Thm. 7.1 has been obtained in [3, Cor. 7], with for S^{4k+1} generalized to a closed contact manifold M with closed Reeb orbits. While both approaches prove the same element $[a^D] \in \pi_1(\text{Diff}_{\eta, str}(M))$ has infinite order, [3] uses the map $a^D : S^1 \rightarrow \text{Diff}_{\eta, str}(M)$ to pull back characteristic classes from $B\text{Diff}_{\eta, str}(M)$ to BS^1 . In contrast, we use the equivalent map $a^L : M \rightarrow LM$, $a^L(m)(\theta) = a^D(\theta)(m)$, to pull back a (nonclosed) Chern-Simons form from LM to M . It would be interesting to understand the relationship between these approaches better. In [6], we reprove [3, Cor. 7] using our approach.

REFERENCES

1. Baues, O. and Kamishima, Y., *Locally homogeneous aspherical Sasaki manifolds*, Differential Geom. Appl. **70** (2020).
2. Blair, D., *Riemannian geometry of contact and symplectic manifolds*, second ed., Progress in Mathematics, vol. 203, Birkhäuser Boston, Ltd., Boston, MA, 2010.
3. Casals, R. and Spáčil, O., *Chern-Weil theory and the group of strict contactomorphisms*, J. Topol. Anal. **8** (2016), no. 1, 59–87.

4. Egi, S., Maeda, Y., and Rosenberg, S., *Isometry groups of circle bundles over integral symplectic manifolds*, <https://arxiv.org/abs/arXiv:2011.01800>.
5. Kobayashi, S. and Nomizu, K., *Foundations of Differential Geometry*, vol. 2, Interscience Publishers, John Wiley & Sons Inc., New York, 1969.
6. Maeda, Y. and Rosenberg, S., *The geometry of loop spaces V: Fundamental groups of geometric transformation groups*, in preparation.
7. Maeda, Y., Rosenberg, S., and Torres-Ardila, F., *The geometry of loop spaces II: Corrections*, <https://arxiv.org/abs/2405.00651>.
8. ———, *The geometry of loop spaces II: Characteristic classes*, Adv. in Math. **287** (2016), 485–518.
9. Tanno, S., *Sasakian manifolds with constant ϕ -holomorphic sectional curvature*, Tohoku Math. J. (2) **21** (1969), 501–507.

TOHOKU FORUM FOR CREATIVITY, TOHOKU UNIVERSITY

Email address: yoshimaeda@tohoku.ac.jp

DEPARTMENT OF MATHEMATICS AND STATISTICS, BOSTON UNIVERSITY

Email address: sr@math.bu.edu