

Edge modes in chiral electron double layers

Klaus Ziegler¹ and Roman Ya. Kezerashvili^{2,3}

¹*Institut für Physik, Universität Augsburg, D-86135 Augsburg, Germany*

²*Physics Department, New York City College of Technology,
The City University of New York, Brooklyn, NY 11201, USA*

³*The Graduate School and University Center,
The City University of New York, New York, NY 10016, USA*

(Dated: December 24, 2024)

We study the quasiparticles in chiral double layers with electron pairing within the framework of the Bogoliubov de Gennes equation. In the presence of an edge it is demonstrated that the quasiparticle modes can be distinguished as edge modes and bulk modes, which appear at different energies. The bulk-edge correspondence is obtained by an analytic continuation from the in-gap edge modes to the bands of bulk modes. By varying the energy we find a transition from localized edge modes to delocalized bulk modes. We calculate the quasiparticle currents, discuss briefly how these currents couple to external currents, and predict how this can be used to control the quasiparticle modes.

Quasiparticle modes of charged particles, such as electrons, create electric currents. They can couple to external currents and external magnetic fields. This offers an opportunity to control quasiparticle modes with macroscopic fields. Moreover, edge currents in a superconductor couple to supercurrents through the Josephson effect. This provides an additional control of quasiparticle modes.

Double layer heterostructures of spatially separated two-dimensional electrons and holes have been a subject of considerable research interest for the last few decades. Inspired by the pairing effect in electron-hole double layers [1–6], we exploit the duality between the electron-hole double layer (EHDL) and the electron double layer (EEDL) to study superconductivity in the latter [7, 8]. Our physical system is a chiral double-layer without interlayer hopping. The electrons in such a system are subject to pairing due to geometric effects [7]. The goal is to study edge modes in a circular geometry with a circular hole on a closed surface or at the center of an infinite double layer. A central question addressed in the present paper concerns the existence of the bulk-edge correspondence for the spectra and the eigenfunctions [9, 10].

The EEDL comprises two spatially separated electronic layers with a positively charged extra layer, which acts as a gate and controls charge neutrality. The latter can either be an external gate or is provided by the positive charges inside the metallic layers. In both cases the entire system preserves charge neutrality. The electrons in the two layers repel each other due to the Coulomb interaction. A geometric constraint enables the electrons to form interlayer Cooper pairs. This is formally linked to the fact that the EEDL is dual to an EHDL [7, 11], where in the latter electron-hole pairs are created by an attractive interlayer Coulomb interaction [1]. The effect can be under-




FIG. 1. Top view of the EEDL. The effective attraction between electrons in the top (red) and the bottom layer (blue) is caused by geometry. The blue (red) square attracts the blue (red) electron in the center to the four red (blue) electrons in the corners.

stood as a geometrically induced attractive interaction between equally charged particle due to the double-layer structure. The Coulomb interaction repels the electrons in each layer, as well as the electrons in different layers. At minimal Coulomb energy we get an ordered structure as illustrated in Fig. 1. The electron at the center of the square is repelled by the four electrons in the corners of the square, which appears as if it is trapped. The same effect exist when we assume an attractive interaction between the center electron and the corner electrons.

The quasiparticles of the EEDL are described by the Bogoliubov de Gennes Hamiltonian of two layers with opposite chiralities [8]

$$H_{\text{EEDL}}(\Delta) = \begin{pmatrix} H & \Delta\sigma_2 \\ \Delta^*\sigma_2 & -H^* \end{pmatrix}, \quad (1)$$

where $H = h_1\sigma_1 + h_2\sigma_2 + h_3\sigma_3$ with Pauli matrices σ_j . $h_{1,2}$ are nearest neighbor tight-binding hopping terms and h_3 is a constant with an additional next-nearest neighbor hopping term. Δ is the spatially uniform pairing order parameter. For

the subsequent discussion we assume a honeycomb lattice with broken time-reversal invariance. The latter can either be caused by a periodic magnetic flux [12] or by a spin texture [13] that provides a single Dirac node in the dispersion. Then we can reduce $h_{1,2} \rightarrow i\partial_{1,2}$ and $h_3 \rightarrow 0$ near the Dirac node in the continuum limit. The Hamiltonian matrix in Eq. (1) can be mapped by a unitary transformation W onto a block-diagonal matrix with two Dirac matrices as (cf. App. A)

$$H_{\text{EEDL}}(\Delta) = W^\dagger H_{\text{Dirac}}(|\Delta|)W, \quad (2)$$

where H_{Dirac} is the 2×2 block diagonal matrix $\text{diag}(h_D(|\Delta|), h_D(-|\Delta|))$ with $h_D(|\Delta|) = i\partial_1\sigma_1 + i\partial_2\sigma_2 + |\Delta|\sigma_3$ and $\Delta = |\Delta|e^{i\phi}$. The unitary matrix W depends on the phase of Δ and is explained in Eq. (A1) of App. A:

$$W = \frac{1}{\sqrt{2}} \begin{pmatrix} e^{-i\phi/2}\sigma_0 & e^{i\phi/2}\sigma_1 \\ -e^{-i\phi/2}\sigma_0 & e^{i\phi/2}\sigma_1 \end{pmatrix}. \quad (3)$$

This mapping reveals that the Hamiltonian $H_{\text{EEDL}}(|\Delta|)$, which describes the coupling of the two layers by the pairing order parameter Δ , is equivalent to two Dirac Hamiltonians $h_D(\pm|\Delta|)$. Thus, the triality of the Hamiltonians H_{EEDL} , H_{EHDL} and H_{Dirac}

$$\begin{array}{ccc} H_{\text{EEDL}} & \longleftrightarrow & H_{\text{EHDL}} \\ & \searrow \quad \swarrow & \\ & H_{\text{Dirac}} & \end{array}$$

enables us to calculate the eigenfunctions of one Hamiltonian and transform them to any of the other Hamiltonians. The duality of H_{EEDL} and H_{EHDL} was used previously to explain pairing in an electronic double layer [7], while the mapping of Eq. (2) yields a connection to the massive Dirac Hamiltonian. For the following it is convenient to determine the eigenfunctions of the Dirac Hamiltonian first, since it separates for the two layers (cf. App. B).

Now we consider a circular geometry, assuming that each layer is either a disk or has a circular hole but is closed otherwise, forming a sphere, a torus, or is an infinite plane. In these cases we have a circular edge. Using polar coordinates $(x, y) = (r \cos \alpha, r \sin \alpha)$, we rewrite the Dirac Hamiltonian for one layer as

$$h_D(|\Delta|) = \begin{pmatrix} |\Delta| & ie^{-i\alpha}(\partial_r - \frac{i}{r}\partial_\alpha) \\ ie^{i\alpha}(\partial_r + \frac{i}{r}\partial_\alpha) & -|\Delta| \end{pmatrix} \quad (4)$$

and solve the eigenvalue problem $h_D(|\Delta|)\phi_E = E\phi_E$. The eigenfunctions for an infinite disk with



FIG. 2. Chiral electronic double layer: circular edges create counterflowing currents.

a circular hole of radius r_0 at the center can be expressed by modified Bessel functions $K_n(cr)$ as

$$\phi_{E,n}(r, \alpha) = A_n \begin{pmatrix} K_n(cr) \\ i\rho K_{n+1}(cr)e^{i\alpha} \end{pmatrix} e^{in\alpha} \quad (5)$$

with the conditions (see Eq. (B5) in App. B)

$$c\rho = E - |\Delta|, \quad -c = \rho(E + |\Delta|). \quad (6)$$

The eigenvalues are degenerate with respect to angular quantum number $n = 0, \pm 1, \dots$, since $h_D(|\Delta|)$ is rotational invariant and the angular momentum operator commutes with $h_D(|\Delta|)$. Subsequently we consider a solution with a fixed n , which is simultaneously an eigenfunction of $h_D(|\Delta|)$ and the angular momentum operator. After the removal of the global phase factor $e^{i(n+1/2)\alpha}$, which does not affect quadratic forms of the eigenfunctions in expectation values, we get from the eigenfunction in Eq. (5)

$$\tilde{\phi}_{E,n}(r, \alpha) = A_n \begin{pmatrix} K_n(cr)e^{-i\alpha/2} \\ i\rho K_{n+1}(cr)e^{i\alpha/2} \end{pmatrix}. \quad (7)$$

This function obeys the spinor relation $\tilde{\phi}_{E,n}(r, \alpha + 2\pi) = -\tilde{\phi}_{E,n}(r, \alpha)$; i.e., the spinor wave function reverses its sign after a full rotation.

The relation in Eq. (6) implies $\rho^2 = (|\Delta| - E)/(|\Delta| + E)$, the energy $E_c = \pm\sqrt{|\Delta|^2 - c^2}$ and $0 \leq c \leq |\Delta|$. ρ^2 is not symmetric with respect to E : it diverges for $E \sim -|\Delta|$ and it vanishes at $E = |\Delta|$. The sign of ρ is fixed by the relations (6), namely $\rho \leq 0$ due to $c \geq 0$. The non-symmetric form of ρ reflects the orthogonality of the wavefunctions for $\pm E$, while $E = 0$ is an exceptional point with two coalescent wavefunctions.

For a full disk of finite radius the solution is analogous but we must replace the modified Bessel function K_n by the modified Bessel function I_n in order to get a finite normalizable eigenfunction for $r \sim 0$ [14]. These solutions represent modes bound to the edge with energies $-|\Delta| \leq E \leq |\Delta|$.

Besides these edge solutions, we also obtain scattering solutions after the analytic continuation $c \rightarrow$




FIG. 3. Bulk and edge spectra for a complex wavenumber k with $\text{Im}(k) = c$.

$\pm ik$ ($k \geq 0$). The different signs do not affect the spectrum $E_k = \pm\sqrt{|\Delta|^2 + k^2}$ but yield different eigenfunctions: the relation [14]

$$\begin{aligned} K_n(-iz) &= \frac{\pi e^{i\pi(n+1)/2}}{2} [J_n(z) + iY_n(z)] \\ &= e^{-in\pi/2 - i\pi/4} \sqrt{2} \frac{e^{iz}}{\sqrt{\pi z}} + O\left(\frac{1}{z}\right), \end{aligned} \quad (8)$$

provides the bulk eigenfunctions in terms of the Bessel functions J_n and Y_n . Thus, the analytic continuation $c \rightarrow \pm ik$ yields a bulk-edge correspondence for the eigenfunctions and the spectrum. There is freedom to interpret k as a function of physical variables. In our example of Dirac particles we use the wavevector \mathbf{k} as $k = |\mathbf{k}|$. In general, the bulk-edge correspondence can connect the same edge mode with different bulk modes, which means that the edge modes are robust under a change of the bulk modes. The two values $\pm ik$ are important, since it provides a superposition of two eigenfunctions to match the boundary conditions. The energies of the bulk solutions $E_k = \pm\sqrt{|\Delta|^2 + k^2}$ represent the usual energy bands of massive Dirac particles in 2D (cf. Fig. 3). The sign of ρ is determined again by the relation (6), after the analytic continuation $c \rightarrow \pm ik$:

$$i\rho = \pm(E - |\Delta|)k = \pm k/(E + |\Delta|). \quad (9)$$

For the upper sign the top band $E \geq |\Delta|$ has $i\rho \geq 0$, while the bottom band $E \leq -|\Delta|$ has $i\rho \leq 0$. For the lower sign we get $i\rho \leq 0$ for the top band and $i\rho \geq 0$ for the bottom band. This reflects the opposite sign of the Chern numbers in the two bands [15].

The edge modes decay exponentially on the scale $1/c = 1/\sqrt{|\Delta|^2 - E^2}$. This scale diverges as E^2

approach the spectral boundary $|\Delta|^2$ of the edge modes. On the other hand, the wavevector of the bulk states vanishes as $k = \sqrt{E^2 - |\Delta|^2}$ for $E^2 \sim |\Delta|^2$. Therefore, $E = \pm|\Delta|$ are singular points in the spectrum, where the edge modes become extended over the entire 2D system. This is reminiscent of a localization-delocalization transition and reflects the critical energy dependence of the edge modes and its transition to bulk modes.

With the solution $\tilde{\phi}_{E,n}$ of the Dirac Hamiltonian in Eq. (7) we return to the eigenvalue problem of the Hamiltonian H_{EEDL} . The coefficients A_n and A'_n of the two chiral layers must be determined through boundary conditions and the normalization. Assuming that the layers are similar, we choose $|A'_n| = |A_n|$ and $A'_n = e^{i\psi} A_n$. Then the transformation in Eq. (2) gives the eigenfunction of H_{EEDL} for the circular edge in the double layer as $e^{i(n+1/2)\alpha} \Psi_{E,n}$ with

$$\Psi_{E,n} = \frac{|A_n|}{\sqrt{2}} \begin{pmatrix} (1 - e^{i\psi})K_n e^{-i\alpha/2} \\ i(\rho - e^{i\psi}\rho^{-1})K_{n+1} e^{i\alpha/2} \\ i(\rho + e^{i\psi}\rho^{-1})K_{n+1} e^{i\alpha/2} \\ (1 + e^{i\psi})K_n e^{-i\alpha/2} \end{pmatrix}. \quad (10)$$

The radial decay as well as the parameters c and ρ are inherited from the Dirac solution in Eq. (7). Therefore, the bulk-edge correspondence is also inherited from the Dirac solution. Moreover, the phase ψ enables us to control the contributions of the Dirac solutions of the top and the bottom layer. An example is $\psi = 0$, which suppresses the zero mode, where $\rho = -1$, in the top layer:

$$\Psi_{0,n} = \sqrt{2}|A_n| \begin{pmatrix} 0 \\ 0 \\ -iK_{n+1} e^{i\alpha/2} \\ K_n e^{-i\alpha/2} \end{pmatrix}. \quad (11)$$

Alternatively, the zero mode in the bottom layer is suppressed for $\psi = \pi$. The actual values of the boundary conditions are determined by the coupling of the double layer to its environment.

Charged quasiparticles, such as electrons, can be observed by the fact that they carry an electric current. This provides also an opportunity to create them by coupling the EEDL to an external current or magnetic field. The current operator for a single Dirac layer reads $j_\mu = (e/i\hbar)[h_D, r_\mu]$, which gives for $E^2 \leq |\Delta|^2$ the expectation value of the edge current density

$$\mathbf{j}_D = |A_n|^2 \rho K_n K_{n+1} \begin{pmatrix} -\sin \alpha \\ \cos \alpha \end{pmatrix}, \quad (12)$$

where $\rho < 0$ implies that the current flows clockwise. According to the relation (6), a sign reverse $|\Delta| \rightarrow -|\Delta|$ implies a sign reverse of ρ , which leads




FIG. 4. Bulk current density in a single chiral disk for $E > |\Delta|$ and $i\rho > 0$. For $E < -|\Delta|$ we get the opposite vorticity. This reflects the different Chern numbers of the top and the bottom band.

to a counter-clockwise current flow. The bulk current density is also represented by a vortex field, as illustrated in Fig. 4, which is a consequence of the angular momentum eigenfunction. The sign change of $i\rho$ when we switch from the top to the bottom band implies a reversal of the vorticity. This reflects the opposite signs of the Chern numbers in the two bands. Since the sign of $i\rho$ depends on the analytic continuation $c \rightarrow ik$ or $c \rightarrow -ik$, the vorticity of the edge currents agree either with the bulk currents of the top or the bottom band.

The quasiparticle current can also be calculated from the expectation value of the current operator $j_\mu = (e/i\hbar)[H_{\text{EEDL}}, r_\mu]$ with respect to $\Psi_{E,n}$. This gives for the edge current densities of the two layers

$$\mathbf{j}_{\uparrow,\downarrow} = C_{\uparrow,\downarrow} \begin{pmatrix} \cos(\alpha + \varphi_{\uparrow,\downarrow}) \\ \sin(\alpha + \varphi_{\uparrow,\downarrow}) \end{pmatrix}, \quad (13)$$

where \uparrow (\downarrow) refers to the top (bottom) layer. The coefficients read

$$C_{\uparrow,\downarrow} = \pm |A_n|^2 |(1 \mp e^{-i\psi})(\rho \mp e^{i\psi} \rho^{-1})| K_n K_{n+1}, \quad (14)$$

where the different signs imply opposite current directions in the two layers. The magnitude of the currents depends on the phase ψ . Moreover, the factor $K_n(cr)K_{n+1}(cr)$ is responsible for the radial decay of the current density. Finally, there is a phase shift:

$$\begin{aligned} \varphi_\uparrow &= \pi/2 + \arg[(1 - e^{-i\psi})(\rho - e^{i\psi} \rho^{-1})], \\ \varphi_\downarrow &= \pi/2 + \arg[(1 + e^{i\psi})(\rho + e^{-i\psi} \rho^{-1})]. \end{aligned} \quad (15)$$

In the special case of $E = 0$ we have $\rho = -1$, $\varphi_\uparrow = \varphi_\downarrow = \pi/2$ and $C_{\uparrow,\downarrow} = \pm 2|A_n|^2(1 \mp \cos \psi)K_n K_{n+1}$. Due to the different signs of $C_{\uparrow,\downarrow}$ the currents flow clockwise in the top layer and anti-clockwise in the bottom layer around the hole (cf. Fig. 2). These expressions indicate that the phase ψ of the boundary conditions can be used to control the edge currents.

The quasiparticle edge current density is balanced with the supercurrent density further away from edge, based on a continuity equation [8, 16–18], while the quasiparticle currents of the bulk eigenfunctions are independent of the supercurrents. Moreover, a macroscopic edge current radiates energy from the system to the environment. Therefore, quasiparticles would either disappear after some time, leading to $A'_n = A_n = 0$ as an equilibrium state, or they are balanced by an external supply of energy, for instance, by a magnetic flux.

In conclusion, we have studied quasiparticle edge modes in a superconducting EEDL. By exploiting the duality of the Hamiltonians H_{EEDL} and H_{Dirac} the simultaneous eigenfunctions of H_{EEDL} and the angular momentum operator were calculated. It turned out that the edge modes in the layers can be controlled separately, they can be switched off individually in the two layers by changing the boundary conditions at the edge. The edge modes carry a current density, which can couple to external fields. This predicts an experimental control of the quasiparticle edge modes. Moreover, an advantage of the superconducting EEDL is that the quasiparticle currents are balanced with the supercurrents as a result of charge conservation. In other words, the edge currents are maintained by the superconducting currents. Finally, there is a bulk-edge correspondence, which is provided by an analytic continuation. This correspondence characterizes two critical energies $E = \pm|\Delta|$, where a transition from edge modes to bulk modes and vice versa appear. These transitions are reminiscent of localization-delocalization transitions for the quasiparticle modes in disordered systems. An application of these ideas to a non-uniform pairing order parameter Δ is left as a project for the future. This could be of interest in the study of the Josephson effect in EEDL systems.

Acknowledgment: We are grateful for inspiring discussions with Andrii Iurov in the early stage of this work.

-
- [1] Yu. E. Lozovik and V. I. Yudson. A new mechanism for superconductivity: pairing between spatially separated electrons and holes. *Soviet Journal of Experimental and Theoretical Physics*, 44:389, 1976.
- [2] V. I. Yudson. Charged “few-electron–single spatially separated hole” complexes in a double quantum well near a metal plate. *Phys. Rev. Lett.*, 77:1564–1567, 1996.
- [3] A. V. Balatsky, Y. N. Joglekar, and P. B. Littlewood. Dipolar superfluidity in electron-hole bilayer systems. *Phys. Rev. Lett.*, 93:266801, 2004.
- [4] D. K. Efimkin, Yu. E. Lozovik, and A. A. Sokolik. Electron-hole pairing in a topological insulator thin film. *Phys. Rev. B*, 86:115436, 2012.
- [5] Z. Wang, D. A. Rhodes, K. Watanabe, T. Taniguchi, J. C. Hone, J. Shan, and K. F. Mak. Evidence of high-temperature exciton condensation in two-dimensional atomic double layers. *Nature*, 574(7776):76–80, 2019.
- [6] M. Zimmerman, R. Rapaport, and S. Gazit. Collective interlayer pairing and pair superfluidity in vertically stacked layers of dipolar excitons. *Proceedings of the National Academy of Sciences*, 119(30):e2205845119, 2022.
- [7] A. Sinner, Yu. E. Lozovik, and K. Ziegler. Pairing transition in a double layer with interlayer Coulomb repulsion. *Physical Review Research*, 2(3):033085, 2020.
- [8] K. Ziegler, A. Sinner, and Yu. E. Lozovik. Anomalous josephson effect of s -wave pairing states in chiral double layers. *Phys. Rev. Lett.*, 128:157001, 2022.
- [9] G. M. Graf and M. Porta. Bulk-edge correspondence for two-dimensional topological insulators. *Communications in Mathematical Physics*, 324(3):851–895, 2013.
- [10] J. Shapiro. The bulk-edge correspondence in three simple cases. *Reviews in Mathematical Physics*, 32(03):2030003, 2020.
- [11] K. Ziegler. Chiral josephson effect in double layers: The role of particle–hole duality. *Contributions to Plasma Physics*, 63(9-10):e202300014, 2023.
- [12] F. D. M. Haldane. Model for a quantum hall effect without landau levels: Condensed-matter realization of the “parity anomaly”. *Phys. Rev. Lett.*, 61:2015–2018, 1988.
- [13] A. Hill, A. Sinner, and K. Ziegler. Valley symmetry breaking and gap tuning in graphene by spin doping. *New Journal of Physics*, 13(3):035023, 2011.
- [14] M. Abramowitz and I. A. Stegun. *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*. Dover Publications, Inc., New York, 1964.
- [15] E. Fradkin. *Field Theories of Condensed Matter Physics*. Field Theories of Condensed Matter Physics. Cambridge University Press, 2013.
- [16] G. E. Blonder, M. Tinkham, and T. M. Klapwijk.

Transition from metallic to tunneling regimes in superconducting microconstrictions: Excess current, charge imbalance, and supercurrent conversion. *Phys. Rev. B*, 25:4515–4532, 1982.

- [17] A. Spuntarelli, P. Pieri, and G.C. Strinati. Solution of the bogoliubov–de gennes equations at zero temperature throughout the bcs–bec crossover: Josephson and related effects. *Physics Reports*, 488(4):111–167, 2010.
- [18] F. Pascucci, S. Conti, D. Neilson, J.s Tempere, and A. Perali. Josephson effect as a signature of electron-hole superfluidity in bilayers of van der waals heterostructures. *Phys. Rev. B*, 106:L220503, 2022.

Appendix A: Mapping of the H_{EEDL} Hamiltonian

We have the mapping $H_{\text{EEDL}} \rightarrow H_D = WH_{\text{EEDL}}W^\dagger$ as

$$\begin{aligned} & \begin{pmatrix} h_1\sigma_1 + h_2\sigma_2 & \Delta\sigma_2 \\ \Delta^*\sigma_2 & h_1\sigma_1 - h_2\sigma_2 \end{pmatrix} \\ & \rightarrow W \begin{pmatrix} h_1\sigma_1 + h_2\sigma_2 & \Delta\sigma_2 \\ \Delta^*\sigma_2 & h_1\sigma_1 - h_2\sigma_2 \end{pmatrix} W^\dagger \\ & = \begin{pmatrix} h_1\sigma_1 + h_2\sigma_2 + |\Delta|\sigma_3 & 0 \\ 0 & h_1\sigma_1 + h_2\sigma_2 - |\Delta|\sigma_3 \end{pmatrix} \end{aligned}$$

with $\Delta = |\Delta|e^{i\phi}$ and

$$W = \frac{1}{\sqrt{2}} \begin{pmatrix} e^{-i\phi/2}\sigma_0 & e^{i\phi/2}\sigma_1 \\ -e^{-i\phi/2}\sigma_0 & e^{i\phi/2}\sigma_1 \end{pmatrix}. \quad (\text{A1})$$

This implies the mapping of the eigenvector Φ_E of H_{Dirac} to the eigenvector Ψ_E of H_{EEDL} as

$$\Psi_E = W^\dagger \Phi_E. \quad (\text{A2})$$

Appendix B: Eigenfunctions of the Dirac Hamiltonian

With the ansatz

$$\Phi_{E,n}(r, \alpha) = \begin{pmatrix} f_n(cr) \\ g_n(cr)e^{i\alpha} \end{pmatrix} e^{in\alpha} \quad (\text{B1})$$

we can write for the eigenvalue problem $h_D\Phi_{E,n} = E\Phi_{E,n}$ the equation

$$\begin{pmatrix} i(cg'_n + \frac{n+1}{r}g_n) \\ i(cf'_n - \frac{n}{r}f_n) \end{pmatrix} = \begin{pmatrix} (E - |\Delta|)f_n \\ (E + |\Delta|)g_n \end{pmatrix}. \quad (\text{B2})$$

With the recurrence relations of the modified Bessel functions [14]

$$K'_0 = -K_1, \quad K'_1 = -K_0 - \frac{1}{cr}K_1$$

$$K'_n = -K_{n-1} - \frac{n}{cr}K_n = -K_{n+1} + \frac{n}{cr}K_n \quad (\text{B3})$$

we get for $f_n = K_n$ and $g_n = i\rho K_{n+1}$ the relations

$$\begin{pmatrix} ic(g'_n + \frac{n+1}{cr}g_n) \\ ic(f'_n - \frac{n}{cr}f_n) \end{pmatrix} = \begin{pmatrix} -c\rho(-K_n) \\ ic(-K_{n+1}) \end{pmatrix}$$

$$= \begin{pmatrix} (E - |\Delta|)K_n \\ (E + |\Delta|)i\rho K_{n+1} \end{pmatrix}, \quad (\text{B4})$$

where the second equation implies

$$c\rho = E - |\Delta|, \quad -c = \rho(E + |\Delta|). \quad (\text{B5})$$

Thus, we have $c^2 = |\Delta|^2 - E^2$ and $\rho^2 = (|\Delta| - E)/(|\Delta| + E)$, independent of n . Combining the eigenfunctions of the Dirac Hamiltonians $h_D(\pm|\Delta|)$, we get as the eigenfunction of H_{Dirac}

$$\Phi_{E,n} = \begin{pmatrix} AK_n(cr) \\ Ai\rho K_{n+1}(cr)e^{i\alpha} \\ A'K_n(cr) \\ A'i\rho^{-1}K_{n+1}(cr)e^{i\alpha} \end{pmatrix} e^{in\alpha}$$

and with Eq. (A2) the eigenfunction of H_{EEDL} as

$$\Psi_{E,n} = \frac{1}{\sqrt{2}} \begin{pmatrix} (A_n - A'_n)K_n \\ i(A_n\rho - A'_n\rho^{-1})e^{i\alpha}K_{n+1} \\ i(A_n\rho + A'_n\rho^{-1})e^{i\alpha}K_{n+1} \\ (A_n + A'_n)K_n \end{pmatrix} e^{in\alpha}. \quad (\text{B6})$$