

UNKNOTTEDNESS OF FREE BOUNDARY MINIMAL SURFACES AND SELF-SHRINKERS

SABINE CHU AND GIADA FRANZ

ABSTRACT. We study unknottedness for free boundary minimal surfaces in a three-dimensional Riemannian manifold with nonnegative Ricci curvature and strictly convex boundary, and for self-shrinkers in the three-dimensional Euclidean space. For doing so, we introduce the concepts of boundary graph for free boundary minimal surfaces and of graph at infinity for self-shrinkers. We prove that these surfaces are unknotted in the sense that any two such surfaces with isomorphic boundary graph or graph at infinity are smoothly isotopic.

1. INTRODUCTION

Let (M^3, g) be a three-dimensional Riemannian manifold, possibly noncompact or with nonempty boundary. Let Σ be a smooth surface that is properly embedded in M , possibly with boundary $\partial\Sigma = \Sigma \cap \partial M$. Recall that Σ is a *minimal surface* if its mean curvature H is equal to zero. Moreover, if Σ is minimal and meets ∂M orthogonally, we call it a *free boundary minimal surface*. Equivalently, Σ is a critical point of the area functional with respect to variations constraining the boundary of the surface to the boundary ∂M .

Lawson in [Law70] proved that minimal surfaces in the three-dimensional sphere S^3 are unknotted. Here, following Lawson's definition, we say that a surface is *unknotted* if there exists an ambient isotopy mapping the surface to a fixed standardly embedded surface with the same topology (see Section 2 for more precise definition). It is then natural to pose the same question for different ambient manifolds M . First, note that Lawson's proof works for minimal surfaces in simply connected three-dimensional manifolds with positive Ricci curvature (which are diffeomorphic to S^3 by result of Hamilton [Ham82, Theorem 1.1]). Moreover, Meeks–Yau showed in [MY92, Theorem 5.1] that complete minimal surfaces in \mathbb{R}^3 with finite topological type are unknotted. The result was then extended to remove the assumption of finite topological type in [FMI97] and [FM08].

Here, we consider two settings that are a priori quite different but actually display interesting similarities. Namely, we are interested in:

- Free boundary minimal surfaces in the unit ball $B^3 \subset \mathbb{R}^3$, or more in general in a compact three-dimensional Riemannian manifold (M^3, g) with nonnegative Ricci curvature and strictly convex boundary. Note that such a manifold M is diffeomorphic to the unit ball B^3 by [FL14, Theorem 2.11].
- Self-shrinkers with finite topology in \mathbb{R}^3 . Recall that these are minimal surfaces in \mathbb{R}^3 with respect to the Gaussian metric $e^{-|x|^2/4}g_{\mathbb{R}^3}$ and they appear as singularity models of the mean curvature flow (see [Hui90, Theorem 3.5]).

Unfortunately, the problem in these settings turns out to be more complicated. Consider for example the free boundary case. Note that in the unit ball $B^3 \subset \mathbb{R}^3$, there exist free boundary minimal surfaces of the same genus and number of boundary components for which there is no ambient diffeomorphism mapping one into the other. For example, the surfaces with three boundary components (and large genus) obtained in [KL21, Theorem 1.1] (see also [Keta, Theorem 1.1]) versus [KKMS, Theorem 1.2] are not ambient diffeomorphic. Note that Kapouleas and Li construct their surfaces by desingularizing the union of the critical catenoid and the equatorial disc, while the surfaces obtained by Karpukhin–Kusner–McGrath–Stern are doublings of the equatorial disc.

This means that these constructions differ in the arrangement of the boundary components on the sphere, as displayed in [Figure 1](#).

A similar behavior is expected from self-shrinkers, for which the ends play the role of the boundary components. Observe that Wang in [[Wan16](#), Theorem 1.1] proved a structure theorem for the ends of a self-shrinker in \mathbb{R}^3 . However, this is not as strong as the result for minimal surfaces in \mathbb{R}^3 , for which the geometric arrangement of the ends is always the same (see e.g. [[MY92](#), Theorem 4.1]).

These considerations suggest that any notion of unknottedness should take into account the arrangement of the boundary components of free boundary minimal surfaces and of the ends of self-shrinkers. To this purpose, following [[Fro92](#)], we introduce the concept of boundary graph for free boundary minimal surfaces and of graph at infinity for self-shrinkers and we proved that, for any two such surfaces with the same genus and isomorphic graph, there exists an ambient diffeomorphism mapping one into the other. In the next two sections we discuss in more detail the two settings separately.

1.1. Free boundary minimal surfaces. In this section, let us assume that (M^3, g) is a compact Riemannian manifold with nonnegative Ricci curvature and nonempty strictly convex boundary. The examples of free boundary minimal surfaces discussed above suggest the concept of *boundary graph* $\text{Graph}(\partial\Sigma, \partial M)$ for a properly embedded surface $\Sigma \subset M$. Essentially, we let the vertex set be the set of connected components of $\partial M \setminus \partial\Sigma$, and connect vertices whose corresponding components are both adjacent to some component of $\partial\Sigma$. See [Definition 2.1](#) for a more detailed definition. For instance, the boundary graphs for the surfaces in [[KL21](#)] and [[KKMS](#)] are shown in the last column of [Figure 1](#). In particular, observe that these two graphs are not isomorphic, therefore there cannot exist an ambient diffeomorphism mapping one surface into the other.




FIGURE 1. Schematic representation of the free boundary minimal surfaces in [[KL21](#)] (top row) and [[KKMS](#)] (bottom row), together with the arrangement of the boundary components on the sphere and the resulting boundary graphs. See [[Sch24](#)] for more precise numerical simulations of these surfaces.

However, given the notion of boundary graph, we are still able to prove that free boundary minimal surfaces in M are unknotted, in the following sense.

Theorem 1.1. *Let Σ, Σ' be (smooth, compact, properly embedded) free boundary minimal surfaces in a compact three-dimensional Riemannian manifold (M^3, g) with nonnegative Ricci curvature and strictly convex boundary. Assume that Σ and Σ' have the same genus and that their boundary graphs $\text{Graph}(\partial\Sigma, \partial M)$ and $\text{Graph}(\partial\Sigma', \partial M)$ are isomorphic. Then, Σ and Σ' are smoothly isotopic in the ambient manifold M^3 .*

In the case when Σ and Σ' have connected boundary or when they are topological annuli, [Theorem 1.1](#) follows from [[Mee81](#), Corollary 4]. There, Meeks proves that all minimal surfaces in B^3 with connected boundary are unknotted, and that all minimal surfaces in B^3 diffeomorphic to an annulus are isotopic to the critical catenoid. More in general, a notion of unknottedness of free boundary minimal surfaces in this setting was already proven in [[CF18](#), Corollary 3.3]. However, Choe–Fraser’s result does not contain a discussion on the boundary graphs. Indeed, what they prove is that a free boundary minimal surface Σ divides M into two handlebodies. Part of the proof is similar, but we decided to add the entire proof for completeness, together with the discussion on the boundary graphs.

Note that the free boundary condition in [Theorem 1.1](#) is necessary. Hall in [[Hal84](#)] showed that, if Σ and Σ' are two minimal surfaces of the same genus in the ball $B^3 \subset \mathbb{R}^3$ with the *same boundary*, they are not necessarily isotopic. In particular, he constructs Σ, Σ' with genus zero and equal boundary where Σ is unknotted and Σ' is knotted.

We conclude the discussion about the free boundary setting with an open question, which naturally arises in view of [Theorem 1.1](#).

Open question. *Which finite graphs can be realized as boundary graphs of free boundary minimal surfaces in the unit ball $B^3 \subset \mathbb{R}^3$ (or more in general in a three-dimensional manifold with non-negative Ricci curvature and strictly convex boundary)? How does the answer change if we fix the genus of the surface?*

Recall that Karpukhin–Kusner–McGrath–Stern proved in [[KKMS](#), Theorem 1.2] that, for every natural numbers $g \geq 0$, $b \geq 1$, there exists a free boundary minimal surface in the unit ball B^3 with genus g , b boundary components, and whose boundary graph is a star. Many other examples with different boundary graphs have also been constructed (see the introduction of [[FKS](#)] for a list of recent results). However, we do not yet have a complete answer to the open question above. What we can say so far is that boundary graphs of surfaces in this setting have to be trees (see [Remark 2.2](#)). Moreover, in the special case of a genus zero surface in B^3 , only very certain graphs can be attained. Indeed, recall that genus zero free boundary minimal surfaces are radial graphs by [[MZ24](#), Theorem 2.1]. This implies that $\text{Graph}(\partial\Sigma, \partial B^3)$ must be a star.

1.2. Self-shrinkers. In this section, our ambient manifold is $(\mathbb{R}^3, e^{-|x|^2/4}g_{\mathbb{R}^2})$. As mentioned above, a minimal surface Σ in this space is a self-shrinker. Inspired by the case of free boundary minimal surfaces, we can define the graph at infinity $\text{Graph}(\partial\Sigma, \infty)$ of a self-shrinker Σ as $\text{Graph}(\Sigma \cap \partial B_R(0), \partial B_R(0))$ for any radius $R > 0$ sufficiently large. Thanks to the structure theorem [[Wan16](#), Theorem 1.1], this notion of graph at infinity is well-defined (see [Section 4](#) for more details).

In analogy to the free boundary case, we can prove the following result.

Theorem 1.2. *Let Σ, Σ' be (smooth, complete, properly embedded) self-shrinkers in \mathbb{R}^3 with finite topological type. Assume that Σ and Σ' have the same genus and that their graphs at infinity $\text{Graph}(\partial\Sigma, \infty)$ and $\text{Graph}(\partial\Sigma', \infty)$ are isomorphic. Then, Σ and Σ' are smoothly isotopic in \mathbb{R}^3 .*

Note that unknottedness of compact self-shrinkers and self-shrinkers with one or two asymptotically conical ends was proven in [[MW20](#), Theorem 1.1], [[Mra24](#), Theorem 1.1], [[Mra](#), Corollary 1.2]. Here, we generalize these results to any (finite) number of ends.

Similarly to the free boundary case, one can wonder which graphs can arise as graphs at infinity of self-shrinkers.

Open question. *Which graphs can be realized as graphs at infinity of self-shrinkers in \mathbb{R}^3 ? How does the answer change if we fix the genus of the surface?*

Again, very little is known in this direction apart from the genus zero case. In fact, the only genus zero self-shrinkers are the plane, the self-shrinking sphere and the self-shrinking cylinder by [Bre16, Theorem 2], for which the graph at infinity has two, one and three vertices, respectively. This suggests that the self-shrinkers case is probably more rigid than the free boundary one. We refer to the introduction of [Ketb] for a discussion on the known examples of self-shrinkers in \mathbb{R}^3 .

1.3. Key ideas of the proofs. Unknottedness of (free boundary) minimal surfaces in an ambient manifold (M^3, g) is closely related to the validity of the Frankel property for minimal surfaces in such manifold. Indeed, thanks to [Fro92, Theorem 2.1], the key step to prove Theorem 1.1 and Theorem 1.2 is to show that a surface Σ as in the statements is a Heegaard splitting of M in the sense of [Fro92, Section 1]. Namely, the inclusions $\pi_1(\Sigma) \hookrightarrow \pi_1(C_i)$ are surjective for $i = 1, 2$, where C_1, C_2 are the two connected components of $M \setminus \Sigma$. It turns out that this is equivalent to proving a Frankel property in the universal Riemannian cover \tilde{C}_i of C_i for $i = 1, 2$, which has similar geometric properties as the ambient manifold M .

The Frankel property for free boundary minimal surfaces in a three-dimensional manifold with nonnegative Ricci curvature and strictly convex boundary was proven by Fraser–Li in [FL14, Lemma 2.4]. For self-shrinkers in \mathbb{R}^3 , the Frankel property was proven in increasing generality in [IPR21], [CM23], and [NZ24]. Here we make use of the robust and general statement [NZ24, Proposition 25] to prove the adaptations needed in our setting. The free boundary setting is treated in Section 3, while the self-shrinker case is discussed in Section 4. In Section 2, we prove some topological preliminaries related to [Fro92] about Heegaard splittings, boundary graphs, and unknottedness.

Acknowledgements. We would like to thank Ailana Fraser and Ursula Hamenstädt for their interest and support. Moreover, we would like to thank Alex Mramor and Shengwen Wang for introducing us to the problem and pointing out reference [Mee81], Keaton Naff for interesting discussions on the Frankel property, and Jonathan Zung for answering our topological questions.

G.F. was partially supported by NSF grant DMS-2405361. Moreover, part of this work was performed while G.F. was in residence at the Simons Laufer Mathematical Sciences Institute (formerly MSRI) during the Fall 2024 semester, supported by NSF grant DMS-1928930.

2. TOPOLOGICAL PRELIMINARIES

In this section, we collect the topological results about unknottedness needed in the paper. In particular, we consider a three-dimensional differentiable manifold M^3 with boundary. We assume that M is diffeomorphic to the closed unit ball $B^3 \subset \mathbb{R}^3$ and we say that M is a *differentiable ball*. We will apply the results in this section to compact Riemannian manifolds with nonnegative Ricci curvature and strictly convex boundary (diffeomorphic to B^3 by [FL14, Theorem 2.11]), or to Euclidean balls $B_R(0) \subset \mathbb{R}^3$. Moreover, unless otherwise stated, we assume that $\Sigma \subset M$ is a smooth, compact, connected surface with boundary, properly embedded in M .

We now start by defining the notion of boundary graph, we then proceed discussing Heegaard splittings and unknottedness.

Definition 2.1 (cf. [Fro92, Section 2]). Let $C = \{\gamma_1, \gamma_2, \dots, \gamma_n\}$ be a set of continuous disjoint Jordan curves lying on a smooth surface S . Define the *boundary graph* $\text{Graph}(C, S)$ as follows. The vertices of $\text{Graph}(C, S)$ correspond to the connected components of $S \setminus C$, and two vertices are connected by an edge if and only if the closure of the corresponding components of $S \setminus C$ both contain some $\gamma_i \in C$.

Remark 2.2. Let M^3 be a three-dimensional differentiable ball and let $\Sigma \subset M$ be a properly embedded surface with $\partial\Sigma = \Sigma \cap \partial M$. Then every boundary component of $\partial\Sigma$ divides ∂M in two connected components. Therefore, the boundary graph $\text{Graph}(\partial\Sigma, \partial M)$ is actually a tree.

Note also that we can lower bound the number of isomorphism classes of trees with n vertices by $\frac{n^{n-2}}{n!}$ (by Cayley’s formula, the number of labelled trees on n vertices is n^{n-2} , and there are $n!$ possible labelings), so the questions brought up in the introduction on the possible finite graphs that can be realized as boundary graphs or graphs at infinity is nontrivial.

Definition 2.3 (cf. [Fro92, Section 1]). We say that a (smooth, compact, connected, properly embedded) surface Σ in a compact three-dimensional differentiable manifold M^3 is a *Heegaard splitting* if $M \setminus \Sigma$ consists of two connected components C_1 and C_2 such that C_1, C_2 are irreducible (i.e., every sphere embedded in C_i bounds a ball), and the inclusion maps $\pi_1(\Sigma) \hookrightarrow \pi_1(C_i)$ are surjective for $i = 1, 2$.

Lemma 2.4. *Let M^3 be a three-dimensional differentiable ball and let Σ be a (smooth, compact, connected, properly embedded) surface in M . Then Σ is a Heegaard splitting of M if and only if the inclusion maps $\pi_1(\Sigma) \hookrightarrow \pi_1(C_i)$ are surjective for $i = 1, 2$.*

Proof. Observe that, if M is a differentiable ball, then Σ is two-sided (by e.g. [CKM17, Lemma C.1]) and thus it divides M into two connected components C_1 and C_2 . Moreover, C_1 and C_2 are irreducible: if C_1 was reducible, then there would exist some sphere in C_1 whose inside enclosed a portion of C_2 (because M is irreducible), but then this would mean there was a connected component of Σ inside the sphere that was disjoint from the rest of the surface, which is a contradiction. \square

Remark 2.5. Fix $i = 1$ or $i = 2$, and let $\Pi: \tilde{C}_i \rightarrow C_i$ be the universal cover of C_i . If $\Pi^{-1}(\Sigma)$ is connected, then $\pi_1(\Sigma) \hookrightarrow \pi_1(C_i)$ is surjective. To see this, consider any loop in C_i and lift it to a loop in \tilde{C}_i . As \tilde{C}_i is simply connected, this is in the same homotopy class as some loop on $\Pi^{-1}(\Sigma)$ (which exists because $\Pi^{-1}(\Sigma)$ is connected). Then we can project down to a loop on Σ , so we have surjectivity.

Note that [Definition 2.3](#) is not the classical definition of a Heegaard splitting, which usually does not require the inclusion maps of the fundamental groups to be surjective. Importantly, this requirement implies that the surface is “unknotted” in some way, as we are about to prove. Indeed, we show that if a surface is a Heegaard splitting then it is smoothly isotopic to a paradigmatic unknotted surface, which looks like a thickened unknotted graph and is rigorously defined as follows.

Definition 2.6 (Model surface). Given a tree T and a number $g \in \mathbb{N}$, we define a *model surface* Σ in B^3 with boundary graph T and genus g as follows (see also [Figure 2](#)).




FIGURE 2. Schematical representation of a model surface with prescribed genus and boundary graph.

Choose a node v_0 to be the root of the tree, and denote by v_1, \dots, v_k its direct children for some $k \in \mathbb{N}$. Let S be a sphere centered in the origin and with radius $1/8$. Moreover, take k disjoint flat discs $D_1, \dots, D_k \subset B^3$ with the same radius and with $\partial D_i = D_i \cap \partial B^3$ for $i = 1, \dots, k$. Now, let r_i be the radial segment connecting the center of D_i with S and let C_i be a small tubular neighborhood of r_i with $\partial C_i \subset D_i \cup S$. Assume that C_1, \dots, C_k have the same radius for $i = 1, \dots, k$. Then, consider the surface Σ_1 obtained as the union of S, D_1, \dots, D_k and C_1, \dots, C_k , after removing the small discs bounded by ∂C_i on D_i and S for $i = 1, \dots, k$.

Now, consider the vertex v_i for some $i = 1, \dots, k$. Let k_i be the number of direct children of v_i , which are at distance 2 from the root v_0 . Consider k_i disjoint flat discs in B^3 with boundary contained in the smallest of the components of $\partial B^3 \setminus \partial D_i$. Moreover, similarly as before, connect the center of each of these discs with D_i with small cylindrical bridges. We choose the discs and the bridges of sufficiently small radius to ensure no self-intersections. Repeat this procedure for each $i = 1, \dots, k$ and let Σ_2 be the surface obtained as the union of Σ_1 and all these new discs and cylindrical bridges.

We then repeat this procedure for the vertices in T at distance 3 from the root v_0 , to get a surface Σ_3 . And so on, we repeat the procedure until each nonroot vertex of T corresponds to some disc connected with a cylindrical bridge to the disc corresponding to its mother in the tree. We let Σ' be the surface that we obtain in this way. Finally, let Σ be the surface obtained by taking the connected sum of Σ' with a standardly embedded genus g surface. In particular, we can assume that the genus g surface is attached to the sphere S and it is small enough so that it does not intersect Σ' . The surface Σ is not smooth at the intersection between the discs and the bridges, but it can be smoothed out easily.

Proposition 2.7. *The model surface in Definition 2.6 is a Heegaard splitting.*

Proof. By Lemma 2.4, it suffices to show that if $\Sigma \subset B^3$ is a model surface of genus g with boundary graph a tree T , then $\pi_1(\Sigma) \hookrightarrow \pi_1(C_i)$ is surjective for $i = 1, 2$, where C_1 and C_2 are the two connected components of $B^3 \setminus \Sigma$.

Now, take two points $p_1 \in C_1$ and $p_2 \in C_2$ such that p_1, p_2 are contained in a small tubular neighborhood of the sphere S . For every cylindrical bridge in Σ , consider a simple loop winding around the bridge and contained in $B^3 \setminus \Sigma$. Note that each loop is either contained in C_1 or C_2 . Then, we can connect each loop with a simple curve to p_1 if the loop is in C_1 , or to p_2 if the loop is in C_2 . Let us denote by $\rho_1 \subset C_1$ and $\rho_2 \subset C_2$ the two graphs obtained in this way. Finally, add to ρ_1 one simple loop in C_1 (based in p_1) winding around each hole of the genus g surface, and do the same with ρ_2 .

Then, C_1 and C_2 can be retracted to the graphs ρ_1 and ρ_2 , respectively. Therefore, the loops in the constructions of ρ_1 and ρ_2 are generators of the fundamental group of C_1 and C_2 , respectively. Moreover, note that every loop in ρ_1 and ρ_2 can be isotoped to a loop in Σ . As a result, $\pi_1(\Sigma) \hookrightarrow \pi_1(C_i)$ is surjective for $i = 1, 2$. \square

Theorem 2.8. *Let Σ be a (smooth, compact, connected, properly embedded) Heegaard splitting in a three-dimensional differentiable ball M^3 . Then Σ is unknotted, meaning that it is smoothly isotopic in M to the model surface from Definition 2.6 with the same genus and the same boundary graph. In particular, every pair of Heegaard splittings with the same genus and the same boundary graph are smoothly isotopic.*

Proof. Let Σ' be the model surface defined in Definition 2.6 with the same genus and same boundary graph as Σ . By Proposition 2.7, Σ' is a Heegaard splitting of M too. Therefore, Frohman in [Fro92, Theorem 2.1] showed that there exists a homeomorphism $\tilde{f}: M \rightarrow M$ such that $\tilde{f}(\Sigma) \rightarrow \Sigma'$.

As a first step, we now want to smooth out \tilde{f} to get a diffeomorphism. Define the surfaces $X = \Sigma \cup \partial M$ and $X' = \Sigma' \cup \partial M$. Thanks to [Mun56, Chapter V], $\tilde{f}|_X: X \rightarrow X'$ is isotopic to a diffeomorphism $h_X: X \rightarrow X'$. This can be seen by applying Munkres' smoothing procedure first to $\tilde{f}|_{\partial M}$, sending $\partial \Sigma$ to $\partial \Sigma'$, and then to $\tilde{f}|_\Sigma$.

By [EK71, Corollary 1.2], \tilde{f} is isotopic to a homeomorphism $h: M \rightarrow M$ that coincides with h_X on X . Moreover, we can assume that h is a diffeomorphism from a tubular neighborhood $N(X)$

of X to a tubular neighborhood $N(X')$ of X' . This can be achieved by interpolating h_X in the tubular neighborhood $N(X)$.

Finally, the map $h: M \rightarrow M$ is isotopic to a diffeomorphism $f: M \rightarrow M$, that coincides with h on $N(X)$, thanks to [Cer59, Theorem 5 and Corollary 1] (which assumes the Smale's conjecture, now proven in Hatcher [Hat83] and Bamler–Kleiner [BK23]).

We are left to prove that Σ and Σ' are smoothly isotopic. Note that, by [Hat83, Appendix 1], the group of diffeomorphisms of B^3 that fix the boundary $\partial B^3 = S^2$ is contractible. Moreover, the group of diffeomorphisms of S^2 has two connected components corresponding respectively to the orientation preserving and orientation reversing diffeomorphisms (see [Sma59]). As a result, the diffeomorphism f is smoothly isotopic to either the identity or to a reflection across a plane. This implies that the surface Σ is smoothly isotopic to either Σ' or the reflection of Σ' across a plane. However, note that it is possible to perform the construction of Σ' in Definition 2.6 in such a way that the model surface is symmetric with respect to the reflection across a plane (by choosing the discs to be centered on such plane). This concludes the proof. \square

3. FREE BOUNDARY MINIMAL SURFACES ARE HEEGAARD SPLITTINGS

In this section, we prove the unknottedness of free boundary minimal surfaces stated in Theorem 1.1. Thanks to the topological preliminaries in the previous section, the theorem is a consequence of the following result.

Theorem 3.1 (cf. [CF18, Lemma 3.1 and Theorem 3.2]). *Let Σ be a (smooth, compact, properly embedded) free boundary minimal surface in a three-dimensional Riemannian manifold M^3 fulfilling the hypotheses of Theorem 1.1. Then Σ is a Heegaard splitting of M .*

Proof. Recall that M is diffeomorphic to a three-dimensional ball by [FL14, Theorem 2.11]. Moreover, note that Σ is connected by the Frankel property for free boundary minimal surfaces in M , proved in [FL14, Lemma 2.4]. Therefore, by Lemma 2.4, to show that Σ is a Heegaard splitting we just need to prove that the inclusion maps $\pi_1(\Sigma) \hookrightarrow \pi_1(C_i)$ are surjective for $i = 1, 2$. Without loss of generality, inspired by [Law70], we let $i = 1$ and consider the universal Riemannian cover $\Pi: \tilde{C}_1 \rightarrow C_1$. By Remark 2.5, it is then sufficient to prove that $\Pi^{-1}(\Sigma)$ is connected.

Assume by contradiction that it is not, and denote by $\Sigma_1, \dots, \Sigma_N$ the connected components of $\Pi^{-1}(\Sigma)$. Then $\Pi^{-1}(\partial C_1) = (\cup_{k=1}^N \Sigma_k) \cup \Gamma$ for $\Gamma = \Pi^{-1}(\partial M \cap \partial C_1)$ (note that Γ may be disconnected). Inspired by [FL14, proof of Lemma 2.4], we define $d_k(x)$ to be the distance from a point $x \in \tilde{C}_1$ to Σ_k , and, for every $k \neq \ell = 1, \dots, N$, we consider

$$D := \inf_{k \neq \ell = 1, \dots, N} \inf_{x \in \tilde{C}_1} d_k(x) + d_\ell(x).$$

Without loss of generality, we can assume that the first infimum is realized for $k = 1, \ell = 2$, as k and ℓ varies in $1, \dots, N$. Consider some minimizing sequence $(x_m)_{m \in \mathbb{N}} \subset \tilde{C}_1$ for $d_1 + d_2$, namely $d_1(x_m) + d_2(x_m) \rightarrow \inf_{x \in \tilde{C}_1} d_1(x) + d_2(x) = D$ as $m \rightarrow \infty$. Fix some $\tilde{p}_0 \in \tilde{C}_1$ and let

$$\mathcal{B}_0 = \{\tilde{p} \in \tilde{C}_1 : d(\tilde{p}, \tilde{p}_0) \leq 2 \operatorname{diam}(C_1)\}.$$

Then, for each x_m , there exists $x'_m \in \mathcal{B}_0$ such that $\Pi(x_m) = \Pi(x'_m)$, i.e., there is an isometry of \tilde{C}_1 mapping x_m to x'_m . Therefore, possibly renaming x_m (to be x'_m), we can assume without loss of generality that the sequence $(x_m)_{m \in \mathbb{N}}$ is contained in \mathcal{B}_0 . This ball is compact, so up to subsequence $(x_m)_{m \in \mathbb{N}}$ converges to some x_0 realizing the infimum, namely

$$d_1(x_0) + d_2(x_0) = \lim_{m \rightarrow \infty} d_1(x_m) + d_2(x_m) = \inf_{x \in \tilde{C}_1} d_1(x) + d_2(x) = D.$$

Therefore, we can apply [NZ24, Proposition 25] on \tilde{C}_1 with $N = \Gamma$, $f = 0$, $\kappa = 0$, $1/\alpha = 0$ (note that \tilde{C}_1 has nonnegative Ricci curvature and Γ is convex in \tilde{C}_1) and we obtain a contradiction. Indeed, observe that \tilde{C}_1 cannot be a product manifold $\Sigma_1 \times [0, d]$ since Γ is strictly convex in \tilde{C}_1 .

So we have proved that Σ_1, Σ_2 cannot be disjoint, meaning that $\Pi^{-1}(\Sigma)$ is connected and $\pi_1(\Sigma) \hookrightarrow \pi_1(C_1)$ is surjective. Thus, this concludes the proof that Σ is a Heegaard splitting. \square

Proof of Theorem 1.1. The result follows from Theorem 3.1 together with Theorem 2.8, from the previous section. \square

4. SELF-SHRINKERS ARE HEEGAARD SPLITTINGS

Let $\Sigma \subset \mathbb{R}^3$ be a self-shrinker of the mean curvature flow. Equivalently, Σ is a minimal surface with respect to the Gaussian metric $e^{-|x|^2/4}g_{\mathbb{R}^3}$ on \mathbb{R}^3 . Let us further assume that Σ has finite topology. Then, by [Wan16, Theorem 1.1], each end of Σ is smoothly asymptotic to either a regular cone or a round cylinder. As a consequence, there exist $R_0 > 0$ such that for every $R \geq R_0$ the graphs $\text{Graph}(\Sigma \cap \partial B_R(0), \partial B_R(0))$ are all isomorphic. Then, it makes sense to define the graph of Σ at infinity, denoted $\text{Graph}(\partial\Sigma, \infty)$, to be one of these isomorphic graphs.

As a result, to study the topology of a self-shrinker is sufficient to look at the portion of the self-shrinker inside a sufficiently large ball. In fact, we prove that the self-shrinker is an Heegaard splitting in every sufficiently large Euclidean ball.

Theorem 4.1. *Let Σ be a (smooth, complete, properly embedded) self-shrinker in \mathbb{R}^3 and take $M^3 = B_R(0)$ for some $R \geq 2\sqrt{2}$. Then $\Sigma \cap M$ is a Heegaard splitting of M .*

Proof. First observe that $\Sigma \cap M$ is connected by the Frankel property for self-shrinkers, proved in [CM23, Corollary 0.3] and in [NZ24, Theorem 3] with the explicit radius $2\sqrt{2}$. Therefore, by Lemma 2.4, we just need to prove that the inclusion maps $\pi_1(\Sigma) \hookrightarrow \pi_1(C_i)$ are surjective for $i = 1, 2$.

Without loss of generality, let $i = 1$ and consider the universal Riemannian cover $\Pi: (\tilde{C}_1, \tilde{g}_0) \rightarrow (C_1, g_{\mathbb{R}^3})$. By Remark 2.5, $\pi_1(\Sigma) \hookrightarrow \pi_1(C_1)$ is surjective if and only if $\Pi^{-1}(\Sigma)$ is connected. Assume by contradiction that $\Pi^{-1}(\Sigma)$ is not connected, and denote by $\Sigma_1, \dots, \Sigma_N$ the connected components of $\Pi^{-1}(\Sigma)$. We will now reach a contradiction by proving a Frankel property for the surfaces $\Sigma_1, \dots, \Sigma_N$ on \tilde{C}_1 , inspired by [NZ24, Section 6.2 and Proposition 25].

Consider the function $\tilde{\rho}: \tilde{C}_1 \rightarrow [0, \infty)$ given by $\tilde{\rho}(y) = R^2 - |\Pi(y)|^2$, where we recall that R is the radius of the ball $B_R(0) = M$. Moreover, let $\tilde{\gamma}: \tilde{C}_1 \rightarrow [0, \infty)$ be defined as $\tilde{\gamma}(y) = |\Pi(y)|^2/4$. Let $\tilde{f}: \tilde{C}_1 \rightarrow (-\infty, \infty]$ be such that $e^{\tilde{f}} = \tilde{\rho}^{-2}$ and consider the metric $\tilde{g}_1 = e^{\tilde{f}}\tilde{g}_0$ on \tilde{C}_1 . Observe that $\tilde{f} = \infty$ on $\Gamma = \Pi^{-1}(\partial M \cap \partial C_1)$ (note that $\Pi^{-1}(\partial C_1) = (\cup_{k=1}^N \Sigma_k) \cup \Gamma$ and Γ may be disconnected), so the idea to add this weight \tilde{f} is to “send the boundary Γ at ∞ ”.

Note that what we are doing is to consider the same functions as in [NZ24, Section 6.2] but lifted to the universal Riemannian cover of \tilde{C}_1 . Indeed, with their notation of ρ, γ, f , we have that $\tilde{\rho} = \rho \circ \Pi$, $\tilde{\gamma} = \gamma \circ \Pi$ and $\tilde{f} = f \circ \Pi$. In particular, as observed in [NZ24, Section 6.2], $\Sigma_1, \dots, \Sigma_N$ are minimal surfaces in $(\tilde{C}_1, \tilde{g}_0)$ with respect to the weight $e^{-\tilde{\gamma}}$ (so we say that they are $\tilde{\gamma}$ -minimal surfaces in $(\tilde{C}_1, \tilde{g}_0)$). Equivalently, $\Sigma_1, \dots, \Sigma_N$ are minimal surfaces in $(\tilde{C}_1, \tilde{g}_1 = e^{\tilde{f}}\tilde{g}_0)$ with respect to the weight $e^{-\tilde{\gamma}-\tilde{f}}$ (say $(\tilde{\gamma} + \tilde{f})$ -minimal surfaces in $(\tilde{C}_1, \tilde{g}_1)$). Moreover, by the computation in [NZ24, Section 6.2], the manifold $(\tilde{C}_1, \tilde{g}_1)$ satisfies that the following (-2) -Bakry-Émery-Ricci curvature is nonnegative:

$$\widetilde{\text{Ric}}_{\tilde{\gamma}+\tilde{f}}^{-2} \geq 0,$$

whenever $R \geq 2\sqrt{2}$.

Now, we define the function $d_k(x)$ to be the distance from a point $x \in \tilde{C}_1$ to Σ_k , for $k = 1, \dots, N$. Similarly to the proof of Theorem 3.1, without loss of generality we get that

$$\inf_{k \neq \ell = 1, \dots, N} \inf_{x \in \tilde{C}_1} d_k(x) + d_\ell(x)$$

is achieved by $k = 1, \ell = 2$ and a point $x_0 \in \tilde{C}_1 \setminus \Gamma$. As a result, we can apply [NZ24, Proposition 25] on $(\tilde{C}_1, \tilde{g}_1, e^{-\tilde{f}})$ with $\kappa = 0$ and $\alpha = -2$ (note that there is no free boundary here), and we obtain that $d_1 + d_2$ is constant and Σ_1, Σ_2 are totally geodesic with respect to the metric $e^{-\tilde{f}}\tilde{g}_1 = \tilde{g}_0 = \Pi^*g_{\mathbb{R}^3}$. By unique continuation, this implies that the initial surface Σ is totally geodesic with

respect to the Euclidean metric on \mathbb{R}^3 , namely it is a plane. However, we know that the only self-shrinking plane is the one passing through the origin and in this case we also know that $\Pi^{-1}(\Sigma)$ is connected. This concludes the proof, showing that $\Pi^{-1}(\Sigma)$ is connected and therefore $\Sigma \cap M$ is a Heegaard splitting of M . \square

Proof of Theorem 1.2. Let Σ and Σ' be two self-shrinkers in \mathbb{R}^3 with the same genus and with isomorphic graphs at infinity. Thanks to the structure theorem [Wan16, Theorem 1.1], there exists $R \geq 2\sqrt{2}$ sufficiently large such that $\Sigma \setminus B_R(0)$ is smoothly isotopic in $\mathbb{R}^3 \setminus B_R(0)$ to the cone over $\Sigma \cap \partial B_R(0)$, through an isotopy fixing $\partial B_R(0)$. Let $H_1: [0, 1] \times \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be such isotopy of \mathbb{R}^3 , which is the identity on $B_R(0)$. Similarly, we can assume R to be also sufficiently large such that there exists a smooth isotopy $H_2: [0, 1] \times \mathbb{R}^3 \rightarrow \mathbb{R}^3$ that is the identity on $B_R(0)$ and maps $\Sigma' \setminus B_R(0)$ to the cone over $\Sigma' \cap \partial B_R(0)$.

If we further assume $R \geq 2\sqrt{2}$, thanks to Theorem 1.2 and Theorem 2.8, there exists a smooth isotopy $H_0: [0, 1] \times \overline{B_R(0)} \rightarrow \overline{B_R(0)}$ of $B_R(0)$ mapping Σ to Σ' . Let us extend H_0 to an isotopy $\tilde{H}_0: [0, 1] \times \mathbb{R}^3 \rightarrow \mathbb{R}^3$, by rescaling in $\mathbb{R}^3 \setminus B_R(0)$. Namely, $\tilde{H}_0(t, x) = \frac{|x|}{R} H_0(t, R \frac{x}{|x|})$ for all $x \in \mathbb{R}^3 \setminus B_R(0)$. Then, the isotopy $H: [0, 1] \times \mathbb{R}^3 \rightarrow \mathbb{R}^3$ defined as $H(t, \cdot) = (H_2(t, \cdot))^{-1} \circ \tilde{H}_0(t, \cdot) \circ H_1(t, \cdot)$ maps Σ to Σ' , which is the desired result. \square

REFERENCES

- [BK23] R. H. Bamler and B. Kleiner, *Ricci flow and diffeomorphism groups of 3-manifolds*, J. Amer. Math. Soc. **36** (2023), no. 2, 563–589.
- [Bre16] S. Brendle, *Embedded self-similar shrinkers of genus 0*, Ann. of Math. (2) **183** (2016), no. 2, 715–728.
- [Cer59] J. Cerf, *Groupes d'automorphismes et groupes de difféomorphismes des variétés compactes de dimension 3*, Bulletin de la Société Mathématique de France **87** (1959), 319–329.
- [CF18] J. Choe and A. Fraser, *Mean curvature in manifolds with Ricci curvature bounded from below*, Comment. Math. Helv. **93** (2018), no. 1, 55–69.
- [CKM17] O. Chodosh, D. Ketover, and D. Maximo, *Minimal hypersurfaces with bounded index*, Invent. Math. **209** (2017), no. 3, 617–664.
- [CM23] T. H. Colding and W. P. Minicozzi II, *A strong Frankel Theorem for shrinkers*, preprint (2023), available at [arXiv:2306.08078](https://arxiv.org/abs/2306.08078).
- [EK71] R. D. Edwards and R. C. Kirby, *Deformations of spaces of imbeddings*, Ann. of Math. (2) **93** (1971), 63–88.
- [FKS] G. Franz, D. Ketover, and M. B. Schulz, *Genus one critical catenoid*, preprint, available at [arXiv:2409.12588](https://arxiv.org/abs/2409.12588).
- [FL14] A. Fraser and M. M.-C. Li, *Compactness of the space of embedded minimal surfaces with free boundary in three-manifolds with nonnegative Ricci curvature and convex boundary*, J. Differential Geom. **96** (2014), no. 2, 183–200.
- [FM08] C. Frohman and W. H. Meeks III, *The topological classification of minimal surfaces in \mathbb{R}^3* , Ann. of Math. (2) **167** (2008), no. 3, 681–700.
- [FMI97] C. Frohman and W. H. Meeks III, *The topological uniqueness of complete one-ended minimal surfaces and Heegaard surfaces in \mathbb{R}^3* , Journal of the American Mathematical Society (1997), 495–512.
- [Fro92] C. Frohman, *Heegaard splittings of the three-ball*, Topology '90 (Columbus, OH, 1990), 1992, pp. 123–131.
- [Hal84] P. Hall, *Two topological examples in minimal surface theory*, J. Differential Geom. **19** (1984), 475–81.
- [Ham82] R. S. Hamilton, *Three-manifolds with positive Ricci curvature*, J. Differential Geometry **17** (1982), no. 2, 255–306.
- [Hat83] A. E. Hatcher, *A proof of the Smale conjecture*, $\text{Diff}(S^3) \simeq \text{O}(4)$, Ann. of Math. (2) **117** (1983), no. 3, 553–607.
- [Hui90] G. Huisken, *Asymptotic behavior for singularities of the mean curvature flow*, J. Differential Geom. **31** (1990), no. 1, 285–299.
- [IPR21] D. Impera, S. Pigola, and M. Rimoldi, *The Frankel property for self-shrinkers from the viewpoint of elliptic PDEs*, J. Reine Angew. Math. **773** (2021), 1–20.
- [Keta] D. Ketover, *Free boundary minimal surfaces of unbounded genus*, preprint, available at [arXiv:1612.08691](https://arxiv.org/abs/1612.08691).
- [Keta] ———, *Self-shrinkers whose asymptotic cones fatten*, preprint, available at [arXiv:2407.01240](https://arxiv.org/abs/2407.01240).
- [KKMS] M. Karpukhin, R. Kusner, P. McGrath, and D. Stern, *Embedded minimal surfaces in S^3 and B^3 via equivariant eigenvalue optimization*, preprint, available at [arXiv:2402.13121](https://arxiv.org/abs/2402.13121).
- [KL21] N. Kapouleas and M. M.-c. Li, *Free boundary minimal surfaces in the unit three-ball via desingularization of the critical catenoid and the equatorial disc*, J. Reine Angew. Math. **776** (2021), 201–254.
- [Law70] H. B. Lawson, *The unknottedness of minimal embeddings*, Inventiones Math. **11** (1970), 183–87.

- [Mee81] W. Meeks, *The topological uniqueness of minimal surfaces in three dimensional Euclidean space*, *Topology* **20** (1981), 389–410.
- [Mra24] A. Mramor, *An unknottedness result for noncompact self-shrinkers*, *J. Reine Angew. Math.* **810** (2024), 189–215.
- [Mra] ———, *An unknottedness result for self shrinkers with multiple ends*, preprint, available at [2011.09373](#).
- [Mun56] J. R. Munkres, *Some applications of triangulation theorems*, ProQuest LLC, Ann Arbor, MI, 1956. Thesis (Ph.D.)—University of Michigan.
- [MW20] A. Mramor and S. Wang, *On the topological rigidity of compact self-shrinkers in \mathbb{R}^3* , *Int. Math. Res. Not. IMRN* **7** (2020), 1933–1941.
- [MY92] W. H. Meeks III and S.-T. Yau, *The topological uniqueness of complete minimal surfaces of finite topological type*, *Topology* **31** (1992), no. 2, 305–316.
- [MZ24] P. McGrath and J. Zou, *On the areas of genus zero free boundary minimal surfaces embedded in the unit 3-ball*, *J. Geom. Anal.* **34** (2024), no. 9, Paper No. 274, 14.
- [NZ24] K. Naff and J. J. Zhu, *Half-space intersection properties for minimal hypersurfaces*, preprint (2024), available at [arXiv:2401.09669](#).
- [Sch24] M. B. Schulz, *Geometric analysis gallery*, 2024. URL: [mbschulz.github.io](#). Last visited on 2024/12/15.
- [Sma59] S. Smale, *Diffeomorphisms of the 2-sphere*, *Proc. Amer. Math. Soc.* **10** (1959), 621–626.
- [Wan16] L. Wang, *Asymptotic structure of self-shrinkers*, preprint (2016), available at [arXiv:1610.04904](#).

SABINE CHU

MIT, Department of Mathematics, 77 Massachusetts Avenue, Cambridge, MA 02139, USA

E-mail address: srchu@mit.edu

GIADA FRANZ

MIT, Department of Mathematics, 77 Massachusetts Avenue, Cambridge, MA 02139, USA

E-mail address: gfranz@mit.edu