THE SUPPORT OF KOSTANT'S WEIGHT MULTIPLICITY FORMULA IS AN ORDER IDEAL IN THE WEAK BRUHAT ORDER

PORTIA X. ANDERSON, ESTHER BANAIAN, MELANIE J. FERRERI, OWEN C. GOFF, KIMBERLY P. HADAWAY, PAMELA E. HARRIS, KIMBERLY J. HARRY, NICHOLAS MAYERS, SHIYUN WANG, AND ALEXANDER N. WILSON

ABSTRACT. For integral weights λ and μ of a classical simple Lie algebra \mathfrak{g} , Kostant's weight multiplicity formula gives the multiplicity of the weight μ in the irreducible representation with highest weight λ , which we denote by $m(\lambda, \mu)$. Kostant's weight multiplicity formula is an alternating sum over the Weyl group of the Lie algebra whose terms are determined via a vector partition function. The Weyl alternation set $\mathcal{A}(\lambda, \mu)$ is the set of Weyl group elements that contribute nontrivially to the multiplicity $m(\lambda, \mu)$. In this article, we prove that Weyl alternation sets are order ideals in the weak Bruhat order of the corresponding Weyl group. Specializing to the Lie algebra $\mathfrak{sl}_{r+1}(\mathbb{C})$, we give a complete characterization of the Weyl alternation sets $\mathcal{A}(\tilde{\alpha}, \mu)$, where $\tilde{\alpha}$ is the highest root and μ is a negative root, answering a question of Harry posed in 2024. We also provide some enumerative results that pave the way for our future work where we aim to prove Harry's conjecture that the q-analog of Kostant's weight multiplicity formula $m_q(\tilde{\alpha}, \mu) = q^{r+j-i+1} + q^{r+j-i} - q^{j-i+1}$ when $\mu = -(\alpha_i + \alpha_{i+1} + \cdots + \alpha_j)$ is a negative root of $\mathfrak{sl}_{r+1}(\mathbb{C})$.

1. INTRODUCTION

In this article, we are concerned with the computation of weight multiplicities using Kostant's weight multiplicity formula. For a classical simple Lie algebra \mathfrak{g} of rank $r \geq 1$, let $\Delta = \{\alpha_1, \alpha_2, \ldots, \alpha_r\}$ denote its set of simple roots, Φ denote its set of roots, Φ^+ denote its set of positive roots, Φ^- denote its set of negative roots, $\tilde{\alpha}$ denote its highest root, and let $\rho = \frac{1}{2} \sum_{\alpha \in \Phi^+} \alpha$. Moreover, let W denote the Weyl group associated to \mathfrak{g} , which is generated by reflections denoted as s_1, s_2, \ldots, s_r , through hyperplanes orthogonal to the simple roots $\alpha_1, \alpha_2, \ldots, \alpha_r$, respectively. If $\sigma \in W$, then $\ell(\sigma)$ denotes the length of σ . Now, letting $L(\lambda)$ denote the irreducible highest weight representation of \mathfrak{g} with highest weight λ , Kostant's weight multiplicity formula gives a way to compute the multiplicity of a weight μ in $L(\lambda)$, denoted $m(\lambda, \mu)$. Using an alternating sum over the Weyl group that involves the Kostant partition function, as given in [17, Theorem on p. 589], one has

(1)
$$m(\lambda,\mu) = \sum_{\sigma \in W} (-1)^{\ell(\sigma)} \wp(\sigma(\lambda+\rho) - \mu - \rho)$$

where \wp denotes Kostant's partition function. That is, \wp is the function from the set of weights to $\mathbb{N} = \{0, 1, 2, 3, \ldots\}$ for which $\wp(\xi)$ is the number of ways the weight ξ can be expressed as a linear combination of the positive roots of \mathfrak{g} with non-negative integer coefficients.

Two major challenges arise in computing weight multiplicities via Kostant's formula:

- 1) The number of terms grows factorially in the rank of the Lie algebra as the order of the Weyl group of a classical Lie algebra is either r! or exponential times r! and
- 2) There are no general closed formulas for the value of the Kostant partition function.

However, in practice, many of the terms are zero and hence contribute trivially to the multiplicity [3]. Hence, this motivates work to characterize and enumerate the Weyl group elements that contribute nontrivially to the multiplicity. This set is referred to as the Weyl alternation set and is defined formally below.

Definition 1.1 (Definition 1.1 in [5]). For λ and μ integral weights of \mathfrak{g} , the Weyl alternation set is

(2)
$$\mathcal{A}(\lambda,\mu) = \{ \sigma \in W \mid \wp(\sigma(\lambda+\rho) - \mu - \rho) > 0 \}.$$

Note, $\sigma \in W$ is in $\mathcal{A}(\lambda, \mu)$ if and only if $\sigma(\lambda + \rho) - \mu - \rho$ can be written as a sum of positive roots. The Weyl alternation sets allows one to reduce the computation of $m(\lambda, \mu)$ to be done over $\mathcal{A}(\lambda, \mu)$ rather than over W, which greatly reduces the computation involved. Work determining Weyl alternation sets for a variety of pairs of weights appears in the literature and includes characterizing Weyl alternation sets in both classical and exceptional Lie algebras [2, 12, 4, 6, 7, 8, 9, 10, 11, 12, 13]. Of note is the work of Harris, which establishes that if $\tilde{\alpha}$ is the highest root of $\mathfrak{sl}_{r+1}(\mathbb{C})$, then the Weyl alternation set $\mathcal{A}(\tilde{\alpha}, 0)$ consists of products of certain commuting simple reflections and hence $|\mathcal{A}(\tilde{\alpha}, 0)| = F_r$, the *r*th Fibonacci number [5, Theorem 1.2]. Harry establishes that if $\tilde{\alpha}$ is the highest root and $\mu = \alpha_i + \alpha_{i+1} + \cdots + \alpha_j$ is a height j - i + 1 positive root of $\mathfrak{sl}_{r+1}(\mathbb{C})$, then the Weyl alternation set $\mathcal{A}(\tilde{\alpha}, \mu)$ again consists of certain products of commuting simple reflections and, hence, $|\mathcal{A}(\tilde{\alpha}, \mu)| = F_i \cdot F_{r-j+1}$ [14, Corollary 2.5].

Motivated by the work of Harris and Harry, in this article we prove general structural results about the Weyl alternation set. For example, we establish the following theorem elucidating the poset structure of $\mathcal{A}(\lambda,\mu)$.

Theorem 3.1. Let λ be an integral dominant weight of a simple Lie algebra \mathfrak{g} with Weyl group W. Then for any weight μ , the Weyl alternation set $\mathcal{A}(\lambda,\mu)$ is a (possibly empty) order ideal in the left and right weak Bruhat orders of W.

Next, we generalize the results of Harris and Harry by replacing simple transpositions with elements having a property called *connected influence* and replacing the commuting condition with a condition called *pairwise independence* (terms defined in Section 3), leading to the following result.

Theorem 3.7. For a dominant integral weight λ of a simple Lie algebra \mathfrak{g} and a weight μ such that $\mathcal{A}(\lambda,\mu)$ is nonempty, there exists a unique subset $S \subseteq \mathcal{A}(\lambda,\mu)$ with $1 \notin S$ such that each $b \in S$ has connected influence and any element $\sigma \in \mathcal{A}(\lambda,\mu)$ can be written as a product of a pairwise independent subset of S. Furthermore, there is a bijection between elements of $\mathcal{A}(\lambda,\mu)$ and pairwise independent subsets of S where each pairwise independent subset corresponds to its product.

Note, in Theorem 3.1, the products could be over possibly empty subsets of S. Using these results, we specialize to the Lie algebra of type A_r where the positive roots are given by $\alpha_{i,j} = \alpha_i + \alpha_{i+1} + \cdots + \alpha_j$ for $1 \leq i \leq j \leq r$. Specifically, in Section 4, we characterize the Weyl alternation sets $\mathcal{A}_r(\tilde{\alpha}, \mu)$, where we use the subscript r to denote the rank, $\tilde{\alpha}$ to denote the highest root, and μ to denote a negative root of $\mathfrak{sl}_{r+1}(\mathbb{C})$. In addition, in Section 5, we prove that the cardinalities of these Weyl alternation sets satisfy the two-term recurrence relation defining the Fibonacci numbers. We remark that the initial values of the recurrence relation depend on μ , and we present some examples in Section 5.

Proposition 5.5. Let $r \ge 2$ and $1 \le i \le j \le r-2$. If $\mu = -\alpha_{i,j}$, then $|\mathcal{A}_r(\tilde{\alpha}, \mu)| = |\mathcal{A}_{r-1}(\tilde{\alpha}, \mu)| + |\mathcal{A}_{r-2}(\tilde{\alpha}, \mu)|.$

With this recurrence in hand, we provide the following generating function for the sizes of Weyl alternation sets for all type A negative roots.

Theorem 5.14. If $\tilde{\alpha}$ is the highest root and $\mu = -\alpha_{i,j}$ with $1 \leq i \leq j \leq r$ is a negative root of the Lie algebra of type A_r , then we have the generating function

$$\sum_{1 \le i \le j \le r} |\mathcal{A}_r(\tilde{\alpha}, -\alpha_{i,j})| x^r s^i t^j = \frac{1}{t(1 - x - x^2)} \left((1 - x)t\mathcal{H}(xt, s) + \mathcal{P}(xt, s) - \frac{xst}{1 - xst - (xst)^2} \right)$$

where

$$\mathcal{H}(x,s) = \frac{xs(x^5s + 3x^4s - xs + x^2 + 2x + 1)}{(1 - x - x^2 - 3x^3 - x^4)(1 - xs - (xs)^2)}$$

and

$$\mathcal{P}(x,s) = \frac{xs(x^4s + 3x^3s + x + 1)}{(1 - x - x^2 - 3x^3 - x^4)(1 - xs - (xs)^2)}.$$

The remainder of the article is organized as follows. In Section 2, we provide the necessary background to make our approach precise. In Section 3, we prove Theorem 3.1 establishing that the Weyl alternation sets are order ideals in the weak Bruhat order. Then, we specialize our study to the Lie algebra of type A, and in Section 4, we characterize all of the Weyl alternation sets $\mathcal{A}(\tilde{\alpha},\mu)$ where $\tilde{\alpha}$ is the highest root and μ is a negative root. Then, in Section 5, we give recursive formulas and generating functions for the cardinalities of the Weyl alternation sets considered in the previous section. We conclude with Section 6 in which we detail our future work where we aim to establish a conjecture of Harry for the q-multiplicity of a negative root in the adjoint representation of $\mathfrak{sl}_{r+1}(\mathbb{C})$.

2. Background

For a simple Lie algebra \mathfrak{g} of rank r, there is an associated set $\Phi \subseteq \mathbb{R}^d$ of vectors called a **root system**. Within Φ , there is a distinguished subset $\Delta = \{\alpha_1, \alpha_2, \ldots, \alpha_r\} \subseteq \Phi$ of **simple roots**, which acts as an integral basis for Φ . The **positive roots** $\Phi^+ \subseteq \Phi$ are those that can be written as a linear combination of the simple roots with non-negative integer coefficients. The data of this root system can be summarized in a graph called a **Coxeter–Dynkin diagram** (sometimes referred to as a Coxeter diagram). This graph has vertices labeled by $1, 2, \ldots, r$, and an edge between i and j if and only if α_i and α_j are not orthogonal. The edge between i and j is usually labeled by a positive integer $m_{i,j}$ for which the angle between α_i and α_j is given by $\frac{\pi}{m_{i,j}}$, but we suppress such labels in this paper.

The Weyl group W associated to \mathfrak{g} is generated by simple reflections s_1, s_2, \ldots, s_r where s_i reflects vectors through the hyperplane orthogonal to α_i . That is,

$$s_i(v) = v - 2\frac{(\alpha_i, v)}{(\alpha_i, \alpha_i)}\alpha_i,$$

where (\cdot, \cdot) is the standard inner product on \mathbb{R}^d . The inner product (\cdot, \cdot) is invariant under the action of the Weyl group and that two reflections s_i and s_j commute if and only if α_i and α_j are orthogonal—that is, if α_i and α_j are non-adjacent in the Coxeter–Dynkin diagram. When viewing the Weyl group as an abstract Coxeter group, this action on the roots coincides with what is called the **geometric representation** of the group.

To the Weyl group W, one can associate two posets of interest here, the "right" and "left weak (Bruhat) orders". Recall that a **poset** (\mathcal{P}, \preceq) consists of a set \mathcal{P} along with a binary relation \preceq between the elements of \mathcal{P} that is reflexive, anti-symmetric, and transitive. For $x, y \in \mathcal{P}$, if $x \preceq y$ and $x \neq y$, then we write $x \prec y$. In the case $x \prec y$ and there exists no $z \in \mathcal{P}$ satisfying $x \prec z \prec y$, then $x \prec y$ is a **covering relation**, denoted $x \prec y$, and we say that y covers x. Covering relations are used to define a visual representation of \mathcal{P} called the **Hasse diagram** – a graph whose vertices correspond to elements of \mathcal{P} and whose edges correspond to covering relations. For further details concerning posets we recommend [19].

Now, writing elements of a Weyl group $\sigma \in W$ as **reduced words**, i.e., minimal length expressions for σ as a product of generators s_i , and denoting the length of such expressions by $\ell(\sigma)$, the posets on W of interest here are defined as follows.

Definition 2.1. The **right weak (Bruhat) order** (W, \leq_R) on the Weyl group W is the transitive closure of the covering relations

$$\sigma \lessdot_R \sigma s_i$$
 whenever $\ell(\sigma s_i) > \ell(\sigma)$.

The left weak (Bruhat) order (W, \leq_L) is similarly defined by covering relations

$$\sigma \lessdot_L s_i \sigma$$
 whenever $\ell(s_i \sigma) > \ell(\sigma)$.

The right (resp. left) weak order can be equivalently given as $\sigma <_R \tau$ (resp. $\sigma <_L \tau$) if in some reduced expression of τ a reduced expression of σ appears as a prefix (resp. suffix).

Example 2.2. Figure 1 illustrates the Hasse diagram of the right weak order on the type A_3 Weyl group.

Definition 2.3. Following [19, p. 282], we define an order ideal to be a subposet I of a poset \mathcal{P} with the property that if $y \in I$ and $x \leq y$, then $x \in I$.

Example 2.4. An example order ideal of the right weak order on the type A_3 Weyl group is provided by $I = \{1, s_1, s_2, s_3, s_1 s_3\}$. This order ideal is generated by the elements s_2 and $s_1 s_3$, the maximal elements in I. In Figure 1, we highlight the order ideal I within the Hasse diagram of the right weak order on the type A_3 Weyl group.

3. Weyl alternation sets are order ideals

In this section, we work with a general simple Lie algebra \mathfrak{g} . We call a weight λ integral if $2\frac{(\lambda,\alpha)}{(\alpha,\alpha)} \in \mathbb{Z}$ for all roots α and **dominant** if $(\lambda, \alpha) \geq 0$ for each positive root α . Our main result establishes that $\mathcal{A}(\lambda, \mu)$ for an integral dominant weight λ is an order ideal.

Theorem 3.1. Let λ be an integral dominant weight of a simple Lie algebra \mathfrak{g} with Weyl group W. Then for any weight μ , the Weyl alternation set $\mathcal{A}(\lambda,\mu)$ is a (possibly empty) order ideal in the left and in the right weak Bruhat orders of W.



FIGURE 1. Hasse diagram of the right weak order on the type A_3 Weyl group with the elements of the order ideal $I = \{1, s_1, s_2, s_3, s_1s_3\}$ boxed and in blue text.

Proof. Suppose that $\mathcal{A}(\lambda,\mu) \neq \emptyset$; otherwise, $\mathcal{A}(\lambda,\mu)$ is vacuously an order ideal. To show that $\mathcal{A}(\lambda,\mu)$ is an order ideal with respect to a partial order \leq , it suffices to establish the following: if $\sigma < \tau$ (that is, τ covers σ in the partial order), then the difference $\sigma(\lambda + \rho) - \tau(\lambda + \rho)$ is a linear combination of positive roots with non-negative integer coefficients. Indeed, this implies that $\sigma(\lambda + \rho) - \rho - \mu$ could be obtained from $\tau(\lambda + \rho) - \rho - \mu$ by adding some number of simple roots with positive coefficients. Consequently, if $\wp(\tau(\lambda + \rho) - \rho - \mu) > 0$, then it would then be the case that $\wp(\sigma(\lambda + \rho) - \rho - \mu) > 0$. For example, a partition of the latter could be obtained by adding these same simple roots to the partition of the former, each as their own part. Hence, $\tau \in \mathcal{A}(\lambda,\mu)$ would imply that $\sigma \in \mathcal{A}(\lambda,\mu)$, i.e., $\mathcal{A}(\lambda,\mu)$ would be an order ideal.

First, we consider the right Bruhat order. Suppose that $\sigma \leq_R \tau$. By definition, $\tau = \sigma s_i$ for some simple reflection $s_i \in W$ and $\ell(\sigma s_i) > \ell(\sigma)$. We now compute

$$\sigma(\lambda+\rho) - \sigma s_i(\lambda+\rho) = \sigma(\lambda+\rho) - \left(\sigma(\lambda+\rho) - 2\frac{(\lambda+\rho,\alpha_i)}{(\alpha_i,\alpha_i)}\sigma(\alpha_i)\right) = 2\frac{(\lambda+\rho,\alpha_i)}{(\alpha_i,\alpha_i)}\sigma(\alpha_i)$$

By [1, Equation 4.25], $\ell(\sigma s_i) > \ell(\sigma)$ implies that $\sigma(\alpha_i) \in \Phi^+$. Because λ and ρ are dominant integral weights, $2\frac{(\lambda+\rho,\alpha_i)}{(\alpha_i,\alpha_i)}$ is a non-negative integer, so the difference $\sigma(\lambda+\rho)-\sigma s_i(\lambda+\rho)$ is indeed a linear combination of simple roots with non-negative integer coefficients

Now, consider the left Bruhat order. Suppose instead that $\sigma \leq_L \tau$. By definition $\tau = s_i \sigma$ and $\ell(s_i \sigma) > \ell(\sigma)$. We now compute

$$\sigma(\lambda+\rho) - s_i \sigma(\lambda+\rho) = \sigma(\lambda+\rho) - \left(\sigma(\lambda+\rho) - 2\frac{(\sigma(\lambda+\rho),\alpha_i)}{(\alpha_i,\alpha_i)}\alpha_i\right)$$
$$= 2\frac{(\sigma(\lambda+\rho),\alpha_i)}{(\alpha_i,\alpha_i)}\alpha_i$$
$$= 2\frac{(\lambda+\rho,\sigma^{-1}(\alpha_i))}{(\sigma^{-1}(\alpha_i),\sigma^{-1}(\alpha_i))}\alpha_i,$$

where the last equality holds because the inner product is invariant under the action of W (for a proof, we point the reader to [1, Equation 4.23]). Moreover, $\ell(s_i\sigma) > \ell(\sigma)$ implies that $\ell(\sigma^{-1}s_i) > \ell(\sigma^{-1})$, and [1, Equation 4.25] tells us that $\sigma^{-1}(\alpha_i)$ is a positive root. Further, λ and ρ are dominant integral weights. Hence, $2\frac{(\lambda+\rho,\sigma^{-1}(\alpha_i))}{(\sigma^{-1}(\alpha_i),\sigma^{-1}(\alpha_i))}$ is a non-negative integer, and the difference $\sigma(\lambda+\rho) - s_i\sigma(\lambda+\rho)$ is indeed a linear combination of simple roots with non-negative integer coefficients.

4



FIGURE 2. The type A_7 Dynkin diagram with some influence and extended influence highlighted.

The following result is essential in our work to characterize Weyl group elements that are not contained in certain Weyl alternation sets.

Corollary 3.2. Let λ be a dominant integral weight of a simple Lie algebra \mathfrak{g} . For any weight μ , if σ is a reduced word appearing contiguously in a reduced expression for $\tau \in \mathcal{A}(\lambda, \mu)$, then $\sigma \in \mathcal{A}(\lambda, \mu)$.

Proof. Suppose $\tau \in \mathcal{A}(\lambda, \mu)$ and σ is a reduced word appearing contiguously in a reduced expression for τ . Then $\tau = \tau_1 \sigma \tau_2$ for some reduced words τ_1 and τ_2 . Note, $\tau_1 \sigma \leq_R \tau$. By Theorem 3.1, we have that $\tau_1 \sigma \in \mathcal{A}(\lambda, \mu)$. As $\sigma \leq_L \tau_1 \sigma$, we similarly have that $\sigma \in \mathcal{A}(\lambda, \mu)$.

Definition 3.3. For an element $\sigma \in W$ of the Weyl group, define the **influence** and **extended influence** of σ , denoted $I(\sigma)$ and $\overline{I}(\sigma)$ respectively, by

 $I(\sigma) = \{i : s_i \text{ is in a reduced word for } \sigma\}, \text{ and}$ $\overline{I}(\sigma) = \{i : i \in I(\sigma) \text{ or } i \text{ is adjacent to some } j \in I(\sigma) \text{ in the Dynkin diagram}\}.$

In type A, the extended influence $\overline{I}(\sigma)$ consists of all i such that $i \in I(\sigma)$ or $i \pm 1 \in I(\sigma)$.

Further, we define the following relations of elements in the Weyl group with respect to their influences and extended influences.

Definition 3.4. We say that $I(\sigma)$ is **connected** if the subgraph of the Dynkin diagram it induces is connected. We say that $\sigma, \tau \in W$ are **independent** if $I(\sigma) \cap \overline{I}(\tau) = \emptyset$. A set $S \subseteq W$ is called **pairwise independent** if any pair of elements $\sigma \neq \tau$ in S are independent.

Example 3.5. Consider the type A_7 Dynkin diagrams in Figure 2 with the influences

$$I(s_1s_2) = I(s_1s_2s_1) = \{1, 2\},$$

$$I(s_4s_6s_7) = \{4, 6, 7\},$$

$$\overline{I}(s_1s_2) = \overline{I}(s_1s_2s_1) = \{1, 2, 3\}, \text{ and}$$

$$\overline{I}(s_4s_6s_7) = \{3, 4, 5, 6, 7\}$$

highlighted. Note, $I(s_1s_2) = I(s_1s_2s_1)$ is connected and $I(s_4s_6s_7)$ is disconnected. The elements s_1s_2 and $s_4s_6s_7$ are independent.

Example 3.6. In Table 1, we consider the influence and extended influence of two elements in type D_5 . Note, $I(s_3s_4) \cap \overline{I}(s_2) = \{3\} \neq \emptyset$, so s_3s_4 and s_2 are not independent.

Lemma 3.7. Let $\sigma, \tau \in W$.

(1) If $\sigma = s_{i_1}s_{i_2}\cdots s_{i_{\ell}}$ is a reduced word, then

$$I(\sigma) = \{i_j : 1 \le j \le \ell\}.$$

- (2) If $I(\sigma) = I(\tau)$ and $\sigma(\alpha_i) = \tau(\alpha_i)$ for all $i \in I(\sigma)$, then $\sigma = \tau$.
- (3) If $\{\sigma_1, \sigma_2, \ldots, \sigma_k\} \subseteq W$ is pairwise independent, then

$$I(\sigma_m) \subseteq I(\sigma_1 \sigma_2 \cdots \sigma_k)$$

for all $1 \leq m \leq k$.



TABLE 1. Examples of influence and extended influence for elements in the Weyl group of type D_5 .

Proof. Property (1): This property follows from [1, Corollary 2.2.3], which states that every reduced expression of σ contains a reduced expression of τ if and only if some reduced expression of σ contains a reduced expression of τ .

Property (2): Suppose $I(\sigma) = I(\tau)$ and let $I = I(\sigma) = I(\tau)$. Then $\sigma, \tau \in W_I$ where W_I is the subgroup of W generated by $\{s_i : i \in I\}$. Since the geometric representation of W_I is isomorphic to the action of W_I on $V_I = \mathbb{R}\{\alpha_i : i \in I\}$ (for a reference we recommend [15, Section 5.5]) and the geometric representation of any finite Coxeter group is faithful, W_I acts faithfully on V_I . Hence, if $\sigma(\alpha_i) = \tau(\alpha_i)$ for all $i \in I$, we have that $\sigma = \tau$.

Property (3): Suppose that $\sigma_1, \sigma_2, \ldots, \sigma_k$ are reduced expressions and the set $\{\sigma_1, \sigma_2, \ldots, \sigma_k\}$ is pairwise independent. If $\sigma_1 \sigma_2 \cdots \sigma_k$ were not a reduced word, we could apply a sequence of nil-moves and braid-moves as in [1, Theorem 3.3.1] to arrive at a reduced expression. Since these operations only apply to strings of generators that are adjacent in the Dynkin diagram, each of these moves must occur within a σ_i . But each σ_i is assumed to be reduced, so $\sigma_1 \sigma_2 \cdots \sigma_k$ is a reduced expression. Hence, if s_i occurs in σ_m for some $m \in [k]$, then it occurs in the product $\sigma_1 \sigma_2 \cdots \sigma_k$ and property (3) follows.

Theorem 3.8. For λ a dominant integral weight of a simple Lie algebra \mathfrak{g} and μ a weight such that $\mathcal{A}(\lambda,\mu)$ is nonempty, there exists a unique subset $S \subseteq \mathcal{A}(\lambda,\mu)$ with $1 \notin S$ such that each $b \in S$ has connected influence and any element $\sigma \in \mathcal{A}(\lambda,\mu)$ can be written as a product of a pairwise independent subset of S. Furthermore, there is a bijection between elements of $\mathcal{A}(\lambda,\mu)$ and pairwise independent subsets of S where each pairwise independent subset corresponds to its product.

Proof. Toward proving uniqueness, suppose that S and S' both satisfy the conditions in the statement of the theorem and let $b \in S$. Since $b \in A(\lambda, \mu)$, we have that $b = b'_1b'_2\cdots b'_k$ for $\{b'_1, b'_2, \ldots, b'_k\}$ a pairwise independent subset of S'. By Lemma 3.7 item (3), we have that $I(b'_m) \subseteq I(b)$ for all $1 \leq m \leq k$. As the influence of all elements of S and S' are assumed to be connected, we must in fact have that $I(b) = I(b'_m)$ for all $1 \leq m \leq k$. The only way that the set $\{b'_1, b'_2, \ldots, b'_k\}$ can be pairwise independent is if it consists of a single element b'_1 . Then $b = b'_1 \in S'$, so $S \subseteq S'$. Similarly, $S' \subseteq S$, and S = S'.

To prove that such a subset always exists, consider the set $S \subseteq \mathcal{A}(\lambda, \mu)$ consisting of all elements b with connected influence such that if $b = b_1b_2$ for b_1 and b_2 independent, then $b_1 = 1$ or $b_2 = 1$. We claim that for any $\sigma \in \mathcal{A}(\lambda, \mu)$, we have that $\sigma = b\sigma'$ where $b \in S$ and the elements b and σ' are independent. Indeed, if σ itself is not already an element of S, then $\sigma = \sigma_1 \sigma_2$ with $\sigma_1 <_R \sigma$ where the elements σ_1 and σ_2 are independent. If $\sigma_1 \notin S$, then we can repeat this process. As the left factor decreases in the right weak order each time we repeat, eventually this process terminates with $\sigma = b\sigma'$ with $b \in S$ and the elements b and σ' independent. Iterating this process yields

$$\sigma = b_1 b_2 \cdots b_k,$$

where the elements b_i and $b_{i+1}b_{i+2}\cdots b_k$ are independent for all $1 \leq i \leq k-1$. By Lemma 3.7, $I(b_j) \subseteq I(b_{i+1}b_{i+2}\cdots b_k)$ for j > i. Consequently,

$$\overline{I}(b_i) \cap I(b_j) \subseteq \overline{I}(b_i) \cap I(b_{i+1}b_{i+2}\cdots b_k) = \emptyset.$$

Hence, we have written σ as a product of a pairwise independent set.

To prove that there is a bijection between pairwise independent subsets of S and elements of $\mathcal{A}(\lambda,\mu)$, we only need to demonstrate that the product of any pairwise independent subset of S is in $\mathcal{A}(\lambda,\mu)$ and that any $\sigma \in \mathcal{A}(\lambda,\mu)$ can be written uniquely as the product of a pairwise independent subset.

Let $B \subseteq S$ be a pairwise independent subset and let $\sigma = \prod_{b \in B} b$. Independent elements of W necessarily commute, so this product is well-defined. For any $i \in [r]$ (where r is the rank of the Lie algebra), the coefficient of α_i in $\sigma(\lambda + \rho) - \rho - \mu$ is either:

- (i) the coefficient of α_i in $b(\lambda + \rho) \rho \mu$ for a unique $b \in B$ with $i \in I(b)$, or
- (ii) the coefficient of α_i in $\lambda \mu$ if no such b exists.

Since in (i), $b \in \mathcal{A}(\lambda, \mu)$ and in (ii), $1 \in \mathcal{A}(\lambda, \mu)$, the coefficient of α_i is non-negative in either case. Hence, $\sigma(\lambda + \rho) - \rho - \mu$ is a non-negative linear combination of simple roots, and $\sigma \in \mathcal{A}(\lambda, \mu)$.

Suppose $\sigma = b_1 b_2 \cdots b_k = b'_1 b'_2 \cdots b'_{k'}$ is written in two ways as a product of pairwise independent subsets of S. Fix $1 \le m \le k$ and consider $i \in I(b_m)$. Necessarily, by Lemma 3.7 item (3), $i \in I(b'_{m'})$ for exactly one $1 \le m' \le k'$, hence k = k'. Since I(b) is connected for all $b \in S$, we must have that $I(b_m) = I(b'_{m'})$. Without loss of generality, we can assume m = m', so $I(b_m) = I(b'_m)$ for all $1 \le m \le k$.

Now for $j \in I(b_m)$, we have that

$$\sigma(\alpha_j) = b_1 b_2 \cdots b_k(\alpha_j) = b_m(\alpha_j), \text{ and} \\ \sigma(\alpha_j) = b'_1 b'_2 \cdots b'_k(\alpha_j) = b'_m(\alpha_j).$$

By Lemma 3.7 item (2), $b_m = b'_m$. Hence, the two products correspond to the same pairwise independent subset of S.

Ongoing, we denote the unique subset of $\mathcal{A}(\lambda,\mu)$ described in Theorem 3.8 by BAS (λ,μ) and refer to its elements as the **basic allowable subwords** for the pair λ and μ .

Remark 3.9. Theorem 3.8 shows that $\mathcal{A}(\lambda, \mu)$ has the structure of an *independence system* or *abstract simplicial complex* because each element of $\mathcal{A}(\lambda, \mu)$ corresponds to a pairwise independent subset of BAS (λ, μ) and the property of being pairwise independent is downward-closed.

The following proof, providing sufficient conditions for a set of words to be the set $BAS(\lambda, \mu)$, generalizes [10, Theorem 4.5].

Proposition 3.10. Let λ be a dominant integral weight and let μ be a weight such that $\mathcal{A}(\lambda, \mu) \neq \emptyset$. Let $S \subseteq \mathcal{A}(\lambda, \mu)$, where $1 \notin S$, be such that

- (C1) any simple transposition $s_i \in \mathcal{A}(\lambda, \mu)$ is contained in S,
- (C2) for each $\sigma \in S$, its influence $I(\sigma)$ is connected, and
- (C3) if σ and τ in S are not independent, the product $\sigma\tau$ falls into one of the following cases: (1) $\sigma\tau \in S$,
 - (2) $\sigma \tau = \nu_1 \nu_2 \cdots \nu_m$ where $\{\nu_1, \nu_2, \dots, \nu_m\} \subseteq S$ is a pairwise independent set and $\ell(\nu_1) + \ell(\nu_2) + \cdots + \ell(\nu_m) < \ell(\sigma) + \ell(\tau)$, or
 - (3) $\sigma\tau$ contains a word not in $\mathcal{A}(\lambda,\mu)$ as a contiguous subword.

Then, $BAS(\lambda, \mu) = S$.

Proof. By Theorem 3.8, it suffices to show that every element $\sigma \in \mathcal{A}(\lambda, \mu)$ can be written as the product of a pairwise independent subset of S.

Let $\sigma \in \mathcal{A}(\lambda, \mu)$. By (C1) and Corollary 3.2, any simple transposition appearing in a reduced expression for σ is contained in S. Thus, we can write σ as a product of elements of S. Choose a product $\sigma = b_1 b_2 \cdots b_k$ with $\ell(b_1) + \ell(b_2) + \cdots + \ell(b_k)$ minimal, and amongst these choose the product with k minimal. Suppose there were two elements b_i and b_j in this product that are not independent. Without loss of generality, we can assume b_i and b_j are adjacent in the product (as an element could be swapped with any independent element adjacent to it). By assumption, there are three possible cases. In case (1), we could replace $b_i b_j$ with a single element from S. Since $\ell(\sigma\tau) \leq \ell(\sigma) + \ell(\tau)$, this replacement must either reduce the length sum, contradicting its minimality or leave the sum of the lengths unchanged, contradicting the assumption that k was minimal among products with that length sum. In case (2), we can replace $b_i b_j$ with a pairwise independent product decreasing the overall length sum, contradicting minimality of the length sum. In case (3), $b_i b_j$ contains a subword not in $\mathcal{A}(\lambda, \mu)$. By Corollary 3.2, this contradicts $\sigma \in \mathcal{A}(\lambda, \mu)$. Hence, any pair b_i, b_j in the product is independent, so $\{b_1, b_2, \ldots, b_k\}$ is a pairwise independent subset of S.

Let $\nu \leq \mu$ denote that $\mu - \nu$ can be written as a linear combination of positive roots with non-negative real coefficients. Next, we relate the sets $BAS(\lambda, \mu)$ and $BAS(\lambda, \nu)$ for a dominant integral weight λ and integral weights μ and ν satisfying $\nu \leq \mu$. To do so, we require the following result.

Lemma 3.11. Let λ, μ, ν be integral weights. If $\nu \leq \mu$, then $\mathcal{A}(\lambda, \mu) \subseteq \mathcal{A}(\lambda, \nu)$.

Proof. First, we claim that if α and β are integral weights and $\alpha \leq \beta$, then $\wp(\alpha) > 0$ implies that $\wp(\beta) > 0$. Indeed, in this case, $\beta = \alpha + \sum_{i=1}^{n} c_i \alpha_i$ where $c_i \in \mathbb{Z}_{\geq 0}$ for each *i*. Consequently, a vector partition of α can be completed to a vector partition of β by adding c_i parts equal to α_i for each *i*.

Now, suppose $\sigma \in \mathcal{A}(\lambda, \mu)$. Then $\wp(\sigma(\lambda + \rho) - \mu - \rho) > 0$. Because $\nu \leq \mu$, we have that $\sigma(\lambda + \rho) - \mu - \rho \leq \sigma(\lambda + \rho) - \nu - \rho$. Hence, $\wp(\sigma(\lambda + \rho) - \nu - \rho) > 0$ and $\sigma \in \mathcal{A}(\lambda, \nu)$, completing the proof.

Proposition 3.12. Let λ be an integral dominant weight, and let μ and ν be two other integral weights such that $\nu \leq \mu$. Then, $BAS(\lambda, \mu) = BAS(\lambda, \nu) \cap \mathcal{A}(\lambda, \mu)$.

Proof. Let $S = BAS(\lambda, \nu) \cap \mathcal{A}(\lambda, \mu)$. We want to show $S = BAS(\lambda, \mu)$ by showing S satisfies the three properties in Proposition 3.10.

- (C1) If $s_i \in \mathcal{A}(\lambda, \mu)$, then $s_i \in \mathcal{A}(\lambda, \nu)$ by Lemma 3.11. By Proposition 3.10, $s_i \in BAS(\lambda, \nu)$. Therefore, $s_i \in S$.
- (C2) Since $S \subseteq BAS(\lambda, \nu)$, we automatically know each $\sigma \in S$ has connected influence.
- (C3) Let σ, τ be a pair of non-independent elements of S. Since $\sigma, \tau \in BAS(\lambda, \nu)$, Proposition 3.10 gives three possibilities for the product $\sigma\tau$.
 - (1) If $\sigma \tau \in BAS(\lambda, \nu)$, then either $\sigma \tau \in S$ or $\sigma \tau \notin \mathcal{A}(\lambda, \mu)$. If we are in the latter case, then $\sigma \tau$ necessarily contains a contiguous subword that is not in $\mathcal{A}(\lambda, \mu)$. Thus, we are either in item (1) or item (3) from Condition (C3) of Proposition 3.10 with respect to S.
 - (2) If $\sigma\tau = \beta_1\beta_2\cdots\beta_m$ where $\{\beta_1,\beta_2,\ldots,\beta_m\} \subseteq BAS(\lambda,\nu)$ is a pairwise independent set and $\ell(\beta_1) + \ell(\beta_2) + \cdots + \ell(\beta_m) < \ell(\sigma) + \ell(\tau)$, then either each $\beta_i \in S$ or there exists $\beta_j \notin \mathcal{A}(\lambda,\mu)$. In the latter case, β_j contains a contiguous subword not in $\mathcal{A}(\lambda,\mu)$, hence $\sigma\tau$ does as well. Thus, we are either in item (2) or (3) from Condition (C3) of Proposition 3.10 with respect to S.
 - (3) If $\sigma\tau$ contains a contiguous subword which is not in $\mathcal{A}(\lambda,\nu)$, then $\sigma\tau$ also contains a contiguous subword which is not in $\mathcal{A}(\lambda,\mu)$ by Lemma 3.11. Therefore, we are in item (3) from Condition (C3) of Proposition 3.10 with respect to S.

Therefore, the product of any pair of non-independent elements $\sigma, \tau \in S$ fits into one of the three items of Condition (C3) of Proposition 3.10 with respect to S.

4. The Lie algebra of type A

In this section, we focus exclusively on the Lie algebra of type A. Let $M_{r+1}(\mathbb{C})$ denote the collection of $(r+1) \times (r+1)$ complex-valued matrices and recall that $\mathfrak{sl}_{r+1}(\mathbb{C}) = \{X \in M_{r+1}(\mathbb{C}) : \operatorname{Tr}(X) = 0\}$ is the Lie algebra of type A_r , with $r \ge 1$. Recall that $\mathfrak{sl}_{r+1}(\mathbb{C})$ is a complex vector space with a bilinear product, called the Lie bracket, defined by the commutator bracket [X,Y] = XY - YX for $X, Y \in \mathfrak{sl}_{r+1}(\mathbb{C})$, which is skew-symmetric and satisfies the Jacobi identity. Given $e_1, e_2, \ldots, e_{r+1}$, the standard basis vectors of \mathbb{R}^{r+1} , for each $i \in [r]$, we define $\alpha_i = e_i - e_{i+1}$. In $\mathfrak{sl}_{r+1}(\mathbb{C})$, the simple roots are $\Delta = \{\alpha_1, \alpha_2, \ldots, \alpha_r\}$, the positive roots are $\Phi^+ = \Delta \cup \{\alpha_i + \alpha_{i+1} + \cdots + \alpha_j : 1 \le i < j \le r\}$, and the negative roots are $-\Phi^+$. The highest root is denoted $\tilde{\alpha} = \alpha_1 + \alpha_2 + \cdots + \alpha_r$ and the negative highest root is henceforth denoted by $-\tilde{\alpha} = -(\alpha_1 + \alpha_2 + \cdots + \alpha_r)$. We recall that the Weyl group W of type A_r is isomorphic to the symmetric group \mathfrak{S}_{r+1} . Our main results characterize and enumerate the elements of the Weyl alternation sets $\mathcal{A}_r(\tilde{\alpha}, \mu)$ when μ is a negative root of $\mathfrak{sl}_{r+1}(\mathbb{C})$. We introduce a subscript r on the Weyl alternation set to specify the rank of the Lie algebra.

The following computations form the backbone of our proofs in this section.

Lemma 4.1. For $\mathfrak{g} = \mathfrak{sl}_{r+1}(\mathbb{C})$, the type A_r simple Lie algebra,

- (1) $s_1(\tilde{\alpha}) = \tilde{\alpha} \alpha_1$ and $s_r(\tilde{\alpha}) = \tilde{\alpha} \alpha_r$,
- (2) for $2 \leq j \leq r-1$, $s_j(\tilde{\alpha}) = \tilde{\alpha}$, and
- (3) for all $1 \le j \le r$, $s_j(\rho) = \rho \alpha_j$.

Proof. The computations are straightforward with (1) and (2) appearing in [10, Lemma 3.1] and (3) in [10, Lemma 3.2]. \Box

Now, we iteratively apply Lemma 4.1 to the elements of W, the Weyl group of the Lie algebra of type A_r .

Lemma 4.2. The action of Weyl group elements on $\tilde{\alpha}$ and ρ satisfy the following.

- (1) If $i_1, i_2, \ldots, i_k \in \{2, 3, \ldots, r-1\}$, then $s_{i_k}s_{i_{k-1}}\cdots s_{i_1}(\tilde{\alpha}) = \tilde{\alpha}$. Otherwise, if i_j is the first occurrence of 1 or r in the list i_1, i_2, \ldots, i_k , then $s_{i_k}s_{i_{k-1}}\cdots s_{i_1}(\tilde{\alpha}) = s_{i_k}s_{i_{k-1}}\cdots s_{i_j}(\tilde{\alpha})$.
- (2) For j < r, $s_j s_{j-1} \cdots s_1(\tilde{\alpha}) = \tilde{\alpha} \sum_{i=1}^j \alpha_i$, and for j > 1, $s_j s_{j+1} \cdots s_r(\tilde{\alpha}) = \tilde{\alpha} \sum_{i=j}^r \alpha_i$.
- (3) For $1 \le i \le r$ and $i + k \le r$, we have $s_i s_{i+1} \cdots s_{i+k}(\rho) = \rho \sum_{j=i}^{i+k} (i+k+1-j)\alpha_j$ and similarly $s_{i+k} \cdots s_{i+1} s_i(\rho) = \rho \sum_{j=i}^{i+k} (j-i+1)\alpha_j$.
- (4) For $1 \le k \le r-1$, $s_k s_{k+1}(\rho) = \rho 2\alpha_k \alpha_{k+1}$ and $s_{k+1} s_k(\rho) = \rho \alpha_k 2\alpha_{k+1}$.
- (5) For $1 \le k \le r 1$, $s_k s_{k+1} s_k(\rho) = \rho 2\alpha_k 2\alpha_{k+1}$.
- (6) For $1 \le k \le r-2$, $s_{k+2}s_ks_{k+1}(\rho) = \rho 2\alpha_k \alpha_{k+1} 2\alpha_{k+2}$ and $s_{k+1}s_{k+2}s_k(\rho) = \rho \alpha_k 3\alpha_{k+1} \alpha_{k+2}$.

Proof. Item (1) follows directly from Lemma 4.1.

We show the first statement of (2) by induction on j. The case when j = 1 follows immediately from Lemma 4.1. Suppose the claim holds for j - 1. Applying our inductive hypothesis, we find that

$$s_j s_{j-1} \cdots s_1 \left(\tilde{\alpha} \right) = s_j \left(\tilde{\alpha} - \sum_{i=1}^{j-1} \alpha_i \right) = s_j \left(\tilde{\alpha} - \alpha_{j-1} - \sum_{i=1}^{j-2} \alpha_i \right) = \tilde{\alpha} - \left(\alpha_{j-1} + \alpha_j \right) - \sum_{i=1}^{j-2} \alpha_i$$

The second statement follows analogously.

Item (3) follows from an inductive argument similar to that used in item (2), and item (4) is a special case of (3). Items (5) and (6) are straightforward calculations using Lemma 4.1. \Box

4.1. Allowable subwords. In this section, we introduce a set of elements of the type A_r Weyl group and verify that each is in $\mathcal{A}(\tilde{\alpha}, -\tilde{\alpha})$. We show in Section 4.3 that these elements are precisely BAS $(\tilde{\alpha}, -\tilde{\alpha})$.

Proposition 4.3. If σ is one of the Weyl group elements

(a) s_k with $1 \le k \le r$, (b) $s_{k+1}s_k$ with $2 \le k \le r-2$, (c) s_ks_{k+1} with $2 \le k \le r-2$, (d) $s_ks_{k+1}s_k$ with $2 \le k \le r-2$, or (e) $s_{k+2}s_ks_{k+1}$ with $2 \le k \le r-3$, then $\sigma \in \mathcal{A}(\tilde{\alpha}, -\tilde{\alpha})$.

Proof. It suffices to check that for any $w \in W$ from the list, if $w(\tilde{\alpha}+\rho) = \tilde{\alpha}+\rho-\sum_{i=1}^{r}c_{i}\alpha_{i}$, then $c_{i} \leq 2$ for all *i*. First, if *w* does not have a subword s_{1} or s_{r} , then $w(\tilde{\alpha}) = \tilde{\alpha}$ by Lemma 4.2 (1). Consequently, when *w* does not have a subword s_{1} or s_{r} , we need only focus on $w(\rho)$, and the desired implication follows by Lemma 4.2 parts (3)-(6). The only cases where we consider s_{1} or s_{r} is when *w* is the simple reflection s_{1} or s_{r} . In these cases, the proof uses the simpler computations in Lemma 4.1. Thus, in all cases, $w(\tilde{\alpha}+\rho)+\tilde{\alpha}-\rho=2\tilde{\alpha}-\sum_{i=1}^{r}c_{i}\alpha_{i}$ with all $c_{i} \leq 2$. As noted above, the result follows.

4.2. Forbidden subwords. In this section, we collect some subwords that cannot appear contiguously in a reduced expression for an element of $\mathcal{A}(\tilde{\alpha}, -\tilde{\alpha})$.

Lemma 4.4. The words

 $s_2 s_1, \quad s_1 s_2, \quad s_{r-1} s_r, \quad and \quad s_r s_{r-1},$

and for any $2 \leq i \leq r - 1$, the words

 $s_{i-1}s_is_{i+1}, \quad s_is_{i-1}s_{i+1}, \quad and \quad s_{i+1}s_is_{i-1}$

are not contained in $\mathcal{A}(\tilde{\alpha}, -\tilde{\alpha})$.

Proof. To show that the listed words are not in $\mathcal{A}(\tilde{\alpha}, -\tilde{\alpha})$, it suffices to check that $w(\tilde{\alpha} + \rho) + \tilde{\alpha} - \rho$ has a negative coefficient when expanded as a linear combination of simple roots, meaning that $\wp(w(\tilde{\alpha} + \rho) + \tilde{\alpha} - \rho)$ is necessarily zero. We denote the leftmost reflection in each word w by s_j . Using Lemma 4.1 and Lemma 4.2, it is straightforward to compute that for any word w listed,

(3)
$$w(\tilde{\alpha}+\rho) + \tilde{\alpha}-\rho = 2\tilde{\alpha} - 3\alpha_j - \sum_{i=1}^{j-1} c_i\alpha_i - \sum_{i=j+1}^r c_i\alpha_i$$

with $c_i \leq 2$ for all *i*. Observe that in Equation (3) the coefficient of α_j is -1. As noted above, the result follows.

Lemma 4.5. Let $k \in [r-3]$. The product of the four simple reflections s_k , s_{k+1} , s_{k+2} , and s_{k+3} in any order is not contained in $\mathcal{A}(\tilde{\alpha}, -\tilde{\alpha})$.

Proof. Let σ be a product of s_k , s_{k+1} , s_{k+2} , and s_{k+3} in some order and suppose that $\sigma \in \mathcal{A}(\tilde{\alpha}, -\tilde{\alpha})$. Because s_{k+3} commutes with s_k and s_{k+1} , we can move s_{k+3} to the beginning or the end of the reduced expression for σ . That is, $\sigma = s_{k+3}\tau$ or $\sigma = \tau s_{k+3}$ where τ is a product of s_k , s_{k+1} , and s_{k+2} in some order. By Lemma 4.4, we must have that $\tau = s_k s_{k+2} s_{k+1}$ for $2 \leq k \leq r-3$. Then either

$$\begin{aligned} \sigma &= s_{k+3}\tau \\ &= s_{k+3}s_ks_{k+2}s_{k+1} \\ &= s_k(s_{k+3}s_{k+2}s_{k+1}) \end{aligned}$$

containing a forbidden $s_{k+3}s_{k+2}s_{k+1}$ or

$$= \tau s_{k+3} = s_k s_{k+2} s_{k+1} s_{k+3} = s_k (s_{k+2} s_{k+1} s_{k+3}),$$

containing a forbidden $s_{k+2}s_{k+1}s_{k+3}$. This contradicts the fact that $\sigma \in \mathcal{A}(\tilde{\alpha}, -\tilde{\alpha})$.

 σ

4.3. Weyl alternation set $\mathcal{A}(\tilde{\alpha}, \mu)$ for μ the negative highest root. In this section, we apply Proposition 3.10 to prove that the set of words given in Section 4.1 are precisely the set $BAS(\tilde{\alpha}, -\tilde{\alpha})$. The following result is analogous to [10, Lemma 4.3], and we modify it to fit our purposes.

Lemma 4.6. Let S be the set of elements listed in Proposition 4.3. Let $\sigma, \tau \in S$ be non-independent elements. Then, the product $\sigma\tau$ falls into one of the following three cases:

- (1) $\sigma \tau \in S$,
- (2) $\sigma \tau = \nu_1 \nu_2 \cdots \nu_m$ where $\{\nu_1, \nu_2, \dots, \nu_m\}$ is a (possibly empty) pairwise independent subset of S and $\ell(\nu_1) + \cdots + \ell(\nu_m) < \ell(\sigma) + \ell(\tau)$, or
- (3) $\sigma\tau$ contains a forbidden subword listed in Lemma 4.4 or Lemma 4.5.

Proof. We proceed via cases. Let σ and τ come from one of the following (not necessarily distinct) collections:

(a) s_k with $1 \le k \le r$,

- (b) $s_{k+1}s_k$ with $2 \le k \le r-2$,
- (c) $s_k s_{k+1}$ with $2 \le k \le r-2$,
- (d) $s_k s_{k+1} s_k$ with $2 \le k \le r-2$, or
- (e) $s_{k+2}s_ks_{k+1}$ with $2 \le k \le r-3$.

For each pair of forms that $\sigma, \tau \in S$ can respectively take, we fix the indices of σ , then consider the range of indices of τ for which σ and τ are not independent.

To reduce the number of cases that need to be handled, consider the automorphism of the Weyl group $f: W \to W$ given by the conjugation $f(\sigma) = w_0 \sigma w_0$ by the longest element w_0 of W. This map f sends s_i to s_{r+1-i} and preserves length. Note additionally that f(S) = S and words of the form (a), (d), and (e) are sent to words of the same form. Words of the form (b) are taken to words of the form (c) by f and vice versa. If σ is a forbidden word as in Lemma 4.4, then so is $f(\sigma)$. As the automorphism f preserves words of the form (a), (d), and (e) the form (a), (d), and (e), while replacing a word of the form (c) with a word of the form (b), we can reduce the number of cases we consider.

Products of the form (a)(a): Let $\sigma = s_k$ for some $1 \le k \le r$. For $\tau = s_j$ with $1 \le j \le r$ to not be independent with σ , we must have $j \in \{k-1, k, k+1\}$.

- Let j = k 1 for $2 \le k \le r$. Then $\sigma\tau = s_k s_{k-1}$. If k = 2 (resp., k = r), then $s_2 s_1$ (resp., $s_r s_{r-1}$) is forbidden by Lemma 4.4. That is, $\sigma\tau$ falls into case (3). Otherwise, $3 \le k \le r 1$ and $s_k s_{k-1} \in S$. That is, $\sigma\tau$ falls into case (1).
- Let j = k for $1 \le k \le r$. Then $\sigma\tau = s_k s_k = 1$, a product of an empty subset of S, with $0 = \ell(1) = \ell(s_k s_k) < \ell(s_k) + \ell(s_k) = 2$. That is, $\sigma\tau$ falls into case (2).
- Let j = k + 1 for $1 \le k \le r 1$. Then $\sigma\tau = s_k s_{k+1}$. If k = 1 (resp., k = r 1), then $s_1 s_2$ (resp., $s_{r-1}s_r$) is forbidden by Lemma 4.4. That is, $\sigma\tau$ falls into case (3). Otherwise, $2 \le k \le r 2$ and $s_k s_{k+1} \in S$. That is, $\sigma\tau$ falls into case (1).

Products of the form (b)(b) (equivalent to (c)(c) via f): Let $\sigma = s_{k+1}s_k$ for some $2 \le k \le r-2$. For $\tau = s_{j+1}s_j$ with $2 \le j \le r-2$ to not be independent with σ , we must have $j \in \{k-2, k-1, k, k+1, k+2\}$.

- Let j = k 2 for $4 \le k \le r 2$. Then $\sigma \tau = (s_{k+1}s_k)(s_{k-1}s_{k-2})$ contains $s_k s_{k-1}s_{k-2}$, forbidden by Lemma 4.4. That is, $\sigma \tau$ falls into case (3).
- Let j = k 1 for $3 \le k \le r 2$. Then $\sigma\tau = (s_{k+1}s_k)(s_ks_{k-1}) = s_{k+1}s_{k-1}$ is a pairwise independent product with $2 = \ell(s_{k+1}) + \ell(s_{k-1}) < \ell(\sigma) + \ell(\tau) = 4$. That is, $\sigma\tau$ falls into case (2).
- Let j = k for $2 \le k \le r 2$. Then $\sigma\tau = (s_{k+1}s_k)(s_{k+1}s_k) = s_{k+1}s_ks_{k+1}s_k = s_{k+1}s_{k+1}s_ks_{k+1} = s_ks_{k+1}$, a basic allowable subword of length 2. That is, $\sigma\tau$ falls into case (2).
- Let j = k + 1 for $2 \le k \le r 3$. Then $\sigma \tau = (s_{k+1}s_k)(s_{k+2}s_{k+1})$ contains $s_{k+1}s_ks_{k+2}$ forbidden by Lemma 4.4. That is, $\sigma \tau$ falls into case (3).
- Let j = k + 2 for $2 \le k \le r 4$. Then $\sigma\tau = (s_{k+1}s_k)(s_{k+3}s_{k+2}) = s_{k+3}s_{k+1}s_ks_{k+2}$ contains $s_{k+1}s_ks_{k+2}$ forbidden by Lemma 4.4. That is, $\sigma\tau$ falls into case (3).

Products of the form (d)(d): Let $\sigma = s_k s_{k+1} s_k$ for some $2 \le k \le r-2$. For $\tau = s_j s_{j+1} s_j$ with $2 \le j \le r-2$ to not be independent with σ , we must have $j \in \{k-2, k-1, k, k+1\}$.

- Let j = k 2 for $4 \le k \le r 2$. Then $\sigma\tau = (s_k s_{k+1} s_k)(s_{k-2} s_{k-1} s_{k-2})$ contains $s_{k+1} s_k s_{k-2} s_{k-1}$, forbidden by Lemma 4.5. That is, $\sigma\tau$ falls into case (3).
- Let j = k 1 for $3 \le k \le r 2$. Then $\sigma\tau = (s_k s_{k+1} s_k)(s_{k-1} s_k s_{k-1})$ contains $s_{k+1} s_k s_{k-1}$, forbidden by Lemma 4.4. That is, $\sigma\tau$ falls into case (3).
- Let j = k for $2 \le k \le r 2$. Then $\sigma\tau = (s_k s_{k+1} s_k)(s_k s_{k+1} s_k) = 1$, a product of an empty subset of S, with $0 = \ell(1) = \ell(s_k s_k) < \ell(s_k) + \ell(s_k) = 2$. That is, $\sigma\tau$ falls into case (2).
- Let j = k+1 for $2 \le k \le r-3$. Then $\sigma\tau = (s_k s_{k+1} s_k)(s_{k+1} s_{k+2} s_{k+1})$ contains $s_k s_{k+1} s_{k+2}$, forbidden by Lemma 4.4. That is, $\sigma\tau$ falls into case (3).

Products of the form (e)(e): Let $\sigma = s_{k+2}s_ks_{k+1}$ for some $2 \le k \le r-3$. For $\tau = s_{j+2}s_js_{j+1}$ with $2 \le j \le r-3$ to not be independent with σ , we must have $j \in \{k-3, k-2, k-1, k, k+1, k+2, k+3\}$.

- Let j = k 3 for $5 \le k \le r 3$. Then $\sigma\tau = (s_{k+2}s_ks_{k+1})(s_{k-1}s_{k-3}s_{k-2})$ contains $s_{k+2}s_ks_{k+1}s_{k-1}$, forbidden by Lemma 4.5. That is, $\sigma\tau$ falls into case (3).
- Let j = k 2 for $4 \le k \le r 3$. Then $\sigma\tau = (s_{k+2}s_ks_{k+1})(s_ks_{k-2}s_{k-1})$ contains $s_{k+1}s_ks_{k-2}s_{k-1}$, forbidden by Lemma 4.5. That is, $\sigma\tau$ falls into case (3).
- Let j = k 1 for $3 \le k \le r 3$. Then $\sigma\tau = (s_{k+2}s_ks_{k+1})(s_{k+1}s_{k-1}s_k) = (s_{k+2})(s_ks_{k-1}s_k)$ is a pairwise independent product with $4 = \ell(s_{k+2}) + \ell(s_ks_{k-1}s_k) < \ell(\sigma) + \ell(\tau) = 6$. That is, $\sigma\tau$ falls into case (2).
- Let j = k for $2 \le k \le r 3$. Then $\sigma\tau = (s_{k+2}s_ks_{k+1})(s_{k+2}s_ks_{k+1})$ contains $s_ks_{k+1}s_{k+2}$, forbidden by Lemma 4.4. That is, $\sigma\tau$ falls into case (3).
- Let j = k + 1 for $2 \le k \le r 4$. Then $\sigma\tau = (s_{k+2}s_ks_{k+1})(s_{k+3}s_{k+1}s_{k+2})$ contains $s_{k+2}s_ks_{k+1}s_{k+3}$, forbidden by Lemma 4.5. That is, $\sigma\tau$ falls into case (3).
- Let j = k+2 for $2 \le k \le r-5$. Then $\sigma\tau = (s_{k+2}s_ks_{k+1})(s_{k+4}s_{k+2}s_{k+3})$ contains $s_{k+1}s_{k+4}s_{k+2}s_{k+3}$, forbidden by Lemma 4.5. That is, $\sigma\tau$ falls into case (3).
- Let j = k+3 for $2 \le k \le r-6$. Then $\sigma\tau = (s_{k+2}s_ks_{k+1})(s_{k+5}s_{k+3}s_{k+4}) = (s_{k+2}s_ks_{k+1})(s_{k+3}s_{k+5}s_{k+4})$ contains $s_{k+2}s_ks_{k+1}s_{k+3}$, forbidden by Lemma 4.5. That is, $\sigma\tau$ falls into case (3).

Products of the form (a)(b) or (b)(a) (equivalent to (a)(c) or (c)(a) via f): Let $\sigma = s_k$ for some $1 \leq k \leq r$. For $\tau = s_{j+1}s_j$ with $2 \leq j \leq r-2$ to not be independent with σ , we must have $j \in \{k-2, k-1, k, k+1\}$.

12 ANDERSON, BANAIAN, FERRERI, GOFF, HADAWAY, HARRIS, HARRY, MAYERS, WANG, AND WILSON

- Let j = k 2 for $4 \le k \le r$. Then $-\sigma \tau = (s_k)(s_{k-1}s_{k-2}) = s_k s_{k-1}s_{k-2}$ is forbidden by Lemma 4.4, and $-\tau \sigma = (s_{k-1}s_{k-2})s_k = s_{k-1}s_{k-2}s_k$ is forbidden by Lemma 4.4. That is, both $\sigma \tau$ and $\tau \sigma$ fall into case (3).
- Let j = k 1 for $3 \le k \le r 1$. Then
 - $-\sigma\tau = (s_k)(s_ks_{k-1}) = s_{k-1} \in S, \text{ and} \\ -\tau\sigma = s_ks_{k-1}s_k \in S.$

That is, both $\sigma\tau$ and $\tau\sigma$ fall into case (1).

- Let j = k for $2 \le k \le r 2$. Then
 - $-\sigma\tau = (s_k)(s_{k+1}s_k) = s_k s_{k+1}s_k \in S, \text{ and}$ $-\tau\sigma = (s_{k+1}s_k)s_k = s_{k+1} \in S.$

That is, both $\sigma\tau$ and $\tau\sigma$ fall into case (1).

- Let j = k + 1 for $1 \le k \le r 3$. Then
 - $-\sigma \tau = (s_k)(s_{k+2}s_{k+1}) = s_k s_{k+2}s_{k+1} = s_{k+2}s_k s_{k+1} \in S$ if $1 < k \le r-3$ and contains the forbidden $s_1 s_2$ when k = 1, and
 - $-\tau\sigma = (s_{k+2}s_{k+1})s_k = s_{k+2}s_{k+1}s_k$ is forbidden by Lemma 4.4.

That is, both $\sigma\tau$ and $\tau\sigma$ fall into case (3) when k = 1, while $\sigma\tau$ falls into case (1) and $\tau\sigma$ falls into case (3) when $1 < k \le r - 3$.

Products of the form (a)(d) or (d)(a): Let $\sigma = s_k$ for some $1 \le k \le r$. For $\tau = s_j s_{j+1} s_j$ with $2 \le j \le r-2$ to not be independent with σ , we must have $j \in \{k-2, k-1, k, k+1\}$.

- Let j = k 2 for $4 \le k \le r$. Then $-\sigma\tau = s_k(s_{k-2}s_{k-1}s_{k-2})$ contains $s_ks_{k-1}s_{k-2}$, forbidden by Lemma 4.4, and $-\tau\sigma = (s_{k-2}s_{k-1}s_{k-2})s_k = (s_{k-1}s_{k-2}s_{k-1})s_k$ contains $s_{k-2}s_{k-1}s_k$, forbidden by Lemma 4.4. That is, both $\sigma\tau$ and $\tau\sigma$ fall into case (3).
- Let j = k 1 for $3 \le k \le r 1$. Then $-\sigma \tau = s_k(s_{k-1}s_ks_{k-1}) = s_k(s_ks_{k-1}s_k) = s_{k-1}s_k \in S$, and $-\tau \sigma = (s_{k-1}s_ks_{k-1})s_k = (s_ks_{k-1}s_k)s_k = s_ks_{k-1} \in S$. That is, both $\sigma \tau$ and $\tau \sigma$ fall into case (1).
- Let j = k for $2 \le k \le r-2$. Then $-\sigma\tau = s_k(s_ks_{k+1}s_k) = s_{k+1}s_k \in S$, and
 - $-\tau\sigma = (s_k s_{k+1} s_k) s_k = s_k s_{k+1} \in S.$
- That is, both $\sigma\tau$ and $\tau\sigma$ fall into case (1).
- Let j = k + 1 for $1 \le k \le r 3$. Then
 - $-\sigma\tau = s_k(s_{k+1}s_{k+2}s_{k+1})$ contains $s_ks_{k+1}s_{k+2}$, forbidden by Lemma 4.4, and
 - $-\tau\sigma = (s_{k+1}s_{k+2}s_{k+1})s_k$ contains $s_{k+2}s_{k+1}s_k$, forbidden by Lemma 4.4.
 - That is, both $\sigma\tau$ and $\tau\sigma$ fall into case (3).

Products of the form (a)(e) or (e)(a): Let $\sigma = s_k$ for some $1 \le k \le r$. For $\tau = s_{j+2}s_js_{j+1}$ with $2 \le j \le r-3$ to not be independent with σ , we must have $j \in \{k-3, k-2, k-1, k, k+1\}$.

- Let j = k 3 for $5 \le k \le r$. Then $-\sigma \tau = (s_k)(s_{k-1}s_{k-3}s_{k-2})$ is forbidden by Lemma 4.5, and $-\tau \sigma = (s_{k-1}s_{k-3}s_{k-2})(s_k)$ is forbidden by Lemma 4.5. That is, both $\sigma \tau$ and $\tau \sigma$ fall into case (3).
- Let j = k 2 for $4 \le k \le r 1$. Then
 - $-\sigma \tau = (s_k)(s_k s_{k-2} s_{k-1}) = s_{k-2} s_{k-1} \in S$, and
 - $-\tau\sigma = (s_k s_{k-2} s_{k-1})(s_k)$ contains $s_{k-2} s_{k-1} s_k$, forbidden by Lemma 4.4.

That is, $\sigma\tau$ falls into case (1) and $\tau\sigma$ falls into case (3).

- Let j = k 1 for $3 \le k \le r 2$. Then
 - $-\sigma\tau = (s_k)(s_{k+1}s_{k-1}s_k)$ contains $s_ks_{k-1}s_{k+1}$, forbidden by Lemma 4.4, and
 - $-\tau\sigma = (s_{k+1}s_{k-1}s_k)(s_k) = s_{k+1}s_{k-1}$ is a product of independent elements of S, with $2 = \ell(s_{k+1}) + \ell(s_{k-1}) < \ell(\tau) + \ell(\sigma) = 4$.
- That is, $\sigma \tau$ falls into case (3) and $\tau \sigma$ falls into case (2).
- Let j = k for $2 \le k \le r 3$. Then
 - $-\sigma \tau = (s_k)(s_{k+2}s_ks_{k+1}) = s_{k+2}s_{k+1} \in S$, and

 $-\tau\sigma = (s_{k+2}s_ks_{k+1})(s_k) = s_{k+2}s_{k+1}s_ks_{k+1}$ contains $s_{k+2}s_{k+1}s_k$, forbidden by Lemma 4.4. That is, $\sigma\tau$ falls into case (1) and $\tau\sigma$ falls into case (3).

• Let j = k + 1 for $1 \le k \le r - 4$. Then

 $- \sigma\tau = (s_k)(s_{k+3}s_{k+1}s_{k+2}) = s_{k+3}s_ks_{k+1}s_{k+2} \text{ contains } s_ks_{k+1}s_{k+2}, \text{ forbidden by Lemma 4.4, and}$ $- \tau\sigma = (s_{k+3}s_{k+1}s_{k+2})(s_k) \text{ contains } s_{k+1}s_ks_{k+2}, \text{ forbidden by Lemma 4.4.}$ That is, both $\sigma\tau$ and $\tau\sigma$ fall into case (3).

Products of the form (b)(c) (equivalent to (c)(b) via f): Let $\sigma = s_{k+1}s_k$ for some $2 \le k \le r-2$. For $\tau = s_j s_{j+1}$ with $2 \le j \le r-2$ to not be independent with σ , we must have $j \in \{k-2, k-1, k, k+1, k+2\}$.

- Let j = k 2 for $4 \le k \le r 2$. Then $\sigma \tau = (s_{k+1}s_k)(s_{k-2}s_{k-1})$ is forbidden by Lemma 4.5. That is, $\sigma \tau$ falls into case (3).
- Let j = k 1 for $3 \le k \le r 2$. Then $\sigma \tau = (s_{k+1}s_k)(s_{k-1}s_k)$ contains $(s_{k+1}s_ks_{k-1})$, forbidden by Lemma 4.4. That is, $\sigma \tau$ falls into case (3).
- Let j = k for $2 \le k \le r 2$. Then $\sigma\tau = (s_{k+1}s_k)(s_ks_{k+1}) = 1$, a product of an empty subset of S, with $0 = \ell(1) < \ell(s_{k+1}s_k) + \ell(s_ks_{k+1}) = 4$. That is $\sigma\tau$ falls into case (2).
- Let j = k + 1 for $2 \le k \le r 3$. Then $\sigma\tau = (s_{k+1}s_k)(s_{k+1}s_{k+2})$ contains $s_ks_{k+1}s_{k+2}$, forbidden by Lemma 4.4. That is, $\sigma\tau$ falls into case (3).
- Let j = k + 2 for $2 \le k \le r 4$. Then $\sigma\tau = (s_{k+1}s_k)(s_{k+2}s_{k+3})$ contains $s_{k+1}s_ks_{k+2}$, forbidden by Lemma 4.4. That is, $\sigma\tau$ falls into case (3).

Products of the form (b)(d) or (d)(b) (equivalent to (c)(d) or (d)(c) via f): Let $\sigma = s_{k+1}s_k$ for some $2 \le k \le r-2$. For $\tau = s_j s_{j+1}s_j$ with $2 \le j \le r-2$ to not be independent with σ , we must have $j \in \{k-2, k-1, k, k+1, k+2\}$.

- Let j = k 2 for $4 \le k \le r 2$. Then $-\sigma \tau = (s_{k+1}s_k)(s_{k-2}s_{k-1}s_{k-2})$ contains $s_{k+1}s_ks_{k-2}s_{k-1}$, forbidden by Lemma 4.5, and $-\tau \sigma = (s_{k-2}s_{k-1}s_{k-2})(s_{k+1}s_k)$ contains $s_{k-1}s_{k-2}s_{k+1}s_k$, forbidden by Lemma 4.5. That is, both $\sigma \tau$ and $\tau \sigma$ fall into case (3).
- Let j = k 1 for $3 \le k \le r 2$. Then $-\sigma \tau = (s_{k+1}s_k)(s_{k-1}s_ks_{k-1})$ contains $s_{k+1}s_ks_{k-1}$, forbidden by Lemma 4.4 and $-\tau \sigma = (s_{k-1}s_ks_{k-1})(s_{k+1}s_k)$ contains $s_ks_{k-1}s_{k+1}$, forbidden by Lemma 4.4. That is, both $\sigma \tau$ and $\tau \sigma$ fall into case (3).
- Let j = k for $2 \le k \le r 2$. Then $-\sigma \tau = (s_{k+1}s_k)(s_ks_{k+1}s_k) = s_k \in S$, and $-\tau \sigma = (s_ks_{k+1}s_k)(s_{k+1}s_k) = s_{k+1} \in S$. That is, both $\sigma \tau$ and $\tau \sigma$ fall into case (1).
- Let j = k + 1 for $2 \le k \le r 3$. Then
 - $-\sigma\tau = (s_{k+1}s_k)(s_{k+1}s_{k+2}s_{k+1}) = s_k s_{k+1} s_k s_{k+2} s_{k+1} \text{ contains } s_{k+1} s_k s_{k+2}, \text{ forbidden by}$
 - Lemma 4.4, and

 $-\tau\sigma = (s_{k+1}s_{k+2}s_{k+1})(s_{k+1}s_k) = s_{k+1}s_{k+2}s_k$ is forbidden by Lemma 4.4.

- That is, both $\sigma\tau$ and $\tau\sigma$ fall into case (3).
- Let j = k + 2 for $2 \le k \le r 4$. Then $-\sigma\tau = (s_{k+1}s_k)(s_{k+2}s_{k+3}s_{k+2})$ contains $s_{k+1}s_ks_{k+2}s_{k+3}$, forbidden by Lemma 4.5, and $-\tau\sigma = (s_{k+2}s_{k+3}s_{k+2})(s_{k+1}s_k)$ contains $s_{k+3}s_{k+2}s_{k+1}s_k$, forbidden by Lemma 4.5. That is, both $\sigma\tau$ and $\tau\sigma$ fall into case (3).

Products of the form (b)(e) or (e)(b) (equivalent to (c)(e) or (e)(c) via f): Let $\sigma = s_{k+1}s_k$ for some $2 \le k \le r-2$. For $\tau = s_{j+2}s_js_{j+1}$ with $2 \le j \le r-3$ to not be independent with σ , we must have $j \in \{k-3, k-2, k-1, k, k+1, k+2\}$.

• Let j = k - 3 for $5 \le k \le r - 2$. Then $-\sigma \tau = (s_{k+1}s_k)(s_{k-1}s_{k-3}s_{k-2})$ contains $s_k s_{k-1}s_{k-3}s_{k-2}$, forbidden by Lemma 4.5, and $-\tau \sigma = (s_{k-1}s_{k-3}s_{k-2})(s_{k+1}s_k) = s_{k-3}s_{k-1}s_{k-2}s_{k+1}s_k$ contains $s_{k-1}s_{k-2}s_{k+1}s_k$, forbidden by Lemma 4.5.

That is, both $\sigma\tau$ and $\tau\sigma$ fall into case (3).

• Let j = k - 2 for $4 \le k \le r - 2$. Then

 $- \sigma \tau = (s_{k+1}s_k)(s_k s_{k-2}s_{k-1}) = (s_{k+1})(s_{k-2}s_{k-1}) \text{ is a pairwise independent product with } 3 = \ell(s_{k+1}) + \ell(s_{k-2}s_{k-1}) < \ell(\sigma) + \ell(\tau) = 5, \text{ and}$

$$-\tau\sigma = (s_k s_{k-2} s_{k-1})(s_{k+1} s_k)$$
 contains $s_k s_{k-2} s_{k-1} s_{k+1}$, forbidden by Lemma 4.5

That is,
$$\sigma\tau$$
 falls into case (2) and $\tau\sigma$ falls into case (3).

• Let j = k - 1 for $3 \le k \le r - 2$. Then

 $-\sigma\tau=(s_{k+1}s_k)(s_{k+1}s_{k-1}s_k)=s_ks_{k+1}s_ks_{k-1}s_k$ contains $s_{k+1}s_ks_{k-1}$, forbidden by Lemma 4.4, and

 $-\tau\sigma = (s_{k+1}s_{k-1}s_k)(s_{k+1}s_k)$ contains $s_{k-1}s_ks_{k+1}$, forbidden by Lemma 4.4.

That is, both $\sigma\tau$ and $\tau\sigma$ fall into case (3).

- Let j = k for $2 \le k \le r 3$. Then $-\sigma \tau = (s_{k+1}s_k)(s_{k+2}s_ks_{k+1})$ contains $s_{k+1}s_ks_{k+2}$, forbidden by Lemma 4.4. $-\tau \sigma = (s_{k+2}s_ks_{k+1})(s_{k+1}s_k) = s_{k+2} \in S$. That is, $\sigma \tau$ falls into case (3) and $\tau \sigma$ falls into case (1).
- Let j = k + 1 for $2 \le k \le r 4$. Then $-\sigma \tau = (s_{k+1}s_k)(s_{k+3}s_{k+1}s_{k+2})$ contains $s_k s_{k+3}s_{k+1}s_{k+2}$, forbidden by Lemma 4.5, and $-\tau \sigma = (s_{k+3}s_{k+1}s_{k+2})(s_{k+1}s_k)$ contains $s_{k+2}s_{k+1}s_k$, forbidden by Lemma 4.4.

That is, both $\sigma\tau$ and $\tau\sigma$ fall into case (3).

Products of the form (d)(e) or (e)(d): Let $\sigma = s_k s_{k+1} s_k$ for some $2 \le k \le r-2$. For $\tau = s_{j+2} s_j s_{j+1}$ with $2 \le j \le r-3$ to not be independent with σ , we must have $j \in \{k-3, k-2, k-1, k, k+1, k+2\}$.

- Let j = k 3 for $5 \le k \le r 2$. Then $-\sigma\tau = (s_k s_{k+1} s_k)(s_{k-1} s_{k-3} s_{k-2})$ contains $s_k s_{k-1} s_{k-3} s_{k-2}$, forbidden by Lemma 4.5, and $-\tau\sigma = (s_{k-1} s_{k-3} s_{k-2})(s_k s_{k+1} s_k)$ contains $s_{k-1} s_{k-3} s_{k-2} s_k$, forbidden by Lemma 4.5. That is, both $\sigma\tau$ and $\tau\sigma$ fall into case (3).
- Let j = k 2 for $4 \le k \le r 2$. Then $-\sigma \tau = (s_k s_{k+1} s_k)(s_k s_{k-2} s_{k-1}) = s_k s_{k+1} s_{k-2} s_{k-1}$ is forbidden by Lemma 4.5, and $-\tau \sigma = (s_k s_{k-2} s_{k-1})(s_k s_{k+1} s_k)$ contains $s_{k-2} s_{k-1} s_k s_{k+1}$, forbidden by Lemma 4.5. That is, both $\sigma \tau$ and $\tau \sigma$ fall into case (3).
- Let j = k 1 for $3 \le k \le r 2$. Then $-\sigma \tau = (s_k s_{k+1} s_k)(s_{k+1} s_{k-1} s_k) = (s_{k+1} s_k s_{k+1})(s_{k+1} s_{k-1} s_k) = s_{k+1} s_k s_{k-1} s_k$ contains $s_{k+1} s_k s_{k-1}$, forbidden by Lemma 4.4, and $-\tau \sigma = (s_{k+1} s_{k-1} s_k)(s_k s_{k+1} s_k) = s_{k+1} s_{k-1} s_{k+1} s_k = s_{k-1} s_k \in S.$ That is, $\sigma \tau$ falls into case (3) and $\tau \sigma$ falls into case (1).
- Let j = k for $2 \le k \le r-3$. Then $-\sigma\tau = (s_k s_{k+1} s_k)(s_{k+2} s_k s_{k+1})$ contains $s_{k+1} s_k s_{k+2}$, forbidden by Lemma 4.4, and $-\tau\sigma = (s_{k+2} s_k s_{k+1})(s_k s_{k+1} s_k) = (s_{k+2} s_k s_{k+1})(s_{k+1} s_k s_{k+1}) = s_{k+2} s_{k+1} \in S$. That is, $\sigma\tau$ falls into case (3) and $\tau\sigma$ falls into case (1).
- Let j = k + 1 for $2 \le k \le r 4$. Then $-\sigma \tau = (s_k s_{k+1} s_k)(s_{k+3} s_{k+1} s_{k+2})$ contains $s_k s_{k+3} s_{k+1} s_{k+2}$, forbidden by Lemma 4.5, and $-\tau \sigma = (s_{k+3} s_{k+1} s_{k+2})(s_k s_{k+1} s_k)$ contains $s_{k+3} s_{k+1} s_{k+2} s_k$, forbidden by Lemma 4.5. That is, both $\sigma \tau$ and $\tau \sigma$ fall into case (3).
- Let j = k + 2 for $2 \le k \le r 5$. Then $-\sigma \tau = (s_{k+1}s_ks_{k+1})(s_{k+4}s_{k+2}s_{k+3})$ contains $s_{k+1}s_{k+4}s_{k+2}s_{k+3}$, forbidden by Lemma 4.5, and $-\tau \sigma = (s_{k+4}s_{k+2}s_{k+3})(s_{k+1}s_ks_{k+1})$ contains $s_{k+4}s_{k+2}s_{k+3}s_{k+1}$, forbidden by Lemma 4.5. That is, both $\sigma \tau$ and $\tau \sigma$ fall into case (3).

This completes the proof.

Next, we characterize the set of basic allowable subwords in the case $\mu = -\tilde{\alpha}$.

Theorem 4.7. The set $BAS(\tilde{\alpha}, -\tilde{\alpha})$ of basic allowable subwords of $\mathcal{A}(\tilde{\alpha}, -\tilde{\alpha})$ consists of

- (a) s_k with $1 \le k \le r$,
- (b) $s_{k+1}s_k$ with $2 \le k \le r-2$,
- (c) $s_k s_{k+1}$ with $2 \le k \le r-2$,
- (d) $s_k s_{k+1} s_k$ with $2 \le k \le r-2$, and
- (e) $s_{k+2}s_ks_{k+1}$ with $2 \le k \le r-3$.

Proof. Let S denote the set of elements given in the statement of Theorem 4.7. We proceed by showing that S satisfies the conditions in Proposition 3.10.

Since S contains all simple transpositions s_k for $1 \le k \le r$, it clearly satisfies (C1). Next, considering Lemma 3.7 (1), we can read off the influence of each element as the indices present on simple transpositions in their reduced expressions given above. As these sets are clearly connected, S satisfies (C2). Finally, Lemma 4.6 establishes that S satisfies (C3). Thus, $S = BAS(\tilde{\alpha}, -\tilde{\alpha})$, as desired. П

4.4. Weyl alternation sets $\mathcal{A}(\tilde{\alpha},\mu)$ for other negative roots μ . In this subsection, we consider other negative roots besides the negative hightest root and apply Proposition 3.12 to obtain the Weyl alternation sets $\mathcal{A}(\tilde{\alpha},\mu)$ from that of $\mathcal{A}(\tilde{\alpha},-\tilde{\alpha})$ found above. Each $\mu \in \Phi^-$ can be written as

$$u = -(\alpha_i + \alpha_{i+1} + \dots + \alpha_j)$$

for $1 \leq i \leq j \leq r$.

For the remainder of this paper, we let

$$\alpha_{i,j} = \sum_{k=i}^{j} \alpha_k$$

for any $1 \le i \le j \le r$ and as expected $\alpha_{i,i} = \alpha_i$. Because the actions of s_1 and s_r on weights are distinct from that of s_k , with $2 \le k \le r-1$, we group the negative roots into three cases: (1) $1 = i \le j < r$; (2) $1 < i \leq j = r$; and (3) $1 < i \leq j < r$. In each case, the corresponding corollary below of Proposition 3.12 is obtained by computing the intersection of $BAS(\tilde{\alpha}, -\tilde{\alpha})$, as characterized in Theorem 4.7, with the Weyl alternation set for the relevant negative root. The proofs are similar to that of Lemma 4.6.

Corollary 4.8. Let $\mu = -\alpha_{1,j}$ for $1 \leq j < r$. The set $BAS(\tilde{\alpha}, \mu)$ of basic allowable subwords of $\mathcal{A}(\tilde{\alpha}, \mu)$ consists of

- (a) s_k with $1 \le k < r$,
- (b) $s_{k+1}s_k$ with $2 \le k \le j-1$,
- (c) $s_k s_{k+1}$ with $2 \le k \le \min(j, r-2)$,
- (d) $s_k s_{k+1} s_k$ with $2 \le k \le j-1$, and (e) $s_{k+2} s_k s_{k+1}$ with $2 \le k \le j-2$.

Corollary 4.9. Let $\mu = -\alpha_{i,r}$ for $1 < i \leq r$. The set $BAS(\tilde{\alpha}, \mu)$ of basic allowable subwords of $\mathcal{A}(\tilde{\alpha}, \mu)$ consists of

- (a) s_k with $1 < k \leq r$,
- (a) s_k with $1 \le k \le r$, (b) $s_{k+1}s_k$ with $\max(i-1,2) \le k \le r-2$, (c) $s_ks_{k+1}s_k$ with $i \le k \le r-2$, and (c) $s_{k+2}s_ks_{k+1}$ with $i \le k \le r-2$, (c) $s_{k+2}s_ks_{k+1}$ with $i \le k \le r-3$.
- (c) $s_k s_{k+1}$ with $i \leq k \leq r-2$,

Corollary 4.10. Let $\mu = -\alpha_{i,j}$ for $1 < i \leq j < r$. The set $BAS(\tilde{\alpha}, \mu)$ of basic allowable subwords of $\mathcal{A}(\tilde{\alpha}, \mu)$ consists of

- (a) s_k with 1 < k < r,
- (d) $s_k s_{k+1} s_k$ with $i \le k \le j-1$, and (e) $s_{k+2} s_k s_{k+1}$ with $i \le k \le j-2$. (b) $s_{k+1}s_k$ with $\max(i-1,2) \le k \le j-1$,
- (c) $s_k s_{k+1}$ with $i \le k \le \min(j, r-2)$,

5. Enumerative results

Having characterized the basic allowable words for Weyl alternation sets $\mathcal{A}_r(\tilde{\alpha},\mu)$ for negative roots μ in the Lie algebra of type A_r , we now provide some enumerative results and generating functions for these sets. To begin, we define the Fibonacci numbers by $F_0 = 0$, $F_1 = 1$, $F_2 = 1$, and for $n \ge 2$, $F_n = F_{n-1} + F_{n-2}$. Harris's work [5, Theorem 1.2] establishes that $|\mathcal{A}_r(\tilde{\alpha}, 0)| = F_r$. We can recover this result from our study of $\mathcal{A}(\tilde{\alpha}, -\tilde{\alpha})$ and Proposition 3.12. More precisely, the only elements from $BAS(\tilde{\alpha}, -\tilde{\alpha})$ that would also be in $\mathcal{A}(\tilde{\alpha}, 0)$ are commuting products of $\{s_i : 2 \leq i \leq r-1\}$, the number of which is a Fibonacci number.

We begin this section by enumerating families of Weyl alternation sets $\mathcal{A}_r(\tilde{\alpha},\mu)$ where μ is the sum of only a few negative simple roots. Throughout this section, we let $BAS_r(\lambda,\mu)$ refer to the basic allowable subwords of $\mathcal{A}_r(\lambda,\mu)$. To prove such results, we require the following lemma. We point the reader to [14, Proposition 2.3 for a proof of the classical combinatorial result.

Lemma 5.1. The number of subsets of [n] that do not contain a pair of consecutive numbers is F_{n+2} .

Our first result considers the case where μ is a negative root of height 1 i.e., $\mu = -\alpha_i$ for some index *i*. Recall that the height of a positive root in type A is the number of simple roots.

Proposition 5.2. Let $r \ge 1$ and fix $1 \le i \le r$.

16 ANDERSON, BANAIAN, FERRERI, GOFF, HADAWAY, HARRIS, HARRY, MAYERS, WANG, AND WILSON

- (1) If i = 1 or i = r, then $|\mathcal{A}_r(\tilde{\alpha}, -\alpha_i)| = F_{r+1}$.
- (2) If r > 2 and $2 \le i \le r 1$, then $|\mathcal{A}_r(\tilde{\alpha}, -\alpha_i)| = F_r + F_{i-1}F_{r-i-1} + F_{i-2}F_{r-i}$.

Proof. For item (1): Suppose i = 1. Then, Corollary 4.8 yields $BAS_r(\tilde{\alpha}, -\alpha_1) = \{s_k : 1 \le k < r\}$, so that $\mathcal{A}_r(\tilde{\alpha}, -\alpha_1)$ consists of commuting products of r - 1 simple transpositions. By Lemma 5.1, the number of such products is F_{r+1} . The proof for i = r is similar.

For item (2): First, suppose i = 2. By Corollary 4.10, $BAS_r(\tilde{\alpha}, -\alpha_2) = \{s_k : 1 < k < r\} \cup \{s_2s_3\}$. If r > 3, the elements of $\mathcal{A}_r(\tilde{\alpha}, -\alpha_2)$ can be split into two sets: those that contain s_2s_3 as a subword and those that do not. The latter can be identified with $\mathcal{A}_r(\tilde{\alpha}, 0)$ so that there are F_r such elements. The former are in bijection with commuting products of $\{s_k : 4 < k < r\}$, of which there are F_{r-3} considering Lemma 5.1. If r = 3, then we only have the former case; since we defined $F_{-1} = 0$, the formula still holds. Thus,

$$|\mathcal{A}_r(\tilde{\alpha}, -\alpha_2)| = F_r + F_{r-3} = F_r + F_1 F_{r-3} + F_0 F_{r-2}$$

as claimed. The case i = r - 1 is analogous.

Next, suppose 2 < i < r-1. By Corollary 4.10, $BAS_r(\tilde{\alpha}, -\alpha_i) = \{s_k : 1 < k < r\} \cup \{s_i s_{i-1}, s_i s_{i+1}\}$. We now can break $\mathcal{A}_r(\tilde{\alpha}, -\alpha_i)$ into three disjoint sets:

- (1) those elements without subwords $s_i s_{i-1}$ and $s_i s_{i+1}$,
- (2) those with subword $s_i s_{i+1}$ and without $s_i s_{i-1}$, and
- (3) those with subword $s_i s_{i-1}$ and without $s_i s_{i+1}$.

Since elements in the Weyl alternation set are products of pairwise independent subsets of the basic allowable subwords, we have no elements in $\mathcal{A}_r(\tilde{\alpha}, -\alpha_i)$ with both $s_i s_{i+1}$ and $s_i s_{i-1}$ as subwords.

The number of elements described by (1) is again F_r . The number of elements described by (2) is equal to the number of pairwise commuting products of $\{s_j : 2 \le j \le i-2\}$ and $\{s_k : i+3 \le k \le r-1\}$, of which there are $F_{i-1}F_{r-i-1}$ considering Lemma 5.1. The number of elements (3) is equal to the number of pairwise commuting products of $\{s_j : 2 \le j \le i-3\}$ and $\{s_k : i+2 \le j \le r-1\}$, of which there are $F_{i-2}F_{r-i}$ considering Lemma 5.1. Summing the contributions of these sets gives us

$$|\mathcal{A}_{r}(\tilde{\alpha}, -\alpha_{i})| = F_{r} + F_{i-1}F_{r-i-1} + F_{i-2}F_{r-i},$$

as desired.

If we set i = 3 in item (2) of Proposition 5.2, we recover the Lucas numbers.

Corollary 5.3. Define the Lucas numbers by $L_0 = 2, L_1 = 1$ and $L_n = L_{n-1} + L_{n-2}$ for all $n \ge 2$. Then, if $r \ge 4$, we have $|\mathcal{A}_r(\tilde{\alpha}, -\alpha_3)| = L_{r-1}$.

Next, we consider the case where μ is a negative root of height 2. As arguments for the cardinalities of subsets of $\mathcal{A}_r(\tilde{\alpha}, -\alpha_i - \alpha_{i+1})$ considered in the proof of the following result are similar to those given in the proof Proposition 5.2, for the sake of brevity, such arguments are omitted.

Proposition 5.4.

- (1) If $r \ge 2$ and i = 1 or i = r 1, then $|\mathcal{A}_r(\tilde{\alpha}, -\alpha_i \alpha_{i+1})| = F_{r+1} + F_{r-3} = 3F_{r-1}$.
- (2) If $r \ge 4$ and $2 \le i \le r-2$, then $|\mathcal{A}_r(\tilde{\alpha}, -\alpha_i \alpha_{i+1})| = F_r + F_{i-2}F_{r-i} + 3F_{i-1}F_{r-i-1} + F_iF_{r-i-2}$.

Proof. For item (1): Suppose $r \ge 2$ and i = 1.

- If r = 2, then $-\alpha_1 \alpha_2 = -\tilde{\alpha}$. By Theorem 4.7, we have that $BAS_2(\tilde{\alpha}, -\tilde{\alpha}) = \{s_1, s_2\}$. Hence, $|\mathcal{A}_2(\tilde{\alpha}, -\tilde{\alpha})| = |\{1, s_1, s_2\}| = 3 = 3F_1$.
- If r = 3, then by Corollary 4.8, we have that $BAS_3(\tilde{\alpha}, -\alpha_1 \alpha_2) = \{s_1, s_2\}$. Hence, $|\mathcal{A}_3(\tilde{\alpha}, -\alpha_1 \alpha_2)| = |\{1, s_1, s_2\}| = 3 = 3F_2$.
- Now for $r \ge 4$, using Corollary 4.8, we have that $BAS_r(\tilde{\alpha}, -\alpha_1 \alpha_2) = \{s_k : 1 \le k < r\} \cup \{s_2s_3\}$. Hence, $\mathcal{A}_r(\tilde{\alpha}, -\alpha_1 - \alpha_2)$ can be split into two sets of elements: those that contain s_2s_3 as a subword and those that do not. The number of those elements not containing s_2s_3 is F_{r+1} because we are choosing non-consecutive indices from the set [r-1]. The number of elements containing s_2s_3 is given by F_{r-3} . Hence $|\mathcal{A}_r(\tilde{\alpha}, -\alpha_1 - \alpha_2)| = F_{r+1} + F_{r-3} = 3F_{r-1}$.

When i = r - 1, the argument is analogous.

For item (2): Suppose $r \ge 4$ and $2 \le i \le r - 2$.

• If 2 < i < r - 2, then by Corollary 4.10, we have that

$$BAS(\tilde{\alpha}, -\alpha_i - \alpha_{i+1}) = \{s_k : 1 < k < r\} \cup \begin{cases} \tau_1 = s_i s_{i-1}, \\ \tau_2 = s_{i+1} s_i, \\ \tau_3 = s_i s_{i+1}, \\ \tau_4 = s_{i+1} s_{i+2}, \\ \tau_5 = s_i s_{i+1} s_i \end{cases}$$

We can partition $\mathcal{A}_r(\tilde{\alpha}, -\alpha_i)$ into the following disjoint sets:

- (1) those elements containing τ_1 ,
- (2) those elements containing τ_2 ,
- (3) those elements containing τ_3 ,
- (4) those elements containing τ_4 ,
- (5) those elements containing τ_5 , and
- (6) those elements without any τ_n for $1 \le n \le 5$.

Observe that whenever an element contains τ_n with $n \in \{1, 2, 3, 4, 5\}$, it cannot contain τ_m with $m \in \{1, 2, 3, 4, 5\} \setminus \{n\}$ as the elements τ_n are not independent.

The sets of elements considered in cases (2), (3), and (5) have the same cardinality. In particular, the elements in each case are in bijection with the number of ways to select non-consecutive indices from $\{2, 3, \ldots, i-2\} \cup \{i+3, i+4, \ldots, r-1\}$, so that there are $F_{i-1}F_{r-i-1}$ such elements. Hence, we have in total over these three cases $3F_{i-1}F_{r-i-1}$ elements in the Weyl alternation set. Similarly, the number of elements considered in case (1) is given by $F_{i-2}F_{r-i}$; the number of elements considered in case (4) is given by F_iF_{r-i-2} ; and, the number of elements considered in (6) is given by F_r .

Therefore,

$$|\mathcal{A}_r(\tilde{\alpha}, -\alpha_i - \alpha_{i+1})| = 3F_{i-1}F_{r-i-1} + F_{i-2}F_{r-i} + F_iF_{r-i-2} + F_r.$$

If i = 2, then the same argument holds as in the previous item, except we do not have a term s_{i-1}s_i since s₁ is not in the support of any element of A(α̃, -α₂ - α₃). Since F₀ = 0 the formula still holds.
If i = r - 2, then the argument is analogous to the case where i = 2.

The prevalence of sums of Fibonacci numbers in the previous results comes from a general behavior concerning cardinalities of these Weyl alternation sets. Ongoing, we fix a pair $1 \leq i \leq j$, let $\mu = -(\alpha_i + \alpha_{i+1} + \cdots + \alpha_j)$, vary r, and compute $\mathcal{A}_r(\tilde{\alpha}, \mu)$. Naturally, this expression only makes sense when $r \geq j$. Next, we show that the cardinality of $\mathcal{A}_r(\tilde{\alpha}, \mu)$ satisfies the Fibonacci recurrence.

Proposition 5.5. Let $r \geq 3$ and $1 \leq i \leq j \leq r-2$. If $\mu = -\alpha_{i,j}$, then

$$|\mathcal{A}_r(\tilde{\alpha},\mu)| = |\mathcal{A}_{r-1}(\tilde{\alpha},\mu)| + |\mathcal{A}_{r-2}(\tilde{\alpha},\mu)|.$$

Proof. First, suppose j < r-2. From Corollary 4.8 and Corollary 4.10, we get that $\text{BAS}_r(\tilde{\alpha}, \mu) = \text{BAS}_{r-1}(\tilde{\alpha}, \mu) \cup \{s_{r-1}\}$, so we can partition the elements of $\mathcal{A}_r(\tilde{\alpha}, \mu)$ into two sets based on whether the elements contain s_{r-1} as a subword or not. The set with the elements not containing s_{r-1} is equivalent to $\mathcal{A}_{r-1}(\tilde{\alpha}, \mu)$. The set of elements containing s_{r-1} are in bijection with $\mathcal{A}_{r-2}(\tilde{\alpha}, \mu)$. Thus, $|\mathcal{A}_r(\tilde{\alpha}, \mu)| = |\mathcal{A}_{r-1}(\tilde{\alpha}, \mu)| + |\mathcal{A}_{r-2}(\tilde{\alpha}, \mu)|$, as claimed.

Now, let j = r - 2. In this case, there are two elements of $BAS_r(\tilde{\alpha}, \mu)$ containing a s_{r-1} in their reduced expressions: s_{r-1} and $s_{r-2}s_{r-1}$. The elements of $\mathcal{A}_r(\tilde{\alpha}, \mu)$ not including either of these as a subword can again be regarded as elements of $\mathcal{A}_{r-1}(\tilde{\alpha}, \mu)$. Hence, there are $|\mathcal{A}_{r-1}(\tilde{\alpha}, \mu)|$ such elements. Now, consider the subset of $\mathcal{A}_r(\tilde{\alpha}, \mu)$ consisting of elements containing the basic allowable subword s_{r-1} . These elements are in bijection with elements of $\mathcal{A}_{r-2}(\tilde{\alpha}, \mu)$ that do not include the basic allowable subword s_{r-2} via the map which removes s_{r-1} . Similarly, the subset of elements in $\mathcal{A}_r(\tilde{\alpha}, -\mu)$ consisting of elements containing the basic allowable subword $s_{r-2}s_{r-1}$ is in bijection with elements of $\mathcal{A}_{r-2}(\tilde{\alpha}, -\mu)$ that include the basic allowable subword s_{r-2} . As elements in $\mathcal{A}_{r-2}(\tilde{\alpha}, \mu)$ either contain s_{r-2} or not, the total number of elements is given by $|\mathcal{A}_{r-2}(\tilde{\alpha}, \mu)|$. Thus, the total number of elements of $\mathcal{A}_r(\tilde{\alpha}, \mu)$ containing s_{r-1} or $s_{r-2}s_{r-1}$ is also given by $|\mathcal{A}_{r-2}(\tilde{\alpha}, \mu)|$. Taking the sum over these cases shows that the desired recurrence still holds.

Figure 3 gives an example of the recursion in Proposition 5.5. To prepare for later computations, we provide a general result concerning the generating function for any sequence defined by the two term homogeneous linear recurrence $a_n = a_{n-1} + a_{n-2}$.

18



FIGURE 3. The Hasse diagram for $\mathcal{A}_7(\tilde{\alpha}, \mu)$ in the left weak Bruhat order with $\mu = -\alpha_{2,4}$. The blue subposet with thick edges is $\mathcal{A}_6(\tilde{\alpha}, \mu)$, and the subposet of boxed values is $\mathcal{A}_5(\tilde{\alpha}, \mu)$. Hence $\mathcal{A}_7(\tilde{\alpha}, \mu)$ decomposes into two pieces: one being $\mathcal{A}_6(\tilde{\alpha}, \mu)$ and the other being a translation of $\mathcal{A}_5(\tilde{\alpha}, \mu)$ by s_6 along the dotted edges.

Lemma 5.6. Let $\{a_i\}_{i\geq 0}$ be a sequence such that $a_i = 0$ for $0 \leq i < r$, $a_r = c$, $a_{r+1} = d$, and for $i \geq r+2$, $a_i = a_{i-1} + a_{i-2}$. Then,

$$\sum_{i\geq 0} a_i x^i = \frac{cx^r + (d-c)x^{r+1}}{1 - x - x^2}$$

Proof. We follow a standard computation method regarding generating functions for recursively defined sequences. Let $A(x) = \sum_{i>0} a_i x^i$. Then, we have

$$\begin{aligned} A(x) &= \sum_{i \ge 0} a_i x^i = \sum_{i \ge r} a_i x^i = c x^r + d x^{r+1} + \sum_{i \ge r+2} a_i x^i = c x^r + d x^{r+1} + \sum_{i \ge r+2} a_{i-1} x^i + \sum_{i \ge r+2} a_{i-2} x^i \\ &= c x^r + d x^{r+1} + x (A(x) - c x^r) + x^2 A(x). \end{aligned}$$

The result now follows after a few manipulations of this equation. Solving for A(x) in the above equation, the result follows.

Now, we turn our attention to tracking the sizes of Weyl alternation sets simultaneously. To prepare for this result, it is advantageous to consider the set

 $P_r^i \coloneqq \{ w \in \mathcal{A}_r(\tilde{\alpha}, -\alpha_{i,r}) : r \text{ is not in the influence of } w \}.$

In other words, if r is not in the influence of $w \in \mathcal{A}(\tilde{\alpha}, -\alpha_{i,r})$, then s_r does not appear in a reduced expression for w. We set $p_r^i \coloneqq |P_r^i|$. Our proof for the following result uses the same technique as in [10, Proposition 4.6].

Lemma 5.7. Given $r \ge i + 4$, the sequence p_r^i satisfies the recurrence

$$p_r^i = p_{r-1}^i + p_{r-2}^i + 3p_{r-3}^i + p_{r-4}^i$$

Proof. To establish the result, we provide a bijection

$$\phi: P_{r-1}^i \cup P_{r-2}^i \cup Y_1 \cup Y_2 \cup Y_3 \cup P_{r-4}^i \to P_r^i$$

where Y_1 , Y_2 , and Y_3 are distinct copies of P_{r-3}^i . First of all, from our description of each Weyl alternation set, we can find an isomorphic copy of P_{r-1}^i in the set P_r^i . This inclusion is just the natural inclusion of the symmetric group on n letters into the symmetric group on n + 1 letters. Thus, if $w \in P_{r-1}^i$, we set $\phi(w) = w$. The cokernel of this map is all $w \in P_r^i$ containing s_{r-1} in their reduced expressions. More specifically, the cokernel consists of all $w \in P_r^i$ containing the subword $s_{r-1}, s_{r-2}s_{r-1}, s_{r-1}s_{r-2}, s_{r-2}s_{r-1}s_{r-2}$, or $s_{r-3}s_{r-1}s_{r-2}$.

Given $w \in P_{r-2}^i$, we set $\phi(w) = ws_{r-1}$. Then $\phi(w) \in P_r^i$ for $w \in P_{r-2}^i$ and the image $\phi(P_{r-2}^i)$ constitutes all elements of P_r^i containing the subword s_{r-1} . Next, if $w \in Y_1$, we set $\phi(w) = ws_{r-2}s_{r-1}$. Thus, $\phi(w) \in P_r^i$ for $w \in Y_1$ and the image $\phi(Y_1)$ constitutes all elements of P_r^i containing the subword $s_{r-2}s_{r-1}$. Similarly, if $w \in Y_2$, we set $\phi(w) = ws_{r-1}s_{r-2}$, and if $w \in Y_3$, we set $\phi(w) = ws_{r-2}s_{r-1}s_{r-2}$. So $\phi(w) \in P_r^i$ for $w \in Y_2$ and Y_3 , the image $\phi(Y_2)$ constitutes all elements of P_r^i containing the subword $s_{r-1}s_{r-2}$, and the image $\phi(Y_3)$ constitutes all elements of P_r^i containing the subword $s_{r-2}s_{r-1}s_{r-2}$. Finally, if $w \in P_{r-4}^i$, we set $\phi(w) = ws_{r-3}s_{r-1}s_{r-2}$. Consequently, $\phi(w) \in P_r^i$ for $w \in P_{r-4}^i$ and the image $\phi(P_{r-4}^i)$ constitutes all elements of P_r^i containing the subword $s_{r-3}s_{r-1}s_{r-2}$. Thus, ϕ is indeed a bijection, which shows that the cardinalities of the sets satisfy the desired recurrence.

Lemma 5.8. If $3 < i \leq r$, then the p_r^i satisfy the recurrence

$$p_r^i = p_{r-1}^{i-1} + p_{r-2}^{i-2}$$

Proof. Since i > 3, we know $\max(i-1,2) = i-1$ and $\max(i-2,2) = i-2$. Therefore, if we take the description of $BAS_{r-1}(\tilde{\alpha}, \alpha_{i-1,r-1}) \setminus \{s_{r-1}\}$ and add one to each index, we have $BAS_r(\tilde{\alpha}, \alpha_{i-1,r-1}) \setminus \{s_2, s_r\}$. By the shift in indices, p_{r-1}^{i-1} enumerates the elements of P_r^i that do not contain s_2 in their reduced expressions.

It remains to enumerate the elements of P_r^i that contain s_2 in their reduced expressions. Since $i \geq 4$, when we remove the word s_2 , the resulting Weyl group element is a pairwise independent product of

- (a) s_k with 3 < k < r,
- (d) $s_k s_{k+1} s_k$ with $i \le k \le r-2$, and (e) $s_{k+2} s_k s_{k+1}$ with $i \le k \le r-3$. (b) $s_{k+1}s_k$ with $\max(i-1,4) \le k \le r-2$, (c) s_ks_{k+1} with $i \le k \le r-2$,

Note, if we subtract 2 from all indices, we have a description of $BAS_{r-2}(\tilde{\alpha}, -\alpha_{i-2,r-2}) \setminus \{s_{r-2}\}$. This means that the number of elements of P_r^i that contain s_2 in their reduced expressions is precisely p_{r-2}^{i-2} , completing the proof.

Lemma 5.9. If $1 \le i < r$, then $|\mathcal{A}_r(\tilde{\alpha}, -\alpha_{i,r-1})| = p_r^i$.

Proof. Let i = 1. Then, comparing the inequalities in Theorem 4.7 to those in Corollary 4.8 with j = r - 1, we have that

$$BAS_r(\tilde{\alpha}, -\alpha_{1,r-1}) = BAS_r(\tilde{\alpha}, -\alpha_{1,r}) \setminus \{s_r\}.$$

Similarly, for 1 < i < r, by comparing the inequalities in Corollary 4.9 to those in Corollary 4.10 with j = r - 1, we have also that

$$BAS_r(\tilde{\alpha}, -\alpha_{i,r-1}) = BAS_r(\tilde{\alpha}, -\alpha_{i,r}) \setminus \{s_r\}.$$

The result then follows from the definition of p_r^i as the number of elements in $\mathcal{A}(\tilde{\alpha}, -\alpha_{i,r})$ without s_r in their reduced expressions and the fact that the only basic allowable subword containing s_r is the element s_r itself.

To prepare for our next results, we set $h_r^i = |\mathcal{A}_r(\tilde{\alpha}, -\alpha_{i,r})|$. Next, we show that the quantities h_r^i and p_r^i are closely related.

Lemma 5.10. If $r \ge 2$ and $1 \le i \le r-1$, then $h_r^i = p_r^i + p_{r-1}^i$.

Proof. To establish the result, we provide a bijection between $P_r^i \cup P_{r-1}^i$ and $\mathcal{A}_r(\tilde{\alpha}, -\alpha_{i,r})$. Given $w \in P_r^i$, we set $\phi(w) = w$. By definition, the image $\phi(P_r^i)$ consists of all $w \in \mathcal{A}_r(\tilde{\alpha}, -\alpha_{i,r})$ without a subword s_r , of which there are p_r^i . Given $w \in P_{r-1}^i$, we set $\phi(w) = ws_r$. One can check that $ws_r \in \mathcal{A}_r(\tilde{\alpha}, -\alpha_{i,r})$, and moreover that all $\tau \in \mathcal{A}_r(\tilde{\alpha}, -\alpha_{i,r})$ with a subword s_r can be written this way. Recall that there are p_{r-1}^i such elements of $\mathcal{A}_r(\tilde{\alpha}, -\alpha_{i,r})$. As these subsets of $\mathcal{A}_r(\tilde{\alpha}, -\alpha_{i,r})$ are disjoint, ϕ defines the desired bijection, and we get $h_r^i = p_r^i + p_{r-1}^i$.

Now that we know recurrences for the sequence $\{p_r^i\}_r$ and a relationship between the sequences $\{p_r^i\}_r$ and $\{h_r^i\}_r$, we can conclude that the two sequences satisfy the same recurrences.

i	p_i^i	p_{i+1}^i	p_{i+2}^{i}	p_{i+3}^{i}	h_i^i	h_{i+1}^i	h_{i+2}^i	h_{i+3}^i
1	1	2	3	8	1	3	5	11
2	1	2	6	12	2	3	8	18
3	2	4	9	20	3	6	13	29
TABLE 2. The values of p_r^i and h_r^i for $i \in \{1, 2, 3\}$ and $i \le r \le i + 2$.								

Corollary 5.11. If $r \ge i + 4$, then

$$h_r^i = h_{r-1}^i + h_{r-2}^i + 3h_{r-3}^i + h_{r-4}^i$$

If $i, r \geq 3$, then

$$h_r^i = h_{r-1}^{i-1} + h_{r-2}^{i-2}.$$

By computing initial conditions using Theorem 3.8 and Corollary 4.9, we provide generating functions for both of these sequences.

Lemma 5.12. Define
$$\mathcal{P}^i(x) = \sum_{r \ge i} p_r^i x^r$$
 and $\mathcal{H}^i(x) = \sum_{r \ge i} h_r^i x^r$.

(1) If i = 1, then

$$\mathcal{P}^{1}(x) = \frac{x + x^{2}}{1 - x - x^{2} - 3x^{3} - x^{4}} \qquad and \qquad \mathcal{H}^{1}(x) = \frac{x + 2x^{2} + x^{3}}{1 - x - x^{2} - 3x^{3} - x^{4}}.$$

(2) If
$$i = 2$$
, then

$$\mathcal{P}^{2}(x) = \frac{x^{2} + x^{3} + 3x^{4} + x^{5}}{1 - x - x^{2} - 3x^{3} - x^{4}} \qquad and \qquad \mathcal{H}^{2}(x) = \frac{2x^{3} + x^{4} + 3x^{5} + x^{6}}{1 - x - x^{2} - 3x^{3} - x^{4}}.$$

(3) If i = 3, then

$$\mathcal{P}^{3}(x) = \frac{2x^{3} + 2x^{4} + 3x^{5} + x^{6}}{1 - x - x^{2} - 3x^{3} - x^{4}} \qquad and \qquad \mathcal{H}^{3}(x) = \frac{3x^{3} + 3x^{4} + 4x^{5} + x^{6}}{1 - x - x^{2} - 3x^{3} - x^{4}}$$

Furthermore, defining $\mathcal{P}(x,s) = \sum_{i \ge 1} \mathcal{P}^i(x) s^i$ and $\mathcal{H}(x,s) = \sum_{i \ge 1} \mathcal{H}^i(x) s^i$, we have

$$\mathcal{P}(x,s) = \frac{xs(x^4s + 3x^3s + x + 1)}{(1 - x - x^2 - 3x^3 - x^4)(1 - xs - (xs)^2)}$$

and

$$\mathcal{H}(x,s) = \frac{xs(x^5s + 3x^4s - xs + x^2 + 2x + 1)}{(1 - x - x^2 - 3x^3 - x^4)(1 - xs - (xs)^2)}.$$

Proof. For the cases i = 1, 2, and 3, reasoning as in the proof of Lemma 5.6, the recursions in Lemma 5.7 and Corollary 5.11 guarantee these generating functions must have the form

$$\frac{ax^{i} + bx^{i+1} + cx^{i+2} + dx^{i+3}}{1 - x - x^{2} - 3x^{3} - x^{4}} = ax^{i} + (a+b)x^{i+1} + (2a+b+c)x^{i+2} + (6a+2b+c+d)x^{i+3} + \cdots$$

with coefficients a, b, c, and d depending on the initial conditions. The formulas for these cases are then obtained by first computing the values of p_r^i and h_r^i for the first four values as shown in Table 2, then solving for the coefficients a, b, c, and d.

We now apply the recursion $p_r^i = p_{r-1}^{i-1} + p_{r-2}^{i-2}$ from Lemma 5.8 for $r \ge i \ge 3$ to obtain a formula for $\mathcal{P}(x,s)$. For $i \ge 3$, we first compute

$$\begin{split} \mathcal{P}^{i}(x) &= \sum_{r \geq i} p_{r}^{i} x^{r} \\ &= \sum_{r \geq i} (p_{r-1}^{i-1} + p_{r-2}^{i-2}) x^{r} \\ &= \sum_{r \geq i} p_{r-1}^{i-1} x^{r} + \sum_{r \geq i} p_{r-2}^{i-2} x^{r} \\ &= x \sum_{r \geq i-1} p_{r}^{i-1} x^{r} + x^{2} \sum_{r \geq i-2} p_{r}^{i-2} x^{r} \\ &= x \mathcal{P}^{i-1}(x) + x^{2} \mathcal{P}^{i-2}(x). \end{split}$$

With this recursion in hand, we compute

$$\begin{aligned} \mathcal{P}(x,s) &= \sum_{i \ge 1} \mathcal{P}^{i}(x)s^{i} \\ &= \mathcal{P}^{1}(x)s + \mathcal{P}^{2}(x)s^{2} + \sum_{i \ge 3} \mathcal{P}^{i}(x)s^{i} \\ &= \mathcal{P}^{1}(x)s + \mathcal{P}^{2}(x)s^{2} + \sum_{i \ge 3} x\mathcal{P}^{i-1}(x)s^{i} + \sum_{i \ge 3} x^{2}\mathcal{P}^{i-2}(x)s^{i} \\ &= \mathcal{P}^{1}(x)s + \mathcal{P}^{2}(x)s^{2} + xs(\mathcal{P}(x,s) - \mathcal{P}^{1}(x)s) + (xs)^{2}\mathcal{P}(x,s) \end{aligned}$$

Now, note $(1 - xs - (xs)^2)\mathcal{P}(x, s) = (1 - xs)\mathcal{P}^1(x)s + \mathcal{P}^2(x)s^2$, and $\mathcal{P}(x, s) = \frac{(1 - xs)\mathcal{P}^1(x)s + \mathcal{P}^2(x)s^2}{1 - xs - (xs)^2}$ $(1 - xs)(x + x^2)s + (x^2 + x^3 + 3x^4 + 3x^4)$

$$= \frac{(1-xs)(x+x^2)s + (x^2+x^3+3x^4+x^5)s^2}{(1-x-x^2-3x^3-x^4)(1-xs-(xs)^2)}$$
$$= \frac{xs(x^4s+3x^3s+x+1)}{(1-x-x^2-3x^3-x^4)(1-xs-(xs)^2)}.$$

We handle $\mathcal{H}(x,s)$ similarly, noting that by Corollary 5.11, we also have that

$$\mathcal{H}^{i}(x) = x\mathcal{H}^{i-1}(x) + x^{2}\mathcal{H}^{i-2}(x)$$

for $i \geq 3$. We follow the above computation until

$$\mathcal{H}(x,s) = \frac{(1-xs)\mathcal{H}^{1}(x)s + \mathcal{H}^{2}(x)s^{2}}{(1-xs - (xs)^{2})}$$

where we insert the initial conditions for $\mathcal{H}^{i}(x)$ to obtain

$$\mathcal{H}(x,s) = \frac{(1-xs)(x+2x^2+x^3)s+(2x^3+x^4+3x^5+x^6)s^2}{(1-x-x^2-3x^3-x^4)(1-xs-(xs)^2)} \\ = \frac{xs(x^5s+3x^4s-xs+x^2+2x+1)}{(1-x-x^2-3x^3-x^4)(1-xs-(xs)^2)}.$$

We remark here that the sequence $\{h_r^1\}$ coincides with [16, <u>A196423</u>]. This sequence also counts the cardinality of the following family of sets. Let X_r denote the number of length r sequences x_1, x_2, \ldots, x_r where $x_i \in \{0, 1, 2\}$ and $x_i = |\{x_j : |j - i| = 1 \text{ and } x_j < x_i\}|$. For example, $X_3 = \{000, 001, 100, 101, 020\}$. We show there is a natural relationship between X_r and $\mathcal{A}_r(\tilde{\alpha}, -\tilde{\alpha})$.

Theorem 5.13. There is a bijection between X_r and $\mathcal{A}_r(\tilde{\alpha}, -\tilde{\alpha})$.

Proof. Observe that, an element of X_r can be decomposed into the following (consecutive) subsequences whose neighbors are all 0's

 $\tau_1 := 1, \quad \tau_2 := 2, \quad \tau_3 := 11, \quad \tau_4 := 12, \quad \tau_5 := 21, \quad \text{and} \quad \tau_6 := 121.$

Notice that τ_1 can only appear at the beginning or end of an element of X_r , and no other subsequence τ_i can appear at the beginning or end of the sequence in X_r . Given $x \in X_r$, define $\psi(x) \in A_r$ in the following way. We begin by setting $\psi(x) = id$. Now, we check which subsequences τ_i are contained in x and at what indices they appear.

- (1) If x has a subsequence τ_1 in position i, then we multiply $\psi(x)$ by s_i . Necessarily, i = 1 or i = r as discussed.
- (2) If x has a subsequence τ_2 in position *i*, then we multiply $\psi(x)$ by s_i .
- (3) If x has a subsequence τ_3 (resp. τ_4, τ_5) in positions i, i + 1, we multiply $\psi(x)$ by $s_i s_{i+1}$ (resp $s_{i+1} s_i$, $s_i s_{i+1}, s_i$).
- (4) If x has a subsequence τ_6 in positions i, i+1, i+2, we multiply $\psi(x)$ by $s_i s_{i+2} s_{i+1}$.

Comparing our method of decomposing an element of X_r with Theorem 4.7 shows that $\psi(x) \in \mathcal{A}_r(\tilde{\alpha}, -\tilde{\alpha})$. Moreover, an inverse to ψ can be constructed in a similar manner, verifying that this map is a bijection. \Box

We conclude this section by providing a generating function that records the cardinalities of all Weyl alternation sets in type A for negative roots.

Theorem 5.14. If $\tilde{\alpha}$ is the highest root and $\mu = -\alpha_{i,j}$ with $1 \leq i \leq j \leq r$ is a negative root of the Lie algebra of type A_r , then we have the generating function

$$\sum_{1 \le i \le j \le r} |\mathcal{A}_r(\tilde{\alpha}, -\alpha_{i,j})| x^r s^i t^j = \frac{1}{t(1 - x - x^2)} \left((1 - x)t\mathcal{H}(xt, s) + \mathcal{P}(xt, s) - \frac{xst}{1 - xst - (xst)^2} \right)$$

where $\mathcal{H}(xt,s)$ and $\mathcal{P}(xt,s)$ are as in Lemma 5.12.

Proof. For a pair fixed $1 \le i \le j$, from Proposition 5.5 and Lemma 5.6, we have

$$\sum_{r \ge j} |\mathcal{A}_r(\tilde{\alpha}, -\alpha_{i,j})| x^r = \frac{h_j^i x^j + (|\mathcal{A}_{j+1}(\tilde{\alpha}, -\alpha_{i,j})| - h_j^i) x^{j+1}}{1 - x - x^2}$$
$$= \frac{h_j^i x^j + (p_{j+1}^i - h_j^i) x^{j+1}}{1 - x - x^2},$$

where, by Lemma 5.10, we can identify $\mathcal{A}_{j+1}(\tilde{\alpha}, -\alpha_{i,j})$ with P_{j+1}^i .

Now, if we fix $i \ge 1$ and sum over all j, we have

$$\sum_{j\geq i} \left(\frac{h_j^i x^j + (p_{j+1}^i - h_j^i) x^{j+1}}{1 - x - x^2}\right) t^j = \frac{1}{1 - x - x^2} \left(\sum_{j\geq i} h_j^i x^j t^j + \sum_{j\geq i} p_{j+1}^i x^{j+1} t^j - \sum_{j\geq i} h_j^i x^{j+1} t^j\right)$$
$$= \frac{1}{1 - x - x^2} \left((1 - x)\mathcal{H}^i(xt) + \frac{\mathcal{P}^i(xt) - p_i^i(xt)^i}{t}\right).$$

Finally, we must take the resulting expression and sum over all $i \ge 1$, yielding

$$\sum_{1 \le i \le j \le r} |\mathcal{A}_r(\tilde{\alpha}, -\alpha_{i,j})| x^r s^i t^j = \sum_{i \ge 1} \frac{1}{1 - x - x^2} \left((1 - x)\mathcal{H}^i(xt) + \frac{\mathcal{P}^i(xt) - p_i^i(xt)^i}{t} \right) s^i$$
$$= \frac{1}{t(1 - x - x^2)} \left((1 - x)t \sum_{i \ge 1} \mathcal{H}^i(xt) s^i + \sum_{i \ge 1} \mathcal{P}^i(xt) s^i - \sum_{i \ge 1} p_i^i(xst)^i \right).$$

We understand the first two summations from Lemma 5.12. We can compute $\sum_{i\geq 1} p_i^i(xt)^i s^i$ by combining Lemma 5.6 and Lemma 5.8. From Lemma 5.12, we have the starting values $p_1^1 = p_2^2 = 1$. Therefore, we can conclude

$$\sum_{1 \le i \le j \le r} |\mathcal{A}_r(\tilde{\alpha}, -\alpha_{i,j})| x^r s^i t^j = \frac{1}{t(1 - x - x^2)} \left((1 - x)t\mathcal{H}(xt, s) + \mathcal{P}(xt, s) - \frac{p_1^1 x s t + (p_2^2 - p_1^1)(xst)^2}{1 - x s t - (xst)^2} \right)$$
$$= \frac{1}{t(1 - x - x^2)} \left((1 - x)t\mathcal{H}(xt, s) + \mathcal{P}(xt, s) - \frac{x s t}{1 - x s t - (xst)^2} \right). \square$$

6. FUTURE WORK

Lusztig introduced the q-analog of Kostant's partition function as the polynomial-valued function $\wp_q(\xi) = c_0 + c_1q + c_2q^2 + \cdots + c_kq^k$, where c_i equals the number of ways to write ξ as a sum of exactly *i* positive roots [18, Proposition 9.2]. Then the q-analog of Kostant's weight multiplicity formula is given by

$$m_q(\lambda,\mu) = \sum_{\sigma \in W} (-1)^{\ell(\sigma)} \wp_q(\sigma(\lambda+\rho) - \mu - \rho).$$

In type A_r , using the Weyl alternation set $\mathcal{A}(\tilde{\alpha}, 0)$, Harris gives a completely combinatorial proof that $m_q(\tilde{\alpha}, 0) = \sum_{i=1}^r q^i$ [5, Proposition 3.3]. Again in type A_r , using the Weyl alternation set $\mathcal{A}(\tilde{\alpha}, \mu)$ with $\mu \in \Phi^+$, Harry shows that $m_q(\tilde{\alpha}, \mu)$ is a power of q [14, Theorem 4.1]. Harry conjectured the following.

Conjecture 6.1. If $\tilde{\alpha}$ is the highest root and $\mu = -\alpha_{i,j}$ with $1 \leq i \leq j \leq r$ is a negative root of the Lie algebra of type A_r , then

$$m_q(\tilde{\alpha},\mu) = q^{r+j-i+1} + q^{r+j-i} - q^{j-i+1}.$$

At this moment, we have a proof of Conjecture 6.1 for $\mu = -\alpha_i$ with $1 \le i \le r$. Given our characterization of the Weyl alternation sets $\mathcal{A}(\tilde{\alpha},\mu)$ for negative roots $\mu \in \Phi^-$, in future work, we aim to use similar techniques to give a full proof of Conjecture 6.1.

Acknowledgments

This material is based on work supported by the National Science Foundation under Grant Number DMS 1916439 while the authors were participating in the Mathematics Research Communities (MRC) 2024 Summer Conference at Beaver Hollow Conference Center in Java Center, New York. We extend our thanks to the American Mathematical Society for their support. E. Banaian was partially supported by Research Project 2 from the Independent Research Fund Denmark (grant no. 1026-00050B). This work was supported by a grant from the Simons Foundation (Travel Support for Mathematicians, P. E. Harris).

References

- A. Björner and F. Brenti. Combinatorics of Coxeter groups, volume 231 of Graduate Texts in Mathematics. Springer, New York, 2005.
- [2] K. Chang, P. E. Harris, and E. Insko. Kostant's weight multiplicity formula and the Fibonacci and Lucas numbers. J. Comb., 11(1):141–167, 2020.
- [3] C. Cochet. Vector partition function and representation theory. Formal Power Series and Algebraic Combinatorics Proceedings. Taormina, Italy, 1009–1020, 2005.
- [4] R. E. Garcia, P. E. Harris, M. Loving, L. Martinez, D. Melendez, J. Rennie, G. Rojas Kirby, and D. Tinoco. On Kostant's weight q-multiplicity formula for sl₄(ℂ). Appl. Algebra Engrg. Comm. Comput., 33(4):353–418, 2022.
- [5] P. E. Harris. On the adjoint representation of \mathfrak{sl}_n and the Fibonacci numbers. C. R. Math. Acad. Sci. Paris, 349(17-18):935–937, 2011.
- [6] P. E. Harris. Computing weight multiplicities. In A primer for undergraduate research, Found. Undergrad. Res. Math., pages 193–222. Birkhäuser/Springer, Cham, 2017.
- [7] P. E. Harris, P. Hollander, D. C. Qin, and M. Rodriguez-Hertz. On Kostant's weight q-multiplicity formula for sp₆(C). Appl. Algebra Engrg. Comm. Comput., 35(2):253–289, 2024.
- [8] P. E. Harris, E. Insko, and M. Omar. The q-analog of Kostant's partition function and the highest root of the simple Lie algebras. Australas. J. Combin., 71:68–91, 2018.
- [9] P. E. Harris, E. Insko, and A. Simpson. Computing weight q-multiplicities for the representations of the simple Lie algebras. Appl. Algebra Engrg. Comm. Comput., 29(4):351–362, 2018.
- [10] P. E. Harris, E. Insko, and L. K. Williams. The adjoint representation of a classical Lie algebra and the support of Kostant's weight multiplicity formula. J. Comb., 7(1):75–116, 2016.
- [11] P. E. Harris and E. L. Lauber. Weight q-multiplicities for representations of $\mathfrak{sp}_4(\mathbb{C})$. Zh. Sib. Fed. Univ. Mat. Fiz., 10(4):494–502, 2017.
- [12] P. E. Harris, M. Loving, J. Ramirez, J. Rennie, G. Rojas Kirby, E. Torres Davila, and F. O. Ulysse. Visualizing the support of Kostant's weight multiplicity formula for the rank-2 Lie algebras. *Involve*, 17(2):183–215, 2024.
- [13] P. E. Harris, M. Rahmoeller, L. Schneider, and A. Simpson. When is the q-multiplicity of a weight a power of q? Electron. J. Combin., 26(4):Paper No. 4.17, 55, 2019.
- [14] K. J. Harry. Computing the q-multiplicity of the positive roots of $\mathfrak{sl}_{r+1}(\mathbb{C})$ and products of Fibonacci numbers, 2024.
- [15] J. E. Humphreys. Reflection groups and Coxeter groups, volume 29 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 1990.
- [16] OEIS Foundation Inc., The On-Line Encyclopedia of Integer Sequences, 2023. Published electronically at https://oeis.org.
- [17] B. Kostant. A formula for the multiplicity of a weight. Proc. Nat. Acad. Sci. U.S.A., 44:588–589, 1958.

- [18] G. Lusztig. Singularities, character formulas, and a q-analog of weight multiplicities, volume 101-102 of Astérisque. Soc. Math. France, Paris, 1983.
- [19] R. P. Stanley. Enumerative combinatorics. Volume 1, volume 49 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, second edition, 2012.

(P. X. Anderson) DEPARTMENT OF MATHEMATICS, CORNELL UNIVERSITY, ITHACA, NY 14853 *Email address*: pxa2@cornell.edu

(E. Banaian) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, RIVERSIDE, RIVERSIDE, CALIFORNIA, 92521 *Email address:* estherbanaian@gmail.com

(M. J. Ferreri) DEPARTMENT OF MATHEMATICS, DARTMOUTH COLLEGE, HANOVER, NH, 03755 *Email address:* melanie.ferreri.gr@dartmouth.edu

(O. C. Goff) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN-MADISON, MADISON, WI 5306 *Email address:* ogoff@wisc.edu

(K. P. Hadaway) DEPARTMENT OF MATHEMATICS, IOWA STATE UNIVERSITY, AMES, IA, 50010 *Email address:* nmayers@ncsu.edu

(P. E. Harris, K. J. Harry) Department of Mathematical Sciences, University of Wisconsin-Milwaukee, Milwaukee, WI 53211

Email address: peharris@uwm.edu kjharry@uwm.edu

(N. Mayers) DEPARTMENT OF MATHEMATICS, NORTH CAROLINA STATE UNIVERSITY, RALEIGH, NC, 27605 *Email address:* nmayers@ncsu.edu

(S. Wang) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MINNESOTA TWIN CITIES, MN, 55455 *Email address:* wang8406@umn.edu

(A. N. Wilson) DEPARTMENT OF MATHEMATICS, OBERLIN COLLEGE, OBERLIN, OH 44074 *Email address:* awilson6@oberlin.edu