

NATURALITY OF SL_n QUANTUM TRACE MAPS FOR SURFACES

HYUN KYU KIM AND ZHIHAO WANG

ABSTRACT. The SL_n -skein algebra of a punctured surface \mathfrak{S} , studied by Sikora, is an algebra generated by isotopy classes of n -webs living in the thickened surface $\mathfrak{S} \times (-1, 1)$, where an n -web is a union of framed links and framed oriented n -valent graphs satisfying certain conditions. For each ideal triangulation λ of \mathfrak{S} , Lê and Yu constructed an algebra homomorphism, called the SL_n -quantum trace, from the SL_n -skein algebra of \mathfrak{S} to a so-called balanced subalgebra of the n -root version of Fock and Goncharov's quantum torus algebra associated to λ . We show that the SL_n -quantum trace maps for different ideal triangulations are related to each other via a balanced n -th root version of the quantum coordinate change isomorphism, which extends Fock and Goncharov's isomorphism for quantum cluster varieties. We avoid heavy computations in the proof, by using the splitting homomorphisms of Lê and Sikora, and a network dual to the n -triangulation of λ studied by Schrader and Shapiro.

CONTENTS

1. Introduction	2
1.1. Overview	2
1.2. Main ideas	5
2. Quantum trace maps for stated SL_n -skein algebras	7
2.1. Reduced stated SL_n -skein algebras	7
2.2. The splitting homomorphism	10
2.3. Quantum trace maps	10
3. Quantum cluster algebras	13
3.1. Classical cluster \mathcal{X} -mutations	14
3.2. Quantum mutations for Fock-Goncharov algebras	14
3.3. Quantum mutations for mutable-balanced n -th root Fock-Goncharov algebras	16
3.4. Coordinate change isomorphisms for change of triangulations	19
4. Compatibility between SL_n -quantum trace maps and coordinate change isomorphisms	26
4.1. Compatibility of the quantum trace map with the balanced quantum coordinate change maps $\Theta_{\lambda\lambda'}^\omega$	26
4.2. Proof of compatibility for the case of quadrilateral \mathbb{P}_4	28
4.3. The restriction of $\Theta_{\lambda\lambda'}^\omega$ to the balanced part	36
4.4. A comparison with coordinate change isomorphisms defined by Lê and Yu	39
References	40

1. INTRODUCTION

1.1. **Overview.** Let \mathfrak{S} be a *punctured surface*, that is, a closed oriented surface minus a non-empty finite set of points called punctures such that every component of \mathfrak{S} contains at least one puncture. For an algebraic group G , one considers the moduli space of G -local systems on \mathfrak{S} , which is identified with the space

$$\mathcal{L}_{G,\mathfrak{S}} = \text{Hom}(\pi_1(\mathfrak{S}), G)/G,$$

which is the quotient of the space $\text{Hom}(\pi_1(\mathfrak{S}), G)$ of group homomorphisms $\pi_1(\mathfrak{S}) \rightarrow G$ by the action of G given by conjugation by an element of G . Here we regard this space as a stack, with the algebra of functions $\mathcal{O}(\mathcal{L}_{G,\mathfrak{S}})$ being the algebra of G -invariant functions $\mathcal{O}(\text{Hom}(\pi_1(\mathfrak{S}), G))^G$; one might call this space the G -character stack for \mathfrak{S} . This space is equipped with a natural symplectic structure per each choice of an invariant bilinear form on G [G84], hence the problem of quantization along this symplectic structure arises. Here we focus on the algebraic aspect of quantization, for which one would like to deform the algebra of regular functions $\mathcal{O}(\mathcal{L}_{G,\mathfrak{S}})$ by a one-parameter family of non-commutative algebras, in an appropriate sense. When $G = \text{SL}_n$, $n \geq 2$, it is well-known that $\mathcal{O}(\mathcal{L}_{\text{SL}_n,\mathfrak{S}})$ is generated by the *trace-of-monodromy* functions f_γ associated to isotopy classes of oriented loops γ in \mathfrak{S} ; the value of f_γ at the equivalence class of a group homomorphism $\rho : \pi_1(\mathfrak{S}) \rightarrow \text{SL}_n$ is $\text{trace}(\rho([\gamma]))$, where $[\gamma]$ is the element of $\pi_1(\mathfrak{S})$ represented by γ . When $n \geq 3$, a complete set of algebraic relations among f_γ is rather complicated, and becomes much simpler when one also takes into consideration more functions f_W , each of which is associated to a combinatorial object W called a *web* or an SL_n -*web*; we call it an n -*web*. Here we consider a model developed by Sikora [S01], in which an n -web is a union of oriented loops and oriented graphs in \mathfrak{S} , where each vertex of the graph is required to be either a source or a sink. See [CKM14] for another model of Cautis, Kamnitzer and Morrison, based on 3-valent graphs with edges labeled by fundamental representations of SL_n , and [P22] for the equivalence of the two models.

A quantum deformed algebra for $\mathcal{O}(\mathcal{L}_{\text{SL}_n,\mathfrak{S}})$ is constructed with the help of 3-dimensional quantum topology. The SL_n -*skein algebra* $\mathcal{S}_\omega(\mathfrak{S}) = \mathcal{S}_\omega^{\text{SL}_n}(\mathfrak{S})$ is an associative algebra generated by the isotopy classes of framed n -webs living in the thickened surface $\mathfrak{S} \times (-1, 1)$, modulo the *skein relations* (7)–(10), where the product of two framed n -webs is given by superposition, i.e. stacking one on top of the other; see §2.1. Here ω appearing in the relations is a formal quantum parameter, which is related to another quantum parameter q by

$$q = \omega^{n^2}.$$

There are various versions of this algebra; the case when $n = 2$ is first studied by Turaev [T89] and Przytycki [P99], and for general n by Sikora [S05] and Cautis-Kamnitzer-Morrison [CKM14], building on the work of Kuperberg [K96] on invariants of tensor product representations of Lie algebras and quantum groups.

When $n = 2$, the SL_2 -character stack $\mathcal{L}_{\text{SL}_2,\mathfrak{S}}$ is closely related to the Teichmüller space of \mathfrak{S} , which is the space of isotopy classes of complex structures (or complete hyperbolic metrics) on \mathfrak{S} . Quantization of various versions of Teichmüller spaces has

been studied since late 1980's, where the first fundamental results were obtained by Kashaev [K98] and by Chekhov and Fock [F97, CF99]; the latter was generalized by Fock and Goncharov [FG09a, FG09b] to quantization of cluster Poisson varieties, which include ordinary and higher Teichmüller spaces of \mathfrak{S} [FG06]. To say briefly, Fock and Goncharov studied a certain stack $\mathcal{X}_{\mathrm{PGL}_n, \mathfrak{S}}$ that parametrizes PGL_n -local systems on \mathfrak{S} together with certain extra discrete data at punctures [FG06]. Per each choice of an *ideal triangulation* λ of \mathfrak{S} , which is a collection of unoriented paths in \mathfrak{S} running between punctures that divide \mathfrak{S} into (ideal) triangles, considered up to isotopy, Fock and Goncharov constructed in [FG06] a special coordinate system on $\mathcal{X}_{\mathrm{PGL}_n, \mathfrak{S}}$. The coordinates are parametrized by the vertices of a special quiver Γ_λ obtained by gluing the ' n -triangulation' of each ideal triangle of λ (see §2.3 and Figure 2). It is also shown that $\mathcal{X}_{\mathrm{PGL}_n, \mathfrak{S}}$ possesses a natural Poisson structure whose Poisson brackets among these special coordinates are encoded by the signed adjacency matrix Q_λ of the quiver Γ_λ which counts the number of arrows between vertices, and that the transition map between the special charts for different ideal triangulations λ and λ' are given by a specific composition of the formulas called the *cluster mutations*. These charts, called the *cluster charts*, together with the transition maps, are quantized by certain non-commutative algebras and maps between them. To each cluster chart for λ is associated the quantum torus algebra $\mathcal{X}_q(\mathfrak{S}, \lambda) = \mathcal{X}_q^{\mathrm{PGL}_n}(\mathfrak{S}, \lambda)$ constructed from Q_λ , which is a free associative algebra generated by the symbols X_v and their inverses X_v^{-1} , with v being the vertices of the quiver Γ_λ , modulo the relations $X_v X_w = q^{2Q_\lambda(v,w)} X_w X_v$. Per each pair of triangulations λ and λ' , is constructed an isomorphism [FG09a, FG09b]

$$\Phi_{\lambda\lambda'}^q : \mathrm{Frac}(\mathcal{X}_q(\mathfrak{S}, \lambda')) \rightarrow \mathrm{Frac}(\mathcal{X}_q(\mathfrak{S}, \lambda)) \quad (1)$$

between the skew-fields of fractions of the quantum torus algebras, as a composition of a special sequence of quantum cluster mutations [BZ05]; this sequence consists of $\frac{1}{6}(n^3 - n)$ mutations [FG06]. These quantum coordinate change isomorphisms recover the classical transition maps as $q \rightarrow 1$, and they satisfy the consistency relation $\Phi_{\lambda\lambda'}^q \Phi_{\lambda'\lambda''}^q = \Phi_{\lambda\lambda''}^q$. One can interpret this as having a consistent system of quantum deformed algebras for the Poisson space $\mathcal{X}_{\mathrm{PGL}_n, \mathfrak{S}}$.

For $n = 2$, the SL_2 -skein algebra $\mathcal{S}_\omega^{\mathrm{SL}_2}(\mathfrak{S})$ which quantizes the space $\mathcal{L}_{\mathrm{SL}_2, \mathfrak{S}}$ and the above system of quantum torus algebras $\mathcal{X}_q^{\mathrm{PGL}_2}(\mathfrak{S}, \lambda)$ which quantize the space $\mathcal{X}_{\mathrm{PGL}_2, \mathfrak{S}}$ were expected to be closely related to each other. It was Bonahon and Wong [BW11] who proved this expectation. One important subtlety is that one needs to consider a square-root version of the quantum torus algebra, $\mathcal{Z}_\omega^{\mathrm{PGL}_2}(\mathfrak{S}, \lambda)$, generated by $Z_v^{\pm 1}$ for vertices v of Γ_λ , with relations $Z_v Z_w = \omega^{2Q_\lambda(v,w)} Z_w Z_v$, in which $\mathcal{X}_q^{\mathrm{PGL}_2}(\mathfrak{S}, \lambda)$ is embedded as $X_v = Z_v^2$ and $q = \omega^4$. Note that, for $n = 2$, the vertices of Γ_λ is in bijection with the edges of the ideal triangulation λ . A Laurent monomial $\prod_{v \in \lambda} Z_v^{a_v} \in \mathcal{Z}_\omega^{\mathrm{PGL}_2}(\mathfrak{S}, \lambda)$, for $a_v \in \mathbb{Z}$, where the product is taken with respect to some chosen order, is said to be SL_2 -balanced if for each ideal triangle of λ , the sum of a_v for the edges v forming this triangle is even. Let $\mathcal{Z}_\omega^{\mathrm{bl}; \mathrm{PGL}_2}(\mathfrak{S}, \lambda)$ be the subalgebra of $\mathcal{Z}_\omega^{\mathrm{PGL}_2}(\mathfrak{S}, \lambda)$ generated by the SL_2 -balanced Laurent monomials; it is easy to see that this balanced square-root subalgebra $\mathcal{Z}_\omega^{\mathrm{bl}; \mathrm{PGL}_2}(\mathfrak{S}, \lambda)$ contains $\mathcal{X}_q^{\mathrm{PGL}_2}(\mathfrak{S}, \lambda)$. Hiatt [H10,

BW11, KLS23] constructed an isomorphism

$$\Theta_{\lambda\lambda'}^{\omega;\text{PGL}_2} : \text{Frac}(\mathcal{Z}_\omega^{\text{bl};\text{PGL}_2}(\mathfrak{S}, \lambda')) \rightarrow \text{Frac}(\mathcal{Z}_\omega^{\text{bl};\text{PGL}_2}(\mathfrak{S}, \lambda))$$

between the skew-fields of fractions of the SL_2 -balanced square-root quantum torus algebras, that extends the previously known $\Phi_{\lambda\lambda'}^q$ which is a composition of quantum cluster mutations, and that satisfies the consistency relation $\Theta_{\lambda\lambda'}^{\omega;\text{PGL}_2} \Theta_{\lambda'\lambda''}^{\omega;\text{PGL}_2} = \Theta_{\lambda\lambda''}^{\omega;\text{PGL}_2}$. Bonahon and Wong [BW11] constructed an algebra homomorphism

$$\text{tr}_\lambda^{\text{SL}_2} : \mathcal{S}_\omega^{\text{SL}_2}(\mathfrak{S}) \rightarrow \mathcal{Z}_\omega^{\text{bl};\text{PGL}_2}(\mathfrak{S}, \lambda) \quad (2)$$

called the SL_2 -quantum trace, per each ideal triangulation λ of \mathfrak{S} , that is compatible with the balanced square-root quantum coordinate change maps $\Theta_{\lambda\lambda'}^{\omega;\text{PGL}_2}$, i.e.

$$\text{tr}_\lambda^{\text{SL}_2} = \Theta_{\lambda\lambda'}^{\omega;\text{PGL}_2} \circ \text{tr}_{\lambda'}^{\text{SL}_2}; \quad (3)$$

this compatibility can be interpreted as the *naturality* under change of ideal triangulations. The Bonahon-Wong SL_2 -quantum trace has been widely used ever since, in several important applications including the construction of Fock-Goncharov quantum duality map [AK17], and the study of representations of SL_2 -skein algebras [BW16].

One major direction of research after Bonahon and Wong's work has been on the SL_n -generalization, i.e. on the SL_n -quantum trace maps

$$\text{tr}_\lambda^{\text{SL}_n} : \mathcal{S}_\omega^{\text{SL}_n}(\mathfrak{S}) \rightarrow \mathcal{Z}_\omega^{\text{bl};\text{PGL}_n}(\mathfrak{S}, \lambda) \subset \mathcal{Z}_\omega^{\text{PGL}_n}(\mathfrak{S}, \lambda). \quad (4)$$

Here $\mathcal{Z}_\omega^{\text{PGL}_n}(\mathfrak{S}, \lambda)$ is the n -th root version of the quantum torus algebra $\mathcal{X}_q^{\text{PGL}_n}(\mathfrak{S}, \lambda)$, generated by $Z_v^{\pm 1}$ for vertices v of the n -triangulation quiver Γ_λ for an ideal triangulation λ , modulo the relation $Z_v Z_w = \omega^{2Q_\lambda(v,w)} Z_w Z_v$. The usual Fock-Goncharov quantum (cluster) torus algebra $\mathcal{X}_q^{\text{PGL}_n}(\mathfrak{S}, \lambda)$ embeds into it as $X_v = Z_v^n$, $q = \omega^{n^2}$. The SL_n -balanced subalgebra $\mathcal{Z}_\omega^{\text{bl};\text{PGL}_n}(\mathfrak{S}, \lambda)$ of $\mathcal{Z}_\omega^{\text{PGL}_n}(\mathfrak{S}, \lambda)$ is generated by $\prod_v Z_v^{a_v}$ for $(a_v)_v \in \mathbb{Z}^{\text{vertices of } \Gamma_\lambda}$ that satisfies a suitable SL_n -analog of the SL_2 -balancedness condition, so that in particular we have $\mathcal{X}_q^{\text{PGL}_n}(\mathfrak{S}, \lambda) \subset \mathcal{Z}_\omega^{\text{bl};\text{PGL}_n}(\mathfrak{S}, \lambda)$. The first developments were for the case $n = 3$ studied by Douglas [D24] and the first author [K20], and later Lê and Yu constructed SL_n -quantum trace maps for all $n \geq 2$ [LY23].

One of the most important sought-for properties of $\text{tr}_\lambda^{\text{SL}_n}$ is the naturality under change of ideal triangulations. The naturality should fit into the framework of quantum cluster algebras, in the sense that $\text{tr}_\lambda^{\text{SL}_n}$ for different ideal triangulations must be compatible with each other via a balanced n -th root quantum coordinate change isomorphism

$$\Theta_{\lambda\lambda'}^{\omega;\text{SL}_n} : \text{Frac}(\mathcal{Z}_\omega^{\text{bl};\text{PGL}_n}(\mathfrak{S}, \lambda')) \rightarrow \text{Frac}(\mathcal{Z}_\omega^{\text{bl};\text{PGL}_n}(\mathfrak{S}, \lambda)) \quad (5)$$

that extends $\Phi_{\lambda\lambda'}^q$ in (1) which is a composition of $\frac{1}{6}(n^3 - n)$ quantum cluster mutations. So one should construct such a map $\Theta_{\lambda\lambda'}^{\omega;\text{SL}_n}$ and show the naturality

$$\text{tr}_\lambda^{\text{SL}_n} = \Theta_{\lambda\lambda'}^{\omega;\text{SL}_n} \circ \text{tr}_{\lambda'}^{\text{SL}_n}. \quad (6)$$

This task is accomplished for $n = 3$ by the first author in [K24]. For general n , Lê and Yu have a partial result in [LY23], where they construct a skew-field isomorphism

$$\overline{\Psi}_{\lambda\lambda'}^X : \mathrm{Frac}(\mathcal{Z}_\omega^{\mathrm{bl};\mathrm{PGL}_n}(\mathfrak{S}, \lambda')) \rightarrow \mathrm{Frac}(\mathcal{Z}_\omega^{\mathrm{bl};\mathrm{PGL}_n}(\mathfrak{S}, \lambda))$$

and show the naturality

$$\mathrm{tr}_\lambda^{\mathrm{SL}_n} = \overline{\Psi}_{\lambda\lambda'}^X \circ \mathrm{tr}_{\lambda'}^{\mathrm{SL}_n}.$$

However, Lê and Yu's isomorphism $\overline{\Psi}_{\lambda\lambda'}^X$ is essentially derived from the naturality equation, and its relationship with the quantum cluster isomorphism $\Phi_{\lambda\lambda'}^q$ is not studied, but only suggested [LY23, §14.4].

Here is the main result of the current paper.

Theorem 1.1 (main theorem). *Let \mathfrak{S} be a punctured surface admitting an ideal triangulation.*

- (1) (§3.4) *For any two ideal triangulations λ and λ' of \mathfrak{S} , there exists a balanced n -th root quantum coordinate change map $\Theta_{\lambda\lambda'}^{\omega;\mathrm{SL}_n}$ as in (5), which is a skew-field isomorphism that extends the map $\Phi_{\lambda\lambda'}^q$ in (1), which in turn is constructed as a composition of $\frac{1}{6}(n^3 - n)$ quantum cluster mutations.*
- (2) (Proposition 3.14) *The consistency equation $\Theta_{\lambda\lambda'}^{\omega;\mathrm{SL}_n} \Theta_{\lambda'\lambda''}^{\omega;\mathrm{SL}_n} = \Theta_{\lambda\lambda''}^{\omega;\mathrm{SL}_n}$ holds for any three ideal triangulations λ , λ' and λ'' of \mathfrak{S} .*
- (3) (Theorems 4.1 and 4.15) *The naturality equation $\mathrm{tr}_\lambda^{\mathrm{SL}_n} = \Theta_{\lambda\lambda'}^{\omega;\mathrm{SL}_n} \circ \mathrm{tr}_{\lambda'}^{\mathrm{SL}_n}$ in (6) holds.*
- (4) (Theorems 4.16) *The map $\Theta_{\lambda\lambda'}^{\omega;\mathrm{SL}_n}$ coincides with Lê-Yu's map $\overline{\Psi}_{\lambda\lambda'}^X$.*

Consequently, Lê and Yu's SL_n -quantum trace is now proved to fit well in the framework of quantum cluster algebras. We expect that this comprises an important step towards an attempt of using the Lê-Yu SL_n -quantum trace to approach the Fock-Goncharov duality conjectures for gauge groups SL_n and PGL_n , classical and quantum [FG06, FG09a]. See [AK17] for $n = 2$ and [K20] for $n = 3$.

1.2. Main ideas. A crucial strategy taken by Bonahon and Wong [BW11] in their study of SL_2 -quantum trace maps is to investigate the compatibility under cutting the surface \mathfrak{S} along an ideal arc, i.e. an edge of an ideal triangulation λ . This yields a more general kind of surface, called a *punctured bordered surface* or a *pb surface*, which is obtained from a compact oriented surface with possibly-empty boundary by removing a non-empty finite set of points called punctures, such that each boundary component of the resulting surface is diffeomorphic to an open interval. The *stated SL_2 -skein algebra* $\mathcal{S}_\omega^{\mathrm{SL}_2}(\mathfrak{S})$ (§2.1) is defined using isotopy classes of framed links and stated framed tangles, which are arcs ending at the boundary walls of $\mathfrak{S} \times (-1, 1)$, where the endpoints lying in each boundary wall are required to have distinct heights, i.e. distinct $(-1, 1)$ -coordinates, equipped with a *state* which assigns a value in $\{1, 2\}$ to each endpoint. We mod out by the skein relations and also by certain extra relations regarding the boundary walls, studied by Lê and collaborators [L18, CL22]. The Fock-Goncharov space $\mathcal{X}_{\mathrm{PGL}_2, \mathfrak{S}}$ has a counterpart space $\mathcal{P}_{\mathrm{PGL}_2, \mathfrak{S}}$ studied by Goncharov and Shen [GS19], equipped with cluster charts and corresponding quantum algebras, with

quantum coordinate change isomorphisms. The statements (2)–(3) still hold for pb surfaces admitting ideal triangulations. Moreover, the compatibility under the cutting of a surface also holds in the following sense. Suppose λ is a triangulation of a pb surface \mathfrak{S} , and let \mathfrak{S}_e be the pb surface obtained from \mathfrak{S} by cutting along an edge e of λ that does not lie in the boundary of \mathfrak{S} . Let λ_e be the induced ideal triangulation of \mathfrak{S}_e . The cutting process yields a natural algebra embedding $\mathcal{Z}_\omega(\mathfrak{S}, \lambda) \rightarrow \mathcal{Z}_\omega(\mathfrak{S}_e, \lambda_e)$, which restricts to the map between balanced subalgebras. The cutting also yields a natural map between the stated SL_2 -skein algebras $\mathcal{S}_\omega^{\mathrm{SL}_2}(\mathfrak{S}) \rightarrow \mathcal{S}_\omega^{\mathrm{SL}_2}(\mathfrak{S}_e)$, called the *splitting homomorphism* which makes the following diagram commute [L18, CL22]

$$\begin{array}{ccc} \mathcal{S}_\omega^{\mathrm{SL}_2}(\mathfrak{S}) & \longrightarrow & \mathcal{Z}_\omega^{\mathrm{bl};\mathrm{PGL}_2}(\mathfrak{S}, \lambda) \\ \downarrow & & \downarrow \\ \mathcal{S}_\omega^{\mathrm{SL}_2}(\mathfrak{S}_e) & \longrightarrow & \mathcal{Z}_\omega^{\mathrm{bl};\mathrm{PGL}_2}(\mathfrak{S}_e, \lambda_e), \end{array}$$

which more or less completely characterizes the SL_2 -quantum trace maps [L18].

We take this strategy using cutting as well. We use the *reduced stated SL_n -skein algebra* $\overline{\mathcal{S}}_\omega^{\mathrm{SL}_n}(\mathfrak{S})$ of Lê-Yu [LY23] and Lê-Sikora [LS21] (see §2.1 of the present paper), defined using framed oriented links and stated framed n -webs in $\mathfrak{S} \times (-1, 1)$ which are graphs having 1-valent vertices lying in boundary walls equipped with state values in $\{1, 2, \dots, n\}$ and sink or source n -valent vertices lying in the interior, with the skein relations (7)–(10) and boundary relations (11)–(14); here ‘reduced’ means that we quotient out a certain ideal. In [LY23, LS21] the splitting homomorphism of these algebras are studied, and the Lê-Yu SL_n -quantum trace is proved to be compatible with the splitting homomorphisms [LY23].

It is well known that any two ideal triangulations are connected by a sequence of elementary moves called *flips*, where a flip changes exactly one edge of an ideal triangulation. If λ and λ' is related by flip at an edge e , we focus on the two ideal triangles in λ adjacent to e , and those in λ' adjacent to e . Using the splitting homomorphisms and the corresponding compatibility, through cutting along all edges not equal to e , the problem of proving the sought-for naturality (6) boils down to proving it in the case when the surface \mathfrak{S} is a 4-gon, i.e. a closed disc with 4 punctures on the boundary, which admits exactly two ideal triangulations. However, as seen in [K24], proving (6) for 4-gon directly would still require a heavy computation, already for $n = 3$.

To bypass the difficult computation, we take advantage of another development. Lê and Yu showed in [LY23, Theorem 10.5] that the value under their SL_n -quantum trace $\mathrm{tr}_\lambda^{\mathrm{SL}_n}$ of a stated arc, which has only 1-valent vertices but no n -valent vertices, in the case when the surface \mathfrak{S} is a triangle, a 3-gon, can be expressed as a sum of Laurent monomials enumerated by paths in a certain directed network (a directed graph) given by dual of the n -triangulation. We note that this network is studied first by Schrader and Shapiro [SS17] (see also [CS23]), who also studied a certain compatibility of this sum under a single ‘mutation’ of the network, which in turn is directly related to the cluster mutation of the n -triangulation quiver Γ_λ of Fock and Goncharov (and Shen).

We extend Lê and Yu's result to 4-gon (Lemma 4.5), and use Schrader and Shapiro's compatibility repeatedly to show the sought-for naturality (6) (Proposition 4.2).

As a result, we show Theorem 1.1 for the reduced stated SL_n -skein algebras for all triangulable pb surfaces. This in particular recovers the results of [K24] by a more conceptual proof.

An interesting future research topic is on the generalization of this non-computational proof of the naturality for more general stated n -webs living in the thickened 4-gon, not just stated arcs. It will become an important step in studying the sought-for quantum Fock-Goncharov duality map for SL_n and PGL_n ; see [KS] for $n = 3$.

Acknowledgments. H.K. (the first author) has been supported by KIAS Individual Grant (MG047204) at Korea Institute for Advanced Study. Z.W. (the second author) is supported by the NTU research scholarship from the Nanyang Technological University (Singapore) and the PhD scholarship from the University of Groningen (The Netherlands). The authors thank Thang Lê for helpful discussions.

2. QUANTUM TRACE MAPS FOR STATED SL_n -SKEIN ALGEBRAS

Suppose that R is a commutative domain with an invertible element $\omega^{\frac{1}{2}}$. Set

$$q = \omega^{n^2}$$

with $q^{\frac{1}{2n^2}} = \omega^{\frac{1}{2}}$. Define the following constants:

$$c_i = (-q)^{n-i} q^{\frac{n-1}{2n}}, \quad \mathfrak{t} = (-1)^{n-1} q^{\frac{n^2-1}{n}}, \quad \mathfrak{a} = q^{\frac{n+1-2n^2}{4}}.$$

2.1. Reduced stated SL_n -skein algebras. A **punctured bordered surface** (or **pb surface** for simplicity) \mathfrak{S} is obtained from a compact oriented surface $\overline{\mathfrak{S}}$ with possibly-empty boundary by removing finitely many points, which are called punctures, such that every boundary component of \mathfrak{S} is diffeomorphic to an open interval.

For any positive integer k , we use \mathbb{P}_k to denote the pb surface obtained from the closed disk by removing k punctures from the boundary. We call \mathbb{P}_1 the **monogon**.

Consider the 3-manifold $\mathfrak{S} \times (-1, 1)$, the thickened surface. For a point $(x, t) \in \mathfrak{S} \times (-1, 1)$, the value t is called the **height** of this point. An **n -web** α in $\mathfrak{S} \times (-1, 1)$ is a disjoint union of oriented closed curves and a directed finite graph properly embedded into $\mathfrak{S} \times (-1, 1)$, satisfying the following requirements:

- (1) α only contains 1-valent or n -valent vertices. Each n -valent vertex is a source or a sink. The set of 1-valent vertices is denoted as $\partial\alpha$, which are called **endpoints** of α . For any boundary component c of \mathfrak{S} , we require that the points of $\partial\alpha \cap (c \times (-1, 1))$ have mutually distinct heights.
- (2) Every edge of the graph is an embedded oriented closed interval in $\mathfrak{S} \times (-1, 1)$.
- (3) α is equipped with a continuous transversal **framing**.
- (4) The set of half-edges at each n -valent vertex is equipped with a cyclic order.
- (5) $\partial\alpha$ is contained in $\partial\mathfrak{S} \times (-1, 1)$ and the framing at these endpoints is given by the positive direction of $(-1, 1)$.

We will consider n -webs up to (ambient) **isotopy** which are continuous deformations of n -webs in their class. The empty n -web, denoted by \emptyset , is also considered as an n -web, with the convention that \emptyset is only isotopic to itself.

A **state** for α is a map $s: \partial\alpha \rightarrow \{1, 2, \dots, n\}$. A **stated n -web** in $\mathfrak{S} \times (-1, 1)$ is an n -web equipped with a state.

We say the (stated) n -web α is in **vertical position** if

- (1) the framing at everywhere is given by the positive direction of $(-1, 1)$,
- (2) α is in general position with respect to the projection $\text{pr}: \mathfrak{S} \times (-1, 1) \rightarrow \mathfrak{S} \times \{0\}$,
- (3) at every n -valent vertex, the cyclic order of half-edges as the image of pr is given by the positive orientation of \mathfrak{S} (drawn counter-clockwise in pictures).

By regarding \mathfrak{S} as $\mathfrak{S} \times \{0\}$, there is a projection $\text{pr}: \mathfrak{S} \times (-1, 1) \rightarrow \mathfrak{S}$. For every (stated) n -web α , we can isotope α to be in vertical position. For each boundary component c of \mathfrak{S} , the heights of $\partial\alpha \cap (c \times (-1, 1))$ determine a linear order on $c \cap \text{pr}(\alpha)$. Then a **(stated) n -web diagram** of α is $\text{pr}(\alpha)$ equipped with the usual over/underpassing information at each double point (called a crossing) and a linear order on $c \cap \text{pr}(\alpha)$ for each boundary component c of \mathfrak{S} .

Let S_n denote the permutation group on the set $\{1, 2, \dots, n\}$.

The **stated SL_n -skein algebra** $\mathcal{S}_\omega(\mathfrak{S})$ of \mathfrak{S} is the quotient module of the R -module freely generated by the set of all isotopy classes of stated n -webs in $\mathfrak{S} \times (-1, 1)$ subject to the relations (7)-(14):

$$q^{\frac{1}{n}} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} - q^{-\frac{1}{n}} \begin{array}{c} \diagdown \diagup \\ \diagup \diagdown \end{array} = (q - q^{-1}) \begin{array}{c} \longrightarrow \\ \longrightarrow \end{array}, \quad (7)$$

$$\begin{array}{c} \longrightarrow \\ \circlearrowleft \end{array} = \mathfrak{t} \begin{array}{c} \longrightarrow \end{array}, \quad (8)$$

$$\begin{array}{c} \circlearrowright \end{array} = (-1)^{n-1} [n] \begin{array}{c} \longrightarrow \\ \longrightarrow \end{array}, \quad \text{where } [n] = \frac{q^n - q^{-n}}{q - q^{-1}} = q^{-n+1} + q^{-n+3} + \dots + q^{n-1}, \quad (9)$$

$$\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} = (-q)^{\frac{n(n-1)}{2}} \cdot \sum_{\sigma \in S_n} (-q^{\frac{1-n}{n}})^{\ell(\sigma)} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \begin{array}{c} \sigma_+ \end{array}, \quad (10)$$

where the ellipse enclosing σ_+ is the minimum crossing positive braid representing a permutation $\sigma \in S_n$ and $\ell(\sigma) = \#\{(i, j) \mid 1 \leq i < j \leq n, \sigma(i) > \sigma(j)\}$ is the length of $\sigma \in S_n$.

$$\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} = \mathfrak{a} \sum_{\sigma \in S_n} (-q)^{\ell(\sigma)} \begin{array}{c} \sigma(n) \\ \longrightarrow \\ \sigma(2) \\ \longrightarrow \\ \sigma(1) \\ \longrightarrow \end{array}, \quad (11)$$

$$\begin{array}{c} \text{shaded area} \\ \text{arc from } i \text{ to } j \\ \text{downward arrow} \end{array} = \delta_{\bar{j}, i} \mathbb{C}_i \begin{array}{c} \text{shaded area} \\ \text{vertical line} \end{array}, \quad (12)$$

$$\begin{array}{c} \text{shaded area} \\ \text{arc from } i \text{ to } j \\ \text{downward arrow} \end{array} = \sum_{i=1}^n (\mathbb{C}_{\bar{i}})^{-1} \begin{array}{c} \text{shaded area} \\ \text{arc from } i \text{ to } \bar{i} \\ \text{downward arrow} \end{array} \quad \text{where } \bar{i} = n + 1 - i, \quad (13)$$

$$\begin{array}{c} \text{shaded area} \\ \text{arc from } i \text{ to } j \\ \text{downward arrow} \end{array} = q^{-\frac{1}{n}} \left(\delta_{j < i} (q - q^{-1}) \begin{array}{c} \text{shaded area} \\ \text{arc from } i \text{ to } j \\ \text{downward arrow} \end{array} + q^{\delta_{i,j}} \begin{array}{c} \text{shaded area} \\ \text{arc from } j \text{ to } i \\ \text{downward arrow} \end{array} \right), \quad (14)$$

where $\delta_{j < i} = \begin{cases} 1 & j < i \\ 0 & \text{otherwise} \end{cases}$, $\delta_{i,j} = \begin{cases} 1 & i = j \\ 0 & \text{otherwise} \end{cases}$, and small white dots represent an arbitrary orientation of the edges (left-to-right or right-to-left), consistent for the entire equation. The black dot represents the opposite orientation. When a boundary edge of a shaded area is directed, the direction indicates the height order of the endpoints of the diagrams on that directed line, where going along the direction increases the height, and the involved endpoints are consecutive in the height order. The height order outside the drawn part can be arbitrary.

The algebra structure for $\mathcal{S}_\omega(\mathfrak{S})$ is given by stacking the stated n -webs, i.e. for any two stated n -webs $\alpha, \alpha' \subset \mathfrak{S} \times (-1, 1)$, the product $\alpha\alpha'$ is defined by stacking α above α' . That is, if $\alpha \subset \mathfrak{S} \times (0, 1)$ and $\alpha' \subset \mathfrak{S} \times (-1, 0)$, we have $\alpha\alpha' = \alpha \cup \alpha'$.

For a boundary puncture p of a pb surface \mathfrak{S} , the **corner arcs** $C(p)_{ij}$ and $\bar{C}(p)_{ij}$ are the stated arcs depicted as in Figure 1. For a boundary puncture p which is not on a monogon component of \mathfrak{S} , set

$$C_p = \{C(p)_{ij} \mid i < j\}, \quad \bar{C}_p = \{\bar{C}(p)_{ij} \mid i < j\}.$$

Each element of $C_p \cup \bar{C}_p$ is called a *bad arc* at p .




FIGURE 1. The left is $C(p)_{ij}$ and the right is $\bar{C}(p)_{ij}$.

For a pb surface \mathfrak{S} ,

$$\overline{\mathcal{S}_\omega(\mathfrak{S})} = \mathcal{S}_\omega(\mathfrak{S}) / I^{\text{bad}}$$

is called the **reduced stated SL_n -skein algebra**, defined in [LY23], where I^{bad} is the two-sided ideal of $\mathcal{S}_\omega(\mathfrak{S})$ generated by all bad arcs.

2.2. The splitting homomorphism. Let \mathfrak{S} be a pb surface. An embedding $e : (0, 1) \rightarrow \mathfrak{S}$ is called an **ideal arc** if both $\bar{e}(0)$ and $\bar{e}(1)$ are punctures, where $\bar{e} : [0, 1] \rightarrow \mathfrak{S}$ is the ‘closure’ of e ; we often consider an ideal arc e up to isotopy, and often identify e with its image. Let e be an ideal arc of a pb surface \mathfrak{S} such that it is contained in the interior of \mathfrak{S} . After cutting \mathfrak{S} along e , we get a new pb surface $\text{Cut}_e(\mathfrak{S})$, which has two copies e_1, e_2 for e such that $\mathfrak{S} = \text{Cut}_e(\mathfrak{S})/(e_1 = e_2)$. We use \mathbf{pr}_e to denote the projection from $\text{Cut}_e(\mathfrak{S})$ to \mathfrak{S} . Suppose that α is a stated n -web diagram in \mathfrak{S} , which is transverse to e . Let s be a map from $e \cap \alpha$ to $\{1, 2, \dots, n\}$, and let h be a linear order on $e \cap \alpha$. Then there is a lift diagram $\alpha(h, s)$ for a stated n -web in $\text{Cut}_e(\mathfrak{S})$. For $i = 1, 2$, the heights of the endpoints of $\alpha(h, s)$ on e_i are induced by h (via \mathbf{pr}_e), and the states of the endpoints of $\alpha(h, s)$ on e_i are induced by s (via \mathbf{pr}_e). Then the splitting homomorphism

$$\mathbb{S}_e : \mathcal{S}_\omega(\mathfrak{S}) \rightarrow \mathcal{S}_\omega(\text{Cut}_e(\mathfrak{S}))$$

is defined by

$$\mathbb{S}_e(\alpha) = \sum_{s: \alpha \cap e \rightarrow \{1, \dots, n\}} \alpha(h, s). \quad (15)$$

Furthermore \mathbb{S}_e is an R -algebra homomorphism [LS21].

We can observe that \mathbb{S}_e sends bad arcs to bad arcs. So it induces an algebra homomorphism from $\overline{\mathcal{S}}_\omega(\mathfrak{S})$ to $\overline{\mathcal{S}}_\omega(\text{Cut}_e(\mathfrak{S}))$, which is still denoted as \mathbb{S}_e . When there is no confusion we will omit the subscript e from \mathbb{S}_e .

2.3. Quantum trace maps. In this subsection we review the materials in [LY23] about the quantum trace map, which is a homomorphism from the stated SL_n -skein algebra of a ‘triangulable’ pb surface \mathfrak{S} to a quantum torus algebra associated to a special quiver built from any choice of an ‘(ideal) triangulation’ of \mathfrak{S} .

To begin with, consider barycentric coordinates for an ideal triangle \mathbb{P}_3 so that

$$\mathbb{P}_3 = \{(i, j, k) \in \mathbb{R}^3 \mid i, j, k \geq 0, i + j + k = n\} \setminus \{(0, 0, n), (0, n, 0), (n, 0, 0)\}, \quad (16)$$

where (i, j, k) (or ijk for simplicity) are the barycentric coordinates. Let $v_1 = (n, 0, 0)$, $v_2 = (0, n, 0)$, $v_3 = (0, 0, n)$. Let e_i denote the edge on $\partial\mathbb{P}_3$ whose endpoints are v_i and v_{i+1} . See Figure 2.




FIGURE 2. Barycentric coordinates ijk and a 4-triangulation with its quiver

The n -**triangulation** of \mathbb{P}_3 is the subdivision of \mathbb{P}_3 into n^2 small triangles using lines $i, j, k = \text{constant}$ integers. For the n -triangulation of \mathbb{P}_3 , the vertices and edges of all small triangles, except for v_1, v_2, v_3 and the small edges adjacent to them, form a quiver $\Gamma_{\mathbb{P}_3}$. An **arrow** is the direction of a small edge defined as follows. If a small edge e is in the boundary $\partial\mathbb{P}_3$ then e has the counterclockwise direction of $\partial\mathbb{P}_3$. If e is interior then its direction is the same with that of a boundary edge of \mathbb{P}_3 parallel to e . Assign weight $\frac{1}{2}$ to any boundary arrow and weight 1 to any interior arrow.

Let $V_{\mathbb{P}_3}$ be the set of all points with integer barycentric coordinates of \mathbb{P}_3 :

$$V_{\mathbb{P}_3} = \{ijk \in \mathbb{P}_3 \mid i, j, k \in \mathbb{Z}\}. \quad (17)$$

Elements of $V_{\mathbb{P}_3}$ are called **small vertices**, and small vertices on the boundary of \mathbb{P}_3 are called the **edge vertices**.

A pb surface \mathfrak{S} is **triangulable** if each component of \mathfrak{S} has at least one puncture, and no component of \mathfrak{S} is one the following cases:

- (1) the monogon \mathbb{P}_1 ,
- (2) the bigon \mathbb{P}_2 ,
- (3) once or twice punctured sphere.

A **triangulation** λ of \mathfrak{S} is a collection of disjoint ideal arcs in \mathfrak{S} with the following properties: (1) any two arcs in λ are not isotopic; (2) λ is maximal under condition (1); (3) every puncture is adjacent to at least two ideal arcs. Our definition of triangulation excludes the so-called self-folded triangles. We will call each ideal arc in λ an **edge** of λ . If an edge is isotopic to a boundary component of \mathfrak{S} , we call such an edge a **boundary edge**. We use \mathbb{F}_λ to denote the set of faces after we cut \mathfrak{S} along all ideal arcs in λ that are not boundary edges. It is well-known that any triangulable surface admits a triangulation.

Suppose that \mathfrak{S} is a triangulable pb surface with a triangulation λ . By cutting \mathfrak{S} along all edges not isotopic to a boundary edge, we have a disjoint union of ideal triangles. Each triangle is called a **face** of λ . Then

$$\mathfrak{S} = \left(\bigsqcup_{\tau \in \mathbb{F}_\lambda} \tau \right) / \sim, \quad (18)$$

where each face τ is regarded as a copy of \mathbb{P}_3 , and \sim is the identification of edges of the faces to recover λ . Each face τ is characterized by a **characteristic map** $f_\tau: \mathbb{P}_3 \rightarrow \mathfrak{S}$, which is a homeomorphism when we restrict f_τ to $\text{Int } \mathbb{P}_3$ or the interior of each edge of \mathbb{P}_3 .

An n -**triangulation** of λ is a collection of n -triangulations of the faces τ which are compatible with the gluing \sim , where compatibility means, for any edges b and b' glued via \sim , the vertices on b and b' are identified. Define

$$V_\lambda = \bigcup_{\tau \in \mathbb{F}_\lambda} V_\tau, \quad V_\tau = f_\tau(V_{\mathbb{P}_3}).$$

The images of the weighted quivers $\Gamma_{\mathbb{P}_3}$ by f_τ form a quiver Γ_λ on \mathfrak{S} obtained by gluing. For any two $v, v' \in V_\lambda$, define

$$a_\lambda(v, v') = \begin{cases} w & \text{if there is an arrow from } v \text{ to } v' \text{ of weight } w, \\ 0 & \text{if there is no arrow between } v \text{ and } v'. \end{cases}$$

Let $Q_\lambda: V_\lambda \times V_\lambda \rightarrow \mathbb{Z}$ be the signed adjacency matrix of the weighted quiver Γ_λ defined by

$$Q_\lambda(v, v') = a_\lambda(v, v') - a_\lambda(v', v). \quad (19)$$

What matters about the quiver Γ_λ is only the signed adjacency matrix Q_λ , so we may tidy up the quiver by canceling or adding the weighted arrows in Γ_λ while maintaining Q_λ , so that between each pair of vertices there is only one arrow (with some weight).

We use $Q_{\mathbb{P}_3}$ to denote Q_λ when $\mathfrak{S} = \mathbb{P}_3$.

The **Fock-Goncharov algebra** is the quantum torus algebra associated to Q_λ , i.e.:

$$\mathcal{X}_q(\mathfrak{S}, \lambda) = \mathbb{T}_q(Q_\lambda) = R\langle X_v^{\pm 1}, v \in V_\lambda \rangle / (X_v X_{v'} = q^{2Q_\lambda(v, v')} X_{v'} X_v \text{ for } v, v' \in V_\lambda). \quad (20)$$

The n -**th root Fock-Goncharov algebra** is the quantum torus algebra defined as follows:

$$\mathcal{Z}_\omega(\mathfrak{S}, \lambda) = \mathbb{T}_\omega(Q_\lambda) = R\langle Z_v^{\pm 1}, v \in V_\lambda \rangle / (Z_v Z_{v'} = \omega^{2Q_\lambda(v, v')} Z_{v'} Z_v \text{ for } v, v' \in V_\lambda). \quad (21)$$

Recall that $q = \omega^{n^2}$. There is an algebra embedding from $\mathcal{X}_q(\mathfrak{S}, \lambda)$ to $\mathcal{Z}_\omega(\mathfrak{S}, \lambda)$ given by

$$X_v \mapsto Z_v^n$$

for $v \in V_\lambda$. We will regard $\mathcal{X}_q(\mathfrak{S}, \lambda)$ as a subalgebra of $\mathcal{Z}_\omega(\mathfrak{S}, \lambda)$. For any integer $k \in \mathbb{Z}$ and $v \in V_\lambda$, we will regard the symbol $(X_v)^{\frac{k}{n}}$ as referring to $(Z_v)^k$.

Let $\mathbf{k}_i: V_{\mathbb{P}_3} \rightarrow \mathbb{Z}$ ($i = 1, 2, 3$) be the functions defined by

$$\mathbf{k}_1(ijk) = i, \quad \mathbf{k}_2(ijk) = j, \quad \mathbf{k}_3(ijk) = k. \quad (22)$$

Let $\mathcal{B}_{\mathbb{P}_3} \subset \mathbb{Z}^{V_{\mathbb{P}_3}}$ be the subgroup generated by $\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3$ and $(n\mathbb{Z})^{V_{\mathbb{P}_3}}$. Elements in $\mathcal{B}_{\mathbb{P}_3}$ are called **balanced**.

A vector $\mathbf{k} \in \mathbb{Z}^{V_\lambda}$ is **balanced** if its pullback $\mathbf{k}_\tau := f_\tau^* \mathbf{k}$ to \mathbb{P}_3 is balanced for every face of λ , where for every face τ and its characteristic map $f_\tau: \mathbb{P}_3 \rightarrow \mathfrak{S}$, the pullback $f_\tau^* \mathbf{k}$ is a vector $V_\lambda \rightarrow \mathbb{Z}$ given by $f_\tau^* \mathbf{k}(v) = \mathbf{k}(f_\tau(v))$. Let

$$\mathcal{B}_\lambda = \text{the subgroup of } \mathbb{Z}^{V_\lambda} \text{ generated by all the balanced vectors.} \quad (23)$$

The **balanced Fock-Goncharov algebra** ([LY23, §11.5]) is defined as the monomial subalgebra

$$\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda) = \text{span}_R \{ Z^{\mathbf{k}} \mid \mathbf{k} \in \mathcal{B}_\lambda \} \subset \mathcal{Z}_\omega(\mathfrak{S}, \lambda), \quad (24)$$

where $Z^{\mathbf{k}}$ is as defined in (29); if $\mathbf{k} = (k_v)_{v \in V_\lambda}$, then $Z^{\mathbf{k}}$ is given by $\omega^{m/2} \prod_{v \in V_\lambda} Z^{k_v}$, where the product is taken with respect to any chosen ordering on V_λ , and m is a suitable integer. It is easy to verify that $\mathcal{X}_q(\mathfrak{S}, \lambda) \subset \mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda)$, and that $\{Z^{\mathbf{k}} \mid \mathbf{k} \in \mathcal{B}_\lambda\}$ is an R -basis of $\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda)$.

Suppose that $v, v' \in V_\lambda$. If v and v' are not contained in the same boundary edge, define $H_\lambda(v, v') = Q_\lambda(v, v')$. If v and v' are contained in the same boundary edge, define

$$H_\lambda(v, v') = \begin{cases} -1 & \text{when } v = v', \\ 1 & \text{when there is an arrow from } v \text{ to } v', \\ 0 & \text{otherwise.} \end{cases}$$

Remark 2.1. Note that Q_λ equals $\frac{1}{2}\bar{Q}_\lambda$ in [LY23] and H_λ equals $-\bar{H}_\lambda$ defined in [LY23, §11.7]. There are typos for the definition of \bar{H}_λ in [LY23, §11.7]; the value of $\bar{H}_\lambda(v, v')$ when v and v' are contained in the same boundary edge should be corrected to

$$\bar{H}_\lambda(v, v') = \begin{cases} 1 & \text{when } v = v', \\ -1 & \text{when there is an arrow from } v \text{ to } v', \\ 0 & \text{otherwise.} \end{cases}$$

In [LY23, §11.7], the authors misplaced the positions for -1 and 1 . We confirmed this with Thang Lê, one of the authors of [LY23].

Lemma 2.2. [LY23, Proposition 11.10] If a vector $\mathbf{k} \in \mathbb{Z}^{V_\lambda}$, considered as a row vector, is balanced, then $\mathbf{k}H_\lambda \in (n\mathbb{Z})^{V_\lambda}$.

Remark 2.3. Proposition 11.10 in [LY23] assumes the condition that \mathfrak{S} contains no interior punctures because this proposition involves a matrix \bar{K}_λ (see [LY23, §11.6]), which is defined only when \mathfrak{S} contains no interior punctures. But the arguments in [LY23, Proposition 11.10] yield the above Lemma 2.2 for general \mathfrak{S} . In [LY23, Proposition 11.10], they also show that

$$\mathbf{k}H_\lambda \in (n\mathbb{Z})^{V_\lambda} \implies \mathbf{k} \text{ is balanced}$$

using the matrix \bar{K}_λ when \mathfrak{S} contains no interior punctures.

Theorem 2.4. [BW11, K20, LY22, LY23] Suppose that \mathfrak{S} is a triangulable pb surface with a triangulation λ . Then the following hold.

(a) There is an algebra homomorphism

$$\bar{\text{tr}}_\lambda: \bar{\mathcal{F}}_\omega(\mathfrak{S}) \rightarrow \mathcal{Z}_\omega(\mathfrak{S}, \lambda),$$

called the **(reduced) quantum trace map**, such that $\text{Im } \bar{\text{tr}}_\lambda \subset \mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda)$.

(b) $\bar{\text{tr}}_\lambda$ is injective when $n = 2$ or when $n = 3$.

(c) $\bar{\text{tr}}_\lambda$ is injective when \mathfrak{S} is a polygon.

3. QUANTUM CLUSTER ALGEBRAS

We recall basics of quantum cluster algebras, obtained by gluing quantum torus algebras associated to cluster seeds along quantum mutations. We refer the readers to [BZ05, FG09a, FG09b]. Then we extend the known quantum mutations to balanced subalgebras for n -th root variables. Such an extension was studied in [H10, BW11] for $n = 2$, and in [K24] for $n = 3$.

3.1. Classical cluster \mathcal{X} -mutations. Fix a non-empty set \mathcal{V} , and a non-empty subset \mathcal{V}_{mut} of \mathcal{V} . The elements of \mathcal{V} , \mathcal{V}_{mut} and $\mathcal{V} \setminus \mathcal{V}_{\text{mut}}$ are called **vertices**, **mutable vertices** and **frozen vertices**, respectively. By a quiver Γ we mean a directed graph whose set of vertices is \mathcal{V} and equipped with weight on the edges so that the weight on each edge is 1 unless the edge connects two frozen vertices, in which case the weight is $\frac{1}{2}$. An edge of weight 1 will be called an **arrow**, and an edge of weight $\frac{1}{2}$ a **half-arrow**. As before, we denote by $Q = (Q(u, v))_{u, v \in \mathcal{V}}$ to denote the signed adjacency matrix of Γ . A quiver is considered up to equivalence, where two quivers are equivalent if they yield the same signed adjacency matrix. In particular, a quiver can be represented by a representative quiver without an oriented cycle of length 1 or 2. Let \mathcal{F} be the field of rational functions over \mathbb{Q} with the set of algebraically independent generators enumerated by \mathcal{V} , say $\mathcal{F} = \mathbb{Q}(\{y_v\}_{v \in \mathcal{V}})$. A **cluster \mathcal{X} -seed** is a pair $\mathcal{D} = (\Gamma, (X_v)_{v \in \mathcal{V}})$, where Γ is a quiver and $\{X_v\}_{v \in \mathcal{V}}$ forms an algebraically independent generating set of \mathcal{F} over \mathbb{Q} . The signed adjacency matrix Q of Γ is called the **exchange matrix** of the seed \mathcal{D} .

Suppose that $k \in \mathcal{V}_{\text{mut}}$. The **mutation** μ_k at the mutable vertex $k \in \mathcal{V}_{\text{mut}}$ is defined to be the process that transforms a seed $\mathcal{D} = (\Gamma, (X_v)_{v \in \mathcal{V}})$ into a new seed $\mu_k(\mathcal{D}) = \mathcal{D}' = (\Gamma', (X'_v)_{v \in \mathcal{V}})$. Here

$$X'_v = \begin{cases} X_v^{-1} & v = k, \\ X_v(1 + X_k^{-\text{sgn}(Q(v, k))})^{-Q(v, k)} & v \neq k, \end{cases}$$

where

$$\text{sgn}(a) = \begin{cases} 1 & a > 0, \\ 0 & a = 0, \\ -1 & a < 0, \end{cases}$$

for $a \in \mathbb{R}$, and Γ' is obtained from Γ by the following procedures: (1) reverse all the arrows incident to the vertex k , (2) for each pair of arrows $k \rightarrow i$ and $j \rightarrow k$ draw an arrow $i \rightarrow j$, (3) delete pairs of arrows $i \rightarrow j$ and $j \rightarrow i$ going in the opposite directions (more precisely, tidy up the quiver while maintaining the signed adjacency matrix). We use $Q' = (Q'(u, v))_{u, v \in \mathcal{V}}$ to denote the signed adjacency matrix of Γ' . Then we have

$$Q'(u, v) = \begin{cases} -Q(u, v) & k \in \{u, v\}, \\ Q(u, v) + \frac{1}{2}(Q(u, k)|Q(k, v)| + |Q(u, k)|Q(k, v)) & k \notin \{u, v\}. \end{cases} \quad (25)$$

It is well-known that $\mu_k(\mu_k(\mathcal{D})) = \mathcal{D}$.

3.2. Quantum mutations for Fock-Goncharov algebras. Suppose that $\mathcal{D} = (\Gamma, (X_v)_{v \in \mathcal{V}})$ is an \mathcal{X} -seed whose exchange matrix is Q . Define the **Fock-Goncharov algebra** associated to \mathcal{D} to be the quantum torus algebra

$$\begin{aligned} \mathcal{X}_q(\mathcal{D}) &= \mathbb{T}_q(Q) \\ &= R\langle X_v^{\pm 1}, v \in \mathcal{V} \rangle / (X_v X_{v'} = q^{2Q(v, v')} X_{v'} X_v \text{ for } v, v' \in \mathcal{V}). \end{aligned}$$

It is equipped with the $*$ -structure, whose $*$ -map $\mathcal{X}_q(\mathcal{D}) \rightarrow \mathcal{X}_q(\mathcal{D})$ is a ring anti-homomorphism sending each generator $X_v^{\pm 1}$ to itself $X_v^{\pm 1}$, and $q^{\pm 1}$ to its inverse $q^{\mp 1}$. We understand $\mathcal{X}_q(\mathcal{D})$ as the quantum algebra associated to the seed \mathcal{D} , which yields the classical algebra associated to \mathcal{D} as $q \rightarrow 1$. In this subsection we recall the quantum version of the mutation formula.

The quantum mutation shall be a non-commutative rational map. So we need to use the skew-field of fractions of $\mathcal{X}_q(\mathcal{D})$, which we denote by $\text{Frac}(\mathcal{X}_q(\mathcal{D}))$; for the existence of $\text{Frac}(\mathcal{X}_q(\mathcal{D}))$, see [C95].

A useful notion is the Weyl-ordered product, which is a certain normalization of a Laurent monomial defined as follows: for any $v_1, \dots, v_r \in \mathcal{V}$ and $a_1, \dots, a_r \in \mathbb{Z}$,

$$[X_{v_1}^{a_1} X_{v_2}^{a_2} \dots X_{v_r}^{a_r}] := q^{-\sum_{i < j} Q(v_i, v_j)} X_{v_1}^{a_1} X_{v_2}^{a_2} \dots X_{v_r}^{a_r} \quad (26)$$

In particular, for $\mathbf{t} = (t_v)_{v \in \mathcal{V}} \in \mathbb{Z}^{\mathcal{V}}$, the following notation for the corresponding Weyl-ordered Laurent monomial will become handy:

$$X^{\mathbf{t}} := \left[\prod_{v \in \mathcal{V}} X_v^{t_v} \right].$$

Due to the property of Weyl-ordered products, note that this element is independent of the choice of the order of the product inside the bracket, and that it is invariant under the $*$ -map.

The following special function is a crucial ingredient.

Definition 3.1 (compact quantum dilogarithm [F95, FK94]). *The quantum dilogarithm for a quantum parameter q is the function*

$$\Psi^q(x) = \prod_{r=1}^{+\infty} (1 + q^{2r-1}x)^{-1}.$$

For our purposes, the infinite product can be understood formally, as what matters for us is only the following:

Lemma 3.2. *Suppose that $xy = q^{2m}yx$ for some $m \in \mathbb{Z}$. Then we have*

$$\text{Ad}_{\Psi^q(x)}(y) = yF^q(x, m),$$

where

$$F^q(x, m) = \prod_{r=1}^{|m|} (1 + q^{(2r-1)\text{sgn}(m)}x)^{\text{sgn}(m)}.$$

For example, $F^q(x, 0) = 1$, $F^q(x, 1) = 1 + qx$, $F^q(x, 2) = (1 + qx)(1 + q^3x)$, $F^q(x, -1) = (1 + q^{-1}x)^{-1}$, $F^q(x, -2) = (1 + q^{-1}x)^{-1}(1 + q^{-3}x)^{-1}$, etc.

Instead of understanding the above as a lemma, one can just understand $\text{Ad}_{\Psi^q(x)}$ as a symbol that sends y to $y \cdot F^q(x, m)$. In particular, if $x = X_k$ and $y = X_v$ are generators of the skew-field $\text{Frac}(\mathcal{X}_q(\mathcal{D}))$, consider the expression

$$\text{Ad}_{\Psi^q(X_k)}(X_v) = X_v \cdot F^q(X_k, Q(k, v)).$$

It is easy to check that $\text{Ad}_{\Psi^q(X_k)}$ extends to a unique skew-field isomorphism from $\text{Frac}(\mathcal{X}_q(\mathcal{D}))$ to itself, which we denote by $\text{Ad}_{\Psi^q(X_k)}$.

Definition 3.3 (quantum \mathcal{X} -mutation for Fock-Goncharov algebras [BZ05, FG09a, FG09b]). *Suppose that $\mathcal{D} = (\Gamma, (X_v)_{v \in \mathcal{V}})$ is an \mathcal{X} -seed, $k \in \mathcal{V}_{\text{mut}}$ is a mutable vertex of Γ , and $\mathcal{D}' = \mu_k(\mathcal{D})$. The quantum mutation map is the isomorphism between the skew-fields*

$$\mu_{\mathcal{D}, \mathcal{D}'}^q = \mu_k^q : \text{Frac}(\mathcal{X}_q(\mathcal{D}')) \rightarrow \text{Frac}(\mathcal{X}_q(\mathcal{D}))$$

given by the composition

$$\mu_k^q = \mu_k^{\sharp q} \circ \mu'_k,$$

where μ'_k is the isomorphism of skew-fields

$$\mu'_k : \text{Frac}(\mathcal{X}_q(\mathcal{D}')) \rightarrow \text{Frac}(\mathcal{X}_q(\mathcal{D}))$$

given by

$$\mu'_k(X'_v) = \begin{cases} X_k^{-1} & \text{if } v = k, \\ [X_v X_k^{[Q(v,k)]_+}] & \text{if } v \neq k. \end{cases} \quad (27)$$

where $[a]_+$ stands for the positive part of a real number a

$$[a]_+ := \max\{a, 0\} = \frac{1}{2}(a + |a|),$$

and $[\sim]$ is the Weyl-ordered product as in (26), while $\mu_k^{\sharp q}$ is the following skew-field automorphism

$$\mu_k^{\sharp q} = \text{Ad}_{\Psi^q(X_k)} : \text{Frac}(\mathcal{X}_q(\mathcal{D})) \rightarrow \text{Frac}(\mathcal{X}_q(\mathcal{D})).$$

Suppose that $\mathcal{D} = (\Gamma, (X_v)_{v \in \mathcal{V}})$ is an \mathcal{X} -seed, $k \in \mathcal{V}_{\text{mut}}$ is a mutable vertex, and $\mathcal{D}' = \mu_k(\mathcal{D})$. We have the following.

Lemma 3.4. *Let $\mathbb{F} = \mathbb{Z}$ or $\mathbb{Z}/m\mathbb{Z}$ for some positive integer $m \geq 2$. Let $\mathcal{V}_0 \subset \mathcal{V}$ contain k , and let $\mathbf{t}' = (t'_v)_{v \in \mathcal{V}} \in \mathbb{Z}^{\mathcal{V}}$ satisfy $\sum_{v \in \mathcal{V}} Q'(u, v)t'_v = 0 \in \mathbb{F}$ for all $u \in \mathcal{V}_0$. Suppose that $\mu'_k((X')^{\mathbf{t}'}) = X^{\mathbf{t}}$, where $\mathbf{t} = (t_v)_{v \in \mathcal{V}} \in \mathbb{Z}^{\mathcal{V}}$. Then we have $\sum_{v \in \mathcal{V}} Q(u, v)t_v = 0 \in \mathbb{F}$ for all $u \in \mathcal{V}_0$.*

This lemma, which will be used later in our proofs, can be verified by straightforward computations using definitions. It can also be viewed as a corollary of [FG09b, Lem.2.7] or [K21b, Lem.3.7(1)], where the core of the necessary computation can be found. Hence we omit a proof of the above lemma.

3.3. Quantum mutations for mutable-balanced n -th root Fock-Goncharov algebras. Suppose that $\mathcal{D} = (\Gamma, (X_v)_{v \in \mathcal{V}})$ is an \mathcal{X} -seed whose exchange matrix is Q . Define the n -th root Fock-Goncharov algebra associated to \mathcal{D} to be

$$\begin{aligned} \mathcal{Z}_\omega(\mathcal{D}) &= \mathbb{T}_\omega(Q) \\ &= R\langle Z_v^{\pm 1}, v \in \mathcal{V} \rangle / (Z_v Z_{v'} = \omega^{2Q(v, v')} Z_{v'} Z_v \text{ for } v, v' \in \mathcal{V}), \end{aligned}$$

equipped with the $*$ -map $\mathcal{Z}_\omega(\mathcal{D}) \rightarrow \mathcal{Z}_\omega(\mathcal{D})$ sending each $Z_v^{\pm 1}$ to $Z_v^{\pm 1}$ and $\omega^{\pm 1}$ to $\omega^{\mp 1}$, into which the Fock-Goncharov algebra $\mathcal{X}_q(\mathcal{D})$ embeds as

$$X_v \mapsto Z_v^n, \quad \forall v \in \mathcal{V}.$$

We use $\text{Frac}(\mathcal{Z}_\omega(\mathcal{D}))$ to denote the skew-field of fractions of $\mathcal{Z}_\omega(\mathcal{D})$. Naturally, $\text{Frac}(\mathcal{X}_q(\mathcal{D}))$ embeds into $\text{Frac}(\mathcal{Z}_\omega(\mathcal{D}))$. The Weyl-ordered product can be defined similarly as before: for any $v_1, \dots, v_r \in \mathcal{V}$ and $a_1, \dots, a_r \in \mathbb{Z}$,

$$[Z_{v_1}^{a_1} \cdots Z_{v_r}^{a_r}] := \omega^{-\sum_{i < j} Q(v_i, v_j)} Z_{v_1}^{a_1} \cdots Z_{v_r}^{a_r}. \quad (28)$$

For $\mathbf{t} = (t_v)_{v \in \mathcal{V}} \in \mathbb{Z}^{\mathcal{V}}$ one defines the Weyl-ordered Laurent monomial:

$$Z^{\mathbf{t}} := \left[\prod_{v \in \mathcal{V}} Z_v^{t_v} \right], \quad (29)$$

which does not depend on the order of the product taken, and which is invariant under the $*$ -map.

The goal of the present subsection is to extend the quantum mutation maps μ_k^q defined on $\text{Frac}(\mathcal{X}_q(\mathcal{D}))$ to the n -root versions on $\text{Frac}(\mathcal{Z}_\omega(\mathcal{D}))$. We will soon see that the extension should be defined on a certain sub-skew-field of $\text{Frac}(\mathcal{Z}_\omega(\mathcal{D}))$, rather than the entire $\text{Frac}(\mathcal{Z}_\omega(\mathcal{D}))$.

Let k be a mutable vertex of Γ and let

$$\mathcal{D}' = \mu_k(\mathcal{D}).$$

Recall that the quantum mutation map $\mu_k^q : \text{Frac}(\mathcal{X}_q(\mathcal{D}')) \rightarrow \text{Frac}(\mathcal{X}_q(\mathcal{D}))$ is presented as the composition $\mu_k^q = \mu_k^{\sharp q} \circ \mu_k'$ of the monomial transformation part $\mu_k' : \text{Frac}(\mathcal{X}_q(\mathcal{D}')) \rightarrow \text{Frac}(\mathcal{X}_q(\mathcal{D}))$ and the automorphism part $\mu_k^{\sharp q} : \text{Frac}(\mathcal{X}_q(\mathcal{D})) \rightarrow \text{Frac}(\mathcal{X}_q(\mathcal{D}))$. We shall extend each of μ_k' and $\mu_k^{\sharp q}$ to its n -root versions.

First, define the skew-field isomorphism

$$\nu_k' : \text{Frac}(\mathcal{Z}_\omega(\mathcal{D}')) \rightarrow \text{Frac}(\mathcal{Z}_\omega(\mathcal{D}))$$

such that the action of ν_k' on the generators is given by the formula (27) for μ_k' with X_v replaced by Z_v :

$$\nu_k'(Z_v) = \begin{cases} Z_k^{-1} & \text{if } v = k, \\ \left[Z_v Z_k^{[Q(v,k)]_+} \right] & \text{if } v \neq k. \end{cases} \quad (30)$$

It is easy to check that ν_k' extends μ_k' . We recall a useful lemma, which is straightforward to verify:

Lemma 3.5 ([K24, Lem.3.19]). *The map ν_k' sends a Weyl-ordered Laurent monomial to a Weyl-ordered Laurent monomial.*

For an n -root extension of the automorphism part $\mu_k^{\sharp q}$, we consider the following subalgebra of $\mathcal{Z}_\omega(\mathcal{D})$.

Definition 3.6. *Define*

$$\mathcal{B}_{\mathcal{D}} = \left\{ \mathbf{t} = (t_v)_{v \in \mathcal{V}} \in \mathbb{Z}^{\mathcal{V}} \mid \sum_{v \in \mathcal{V}} Q(u, v) t_v = 0 \in \mathbb{Z}/n\mathbb{Z} \text{ for all } u \in \mathcal{V}_{\text{mut}} \right\},$$

and define $\mathcal{Z}_\omega^{\text{mbl}}(\mathcal{D})$ to be the R -submodule of $\mathcal{Z}_\omega(\mathcal{D})$ spanned by $Z^{\mathbf{t}}$ for $\mathbf{t} \in \mathcal{B}_{\mathcal{D}}$. We call $\mathcal{Z}_\omega^{\text{mbl}}(\mathcal{D})$ the **mutable-balanced subalgebra** of $\mathcal{Z}_\omega(\mathcal{D})$.

Note that $\mathcal{B}_{\mathcal{D}}$ is a subgroup of $\mathbb{Z}^{\mathcal{V}}$. So $\mathcal{Z}_{\omega}^{\text{mbl}}(\mathcal{D})$ is indeed an R -subalgebra of $\mathcal{Z}_{\omega}(\mathcal{D})$. Since $n\mathbb{Z}^{\mathcal{V}} \subset \mathcal{B}_{\mathcal{D}} \subset \mathbb{Z}^{\mathcal{V}}$, we have

$$\mathcal{X}_q(\mathcal{D}) \subset \mathcal{Z}_{\omega}^{\text{mbl}}(\mathcal{D}) \subset \mathcal{Z}_{\omega}(\mathcal{D}).$$

The mutable-balanced subalgebra $\mathcal{Z}_{\omega}^{\text{mbl}}(\mathcal{D})$ is precisely the subalgebra of the n -th root Fock-Goncharov algebra $\mathcal{Z}_{\omega}(\mathcal{D})$ to which we can extend the automorphism-part map $\mu_k^{\sharp q}$ for $\mathcal{X}_q(\mathcal{D})$. More precisely, just as for $\mathcal{X}_q(\mathcal{D})$, one needs to consider the skew-field of fractions $\text{Frac}(\mathcal{Z}_{\omega}^{\text{mbl}}(\mathcal{D}))$, which does make sense because $\mathcal{Z}_{\omega}^{\text{mbl}}(\mathcal{D})$ is a quantum torus algebra [C95], which in turn follows from the fact that $\mathcal{B}_{\mathcal{D}} \subset \mathbb{Z}^{\mathcal{V}}$ is a finitely generated free \mathbb{Z} -module.

Lemma 3.7. *There exists a unique skew-field isomorphism*

$$\nu_k^{\sharp\omega} := \text{Ad}_{\Psi^q(X_k)} : \text{Frac}(\mathcal{Z}_{\omega}^{\text{mbl}}(\mathcal{D})) \rightarrow \text{Frac}(\mathcal{Z}_{\omega}^{\text{mbl}}(\mathcal{D}))$$

such that for each $\mathbf{t} = (t_v)_{v \in \mathcal{V}} \in \mathcal{B}_{\mathcal{D}}$, we have

$$\nu_k^{\sharp\omega}(Z^{\mathbf{t}}) = Z^{\mathbf{t}} F^q(X_k, m),$$

where $m = \frac{1}{n} \sum_{v \in \mathcal{V}} Q(k, v)t_v$.

Proof. First, for $\mathbf{t} \in \mathcal{B}_{\mathcal{D}}$, we have $m \in \mathbb{Z}$, so that the expression $F^q(X_k, m)$ is well-defined. It is straightforward to check that $\nu_k^{\sharp\omega}$ extends to a unique algebra homomorphism. Similar arguments show that there is a unique skew-field homomorphism

$$\text{Ad}_{\Psi^q(X_k)^{-1}} : \text{Frac}(\mathcal{Z}_{\omega}^{\text{mbl}}(\mathcal{D})) \rightarrow \text{Frac}(\mathcal{Z}_{\omega}^{\text{mbl}}(\mathcal{D}))$$

such that $\text{Ad}_{\Psi^q(X_k)^{-1}}(Z^{\mathbf{t}}) = Z^{\mathbf{t}} F^q(X_k, -m)$ for each $\mathbf{t} = (t_v)_{v \in \mathcal{V}} \in \mathcal{B}_{\mathcal{D}}$, with $m = \frac{1}{n} \sum_{v \in \mathcal{V}} Q(k, v)t_v$. It is easy to see that $\nu_k^{\sharp\omega} = \text{Ad}_{\Psi^q(X_k)}$ and $\text{Ad}_{\Psi^q(X_k)^{-1}}$ are inverses to each other. \square

One can observe that $\text{Frac}(\mathcal{X}_q(\mathcal{D})) \subset \text{Frac}(\mathcal{Z}_{\omega}^{\text{mbl}}(\mathcal{D}))$, and that $\nu_k^{\sharp\omega}$ indeed extends $\mu_k^{\sharp q}$ defined in Def.3.3, as already hinted by the fact that we denoted both by the symbol $\text{Ad}_{\Psi^q(X_k)}$ in eq.(3.2).

The following lemma, which follows from Lemma 3.4, says that the skew-fields of fractions of the mutable-balanced subalgebras are compatible with the monomial-transformation isomorphism $\nu_k^{\sharp\omega}$.

Lemma 3.8. *The map $\text{Frac}(\mathcal{Z}_{\omega}(\mathcal{D}')) \xrightarrow{\nu_k'} \text{Frac}(\mathcal{Z}_{\omega}(\mathcal{D}))$ defined in (30) restricts to a skew-field isomorphism $\text{Frac}(\mathcal{Z}_{\omega}^{\text{mbl}}(\mathcal{D}')) \xrightarrow{\nu_k'} \text{Frac}(\mathcal{Z}_{\omega}^{\text{mbl}}(\mathcal{D}))$. \square*

We define the quantum mutation ν_k^{ω} for the n -th root Fock-Goncharov algebras, or more precisely for the skew-fields of fractions of their mutable-balanced subalgebras, to be the composition

$$\nu_k^{\omega} := \nu_k^{\sharp\omega} \circ \nu_k' : \text{Frac}(\mathcal{Z}_{\omega}^{\text{mbl}}(\mathcal{D}')) \xrightarrow{\nu_k'} \text{Frac}(\mathcal{Z}_{\omega}^{\text{mbl}}(\mathcal{D})) \xrightarrow{\nu_k^{\sharp\omega}} \text{Frac}(\mathcal{Z}_{\omega}^{\text{mbl}}(\mathcal{D})).$$

Lemma 3.9. *The map ν_k^{ω} extends μ_k^q . \square*

One useful property is:

Lemma 3.10 ([K24, Lem.3.27]). *The map ν_k^ω preserves the $*$ -structures.*

In order to apply to our situation, consider a triangulable pb surface \mathfrak{S} with a triangulation λ . Letting \mathcal{V} be V_λ and \mathcal{V}_{mut} be the subset of V_λ consisting of the vertices contained in the interior of \mathfrak{S} , one obtains a cluster \mathcal{X} -seed $\mathcal{D}_\lambda := (\Gamma_\lambda, (X_v)_{v \in V_\lambda})$. We denote the n -root Fock-Goncharov algebra and its mutable-balanced subalgebra as

$$\mathcal{Z}_\omega(\mathfrak{S}, \lambda) = \mathcal{Z}_\omega(\mathcal{D}_\lambda), \quad \mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}, \lambda) = \mathcal{Z}_\omega^{\text{mbl}}(\mathcal{D}_\lambda).$$

On the other hand, recall the balanced subalgebra $\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda)$ of $\mathcal{Z}_\omega(\mathfrak{S}, \lambda)$ defined in (24).

Lemma 3.11. *We have $\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda) \subset \mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}, \lambda)$.*

Proof. It suffices to show $\mathcal{B}_\lambda \subset \mathcal{B}_{\mathcal{D}_\lambda}$. Suppose that $\mathbf{t} = (t_v)_{v \in V_\lambda} \in \mathcal{B}_\lambda$ and $k \in \mathcal{V}_{\text{mut}}$. Lemma 2.2 implies that $\sum_{v \in V_\lambda} t_v H_\lambda(v, k) = 0 \in \mathbb{Z}/n\mathbb{Z}$. From the definition of H_λ , it follows that $H_\lambda(v, k) = Q_\lambda(v, k) = -Q_\lambda(k, v)$ for any $v \in V_\lambda$. Then we have

$$\sum_{v \in V_\lambda} Q_\lambda(k, v) t_v = 0 \in \mathbb{Z}/n\mathbb{Z}.$$

This shows that $\mathbf{t} \in \mathcal{B}_{\mathcal{D}_\lambda}$. □

Remark 3.12. *When the triangulable pb surface \mathfrak{S} has empty boundary, one would obtain $\mathcal{B}_\lambda = \mathcal{B}_{\mathcal{D}_\lambda}$ and hence $\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda) = \mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}, \lambda)$ if one could prove the converse of Lemma 2.2, upgrading it to an if-and-only-if statement. One can try to come up with a modified version of $\mathcal{B}_{\mathcal{D}}$ for a general cluster \mathcal{X} -seed \mathcal{D} that involves additional equations about the frozen vertices, so that it matches \mathcal{B}_λ in the case of a seed coming from a triangulation λ of a triangulable pb surface \mathfrak{S} with boundary as well. We do not undertake these tasks in the present paper. See [K24, Rem.3.25] for a related question.*

Eventually, in the next section §4 we will be applying the n -th root quantum mutations ν_k^ω to the balanced subalgebras $\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda) \subset \mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}, \lambda)$, or more precisely to their skew-fields of fractions $\text{Frac}(\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda))$, which make sense since $\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda)$ is a quantum torus algebra, which in turn holds because $\mathcal{B}_\lambda \subset \mathcal{B}_{\mathcal{D}_\lambda}$ is a finitely generated free \mathbb{Z} -module.

Still in this section, we will work on the mutable-balanced algebras $\mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}, \lambda)$ and the skew-fields of fractions thereof, for triangulations λ of a triangulable pb surface \mathfrak{S} . In the following subsection, we will compose the maps ν_k^ω studied in the present subsection to construct a coordinate change isomorphism between $\text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}, \lambda))$ for different triangulations λ .

3.4. Coordinate change isomorphisms for change of triangulations. Let \mathfrak{S} be a triangulable pb surface with a triangulation λ . In the last subsection we considered a particular cluster \mathcal{X} -seed $\mathcal{D}_\lambda = (\Gamma_\lambda, (X_v)_{v \in V_\lambda})$, associated to a special quiver Γ_λ , with certain vertex sets $\mathcal{V} = V_\lambda$ and \mathcal{V}_{mut} . This cluster \mathcal{X} -seed comes from a special coordinate chart on a certain moduli space $\mathcal{P}_{\text{PGL}_n, \mathfrak{S}}$ studied by Fock and Goncharov [FG06], and by Goncharov and Shen [GS19]. This space parametrizes PGL_n -local systems on \mathfrak{S} together with some extra data at the punctures and the boundary

of \mathfrak{S} . The stated SL_n -skein algebra $\mathcal{S}_\omega(\mathfrak{S})$ plays a crucial role in understanding the deformation quantization of this space. In [FG06, GS19], the authors construct a coordinate chart for this space per each triangulation λ of \mathfrak{S} , whose coordinate variables are denoted by X_v , enumerated by the vertices v of the quiver Γ_λ . They further show that these special charts for different triangulations are related to each other by a special sequence of mutations. In the present paper we do not recall the definition of the moduli space $\mathcal{P}_{\mathrm{PGL}_n, \mathfrak{S}}$ nor the construction of the coordinates, but only recall the sequence of mutations.

We begin with the elementary change of triangulations, namely with the case when λ and λ' are triangulations of \mathfrak{S} that differ exactly by one edge, say e . We say that each of these triangulations are obtained from each other by a **flip** at the edge e . Consider the ideal quadrilateral formed by the two ideal triangles of λ sharing e (it may not actually be a quadrilateral, but that doesn't matter here). Let $V_{\lambda;e}$ be the set of all vertices of V_λ that either lie in the interior of one of these two triangles or lie on e ; so $V_{\lambda;e}$ consists of $(n-1)^2$ vertices. For each $i = 0, 1, 2, \dots, n-2$, we will define a subset $V_{\lambda;e}^{(i)}$ of $V_{\lambda;e}$. Draw this ideal quadrilateral for λ as in Figure 3, so that e is the ‘middle vertical line’, and after flipping at e , the flipped e would be the ‘middle horizontal line’. As in Figure 3, for each vertex v in $V_{\lambda;e}$ one can define the vertical distance $d_{\mathrm{vert}}(v) \in \mathbb{N}$ from the middle horizontal line, and the horizontal distance $d_{\mathrm{hori}}(v) \in \mathbb{N}$ from the middle vertical line. Define

$$V_{\lambda;e}^{(i)} := \left\{ v \in V_{\lambda;e} \mid \begin{array}{l} d_{\mathrm{vert}}(v) \leq i, \quad d_{\mathrm{vert}}(v) \equiv i \pmod{2}, \\ d_{\mathrm{hori}}(v) \leq n-2-i, \quad d_{\mathrm{hori}}(v) \equiv n-2-i \pmod{2} \end{array} \right\}.$$

See Figure 3; so each $V_{\lambda;e}^{(i)}$ can be viewed as forming the grid points of a rectangle, consisting of $(i+1)(n-i-1)$ points. Notice that these sets for different i are not necessarily disjoint with each other. For example, if $n \geq 4$, then $V_{\lambda;e}^{(0)}$ and $V_{\lambda;e}^{(2)}$ have $n-3$ vertices in common.

The sought-for mutation sequence of Fock, Goncharov and Shen consists of first mutations at all vertices of $V_{\lambda;e}^{(0)}$ in any order, then mutations at all vertices of $V_{\lambda;e}^{(1)}$ in any order, etc., then lastly mutations at all vertices of $V_{\lambda;e}^{(n-2)}$ in any order. Regardless of the orders chosen, this mutation sequence connects the coordinate chart \mathcal{D}_λ of $\mathcal{P}_{\mathrm{PGL}_n, \mathfrak{S}}$ for λ to the chart $\mathcal{D}_{\lambda'}$ for λ' . In particular, the length of this mutation sequence is $\sum_{i=0}^{n-2} (i+1)(n-1-i) = \sum_{j=1}^{n-1} j(n-j) = \frac{1}{6}(n^3 - n) =: r$. Let us denote the corresponding sequence of vertices as

$$v_1, v_2, \dots, v_r.$$

So, the first $n-1$ of them would be the elements of $V_{\lambda;e}^{(0)}$, and we have

$$\mathcal{D}_{\lambda'} = \mu_{v_r} \cdots \mu_{v_2} \mu_{v_1}(\mathcal{D}_\lambda) \tag{31}$$

and in particular,

$$\Gamma_{\lambda'} = \mu_{v_r} \cdots \mu_{v_2} \mu_{v_1}(\Gamma_\lambda).$$




FIGURE 3. Vertices of the n -triangulation quiver involved in the flip of triangulations; the above example picture is for the case $n = 4$

At the quantum level, we first define the quantum coordinate change isomorphism for the skew-fields of fractions of the usual Fock-Goncharov algebras

$$\Phi_{\lambda\lambda'}^q := \mu_{v_1}^q \circ \cdots \circ \mu_{v_r}^q : \text{Frac}(\mathcal{X}_q(\mathfrak{S}, \lambda')) \rightarrow \text{Frac}(\mathcal{X}_q(\mathfrak{S}, \lambda)). \quad (32)$$

Note that the order of the mutation sequence seems to be reversed because of the contravariant nature of the classical and quantum coordinate change maps. We then define the n -root balanced version of the quantum coordinate change isomorphism to be

$$\Theta_{\lambda\lambda'}^\omega := \nu_{v_1}^\omega \circ \cdots \circ \nu_{v_r}^\omega : \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}, \lambda')) \rightarrow \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}, \lambda)). \quad (33)$$

Suppose now that λ and λ' are any two triangulations of \mathfrak{S} . A **triangulation sweep** connecting λ and λ' is a sequence of triangulations $\Lambda = (\lambda_1, \dots, \lambda_m)$ such that $\lambda_1 = \lambda$, $\lambda_m = \lambda'$, and λ_{i+1} is obtained from λ_i by a flip for each $1 \leq i \leq m-1$. It is well known that for any λ and λ' , there exists a triangulation sweep Λ connecting λ and λ' ([L09]). For any such triangulation sweep Λ , we define the corresponding quantum coordinate change isomorphisms

$$\Phi_\Lambda^q := \Phi_{\lambda_1\lambda_2}^q \circ \cdots \circ \Phi_{\lambda_{m-1}\lambda_m}^q : \text{Frac}(\mathcal{X}_q(\mathfrak{S}, \lambda')) \rightarrow \text{Frac}(\mathcal{X}_q(\mathfrak{S}, \lambda)), \quad (34)$$

$$\Theta_\Lambda^\omega := \Theta_{\lambda_1\lambda_2}^\omega \circ \cdots \circ \Theta_{\lambda_{m-1}\lambda_m}^\omega : \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}, \lambda')) \rightarrow \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}, \lambda)). \quad (35)$$

Proposition 3.13 ([BZ05, FG09a, KN11]; see also [K21a, K24]). Φ_Λ^q depends only on λ and λ' .

In order to prove this proposition, one should show that any relation satisfied by classical coordinate change maps for flips, say by the maps $\Phi_{\lambda\lambda'}^1$, such as the pentagon relation, is also satisfied by the quantum counterparts $\Phi_{\lambda\lambda'}^q$. It suffices to show that any relation satisfied by classical mutation maps μ_k^1 is satisfied by the quantum counterparts μ_k^q . One could use the decomposition $\mu_k^q = \mu_k^{\sharp q} \circ \mu'_k$ for each quantum mutation μ_k^q . Then, for a sought-for equation among μ_k^q , one can move all μ'_k to the left (or right), with the help of (27). By also exploiting a slight variant of the decomposition $\mu_k^q = \mu_k^{\sharp q} \circ \mu'_k$ (a ‘signed’ decomposition), one can ensure that the problem boils down to proving an equation among μ'_k ’s only, and an equation among $\mu_k^{\sharp q}$ ’s only. We refer the readers to [KN11, K21a] how this argument works. Essentially the same argument works for the balanced n -th root quantum mutations, yielding the following.

Proposition 3.14. Θ_Λ^ω depends only on λ and λ' . □

Therefore, when we do not have to keep track of a specific triangulation sweep, we can write

$$\Phi_{\lambda\lambda'}^q = \Phi_\Lambda^q \quad \text{and} \quad \Theta_{\lambda\lambda'}^\omega = \Theta_\Lambda^\omega.$$

One obvious remark:

Remark 3.15. For any two triangulations λ and λ' , the balanced n -th root quantum coordinate change map $\Theta_{\lambda\lambda'}^\omega$ extends the quantum coordinate change map $\Phi_{\lambda\lambda'}^q$.

A crucial property we shall develop and use about the balanced quantum coordinate change maps is the compatibility with cutting and gluing of a pb surface along an ideal arc.

Still, let \mathfrak{S} be a triangulable pb surface with a triangulation λ . In principle, one can cut the surface \mathfrak{S} along any ideal arc, but here let us assume that e is a non-boundary edge of the triangulation λ . As in §2.2, we denote by $\text{Cut}_e(\mathfrak{S})$ the pb surface obtained from \mathfrak{S} by cutting along edge e , and denote the projection map by

$$\mathbf{pr}_e : \text{Cut}_e(\mathfrak{S}) \rightarrow \mathfrak{S},$$

so that $\mathbf{pr}_e^{-1}(e)$ consists of two ideal arcs e', e'' of $\text{Cut}_e\mathfrak{S}$. In §2.2 we studied the induced splitting homomorphism \mathbb{S}_e between the stated SL_n -skein algebras and that between the reduced stated SL_n -skein algebras; here we will need the latter

$$\mathbb{S}_e : \overline{\mathcal{F}}_\omega(\mathfrak{S}) \rightarrow \overline{\mathcal{F}}_\omega(\text{Cut}_e(\mathfrak{S})).$$

We further need to investigate a ‘splitting map’ between the (n -th root) Fock-Goncharov algebras. Note that $\lambda_e = (\lambda \setminus \{e\}) \cup \{e', e''\}$ is a triangulation of $\text{Cut}_e(\mathfrak{S})$. We say λ_e is induced from λ . For a small vertex v in e , i.e. $v \in V_\lambda$ lying in e , we have $\mathbf{pr}_e^{-1}(v) = \{v', v''\}$, where v' and v'' are small vertices of λ' . There is an algebra embedding

$$\mathcal{S}_e : \mathcal{Z}_\omega(\mathfrak{S}, \lambda) \rightarrow \mathcal{Z}_\omega(\text{Cut}_e(\mathfrak{S}), \lambda_e) \tag{36}$$

defined on the generators Z_v , $v \in V_\lambda$, by

$$\mathcal{S}_e(Z_v) = \begin{cases} Z_v & \text{if } v \text{ is not contained in } e, \\ [Z_{v'}Z_{v''}] & \text{if } v \text{ is contained in } e \text{ and } \mathbf{pr}_e^{-1}(v) = \{v', v''\}, \end{cases} \tag{37}$$

where $[\sim]$ is the Weyl-ordered product (28); since we are not allowing self-folded triangles in this paper, in fact $Z_{v'}$ commutes with $Z_{v''}$ in the second case, and hence $\mathcal{S}_e(Z_v) = Z_{v'}Z_{v''}$.

The following statement is about the compatibility of the quantum trace map $\overline{\text{tr}}_\lambda$ (Theorem 2.4) with the splitting homomorphisms.

Theorem 3.16 ([LY23]). *The following diagram commutes*

$$\begin{array}{ccc} \overline{\mathcal{F}}_\omega(\mathfrak{S}) & \xrightarrow{\mathbb{S}_e} & \overline{\mathcal{F}}_\omega(\text{Cut}_e(\mathfrak{S})) \\ \downarrow \overline{\text{tr}}_\lambda & & \downarrow \overline{\text{tr}}_{\lambda_e} \\ \mathcal{Z}_\omega(\mathfrak{S}, \lambda) & \xrightarrow{\mathcal{S}_e} & \mathcal{Z}_\omega(\text{Cut}_e(\mathfrak{S}), \lambda_e), \end{array}$$

where \mathbb{S}_e is the splitting homomorphism defined in subsection 2.2.

Lemma 3.17. *The algebra embedding $\mathcal{S}_e: \mathcal{Z}_\omega(\mathfrak{S}, \lambda) \rightarrow \mathcal{Z}_\omega(\text{Cut}_e(\mathfrak{S}), \lambda_e)$ restricts to an algebra embedding from $\mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}, \lambda)$ to $\mathcal{Z}_\omega^{\text{mbl}}(\text{Cut}_e(\mathfrak{S}), \lambda_e)$.*

Proof. For any $\mathbf{t} \in \mathbb{Z}^{V_\lambda}$, we have $Z^{\mathbf{t}} \in \mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}, \lambda)$ if and only if, for each mutable vertex $k \in V_\lambda$ (i.e. k is not contained in the boundary), there exists an integer m such that $Z^{\mathbf{t}}X_k = q^{2m}X_kZ^{\mathbf{t}}$ (see Definition 3.6).

Suppose that $k' \in V_{\lambda_e}$ is a mutable vertex, and that $\mathbf{t}' \in \mathbb{Z}^{V_\lambda}$ satisfies $Z^{\mathbf{t}'} \in \mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}, \lambda)$. Since k is also a mutable vertex for V_λ , the above discussion shows that there exists an integer m' such that $Z^{\mathbf{t}'}X_{k'} = q^{2m'}X_{k'}Z^{\mathbf{t}'}$. Since $k' \in V_{\lambda_e}$ is a mutable vertex, we have $\mathcal{S}_e(X_{k'}) = X_{k'}$. This implies that $\mathcal{S}_e(Z^{\mathbf{t}'})X_{k'} = q^{2m'}X_{k'}\mathcal{S}_e(Z^{\mathbf{t}'}) \in \mathcal{Z}_\omega(\text{Cut}_e(\mathfrak{S}), \lambda_e)$. Thus we have $\mathcal{S}_e(Z^{\mathbf{t}'}) \in \mathcal{Z}_\omega^{\text{mbl}}(\text{Cut}_e(\mathfrak{S}), \lambda_e)$. \square

Suppose that λ and λ' are two triangulations and that λ' is obtained from λ by performing a flip at an edge e_1 . Let $e \in \lambda \setminus \{e_1\}$. Suppose that λ_e (resp. λ'_e) is the triangulation of $\text{Cut}_e(\mathfrak{S})$ induced by λ (resp. λ'). Then λ'_e is obtained from λ_e by performing a flip at e_1 . Let v_1, \dots, v_r be a sequence of mutable vertices of Γ_λ such that $\mathcal{D}_{\lambda'} = \mu_{v_r} \cdots \mu_{v_1}(\mathcal{D}_\lambda)$, in the sense of (31). Since $e \neq e_1$, each $v_i = \mathbf{pr}_e^{-1}(v_i) \in V_{\lambda_e}$ is contained in the interior of $\text{Cut}_e(\mathfrak{S})$. We claim that

$$\mathcal{D}_{\lambda'_e} = \mu_{v_r} \cdots \mu_{v_1}(\mathcal{D}_{\lambda_e}) \tag{38}$$

holds. Consider the subsurface of \mathfrak{S} formed by the two ideal triangles of λ having e_1 as a side. Say that the three edges of one triangle are e_1, e_2, e_3 in the clockwise order, while those of the other are e_1, e_4, e_5 in the clockwise order. If e appears at most once in the sequence e_2, e_3, e_4, e_5 , then (38) is easy to see. In case $e = e_2 = e_4$ or $e = e_3 = e_5$, then one has to be careful about variables associated to vertices lying in e when checking (38), but one can still check the equality. The case $e_2 = e_3$ or $e_4 = e_5$ is excluded, because then λ has a self-folded triangle. The case $e_2 = e_5$ or $e_3 = e_4$ is excluded, because then λ' has a self-folded triangle. We thus have (38) indeed.

The following lemma can be viewed as a balanced n -th root quantum version of (38). See [K24, Prop.3.34] for the case when $n = 3$, the proof of which we closely follow in our proof below.

Lemma 3.18. *The following diagram commutes*

$$\begin{array}{ccc} \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}, \lambda')) & \xrightarrow{\mathcal{S}_e} & \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\text{Cut}_e(\mathfrak{S}), \lambda'_e)) \\ \downarrow \Theta_{\lambda\lambda'}^\omega & & \downarrow \Theta_{\lambda_e\lambda'_e}^\omega \\ \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}, \lambda)) & \xrightarrow{\mathcal{S}_e} & \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\text{Cut}_e(\mathfrak{S}), \lambda_e)). \end{array}$$

Proof. Let $\mathcal{D}_0 := \mathcal{D}_\lambda$, and for each $i = 1, \dots, r$ let $\mathcal{D}_i = \mu_{v_i}(\mathcal{D}_{i-1})$ be the i -th seed in the sequence of seeds connecting \mathcal{D}_λ and $\mathcal{D}_{\lambda'}$. In particular, $\mathcal{D}_r = \mathcal{D}_{\lambda'}$. Likewise for the cut surface $\text{Cut}_e(\mathfrak{S})$, let $\mathcal{D}_{0,e} := \mathcal{D}_{\lambda_e}$, and $\mathcal{D}_{i,e} := \mu_{v_i}(\mathcal{D}_{i-1,e})$ for $i = 1, \dots, r$. For each $1 \leq i \leq r$ we have the algebra embedding

$$\mathcal{S}_e : \mathcal{Z}_\omega(\mathcal{D}_i) \rightarrow \mathcal{Z}_\omega(\mathcal{D}_{i,e})$$

from (36). We first claim that this restricts to an algebra embedding between the mutable-balanced subalgebras

$$\mathcal{S}_e : \mathcal{Z}_\omega^{\text{mbl}}(\mathcal{D}_i) \rightarrow \mathcal{Z}_\omega^{\text{mbl}}(\mathcal{D}_{i,e}). \quad (39)$$

Indeed, let $\mathbf{t} = (t_v)_{v \in \mathcal{V}} \in \mathcal{B}_{\mathcal{D}_i} \subset \mathbb{Z}^\mathcal{V}$, so that $\sum_{v \in \mathcal{V}} Q_i(u, v)t_v \in n\mathbb{Z}$ for all mutable vertices u of \mathcal{V} , i.e. vertices of V_λ not lying on the boundary of \mathfrak{S} , where Q_i is the exchange matrix of the seed \mathcal{D}_i . Then

$$\mathcal{S}_e(\mathbf{Z}^{\mathbf{t}}) = \mathbf{Z}^{\mathbf{t}'} \quad (40)$$

with $\mathbf{t}' = (t'_v)_{v \in \mathcal{V}_e} \in \mathbb{Z}^{\mathcal{V}_e}$, where $\mathcal{V}_e = V_{\lambda_e}$; if $v \in \mathcal{V}$ does not lie on e so that $\mathbf{pr}_e^{-1}(v) = \{v\}$ then $t'_v = t_v$, and if $v \in \mathcal{V}$ lies on e so that $\mathbf{pr}_e^{-1}(v) = \{v', v''\}$ then $t'_{v'} = t'_{v''} = t_v$. If $u \in \mathcal{V}_e$ is mutable, then u does not lie in the boundary of $\text{Cut}_e(\mathfrak{S})$, hence $\mathbf{pr}_e^{-1}(u) = \{u\}$ where u is a mutable vertex of $\mathcal{V} = V_\lambda$ not lying on e . We would like to show that

$$\sum_{v \in \mathcal{V}_e} Q_{i,e}(u, v)t'_v \in n\mathbb{Z}$$

holds, where $Q_{i,e}$ is the exchange matrix of $\mathcal{D}_{i,e}$. For each $v \in \mathcal{V}$, observe that

$$Q_i(u, v) = \sum_{w \in \mathbf{pr}_e^{-1}(v)} Q_{i,e}(u, w). \quad (41)$$

Since $\mathbf{pr}_e : \mathcal{V}_e \rightarrow \mathcal{V}$ is surjective, and since $t'_w = t_{\mathbf{pr}_e(w)}$ for each $w \in \mathcal{V}_e$, it follows that

$$\sum_{v \in \mathcal{V}} Q_i(u, v)t_v = \sum_{w \in \mathcal{V}_e} Q_{i,e}(u, w)t'_w. \quad (42)$$

The left hand side belongs to $n\mathbb{Z}$, it follows that so does the right hand side, as desired.

We still denote by \mathcal{S}_e the map between the skew-fields of fractions $\text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathcal{D}_i)) \rightarrow \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathcal{D}_{i,e}))$ induced by $\mathcal{S}_e : \mathcal{Z}_\omega^{\text{mbl}}(\mathcal{D}_i) \rightarrow \mathcal{Z}_\omega^{\text{mbl}}(\mathcal{D}_{i,e})$. We now claim that the

following diagram commutes:

$$\begin{array}{ccc} \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathcal{D}_i)) & \xrightarrow{\mathcal{S}_e} & \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathcal{D}_{i,e})) \\ \nu_{v_i}^\omega \downarrow & & \downarrow \nu_{v_i}^\omega \\ \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathcal{D}_{i-1})) & \xrightarrow{\mathcal{S}_e} & \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathcal{D}_{i-1,e})). \end{array} \quad (43)$$

To show that the diagram in (43) commutes, it suffices to show that the following two diagrams commute:

$$\begin{array}{ccc} \text{Frac}(\mathcal{Z}_\omega(\mathcal{D}_i)) & \xrightarrow{\mathcal{S}_e} & \text{Frac}(\mathcal{Z}_\omega(\mathcal{D}_{i,e})) \\ \nu_{v_i}' \downarrow & & \downarrow \nu_{v_i}' \\ \text{Frac}(\mathcal{Z}_\omega(\mathcal{D}_{i-1})) & \xrightarrow{\mathcal{S}_e} & \text{Frac}(\mathcal{Z}_\omega(\mathcal{D}_{i-1,e})) \end{array} \quad (44)$$

and

$$\begin{array}{ccc} \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathcal{D}_{i-1})) & \xrightarrow{\mathcal{S}_e} & \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathcal{D}_{i-1,e})) \\ \nu_{v_i}^{\#\omega} \downarrow & & \downarrow \nu_{v_i}^{\#\omega} \\ \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathcal{D}_{i-1})) & \xrightarrow{\mathcal{S}_e} & \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathcal{D}_{i-1,e})). \end{array} \quad (45)$$

Note that the first diagram restricts to the maps between the mutable-balanced subalgebras (or the skew-fields of fractions thereof), since we showed that both ν_{v_i}' and \mathcal{S}_e restrict to the mutable-balanced subalgebras.

Let us temporarily denote the generators of $\mathcal{Z}_\omega(\mathcal{D}_i)$ or $\mathcal{Z}_\omega(\mathcal{D}_{i,e})$ by Z'_v , and those of $\mathcal{Z}_\omega(\mathcal{D}_{i-1})$ or $\mathcal{Z}_\omega(\mathcal{D}_{i-1,e})$ by Z_v . For any $v \in \mathcal{V}$ such that v is not contained in e , we have $\mathbf{pr}_e^{-1}(v) = \{v\}$ (also $\mathbf{pr}_e^{-1}(v_i) = \{v_i\}$), so $Q_i(v, v_i) = Q_{i,e}(v, v_i)$ (cf.(41)), $\mathcal{S}_e(Z'_v) = Z'_v$ and $\mathcal{S}_e(Z'_{v_i}) = Z'_{v_i}$ ((37)). In view of (30), we can observe

$$\mathcal{S}_e(\nu_{v_i}'(Z'_v)) = \nu_{v_i}'(\mathcal{S}_e(Z'_v)).$$

Now let $v \in \mathcal{V}$ be a vertex that lies in e ; then $(\mathbf{pr}_e)^{-1}(v) = \{v', v''\}$, for some distinct $v', v'' \in \mathcal{V}_e$. From (37) and (30), together with Lemma 3.5, we have

$$\begin{aligned} \nu_{v_i}'(\mathcal{S}_e(Z'_v)) &= \nu_{v_i}'([Z'_{v'} Z'_{v''}]) = [Z_{v'} Z_{v_i}^{[Q_{i-1,e}(v', v_i)]_+} Z_{v''} Z_{v_i}^{[Q_{i-1,e}(v'', v_i)]_+}], \\ \mathcal{S}_e(\nu_{v_i}'(Z'_v)) &= \mathcal{S}_e([Z_v Z_{v_i}^{[Q_{i-1}(v, v_i)]_+}]) = [Z_{v'} Z_{v''} Z_{v_i}^{[Q_{i-1}(v, v_i)]_+}]. \end{aligned}$$

From (41) one has

$$Q_{i-1}(v, v_i) = Q_{i-1,e}(v', v_i) + Q_{i-1,e}(v'', v_i).$$

When $n > 2$, we claim that at least one of $Q_{i-1,e}(v', v_i)$ and $Q_{i-1,e}(v'', v_i)$ is zero. As observed in the proof of [K24, Prop.3.34], the only possible case when both are nonzero is when e and e_1 shares a puncture in the interior of the surface, such that the valence of λ at this puncture is 2. However, then flipping at e would yield a self-folded triangle (in λ'), so this case is excluded. When $n = 2$, it is easy to check that $Q_{i-1,e}(v', v_i)Q_{i-1,e}(v'', v_i) \geq 0$. This shows $[Q_{i-1}(v, v_i)]_+ = [Q_{i-1,e}(v', v_i)]_+ +$

$[Q_{i-1,e}(v'', v_i)]_+$ and therefore $\nu'_{v_i}(\mathcal{S}_e(Z'_v)) = \mathcal{S}_e(\nu'_{v_i}(Z'_v))$. Thus the diagram in (44) commutes.

For $\mathbf{t} = (t_v)_{v \in \mathcal{V}} \in \mathcal{B}_{\mathcal{D}_{i-1}}$, suppose that $\mathcal{S}_e(Z^{\mathbf{t}}) = Z^{\mathbf{t}'}$, where $\mathbf{t}' = (t'_v)_{v \in \mathcal{V}_e} \in \mathcal{B}_{\mathcal{D}_{i-1,e}}$. Lemma 3.7 implies that

$$\begin{aligned} \nu_{v_i}^{\sharp\omega}(\mathcal{S}_e(Z^{\mathbf{t}})) &= Z^{\mathbf{t}'} F^q(X_{v_i}, \frac{1}{n} \sum_{v \in \mathcal{V}_e} Q_{i-1,e}(v_i, v) t'_v), \\ \mathcal{S}_e(\nu_{v_i}^{\sharp\omega}(Z^{\mathbf{t}})) &= \mathcal{S}_e(Z^{\mathbf{t}'} F^q(X_{v_i}, \frac{1}{n} \sum_{v \in \mathcal{V}} Q_{i-1}(v_i, v) t_v)) = Z^{\mathbf{t}'} F^q(X_{v_i}, \frac{1}{n} \sum_{v \in \mathcal{V}} Q_{i-1}(v_i, v) t_v). \end{aligned}$$

From (42) we have

$$\sum_{v \in \mathcal{V}_e} Q_{i-1,e}(v_i, v) t'_v = \sum_{v \in \mathcal{V}} Q_{i-1}(v_i, v) t_v.$$

Thus we have $\nu_{v_i}^{\sharp\omega}(\mathcal{S}_e(Z^{\mathbf{t}})) = \mathcal{S}_e(\nu_{v_i}^{\sharp\omega}(Z^{\mathbf{t}}))$, and hence the diagram in (45) commutes, finishing the proof. \square

4. COMPATIBILITY BETWEEN SL_n -QUANTUM TRACE MAPS AND COORDINATE CHANGE ISOMORPHISMS

4.1. Compatibility of the quantum trace map with the balanced quantum coordinate change maps $\Theta_{\lambda\lambda'}^\omega$. Let \mathfrak{S} be a triangulable pb surface. For each triangulation λ of \mathfrak{S} , we have a quantum trace homomorphism

$$\overline{\text{tr}}_\lambda : \overline{\mathcal{F}}_\omega(\mathfrak{S}) \rightarrow \mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda) \subset \mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}, \lambda) \subset \mathcal{Z}_\omega(\mathfrak{S}, \lambda),$$

from Theorem 2.4; see Lemma 3.11 for $\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda) \subset \mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}, \lambda)$. Meanwhile, for any pair of triangulations λ and λ' , we developed the balanced n -th root quantum coordinate change isomorphism

$$\Theta_{\lambda\lambda'}^\omega : \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}, \lambda')) \rightarrow \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}, \lambda))$$

in the previous section, which extends the previously known quantum coordinate change isomorphism $\Phi_{\lambda\lambda'}^q : \text{Frac}(\mathcal{X}_q(\mathfrak{S}, \lambda')) \rightarrow \text{Frac}(\mathcal{X}_q(\mathfrak{S}, \lambda))$ coming from the theory of quantum cluster algebras.

In the present subsection we show that the quantum trace maps for different triangulations are compatible with each other via the isomorphisms $\Theta_{\lambda\lambda'}^\omega$. This compatibility, formulated as the following theorem, constitutes the core part of the main result of the present paper.

Theorem 4.1 (compatibility of quantum trace with mutable-balanced n -root quantum coordinate change maps $\Theta_{\lambda\lambda'}^\omega$). *Let \mathfrak{S} be a triangulable pb surface. For any two triangulations λ and λ' of \mathfrak{S} , the following diagram commutes:*

$$\begin{array}{ccc} & \overline{\mathcal{F}}_\omega(\mathfrak{S}) & \\ \overline{\text{tr}}_{\lambda'} \swarrow & & \searrow \overline{\text{tr}}_\lambda \\ \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}, \lambda')) & \xrightarrow{\Theta_{\lambda\lambda'}^\omega} & \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}, \lambda)). \end{array} \quad (46)$$

That is, we have

$$\overline{\text{tr}}_\lambda = \Theta_{\lambda\lambda'}^\omega \circ \overline{\text{tr}}_{\lambda'}. \quad (47)$$

Here is how we break down the proof. First, from Proposition 3.14 we immediately obtain the consistency among the quantum coordinate change isomorphisms; that is, for any triangulations λ , λ' and λ'' , we have

$$\Theta_{\lambda\lambda'}^\omega \circ \Theta_{\lambda'\lambda''}^\omega = \Theta_{\lambda\lambda''}^\omega. \quad (48)$$

Now, since any two triangulations are connected by a sequence of flips, it suffices to prove Theorem 4.1 for the case when λ and λ' are related by a single flip at an edge, say e_1 .

There are two triangles of λ having e_1 as a side. Let one of them have sides e_1, e_2, e_3 in the clockwise order, and the other have sides e_1, e_4, e_5 in the clockwise order. Cutting the surface \mathfrak{S} along all the edges in the set $\{e_2, e_3, e_4, e_5\}$ (whose cardinality may be less than 4) that do not lie in the boundary of \mathfrak{S} yields a triangulable pb surface, the connected component of which that contains (the image of) e_1 is isomorphic to \mathbb{P}_4 , the ideal quadrilateral, the 4-gon, i.e. the closed disc minus four punctures at the boundary. By a repeated application of Lemma 3.18 about the compatibility between the coordinate change maps $\Theta_{\lambda\lambda'}$ and the splitting homomorphisms of skein algebras, the problem boils down to proving Theorem 4.1 in the case when the surface \mathfrak{S} is the 4-gon \mathbb{P}_4 (we will justify this ‘boiling down’ process). So we shall first show the following:

Proposition 4.2. *Theorem 4.1 holds when \mathfrak{S} is the 4-gon \mathbb{P}_4 , where λ, λ' are the triangulations illustrated in Figure 4.*




FIGURE 4. Two triangulations λ and λ' of \mathbb{P}_4

In fact, the core computational part of the proof of the entire Theorem 4.1, hence the main difficulty thereof, lies in this case of \mathbb{P}_4 , i.e. in the proof of Proposition 4.2, which we postpone until the next subsection. At the moment, we explain how Proposition 4.2 implies Theorem 4.1, i.e. justify the ‘boiling down’ process.

Proof of Theorem 4.1, using Proposition 4.2. Let \mathfrak{S} be a triangulable pb surface, and λ and λ' be triangulations of \mathfrak{S} . As mentioned above, it suffices to show the statement, (47), in the case when λ and λ' are related by a flip at an edge, say e_1 .

Define

$$E = \{e \in \lambda \mid e \neq e_1 \text{ and } e \text{ is not a boundary edge}\}. \quad (49)$$

We use $\mathfrak{S}_E := \text{Cut}_E(\mathfrak{S})$ to denote the pb surface obtained from \mathfrak{S} by cutting \mathfrak{S} along all edges in E . Then the connected components of \mathfrak{S}_E consist of exactly one \mathbb{P}_4 and

some number of \mathbb{P}_3 . We use λ_E (resp. λ'_E) to denote the triangulation of \mathfrak{S}_E induced by λ (resp. λ').

Then λ'_E is obtained from λ_E by a flip at the edge e_1 . Consider the following diagram

$$\begin{array}{ccccc}
\text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}, \lambda')) & \xrightarrow{\Theta_{\lambda\lambda'}^\omega} & \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}, \lambda)) & & \\
\downarrow \mathcal{S}_E & \swarrow \overline{\text{tr}}_{\lambda'} & \nearrow \overline{\text{tr}}_\lambda & & \downarrow \mathcal{S}_E \\
& \mathcal{S}_\omega(\mathfrak{S}) & & & \\
& \downarrow \mathbb{S}_E & & & \\
& \mathcal{S}_\omega(\mathfrak{S}_E) & & & \\
& \swarrow \overline{\text{tr}}_{\lambda'_E} & \searrow \overline{\text{tr}}_{\lambda_E} & & \\
\text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}_E, \lambda'_E)) & \xrightarrow{\Theta_{\lambda_E\lambda'_E}^\omega} & \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}_E, \lambda_E)) & &
\end{array} \tag{50}$$

where all the vertical arrows are the algebra homomorphisms induced by the splitting maps (for cutting along all the edges in E), as in Theorem 3.16; note that one can cut the edges in any order, as the splitting maps for different edges of a triangulation commute with each other. Observe that the two long vertical arrows in the above diagram are injective. Theorem 3.16 implies that each of the left and the right trapezoids of this diagram commutes. Proposition 4.2 implies that the lower triangle in this diagram commutes. Lemma 3.18 implies that the outermost square of this diagram commutes.

One can now show that the upper triangle also commutes. Let us elaborate. Pick any $u \in \mathcal{S}_\omega(\mathfrak{S})$. Then

$$\begin{aligned}
\mathcal{S}_E(\Theta_{\lambda\lambda'}^\omega(\overline{\text{tr}}_{\lambda'}(u))) &= \Theta_{\lambda\lambda'}^\omega(\mathcal{S}_E(\overline{\text{tr}}_{\lambda'}(u))) \quad (\because \text{outermost square}) \\
&= \Theta_{\lambda\lambda'}^\omega(\overline{\text{tr}}_{\lambda'_E}(\mathbb{S}_E(u))) \quad (\because \text{left trapezoid}) \\
&= \overline{\text{tr}}_{\lambda_E}(\mathbb{S}_E(u)) \quad (\because \text{lower triangle}) \\
&= \mathcal{S}_E(\overline{\text{tr}}_\lambda(u)) \quad (\because \text{right trapezoid}).
\end{aligned}$$

Since \mathcal{S}_E (for the right vertical arrow) is injective, we get $\Theta_{\lambda\lambda'}^\omega(\overline{\text{tr}}_{\lambda'}(u)) = \overline{\text{tr}}_\lambda(u)$, as desired. \square

4.2. Proof of compatibility for the case of quadrilateral \mathbb{P}_4 . In the present subsection we prove Proposition 4.2. We shall verify that (47) holds when applied to generating elements of the reduced stated SL_n -skein algebra $\mathcal{S}_\omega(\mathbb{P}_4)$. Consider the three corner arcs a, b, c in \mathbb{P}_4 as in Figure 5. For $i, j \in \{1, 2, \dots, n\}$, we use a_{ij} to denote the stated web diagram such that the state of the starting point (resp. endpoint) of a is i (resp. j). Similarly, we define b_{ij} and c_{ij} . We have the following.

Lemma 4.3. [LY23, Theorem 6.1(c)] *The stated SL_n -skein algebra $\mathcal{S}_\omega(\mathbb{P}_4)$ is generated by a_{ij}, b_{ij}, c_{ij} for $i, j \in \{1, \dots, n\}$.*

In view of the definition of the bad arcs and reduced stated SL_n -skein algebra, we get:




FIGURE 5. Three corner arcs a, b, c in \mathbb{P}_4 . Four edges of \mathbb{P}_4 are labeled by e_2, e_3, e_4, e_5

Lemma 4.4. *The reduced stated SL_n -skein algebra $\overline{\mathcal{S}}_\omega(\mathbb{P}_4)$ is generated by a_{ij}, b_{ij}, c_{ij} for $i, j \in \{1, \dots, n\}$ with $i \geq j$.*


Thus it suffices to check (47) when applied to these generators. See [K24, Lem.4.2] for another argument that allows us to focus only on these corner arcs in \mathbb{P}_4 .

We now explain our proof of (47) for the generator $a_{ij} \in \overline{\mathcal{S}}_\omega(\mathbb{P}_4)$, with $i \geq j$. Proof by a direct computation would be quite involved; see [K24] for such a proof, for the case $n = 3$. Instead, we adapt the methodology using paths lying in a certain ‘network’, which is developed initially by Schrader and Shapiro in [SS17] and further investigated in [CS23, LY23]. This methodology, which we shall now briefly recall, will let us bypass difficult computations.

Choose any triangulation of \mathbb{P}_4 . For each triangle t of this triangulation, consider its n -triangulation, which subdivides the triangle t into n^2 small triangles. Consider a graph dual to this n -triangulation of t , such that the set of all edges of this graph is in bijection with the set of all edges of the n -triangulation. In particular, for each small triangle there is a vertex of this graph, and for each boundary edge of the n -triangulation there is a vertex; see the blue graph in Figure 6. These graphs for different triangles can naturally be glued together.

Now we give orientations to the edges of this graph in a particular way that depends on the choice of a distinguished side of each (big) triangle; we call the resulting oriented graph a **network**. There is a left-turn version and a right-turn version; here we only need to consider the left-turn version. See [SS17, CS23, LY23] for the precise recipe. In particular, in each (big) triangle, the orientations of the edges meeting the distinguished side of the triangle are ‘outward’, while those of the edges meeting the other sides are ‘inward’. Figure 6 presents examples of (left-turn) networks \mathcal{N} and \mathcal{N}' for the two triangulations λ and λ' of \mathbb{P}_4 , where e_3 is the ‘outward’ side, and e_2, e_4, e_5 are the ‘inward’ sides, where we are using the same labels for the four boundary arcs of \mathbb{P}_4 as in Figure 4. Although Figure 6 is only about the case when $n = 4$, we believe that the readers can easily see how it would work for general n . We label the vertices of these networks lying on the bottom side e_2 as $\alpha_1, \dots, \alpha_n$, and those lying on the left side e_3 as β_1, \dots, β_n , as appearing in Figure 6.

A **path** in a network is a concatenation of oriented edges of the network. For $1 \leq i, j \leq n$, let $\mathcal{N}(ij)$ (resp. $\mathcal{N}'(ij)$) be the set of all paths in the network \mathcal{N} (resp. \mathcal{N}') from α_i to β_j . For example, $\mathcal{N}(11)$ consists of a single path, which is made of

FIGURE 6. (left-turn) Networks for triangulations of \mathbb{P}_4 ; when $n = 4$

three oriented edges. As can be verified by inspection in Figure 6, $\mathcal{N}(21)$ consists of two paths, if $n \geq 2$; one path is made of six oriented edges, and the other path of eight oriented edges.

We shall soon see that the role of paths is that the value of a stated arc under the quantum trace map can be presented as a sum over all paths starting and ending with specified boundary vertices of a suitable network. The summand corresponding to a path p is a balanced Weyl-ordered Laurent monomial $Z^{\mathbf{t}}$ (see (29)). To describe the exponent \mathbf{t} , we need some definitions.

For any path p in $\mathcal{N}(ij)$ (resp. $\mathcal{N}'(ij)$), define the element $\mathbf{k}(p, \lambda) = (k_v(p))_{v \in V_\lambda} \in \mathbb{Z}^{V_\lambda}$ (resp. $\mathbf{k}(p, \lambda') = (k_v(p))_{v \in V_{\lambda'}} \in \mathbb{Z}^{V_{\lambda'}}$) as

$$k_v(p) = \begin{cases} 1 & \text{if } v \in V_\lambda \text{ (resp. } v \in V_{\lambda'}) \text{ lies strictly on the left of the path } p, \\ 0 & \text{otherwise,} \end{cases}$$

where the elements of V_λ (resp. $V_{\lambda'}$) are naturally thought of as vertices of the n -triangulation of the triangles of λ (resp. λ') used in the definition of the network \mathcal{N} (resp. \mathcal{N}'); see Figure 6.


Note that for each $i = 1, 2, \dots, n$, $\mathcal{N}(ii)$ (resp. $\mathcal{N}'(ii)$) consists of a single path, say p_i (resp. p'_i). Now define

$$\mathbf{k} := \sum_{1 \leq i \leq n} \mathbf{k}(p_i, \lambda) \in \mathbb{Z}^{V_\lambda}, \quad \mathbf{k}' := \sum_{1 \leq i \leq n} \mathbf{k}(p'_i, \lambda') \in \mathbb{Z}^{V_{\lambda'}}.$$



Lemma 4.5 (quantum trace as sum over paths in a network). *One has*

- (1) $\overline{\text{tr}}_\lambda(a_{ij})$ equals $\sum_{p \in \mathcal{N}(ij)} [X^{\mathbf{k}(p, \lambda)} Z^{\mathbf{k}}] \in \mathcal{Z}_\omega^{\text{bl}}(\mathbb{P}_4, \lambda)$.
- (2) $\overline{\text{tr}}_{\lambda'}(a_{ij})$ equals $\sum_{p' \in \mathcal{N}'(ij)} [X^{\mathbf{k}(p', \lambda')} Z^{\mathbf{k}'}] \in \mathcal{Z}_\omega^{\text{bl}}(\mathbb{P}_4, \lambda')$.

Proof. (1) The edges of the network \mathcal{N} meeting the diagonal ideal arc of λ are oriented toward the lower-left direction as seen in Figure 6; so a path in \mathcal{N} starting at a vertex

in the left triangle  can never leave this triangle. Therefore each path $p \in \mathcal{N}(ij)$ is completely contained in the left triangle. Then the sought-for result follows directly from [LY23, Theorem 10.5], which expresses the quantum trace for a triangle \mathbb{P}_3 as a sum over paths in a network.

(2) A path $p \in \mathcal{N}'(ij)$ necessarily meets both triangles of λ' , so [LY23, Theorem 10.5] does not directly apply. Denote by e'_1 the diagonal ideal arc of λ' . For each $i = 1, \dots, n$, recall that there is only one path in $\mathcal{N}'(ii)$, which we denoted by p'_i ; this path p'_i intersects e'_1 at a single point, which we denote by γ_i as in Figure 6.

We now cut \mathbb{P}_4 along e'_1 . Note that $\text{Cut}_{e'_1}(\mathbb{P}_4)$ consists of two copies of triangles \mathbb{P}_3 . The triangle  containing α_i (resp. the triangle  containing β_j) is denoted by \mathbb{P}_3^α (resp. \mathbb{P}_3^β). We use $\mathbf{pr}_{e'_1}$ to denote the projection from $\text{Cut}_{e'_1}(\mathbb{P}_4)$ to \mathbb{P}_4 . For each $i = 1, \dots, n$, note that $(\mathbf{pr}_{e'_1})^{-1}(\gamma_i)$ consists of two elements. The one contained in \mathbb{P}_4^α (resp. \mathbb{P}_4^β) is denoted by w_i^α (resp. w_i^β). The network \mathcal{N}' for \mathbb{P}_4 induces a network $(\mathbf{pr}_{e'_1})^{-1}(\mathcal{N}')$ for $\text{Cut}_{e'_1}(\mathbb{P}_4)$, which has two components. The component contained in \mathbb{P}_3^α (resp. \mathbb{P}_3^β) is denoted by \mathcal{N}^α (resp. \mathcal{N}^β). Then \mathcal{N}^α (resp. \mathcal{N}^β) contains vertices w_i^α (resp. w_i^β) for $i = 1, \dots, n$. The preimage $(\mathbf{pr}_{e'_1})^{-1}(a)$ in $\text{Cut}_{e'_1}(\mathbb{P}_4)$ of the oriented arc a in \mathbb{P}_4 consists of two oriented arcs. The one contained in \mathbb{P}_3^α (resp. \mathbb{P}_3^β) is denoted by a^α (resp. a^β). For each $i, j \in \{1, \dots, n\}$, similarly as a_{ij} , we can define stated arcs a_{ij}^α (resp. a_{ij}^β) in \mathbb{P}_3^α (resp. \mathbb{P}_3^β), whose state values at the initial and the terminal endpoints are i and j respectively.

The definition of the splitting homomorphism $\mathbb{S}_{e'_1}$ in equation (15) shows that

$$\mathbb{S}_{e'_1}(a_{ij}) = \sum_{1 \leq t \leq n} a_{it}^\alpha \otimes a_{tj}^\beta \in \overline{\mathcal{F}}_\omega(\mathbb{P}_3^\alpha) \otimes_R \overline{\mathcal{F}}_\omega(\mathbb{P}_3^\beta) = \overline{\mathcal{F}}_\omega(\text{Cut}_{e'_1}(\mathbb{P}_4)).$$

As before, let $\lambda'_{e'_1}$ denote the triangulation of $\text{Cut}_{e'_1}(\mathbb{P}_4) = \mathbb{P}_3^\alpha \sqcup \mathbb{P}_3^\beta$ induced by λ' ; that is, $\lambda'_{e'_1} = \lambda^\alpha \sqcup \lambda^\beta$, where λ^α (resp. λ^β) is the unique triangulation of \mathbb{P}_3^α (resp. \mathbb{P}_3^β). Hence one can express $\overline{\text{tr}}_{\lambda'_{e'_1}} : \overline{\mathcal{F}}_\omega(\text{Cut}_{e'_1}(\mathbb{P}_4)) \rightarrow \mathcal{Z}_\omega(\text{Cut}_{e'_1}(\mathbb{P}_4), \lambda'_{e'_1})$ as $\overline{\text{tr}}_{\lambda^\alpha} \otimes \overline{\text{tr}}_{\lambda^\beta} : \overline{\mathcal{F}}_\omega(\mathbb{P}_3^\alpha) \otimes_R \overline{\mathcal{F}}_\omega(\mathbb{P}_3^\beta) \rightarrow \mathcal{Z}_\omega(\mathbb{P}_3^\alpha, \lambda^\alpha) \otimes_R \mathcal{Z}_\omega(\mathbb{P}_3^\beta, \lambda^\beta)$. For each of $\overline{\text{tr}}_{\lambda^\alpha}$ and $\overline{\text{tr}}_{\lambda^\beta}$ we apply [LY23, Theorem 10.5]. That is, note that

$$\begin{aligned} \overline{\text{tr}}_{\lambda'_{e'_1}}(\mathbb{S}_{e'_1}(a_{ij})) &= (\overline{\text{tr}}_{\lambda^\alpha} \otimes \overline{\text{tr}}_{\lambda^\beta}) \left(\sum_{1 \leq t \leq n} a_{it}^\alpha \otimes a_{tj}^\beta \right) \\ &= \sum_{1 \leq t \leq n} \overline{\text{tr}}_{\lambda^\alpha}(a_{it}^\alpha) \otimes \overline{\text{tr}}_{\lambda^\beta}(a_{tj}^\beta) \\ &= \sum_{1 \leq t \leq n} \left(\sum_{p \in \mathcal{N}^\alpha(it)} [X^{\mathbf{k}(p, \lambda^\alpha)} Z^{\mathbf{k}^\alpha}] \otimes \sum_{p \in \mathcal{N}^\beta(tj)} [X^{\mathbf{k}(p, \lambda^\beta)} Z^{\mathbf{k}^\beta}] \right), \end{aligned} \quad (51)$$

$$(\because [\text{LY23, Theorem 10.5}]) \quad (52)$$

where

$$\mathcal{N}^\alpha(it) = \{\text{paths in } \mathcal{N}^\alpha \text{ from } \alpha_i \text{ to } \gamma_t^\alpha\},$$

$$\begin{aligned}
\mathcal{N}^\beta(tj) &= \{\text{paths in } \mathcal{N}^\beta \text{ from } \gamma_t^\beta \text{ to } \beta_j\}, \\
\mathbf{k}(p, \lambda^\alpha) &= (k_v(p))_{v \in V_{\lambda^\alpha}}, \quad k_v(p) = \begin{cases} 1 & \text{if } v \in \lambda^\alpha \text{ lies on the left of } p, \\ 0 & \text{otherwise,} \end{cases} \\
\mathbf{k}(p, \lambda^\beta) &= (k_v(p))_{v \in V_{\lambda^\beta}}, \quad k_v(p) = \begin{cases} 1 & \text{if } v \in \lambda^\beta \text{ lies on the left of } p, \\ 0 & \text{otherwise,} \end{cases} \\
\mathbf{k}^\alpha &= \sum_{i=1}^n \mathbf{k}_{(p_i)^\alpha}, \quad \text{where } (p_i)^\alpha \text{ is the only constituent path of } \mathcal{N}^\alpha(ii), \\
\mathbf{k}^\beta &= \sum_{i=1}^n \mathbf{k}_{(p_i)^\beta}, \quad \text{where } (p_i)^\beta \text{ is the only constituent path of } \mathcal{N}^\beta(ii).
\end{aligned}$$

The natural concatenation yields a map

$$\bigsqcup_{1 \leq t \leq n} (\mathcal{N}^\alpha(it) \times \mathcal{N}^\beta(tj)) \rightarrow \mathcal{N}'(ij), \quad (53)$$

which is easily seen to be an injection. It is also surjective, as any path p' in $\mathcal{N}'(ij)$ must meet the diagonal ideal arc e'_1 of λ' exactly once, say at γ_t ; then one can see that p' is the concatenation of a uniquely determined path in $\mathcal{N}^\alpha(it)$, say $(p')^\alpha$, and a uniquely determined path in $\mathcal{N}^\beta(tj)$, say $(p')^\beta$.

Recall from (36) the algebra embedding

$$\mathcal{S}_{e'_1} : \mathcal{Z}_\omega(\mathbb{P}_4, \lambda') \rightarrow \mathcal{Z}_\omega(\text{Cut}_{e'_1}(\mathbb{P}_4), \lambda'_{e'_1}) \cong \mathcal{Z}_\omega(\mathbb{P}_3^\alpha, \lambda^\alpha) \otimes_R \mathcal{Z}_\omega(\mathbb{P}_3^\beta, \lambda^\beta).$$

For any path p' in $\mathcal{N}'(ij)$ we can observe

$$\mathcal{S}_{e'_1}[X^{\mathbf{k}(p', \lambda')} Z^{\mathbf{k}'}] = [X^{\mathbf{k}((p')^\alpha, \lambda^\alpha)} Z^{\mathbf{k}^\alpha}] \otimes [X^{\mathbf{k}((p')^\beta, \lambda^\beta)} Z^{\mathbf{k}^\beta}], \quad (54)$$

which one can see from the definition of $\mathcal{S}_{e'_1}$ in (37) to hold up to a multiplicative constant that is an integer power of $\omega^{1/2}$. One can further observe that the multiplicative constant is 1, by showing that $\mathcal{S}_{e'_1}$ sends a Weyl-ordered Laurent monomial to a Weyl-ordered Laurent monomial, which in turn is straightforward to prove, e.g. by the same argument as in the proof of [K24, Lem.3.19(3)–(5)].

Now, for $i, j \in \{1, \dots, n\}$, take the sum of (54) over all paths p' in $\mathcal{N}'(ij)$:

$$\begin{aligned}
\mathcal{S}_{e'_1} \left(\sum_{p' \in \mathcal{N}'(ij)} [X^{\mathbf{k}(p', \lambda')} Z^{\mathbf{k}'}] \right) &= \sum_{p' \in \mathcal{N}'(ij)} ([X^{\mathbf{k}((p')^\alpha, \lambda^\alpha)} Z^{\mathbf{k}^\alpha}] \otimes [X^{\mathbf{k}((p')^\beta, \lambda^\beta)} Z^{\mathbf{k}^\beta}]) \\
&= \sum_{1 \leq t \leq n} \left(\sum_{p \in \mathcal{N}^\alpha(it)} [X^{\mathbf{k}(p, \lambda^\alpha)} Z^{\mathbf{k}^\alpha}] \right) \otimes \left(\sum_{p \in \mathcal{N}^\beta(tj)} [X^{\mathbf{k}(p, \lambda^\beta)} Z^{\mathbf{k}^\beta}] \right),
\end{aligned}$$

where the latter equality is due to the bijection in (53). We thus have

$$\begin{aligned}
\mathcal{S}_{e'_1} \left(\sum_{p' \in \mathcal{N}'(ij)} [X^{\mathbf{k}(p', \lambda')} Z^{\mathbf{k}'}] \right) &= \overline{\text{tr}}_{\lambda'_{e'_1}}(\mathcal{S}_{e'_1}(a_{ij})) \quad (\because (51)) \\
&= \mathcal{S}_{e'_1}(\overline{\text{tr}}_{\lambda'}(a_{ij})) \quad (\because \text{Theorem 3.16}).
\end{aligned}$$

Since \mathcal{S}_{e_1} is injective, it follows that $\sum_{p' \in \mathcal{N}'(ij)} [X^{\mathbf{k}(p', \lambda')} Z^{\mathbf{k}'}]$ equals $\overline{\text{tr}}_{\lambda'}(a_{ij})$, as desired. \square

Remark 4.6. *Using the same technique, Lemma 4.5 can be generalized to calculate the image of the stated corner arcs in any triangulable pb surface under the quantum trace map.*

Keep in mind that our goal is to show the following form of compatibility:

$$\Theta_{\lambda\lambda'}^\omega(\overline{\text{tr}}_{\lambda'}(a_{ij})) = \overline{\text{tr}}_\lambda(a_{ij}).$$

The reason why we invoked the description of the value of the quantum trace in terms of sum over paths in a network is because Schrader and Shapiro showed a form of compatibility of such sums over paths with mutations, namely [SS17, Proposition 4.2]. Each cluster mutation corresponds to mutation of a network (see [SS17, Figure 9]), and the sequence of cluster mutations realizing the flip $\lambda \rightsquigarrow \lambda'$ along the ideal arc e_1 in \mathbb{P}_4 (as in (31)) transforms the network \mathcal{N} into the network \mathcal{N}' . Applying [SS17, Proposition 4.2] to this mutation sequence, we obtain the following:

Lemma 4.7. *For each $i, j \in \{1, \dots, n\}$, one has*

$$\Phi_{\lambda\lambda'}^q\left(\sum_{p' \in \mathcal{N}'(ij)} X^{\mathbf{k}(p', \lambda')}\right) = \sum_{p \in \mathcal{N}(ij)} X^{\mathbf{k}(p, \lambda)}. \quad \square$$

We stress that this lemma settles the most involved part of the proof of our sought-for Proposition 4.2. That is, a direct computational proof of this lemma, for each fixed n , would require an enormous effort; see [K24] for a version of such a proof for $n = 3$. In a sense, the advantage of using the network description of quantum trace mainly lies in this lemma.

We now perform the remaining steps to prove Proposition 4.2.

Lemma 4.8. *One has*

$$\Theta_{\lambda\lambda'}^\omega(Z^{\mathbf{k}'}) = Z^{\mathbf{k}}.$$

Note that $\mathcal{N}'(ii) = \{p'_i\}$ and $\mathcal{N}(ii) = \{p_i\}$, so from Lemma 4.7 applied to $i = j$ we get

$$\Phi_{\lambda\lambda'}^q(X^{\mathbf{k}(p'_i, \lambda')}) = X^{\mathbf{k}(p_i, \lambda)}, \quad (55)$$

hence $\Phi_{\lambda\lambda'}^q(X^{\mathbf{k}'}) = X^{\mathbf{k}}$ holds up to a multiplicative constant that is an integer power of $q^{1/2}$. One can show that this constant is 1, as we shall see later. However, $\Phi_{\lambda\lambda'}^q(X^{\mathbf{k}'}) = X^{\mathbf{k}}$ does not directly imply the sought-for $\Theta_{\lambda\lambda'}^\omega(Z^{\mathbf{k}'}) = Z^{\mathbf{k}}$, although these two are consistent with each other, in view of Remark 3.15. We give a direct proof of Lemma 4.8 below.

Note that any ground ring R is an algebra over $\mathbb{Z}[\omega^{\pm\frac{1}{2}}]$. Then we can use the functor $- \otimes_{\mathbb{Z}[\omega^{\pm\frac{1}{2}}]} R$ to change the ground ring from $\mathbb{Z}[\omega^{\pm\frac{1}{2}}]$ to R . Due to this functor, we assume that the ground ring is $\mathbb{Z}[\omega^{\pm\frac{1}{2}}]$ in the following proof.

Proof of Lemma 4.8. Denote by \mathcal{D}_λ and $\mathcal{D}_{\lambda'}$ the cluster seeds corresponding to λ and λ' , and that the sequence of mutations at the vertices v_1, v_2, \dots, v_r turns \mathcal{D}_λ to $\mathcal{D}_{\lambda'}$, as in (31). Like we did in the proof of Lemma 3.18, let $\mathcal{D}_0 = \mathcal{D}_\lambda$, and let $\mathcal{D}_k = \mu_{v_k}(\mathcal{D}_{k-1})$ for each $k = 1, \dots, r$, so that $\mathcal{D}_r = \mathcal{D}_{\lambda'}$ in particular. Denote by \mathcal{V} the underlying vertex sets of all these seeds \mathcal{D}_k , $k = 0, 1, \dots, r$, which are naturally identified with each other; in particular, $\mathcal{V} = V_\lambda = V_{\lambda'}$. Let \mathcal{V}_{mut} be the vertices lying in the interior of \mathbb{P}_4 . Let Q_k be the exchange matrix of \mathcal{D}_k .

Let $i \in \{1, \dots, n\}$. Define $\mathbf{t}^{(k)} = (t_v^{(k)})_{v \in \mathcal{V}} \in \mathbb{Z}^{\mathcal{V}}$ by the following equations:

$$\mathbf{t}^{(r)} = \mathbf{k}(p'_i, \lambda'), \quad \nu'_{v_k}(Z^{\mathbf{t}^{(k)}}) = Z^{\mathbf{t}^{(k-1)}}, \quad k = r, r-1, \dots, 2, 1.$$

More explicitly, in view of (30), the second equation says that

$$t_{v_k}^{(k-1)} = -t_{v_k}^{(k)} + \sum_{v \in \mathcal{V}} [Q_{(k-1)}(v, v_k)]_+ t_v^{(k)}, \quad \text{and} \quad t_v^{(k-1)} = t_v^{(k)} \quad \text{for all } v \neq v_k. \quad (56)$$

For the seed $\mathcal{D}_{\lambda'}$, it is straightforward to observe that

$$\sum_{v \in \mathcal{V}} Q_r(u, v) t_v^{(r)} = 0 \quad \text{for all } u \in \mathcal{V}_{\text{mut}}. \quad (57)$$

One can translate this condition into $Z_u Z^{\mathbf{t}^{(r)}} = Z^{\mathbf{t}^{(r)}} Z_u \in \mathcal{Z}_\omega(\mathcal{D}^{(r)})$, $\forall u \in \mathcal{V}_{\text{mut}}$. By applying Lemma 3.4 to (57) inductively, one obtains:

$$\text{For each } k = r, r-1, \dots, 1, \text{ one has } \sum_{v \in \mathcal{V}} Q_{k-1}(u, v) t_v^{(k-1)} = 0, \quad \forall u \in \mathcal{V}_{\text{mut}}. \quad (58)$$

Since $\nu_{v_k}^\omega = \nu_{v_k}^{\sharp\omega} \circ \nu'_{v_k}$, in view of the action of $\nu_{v_k}^{\sharp\omega}$ as written in Lemma 3.7, it is easy to see that (58) implies:

$$\text{For each } k = r, r-1, \dots, 1, \text{ one has } \nu_{v_k}^\omega(Z^{\mathbf{t}^{(k)}}) = Z^{\mathbf{t}^{(k-1)}}.$$

Thus $\Theta_{\lambda\lambda'}^\omega(Z^{\mathbf{t}^{(r)}}) = \nu_{v_r}^\omega \dots \nu_{v_1}^\omega(Z^{\mathbf{t}^{(r)}}) = Z^{\mathbf{t}^{(0)}}$.

On the other hand, a verbatim proof, with all $\nu_i^?$ replaced by $\mu_i^?$ and $Z^?$ by $X^?$, shows $\Phi_{\lambda\lambda'}^q(X^{\mathbf{t}^{(r)}}) = X^{\mathbf{t}^{(0)}}$. Since $\mathbf{t}^{(r)} = \mathbf{k}(p'_i, \lambda')$, now (55) (together with injectivity of $\Phi_{\lambda\lambda'}^q$) tells us that $\mathbf{t}^{(0)} = \mathbf{k}(p_i, \lambda)$. Hence we showed that

$$\Theta_{\lambda\lambda'}^\omega(Z^{\mathbf{k}(p'_i, \lambda')}) = Z^{\mathbf{k}(p_i, \lambda)}$$

holds for all $i = 1, \dots, n$.

From (57) it follows that $Z^{\mathbf{k}(p'_i, \lambda')}$, $i = 1, \dots, n$, mutually commute with each other, hence $Z^{\mathbf{k}(p'_1, \lambda')} Z^{\mathbf{k}(p'_2, \lambda')} \dots Z^{\mathbf{k}(p'_n, \lambda')} = Z^{\mathbf{k}(p'_1, \lambda') + \dots + \mathbf{k}(p'_n, \lambda')}$ (see e.g. [K24, Lem.3.20(A)]) which equals $Z^{\mathbf{k}'}$. Similarly, from (58) for $k = 1$ we see that $Z^{\mathbf{t}^{(0)}} = Z^{\mathbf{k}(p_i, \lambda)}$, $i = 1, \dots, n$, mutually commute with each other, hence

$$Z^{\mathbf{k}(p_1, \lambda)} \dots Z^{\mathbf{k}(p_n, \lambda)} = Z^{\mathbf{k}(p_1, \lambda) + \dots + \mathbf{k}(p_n, \lambda)} = Z^{\mathbf{k}}.$$

Since $\Theta_{\lambda\lambda'}^\omega$ is a homomorphism, it follows that $\Theta_{\lambda\lambda'}^\omega(Z^{\mathbf{k}'}) = Z^{\mathbf{k}}$ holds, as desired. \square

We now assemble Lemmas 4.7 and 4.8. We first perform a necessary preliminary computation. For $i, j \in \{1, \dots, n\}$, define $\mathbf{t}_{ij} = ((t_{ij})_v)_{v \in \mathcal{V}} \in \mathbb{Z}^{\mathcal{V}}$ as

$$(t_{ij})_v = \begin{cases} n & \text{if } v = \alpha_l \text{ for } 1 \leq l \leq i, \text{ or } v = \beta_m \text{ for } 1 \leq m \leq j, \\ 0 & \text{otherwise,} \end{cases}$$

keeping the identification $\mathcal{V} = V_\lambda = V_{\lambda'}$ in mind. Write $\mathbf{k} = (k_v)_{v \in V_\lambda}$ and $\mathbf{k}' = (k'_v)_{v \in V_{\lambda'}}$; recall $\mathbf{k}(p, \lambda) = (k_v(p))_{v \in V_\lambda}$ and $\mathbf{k}(p', \lambda') = (k_v(p'))_{v \in V_{\lambda'}}$. For each $p \in \mathcal{N}(ij)$, from (57) it follows that

$$\sum_{v \in \mathcal{V}} \sum_{u \in V_\lambda} n Q_\lambda(v, u) k_v(p) k_u = \sum_{v \in \{\alpha_l, \beta_m | 1 \leq l \leq i, 1 \leq m \leq j\}} \sum_{u \in V_\lambda} Q_\lambda(v, u) (t_{ij})_v k_u =: m_{ij}.$$

For each $p' \in \mathcal{N}'(ij)$, from (58) (for $k = 1$) it follows that

$$\sum_{v \in \mathcal{V}} \sum_{u \in V_{\lambda'}} n Q_{\lambda'}(v, u) k_v(p') k'_u = \sum_{v \in \{\alpha_l, \beta_m | 1 \leq l \leq i, 1 \leq m \leq j\}} \sum_{u \in V_{\lambda'}} Q_{\lambda'}(v, u) (t'_{ij})_v k'_u =: m'_{ij}.$$

Note that the integers m_{ij} and m'_{ij} do not depend on p or p' , respectively. We will eventually be able to show that $m_{ij} = m'_{ij}$. At the moment, we have

$$\overline{\text{tr}}_\lambda(a_{ij}) \stackrel{\text{Lemma 4.5(1)}}{=} \sum_{p \in \mathcal{N}(ij)} [X^{\mathbf{k}(p, \lambda)} Z^{\mathbf{k}}] = \omega^{-m_{ij}} \sum_{p \in \mathcal{N}(ij)} X^{\mathbf{k}(p, \lambda)} Z^{\mathbf{k}}, \quad (59)$$

$$\overline{\text{tr}}_{\lambda'}(a_{ij}) \stackrel{\text{Lemma 4.5(2)}}{=} \sum_{p' \in \mathcal{N}'(ij)} [X^{\mathbf{k}(p', \lambda')} Z^{\mathbf{k}'}] = \omega^{-m'_{ij}} \sum_{p' \in \mathcal{N}'(ij)} X^{\mathbf{k}(p', \lambda')} Z^{\mathbf{k}'}. \quad (60)$$

Observe now that

$$\begin{aligned} \Theta_{\lambda \lambda'}^\omega(\overline{\text{tr}}_{\lambda'}(a_{ij})) &= \Theta_{\lambda \lambda'}^\omega(\omega^{-m'_{ij}} \sum_{p' \in \mathcal{N}'(ij)} X^{\mathbf{k}(p', \lambda')} Z^{\mathbf{k}'}) \quad (\because (60)) \\ &= \omega^{-m'_{ij}} \Phi_{\lambda \lambda'}^q(\sum_{p' \in \mathcal{N}'(ij)} X^{\mathbf{k}(p', \lambda')} Z^{\mathbf{k}'}) \Theta_{\lambda \lambda'}^\omega(Z^{\mathbf{k}'}) \quad (\because \text{Remark 3.15}) \\ &= \omega^{-m'_{ij}} \sum_{p \in \mathcal{N}(ij)} X^{\mathbf{k}(p, \lambda)} Z^{\mathbf{k}} \quad (\because \text{Lemmas 4.7 and 4.8}) \\ &= \omega^{m_{ij} - m'_{ij}} \overline{\text{tr}}_\lambda(a_{ij}). \quad (\because (59)) \end{aligned}$$

From Lemma 4.5, both $\overline{\text{tr}}_{\lambda'}(a_{ij}) \in \mathcal{Z}_\omega(\mathbb{P}_4, \lambda')$ and $\overline{\text{tr}}_\lambda(a_{ij}) \in \mathcal{Z}_\omega(\mathbb{P}_4, \lambda)$ are sum of Weyl-ordered Laurent monomials, hence they are invariant under the $*$ -maps. By Lemma 3.10 it follows that $\Theta_{\lambda \lambda'}^\omega$ preserves the $*$ -structures, hence sends $*$ -invariant elements to $*$ -invariant elements. It then follows that $\omega^{m_{ij} - m'_{ij}} = 1$. We thus obtain the sought-for compatibility $\Theta_{\lambda \lambda'}^\omega(\overline{\text{tr}}_{\lambda'}(a_{ij})) = \overline{\text{tr}}_\lambda(a_{ij})$ for the stated arcs a_{ij} .

The proofs for the remaining generators b_{ij} and c_{ij} are similar, and we omit them. In view of Lemma 4.4, the compatibility $\Theta_{\lambda \lambda'}^\omega(\overline{\text{tr}}_{\lambda'}(u)) = \overline{\text{tr}}_\lambda(u)$ holds for all elements $u \in \overline{\mathcal{S}}_\omega(\mathbb{P}_4)$, hence Proposition 4.2 is proved. As explained in §4.1, this finishes the proof of Theorem 4.1.

4.3. The restriction of $\Theta_{\lambda\lambda'}^\omega$ to the balanced part. Theorem 4.1 states that the quantum trace maps $\overline{\text{tr}}_\lambda$ and $\overline{\text{tr}}_{\lambda'}$ for different triangulations are compatible with each other through the quantum coordinate change map $\Theta_{\lambda\lambda'}^\omega : \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}, \lambda')) \rightarrow \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}, \lambda))$ between the skew-fields of fractions of the mutable-balanced algebras. Meanwhile, in view of Theorem 2.4(a), the image of the quantum trace maps $\overline{\text{tr}}_{\lambda'}$ and $\overline{\text{tr}}_\lambda$ lie in $\text{Frac}(\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda'))$ and $\text{Frac}(\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda))$, the skew-fields of fractions of the balanced algebras. In the present subsection we show that $\Theta_{\lambda\lambda'}^\omega$ restricts to the map between these balanced fraction skew-fields.

Proposition 4.9. *Let \mathfrak{S} be a triangulable pb surface, and λ and λ' be triangulations of \mathfrak{S} . The quantum coordinate change isomorphism $\Theta_{\lambda\lambda'}^\omega : \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}, \lambda')) \rightarrow \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}, \lambda))$ established in §3.4 restricts to an isomorphism*

$$(\Theta_{\lambda\lambda'}^\omega)^{\text{bl}} : \text{Frac}(\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda')) \rightarrow \text{Frac}(\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda)).$$

Since $\Theta_{\lambda'\lambda}^\omega : \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}, \lambda)) \rightarrow \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}, \lambda'))$ is the inverse of $\Theta_{\lambda\lambda'}^\omega$ (which follows from (48), $\Theta_{\lambda\lambda}^\omega = \text{Id}$ and $\Theta_{\lambda'\lambda'}^\omega = \text{Id}$), it suffices to show that $\Theta_{\lambda\lambda'}^\omega$ does restrict to a homomorphism $\text{Frac}(\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda')) \rightarrow \text{Frac}(\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda))$.

Our strategy to show this is to use the splitting homomorphisms \mathcal{S}_e for the skew-fields of fractions of the mutable-balanced algebras, as in (39). We first show:

Lemma 4.10. *Let \mathfrak{S} be a triangulable pb surface, and λ be a triangulation of \mathfrak{S} . Let e be an edge of λ that is not a boundary edge. Let $\text{Cut}_e(\mathfrak{S})$ be the pb surface obtained from \mathfrak{S} by cutting along e , and let λ_e be the triangulation of $\text{Cut}_e(\mathfrak{S})$ induced by λ . Then, the splitting homomorphism $\mathcal{S}_e : \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}, \lambda)) \rightarrow \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\text{Cut}_e(\mathfrak{S}), \lambda_e))$ in (39) restricts to the homomorphism*

$$\mathcal{S}_e^{\text{bl}} : \text{Frac}(\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda)) \rightarrow \text{Frac}(\mathcal{Z}_\omega^{\text{bl}}(\text{Cut}_e(\mathfrak{S}), \lambda_e)).$$

Proof. Let $Z^{\mathbf{t}} \in \mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda) \subset \mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}, \lambda) \subset \mathcal{Z}_\omega(\mathfrak{S}, \lambda)$, with $\mathbf{t} = (t_v)_{v \in V_\lambda} \in \mathcal{B}_\lambda$, where $\mathcal{B}_\lambda \subset \mathbb{Z}^{V_\lambda}$ is as defined in (23). From (40) we have $\mathcal{S}_e(Z^{\mathbf{t}}) = Z^{\mathbf{t}'}$, with $\mathbf{t}' = (t'_v)_{v \in V_{\lambda_e}} \in \mathbb{Z}^{V_{\lambda_e}}$ given by the equation $t'_v = t_{\mathbf{pr}_e(v)}$, where \mathbf{pr}_e stands for the gluing projection map $\text{Cut}_e(\mathfrak{S}) \rightarrow \mathfrak{S}$ and also the induced map $V_{\lambda_e} \rightarrow V_\lambda$. Thus, for each ideal triangle τ of λ , if we denote by τ' the corresponding unique ideal triangle of λ_e , then \mathbf{pr}_e induces a bijection from $V_{\lambda_e} \cap \tau'$ to $V_\lambda \cap \tau$, and we have $t'_v = t_{\mathbf{pr}_e(v)}$ for all $v \in V_{\lambda_e} \cap \tau'$. In view of (23), the balancedness condition is described for each ideal triangle, and hence one can observe that the balancedness of \mathbf{t} (with respect to λ) implies that of \mathbf{t}' (with respect to λ_e). So $\mathcal{S}_e(Z^{\mathbf{t}}) = Z^{\mathbf{t}'} \in \mathcal{Z}_\omega^{\text{mbl}}(\text{Cut}_e(\mathfrak{S}), \lambda_e)$ lies in $\mathcal{Z}_\omega^{\text{bl}}(\text{Cut}_e(\mathfrak{S}), \lambda_e)$. \square

By using the splitting homomorphisms, we will see that proving Proposition 4.9 boils down to the case when \mathfrak{S} is a disjoint union of polygons. To settle the polygon case, we invoke the following lemma:

Lemma 4.11. [LY23, Theorem 11.7 and section 14.2] *Suppose that the pb surface \mathfrak{S} is a polygon \mathbb{P}_k with $k \geq 3$, and λ is a triangulation of \mathfrak{S} . Then the quantum trace map $\overline{\text{tr}}_\lambda : \overline{\mathcal{F}}_\omega(\mathfrak{S}) \rightarrow \mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda)$ induces an isomorphism*

$$\text{Frac}(\overline{\text{tr}}_\lambda) : \text{Frac}(\overline{\mathcal{F}}_\omega(\mathfrak{S})) \rightarrow \text{Frac}(\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda))$$

between the corresponding skew-fields of fractions.

Proposition 4.12. *Proposition 4.9 holds when \mathfrak{S} is a disjoint union of polygons.*

Proof. It is straightforward to see that it suffices to show the statement when \mathfrak{S} is a (single) polygon \mathbb{P}_k , with $k \geq 3$. Our Theorem 4.1 yields $\Theta_{\lambda\lambda'}^\omega(\text{Im}(\overline{\text{tr}}_{\lambda'})) \subset \text{Im}(\overline{\text{tr}}_\lambda)$. So Lemma 4.11 implies that $\Theta_{\lambda\lambda'}^\omega$ restricts to a homomorphism $\text{Frac}(\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda')) \rightarrow \text{Frac}(\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda))$. \square

We are now ready to prove Proposition 4.9.

Proof of Proposition 4.9. By the consistency (48), it suffices to prove the case when λ and λ' are related by a flip at an edge, say e . As in the proof of Theorem 4.1, we cut along all edges in E defined as in (49), i.e. along all non-boundary edges of λ not equal to e . The resulting pb surface is denoted by $\mathfrak{S}_E := \text{Cut}_E(\mathfrak{S})$, which is a disjoint union of a single copy of \mathbb{P}_4 which contains e and some number of copies of triangles \mathbb{P}_3 . Let λ_E and λ'_E be the triangulations of \mathfrak{S}_E induced by λ and λ' . Consider the following diagram

$$\begin{array}{ccccc}
 \text{Frac}(\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda')) & \overset{\exists(\Theta_{\lambda\lambda'}^\omega)^{\text{bl}}?}{\dashrightarrow} & & & \text{Frac}(\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda)) \\
 \downarrow \mathcal{S}_E^{\text{bl}} & \searrow \iota & & \swarrow \iota & \downarrow \mathcal{S}_E^{\text{bl}} \\
 & \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}, \lambda')) & \xrightarrow{\Theta_{\lambda\lambda'}^\omega} & \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}, \lambda)) & \\
 & \downarrow \mathcal{S}_E & & \downarrow \mathcal{S}_E & \\
 & \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}_E, \lambda'_E)) & \xrightarrow{\Theta_{\lambda_E\lambda'_E}^\omega} & \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}_E, \lambda_E)) & \\
 \downarrow \mathcal{S}_E^{\text{bl}} & \swarrow \iota & & \swarrow \iota & \downarrow \mathcal{S}_E^{\text{bl}} \\
 \text{Frac}(\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}_E, \lambda'_E)) & \xrightarrow{(\Theta_{\lambda_E\lambda'_E}^\omega)^{\text{bl}}} & & & \text{Frac}(\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}_E, \lambda_E))
 \end{array} \tag{61}$$

The inner square of the above diagram is the outermost square of the diagram in (50), hence is commutative. The four diagonal arrows denoted by ι are the natural embeddings. So the left and the right trapezoids are commutative, by Lemma 4.10. The bottom horizontal map comes from Proposition 4.12; so the bottom trapezoid commutes. The goal is to show that there exists the top horizontal map $(\Theta_{\lambda\lambda'}^\omega)^{\text{bl}}$ that makes the top trapezoid commutes; since ι are injective, such a map $(\Theta_{\lambda\lambda'}^\omega)^{\text{bl}}$ is unique if it exists. From now on, we may omit writing ι if it's clear from the context.

Let $Z^{\mathbf{t}'} \in \mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda') \subset \text{Frac}(\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda'))$, belonging to the upper left corner of the diagram; that is, $\mathbf{t}' = (t'_v)_{v \in V_{\lambda'}} \in \mathcal{B}_{\lambda'} \subset \mathbb{Z}^{V_{\lambda'}}$, where $\mathcal{B}_{\lambda'}$ is defined in (23). As $Z^{\mathbf{t}'} = \iota(Z^{\mathbf{t}'}) \in \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}, \lambda'))$, we may apply $\Theta_{\lambda\lambda'}^\omega$ and get an element $\Theta_{\lambda\lambda'}^\omega(Z^{\mathbf{t}'}) \in \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathfrak{S}, \lambda))$. All we need to check is whether $\Theta_{\lambda\lambda'}^\omega(Z^{\mathbf{t}'})$ lies in $\text{Frac}(\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda))$.

Recall from (31) again that the cluster seeds \mathcal{D}_λ and $\mathcal{D}_{\lambda'}$ are connected by the sequence of mutations at the vertices v_1, \dots, v_r . As in the proof of Lemma 4.8 (also Lemma 3.18), let $\mathcal{D}_0 = \mathcal{D}_\lambda$, and $\mathcal{D}_k = \mu_{v_k}(\mathcal{D}_{k-1})$ for $k = 1, \dots, r$. In particular, $\mathcal{D}_r = \mathcal{D}_{\lambda'}$. Let $\mathbf{t}^{(r)} := \mathbf{t}'$, and define $\mathbf{t}^{(k)}$, $k = r-1, \dots, 2, 1$, recursively by the equation $\nu'_{v_k}(Z^{\mathbf{t}^{(k)}}) = Z^{\mathbf{t}^{(k-1)}}$; see (56) for the formula of $\mathbf{t}^{(k-1)}$ in terms of

$\mathbf{t}^{(k)}$. Note that $\nu_{v_k}^\omega(Z^{\mathbf{t}^{(k)}}) = \nu_{v_k}^{\sharp\omega}(\nu_{v_k}'(Z^{\mathbf{t}^{(k)}})) = \nu_{v_k}^{\sharp\omega}(Z^{\mathbf{t}^{(k-1)}}) = Z^{\mathbf{t}^{(k-1)}} \cdot (*)_{k-1}$ for some nonzero element $(*)_{k-1}$ of $\text{Frac}(\mathcal{X}_q(\mathcal{D}_{k-1}))$, due to Lemma 3.7. Note also that the isomorphism $\nu_{v_k}^\omega : \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathcal{D}_k)) \rightarrow \text{Frac}(\mathcal{Z}_\omega^{\text{mbl}}(\mathcal{D}_{k-1}))$ restricts to an isomorphism $\Phi_{v_k}^q : \text{Frac}(\mathcal{X}_q(\mathcal{D}_k)) \rightarrow \text{Frac}(\mathcal{X}_q(\mathcal{D}_{k-1}))$ (Lemma 3.9). Thus we arrive at

$$\Theta_{\lambda\lambda'}^\omega(Z^{\mathbf{t}^{(r)}}) = \nu_{v_r}^\omega \cdots \nu_{v_1}^\omega(Z^{\mathbf{t}^{(r)}}) = \nu_{v_r}^\omega \cdots \nu_{v_2}^\omega(Z^{\mathbf{t}^{(r-1)}} \cdot (*)_{r-1}) = \cdots = Z^{\mathbf{t}^{(0)}} \cdot (*),$$

where $\mathbf{t}^{(0)} \in \mathbb{Z}^{V_\lambda}$ and $(*)$ is a nonzero element of $\text{Frac}(\mathcal{X}_q(\mathcal{D}_0)) = \text{Frac}(\mathcal{X}_q(\mathfrak{S}, \lambda))$. It remains to show that $\mathbf{t}^{(0)}$ is balanced, i.e. lies in \mathcal{B}_λ .

We now apply \mathcal{S}_{λ_0} to $Z^{\mathbf{t}^{(0)}} \cdot (*)$. Then, from the diagram,

$$\begin{aligned} \mathcal{S}_E(\Theta_{\lambda\lambda'}^\omega(\iota(Z^{\mathbf{t}'}))) &= \Theta_{\lambda_E\lambda'_E}^\omega(\mathcal{S}_E(\iota(Z^{\mathbf{t}'}))) \quad (\because \text{inner square}) \\ &= \Theta_{\lambda_E\lambda'_E}^\omega(\iota(\mathcal{S}_E^{\text{bl}}(Z^{\mathbf{t}'}))) \quad (\because \text{left trapezoid}) \\ &= \iota((\Theta_{\lambda_E\lambda'_E}^\omega)^{\text{bl}}(\mathcal{S}_E^{\text{bl}}(Z^{\mathbf{t}'}))) \quad (\because \text{bottom trapezoid}) \end{aligned}$$

Note that $\mathcal{S}_E^{\text{bl}}(Z^{\mathbf{t}'})$ is of the form $Z^{\mathbf{t}''}$ with $\mathbf{t}'' \in \mathcal{B}_{\bar{\lambda}'} \subset \mathbb{Z}^{V_{\bar{\lambda}'}}$, so by the similar arguments as $\Theta_{\lambda\lambda'}^\omega$ we see that $\iota((\Theta_{\lambda_E\lambda'_E}^\omega)^{\text{bl}}(\mathcal{S}_E^{\text{bl}}(Z^{\mathbf{t}'})))$ is of the form $Z^{\mathbf{t}'''} \cdot (**)$, with $\mathbf{t}''' \in \mathbb{Z}^{V_{\bar{\lambda}'}}$ and $0 \neq (**) \in \text{Frac}(\mathcal{X}_q(\mathfrak{S}', \bar{\lambda}'))$. As the image of $(\Theta_{\lambda_E\lambda'_E}^\omega)^{\text{bl}}$ lies in $\text{Frac}(\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}', \bar{\lambda}'))$, and since $\text{Frac}(\mathcal{X}_q(\mathfrak{S}', \lambda))$ is contained in $\text{Frac}(\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}', \bar{\lambda}'))$, it follows that $Z^{\mathbf{t}'''}$ lies in $\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}', \bar{\lambda}')$, i.e. \mathbf{t}''' lies in $\mathcal{B}_{\bar{\lambda}'}$. On the other hand, $\mathcal{S}_E(\Theta_{\lambda\lambda'}^\omega(\iota(Z^{\mathbf{t}'}))) = \mathcal{S}_E(Z^{\mathbf{t}^{(0)}}) \cdot \mathcal{S}_E((*))$; here $\mathcal{S}_E(Z^{\mathbf{t}^{(0)}}) = Z^{\mathbf{t}''''}$ for some $\mathbf{t}'''' \in \mathbb{Z}^{V_{\bar{\lambda}'}}$ and $\mathcal{S}_E((*))$ is a nonzero element of $\text{Frac}(\mathcal{X}_q(\mathfrak{S}', \bar{\lambda}'))$. By the equality $Z^{\mathbf{t}''''} \cdot \mathcal{S}_E((*)) = Z^{\mathbf{t}'''} \cdot (**)$, it follows that \mathbf{t}'''' also belongs to $\mathcal{B}_{\bar{\lambda}'}$. We use the arguments in the proof of Lemma 4.10. Since the balancedness condition for a Laurent monomial is described for each triangle in a triangulation, and since the splitting homomorphism \mathcal{S}_E preserves this condition, it follows that the balancedness of $Z^{\mathbf{t}''''} = \mathcal{S}_E(Z^{\mathbf{t}^{(0)}})$ implies that of $Z^{\mathbf{t}^{(0)}}$, i.e. we get $\mathbf{t}^{(0)} \in \mathcal{B}_\lambda$, as desired. This finishes the proof of Proposition 4.9. \square

The two remarks below easily follow.

Remark 4.13. *The balanced quantum coordinate change isomorphism $(\Theta_{\lambda\lambda'}^\omega)^{\text{bl}} : \text{Frac}(\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda')) \rightarrow \text{Frac}(\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda))$ obtained in Proposition 4.9 extends the map $\Phi_{\lambda\lambda'}^q : \text{Frac}(\mathcal{X}_q(\mathfrak{S}, \lambda')) \rightarrow \text{Frac}(\mathcal{X}_q(\mathfrak{S}, \lambda))$. (see Remark 3.15.)*

Remark 4.14. *The consistency equation holds (see (48)):*

$$(\Theta_{\lambda\lambda'}^\omega)^{\text{bl}} \circ (\Theta_{\lambda'\lambda''}^\omega)^{\text{bl}} = (\Theta_{\lambda\lambda''}^\omega)^{\text{bl}}$$

By collecting what we have proved, we obtain the following version of Theorem 4.1 written this time with respect to the coordinate change maps $(\Theta_{\lambda\lambda'}^\omega)^{\text{bl}}$ between the skew-fields of fractions of balanced algebras.

Theorem 4.15 (compatibility of quantum trace with balanced n -root quantum coordinate change maps $(\Theta_{\lambda\lambda'}^\omega)^{\text{bl}}$). *Let \mathfrak{S} be a triangulable pb surface. For any two*

triangulations λ and λ' of \mathfrak{S} , the following diagram commutes:

$$\begin{array}{ccc}
 & \overline{\mathcal{F}}_\omega(\mathfrak{S}) & \\
 \overline{\text{tr}}_{\lambda'} \swarrow & & \searrow \overline{\text{tr}}_\lambda \\
 \text{Frac}(\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda')) & \xrightarrow{(\Theta_{\lambda\lambda'}^\omega)^{\text{bl}}} & \text{Frac}(\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda)).
 \end{array} \tag{62}$$

That is, we have

$$\overline{\text{tr}}_\lambda = (\Theta_{\lambda\lambda'}^\omega)^{\text{bl}} \circ \overline{\text{tr}}_{\lambda'}. \tag{63}$$

4.4. A comparison with coordinate change isomorphisms defined by Lê and Yu. Suppose that \mathfrak{S} is a triangulable pb surface and λ, λ' are two triangulations of \mathfrak{S} . Lê and Yu also constructed a coordinate change isomorphism [LY23, Theorem 14.2]

$$\overline{\Psi}_{\lambda\lambda'}^X : \text{Frac}(\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda')) \rightarrow \text{Frac}(\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda))$$

such that the following diagram commutes:

$$\begin{array}{ccc}
 & \overline{\mathcal{F}}_\omega(\mathfrak{S}) & \\
 \overline{\text{tr}}_{\lambda'} \swarrow & & \searrow \overline{\text{tr}}_\lambda \\
 \text{Frac}(\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda')) & \xrightarrow{\overline{\Psi}_{\lambda\lambda'}^X} & \text{Frac}(\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda)).
 \end{array} \tag{64}$$

For any three triangulations $\lambda, \lambda', \lambda''$ of \mathfrak{S} , the consistency equation holds:

$$\overline{\Psi}_{\lambda\lambda'}^X \circ \overline{\Psi}_{\lambda'\lambda''}^X = \overline{\Psi}_{\lambda\lambda''}^X. \tag{65}$$

Here we briefly recall the construction of $\overline{\Psi}_{\lambda\lambda'}^X$ of Lê and Yu. As we mentioned in Lemma 4.11, in [LY23] Lê and Yu proved that the quantum trace map

$$\overline{\text{tr}}_\lambda : \overline{\mathcal{F}}_\omega(\mathfrak{S}) \rightarrow \mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda)$$

induces an isomorphism

$$\text{Frac}(\overline{\text{tr}}_\lambda) : \text{Frac}(\overline{\mathcal{F}}_\omega(\mathfrak{S})) \rightarrow \text{Frac}(\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda))$$

when \mathfrak{S} is a polygon. They define $\overline{\Psi}_{\lambda\lambda'}^X$ to be $\text{Frac}(\overline{\text{tr}}_\lambda) \circ \text{Frac}(\overline{\text{tr}}_{\lambda'})^{-1}$ when \mathfrak{S} is a polygon. Then they generalize the definition of $\overline{\Psi}_{\lambda\lambda'}^X$ by cutting \mathfrak{S} into a collection of polygons ([LY23, §14.3]).

We show that the Lê-Yu coordinate change map $\overline{\Psi}_{\lambda\lambda'}^X$ coincides with the balanced quantum cluster coordinate change map $(\Theta_{\lambda\lambda'}^\omega)^{\text{bl}}$ which we studied in the present paper. This answers the expectation stated in [LY23, §14.4], which provided the very motivation for the present paper.

Theorem 4.16 (Lê-Yu quantum coordinate change equals balanced quantum cluster coordinate change). *Suppose that \mathfrak{S} is a triangulable pb surface and λ, λ' are two triangulations of \mathfrak{S} . We have*

$$\overline{\Psi}_{\lambda\lambda'}^X = (\Theta_{\lambda\lambda'}^\omega)^{\text{bl}}.$$

Proof. Because of Remark 4.14 and equation (65), we can assume that λ' is obtained from λ by a flip on an edge $e \in \lambda$. When \mathfrak{S} is a polygon, Lemma 4.11 implies the theorem. That is, Lemma 4.11 implies that an isomorphism $\overline{\Psi}_{\lambda\lambda'}^X : \text{Frac}(\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda')) \rightarrow \text{Frac}(\mathcal{Z}_\omega^{\text{bl}}(\mathfrak{S}, \lambda))$ making the diagram in (64) is unique (see [LY23, Cor.14.3] for such a uniqueness), hence equals $(\Theta_{\lambda\lambda'}^\omega)^{\text{bl}}$, in view of Theorem 4.15.

When \mathfrak{S} is a general triangulable pb surface, we can use the trick as in the proofs of Theorem 4.1 and Proposition 4.9. We cut \mathfrak{S} into a \mathbb{P}_4 containing e and some \mathbb{P}_3 . Consider the outermost square of the diagram in (61), which is commutative. This diagram still commutes if we replace the top horizontal map by $\overline{\Psi}_{\lambda\lambda'}^X$ and the bottom horizontal map by $\overline{\Psi}_{\lambda_E\lambda'_E}^X$, because the definition of the Lê-Yu maps $\overline{\Psi}_{\lambda\lambda'}^X$ (see [LY23, equation (253)]) implies that they are compatible with the splitting homomorphisms. From the polygon case, we see that the original bottom horizontal map $(\Theta_{\lambda_E\lambda'_E}^\omega)^{\text{bl}}$ coincides with $\overline{\Psi}_{\lambda_E\lambda'_E}^X$.

Since the right vertical map $\mathcal{S}_E^{\text{bl}}$ in this original square diagram are injective, it follows that the top horizontal map that makes this diagram commutative is unique, if exists. By this uniqueness, it follows that $(\Theta_{\lambda\lambda'}^\omega)^{\text{bl}}$ coincides with $\overline{\Psi}_{\lambda\lambda'}^X$, as desired. \square

Remark 4.17. *The proof of Theorem 4.16 implies that $\overline{\Psi}_{\lambda\lambda'}^X = (\Theta_{\lambda\lambda'}^\omega)^{\text{bl}}$ as long as $\overline{\Psi}_{\lambda'\lambda}^X$ makes the diagram (64) commute and satisfies the equation (65). This shows the uniqueness of $\overline{\Theta}_{\lambda\lambda'}^\omega$ with properties stated in Theorem 4.1 and Remark 4.14.*

REFERENCES

- [AK17] D. G. L. Allegretti and H. Kim, *A duality map for quantum cluster varieties from surfaces*, Adv. Math. **306** (2017), 1164–1208.
- [BZ05] A. Berenstein and A. Zelevinsky, *Quantum cluster algebras*, Adv. Math. **195**(2) (2005), 405–455.
- [BW11] F. Bonahon and H. Wong, *Quantum traces for representations of surface groups in $SL_2(\mathbb{C})$* , Geom. Topol. **15**(3) (2011), 1569–1615.
- [BW16] F. Bonahon and H. Wong, *Representations of the Kauffman bracket skein algebra I: invariants and miraculous cancellations*, Invent. Math. **204** (2016), 195–243.
- [CKM14] S. Cautis, J. Kamnitzer, and S. Morrison, *Webs and quantum skew Howe duality*, Math. Ann. **360** (2014), 351–390.
- [CF99] L. Chekhov and V. V. Fock, *A quantum Teichmüller space*, Theor. Math. Phys. **120** (1999) 511–528.
- [CS23] L. O. Chekhov and M. Shapiro, *Log-canonical coordinates for symplectic groupoid and cluster algebras*, Int. Math. Res. Notices **2023** no. 11 (2023), 9565–9652.
- [C95] P. M. Cohn, *Skew fields: theory of general division rings*, Encyclopedia of Mathematics and its Applications **57**, Cambridge Univ. Press, 1995.
- [CL22] F. Costantino and T. T. Q. Lê, *Stated skein algebras of surfaces*, J. Eur. Math. Soc. **24**(12) (2022), 4063–4142.
- [D24] D. C. Douglas, *Quantum traces for $SL_n(\mathbb{C})$: The case $n = 3$* . J. Pure Appl. Algebra **228**(7) (2024), article no. 107652.
- [F95] L. D. Faddeev, *Discrete Heisenberg-Weyl group and modular group*, Lett. Math. Phys. **34** (1995), 249–254.

- [FK94] L. D. Faddeev and R. M. Kashaev, *Quantum dilogarithm*, Mod. Phys. Lett. **A9** (1994) 427–434.
- [F97] V. V. Fock, *Dual Teichmüller spaces*, preprint (1997). arXiv:hep-th/9702018
- [FG06] V. V. Fock and A. B. Goncharov, *Moduli spaces of local systems and higher Teichmüller theory*, Publ. Math. Inst. Hautes Études Sci. **103** (2006), 1–211.
- [FG09a] V. V. Fock and A. B. Goncharov, *Cluster ensembles, quantization and the dilogarithm*, Ann. Sci. Éc. Norm. Supér. (4) **42** (2009), no. 6, 865–930.
- [FG09b] V. V. Fock and A. B. Goncharov, *The quantum dilogarithm and representations of the quantum cluster varieties*, Invent. Math. **175**(2) (2009), 223–286.
- [G84] W. M. Goldman, *The symplectic nature of fundamental groups of surfaces*, Adv. Math. **54** (1984), 200–225.
- [GS19] A. B. Goncharov and L. Shen, *Quantum geometry of moduli spaces of local systems and representation theory*, arXiv:1904.10491
- [H10] C. Hiatt, *Quantum traces in quantum Teichmüller theory*, Algebr. Geom. Topol. **10**(3) (2010), 1245–1283.
- [K98] R. M. Kashaev, *Quantization of Teichmüller spaces and the quantum dilogarithm*, Lett. Math. Phys. **43** (1998) 105–115.
- [KN11] R. M. Kashaev and T. Nakanishi, *Classical and quantum dilogarithm identities*, SIGMA Symmetry Integrability Geom. Methods Appl. **7** (2011), article no. 102
- [K20] H. Kim, *SL_3 -laminations as bases for PGL_3 cluster varieties for surfaces*, arXiv:2011.14765 (2020).
- [K21a] H. Kim, *Phase constants in the Fock-Goncharov quantum cluster varieties*, Anal.Math. Phys. **11** (2021), no. 1, article no. 2
- [K21b] H. Kim, *Irreducible self-adjoint representations of quantum Teichmüller space and the phase constants*, J. Geom. Phys. **162** (2021), 104103.
- [K24] H. Kim, *Naturality of SL_3 quantum trace maps for surfaces*, Quantum Topol., online first (2024), arXiv:2104.06286.
- [KLS23] H. Kim, T. T. Q. Lê, and M. Son, *SL_2 quantum trace in quantum Teichmüller theory via writhe*, Algebr. Geom. Topol. **23**(1) (2023), 339–418.
- [KS] H. Kim and L. Shen, *On the canonicity of SL_3 - PGL_3 duality maps for cluster varieties for surfaces* (tentative title), in preparation.
- [K96] G. Kuperberg, *Spiders for Rank 2 Lie Algebras*, Commun. Math. Phys. **180** (1996), 109–151.
- [L09] D. Labardini-Fragoso, *Quivers with potentials associated to triangulated surfaces*, Proceedings of the London Mathematical Society **98**(3) (2009): 797–839.
- [L18] T. T. Q. Lê, *Triangular decomposition of skein algebras*, Quantum Topol. **9**(3) (2018), 591–632.
- [LS21] T. T. Q. Lê, A. S. Sikora, *Stated $SL(n)$ -Skein Modules and Algebras*, arXiv:2201.00045 (2021).
- [LY23] T. T. Q. Lê, T. Yu, *Quantum traces for SL_n -skein algebras*, arXiv:2201.00045 (2023).
- [LY22] T. T. Q. Lê, T. Yu, *Quantum traces and embeddings of stated skein algebras into quantum tori*, Selecta Math. (N.S.) **28** (2022), no. 4, Paper No. 66, 48 pp.
- [P22] A. Poudel, *A comparison between SL_n spider categories*, arXiv:2210.09289
- [P99] J. Przytycki, *Fundamentals of Kauffman bracket skein modules*, Kobe J. Math. **16** (1999) 45–66.
- [SS17] G. Schrader, and A. Shapiro *Continuous tensor categories from quantum groups I: algebraic aspects*, arXiv:1708.08107 (2017).
- [S01] A. S. Sikora, *SL_n -character varieties as spaces of graphs*, T. Am. Math. Soc. **35** no.7 (2001), 2773–2804.
- [S05] A. S. Sikora, *Skein theory for $SU(n)$ -quantum invariants*, Alg. Geom. Topol. **5** (2005), 865–897.

- [T89] V. G. Turaev, “Algebra of loops on surfaces, algebra of knots, and quantization” in *Braid group, knot theory and statistical mechanics*, Adv. Ser. Math. Phys. **9**, World Sci. Publ., Teaneck, NJ, 1989. pp.59–95.

SCHOOL OF MATHEMATICS, KOREA INSTITUTE FOR ADVANCED STUDY (KIAS), 85 HOEGI-RO,
DONGDAEMUN-GU, SEOUL 02455, REPUBLIC OF KOREA

Email address: `hkim@kias.re.kr`

ZHIHAO WANG, SCHOOL OF PHYSICAL AND MATHEMATICAL SCIENCES, NANYANG TECHNO-
LOGICAL UNIVERSITY, 21 NANYANG LINK SINGAPORE 637371

Email address: `ZHIHA0003@e.ntu.edu.sg`

UNIVERSITY OF GRONINGEN, BERNOULLI INSTITUTE, 9700 AK GRONINGEN, THE NETHER-
LANDS

Email address: `wang.zhihao@rug.nl`