## JONES–WENZL PROJECTIONS OF TYPE D AND DYCK TILINGS

KEIICHI SHIGECHI

Abstract. We study the relation between a coefficient of an element of the Jones–Wenzl projection in the Temperley–Lieb algebra of type  $D$  and an enumeration of Dyck tilings. The coefficient can be non-recursively expressed as an enumerative generating function of Dyck tilings by considering the generalized Hermite histories, which we call bi-colored vertical Hermite histories, on the tilings.

## 1. INTRODUCTION

The Jones–Wenzl projection [\[6,](#page-29-0) [8,](#page-29-1) [18\]](#page-29-2) is a special element of the Temperley–Lieb algebra [\[17\]](#page-29-3). This projection appears in many branches of mathematics and mathematical physics: it gives a way to calculate the Jones polynomial of a knot [\[7,](#page-29-4) [8\]](#page-29-1), graphical calculus of the canonical bases in tensor products of the quantum group  $U_q(\mathfrak{sl}_2)$  [\[4\]](#page-29-5), and Soergel calculus [\[2\]](#page-29-6). In this paper, we study the Jones–Wenzl projections of type  $D$  in terms of combinatorial objects called Dyck tilings. We prove that a coefficient in the Jones–Wenzl projection of type  $D$  can be non-recursively expressed as the generating function of Dyck tilings.

The Jones–Wenzl projections of type A were introduced by V. F. R. Jones in  $[6]$ , and the recursive formula was given by H. Wenzl in [\[18\]](#page-29-2). This recursive formula was further simplified by I. B. Frenkel and M. G. Khovanov in [\[4\]](#page-29-5), and a graphical interpretation of the same recursive formula was given by S. Morrison in [\[11\]](#page-29-7). In [\[13\]](#page-29-8), the author gave a combinatorial interpretation of the Jones–Wenzl projections of types A and B. More concretely, the coefficient of an element in the projection is given by the generating function of combinatorial objects called Dyck tilings. In [\[1\]](#page-29-9), J. Baine gives another interpretation of the coefficients of the projection in terms of Kazhdan–Lusztig polynomials, equivalently, the graded ranks of Soergel modules, for a wide class of types which includes types A and  $B$ , but not type  $D$ .

A Dyck tiling is a tiling of a region between two Dyck paths by use of Dyck tiles. Dyck tilings first appeared in the study of the Kazhdan–Lusztig polynomials for the Grassmannian permutations in [\[14\]](#page-29-10), and independently in the study of double-dimer models [\[9\]](#page-29-11). We have two types of Dyck tilings (type I and type II in  $[14]$ ), and we make use of only cover-inclusive Dyck tilings (corresponding to type I in  $[14]$ ) in this paper. In  $[10]$ , basic tools such as Dyck tiling strips, Dyck tiling ribbons, and Hermite histories, were established to study Dyck tilings. Among these tools, we generalize and make use of Hermite histories, which we call vertical Hermite histories, to give a combinatorial interpretation of the coefficients of elements in the Jones–Wenzl projection.

We start with the recursive relation for the Jones–Wenzl projection obtained by P. Sentinelli in  $[12]$ , which is the type D analogue of the Wenzl's recursive formula. Then, we simplify the recursive relation further along the spirit of Frenkel–Khovanov [\[4\]](#page-29-5) and Morrison [\[11\]](#page-29-7). By applying the method used by Morrison, we obtain two types of recursive relations in terms of the diagrams according to the parity. One of the main results of this paper is that these simplified recursive formulas can be interpreted in terms of Dyck tilings by imposing a simple condition on them, or by introducing a new combinatorial description of Hermite histories which we call bi-colored vertical

Date: December 24, 2024.

Hermite histories. The advantage of our method is that a coefficient in the Jones–Wenzl projections is given by an enumerative generating function of Dyck tilings.

It is well-known that the elements in the Temperley–Lieb algebra of type A correspond to  $n+1$ strand Temperley–Lieb diagrams which are diagrams with  $n+1$  marked points on the top and the bottom, with  $n+1$  non-intersecting strands connecting these marked points. Similarly, the elements in the Tempereley–Lieb algebra of type D correspond to the  $n+1$ -strand Temperley–Lieb diagrams with dots. The diagrams are classified into two classes according to the parity of the number of dots in the diagram. The coefficients of the elements in the projection behave differently according to the parity. Therefore, we have two types of recursive formulas for the coefficients depending on the parity. We discuss these two cases in what follows.

The Temperley–Lieb algebra  $TL_{n+1}^D$  of type D has the set of generators  $\{E_0, E_1, \ldots, E_n\}$  with some relations. If the number of dots in a diagram is even, this means that the algebraic representation of the diagram does not contain the product  $E_0E_1$  of the two generators. This condition is equivalent to considering the case where the product satisfies  $E_0E_1 = 0$ . Then, this case can be essentially reduced to the the type B case (see e.g.  $[16]$ ). Proposition [3.7](#page-9-0) also shows this fact since the form of the recursive relation for the projections of type D (under the condition  $E_0E_1 = 0$ ) is the same as types A and B which are studied in  $[13]$ . From these facts, the coefficients are expressed in terms of the generating functions of Dyck tilings which satisfy a simple condition (Theorem [4.9\)](#page-17-0).

If a diagram contains an odd number of dots, the algebraic representation of the diagram contains the product  $E_0E_1 \neq 0$ . The existence of the product  $E_0E_1$  is the main difficulty to obtain a combinatorial interpretation. Below, we explain two difficulties coming from the product.

Firstly, in [\[5\]](#page-29-15), R. M. Green introduced a graphical representation of the Temperley–Lieb algebra of type D. In this representation, we have to introduce an equivalence of the diagrams corresponding to  $E_1E_0$  and  $E_0E_1$  which naively give different diagrams although we have  $E_1E_0 = E_0E_1$ . Instead of the equivalence relations, we introduce a similar, but more straightforward graphical representation. When a diagram contains the product  $E_0E_1$ , we give a unique dot in the diagram. This is welldefined since if the diagram contains the product  $E_0E_1$ , then it does not contain any other  $E_0$  and  $E_1$  in its algebraic representation. We regard the unique dot as a landmark which indicates that the diagram contains the product  $E_0E_1$ . A diagram satisfies simple conditions (P1) and (P2) as in Section [2.2.](#page-2-0)

Secondly, if we derive the recurrence relation for a diagram with a unique dot, then this recurrence relation contains minus signs (Proposition [3.10\)](#page-11-0). The minus signs prevent us to make a combinatorial interpretation since we want to connect a coefficient in the projections with a generating function which enumerates combinatorial objects without minus signs. To overcome this point, we study the relation between the elements of type  $A$  and type  $D$ . If we take an appropriate linear combination of elements of type  $D$ , then this linear combination satisfies the recursive relation for type  $A$  (Section [3.4\)](#page-11-1). This allows us to rewrite the recursive relation into another recursive relation which does not have minus signs. This new recursive relation naturally gives a combinatorial description of an element in the projection in terms of Dyck tilings satisfying simple conditions  $(Q1)$  to  $(Q4)$  in Section [4.5.](#page-19-0)

By combining the above two mentioned prescriptions, the recurrence relation becomes minus sign free, however, it involves the generating function of not only type  $D$ , but also type  $A$  (Proposition [3.15\)](#page-13-0). We introduce new combinatorial objects on a Dyck tiling, which we call bi-colored vertical Hermite histories. The bi-colored vertical Hermite histories fit well to this situation, and give a combinatorial interpretation of the coefficients in the projections (Theorem [4.16\)](#page-20-0). A coefficient is expressed as a generating function of Dyck tilings using the bi-colored vertical Hermite histories. Especially, this combinatorial description is enumerative and non-recursive.

The paper is organized as follows. In Section [2,](#page-2-1) we introduce the Temperley–Lieb algebra of type D, the graphical representation, and Jones–Wenzl projections. We simplify the recursive relation of the projections, and give diagrammatic recursive relations in Section [3.](#page-6-0) In Section [4,](#page-14-0) after introducing Dyck tilings and vertical Hermite histories, we give a combinatorial interpretation of Jones–Wenzl projections in terms of bi-colored vertical Hermite histories. Some explicit expressions of the coefficients in the Jones–Wenzl projections are given in Section [5.](#page-23-0) In Appendix [A,](#page-28-0) we give the explicit expression for the Jones–Wenzl projection of type D of rank four.

**Notation.** We will use the standard notation for q-integers:  $[n] := (q^n - q^{-n})/(q - q^{-1})$ .

<span id="page-2-1"></span>Acknowledgement. The author gratefully acknowledges Byung-Hak Hwang for collaboration at the early stage of this work and valuable comments on the manuscript.

2. Temperley–Lieb algebra of type D

2.1. **Definition.** The Dynkin diagram of type  $D$  is depicted as follows:



We take the convention that the Dynkin diagram of type  $D_{n+1}$  is of rank  $n+1$ , and the labels of the nodes are  $0, 1, 2, ..., n$ . We have  $D_1 = A_1, D_2 = A_1 \times A_1$ , and  $D_3 = A_3$ .

The Temperley–Lieb algebra  $TL_{n+1}^D$  of type D is a  $\mathbb{C}(q)$ -algebra with generators  $\{E_0, E_1, \ldots, E_n\}$ which satisfy the following relations:

$$
E_i^2 = -[2]E_i, \t\forall i \in \{0, 1, ..., n\}
$$
  
\n
$$
E_i E_j E_i = E_i, \t\text{if } i \text{ and } j \text{ are adjacent in the Dynkin diagram } D_{n+1},
$$
  
\n
$$
E_i E_j = E_j E_i, \t\text{if } i \text{ and } j \text{ are not adjacent in the Dynkin diagram } D_{n+1}.
$$

The Temperley–Lieb algebra  $TL_{n+1}^A$  of type A is generated by the subset  $\{E_1, E_2, \ldots, E_n\}$  or  $\{E_0, E_2, \ldots, E_n\}$  of generators. It is well-known that the dimension of  $\mathrm{TL}_n^A$  is given by the Catalan number  $C_n := \frac{1}{n+1} \binom{2n}{n}$  $n_n^{2n}$  = 1, 2, 5, 14, .... Similarly, the dimension of TL<sub>n</sub><sup>D</sup> is given by

<span id="page-2-2"></span>(2.1) 
$$
\dim TL_n^D = \frac{n+3}{2}C_n - 1.
$$

The dimension of  $TL_n^D$  is equal to the number of fully commutative elements of type D studied in  $[3, 5, 15]$  $[3, 5, 15]$  $[3, 5, 15]$  $[3, 5, 15]$ . In Section [4.2,](#page-15-0) we derive the formula  $(2.1)$  by use of the correspondence between a Temperley–Lieb diagram and another diagram called a chord diagram.

Let  $w := (s_1, \ldots, s_l)$  be a sequence of integers such that  $s_i \in [0, n]$ . We call  $E_w := E_{s_1} \ldots E_{s_l}$  in  $TL_{n+1}^D$  a word. A word  $E_w$  is said to be a reduced expression if  $E_w$  can not be written with less than l generators. The integer l is the length of  $E_w$ .

<span id="page-2-0"></span>2.2. Graphical representation. We first recall the graphical representation of  $TL_{n+1}^A$ . An  $n+1$ strand Temperley–Lieb diagram is a diagram with  $n+1$  stands such that it has  $n+1$  marked points on the top and on the bottom, and  $n + 1$  arcs joining these marked points are non-intersecting. An arc joining two marked points on the top (resp. on the bottom) is called a *cup* (resp. a *cap*). A cup c is called inner-most cup if there is no smaller cup inside c. A cup is called outer-most if

there is no larger cup outside c and no vertical strands left to  $c$ . An inner-most or outer-most cap is similarly defined. We also say that a vertical strand which connects the right-most points on the top and bottom is an inner-most cap. A vertical strand which is left-most in a diagram is said to be an outer-most cap.

An example of Temperley–Lieb diagram is given in Figure [2.1.](#page-3-0) This diagram has three caps and



<span id="page-3-0"></span>Figure 2.1. An example of 8-strand Temperley–Lieb diagram

three cups. The cup connecting the first and second points is both inner-most and outer-most. The cup connecting the fourth and seventh points is neither inner-most nor outer-most. The cup connecting the fifth and sixth points is inner-most. We have two vertical strands which connect points on the bottom and on the top. The left-most vertical strand is an outer-most cap.

Let  $D_1$  and  $D_2$  be n+1-strand Temperley–Lieb diagrams of type A. Then, the product  $D = D_1D_2$ of  $D_1$  and  $D_2$  is given by putting  $D_1$  on the top of  $D_2$ . If there is a loop in D, we give a factor  $-[2]$ and delete the loop from the diagram. For example, two diagrams  $D_1$  and  $D_2$ 



give the diagram  $D = D_1 D_2$ :



To consider  $TL_{n+1}^D$ , we generalize the notion of the  $n+1$ -strand Temperley–Lieb diagrams. For type  $D$ , we allow to put a "dot" on a cup, a cap and a vertical strand. A diagram with dots satisfies the following properties:

(P1) The number of dots in a diagram is one, or an even non-negative integer. A cap, a cup, or a vertical strand contains at most one dot.

(P2) We consider the following three cases:

- (a) The number of dots is even. A dot is on an outer-most cap, outer-most cup or a left-most vertical strand.
- (b) The number of dots is one. A dot is on a cap or on a vertical strand, which connects the left-most marked point on the bottom and another marked point.
- (c) The diagram consisting of  $n + 1$  vertical strands (without caps and cups) does not have a dot.

We do not consider a diagram  $D$  such that the number of dots in  $D$  is an odd number greater than one.

Following [\[5\]](#page-29-15), we introduce a graphical representation for  $TL_{n+1}^D$ . The generator  $E_i$ ,  $1 \leq i \leq n$ is depicted as

$$
E_i = \left| \begin{array}{c} \cdots \\ \cdots \\ \cdots \end{array} \right| \left| \begin{array}{c} \diagdown \\ \diagdown \\ \diagdown \cdots \end{array} \right| \left| \begin{array}{c} \cdots \\ \cdots \\ \cdots \end{array} \right|
$$

and the generator  $E_0$  is depicted as

$$
E_0=\begin{array}{c|c} \cdot\cdot\cdot & \cdot \\ \hline 1 & 2 & 3 & n+1 \end{array}
$$

The identity 1 corresponds to the diagram with  $n + 1$  vertical strands without dots. The property (P2c) implies that there is no diagram which has the same connectivity as 1 and has a dot. Note that only the generator  $E_0$  has two dots, and other generators do not have dots in the graphical representation. The generators  $E_1$  and  $E_0$  have the same connectivity of marked points, and we distinguish between  $E_0$  and  $E_1$  by the existence of dots.

In the case of type A, we have a graphical calculation of an element  $E_w$  which may not be a reduced expression. If w is not a reduced expression, we use the algebraic relations  $E_iE_{i\pm 1}E_i =$  $E_i$  and  $E_i^2 = -[2]E_i$  to obtain a reduced expression. The former relation does not change the connectivity of a diagram, but the latter relation produces a loop. The weight  $-[2]$  of a loop corresponds to the factor in the relation. In the case of type  $D$ , we impose the properties (P1) and  $(P2)$  on a diagram. Due to these properties, we can not have a graphical calculation of  $E_w$  which is not a reduced expression. In what follows, we consider the graphical representation of a reduced expression in  $TL_{n+1}^D$ .

An element of  $TL_{n+1}^D$  is given by a product of generators. Suppose that a reduced expression  $E_w := E_{s_1} \dots E_{s_m}, s_i \in [0, n]$  is written as a product  $E_w = E_{w_1} E_{w_2}$  of two reduced expressions  $E_{w_1}$  and  $E_{w_2}$ . In other words, we have  $l(w) = l(w_1) + l(w_2)$ . Let D, D<sub>1</sub> and D<sub>2</sub> be diagrams corresponding to w,  $w_1$  and  $w_2$ . Then, the diagram D is obtained from  $D_1$  and  $D_2$  by putting  $D_1$ on the top of  $D_2$ , and by applying the rule (P2b) to D if D has an odd number of dots.

The graphical representation of generators implies that we have a loop without dots if we have  $E_i^2 = -[2]E_i, i \ge 1$ , in its algebraic representation. As far as we consider a reduced expression, we do not have such a loop. Similarly, we have no loops with two dots since such a loop appears if and only if the expression contains  $E_0^2 = -[2]E_0$ . A vertical strand, a cup, or a cap contains two dots if  $E_w$  contains the product  $E_0E_2E_0$  in its algebraic representation. Since we consider only reduced expressions, we do not have such a vertical strand, a cup or a cap. From these, it is clear that a vertical strand, a cup or a cap contains at most one dot.

We may have a loop with a single dot in the diagram D. Such a loop comes from the product of two generators  $E_0$  and  $E_1$  in the algebraic representation. In this case, we simply delete the loop with a dot from the diagram. Then, since a deletion of a loop with a dot implies that we have an odd number of dots in the diagram, we change the position of the dots such that it satisfies the condition (P2b). In fact, if a reduced expression  $E_w$  contains the product  $E_0E_1$  (or equivalently  $E_1E_0$ , we have no other  $E_0$  and  $E_1$  in its algebraic representation. This means that if the number of dots in a diagram is odd, then we have a unique dot which comes from the product  $E_0E_1$ . Therefore, the property (P2b) is well-defined.

In this way, we obtain a graphical representation D of a reduced word  $E_w$  in  $TL_{n+1}^D$ .

Remark 2.2. Three remarks are in order:

- (1) Consider the product  $E_0E_1$  of two generators. By the defining relations, we have  $E_0E_1 =$  $E_1E_0$ . If we apply a naive representation by diagrams,  $E_0E_1$  gives a diagram with a dotted cup and a loop with a single dot. On the other hand,  $E_1E_0$  gives a diagram with a dotted cap and a loop with a single dot. However, the two products  $E_1E_0$  and  $E_0E_1$  define the same element in  $\mathrm{TL}_{n+1}^D$ , we need to introduce an equivalence relation on diagrams for  $E_1E_0$ and  $E_0E_1$ . This is achieved in [\[5\]](#page-29-15).
- (2) In our convention, to distinguish a diagram with a single dot from diagrams with an even number of dots, we introduce the rule (P2b). If an element E' in  $TL_{n+1}^D$  contains the product  $E_0E_1$ , then it is easy to see that E' contains no other  $E_0$  or  $E_1$  if E' is a reduced expression. This means that if the number of dots is odd in a diagram D, then D contains a unique dot since we have the generator  $E_0$  and we delete one of the two dots by deleting a loop with a single dot. We regard a unique dot as a sign of the parity of the number of dots.
- (3) By the property (P2b), we depict

$$
E_0E_1=\begin{array}{c|c}\n\diagdown\ddots\\
1&2&3&n+1\n\end{array}
$$

The product  $E_0E_1$  has the same connectivity of the strands as  $E_0$  and  $E_1$ , and has a unique dot on the unique cap.

.

<span id="page-5-0"></span>**Example 2.3.** The graphical representations of  $A = E_2 E_0 E_1$  and  $B = E_0 E_2 E_3 E_1 E_2 E_0$  in  $TL_4^D$ are given by



2.3. Jones–Wenzl projections. We summarize the properties which characterize the Jones– Wenzl projection of types A and D. The Jones–Wenzl projection  $P_n$ ,  $0 \leq n$ , is a unique element in  $TL_{n+1}^A$  such that

- $(1)$   $P_n \neq 0;$
- (2)  $P_n P_i = P_i P_n = P_n$  for  $0 \le i \le n$ ;
- (3)  $E_i P_n = P_n E_i = 0$  for  $1 \le i \le n$ .

These three conditions uniquely fix the element  $P_n$  for  $0 \leq n$ .

Similarly, the Jones–Wenzl projection  $Q_n$ ,  $0 \leq n$ , of type D is a unique element in  $TL_{n+1}^D$  such that

(1) 
$$
Q_n \neq 0
$$
;  
(2)  $Q_n Q_i = Q_i Q_n = Q_n$  for  $0 \leq i \leq n$ ;

(3)  $E_i Q_n = Q_n E_i = 0$  for  $0 \le i \le n$ .

The second conditions for  $P_n$  and  $Q_n$  insure that the coefficient of the identity 1 is one.

**Example 2.4.** We give first few explicit expressions of  $P_n$  and  $Q_n$ :

$$
P_0 = 1, \qquad P_1 = 1 + \frac{1}{[2]} E_1,
$$
  

$$
P_2 = 1 + \frac{[2]}{[3]} (E_1 + E_2) + \frac{1}{[3]} (E_1 E_2 + E_2 E_1),
$$

and

$$
Q_0 = 1 + \frac{1}{[2]} E_0, \qquad Q_1 = 1 + \frac{1}{[2]} (E_0 + E_1) + \frac{1}{[2]^2} E_0 E_1,
$$
  
\n
$$
Q_2 = 1 + \frac{[3]}{[4]} (E_0 + E_1) + \frac{[2]^2}{[4]} E_2 + \frac{[2]}{[4]} (E_0 E_2 + E_2 E_0 + E_1 E_2 + E_2 E_1)
$$
  
\n
$$
+ \frac{[2]^3}{[4][3]} E_0 E_1 + \frac{1}{[4]} (E_0 E_2 E_1 + E_1 E_2 E_0)
$$
  
\n
$$
+ \frac{[2]^2}{[4][3]} (E_0 E_1 E_2 + E_2 E_0 E_1) + \frac{[2]}{[4][3]} E_2 E_0 E_1 E_2.
$$

Since  $D_3 = A_3$ , the projection  $P_3$  is expressed in terms of  $Q_2$  by replacing  $E_0$  with  $E_3$ . The explicit expression of  $Q_3$  is given in Appendix [A.](#page-28-0)

## 3. Recurrence relations for Jones–Wenzl projections

<span id="page-6-0"></span>3.1. **Type** A. In [\[18\]](#page-29-2), Wenzl gave a recurrence relation for  $P_n$ :

$$
P_n = P_{n-1} + \frac{[n]}{[n+1]} P_{n-1} E_n P_{n-1},
$$

with the initial condition  $P_0 = 1$ .

In  $[4, 11]$  $[4, 11]$ , the above recurrence relation is further simplified. For this, we define the elements in  $\mathrm{TL}_{n+1}^A$  by  $g_{n,i}^A = E_n E_{n-1} \dots E_i$  for  $1 \leq i \leq n$ , and  $g_{n,n+1}^A = 1$ . Note that the diagram corresponds to  $g_{n,n+1}^A$  is the diagram consisting of  $n+1$  vertical strands. Then, we have the following proposition:

**Proposition 3.1** ([\[11,](#page-29-7) Proposition 3.3]). The coefficient of  $g_{n,i}^A$ ,  $1 \le i \le n+1$ , in  $P_n$  is  $\frac{[i]}{[n+1]}$ , and  $P_n$  satisfies the recurrence relation:

$$
P_n = P_{n-1} \left( \sum_{i=1}^{n+1} \frac{[i]}{[n+1]} g_{n,i}^A \right).
$$

3.2. Type D. To state the recurrence relation for  $Q_n$ , we introduce some notations. We define the element  $E_{w_n}$  in  $TL_{n+1}^D$  by

$$
E_{w_n} := E_n E_{n-1} \cdots E_2 E_0 E_1 E_2 \cdots E_n,
$$

and define two coefficients

$$
A_n := \begin{cases} \frac{1}{[2]}, & \text{if } n = 1, \\ \frac{[n][2n-2]}{[2n][n-1]}, & \text{otherwise} \end{cases}, \qquad B_n := \frac{[n]}{[2n][n+1]}.
$$

<span id="page-6-2"></span>**Proposition 3.2** ([\[12,](#page-29-13) Proposition 4.8]). The generalized Jones–Wenzl projection  $Q_n$  satisfies the following recurrence relation:

<span id="page-6-1"></span>(3.1) 
$$
Q_n = Q_{n-1} + A_n Q_{n-1} E_n Q_{n-1} + B_n Q_{n-1} E_{w_n} Q_{n-1}.
$$

To simplify Eq. [\(3.1\)](#page-6-1) further, we first define some elements in  $TL_{n+1}^D$ . For  $i = 1, 2, ..., n+1$  and  $j = 0, 1, \ldots, n$ , we define

$$
g_{n,i} := \begin{cases} 1, & \text{if } i = n+1, \\ E_n E_{n-1} \cdots E_i, & \text{if } i \le n, \end{cases}
$$

$$
h_{n,j} := \begin{cases} E_n E_{n-1} \cdots E_2 E_0, & \text{if } j = 0, \\ E_n E_{n-1} \cdots E_2 E_0 E_1 E_2 \cdots E_j, & \text{if } j \ge 1. \end{cases}
$$

<span id="page-7-0"></span>Note that we have  $h_{n,n} = E_{w_n}$ .

**Proposition 3.3.** The projections  $Q_n$  satisfy the following recurrence relation:

<span id="page-7-1"></span>(3.2) 
$$
Q_{n+1} = Q_n \left( \sum_{i=1}^{n+2} \mathrm{coef}(g_{n+1,i}) g_{n+1,i} + \sum_{j=0}^{n+1} \mathrm{coef}(h_{n+1,j}) h_{n+1,j} \right),
$$

where the two coefficients are given by

<span id="page-7-3"></span>(3.3) 
$$
\operatorname{coef}(g_{n,i}) := \begin{cases} \frac{[n]}{[2n]}, & if \ i = 1, \\ \frac{[n][2i-2]}{[2n][i-1]}, & otherwise, \end{cases} \qquad \operatorname{coef}(h_{n,j}) := \frac{[n][n+1-j]}{[2n][n+1]},
$$

for  $1 \le i \le n+1$  and  $0 \le j \le n$ .

Before proceeding with the proof of Proposition [3.3,](#page-7-0) we write down the actions of  $E_{n+1}$  and  $E_{w_{n+1}}$  on  $g_{n,i}$  and  $h_{n,j}$ .

<span id="page-7-2"></span>Lemma 3.4. We have

$$
E_{n+1}g_{n,i} = g_{n+1,i}, \t for 1 \le i \le n+1,
$$
  
\n
$$
E_{n+1}h_{n,j} = h_{n+1,j}, \t for 0 \le j \le n,
$$
  
\n
$$
E_{w_{n+1}}g_{n,i} = h_{n+1,i}, \t for 1 \le i \le n+1,
$$
  
\n
$$
E_{w_{n+1}}h_{n,0} = h_{n+1,1},
$$
  
\n
$$
E_{w_{n+1}}h_{n,j} = -[2]h_{n+1,j}, \t for 1 \le j \le n.
$$

*Proof.* We calculate them by use of the definitions of  $g_{n,i}$  and  $h_{n,j}$ , and the defining relations of  $TL_{n+1}^D$ .  $D$ <sub>n+1</sub>.

*Proof of Proposition [3.3.](#page-7-0)* We prove the statement by induction on n following the method developed in [\[11\]](#page-29-7). For  $n = 1$ , one can show Eq. [\(3.2\)](#page-7-1) by a simple calculation. We assume that Eq. (3.2) holds up to  $n-1$ .

Let  $\mathcal{K}_n$  be the linear span of the elements  $g_{n,i}$  and  $h_{n,j}$ , and  $\mathcal{K}_n^{\perp}$  the linear span of the other elements. Then, an element  $D \in \mathcal{K}_n^{\perp}$  can be written as  $D = E_j D'$  with some  $j \in \{0, 1, 2, \ldots, n-1\}$ and D'. This implies that  $Q_n E_{n+1} \mathcal{K}_n^{\perp} = 0$  since we have  $E_{n+1} E_j = E_j E_{n+1}$  and  $Q_n E_j = 0$ by definition of  $Q_n$ . The projection  $Q_n$  is decomposed as  $Q_n = Q_n^{\mathcal{K}} + Q_n^{\mathcal{K}^{\perp}}$  with  $Q_n^{\mathcal{K}} \in \mathcal{K}_n$  and  $Q_n^{\mathcal{K}^{\perp}} \in \mathcal{K}_n^{\perp}$ . We have

$$
Q_n E_{n+1} Q_n = Q_n E_{n+1} (Q_n^{\mathcal{K}} + Q_n^{\mathcal{K}^\perp}) = Q_n E_{n+1} Q_n^{\mathcal{K}},
$$
  

$$
Q_n E_{w_{n+1}} Q_n = Q_n E_{w_{n+1}} Q_n^{\mathcal{K}}.
$$

By combining these together, we have

<span id="page-8-0"></span>(3.4) 
$$
Q_{n+1} = Q_n + A_{n+1}Q_n E_{n+1} Q_n + B_{n+1}Q_n E_{w_{n+1}} Q_n,
$$

$$
= Q_n + A_{n+1}Q_n E_{n+1} Q_n^{\mathcal{K}} + B_{n+1}Q_n E_{w_{n+1}} Q_n^{\mathcal{K}}.
$$

Since  $Q_n^{\mathcal{K}} \in \mathcal{K}_n$ ,  $Q_n$  has an expression of the following form:

<span id="page-8-1"></span>(3.5) 
$$
Q_n^{\mathcal{K}} = \sum_i \text{coef}'(g_{n,i}) g_{n,i} + \sum_j \text{coef}'(h_{n,j}) h_{n,j}.
$$

with some coefficients coef' $(g_{n,i})$  and coef' $(h_{n,j})$ . From Eqs. [\(3.4\)](#page-8-0) and [\(3.5\)](#page-8-1), Lemma [3.4](#page-7-2) and the induction hypothesis, we obtain recursive formulas for  $\text{coef}'(g_{n,i})$ ,  $\text{coef}'(h_{n,j})$ ,  $\text{coef}(g_{n,i})$  and  $\mathrm{coef}(h_{n,j})$ . These recursive formulas can be solved as

$$
\mathrm{coef}(g_{n,i}) = \mathrm{coef}'(g_{n,i}) = \frac{[n][2i-2]}{[2n][i-1]}, \qquad \mathrm{coef}(h_{n,j}) = \mathrm{coef}'(h_{n,j}) = \frac{[n][n+1-j]}{[2n][n+1]},
$$

which implies Eq.  $(3.3)$ . By substituting these into Eq.  $(3.4)$  and rearranging the terms, we obtain Eq.  $(3.2)$ , which completes the proof.

<span id="page-8-3"></span>3.3. Recurrence relation via diagrams. To derive the recurrence relation via diagrams, we first consider the graphical representation of the elements  $g_{n,i}$  and  $h_{n,j}$  with  $1 \leq i \leq n+1$  and  $0 \leq j \leq n$ . The element  $g_{n,i}$ ,  $1 \leq i \leq n$  has a cup connecting the right-most point with the point next to it on the top, a cap connecting the *i*-th point with the  $i + 1$ -th point on the bottom, and  $n-1$  vertical strands. The element  $g_{n,n+1}$  consists of  $n+1$  vertical strands. Similarly, the element  $h_{n,j}, 1 \leq j \leq n$ , has a cup connecting the right-most point with the point next to it on the top, a cap connecting the j-th point with the j + 1-th point on the bottom, and  $n-1$  vertical strands. Further, a cap or a strand which contains the left-most point in the bottom has a dot. The element  $h_{n,0}$  has the same connectivity as  $g_{n,1}$ , and the unique cap and the left-most vertical strands have a dot. These elements  $g_{n,i}$  and  $h_{n,j}$  satisfy the properties (P1) and (P2). In Figure [3.5,](#page-8-2) we depict the elements  $g_{3,i}$  and  $h_{3,i}$ .



<span id="page-8-2"></span>FIGURE 3.5. The graphical representation of  $g_{3,i}$  and  $h_{3,i}$ .

We recall the recurrence relation via diagrams for  $TL_{n+1}^A$ . In [\[11\]](#page-29-7) (see also [\[4\]](#page-29-5)), Morrison gave a recursive formula for the coefficients of the elements, equivalently for  $n + 1$ -strand Temperley–Lieb diagrams.

By definition, note that the vertical strand which connects the right-most points on the top and on the bottom is called an inner-most cap. Any  $n+1$ -strand Temperley–Lieb diagram has at least one inner-most cap. This property allows us to have a recurrence relation via diagrams.

Let D be an  $n + 1$ -strand Temperley–Lieb diagram. Suppose that an inner-most cap c in D connects the *i*-th bottom point from left and  $i + 1$ -th bottom point from left for  $1 \leq i \leq n$ . Then, the position of c is defined as i. If the right-most points on the top and on the bottom in  $D$  are connected by a vertical strand, the position of this inner-most cap is defined as  $n + 1$ .

<span id="page-9-3"></span>We first recall the recurrence relation the coefficients of the projection for type A.

**Proposition 3.6** (Proposition 4.1 in [\[11\]](#page-29-7)). Let D be an  $n+1$ -strand Temperley–Lieb diagram. Let  $I(D) \subseteq [1, n + 1]$  be the set of positions of inner-most caps in D, and  $D_i$  the n-strand Temperley-Lieb diagram obtained from D by removing the *i*-th inner-most cap. The coefficient  $\text{coef}_n^A(D)$  of D in  $P_n$  satisfies the following recurrence relation:

<span id="page-9-2"></span>(3.6) 
$$
\qquad \qquad \text{coef}_n^A(D) = \sum_{i \in I(D)} \frac{[i]}{[n+1]} \text{coef}_{n-1}^A(D_i).
$$

In what follows, we will derive the recurrence relation via diagrams for  $TL_{n+1}^D$ . Let D be an  $n + 1$ -strand Tempereley–Lieb diagram possibly with dots. For convenience, we simply write the coefficient coef<sub>n+1</sub>(D) of D in  $Q_{n+1}$  as D.

First, we consider the case where D contains an even number of dots. We define two sets  $I^{\bullet}(D)$ and  $I^{\circ}(D)$ . We define the set  $I^{\bullet}(D) := \{1\}$  if there exists a dotted inner-most cap at the position one and  $I^{\bullet}(D) = \emptyset$  otherwise. Similarly, the set  $I^{\circ}(D)$  is defined as the set of the positions of inner-most caps without a dot. Then, we define  $I(D) := I^{\bullet}(D) \cup I^{\circ}(D)$ .

<span id="page-9-0"></span>**Proposition 3.7.** Suppose that D has a diagram consisting of  $n + 1$  strands with an even number of dots. Let  $I(D)$  be the set as above. We denote by  $D_i$ , the n-strand diagram obtained from D by removing the *i*-th inner-most cap. Further, if  $i = 1 \in I^{\bullet}(D)$ , we add a dot on a cap or a vertical strand which contains the left-most point at the bottom in  $D_i$ . Then, we have

<span id="page-9-1"></span>(3.7) 
$$
D = \sum_{i \in I(D)} \mathrm{coef}(g_{n,i}) D_i.
$$

We can recursively apply Eq. [\(3.7\)](#page-9-1) to the diagram  $D_i$  since  $D_i$  has an even number of dots.

Remark 3.8. Four remarks are in order:

- (1) The diagram  $D$  has an even number of dots. Similarly, a diagram  $D_i$  also has an even number of dots. This is because we add a dot to a diagram after the removal of an innermost cap with a dot at position one.
- (2) Suppose we have two dots on a cap or a vertical strand after we add a dot on the cap or the vertical strand. Then, we delete these two dots.
- (3) If an inner-most cap at position  $i \neq 1$  has a dot, this inner-most cap cannot be removed to produce a diagram of smaller size. This comes from the fact that we have no generator which has the same connectivity as  $E_i$ ,  $i \geq 2$ , and has a dotted cap and a dotted cup. This is the reason why we have  $I^{\bullet}(D) = \{1\}$  if there is a dotted cap at position one, and  $I^{\bullet}(D) = \emptyset$ otherwise.
- (4) Suppose that  $1 \in I(D)$ . Then, we have an inner-most cap with or without a dot at the position one. Since we have  $\text{coef}(h_{n,0}) = \text{coef}(g_{n,1})$  by definition, we do not need to distinguish these two coefficients.

*Proof of Proposition [3.7.](#page-9-0)* Recall we have the recurrence relation  $(3.2)$  in Proposition [3.3.](#page-7-0) Let D be a diagram with  $n + 1$  strands and with an even number of dots. This implies that the algebraic representation of D does not contain the product  $E_0E_1$ . Recall that the elements  $h_{n,j}$  with  $j \geq 1$ contain the product  $E_0E_1$ . If we multiply  $h_{n,j}$  by any  $g_{m,i'}$  or  $h_{m,j'}$  with  $m \leq n-1$ , the result

also contains the product of  $E_0E_1$ . This fact and Eq. [\(3.2\)](#page-7-1) imply that D is expressed as a product  $D'g_{n,i}$  or  $D''h_{n,0}$  such that  $D'$  or  $D''$  is a diagram of n strands and  $1 \le i \le n+1$ . Here,  $D'$  or  $D''$ is a diagram with n strands and when we consider the product with  $g_{n,i}$  or  $h_{n,0}$ , we add a vertical strand to the right of  $D'$  or  $D''$  such that  $D'$  or  $D''$  and the vertical strand form a diagram with  $n+1$  strands. By the same argument as the case of type A (see [\[11\]](#page-29-7) for details), the connectivity of a diagram D' is the same as that of  $D_i$ . In the case  $D = D' g_{n,i}$ , it is obvious that the position of dots in D' is the same as  $D_i$ . From these, we have  $D' = D_i$ . If  $D = D'' h_{n,0}$ , we have to add a dot on the cap or the vertical strand which contains the left-most bottom point after deleting the left-most cap, since the graphical representation of  $h_{n,0}$  has a vertical strand with a dot which contains the left-most top point. Since each cap or a vertical strand has at most one dot, if there exist two dots on a cap or a vertical strand, then we delete these two dots. From these, we have  $D'' = D_1$  if  $D = D''h_{n,0}$  and  $1 \in I^{\bullet}(D)$ . Finally, we have  $\text{coef}(g_{n,1}) = \text{coef}(h_{n,0})$ . By combining these observations together, we have Eq.  $(3.7)$ .

**Example 3.9.** We consider the coefficient of  $E_1E_2E_3E_0$  in  $P_3$ . We calculate

$$
\begin{array}{c}\n\bigvee \left(\begin{array}{c}\n\searrow \\
\hline\n\end{array}\right) = \frac{[3]}{[6]} \cdot \bigvee \left(\begin{array}{c}\n\searrow \\
\hline\n\end{array}\right) + \frac{[3][4]}{[6][2]} \cdot \bigvee \left(\begin{array}{c}\n\searrow \\
\hline\n\end{array}\right) \\
= \left(\frac{[3]}{[6]} + \frac{[3][4]}{[6][2]}\right) \frac{[2]}{[4]} \cdot \bigvee \left(\begin{array}{c}\n\searrow \\
\hline\n\end{array}\right) \\
= \frac{[3]^2[2]}{[6][4]} \cdot \frac{1}{[2]} \cdot \left|\begin{array}{c}\n\searrow \\
\hline\n\end{array}\right| \\
= \frac{[3]^2}{[6][4]}.\n\end{array}
$$

Note that we are not allowed to remove the vertical strand with a dot at the position 3 (in the first row). In the calculation of the coefficient of  $E_1E_2E_3E_0$ , we have only diagrams which have an even number of dots.

Compare the result with Example [4.10.](#page-18-0)

Secondly, we consider the case where a diagram D contains a single dot, which implies that D corresponds to an element containing the product  $E_0E_1$  in its algebraic representation. The connectivity of the diagram D is different from 1 by the property (P2c). Let  $I(D) \subseteq [1, n+1]$ be the set of positions of inner-most caps possibly with a dot. If  $n + 1 \in I(D)$ , this means that a diagram has an inner-most cap as a vertical strand connecting the right-most marked points on the top and bottom. Given  $i \in I(D)$ , we denote by  $D_i$  the diagram obtained from D by deleting the inner-most cap at the position *i*. The size of  $D_i$  is one smaller than that of D. We have two cases: 1)  $D_i \neq 1$ , and 2)  $D_i = 1$ .

Case 1). We define  $\mathcal{D}_{i}^{e}$  as the set of diagrams D such that D has the same connectivity of caps, cups and vertical strands as  $D_i$ , and has an even number of dots. The positions of dots in the diagram  $D' \in \mathcal{D}_i^e$  satisfy the property (P2a). Similarly, we denote by  $D_i^{\text{odd}}$  the unique diagram which has the same connectivity of caps, cups and vertical strands as  $D_i$ , and has a unique dot satisfying the property (P2b).

Case 2). Since  $D_i = 1$ , we define  $\mathcal{D}_i^e := \{1\}$  and  $D_i^{\text{odd}} := \emptyset$  by the property (P2c).

<span id="page-11-0"></span>**Proposition 3.10.** Let D be a diagram with a unique dot, and  $I(D)$ ,  $\mathcal{D}_i^e$ , and  $D_i^{\text{odd}}$  be as above. Then, we have

<span id="page-11-2"></span>(3.8) 
$$
D = \sum_{i \in I(D)} \mathrm{coef}(h_{n,i}) \left( \sum_{D' \in \mathcal{D}_i^e} D' - [2]D_i^{\mathrm{odd}} \right) + \sum_{i \in I(D)} 2^{\delta(i=1)} \mathrm{coeff}(g_{n,i}) D_i^{\mathrm{odd}},
$$

<span id="page-11-3"></span>where  $\delta(P)$  is the Kronecker delta function, namely,  $\delta(P) = 1$  if P is true and  $\delta(P) = 0$  otherwise.

Remark 3.11. Two remarks are in order:

- (1) The factor  $2^{\delta(i=1)}$  comes from the fact that we have  $\text{coef}(h_{n+1,0}) = \text{coef}(g_{n+1,1}),$  and  $I(D) \subseteq$ [1, n + 1]. The set  $I(D)$  of the positive integers cannot detect the generator  $h_{n+1,0}$  since  $0 \notin I(D)$ .
- (2) The factor -[2] in front of  $D_i^{\text{odd}}$  comes from the fact that if we multiply  $h_{n,i}$  with  $h_{m,j}$  such that  $i, j \neq 0$ , then this product gives  $-[2]h'$  with some  $h' \in TL_n^D$ . In other words, both  $h_{n,i}$  and  $h_{m,j}$  contain the product  $E_0E_1$ , and we use  $E_1^2 = -[2]E_1$  or  $E_0^2 = -[2]E_0$  in the calculation of the product.

*Proof of Proposition [3.10.](#page-11-0)* We rewrite the recurrence relation  $(3.2)$  in Proposition [3.3](#page-7-0) in terms of diagrams. Recall that the diagram  $D$  has a unique single dot which implies that the element corresponding to D contains the product  $E_0E_1$ .

If we have  $D = D'g_{n,i}$  or  $D = D''h_{n,0}$  where  $D'$  or  $D''$  is an element in  $Q_{n-1}, D'$  or  $D''$  has to contain the product  $E_0E_1$  since  $g_{n,i}$  or  $h_{n,0}$  does not contain the product  $E_0E_1$ . Further we have  $\text{coef}(g_{n,1}) = \text{coef}(h_{n,0})$ . This gives the factor  $2^{\delta(i=1)}$ . By the same argument as the proof of Proposition [3.7,](#page-9-0) D' or  $D''$  has the same connectivity as  $D_i^{\text{odd}}$ . The number of dots in D' or D" is one since it contains  $E_0E_1$ . Then, the diagram D' or D'' satisfies the property (P2b). From these, we have the second sum in the right hand side of Eq. [\(3.8\)](#page-11-2).

Suppose that we have  $D = D'h_{n,i}$  with  $1 \leq i \leq n$  where  $D'$  is an element in  $Q_{n-1}$ . Then, since  $h_{n,i}$  contains the product  $E_0E_1$ , we have two cases for  $D'$ : a) the diagram  $D'$  has an even number of dots, and b)  $D'$  has a unique single dot.

- (a) Again, by the same argument as the proof of Proposition [3.7,](#page-9-0) the connectivity of  $D'$  is the same as diagrams in  $\mathcal{D}_{i}^{e}$ . Therefore, it is enough to show that a diagram  $D' \in \mathcal{D}_{e}^{i}$  satisfies  $D = D'h_{n,i}$ . However, the facts that D and  $h_{n,i}$  contain a single dot, and D' contains an even number of dots, imply that if  $D' \in \mathcal{D}_{e}^{i}$ , then  $D = D'h_{n,i}$ . From this observation, we have the term  $\sum_{i \in I(D)} \text{coef}(h_{n,i}) \sum_{D' \in \mathcal{D}_i^e} D'.$
- (b) Since  $D'$  has a unique single dot, the diagram is uniquely fixed if the connectivity of caps, cups and vertical strands is fixed. The condition that  $D = D'h_{n,i}$  implies that the connectivity of D' is the same as that of  $D_i^{\text{odd}}$ . Then, it is easy to see that  $D' = D_i^{\text{odd}}$  since  $D'$  contains a unique dot. The element corresponding to  $D_i^{\text{odd}}$  contains the product  $E_0E_1$ . By the second remark in Remark [3.11,](#page-11-3)  $D_i^{\text{odd}} h_{n,i}$  yields the factor  $-[2]$ . By combining these observations, we have the term  $-[2] \sum_{i \in I(D)} \text{coef}(h_{n,i}) D_i^{\text{odd}}.$

From the two cases (a) and (b), we have the first sum in the right hand side of Eq. [\(3.8\)](#page-11-2). This completes the proof.  $\Box$ 

<span id="page-11-1"></span>3.4. Relation between types  $A$  and  $D$ . In this section, we will show that a linear combination of the diagrams in  $TL_{n+1}^D$  satisfies the recurrence relation of a diagram in  $TL_{n+1}^A$ . This linear combination of the diagrams of type  $D$  plays a central role when we interpret the recursive relation [\(3.8\)](#page-11-2) in terms of the Dyck tilings.

Let D be an  $n+1$ -strand Temperley-Lieb diagram for  $TL_{n+1}^D$ . The diagram may have dots. We denote by  $D^A$  the diagram obtained from D by deleting all dots. In other words, we focus on the only connectivity of  $\overline{D}$  and ignore the dots. This operation naturally defines the forgetful map For :  $D \mapsto D^A$ . Let  $\mathcal{D}^e(D^A)$  be the set of diagrams D such that  $\text{For}(D) = D^A$  and they have an even number of dots satisfying (P2a). We have a unique diagram which has an odd number of dots and has the same connectivity as  $D^A$  if  $D^A \neq 1$ . Therefore, we denote by  $D^{\text{odd}}$  this unique diagram. As a consequence, the diagram  $D^{odd}$  has a single dot, and it satisfies (P2b). Here, the superscript A in  $D^A$  stands for type A, and the superscript odd in  $D^{\text{odd}}$  stands for a diagram of type D with a single dot.

**Definition 3.12.** Let  $D^A$ ,  $\mathcal{D}^e(D^A)$  and  $D^{\text{odd}}$  be the diagram given as above. We define the following linear combination:

<span id="page-12-0"></span>(3.9) 
$$
D^{A} := \sum_{D \in \mathcal{D}^{e}(D^{A})} D - [2] D^{\text{odd}}.
$$

The linear combination  $D^A$  in Eq. [\(3.9\)](#page-12-0) can be regarded as an element of Temperely–Lieb algebra of type A by the next proposition.

<span id="page-12-1"></span>**Proposition 3.13.** The linear combination  $D^A$  satisfies the recurrence relation [\(3.6\)](#page-9-2) of type A in Proposition [3.6.](#page-9-3)

*Proof.* If  $D^A = 1$ , we have  $\mathcal{D}^e(D^A) = \{1\}$  and  $D^{odd} = \emptyset$ . In this case, the claim is trivial. For  $D^A \neq 1$ , it is enough to show that the linear combination [\(3.9\)](#page-12-0) satisfies the recurrence relation  $(3.6)$  of type A by use of the recurrence relations  $(3.7)$  and  $(3.8)$ . We compute the coefficient of  $D_i^A$ ,  $i \in I(D^A)$ , where  $D_i^A$  is the diagram of type A obtained from  $D^A$  by removing the inner-most cap at position *i*. We consider the three cases: the first one is  $i = n + 1$ , the second one is  $1 \le i \le n$ , and the third one is  $i = 1$ .

Firstly, suppose that  $n+1 \in I(D^A)$ . The diagram  $D^A$  has a vertical strand which connects the two right-most marked points on the top and bottom. This vertical strand is inner-most by definition. We calculate the contributions of the right hand side of Eq.  $(3.9)$  to  $D_{n+1}^A$ . When  $D \in \mathcal{D}^e(D^A)$ , and the  $n + 1$ -st inner-most cap does not have a dot, we have a coefficient coef $(g_{n+1,n+1}) = 1$ . Further, the new diagram (obtained from  $D$  by removing the right-most vertical strand) contains an even number of dots. We have all diagrams with an even number of dots and with the same connectivity as  $D_{n+1}^A$ . If D has a dot at the  $n+1$ -st inner-most cap, this does not contribute to  $D_{n+1}^A$  since we have a recurrence relation [\(3.7\)](#page-9-1). We apply Eq. [\(3.8\)](#page-11-2) to the diagram  $D^{\text{odd}}$ . Note that the coefficient of  $h_{n,n+1}$  is equal to zero since it is not defined. For the second term, we have  $\text{coef}(g_{n,n+1})D_{n+1}^{\text{odd}}$ . By combining these with Eq.  $(3.9)$ , we have the linear combination of the diagrams for  $D_{n+1}^A$  with the coefficient 1.

Secondly, suppose that  $1 \lt i \leq n$  and  $i \in I(D^A)$ . We apply the recurrence relations [\(3.7\)](#page-9-1) and  $(3.8)$  to Eq.  $(3.9)$ . Explicitly, we have

$$
\begin{split}\n&\cosh(g_{n,i})\sum_{D\in\mathcal{D}^e(D_i^A)}D-[2]\left(\cosh(h_{n,i})\sum_{D\in\mathcal{D}^e(D_i^A)}D-[2]\mathrm{coeff}(h_{n,i})D_i^{\mathrm{odd}}+\mathrm{coeff}(g_{n,i})D_i^{\mathrm{odd}}\right), \\
&=\frac{[i]}{[n+1]}\left(\sum_{D\in\mathcal{D}^e(D_i^A)}D-[2]D_i^{\mathrm{odd}}\right), \\
&=\frac{[i]}{[n+1]}D_i^A,\n\end{split}
$$

where  $D \in \mathcal{D}^e(D_i^A)$  (resp.  $D_i^{\text{odd}}$ ) are the diagrams of even (resp. odd) number of dots such that For $(D) = D_i^A$  (resp. For $(D_i^{\text{odd}}) = D_i^A$ ).

Finally, suppose that  $i = 1 \in I(D^A)$ . By applying the recurrence relations [\(3.7\)](#page-9-1) and [\(3.8\)](#page-11-2) to Eq.  $(3.9)$ , we obtain

<span id="page-13-1"></span>
$$
2 \cdot \text{coef}(g_{n,1}) \sum_{D \in \mathcal{D}^e} D - [2] \left( \text{coef}(h_{n,1}) \sum_{D \in \mathcal{D}^e} D - [2] \text{coef}(h_{n,1}) D_1^{\text{odd}} + 2 \cdot \text{coef}(g_{n,1}) D_1^{\text{odd}} \right),
$$
  
(3.10)  

$$
= \frac{1}{[n+1]} \left( \sum_{D \in \mathcal{D}^e} D - [2] D_1^{\text{odd}} \right),
$$
  

$$
= \frac{1}{[n+1]} D_1^A.
$$

where  $\mathcal{D}^e := \mathcal{D}^e(D_1^A)$ . The first factor 2 in Eq. [\(3.10\)](#page-13-1) comes from the fact that we have two generators  $E_1E_2E_0$  and  $E_0E_2E_1$ , whose diagrams are the same as  $E_1$  and  $E_0$  if we ignore the dots.

Form these, the linear combination  $(3.9)$  satisfies the recurrence relation  $(3.6)$ .

Example 3.14. Let  $D^A$  be a diagram without dots:

$$
D^A = \bigcup_{\bigcap_{i=1}^n} \ \ \bigg| \ .
$$

Then, we consider the following linear combination in  $TL_3^D$ :

<span id="page-13-2"></span>(3.11)  $D^A = E_1 + E_0 + E_1 E_2 E_0 + E_0 E_2 E_1 - [2] E_0 E_1.$ 

Here, we make use of the algebraic representations of the diagrams. By a simple calculation using Eq.  $(3.6)$ , we have

$$
\operatorname{coef}^A(D^A) = \frac{[2]}{[3]},
$$

where  $\text{coef}^A(D)$  is the coefficient of the diagram D in the type A case. By a similar calculation, the right hand side of Eq.  $(3.11)$  gives

$$
\frac{[3]}{[4]} + \frac{[3]}{[4]} + \frac{1}{[4]} + \frac{1}{[4]} - [2]\frac{[2]^3}{[4][3]} = \frac{[2]}{[3]}.
$$

3.5. Recurrence relation via diagrams with a unique dot. By combining the result in Section [3.3](#page-8-3) with that in Section [3.4,](#page-11-1) we rewrite the recurrence relation [\(3.8\)](#page-11-2) in terms of the diagrams of types A and D.

Let D be the diagram with a unique dot. Let  $D_i$  be the diagram obtained from D by removing an inner-most cap at position *i*. We define  $D_i^A := \text{For}(D_i)$  where For is the forgetful map. Similarly,  $D_i^{\text{odd}}$  is a diagram with a unique dot such that  $\text{For}(D_i^{\text{odd}}) = D_i$ .

<span id="page-13-0"></span>The following is a direct consequence of Propositions [3.10](#page-11-0) and [3.13.](#page-12-1)

**Proposition 3.15.** Let D,  $D_i^A$  and  $D_i^{\text{odd}}$  be diagrams as above. Then, we have

<span id="page-13-3"></span>(3.12) 
$$
D = \sum_{i \in I(D)} \text{coef}(h_{n,i}) D_i^A + \sum_{i \in I(D)} 2^{\delta(i=1)} \text{coef}(g_{n,i}) D_i^{\text{odd}}.
$$

Note that the recurrence relation  $(3.8)$  is not minus sign free, *i.e.*, we have a term with minus sign  $-[2]D_i^{\text{odd}}$ . However, the recurrence relation  $(3.12)$  is a minus sign free because of Proposition [3.13.](#page-12-1)

The coefficient of D in  $Q_n$  is expressed as the linear combination of  $D_i^A$  and  $D_i^{\text{odd}}$  with positive coefficients. However, the recurrence relation contains the diagrams not only of type D but also of type A. This form plays a central role when we interpret the recurrence relation [\(3.12\)](#page-13-3) in terms of the Dyck tilings in the next section.

## 4. Dyck tilings

## <span id="page-14-0"></span>4.1. **Definition.** We briefly introduce the notion of cover-inclusive Dyck tilings following [\[10,](#page-29-12) [14\]](#page-29-10).

A Dyck path of size n is a lattice path from  $(0, 0)$  to  $(2n, 0)$  such that it consists of up steps  $(1, 1)$ and down steps  $(1, -1)$  and it never goes below the horizontal line  $y = 0$ . By its definition, a Dyck path has  $n$  up steps and  $n$  down steps. For example, we have five Dyck paths of size three:

## UUUDDD UUDUDD UUDDUD UDUUDD UDUDUD

where U (resp. D) stands for an up (resp. down) step. We call the Dyck path  $U^nD^n$  the top Dyck path of size n. Let  $\lambda, \mu$  be two Dyck paths. We write  $\lambda \leq \mu$  if  $\mu$  is above  $\lambda$ .

A ribbon is a connected skew shape which does not contain a 2-by-2 rectangle. A ribbon is called a Dyck tile if the centers of the unit cells in the ribbon form a Dyck path. A unit cell is a Dyck tile of size zero, and the size of a Dyck tile  $d$  is the size of the Dyck path which characterizes  $d$ . We call a Dyck tile of size zero a trivial Dyck tile, and other Dyck tiles non-trivial Dyck tiles.

Let  $\lambda, \mu$  be two Dyck paths such that  $\lambda \leq \mu$ . We consider the tiling of the region above  $\lambda$ and below  $\mu$  by Dyck tiles. In this paper, we consider only cover-inclusive Dyck tilings. Here, "cover-inclusiveness" means that if we move a Dyck tile downward by  $(0, -2)$ , then it is contained in another Dyck tile or strictly below the path  $\lambda$ . We give an example of a cover-inclusive Dyck



<span id="page-14-1"></span>FIGURE 4.1. An example of a cover-inclusive Dyck tiling of size 8

tiling in Figure [4.1.](#page-14-1) We have fifteen Dyck tiles, and three non-trivial Dyck tiles.

Remark 4.2. The cover-inclusive Dyck tilings correspond to Dyck tilings which satisfy the rule I in [\[14\]](#page-29-10).

A Dyck tile d consists of several unit squares. Given a unit box s, we denote by  $h(s)$  the height of the center of s.

Definition 4.3. We define the height of d by

$$
h(d) := \min\{h(s) : s \in d\}.
$$

In Figure [4.1,](#page-14-1) the heights of three non-trivial Dyck tiles are one, two, and four.

<span id="page-15-0"></span>4.2. Correspondence between a Dyck path and a Temperley–Lieb diagram. In this section, we give a correspondence between a Dyck path of size  $n$  and an n-strand Temperley–Lieb diagram  $D$  possibly with dots. We fold an *n*-strand Temperley–Lieb diagram  $D$  down to the right such that 2n points are in line. Then, we obtain a graph  $\mu$  with 2n points such that a point is connected to another point by a cap. We have n caps in the graph  $\mu$ . Note that the strands in D are non-intersecting, therefore the caps in  $\mu$  are also non-intersecting. Obviously, we have a one-to-one correspondence between a strand in D and a cap in  $\mu$ . A cap in  $\mu$  has a dot if the corresponding strand in  $D$  has a dot. If we replace the left end point of a cap with  $U$ , and the right end point of a cap with D, then we have a Dyck path. The Dyck path which consists of possibly dotted caps corresponds to the diagram D.

In Figure [4.4,](#page-15-1) we give the Dyck path with a dot which corresponds to the diagram A in Example [2.3.](#page-5-0) The corresponding Dyck path is *UDUUDUDD* and the left-most cap has a dot.



<span id="page-15-1"></span>Figure 4.4. A Dyck path corresponding to a 4-strand Temperley–Lieb diagram.

We call a graph consisting of  $2n$  points and n non-crossing caps with or without dots a *chord* diagram. The number of chord diagrams consisting of  $n$  caps without dots is given by the Catalan number  $C_n$ . Similarly, the number of chord diagrams of n caps with dots which correspond to the *n* strand Temperley–Lieb diagrams of type *D* is given by dim  $TL_p^D$ . This is easily obtained since the number of chord diagrams with an even number of dots is  $\frac{1}{2} \binom{2n}{n}$  $\binom{2n}{n}$ , and those with an odd number of dots is  $C_n - 1$ . Here, we use the property (P2c), which implies that the chord diagram corresponding to the identity 1 cannot have a dot. As a consequence, the number of chord diagrams with dots is given by

$$
\frac{1}{2} \begin{bmatrix} 2n \\ n \end{bmatrix} + C_n - 1 = \dim \mathrm{TL}_n^D.
$$

4.3. Hermite histories. We introduce the notion of Hermite histories on a cover-inclusive Dyck tiling: horizontal and vertical. (Horizontal Hermite histories are introduced in [\[10,](#page-29-12) Section 3].) We



<span id="page-15-2"></span>Figure 4.5. Horizontal Hermite history (left) and vertical Hermite history (right).

draw a line in Dyck tiles in a Dyck tiling as follows. The boundary of a Dyck tile consists of up steps  $(\angle)$  and down steps  $(\angle)$ . For a horizontal Hermite history, we write a line from the left-most up step to the right-most up step in a Dyck tile. We concatenate the lines in Dyck tiles if it is possible to extend. A line passing through Dyck tiles is called a horizontal trajectory. Similarly, for a vertical Hermite history, we write a line from the left-most down step to the right-most down step in a Dyck tile. We extend the lines if possible. A line obtained as above is called a vertical trajectory. We emphasize that the length of a trajectory is larger than zero. One can consider

trajectories of length zero in an Hermite history (see Remark [4.7\)](#page-16-0). Figure [4.5](#page-15-2) gives examples of horizontal and vertical Hermite histories. We have three horizontal trajectories in the left figure, and two vertical trajectories in the right figure. These trajectories have length larger than zero.

A horizontal Hermite history can be obtained as a mirror image of a vertical Hermite history, and vice versa.

<span id="page-16-1"></span>**Proposition 4.6.** Let T be a Dyck tiling of size n above  $\lambda$  and below  $\mu$ . A horizontal or vertical Hermite history  $H(T)$  of T has the following properties:

- (a)  $H(T)$  contains at most n 1 trajectories with non-zero length.
- (b) All trajectories are non-crossing.
- (c) Any trajectory starts from an up step in  $\mu$  for the horizontal Hermite history. Similarly, all trajectories ends at a down step in µ for the vertical Hermite history.

Proof. We prove only for a vertical Hermite history since the horizontal case can be proven in a similar way.

(a) By construction of a trajectory, an down step in  $\lambda$  is connected to an down step in  $\mu$  by a trajectory. Let d be the right-most down step in  $\lambda$ . Since  $\lambda$  and  $\mu$  share the same down step  $d$  at their right ends, there is no trajectory with non-zero length which starts from  $d$ . If we take  $\lambda = (UD)^n$  and  $\mu = U^n D^n$ , and T does not have non-trivial Dyck tiles, then it is obvious that we have  $n-1$  trajectories with non-zero length. Since we have n down steps in  $\lambda$  and  $\mu$ , the maximal number of trajectories with non-zero is  $n-1$ .

(b) Any Dyck tile t in T has a line inside t. The right-most down step in t is attached to the left-most down step of another Dyck tile, or to a down step in  $\mu$ . This property insures that all trajectories are non-crossing.

(c) A down step d in  $\mu$  is a down step of a Dyck tile t, or a down step of  $\lambda$ . If d is the right-most down step of  $t$ , then there exists a trajectory which ends at  $d$ . Otherwise, there is no trajectory which ends at d and has non-zero legnth. Recall that we consider a cover-inclusive Dyck tiling  $T$ . Especially, if two Dyck tiles  $t_1$  is on the top of another Dyck tile  $t_2$ , the size of  $t_1$  is weakly smaller than that of  $t_2$ . This means that the right end of the trajectory containing  $t_2$  is right to the right end of the trajectory containing  $t_1$ . Recall we concatenate lines in Dyck tiles to obtain a trajectory. If a trajectory is below another trajectory, the right end of the former is right to the latter. These implies that the right end of a trajectory ends at a down step in  $\mu$ .

<span id="page-16-0"></span>**Remark 4.7.** Suppose that we have  $m \leq n-2$  trajectories in a vertical Hermite history  $H(T)$ . By the proof of (c) in Proposition [4.6,](#page-16-1) there exists a down step d in  $\mu$  such that there exists no trajectory ending at d. If we regard this situation as a trajectory of length zero, we have n trajectories in total. Especially, if  $n = 1$ , we have no trajectories of length larger than zero. This situation can be interpreted as follows: there is a trajectory of length zero starting from the unique down step in  $\lambda$  to the unique down step in  $\mu$ . Therefore, if we allow the trajectories of length zero, then any Hermite history contains n trajectories.

4.4. Dyck tilings and the diagram with an even number of dots. Let D be an  $n+1$ -strand Temperley–Lieb diagram with an even number of dots. Then, by unfolding the diagram  $D$ , we have a chord diagram C of size  $2n+2$ . Since D does not contain the product  $E_0E_1$  in the algebraic representation, the chord diagram  $C$  also has an even number of dots. Since  $D$  contains only an even number of dots, D satisfies the recurrence relation [\(3.7\)](#page-9-1).

First, suppose that the Jones–Wenzl projection  $R_n$  for a Temperley–Lieb algebra has the form

$$
R_n = R_{n-1} \left( \sum_{i \in I(D)} \frac{\langle i \rangle}{\langle n+1 \rangle} D_i \right),
$$

where  $R_n$  is the projection,  $D_i$  is the diagram obtained from D by removing the inner-most cap at position i and  $\langle n \rangle$  is a function of n. The projections for types A and B, and type D with  $E_0E_1 = 0$ have this form. In type A, we have  $\langle n \rangle$  is simply  $[n]$ , and  $\langle n \rangle$  is simply  $[n]$ , for type B (see [\[13\]](#page-29-8) for a definition of  $[n]_s$ ). In type D (an even number of dots),  $\langle n \rangle$  is  $\frac{[2(n-1)]}{[n-1]}$ . Once again, note that this works well only in the case of the diagrams of type  $D$  with an even number of dots. If the diagram contains a unique dot, then the recurrence relation  $(3.8)$  contains the coefficients

$$
\mathrm{coef}(h_{n,i}) = \frac{[n][n+1-i]}{[2n][n+1]},
$$

for  $1 \leq i \leq n$ , and these coefficients cannot be written as  $\langle i \rangle / \langle n + 1 \rangle$ .

Secondly, we have the similarity between type  $D$  (an even number of dots) and type  $B$ . In type B, we cannot remove the inner-most cap at the position  $i > 1$  if it has a dot [\[13\]](#page-29-8). Similarly in type  $D$ , the diagram of  $E_0$  has two dots, and we are not allowed to have a diagram which has the same connectivity as  $E_i$ ,  $i > 1$ , and has two dots on a cap and a cup. Therefore, in type D, we cannot remove the inner-most cap with a dot if its position is  $i > 1$ .

The second observation can be easily translated in terms of Dyck tilings. Some Dyck tilings are not allowed in type B, and such Dyck tilings are also not allowed in type D as well.

We can apply the same algorithm for type B studied in  $[13]$  for these diagrams of type D. When a diagram D contains a cap with a dot, this cap corresponds to a pair of an up step and a down step in the chord diagram  $C$  corresponding to  $D$ . Let  $D$  be a Dyck tiling whose bottom Dyck path is the same as C, and top Dyck path is  $U^{n+1}D^{n+1}$ . Let c be a dotted cap in a chord diagram, and  $x_l$  (resp.  $x_r$ ) the position of the left (resp. right) end of c in the chord diagram C. Then, the size of the cap c is defined to be  $l(c) := (x_r - x_l + 1)/2$ .

**Definition 4.8.** Let  $\mu$  be the Dyck path. The generating function of Dyck tilings above  $\mu$  and below the top path is defined to be

$$
Z(\mu) := \sum_{T} \prod_{d \in T} \frac{[h(d)][2h(d) - 2]}{[2h(d)][h(d) - 1]},
$$

where  $d \in T$  is a Dyck tile in a Dyck tiling T, we define  $\frac{[2h(d)-2]}{[h(d)-1]} := 1$  if  $h(d) = 1$ , and the sum  $\sum'$ is taken over all Dyck tilings such that there is no Dyck tile of size  $l(c)$  above the dotted cap c.

<span id="page-17-0"></span>Then, we have the following theorem.

**Theorem 4.9.** Let D be a diagram with an even number of dots, and  $\mu$  the Dyck path corresponding to D. The coefficient coef(D) of D in  $Q_n$  is given by

$$
\mathrm{coef}(D) = Z(\mu).
$$

*Proof.* We prove that the generating function  $Z(\mu)$  satisfies the recurrence relation [\(3.7\)](#page-9-1). Fix a Dyck tiling D above  $\mu$  and below the top path  $U^{n+1}D^{n+1}$  such that there is no Dyck tile of size  $l(c)$ above a dotted cap c. Let d be the left-most down step in the top path, and consider a trajectory  $T_d$  in the vertical Hermite history of D which contains the down step d. We have two cases: 1)  $T_d$ connects the down step d and a step in the bottom Dyck path  $\mu$ , and 2)  $T_d$  does not touch a down step in  $\mu$ .

(1) The trajectory  $T_d$  consists of only trivial Dyck tiles. Let  $h_{low}$  be the height of the lowest unit cell in  $T_d$ . Then, the product of the weights of the Dyck tiles in  $T_d$  is given by

<span id="page-18-1"></span>(4.1) 
$$
\frac{[n][2(h_{low}-1)]}{[2n][h_{low}-1]}
$$

if  $h_{low} \geq 2$  and  $[n]/[2n]$  if  $h_{low} = 1$ . Since  $T_d$  touches a step in the bottom Dyck path  $\mu$ , there is an inner-most cap at position  $h_{low}$ . To remove the cap at position  $h_{low}$  corresponds to give the weight [\(4.1\)](#page-18-1), and to delete the trajectory  $T_d$ . If  $h_{low} = 1$ , we remove the cap at position one. If  $h_{low} \geq 2$ , the cap at position  $h_{low}$  does not have a dot since this cap is not outer-most. By removing an inner-most cap, we have a Dyck tiling of smaller size.

(2) The trajectory  $T_d$  consists of only trivial Dyck tiles, however, the low end of  $T_d$  touches a down step of non-trivial Dyck tile  $d_1$ . As in the case of (1), the product of the weights of the Dyck tiles in  $T_d$  is given by Eq. [\(4.1\)](#page-18-1). By the same argument as (1), to remove the cap at position  $h_{low}$  corresponds to give the weight, to delete the trajectory  $T_d$  and to shrink the size of  $d_1$  by one. Suppose that we have a dotted cap c and there is a non-trivial Dyck tile of size  $l(c)$  above c. The above observation implies that the trajectory  $T<sub>d</sub>$  detects the inner-most dotted cap at position  $h_{low}$ . However, we are not allowed to remove the inner-most dotted cap at position  $h_{low} \geq 2$ . This is avoidable if we consider only Dyck tilings such that there is no Dyck tile of size  $l(c)$  above c. In fact, the generating function  $Z(\mu)$  satisfies this condition.

Suppose that the bottom path of the Dyck tiling corresponds to the diagram D. By the correspondence between a Temperley–Lieb diagram and a chord diagram, the Dyck tiling of smaller size corresponds to  $D_i$  in Eq. [\(3.7\)](#page-9-1). By combining (1), (2) and the above observation, it is clear that  $Z(\mu)$  satisfies the recurrence relation [\(3.7\)](#page-9-1).

<span id="page-18-0"></span>**Example 4.10.** We consider the coefficient of  $E_1E_2E_3E_0$  in  $P_3$ .



We have two admissible Dyck tilings.



Then, the coefficient of this diagram is given by

$$
\frac{[3]}{[6]} \frac{[2]}{[4]} \frac{1}{[2]} + \frac{[3]}{[6]} \frac{1}{[2]} = \frac{[3]^2}{[6][4]}.
$$

There are two non-admissible diagrams:



The Dyck tile with  $\star$  violates the condition. The dotted cap below  $\star$  has a Dyck tile of size 1.

<span id="page-19-0"></span>4.5. The diagram with a unique dot. We introduce two colors on a vertical Hermite history. Recall that a line in an Hermite history is called a trajectory. In other words, an Hermite history consists of several trajectories. As in Remark [4.7,](#page-16-0) we have n trajectories in an Hermite history if we allow a trajectory of length zero.

We consider the following conditions on a vertical Hermite history.

- (Q1) There exists a unique trajectory with the color green. The length of this trajectory is non-zero.
- (Q2) The trajectories left to the green trajectory are all red. If the left-most trajectory is green, then there is no red trajectories with length non-zero.
- (Q3) There are no trajectories on Dyck tiles right to the green trajectory. That is, we have a Dyck tiling  $T$  right to the green trajectory but remove all the trajectories on the Dyck tiles in  $T$ .

The condition (Q3) implies that there may be a region without trajectories. Such a region may be empty.

We impose one more condition on the vertical Hermite history. By Remark [4.7,](#page-16-0) we have  $n$ vertical trajectories in a vertical Hermite history if we include trajectories of length zero. Fix a Dyck tiling and its vertical Hermite history. We denote by  $T_{\geq 0}$  the set of vertical trajectories of size weakly larger than zero. Suppose that the unique green trajectory  $t$  is the  $m$ -th vertical trajectory from left in  $T_{\geq 0}$ . Let  $t_i$ ,  $1 \leq i \leq n-m$ , be the  $m+i$ -th vertical trajectory from left in  $T_{\geq 0}$ . We denote by  $n_i$  the number of Dyck tiles in the trajectory  $t_i$ . These  $n - m$  vertical trajectories are removed by (Q3). We consider the following condition:

(Q4)  $n_1 \geq n_2 \geq \ldots \geq n_{n-m}$ .

Note that  $n_{n-m} = 0$  in (Q4) since the lower Dyck path and the top Dyck path in a Dyck tiling share the same right-most down step and we have a trajectory of length zero.

**Definition 4.11.** We call a vertical Hermite history which satisfies the conditions  $(Q1)$  to  $(Q4)$  a bi-colored vertical Hermite history.

We first study the condition (Q4). Fix a Dyck tiling T of size n above a Dyck path  $\lambda$  and below the top Dyck path, and its bi-colored vertical Hermite history  $V$ . Suppose that the m-th trajectory from left in  $V$  is the unique green trajectory. We introduce the operation which we call a deletion of T. Let d be the left-most down step of the lowest Dyck tile in the first trajectory. Let  $\lambda_{i+1}$  be the down step of  $\lambda$  which is obtained by moving d downward. Since  $\lambda_{i+1}$  is a down step and d is in the first trajectory, the step  $\lambda_i$  is an up step in  $\lambda$ . We delete all the Dyck tiles in the first vertical trajectory and shrink all the Dyck tiles above  $\lambda_i$  and  $\lambda_{i+1}$ . The newly obtained Dyck tiling has a lower Dyck path  $\lambda'$  and the upper Dyck path  $U^{n-1}D^{n-1}$ . Here, the Dyck path  $\lambda'$  is obtained from  $\lambda$  by deleting  $\lambda_i$  and  $\lambda_{i+1}$ . This operation is visualized in Figure [4.12.](#page-19-1) We call this operation a



<span id="page-19-1"></span>Figure 4.12. A deletion of a Dyck tiling

deletion of T. We perform the deletion of T  $m$  times. Then, we obtain a Dyck tiling  $T'$  of size  $n - m$ . The condition (Q4) is equivalent to the following condition on  $T'$ :

(Q4′ ) T ′ consists of only trivial Dyck tiles.

This condition captures the property of  $(Q4)$  in terms of a Dyck tiling. We make use of this equivalence between  $(Q4)$  and  $(Q4')$  in the proof of Theorem [4.16](#page-20-0) below.

**Remark 4.13.** The equivalence between  $(Q_4)$  and  $(Q_4)$  holds if and only if the upper path of a Dyck tiling is the top Dyck path  $U^nD^n$ . Let us consider a Dyck tiling whose upper path is the top Dyck path. Then, by Proposition [4.6,](#page-16-1) any vertical trajectory ends at a down step in the upper path of a Dyck tiling. Since the upper path of a Dyck tiling is the top Dyck path, it is obvious that  $n_i < n_{i+1}$  gives a Dyck tiling with non-trivial Dyck tiles. A non-trivial Dyck tile appears only in this way.

Example 4.14. Consider the Dyck tiling in Figure [4.5.](#page-15-2) We have the following unique vertical Hermite history with two colors.



The left-most vertical trajectory (which is a red trajectory in the figure above) in the Dyck tiling cannot be green. If the first vertical trajectory is green, then we have  $(n_1, n_2, n_3) = (0, 1, 0)$ , which violates the condition  $(Q_4)$ .

Let T be a Dyck tiling with bi-colored trajectories satisfying the conditions  $(Q1)$  to  $(Q4)$ , and  $T^A$  be the Dyck tiling of the region without trajectories in T. We are allowed to consider the Dyck tilings such that it may have larger Dyck tiles than the Dyck tiles in  $T^A$ . The Dyck tiles are within the region without trajectories in T. Then, we define the weight  $wt(T^A)$  of  $T^A$  by

<span id="page-20-1"></span>(4.2) 
$$
\text{wt}(T^A) := \sum_{T' \in \mathcal{T}^A} \prod_{d \in T'} \frac{[h(d)]}{[h(d) + 1]},
$$

where  $\mathcal{T}^A$  is the set of Dyck tilings in the region without trajectories in T, and d is a Dyck tile in  $T' \in \mathcal{T}^A$ .

Let  $t_r$  be a red vertical trajectory, and we define the top and bottom heights  $h_t(t_r)$  and  $h_b(t_r)$ of  $t_r$  by the height of the top Dyck tile and the bottom Dyck tile in  $t_r$  respectively. Similarly,  $t_q$ be a green vertical trajectory. The heights  $h_t(t_q)$  and  $h_b(t_q)$  are defined in a similar way.

Let  $\lambda \leq \mu$  be two Dyck paths. Let  $\mathcal{D}(\lambda, \mu)$  be the set of bi-colored vertical Hermite histories of Dyck tilings above  $\lambda$  and below  $\mu$ . Suppose that we have a Dyck tiling  $T^A$  in the region without trajectories in the vertical Hermite history.

Definition 4.15. We define

$$
(4.3) \quad Z'(\lambda,\mu) := \sum_{D \in \mathcal{D}(\lambda,\mu)} 2^{N(t_r)} \text{wt}(T^A) \prod_{t_r} \frac{[h_t(t_r)][2(h_b(t_r)-1)]}{[2h_t(t_r)][h_b(t_r)-1]} \prod_{t_g} \frac{[h_t(t_g)][h_t(t_g)+1-h_b(t_g)]}{[2h_t(t_g)][h_t(t_g)+1]},
$$

where  $N(t_r)$  be the number of red trajectories  $t_r$  such that  $h_b(t_r) = 1$ . When  $h_b(t_r) = 1$ , we define  $\frac{[2(h_b(t_r)-1)]}{[h_b(t_r)-1]}:=1.$ 

Let  $D^{odd}$  be an  $n + 1$ -strand diagram with a single dot, and  $\lambda$  is the Dyck path corresponding to the diagram  $\text{For}(D^{\text{odd}})$ . Then, we have the following theorem which is one of the main results.

<span id="page-20-0"></span>**Theorem 4.16.** Let  $Z'(\lambda, \mu)$  be as above, and  $\mu_0$  the top Dyck path of size  $n + 1$ . The coefficient  $\text{coef}(D^{\text{odd}})$  of the diagram  $D^{\text{odd}}$  is given by

<span id="page-20-2"></span>(4.4) 
$$
\operatorname{coef}(D^{\text{odd}}) = Z'(\lambda, \mu_0).
$$

*Proof.* We will show that the generating function  $Z'(\lambda, \mu_0)$  satisfies the recurrence relation [\(3.12\)](#page-13-3) in Proposition [3.15](#page-13-0) if a bi-colored vertical Hermite history for  $Z'(\lambda, \mu_0)$  satisfies the conditions from (Q1) to (Q4). As in the proof of Theorem [4.9,](#page-17-0) we consider the trajectory  $T_d$  which is left-most in the bi-colored vertical Hermite history. We first show that a bi-colored vertical Hermite history satisfies the conditions (Q1) to (Q3). We consider two cases: 1)  $T_d$  is a green trajectory, and 2)  $T_d$ is a red trajectory.

- (1) Since the color of  $T_d$  is green, we have no green or red trajectories right to  $T_d$ . We may have a Dyck tiling without colored trajectories right to  $T<sub>d</sub>$ . From Eq. [\(3.12\)](#page-13-3), this corresponds to the case such that  $T_d$  gives the factor coef( $h_{n,i}$ ), and the region without colored trajectories corresponds to the generating function of Dyck tilings of type A. This is achieved by choosing  $i \in I(D)$ , and the first sum in the right hand side of Eq. [\(3.12\)](#page-13-3) is given.
- (2) Since the color of  $T_d$  is red, we have red trajectories and a unique right-most green trajectory right to  $T_d$ . The recurrence relation [\(3.12\)](#page-13-3) implies that if we have a coefficient coef( $g_{n,i}$ ), then we have a diagram  $D_i^{\text{odd}}$ . Since this diagram  $D_i^{\text{odd}}$  contains a unique dot, we apply the recurrence relation  $(3.12)$  once again. This procedure implies that unless we have a coefficient coef( $g_{n,i}$ ), we recursively apply the recurrence relation [\(3.12\)](#page-13-3). This stops when we have  $\text{coef}(h_{n,i})$  and produces the diagram  $D_i^A$ . This means that there exists a unique green trajectory  $T_q$ , and the trajectories left to  $T_q$  are all red. Then, the region right to the trajectory  $T_q$  does not have colored trajectories. We have a factor 2 when  $i = 1 \in I(D)$ , which comes from the fact that we have  $\text{coef}(h_{n,0}) = \text{coef}(g_{n,1})$  in Eq. [\(3.2\)](#page-7-1) in Proposition [3.3.](#page-7-0) By combining these together, we have the second term in Eq. [\(3.12\)](#page-13-3).

From these, a bi-colored vertical Hermite history in the generating function satisfies the conditions  $(Q1)$  to  $(Q3)$ .

Below, we show that a bi-colored vertical Hermite history satisfies the condition (Q4). Consider the first sum in the recurrence relation [\(3.12\)](#page-13-3). Since  $D_i^A$  is obtained from D by removing an innermost cap, the size of  $D_i^A$  is one smaller than that of D. Recall that the diagram  $D_i^A$  is of type A. If the lower Dyck path corresponding to  $D_i^A$  is  $\lambda$ , we have to consider all the Dyck tilings above  $\lambda$ and below the top Dyck path.

The expression  $(4.2)$  implies that we need to have the Dyck tiling T which consists of only trivial Dyck tiles. If the top and the bottom Dyck paths are given, we have a unique Dyck tiling with only trivial Dyck tiles, and other Dyck tilings are with non-trivial Dyck tiles. Such other Dyck tilings are obtained from T by merging Dyck tiles into a larger Dyck tile.

Suppose that a Dyck tiling T' has  $m \geq 0$  red trajectories and a unique green trajectory. Then, since the size of the Dyck tiling  $T'$  is  $n+1$ , the size of the region without colored trajectories is to be  $n + 1 - (m + 1) = n - m$ . The observation in the previous paragraph implies that if we perform the deletion of the Dyck tiling  $T'$   $m + 1$  times, then we need to have a Dyck tiling which consists of only trivial Dyck tiles. This is nothing but  $(Q4')$ . By the equivalence between  $(Q4)$  and  $(Q4')$ , a bi-colored vertical Hermite history satisfies the condition  $(Q_4)$ . On the contrary, suppose that a bi-colored vertical Hermite history violates the condition (Q4). Then, the region without colored trajectories consists of a Dyck tiling with a non-trivial Dyck tile after the  $m + 1$  deletions. In this case, we can not have the Dyck tiling consisting T of only trivial Dyck tiles when we calculate the weight  $(4.2)$  of  $D_i^A$ . From these observations, it is clear that bi-colored vertical Hermite histories in the generating function contain only the trajectories which satisfy the condition (Q4).

As a consequence, a bi-colored vertical Hermite history in  $Z'(\lambda, \mu_0)$  satisfies the conditions from (Q1) to (Q4), and  $Z'(\lambda, \mu_0)$  also satisfies the recurrence relation [\(3.12\)](#page-13-3). These imply Eq. [\(4.4\)](#page-20-2), which completes the proof.  $\Box$  **Example 4.17.** We calculate the coefficient of  $E_0E_1$  in  $Q_3$ . The diagram for  $E_0E_1$  and its corresponding top and bottom Dyck paths are as follows.



We have six bi-colored vertical Hermite histories.



The sum of the weights of the above diagrams is given by

$$
\frac{[3]^2}{[6][4]} \frac{[2]}{[3]} \frac{1}{[2]} + 2 \frac{[3]}{[6]} \frac{[2]}{[4][3]} \frac{1}{[2]} + 2 \frac{[3]}{[6]} \frac{[2][2]}{[4][1]} \frac{1}{[2]^2} + \frac{[2]^2}{[4][3]} \frac{1}{[2]} + 2 \frac{[2]}{[4]} \frac{1}{[2]^2} + \frac{1}{[2]^2} = \frac{[3]^3}{[6][4]}.
$$

<span id="page-22-0"></span>**Example 4.18.** We calculate the coefficient of  $E_0E_1E_3$  in  $Q_3$ . The diagram for  $E_0E_1E_3$  and its corresponding top and bottom Dyck paths are given by



We have eight bi-colored vertical Hermite histories.



The sum of the weight is given by

<span id="page-22-1"></span>(4.5)

$$
\frac{[3]^2}{[6][4]} \left( \frac{1}{[3][2]} + \frac{1}{[2]} \right) + 2 \frac{[3]}{[6]} \frac{[2]^2}{[4][3]} \frac{1}{[2]} + 2^2 \frac{[3]}{[6]} \frac{[2]}{[4]} \frac{1}{[2]^2} \n+ \frac{[3]}{[6][4]} \left( \frac{1}{[3][2]} + \frac{1}{[2]} \right) + \frac{[3][4]}{[6][2]} \frac{[2]^2}{[4][3]} \frac{1}{[2]} + 2 \frac{[3][4]}{[6][2]} \frac{[2]}{[4]} \frac{1}{[2]^2} \n+ 2 \frac{[3]}{[6]} \frac{1}{[2]^2} + \frac{[3][4]}{[6][2]} \frac{1}{[2]^2} \n= \frac{[2]^3}{[6][4]} \left( [2]^2 + 1 \right).
$$

There are two non-admissible bi-colored Hermite histories of Dyck tilings:

<span id="page-23-1"></span>

The both bi-colored Hermite histories have three (removed) trajectories right to the green trajectory. Then, the numbers of Dyck tiles in these three trajectories are  $(n_1, n_2, n_3) = (0, 1, 0)$  from left to right if we apply the deletion of the Dyck tilings. They are not admissible since  $n_1 < n_2$  violates  $(Q_4)$ .

In what follows, we consider the practical reason why the condition  $(Q_4)$  on a bi-colored vertical Hermite history is necessary with an example.

For simplicity, we consider the two non-admissible Dyck tilings in Eq. [\(4.6\)](#page-23-1) of Example [4.18.](#page-22-0) The left non-admissible Dyck tiling: The green vertical trajectory corresponds to the coefficient  $\mathrm{coef}(h_{3,1})$ . Then, diagrammatically, we calculate

<span id="page-23-2"></span>(4.7) 
$$
\bigcup_{D} \bigcup_{D} \longrightarrow \mathrm{coef}(h_{3,1}) \cdot \bigcup_{A},
$$

where the green cap corresponds to the green trajectory. The right-hand side of Eq.  $(4.7)$  implies that we have a diagram of type A with three strands. We compare this situation with the left Dyck tiling in Eq. [\(4.6\)](#page-23-1). This Dyck tiling contains three trivial Dyck tiles on the green trajectory, and a unique non-trivial Dyck tile of size 1. This non-trivial Dyck tile has no trajectory, and this means that this tile corresponds to type A. The region of type A corresponds to a single tile at height 1 since this tile is a non-trivial Dyck tile. By shrinking the Dyck tile, it is easy to see that this Dyck tile corresponds to the following diagram of type A:

A

This contains only two strands and does not fit to the situation in Eq. [\(4.7\)](#page-23-2).

The right Dyck tiling in Eq. [\(4.6\)](#page-23-1) is not admissible in the same reason.

On the contrary, we consider the left-most admissible Dyck tiling in the second row in Eq. [\(4.5\)](#page-22-1). The green trajectory corresponds to  $\text{coef}(h_{3,3})$ . As a diagram, we calculate

<span id="page-23-3"></span>(4.8) 
$$
\bigcup_{D} \bigcup_{D} \text{coeff}(h_{3,3}) \cdot \bigcup_{A} \Big|_{A}.
$$

The Dyck tiling contains one Dyck tile with a green trajectory, and three Dyck tiles without trajectories. These three Dyck tiles consist of two Dyck tiles of height 1 and one Dyck tiles of height 2. This diagram of type A indeed corresponds to the diagram in the right-hand side of Eq. [\(4.8\)](#page-23-3). Therefore, this Dyck tiling is admissible.

## 5. Some explicit expressions of the coefficients

<span id="page-23-0"></span>In this section, we give some explicit expressions of the coefficients of the elements in Jones–Wenzl projections. We write  $\operatorname{coeff}^n(A)$  as a coefficient of A in the projection  $Q_n$ .

We first study the symmetry of the coefficients of the diagram in the Jones–Wenzl projection  $Q_n$ .

Let D be an  $n+1$ -strand Temperley–Lieb diagram in  $TL_{n+1}^D$ . The diagram D has an algebraic representation, that is, D corresponds to the word  $E_D := E_{s_1} \cdots E_{s_m}$  where  $E_D$  is a reduced expression. Given a word  $E_D$ , we introduce a transposed word  $E_D := E_{s_m} \cdots E_{s_1}$ .

<span id="page-24-4"></span>**Proposition 5.1.** Let  $E_D$  and  $\overline{E_D}$  be words as above. Then, we have  $\text{coef}^n(E_D) = \text{coef}^n(\overline{E_D})$ . Further, the coefficient of D in  $Q_n$  is equal to that of  $\overline{D}$ , where  $\overline{D}$  is the upside down diagram obtained from D (We change the position of a unique dot in  $\overline{D}$  to satisfy the property (P2b) if  $\overline{D}$ has a unique dot.).

*Proof.* The recurrence relation  $(3.2)$  in Proposition [3.3](#page-7-0) can be rewritten as

$$
Q_{n+1} = \left(\sum_{i=1}^{n+2} \mathrm{coeff}(g_{n+1,i})\overline{g_{n+1,i}} + \sum_{j=0}^{n+1} \mathrm{coeff}(h_{n+1,j})\overline{h_{n+1,j}}\right) Q_{n-1}.
$$

This implies that  $\mathrm{coef}(E_D) = \mathrm{coef}(\overline{E_D})$ . Since the diagram  $\overline{D}$  is nothing but a diagram for  $\overline{E_D}$ , the second statement follows.  $\Box$ 

Let  $E_D := E_{s_1} \dots E_{s_m} \in TL_{n+1}^D$  be a word of  $E_i$ ,  $0 \le i \le n$ . We define an involution  $\theta : E_D \mapsto$  $E_{D'}$  where  $E_{D'}$  is obtained from  $E_D$  by replacing  $E_0$  by  $E_1$  and  $E_1$  by  $E_0$  in  $E_D$ .

<span id="page-24-3"></span>**Proposition 5.2.** We have  $\mathrm{coef}^n(E_D) = \mathrm{coef}^n(E_{D'})$ .

*Proof.* Note that  $\theta$  is an involution of the Dynkin diagram  $D_{n+1}$  such that we exchange the labels 0 and 1 and fix other labels. The statement immediately follows from this fact.

We introduce the following summation formula which is used later.

<span id="page-24-1"></span>Lemma 5.3. We have

$$
\sum_{k=j+1}^{n} \frac{[k][k-1]}{[2k][2(k-1)]} = \frac{[n][j][n-j]}{[2n][2j]}.
$$

*Proof.* By induction on *n*, one can easily prove the claim.

We first consider some cases where the coefficient is expressed by a simple fraction of  $q$ -integers. We first consider the reduced expressions without the product  $E_0E_1$  in its algebraic representation.

<span id="page-24-2"></span>**Proposition 5.4.** Let i and j be positive integers such that  $1 \leq i \leq j \leq n$ . We have

<span id="page-24-0"></span>(5.1) 
$$
\operatorname{coef}^n(E_j E_{j-1} \dots E_i) = \begin{cases} \frac{[n][n+1]}{[2n][2]}, & \text{if } i = j = 1, \\ \frac{[n][2(i-1)][n+1-j]}{[2n][i-1]}, & \text{otherwise.} \end{cases}
$$

*Proof.* We prove the second cases in Eq.  $(5.1)$  by induction on n since the first case can be proven in a similar way. By use of Eq. [\(3.7\)](#page-9-1) in Proposition [3.7](#page-9-0) and the induction hypothesis, we have

$$
\begin{split} \text{coef}^n(E_j \cdots E_i) &= \frac{[n][2(i-1)]}{[2n][i-1]} \text{coef}^{n-1}(E_{n-1}E_{n-2}\cdots E_j) + \text{coef}^{n-1}(E_j \cdots E_i), \\ &= \frac{[n][n-1][2(i-1)][2(j-1)]}{[2n][2(n-1)][i-1][j-1]} + \text{coef}^{n-1}(E_{n-1}E_{n-2}\cdots E_j), \\ &= \left(\sum_{k=j+1}^n \frac{[k][k-1]}{[2k][2(k-1)]}\right) \frac{[2(i-1)][2(j-1)]}{[i-1][j-1]} + \frac{[j][2(j-1)]}{[2j][j-1]}, \\ &= \frac{[n][2(i-1)][n+1-j]}{[2n][i-1]}, \end{split}
$$

where we have used Lemma [5.3.](#page-24-1)

<span id="page-25-4"></span>In what follows, we consider the coefficients of elements which contains a product  $E_0E_1$ . **Proposition 5.5.** Let  $D = E_j E_{j-1} \cdots E_1 E_0 E_2 \cdots E_i$  with  $j \ge 1$  and  $i \ge 2$ . We have

<span id="page-25-0"></span>(5.2) 
$$
\qquad \qquad \text{coef}^n(D) = \frac{[n][n+1-i][n+1-j]}{[2n][n+1]}.
$$

Further, we have

<span id="page-25-3"></span>(5.3) 
$$
\operatorname{coef}^n(E_0E_1) = \frac{[n]^3}{[2n][n+1]}.
$$

*Proof.* We prove Eq. [\(5.2\)](#page-25-0) by induction on n. For  $n = 1, 2$ , one can show that Eq. (5.2) holds by a simple calculation.

We make use of the recurrence relation  $(3.1)$  in Proposition [3.2.](#page-6-2) We calculate the contributions to coef<sup>n</sup>(D) from the three terms in the right hand side of Eq.  $(3.1)$ . From the first term, we have  $\mathrm{coef}^{n-1}(D)$ . Therefore, by induction hypothesis, we have

<span id="page-25-1"></span>(5.4) 
$$
\qquad \qquad \text{coef}^{n-1}(D) = \frac{[n-1][n-j][n-i]}{[2n-2][n]}.
$$

From the second term, we have three cases:

- (1)  $D = (E_j \cdots E_1 E_0 E_2 \cdots E_{n-1}) \cdot E_n \cdot (E_{n-1} \cdots E_i),$
- (2)  $D = (E_j \cdots E_1 E_0 E_2 \cdots E_{n-1}) \cdot E_n \cdot (E_{n-1} \cdots E_1 E_0 E_2 \cdots E_i)$
- (3)  $D = (E_j E_{j+1} \cdots E_{n-1}) \cdot E_n \cdot (E_{n-1} \cdots E_1 E_0 E_2 \cdots E_i),$

In the case of (2), we have an extra factor  $-[2]$ . By Eq. [\(5.1\)](#page-24-0), the three cases (1) to (4) contribute to  $Q_n$  as

<span id="page-25-2"></span>(5.5)  

$$
\frac{[n][2n-2]}{[2n][n-1]} \left\{ \frac{[n-1][n-j]}{[2n-2][n]} \left( \frac{[n-1][2i-2]}{[2n-2][i-1]} - [2] \frac{[n-1][n-i]}{[2n-2][n]} \right) + \frac{[n-1][n-i]}{[2n-2][n]} \frac{[n-1][2j-2]}{[2n-2][j-1]} \right\}.
$$

Let  $D_{j,i} := E_j E_{j-1} \cdots E_1 E_0 E_2 \cdots E_i$ . From the third term in Eq. [\(3.1\)](#page-6-1), we have  $D = D' E_{w_n} D''$ where D is either  $E_jE_{j+1}\cdots E_{n-1}$  or  $D_{j,n-1}$ , and D'' is either  $E_{n-1}E_{n-2}\cdots E_i$  or  $D_{n-1,i}$ . Note

that if we choose  $D_{j,n-1}$  or  $D_{n-1,i}$ , then  $D'E_{w_n}D''$  has a factor  $-[2]$ . Therefore, the contribution to  $Q_n$  is given by

<span id="page-26-0"></span>
$$
(5.6) \qquad \frac{[n]}{[2n][n+1]} \left( \frac{[n-1][2j-2]}{[2n-2][j-1]} - [2] \frac{[n-1][n-j]}{[2n-2][n]} \right) \left( \frac{[n-1][2i-2]}{[2n-2][i-1]} - [2] \frac{[n-1][n-i]}{[2n-2][n]} \right).
$$

By taking the sum of Eqs.  $(5.4)$ ,  $(5.5)$  and  $(5.6)$ , we obtain Eq.  $(5.2)$ . Equation  $(5.3)$  can be shown in a similar way.  $\Box$ 

In some cases, the coefficient of an element in  $Q_n$  is given by a simple fraction of q-integers as we have seen Propositions [5.4](#page-24-2) and [5.5.](#page-25-4) However, in general, the coefficients are the sum of fractions of q-integers. A typical example of such coefficients is  $\text{coef}^n(E_0E_1E_3)$ .

<span id="page-26-1"></span>**Proposition 5.6.** The coefficient  $\text{coef}^n(E_0E_1E_3)$  is given by

(5.7) 
$$
\operatorname{coef}^n(E_0E_1E_3) = \frac{[n-1]^3[n-2][2]^2}{[2n-2][n+1][n]} + 2\frac{[n][n-1]^3[n-2][2]}{[2n][2n-2][n+1]}.
$$

<span id="page-26-3"></span>Before proceeding to the proof of Proposition [5.6,](#page-26-1) we will calculate the necessary coefficients.

Lemma 5.7. We have

$$
\mathrm{coeff}^n(E_1E_2E_0) = \mathrm{coeff}^n(E_0E_2E_1) = \frac{[n][n-1]}{[2n][2]}.
$$

*Proof.* We make use of Eq.  $(3.2)$  in Proposition [3.3.](#page-7-0) Since the word  $E_0E_2E_1$  is rewritten as

$$
E_0E_2E_1 = E_0E_2E_3\cdots E_{n-1} \cdot E_nE_{n-1}\cdots E_1,
$$

we have

$$
\begin{aligned}\n\text{coef}^n(E_0 E_2 E_1) &= \text{coef}^{n-1}(E_0 E_2 \cdots E_{n-1}) \text{coef}^n(g_{n,1}) + \text{coef}^{n-1}(E_0 E_2 E_1), \\
&= \frac{[n][n-1]}{[2n][2n-2]} + \text{coef}^{n-1}(E_0 E_1 E_2), \\
&= \sum_{k=2}^n \frac{[k][k-1]}{[2k][2k-2]}, \\
&= \frac{[n][n-1]}{[2n][2]},\n\end{aligned}
$$

where we have used Lemma [5.3.](#page-24-1) By applying  $\theta$  on  $E_0E_2E_1$ , and Proposition [5.2,](#page-24-3) we have the equality  $\text{coef}^n(E_0E_2E_1) = \text{coef}^n(E_1E_2E_0)$ , which completes the proof.

<span id="page-26-4"></span>Lemma 5.8. We have

<span id="page-26-2"></span>(5.8) 
$$
\operatorname{coef}^n(E_n E_{n-1} \cdots E_3 E_1 E_0) = \frac{[n-1]^2[2]}{[2n][n+1]} + \frac{[n-1]^3[2]^2}{[2n][2n-2]}.
$$

*Proof.* We make use of Eq. [\(3.2\)](#page-7-1). The word  $E_nE_{n-1}\cdots E_3E_1E_0$  can be expressed as products of two words  $H_1H_2$  where  $H_1$  is in  $Q_{n-1}$  and  $H_2$  is either  $g_{n,i}$  or  $h_{n,j}$ . More precisely, we have if  $H_2$ is either  $h_{n,1}$  or  $h_{n,3}$ , then  $H_1$  is either  $E_0$ ,  $E_1$ ,  $E_1E_0$ ,  $E_1E_2E_0$  or  $E_0E_2E_1$ . Similarly, if  $H_2$  is either  $g_{n,1}, h_{n,0}$  or  $g_{n,3}$ , then  $H_1$  is  $E_0E_1$ . By taking the sum of all contributions, we have Eq. [\(5.8\)](#page-26-2).

<span id="page-26-5"></span>Lemma 5.9. We have

$$
\mathrm{coeff}^n(E_n E_{n-1} \cdots E_3 E_1) = \mathrm{coeff}^n(E_n E_{n-1} \cdots E_3 E_0) = \frac{[n]^2 [n-1][3]}{[2n][2n-2][2]}.
$$

*Proof.* The word  $E_n \cdots E_3 E_1$  can be expressed as a product of two words  $H_1 H_2$  where  $H_1 = E_1$ and  $H_2$  is either  $g_{n,3}$  or  $g_{n,1}$ . By Eq. [\(3.2\)](#page-7-1), we have

$$
\mathrm{coeff}^n(E_n\cdots E_3E_1)=\mathrm{coeff}^{n-1}(E_1)(\mathrm{coeff}(g_{n,3})+\mathrm{coeff}(g_{n,1}))=\frac{[n]^2[n-1][3]}{[2n][2n-2][2]},
$$

<span id="page-27-2"></span>where we have used Eq.  $(5.1)$ .

Lemma 5.10. We have

<span id="page-27-0"></span>(5.9) 
$$
\operatorname{coef}^n(E_0 E_n E_{n-1} \cdots E_1) = \operatorname{coef}^n(E_1 E_n E_{n-1} \cdots E_2 E_0) = \frac{[n]^2 [n-1]}{[2n][2n-2][2]}.
$$

*Proof.* The word  $E_0E_nE_{n-1}\cdots E_1$  is uniquely written as a product of two words  $H_1H_2$  where  $H_1 = E_0$  and  $H_2 = g_{n,1}$ . By Eq. [\(3.2\)](#page-7-1),  $\operatorname{coeff}^n(E_0 g_{n,1}) = \operatorname{coeff}^{n-1}(E_0) \operatorname{coeff}^n(g_{n,1})$ . From Propositions [5.1](#page-24-4) and [5.4,](#page-24-2) and the definition of  $\operatorname{coef}^n(g_{n,1})$ , we have Eq. [\(5.9\)](#page-27-0).

<span id="page-27-3"></span>Lemma 5.11. We have

<span id="page-27-1"></span>(5.10) 
$$
\operatorname{coef}^n(E_0E_2E_1g_{n,3}) = \operatorname{coef}^n(E_1E_2E_0g_{n,3}) = \frac{[n][n-1][n-2][3]}{[2n][2n-2][2]}.
$$

*Proof.* Note that the word  $E_0E_2E_1g_{n,3}$  can be also expressed as  $E_0E_2E_1g_{n,1}$ . Therefore, we have two expressions for  $E_0E_2E_1g_{n,3}$ . By Eq. [\(3.2\)](#page-7-1) and Lemma [5.7,](#page-26-3) we have Eq. [\(5.10\)](#page-27-1).

<span id="page-27-4"></span>Lemma 5.12. We have

$$
\mathrm{coeff}^n(E_1E_2E_0g_{n,1}) = \mathrm{coeff}^n(E_0E_2E_1h_{n,0}) = \frac{[n][n-1][n-2]}{[2n][2n-2][2]}.
$$

*Proof.* Since there is no other ways to describe the word  $E_1E_2E_0g_{n,1}$  in terms of a product of words  $H_1$  and  $H_2$  such that  $H_1$  is in  $Q_{n-1}$  and  $H_2$  is either  $g_{n,i}$  or  $h_{n,j}$ , we have

$$
\mathrm{coeff}^n(E_1E_2E_0g_{n,1}) = \mathrm{coeff}^{n-1}(E_1E_2E_0)\mathrm{coeff}(g_{n,1}) = \frac{[n][n-1][n-2]}{[2n][2n-2][2]},
$$

where we have used Lemma [5.7](#page-26-3) and Eq.  $(3.3)$ .

We are ready to prove Proposition [5.6.](#page-26-1)

*Proof of Proposition [5.6.](#page-26-1)* We rewrite  $E_0E_1E_3$  as a product  $H_1H_2$  of two words  $H_1$  and  $H_2$  such that  $H_1$  is in  $Q_{n-1}$  and  $H_2$  is either  $g_{n,i}$  or  $h_{n,j}$ . Below, we list up all the possibilities of a pair of two elements  $H_1$  and  $H_2$ .

Suppose that  $H_1 = E_0E_1\overline{g_{n-1,3}}$ . Then,  $H_2$  is either  $g_{n,1}$ ,  $h_{n,0}$  or  $g_{n,3}$ .

Similarly, if  $H_2$  is either  $h_{n,1}$  or  $h_{n,3}$ , then  $H_1$  is either  $E_0\overline{g_{n-1,3}}$ ,  $E_1\overline{g_{n-1,3}}$ ,  $E_0E_1\overline{g_{n-1,3}}$ ,  $\overline{g_{n-1,1}}E_0$ ,  $h_{n-1,0}E_1, \overline{g_{n-1,3}}E_1E_2E_0, \overline{g_{n-1,3}}E_0E_2E_1, \overline{g_{n-1,1}}E_0E_2E_1, \text{ or } \overline{h_{n-1,0}}E_1E_2E_0.$ 

Finally, if  $H_2$  is the identity, then  $H_1 = E_0 E_1 E_3$ .

Note that if  $H_1 = E_0 E_1 \overline{g_{n-1,3}}$  and  $H_2 = h_{n,1}$  or  $h_{n,3}$ , then we have  $H_1 H_2 = -[2]E_0 E_1 E_3$ .

$$
\begin{split} \mathrm{coeff}^{n}(E_{0}E_{1}E_{3}) &= \left(\frac{[n-2]^{2}[2]}{[2n-2][n]} + \frac{[n-2]^{3}[2]^{2}}{[2n-2][2n-4]}\right) \\ &\times \left(2\frac{[n]}{[2n]} + \frac{[n][4]}{[2n][2]} - [2]\frac{[n]^{2}}{[2n][n+1]} - [2]\frac{[n][n-2]}{[2n][n+1]}\right) \\ &+ 2\left(\frac{[n]^{2}}{[2n][n+1]} + \frac{[n][n-2]}{[2n][n+1]}\right) \\ &\times \left(\frac{[n-1]^{2}[n-2][3]}{[2n-2][2n-4][2]} + \frac{[n-1]^{2}[n-2]}{[2n-2][2n-4][2]} + \frac{[n-1][n-2][n-3]}{[2n-2][2n-4][2]} + \frac{[n-1][n-2][n-3]}{[2n-2][2n-4][2]}\right) \\ &+ \mathrm{coeff}^{-1}(E_{0}E_{1}E_{3}), \\ &= \sum_{k=2}^{n} \left(\frac{[k-2]^{2}[2]^{3}}{[2k-2][k][k+1]} + \frac{[k-2]^{3}[2]^{4}}{[2k-2][2k-4][k+1]} + 2\frac{[k][k-1]^{2}[k-2]^{2}[2]^{3}}{[2k][2k-2][2k-4][k+1]}\right), \\ &= \frac{[n-1]^{3}[n-2][2]^{2}}{[2n-2][n+1][n]} + 2\frac{[n][n-1]^{3}[n-2][2]}{[2n][2n-2][n+1]}, \end{split}
$$

<span id="page-28-0"></span>where we have used Lemmas [5.7,](#page-26-3) [5.8,](#page-26-4) [5.9,](#page-26-5) [5.10,](#page-27-2) [5.11,](#page-27-3) [5.12](#page-27-4) and Proposition [5.1.](#page-24-4)  $\Box$ 

# APPENDIX A. THE JONES–WENZL PROJECTION  $Q_3$

The Jones–Wenzl projection  $Q_3$  for  $TL_4^D$  is given as follows:

By Eq.  $(3.2)$ , we have

$$
Q_3 = 1 + \frac{[3][4]}{[6][2]}(E_0 + E_1 + E_3) + \frac{[3][2]^2}{[6]}E_2 + \frac{[3]^3}{[6][4]}(E_1E_0 + E_3E_0 + E_3E_1)
$$
  
+ 
$$
\frac{[3][2]}{[6]}(E_2E_0 + E_2E_1 + E_0E_2 + E_1E_2 + E_3E_2 + E_2E_3)
$$
  
+ 
$$
\frac{[3]^2[2]}{[6][4]}(E_2E_1E_0 + E_2E_3E_0 + E_2E_3E_1 + E_1E_0E_2 + E_3E_0E_2 + E_3E_1E_2)
$$
  
+ 
$$
\frac{[3]}{[6]}(E_1E_2E_0 + E_3E_2E_0 + E_0E_2E_1 + E_3E_2E_1 + E_0E_2E_3 + E_1E_2E_3)
$$
  
+ 
$$
\frac{[2]^3([2]^2 + 1)}{[6][4]}E_3E_1E_0
$$
  
+ 
$$
\frac{[3]^2}{[6][4]}(E_3E_2E_1E_0 + E_3E_1E_2E_0 + E_1E_2E_3E_0 + E_0E_3E_2E_1 + E_0E_2E_3E_1 + E_1E_0E_2E_3)
$$
  
+ 
$$
\frac{[3]^2}{[6][4]}(E_2E_1E_0E_2 + E_2E_3E_0E_2 + E_2E_3E_1E_2)
$$
  
+ 
$$
\frac{[2]^2([2]^2 + 1)}{[6][4]}(E_2E_3E_1E_0 + E_3E_1E_0E_2)
$$
  
+ 
$$
\frac{[3][2]}{[6][4]}(E_2E_3E_1E_2 + E_2E_0E_3E_2E_1 + E_3E_2E_1E_0E_2 + E_1E_2E_3E_0E_2 + E_0E_2E_3E_1E_2 + E_2E_3E_1E_2 + E_2E_2E_3E_1E_2 + E_2E_2E_3E_1E_2 + E_2E_2E_3E_1E_2 + E_2E_2E_3E_1E_2 + E_2E
$$

+ 
$$
\frac{[2]([2]^2+1)}{[6][4]}E_2E_3E_1E_0E_2
$$
  
+ 
$$
\frac{[3]}{[6][4]}(E_0E_2E_3E_1E_2E_0 + E_1E_2E_3E_0E_2E_1 + E_3E_2E_1E_0E_2E_3).
$$

## **REFERENCES**

- <span id="page-29-9"></span><span id="page-29-6"></span>[1] J. Baine, On the coefficients in the Jones–Wenzl idemotent, preprint (2024), 13 pages,  $arXiv:2406.06333$ .
- <span id="page-29-16"></span>[2] B. Elias and G. Williamson, Soergel Calculus, Represent. Theory 20 (2016), 295–374, [arXiv:1309.0865](http://arxiv.org/abs/1309.0865), [doi](http://dx.doi.org/10.1090/ert/481).
- <span id="page-29-5"></span>[3] C. K. Fan, A Hecke algebra quotient and some combinatorial applications, J. Algebraic Combin. 5 (1996), no. 3, 175–189, [doi](http://dx.doi.org/10.1023/A:1022443327568).
- [4] I. B. Frenkel and M. G. Khovanov, Canonical bases in tensor products and graphical calculus for  $U_q(\mathfrak{sl}_2)$ , Duke Math. J. 87 (1997), no. 3, 409–480.
- <span id="page-29-15"></span><span id="page-29-0"></span>[5] R. M. Green, Generalized Temperley–Lieb algebras and decorated tangles, J. Knot Theory Ramifications 7 (1998), 155–171, [arXiv:q-alg/9712018](http://arxiv.org/abs/q-alg/9712018).
- <span id="page-29-4"></span>[6] V. F. R. Jones, *Index for subfactors*, Invent. Math. **72** (1983), 1–25.
- <span id="page-29-1"></span>[7] , Hecke algebra representations of braid groups and link polynomials, Ann. of Math. 126 (1987), no. 2, 335–388.
- <span id="page-29-11"></span>[8] , On knot invariants related to some statistical mechanical models, Pacific J. Math. 137 (1989), no. 2, 311–334.
- [9] R. W. Kenyon and D. B. Wilson, Double-dimer pairings and skew Young diagrams, Electron. J. Combin. 18 (2011), no. 1, P130, [arXiv:1007.2006](http://arxiv.org/abs/1007.2006).
- <span id="page-29-12"></span>[10] J. S. Kim, K. Mészáros, G. Panova, and D. B. Wilson, Dyck tilings, increasing trees, descents and inversions, J. Combin. Theory Ser. A 122 (2014), 9-27,  $arXiv:1205.6578$ .
- <span id="page-29-13"></span><span id="page-29-7"></span>[11] S. Morrison, A formula for the Jones–Wenzl projections, Proc. Centre Math. Appl. (2017), 367–378, [arXiv:1503.00384](http://arxiv.org/abs/1503.00384).
- [12] P. Sentinelli, The Jones-Wenzl idempotent of a generalized Temperley-Lieb algebra, Journal of Algebra 528 (2019), 505–524, [doi](http://dx.doi.org/https://doi.org/10.1016/j.jalgebra.2019.03.008).
- <span id="page-29-10"></span><span id="page-29-8"></span>[13] K. Shigechi, Jones–Wenzl projections and Dyck tilings: type A and B, preprint (2024), 12 pages, [arXiv:2402.09887](http://arxiv.org/abs/2402.09887).
- [14] K. Shigechi and P. Zinn-Justin, Path representation of maximal parabolic Kazhdan–Lusztig polynomials, J. Pure Appl. Alegebra 216 (2012), no. 11, 2533-2548, [arXiv:1001.1080](http://arxiv.org/abs/1001.1080).
- <span id="page-29-17"></span><span id="page-29-14"></span>[15] J. R. Stembridge, On the Fully Commutative Elements of Coxeter Groups, J. Algebr. Comb. 5 (1996), 353–385, [doi](http://dx.doi.org/10.1007/BF00193185).
- [16] C. Stroppel and Z. Wojciechowski, *Diagrammatics for the smallest quantum coideal and Jones–Wenzl projectors*, preprint (2024), 20 pages, [arXiv:2406.12132](http://arxiv.org/abs/2406.12132).
- <span id="page-29-3"></span>[17] H. N. V. Temperley and E. H. Lieb, Relations between the 'percolation' and 'coloring' problem and other graphtheoretical problems associated with regular planar lattices: some exact results for the 'percolation problem', Proc. Roy. Soc. Lond. A 322 (1971), 251–280.
- <span id="page-29-2"></span>[18] H. Wenzl, On sequences of projections, C. R. Math. Rep. Acad. Sci. Canada 9 (1987), no. 1, 5–9.

Email address: k1.shigechi AT gmail.com