Polynomial and analytic methods for classifying complexity of planar graph homomorphisms

Jin-Yi Cai As jyc@cs.wisc.edu ama

Ashwin Maran amaran@wisc.edu

Abstract

We introduce some polynomial and analytic methods in the classification program for the complexity of planar graph homomorphisms. These methods allow us to handle infinitely many lattice conditions and isolate the new P-time tractable matrices represented by tensor products of matchgates. We use these methods to prove a complexity dichotomy for 4×4 matrices that says Valiant's holographic algorithm is universal for planar tractability in this setting.

Contents

1	Introduction	1
2	Preliminaries 2.1 Model of Computation 2.2 Definitions	2 2 4
3	Reduction from the Potts Model	5
4	Lattice on Eigenvalues	8
5	Proof of Hardness	13
6	Lattice on Diagonal Entries	18
7	Confluence and Pairwise Order Independence	23
8	Diagonal and Pairwise Order Distinctness	33
9	Hardness for Diagonal Distinct 4 x 4 matrices	45
10	Hardness for non-Diagonal Distinct 4 x 4 matrices	51
11	Forms (I), (III), (IV) and (VI) 11.1 Form (I)	60 60 68 72 88
12	Dichotomy of 4 x 4 matrices	93

1 Introduction

Given graphs G and H, a mapping from V(G) to V(H) is called a homomorphism if edges of G are mapped to edges of H. This is put in a more general or quantitative setting by the notion of a partition function. Let $M = (M_{ij})$ be a $q \times q$ symmetric matrix. In this paper we consider non-negative arbitrary real entries $M_{ij} \in \mathbb{R}_{\geq 0}$; if $M_{ij} \in \{0, 1\}$, then M is the unweighted adjacency matrix of a graph $H = H_M$. Given M, the partition function $Z_M(G)$ for any input undirected multi-graph G = (V, E) is defined as

$$Z_M(G) = \sum_{\sigma: V \to [q]} \prod_{(u,v) \in E} M_{\sigma(u)\sigma(v)}.$$

Obviously isomorphic graphs $G \cong G'$ have the same value $Z_M(G) = Z_M(G')$, and thus every M defines a graph property $Z_M(\cdot)$. For a 0-1 matrix M, $Z_M(G)$ counts the number of homomorphisms from G to H. Graph homomorphism (GH) encompasses a great deal of graph properties [Lov12].

Each M defines a computational problem, denoted by GH(M): given an input graph G compute $Z_M(G)$. The complexity of GH(M) has been a major focus of research. A number of increasingly general complexity dichotomy theorems have been achieved. Dyer and Greenhill [DG00] proved that for any 0-1 symmetric matrix M, computing $Z_M(G)$ is either in P-time or is #P-complete. Bulatov and Grohe [BG05] found a complete classification for GH(M) for all nonnegative matrices M. Goldberg, Grohe, Jerrum, and Thurley [Gol+10] then proved a dichotomy for all real-valued matrices M. Finally, Cai, Chen, and Lu [CCL13] established a dichotomy for all complex valued matrices M. We also note that graph homomorphism can be viewed as a special case of counting CSP. For counting CSP, a series of results established a complexity dichotomy for any set of constraint functions \mathcal{F} , going from 0-1 valued [Bul13; DR10; DR11; DR13] to nonnegative rational valued [Bul+12], to nonnegative real valued [CCL16], to all complex valued functions [CC17].

Parallel to this development, Valiant [Val08] introduced holographic algorithms. It is well known that counting the number of perfect matchings (#PM) is #P-complete [Val79]. On the other hand, since the 60's, there has been a famous FKT algorithm [Kas61; TF61; Kas63; Kas67] that can compute #PM on planar graphs in P-time. Valiant's holographic algorithms greatly extended its reach, in fact so much so that a most intriguing question arises: Is this a *universal* algorithm that every counting problem expressible as a sum-of-products that can be solved in P-time on planar graphs (but #P-hard in general) is solved by this method alone? Such a universality statement must appear to be extraordinarily, if not overly, ambitious.

After a series of work [CLX09; CGW16; CF19; Bac17; Bac18; YF22; FYY19; FY14; CF19] it was established that for every set of complex valued constraint functions \mathcal{F} on the Boolean domain (i.e., q = 2) there is a 3-way exact classification for #CSP(\mathcal{F}): (1) P-time solvable, (2) P-time solvable over planar graphs but #P-hard over general graphs, (3) #P-hard over planar graphs. Moreover, category (2) consists of precisely those problems that can be solved by Valiant's holographic algorithm using FKT. Cai and Maran [CM23] showed that for GH the same 3-way exact classification holds even on the domain q = 3, and category (2) again consists of precisely those problems that can be solved by Valiant's holographic algorithm using FKT. So far little is known for higher domain problems (q > 3) on this universality question. Let P1-GH(M) denote the problem GH(M) when the input graphs are restricted to planar graphs. Planar GH is also intimately related to quantum isomorphism of graphs, a relaxation of classical isomorphism [Ats+19]. It is known that graphs H and H' are quantum isomorphic iff there is a perfect winning strategy in a two-player graph isomorphism game in which the players share and perform measurements on an entangled quantum state. This is also equivalent to the existence of a quantum permutation matrix transforming H to H' [LMR20]. Let M and M' be the adjacency matrices of H and H'. Mančinska and Roberson [MR20] proved that H and H' are quantum isomorphic iff $Z_M(G) = Z_{M'}(G)$ for every planar graph G, i.e., H and H' define the same Planar GH problem. Furthermore, a fascinating consequence of this line of work is that it is undecidable whether P1-GH(M) = P1-GH(M') [MR20], which hints at the difficulty that we face in this paper.

Our goal is to classify the complexity of Pl-GH(M), i.e., when the input G is restricted to be planar for $Z_M(G)$. (The underlying graph H_M is not restricted to planar graphs.) We want to classify the problems Pl-GH(M): What is the computational complexity of $Z_M(G)$ from planar input graphs G? We present some strong polynomial and analytic techniques that will help us approach this problem. We demonstrate the power of these techniques by giving a complete classification of the complexity of Pl-GH(M) for all non-negative real valued full rank 4×4 matrices M. The full rank 4×4 case is particularly important as it is the first case where tensor product of matchgates [Val08; Val01] defines new P-time tractable problems. We prove that an exact classification according to the three categories above holds for this class, and a holographic reduction to FKT remains a *universal* algorithm for category (2).

2 Preliminaries

2.1 Model of Computation

The Turing machine model is naturally suited to the study of computation over discrete structures such as integers or graphs. When $M \in \mathbb{R}^{q \times q}$, for Pl-GH(M) one usually restricts M to be a matrix with only algebraic numbers. This is strictly for the consideration of the model of computation, even though allowing all real-valued matrices would be more natural.

There is a formal (albeit nonconstructive) method to treat Pl-GH(M) for arbitrary real-valued matrices M and yet stay strictly within the Turing machine model in terms of bit-complexity. In this paper, because our proof depends heavily on analytic argument with continuous functions on the real line, this logical formal view becomes necessary.

To begin with, we recall a theorem from field theory: Every extension field \mathbf{F} over \mathbb{Q} by a finite set of real numbers is a finite algebraic extension \mathbf{E}' of a certain purely transcendental extension field \mathbf{E} over \mathbb{Q} , which has the form $\mathbf{E} = \mathbb{Q}(X_1, \ldots, X_m)$ where $m \ge 0$ and X_1, \ldots, X_m are algebraically independent [Jac85] (Theorem 8.35, p. 512). \mathbf{F} is said to have a finite transcendence degree m over \mathbb{Q} . It is known that m is uniquely defined for \mathbf{F} . Since char $\mathbb{Q} = 0$, the finite algebraic extension \mathbf{E}' over \mathbf{E} is actually simple, $\mathbf{E}' = \mathbf{E}(\beta)$ for some β , and it is specified by a minimal polynomial in $\mathbf{E}[X]$. Now given a real matrix M, let $\mathbf{F} = \mathbb{Q}(M)$ be the extension field by adjoining the entries of M. We consider M is fixed for the problem P1-GH(M), and thus we may assume (nonconstructively) that the form $\mathbf{F} = \mathbf{E}(\beta)$ and $\mathbf{E} = \mathbb{Q}(X_1, \dots, X_m)$ are given. (This means, among other things, that the minimal polynomial of β over \mathbf{E} is given, and all arithmetic operations can be performed on \mathbf{F} .)

Now, the computational problem Pl-GH(M) is the following: Given a planar G, compute $Z_M(G)$ as an element in \mathbf{F} (which is expressed as a polynomial in β with coefficients in \mathbf{E}). More concretely, we can show that this is equivalent to the following problem COUNT(M): The input is a pair (G, x), where G = (V, E) is a planar graph and $x \in \mathbf{F}$. The output is

$$\#_M(G,x) = \left| \left\{ \sigma : V \to [q] : \prod_{(u,v) \in E} m_{\sigma(u),\sigma(v)} = x \right\} \right|,$$

a non-negative integer. Note that, in this definition, we are basically combining terms with the same product value in the definition of $Z_M(G)$.

Let n = |E|. Define X to be the set of all possible product values appearing in $Z_M(G)$:

$$X = \left\{ \prod_{i,j \in [q]} m_{ij}^{k_{ij}} \mid \text{integers } k_{ij} \ge 0 \text{ and } \sum_{i,j \in [q]} k_{ij} = n \right\}.$$
 (1)

There are $\binom{n+q^2-1}{q^2-1} = n^{O(1)}$ many integer sequences $(k_{i,j})$ such that $k_{i,j} \ge 0$ and $\sum_{i,j \in [q]} k_{i,j} = n$. X is defined as a set, not a multi-set. After removing repeated elements the cardinality |X| is also polynomial in n. For fixed and given **F** the elements in X can be enumerated in polynomial time in n. (It is important that **F** and q are all treated as fixed constants.) It then follows from the definition that $\#_M(G, x) = 0$ for any $x \notin X$. This gives us the following relation:

$$Z_M(G) = \sum_{x \in X} x \cdot \#_M(G, x), \quad ext{for any graph } G,$$

and thus, $Pl-GH(M) \leq COUNT(M)$.

For the other direction, we construct, for any $p \in [|X|]$ (recall that |X| is polynomial in n), a planar graph T_pG from G by replacing every edge of G with p parallel edges. Then,

$$Z_M(T_pG) = \sum_{x \in X} x^p \cdot \#_M(G, x), \text{ for any graph } G.$$

This is a Vandermonde system; it has full rank since elements in X are distinct by definition. So by querying Pl-GH(M) for the values of $Z_M(T_pG)$, we can solve it in polynomial time and get $\#_M(G, x)$ for every non-zero $x \in X$. To obtain $\#_M(G, 0)$ (if $0 \in X$), we note that

$$\sum_{x \in X} \#_M(G, x) = q^{|V|}.$$

This gives us a polynomial-time reduction and thus, $COUNT(M) \leq Pl-GH(M)$. We have proved

Lemma 1. For any fixed $M \in \mathbb{R}^{q \times q}$, $Pl-GH(M) \equiv COUNT(M)$.

Thus, Pl-GH(M) can be identified with the problem of producing those polynomially many integer coefficients in the canonical expression for $Z_M(G)$ as a sum of (distinct) terms from X.

This formalistic view has the advantage that we can treat the complexity of Pl-GH(M) for general M, and not restricted to algebraic numbers. Thus, numbers such as e or π need not be excluded. More importantly, in this paper this generality is essential, due to the proof technique that we employ. Furthermore, once freed from this restriction we in fact explicitly use transcendental numbers as a tool in our proof (see Lemma 16). In short, in this paper, treating the complexity of Pl-GH(M) for general real M is not a *bug* but a *feature*.

However, we note that this treatment has the following subtlety. For the computational problem P1-GH(M) the formalistic view demands that \mathbf{F} be specified in the form $\mathbf{F} = \mathbf{E}(\beta)$. Such a form exists, and its specification is of constant size when measured in terms of the size of the input graph G. However, in reality many basic questions for transcendental numbers are unknown. For example, it is still unknown whether $e + \pi$ or $e\pi$ are rational, algebraic irrational or transcendental, and it is open whether $\mathbb{Q}(e,\pi)$ has transcendence degree 2 (or 1) over \mathbb{Q} , i.e., whether e and π are algebraically independent. The formalistic view here non-constructively assumes this information is given for \mathbf{F} . A polynomial time reduction $\Pi_1 \leq \Pi_2$ from one problem to another in this setting merely implies that the *existence* of a polynomial time algorithm for Π_2 logically implies the *existence* of a polynomial time algorithm for Π_1 . We do not actually obtain such an algorithm constructively.

This logical detour not withstanding, if a reader is interested only in the complexity of Pl-GH(M) for integer matrices M, then the complexity dichotomy proved in this paper holds according to the standard definition of Pl-GH(M) for integral M in terms of the model of computation; the fact that this is proved in a broader setting for all real matrices M is irrelevant. This is akin to the situation in analytic number theory, where one might be interested in a question strictly about the ordinary integers, but the theorems are proved in a broader setting of analysis.

2.2 Definitions

As we will refer to various different types of matrices throughout this paper, it will be helpful to establish some notation. Given a positive integer $q \ge 1$, we let $\operatorname{Sym}_q(X)$ denote the set of $q \times q$ symmetric matrices such that each entry of the matrix is from the set $X \subset \mathbb{R}$. For example, with $X = \mathbb{R}$ (respectively, $\mathbb{R}_{\ge 0}$, or $\mathbb{R}_{\ne 0}$) $\operatorname{Sym}_q(X)$ denotes $q \times q$ symmetric matrices with real (respectively, non-negative, or non-zero) entries. Similarly, we let $\operatorname{Sym}_q^{\mathsf{F}}(X)$ denote the set of $q \times q$ full rank symmetric matrices such that each entry of the matrix is from the set $X \subset \mathbb{R}$, and let $\operatorname{Sym}_q^{\mathsf{pd}}(X)$ denote the set of $q \times q$ positive definite symmetric matrices with entries from X.

Now, consider some $M \in \text{Sym}_q(\mathbb{R})$ with entries $M_{ij} \in \mathbb{R}$ for $i, j \in [q]$. Given a planar, undirected multi-graph G = (V, E), we can perform certain elementary operations (that preserve planarity) on the graph G to transform it into a new graph G', such that $Z_M(G') = Z_{M'}(G)$ for some matrix M'. For most of this paper we will use two such operations, thickening and stretching.

From any planar multi-graph G = (V, E), and a positive integer n, we can construct the planar multi-graph T_nG , by replacing every edge in G with n parallel edges between the same vertices. This process is called *thickening*. Clearly $Z_M(T_nG) = Z_{T_nM}(G)$, where $T_nM \in \text{Sym}_q(\mathbb{R})$ with entries $((M_{i,j})^n)$ for $i, j \in [q]$. In particular, Pl-GH $(T_nM) \leq \text{Pl-GH}(M)$ for all $n \geq 1$.

Similarly, from any planar multi-graph G = (V, E), and a positive integer n, we can construct the planar multi-graph $S_n G$ by replacing every edge $e \in E$ with a path of length n. This process is called *stretching*. It is also easily seen that $Z_M(S_nG) = Z_{S_nM}(G)$, where $S_nM = M^n$, the *n*-th power of M. So, we also have $\text{Pl-GH}(S_nM) \leq \text{Pl-GH}(M)$ for all $n \geq 1$.

3 Reduction from the Potts Model

We now consider the thickening operation more closely. For $m, n \ge 1$, let

$$\mathcal{P}_m(n) = \left\{ \mathbf{x} = (x_i)_{i \in [m]} \in \mathbb{Z}^m \mid (\forall i \in [m]) \ [x_i \ge 0] \text{ and } \sum_{i \in [m]} x_i = n \right\}.$$

We note that given any graph G = (V, E),

$$Z_M(T_nG) = Z_{T_nM}(G) = \sum_{x \in X(G)} x^n \cdot \#_M(G, x)$$
⁽²⁾

where

$$X(G) = \left\{ \prod_{i,j \in [q]} M_{ij}^{k_{ij}} \mid \mathbf{k} = (k_{ij})_{i,j \in [q]} \in \mathcal{P}_{q^2}(|E|) \right\},$$
(3)

and

$$\#_M(G,x) = \left| \left\{ \sigma : V \to [q] : \prod_{(u,v) \in E} M_{\sigma(u),\sigma(v)} = x \right\} \right|.$$

$$\tag{4}$$

Note that given any $x \in X(G)$, $\#_M(G, x)$ does not depend on n, but depends only on the entries of the matrix M. We will deal with this dependence now.

Definition 2. Let $\mathcal{A} \subseteq \mathbb{R}_{\neq 0}$ be a set of non-zero real numbers. A finite set $\{g_t\}_{t\in[d]} \in (\mathbb{R}_{>1})^d$, for some integer $d \geq 0$, is called a generating set of \mathcal{A} if for every $a \in \mathcal{A}$, there exists a unique $(e_0, e_1, \ldots, e_d) \in \{0, 1\} \times \mathbb{Z}^d$ such that $a = (-1)^{e_0} g_1^{e_1} \cdots g_d^{e_d}$.

Remark. The uniqueness of the exponents implies the following property of $\{g_t\}_{t\in[d]}$: Whenever $\prod_{t\in[d]} g_t^{a_t} = \prod_{t\in[d]} g_t^{b_t}$ we have $a_t = b_t$ for all $t\in[d]$, i.e., any such expression has unique exponents.

Lemma 3. Every finite set $\mathcal{A} \subset \mathbb{R}_{\neq 0}$ of non-zero real numbers has a generating set.

Proof. Consider the multiplicative group \mathcal{G} generated by the positive real numbers $\{|a| : a \in \mathcal{A}\}$. It is a subgroup of the multiplicative group $(\mathbb{R}_{>0}, \cdot)$. Since \mathcal{A} is finite, and $(\mathbb{R}_{>0}, \cdot)$ is torsion-free, the



Figure 1: A graph G, the thickened graph T_3G , and the stretched graph S_2G .

group \mathcal{G} is a finitely generated free Abelian group, and thus isomorphic to \mathbb{Z}^d for some $d \ge 0$. Let f be this isomorphism from \mathbb{Z}^d to the multiplicative group. By flipping ± 1 in \mathbb{Z} we may assume that this isomorphism maps the basis elements of \mathbb{Z}^d to some elements $\{g_t\}_{t\in[d]}$ such that $g_t > 1$ for all $t \in [d]$. The set $\{g_t\}_{t\in[d]}$ is a generating set.

We now use Lemma 3 to find a generating set for the entries $(M_{ij})_{i,j\in[q]}$ of any matrix $M \in$ Sym_q($\mathbb{R}_{\neq 0}$). Note that this generating set need not be unique. However, with respect to a fixed generating set, for any M_{ij} , there are unique integers $e_{ij0} \in \{0,1\}$, and $e_{ij1}, \ldots, e_{ijd} \in \mathbb{Z}$, such that

$$M_{ij} = (-1)^{e_{ij0}} \cdot g_1^{e_{ij1}} \cdots g_d^{e_{ijd}}.$$
 (5)

Remark. It should be noted that since M is symmetric, $M_{ij} = M_{ji}$ for all $i, j \in [q]$. The uniqueness of the integers e_{ijt} in Eq. (5) then implies that for all $i, j \in [q]$, $e_{ijt} = e_{jit}$ for all $t \in [d]$.

Lemma 4. Let $M \in Sym_q(\mathbb{R}_{\neq 0})$ with a generating set $\{g_t\}_{t \in [d]}$ for its entries. There exists an $N = cM \in Sym_q(\mathbb{R}_{\neq 0})$, for some $c \in \mathbb{R}_{>0}$, such that $\mathsf{Pl-GH}(N) \equiv \mathsf{Pl-GH}(M)$, and $\{g_t\}_{t \in [d]}$ is also a generating set for the entries of N, with unique integers $e_{ij0} \in \{0,1\}$ and $e_{ijt} \in \mathbb{Z}_{\geq 0}$ satisfying

$$N_{ij} = (-1)^{e_{ij0}} \cdot g_1^{e_{ij1}} \cdots g_d^{e_{ijd}}.$$
 (6)

Proof. For any $c \neq 0$, and any planar graph G = (V, E), we have

$$Z_{cM}(G) = \sum_{\sigma: V \to [q]} \prod_{(u,v) \in E} (c \cdot M)_{\sigma(u)\sigma(v)} = c^{|E|} \cdot \sum_{\sigma: V \to [q]} \prod_{(u,v) \in E} M_{\sigma(u)\sigma(v)} = c^{|E|} Z_M(G).$$

Therefore, $\text{Pl-GH}(M) \equiv \text{Pl-GH}(cM)$ for all $c \neq 0$. As the entries of M are generated by $\{g_t\}_{t \in [d]}$, we have unique integers $e'_{ij0} \in \{0, 1\}$, and $e'_{ijt} \in \mathbb{Z}$, such that

$$M_{ij} = (-1)^{e'_{ij0}} g_1^{e'_{ij1}} \cdots g_d^{e'_{ijd}}$$

Now let $c = (g_1 \cdots g_d)^{-e} \in \mathbb{R}_{>0}$, where $e = \min_{i,j \in [q], t \in [d]} \{e'_{ijt}\}$.

Clearly, $\{g_t\}_{t\in[d]}$ is also a generating set for the entries of N. If we let $e_{ij0} = e'_{ij0}$, and $e_{ijt} = e'_{ijt} - e$ for all $i, j \in [q]$, and $t \in [d]$, we see that $e_{ijt} \ge 0$, and that

$$(cM)_{ij} = (-1)^{e_{ij0}} \cdot g_1^{e_{ij1}} \cdots g_d^{e_{ijd}}$$

for all $i, j \in [d]$.

Remark. For P1-GH(M), given by $M \in Sym(\mathbb{R}_{\neq 0})$ with generating set $\{g_t\}_{t \in [d]}$, Lemma 4 allows us to replace M with the matrix N = cM whose entries are generated by $\{g_t\}_{t \in [d]}$ such that $e_{ijt} \geq 0$ for all $i, j \in [q]$ and $t \in [d]$. In the following we will often make this substitution when convenient.

Definition 5. Given PI-GH(M) defined by $M \in Sym_q(\mathbb{R}_{\neq 0})$ with entries $(M_{ij})_{i,j\in q}$, we assume a generating set $\{g_t\}_{t\in[d]}$ is chosen and the replacement of N = cM in Lemma 4 has been made so that the integers $e_{ijt} \geq 0$ in Eq. (5). We define the function $\mathcal{T}_M : \mathbb{R}^d \to Sym_q(\mathbb{R})$ such that

$$\mathcal{T}_M(\mathbf{p})_{ij} = (-1)^{e_{ij0}} \cdot p_1^{e_{ij1}} \cdots p_d^{e_{ijd}}$$

is a signed monomial in $\mathbf{p} = (p_1, \ldots, p_d)$ for all $i, j \in [q]$.

	_	_	_	

Lemma 6. Let $M \in Sym_q(\mathbb{R}_{\neq 0})$, with entries $(M_{ij})_{i,j\in q}$ generated by some $\{g_t\}_{t\in [d]}$. Then, Pl-GH $(\mathcal{T}_M(\mathbf{p})) \leq$ Pl-GH(M) for all $\mathbf{p} \in \mathbb{R}^d$.

Proof. Replacing M by N = cM as in Lemma 4, we may assume $\mathcal{T}_M(\mathbf{p})$ is defined in Definition 5 with all $e_{ijt} \ge 0$ in Eq. (5). For any $n \ge 1$, and any given graph G = (V, E),

$$Z_M(T_nG) = \sum_{x \in X(G)} x^n \cdot \#_M(G, x), \tag{7}$$

where X(G) and $\#_M(G, x)$ are given in Eqs. (3) and (4). By definition X(G) is a set, and so each $x \in X(G)$ is distinct, and $|X(G)| \leq |E|^{O(1)}$. By oracle access to Pl-GH(M) we can get $Z_M(T_nG)$ for $n \in [|X(G)|]$. Then Eq. (7) is a full rank Vandermonde system of linear equations, which can be solved in polynomial time to find $\#_M(G, x)$ for all $x \in X(G)$.

Now, let us consider the set X(G) more closely. Given any $x \in X(G)$, we see that $x = \prod M_{ij}^{k_{ij}}$ for some (not necessarily unique) $\mathbf{k} \in \mathcal{P}_{q^2}(|E|)$. Since $\{g_t\}_{t \in [d]}$ is a generating set for the entries of M, any $x \in X(G)$ can be represented uniquely as

$$x = (-1)^{e_0^x} g_1^{e_1^x} \cdots g_d^{e_d^x},$$

with exponents $e_0^x \in \{0, 1\}$, and $e_1^x, \ldots, e_d^x \in \mathbb{Z}_{\geq 0}$.

Fix any $\mathbf{p} = (p_1, \ldots, p_d) \in \mathbb{R}^d$. We define the function $\widehat{y} : X(G) \to \mathbb{R}$, such that

$$\hat{y}(x) = (-1)^{e_0^x} \cdot p_1^{e_1^x} \cdots p_d^{e_d^x}$$

By definition of X(G), we see that for any $x \in X(G)$, there exist $\mathbf{k} \in \mathcal{P}_{q^2}(|E|)$, such that

$$x = \prod_{i,j \in [q]} M_{ij}^{k_{ij}} = (-1)^{\sum_{i,j \in [q]} k_{ij} e_{ij0}} \cdot g_1^{\sum_{i,j \in [q]} k_{ij} e_{ij1}} \cdots g_d^{\sum_{i,j \in [q]} k_{ij} e_{ijd}}$$

By definition of \hat{y} , this means that $\hat{y}(x) = \prod_{i,j \in [q]} \mathcal{T}_M(\mathbf{p})_{ij}^{k_{ij}}$. Now, let

$$Y(G) = \left\{ \prod_{i,j\in[q]} \mathcal{T}_M(\mathbf{p})_{ij}^{k_{ij}} \mid \mathbf{k} = (k_{ij})_{i,j\in[q]} \in \mathcal{P}_{q^2}(|E|) \right\}.$$

Consider any $\sigma: V \to [q]$. For a given σ , we define $\mathbf{k} \in \mathcal{P}_{q^2}(|E|)$ such that $k_{ij} = |\{(u, v) \in E : \sigma(u) = i, \sigma(v) = j\}|$, for all $i, j \in [q]$. Then

$$\widehat{y}\left(\prod_{(u,v)\in E} M_{\sigma(u)\sigma(v)}\right) = \widehat{y}\left(\prod_{i,j\in[q]} M_{ij}^{k_{ij}}\right) = \prod_{i,j\in[q]} \mathcal{T}_M(\mathbf{p})_{ij}^{k_{ij}} = \prod_{(u,v)\in E} \mathcal{T}_M(\mathbf{p})_{\sigma(u)\sigma(v)}$$

This implies that for any $y \in Y(G)$,

$$\left\{\sigma: V \to [q] : \prod_{(u,v) \in E} \mathcal{T}_M(\mathbf{p})_{\sigma(u),\sigma(v)} = y\right\} = \bigsqcup_{\substack{x \in X(G):\\ \widehat{y}(x) = y}} \left\{\sigma: V \to [q] : \prod_{(u,v) \in E} M_{\sigma(u),\sigma(v)} = x\right\}.$$

Therefore, for any $y \in Y(G)$,

$$\#_{\mathcal{T}_{M}(\mathbf{p})}(G, y) = \left| \left\{ \sigma : V \to [q] : \prod_{(u,v) \in E} \mathcal{T}_{M}(\mathbf{p})_{\sigma(u),\sigma(v)} = y \right\} \right|$$
$$= \sum_{x \in X(G): \ \widehat{y}(x) = y} \left| \left\{ \sigma : V \to [q] : \prod_{(u,v) \in E} M_{\sigma(u),\sigma(v)} = x \right\} \right| = \sum_{x \in X(G): \ \widehat{y}(x) = y} \#_{M}(G, x).$$

Having already obtained $\#_M(G, x)$ for all $x \in X(G)$, we can compute in polynomial time

$$\sum_{x \in X(G)} \widehat{y}(x) \cdot \#_M(G, x) = \sum_{y \in Y(G)} y \cdot \sum_{x \in X(G): \widehat{y}(x) = y} \#_M(G, x) = \sum_{y \in Y(G)} y \cdot \#_{\mathcal{T}_M(\mathbf{p})}(G, y) = Z_{\mathcal{T}_M(\mathbf{p})}(G).$$

Therefore, $\operatorname{Pl-GH}(\mathcal{T}_M(\mathbf{p})) \leq \operatorname{Pl-GH}(M)$.

We will now need the following theorem and corollary from [Ver05]:

Theorem 7. For $x, y \in \mathbb{C}$, evaluating the Tutte polynomial at (x, y) is #P-hard over planar graphs unless $(x - 1)(y - 1) \in \{1, 2\}$ or $(x, y) \in \{(1, 1), (-1, -1), (\omega, \omega^2), (\omega^2, \omega)\}$, where $\omega = e^{2\pi i/3}$. In each exceptional case, the problem is in polynomial time.

Corollary 8. The q-state Potts Model Pl-GH(Potts_q(x)) is #P-hard for any integer $q \ge 3$, and real $x \ne 1$, where Potts_q(x) $\in Sym_q(\mathbb{R})$ is the matrix with entries (Potts_q(x)_{ij}) such that Potts_q(x)_{ij} = 1 if $i \ne j$, and Potts_q(x)_{ij} = x otherwise.

Note that $Pl-GH(Potts_q(0))$ is the problem of counting vertex coloring with q colors on planar graphs. Theorem 7 allows us to prove our first hardness result.

Lemma 9. Let $M \in Sym_q(\mathbb{R}_{\neq 0})$, and let \mathcal{T}_M be the function defined in Definition 5. Furthermore, assume for all $i \in [q]$ there exists some (not necessarily distinct) $t(i) \in \{1, \ldots, d\}$, such that $e_{iit(i)} > 0$, and $e_{jkt(i)} = 0$ for all $j \neq k$. Then Pl-GH(M) is #P-hard.

Proof. We apply Lemma 6. Let $\mathbf{p}^* \in \mathbb{R}^d$ defined by $p_t^* = 0$ for all $t \in [d]$, such that t = t(i) for some $i \in [q]$, and $p_t^* = 1$ for all other $t \in [d]$. Then, we see that $\mathcal{T}_M(\mathbf{p}^*)_{ii} = 0$ for all $i \in [q]$, and $\mathcal{T}_M(\mathbf{p}^*)_{ij} = \pm 1$ for all $i \neq j \in [q]$. Therefore, $T_2(\mathcal{T}_M(\mathbf{p}^*)) = \mathsf{Potts}_q(0)$.

From Lemma 6, we get $Pl-GH(\mathcal{T}_M(\mathbf{p}^*)) \leq Pl-GH(M)$. Therefore,

$$\operatorname{Pl-GH}(\operatorname{Potts}_q(0)) \leq \operatorname{Pl-GH}(\mathcal{T}_M(\mathbf{p}^*)) \leq \operatorname{Pl-GH}(M).$$

It follows from Corollary 8 that Pl-GH(M) is #P-hard.

4 Lattice on Eigenvalues

In this section we focus on full ranked matrices. Using stretching, we shall prove the hardness of a more interesting class of matrices than we were able to do in Lemma 9. Consider any $M \in \text{Sym}_q(\mathbb{R})$. There exists some (not necessarily unique) real orthogonal matrix H, and a real diagonal matrix $D = \text{diag}(\lambda_1, \ldots, \lambda_q)$ such that

$$M = HDH^{\mathrm{T}},$$

where $(\lambda_1, ..., \lambda_q)$ are the eigenvalues of M and the columns of H are the corresponding eigenvectors. In the rest of the paper, when we refer to $M = HDH^{\mathsf{T}}$, it is to be understood that we refer to such an orthogonal matrix H, and diagonal matrix D.

From the decomposition $M = HDH^{\mathsf{T}}$, we have $M^n = HD^nH^{\mathsf{T}}$, and

$$(M^n)_{ij} = (H_{i1}H_{j1})\lambda_1^n + \dots + (H_{iq}H_{jq})\lambda_q^n.$$

It follows that

$$Z_{M^n}(G) = \sum_{\mathbf{k}\in\mathcal{P}_q(|E|)} c_H(G,\mathbf{k}) \cdot \left(\lambda_1^{k_1}\cdots\lambda_q^{k_q}\right)^n,\tag{8}$$

where

$$c_H(G, \mathbf{k}) = \sum_{\sigma: V \to [q]} \left(\sum_{\substack{E_1 \sqcup \cdots \sqcup E_q = E \\ |E_i| = k_i}} \left(\prod_{i \in [q]} \prod_{(u,v) \in E_i} H_{\sigma(u)i} H_{\sigma(v)i} \right) \right)$$

depends only on G and the orthogonal matrix H, but not on D.

Before we can analyze Eq. (8) in greater detail, we will need a few more definitions.

Definition 10. Let $\mathcal{A} = (a_1, \ldots, a_n)$ be a tuple of non-zero real numbers (not necessarily distinct). The lattice of \mathcal{A} consists of the set defined as

$$\mathcal{L}(\mathcal{A}) = \left\{ (x_1, \dots, x_n) \in \mathbb{Z}^n \mid \sum_{i=1}^n x_i = 0, \prod_{i=1}^n a_i^{x_i} = 1 \right\},\$$

with addition in \mathbb{Z}^n .

Lemma 11. Let $\mathcal{A} = (a_1, \ldots, a_n)$ be a tuple of non-zero real numbers. The set $\mathcal{L}(\mathcal{A})$ forms a lattice and is isomorphic to \mathbb{Z}^d for some unique integer $0 \le d \le n$.

Proof. We note that $\mathcal{L}(\mathcal{A})$ is a subgroup of the finitely generated (discrete) Abelian group \mathbb{Z}^n . So, $\mathcal{L}(\mathcal{A})$ is itself a finitely generated (discrete) Abelian group, and is torsion-free. It follows that it is a lattice and is isomorphic to \mathbb{Z}^d for some unique $0 \le d \le n$.

Definition 12. Let $\mathcal{A} = (a_1, \ldots, a_n)$ be a tuple of non-zero real numbers. The lattice dimension of \mathcal{A} , denoted by dim($\mathcal{L}(\mathcal{A})$), is the unique integer $d \geq 0$ such that $\mathcal{L}(\mathcal{A}) \cong \mathbb{Z}^d$. A set $\{\mathbf{x}_1, \ldots, \mathbf{x}_d\} \subset \mathcal{L}(\mathcal{A})$, is called a lattice basis of \mathcal{A} if there exists an isomorphism from $\mathcal{L}(\mathcal{A})$ to \mathbb{Z}^d that maps this to a basis of \mathbb{Z}^d .

Remark. Here we note the known fact that if $\mathbb{Z}^d \cong \mathbb{Z}^{d'}$, then d = d'. This fact, together with Lemma 11 guarantees that the lattice dimension of any given tuple of non-zero reals is well-defined.

With the help of these new definitions, we can now go back to studying the effects of the stretching gadget.

Lemma 13. Let $M \in Sym_q^{\mathsf{F}}(\mathbb{R})$, such that $M = HDH^{\mathsf{T}}$, where $D = \operatorname{diag}(\lambda_1, \ldots, \lambda_q)$. Then $\mathsf{Pl-GH}(H\Delta H^{\mathsf{T}}) \leq \mathsf{Pl-GH}(M)$ for any diagonal matrix $\Delta = \operatorname{diag}(\Delta_1, \ldots, \Delta_q)$, such that $\Delta_i \in \mathbb{R}_{\neq 0}$ for all $i \in [q]$, and $\mathcal{L}(\lambda_1, \ldots, \lambda_q) \subseteq \mathcal{L}(\Delta_1, \ldots, \Delta_q)$. *Proof.* We recall that Eq. (8) states that for any graph G = (V, E),

$$Z_M(S_nG) = \sum_{\mathbf{k}\in\mathcal{P}_q(|E|)} c_H(G,\mathbf{k}) \cdot \left(\lambda_1^{k_1}\cdots\lambda_q^{k_q}\right)^n,$$

where

$$c_H(G, \mathbf{k}) = \sum_{\sigma: V \to [q]} \left(\sum_{\substack{E_1 \sqcup \cdots \sqcup E_q = E \\ |E_i| = k_i}} \left(\prod_{i \in [q]} \prod_{(u,v) \in E_i} H_{\sigma(u)i} H_{\sigma(v)i} \right) \right).$$

We now define the set

$$\Lambda_D(G) = \left\{ \prod_{i \in [q]} \lambda_i^{k_i} \middle| \mathbf{k} \in \mathcal{P}_q(|E|) \right\}.$$

For each $\mu \in \Lambda_D(G)$, we then define

$$X_D(G,\mu) = \left\{ \mathbf{k} \in \mathcal{P}_q(|E|) \middle| \prod_{i \in [q]} \lambda_i^{k_i} = \mu \right\} \text{ and } c_{H,D}(G,\mu) = \sum_{\mathbf{k} \in X_D(G,\mu)} c_H(G,\mathbf{k}).$$

Putting everything together, we see that

$$Z_M(S_nG) = \sum_{\mu \in \Lambda_D(G)} c_{H,D}(G,\mu) \cdot \mu^n.$$

Since $\Lambda_D(G)$ is a set, each $\mu \in \Lambda_D(G)$ is distinct, and $|\Lambda_D(G)| \leq |E|^{O(1)}$. With oracle access to P1-GH(M) we can get $Z_M(S_nG)$ for $n \in [|\Lambda_D(G)|]$. Then we have a full rank Vandermonde system of linear equations, which can be solved in polynomial time to find $c_{H,D}(G,\mu)$ for all $\mu \in \Lambda_D(G)$.

Now, we consider

$$Z_{H\Delta H^{\mathsf{T}}}(G) = \sum_{\nu \in \Lambda_{\Delta}(G)} c_{H,\Delta}(G,\nu) \cdot \nu.$$

Consider $\mathbf{k}, \mathbf{l} \in X_D(G, \mu)$ for some $\mu \in \Lambda_D(G)$. By definition, this implies that

$$\prod_{i \in [q]} \lambda_i^{k_i} = \prod_{i \in [q]} \lambda_i^{l_i} = \mu$$

Therefore, $(k_1 - l_1, \ldots, k_q - l_q) \in \mathcal{L}(\lambda_1, \ldots, \lambda_q) \subseteq \mathcal{L}(\Delta_1, \ldots, \Delta_q)$ by our choice of Δ . So, it follows that there exists some $\nu \in \Lambda_{\Delta}(G)$ such that

$$\prod_{i \in [q]} \Delta_i^{k_i} = \prod_{i \in [q]} \Delta_i^{l_i} = \nu.$$

Therefore, given any $\mu \in \Lambda_D(G)$, there exists some $\nu \in \Lambda_\Delta(G)$, such that $X_D(G,\mu) \subseteq X_\Delta(G,\nu)$. Now, we consider some $\nu \in \Lambda_\Delta(G)$. Let $\mathbf{k} \in X_\Delta(G,\nu)$. Then, we let $\mu = \prod_{i \in [q]} \lambda_i^{k_i}$. We see that $\mathbf{k} \in X_D(G,\mu)$. This implies that given any $\nu \in \Lambda_\Delta(G)$, there exist some $\mu_1, \ldots, \mu_t \in \Lambda_D(G)$, such that

$$X_{\Delta}(G,\nu) = X_D(G,\mu_1) \sqcup \cdots \sqcup X_D(G,\mu_t).$$

Now, we know that

$$c_{H,\Delta}(G,\nu) = \sum_{\mathbf{k}\in X_{\Delta}(\nu)} c_{H}(G,\mathbf{k}) = \sum_{i\in[t]} \left(\sum_{\mathbf{l}\in X_{D}(\mu_{i})} c_{H}(G,\mathbf{l})\right) = \sum_{i\in[t]} c_{H,D}(G,\mu_{i}).$$

Note that we have already computed $c_{H,D}(G,\mu)$ for all $\mu \in \Lambda_D(G)$. Therefore, we can compute $c_{H,\Delta}(G,\nu)$ for each $\nu \in \Lambda_{\Delta}(G)$, and therefore, $Z_{H\Delta H^{\mathsf{T}}}(G)$ can also be computed. Therefore, $\mathsf{Pl-GH}(H\Delta H^{\mathsf{T}}) \leq \mathsf{Pl-GH}(M)$.

Corollary 14. If $M = HDH^{\mathsf{T}} \in Sym_q^{\mathsf{F}}(\mathbb{R})$ is such that its eigenvalues $(\lambda_1, \ldots, \lambda_q)$ have lattice dimension zero, then $\mathsf{Pl-GH}(H\Delta H^{\mathsf{T}}) \leq \mathsf{Pl-GH}(M)$ for any diagonal matrix $\Delta = diag(\Delta_1, \ldots, \Delta_q)$ such that $\Delta_i \in \mathbb{R}_{\neq 0}$ for all $i \in [q]$.

Proof. If the lattice dimension of $(\lambda_1, \ldots, \lambda_q)$ is zero, it implies that $\mathcal{L}(\lambda_1, \ldots, \lambda_q) = \{\mathbf{0}\}$. Therefore, $\mathcal{L}(\lambda_1, \ldots, \lambda_q) \subseteq \mathcal{L}(\Delta_1, \ldots, \Delta_q)$ for any $\Delta = (\Delta_1, \ldots, \Delta_q)$ such that $\Delta_i \in \mathbb{R}_{\neq 0}$. The result then follows from Lemma 13.

We will now prove that there exists some Δ such that $Pl-GH(H\Delta H^T)$ is #P-hard.

Lemma 15. Let $\mathcal{A} = \{a_1, \ldots, a_n\}$, and $\mathcal{B} = \{b_1, \ldots, b_m\}$ be finite sets of positive real numbers. There exists some $\kappa \in \mathbb{R}$ such that $\kappa + a_i > 1$ for all $i \in [n]$, κ is transcendental to the field $\mathbf{F} = \mathbb{Q}(\mathcal{B})$, and for all $(e_1, \ldots, e_n) \in \mathbb{Z}^n$,

$$(\kappa + a_1)^{e_1} \cdots (\kappa + a_n)^{e_n} = 1 \implies (e_1, \dots, e_n) = \mathbf{0}$$

Proof. For each $\mathbf{e} = (e_1, \ldots, e_n) \neq \mathbf{0} \in \mathbb{Z}^n$, we define the polynomial $f_{\mathbf{e}} : \mathbb{R} \to \mathbb{R}$ as

$$f_{\mathbf{e}}(x) = \prod_{i \in [n]: e_i > 0} (x + a_i)^{e_i} - \prod_{i \in [n]: e_i < 0} (x + a_i)^{-e_i}.$$

We can see that no such $f_{\mathbf{e}}$ is the zero polynomial as each $a_i \in \mathcal{A}$ is a distinct element in a unique factorization domain. Therefore, for each $f_{\mathbf{e}}$, the set $\emptyset_{\mathbf{e}} = \{x \in \mathbb{R} : f_{\mathbf{e}}(x) = 0\}$ is finite. We also see that the set $Z \subset \mathbb{R}$ of all the algebraic numbers over the field $\mathbf{F} = \mathbb{Q}(\mathcal{B})$ is countable, since \mathcal{B} is finite. Therefore, $(\bigcup_{\mathbf{e}} \emptyset_{\mathbf{e}}) \cup Z$ is a countable set. Therefore, we can pick some $\kappa \in \mathbb{R} \setminus ((\bigcup_{\mathbf{e}} \emptyset_{\mathbf{e}}) \cup Z)$ such that $\kappa + a_i > 1$ for all $i \in [n]$. This κ satisfies all the requirements of the lemma.

Lemma 16. Let $M = HDH^{\mathsf{T}} \in Sym_q^{\mathsf{F}}(\mathbb{R}_{\neq 0})$ (for $q \geq 3$). Then there exists a diagonal matrix $\Delta = D + \kappa I$ for some $\kappa \in \mathbb{R}$, such that $\mathsf{Pl-GH}(H\Delta H^{\mathsf{T}})$ is #P-hard.

Proof. Let $(M_{ij})_{i,j\in[q]}$ be the entries of the matrix M generated by some $\{g_t\}_{t\in[d]}$. We will replace M with the matrix N guaranteed by Lemma 4. So, we may assume that $e_{ijt} \ge 0$ for all $i, j \in [q]$, and $t \in [d]$. Let $\mathcal{A} = \{M_{ii} : i \in [q]\}$ be the set of diagonal elements (with duplicates removed). Without loss of generality, we let $\mathcal{A} = \{M_{11}, \ldots, M_{rr}\}$ for some $1 \le r \le q$. Let $\mathcal{B} = \{g_t\}_{t\in[d]}$. With this choice of \mathcal{A} and \mathcal{B} , we can let $\kappa \in \mathbb{R}$ be the number whose existence is guaranteed by Lemma 15. We will now let $\Delta = D + \kappa I$, a diagonal matrix with all diagonal elements positive.

Clearly, $H\Delta H^{\mathsf{T}} = M + \kappa I \in \operatorname{Sym}_{q}^{\mathsf{F}}(\mathbb{R}_{\neq 0})$. All its non-diagonal entries are the same as that of M, and can be represented as a product of non-negative integer powers of the generating set $\{g_t\}_{t\in[d]}$, up to a \pm sign. The diagonal entries of $H\Delta H^{\mathsf{T}}$ can also be trivially represented as non-negative integer powers of the set $\{\kappa + M_{11}, \ldots, \kappa + M_{rr}\}$. We will now show that each entry of $H\Delta H^{\mathsf{T}}$ can in fact be expressed *uniquely* as a product of integer powers of $\{g_t\}_{t\in[d]} \cup \{\kappa + M_{tt} : t \in [r]\}$, up to a \pm sign. To show that, we only need to prove that for any $(e_1, \ldots, e_d, e'_1, \ldots, e'_r) \in \mathbb{Z}^{d+r}$, if

$$g_1^{e_1} \cdots g_d^{e_d} \cdot (\kappa + M_{11})^{e_1'} \cdots (\kappa + M_{rr})^{e_r'} = 1,$$
(9)

then $(e_1, \ldots, e_d, e'_1, \ldots, e'_r) = \mathbf{0}$.

First, we assume $(e_1, \ldots, e_d) \neq \mathbf{0}$. In Eq. (9), since $\{g_t\}_{t \in [d]}$ is a generating set for M, and $(e_1, \ldots, e_d) \neq \mathbf{0}$, we have $\prod_{t \in [d]} g_t^{e_t} \neq 1$. Therefore, $(e'_1, \ldots, e'_r) \neq \mathbf{0}$ by Eq. (9). Separating out positive and negative e'_t 's, we have

$$\left(\prod_{t \in [d]} g_t^{e_t}\right) \cdot \prod_{t \in [r]: e_t' > 0} (\kappa + M_{tt})^{e_t'} = \prod_{t \in [r]: e_t' < 0} (\kappa + M_{tt})^{-e_t'}.$$
(10)

Since $(e'_1, \ldots, e'_r) \neq \mathbf{0}$, at least one side is a non-constant polynomial in κ over the field $\mathbf{F} = \mathbb{Q}(\{g_t\}_{t \in [d]})$, yet both sides have different leading coefficients. This contradicts our assumption that κ is transcendental to \mathbf{F} . Therefore, we must have $(e_1, \ldots, e_d) = \mathbf{0}$. But then, $\prod_{t \in [d]} g_t^{e_t} = 1$, which implies that

$$\prod_{t \in [r]} (\kappa + M_{tt})^{e'_t} = 1.$$

Lemma 15 implies that for κ , this is only possible if $(e'_1, \ldots, e'_r) = \mathbf{0}$, so $(e_1, \ldots, e_d, e'_1, \ldots, e'_r) = \mathbf{0}$.

This proves that $\{g_t\}_{t\in[d]} \cup \{\kappa + M_{tt} : t \in [r]\}$ is a generating set of the entries of $H\Delta H^{\mathsf{T}}$. Importantly, if we let e_{ijt} and e'_{ijt} be the unique integers such that

$$(H\Delta H^{\mathsf{T}})_{ij} = (-1)^{e_{ij0}} g_1^{e_{ij1}} \cdots g_d^{e_{ijd}} \cdot (\kappa + M_{11})^{e'_{ij1}} \cdots (\kappa + M_{r'})^{e'_{ijr}},$$

we see that $e_{ijt}, e'_{ijt'} \ge 0$ for all $i, j \in [q], t \in [d]$, and $t' \in [r]$. Moreover, for every $i \in [q]$, we know that there exists some $t(i) \in [r]$ such that $e'_{iit(i)} = 1$, and $e'_{jkt(i)} = 0$ for all $j \ne k$. So, from Lemma 9, we conclude that $Pl-GH(H\Delta H^T)$ is #P-hard.

Corollary 14 and Lemma 16 immediately allow us to prove the following theorem.

Theorem 17. If $M \in Sym_q^{\mathsf{F}}(\mathbb{R}_{\neq 0})$ (for $q \geq 3$) has eigenvalues with lattice dimension 0, then $\mathsf{Pl-GH}(M)$ is #P-hard.

With a little more effort however, Lemma 16 allows us to prove a slightly stronger version of Theorem 17. Given $i \neq j \in [q]$, we define $\delta_{ij} \in \mathbb{Z}^q$ such that

$$\delta_{ij}(k) = \begin{cases} 1 & \text{if } k = i, \\ -1 & \text{if } k = j, \\ 0 & \text{otherwise} \end{cases}$$

We then let

$$\mathcal{D} = \{\delta_{ij} : i \neq j \in [q]\} \tag{11}$$

Theorem 18. If $M \in Sym_q^{\mathsf{F}}(\mathbb{R}_{\neq 0})$ (for $q \geq 3$) has eigenvalues $(\lambda_1, \ldots, \lambda_q)$ with a lattice basis \mathcal{B} such that $\mathcal{B} \subseteq \mathcal{D}$, then $\mathsf{Pl-GH}(M)$ is #P-hard.

Proof. From Lemma 16, we know that given $M = HDH^{\mathsf{T}}$, there exists a diagonal matrix $\Delta = D + \kappa I$ for some $\kappa \in \mathbb{R}$ such that Pl - $\mathsf{GH}(H\Delta H^{\mathsf{T}})$ is $\#\mathsf{P}$ -hard. Now, if $\delta_{ij} \in \mathcal{B}$ for some i < j, we know that $\lambda_i^{+1} \cdot \lambda_j^{-1} = 1$. This implies that $\lambda_i = \lambda_j$. Therefore, $\lambda_i + \kappa = \lambda_j + \kappa$. So, $\Delta_i^{+1} \cdot \Delta_j^{-1} = 1$. Therefore, $\delta_{ij} \in \mathcal{L}(\Delta_1, \ldots, \Delta_q)$. Since this is true for all $\delta_{ij} \in \mathcal{B}$, it follows that $\mathcal{B} \subseteq \mathcal{L}(\Delta_1, \ldots, \Delta_q)$. Therefore, $\mathcal{L}(\lambda_1, \ldots, \lambda_q) \subseteq \mathcal{L}(\Delta_1, \ldots, \Delta_q)$. Now, Lemma 13 implies that Pl - $\mathsf{GH}(H\Delta H^{\mathsf{T}}) \leq \mathsf{Pl}$ - $\mathsf{GH}(M)$. By our choice of Δ , this implies that Pl - $\mathsf{GH}(M)$ is #P-hard.

Remark. It should be noted that the identity matrix $I \in Sym_q^{\mathsf{F}}(\mathbb{R})$ has eigenvalues $(1, \ldots, 1)$, which trivially has a lattice basis $\mathcal{B} = \{\delta_{1j} : 2 \leq j \leq q\} \subseteq \mathcal{D}$. However, we are not claiming that Pl -GH(I) is #P-hard (in fact the problem Pl -GH(I) is clearly in P). This is because the identity matrix does not satisfy the crucial property that all entries of the matrix belong to $\mathbb{R}_{\neq 0}$. In general, Theorem 18 implies that if $M \in Sym_q^{\mathsf{F}}(\mathbb{R})$ has eigenvalues with a lattice basis $\mathcal{B} \subseteq \mathcal{D}$, then either Pl -GH(M) is #P-hard, or M is guaranteed to have zero entries.

5 Proof of Hardness

We will now try to understand the lattice $\mathcal{L}(\lambda_1, \ldots, \lambda_q)$ of the eigenvalues of matrices $M \in \text{Sym}_q^F(\mathbb{R}_{>0})$ which do not satisfy the conditions of Theorem 18. Let

$$\chi_q = \left\{ \mathbf{x} = (x_1, \dots, x_q) \in \mathbb{Z}^q \middle| x_1 + \dots + x_q = 0 \right\}$$

Consider some $\mathbf{0} \neq \mathbf{x} \in \chi_q$. We will use this \mathbf{x} to define a polynomial $\phi_{\mathbf{x}} : \mathbb{R}^q \to \mathbb{R}$ as

$$\phi_{\mathbf{x}}(\alpha_1,\ldots,\alpha_q) = \prod_{i \in [q]: x_i > 0} \alpha_i^{x_i} - \prod_{i \in [q]: x_i < 0} \alpha_i^{-(x_i)}.$$
(12)

Since $\mathbf{x} \neq \mathbf{0}$, we can see that $\phi_{\mathbf{x}}$ is not the trivial constant zero function. Moreover, since $\sum_{i \in [q]} x_i = 0$, we see that

$$\sum_{i \in [q]: x_i > 0} x_i = \sum_{i \in [q]: x_i < 0} -(x_i).$$

So, we see that $\phi_{\mathbf{x}}$ is a homogeneous polynomial. Moreover, by construction, we see that

$$\mathbf{x} \in \mathcal{L}(\alpha_1, \dots, \alpha_q) \iff \phi_{\mathbf{x}}(\alpha_1, \dots, \alpha_q) = 0.$$

We will now define the polynomial $\Phi_{\mathbf{x}} : \mathbb{R}^q \to \mathbb{R}$ as follows:

$$\Phi_{\mathbf{x}}(\alpha_1,\ldots,\alpha_q) = \prod_{\sigma \in S_q} \phi_{\mathbf{x}}(\alpha_{\sigma(1)},\ldots,\alpha_{\sigma(q)}),$$
(13)

where S_q is the symmetric group over [q]. By construction, we see that $\Phi_{\mathbf{x}}$ is a symmetric homogeneous polynomial. In the next section, we will explore this polynomial in greater detail, and exploit its symmetry to prove some useful results. For now, the most important property of this polynomial is that if $\mathbf{x} \in \mathcal{L}(\alpha)$ for some $\alpha = (\alpha_1, \ldots, \alpha_q)$, then $\Phi_{\mathbf{x}}(\alpha) = 0$. However, because we constructed $\Phi_{\mathbf{x}}$ to be symmetric, the converse does not quite hold.

Lemma 19. Let $\mathbf{x}, \mathbf{y} \in \chi_q$. Let $\sigma \in S_q$ such that $x_i = y_{\sigma(i)}$ for all $i \in [q]$. Then, $\Phi_{\mathbf{x}} = \Phi_{\mathbf{y}}$.

Proof. By definition, given any $\alpha \in \mathbb{R}^q$, we know that

$$\phi_{\mathbf{x}}(\alpha_{\sigma(1)}, \dots, \alpha_{\sigma(q)}) = \prod_{i \in [q]: \ x_i > 0} \alpha_{\sigma(i)}^{x_i} - \prod_{i \in [q]: \ x_i < 0} \alpha_{\sigma(i)}^{-(x_i)}$$
$$= \prod_{i \in [q]: \ y_{\sigma(i)} > 0} \alpha_{\sigma(i)}^{y_{\sigma(i)}} - \prod_{i \in [q]: \ y_{\sigma(i)} < 0} \alpha_{\sigma(i)}^{-(y_{\sigma(i)})}$$
$$= \prod_{j \in [q]: \ y_j > 0} \alpha_j^{y_j} - \prod_{j \in [q]: \ y_j < 0} \alpha_j^{-(y_j)}$$
$$= \phi_{\mathbf{y}}(\alpha_1, \dots, \alpha_q).$$

Therefore,

$$\Phi_{\mathbf{y}}(\alpha_1, \dots, \alpha_q) = \prod_{\sigma' \in S_q} \phi_{\mathbf{y}}(\alpha_{\sigma'(1)}, \dots, \alpha_{\sigma'(q)})$$
$$= \prod_{\sigma' \in S_q} \phi_{\mathbf{x}}(\alpha_{\sigma(\sigma'(1))}, \dots, \alpha_{\sigma(\sigma'(q))})$$
$$= \prod_{\tau \in S_q} \phi_{\mathbf{x}}(\alpha_{\tau(1)}, \dots, \alpha_{\tau(q)})$$
$$= \Phi_{\mathbf{x}}(\alpha_1, \dots, \alpha_q).$$

-		

Lemma 19 implies that if $\mathbf{x} \in \mathcal{L}(\lambda_1, \ldots, \lambda_q)$, then $\Phi_{\mathbf{y}}(\lambda_1, \ldots, \lambda_q) = 0$ for all $\mathbf{y} \in \chi_q$ such that $y_{\sigma(i)} = x_i$ for some $\sigma \in S_q$. To navigate through some of the problems introduced by this, we will now introduce some notation.

Notation 20. Let $X \subset \chi_q$. Then, given any $\sigma \in S_q$, we define

$$X^{\sigma} = \{ \mathbf{x} \in \chi_q : (x_{\sigma(1)}, \dots, x_{\sigma(q)}) \in X \}.$$

We also define

$$\overline{X} = \bigcup_{\sigma \in S_q} X^{\sigma}$$

Lemma 21. Let $\alpha = (\alpha_1, \ldots, \alpha_q)$ be non-zero reals. Then,

$$\mathbf{x} \in \overline{\mathcal{L}(\alpha)} \iff \Phi_{\mathbf{x}}(\alpha) = 0.$$

Proof. If $\mathbf{x} \in \overline{\mathcal{L}(\alpha)}$, we know that there exists some $\sigma \in S_q$ such that $\mathbf{x} \in \mathcal{L}(\alpha)^{\sigma}$. If we now define $\mathbf{y} \in \chi_q$, such that $y_i = x_{\sigma(i)}$ for all $i \in [q]$, we see that $\mathbf{y} \in \mathcal{L}(\alpha)$. Now, Lemma 19 implies that $\Phi_{\mathbf{x}}(\alpha) = \Phi_{\mathbf{y}}(\alpha) = 0$.

If $\Phi_{\mathbf{x}}(\alpha) = 0$, then we see that $\phi_{\mathbf{x}}(\alpha_{\sigma(1)}, \ldots, \alpha_{\sigma(q)}) = 0$ for some $\sigma \in S_q$. Therefore, $\phi_{\mathbf{y}}(\alpha) = 0$, where $y_i = x_{\sigma^{-1}(i)}$ for all $i \in [q]$. So, $\mathbf{y} \in \mathcal{L}(\alpha)$, which implies that $\mathbf{x} \in \mathcal{L}(\alpha)^{\sigma^{-1}} \subset \overline{\mathcal{L}(\alpha)}$.

We recall from Eq. (13) that given any $\mathbf{0} \neq \mathbf{x} \in \chi_q$, the polynomial $\Phi_{\mathbf{x}}$ is a symmetric, homogeneous polynomial. We now recall the Fundamental Theorem of Symmetric Polynomials, stated below:

Theorem 22. Let $\Phi(\alpha_1, \ldots, \alpha_q)$ be a symmetric polynomial on q variables. Let e_i for $i \in [q]$ represent the elementary symmetric polynomials, such that

$$e_i(\alpha_1,\ldots,\alpha_q) = \sum_{1 \le t_1 < \cdots < t_i \le q} \alpha_{t_1} \cdots \alpha_{t_i}.$$

Then, there exists a polynomial function $\widehat{\Phi} : \mathbb{R}^q \to \mathbb{R}$ such that

$$\Phi(\alpha_1,\ldots,\alpha_q) = \widehat{\Phi}\Big(e_1(\alpha_1,\ldots,\alpha_q),\ldots,e_q(\alpha_1,\ldots,\alpha_q)\Big).$$

Now, consider some $\mathbf{0} \neq \mathbf{x} \in \chi_q$, and $M \in \operatorname{Sym}_q(\mathbb{R})$ with eigenvalues $(\lambda_1, \ldots, \lambda_q)$. Theorem 22 implies that there exists a polynomial $\widehat{\Phi}_{\mathbf{x}}$ such that

$$\Phi_{\mathbf{x}}(\lambda_1,\ldots,\lambda_q) = \widehat{\Phi_{\mathbf{x}}}\Big(e_1(\lambda_1,\ldots,\lambda_q),\ldots,e_q(\lambda_1,\ldots,\lambda_q)\Big).$$

We will now use the fact that these $e_i(\lambda_1, \ldots, \lambda_q)$ are all just coefficients of the characteristic polynomial of M. Thus, each e_i can be written as a homogeneous polynomial of the entries of M.

Theorem 23. Let $M \in Sym_q(\mathbb{R})$ with eigenvalues $(\lambda_1, \ldots, \lambda_q)$. For any elementary symmetric polynomial e_k for $k \in [q]$, there exists a homogeneous polynomial $s_k : Sym_q(\mathbb{R}) \to \mathbb{R}$ of degree k such that

$$e_k(\lambda_1,\ldots,\lambda_q) = s_k\left((M_{ij})_{i\leq j\in[q]}\right).$$

We will finally define the polynomial $\Psi_{\mathbf{x}} : \operatorname{Sym}_q(\mathbb{R}) \to \mathbb{R}$ as

$$\Psi_{\mathbf{x}}(M) = \widehat{\Phi_{\mathbf{x}}}\left(s_1\left((M_{ij})_{i \le j \in [q]}\right), \dots, s_q\left((M_{ij})_{i \le j \in [q]}\right)\right).$$
(14)

Definition 24. A function $F : Sym_q(\mathbb{R}) \to \mathbb{R}$ is called a $Sym_q(\mathbb{R})$ -polynomial if it is a homogeneous polynomial in the entries of the matrix.

Remark. Note that we require $Sym_q(\mathbb{R})$ -polynomials to be homogeneous. If F is a $Sym_q(\mathbb{R})$ polynomial of degree d, then $F(cM) = c^d \cdot F(M)$. In particular, this would allow us to conclude that F(M) = 0 if and only if F(cM) = 0. So, we can safely replace any M with N = cM as guaranteed
by Lemma 4 without changing whether F(M) = 0 for any $Sym_a(\mathbb{R})$ polynomial F.

Clearly, given any $\mathbf{0} \neq \mathbf{x} \in \chi_q$, we see that $\Psi_{\mathbf{x}}(M)$ is polynomial in the entries of M. Moreover, we note that for each $i \in [q]$, e_i and s_i are homogeneous polynomials of the same degree. Since $\Phi_{\mathbf{x}}$ is a homogeneous polynomial, by construction of $\widehat{\Phi}$, it follows that $\Psi_{\mathbf{x}}$ is also a homogeneous polynomial, of the same degree as $\Phi_{\mathbf{x}}$. In summation, we have managed to prove the following lemma.

Lemma 25. Let $\mathbf{0} \neq \mathbf{x} \in \chi_q$. There exists a $Sym_q(\mathbb{R})$ -polynomial $\Psi_{\mathbf{x}}$, such that given any $M \in Sym_q^{\mathsf{F}}(\mathbb{R})$ with eigenvalues $(\lambda_1, \ldots, \lambda_q)$,

$$\Psi_{\mathbf{x}}(M) = 0 \iff \mathbf{x} \in \overline{\mathcal{L}(\lambda_1, \dots, \lambda_q)}.$$

The proof immediately follows from the construction of $\Psi_{\mathbf{x}}$, and Lemma 21. Before we can proceed further, we will need to establish a few important properties of $\overline{\mathcal{L}(\lambda_1,\ldots,\lambda_q)}$.

Lemma 26. Let $\mathcal{L}_1 = \mathcal{L}(\alpha_1, \ldots, \alpha_q)$ and $\mathcal{L}_2 = \mathcal{L}(\beta_1, \ldots, \beta_q)$ be lattices such that $\beta_i > 0$ for all $i \in [q]$. Let $d_1 = \dim(\mathcal{L}_1)$ and $d_2 = \dim(\mathcal{L}_2)$. If $\overline{\mathcal{L}_2} \subsetneq \overline{\mathcal{L}_1}$, then $d_2 < d_1$.

Proof. We will first define the rational span of a lattice \mathcal{L} of dimension d with a lattice basis $\mathcal{B} = \{\mathbf{y}_1, \dots, \mathbf{y}_d\}$ to be

$$\operatorname{Qsp}(\mathcal{L}) = \{c_1 \cdot \mathbf{y}_1 + \dots + c_d \cdot \mathbf{y}_d \mid c_1, \dots, c_d \in \mathbb{Q}\}$$

It is easily seen that $\operatorname{Qsp}(\mathcal{L})$ is a \mathbb{Q} -vector space of the same dimension d, as the lattice dimension of \mathcal{L} . We also note that given any $\sigma \in S_q$, \mathcal{L}^{σ} is also a lattice of dimension d, with \mathcal{B}^{σ} as a lattice basis. Therefore,

$$\overline{\operatorname{Qsp}(\mathcal{L})} = \bigcup_{\sigma \in S_q} \operatorname{Qsp}(\mathcal{L}^{\sigma})$$

for any lattice \mathcal{L} .

Now, let us consider any $\mathbf{y} \in \overline{\operatorname{Qsp}(\mathcal{L}_2)}$. There exists some $\sigma \in S_q$, such that $\mathbf{y} \in \operatorname{Qsp}(\mathcal{L}_2^{\sigma})$. So, there exists some $c \in \mathbb{Z}_{>0}$, such that $c \cdot \mathbf{y} \in \mathcal{L}_2^{\sigma} \subset \overline{\mathcal{L}_2}$. From our assumptions, we know that $\overline{\mathcal{L}_2} \subset \overline{\mathcal{L}_1}$. Therefore, there exists some $\tau \in S_q$ such that $c \cdot \mathbf{y} \in \mathcal{L}_1^{\tau}$. So, $\mathbf{y} \in \operatorname{Qsp}(\mathcal{L}_1^{\tau}) \subset \overline{\operatorname{Qsp}(\mathcal{L}_1)}$. Therefore, we have shown that

$$\overline{\operatorname{Qsp}(\mathcal{L}_2)} \subseteq \overline{\operatorname{Qsp}(\mathcal{L}_1)}.$$

Hence,

$$\operatorname{Qsp}(\mathcal{L}_2) \subseteq \overline{\operatorname{Qsp}(\mathcal{L}_2)} \subseteq \overline{\operatorname{Qsp}(\mathcal{L}_1)} = \bigcup_{\sigma \in S_q} \operatorname{Qsp}(\mathcal{L}_1^{\sigma})$$

Since this is a finite union of Q-vector spaces, we know that there must be some $\tau \in S_q$, such that $\operatorname{Qsp}(\mathcal{L}_2) \subseteq \operatorname{Qsp}(\mathcal{L}_1^{\tau})$. In particular, this implies that $d_2 = \dim(\operatorname{Qsp}(\mathcal{L}_2)) \leq \dim(\operatorname{Qsp}(\mathcal{L}_1^{\tau})) = d_1$. Now, for a contradiction assume $d_2 = d_1$. This implies that in fact, $\operatorname{Qsp}(\mathcal{L}_2) = \operatorname{Qsp}(\mathcal{L}_1^{\tau})$.

Now, we recall that there exists some $\mathbf{x} \in \overline{\mathcal{L}_1} \setminus \overline{\mathcal{L}_2}$. Being outside \mathcal{L}_2 , clearly $\mathbf{x} \neq \mathbf{0}$. There exists some $\sigma \in S_q$, such that $\mathbf{x} \in \mathcal{L}_1^{\sigma}$. If we let $\mathbf{y} \in \chi_q$ such that $y_i = x_{\tau^{-1}(\sigma(i))}$ for all $i \in [q]$, we see that $\mathbf{y} \in \mathcal{L}_1^{\tau} \subset \operatorname{Qsp}(\mathcal{L}_1^{\tau}) = \operatorname{Qsp}(\mathcal{L}_2)$. Therefore, there exists some $c \in \mathbb{Z}_{>0}$, such that $c \cdot \mathbf{y} \in \mathcal{L}_2$. We recall that $\mathcal{L}_2 = \mathcal{L}(\beta_1, \ldots, \beta_q)$. Therefore, by definition, this implies that

$$\beta_1^{c \cdot y_1} \cdots \beta_q^{c \cdot y_q} = \left(\beta_1^{y_1} \cdots \beta_q^{y_q}\right)^c = 1 \implies \beta_1^{y_1} \cdots \beta_q^{y_q} = \pm 1$$

Since $\beta_i > 0$ for all $i \in [q]$, it must be the case that $\beta_1^{y_1} \cdots \beta_q^{y_q} = 1$, which means that $\mathbf{y} \in \mathcal{L}_2$. By the construction of \mathbf{y} , it follows that $\mathbf{x} \in \mathcal{L}_2^{\tau^{-1}\sigma} \subset \overline{\mathcal{L}}_2$. But this contradicts our assumption about \mathbf{x} that $\mathbf{x} \notin \overline{\mathcal{L}}_2$. Therefore, our assumption that $d_2 = d_1$ must be false.

We will now see how Lemma 26 allows us to prove the #P-hardness of Pl-GH(M). We will need just one more definition.

Definition 27. Let $M \in Sym_q^{\mathbf{F}}(\mathbb{R}_{\neq 0})$ have eigenvalues $(\lambda_1, \ldots, \lambda_q)$. Let \mathcal{F} be a countable set of $Sym_q(\mathbb{R})$ -polynomials. A matrix $N \in Sym_q^{\mathbf{F}}(\mathbb{R}_{\neq 0})$ is called a reduct of (M, \mathcal{F}) if $\mathsf{Pl-GH}(N) \leq \mathsf{Pl-GH}(M)$, and $F(N) \neq 0$ for all $F \in \mathcal{F}$. We denote the set of reducts of (M, \mathcal{F}) by

$$\mathfrak{R}(M,\mathcal{F}) = \Big\{ N \in Sym_q^{\mathsf{F}}(\mathbb{R}_{\neq 0}) \Big| \ N \ is \ a \ reduct \ of \ (M,\mathcal{F}) \Big\}.$$

Consider any $M \in \operatorname{Sym}_q^{pd}(\mathbb{R}_{\neq 0})$, with eigenvalues $(\lambda_1, \ldots, \lambda_q)$. We define

$$\mathcal{F}_M = \Big\{ \Psi_{\mathbf{y}} : \mathbf{0} \neq \mathbf{y} \in \chi_q, \Psi_{\mathbf{y}}(M) \neq 0 \Big\}.$$
(15)

Since χ_q is a countable set, it follows that \mathcal{F}_M is also a countable set of $\operatorname{Sym}_q(\mathbb{R})$ -polynomials. From Lemma 25, we see that \mathcal{F}_M consists of precisely the polynomials indexed by vectors in $\chi_q \setminus \overline{\mathcal{L}(\lambda_1, \ldots, \lambda_q)}$.

Now, if the eigenvalues $(\lambda_1, \ldots, \lambda_q)$ of M have a lattice basis \mathcal{B} such that $\mathcal{B} \subset \mathcal{D}$, we know from Theorem 18 that Pl-GH(M) is #P-hard. Otherwise, for any lattice basis \mathcal{B} of the eigenvalues, there exists some $\mathbf{x} \in \mathcal{B} \setminus \mathcal{D}$. Our goal will be to prove that there exists some

$$N_1 \in \mathfrak{R}(M, \mathcal{F}_M \cup \{\Psi_{\mathbf{x}}\}) \cap \operatorname{Sym}_q^{\mathsf{pd}}(\mathbb{R}_{\neq 0}).$$

If such an N_1 exists, let its eigenvalues be (μ_1, \ldots, μ_q) . From Definition 27, for all $\mathbf{0} \neq \mathbf{y} \in \chi_q$, if $\Psi_{\mathbf{y}}(M) \neq 0$, then we also have $\Psi_{\mathbf{y}}(N_1) \neq 0$. In other words, for all $\mathbf{0} \neq \mathbf{y} \in \chi_q$, $\Psi_{\mathbf{y}}(N_1) = 0$ implies that $\Psi_{\mathbf{y}}(M) = 0$. So, Lemma 25 now implies that

$$\overline{\mathcal{L}(\mu_1,\ldots,\mu_q)}\subseteq\overline{\mathcal{L}(\lambda_1,\ldots,\lambda_q)}.$$

Moreover, by our choice of N_1 , we know that $\Psi_{\mathbf{x}}(N_1) \neq 0$, i.e., $\mathbf{x} \notin \overline{\mathcal{L}(\mu_1, \ldots, \mu_q)}$, again by Lemma 25. But by our choice of \mathbf{x} , we know that $\mathbf{x} \in \mathcal{B} \subset \mathcal{L}(\lambda_1, \ldots, \lambda_q) \subseteq \overline{\mathcal{L}(\lambda_1, \ldots, \lambda_q)}$. Therefore, we see that in fact,

$$\overline{\mathcal{L}(\mu_1,\ldots,\mu_q)} \subsetneq \overline{\mathcal{L}(\lambda_1,\ldots,\lambda_q)}.$$

Therefore, Lemma 26 tells us that

$$\dim(\mathcal{L}(\mu_1,\ldots,\mu_q)) < \dim(\mathcal{L}(\lambda_1,\ldots,\lambda_q)).$$

Now we can repeat this process with N_1 in place of M. Since the lattice dimension of $\mathcal{L}(\lambda_1, \ldots, \lambda_q)$ is some finite $d \in \mathbb{Z}_{\geq 0}$, we can only repeat this process k times, for some $k \leq d$, until we find some $N_k \in \operatorname{Sym}_q^{pd}(\mathbb{R}_{\neq 0})$ such that it has a basis $\mathcal{B} \subset \mathcal{D}$, and thus $\operatorname{Pl-GH}(N_k)$ is #P-hard by Theorem 18. Furthermore,

$$\operatorname{Pl-GH}(N_k) \leq \cdots \leq \operatorname{Pl-GH}(N_1) \leq \operatorname{Pl-GH}(M).$$

This would then allow us to prove that Pl-GH(M) is #P-hard.

In other words, given a matrix $M \in \operatorname{Sym}_{q}^{pd}(\mathbb{R}_{\neq 0})$, and any non-zero $\mathbf{x} \in \chi_q \setminus \mathcal{D}$, our goal will be to prove that there exists some $N \in \mathfrak{R}(M, \mathcal{F}_M \cup \{\Psi_{\mathbf{x}}\}) \cap \operatorname{Sym}_{q}^{pd}(\mathbb{R}_{\neq 0})$. In subsequent sections, we will find larger and larger classes of matrices M and \mathbf{x} for which we can find such an N.

6 Lattice on Diagonal Entries

Let $M \in \operatorname{Sym}_q^{\mathsf{F}}(\mathbb{R}_{\neq 0})$, and let \mathcal{F} be a countable set of $\operatorname{Sym}_q(\mathbb{R})$ -polynomials. Since our goal is to show that for some choices of M and $\mathcal{F}, \mathfrak{R}(M, \mathcal{F}) \neq \emptyset$, we will first prove some sufficient conditions for this set to be non-empty. We will recall from Lemma 3 that given any $M \in \operatorname{Sym}_q(\mathbb{R}_{\neq 0})$, there exists a generating set $\{g_1, \ldots, g_d\}$ for the entries of M. We also recall that Lemma 4 allows us to replace this matrix M with a matrix N = cM satisfying $\operatorname{Pl-GH}(M) \equiv \operatorname{Pl-GH}(N)$ and such that the entries of N are generated by $\{g_t\}_{t\in[d]}$ with exponents $e_{ijt} \geq 0$ for all $i, j \in [q]$, and $t \in [d]$. Therefore, Lemma 6 tells us that $\operatorname{Pl-GH}(\mathcal{T}_M(\mathbf{p})) \leq \operatorname{Pl-GH}(M)$ for all $\mathbf{p} \in \mathbb{R}^d$, where \mathcal{T}_M is as defined in Definition 5, and satisfies $\mathcal{T}_M(g_1, \ldots, g_d) = M$.

Lemma 28. Let $M \in Sym_q^{\mathsf{F}}(\mathbb{R}_{\neq 0})$ be a matrix, whose entries are generated by $\{g_t\}_{t\in[d]}$. Let \mathcal{F} be a countable set of $Sym_q(\mathbb{R})$ -polynomials, such that for each $F \in \mathcal{F}$, there exists some $\mathbf{p}_F \in \mathbb{R}^d$, such that $F(\mathcal{T}_M(\mathbf{p}_F)) \neq 0$. Then, there exists $N = \mathcal{T}_M(\mathbf{p}^*) \in \mathfrak{R}(M, \mathcal{F}) \cap Sym_q^{\mathsf{F}}(\mathbb{R}_{\neq 0})$ for some $\mathbf{p}^* \in \mathbb{R}^d$. Moreover,

- if $M \in Sym_a^{\mathbf{F}}(\mathbb{R}_{>0})$, we can ensure that $N \in Sym_a^{\mathbf{F}}(\mathbb{R}_{>0})$,
- if $M \in Sym_q^{pd}(\mathbb{R}_{\neq 0})$, we can ensure that $N \in Sym_q^{pd}(\mathbb{R}_{\neq 0})$,
- if $M \in Sym_a^{pd}(\mathbb{R}_{>0})$, we can ensure that $N \in Sym_a^{pd}(\mathbb{R}_{>0})$.

Proof. We will first replace M with cM as guaranteed by Lemma 4, for some c > 0. Since $M \in \operatorname{Sym}_q^{\mathsf{F}}(\mathbb{R}_{\neq 0})$, it follows that $cM \in \operatorname{Sym}_q^{\mathsf{F}}(\mathbb{R}_{\neq 0})$ as well. If $M \in \operatorname{Sym}_q^{\mathsf{F}}(\mathbb{R}_{>0})$ (similarly, $\operatorname{Sym}_q^{\mathsf{pd}}(\mathbb{R}_{\neq 0})$ and $\operatorname{Sym}_q^{\mathsf{pd}}(\mathbb{R}_{>0})$), then so is cM. We note from the definition of \mathcal{T}_M in Definition 5, that the entries of the matrix $\mathcal{T}_M(\mathbf{p})$ are all polynomials in \mathbf{p} . Since each $F \in \mathcal{F}$ is a $\operatorname{Sym}_q(\mathbb{R})$ -polynomial, it follows that $F(\mathcal{T}_M(\mathbf{p}))$ is a polynomial in \mathbf{p} for all $F \in \mathcal{F}$. Moreover, by our choice of \mathcal{F} , we know that for each $F \in \mathcal{F}$, there exists some $\mathbf{p}_F \in \mathbb{R}^d$ such that $F(\mathcal{T}_M(\mathbf{p})) \neq 0$. We also note that $\det(\mathcal{T}_M(g_1,\ldots,g_d)) = \det(M) \neq 0$, since M is full rank. Therefore, $F(\mathcal{T}_M(\mathbf{p})) : \mathbb{R}^d \to \mathbb{R}$ is a non-zero polynomial for all $F \in \mathcal{F} \cup \{\det\}$.

Let us now assume that M is positive definite. So, the eigenvalues $(\lambda_1, \ldots, \lambda_q)$ of $\mathcal{T}_M(g_1, \ldots, g_d) = M$ will all be positive. We note that each entry of the matrix $\mathcal{T}_M(\mathbf{p})$ is a continuous function of \mathbf{p} . So, each of the coefficients of the characteristic polynomial of $\mathcal{T}_M(\mathbf{p})$ is a continuous function of \mathbf{p} . Since the eigenvalues of $\mathcal{T}_M(\mathbf{p})$ are simply the roots of the characteristic polynomial, we can use the well-known fact that the roots of a polynomial are continuous in the coefficients [HM87] to see that eigenvalues of $\mathcal{T}_M(\mathbf{p})$ are also continuous as functions of \mathbf{p} . In other words, there exist open intervals I_1, \ldots, I_d such that $g_t \in I_t$ for all $t \in [d]$, and $\mathcal{T}_M(\mathbf{p})$ is positive definite for all $\mathbf{p} \in (I_1 \times \cdots \times I_d) = U$. Since $g_t > 1$ for all $t \in [q]$, we may further assume that $U \subset (\mathbb{R}_{>1})^d$. By

construction, we note that U has positive measure. If on the other hand, M is not positive definite, we can simply let $U = (\mathbb{R}_{>1})^d$.

We note that for each $F \in \mathcal{F} \cup \{\det\}$, the set $\emptyset_F = \{\mathbf{p} : F(\mathcal{T}_M(\mathbf{p})) = 0\}$ has measure 0, since $F(\mathcal{T}_M(\mathbf{p}))$ is a non-zero polynomial. The measure of a countable union of measure 0 sets is also 0. Therefore, there exists some $\mathbf{p}^* = (p_1^*, \ldots, p_d^*) \in U \setminus \bigcup_F \emptyset_F$. If we let $N = \mathcal{T}_M(\mathbf{p}^*)$, we now see that $F(N) \neq 0$ for all $F \in \mathcal{F} \cup \{\det\}$. In particular, this implies that $\det(N) \neq 0$. Moreover, since $\mathbf{p}^* \in U \subseteq (\mathbb{R}_{>1})^d$, we see that $N \in \operatorname{Sym}_q^F(\mathbb{R}_{\neq 0})$. So, $N \in \mathfrak{R}(M, \mathcal{F}) \cap \operatorname{Sym}_q^F(\mathbb{R}_{\neq 0})$ is the required matrix.

Moreover, if $M \in \operatorname{Sym}_q(\mathbb{R}_{>0})$, we know from Eq. (5) that $e_{ij0} = 0$ for all $i, j \in [q]$. Since $\mathbf{p}^* \in U$, this implies that $N \in \operatorname{Sym}_q(\mathbb{R}_{>0})$ as well. Finally, if $M \in \operatorname{Sym}_q^{pd}(\mathbb{R}_{\neq 0})$, we note that by our choice of $U, N \in \operatorname{Sym}_q^{pd}(\mathbb{R}_{\neq 0})$ as well. Therefore, we see that this N satisfies all the requirements of the lemma.

Corollary 29. Let $M \in Sym_q^{pd}(\mathbb{R}_{>0})$. Let \mathcal{F} be a countable set of $Sym_q(\mathbb{R})$ -polynomials such that $F(M) \neq 0$ for all $F \in \mathcal{F}$. Let $\mathbf{0} \neq \mathbf{x} \in \chi_q$ such that $\Psi_{\mathbf{x}}(T_n M) \neq 0$ for some integer $n \geq 1$. Then, there exists some $N \in \mathfrak{R}(M, \mathcal{F} \cup \mathcal{F}_M \cup \{\Psi_{\mathbf{x}}\}) \cap Sym_q^{pd}(\mathbb{R}_{>0})$.

Proof. Let $\{g_t\}_{t\in[d]}$ be a generating set of the entries of M. As we have already seen, we may replace M with $c \cdot M$ as guaranteed by Lemma 4. We have seen in the remark following Definition 24 that F(M) = 0 if and only if F(cM) = 0. Therefore, $F(c \cdot M) \neq 0$ for all $F \in \mathcal{F}$. By our choice of \mathcal{F} , and the definition of \mathcal{F}_M in Eq. (15), we see that for all $F \in \mathcal{F} \cup \mathcal{F}_M$, $F(\mathcal{T}_M(g_1, \ldots, g_d)) = F(M) \neq 0$. Moreover, we also see that since $M \in \operatorname{Sym}_q(\mathbb{R}_{>0})$, $\Psi_{\mathbf{x}}(\mathcal{T}_M((g_1)^n, \ldots, (g_d)^n)) = \Psi_{\mathbf{x}}(\mathcal{T}_n M) \neq 0$ as given. Let $\mathcal{F}' = \mathcal{F} \cup \mathcal{F}_M \cup \{\Psi_{\mathbf{x}}\}$. It is a countable set of $\operatorname{Sym}_q(\mathbb{R})$ -polynomials with the property that for every $F \in \mathcal{F}'$ there exists some $\mathbf{p}_F \in \mathbb{R}^d$ such that $F(\mathcal{T}_M(\mathbf{p}_F)) \neq 0$. We are also given that $M \in \operatorname{Sym}_q^{\mathrm{pd}}(\mathbb{R}_{>0})$. Therefore, Lemma 28 implies that there exists some $N \in \mathfrak{R}(M, \mathcal{F}') \cap \operatorname{Sym}_q^{\mathrm{pd}}(\mathbb{R}_{>0})$.

Now we will consider $M \in \operatorname{Sym}_q^{\operatorname{pd}}(\mathbb{R}_{>0})$ and $\mathbf{0} \neq \mathbf{x} \in \chi_q$, such that $\Psi_{\mathbf{x}}(T_n M) = 0$ for all n. To better understand these matrices, it will be helpful for us to study the function $\Psi_{\mathbf{x}} : \operatorname{Sym}_q(\mathbb{R}) \to \mathbb{R}$ better. We already know that the function $\Psi_{\mathbf{x}}$ is a homogeneous polynomial of some degree $d \geq 1$.

Now, we will try to understand the individual terms of this polynomial. Recall that we have defined

$$\mathcal{P}_{q^2}(d) = \left\{ (k_{ij})_{i,j \le q} \in \mathbb{Z}^{q^2} \mid k_{ij} \ge 0, \sum_{i,j \in [q]} k_{ij} = d \right\}.$$

Given any $\mathbf{k} \in \mathcal{P}_{q^2}(d)$, we will now define $m_{\mathbf{k}} : \operatorname{Sym}_q(\mathbb{R}) \to \mathbb{R}$, such that

$$m_{\mathbf{k}}(M) = \prod_{i,j \in [q]} M_{ij}^{k_{ij}}$$

As $\Psi_{\mathbf{x}}$ had degree d, each $m_{\mathbf{k}}$ represents a monomial term in $\Psi_{\mathbf{x}}$ (with possibly a 0 coefficient). We then let

$$\mathcal{M} = \left\{ m_{\mathbf{k}} \, \middle| \, \mathbf{k} \in \mathcal{P}_{q^2}(d) \right\},\,$$

and we can now express $\Psi_{\mathbf{x}}$ as

$$\Psi_{\mathbf{x}}(M) = \sum_{m \in \mathcal{M}} c_m(\mathbf{x}) m(M),$$

where $c_m(\mathbf{x}) \in \mathbb{R}$ for all $m \in \mathcal{M}$.

Lemma 30. Let $\mathbf{0} \neq \mathbf{x} \in \chi_q$. There exists $\epsilon > 0$, such that given any matrix $M \in Sym_q(\mathbb{R})$ satisfying the conditions that $|M_{ij} - I_{ij}| < \epsilon$ for all $i, j \in [q]$, and $\Psi_{\mathbf{x}}(T_n M) = 0$ for all $n \ge 1$, then

$$\Psi_{\mathbf{x}}(diag(M)) = 0,$$

where $diag(M) \in Sym_{q}(\mathbb{R})$ is the diagonal matrix such that

$$diag(M)_{ij} = \begin{cases} M_{ii} & \text{if } i = j, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. We will let $\epsilon = 1/3d$, and assume that $|M_{ij} - I_{ij}| < \epsilon$ for all $i, j \in [q]$. By construction of \mathcal{M} , we see that

$$\Psi_{\mathbf{x}}(T_n M) = \sum_{m \in \mathcal{M}} c_m(\mathbf{x}) \left(m(M) \right)^n.$$

Note that each $m \in \mathcal{M}$ is $m_{\mathbf{k}}$ for some $\mathbf{k} \in \mathcal{P}_{q^2}(d)$, where d is the degree of $\Psi_{\mathbf{x}}$. Now, we let

$$\mathcal{M}(M) = \left\{ m(M) \mid m \in \mathcal{M} \right\}$$

be the set of all values of the monomial terms in \mathcal{M} when evaluated at M. For all the values $v \in \mathcal{M}(M)$, we let

$$c_{v,M}(\mathbf{x}) = \sum_{m \in \mathcal{M}: m(M) = v} c_m(\mathbf{x}).$$

So, we see that

$$\Psi_{\mathbf{x}}(T_n M) = \sum_{v \in \mathcal{M}(M)} c_{v,M}(\mathbf{x}) \cdot v^n.$$

We can see that $|\mathcal{M}(M)|$ is O(1). So, in particular, the equations $\Psi_{\mathbf{x}}(T_n M) = 0$ form a full rank Vandermonde system of linear equations. This implies that $c_{v,M}(\mathbf{x}) = 0$ for all $v \in \mathcal{M}(M)$.

Setting that aside for a moment, we also note that by construction of $\operatorname{diag}(M)$,

$$m_{\mathbf{k}}(\operatorname{diag}(M)) = \begin{cases} m_{\mathbf{k}}(M) & \text{if } k_{ij} = 0 \ \forall \ i \neq j, \\ 0 & \text{otherwise.} \end{cases}$$

Therefore, if we define

diag
$$(\mathcal{M}) = \left\{ m_{\mathbf{k}} \mid \mathbf{k} \in \mathcal{P}_{q^2}(d), \sum_{i \in [q]} k_{ii} = d \right\},\$$

we see that $\operatorname{diag}(\mathcal{M}) \subset \mathcal{M}$, and

$$\Psi_{\mathbf{x}}(\operatorname{diag}(M)) = \sum_{m \in \operatorname{diag}(\mathcal{M})} c_m(\mathbf{x}) m(M).$$

So, if we let

$$\operatorname{diag}(\mathcal{M})(M) = \left\{ m(M) \mid m \in \operatorname{diag}(\mathcal{M}) \right\}, \quad \text{and} \quad c_{v,\operatorname{diag}(M)}(\mathbf{x}) = \sum_{m \in \operatorname{diag}(\mathcal{M}): \ m(M) = v} c_m(\mathbf{x})$$

for all $v \in \operatorname{diag}(\mathcal{M})(M)$, then we see that

$$\Psi_{\mathbf{x}}(\operatorname{diag}(M)) = \sum_{v \in \operatorname{diag}(\mathcal{M})(M)} c_{v,\operatorname{diag}(M)}(\mathbf{x}) \cdot v.$$

Let us now consider $m_{\mathbf{k}} \in \operatorname{diag}(\mathcal{M})$. We note that by our choice of ϵ and M, $m_{\mathbf{k}}(M) > (1-\epsilon)^d$. On the other hand, let us now consider some $m_{\mathbf{k}'} \in \mathcal{M} \setminus \operatorname{diag}(\mathcal{M})$. We note that by our choice of M, $|m_{\mathbf{k}'}(M)| < (1+\epsilon)^{d-1} \cdot \epsilon$. Therefore,

$$\frac{|m_{\mathbf{k}'}(M)|}{|m_{\mathbf{k}}(M)|} < \frac{(1+\epsilon)^{d-1} \cdot \epsilon}{(1-\epsilon)^d} = \left(\frac{1+\epsilon}{1-\epsilon}\right)^d \cdot \frac{\epsilon}{1+\epsilon}$$

Now, we note that

$$\left(\frac{1+\epsilon}{1-\epsilon}\right)^d = \left(1+\frac{2\epsilon}{1-\epsilon}\right)^d < \left(e^{\left(\frac{2\epsilon}{1-\epsilon}\right)}\right)^d.$$

From our choice of $\epsilon = 1/3d$, we know that $\epsilon \leq 1/3$ since $d \geq 1$. Therefore, $1 - \epsilon \geq 2/3$. So,

$$\frac{2\epsilon}{1-\epsilon} \le 3\epsilon \le \frac{1}{d}$$

So,

$$\frac{|m_{\mathbf{k}'}(M)|}{|m_{\mathbf{k}}(M)|} < \left(\frac{1+\epsilon}{1-\epsilon}\right)^d \cdot \frac{\epsilon}{1+\epsilon} < \frac{e}{3} < 1.$$

This proves that if $v \in \operatorname{diag}(\mathcal{M})(M)$, then there cannot exist any $m \in \mathcal{M} \setminus \operatorname{diag}(\mathcal{M})$, such that v = m(M). Therefore,

$$c_{v,\operatorname{diag}(M)}(\mathbf{x}) = \sum_{m \in \operatorname{diag}(\mathcal{M}): \ m(M) = v} c_m(\mathbf{x}) = \sum_{m \in \mathcal{M}: \ m(M) = v} c_m(\mathbf{x}) = c_{v,M}(\mathbf{x})$$

for all $v \in \operatorname{diag}(\mathcal{M})(M)$. But we already know that $c_{v,M}(\mathbf{x}) = 0$ for all $v \in \mathcal{M}(M)$. Therefore, $c_{v,\operatorname{diag}(M)}(\mathbf{x}) = 0$ for all $v \in \operatorname{diag}(\mathcal{M})(M)$. Therefore,

$$\Psi_{\mathbf{x}}(\operatorname{diag}(M)) = \sum_{v \in \operatorname{diag}(\mathcal{M})(M)} c_{v,\operatorname{diag}(M)}(\mathbf{x}) \cdot v = 0.$$

In order to make effective use of Lemma 30, we will need one more lemma that provides a sufficient condition to prove that $\mathfrak{R}(M, \mathcal{F})$ is not empty.

Definition 31. Let $M = HDH^{\mathsf{T}} \in Sym_q^{\mathsf{pd}}(\mathbb{R})$. We now define the function $\mathcal{S}_M : \mathbb{R} \to Sym_q^{\mathsf{pd}}(\mathbb{R})$, such that

$$\mathcal{S}_M(\theta) = HD^{\theta}H^{\mathsf{T}}.$$

Remark. Let $M = HDH^{\mathsf{T}} \in Sym_q^{\mathsf{pd}}(\mathbb{R})$ have the eigenvalues $(\lambda_1, \ldots, \lambda_q)$. Since these are all positive, D^{θ} is well-defined for all $\theta \in \mathbb{R}$, as the diagonal matrix such that $(D^{\theta})_{ii} = (\lambda_i)^{\theta}$. Consequently, S_M is well-defined. Moreover, we note that

$$\mathcal{S}_M(\theta)_{ij} = (H_{i1}H_{j1})e^{\theta \cdot \log(\lambda_1)} + \dots + (H_{iq}H_{jq})e^{\theta \cdot \log(\lambda_q)},$$

for all $i, j \in [q]$. Therefore, $S_M(\theta)_{ij}$ is a real analytic function in θ for all $i, j \in [q]$.

Lemma 32. Let $M = HDH^{\mathsf{T}} \in Sym_q^{\mathsf{pd}}(\mathbb{R}_{\neq 0})$. Let \mathcal{F} be a countable set of $Sym_q(\mathbb{R})$ -polynomials, such that for each $F \in \mathcal{F}$, there exists some $\theta_F \in \mathbb{R}$ such that $F(\mathcal{S}_M(\theta_F)) \neq 0$. Then, given any interval $(a,b) \subset \mathbb{R}$, there exists some $\theta^* \in (a,b)$ such that $\mathcal{S}_M(\theta^*) \in \mathfrak{R}(M,\mathcal{F}) \cap Sym_q^{\mathsf{pd}}(\mathbb{R}_{\neq 0})$.

Proof. We will let

$$\mathcal{A} = \Big\{ \mathcal{S}_M(\theta)_{ij} : i, j \in [q] \Big\} \cup \Big\{ F(\mathcal{S}_M(\theta)) : F \in \mathcal{F} \Big\}.$$

Since each $F \in \mathcal{F}$ is polynomial in the entries of the input matrix, and since the entries of the matrix $\mathcal{S}_M(\theta)$ are all real analytic functions of θ , we see that each function in \mathcal{A} is a real valued analytic function of θ . We also know that $(\mathcal{S}_M(1))_{ij} = (HDH^T)_{ij} = M_{ij} \neq 0$ for all $i, j \in [q]$, and that $F(\mathcal{S}_M(\theta_F)) \neq 0$ for all $F \in \mathcal{F}$. So, each function in \mathcal{A} is a non-zero analytic function.

Since all the functions in \mathcal{A} are non-zero real analytic functions, the Identity Theorem for Real Analytic Functions ([KP02], Corollary 1.2.7) implies that the set of zeros of any of these functions does not have an accumulation point. In particular, each of these functions has only finitely many zeros within (a, b). Since the countable union of finite sets is only countable, there exists some $a < \theta^* < b$ such that if we let $N = \mathcal{S}_M(\theta^*)$, then $N_{ij} \neq 0$ for all $i, j \in [q]$, and $F(N) \neq 0$ for all $F \in \mathcal{F}$. We may also assume that $\theta^* \neq 0$.

We also note that since $\theta^* \neq 0$, and $\lambda_i > 0$ for all $i \in [q]$,

$$\lambda_1^{y_1} \cdots \lambda_q^{y_q} = 1 \iff (\lambda_1^{y_1} \cdots \lambda_q^{y_q})^{\theta^*} = 1 \iff (\lambda_1^{\theta^*})^{y_1} \cdots (\lambda_q^{\theta^*})^{y_q} = 1,$$

for all $\mathbf{y} \in \chi_q$. This implies that $\mathcal{L}(\lambda_1, \dots, \lambda_q) = \mathcal{L}(\lambda_1^{\theta^*}, \dots, \lambda_q^{\theta^*})$. Now, Lemma 13 implies that $\mathsf{P1-GH}(N) \equiv \mathsf{P1-GH}(M)$. Therefore, $N = \mathcal{S}_M(\theta^*) \in \mathfrak{R}(M, \mathcal{F}) \cap \operatorname{Sym}_q^{\mathsf{pd}}(\mathbb{R}_{\neq 0})$ is the required matrix.

We will now use Lemma 32 to prove a useful corollary.

Corollary 33. Let $M \in Sym_q^{pd}(\mathbb{R}_{>0})$. Let \mathcal{F} be a countable set of $Sym_q(\mathbb{R})$ -polynomials, such that for each $F \in \mathcal{F}$, there exists some $\theta_F \in \mathbb{R}$ such that $F(\mathcal{S}_M(\theta_F)) \neq 0$. Then, there exists some $N = \mathcal{S}_M(\theta^*) \in \mathfrak{R}(M, \mathcal{F}) \cap Sym_q^{pd}(\mathbb{R}_{>0})$.

Proof. We note that $M_{ij} > 0$ for all $i, j \in [q]$. We also note that \mathcal{S}_M defined in Definition 31 is continuous as a function of θ . Since $\mathcal{S}_M(1) = M$, there exists some a < 1 < b such that $\mathcal{S}_M(\theta)_{ij} > 0$ for all $i, j \in [q]$, and $\theta \in (a, b)$. We can apply Lemma 32 with this choice of a < b to find the required $N = \mathcal{S}_M(\theta^*) \in \mathfrak{R}(M, \mathcal{F}) \cap \operatorname{Sym}_q^{pd}(\mathbb{R}_{>0})$.

Theorem 34. Let $M \in Sym_q^{pd}(\mathbb{R}_{>0})$. Let \mathcal{F} be a countable set of $Sym_q(\mathbb{R})$ -polynomials, such that $F(M) \neq 0$ for all $F \in \mathcal{F}$. Let $\mathbf{0} \neq \mathbf{x} \in \chi_q$. If $\Phi_{\mathbf{x}}(M_{11}, \ldots, M_{qq}) \neq 0$, there exists some $N \in \mathfrak{R}(M, \mathcal{F} \cup \mathcal{F}_M \cup \{\Psi_{\mathbf{x}}\}) \cap Sym_q^{pd}(\mathbb{R}_{>0})$.

Proof. We consider $S_M : \mathbb{R} \to \text{Sym}_q^{\mathsf{F}}(\mathbb{R})$ as defined in Definition 31. We see that $F(S_M(1)) = F(M) \neq 0$ for all $F \in \mathcal{F} \cup \mathcal{F}_M$. We also see that the function $\zeta : N \mapsto \Psi_{\mathbf{x}}(\text{diag}(N))$ is a $\text{Sym}_q(\mathbb{R})$ -polynomial. We note that since diag(N) is a diagonal matrix, its eigenvalues are precisely the diagonal values of diag(N), i.e., (N_{11}, \ldots, N_{qq}) . Therefore, from the construction of $\Phi_{\mathbf{x}}$ and $\Psi_{\mathbf{x}}$, we know that

$$\Psi_{\mathbf{x}}(\operatorname{diag}(N)) = 0 \iff \Phi_{\mathbf{x}}(N_{11}, \dots, N_{qq}) = 0$$

for all $N \in \text{Sym}_{q}^{\mathsf{F}}(\mathbb{R})$. In particular, since $\Phi_{\mathbf{x}}(\mathcal{S}_{M}(1)_{11}, \ldots, \mathcal{S}_{M}(1)_{qq}) = \Phi_{\mathbf{x}}(M_{11}, \ldots, M_{qq}) \neq 0$ by our choice of M, we see that $\zeta(\mathcal{S}_{M}(1)) \neq 0$. So, if we let

$$\mathcal{F}' = \mathcal{F} \cup \mathcal{F}_M \cup \{\zeta\},\$$

we see that for each $F \in \mathcal{F}'$, there exists some $\theta_F \in \mathbb{R}$ such that $F(\mathcal{S}_M(\theta_F)) \neq 0$.

We now note that $S_M(0) = HD^0H^T = HH^T = I$. Given $\mathbf{x} \in \chi_q$, we will now let $\epsilon > 0$ be the number whose existence is guaranteed by Lemma 30. Since $S_M(\theta)$ is continuous as a function of θ , we know that there exists some $\delta > 0$, such that for all $0 < \theta < \delta$, for all $i, j \in [q], |S_M(\theta)_{ij} - I_{ij}| < \epsilon$. We can now use Lemma 32 to find $0 < \theta^* < \delta$ such that $M' = S_M(\theta^*) \in \mathfrak{R}(M, \mathcal{F}')$. It should be stressed here that the entries of M' may be negative.

Since $\zeta \in \mathcal{F}'$, we know that $\zeta(M') = \Psi_{\mathbf{x}}(\operatorname{diag}(M')) \neq 0$. Moreover, due to our choice of $\theta^* < \delta$, Lemma 30 allows us to conclude that there exists some $n \geq 1$ such that $\Psi_{\mathbf{x}}(T_nM') \neq 0$. We can now define the $\operatorname{Sym}_q(\mathbb{R})$ -polynomial $\xi : \operatorname{Sym}_q(\mathbb{R}) \to \mathbb{R}$ such that $\xi(N) = \Psi_{\mathbf{x}}(T_nN)$. Since $n \geq 1$ is an integer, the entries of $T_n(N)$ are all homogeneous polynomials in the entries of the matrix N, and therefore, ξ is a $\operatorname{Sym}_q(\mathbb{R})$ -polynomial. We have seen that $\xi(\mathcal{S}_M(\theta^*)) \neq 0$. Therefore, since $M \in \operatorname{Sym}_q^{\operatorname{pd}}(\mathbb{R}_{>0})$, we may use Corollary 33 to find a new $M'' \in \mathfrak{R}(M, \mathcal{F} \cup \mathcal{F}_M \cup \{\xi\}) \cap \operatorname{Sym}_q^{\operatorname{pd}}(\mathbb{R}_{>0})$. Since $\xi(M'') \neq 0$, we see that $\Psi_{\mathbf{x}}(T_nM'') \neq 0$.

But then, Corollary 29 implies that there exists some $N \in \mathfrak{R}(M'', \mathcal{F} \cup \mathcal{F}_M \cup \{\Psi_{\mathbf{x}}\}) \cap \operatorname{Sym}_q^{pd}(\mathbb{R}_{>0})$. Since $\operatorname{Pl-GH}(M'') \leq \operatorname{Pl-GH}(M)$, this means that $N \in \mathfrak{R}(M, \mathcal{F} \cup \mathcal{F}_M \cup \{\Psi_{\mathbf{x}}\}) \cap \operatorname{Sym}_q^{pd}(\mathbb{R}_{>0})$ is the required matrix.

7 Confluence and Pairwise Order Independence

In this section, we will prove the #P-hardness of P1-GH(M) for a subset of the matrices $M \in \text{Sym}_q^{\text{pd}}(\mathbb{R}_{>0})$. We will do so by identifying a large class of matrices $M \in \text{Sym}_q^{\text{pd}}(\mathbb{R}_{>0})$, and a large class of $\mathbf{x} \in \chi_q$, such that $\Phi_{\mathbf{x}}(M_{11}, \ldots, M_{qq}) \neq 0$.

Definition 35. Let I, J be finite sets. Let $\mathbf{x} = (x_i)_{i \in I} \in \mathbb{Z}_{>0}^{|I|}$, and $\mathbf{y} = (y_j)_{j \in J} \in \mathbb{Z}_{>0}^{|J|}$. We say that (\mathbf{x}, \mathbf{y}) is a confluence if it satisfies the following two conditions:

- 1. $\sum_{i \in I} x_i = \sum_{j \in J} y_j$, and
- 2. for any $S_1, S_2 \subseteq I$, and $T_1, T_2 \subseteq J$,

$$\left|\sum_{i\in S_1} x_i = \sum_{j\in T_1} y_j \quad and \quad \sum_{i\in S_2} x_i = \sum_{j\in T_2} y_j\right| \implies \sum_{i\in S_1\cap S_2} x_i = \sum_{j\in T_1\cap T_2} y_j.$$

As all $x_i, y_j > 0$, in a *confluence* clearly $I = \emptyset$ if and only if $J = \emptyset$. We will only be interested in nonempty I and J. While Definition 35 is concise, we will now prove that it has an alternate equivalent definition, with some useful properties.

Lemma 36. Consider (\mathbf{x}, \mathbf{y}) , where $\mathbf{x} = (x_i)_{i \in I} \in \mathbb{Z}_{>0}^{|I|}$, and $\mathbf{y} = (y_i)_{j \in J} \in \mathbb{Z}_{>0}^{|J|}$ for nonempty finite sets I, and J. (\mathbf{x}, \mathbf{y}) is a confluence if and only if there exist partitions $I = S_1 \sqcup \cdots \sqcup S_r$ and $J = T_1 \sqcup \cdots \sqcup T_r$ for some $r \ge 1$, such that for any $S \subseteq I$, $T \subseteq J$,

$$\sum_{i \in S} x_i = \sum_{j \in T} y_j \iff \left\lfloor S = \bigsqcup_{a \in P} S_a \text{ and } T = \bigsqcup_{a \in P} T_a, \text{ for some } P \subseteq [r] \right\rfloor.$$
(16)

Furthermore, the paired partition $\{(S_1, T_1), \ldots, (S_r, T_r)\}$ is unique up to the order of the pairs.

Proof. We will first assume that $I = S_1 \sqcup \cdots \sqcup S_r$, and $J = T_1 \sqcup \cdots \sqcup T_r$ satisfies the condition Eq. (16). If we let P = [r], we immediately get that $\sum_{i \in I} x_i = \sum_{j \in J} y_j$. Moreover, for any $a \in [r]$, let $P = \{a\}$ then $\sum_{i \in S_a} x_i = \sum_{j \in T_a} y_j$.

Now if we consider any $S_1, S_2 \subseteq I$, and $T_1, T_2 \subseteq J$ such that

$$\sum_{i \in S_1} x_i = \sum_{j \in T_1} y_j \quad \text{and} \quad \sum_{i \in S_2} x_i = \sum_{j \in T_2} y_j,$$

we know that there exist some $P_1, P_2 \subseteq [r]$ such that $S_1 = \bigsqcup_{a \in P_1} S_a, T_1 = \bigsqcup_{a \in P_1} T_a, S_2 = \bigsqcup_{a \in P_2} S_a$, and $T_2 = \bigsqcup_{a \in P_2} T_a$. Therefore, if we let $P_3 = P_1 \cap P_2$, we see that $S_1 \cap S_2 = \bigsqcup_{a \in P_3} S_a$, and $T_1 \cap T_2 = \bigsqcup_{a \in P_3} T_a$. Therefore,

$$\sum_{i \in S_1 \cap S_2} x_i = \sum_{a \in P_3} \sum_{i \in S_a} x_i = \sum_{a \in P_3} \sum_{j \in T_a} y_j = \sum_{j \in T_1 \cap T_2} y_j.$$

This proves that (\mathbf{x}, \mathbf{y}) is a confluence.

Conversely, assume that (\mathbf{x}, \mathbf{y}) is a confluence. Let

$$\mathcal{C} = \left\{ (S,T) : S \subseteq I, T \subseteq J, \quad \sum_{i \in S} x_i = \sum_{j \in T} y_j \right\}, \text{ and}$$
$$\mathcal{M} = \left\{ (S,T) \in \mathcal{C} : (S,T) \neq (\emptyset,\emptyset) \text{ and } (\forall S' \subsetneq S, \forall T' \subsetneq T) \left[(S',T') \in \mathcal{C} \implies S' = T' = \emptyset \right] \right\}$$

be the minimal members of C. Since C and consequently \mathcal{M} are finite sets, we may assume that $\mathcal{M} = \{(S_1, T_1), \ldots, (S_r, T_r)\}$ for some $r \geq 1$. We will now show that $\{S_1, \ldots, S_r\}$, and $\{T_1, \ldots, T_r\}$ are the required partitions of I and J respectively.

For $a \neq b \in [r]$, we first consider $(S_a, T_a), (S_b, T_b) \in \mathcal{M}$. By definition of $\mathcal{M} \subseteq \mathcal{C}$, we know that $\sum_{i \in S_a} x_i = \sum_{j \in T_a} y_j$, and $\sum_{i \in S_b} x_i = \sum_{j \in T_b} y_j$. Since (\mathbf{x}, \mathbf{y}) is a confluence, this implies that $\sum_{i \in S_a \cap S_b} x_i = \sum_{j \in T_a \cap T_b} y_j$. But $(S_a \cap S_b) \subseteq S_a$, and $(T_a \cap T_b) \subseteq T_a$. Since $(S_a, T_a) \neq (S_b, T_b)$, we may assume without loss of generality that $(S_a \cap S_b) \subseteq S_a$. But then, since $x_i \in \mathbb{Z}_{>0}$ for all $i \in I$, we see that

$$\sum_{j \in T_a} y_j = \sum_{i \in S_a} x_i > \sum_{i \in S_a \cap S_b} x_i = \sum_{j \in T_a \cap T_b} y_j.$$

Therefore, $(T_a \cap T_b) \subsetneq T_a$ as well. But by our definition of \mathcal{M} , since $(S_a \cap S_b, T_a \cap T_b) \in \mathcal{C}$, and $(S_a, T_a) \in \mathcal{C}$, it must be the case that $S_a \cap S_b = T_a \cap T_b = \emptyset$. Therefore, $\{S_1, \ldots, S_r\}$ and $\{T_1, \ldots, T_r\}$ are both pairwise disjoint.

Since $(S_a, T_a) \in \mathcal{M} \subseteq \mathcal{C}$ for all $a \in [r]$, it is trivial to see that given any $P \subseteq [r]$,

$$\sum_{a \in P} \sum_{i \in S_a} x_i = \sum_{a \in P} \sum_{j \in T_a} y_j, \text{ and therefore, } \left(\bigcup_{a \in P} S_a, \bigcup_{a \in P} T_a \right) \in \mathcal{C}.$$

Now, using induction, we will show that given any $(S,T) \in \mathcal{C}$, we can find some $P \subseteq [r]$, such that $S = \bigsqcup_{a \in P} S_a$, and $T = \bigsqcup_{a \in P} T_a$. This is trivially true when |S| + |T| = 0. Let us now assume that it is true when |S| + |T| < k for some k > 0. Now, we consider some $(S,T) \in \mathcal{C}$ such that |S| + |T| = k. If $(S,T) \in \mathcal{M}$, we are already done. Otherwise, by definition of \mathcal{M} , we know that there exist some $(S',T') \in \mathcal{C}$ such that $S' \subsetneq S$, and $T' \subsetneq T$, but $S' \neq \emptyset$, and $T' \neq \emptyset$. But we note that

$$\sum_{i \in S \setminus S'} x_i = \sum_{i \in S} x_i - \sum_{i \in S'} x_i = \sum_{j \in T} y_j - \sum_{j \in T'} y_j = \sum_{j \in T \setminus T'} y_j,$$

which implies that $(S \setminus S', T \setminus T') \in \mathcal{C}$. We note that 0 < |S'| + |T'| < k. Therefore, $|S \setminus S'| + |T \setminus T'| = (|S|+|T|) - (|S'|+|T'|) < k$. Therefore, our induction hypothesis implies that there exist $P_1, P_2 \subseteq [r]$, such that $S' = \bigcup_{a \in P_1} S_a$, $T' = \bigcup_{a \in P_1} T_a$, $(S \setminus S') = \bigcup_{b \in P_2} S_b$, and $(T \setminus T') = \bigcup_{b \in P_2} T_b$. Therefore, there exists $P = P_1 \cup P_2 \subseteq [r]$ such that

$$S = \bigcup_{a \in P} S_a$$
, and $T = \bigcup_{a \in P} T_a$.

This completes the induction. In particular, since (\mathbf{x}, \mathbf{y}) is a confluence, we know that $(I, J) \in \mathcal{C}$. This then means that there exists some $P \subseteq [r]$ such that $I = \bigsqcup_{a \in P} S_a$, and $J = \bigsqcup_{a \in P} T_a$. Recall that $S_a \cap S_b = \emptyset$ for $a \neq b$. Each $(S_a, T_a) \in \mathcal{M}$ satisfies $S_a \neq \emptyset$. If there is some $a' \notin P$ then $S_{a'} = S_{a'} \cap I = S_{a'} \cap [\bigcup_{a \in P} S_a] = \emptyset$, a contradiction. This implies that P = [r]. This proves that $I = S_1 \sqcup \cdots \sqcup S_r$, and $J = T_1 \sqcup \cdots \sqcup T_r$. This finishes the proof of the existence of the paired partition $\mathcal{M} = \{(S_1, T_1), \ldots, (S_r, T_r)\}$.

For uniqueness, let $\mathcal{M}' = \{(S'_1, T'_1), \dots, (S'_{\rho}, T'_{\rho})\}$ be another such paired partition. We first note that for \mathcal{M} , if $a \neq b \in [r]$ then $\sum_{i \in S_a} x_i \neq \sum_{i \in S_b} x_i$. Otherwise, by the confluence property, $\sum_{j \in T_a} y_j = \sum_{i \in S_a} x_i = \sum_{i \in S_b} x_i$ implies $0 < \sum_{j \in T_a} y_j = \sum_{j \in T_a \cap T_a} y_j = \sum_{i \in S_a \cap S_b} x_i = 0$, a contradiction. The same is true for \mathcal{M}' .

For any $(S_a, T_a) \in \mathcal{M}$, as $\sum_{i \in S_a} x_i = \sum_{j \in T_a} y_j$, there exists $P' \subseteq [\rho]$ such that $S_a = \bigcup_{c \in P'} S'_c$. As $S_a \neq \emptyset$, we have $P' \neq \emptyset$. Pick any $c \in P'$. Then by the same argument there exists $P \subseteq [r]$ such that $S'_c = \bigcup_{b \in P} S_b$. As $\{S_a : a \in [r]\}$ are pairwise disjoint and nonempty, P' must be a singleton set $\{c\}$ and then in turn P for this c is also a singleton set $\{b\}$. This sets up a mapping sending $a \in [r]$ to $c \in [\rho]$. Since $\sum_{i \in S_a} x_i = \sum_{i \in S'_c} x_i = \sum_{i \in S_b} x_i$, we must have a = b. Hence the mapping sending a to c is 1-1. Switching the role of \mathcal{M} and \mathcal{M}' shows that $r = \rho$ and the two paired partitions are in 1-1 correspondence. This proves uniqueness.

Now that we have defined confluences and understood their properties better, we will now relate them to $\mathbf{x} \in \chi_q$, and see how they are relevant to our problem at hand.

Definition 37. Let $\mathbf{0} \neq \mathbf{x} \in \chi_q$. We will define $I^+ = \{i \in [q] : x_i > 0\}$, and $I^- = \{i \in [q] : x_i < 0\}$. Then, we let $\mathbf{x}^+ = (x_i)_{i \in I^+}$, and $\mathbf{x}^- = (-x_i)_{i \in I^-}$. We say that \mathbf{x} is confluent if $(\mathbf{x}^+, \mathbf{x}^-)$ is a confluence.

If \mathbf{x} is confluent, let $I^+ = S_1 \sqcup \cdots \sqcup S_r$ and $I^- = T_1 \sqcup \cdots \sqcup T_r$ be the partition guaranteed by Lemma 36. We say $((S_1, T_1), \ldots, (S_r, T_r))$ is the confluence basis of \mathbf{x} . The confluence basis is uniquely defined, up to the order of the pairs.

An example **x** that is *non*-confluent is $(1, 1, -1, -1) \in \chi_4$. This will play a pivotal role later in isolating tensor products. Before we can study confluent **x**, we will need a few more definitions.

Definition 38. We say that a polynomial $f : \mathbb{R}^d \to \mathbb{R}$ is a 0-1 polynomial with q terms if it can be expressed as

$$f(p_1,\ldots,p_d) = \sum_{i \in [q]} \prod_{t \in [d]} p_t^{x_{i,t}},$$

where each monomial $\prod_{t \in [d]} p_t^{x_{i,t}}$ is distinct, and has a coefficient 1.

Definition 39. Two polynomials $f, g : \mathbb{R}^d \to \mathbb{R}$ are said to be proportional to each other, if there exist integers $x_1, \ldots, x_d \ge 0$ and $y_1, \ldots, y_d \ge 0$ such that for all $\mathbf{p} \in \mathbb{R}^d$,

$$p_1^{x_1} \cdots p_d^{x_d} \cdot f(\mathbf{p}) = p_1^{y_1} \cdots p_d^{y_d} \cdot g(\mathbf{p}).$$

We will first focus on 0-1 polynomials $f : \mathbb{R} \to \mathbb{R}$ in one variable with q terms. Such polynomials must be of the form

$$f(p) = p^{x_1} + p^{x_2} + \dots + p^{x_q}$$

where $0 \le x_1 < x_2 < \cdots < x_q$.

Lemma 40. Let $\mathbf{0} \neq \mathbf{x} \in \chi_q$ be confluent, with the confluence basis $((S_1, T_1), \ldots, (S_r, T_r))$. Let $f_1, \ldots, f_q : \mathbb{R} \to \mathbb{R}$ be 0-1 polynomials with q terms such that $\phi_{\mathbf{x}}(f_1(p), \ldots, f_q(p)) = 0$ for all $p \in \mathbb{R}$. Then, given any $i, j \in [q]$, such that $i, j \in S_a \cup T_a$ for some $a \in [r]$, we have that f_i and f_j are proportional to each other.

Proof. Since f_1, \ldots, f_q are all 0-1 polynomials with q terms, we may assume that

$$f_i(p) = p^{z_{i,1}} + \dots + p^{z_{i,q}}$$

for integers $0 \le z_{i,1} < \cdots < z_{i,q}$ for all $i \in [q]$. We will now let $d_i = z_{i,1}$, and $y_{i,j} = z_{i,j} - d_i$ for all $i, j \in [q]$. If we now let

$$g_i(p) = p^{y_{i,1}} + \dots + p^{y_{i,q}},$$

we see that $f_i(p) = p^{d_i}g_i(p)$ for all $i \in [q]$. By construction, we also see that $y_{i,1} = 0$ for all $i \in [q]$. Therefore, for any $i \neq j \in [q]$, we have $g_i = g_j$ if and only if f_i and f_j are proportional to each other.

Since $\phi_{\mathbf{x}}(f_1(p), \ldots, f_q(p)) = 0$, we see that

$$\prod_{i \in I^+} p^{x_i \cdot d_i} \cdot g_i(p)^{x_i} = \prod_{i \in I^+} f_i(p)^{x_i} = \prod_{i \in I^-} f_i(p)^{-x_i} = \prod_{i \in I^-} p^{-x_i \cdot d_i} \cdot g_i(p)^{-x_i}.$$

By our construction of g_i , we see that $p \nmid g_i$ for all $i \in [q]$. Therefore,

$$\prod_{i \in I^+} p^{x_i \cdot d_i} = \prod_{i \in I^-} p^{-x_i \cdot d_i}, \text{ and } \prod_{i \in I^+} g_i(p)^{x_i} = \prod_{i \in I^-} g_i(p)^{-x_i}.$$

For any $i \in [q]$, if $x_i \neq 0$ then either $i \in I^+$ or $i \in I^-$. For any $i \in I^+ \cup I^-$, there exists a unique $a \in [r]$, and we will denote it by $\sigma(i)$, such that $i \in S_a$ or $i \in T_a$, i.e., $i \in S_{\sigma(i)} \cup T_{\sigma(i)}$.

We will now prove by induction that given any $i, j \in S_a \cup T_a$ for some $a \in [r]$, i.e., $\sigma(i) = \sigma(j)$, we have

$$g_i = g_j.$$

For all $i \in [q]$, given any $1 \le t \le q$, we define a truncated version of g_i ,

$$g_i|_t(p) = p^{y_{i,1}} + \dots + p^{y_{i,t}}$$

By the construction of g_i , we know that $y_{i,1} = 0$ for all $i \in [q]$. So, $g_i|_1(p) = p^{y_{i,1}} = 1$ for all $i \in [q]$. Now, for each $r \leq k \leq rq$, we define the following:

Statement (k): There exist integers $1 \le t_1, \ldots, t_r \le q$ such that $t_1 + \cdots + t_r = k$, and some polynomials h_1, \ldots, h_r such that $g_i|_{t_{\sigma(i)}} = h_{\sigma(i)}$ for all $i \in I^+ \cup I^-$.

We will momentarily assume the following claim, which we shall prove shortly.

Claim 41. If **Statement** (k) is true for some $r \le k < rq$, then there exists some $k < k' \le rq$, such that **Statement** (k') is true.

We have already seen that when $t_1 = \cdots = t_r = 1$, and $h_1 = \cdots = h_r = 1$, $g_i|_{t_{\sigma(i)}} = 1 = h_{\sigma(i)}$ for all $i \in [q]$. Therefore, **Statement** (r) is true. So, **Claim 41** implies that **Statement** (rq) is true. In this case with k = rq, based on the bound on the t_a 's, all $t_a = q$. Then $g_i = g_i|_q = g_i|_{t_{\sigma(i)}} = h_{\sigma(i)}$. Hence if $\sigma(i) = \sigma(j)$ then $g_i = g_j$. From the construction of g_i , it now follows that f_i and f_j are proportional to each other, if $i, j \in S_a \cup T_a$ for some $a \in [r]$.

We will now finish the proof of Lemma 40 by proving Claim 41.

Proof of Claim 41. Let us assume that there exist $1 \leq t_1, \ldots, t_r \leq q$, as well as polynomials h_1, \ldots, h_r that satisfy **Statement** (k) for some $r \leq k < rq$. We will consider the following equation

$$\prod_{i \in I^+} g_i(p)^{x_i} - \prod_{i \in I^+} \left(g_i |_{t_{\sigma(i)}}(p) \right)^{x_i} = \prod_{i \in I^-} g_i(p)^{-x_i} - \prod_{i \in I^-} \left(g_i |_{t_{\sigma(i)}}(p) \right)^{-x_i}.$$
(17)

We can see that Eq. (17) is true since we know that

$$\prod_{i \in I^+} g_i(p)^{x_i} = \prod_{i \in I^-} g_i(p)^{-x_i},$$

and the induction hypothesis tells us that since \mathbf{x} is confluent,

$$\prod_{i \in I^+} \left(g_i |_{t_{\sigma(i)}}(p) \right)^{x_i} = \prod_{a \in [r]} h_a^{\sum_{i \in S_a} x_i} = \prod_{a \in [r]} h_a^{\sum_{i \in T_a} (-x_i)} = \prod_{i \in I^-} \left(g_i |_{t_{\sigma(i)}}(p) \right)^{-x_i}$$

Since we have assumed that $t_1 + \cdots + t_r = k < rq$, we see that there exists some $a \in [r]$ such that $t_a < q$. Since $\sigma : I^+ \cup I^- \to [r]$ is onto, Eq. (17) is not the trivial 0 = 0 equation. Now, we will consider the least degree term of the LHS and RHS of Eq. (17). If we focus on the LHS, we notice that

$$\prod_{i \in I^+} g_i(p)^{x_i} = \sum_{k_{i,j}: \ (\forall i \in I^+, \forall j \in [x_i])[1 \le k_{i,j} \le q]} p^{\sum_{i \in I^+} \left(y_{i,k_{i,1}} + \dots + y_{i,k_{i,x_i}} \right)}$$

However, any term $p^{\sum_{i \in I^+} (y_{i,k_{i,1}} + \dots + y_{i,k_{i,x_i}})}$ for which $k_{i,j} \leq t_{\sigma(i)}$ for all $i \in I^+$ and $j \in [x_i]$ would appear in

$$\prod_{i\in I^+} \left(g_i|_{t_{\sigma(i)}}(p)\right)^x$$

as well, and get cancelled in the LHS of Eq. (17). Therefore, to be a candidate for the least degree term remaining, it must be obtained by choosing one least degree remaining term $p^{y_{i,t_{\sigma(i)}+1}}$ from one factor $g_i(p)$ and the term $p^{y_{j,1}} = p^0 = 1$ for all other factor polynomials. Furthermore, all such product terms from $\prod_{i \in I^+} g_i(p)^{x_i}$ are not cancelled from $\prod_{i \in I^+} \left(g_i|_{t_{\sigma(i)}}(p)\right)^{x_i}$ in Eq. (17), since all terms from this second product have already been used to cancel the corresponding terms from the first product. Hence, this least degree term has degree

$$d^+ = \min_{i \in I^+} \left(y_{i, t_{\sigma(i)} + 1} \right),$$

and this occurs exactly x_i times for each minimizer *i* that achieves this minimum. Summing up, the least degree term of the LHS of Eq. (17) is therefore

$$\sum_{\substack{i \in I^+:\\ h_{i,t_{\sigma(i)}+1}=d^+}} x_i \cdot p^{d^+}$$

IJ

Similarly, if we let

$$d^- = \min_{i \in I^-} \left(y_{i, t_{\sigma(i)} + 1} \right),$$

we find that the least degree term of the RHS of Eq. (17) is

$$\sum_{\substack{i\in I^-:\\y_{i,t_{\sigma(i)}+1}=d^-}} (-x_i)\cdot p^d$$

Since the least degree terms of the LHS and the RHS have to be identical, this implies that $d^+ = d^-$, and also that $\sum_{i \in S} x_i = \sum_{i \in T} (-x_i)$, where $S = \{i \in I^+ : y_{i,t_{\sigma(i)}+1} = d^+\}$, and $T = \{i \in I^- : y_{i,t_{\sigma(i)}+1} = d^-\}$. Since **x** is confluent, by definition, this means that there exists some $P \subseteq [r]$ such that $S = \bigsqcup_{a \in P} S_a$, and $T = \bigsqcup_{a \in P} T_a$. But this means that $y_{i,t_{\sigma(i)}+1} = d^+ = d^-$ for all $i \in \bigcup_{a \in P} (S_a \cup T_a)$.

In other words, we have shown that there exist integers $1 \leq t'_1, \ldots, t'_r \leq q$ such that $t'_a = t_a + 1$ for $a \in P$, and $t'_a = t_a$ for $a \notin P$, as well as polynomials h'_1, \ldots, h'_r such that $g_i|_{t'_{\sigma(i)}} = h'_{\sigma(i)}$ for all $i \in I^+ \cup I^-$. Since $k' = t'_1 + \cdots + t'_r > t_1 + \cdots + t_r = k$, this finishes the proof of Claim 41.

We will now extend Lemma 40 to be applicable for all 0-1 polynomials $f_i : \mathbb{R}^d \to \mathbb{R}$.

Lemma 42. Let $\mathbf{0} \neq \mathbf{x} \in \chi_q$ be confluent, with the confluence basis $((S_1, T_1), \ldots, (S_r, T_r))$. Let $f_1, \ldots, f_q : \mathbb{R}^d \to \mathbb{R}$ be 0-1 polynomials with q terms such that $\phi_{\mathbf{x}}(f_1(\mathbf{p}), \ldots, f_q(\mathbf{p})) = 0$ for all $\mathbf{p} \in \mathbb{R}^d$. Then, given any $i, j \in [q]$, such that $i, j \in S_a \cup T_a$ for some $a \in [r]$, we have that f_i and f_j are proportional to each other.

Proof. From Definition 38, we may assume that

$$f_j(\mathbf{p}) = \sum_{i \in [q]} \prod_{t \in [d]} p_t^{x_{i,t,j}}$$

for all $j \in [q]$. We let $\mathfrak{m} = \max_{i \in [q], t \in [d], j \in [q]}(x_{i,j,t})$. For all $j \in [q]$, we know from Definition 38 that the following sequences are pairwise distinct

$$(x_{1,1,j}, x_{1,2,j}, \dots, x_{1,d,j}) \neq (x_{2,1,j}, x_{2,2,j}, \dots, x_{2,d,j}) \neq \dots \neq (x_{q,1,j}, x_{q,2,j}, \dots, x_{q,d,j}).$$

Therefore for any $m > 2\mathfrak{m}$, if we let

$$z_{i,j} = \sum_{t \in [d]} m^t \cdot x_{i,t,j},$$

we see that for all $j \in [q]$ they are pairwise distinct, $z_{1,j} \neq \cdots \neq z_{q,j}$. Moreover,

$$f_j(p^m, p^{m^2}, \dots, p^{m^d}) = \sum_{i \in [q]} p^{z_{i,j}}$$

for all $j \in [q]$. We can now define $g_j : \mathbb{R} \to \mathbb{R}$ as

$$g_j(p) = f_j(p^m, p^{m^2}, \dots, p^{m^d}).$$

By our choice of $m > 2\mathfrak{m}$, we see that each g_j is a 0-1 polynomial with (exactly) q terms, i.e., no terms get combined.

By hypothesis,

$$\phi_{\mathbf{x}}(f_1(\mathbf{p}),\ldots,f_q(\mathbf{p}))=0$$

for all $\mathbf{p} \in \mathbb{R}^d$. Therefore,

$$\phi_{\mathbf{x}}(g_1(p),\ldots,g_q(p))=0$$

for all $p \in \mathbb{R}$. So, Lemma 40 tells us that given any $i, j \in [q]$ such that $i, j \in S_a \cup T_a$ for some $a \in [r]$, we have that g_i and g_j are proportional to each other. But from our construction of g_i and g_j , this means that there exists some $d_{i,j} \in \mathbb{Z}$ such that

$$\frac{p^{z_{1,i}} + \dots + p^{z_{q,i}}}{p^{z_{1,j}} + \dots + p^{z_{q,j}}} = \frac{g_i(p)}{g_j(p)} = p^{d_{i,j}}.$$

Therefore, there exists some $\sigma \in S_q$ such that

$$p^{z_{1i}-z_{\sigma(1)j}} = p^{z_{2i}-z_{\sigma(2)j}} = \dots = p^{z_{qi}-z_{\sigma(q)j}} = p^{d_{i,j}}$$

From the construction of z_{1i}, \ldots, z_{qi} , and z_{1j}, \ldots, z_{qj} , this implies that

$$\sum_{t \in [d]} m^t \cdot (x_{1,t,i} - x_{\sigma(1),t,j}) = \sum_{t \in [d]} m^t \cdot (x_{2,t,i} - x_{\sigma(2),t,j}) = \dots = \sum_{t \in [d]} m^t \cdot (x_{q,t,i} - x_{\sigma(q),t,j}).$$

We note that $-\mathfrak{m} \leq (x_{1ti} - x_{\sigma(1)tj}), \ldots, (x_{qti} - x_{\sigma(q)tj}) \leq \mathfrak{m}$ for all $t \in [d]$. So, our choice of $m > 2\mathfrak{m}$ implies that for all $t \in [d]$ there exists some $r_t \in \mathbb{Z}$ such that,

$$(x_{1,t,i} - x_{\sigma(1),t,j}) = \dots = (x_{q,t,i} - x_{\sigma(q),t,j}) = r_t.$$

Therefore,

$$\prod_{t \in [d]} p_t^{x_{1,t,i}} = \prod_{t \in [d]} p_t^{r_t} \cdot p_t^{x_{\sigma(1),t,j}}, \quad \dots, \quad \prod_{t \in [d]} p_t^{x_{q,t,i}} = \prod_{t \in [d]} p_t^{r_t} \cdot p_t^{x_{\sigma(q),t,j}}.$$

So,

$$f_i(\mathbf{p}) = \prod_{t \in [d]} p_t^{x_{1,t,i}} + \dots + \prod_{t \in [d]} p_t^{x_{q,t,i}} = \prod_{t \in [d]} p_t^{r_t} \cdot \left(\prod_{t \in [d]} p_t^{x_{1,t,j}} + \dots + \prod_{t \in [d]} p_t^{x_{q,t,j}} \right) = \prod_{t \in [d]} p_t^{r_t} \cdot f_j(\mathbf{p}).$$

This proves that f_i and f_j are proportional to each other.

We are now finally ready to study how confluent $\mathbf{x} \in \chi_q$ interact with matrices $M \in \text{Sym}_q^{\mathsf{F}}(\mathbb{R}_{\neq 0})$. We will initially restrict ourselves to a very special family of matrices.

Definition 43. Let $M \in Sym_q(\mathbb{R})$. We say that M is a row full matrix if the elements of each row of M are pairwise distinct:

$$\forall i \in [q], \ M_{i1} \neq M_{i2} \neq \cdots \neq M_{iq}.$$

Definition 44. Let $M \in Sym_q(\mathbb{R})$. We say that the rows $i \neq j$ of M are order dependent if there exists some $\sigma \in S_q$ such that $(M_{i1}, M_{i2}, \ldots, M_{iq})$ and $(M_{j\sigma(1)}, M_{j\sigma(2)}, \ldots, M_{j\sigma(q)})$ are linearly dependent. We say that $M \in Sym_q(\mathbb{R})$ is pairwise order independent (or p.o. independent) if no rows $i \neq j$ of M are order dependent.

Lemma 45. There exist $Sym_q(\mathbb{R})$ -polynomials $\rho_{\texttt{full}}$ and $\rho_{\texttt{indep}}$ such that given any $M \in Sym_q(\mathbb{R})$, $\rho_{\texttt{full}}(M) \neq 0$ if and only if M is row full, and $\rho_{\texttt{indep}}(M) \neq 0$ if and only if M is p.o. independent.

Proof. For all $i \in [q]$, we can define the $\text{Sym}_q(\mathbb{R})$ -polynomial $\rho_{\texttt{full},i}(M) = \prod_{j_1 \neq j_2 \in [q]} (M_{ij_1} - M_{ij_2})$. We then define $\rho_{\texttt{full}}$ such that

$$\rho_{\texttt{full}}(M) = \prod_{i \in [q]} \rho_{\texttt{full},i}(M)$$

We can easily see that for any $M \in \text{Sym}_q(\mathbb{R})$, $\rho_{\text{full}}(M) \neq 0$ if and only if M is a row full matrix. We can also see that $\rho_{\text{full}}(M)$ is a homogeneous polynomial in the entries of M. So, it is a $\text{Sym}_q(\mathbb{R})$ -polynomial.

We know that given any two vectors $\mathbf{a}, \mathbf{b} \in \mathbb{R}^q$, the Gram determinant $\mathfrak{g} : \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R}$:

$$\mathfrak{g}(\mathbf{a},\mathbf{b}) = \det egin{bmatrix} \langle \mathbf{a},\mathbf{a}
angle & \langle \mathbf{a},\mathbf{b}
angle \\ \langle \mathbf{b},\mathbf{a}
angle & \langle \mathbf{b},\mathbf{b}
angle \end{bmatrix}$$

is a homogeneous polynomial, where $\langle \cdot, \cdot \rangle$ denotes inner product, such that $\mathfrak{g}(\mathbf{a}, \mathbf{b}) \neq 0$ if and only if **a** and **b** are linearly independent. We will now define the function $\rho_{\mathtt{indep},ij}$ for $i \neq j \in [q]$ as

$$\rho_{\mathtt{indep},ij}(M) = \prod_{\sigma \in S_q} \mathfrak{g}\Big((M_{i1}, \dots, M_{iq}), (M_{j\sigma(1)}, \dots, M_{j\sigma(q)})\Big),\tag{18}$$

and the function ρ_{indep} as

$$\rho_{\text{indep}}(M) = \prod_{i \neq j \in [q]} \rho_{\text{indep}, ij}(M).$$
(19)

From the construction, we can easily see that $\rho_{indep}(M) \neq 0$ if and only if M is p.o. independent. We can also see that $\rho_{indep}(M)$ is a homogeneous polynomial in the entries of M. So, it is a $\text{Sym}_q(\mathbb{R})$ -polynomial.

We will now study how confluent $\mathbf{x} \in \chi_q$ interact with row full, p.o. independent matrices in $\operatorname{Sym}_q^{\operatorname{pd}}(\mathbb{R}_{>0})$.

Theorem 46. Let $M \in Sym_q^{pd}(\mathbb{R}_{>0})$ be a row full, p.o. independent matrix. Let \mathcal{F} be a countable set of $Sym_q(\mathbb{R})$ -polynomials such that $F(M) \neq 0$ for all $F \in \mathcal{F}$. Given any confluent $\mathbf{0} \neq \mathbf{x} \in \chi_q$, there exists some row full, p.o. independent $N \in \mathfrak{R}(M, \mathcal{F} \cup \{\Psi_{\mathbf{x}}\}) \cap Sym_q^{pd}(\mathbb{R}_{>0})$.

Proof. We may assume that the entries of M are generated by $\{g_t\}_{t\in[d]}$. We recall that we may replace M with some cM as in Lemma 4. So, we know that for $i, j \in [q]$, there exist unique integers $e_{ij0} \in \{0, 1\}$, and $e_{ij1}, \ldots, e_{ijd} \ge 0$, such that

$$M_{ij} = (-1)^{e_{ij0}} \cdot g_1^{e_{ij1}} \cdots g_d^{e_{ijd}}.$$

Since, $M_{ij} > 0$ for all $i, j \in [q]$ by our choice of M, we see that $e_{ij0} = 0$ for all $i, j \in [q]$. So, from Definition 5, we see that

$$\mathcal{T}_M(\mathbf{p})_{ij} = p_1^{e_{ij1}} \cdots p_d^{e_{ijd}}$$

is a polynomial in **p** for all $i, j \in [q]$. So, we see that for all $i \in [q]$,

$$(\mathcal{T}_M(\mathbf{p})^2)_{ii} = \sum_{j \in [q]} (\mathcal{T}_M(\mathbf{p})_{ij})^2 = \sum_{j \in [q]} \prod_{t \in [d]} p_t^{2e_{ijt}}.$$

Since M is a row full matrix, we know that for all $i \in [q]$, $M_{i1} \neq M_{i2} \neq \cdots \neq M_{iq}$ are pairwise distinct. Since $\{g_t\}_{t\in[d]}$ is a generating set, this implies that for each $i \in [q]$, $(e_{i11}, e_{i12}, \ldots, e_{i1d}) \neq$ $(e_{i21}, e_{i22}, \ldots, e_{i2d}) \neq \cdots \neq (e_{iq1}, e_{iq2}, \ldots, e_{iqd})$ are also pairwise distinct. Therefore, for each $i \in [q]$, we see that $(\mathcal{T}_M(\mathbf{p})^2)_{ii}$ is a 0-1 polynomial with exactly q terms. Since \mathbf{x} is confluent, we know that it has a confluence basis $((S_1, T_1), \ldots, (S_r, T_r))$. Since $r \geq 1$, and $S_1, T_1 \neq \emptyset$, we may pick some $i \in S_1$, and $j \in T_1$. We will now assume that $\phi_{\mathbf{x}} ((\mathcal{T}_M(\mathbf{p})^2)_{11}, \ldots, (\mathcal{T}_M(\mathbf{p})^2)_{qq}) = 0$ for all $\mathbf{p} \in \mathbb{R}^d$. Lemma 42 now implies that by our choice of $i, j \in [q], (\mathcal{T}_M(\mathbf{p})^2)_{ii}$ and $(\mathcal{T}_M(\mathbf{p})^2)_{jj}$ are proportional to each other. But this means that there exists some $c_1, \ldots, c_d \in \mathbb{Z}$ such that

$$\frac{\prod_{t \in [d]} p_t^{2e_{i1t}} + \dots + \prod_{t \in [d]} p_t^{2e_{iqt}}}{\prod_{t \in [d]} p_t^{2e_{j1t}} + \dots + \prod_{t \in [d]} p_t^{2e_{jqt}}} = \prod_{t \in [d]} p_t^{c_t}.$$

Therefore, there exists some $\sigma \in S_q$ such that

$$\prod_{t \in [d]} p_t^{2e_{i1t}} = \prod_{t \in [d]} p_t^{c_t} \cdot \prod_{t \in [d]} p_t^{2e_{j\sigma(1)t}}, \quad \dots, \quad \prod_{t \in [d]} p_t^{2e_{iqt}} = \prod_{t \in [d]} p_t^{c_t} \cdot \prod_{t \in [d]} p_t^{2e_{j\sigma(q)t}}$$

as an identity in **p**. In particular, when we evaluate these expressions at (g_1, \ldots, g_d) , we see that there exists a constant $c = \prod_{t \in [d]} g_t^{c_t} \in \mathbb{R}$ such that

$$((M_{i1})^2, \dots, (M_{iq})^2) = c \cdot ((M_{j\sigma(1)})^2, \dots, (M_{j\sigma(q)})^2)$$

Since $M_{ij} > 0$ for all $i, j \in [q]$ by our choice of M, this implies that in fact

$$(M_{i1},\ldots,M_{iq}) = \sqrt{c} \cdot (M_{j\sigma(1)},\ldots,M_{j\sigma(q)}).$$

This contradicts our assumption that M is p.o. independent, since $i \neq j$. Therefore, our assumption that $\phi_{\mathbf{x}} \left(\mathcal{T}_M(\mathbf{p})^2 \right)_{11}, \ldots, \left(\mathcal{T}_M(\mathbf{p})^2 \right)_{qq} \right) = 0$ must be false.

Since our only restriction on \mathbf{x} was that it be confluent, $\phi_{\mathbf{y}}((\mathcal{T}_M(\mathbf{p})^2)_{11}, \ldots, (\mathcal{T}_M(\mathbf{p})^2)_{qq})$ must be a non-zero polynomial in \mathbf{p} for all confluent $\mathbf{y} \in \chi_q$. In particular, given any $\sigma \in S_q$, we can construct $\mathbf{y} \in \chi_q$ such that $y_i = x_{\sigma(i)}$ for all $i \in [q]$. Since \mathbf{x} is confluent, it is trivial to see that \mathbf{y} is also confluent. Therefore, it follows from the construction of $\Phi_{\mathbf{x}}$ that

$$\Phi_{\mathbf{x}}((\mathcal{T}_M(\mathbf{p})^2)_{11},\ldots,(\mathcal{T}_M(\mathbf{p})^2)_{qq}) = \prod_{\substack{\mathbf{y}\in\chi_q:\exists \ \sigma\in S_q\\\forall \ i\in[q],y_i=x_{\sigma(i)}}} \phi_{\mathbf{y}}((\mathcal{T}_M(\mathbf{p})^2)_{11},\ldots,(\mathcal{T}_M(\mathbf{p})^2)_{qq})$$

is a product of non-zero polynomials, and is therefore, a non-zero polynomial. So, there exists some $\mathbf{p}_{\mathbf{x}} \in \mathbb{R}^d$ such that $\Phi_{\mathbf{x}}((\mathcal{T}_M(\mathbf{p}_{\mathbf{x}})^2)_{11}, \ldots, (\mathcal{T}_M(\mathbf{p}_{\mathbf{x}})^2)_{qq}) \neq 0.$

We will now define $\zeta : N \mapsto \Phi_{\mathbf{x}}((N^2)_{11}, \ldots, (N^2)_{qq})$ for all $N \in \operatorname{Sym}_q(\mathbb{R})$. We can see that since the entries of the matrix N^2 are homogeneous polynomials in the entries of the matrix N, ζ is in fact a $\operatorname{Sym}_q(\mathbb{R})$ -polynomial. From Lemma 45, we also know that there exist $\operatorname{Sym}_q(\mathbb{R})$ polynomials ρ_{full} , and ρ_{indep} . We will now let $\mathcal{F}' = \mathcal{F} \cup \{\zeta, \rho_{\mathsf{full}}, \rho_{\mathsf{indep}}\}$. We know that for every $F \in \mathcal{F}$, $F(\mathcal{T}_M(g_1, \ldots, g_d)) = F(M) \neq 0$. We have also seen that $\zeta(\mathcal{T}_M(\mathbf{p}_{\mathbf{x}})) \neq 0$. Since M is row full and p.o. independent, we also see that $\rho_{\mathsf{full}}(\mathcal{T}_M(g_1, \ldots, g_d)) = \rho_{\mathsf{full}}(M) \neq 0$, and $\rho_{\mathsf{indep}}(\mathcal{T}_M(g_1, \ldots, g_d)) = \rho_{\mathsf{indep}}(M) \neq 0$. Therefore, Lemma 28 tells us that there exists some $M' \in \mathfrak{R}(M, \mathcal{F}') \cap \operatorname{Sym}_q^{\mathsf{pd}}(\mathbb{R}_{>0})$.

Therefore, $F(M') \neq 0$ for all $F \in \mathcal{F}$, and $\Phi_{\mathbf{x}}(((M')^2)_{11}, \ldots, ((M')^2)_{qq}) \neq 0$. We will now define $\xi : N \mapsto \Phi_{\mathbf{x}}(N_{11}, \ldots, N_{qq})$ for all $N \in \operatorname{Sym}_q(\mathbb{R})$, and let $\mathcal{F}'' = \mathcal{F} \cup \{\rho_{\mathsf{full}}, \rho_{\mathsf{indep}}, \xi\}$. From our construction of M', we know that $F(M') \neq 0$ for all $F \in \mathcal{F} \cup \{\rho_{\mathsf{full}}, \rho_{\mathsf{indep}}\}$, and $\xi((M')^2) = \zeta(M') \neq 0$. (We note that as $M' \in \operatorname{Sym}_q^{\mathsf{pd}}(\mathbb{R}_{>0})$, we have $(M')^2 = \mathcal{S}_{M'}(2)$.) Therefore, Corollary 33 implies that there exists some $M'' \in \mathfrak{R}(M', \mathcal{F}'') \cap \operatorname{Sym}_q^{\mathsf{pd}}(\mathbb{R}_{>0})$. Since $\mathsf{Pl-GH}(M') \leq \mathsf{Pl-GH}(M)$, this means that $M'' \in \mathfrak{R}(M, \mathcal{F}'')$. Finally, since $\Phi_{\mathbf{x}}((M'')_{11}, \ldots, (M'')_{qq}) = \xi(M'') \neq 0$, Theorem 34 implies that there exists some $N \in \mathfrak{R}(M'', \mathcal{F} \cup \{\rho_{\mathsf{full}}, \rho_{\mathsf{indep}}, \Psi_{\mathbf{x}}\}) \cap \operatorname{Sym}_q^{\mathsf{pd}}(\mathbb{R}_{>0})$. This N is our required matrix

$$N \in \mathfrak{R}(M, \mathcal{F} \cup \{\rho_{\rho_{\mathtt{full}}, \mathtt{indep}}, \Psi_{\mathbf{x}}\}) \cap \operatorname{Sym}_{q}^{\mathtt{pd}}(\mathbb{R}_{>0}).$$

8 Diagonal and Pairwise Order Distinctness

We will now try to understand the requirement that M be a p.o. independent matrix.

Definition 47. Let $M \in Sym_q(\mathbb{R})$. We say that the rows $i \neq j$ of the matrix M are order identical, if there exists some $\sigma \in S_q$ such that

$$(M_{i1},\ldots,M_{iq})=(M_{j\sigma(1)},\ldots,M_{j\sigma(q)}).$$

We say that $M \in Sym_q(\mathbb{R})$ is pairwise order distinct (or p.o. distinct) if no two rows $i \neq j$ of M are order identical.

Lemma 48. There exists a $Sym_q(\mathbb{R})$ -polynomial ρ_{dist} , such that for any $M \in Sym_q(\mathbb{R})$, $\rho_{dist}(M) \neq 0$ if and only if M is p.o. distinct.

Proof. We know that given any two vectors $\mathbf{a}, \mathbf{b} \in \mathbb{R}^q$,

$$\mathbf{a} = \mathbf{b} \iff \sum_{k \in [q]} (a_k - b_k)^2 = 0.$$

So, we can define the function $\rho_{\mathtt{dist},ij}$ for $i \neq j \in [q]$ as

$$\rho_{\texttt{dist},ij}(M) = \prod_{\sigma \in S_q} \sum_{k \in [q]} (M_{ik} - M_{j\sigma(k)})^2,$$

and the function ρ_{dist} as

$$\rho_{\texttt{dist}}(M) = \prod_{i \neq j \in [q]} \rho_{\texttt{dist}, ij}(M)$$

From the construction, we can easily see that $\rho_{\mathtt{dist}}(M) \neq 0$ if and only if M is p.o. distinct. We can also see that $\rho_{\mathtt{dist}}(M)$ is a homogeneous polynomial in the entries of M. So, it is a $\operatorname{Sym}_q(\mathbb{R})$ -polynomial.

Lemma 49. Let $M \in Sym_q^{pd}(\mathbb{R}_{>0})$ be a row full, p.o. distinct matrix. Let \mathcal{F} be a countable set of $Sym_q(\mathbb{R})$ -polynomials such that $F(M) \neq 0$ for all $F \in \mathcal{F}$. Given any $i \neq j \in [q]$, there exists some row full, p.o. distinct $N \in \mathfrak{R}(M, \mathcal{F} \cup \{\rho_{indep,ij}\}) \cap Sym_q^{pd}(\mathbb{R}_{>0})$, where $\rho_{indep,ij}$ is as defined in Eq. (18) in the proof of Lemma 45.

Proof. Without loss of generality, we may assume that i = 1, and j = 2. We may also assume that $M = HDH^{\mathsf{T}}$. We will consider the function \mathcal{S}_M defined in Definition 31. If there exists some θ such that $\rho_{\mathsf{indep},12}(\mathcal{S}_M(\theta)) \neq 0$, then we can immediately use Corollary 33 to find some $N \in \mathfrak{R}(M, \mathcal{F} \cup \{\rho_{\mathsf{full}}, \rho_{\mathsf{indep},12}\}) \cap \operatorname{Sym}_q^{\mathsf{pd}}(\mathbb{R}_{>0})$, and we would be done. Let us now assume that $\rho_{\mathsf{indep},12}(\mathcal{S}_M(\theta)) = 0$ for all $\theta \in \mathbb{R}$.

Since M is row full and p.o. distinct, it follows that $\rho_{full}(\mathcal{S}_M(1)) = \rho_{full}(M) \neq 0$, and $\rho_{dist}(\mathcal{S}_M(1)) = \rho_{dist}(M) \neq 0$. We will now define $\xi : N \mapsto \rho_{full}(T_2N)$. Since the entries of the matrix T_2N are homogeneous polynomials in the entries of N, ξ is clearly a $\operatorname{Sym}_q(\mathbb{R})$ -polynomial. Since $M \in \operatorname{Sym}_q(\mathbb{R}_{>0})$ is a row full matrix, we note T_2M is also a row full matrix. So, $\xi(\mathcal{S}_M(1)) = \rho_{full}(T_2M) \neq 0$.

Note that $\mathcal{S}_M(0) = I$. Since $\mathcal{S}_M(\theta)_{ij}$ is a continuous function of θ , we know that there exists some $\delta > 0$ such that $|\mathcal{S}_M(\theta)_{ij} - I_{ij}| < \frac{1}{3}$ for all $0 < \theta < \delta$, for all $i, j \in [q]$. With this choice of δ , Lemma 32 allows us to find some $M' = HD^{\theta'}H^{\mathsf{T}} = \mathcal{S}_M(\theta') \in \mathfrak{R}(M, \mathcal{F} \cup \{\rho_{\mathsf{full}}, \rho_{\mathsf{dist}}, \xi\}) \cap \operatorname{Sym}_q^{\mathsf{pd}}(\mathbb{R}_{\neq 0})$, such that $0 < \theta' < \delta$. We should stress here that M' may have negative entries.

From the definition of $\rho_{indep,12}$, we see that

$$\rho_{\texttt{indep},12}(\mathcal{S}_M(\theta)) = \prod_{\sigma \in S_q} \mathfrak{g}\Big((\mathcal{S}_M(\theta)_{11}, \dots, \mathcal{S}_M(\theta)_{1q}), (\mathcal{S}_M(\theta)_{2\sigma(1)}, \dots, \mathcal{S}_M(\theta)_{2\sigma(q)})\Big).$$

Since $\rho_{indep,12}(\mathcal{S}_M(\theta))$ is identically 0, and is the product of real analytic functions, it follows that there must be some $\sigma \in S_q$ such that $\mathfrak{g}\left((\mathcal{S}_M(\theta)_{11},\ldots,\mathcal{S}_M(\theta)_{1q}),(\mathcal{S}_M(\theta)_{2\sigma(1)},\ldots,\mathcal{S}_M(\theta)_{2\sigma(q)})\right) = 0$ for all θ . This implies that for all $i, j \in [q], M'_{1i} \cdot M'_{2\sigma(j)} = M'_{1j} \cdot M'_{2\sigma(i)}$.

From our construction of M', we have $M'_{11}, M'_{22} > 2/3$, and $|M'_{ij}| < 1/3$ for all $i \neq j$. Therefore, given any $t_1 \neq 1, t_2 \neq 2 \in [q]$, we see that $M'_{11} > |M'_{1t_1}|$ and $M'_{22} > |M'_{2t_2}|$. Therefore, $|M'_{11}| \cdot |M'_{22}| > |M'_{1t_1}| \cdot |M'_{2t_2}|$. But we know that for all $j \in [q]$, $|M'_{11}| \cdot |M'_{2\sigma(j)}| = |M'_{1j}| \cdot |M'_{2\sigma(1)}|$. In particular, it must also be true for $j = \sigma^{-1}(2)$. This is only possible if j = 1 or $\sigma(1) = 2$. These are equivalent, namely $\sigma(1) = 2$. Now, let us assume that there exists some arbitrary $\sigma' \in S_q$, such that $\mathfrak{g}\Big((M'_{11}, \ldots, M'_{1q}), (M'_{2\sigma'(1)}, \ldots, M'_{2\sigma'(q)})\Big) = 0$. By the same argument, we have $\sigma'(1) = 2$. But this implies that, for all $t \in [q]$,

$$\frac{|M'_{1t}|}{|M'_{2\sigma'(t)}|} = \frac{|M'_{11}|}{|M'_{22}|} = \frac{|M'_{1t}|}{|M'_{2\sigma(t)}|}.$$

(The ratios are well defined as $M' \in \operatorname{Sym}_q^{pd}(\mathbb{R}_{\neq 0})$.) By construction, M' satisfies the property that $\xi(M') = \rho_{full}(T_2M') \neq 0$. Therefore, T_2M' is row full, and thus $|M'_{2\sigma'(t)}| = |M'_{2\sigma(t)}|$ implies that $\sigma'(t) = \sigma(t)$, for all $t \in [q]$. Therefore, we conclude that

$$\left((\forall \ \theta \in \mathbb{R}) \left[\mathfrak{g} \left((\mathcal{S}_M(\theta)_{11}, \dots, \mathcal{S}_M(\theta)_{1q}), (\mathcal{S}_M(\theta)_{2\sigma'(1)}, \dots, \mathcal{S}_M(\theta)_{2\sigma'(q)}) \right) = 0 \right] \right) \implies \sigma' = \sigma.$$

Therefore, we can define $\zeta_{\tau} : N \mapsto \mathfrak{g}((N_{11}, \ldots, N_{1q}), (N_{2\tau(1)}, \ldots, N_{2\tau(q)}))$ for all $\tau \in S_q$. We can now use Corollary 33 to find $M'' \in \mathfrak{R}(M, \mathcal{F} \cup \{\rho_{\mathtt{full}}, \rho_{\mathtt{dist}}\} \cup \{\zeta_{\tau} : \tau \neq \sigma \in S_q\}) \cap \operatorname{Sym}_q^{\mathtt{pd}}(\mathbb{R}_{>0}).$

Now, let the entries of this matrix M'' be generated by some $\{g_1, \ldots, g_d\}$. Recall that we may assume by replacing M'' with some cM'', that there exist unique integers e_{ijt} for that for all $i, j \in [q]$,

$$(M'')_{ij} = (-1)^{e_{ij0}} g_1^{e_{ij1}} \cdots g_d^{e_{ijd}},$$

where $e_{ij0} \in \{0, 1\}$ and $e_{ijt} \ge 0$ for all $t \in [d]$, for all $i, j \in [q]$. Since $M'' \in \text{Sym}_q(\mathbb{R}_{>0})$, we see that $e_{ij0} = 0$ for all $i, j \in [q]$. Moreover, M'' is a row full matrix by construction. Therefore, for each $i \in [q]$, we have pairwise distinct tuples

$$(e_{i11}, e_{i12}, \dots, e_{i1d}) \neq (e_{i21}, e_{i22}, \dots, e_{i2d}) \neq \dots \neq (e_{iq1}, e_{iq2}, \dots, e_{iqd}).$$

Let $\mathfrak{m} = \max_{i,j \in [q], t \in [d]}(e_{ijt})$. Pick any integer $m > \mathfrak{m}$, and let

$$z_{ij} = \sum_{t \in [d]} m^t \cdot e_{ijt}.$$
We see that because of our choice of $m, z_{i1} \neq z_{i2} \neq \cdots \neq z_{iq}$ are pairwise distinct for all $i \in [q]$. We also observe that for all $i, j \in [q]$,

$$\mathcal{T}_{M''}(p^m,\ldots,p^{m^d})_{ij}=p^{z_{ij}}.$$

Moreover, we may assume that $\zeta_{\sigma}(\mathcal{T}_{M''}(p^m,\ldots,p^{m^d})) = 0$ for all $p \in \mathbb{R}$, since otherwise, we would be able to use Lemma 28 to find the required $N \in \mathfrak{R}(M'', \mathcal{F} \cup \{\rho_{\mathtt{full}}, \rho_{\mathtt{dist}}\} \cup \{\zeta_{\tau} : \tau \in S_q\}) \cap$ $\operatorname{Sym}_q^{\mathtt{pd}}(\mathbb{R}_{>0}) = \mathfrak{R}(M'', \mathcal{F} \cup \{\rho_{\mathtt{full}}, \rho_{\mathtt{dist}}, \rho_{\mathtt{indep}, 12}\}) \cap \operatorname{Sym}_q^{\mathtt{pd}}(\mathbb{R}_{>0})$. Therefore, the two row vectors

$$(p^{z_{11}}, \dots, p^{z_{1q}})$$
 and $(p^{z_{2\sigma(1)}}, \dots, p^{z_{2\sigma(q)}})$

are linearly dependent for all p, and so one is a scalar multiple of the other for every p. This scalar multiple is a function of p, and is obtainable as a ratio of two monomials, and thus is itself a power of p. Therefore, there exists some integer $c \in \mathbb{Z}$ such that $z_{11} - z_{2\sigma(1)} = \cdots = z_{1q} - z_{2\sigma(q)} = c$. We will now define one more function $\mathcal{T}_M^* : \mathbb{R} \to \operatorname{Sym}_q(\mathbb{R})$ such that

$$\mathcal{T}_M^*(p) = (\mathcal{T}_{M''}(p^m, \dots, p^{m^d}))^2.$$

Let us first assume that $\zeta_{\sigma}(\mathcal{T}_{M}^{*}(p)) = 0$ for all $p \in \mathbb{R}$. This implies that

$$(\mathcal{T}_M^*(p)^2)_{11} \cdot (\mathcal{T}_M^*(p)^2)_{2\sigma(t)} = (\mathcal{T}_M^*(p)^2)_{2\sigma(1)} \cdot (\mathcal{T}_M^*(p)^2)_{13}$$

for all $t \in [q]$. We recall that $\sigma(1) = 2$. Therefore, for all $t \in [q]$, we will define $f_t : \mathbb{R} \to \mathbb{R}$ as $f_t(p) = (\mathcal{T}^*_M(p)^2)_{11} \cdot (\mathcal{T}^*_M(p)^2)_{2\sigma(t)} - (\mathcal{T}^*_M(p)^2)_{22} \cdot (\mathcal{T}^*_M(p)^2)_{1t}$, and we find that

$$f_t(p) = \left(\sum_{i \in [q]} p^{2z_{i1}}\right) \cdot \left(\sum_{i \in [q]} p^{z_{i2} + z_{i\sigma(t)}}\right) - \left(\sum_{i \in [q]} p^{2z_{i2}}\right) \cdot \left(\sum_{i \in [q]} p^{z_{i1} + z_{it}}\right)$$
$$= \left(\sum_{i,j \in [q]} p^{2z_{i1} + z_{j2} + z_{j\sigma(t)}}\right) - \left(\sum_{i,j \in [q]} p^{2z_{i2} + z_{j1} + z_{jt}}\right).$$

Since we know that $f_t(p) = 0$ for all $p \in \mathbb{R}$, it follows that $\frac{df_t}{dp}(p) = 0$ for all $p \in \mathbb{R}$ as well. Specifically, we note that

$$\begin{aligned} \frac{df_t}{dp}(1) &= \left(\sum_{i,j\in[q]} 2z_{i1} + z_{j2} + z_{j\sigma(t)}\right) - \left(\sum_{i,j\in[q]} 2z_{i2} + z_{j1} + z_{jt}\right) \\ &= q \left(2\sum_{i\in[q]} (z_{i1}) + \sum_{i\in[q]} (z_{i2}) + \sum_{i\in[q]} (z_{i\sigma(t)})\right) - q \left(2\sum_{i\in[q]} (z_{i2}) + \sum_{i\in[q]} (z_{i1}) + \sum_{i\in[q]} (z_{it})\right) \\ &= q \left(\sum_{i\in[q]} (z_{i1}) - \sum_{i\in[q]} (z_{i2}) + \sum_{i\in[q]} (z_{i\sigma(t)}) - \sum_{i\in[q]} (z_{it})\right).\end{aligned}$$

We recall that $z_{11} - z_{2\sigma(1)} = \cdots = z_{1q} - z_{2\sigma(q)} = c$ for some integer $c \in \mathbb{Z}$. This means that $\sum_{i \in [q]} (z_{i1}) - \sum_{i \in [q]} (z_{i2}) = cq$. Therefore, we have found that

$$\sum_{i \in [q]} (z_{it}) - \sum_{i \in [q]} (z_{i\sigma(t)}) = cq$$

Since this is true for all $t \in [q]$, by the same argument, we also get that

$$\sum_{i \in [q]} (z_{i\sigma(t)}) - \sum_{i \in [q]} (z_{i\sigma(\sigma(t))}) = cq.$$

This can be repeated. Since $\sigma \in S_q$, we know that there exists some $2 \le n \le q$ such that $\sigma^n(1) = 1$. So, we have shown that

$$0 = \sum_{k=1}^{n} \left(\sum_{i \in [q]} (z_{i\sigma^{k-1}(1)}) - \sum_{i \in [q]} (z_{i\sigma^{k}(1)}) \right) = cnq.$$

Since n > 0 and q > 0, this implies that c = 0. But from our construction of z_{ij} , this implies that $e_{1jt} = e_{2\sigma(j)t}$, for all $j \in [q]$ and $t \in [d]$. This implies that $M''_{1j} = M''_{2\sigma(j)}$ for all $j \in [q]$. This contradicts our assumption that $\rho_{dist}(M'') \neq 0$. Therefore, our assumption that $\zeta_{\sigma}(\mathcal{T}^*_M(p)) = 0$ for all $p \in \mathbb{R}$ must be false. We can therefore use Lemma 28 to find $M''' \in \mathfrak{R}(M'', \mathcal{F} \cup \{\zeta_{\tau} : \tau \neq \sigma\} \cup \{\rho_{full}, \rho_{dist}, \zeta^*\}) \cap \operatorname{Sym}_q^{pd}(\mathbb{R}_{>0})$, where $\zeta^* : N \mapsto \zeta_{\sigma}(N^2)$. Since $\zeta_{\tau}(M''') \neq 0$, for all $\tau \neq \sigma$, and $\zeta_{\sigma}((M''')^2) \neq 0$ we can then use Corollary 33 to find the required $N \in \mathfrak{R}(M'', \mathcal{F} \cup \{\rho_{full}, \rho_{dist}, \rho_{indep, 12}\}) \cap \operatorname{Sym}_q^{pd}(\mathbb{R}_{>0})$.

Corollary 50. Let $M \in Sym_q^{pd}(\mathbb{R}_{>0})$ be a row full, p.o. distinct matrix. Let \mathcal{F} be a countable set of $Sym_q(\mathbb{R})$ -polynomials such that $F(M) \neq 0$ for all $F \in \mathcal{F}$. There exists some row full, p.o. independent $N \in \mathfrak{R}(M, \mathcal{F}) \cap Sym_q^{pd}(\mathbb{R}_{>0})$.

Proof. Since M is row full, and p.o. distinct, Lemma 49 implies that there exists some $M^{1,2} \in \mathfrak{R}(M, \mathcal{F} \cup \{\rho_{\texttt{full}}, \rho_{\texttt{dist}}, \rho_{\texttt{indep}, 12}\}) \cap \operatorname{Sym}_q^{\texttt{pd}}(\mathbb{R}_{>0})$. We can now use Lemma 49 again to find $M^{1,3} \in \mathfrak{R}(M^{1,2}, \mathcal{F} \cup \{\rho_{\texttt{full}}, \rho_{\texttt{dist}}, \rho_{\texttt{indep}, 12}, \rho_{\texttt{indep}, 13}\}) \cap \operatorname{Sym}_q^{\texttt{pd}}(\mathbb{R}_{>0})$ which is contained in

$$\mathfrak{R}(M, \mathcal{F} \cup \{\rho_{\mathtt{full}}, \rho_{\mathtt{dist}}, \rho_{\mathtt{indep}, 12}, \rho_{\mathtt{indep}, 13}\}) \cap \mathrm{Sym}_q^{\mathtt{pd}}(\mathbb{R}_{>0})$$

By repeating this up to $\binom{q}{2}$ times, we can find the required

$$N \in \Re(M, \mathcal{F} \cup \{\rho_{\texttt{full}}, \rho_{\texttt{dist}}\} \cup \{\rho_{\texttt{indep}, ij} : i \neq j \in [q]\}) \cap \operatorname{Sym}_q^{\texttt{pd}}(\mathbb{R}_{>0}) = \\ \Re(M, \mathcal{F} \cup \{\rho_{\texttt{full}}, \rho_{\texttt{dist}}, \rho_{\texttt{indep}}\}) \cap \operatorname{Sym}_q^{\texttt{pd}}(\mathbb{R}_{>0}),$$

where ρ_{indep} is defined in Eq. (19).

Lemma 51. Let $M \in Sym_q^{pd}(\mathbb{R}_{>0})$ be a row full, p.o. distinct matrix. If \mathbf{x} is confluent for all $\mathbf{0} \neq \mathbf{x} \in \chi_q$ such that $\Psi_{\mathbf{x}}(M) = 0$, then Pl-GH(M) is #P-hard.

Proof. Let \mathcal{F}_M be as defined in Eq. (15). Let $M' \in \mathfrak{R}(M, \mathcal{F}_M) \cap \operatorname{Sym}_q^{pd}(\mathbb{R}_{>0})$ be the row full, p.o. independent matrix whose existence is guaranteed by Corollary 50.

Let \mathcal{L}_M be the lattice formed by the eigenvalues of M, and let $\mathcal{L}_{M'}$ be the lattice formed by the eigenvalues of M'. By our assumption about M, we know that all $\mathbf{0} \neq \mathbf{x} \in \mathcal{L}_M$ are confluent. From the definition of $\overline{\mathcal{L}}_M$, we see that this immediately implies that all $\mathbf{0} \neq \mathbf{x} \in \overline{\mathcal{L}}_M$ are confluent. By our construction of M', and our choice of \mathcal{F}_M as defined in Eq. (15), we see that $\overline{\mathcal{L}}_{M'} \subseteq \overline{\mathcal{L}}_M$. So, all $\mathbf{0} \neq \mathbf{x} \in \overline{\mathcal{L}}_{M'}$ are confluent.

Let $d = \dim \mathcal{L}_{M'}$, the lattice dimension of $\mathcal{L}_{M'}$. If d > 0, we can pick some $\mathbf{x}_1 \neq \mathbf{0}$ from this lattice. We know that \mathbf{x}_1 is confluent. Since M' is row full and p.o. independent, we can use **Theorem 46** to find some $N_1 \in \mathfrak{R}(M', \mathcal{F}_{M'} \cup \{\Psi_{\mathbf{x}_1}\}) \cap \operatorname{Sym}_q^{pd}(\mathbb{R}_{>0})$. By Lemma 25, $\mathbf{x}_1 \notin \overline{\mathcal{L}_{N_1}}$, where \mathcal{L}_{N_1} is the lattice formed by the eigenvalues of N_1 . We also have $\overline{\mathcal{L}_{N_1}} \subsetneq \overline{\mathcal{L}_{M'}}$. Hence, by Lemma 26, $\dim \mathcal{L}_{N_1} < d$. If the lattice dimension of \mathcal{L}_{N_1} is non-zero, we can repeat this process by picking some $\mathbf{0} \neq \mathbf{x}_2 \in \mathcal{L}_{N_1}$, and then using Theorem 46 once again to find some $N_2 \in \mathfrak{R}(N_1, \mathcal{F}_{N_1} \cup \{\Psi_{\mathbf{x}_1}, \Psi_{\mathbf{x}_2}\}) \cap$ $\operatorname{Sym}_q^{pd}(\mathbb{R}_{>0}) \subseteq \mathfrak{R}(M', \mathcal{F}_{N_1} \cup \{\Psi_{\mathbf{x}_1}, \Psi_{\mathbf{x}_2}\}) \cap \operatorname{Sym}_q^{pd}(\mathbb{R}_{>0})$. We can therefore repeat this process up to d times to find some matrix $N \in \operatorname{Sym}_q^{\mathbf{F}}(\mathbb{R}_{>0})$ such that $\operatorname{Pl-GH}(N) \leq \operatorname{Pl-GH}(M') \leq \operatorname{Pl-GH}(M)$, and the lattice dimension of \mathcal{L}_N is 0. Theorem 18 then proves that $\operatorname{Pl-GH}(N)$ is #P-hard, which implies that $\operatorname{Pl-GH}(M)$ is also #P-hard.

We will now prove that the requirement that the matrix M be p.o. distinct is sufficient, and that the row fullness requirement is redundant. In order to do that, we will make use of an intermediary condition.

Definition 52. Let $M \in Sym_q(\mathbb{R})$. We say that it is diagonal distinct, if $M_{11} \neq M_{22} \neq \cdots \neq M_{qq}$ are pairwise distinct.

Lemma 53. There exists a $Sym_q(\mathbb{R})$ -polynomial ρ_{diag} such that for any $M \in Sym_q(\mathbb{R})$, $\rho_{diag}(M) \neq 0$ if and only if M is diagonal distinct.

Proof. We will define $\rho_{\text{diag}} : \text{Sym}_{q}(\mathbb{R}) \to \mathbb{R}$ such that

$$\rho_{\text{diag}}(N) = \prod_{i \neq j \in [q]} (N_{ii} - N_{jj}).$$

It is easily seen that $\rho_{\text{diag}}(N)$ is a $\text{Sym}_q(\mathbb{R})$ -polynomial such that $\rho_{\text{diag}}(N) \neq 0$ if and only if N is diagonal distinct.

Lemma 54. Let $M \in Sym_q^{pd}(\mathbb{R}_{>0})$ be a row full, p.o. distinct matrix. Let \mathcal{F} be a family of $Sym_q(\mathbb{R})$ polynomial, such that $F(M) \neq 0$ for all $F \in \mathcal{F}$. There exists some diagonal distinct $N \in \mathfrak{R}(M, \mathcal{F}) \cap$ $Sym_q^{pd}(\mathbb{R}_{>0})$.

Proof. If M were itself diagonal distinct, we would be done, so let us assume otherwise. We may assume that $M_{ii} = M_{jj}$ for some $i \neq j$. We assume that the entries of M are generated by some $\{g_t\}_{t\in[d]}$. As before, we may use Lemma 4 to replace M with some cM, and consider the function \mathcal{T}_M as defined in Definition 5. Since $M \in \text{Sym}_q(\mathbb{R}_{>0})$, we see that $e_{ij0} = 0$ for all $i, j \in [q]$. We will first assume that $\rho_{\text{diag}}(\mathcal{T}_M(\mathbf{p})^2) = 0$ for all $\mathbf{p} \in \mathbb{R}^d$. Now, we let $\mathfrak{m} = \max_{i,j \in [q], t \in [d]} (e_{ijt})$, pick some $m > \mathfrak{m}$, and define $z_{ij} = \sum_{t \in [d]} m^t \cdot e_{ijt}$ for all $i, j \in [q]$. We note that

$$\mathcal{T}_M(p^m, p^{m^2}, \dots, p^{m^d})_{ij} = p^{z_{ij}}$$

for all $i, j \in [q]$, and because of our choice of $m, z_{ij} = z_{i'j'}$ if and only if $(e_{ij1}, e_{ij2}, \ldots, e_{ijd}) = (e_{i'j'1}, e_{i'j'2}, \ldots, e_{i'j'd})$. We recall that by our assumption, $\rho_{\text{diag}}(\mathcal{T}_M(p^m, \ldots, p^{m^d})^2) = 0$, for all $p \in \mathbb{R}$. But that implies that for some $i \neq j \in [q], (\mathcal{T}_M(p^m, \ldots, p^{m^d})^2)_{ii} = (\mathcal{T}_M(p^m, \ldots, p^{m^d})^2)_{jj}$. This implies that

$$p^{2z_{i1}} + p^{2z_{i2}} + \dots + p^{2z_{iq}} = p^{2z_{j1}} + p^{2z_{j2}} + \dots + p^{2z_{jq}}$$

for all $p \in \mathbb{R}$. This implies that there exists some $\sigma \in S_q$ such that $z_{it} = z_{j\sigma(t)}$ for all $t \in [q]$. But from our construction of z_{ij} , this implies that $M_{it} = M_{j\sigma(t)}$ for all $t \in [q]$. This contradicts our assumption that M is p.o. distinct. Therefore, our assumption that $\rho_{\text{diag}}(\mathcal{T}_M(\mathbf{p})^2) = 0$ for all $\mathbf{p} \in \mathbb{R}^d$ must be false.

So, $\rho_{\text{diag}}(\mathcal{T}_M(\mathbf{p}^*)^2) \neq 0$ for some $\mathbf{p}^* \in \mathbb{R}^d$. But then, Lemma 28 allows us to first find $M' \in \text{Sym}_q^{\text{pd}}(\mathbb{R}_{>0})$ such that $F(M') \neq 0$ for all $F \in \mathcal{F}$, and $\rho_{\text{diag}}((M')^2) \neq 0$. Then, Corollary 33 allows us to find the required $N \in \mathfrak{R}(M, \mathcal{F} \cup \{\rho_{\text{diag}}\}) \cap \text{Sym}_q^{\text{pd}}(\mathbb{R}_{>0})$.

Somewhat informally, we have proved that

row full
$$+$$
 p.o. distinct \implies diagonal distinct.

We will prove this is actually an equivalence. To prove the other direction of this equivalence will require us to set up some more machinery. Specifically, we need to make use of a new edge gadget.

Given any graph G = (V, E), and any $n \in \mathbb{Z}_{>0}$, we will construct the graph $R_n G$ by replacing each edge of G with a path of length 3, and then replacing the middle edge of each such path with n parallel edges. Clearly, this gadget preserves planarity. Moreover, we note that for any $M \in \text{Sym}_q(\mathbb{R})$, given any graph G = (V, E),

$$Z_M(R_nG) = \sum_{\sigma:V \to [q]} \prod_{(u,v) \in E} \sum_{a,b \in [q]} M_{\sigma(u)a}(M_{ab})^n M_{\sigma(v)b} = \sum_{\sigma:V \to [q]} \prod_{(u,v) \in E} (R_nM)_{\sigma(u)\sigma(v)},$$



(a) The edge gadget R_3 . (b) An

(b) An example graph G, and the graph R_3G .

Figure 2: The edge gadget R_n

where $R_n(M) \in \text{Sym}_q(\mathbb{R})$ such that for all $i, j \in [q]$,

$$(R_n M)_{ij} = \sum_{a,b \in [q]} (M_{ab})^n \cdot M_{ia} M_{jb}.$$

Definition 55. Let $M \in Sym_q(\mathbb{R}_{>0})$. We can now define a function $\mathcal{R}_M : \mathbb{R} \to Sym_q(\mathbb{R}_{>0})$, such that for all $i, j \in [q]$,

$$\mathcal{R}_M(\theta)_{ij} = \sum_{a,b \in [q]} (M_{ab})^{\theta} \cdot M_{ia} M_{jb}.$$

In particular, for any $M \in \text{Sym}_q(\mathbb{R}_{>0})$ and integer $n \ge 1$,

$$\mathcal{R}_M(n) = R_n M. \tag{20}$$

Lemma 56. Let $M \in Sym_q(\mathbb{R}_{>0})$. Then, $\operatorname{Pl-GH}(\mathcal{R}_M(\theta)) \leq \operatorname{Pl-GH}(M)$ for all $\theta \in \mathbb{R}$.

Proof. We know that for any $n \ge 1$, and given any planar graph G = (V, E),

$$Z_M(R_nG) = \sum_{\sigma: V \to [q]} \prod_{(u,v) \in E} \sum_{a,b \in [q]} M_{\sigma(u)a}(M_{ab})^n M_{\sigma(v)b}$$

We recall that

$$\mathcal{P}_m(n) = \left\{ \mathbf{x} = (x_1, \dots, x_m) \in \mathbb{Z}^m \mid (\forall i \in [m]) \ [x_i \ge 0], \text{ and } x_1 + \dots + x_m = n \right\}.$$

We define the *set* (with no duplicates)

$$X_M(G) = \left\{ \prod_{i,j \in [q]} M_{ij}^{k_{ij}} \middle| \mathbf{k} \in \mathcal{P}_{q^2}(|E|) \right\}.$$

For each $\mathbf{k} \in \mathcal{P}_{q^2}(|E|)$, we will define $x_M(\mathbf{k}) = \prod_{i,j \in [q]} M_{ij}^{k_{ij}}$, and

$$Y_M(\mathbf{k}) = \left\{ (\mathbf{a}, \mathbf{b}) \in [q]^{|E|} \mid (\forall i, j \in [q]) \ [k_{ij} = |\{t : a_t = i, b_t = j\}|] \right\}.$$

Then, for each $x \in X_M(G)$, we define

$$c_M(G, x) = \sum_{\sigma: V \to [q]} \left(\sum_{\substack{\mathbf{k} \in \mathcal{P}_{q^2}(|E|):\\ x_M(\mathbf{k}) = x}} \left(\sum_{\substack{\mathbf{a}, \mathbf{b} \in Y_M(\mathbf{k})}} \left(\prod_{(u,v) \in E} M_{\sigma(u)a_u} M_{\sigma(v)b_v} \right) \right) \right).$$

Finally, we note that

$$Z_M(R_nG) = \sum_{x \in X_M(G)} x^n \cdot c_M(G, x).$$

Since each $x \in X_M(G)$ is distinct, and since $|X_M(G)| \leq |E|^{O(1)}$, we can use oracle access to P1-GH(M) to compute $Z_M(R_nG)$ for $n \in [|X_M(G)|]$. We will then have a full rank Vandermonde system of linear equations, which can be solved in polynomial time to find $c_M(G, x)$ for all $x \in X_M(G)$.

We will now note that each $x \in X_M(G)$ is the product of entries of the matrix M. Since $M \in \text{Sym}_q(\mathbb{R}_{>0})$, this implies that every element of $X_M(G)$ is positive. Moreover, we see that for every $\mathbf{k} \in \mathcal{P}_{q^2}(|E|)$, and for every $\theta \in \mathbb{R}$,

$$(x_M(\mathbf{k}))^{\theta} = \prod_{i,j \in [q]} (M_{ij}^{k_{ij}})^{\theta} = \prod_{i,j \in [q]} (M_{ij}^{\theta})^{k_{ij}}$$

is well defined. Therefore, for any $\theta \in \mathbb{R}$, we can compute the quantity

$$\sum_{x \in X_M(G)} x^{\theta} \cdot c_M(G, x) = \sum_{\sigma: V \to [q]} \prod_{(u,v) \in E} \sum_{a,b \in [q]} M_{\sigma(u)a}(M_{ab})^{\theta} M_{\sigma(v)b} = \sum_{\sigma: V \to [q]} \prod_{(u,v) \in E} \mathcal{R}_M(\theta)_{\sigma(u)\sigma(v)}$$

in polynomial time. This proves that $Pl-GH(\mathcal{R}_M(\theta)) \leq Pl-GH(M)$ for all $\theta \in \mathbb{R}$.

Lemma 57. Let $M \in Sym_q^{pd}(\mathbb{R}_{>0})$. Let \mathcal{F} be a countable set of $Sym_q(\mathbb{R})$ -polynomial, such that for all $F \in \mathcal{F}$, there exists some $\theta_F \in \mathbb{R}$ such that $F(\mathcal{R}_M(\theta_F)) \neq 0$. Then, there exists some $N = \mathcal{R}_M(\theta^*) \in \mathfrak{R}(M, \mathcal{F}) \cap Sym_q^{pd}(\mathbb{R}_{>0})$.

Proof. Since each $F \in \mathcal{F}$ is polynomial in the entries of the input matrix, and since each entry of the matrix $\mathcal{R}_M(\theta)$ is a real analytic function of θ , we see that $F(\mathcal{R}_M(\theta))$ is a real analytic function of θ for all $F \in \mathcal{F}$. Moreover, since $F(\mathcal{R}_M(\theta_F)) \neq 0$ for all $F \in \mathcal{F}$, we see that these are non-zero analytic functions.

From the definition of $\mathcal{R}_M(\theta)$, we can see that for all $i, j \in [q]$, $\mathcal{R}_M(\theta)_{ij} > 0$. We also note that $\mathcal{R}_M(1) = M^3$, and since M is positive definite, it follows that M^3 is also positive definite. We note that the eigenvalues of the matrix $\mathcal{R}_M(\theta)$ are continuous functions of the entries of the matrix, and consequently, are continuous functions of θ . This implies that there exists some interval $(a, b) \subset \mathbb{R}$ such that $1 \in (a, b)$, and $\mathcal{R}_M(\theta) \in \operatorname{Sym}_q^{pd}(\mathbb{R}_{>0})$.

Now, since we know that for each $F \in \mathcal{F}$, $F(\mathcal{R}_M(\theta))$ is a non-zero real analytic function, we know that for each $F \in \mathcal{F}$, there are only finitely many zeros within the interval (a, b) ([KP02], Corollary 1.2.7). Since a countable union of finite sets is countable, there must exist some $\theta^* \in (a, b)$ such that $F(\mathcal{R}_M(\theta^*)) \neq 0$. From our choice of (a, b), we also see that $\mathcal{R}_M(\theta^*) \in \operatorname{Sym}_q^{pd}(\mathbb{R}_{>0})$. So, it is the required matrix.

We can now use this gadget to first show that if $M \in \operatorname{Sym}_q^{\operatorname{pd}}(\mathbb{R}_{>0})$ is diagonal distinct, we can find a row full and diagonal distinct $N \in \operatorname{Sym}_q^{\operatorname{pd}}(\mathbb{R}_{>0})$ such that $\operatorname{Pl-GH}(N) \leq \operatorname{Pl-GH}(M)$. Later, we will show that if $M \in \operatorname{Sym}_q^{\operatorname{pd}}(\mathbb{R}_{>0})$ is both row full and diagonal distinct, then we can in fact find a row full, p.o. distinct $N \in \operatorname{Sym}_q^{\operatorname{pd}}(\mathbb{R}_{>0})$.

Lemma 58. Let $M \in Sym_q^{pd}(\mathbb{R}_{>0})$ be a diagonal distinct matrix. Let \mathcal{F} be a countable set of $Sym_q(\mathbb{R})$ -polynomials such that $F(M) \neq 0$ for all $F \in \mathcal{F}$. Then there exists a row full, diagonal distinct $N \in \mathfrak{R}(M, \mathcal{F}) \cap Sym_q^{pd}(\mathbb{R}_{>0})$.

Proof. Since $S_M(0) = I$, there exists some $\delta > 0$ such that $|S_M(\theta)_{ij} - I_{ij}| \leq \frac{1}{3}$ for all $i, j \in [q]$, and $0 < \theta < \delta$. Therefore, we can use Lemma 32 to find $M' = S_M(\theta^*) \in \mathfrak{R}(M, \{\rho_{\text{diag}}\}) \cap \operatorname{Sym}_q^{pd}(\mathbb{R}_{\neq 0})$, for some $\theta^* \in \mathbb{R}$, such that $|(M')_{ij} - I_{ij}| \leq \frac{1}{3}$ for all $i, j \in [q]$.

We can now consider $R_n(M')$ for $n \ge 1$. From the definition, we see that

$$R_n(M')_{ij} = \sum_{a,b \in [q]} (M'_{ab})^n \cdot (M')_{ia} (M')_{jb}.$$

We may now let $X = \{(M')_{ab} : a, b \in [q]\}$, and for each $x \in X$, define

$$c_{ij}(x) = \sum_{a,b \in [q]:(M')_{ab} = x} (M')_{ia} (M')_{jb}.$$

So, we can express $R_n(M')_{ij}$ as

$$R_n(M')_{ij} = \sum_{x \in X} x^n \cdot c_{ij}(x)$$

for all $i, j \in [q]$. Therefore, for any $i \in [q]$, and $j \neq j' \in [q]$, we see that

$$R_n(M')_{ij} - R_n(M')_{ij'} = \sum_{x \in X} x^n \cdot (c_{ij}(x) - c_{ij'}(x)).$$

Since each $x \in X$ is distinct, and |X| = O(1), this forms a full rank Vandermonde system of linear equations. If $R_n(M')_{ij} - R_n(M')_{ij'} = 0$ for all $n \ge 1$, that would imply that $c_{ij}(x) - c_{ij'}(x) = 0$ for all $x \in X$.

In particular, we note that by our choice of M', $\rho_{\text{diag}}(M') \neq 0$. So, $(M')_{jj}$ is not equal to any other diagonal entry of M'. By our choice of M', we also know that the diagonal entries of M'have a higher absolute value than any non-diagonal entry. Therefore, $(M')_{ab} = (M')_{jj}$ implies that (a, b) = (j, j). So,

$$c_{ij}((M')_{jj}) - c_{ij'}((M')_{jj}) = (M')_{ij}(M')_{jj} - (M')_{ij}(M')_{jj'} = 0.$$

Since $M' \in \text{Sym}_q(\mathbb{R}_{\neq 0})$, we know that $(M')_{ij} \neq 0$. So, we have shown that $(M')_{jj} = (M')_{jj'}$, which contradicts our assumption that the diagonal entries of M' have a higher absolute value than any non-diagonal entry. This implies that it cannot be possible that $R_n(M')_{ij} - R_n(M')_{ij'} = 0$ for all $n \geq 1$.

Therefore, given any $i, j \neq j' \in [q]$, there exists some $n_{i,j,j'} \in \mathbb{Z}_{>0}$ such that $(R_{n_{i,j,j'}}(M'))_{ij} - (R_{n_{i,j,j'}}(M'))_{ij'} \neq 0$. We can now define the function $\zeta_{i,j,j'} : N \mapsto (R_{n_{i,j,j'}}N)_{ij} - (R_{n_{i,j,j'}}N)_{ij'}$. From the definition of $R_{n_{i,j,j'}}$ in Definition 55, we can see that $\zeta_{i,j,j'}$ is a $\operatorname{Sym}_q(\mathbb{R})$ -polynomial.

Recall that by construction, $M' = S_M(\theta^*)$ for some $\theta^* \in \mathbb{R}$. So, we have seen that for all $i, j \neq j' \in [q], \zeta_{i,j,j'}(S_M(\theta^*)) \neq 0$. We can now let

$$\mathcal{F}' = \left\{ F' : N \mapsto F(N^3) \mid F \in \mathcal{F} \cup \{\rho_{\text{diag}}\} \right\} \cup \{\zeta_{i,j,j'}\}_{i,j \neq j' \in [q]}.$$

We note that $(\mathcal{S}_M(\frac{1}{3}))^3 = M$, and $F(M) \neq 0$ for all $F \in \mathcal{F} \cup \{\rho_{\mathtt{diag}}\}$, and $\mathcal{S}_M(\theta^*) = M'$. Therefore, Corollary 33 allows us to find $M'' \in \mathfrak{R}(M, \mathcal{F}') \cap \operatorname{Sym}_q^{\mathtt{pd}}(\mathbb{R}_{>0})$. Since $M'' \in \operatorname{Sym}_q(\mathbb{R}_{>0})$, we can now consider the function $\mathcal{R}_{M''}$. We see that by construction of M'', $\mathcal{R}_{M''}(1) = (M'')^3$. So, for all $F \in \mathcal{F} \cup \{\rho_{\text{diag}}\}$, we see that $F(\mathcal{R}_{M''}(1)) = F((M'')^3) \neq 0$ by the fact that $M'' \in \mathfrak{R}(M, \mathcal{F}')$.

Finally, consider the functions $\xi_{i,j,j'}: N \mapsto N_{ij} - N_{ij'}$. Clearly $\xi_{i,j,j'}$ are $\operatorname{Sym}_q(\mathbb{R})$ -polynomials. From our choice of \mathcal{F}' , and $\mathcal{R}_{M''}(n_{i,j,j'}) = R_{n_{i,j,j'}}(M'')$ by Eq. (20), we know that for every $i, j \neq j' \in [q]$, there exists $n_{i,j,j'}$, such that

$$\xi_{i,j,j'}(\mathcal{R}_{M''}(n_{i,j,j'})) = \xi_{i,j,j'}(R_{n_{i,j,j'}}(M'')) = (R_{n_{i,j,j'}}(M''))_{ij} - (R_{n_{i,j,j'}}(M''))_{ij'} = \zeta_{i,j,j'}(M'') \neq 0.$$

Therefore, we apply Lemma 57 to the function set $\mathcal{F} \cup \{\rho_{\mathtt{diag}}\} \cup \{\xi_{i,j,j'} : i \in [q], j \neq j' \in [q]\}$, to get a row full, diagonal distinct $N \in \mathfrak{R}(M'', \mathcal{F}) \cap \operatorname{Sym}_q^{\mathtt{pd}}(\mathbb{R}_{>0})$. Note that $\rho_{\mathtt{full}} = \prod_{i \in [q]} \prod_{j \neq j' \in [q]} \xi_{i,j,j'}$ Since $\mathtt{Pl-GH}(M'') \leq \mathtt{Pl-GH}(M)$, this is the required matrix.

Lemma 59. Let $M \in Sym_q^{pd}(\mathbb{R}_{>0})$ be a diagonal distinct matrix. Let \mathcal{F} be a countable set of $Sym_q(\mathbb{R})$ -polynomials such that $F(M) \neq 0$ for all $M \in \mathcal{F}$. There exists some row full, p.o. distinct $N \in \mathfrak{R}(M, \mathcal{F}) \cap Sym_q^{pd}(\mathbb{R}_{>0})$.

Proof. We will first use Lemma 58 to find a row full, diagonal distinct matrix

$$M' \in \mathfrak{R}(M, \mathcal{F} \cup \{\rho_{\texttt{full}}, \rho_{\texttt{diag}}\}) \cap \operatorname{Sym}_{q}^{\texttt{pd}}(\mathbb{R}_{>0})$$

We may assume that $M' = HDH^{\mathsf{T}}$. Since $\mathcal{S}_{M'}(0) = I$, we know that there exists some $\delta > 0$ such that $|\mathcal{S}_{M'}(\theta)_{ij} - I_{ij}| < \frac{1}{3}$ for all $i, j \in [q]$, for all $0 < \theta < \delta$. We can now use Lemma 32 to find some $M'' = HD^{\theta^*}H^{\mathsf{T}} \in \mathfrak{R}(M', \mathcal{F} \cup \{\rho_{\mathsf{diag}}, \rho_{\mathsf{full}}\})$, where $0 < \theta^* < \delta$. From our choice of θ^* , we can see that $(M'')_{ii} > (M'')_{jk}$ for all $i \in [q]$, and $j \neq k \in [q]$. Moreover, since (M'') is diagonal distinct, we also see that $(M'')_{ii} \neq (M'')_{jj}$ for any $i \neq j$. Therefore, given any $i \neq j$, we see that $(M'')_{ii} \neq (M'')_{jk}$ for all $k \in [q]$. In particular, this implies that there does not exist any $\sigma \in S_q$ such that $((M'')_{i1}, \ldots, (M'')_{iq}) = ((M'')_{j\sigma(1)}, \ldots, (M'')_{j\sigma(q)})$. This proves that $\rho_{\mathsf{dist}}(HD^{\theta^*}H^{\mathsf{T}}) \neq 0$. Now, Corollary 33 allows us to find the required $N \in \mathfrak{R}(M', \mathcal{F} \cup \{\rho_{\mathsf{full}}, \rho_{\mathsf{dist}}\}) \cap \operatorname{Sym}_q^{\mathsf{pd}}(\mathbb{R}_{>0})$.

Informally, Lemma 58 and Lemma 59 together imply the following:

diagonal distinct \implies row full + p.o. distinct.

Together with Lemma 54, this means that

diagonal distinct \iff row full + p.o. distinct.

Lemma 51 now lets us immediately prove the following.

Corollary 60. Let $M \in Sym_q^{pd}(\mathbb{R}_{>0})$ be a diagonal distinct matrix. If \mathbf{x} is confluent for all $\mathbf{0} \neq \mathbf{x} \in \chi_q$ such that $\Psi_{\mathbf{x}}(M) = 0$, then Pl-GH(M) is #P-hard.

Our next goal will be to prove the following:

diagonal distinct \iff p.o. distinct.

We note that Lemma 59 already implies that

diagonal distinct \implies p.o. distinct.

So, we only need to show that the converse is also true.

Lemma 61. Fix any $i \neq j \in [q]$. Let $M \in Sym_q^{pd}(\mathbb{R}_{>0})$ be a matrix such that rows i and j are not order identical. Let \mathcal{F} be a countable set of $Sym_q(\mathbb{R})$ -polynomials such that $F(M) \neq 0$ for all $F \in \mathcal{F}$. Then there exists some $N \in \mathfrak{R}(M, \mathcal{F}) \cap Sym_q^{pd}(\mathbb{R}_{>0})$, such that $N_{ii} \neq N_{jj}$.

Proof. We can assume without loss of generality that i = 1 and j = 2. We will define $\xi : N \mapsto N_{11} - N_{22}$. If there is some θ such that $\xi(\mathcal{S}_M(\theta)) \neq 0$, then we are done by Corollary 33. So we may assume that $\xi(\mathcal{S}_M(\theta)) = 0$ for all $\theta \in \mathbb{R}$. We also recall from Lemma 48 that there exists a $\operatorname{Sym}_q(\mathbb{R})$ -polynomial $\rho_{\operatorname{dist},12}$ such that $\rho_{\operatorname{dist},12}(N) \neq 0$ on any matrix N if and only if rows 1 and 2 of N are not order identical. We will now define $\zeta : N \mapsto \rho_{\operatorname{dist},12}(T_4N)$. Since $\rho_{\operatorname{dist},12}(M) \neq 0$, and $M \in \operatorname{Sym}_q(\mathbb{R}_{>0})$, it follows that $\zeta(M) = \rho_{\operatorname{dist},12}(T_4M) \neq 0$ as well.

We will now let $\mathcal{F}' = \{F' : N \mapsto F(N^3) \mid F \in \mathcal{F}\}$. We see that $F'(\mathcal{S}_M(\frac{1}{3})) = F(\mathcal{S}_M(1)) = F(M) \neq 0$ for all $F' \in \mathcal{F}'$. We also know that $\zeta(\mathcal{S}_M(1)) = \zeta(M) \neq 0$. Since $\mathcal{S}_M(0) = I$, we know that there exists some $\delta > 0$ such that $|\mathcal{S}_M(\theta)_{ij} - I_{ij}| < \frac{1}{3}$ for all $0 < \theta < \delta$, for all $i, j \in [q]$. So, we can use Lemma 32 to find $M' = \mathcal{S}_M(\theta^*) \in \mathfrak{R}(M, \mathcal{F}' \cup \{\zeta\}) \cap \operatorname{Sym}_q^{pd}(\mathbb{R}_{\neq 0})$ where $0 < \theta^* < \delta$.

We know that $\xi(M') = 0$. So, we see that $(M')_{11} = (M')_{22}$. We will now consider $R_n(T_mM')$ for $n \ge 1$ and $m \ge 1$. Let us first assume that $\xi(R_n(T_mM')) = 0$ for all $n, m \ge 1$. We will now define an equivalence relation on [q] such that $i \sim j$ if and only if $(R_n(T_mM'))_{ii} = (R_n(T_mM'))_{jj}$ for all $n, m \ge 1$. By our assumption, we see that $1 \sim 2$. This equivalence relation defines a partition $[q] = S_1 \sqcup \cdots \sqcup S_r$ for some r < q. We may assume without loss of generality that $1, 2 \in S_1$. We will now fix some odd $m \ge 1$, and consider $\xi(R_n(T_mM'))$ for all $n \ge 1$. We know that

$$\xi(R_n(T_mM')) = \sum_{a,b \in [q]} (M'_{ab})^{nm} \cdot ((M'_{1a})^m (M'_{1b})^m - (M'_{2a})^m (M'_{2b})^m)$$

We now let $X = \{(M'_{ab}) : a, b \in [q]\}$, and

$$c_{m,11}(x) = \sum_{a,b \in [q]:M'_{ab} = x} (M'_{1a}M'_{1b})^m, \quad \text{and} \quad c_{m,22}(x) = \sum_{a,b \in [q]:M'_{ab} = x} (M'_{2a}M'_{2b})^m,$$

So, we see that

$$\xi(R_n(T_mM')) = \sum_{x \in X} x^{mn} \cdot (c_{m,11}(x) - c_{m,22}(x)).$$

Since m is odd by our choice, we see that $x^m = (x')^m$ for $x, x' \in X$ implies that x = x'. So, if $\xi(R_n(T_mM')) = 0$ for all $n \ge 1$, this forms a full rank Vandermonde system of linear equations of size O(1). This implies that $c_{m,11}(x) - c_{m,22}(x) = 0$ for all $x \in X$.

Now, we note that by our construction of M', we ensured that every diagonal entry is greater than the absolute values of all non-diagonal entries. Now, given any $i \in [q]$, we know that since $[q] = S_1 \sqcup \cdots S_r$, there exists some $t \in [r]$ such that $i \in S_t$. We now see that

$$c_{m,11}((M')_{ii}) - c_{m,22}((M')_{ii}) = \sum_{j \in S_t} \left((M'_{1j})^{2m} - (M'_{2j})^{2m} \right).$$

Since we can do this for all $i \in [q]$, we see that for all $t \in [r]$,

$$\sum_{j \in S_t} (M'_{1j})^{2m} = \sum_{j \in S_t} (M'_{2j})^{2m}.$$

So,

$$\sum_{j \in [q]} (M'_{1j})^{2m} = \sum_{t \in [r]} \sum_{j \in S_t} (M'_{1j})^{2m} = \sum_{t \in [r]} \sum_{j \in S_t} (M'_{2j})^{2m} = \sum_{j \in [q]} (M'_{2j})^{2m}.$$

Since this is true for all odd m = 2k - 1 with $k \ge 1$, we have a Vandermonde system of equations of the form

$$\begin{pmatrix} (M_{11}')^2 & (M_{12}')^2 & \cdots & (M_{1q}')^2 & (M_{21}')^2 & (M_{22}')^2 & \cdots & (M_{2q}')^2 \\ (M_{11}')^6 & (M_{12}')^6 & \cdots & (M_{1q}')^6 & (M_{12}')^6 & (M_{22}')^6 & \cdots & (M_{2q}')^6 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ (M_{11}')^{4k+2} & (M_{12}')^{4k+2} & \cdots & (M_{1q}')^{4k+2} & (M_{12}')^{4k+2} & (M_{22}')^{4k+2} & \cdots & (M_{2q}')^{4k+2} \end{pmatrix} \times \begin{pmatrix} 1 \\ 1 \\ \vdots \\ 1 \\ -1 \\ -1 \\ \vdots \\ -1 \end{pmatrix} = \begin{pmatrix} 0 \\ \vdots \\ 0 \end{pmatrix}$$

This is only possible if the following condition is satisfied: For any v, let $n_v = |\{i \mid |M'_{1i}| = v\}|$ and $n'_v = |\{i \mid |M'_{2i}| = v\}|$. Then $n_v = n'_v$ for all v. This can be seen by first ordering the elements of the following multisets by magnitude

$$\{|M'_{11}|, |M'_{12}|, \dots |M'_{1q}|\}, \text{ and } \{|M'_{21}|, |M'_{22}|, \dots |M'_{2q}|\},\$$

and then taking a sufficiently large k. Thus the entries of $|M'_{1i}|$ and $|M'_{2i}|$ can be matched in a 1-1 correspondence. Hence, there exists some $\sigma \in S_q$ such that $(M'_{1i})^4 = (M'_{2\sigma(i)})^4$ for all $i \in [q]$. But from our construction of M', we ensured that $\zeta(M') \neq 0$, where ζ was defined such that $\zeta : N \mapsto \rho_{\text{dist},12}(T_4N)$. This contradiction therefore implies that it is not possible that $\xi(R_n(T_m(M'))) = 0$ for all $n, m \geq 1$.

So, we may assume that there exists some $n \geq 1$ and $m \geq 1$ such that $\xi(R_n(T_mM')) \neq 0$. We can now define $\xi' : N \mapsto \xi(R_n(T_mN))$, and we see that $\xi'(\mathcal{S}_M(\theta^*)) \neq 0$. So, Corollary 33 allows us to find some $M'' \in \mathfrak{R}(M, \mathcal{F}' \cup \{\xi'\}) \cap \operatorname{Sym}_q^{pd}(\mathbb{R}_{>0})$. We can assume that the entries of M'' are generated by some $\{g''_t\}_{t\in[d]}$. We also know that using Lemma 4, we can replace M'' with some cM''. If we now let $\xi'' : N \mapsto (R_n(N))_{11} - (R_n(N))_{22}$, we can see that since $M'' \in \operatorname{Sym}_q(\mathbb{R}_{>0}), \ \xi''(\mathcal{T}_{M''}((g''_1)^m, \dots, (g''_d)^m)) = \xi''(T_mM'') = \xi'(M'') \neq 0$. So, Lemma 28 allows us to find some $M''' \in \mathfrak{R}(M'', \mathcal{F}' \cup \{\xi''\}) \cap \operatorname{Sym}_q^{pd}(\mathbb{R}_{>0})$. We note that since $M''' \in \operatorname{Sym}_q(\mathbb{R}_{>0}), \ \xi(\mathcal{R}_{M'''}(n)) = \xi(R_n(M''')) = \xi''(M''') \neq 0$. Also, $\mathcal{R}_{M'''}(1) = (M''')^3$, and so, for all $F \in \mathcal{F}$, Finally, Lemma 57 allows us to find the required $N \in \mathfrak{R}(M''', \mathcal{F} \cup \{\xi\}) \cap \operatorname{Sym}_q^{pd}(\mathbb{R}_{>0})$.

We can repeatedly apply Lemma 61 to all distinct pairs $i \neq j \in [q]$, while each time incorporating an additional polynomial ξ in the set \mathcal{F} that represents the last pair that is ensured to be not order identical. More formally, we can prove the following corollary. **Corollary 62.** Let $M \in Sym_q^{pd}(\mathbb{R}_{>0})$ be a p.o. distinct matrix, that is not necessarily row full. Let \mathcal{F} be a countable set of $Sym_q(\mathbb{R})$ -polynomials such that $F(M) \neq 0$ for all $F \in \mathcal{F}$. There exists some diagonal distinct $N \in \mathfrak{R}(M, \mathcal{F}) \cap Sym_q^{pd}(\mathbb{R}_{>0})$.

Proof. We can define the $\operatorname{Sym}_q(\mathbb{R})$ -polynomial $\rho_{\operatorname{diag},ij}$ such that $\rho_{\operatorname{diag},ij}(N) : N \mapsto N_{ii} - N_{jj}$. Since M is p.o. distinct, Lemma 61 allows us to find $M^{(12)} \in \mathfrak{R}(M, \mathcal{F} \cup \{\rho_{\operatorname{dist}}, \rho_{\operatorname{diag},12}\}) \cap \operatorname{Sym}_q^{\operatorname{pd}}(\mathbb{R}_{>0})$. We can repeat this process again with $M^{(12)}$ to find $M^{(13)} \in \mathfrak{R}(M^{(12)}, \mathcal{F} \cup \{\rho_{\operatorname{dist}}, \rho_{\operatorname{diag},12}, \rho_{\operatorname{diag},13}\}) \cap \operatorname{Sym}_q^{\operatorname{pd}}(\mathbb{R}_{>0})$.

After repeating this for all $i \neq j \in [q]$, we obtain $N \in \mathfrak{R}(M, \mathcal{F} \cup \{\rho_{\mathtt{dist}}\} \cup \{\rho_{\mathtt{diag},ij} : i \neq j \in [q]\}) \cap \operatorname{Sym}_q^{\mathtt{pd}}(\mathbb{R}_{>0})$. So, we see that $N \in \mathfrak{R}(M, \mathcal{F}_M \cup \{\rho_{\mathtt{diag}}\}) \cap \operatorname{Sym}_q^{\mathtt{pd}}(\mathbb{R}_{>0})$ is the required matrix.

This finishes our proof that

row full + p.o. distinct
$$\iff$$
 diagonal distinct \iff p.o. distinct.

We can therefore also prove the following corollary.

Corollary 63. Let $M \in Sym_q^{pd}(\mathbb{R}_{>0})$ be a p.o. distinct matrix. If \mathbf{x} is confluent for all $\mathbf{0} \neq \mathbf{x} \in \chi_q$ such that $\Psi_{\mathbf{x}}(M) = 0$, then Pl-GH(M) is #P-hard.

Proof. Corollary 62 implies that there exists some diagonal distinct $N \in \mathfrak{R}(M, \mathcal{F}_M) \cap \operatorname{Sym}_q^{pd}(\mathbb{R}_{>0})$. Corollary 60 then proves that $\operatorname{Pl-GH}(N)$ is $\#\operatorname{P-hard}$, which then implies that $\operatorname{Pl-GH}(M)$ is also $\#\operatorname{P-hard}$.

9 Hardness for Diagonal Distinct 4×4 matrices

We are now ready to use all this quite elaborate machinery that we have built to prove a complexity dichotomy in the case where q = 4.

Lemma 64. Let $\lambda_1, \ldots, \lambda_4 > 0$. If $\overline{\mathcal{L}(\lambda_1, \ldots, \lambda_4)}$ contains any $\mathbf{x} \neq \mathbf{0}$ that is non-confluent, then such an \mathbf{x} must satisfy $|x_1| = |x_2| = |x_3| = |x_4|$, and in this case, $(1, 1, -1, -1) \in \overline{\mathcal{L}(\lambda_1, \ldots, \lambda_4)}$.

Proof. By definition $\mathbf{0} \neq \mathbf{x} \in \chi_q$, and $\sum_i x_i = 0$. Let $I^+ = \{i \in [4] : x_i > 0\}$, and $I^- = \{i \in [4] : x_i < 0\}$. Since $\mathbf{x} \neq \mathbf{0}$, we know that both $I^+, I^- \neq \emptyset$. If $|I^+| = 1$ or $|I^-| = 1$, \mathbf{x} must be confluent, by Lemma 36, since the only choice for nonempty $S \subseteq I^+$ and $T \subseteq I^-$ such that $\sum_{i \in S} x_i = \sum_{j \in T} (-x_j)$ is in fact $S = I^+$ and $T = I^-$. This leaves us to consider as the only non-trivial case, the scenario where $|I^+| = |I^-| = 2$. We note that if $\mathbf{x} \in \overline{\mathcal{L}(\lambda_1, \ldots, \lambda_4)}$, then $\mathbf{y} \in \overline{\mathcal{L}(\lambda_1, \ldots, \lambda_4)}$ as well for any \mathbf{y} such that $y_i = x_{\sigma(i)}$ for some $\sigma \in S_4$. So, we may assume without loss of generality that $I^+ = \{1, 4\}$, and $I^- = \{2, 3\}$.

Since **x** is not confluent, there must be $S_1, S_2 \subseteq \{1, 4\}$, and $T_1, T_2 \subseteq \{2, 3\}$ such that

$$\sum_{i \in S_1} x_i = \sum_{j \in T_1} (-x_j) \text{ and } \sum_{i \in S_2} x_i = \sum_{j \in T_2} (-x_j) \text{ but } \sum_{i \in S_1 \cap S_2} x_i \neq \sum_{j \in T_1 \cap T_2} (-x_j).$$

Since all $x_i \neq 0$, if one of S_1, S_2, T_1 or $T_2 = \emptyset$, so is the corresponding set, violating the last inequality. So, we may rule out any empty sets, and also the case where $S_1 = S_2 = \{1, 4\}$ and the case where $T_1 = T_2 = \{2, 3\}$. Without loss of generality, $S_1 \neq \{1, 4\}$, and we may assume that $S_1 = \{1\}$. This implies that $T_1 \neq \{2, 3\}$ since we know that $x_1 < x_1 + x_4 = (-x_2) + (-x_3)$. Once again, without loss of generality, we may assume that $T_1 = \{2\}$.

Now, if $S_2 = \{1, 4\}$, that would force $T_2 = \{2, 3\}$ since $x_1 + x_4 > (-x_2)$ and $x_1 + x_4 > (-x_3)$. But then, $S_1 \cap S_2 = S_1$ and $T_1 \cap T_2 = T_1$, which implies that $\sum_{i \in S_1 \cap S_2} x_i = \sum_{j \in T_1 \cap T_2} (-x_j)$, a contradiction. Therefore, $S_2 \neq \{1, 4\}$. So the only possibilities are $S_2 = \{1\}$ or $S_2 = \{4\}$. By symmetry, the only possibilities for T_2 are $T_2 = \{2\}$ or $T_2 = \{3\}$.

If $S_2 = \{1\}$, then we claim $T_2 = \{3\}$, for otherwise $T_2 = \{2\}$ and that would lead to an equality $\sum_{i \in S_1 \cap S_2} x_i = x_1 = -x_2 = \sum_{j \in T_1 \cap T_2} (-x_j)$, a contradiction. Then we have $x_1 = -x_2$ and $x_1 = -x_3$ which leads to $x_4 = (x_1 + x_4) - x_1 = (-x_2 + -x_3) - x_1 = x_1$. This proves that $|x_1| = |x_2| = |x_3| = |x_4|$, and also that $(x, -x, -x, x) \in \mathcal{L}(\lambda_1, \dots, \lambda_4)^{\sigma}$ for some $x \in \mathbb{Z}_{>0}$, and $\sigma \in S_4$. This means that $(\lambda_{\sigma(1)})^x (\lambda_{\sigma(2)})^{-x} (\lambda_{\sigma(3)})^{-x} (\lambda_{\sigma(4)})^x = 1$. Since $\lambda_1, \dots, \lambda_4 > 0$, this implies that $(\lambda_{\sigma(1)}) (\lambda_{\sigma(2)})^{-1} (\lambda_{\sigma(3)})^{-1} (\lambda_{\sigma(4)}) = 1$. Therefore, $(1, -1, -1, 1) \in \overline{\mathcal{L}(\lambda_1, \dots, \lambda_4)}$.

Finally, if $S_2 = \{4\}$, then $S_1 \cap S_2 = \emptyset$. Then $T_2 = \{2\}$, for otherwise $T_2 = \{3\}$, would give $T_1 \cap T_2 = \emptyset$ and then we would have an equality $\sum_{i \in S_1 \cap S_2} x_i = 0 = \sum_{j \in T_1 \cap T_2} (-x_j)$, a contradiction. Then we have $x_1 = -x_2$ and $x_4 = -x_2$, which also leads to $-x_3 = x_1$ as above. Hence, even in this case, we find that $|x_1| = |x_2| = |x_3| = |x_4|$, and that (x, -x, -x, x) in $\overline{\mathcal{L}(\lambda_1, \ldots, \lambda_4)}$, for some $x \in \mathbb{Z}_{>0}$. This once again implies that $(1, -1, -1, 1) \in \overline{\mathcal{L}(\lambda_1, \ldots, \lambda_4)}$.

Our goal will be to now show that given a diagonal distinct matrix $M \in \text{Sym}_4^{\text{pd}}(\mathbb{R}_{>0})$, we can find some $N \in \mathfrak{R}(M, \mathcal{F}_M)$ with eigenvalues (μ_1, \ldots, μ_4) such that $(1, -1, -1, 1) \notin \overline{\mathcal{L}}(\mu_1, \ldots, \mu_4)$. Corollary 60 then implies that P1-GH(N) is #P-hard, which means that P1-GH(M) is also #P-hard. On the other hand, if we are unable to find such a matrix N, then we will show that M is isomorphic to $A \otimes B$ for some $A, B \in \text{Sym}_2(\mathbb{R}_{>0})$. Here for any $\sigma \in S_4$ we define the matrix M^{σ} to be such that $(M^{\sigma})_{ij} = M_{\sigma(i)\sigma(j)}$. We say that M^{σ} is isomorphic to M.

Lemma 65. There exist $Sym_4(\mathbb{R})$ -polynomials, $\varrho_{\texttt{tensor}}$ and $\rho_{\texttt{tensor}}$ such that given any $M \in Sym_4(\mathbb{R}_{\neq 0})$,

- $\varrho_{\text{tensor}}(M) = 0$ if and only if $M = A \otimes B$ for some $A, B \in Sym_2(\mathbb{R})$, and
- $\rho_{\text{tensor}}(M) = 0$ if and only if M is isomorphic to $A \otimes B$ for some $A, B \in Sym_2(\mathbb{R})$.

Proof. We will first define ρ_{tensor} as

$$\varrho_{\texttt{tensor}}(N) = (N_{14} - N_{23})^4 + (N_{11}N_{44} - N_{22}N_{33})^2 + \sum_{i \in [4]} (N_{i1}N_{i4} - N_{i2}N_{i3})^2.$$

We can see that ρ_{tensor} is a homogeneous polynomial in the entries of the matrix. So, it is a $\text{Sym}_4(\mathbb{R})$ -polynomial. Note that if we index the rows and columns not from $\{1, 2, 3, 4\}$ but rather from $D = \{00, 01, 10, 11\}$, the polynomial $\rho_{\text{tensor}}(N)$ takes the form

$$\varrho_{\texttt{tensor}}(N) = (N_{00,11} - N_{01,10})^4 + (N_{00,00}N_{11,11} - N_{01,01}N_{10,10})^2 + \sum_{ab\in D} (N_{ab,00}N_{ab,11} - N_{ab,01}N_{ab,10})^2.$$

From this form, it is clear that $\rho_{tensor}(N)$ is invariant if we flip the bit a or b in the matrix N.

We see that $\rho_{tensor}(M) = 0$ if and only if $M_{14} = M_{23}$, $M_{11}M_{44} - M_{22}M_{33}$, and

$$M_{11}M_{14} = M_{12}M_{13}, \quad M_{12}M_{24} = M_{22}M_{23}, \quad M_{13}M_{34} = M_{23}M_{33}, \quad M_{14}M_{44} = M_{24}M_{34}$$

Let $M \in \text{Sym}_4(\mathbb{R}_{\neq 0})$. We will now let $x = \frac{M_{13}}{M_{11}}$, and $y = \frac{M_{33}}{M_{11}}$. (As we assume the entries of M are non-zero, the divisions are well defined.) Since $M_{11}M_{14} = M_{12}M_{13}$, this means that $M_{14} = x \cdot M_{12}$. Similarly, since $M_{11}M_{44} = M_{22}M_{33}$, we see that $M_{44} = y \cdot M_{22}$. Since (by symmetry) $M_{23} = M_{14} = x \cdot M_{12}$, and $M_{12}M_{24} = M_{22}M_{23}$, we see that $M_{24} = x \cdot M_{22}$. Since $M_{33} = y \cdot M_{11}$, and $M_{13}M_{34} = M_{23}M_{33}$, we see that $M_{34} = \frac{xy \cdot M_{11}M_{12}}{x \cdot M_{11}} = y \cdot M_{12}$. Putting everything together, we see that

$$M = \begin{pmatrix} M_{11} & M_{12} & xM_{11} & xM_{12} \\ M_{12} & M_{22} & xM_{12} & xM_{22} \\ xM_{11} & xM_{12} & yM_{11} & yM_{12} \\ xM_{12} & xM_{22} & yM_{12} & yM_{22} \end{pmatrix} = \begin{pmatrix} 1 & x \\ x & y \end{pmatrix} \otimes \begin{pmatrix} M_{11} & M_{12} \\ M_{12} & M_{22} \end{pmatrix}.$$

So, we see that $\rho_{\texttt{tensor}}(M) = 0$ implies that $M = A \otimes B$ for some $A, B \in \text{Sym}_2(\mathbb{R})$.

Now, let us assume that $M = A \otimes B$ for some $A, B \in \text{Sym}_2(\mathbb{R})$. So, we see that

$$M = \begin{pmatrix} A_{11}B_{11} & A_{11}B_{12} & A_{12}B_{11} & A_{12}B_{12} \\ A_{11}B_{12} & A_{11}B_{22} & A_{12}B_{12} & A_{12}B_{22} \\ A_{12}B_{11} & A_{12}B_{12} & A_{22}B_{11} & A_{22}B_{12} \\ A_{12}B_{12} & A_{12}B_{22} & A_{22}B_{12} & A_{22}B_{22} \end{pmatrix}$$

But now we can verify that $\rho_{\text{tensor}}(M) = 0$. This proves that $\rho_{\text{tensor}}(M) = 0$ if and only if $M = A \otimes B$ for some $A, B \in \text{Sym}_2(\mathbb{R})$.

Given any $M \in \text{Sym}_4(\mathbb{R}_{\neq 0})$, and $\sigma \in S_4$, we can define M^{σ} to be the matrix such that $(M^{\sigma})_{ij} = M_{\sigma(i)\sigma(j)}$. Matrices isomorphic to M take the form M^{σ} under a simultaneous row and column permutation by some σ . Now, we can define

$$\rho_{\texttt{tensor}}(M) = \prod_{\sigma \in S_4} \varrho_{\texttt{tensor}}(M^{\sigma}).$$

We can see that for each $\sigma \in S_4$, $\rho_{\texttt{tensor}}(M^{\sigma})$ is a homogeneous polynomial in the entries of M. So, $\rho_{\texttt{tensor}}$ is a $\text{Sym}_q(\mathbb{R})$ -polynomial, such that $\rho_{\texttt{tensor}}(M) = 0$ implies that M is isomorphic to $A \otimes B$ for some $A, B \in \text{Sym}_2(\mathbb{R})$. Similarly, if M is isomorphic to some $A \otimes B$, it follows that there exists some $\sigma \in S_4$ such that $M^{\sigma} = A \otimes B$. This implies that $\rho_{\texttt{tensor}}(M) = 0$.

Remark. For $M \in Sym_4(\mathbb{R}_{>0})$, being expressible as $M = A \otimes B$ for some $A, B \in Sym_2(\mathbb{R})$ is equivalent to being expressible as $M = A \otimes B$ for some $A, B \in Sym_2(\mathbb{R}_{>0})$. This is because if A or B have a zero entry, there would be a zero entry in M as well, and if A and B have any entries a and b of the opposite signs, then ab < 0 would be an entry of M. Then all entries of A have the same sign, as well as that of B, and their signs are the same. Finally, if they are both -, then replace A and B by -A and -B. We will now show that given a diagonal distinct matrix $M \in \operatorname{Sym}_{4}^{\operatorname{pd}}(\mathbb{R}_{>0})$, if we cannot find some $N \in \mathfrak{R}(M, \mathcal{F}_M)$ such that $\Psi_{(1,-1,-1,1)}(N) \neq 0$, then M needs to satisfy more and more conditions, until we are able to prove that $\rho_{\operatorname{tensor}}(M) = 0$. We will start with a lemma that is applicable for all $M \in \operatorname{Sym}_{4}^{\operatorname{pd}}(\mathbb{R}_{>0})$, and not just diagonally distinct matrices.

Lemma 66. Let $M \in Sym_4^{pd}(\mathbb{R}_{>0})$. Let \mathcal{F} be a countable set of $Sym_4(\mathbb{R})$ -polynomials such that $F(M) \neq 0$ for all $F \in \mathcal{F}$. Let ξ be the $Sym_4(\mathbb{R})$ -polynomial such that $\xi : N \mapsto \phi_{(1,-1,-1,1)}(N_{11},\ldots,N_{44})$. Then there exists some $N \in \mathfrak{R}(M, \mathcal{F} \cup \{\xi\}) \cap Sym_4^{pd}(\mathbb{R}_{>0})$, unless $M_{11} \cdot M_{44} = M_{22} \cdot M_{33}$, and $M_{14} = M_{23}$.

Proof. If $\xi(M) \neq 0$, then we are done already, so we may assume that $\xi(M) = 0$. This immediately implies that $M_{11} \cdot M_{44} = M_{22} \cdot M_{33}$. So, our goal now is just to show that if no $N \in \mathfrak{R}(M, \mathcal{F} \cup \{\xi\}) \cap \operatorname{Sym}_{4}^{pd}(\mathbb{R}_{>0})$ can be found, then $M_{14} = M_{23}$.

Let the entries of M be generated by some $\{g_t\}_{t\in[d]}$. We know from Lemma 4 that we can replace M with some $c \cdot M$ such that the entries of M are generated by $\{g_t\}_{t\in[d]}$ with $e_{ijt} \geq 0$ for all $i, j \in [4]$, and $t \in [d]$. Since $M \in \text{Sym}_4(\mathbb{R}_{>0})$, we also know that $e_{ij0} = 0$ for all $i, j \in [4]$. We will now let $\mathfrak{m} = \max_{i, j \in [4], t \in [d]}(e_{ijt})$, and pick some $m > \mathfrak{m}$. We then let

$$z_{ij} = \sum_{t \in [d]} m^t e_{ijt}$$

for all $i, j \in [4]$. We will now define $\mathcal{T}_M^* : \mathbb{R} \to \operatorname{Sym}_4(\mathbb{R})$ such that

(

$$\mathcal{T}_M^*(p)_{ij} = \mathcal{T}_M(p^m, \dots, p^{m^d})_{ij} = p^{z_{ij}}$$

for all $i, j \in [4]$. We will now define the Sym₄(\mathbb{R})-polynomial ξ_2 such that $\xi_2 : N \mapsto \xi(N^2)$.

If $\xi(\mathcal{T}_M^*(p)) \neq 0$ for some $p \in \mathbb{R}$, then Lemma 28 allows us to find some $N \in \mathfrak{R}(M, \mathcal{F} \cup \{\xi\}) \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{>0})$, and we will be done. Similarly, if $\xi_2(\mathcal{T}_M^*(p)) \neq 0$ for some $p \in \mathbb{R}$, we can first use Lemma 28 allows us to find some $M' \in \mathfrak{R}(M, \mathcal{F} \cup \{\xi_2\}) \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{>0})$. Then, since $F(M') \neq 0$ for all $F \in \mathcal{F}$, and $\xi((M')^2) \neq 0$, Corollary 33 allows us to once again find the required $N \in \mathfrak{R}(M'', \mathcal{F} \cup \{\xi\}) \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{>0})$.

On the other hand, let us assume that $\xi(\mathcal{T}_M^*(p)) = 0$, and $\xi_2(\mathcal{T}_M^*(p)) = 0$ for all $p \in \mathbb{R}$. We will now define $\zeta, \zeta_2 : \mathbb{R} \to \mathbb{R}$ such that $\zeta(p) = \xi(\mathcal{T}_M^*(p))$, and $\zeta_2(p) = \xi_2(\mathcal{T}_M^*(p))$ for all $p \in \mathbb{R}$. We note that

$$\begin{aligned} \zeta(p) &= \mathcal{T}_{M}^{*}(p)_{11} \cdot \mathcal{T}_{M}^{*}(p)_{44} - \mathcal{T}_{M}^{*}(p)_{22} \cdot \mathcal{T}_{M}^{*}(p)_{33} = p^{z_{11}+z_{44}-z_{22}-z_{33}}, \text{ and} \\ \zeta_{2}(p) &= (\mathcal{T}_{M}^{*}(p)^{2})_{11} \cdot (\mathcal{T}_{M}^{*}(p)^{2})_{44} - (\mathcal{T}_{M}^{*}(p)^{2})_{22} \cdot (\mathcal{T}_{M}^{*}(p)^{2})_{33} \\ &= \left(\sum_{i \in [4]} p^{2z_{1i}}\right) \cdot \left(\sum_{j \in [4]} p^{2z_{4j}}\right) - \left(\sum_{i \in [4]} p^{2z_{2i}}\right) \cdot \left(\sum_{j \in [4]} p^{2z_{3j}}\right) \\ &= \sum_{i,j \in [4]} \left(p^{2z_{1i}+2z_{4j}} - p^{2z_{2i}+2z_{3j}}\right). \end{aligned}$$

From our assumptions above, we know that $\zeta(p) = 0$, and $\zeta_2(p) = 0$ for all $p \in \mathbb{R}$. Therefore $\frac{d}{dp}(\zeta(p)) = 0$, and $\frac{d}{dp}(\zeta_2(p)) = 0$ for all $p \in \mathbb{R}$. Specifically, when evaluated at p = 1, we see that

$$\frac{d\zeta}{dp}(1) = z_{11} + z_{44} - z_{22} - z_{33} = 0.$$

This implies that $z_{11} + z_{44} = z_{22} + z_{33}$. Similarly,

$$\frac{d\zeta_2}{dp}(1) = \sum_{i,j \in [4]} (2z_{i1} + 2z_{4j} - 2z_{2i} - 2z_{3j})$$

= $(2z_{11} + 2z_{12} + 2z_{13} + 2z_{14}) + (2z_{14} + 2z_{24} + 2z_{34} + 2z_{44})$
 $- (2z_{12} + 2z_{22} + 2z_{23} + 2z_{24}) - (2z_{13} + 2z_{23} + 2z_{33} + 2z_{34})$
= $2z_{11} + 4z_{14} + 2z_{44} - 2z_{22} - 4z_{23} - 2z_{33} = 0.$

Therefore, $2(z_{11} + z_{44} - z_{22} - z_{33}) + 4(z_{14} - z_{23}) = 0$. Since we already saw that $z_{11} + z_{44} = z_{22} + z_{33}$, this implies that $z_{14} = z_{23}$.

We recall that by our choice of $m > \mathfrak{m}$, and construction of z_{ij} , $z_{14} = z_{23}$ implies that $(e_{141}, \ldots, e_{14d}) = (e_{231}, \ldots, e_{23d})$. But this means that $M_{14} = M_{23}$, which finishes the proof.

Lemma 67. Let $M \in Sym_4^{pd}(\mathbb{R}_{>0})$ be a diagonal distinct matrix. Let \mathcal{F} be a countable set of $Sym_4(\mathbb{R})$ -polynomials such that $F(M) \neq 0$ for all $F \in \mathcal{F}$. Let ξ be the $Sym_4(\mathbb{R})$ -polynomial such that $\xi : N \mapsto \phi_{(1,-1,-1,1)}(N_{11},\ldots,N_{44})$. Then there exists some diagonal distinct $N \in \mathfrak{R}(M, \mathcal{F} \cup \{\xi\}) \cap Sym_4^{pd}(\mathbb{R}_{>0})$, unless $M = A \otimes B$ for some $A, B \in Sym_2(\mathbb{R}_{>0})$.

Proof. We will assume that $M \neq A \otimes B$ for any $A, B \in \operatorname{Sym}_2(\mathbb{R}_{>0})$. From Lemma 65, we know that $\rho_{\text{tensor}}(M) \neq 0$. So we can add ρ_{diag} and ρ_{tensor} to \mathcal{F} assumed for M. Our goal will be to show that we can construct some diagonal distinct $N' \in \mathfrak{R}(M, \mathcal{F}) \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{>0})$ such that $N'_{11} \cdot N'_{44} \neq N'_{22} \cdot N'_{33}$ or $N'_{14} \neq N'_{23}$. Then, Lemma 66 would allow us to find the required $N \in$ $\mathfrak{R}(N', \mathcal{F} \cup \{\rho_{\text{diag}}, \xi\}) \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{>0})$.

We recall that since $S_M(0) = I$, there exists some $\delta > 0$ such that $|S_M(\theta)_{ij} - I_{ij}| < \frac{1}{3}$ for all $0 < \theta < \delta$ for all $i, j \in [4]$. We will now let

$$\mathcal{F}' = \{ F' : N \mapsto F(N^3) \mid F \in \mathcal{F} \}.$$

We note that $F'(\mathcal{S}_M(\frac{1}{3})) \neq 0$ for all $F' \in \mathcal{F}'$, and $\rho_{\text{diag}}(\mathcal{S}_M(1)) = \rho_{\text{diag}}(M) \neq 0$, and $\rho_{\text{tensor}}(\mathcal{S}_M(1)) = \rho_{\text{tensor}}(M) \neq 0$. We can therefore use Lemma 32 to find $M' = \mathcal{S}_M(\theta^*) \in \mathfrak{R}(M, \mathcal{F}' \cup \{\rho_{\text{diag}}, \rho_{\text{tensor}}\}) \cap \text{Sym}_4^{pd}(\mathbb{R}_{\neq 0})$ for some $0 < \theta^* < \delta$.

If $(M')_{14} \neq (M')_{23}$ or $M'_{11}M'_{44} \neq M'_{22}M'_{33}$, we will be done. So, we may assume otherwise. We will first assume that $(R_n(M'))_{14} - (R_n(M'))_{23} = 0$ for all $n \geq 1$. We note that

$$R_n(M')_{14} - R_n(M')_{23} = \sum_{a,b \in [4]} (M'_{ab})^n \cdot (M'_{1a}M'_{4b} - M'_{2a}M'_{3b}).$$

We may now let $X = \{(M')_{ab} : a, b \in [4]\}$, and for each $x \in X$, define

$$c_{14}(x) = \sum_{a,b \in [4]: (M')_{ab} = x} (M')_{1a}(M')_{4b}, \text{ and } c_{23}(x) = \sum_{a,b \in [4]: (M')_{ab} = x} (M')_{2a}(M')_{3b}.$$

So, we see that

$$R_n(M')_{14} - R_n(M')_{23} = \sum_{x \in X} x^n \cdot (c_{14}(x) - c_{23}(x)).$$

Since each $x \in X$ is distinct, and $|X| \leq O(1)$, this forms a full rank Vandermonde system of linear equations. Since we have assumed that $(R_n(M'))_{14} - (R_n(M'))_{23} = 0$ for all $n \geq 1$, this implies that $c_{14}(x) - c_{23}(x) = 0$ for all $x \in X$.

Now, by our choice of M', we know that all the diagonal terms M' of are distinct. Also any diagonal element of $M' = S_M(\theta^*)$ is greater than the absolute value of any off diagonal element, by our choice of $0 < \theta^* < \delta$. Therefore, $(M')_{ii} = (M')_{ab}$ implies that (a, b) = (i, i) for all $i \in [4]$. This implies that

$$c_{14}((M')_{ii}) - c_{23}((M')_{ii}) = (M')_{1i}(M')_{4i} - (M')_{2i}(M')_{3i} = 0$$

for all $i \in [4]$. We also already know that $(M')_{14} = (M')_{23}$, and $M'_{11}M'_{44} = M'_{22}M'_{33}$. But this implies that $\rho_{\texttt{tensor}}(M') = 0$, which is a contradiction to $M' \in \mathfrak{R}(M, \mathcal{F}' \cup \{\rho_{\texttt{diag}}, \rho_{\texttt{tensor}}\})$.

This means that there exists some $n^* \geq 1$, such that $(R_{n^*}(M'))_{14} - R_{n^*}(M'))_{23} \neq 0$. This implies that we can define the $\operatorname{Sym}_4(\mathbb{R})$ -polynomial $\xi_{n^*} : N \mapsto (R_{n^*}(N))_{14} - (R_{n^*}(N))_{23}$, and we see that $\xi_{n^*}(\mathcal{S}_M(\theta^*)) = \xi_{n^*}(M') \neq 0$. So, using $\mathcal{S}_M(\frac{1}{3})$, Corollary 33 allows us to first find some $M'' \in \mathfrak{R}(M, \mathcal{F}' \cup \{\rho_{\operatorname{diag}}^*, \xi_{n^*}\}) \cap \operatorname{Sym}_4^{\operatorname{pd}}(\mathbb{R}_{>0})$, where $\rho_{\operatorname{diag}}^*(N) = \rho_{\operatorname{diag}}(N^3)$. By construction, we see that $F(\mathcal{R}_{M''}(1)) \neq 0$ for all $F \in \mathcal{F}$, and $\rho_{\operatorname{diag}}(\mathcal{R}_{M''}(1)) \neq 0$, as $\mathcal{R}_{M''}(1) = (M'')^3$. Moreover, we see that $(\mathcal{R}_{M''}(n^*))_{14} - (\mathcal{R}_{M''}(n^*))_{23} \neq 0$. Therefore, Lemma 57 allows us to find some $N' \in$ $\mathfrak{R}(M'', \mathcal{F} \cup \{\rho_{\operatorname{diag}}\}) \cap \operatorname{Sym}_4^{\operatorname{pd}}(\mathbb{R}_{>0})$, such that $N'_{14} \neq N'_{23}$. Lemma 66 then allows us to find the required $N \in \mathfrak{R}(N', \mathcal{F} \cup \{\rho_{\operatorname{diag}}, \xi\}) \cap \operatorname{Sym}_4^{\operatorname{pd}}(\mathbb{R}_{>0})$.

We are now ready to state a hardness criterion that applies to all Pl-GH(M) where M is diagonal distinct.

Theorem 68. Let $M \in Sym_4^{pd}(\mathbb{R}_{>0})$ be a diagonal distinct matrix. Then Pl-GH(M) is #P-hard unless M is isomorphic to $A \otimes B$ for some $A, B \in Sym_2(\mathbb{R}_{>0})$.

Proof. Let us assume that M is not isomorphic to any $A \otimes B$ for any $A, B \in \text{Sym}_2(\mathbb{R})$. (Note that for $M \in \text{Sym}_4(\mathbb{R}_{>0})$, M is isomorphic to $A \otimes B$ for some $A, B \in \text{Sym}_2(\mathbb{R})$ is equivalent to being isomorphic to $A \otimes B$ for some $A, B \in \text{Sym}_2(\mathbb{R}_{>0})$.) So, Lemma 65 tells us that $\rho_{\texttt{tensor}}(M) \neq 0$. We will now construct

$$\mathcal{F} = \mathcal{F}_M \cup \{ \rho_{\texttt{diag}}, \rho_{\texttt{tensor}} \},$$

where \mathcal{F}_M is as defined in Eq. (15). We can see that $F(M) \neq 0$ for all $F \in \mathcal{F}$. We recall from Eq. (12) that

$$\Phi_{(1,-1,-1,1)}(N_{11},\ldots,N_{44}) = \prod_{\sigma \in S_4} \phi_{(1,-1,-1,1)}(N_{\sigma(1)\sigma(1)},\ldots,N_{\sigma(4)\sigma(4)}).$$

So, we will define the $\text{Sym}_4(\mathbb{R})$ -polynomials ξ_{σ} for $\sigma \in S_4$ as

$$\xi_{\sigma}(N) = \phi_{(1,-1,-1,1)}(N_{\sigma(1)\sigma(1)}, \dots, N_{\sigma(4)\sigma(4)}),$$

and the Sym₄(\mathbb{R})-polynomial ξ such that $\xi : N \mapsto \prod_{\sigma \in S_4} \xi_{\sigma}(N)$.

For each $\sigma \in S_4$, we may define $M^{\sigma} \in \text{Sym}_4^{\text{pd}}(\mathbb{R}_{>0})$ as in Lemma 65 to be such that $(M^{\sigma})_{ij} = M_{\sigma(i)\sigma(j)}$. We will also let $S_4 = \{\sigma_1, \ldots, \sigma_{24}\}$. Now, Lemma 67 implies that since $\rho_{\text{tensor}}(M^{\sigma_1}) \neq 0$,

we can find some $M_1 \in \mathfrak{R}(M, \mathcal{F} \cup \{\xi_{\sigma_1}\}) \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{>0})$. Since $F(M_1) \neq 0$ for all $F \in \mathcal{F}$, starting with M_1 instead of M, we can repeat this to find $M_2 \in \mathfrak{R}(M_1, \mathcal{F} \cup \{\xi_{\sigma_1}, \xi_{\sigma_2}\}) \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{>0})$.

After repeating this process for all $\sigma \in S_4$, we end up with some

$$M' = M_{24} \in \mathfrak{R}(M, \mathcal{F} \cup \{\xi\}) \cap \operatorname{Sym}_4^{\operatorname{pd}}(\mathbb{R}_{>0}).$$

Since $\Phi_{(1,-1,-1,1)}(M'_{11},\ldots,M'_{44}) \neq 0$, Theorem 34 allows us to find some $N \in \mathfrak{R}(M',\mathcal{F}_M \cup \{\rho_{\mathtt{diag}},\Psi_{(1,-1,-1,1)}\}) \cap \mathrm{Sym}_4^{\mathtt{pd}}(\mathbb{R}_{>0}).$

Let (μ_1, \ldots, μ_4) be the eigenvalues of N. Let \mathcal{L}_N be the lattice formed by these eigenvalues. By construction, we see that $(1, -1, -1, 1) \notin \overline{\mathcal{L}_N}$. So, from Lemma 64, we see that all $\mathbf{0} \neq \mathbf{x} \in \overline{\mathcal{L}_N}$ are confluent. Since N is diagonal distinct, Corollary 60 then proves that $\mathsf{Pl-GH}(N)$ is $\#\mathsf{P}$ -hard. Since $\mathsf{Pl-GH}(M)$, we see that $\mathsf{Pl-GH}(M)$ is also $\#\mathsf{P}$ -hard.

10 Hardness for non-Diagonal Distinct 4×4 matrices

In this section, we will deal with the matrices $M \in \text{Sym}_4^{\text{pd}}(\mathbb{R}_{>0})$ that are not diagonal distinct. We will first show that in this case, there are only finitely many forms that the matrix M must take. Then, we will show that for each such form, Pl-GH(M) is #P-hard, unless M is isomorphic to $A \otimes B$ for some $A, B \in \text{Sym}_2(\mathbb{R}_{\neq 0})$.

Lemma 69. Let $M \in Sym_4^{pd}(\mathbb{R}_{>0})$ such that $M_{11} = M_{22}$, $M_{22} \neq M_{33} \neq M_{44}$ are pairwise distinct, and $\rho_{tensor}(M) \neq 0$. Unless M is of one of the two forms below (Forms (I) or (II)), Pl-GH(M) is #P-hard.

Proof. We apply Lemma 61 to M. Suppose rows 1 and 2 are not order identical. The pairwise distinctness of $\{M_{22}, M_{33}, M_{44}\}$ and of $\{M_{11}, M_{33}, M_{44}\}$ are conditions expressible as nonvanishing of $\operatorname{Sym}_q(\mathbb{R})$ -polynomials and thus maintained by Lemma 61, and hence its conclusion $N_{11} \neq N_{22}$ in addition to the above pairwise distinctness conditions gives diagonal distinctness. Therefore, from Lemma 61, unless rows 1 and 2 are order identical, we get some diagonal distinct $N \in \mathfrak{R}(M, \mathcal{F}_M \cup \{\rho_{\text{tensor}}\}) \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{>0})$. Theorem 68 then proves that $\operatorname{Pl-GH}(N)$, and consequently, $\operatorname{Pl-GH}(M)$ are #P-hard.

So, we may assume rows 1 and 2 are order identical. There exists some $\sigma \in S_4$ such that $M_{1i} = M_{2\sigma(i)}$ for all $i \in [4]$. We are given $M_{11} = M_{22}$. We also have $M_{12} = M_{21}$ by M being symmetric. From the identical multisets $\{M_{11}, M_{12}, M_{13}, M_{14}\}$ and $\{M_{21}, M_{22}, M_{23}, M_{24}\}$ if we

remove the element pairs $M_{11} = M_{22}$ and $M_{12} = M_{21}$, we still have an equal multiset $\{M_{13}, M_{14}\} = \{M_{23}, M_{24}\}$. So, we may assume that $(\sigma(1) = 2, \sigma(2) = 1, \text{ and}) \sigma(3) = 3 \text{ or } \sigma(3) = 4$. Now, if $\sigma(3) = 3$, that means that $\sigma(4) = 4$, which forces $M_{13} = M_{23}$, and $M_{14} = M_{24}$. In that case, we see that M is of Form (I) above. On the other hand, if $\sigma(3) = 4$, that means $\sigma(4) = 3$, which forces M to be of Form (II) above.

Remark. When we say that a matrix is of Form (I), we do not require that $M_{13} \neq M_{14}$ for example. In general, for all the Forms that we will describe, we allow the possibility that some of the values denoted by distinct symbols may be equal to each other. All we require is that entries denoted by the same symbol are equal. So, it may be possible that a matrix is of more than one Form.

Lemma 70. Let $M \in Sym_4^{pd}(\mathbb{R}_{>0})$ such that $M_{11} = M_{22} \neq M_{33} = M_{44}$, and $\rho_{tensor}(M) \neq 0$. Unless M is isomorphic to a matrix of the Form (I) or Form (II), or is of the form below (Form (III)), Pl-GH(M) is #P-hard.

(M_{11})	M_{12}	M_{13}	M_{14}		(a)	x	y	z
M_{12}	M_{11}	M_{14}	M_{13}	=	x	a	z	y
M_{13}	M_{14}	M_{33}	M_{34}		y	z	b	t
M_{14}	M_{13}	M_{34}	M_{33}		$\langle z \rangle$	y	t	b)

Form	(III)
Form	(111)

Proof. Let us first assume that it is not the case that rows 1 and 2 are order identical, and that rows 3 and 4 are order identical. By symmetry, we may assume that rows 3 and 4 are not order identical. We will also assume that M is not isomorphic to a matrix of Form (I) or of Form (II). So, if we define $\zeta : N \mapsto (N_{11} - N_{33})(N_{11} - N_{44})(N_{22} - N_{33})(N_{22} - N_{44}), \zeta_1 : N \mapsto (N_{13} - N_{23})^2 + (N_{14} - N_{24})^2$, and $\zeta_2 : N \mapsto (N_{13} - N_{24})^2 + (N_{14} - N_{23})^2$, we see that $\zeta(M) \neq 0, \zeta_1(M) \neq 0$, and $\zeta_2(M) \neq 0$. From Lemma 61, we know that since rows 3 and 4 are not order identical, we can construct some $N \in \Re(M, \mathcal{F}_M \cup \{\rho_{\text{tensor}}, \zeta, \zeta_1, \zeta_2\}) \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{>0})$ such that $N_{22} \neq N_{33} \neq N_{44}$, and $N_{11} \neq N_{33} \neq N_{44}$. If $N_{11} \neq N_{22}$, then N is diagonal distinct, and Theorem 68 implies that Pl-GH(N) \leq Pl-GH(M) is #P-hard. Moreover, by our construction of N, we ensured that even if $N_{11} = N_{22}$, N is not of Form (I). So, we can see from Lemma 69 that Pl-GH(N) \leq Pl-GH(M) must be #P-hard.

So, we may assume that rows 1 and 2 are order identical, and that rows 3 and 4 are also order identical. So, there must exist some $\sigma_1, \sigma_2 \in S_4$ such that $M_{1i} = M_{2\sigma_1(i)}$, and $M_{3i} = M_{4\sigma_2(i)}$ for all $i \in [4]$. We already know that $M_{11} = M_{22}$, and that $M_{12} = M_{21}$. So, we may assume that $\sigma_1(1) = 2$, and $\sigma_1(2) = 1$. Similarly, we may assume that $\sigma_2(3) = 4$, and $\sigma_2(4) = 3$. Now, if $\sigma_1(3) = 3$, that means that $\sigma_1(4) = 4$, which forces $M_{13} = M_{23}$, and $M_{14} = M_{24}$. But then irrespective of whether $\sigma_2(1) = 1$ or $\sigma_2(1) = 2$, we see that $M_{31} = M_{4\sigma_2(1)}$ which implies that $M_{13} = M_{14} = M_{23} = M_{24}$. In that case, we see that M is of Form (III) above (with $M_{13} = M_{14}$). On the other hand, if $\sigma_1(3) = 4$, that means $\sigma_1(4) = 3$, which forces M to be of Form (III) above.

Lemma 71. Let $M \in Sym_4^{pd}(\mathbb{R}_{>0})$ such that $M_{11} = M_{22} = M_{33} \neq M_{44}$, and $\rho_{tensor}(M) \neq 0$. Unless M is isomorphic to a matrix of Form (I) or Form (II), or is of the two forms below, Pl-GH(M) is #P-hard.

Proof. We will first suppose rows 1, 2 and 3 are not all order identical to each other. By the symmetry of our assumptions so far among $\{1, 2, 3\}$ in this lemma, without loss of generality suppose rows 2 and 3 are not order identical. We can also assume that M is not isomorphic to a matrix of Form (I) or Form (II). We can now define $\zeta : N \mapsto (N_{11} - N_{44})(N_{22} - N_{44})(N_{33} - N_{44})$ $\zeta_1 : N \mapsto (N_{13} - N_{23})^2 + (N_{14} - N_{24})^2, \ \zeta_2 : N \mapsto (N_{13} - N_{24})^2 + (N_{14} - N_{23})^2, \ \zeta_3 : N \mapsto (N_{13} - N_{24})^2 + (N_{14} - N_{23})^2, \ \zeta_3 : N \mapsto (N_{13} - N_{24})^2 + (N_{14} - N_{23})^2, \ \zeta_3 : N \mapsto (N_{13} - N_{24})^2 + (N_{14} - N_{23})^2, \ \zeta_3 : N \mapsto (N_{13} - N_{24})^2 + (N_{14} - N_{23})^2, \ \zeta_3 : N \mapsto (N_{13} - N_{24})^2 + (N_{14} - N_{23})^2, \ \zeta_3 : N \mapsto (N_{13} - N_{24})^2 + (N_{14} - N_{23})^2, \ \zeta_3 : N \mapsto (N_{13} - N_{24})^2 + (N_{14} - N_{23})^2, \ \zeta_3 : N \mapsto (N_{13} - N_{24})^2 + (N_{14} - N_{23})^2, \ \zeta_3 : N \mapsto (N_{13} - N_{24})^2 + (N_{14} - N_{23})^2, \ \zeta_3 : N \mapsto (N_{13} - N_{24})^2 + (N_{14} - N_{23})^2, \ \zeta_3 : N \mapsto (N_{13} - N_{24})^2 + (N_{14} - N_{23})^2 + (N_{14} - N_{24})^2 + (N_{14} -$ $(N_{12} - N_{23})^2 + (N_{14} - N_{34})^2$, and $\zeta_4 : N \mapsto (N_{12} - N_{34})^2 + (N_{14} - N_{23})^2$. (Here, given $M_{11} = M_{22}$, being not in Form (I) or Form (II) implies that $\zeta_1(M) \neq 0$ and $\zeta_2(M) \neq 0$. Note that ζ_3 and ζ_4 are obtained from ζ_1 and ζ_2 , respectively, by exchanging rows and columns indexed by 2 and 3.) We see that $\zeta(M) \neq 0$, and $\zeta_i(M) \neq 0$ for all $i \in [4]$. By Lemma 61, we can obtain some $N \in \mathfrak{R}(M, \mathcal{F}_M \cup \{\rho_{\texttt{tensor}}, \zeta, \zeta_1, \zeta_2, \zeta_3, \zeta_4\}) \cap \operatorname{Sym}_4^{\mathsf{pd}}(\mathbb{R}_{>0}) \text{ satisfying } N_{22} \neq N_{33}. \text{ If } N_{11} \neq N_{22} \neq N_{33} \neq N_{33} = N_{33} + N_{3$ N_{44} are all pairwise distinct, then Theorem 68 implies that $Pl-GH(N) \leq Pl-GH(M)$ is #P-hard. If however, $N_{11} = N_{22}$, since $\zeta(N) \neq 0$, it must be the case that $N_{11} = N_{22}$, but $N_{22} \neq N_{33} \neq N_{44}$ are pairwise distinct. So, Lemma 69 implies that $Pl-GH(N) \leq Pl-GH(M)$ is #P-hard unless N is of Form (I) or (II). But, by construction, $\zeta_1(N) \neq 0$, and $\zeta_2(N) \neq 0$. So, N is not of Form (I) or (II), which proves that $Pl-GH(N) \leq Pl-GH(M)$ is #P-hard. Similarly, if $N_{11} = N_{33}$, then since $\zeta(N) \neq 0$, we see that $N_{22} \neq N_{33} \neq N_{44}$ are pairwise distinct. Moreover, since $\zeta_3(N) \neq 0$, and $\zeta_4(N) \neq 0$, we see that if we switch rows and columns $2 \leftrightarrow 3$ in N, we have a matrix N' that is isomorphic to N that satisfies the condition that $(N')_{11} = (N')_{22}$, but $(N')_{22} \neq (N')_{33} \neq (N')_{44}$ are pairwise distinct, and the condition that $\zeta_3(N) \neq 0$ and $\zeta_4(N) \neq 0$ translates to N' is not of Form (I) or (II). So, Lemma 69 implies that $Pl-GH(N') \equiv Pl-GH(N) \leq Pl-GH(M)$ is #P-hard.

So, rows 1, 2 and 3 are all order identical to each other, and there exist some $\sigma_1, \sigma_2 \in S_4$ such that $M_{1i} = M_{2\sigma_1(i)}$, and $M_{1i} = M_{3\sigma_2(i)}$ for all $i \in [4]$. We already know that $M_{11} = M_{22}$, and that $M_{12} = M_{21}$. So, we may assume that $\sigma_1(1) = 2$, and $\sigma_1(2) = 1$. Similarly, we may assume that $\sigma_2(1) = 3$, and $\sigma_2(3) = 1$. So far we have the following

$$\begin{pmatrix} a & x & y & * \\ x & a & * & * \\ y & * & a & * \\ * & * & * & * \end{pmatrix}$$

There are two possibilities: $\sigma_1(3) = 3$ or $\sigma_1(3) = 4$. Suppose $\sigma_1(3) = 3$, then $\sigma_1(4) = 4$, which

forces $M_{13} = M_{23}$, and $M_{14} = M_{24}$, and we have the following

$$\begin{pmatrix} a & x & y & z \\ x & a & y & z \\ y & y & a & * \\ z & z & * & * \end{pmatrix}$$

Since rows 1 and 3 are order identical, the y entry at M_{31} is either y = x or y = z. The two cases are listed in Form (IV) and Form (V) (with y = z) respectively.

Now we may assume $\sigma_1(3) = 4$. Then we have the following

$$\begin{pmatrix} a & x & y & z \\ x & a & z & y \\ y & z & a & * \\ z & y & * & * \end{pmatrix}$$

Again since row 1 and 3 are order identical, the M_{34} entry must be x, this gives Form (V).

Lemma 72. Let $M \in Sym_4^{pd}(\mathbb{R}_{>0})$ such that $M_{11} = M_{22} = M_{33} = M_{44}$, and $\rho_{tensor}(M) \neq 0$. Unless M is isomorphic to a matrix of Forms (I) - (V), or is of the form below, Pl-GH(M) is #P-hard.

$$\begin{pmatrix} M_{11} & M_{12} & M_{13} & M_{14} \\ M_{12} & M_{11} & M_{14} & M_{13} \\ M_{13} & M_{14} & M_{11} & M_{12} \\ M_{14} & M_{13} & M_{12} & M_{11} \end{pmatrix} = \begin{pmatrix} a & x & y & z \\ x & a & z & y \\ y & z & a & x \\ z & y & x & a \end{pmatrix}$$

Form	(VI)
------	------

Proof. We first suppose rows 1, 2, 3 and 4 are not all order identical to each other. By symmetry, without loss of generality suppose rows 3 and 4 are not order identical. We may also assume that M is not of Forms (I) - (V). We can now define the $\text{Sym}_4(\mathbb{R})$ -polynomial ζ_1 such that $\zeta_1(N) = 0$ if and only if N is isomorphic to a matrix of Form (I). Similarly, we can define ζ_2 such that $\zeta_2(N) = 0$ if and only if N is isomorphic to a matrix of Form (II). We can also define $\zeta_3, \zeta_4, \zeta_5$ similarly for Forms (III) - (V). By our assumption, $\zeta_i(M) \neq 0$ for all $i \in [5]$.

Since rows 3 and 4 are not order identical, by Lemma 61, we can obtain some $N \in \mathfrak{R}(M, \mathcal{F}_M \cup \{\rho_{\mathsf{tensor}}, \zeta_1, \ldots, \zeta_5\}) \cap \operatorname{Sym}_4^{\mathsf{pd}}(\mathbb{R}_{>0})$ satisfying $N_{33} \neq N_{44}$. By our construction, we also know that N is not isomorphic to any matrix of Forms (I) - (V). Now, the set of diagonal elements of N (after removal of duplicates) has cardinality 2 or 3 or 4. The cardinality 4 case is diagonal distinct, and then Theorem 68 allows us to prove that P1-GH(M) is #P-hard. The cases of cardinality 2 or 3 fall into the cases of Lemma 69, Lemma 70 or Lemma 71, and in either case, we see that P1-GH(N) \leq P1-GH(M) is #P-hard, since N is not isomorphic to any matrix of Form (I) - (V).

So suppose rows 1, 2, 3 and 4 are all order identical to each other. We have the following setting

$$\begin{pmatrix} a & x & y & z \\ x & a & * & * \\ y & * & a & * \\ z & * & * & a \end{pmatrix}$$

As in the proof of Lemma 71, there are two possibilities: $\sigma_1(3) = 3$ or $\sigma_1(3) = 4$. Suppose $\sigma_1(3) = 3$, then $\sigma_1(4) = 4$, which forces $M_{13} = M_{23}$, and $M_{14} = M_{24}$, and we have the following

$$\begin{pmatrix} a & x & y & z \\ x & a & y & z \\ y & y & a & * \\ z & z & * & a \end{pmatrix}$$

Then y from the 3rd row must match either x or z. If y = x, then the 3rd row is (y, y, a, z) and then the 4th row is (z, z, z, a), and thus x = y = z, and we have Form (VI) (with x = y = z). If y = z then we have

$$\begin{pmatrix} a & x & y & y \\ x & a & y & y \\ y & y & a & * \\ y & y & * & a \end{pmatrix}$$

Then $M_{34} = x$ and we have Form (VI) (with y = z). Finally if $\sigma_1(3) = 4$, we also have Form (VI).

Now that we have listed all the different Forms that the matrices must take, we will show one by one that for each of the Forms, matrices M belonging to that Form must either be isomorphic to some $A \otimes B$, or that Pl-GH(M) is #P-hard. We will first show that Form (II) and Form (V) can actually be reduced to one of the other forms.

Lemma 73. Let $M \in Sym_4^{pd}(\mathbb{R}_{>0})$ be of Form (II), such that $M_{11} = M_{22}$ and $M_{22} \neq M_{33} \neq M_{44}$ are pairwise distinct, and $\rho_{tensor}(M) \neq 0$. Then, either Pl-GH(M) is #P-hard, or M is also isomorphic to a matrix of Form (I).

Proof. We first let $\zeta : N \mapsto (N_{11} - N_{33})(N_{11} - N_{44})(N_{22} - N_{33})(N_{22} - N_{44})(N_{33} - N_{44})$. We note that $\zeta(M) \neq 0$ by our choice of M. Moreover, we note that if $\zeta(N) \neq 0$, then N cannot be isomorphic to any matrix of Form (III), (IV), (V), or (VI), as there are at least three distinct diagonal elements. We have already seen that there exists some $\text{Sym}_4(\mathbb{R})$ -polynomial ζ_1 such that $\zeta_1(N) = 0$ if and only if N is isomorphic to a matrix of Form (I). Similarly, there exist $\text{Sym}_4(\mathbb{R})$ polynomials ζ_2, \ldots, ζ_6 for Forms (II) - (VI), respectively. By our assumption about M, we have $\zeta_i(M) \neq 0$ for all $i \in \{3, 4, 5, 6\}$. We are given that M has Form (II), thus $\zeta_2(M) = 0$. We will assume $\zeta_1(M) \neq 0$, and show that P1-GH(M) is #P-hard.

We will now define $\xi : N \mapsto (N_{14} - N_{23})^2 + (N_{13} - N_{24})^2$. Since *M* has Form (II), we have $\xi(M) = 0$. Moreover, for any N, $\zeta(N) \neq 0$ and $\xi(N) \neq 0$ implies that *N* is not isomorphic

to any matrix of Form (II). Indeed, to be isomorphic to a matrix of Form (II) and having the pairwise distinctness of $\{M_{11}, M_{33}, M_{44}\}$ and of $\{M_{22}, M_{33}, M_{44}\}$ given by $\zeta(N) \neq 0$ we only need to consider N under possibly the permutations $1 \leftrightarrow 2$ and $3 \leftrightarrow 4$; but these permutations do not change $\xi(N) \neq 0$. In other words, $[\zeta(N) \neq 0] \land [\xi(N) \neq 0]$ implies that $\zeta_2(N) \neq 0$. So, if there exists some $\theta \in \mathbb{R}$ such that $\xi(\mathcal{S}_M(\theta)) \neq 0$, we will be able to immediately find $N \in \mathfrak{R}(M, \{\rho_{\texttt{tensor}}, \zeta, \zeta_1, \xi\}) \cap \operatorname{Sym}_4^{\mathsf{pd}}(\mathbb{R}_{>0}) \subseteq \mathfrak{R}(M, \{\rho_{\texttt{tensor}}, \zeta, \zeta_1, \zeta_2\}) \cap \operatorname{Sym}_4^{\mathsf{pd}}(\mathbb{R}_{>0})$, by Corollary 33. Since $\zeta(N) \neq 0$, we see that N must either be diagonal distinct, or $N_{11} = N_{22} \neq N_{33} \neq N_{44}$ pairwise distinct. But since $\zeta_i(N) \neq 0$ for $i \in \{1, 2\}$, we see from Theorem 68, and Lemma 69, that $\mathsf{Pl-GH}(N) \leq \mathsf{Pl-GH}(M)$ is #P-hard. So, we may assume $\xi(\mathcal{S}_M(\theta)) = 0$ for all $\theta \in \mathbb{R}$.

We will now let $\mathcal{F}' = \{F' : N \mapsto F(N^3) \mid F \in \{\rho_{\texttt{tensor}}, \zeta, \zeta_1\}\}$. Since $\mathcal{S}_M(0) = I$, we know that there exists some $\delta > 0$ such that $|\mathcal{S}_M(\theta)_{ij} - I_{ij}| < \frac{1}{3}$ for all $i, j \in [4]$, for all $0 < \theta < \delta$. We can now use Lemma 32 to find some $M' = \mathcal{S}_M(\theta^*) \in \mathfrak{R}(M, \mathcal{F}' \cup \{\rho_{\texttt{tensor}}, \zeta, \zeta_1\}) \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{\neq 0})$ such that $0 < \theta^* < \delta$. We note that by our assumption, $\xi(M') = 0$.

Now, since $\zeta(M') \neq 0$, we see that $(M')_{11} \neq (M')_{33} \neq (M')_{44}$ and $(M')_{22} \neq (M')_{33} \neq (M')_{44}$ are both pairwise distinct. If $(M')_{11} \neq (M')_{22}$, M' would be diagonal distinct, and Pl-GH $(M') \leq$ Pl-GH(M) is #P-hard, due to Theorem 68. On the other hand, suppose $(M')_{11} = (M')_{22}$. If M'is not of Form (II), we see that since $\zeta_1(M') \neq 0$, Pl-GH $(M') \leq$ Pl-GH(M) is #P-hard, due to Lemma 69. So, we may assume that M' is of Form (II), such that $(M')_{11} = (M')_{22} \neq (M')_{33} \neq$ $(M')_{44}$ are pairwise distinct, and $\xi(M') = 0$.

We will now consider $\mathcal{R}_{M'}(n)$ for all $n \geq 1$. Let us first assume that $\xi(R_n(M')) = 0$ for all $n \geq 1$. We note that

$$(R_n(M'))_{14} - (R_n(M'))_{23} = \sum_{a,b \in [4]} (M'_{ab})^n \cdot (M'_{1a}M'_{4b} - M'_{2a}M'_{3b}).$$

We will let $X = \{M'_{ab} : a, b \in [4]\}$, and define $c_{ij}(x) = \sum_{a,b \in [4]: M'_{ab} = x} (M'_{ia}M'_{jb})$ for all $i, j \in [4]$. So, we see that

$$(R_n(M'))_{14} - (R_n(M'))_{23} = \sum_{x \in X} x^n \cdot (c_{14}(x) - c_{23}(x)).$$

Since each $x \in X$ is distinct, the equations $\xi(R_n(M')) = 0$ form a full rank Vandermonde system of size O(1). This implies that $c_{14}(x) - c_{23}(x) = 0$ for all $x \in X$. But since $\zeta(M') \neq 0$, and $0 < \theta^* < \delta$ in the definition of M', we ensured that $M'_{44} = M'_{ab}$ if and only if (a, b) = (4, 4), and we have also ensured that $M'_{33} = M'_{ab}$ if and only if (a, b) = (3, 3). Now, we see that

$$c_{14}(M'_{44}) - c_{23}(M'_{44}) = M'_{14}M'_{44} - M'_{24}M'_{34}.$$
$$c_{14}(M'_{33}) - c_{23}(M'_{33}) = M'_{13}M'_{34} - M'_{23}M'_{33}.$$

Since $\xi(M') = \xi(\mathcal{S}_M(\theta^*)) = 0$, we also know that $M'_{24} = M'_{13}$ and $M'_{23} = M'_{14}$. So, we also have

$$c_{14}(M'_{44}) - c_{23}(M'_{44}) = M'_{14}M'_{44} - M'_{13}M'_{34}.$$
$$c_{14}(M'_{33}) - c_{23}(M'_{33}) = M'_{13}M'_{34} - M'_{14}M'_{33}.$$

Then, it follows that

$$M_{14}'M_{44}' = M_{13}'M_{34}' = M_{14}'M_{33}'$$

which implies that $M'_{33} = M'_{44}$, since $M'_{14} \neq 0$. But by our construction of M', we have $\zeta(M') \neq 0$. This is a contradiction.

Therefore, our assumption that $\xi(R_n(M')) = 0$ for all $n \ge 1$ must be false. Let $\xi \circ R_n$ denote the composition of ξ and R_n . There exists some $n^* \ge 1$ such that $(\xi \circ R_{n^*})(M') = (\xi \circ R_{n^*})(\mathcal{S}_M(\theta^*)) \ne 0$. So, Corollary 33 allows us to find some $M'' \in \mathfrak{R}(M, \mathcal{F}' \cup \{(\xi \circ R_{n^*})\}) \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{>0})$. Since $F(\mathcal{R}_{M''}(1)) = F((M'')^3) \ne 0$ for all $F \in \{\rho_{\texttt{tensor}}, \zeta, \zeta_1\}$, and $\xi(\mathcal{R}_{M''}(n^*)) \ne 0$, Lemma 57 allows us to find some $N \in \mathfrak{R}(M, \{\rho_{\texttt{tensor}}, \zeta, \zeta_1, \xi\}) \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{>0}) \subseteq \mathfrak{R}(M, \{\rho_{\texttt{tensor}}, \zeta, \zeta_1, \zeta_2\}) \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{>0})$.

Since $\zeta(N) \neq 0$, we see that N must either be diagonal distinct, or $N_{11} = N_{22} \neq N_{33} \neq N_{44}$ pairwise distinct. If N is diagonal distinct, Theorem 68 implies that $Pl-GH(N) \leq Pl-GH(M)$ is #P-hard. Otherwise, since $\zeta_i(N) \neq 0$ for all $i \in \{1, 2\}$, Lemma 69 implies that $Pl-GH(N) \leq Pl-GH(M)$ is #P-hard.

Lemma 74. Let $M \in Sym_4^{pd}(\mathbb{R}_{>0})$ be of Form (V), such that $M_{11} = M_{22} = M_{33} \neq M_{44}$, and $\rho_{tensor}(M) \neq 0$. Then, either Pl-GH(M) is #P-hard, or M is also isomorphic to a matrix of Form (I), or (IV).

Proof. We will first let $\zeta : N \mapsto (N_{11} - N_{44})(N_{22} - N_{44})(N_{33} - N_{44})$. By our choice of M, we see that $\zeta(M) \neq 0$. Moreover, we note that $\zeta(N) \neq 0$ implies that N cannot be isomorphic to any matrix of Form (III) or (VI), since in these two Forms, every diagonal element coincides with another diagonal element. We have already seen that there exist $\text{Sym}_4(\mathbb{R})$ -polynomials ζ_1, \ldots, ζ_6 such that $\zeta_i(N) = 0$, for $i \in [6]$, if and only if N is isomorphic to a matrix of the Form (I) - (VI), respectively. From our choice of M, we note that since $M_{11} = M_{22} = M_{33} \neq M_{44}$, it immediately follows that $\zeta_3(M) \neq 0$, and $\zeta_6(M) \neq 0$. We are given that M is in Form (V) and hence $\zeta_5(M) = 0$. We will assume $\zeta_i(M) \neq 0$ for all $i \in \{1, 4\}$, and show that P1-GH(M) is #P-hard.

We will now define $\xi : N \mapsto (N_{14} - N_{23})^2 + (N_{13} - N_{24})^2 + (N_{12} - N_{34})^2$. We see that since M is of Form (V), $\xi(M) = 0$. Moreover, if $\zeta(N) \neq 0$, and $\xi(N) \neq 0$, that implies that N is not isomorphic to any matrix of Form (V). Indeed, since $\zeta(N) \neq 0$, N_{44} is distinct from the other diagonal elements, and so the only permutations that could be isomorphisms of N with a matrix of Form (V) must fix 4. However the group S_3 of permutations on $\{1, 2, 3\}$ is generated by $1 \leftrightarrow 2$ and $1 \leftrightarrow 3$, both of which keep $\xi(N)$ invariant. In other words, $[\zeta(N) \neq 0] \wedge [\xi(N) \neq 0]$ implies that $\zeta_5(N) \neq 0$. So, if there exists some $\theta \in \mathbb{R}$ such that $\xi(\mathcal{S}_M(\theta)) \neq 0$, we will be able to immediately find $N \in \mathfrak{R}(M, \{\rho_{\texttt{tensor}}, \zeta, \zeta_1, \zeta_4, \xi\}) \cap \operatorname{Sym}_4^{\mathsf{pd}}(\mathbb{R}_{>0}) \subseteq \mathfrak{R}(M, \{\rho_{\texttt{tensor}}, \zeta, \zeta_1, \zeta_4, \zeta_5\}) \cap \operatorname{Sym}_4^{\mathsf{pd}}(\mathbb{R}_{>0})$, by Corollary 33. Since $\zeta(N) \neq 0$, we see that $N_{11}, N_{22}, N_{33} \neq N_{44}$. If N is diagonal distinct, then Theorem 68 implies that $Pl-GH(N) \leq Pl-GH(M)$ is #P-hard. Otherwise, if N_{11}, N_{22}, N_{33} are not all identical, then from Lemma 69, we see that unless N is isomorphic to a matrix in Form (I) or (II), $\text{Pl-GH}(N) \leq \text{Pl-GH}(M)$ is #P-hard. But, we know that $\zeta_1(N) \neq 0$. So, N cannot be isomorphic to a matrix in Form (I). Moreover, from Lemma 73, we know that even if N is isomorphic to a matrix in Form (II), since it is not isomorphic to any matrix of Form (I), $Pl-GH(N) \leq Pl-GH(M)$ is #P-hard. This leaves only one possibility that $N_{11} = N_{22} = N_{33} \neq N_{44}$. Again, Lemma 71 implies that unless N is isomorphic to a matrix in Form (I), (II), (IV), or (V), $Pl-GH(N) \leq Pl-GH(M)$ is #P-hard.

But, by our construction of N, $\zeta_i(N) \neq 0$ for all $i \in \{1, 4, 5\}$. Similarly, even if $\zeta_2(N) = 0$, since $\zeta_1(N) \neq 0$, Lemma 73 implies that Pl-GH $(N) \leq$ Pl-GH(M) is #P-hard. Therefore, we see that when $\xi(\mathcal{S}_M(\theta)) \neq 0$ for some $\theta \in \mathbb{R}$, Pl-GH(M) is #P-hard. So, we may assume $\xi(\mathcal{S}_M(\theta)) = 0$ for all $\theta \in \mathbb{R}$.

We will now let $\mathcal{F}' = \{F' : N \mapsto F(N^3) \mid F \in \{\rho_{\texttt{tensor}}, \zeta, \zeta_1, \zeta_4\}\}$. Since $\mathcal{S}_M(0) = I$, we know that there exists some $\delta > 0$ such that $|\mathcal{S}_M(\theta)_{ij} - I_{ij}| < \frac{1}{3}$ for all $i, j \in [4]$, for all $0 < \theta < \delta$. We can now use Lemma 32 to find some $M' = \mathcal{S}_M(\theta^*) \in \mathfrak{R}(M, \mathcal{F}' \cup \{\rho_{\texttt{tensor}}, \zeta, \zeta_1, \zeta_4\}) \cap \text{Sym}_4^{pd}(\mathbb{R}_{\neq 0})$ such that $0 < \theta^* < \delta$. We note that by our assumption, $\xi(M') = 0$.

Now, since $\zeta(M') \neq 0$, we see that $(M')_{44}$ is distinct from each of $(M')_{11}, (M')_{22}, (M')_{33}$. If $(M')_{11} \neq (M')_{22} \neq (M')_{33}$ are pairwise distinct, M' would be diagonal distinct, and Pl-GH $(M') \leq$ Pl-GH(M) is #P-hard, due to Theorem 68. On the other hand, if the set $\{(M')_{11}, (M')_{22}, (M')_{33}\}$ has cardinality 2, we know from Lemma 69 that Pl-GH $(M') \leq$ Pl-GH(M) is #P-hard, unless M' is isomorphic to a matrix in Form (I) or (II). But since $\zeta_1(M') \neq 0$, we know that M' is not isomorphic to a matrix in Form (I). Moreover, from Lemma 73, we also know that even if $\zeta_2(M') = 0$, since $\zeta_1(M') \neq 0$, Pl-GH $(M') \leq$ Pl-GH(M') is #P-hard, unless M' is isomorphic to a matrix in Form Lemma 71, we see that Pl-GH $(M') \leq$ Pl-GH(M) is #P-hard, unless M' is isomorphic to a matrix in Form (I), (II), (IV), or (V). Since $\zeta_i(M') \neq 0$ for $i \in \{1,4\}$, we see that Forms (I), or (IV) are ruled out for M'. Moreover, since $\zeta_1(M') \neq 0$, we see from Lemma 73 that even if $\zeta_2(M') = 0$, Pl-GH $(M') \leq$ Pl-GH(M) is #P-hard. So, we may assume that M' is of Form (V), such that $(M')_{11} = (M')_{22} = (M')_{33} \neq (M')_{33} \in (M')_{44}$, and $\xi(M') = 0$.

We will now consider $\mathcal{R}_{M'}(n)$ for all $n \geq 1$. Let us first assume that $\xi(R_n(M')) = 0$ for all $n \geq 1$. We note that

$$(R_n(M'))_{14} - (R_n(M'))_{23} = \sum_{a,b \in [4]} (M'_{ab})^n \cdot (M'_{1a}M'_{4b} - M'_{2a}M'_{3b}).$$

We will let $X = \{M'_{ab} : a, b \in [4]\}$, and define $c_{ij}(x) = \sum_{a,b \in [4]: M'_{ab} = x} (M'_{ia}M'_{jb})$ for all $i, j \in [4]$. So, we see that

$$(R_n(M'))_{14} - (R_n(M'))_{23} = \sum_{x \in X} x^n \cdot (c_{14}(x) - c_{23}(x)).$$

Since each $x \in X$ is distinct, the equations $\xi(R_n(M')) = 0$ form a full rank Vandermonde system of size O(1). This implies that $c_{14}(x) - c_{23}(x) = 0$ for all $x \in X$. But since $\zeta(M') \neq 0$, and $0 < \theta^* < \delta$ in the definition of M', we ensured that $M'_{44} = M'_{ab}$ if and only if (a, b) = (4, 4). Moreover, from our construction of M', we have also ensured that $\{(a, b) : M'_{ab} = M'_{33}\} = \{(1, 1), (2, 2), (3, 3)\}$. Now, we see that

$$c_{14}(M'_{44}) - c_{23}(M'_{44}) = M'_{14}M'_{44} - M'_{24}M'_{34}$$

Since $\xi(M') = \xi(\mathcal{S}_M(\theta^*)) = 0$, we also know that $M'_{24} = M'_{13}$, $M'_{23} = M'_{14}$, and $M'_{12} = M'_{34}$. So, we also have

$$c_{14}(M'_{44}) - c_{23}(M'_{44}) = M'_{14}M'_{44} - M'_{13}M'_{34}$$

Moreover, we also see that

$$c_{14}(M'_{33}) - c_{23}(M'_{33}) = M'_{11}M'_{14} + M'_{12}M'_{24} + M'_{13}M'_{34} - M'_{12}M'_{13} - M'_{22}M'_{23} - M'_{23}M'_{33} = M'_{13}M'_{34} - M'_{23}M'_{33} - M'_{23}M'_{33} - M'_{23}M'_{33} = M'_{13}M'_{34} - M'_{23}M'_{33} - M'_{23}M'_{33} - M'_{23}M'_{33} = M'_{13}M'_{34} - M'_{23}M'_{33} - M'_{23}M'_{33} = M'_{13}M'_{34} - M'_{23}M'_{33} - M'_{23}M'_{33} = M'_{13}M'_{34} - M'_{23}M'_{33} - M'$$

since $M'_{22}M'_{23} = M'_{11}M'_{14}$, and $M'_{12}M'_{13} = M'_{12}M'_{24}$. Then, it follows that

$$M_{14}'M_{44}' = M_{13}'M_{34}' = M_{14}'M_{33}',$$

which implies that $M'_{33} = M'_{44}$, since $M'_{14} \neq 0$. But by our construction of M', we have $\zeta(M') \neq 0$. This is a contradiction.

Therefore, our assumption that $\xi(R_n(M')) = 0$ for all $n \ge 1$ must be false. Let $\xi \circ R_n$ denote the composition of ξ and R_n . There exists some $n^* \ge 1$ such that $(\xi \circ R_{n^*})(M') = (\xi \circ R_{n^*})(\mathcal{S}_M(\theta^*)) \ne 0$. So, Corollary 33 allows us to find some $M'' \in \mathfrak{R}(M, \mathcal{F}' \cup \{(\xi \circ R_{n^*})\}) \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{>0})$. Since $F(\mathcal{R}_{M''}(1)) = F((M'')^3) \ne 0$ for all $F \in \{\rho_{\texttt{tensor}}, \zeta, \zeta_1, \zeta_4\}$, and $\xi(\mathcal{R}_{M''}(n^*)) \ne 0$, Lemma 57 allows us to find some $N \in \mathfrak{R}(M, \{\rho_{\texttt{tensor}}, \zeta, \zeta_1, \zeta_4, \xi\}) \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{>0}) \subseteq \mathfrak{R}(M, \{\rho_{\texttt{tensor}}, \zeta, \zeta_1, \zeta_4, \xi\}) \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{>0})$.

Once again, in view of $\zeta(N) \neq 0$, we consider the cardinality of the set $\{N_{11}, N_{22}, N_{33}\}$. If the cardinality is 3, then N is diagonal distinct, and Theorem 68 implies that $\text{Pl-GH}(N) \leq \text{Pl-GH}(M)$ is #P-hard. If the cardinality is 2, then since $\zeta_1(N) \neq 0$, by Lemma 69 and Lemma 73, $\text{Pl-GH}(N) \leq \text{Pl-GH}(M)$ is #P-hard. If the cardinality is 1, then since $\zeta_4(N) \neq 0$ and $\zeta_5(N) \neq 0$, and also $\zeta_1(N) \neq 0$, by Lemma 71 and by Lemma 73, $\text{Pl-GH}(N) \leq \text{Pl-GH}(M)$ is #P-hard.

Lemma 69, Lemma 70, Lemma 71, Lemma 72, Lemma 73, and Lemma 74 together imply the following theorem.

Theorem 75. Let $M \in Sym_4^{pd}(\mathbb{R}_{>0})$ be a non-diagonally distinct matrix such that $\rho_{tensor}(M) \neq 0$. Let \mathcal{F} be a countable set of $Sym_4(\mathbb{R})$ -polynomials such that $F(M) \neq 0$ for all $F \in \mathcal{F}$. Unless M is isomorphic to a matrix of Form (I), (III), (IV), or (VI), Pl-GH(M) is #P-hard. Moreover,

- 1. If M is isomorphic to a matrix of Form (I), but not isomorphic to any matrix of Form (III), (IV), or (VI), then either Pl-GH(M) is #P-hard, or there exists some $N \in \mathfrak{R}(M, \mathcal{F}) \cap$ Sym₄^{pd}($\mathbb{R}_{>0}$) such that N is isomorphic to a matrix of Form (I), and $N_{11} = N_{22}$, but $N_{22} \neq$ $N_{33} \neq N_{44}$ are pairwise distinct.
- 2. If M is isomorphic to a matrix of Form (III), but not isomorphic to any matrix of Form (VI), then either Pl-GH(M) is #P-hard, or there exists some $N \in \mathfrak{R}(M, \mathcal{F}) \cap Sym_4^{pd}(\mathbb{R}_{>0})$ such that N is isomorphic to a matrix of Form (III), and $N_{11} = N_{22} \neq N_{33} = N_{44}$.
- 3. If M is isomorphic to a matrix of Form (IV), but not isomorphic to any matrix of Form (VI), then either Pl-GH(M) is #P-hard, or there exists some $N \in \mathfrak{R}(M, \mathcal{F}) \cap Sym_4^{pd}(\mathbb{R}_{>0})$ such that N is isomorphic to a matrix of Form (IV), and $N_{11} = N_{22} = N_{33} \neq N_{44}$.

Remark. In the statement of Theorem 75, a 4th item not listed explicitly but is logically implied is that M can be isomorphic to a matrix of Form (VI). The enumeration is in a reverse order of the last appearance of (I), (III), (IV), or (VI), when M is isomorphic to a matrix of Form (I), (III), (IV), or (VI). After the implicit 4th item of Form (VI), item 3 is when M is in Form (IV) but not (VI), item 2 is when M is in Form (III) but not (IV) or (VI), and item 1 is when M is in Form (I) but not (III) or (IV) or (VI). We note that if a matrix M is both isomorphic to a matrix of Form (III), and also isomorphic to a matrix of Form (IV), then in fact all diagonal entries are equal, and all off diagonal entries are equal. To see this, in Form (IV) there are three equal diagonal entries and in Form (III) they come in two equal pairs (a, a) and (b, b). Thus a = b and all diagonal entries are equal. For the off diagonal entries, define two graphs K and K', both a copy of K₄, but with labeled edges according to Forms (III) and (IV) respectively. In K the edges will be labeled with t, x, y and z, and the list of incident edges for the 4 vertices is (tyz, tyz, xyz, xyz). In K' we will label them x' and z', and the list of incident edges is (x'x'z', x'x'z', x'x'z', z'z'z'). Being isomorphic, from z'z'z' we have y = z and either t = y or x = y. Hence there are two equal triples in K, which implies that x' = z' in K', and hence all labels are equal. Thus, such a matrix M is in the Potts model, which is a special case of Form (VI). Consequently, in the 2nd (respectively, the 3rd) item, when we assume that M is isomorphic to a matrix of Form (III) (respectively, Form (IV)), but not isomorphic to any matrix of Form (VI), it follows that M is not isomorphic to any matrix of Form (IV) (respectively, Form (III)) either. (In particular, in item 2, it is not explicitly stated that M is not in Form (IV), as being in Form (III) but not Form (VI) already implies this.)

11 Forms (I), (III), (IV) and (VI)

Following Theorem 75, we now only need to deal with the matrices of the Forms (I), (III), (IV), and (VI), as they are in Theorem 75. We will deal with these Forms, one by one.

Remark. Due to Theorem 75, from this point onwards, when we refer to a matrix M of Form (I), we may assume that $M_{11} = M_{22}$, but $M_{22} \neq M_{33} \neq M_{44}$ are pairwise distinct. Similarly, if we refer to a matrix of Form (III), we may assume that $M_{11} = M_{22} \neq M_{33} = M_{44}$, and if we refer to a matrix of Form (IV) we may also assume that it satisfies $M_{11} = M_{22} = M_{33} \neq M_{44}$.

11.1 Form (I)

We will now deal with matrices of Form (I) such that $M_{11} = M_{22}$, and $M_{22} \neq M_{33} \neq M_{44}$ are pairwise distinct. Our strategy will be to show that, if the matrix $M \in \text{Sym}_4^{\text{pd}}(\mathbb{R}_{>0})$ is of Form (I) such that $\rho_{\text{tensor}}(M) \neq 0$, then for any $\mathbf{0} \neq \mathbf{x} \in \chi_4$, we have $\Psi_{\mathbf{x}}(M) \neq 0$. Before we can jump into it, we will prove that we may assume that the matrix M of Form (I) has some additional structural properties.

$$\begin{pmatrix} a & x & y & z \\ x & a & y & z \\ y & y & b & t \\ z & z & t & c \end{pmatrix}$$

Figure 7: Form (I)

Lemma 76. Let $M \in Sym_4^{pd}(\mathbb{R}_{>0})$ be of Form (I) such that $\rho_{tensor}(M) \neq 0$, where $M_{11} = M_{22}$, but $M_{22} \neq M_{33} \neq M_{44}$ are pairwise distinct. Let \mathcal{F} be a countable set of $Sym_4(\mathbb{R})$ -polynomials such that $F(M) \neq 0$ for all $F \in \mathcal{F}$. Then, either Pl-GH(M) is #P-hard, or there exists some $N \in \mathfrak{R}(M, \mathcal{F} \cup \{\rho_{\texttt{tensor}}\}) \cap Sym_4^{pd}(\mathbb{R}_{>0})$ that is of Form (I), such that $N_{11} \neq N_{33} \neq N_{44}$ and $N_{22} \neq N_{33} \neq N_{44}$ are both pairwise distinct, $N_{12} \neq N_{13} \neq N_{14}$ are pairwise distinct, $N_{34} \neq N_{13}, N_{14}$, and $N_{ii} \neq N_{jk}$ for all $i \in [4]$ and for all $j \neq k \in [4]$.

Remark. The matrix N of Form (I) from the conclusion of Lemma 76 stipulates that all diagonal elements are distinct except $M_{11} = M_{22}$, and are distinct from all off diagonal elements. And all off diagonal elements denoted by distinct letters in Fig. 7 are distinct except possibly x = t.

Proof. We will first define $\zeta : N \mapsto (N_{11} - N_{33})(N_{11} - N_{44})(N_{22} - N_{33})(N_{22} - N_{44})(N_{33} - N_{44}),$ and $\zeta' : N \mapsto \prod_{i \in [4], j \neq k \in [4]} (N_{ii} - N_{jk})$. We note that by our choice of M, $\zeta(M) \neq 0$. While we cannot claim that $\zeta'(M) \neq 0$, we note that $\zeta'(I) \neq 0$. We will let $\mathcal{F}' = \{F' : N \mapsto F(N^3) \mid F \in \mathcal{F} \cup \{\rho_{\texttt{tensor}}, \zeta, \zeta'\}\}$. We will now consider $\mathcal{S}_M(\theta)$. We know that $F'(\mathcal{S}_M(\theta)) \neq 0$ for all $F' \in \mathcal{F}'$ for some $\theta \in \{\frac{1}{3}, 0\}$. We also know that there exists some $\delta > 0$ such that $|\mathcal{S}_M(\theta)_{ij} - I_{ij}| < \frac{1}{3}$ for all $i, j \in [4]$, and $0 < \theta < \delta$. We can therefore use Lemma 32 to find some $M' = \mathcal{S}_M(\theta^*) \in \mathfrak{R}(M, \mathcal{F}' \cup \{\zeta, \zeta'\}) \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{\neq 0})$ for some $0 < \theta^* < \delta$.

We will now consider $R_n(M')$ for all $n \ge 1$. We will let $X = \{M'_{ab} : a, b \in [4]\}$, and for each $x \in X$, and $i, j \in [4]$, define $c_{ij}(x) = \sum_{a,b \in [4]: M'_{ab} = x} (M'_{ia}M'_{jb})$. Then for any $i \ne j \in \{2, 3, 4\}$, we see that

$$(R_n(M'))_{1i} - (R_n(M'))_{1j} = \sum_{x \in X} x^n \cdot (c_{1i}(x) - c_{1j}(x))$$

Let us assume that for some $i \neq j \in \{2, 3, 4\}$, $(R_n(M'))_{1i} - (R_n(M'))_{1j} = 0$ for all $n \geq 1$. In that case, we see that we have a full rank Vandermonde system of linear equations, which implies that $c_{1i}(x) - c_{1j}(x) = 0$ for all $x \in X$.

From our construction of M', we know that $M'_{ab} = M'_{33}$ implies that (a, b) = (3, 3), and that $M'_{ab} = M'_{44}$ implies that (a, b) = (4, 4). This is because by choosing $0 < \theta^* < \delta$ our M' is close to I, and thus diagonal elements are all distint from off diagonal elements, and $\zeta(M') \neq 0$ separates M'_{33} and M'_{44} from each other and also from the other diagonal elements. So, we see that

$$c_{1i}(M'_{33}) - c_{1j}(M'_{33}) = M'_{13}M'_{i3} - M'_{13}M'_{j3}, \text{ and } c_{1i}(M'_{44}) - c_{1j}(M'_{44}) = M'_{14}M'_{i4} - M'_{14}M'_{j4}.$$

We note that by our choice of $i \neq j \in \{2,3,4\}$, at least one of $i, j \in \{3,4\}$. Without loss of generality, we may assume that it is *i*. If i = 3, that implies that $M'_{13}M'_{33} = M'_{13}M'_{j3}$ for some $j \neq 3$, and if i = 4, that implies that $M'_{14}M'_{44} = M'_{14}M'_{j4}$ for some $j \neq 4$. As $M'_{13}, M'_{14} \neq 0$, in either case, we get a contradiction, since by our choice of M', no diagonal entry can be equal to a non-diagonal entry. Therefore, for each $i \neq j \in \{2,3,4\}$, there exists some $n_{ij} \geq 1$ such that $(R_{n_{ij}}(M'))_{1i} - (R_{n_{ij}}(M'))_{1j} \neq 0$.

Similarly, for $i \in \{3, 4\}$, we see that for all $n \ge 1$,

$$(R_n(M'))_{1i} - (R_n(M'))_{34} = \sum_{x \in X} x^n \cdot (c_{1i}(x) - c_{34}(x)).$$

Let us assume that for some $i \in \{3, 4\}$, $(R_n(M'))_{1i} - (R_n(M'))_{34} = 0$ for all $n \ge 1$. Once again, since this forms a full rank Vandermonde system of linear equations, we see that $c_{1i}(x) - c_{34}(x) = 0$ for all $x \in X$. If i = 3, we will consider $c_{13}(M'_{44}) - c_{34}(M'_{44}) = M'_{14}M'_{34} - M'_{34}M'_{44} = 0$, which implies that $M'_{14} = M'_{44}$, and if i = 4, we consider $c_{14}(M'_{33}) - c_{34}(M'_{33}) = M'_{13}M'_{34} - M'_{33}M'_{34} = 0$, which implies that $M'_{13} = M'_{33}$. Since neither of these are possible, we conclude that for each $i \in \{3, 4\}$, there exists some $n'_i \ge 1$ such that $(R_{n'_i}(M'))_{1i} - (R_{n'_i}(M'))_{34} \ne 0$.

For each $i \neq j \in \{2,3,4\}$, we can now construct a $\operatorname{Sym}_4(\mathbb{R})$ -polynomial $\xi_{ij} : N \mapsto (R_{n_{ij}}(N))_{1i} - (R_{n_{ij}}(N))_{1j}$. For each $i \in \{3,4\}$, we can also construct $\xi'_i : N \mapsto (R_{n'_i}(N))_{1i} - (R_{n'_i}(N))_{34}$. We have seen that $\xi_{ij}(\mathcal{S}_M(\theta^*)) \neq 0$ for all $i \neq j \in \{2,3,4\}$, and $\xi'_i(\mathcal{S}_M(\theta^*)) \neq 0$, for all $i \in \{3,4\}$. So, we can use Corollary 33 to find some $M'' \in \mathfrak{R}(M, \mathcal{F}' \cup \{\xi_{23}, \xi_{24}, \xi_{34}, \xi'_3, \xi'_4\}) \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{>0})$. But then, we see that $F(\mathcal{R}_{M''}(1)) \neq 0$ for all $F \in \mathcal{F} \cup \{\rho_{\text{tensor}}, \zeta, \zeta'\}$. Furthermore, let $\xi_{ij}^o : N \mapsto N_{1i} - N_{1j}$ for all $i \neq j \in \{2,3,4\}$, and $\xi'_i(\mathcal{R}_{M''}(n_{ij})) \neq 0$ for all $i \in \{3,4\}$. Then $\xi_{ij}^o(\mathcal{R}_{M''}(n_{ij})) \neq 0$ for all $i \neq j \in \{2,3,4\}$, and $\xi'_i(\mathcal{R}_{M''}(n'_i)) \neq 0$ for all $i \in \{3,4\}$. So, Lemma 57 allows us to find the required $N \in \mathfrak{R}(M, \mathcal{F} \cup \{\rho_{\text{tensor}}, \zeta, \zeta', \xi_{23}^o, \xi_{24}^o, \xi_{34}^o, \xi_{4}^{\prime o}\}) \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{>0})$.

Since $\zeta(N) \neq 0$, we see that $N_{11} \neq N_{33} \neq N_{44}$, and $N_{22} \neq N_{33} \neq N_{44}$ are both pairwise distinct. If $N_{11} \neq N_{22}$, then N is diagonal distinct, and since $\rho_{\texttt{tensor}}(N) \neq 0$, Theorem 68 allows us to conclude that Pl-GH(M) is #P-hard. Otherwise, if N is not of Form (I), then Lemma 69 and Lemma 73 tell us once again that Pl-GH(N) is #P-hard. So, the only other possibility is that N is of Form (I) such that $N_{12} \neq N_{13} \neq N_{14}$ are pairwise distinct (because $\xi_{23}^o(N) \neq 0, \xi_{24}^o(N) \neq 0, \xi_{34}^o(N) \neq 0$), and $N_{34} \neq N_{13}, N_{14}$ (because $\xi_{33}^{\prime o}(N) \neq 0, \xi_{4}^{\prime o}(N) \neq 0$), and $N_{ii} \neq N_{jk}$ for all $i \in [4]$ and for all $j \neq k \in [4]$ (because $\zeta'(N) \neq 0$).

Lemma 76 allows us to claim that for any $M \in \text{Sym}_4^{\text{pd}}(\mathbb{R}_{>0})$ of Form (I) (such that $\rho_{\text{tensor}}(M) \neq 0$), we may assume that $M_{12} \neq M_{13} \neq M_{14}$ are pairwise distinct, and that $M_{34} \neq M_{13}, M_{14}$. Ideally, we would like to be able to say that we can assume that $M_{12} \neq M_{34}$ as well. That is however, not true in general. As it turns out, it is possible that $M_{12} = M_{34}$, but in that case, we will be able to show that $\rho_{\text{tensor}}(M) = 0$.

Lemma 77. Let $M \in Sym_4^{pd}(\mathbb{R}_{>0})$ be of Form (I) such that $\rho_{tensor}(M) \neq 0$. Let \mathcal{F} be a countable set of $Sym_4(\mathbb{R})$ -polynomials such that $F(M) \neq 0$ for all $F \in \mathcal{F}$. Then, either Pl-GH(M) is #P-hard, or there exists some $N \in \mathfrak{R}(M, \mathcal{F} \cup \{\rho_{tensor}\}) \cap Sym_4^{pd}(\mathbb{R}_{>0})$ that is of Form (I), such that either $N_{12} \neq N_{34}$, or $(N_{11})^2 \neq N_{33}N_{44}$.

Proof. We will first define $\zeta : N \mapsto (N_{11} - N_{33})(N_{11} - N_{44})(N_{22} - N_{33})(N_{22} - N_{44})(N_{33} - N_{44})$ as in the proof of Lemma 76. From our choice of M, we note that $\zeta(M) \neq 0$. We will now let $\mathcal{F}' = \{F' : N \mapsto F(N^3) \mid F \in \mathcal{F} \cup \{\rho_{\texttt{tensor}}, \zeta\}\}$. We note that $F'(\mathcal{S}_M(\frac{1}{3})) \neq 0$ for all $F' \in \mathcal{F}'$. We also know that there exists some $\delta > 0$ such that for all $0 < \theta < \delta$, $|\mathcal{S}_M(\theta)_{ij} - I_{ij}| < \frac{1}{3}$. We can now use Lemma 32 to find $M' = \mathcal{S}_M(\theta^*) \in \mathfrak{R}(M, \mathcal{F}' \cup \{\zeta\}) \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{\neq 0})$, such that $0 < \theta^* < \delta$. Let us assume that M' is not of Form (I). Since $\zeta(M') \neq 0$, we know that $(M')_{11} \neq (M')_{33} \neq (M')_{44}$ are pairwise distinct, and $(M')_{22} \neq (M')_{33} \neq (M')_{44}$ are also pairwise distinct. If $(M')_{11} \neq (M')_{22}$, then M' is diagonal distinct, and Theorem 68 implies that P1-GH $(M') \leq$ P1-GH(M) is #P-hard. Otherwise, Lemma 69 implies that unless M' is of Form (I), then Lemma 73 implies that P1-GH $(M') \leq$ P1-GH(M) is #P-hard. Finally, if M' has Form (II), and does not have Form (I), then Lemma 73 implies that P1-GH $(M') \leq$ P1-GH $(M') \leq$ P1-GH(M') is $(M'_{11})^2 \neq M'_{33}M'_{44}$, we can immediately use Corollary 33 to find the required N. So, we may assume otherwise: $M'_{12} = M'_{34}$ and $(M'_{11})^2 = M'_{33}M'_{44}$. In terms of notations in Fig. 7, we have x = t and $a^2 = bc$.

We will now consider $R_n(M')$ for all $n \ge 1$. We will let $X = \{(M')_{ab} : a, b \in [4]\}$, and define $c_{ij}(x) = \sum_{a,b \in [4]: (M')_{ab} = x} M'_{ia}M'_{jb}$. We note that

$$R_n(M')_{12} - R_n(M')_{34} = \sum_{x \text{ in } X} x^n \cdot (c_{12}(x) - c_{34}(x)).$$

If $R_n(M')_{12} - R_n(M')_{34} = 0$ for all $n \ge 1$, then we have a Vandermonde system of linear equations of size O(1). That implies that $c_{12}(x) - c_{34}(x) = 0$ for all $x \in X$.

From our construction of M', we know that $M'_{ab} = M'_{33}$ implies that (a, b) = (3, 3), $M'_{ab} = M'_{44}$ implies that (a, b) = (4, 4), and that $M'_{ab} = M'_{11}$ implies that $(a, b) \in \{(1, 1), (2, 2)\}$. So, we see that

$$c_{12}(M'_{33}) - c_{34}(M'_{33}) = M'_{13}M'_{13} - M'_{33}M'_{34},$$

$$c_{12}(M'_{44}) - c_{34}(M'_{44}) = M'_{14}M'_{14} - M'_{34}M'_{44},$$

$$c_{12}(M'_{11}) - c_{34}(M'_{11}) = M'_{11}M'_{12} + M'_{12}M'_{11} - M'_{13}M'_{14} - M'_{13}M'_{14}$$

$$= 2M_{11}M'_{12} - 2M'_{13}M'_{14},$$

and they all equal to 0. (Here we used the fact that $M'_{23} = M'_{13}, M'_{24} = M'_{14}, M'_{11} = M'_{22}$ in Form (I).) In terms of the notations in Fig. 7 (with x = t, and $a^2 = bc$), we have $y^2 = bx$, $z^2 = cx$, and ax = yz. Together with x = t and $a^2 = bc$, we can verify that $\rho_{\text{tensor}}(M') = 0$, where ρ_{tensor} is defined in Lemma 65. (We use the switching $2 \leftrightarrow 4$ in ρ_{tensor} to ρ_{tensor} .)

But by construction of M', we ensured that $\rho_{\text{tensor}}(M') \neq 0$. Therefore, there must exist some $n \geq 1$ such that $R_n(M')_{12} - R_n(M')_{34} \neq 0$. So, if we define $\xi : N \mapsto R_n(N)_{12} - R_n(N)_{34}$, we see that $\xi(\mathcal{S}_M(\theta^*)) = \xi(M') \neq 0$. Now, we can use Corollary 33 to find $M'' \in \mathfrak{R}(M, \mathcal{F} \cup \{\xi\}) \cap \text{Sym}_4^{\text{pd}}(\mathbb{R}_{>0})$. Finally, Lemma 57 allows us to find $N \in \mathfrak{R}(M, \mathcal{F} \cup \{\rho_{\text{tensor}}, \zeta, \xi^o\}) \cap \text{Sym}_4^{\text{pd}}(\mathbb{R}_{>0})$, where $\xi^o : N \mapsto N_{12} - N_{34}$. Since $\zeta(N) \neq 0$, N may either be diagonal distinct, or of the form $N_{11} = N_{22}$ with $N_{22} \neq N_{33} \neq N_{44}$ pairwise distinct. If N is diagonal distinct, then Theorem 68 implies that P1-GH(N) \leq P1-GH(M) is #P-hard. Otherwise, Lemma 69 implies that unless N is either of Form (I), P1-GH(N) \leq P1-GH(M) is #P-hard. Since $\xi^o(N) \neq 0$, we know that $N_{12} \neq N_{34}$. So, N is the required matrix.

Remark. We note that in the statement of Lemma 77, we claim to be able to find some N such that $N_{12} \neq N_{34}$, or $(N_{11})^2 \neq N_{33}N_{44}$, however, at the end of the proof, we were able to find some N such that $N_{12} \neq N_{34}$. So, it appears as if we could eliminate the option that $(N_{11})^2 \neq N_{33}N_{44}$ from the statement of the lemma. However, we cannot do that. We note that in the proof of Lemma 77, when we constructed M', if $M'_{12} = M'_{34}$, but $(M'_{11})^2 \neq M'_{33}M'_{44}$, we could then have produced some N such that $N_{12} = N_{34}$, but $(N_{11})^2 \neq M'_{33}M'_{44}$.

We will now show that given a matrix $M \in \operatorname{Sym}_{4}^{\operatorname{pd}}(\mathbb{R}_{>0})$ of Form (I), and any $\mathbf{0} \neq \mathbf{x} \in \chi_{4}$ of support size > 2, we can find some $N \in \mathfrak{R}(M, \mathcal{F}_{M} \cup \{\phi_{\mathbf{x}}\}) \cap \operatorname{Sym}_{4}^{\operatorname{pd}}(\mathbb{R}_{>0})$. We note that if $\mathbf{0} \neq \mathbf{x} \in \chi_{4}$ has support size ≤ 2 , it must be of the form $\mathbf{x} = c \cdot \delta_{ij}$ for some $\delta_{ij} \in \mathcal{D}$ (as defined in Eq. (11)). We will show later that our ability to use Theorem 18 is not hindered by such $\mathbf{x} \in \chi_4$, so we do not have to worry about them.

Lemma 78. Let $M \in Sym_4^{pd}(\mathbb{R}_{>0})$ be of Form (I) such that $\rho_{tensor}(M) \neq 0$. Let \mathcal{F} be a countable set of $Sym_4(\mathbb{R})$ -polynomials such that $F(M) \neq 0$ for all $F \in \mathcal{F}$. Let $\xi_{\mathbf{x}}$ be a $Sym_4(\mathbb{R})$ -polynomial such that $\xi_{\mathbf{x}} : N \mapsto \phi_{\mathbf{x}}(N_{11}, \ldots, N_{44})$ for any $\mathbf{0} \neq \mathbf{x} \in \chi_4$ of support size greater than 2. Then, either Pl-GH(M) is #P-hard, or there exists some $N \in \mathfrak{R}(M, \mathcal{F} \cup \{\rho_{tensor}, \xi_{\mathbf{x}}\}) \cap Sym_4^{pd}(\mathbb{R}_{>0})$ of Form (I).

Proof. We will first use Lemma 77 to find $M' \in \mathfrak{R}(M, \mathcal{F} \cup \{\rho_{\texttt{tensor}}\}) \cap \operatorname{Sym}_{4}^{\mathsf{pd}}(\mathbb{R}_{>0})$ that is of Form (I), such that either $(M')_{12} \neq (M')_{34}$, or $((M')_{11})^2 \neq (M')_{33}(M')_{44}$. (In Fig. 7, this is $x \neq t$ or $a^2 \neq bc$.) We now define $\zeta : N \mapsto (N_{11} - N_{33})(N_{11} - N_{44})(N_{22} - N_{33})(N_{22} - N_{44})(N_{33} - N_{44})$. We can now construct a matrix $(M')^{\sigma}$ by permuting both the rows and columns of M by some $\sigma \in S_4$ (that switches $2 \leftrightarrow 4$). Then we see that the condition that either $(M')_{12} \neq (M')_{34}$, or $((M')_{11})^2 \neq (M')_{33}(M')_{44}$ means exactly that either $(M')_{14}^{\sigma} \neq (M')_{23}^{\sigma}$, or $(M')_{11}^{\sigma}(M')_{44}^{\sigma} \neq (M')_{22}^{\sigma}(M')_{33}^{\sigma}$. (By Form (I), $M'_{11} = M'_{22}$, which gives $(M')_{11}^{\sigma} = (M')_{44}^{\sigma}$.) This is precisely the condition for Lemma 66 that allows us to find some $(M'')^{\sigma} \in \mathfrak{R}((M')^{\sigma}, \mathcal{F}^{\sigma} \cup \{\xi_{(1,-1,-1,1)}\}) \cap \operatorname{Sym}_{4}^{\mathsf{pd}}(\mathbb{R}_{>0})$, where

$$\mathcal{F}^{\sigma} = \left\{ F^{\sigma} : N \mapsto F(N^{\sigma^{-1}}) \mid F \in \mathcal{F} \cup \{\rho_{\texttt{tensor}}, \zeta\} \right\}$$

If we now look at (M'') (by switching back $2 \leftrightarrow 4$), we see that $\xi_{(1,1,-1,-1)}(M'') \neq 0$, and that $F(M'') \neq 0$ for all $F \in \mathcal{F} \cup \{\rho_{\texttt{tensor}}, \zeta\}$. From the construction of ζ , we can see that if $M''_{11} \neq M''_{22}$, then M'' is diagonal distinct, and Theorem 68 implies that $\texttt{Pl-GH}(M'') \leq \texttt{Pl-GH}(M)$ is #P-hard. On the other hand, if $M''_{11} = M''_{22}$, then Lemma 69, and Lemma 73 together imply that unless M'' is of Form (I), then $\texttt{Pl-GH}(M'') \leq \texttt{Pl-GH}(M)$ is #P-hard. So, we may assume that M'' is in fact, of Form (I).

Finally, we can use Lemma 76, to find $M''' \in \mathfrak{R}(M'', \mathcal{F} \cup \{\rho_{\texttt{tensor}}, \xi_{(1,1,-1,-1)}, \zeta\}) \cap \text{Sym}_{jk}^{pd}(\mathbb{R}_{>0})$ such that $M''_{12} \neq M''_{13} \neq M''_{14}$ are pairwise distinct, $M''_{34} \neq M''_{13}, M''_{14}$, and that $M''_{ii} \neq M''_{jk}$ for all $i \in [4]$, and $j \neq k \in [4]$. We note that since $\zeta(M''') \neq 0$, if M''' is not of Form (I), then Theorem 75 implies that Pl-GH(M) is #P-hard. So, we may assume that M''' is of Form (I). For convenience, we will rename this M''' as M.

Let us consider any $\mathbf{0} \neq \mathbf{x} = (x_1, x_2, x_3, x_4) \in \chi_4$. If $\xi_{\mathbf{x}}(M) = \phi_{\mathbf{x}}(M_{11}, \dots, M_{44}) \neq 0$, then we are already done. So we may assume that $\xi_{\mathbf{x}}(M) = 0$. Let us identify those \mathbf{x} for which $\xi_{\mathbf{x}}(M) = 0$. We already know that $\xi_{(1,1,-1,-1)}(M) \neq 0$. For any \mathbf{x} such that $x_1 + x_2 = 2c$, and $x_3 = x_4 = -c$, for any non-zero $c \in \mathbb{Z}$, since we have $M_{11} = M_{22}$, this implies that $\xi_{\mathbf{x}}(M) = (\xi_{(1,1,-1,-1)}(M))^c \neq 0$. We will now make use of the following claim, which we shall prove shortly.

Claim 79. Let $\alpha_1, \ldots, \alpha_4 \in \mathbb{R}$, such that $\alpha_1 = \alpha_2$, and $|\alpha_2| \neq |\alpha_3| \neq |\alpha_4|$ are pairwise distinct. If $\phi_{\mathbf{x}}(\alpha_1, \ldots, \alpha_4) = 0$ for some $\mathbf{x} \in \chi_4$ with support size greater than 2, then $x_3 \neq 0$, $x_4 \neq 0$, and $x_1 + x_2 \neq 0$.

Since $M \in \text{Sym}_4(\mathbb{R}_{>0})$ and is of Form (I), we see that $M_{11} = M_{22}$, and $|M_{22}| \neq |M_{33}| \neq |M_{44}|$ are pairwise distinct. So, Claim 79 lets us conclude that $\xi_{\mathbf{x}}(M) = 0$ implies $x_3, x_4 \neq 0$, and $x_1 + x_2 \neq 0$.

We will now let the entries of M be generated by some $\{g_t\}_{t\in[d]}$. We may assume that M is replaced with some cM as in Lemma 4 such that $e_{ijt} \ge 0$ for all $i, j \in [4]$, and $t \in [d]$. Now, we will let $\mathfrak{m} = \max_{i,j\in[4],t\in[d]}(e_{ijt})$, pick some $m > \mathfrak{m}$, and define $z_{ij} = \sum_{t\in[d]} m^t \cdot e_{ijt}$ for all $i, j \in [4]$. We will now define $\mathcal{T}_M^* : \mathbb{R} \to \operatorname{Sym}_4(\mathbb{R})$ such that

$$\mathcal{T}_M^*(p)_{ij} = \mathcal{T}_M(p^m, p^{m^2}, \dots, p^{m^d})_{ij} = p^{z_{ij}}.$$

From our choice of m, we see that $z_{ij} = z_{i'j'}$ if and only if $(e_{ij1}, \ldots, e_{ijd}) = (e_{i'j'1}, \ldots, e_{i'j'd})$. So, from our choice of M, we see that in fact, $z_{12} \neq z_{13} \neq z_{14}$, $z_{34} \neq z_{13}$, z_{14} , and $z_{ii} \neq z_{ij}$ for all $i \neq j \in [4]$.

We will now consider $\xi_{\mathbf{x}}((\mathcal{T}_M^*(p))^2)$ for all $p \in \mathbb{R}$. Since $x_1 + x_2 \neq 0$, we may assume without loss of generality that $x_1 + x_2 > 0$. We may also assume by symmetry that $x_3 \geq x_4$. There are now two possibilities we have to deal with: x_3 may be positive, in which case x_4 must be negative (since $x_1 + \cdots + x_4 = 0$), or both x_3 and x_4 may be negative. (By Claim 79, $x_3, x_4 \neq 0$.)

We will first deal with the case where $x_3 > 0$. In this case, we see that

$$\xi_{\mathbf{x}}(\mathcal{T}_{M}^{*}(p)^{2}) = (p^{2z_{11}} + p^{2z_{12}} + p^{2z_{13}} + p^{2z_{14}})^{x_{1}+x_{2}} \cdot (2p^{2z_{13}} + p^{2z_{33}} + p^{2z_{34}})^{x_{3}} - (2p^{2z_{14}} + p^{2z_{34}} + p^{2z_{44}})^{x_{1}+x_{2}+x_{3}}.$$

Here we used the fact that $\mathcal{T}_{M}^{*}(p)$ has Form (I) (because M does) and thus $(\mathcal{T}_{M}^{*}(p)^{2})_{11} = (\mathcal{T}_{M}^{*}(p)^{2})_{22}$. We note that by our choice of z_{ij} , the exponents $2z_{11} \neq 2z_{12} \neq 2z_{13} \neq 2z_{14}$ are pairwise distinct, $2z_{13} \neq 2z_{33} \neq 2z_{34}$ are pairwise distinct, and $2z_{14} \neq 2z_{34} \neq 2z_{44}$ are pairwise distinct. So, the following claim (which we shall prove soon) allows us to claim that $\xi_{\mathbf{x}}(\mathcal{T}_{M}^{*}(p)^{2})$ is not the zero polynomial.

Claim 80. Let $n, m \in \mathbb{Z}_{>0}$, and $y_{11} < y_{12} < y_{13} < y_{14}$, $y_{21} < y_{22} < y_{23}$, and $y_{31} < y_{32} < y_{33} \in \mathbb{Z}_{>0}$. Assume the multiset $\{\alpha_1, \alpha_2, \alpha_3\} = \{\beta_1, \beta_2, \beta_3\} = \{2, 1, 1\}$. Define the polynomial $f : \mathbb{R} \to \mathbb{R}$ such that

$$f(p) = (p^{y_{11}} + p^{y_{12}} + p^{y_{13}} + p^{y_{14}})^n \cdot (\alpha_1 p^{y_{21}} + \alpha_2 p^{y_{22}} + \alpha_3 p^{y_{23}})^m - (\beta_1 p^{y_{31}} + \beta_2 p^{y_{32}} + \beta_3 p^{y_{33}})^{n+m},$$

then f(p) is not the zero polynomial.

On the other hand, if $x_3 < 0$, we see that

$$\xi_{\mathbf{x}}(\mathcal{T}_{M}^{*}(p)^{2}) = (p^{2z_{11}} + p^{2z_{12}} + p^{2z_{13}} + p^{2z_{14}})^{x_{1}+x_{2}} - (2p^{2z_{13}} + p^{2z_{33}} + p^{2z_{34}})^{-x_{3}} \cdot (2p^{2z_{14}} + p^{2z_{34}} + p^{2z_{44}})^{x_{1}+x_{2}+x_{3}}.$$

We will now make use of the following claim, which we shall prove shortly.

Claim 81. Let $n, m \in \mathbb{Z}_{>0}$, and $y_{11} < y_{12} < y_{13} < y_{14}$, $y_{21} < y_{22} < y_{23}$, and $y_{31} < y_{32} < y_{33} \in \mathbb{Z}_{>0}$. Assume the multiset $\{\alpha_1, \alpha_2, \alpha_3\} = \{\beta_1, \beta_2, \beta_3\} = \{2, 1, 1\}$. Define the polynomial $f : \mathbb{R} \to \mathbb{R}$ such that

$$f(p) = (p^{y_{11}} + p^{y_{12}} + p^{y_{13}} + p^{y_{14}})^{n+m} - (\alpha_1 p^{y_{21}} + \alpha_2 p^{y_{22}} + \alpha_3 p^{y_{23}})^n \cdot (\beta_1 p^{y_{31}} + \beta_2 p^{y_{32}} + \beta_3 p^{y_{33}})^m,$$

then f(p) is not the zero polynomial, unless n = m.

Claim 81 implies that if $\xi_{\mathbf{x}}(\mathcal{T}_M^*(p)^2)$ is the zero polynomial only if $-x_3 = x_1 + x_2 + x_3$. This implies that if $x_3 = -c$, then $x_1 + x_2 = 2c$, and $x_4 = -c$. In other words, $\mathbf{x} = (x_1, x_2, -c, -c)$ for some $c \in \mathbb{Z}_{\neq 0}$, where $x_1 + x_2 = 2c$. But by our construction of M, we already ensured that for such $\mathbf{x} \in \chi_4, \xi_{\mathbf{x}}(M) \neq 0$. So, we see that if $\xi_{\mathbf{x}}(M) = 0$, then $\xi_{\mathbf{x}}(\mathcal{T}_M^*(p)^2)$ is not the zero polynomial.

So, we see that in either case, when $\mathbf{x} \in \chi_4$ has a support size greater than 2, if $\xi_{\mathbf{x}}(M) = 0$, there exists some $p^* \in \mathbb{R}$ such that $\xi_{\mathbf{x}}((\mathcal{T}_M^*(p^*))^2) \neq 0$. We will now define $\xi_{2,\mathbf{x}} : N \mapsto \phi_{\mathbf{x}}(N^2)$, and $\zeta : N \mapsto (N_{11} - N_{33})(N_{11} - N_{44})(N_{22} - N_{33})(N_{22} - N_{44})(N_{33} - N_{44})$. So, we can use Lemma 28 to find some $M' \in \mathfrak{R}(M, \mathcal{F} \cup \{\rho_{\texttt{tensor}}, \xi_{2,\mathbf{x}}, \zeta\}) \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{>0})$. Since $\xi_{2,\mathbf{x}}(M') = \xi_{\mathbf{x}}((M')^2) \neq 0$, Corollary 33 allows us to find the required $N \in \mathfrak{R}(M', \mathcal{F} \cup \{\rho_{\texttt{tensor}}, \xi_{\mathbf{x}}, \zeta\}) \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{>0})$. If N is diagonal distinct, Theorem 68 proves that $\operatorname{Pl-GH}(M)$ is $\#\operatorname{P-hard}$. Otherwise, Theorem 75 implies that either $\operatorname{Pl-GH}(N) \leq \operatorname{Pl-GH}(M)$ is $\#\operatorname{P-hard}$, or N is of Form (I) as required. \square

We will now finish the proof of Lemma 78 by proving Claim 79, Claim 80 and Claim 81.

Proof of Claim 79. For a contradiction, suppose $x_3 = 0$ or $x_4 = 0$ or $x_1 + x_2 = 0$. First suppose $x_3 = 0$. If $x_1 + x_2 = 0$, that would imply that $x_4 = 0$ as well, which is not possible, since the support size of **x** is greater than 2 by assumption. So, without loss of generality, we may assume that $x_1 + x_2 > 0$, by replacing **x** with $-\mathbf{x}$. Now,

$$\phi_{\mathbf{x}}(\alpha_1,\ldots,\alpha_4) = (\alpha_1)^{x_1+x_2} - (\alpha_4)^{x_1+x_2}.$$

Since $|\alpha_1| \neq |\alpha_4|$, we see that it is not possible that $\phi_{\mathbf{x}}(\alpha_1, \ldots, \alpha_4) = 0$. So $x_3 = 0$ is impossible. By symmetry, we see that $x_4 = 0$ is also impossible.

Now, let $x_1 + x_2 = 0$. This implies that $x_3 + x_4 = 0$ as well. We may assume without loss of generality that $x_3 > 0$, since we already know that $x_3 \neq 0$. Once again, we see that $\phi_{\mathbf{x}}(\alpha_1, \ldots, \alpha_4) = 0$ would imply that $(\alpha_3)^{x_3} = (\alpha_4)^{x_3}$, which is also known to be not possible since $|\alpha_3| \neq |\alpha_4|$. This finishes the proof of Claim 79.

Proof of Claim 80. For convenience, we will define the polynomials t_1, t_2, t_3 such that

$$t_1(p) = (p^{y_{11}} + p^{y_{12}} + p^{y_{13}} + p^{y_{14}}),$$

$$t_2(p) = (\alpha_1 p^{y_{21}} + \alpha_2 p^{y_{22}} + \alpha_3 p^{y_{23}}), \text{ and}$$

$$t_3(p) = (\beta_1 p^{y_{31}} + \beta_2 p^{y_{32}} + \beta_3 p^{y_{33}}).$$

Then, we see that $f(p) = t_1(p)^n \cdot t_2(p)^m - t_3(p)^{n+m}$.

Let us assume that f(p) = 0 for all $p \in \mathbb{R}$. This means that

$$t_1(p)^n \cdot t_2(p)^m = t_3(p)^{n+m}$$

for all $p \in \mathbb{R}$. Now, if $\beta_1 = 2$, that means that the least degree term of the RHS has a coefficient of 2^{n+m} . But the least degree term of the LHS can only be $\alpha_1^m \in \{1, 2^m\}$. We are given n > 0. In either case, it is not equal to 2^{n+m} , and we get a contradiction. So, we find that $\beta_1 = 1$. Now, if $\alpha_1 = 2$, we get a similar contradiction (using m > 0), as the least degree term of the RHS will be 1, and not equal to 2^m . So, we see that $\alpha_1 = \beta_1 = 1$. By reasoning about the highest degree term instead of the least degree term, we can also see that $\alpha_3 = \beta_3 = 1$. This means that $\alpha_2 = \beta_2 = 2$. Now, $p^{ny_{11}+my_{21}}$, and $p^{(n+m)y_{31}}$ are the least degree terms of the LHS and RHS respectively. So, they must be equal. This means that $ny_{11} + my_{21} = (n+m)y_{31}$. It also means that

$$t_1(p)^n \cdot t_2(p)^m - p^{ny_{11} + my_{21}} = t_3(p)^{n+m} - p^{(n+m)y_{31}}.$$
(21)

The least degree term of the RHS of Eq. (21) will now be $2(n+m)p^{(n+m-1)y_{31}+y_{32}}$ (here the coefficient 2 is β_2), while the least degree term of the LHS may be either $np^{(n-1)y_{11}+y_{12}+my_{21}}$ or $2mp^{ny_{11}+(m-1)y_{21}+y_{22}}$, or their sum (if the degrees are the same). However, in either case, we find that the coefficient of the least degree term of the LHS is $\leq n + 2m < 2(n+m)$. This implies that Eq. (21) is not true. So, our assumption that f(p) = 0 for all $p \in \mathbb{R}$ must be false.

Proof of Claim 81. For convenience, we will define the polynomials t_1, t_2, t_3 such that

$$t_1(p) = (p^{y_{11}} + p^{y_{12}} + p^{y_{13}} + p^{y_{14}}),$$

$$t_2(p) = (\alpha_1 p^{y_{21}} + \alpha_2 p^{y_{22}} + \alpha_3 p^{y_{23}}), \text{ and}$$

$$t_3(p) = (\beta_1 p^{y_{31}} + \beta_2 p^{y_{32}} + \beta_3 p^{y_{33}}).$$

Then, we see that $f(p) = t_1(p)^{n+m} - t_2(p)^n \cdot t_3(p)^m$.

Let us assume that f(p) = 0 for all $p \in \mathbb{R}$. This means that

$$t_1(p)^{n+m} = t_2(p)^n \cdot t_3(p)^m$$

for all $p \in \mathbb{R}$. We see that the coefficient of the least degree term of the LHS is 1. However, if $\alpha_1 = 2$, or $\beta_1 = 2$, the coefficient of the least degree term of the LHS would be either 2^n or 2^m or 2^{n+m} . Since this is not possible, as n, m > 0, we may conclude that $\alpha_1 = \beta_1 = 1$. Similarly, by reasoning about the highest degree terms of the LHS and the RHS, we can see that $\alpha_3 = \beta_3 = 1$. This means that $\alpha_2 = \beta_2 = 2$.

We note that the least degree term of the LHS is $p^{(n+m)y_{11}}$, and the least degree term of the RHS is $p^{ny_{21}+my_{31}}$. These terms must therefore be equal. This also means that

$$t_1(p)^{n+m} - p^{(n+m)y_{11}} = t_2(p)^n t_3(p)^m - p^{ny_{21}+my_{31}}.$$
(22)

The least degree term of the LHS of Eq. (22) will now be $(n+m)p^{(n+m-1)y_{11}+y_{12}}$, while the least degree term of the RHS will be either $2np^{(n-1)y_{21}+y_{22}+my_{31}}$ or $2mp^{ny_{21}+(m-1)y_{31}+y_{32}}$, or their sum (if the degrees are the same). We observe the coefficients cannot be the same if the degrees are the same, since 2n+2m > n+m. So, the only remaining possibility is that 2n = n+m, or 2m = n+m. In either case, we see that unless n = m, it is not possible that f(p) = 0 for all $p \in \mathbb{R}$.

We are finally ready to prove that if M is of Form (I) and not isomorphic to a tensor product then Pl-GH(M) is #P-hard.

Lemma 82. Let $M \in Sym_4^{pd}(\mathbb{R}_{>0})$ be of Form (I) such that $\rho_{tensor}(M) \neq 0$. Then, Pl-GH(M) is #P-hard.

Proof. Let $(\lambda_1, \ldots, \lambda_4)$ be the eigenvalues of M. Let \mathcal{B} be a lattice basis of the lattice $\mathcal{L}(\lambda_1, \ldots, \lambda_4)$. Let us assume that $c \cdot \delta_{ij} \in \mathcal{B}$ for some $c \in \mathbb{Z}$, and some $i \neq j \in [4]$. Being part of a basis, $c \neq 0$. This implies that $(\lambda_i)^c (\lambda_j)^{-c} = 1$. Since $(\lambda_1, \ldots, \lambda_4)$ are all positive, this implies that $\lambda_i \lambda_j^{-1} = 1$. In other words, we see that $\delta_{ij} \in \mathcal{L}(\lambda_1, \ldots, \lambda_4)$. Since \mathcal{B} is a lattice basis, we have $c = \pm 1$. So, we may replace all such $c \cdot \delta_{ij}$ with δ_{ij} in \mathcal{B} and still have a lattice basis.

Now, let us assume that there exists some $\mathbf{x} \in \mathcal{B} \setminus \mathcal{D}$. (Recall that \mathcal{D} is defined in Eq. (11).) As $\mathbf{0} \neq \mathbf{x} \in \chi_4$, we know that the support of this \mathbf{x} must be greater than 2. Now, for each $\sigma \in S_4$, we can construct \mathbf{x}^{σ} such that $(\mathbf{x}^{\sigma})_i = x_{\sigma(i)}$. We will let $S_4 = \{\sigma_1, \ldots, \sigma_{24}\}$, and define $\xi_i : N \mapsto \phi_{\mathbf{x}^{\sigma_i}}(N_{11}, \ldots, N_{44})$ for $i \in [24]$. We can now use Lemma 78 to find $M_1 \in \mathfrak{R}(M, \mathcal{F}_M \cup \{\rho_{\mathsf{tensor}}, \xi_1\}) \cap \operatorname{Sym}_4^{\mathsf{pd}}(\mathbb{R}_{>0})$ where \mathcal{F}_M is as defined in Eq. (15). We can now repeat this process with M_1 in place of M to find $M_2 \in \mathfrak{R}(M_1, \mathcal{F}_M \cup \{\rho_{\mathsf{tensor}}, \xi_1, \xi_2\}) \cap \operatorname{Sym}_4^{\mathsf{pd}}(\mathbb{R}_{>0})$.

After repeating this for all $i \in [24]$, we can find $M' = M_{24} \in \mathfrak{R}(M, \mathcal{F}_M \cup \{\rho_{\texttt{tensor}}, \xi_1, \ldots, \xi_{24}\}) \cap$ $\operatorname{Sym}_4^{pd}(\mathbb{R}_{>0})$. So, we see that $M' \in \mathfrak{R}(M, \mathcal{F}_M \cup \{\rho_{\texttt{tensor}}, \xi\}) \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{>0})$, where $\xi : N \mapsto \Phi_{\mathbf{x}}(N_{11}, \ldots, N_{44})$. Now, Theorem 34 implies that we can find some $N_1 \in \mathfrak{R}(M, \mathcal{F}_M \cup \{\rho_{\texttt{tensor}}, \Psi_{\mathbf{x}}\}) \cap$ $\operatorname{Sym}_4^{pd}(\mathbb{R}_{>0})$.

If the eigenvalues of N_1 have no lattice basis $\mathcal{B} \subseteq \mathcal{D}$, we can now repeat this whole process with N_1 instead of M. From Lemma 26, we know that after repeating this process at most 4 times, we will have some $N \in \text{Sym}_4^{\text{pd}}(\mathbb{R}_{>0})$, such that $\text{Pl-GH}(N) \leq \text{Pl-GH}(M)$, and the eigenvalues of N have a lattice basis $\mathcal{B} \subseteq \mathcal{D}$. Now, Theorem 18 proves that $\text{Pl-GH}(N) \leq \text{Pl-GH}(M)$ is #P-hard.

11.2 Form (IV)

We will postpone our treatment of Form (III) for a while, and deal with matrices of Form (IV) first.

$$\begin{pmatrix} a & x & x & z \\ x & a & x & z \\ x & x & a & z \\ z & z & z & b \end{pmatrix}$$

Figure 8: Form (IV)

Lemma 83. Let $M \in Sym_4^{pd}(\mathbb{R}_{>0})$ be of Form (IV) such that $\rho_{tensor}(M) \neq 0$. Let \mathcal{F} be a countable set of $Sym_4(\mathbb{R})$ -polynomials such that $F(M) \neq 0$ for all $F \in \mathcal{F}$. Let $\mathbf{0} \neq \mathbf{x} \in \chi_4$ have support size greater than 2. Then, either Pl-GH(M) is #P-hard, or there exists some $N \in \mathfrak{R}(M, \mathcal{F} \cup \{\rho_{tensor}, \Psi_{\mathbf{x}}\}) \cap Sym_4^{pd}(\mathbb{R}_{>0})$ of Form (IV).

Proof. We will first define $\zeta : N \mapsto (N_{11} - N_{44})(N_{22} - N_{44})(N_{33} - N_{44})$. By our choice of M, we know that $\zeta(M) \neq 0$. We will now consider $\mathcal{S}_M(\theta)$. We know that there exists some $\delta > 0$ such that $|\mathcal{S}_M(\theta)_{ij} - I_{ij}| < \frac{1}{3}$ for all $i, j \in [4]$, for all $0 < \theta < \delta$. We can now use Lemma 32 to find some $M' = \mathcal{S}_M(\theta^*) \in \mathfrak{R}(M, \mathcal{F} \cup \{\rho_{\texttt{tensor}}, \zeta\}) \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{\neq 0})$, such that $0 < \theta^* < \delta$.

By construction of M', we know that $M'_{44} \neq M'_{11}, M'_{22}, M'_{33}$. If it is not the case that $M'_{11} = M'_{22} = M'_{33}$, then using Lemma 69, we can find some $M'' \in \mathfrak{R}(M, \mathcal{F} \cup \{\rho_{\texttt{tensor}}\} \cap \operatorname{Sym}_{4}^{\mathsf{pd}})(\mathbb{R}_{>0})$

that is diagonal distinct, or isomorphic to a matrix of Form (I). If M'' is diagonal distinct, then Theorem 68 tells us that Pl-GH(M'') and therefore Pl-GH(M) are #P-hard, If it is isomorphic to a matrix of Form (I), then Lemma 82 similarly implies that Pl-GH(M) is #P-hard. So, we may assume that $(M')_{11} = (M')_{22} = (M')_{33} \neq (M')_{44}$. Theorem 75 now tells us that unless M' is of Form (IV), Pl-GH(M) is again #P-hard. So, we may assume that M' is of Form (IV).

We will now consider $T_{2n}(M')$ for all $n \ge 1$. We let

$$T_{2n}(M') = \begin{pmatrix} (z_{11})^n & (z_{12})^n & (z_{12})^n & (z_{14})^n \\ (z_{12})^n & (z_{11})^n & (z_{12})^n & (z_{14})^n \\ (z_{12})^n & (z_{12})^n & (z_{11})^n & (z_{14})^n \\ (z_{14})^n & (z_{14})^n & (z_{14})^n & (z_{44})^n \end{pmatrix}$$

where $z_{11} = (M'_{11})^2$, $z_{12} = (M'_{12})^2$, $z_{14} = (M'_{14})^2$, and $z_{44} = (M'_{44})^2$. As squares of non-zero numbers they are all positive. It can be verified that the eigenvalues of $T_{2n}(M')$ are (with multiplicity):

$$\lambda_{1}(T_{2n}(M')) = (z_{11})^{n} - (z_{12})^{n},$$

$$\lambda_{2}(T_{2n}(M')) = (z_{11})^{n} - (z_{12})^{n},$$

$$\lambda_{3}(T_{2n}(M')) = \frac{1}{2} \left((z_{11})^{n} + 2(z_{12})^{n} + (z_{44})^{n} - \sqrt{((z_{11})^{n} + 2(z_{12})^{n} - (z_{44})^{n})^{2} + 12((z_{14})^{n})^{2}} \right),$$

$$\lambda_{4}(T_{2n}(M')) = \frac{1}{2} \left((z_{11})^{n} + 2(z_{12})^{n} + (z_{44})^{n} + \sqrt{((z_{11})^{n} + 2(z_{12})^{n} - (z_{44})^{n})^{2} + 12((z_{14})^{n})^{2}} \right).$$

(23)

From our construction of M', we know that $z_{11} > z_{12}$. So, we can see that

$$\lim_{n \to \infty} \frac{\lambda_1(T_{2n}M')}{(z_{11})^n} = \lim_{n \to \infty} \frac{\lambda_2(T_{2n}M')}{(z_{11})^n} = \lim_{n \to \infty} \frac{(z_{11})^n - (z_{12})^n}{(z_{11})^n} = 1$$

We also note that by our construction of M', $z_{11} \neq z_{44}$. We shall prove the following claim shortly: Claim 84. If $z_{11} > z_{44}$, then

$$\lim_{n \to \infty} \frac{\lambda_3(T_{2n}M')}{(z_{44})^n} = \lim_{n \to \infty} \frac{\lambda_4(T_{2n}M')}{(z_{11})^n} = 1.$$

If $z_{11} < z_{44}$, then

$$\lim_{n \to \infty} \frac{\lambda_3(T_{2n}M')}{(z_{11})^n} = \lim_{n \to \infty} \frac{\lambda_4(T_{2n}M')}{(z_{44})^n} = 1.$$

So, we see that for large enough values of n, $\lambda_i(T_{2n}M') > 0$ for all $i \in [4]$. We can now define the function $\varphi_{\mathbf{x}} : (\mathbb{R}_{\neq 0})^4 \to \mathbb{R}$, such that $\varphi_{\mathbf{x}}(\alpha_1, \ldots, \alpha_4) = \prod_{i \in [4]} (\alpha_i)^{x_i}$. For large enough values of n, we see that $\varphi_{\mathbf{x}}(\lambda_1(T_{2n}M'), \ldots, \lambda_4(T_{2n}M'))$ is well-defined. Moreover, we note that

$$\phi_{\mathbf{x}}(\lambda_1(T_{2n}M'),\ldots,\lambda_4(T_{2n}M'))=0\iff \varphi_{\mathbf{x}}(\lambda_1(T_{2n}M'),\ldots,\lambda_4(T_{2n}M'))=1.$$

We will now study the behavior of the function $\phi_{\mathbf{x}}(\lambda_1(T_{2n}M'),\ldots,\lambda_4(T_{2n}M'))$, by studying the function $\varphi_{\mathbf{x}}(\lambda_1(T_{2n}M'),\ldots,\lambda_4(T_{2n}M'))$.

We first note from Eq. (23) that $\lambda_1(T_{2n}M') = \lambda_2(T_{2n}M')$ for all $n \geq 1$. We also note that $\lambda_4(T_{2n}M') > \lambda_1(T_{2n}M'), \lambda_3(T_{2n}M')$ for large enough n. If $\lambda_1(T_{2n}M') = \lambda_3(T_{2n}M')$, that implies that

$$(z_{44})^n - (z_{11})^n + 4(z_{12})^n = \sqrt{((z_{11})^n + 2(z_{12})^n - (z_{44})^n)^2 + 12((z_{14})^n)^2}.$$

Squaring both sides, we find that

$$((z_{44})^n - (z_{11})^n)^2 + 16(z_{12})^{2n} + 8(z_{12})^n ((z_{44})^n - (z_{11})^n) = ((z_{44})^n - (z_{11})^n)^2 + 4(z_{12})^{2n} - 4(z_{12})^n ((z_{44})^n - (z_{11})^n) + 12(z_{14})^{2n}.$$

On rearranging terms, we find that

$$12(z_{12})^{2n} + 12(z_{12}z_{44})^n = 12(z_{14})^{2n} + 12(z_{12}z_{11})^n$$

If we write this as $A^n + B^n = C^n + D^n$, where $A = (z_{12})^2$, $B = z_{12}z_{44}$, $C = (z_{14})^2$ and $D = z_{12}z_{11}$, then we know that A < D and $B \neq D$. (1) If $C \geq D$, then the RHS has order C^n . As $A < D \leq C$, to match the leading order, B = C (in this case, C = D is impossible). But then $A^n = D^n$, contradicting A < D. (2) If C < D, then the RHS has order D^n . But this cannot be since A < D and $B \neq D$. So, this equation cannot be true for large enough values of n. So, we see that $\lambda_1(T_{2n}M') = \lambda_2(T_{2n}M')$ and $|\lambda_1(T_{2n}M')| \neq |\lambda_3(T_{2n}M')| \neq |\lambda_4(T_{2n}M')|$ are pairwise distinct for large values of n. So, Claim 79 immediately implies that if $x_3 = 0$, or $x_4 = 0$, or $x_1 + x_2 = 0$, then $\phi_{\mathbf{x}}(\lambda_1(T_{2n}M'), \dots, \lambda_4(T_{2n}M')) \neq 0$ for large enough n.

We will now assume that $x_3 \neq 0$, and $x_4 \neq 0$. Now, we find that when $z_{11} > z_{44}$,

$$\varphi_{\mathbf{x}}(\lambda_1(T_{2n}M'),\ldots,\lambda_4(T_{2n}M')) = \left((z_{11})^{n(x_1+x_2+x_4)}(z_{44})^{n(x_3)}\right) \cdot \left(\frac{\lambda_1(T_{2n}M')}{(z_{11})^n}\right)^{x_1} \cdot \left(\frac{\lambda_2(T_{2n}M')}{(z_{11})^n}\right)^{x_2} \cdot \left(\frac{\lambda_3(T_{2n}M')}{(z_{44})^n}\right)^{x_3} \cdot \left(\frac{\lambda_4(T_{2n}M')}{(z_{11})^n}\right)^{x_4}$$

So, we see that

$$\lim_{n \to \infty} \varphi_{\mathbf{x}}(\lambda_1(T_{2n}M'), \dots, \lambda_4(T_{2n}M')) = \lim_{n \to \infty} \left(\frac{z_{44}}{z_{11}}\right)^{nx_5}$$

which is either 0 or ∞ depending on whether $x_3 > 0$, or $x_3 < 0$. Similarly, when $z_{11} < z_{44}$, we find that

$$\varphi_{\mathbf{x}}(\lambda_1(T_{2n}M'),\dots,\lambda_4(T_{2n}M')) = \left((z_{11})^{n(x_1+x_2+x_3)}(z_{44})^{n(x_4)}\right) \cdot \left(\frac{\lambda_1(T_{2n}M')}{(z_{11})^n}\right)^{x_1} \cdot \left(\frac{\lambda_2(T_{2n}M')}{(z_{11})^n}\right)^{x_2} \cdot \left(\frac{\lambda_3(T_{2n}M')}{(z_{11})^n}\right)^{x_3} \cdot \left(\frac{\lambda_4(T_{2n}M')}{(z_{44})^n}\right)^{x_4}$$

So, we see that

$$\lim_{n \to \infty} \varphi_{\mathbf{x}}(\lambda_1(T_{2n}M'), \dots, \lambda_4(T_{2n}M')) = \lim_{n \to \infty} \left(\frac{z_{44}}{z_{11}}\right)^{nx_4}$$

which is either 0 or ∞ depending on whether $x_4 < 0$, or $x_4 > 0$.
So, we see that in either case, $\lim_{n\to\infty} \varphi_{\mathbf{x}}(\lambda_1(T_{2n}M'),\ldots,\lambda_4(T_{2n}M'))$ is either 0 or ∞ , and so it is away from 1. This proves that for all $\mathbf{x} \in \chi_4$ of support size greater than 2, for large enough $n \geq 1, \ \phi_{\mathbf{x}}(\lambda_1(T_{2n}M'),\ldots,\lambda_4(T_{2n}M')) \neq 0$. In particular, given any $\sigma \in S_4$, we can see that $\phi_{\mathbf{y}}(\lambda_1(T_{2n}M'),\ldots,\lambda_4(T_{2n}M')) \neq 0$ for all \mathbf{y} such that $y_{\sigma(i)} = x_i$, for some large enough value of n. This proves there exists some n^* , such that

$$\Phi_{\mathbf{x}}(\lambda_1(T_{2n^*}M'),\ldots,\lambda_4(T_{2n^*}M')) = \Psi_{\mathbf{x}}(T_{2n^*}M') \neq 0.$$

We will now define the $\operatorname{Sym}_4(\mathbb{R})$ -polynomial $\xi : N \mapsto \Psi_{\mathbf{x}}(T_{2n^*}N)$. Since $M' = \mathcal{S}_M(\theta^*)$ by construction, we see that $\xi(\mathcal{S}_M(\theta^*)) \neq 0$. So, we can use Corollary 33 to find some $M'' \in \mathfrak{R}(M, \mathcal{F} \cup \{\rho_{\texttt{tensor}}, \zeta, \xi\}) \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{>0})$. We now see that $\Psi_{\mathbf{x}}(T_{2n^*}M'') = \xi(M'') \neq 0$. So, Corollary 29 allows us to find the required $N \in \mathfrak{R}(M, \mathcal{F} \cup \{\rho_{\texttt{tensor}}, \zeta, \Psi_{\mathbf{x}}\}) \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{>0})$.

Proof of Claim 84. First, we note that $\lambda_3(T_{2n}M')$ and $\lambda_4(T_{2n}M')$ are symmetric if we exchange z_{11} and z_{44} . So, it will be sufficient for us to prove this claim when $z_{11} > z_{44}$. We note that

$$(z_{11})^n + 2(z_{12})^n + (z_{44})^n = (z_{11})^n \left(1 + 2\left(\frac{z_{12}}{z_{11}}\right)^n + \left(\frac{z_{44}}{z_{11}}\right)^n\right).$$

So,

$$\lim_{n \to \infty} \frac{(z_{11})^n + 2(z_{12})^n + (z_{44})^n}{(z_{11})^n} = 1.$$

We also note that

$$\sqrt{((z_{11})^n + 2(z_{12})^n - (z_{44})^n)^2 + 12((z_{14})^n)^2} = (z_{11})^n \cdot \sqrt{\left(1 + 2\left(\frac{z_{12}}{z_{11}}\right)^n - \left(\frac{z_{44}}{z_{11}}\right)^n\right)^2 + 12\left(\left(\frac{z_{14}}{z_{11}}\right)^n\right)^2}.$$

So,

$$\lim_{n \to \infty} \frac{\sqrt{((z_{11})^n + 2(z_{12})^n - (z_{44})^n)^2 + 12((z_{14})^n)^2}}{(z_{11})^n} = 1$$

This implies that

$$\lim_{n \to \infty} \frac{\lambda_4(T_2 M')}{(z_{11})^n} = 1.$$

We now note that

$$\begin{split} \lambda_{3}(T_{2n}M') &= \frac{1}{2} \left(\frac{((z_{11})^{n} + 2(z_{12})^{n} + (z_{44})^{n})^{2} - ((z_{11})^{n} + 2(z_{12})^{n} - (z_{44})^{n})^{2} - 12((z_{14})^{n})^{2}}{(z_{11})^{n} + 2(z_{12})^{n} + (z_{44})^{n} + \sqrt{((z_{11})^{n} + 2(z_{12})^{n} - (z_{44})^{n})^{2} + 12((z_{14})^{n})^{2}}} \right) \\ &= \frac{1}{2} \left(\frac{4(z_{44})^{n}((z_{11})^{n} + (2z_{12})^{n}) - 12(z_{14})^{2n}}{(z_{11})^{n} + 2(z_{12})^{n} + (z_{44})^{n} + \sqrt{((z_{11})^{n} + 2(z_{12})^{n} - (z_{44})^{n})^{2} + 12((z_{14})^{n})^{2}}} \right) \\ &= \frac{(z_{44})^{n}}{2} \left(\frac{4 + 8\left(\frac{z_{12}}{z_{11}}\right)^{n} - 12\left(\frac{(z_{14})^{2}}{z_{44}z_{11}}\right)^{n}}{1 + 2\left(\frac{z_{12}}{z_{11}}\right)^{n} + \left(\frac{z_{44}}{z_{11}}\right)^{n} + \sqrt{\left(1 + 2\left(\frac{z_{12}}{z_{11}}\right)^{n} - \left(\frac{z_{44}}{z_{11}}\right)^{n}\right)^{2} + 12\left(\left(\frac{z_{14}}{z_{11}}\right)^{n}\right)^{2}}} \right). \end{split}$$

We already know that

So,

$$\lim_{n \to \infty} \left(1 + 2\left(\frac{z_{12}}{z_{11}}\right)^n + \left(\frac{z_{44}}{z_{11}}\right)^n + \sqrt{\left(1 + 2\left(\frac{z_{12}}{z_{11}}\right)^n - \left(\frac{z_{44}}{z_{11}}\right)^n\right)^2 + 12\left(\left(\frac{z_{14}}{z_{11}}\right)^n\right)^2\right)} = 2.$$
$$\lim_{n \to \infty} \frac{\lambda_3(T_{2n}M')}{(z_{44})^n} = \frac{1}{2}\left(\frac{4}{2}\right) = 1.$$

We can now prove that Pl-GH(M), when M is of Form (IV) that is not isomorphic to a tensor product, must also be #P-hard.

Lemma 85. Let $M \in Sym_4^{pd}(\mathbb{R}_{>0})$ be of Form (IV) such that $\rho_{tensor}(M) \neq 0$. Then, Pl-GH(M) is #P-hard.

Proof. The proof is very similar to the proof of Lemma 82. Let $(\lambda_1, \ldots, \lambda_4)$ be the eigenvalues of M. Let \mathcal{B} be a lattice basis of the lattice $\mathcal{L}(\lambda_1, \ldots, \lambda_4)$. If there is some $c \cdot \delta_{ij} \in \mathcal{B}$ for some non-zero $c \in \mathbb{Z}$, we may replace all such $c \cdot \delta_{ij}$ with δ_{ij} in \mathcal{B} and still have a lattice basis.

Now, let us assume that there exists some $\mathbf{x} \in \mathcal{B} \setminus \mathcal{D}$. We know that the support of this \mathbf{x} must be greater than 2. Unless Pl-GH(M) is #P-hard, we can now use Lemma 83 to find some $N_1 \in \mathfrak{R}(M, \mathcal{F}_M \cup \{\rho_{\texttt{tensor}}, \Psi_{\mathbf{x}}\} \cap \text{Sym}_4^{pd}(\mathbb{R}_{>0})$ of Form (IV).

If the eigenvalues of N_1 have no lattice basis $\mathcal{B} \subseteq \mathcal{D}$, we can now repeat this whole process with N_1 instead of M. From Lemma 26, we know that after repeating this process finitely many times, we will have some $N \in \text{Sym}_4^{\text{pd}}(\mathbb{R}_{>0})$, such that $\text{Pl-GH}(N) \leq \text{Pl-GH}(M)$, and the eigenvalues of N have a lattice basis $\mathcal{B} \subseteq \mathcal{D}$. Now, Theorem 18 proves that $\text{Pl-GH}(N) \leq \text{Pl-GH}(M)$ is #P-hard. \Box

11.3 Form (III)

We will now deal with matrices of Form (III). We will once again need some setup, by proving that the matrix M of Form (III) may be assumed to have some additional structure. Note that by a simultaneous permutation we can permute the rows and columns of M such that $M_{11} = M_{22} > M_{33} = M_{44}$, and that $M_{13} \ge M_{14}$. For the rest of this section, we will assume this is the case.

$$\begin{pmatrix} a & x & y & z \\ x & a & z & y \\ y & z & b & t \\ z & y & t & b \end{pmatrix}$$

Figure 9: Form (III)

Lemma 86. Let $M \in Sym_4^{pd}(\mathbb{R}_{>0})$. Let \mathcal{F} be a countable set of $Sym_4(\mathbb{R})$ -polynomials such that $F(M) \neq 0$ for all $F \in \mathcal{F}$. Then there exists some $N \in \mathfrak{R}(M, \mathcal{F}) \cap Sym_4^{pd}(\mathbb{R}_{>0})$, such that $N_{ii} > N_{jk}$ for all $i \in [4]$, and $j \neq k \in [4]$.

Proof. We may assume that the entries of M are generated by some $\{g_t\}_{t\in[d]}$. We can use Lemma 4 to replace M with some other cM such that $e_{ijt} \geq 0$ for all $i, j \in [4], t \in [d]$. We will now let $\mathcal{F}' = \{F' : N \mapsto F(T_2N) : F \in \mathcal{F}\}$. We note that since $F(M) \neq 0$ for all $F \in \mathcal{F}, F'(\mathcal{T}_M(g_1^{1/2}, \ldots, g_d^{1/2})) \neq 0$ for all $F' \in \mathcal{F}'$. So, Lemma 28 allows us to pick some $M' \in \mathfrak{R}(M, \mathcal{F}') \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{>0})$.

We will now consider $\mathcal{S}_{M'}(\theta)$. We know that there exists some $\delta > 0$ such that $|\mathcal{S}_{M'}(\theta)_{ij} - I_{ij}| < \frac{1}{3}$ for all $i, j \in [4]$, for all $0 < \theta < \delta$. So, we can use Lemma 32 to find some $M'' = \mathcal{S}_{M'}(\theta^*) \in \mathfrak{R}(M', \mathcal{F}') \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{\neq 0})$, such that $0 < \theta^* < \delta$.

We will now let $N = T_2(M'')$. Clearly, $N \in \text{Sym}_4(\mathbb{R}_{>0})$. By our choice of M'', we know that $N_{ii} > \frac{4}{9}$, and $N_{jk} < \frac{1}{9}$ for all $i \in [4]$, and $j \neq k \in [4]$. By the **Gershgorin Circle Theorem** [Ger31], we know that if we let $r_i = \sum_{j \in [4] \setminus \{i\}} |N_{ij}|$, then all the eigenvalues of N lie within one of the intervals $[N_{ii} - r_i, N_{ii} + r_i]$. But as we have seen, $N_{ii} - r_i > (\frac{4}{9} - 3 \cdot \frac{1}{9}) > 0$ for all $i \in [4]$. This implies that all eigenvalues of N are positive. So, $N \in \text{Sym}_4^{\text{pd}}(\mathbb{R}_{>0})$. Moreover, by our choice of \mathcal{F}' , we know that since $F'(M'') \neq 0$ for all $F' \in \mathcal{F}'$, $F(N) \neq 0$ for all $F \in \mathcal{F}$. So, $N \in \mathfrak{R}(M, \mathcal{F}) \cap \text{Sym}_4^{\text{pd}}(\mathbb{R}_{>0})$ is the required matrix.

As it turns out, it is possible for matrices of Form (III) to be isomorphic to $A \otimes B$ for some $A, B \in \text{Sym}_2(\mathbb{R})$. We will now show that if $\rho_{\texttt{tensor}}(M) \neq 0$, then M has sufficient structure, that we can prove that Pl-GH(M) is #P-hard.

Lemma 87. Let $M \in Sym_4^{pd}(\mathbb{R}_{>0})$ be of Form (III) such that $\rho_{tensor}(M) \neq 0$. Let \mathcal{F} be a countable set of $Sym_4(\mathbb{R})$ -polynomials such that $F(M) \neq 0$ for all $F \in \mathcal{F}$. Then, either Pl-GH(M) is #P-hard, or there exists some $N \in \mathfrak{R}(M, \mathcal{F} \cup \{\rho_{tensor}\}) \cap Sym_4^{pd}(\mathbb{R}_{>0})$ of Form (III), such that $N_{12} \neq N_{34}$.

Proof. If $M_{12} \neq M_{34}$, we are already done, so we may assume otherwise. We will let $\zeta : N \mapsto (N_{11} - N_{33})(N_{11} - N_{44})(N_{22} - N_{33})(N_{22} - N_{44})$. We know that $\zeta(M) \neq 0$. We will let $\mathcal{F}' = \{F' : N \mapsto F(N^3) \mid F \in \mathcal{F} \cup \{\rho_{\mathsf{tensor}}, \zeta\}\}$. We also know that there exists some $\delta > 0$ such that for all $0 < \theta < \delta$, $|\mathcal{S}_M(\theta)_{ij} - I_{ij}| < \frac{1}{3}$. We can now use Lemma 32 to find $M' = \mathcal{S}_M(\theta^*) \in \mathfrak{R}(M, \mathcal{F} \cup \mathcal{F}' \cup \{\zeta, \rho_{\mathsf{tensor}}\}) \cap \operatorname{Sym}_4^{\mathsf{pd}}(\mathbb{R}_{>0})$, for some $0 < \theta^* < \delta$. If M' is not of Form (III), that would imply that it is either diagonal distinct, or isomorphic to a matrix of Form (I) or Form (II), or satisfies $M'_{11} = M'_{22} \neq M'_{33} = M'_{44}$, but is not of Form (III). In either case, since $\rho_{\mathsf{tensor}}(M') \neq 0$, we see that P1-GH(M') \leq P1-GH(M) is #P-hard, due to either Theorem 68, or Lemma 73 and Lemma 82, or Lemma 70. So, we may assume that M' is of Form (III).

We will now consider $(R_n M')_{12} - (R_n M')_{34}$ for all $n \ge 1$. We let $X = \{(M')_{ab} : a, b \in [4]\}$, and $c_{ij}(x) = \sum_{a,b \in [4]: M'_{ab} = x} (M')_{ia} (M')_{jb}$. We see that

$$(R_n M')_{12} - (R_n M')_{34} = \sum_{x \in X} x^n \cdot (c_{12}(x) - c_{34}(x))$$

for all $n \ge 1$. So, the equations $(R_n M')_{12} - (R_n M')_{34} = 0$ form a full rank Vandermonde system of equations of size O(1). This implies that $c_{12}(x) - c_{34}(x) = 0$ for all $x \in X$. By construction of M', we know that $(M')_{ab} = (M')_{11}$ implies that $(a, b) \in \{(1, 1), (2, 2)\}$, and that $(M')_{ab} = (M'')_{33}$ implies that $(a, b) = \{(3, 3), (4, 4)\}$. So, we see that (using the fact that M' has Form (III))

$$c_{12}((M')_{11}) - c_{34}((M')_{11}) = 2(M')_{11}(M')_{12} - 2(M')_{13}(M')_{14} = 0$$
, and

$$c_{12}((M')_{33}) - c_{34}((M')_{33}) = 2(M')_{13}(M')_{14} - 2(M')_{33}(M')_{34} = 0.$$

Hence, $(M')_{11}(M')_{12} = (M')_{33}(M')_{34}$. Since we know that $(M')_{11} \neq (M')_{33}$, this implies that $(M')_{12} \neq (M')_{34}$.

So, if $(R_nM')_{12} - (R_nM')_{34} = 0$ for all $n \ge 1$, then $M' \in \mathfrak{R}(M, \mathcal{F} \cup \{\rho_{\texttt{tensor}}\}) \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{>0})$ is the required matrix. On the other hand, suppose there exists some $n \ge 1$ such that $(R_nM')_{12} - (R_nM')_{34} \ne 0$. In that case, we will construct $\xi' : N \mapsto (R_nN)_{12} - (R_nN)_{34}$. We see that $\xi'(\mathcal{S}_M(\theta^*)) \ne 0$. So, Corollary 33 lets us find some $M'' \in \mathfrak{R}(M, \mathcal{F}' \cup \{\xi'\}) \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{>0})$. Now, if we let $\xi : N \mapsto N_{12} - N_{34}$, we see that $\xi(\mathcal{R}_{M''}(n)) \ne 0$. We also see that $F(\mathcal{R}_{M''}(1)) = F((M'')^3) \ne 0$ for all $F \in \mathcal{F} \cup \{\rho_{\texttt{tensor}}, \zeta\}$. So, we can use Lemma 57 to find some $N \in \mathfrak{R}(M, \mathcal{F} \cup \{\rho_{\texttt{tensor}}, \zeta, \xi\}) \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{>0})$.

Since $\zeta(N) \neq 0$, if N is not of Form (III), then it must either be diagonal distinct, or isomorphic to a matrix of Form (I) or Form (II), or satisfies $N_{11} = N_{22} \neq N_{33} = N_{44}$, but is not of Form (III). In any case, since $\rho_{\texttt{tensor}}(N) \neq 0$, we see that $\texttt{Pl-GH}(N) \leq \texttt{Pl-GH}(M)$ is #P-hard, due to either **Theorem 68**, or Lemma 73 and Lemma 82, or Lemma 70. On the other hand, if N is of Form (III), it is the required matrix.

Lemma 88. Let $M \in Sym_4^{pd}(\mathbb{R}_{>0})$ be of Form (III) such that $\rho_{tensor}(M) \neq 0$. Let \mathcal{F} be a countable set of $Sym_4(\mathbb{R})$ -polynomials such that $F(M) \neq 0$ for all $F \in \mathcal{F}$. Then, either Pl-GH(M) is #P-hard, or there exists some $N \in \mathfrak{R}(M, \mathcal{F} \cup \{\rho_{tensor}\}) \cap Sym_4^{pd}(\mathbb{R}_{>0})$ of Form (III), such that $N_{11}N_{33}N_{12}N_{34} - (N_{13}N_{14})^2 \neq 0$.

Proof. We let $\xi : N \mapsto (N_{11}N_{33}N_{12}N_{34}) - (N_{13}N_{14})^2$. If $\xi(M) \neq 0$, we are already done, so we may assume otherwise. We also let $\zeta : N \mapsto (N_{11} - N_{33})(N_{11} - N_{44})(N_{22} - N_{33})(N_{22} - N_{44})$. We note that $\zeta(M) \neq 0$. We will first use Lemma 87 to find $M' \in \Re(M, \mathcal{F} \cup \{\rho_{\text{tensor}}, \zeta\}) \cap \text{Sym}_4^{\text{pd}}(\mathbb{R}_{>0})$ of Form (III) such that $(M')_{12} \neq (M')_{34}$. We can now use Lemma 86 to find $M'' \in \Re(M', \mathcal{F} \cup \{\rho_{\text{tensor}}, \zeta\}) \cap$ $\text{Sym}_4^{\text{pd}}(\mathbb{R}_{>0})$, such that $(M'')_{ii} > (M'')_{jk}$ for all $i \in [4]$, and $j \neq k \in [4]$. Since $\zeta(M'') \neq 0$, if M'' is not of Form (III), then it must either be diagonal distinct, or isomorphic to a matrix of Form (I) or Form (II), or satisfies $(M'')_{11} = (M'')_{22} \neq (M'')_{33} = (M'')_{44}$, but is not of Form (III). In any case, since $\rho_{\text{tensor}}(M'') \neq 0$, we see that P1-GH $(N) \leq$ P1-GH(M) is #P-hard, due to either Theorem 68, or Lemma 73 and Lemma 82, or Lemma 70. On the other hand, we may assume that M'' is of Form (III). For convenience, we may rename this M'' as M. We note that we can also simultaneously permute the rows and columns of M (first by a possible switch $\{1, 2\} \leftrightarrow \{3, 4\}$ and then a possible flip $3 \leftrightarrow 4$) such that $M_{11} > M_{33}$, and $M_{13} \geq M_{14}$ without loss of generality.

We may assume that the entries of M are generated by some $\{g_t\}_{t\in[d]}$. We can also replace this M with some cM as guaranteed by Lemma 4 such that $e_{ijt} \ge 0$ for all $i, j \in [4]$, and $t \in [d]$. We will now define a function $\widehat{\mathcal{T}_M} : \mathbb{R}^{d+1} \to \operatorname{Sym}_4(\mathbb{R})$ such that

$$\widehat{\mathcal{T}}_M(p, z_1, \dots, z_d)_{ij} = \mathcal{T}_M(p^{z_1}, \dots, p^{z_d})_{ij}$$
(24)

for all $i, j \in [4]$. By our choice of M, we know that $\widehat{\mathcal{T}}_M(e, \log(g_1), \ldots, \log(g_d)) = M$. We note that $\widehat{\mathcal{T}}_M(p, z_1, \ldots, z_d)_{ij}$ is continuous as a function of all its variables, for all $i, j \in [4]$. So, there exist non-empty intervals $I_1, \ldots, I_d \subseteq \mathbb{R}_{>0}$ such that for all $(z_1, \ldots, z_d) \in I_1 \times \cdots \times I_d$, $\widehat{\mathcal{T}}_M(e, z_1, \ldots, z_d)$

also satisfies the properties of M that

$$M_{11} > M_{33}, M_{13} \ge M_{14}, M_{12} \ne M_{34}, \text{ and } M_{ii} > M_{jk}$$

for all $i \in [4]$, for $j \neq k \in [4]$. But now, we can pick some $(z_1^*, \ldots, z_d^*) \in \mathbb{Q}^d \cap (I_1 \times \cdots \times I_d)$. Since $z_t^* > 0$ for all $t \in [d]$, there exists some $Z^* \in \mathbb{Z}_{>0}$, such that $Z \cdot z_t^* \in \mathbb{Z}_{>0}$ for all $t \in [d]$. We will now define $\mathcal{T}_M^* : \mathbb{R} \to \text{Sym}_4(\mathbb{R})$ such that

$$\mathcal{T}_M^*(p)_{ij} = \widehat{\mathcal{T}_M}(p, Z^* \cdot z_i^*, \dots, Z^* \cdot z_d^*)_{ij} = p^{z_{ij}}$$

$$\tag{25}$$

for some $z_{ij} = Z^* \sum_{t \in [d]} e_{ijt} z_t^* \in \mathbb{Z}_{\geq 0}$ for all $i, j \in [4]$. We also see that since $\mathcal{T}_M^*(e)_{ii} > \mathcal{T}_M^*(e)_{jk}$ for all $i \in [4]$, and $j \neq k \in [4]$, it must be the case that $z_{ii} > z_{jk}$ for all $i \in [4]$, and $j \neq k \in [4]$. If $\xi(\mathcal{T}_M^*(p)) \neq 0$ for some $p \in \mathbb{R}$, then from the construction of \mathcal{T}_M^* , we see that there must exist some $\mathbf{p} \in \mathbb{R}^d$ such that $\xi(\mathcal{T}_M(\mathbf{p})) = \xi(\mathcal{T}_M^*(p)) \neq 0$. Therefore, Lemma 28 allows us to find some $N \in \mathfrak{R}(M, \mathcal{F} \cup \{\rho_{\mathsf{tensor}}, \zeta, \xi\}) \cap \operatorname{Sym}_4^{\mathsf{pd}}(\mathbb{R}_{>0})$. Since $\zeta(N) \neq 0$, if N is not of Form (III), it must either be diagonal distinct, or isomorphic to a matrix of Form (I) or (II), or it should satisfy the equation $N_{11} = N_{22} \neq N_{33} = N_{44}$, but not be of Form (III). In any of these cases, since $\rho_{\mathsf{tensor}}(N) \neq 0$, it follows from Theorem 68, or Lemma 73 and Lemma 82, or Lemma 70, that P1-GH(N) \leq P1-GH(M) is #P-hard. On the other hand, if N is of Form (III), then we note that it is the required matrix, and we are done.

We may now assume that $\xi(\mathcal{T}_M^*(p)) = 0$ for all $p \in \mathbb{R}$. This means that $p^{z_{11}+z_{33}+z_{12}+z_{34}} = p^{2z_{13}+2z_{14}}$ for all $p \in \mathbb{R}$. So, we see that

$$z_{13} + z_{14} = \frac{(z_{11} + z_{12}) + (z_{33} + z_{34})}{2},$$
(26)

i.e., $z_{13} + z_{14}$ is the average of the other two sums. In particular, we have three possibilities,

Case 0. $z_{11} + z_{12} = z_{13} + z_{14} = z_{33} + z_{34}$; or

Case 1. $z_{11} + z_{12} > z_{13} + z_{14} > z_{33} + z_{34}$; or

Case 2. $z_{11} + z_{12} < z_{13} + z_{14} < z_{33} + z_{34}$.

We will now consider

$$\xi((\mathcal{T}_{M}^{*}(p))^{2}) = ((\mathcal{T}_{M}^{*}(p))^{2})_{11}((\mathcal{T}_{M}^{*}(p))^{2})_{33}((\mathcal{T}_{M}^{*}(p))^{2})_{12}((\mathcal{T}_{M}^{*}(p))^{2})_{34} - (((\mathcal{T}_{M}^{*}(p))^{2})_{13}((\mathcal{T}_{M}^{*}(p))^{2})_{14})^{2}.$$

We will show that this function in p is not identically 0. For a contradiction, we assume that $\xi(((\mathcal{T}_M^*(p))^2)) = 0$ for all $p \in \mathbb{R}$. Thus, we have for all $p \in \mathbb{R}$

$$((\mathcal{T}_{M}^{*}(p))^{2})_{11}((\mathcal{T}_{M}^{*}(p))^{2})_{33}((\mathcal{T}_{M}^{*}(p))^{2})_{12}((\mathcal{T}_{M}^{*}(p))^{2})_{34} = \left(((\mathcal{T}_{M}^{*}(p))^{2})_{13}((\mathcal{T}_{M}^{*}(p))^{2})_{14}\right)^{2}$$
(27)

where (using the Form (III))

$$\begin{split} &((\mathcal{T}_{M}^{*}(p))^{2})_{11} = p^{2z_{11}} + p^{2z_{12}} + p^{2z_{13}} + p^{2z_{14}}, \\ &((\mathcal{T}_{M}^{*}(p))^{2})_{33} = p^{2z_{13}} + p^{2z_{14}} + p^{2z_{33}} + p^{2z_{34}}, \\ &((\mathcal{T}_{M}^{*}(p))^{2})_{12} = 2p^{z_{11}+z_{12}} + 2p^{z_{13}+z_{14}}, \\ &((\mathcal{T}_{M}^{*}(p))^{2})_{34} = 2p^{z_{13}+z_{14}} + 2p^{z_{33}+z_{34}}, \\ &((\mathcal{T}_{M}^{*}(p))^{2})_{13} = p^{z_{11}+z_{13}} + p^{z_{12}+z_{14}} + p^{z_{13}+z_{33}} + p^{z_{14}+z_{34}}, \\ &((\mathcal{T}_{M}^{*}(p))^{2})_{14} = p^{z_{11}+z_{14}} + p^{z_{12}+z_{13}} + p^{z_{13}+z_{34}} + p^{z_{14}+z_{33}}. \end{split}$$

Step 1. The leading degree term of the LHS of Eq. (27) is either 4p<sup>3z₁₁+2z₃₃+z₁₂+z₁₃+z₁₄ (in the enumerated Case 1. of Eq. (26)) or 4p<sup>2z₁₁+3z₃₃+z₃₄+z₁₃+z₁₄ (in Case 2.) or has coefficient 16 (in Case 0).
</sup></sup>

By our choice of z_{ij} , we know that $z_{11} > z_{33} > z_{ij}$ for all $i \neq j \in [4]$. So, $p^{2z_{11}}$ is the leading degree term of

$$((\mathcal{T}_M^*(p))^2)_{11} = p^{2z_{11}} + p^{2z_{12}} + p^{2z_{13}} + p^{2z_{14}},$$

and $p^{2z_{33}}$ is the leading degree term of

$$((\mathcal{T}_M^*(p))^2)_{33} = p^{2z_{13}} + p^{2z_{14}} + p^{2z_{33}} + p^{2z_{34}}$$

Now, if the two terms of $((\mathcal{T}_M^*(p))^2)_{12}$ collapse into one term, that implies that $z_{11} + z_{12} = z_{13} + z_{14}$. But then, Eq. (26) implies that $z_{13} + z_{14} = z_{33} + z_{34}$ as well. This would imply that $((\mathcal{T}_M^*(p))^2)_{12} = 4p^{z_{11}+z_{12}} = ((\mathcal{T}_M^*(p))^2)_{34}$. This means that the leading degree term of the LHS of Eq. (27) will be

$$16p^{4z_{11}+2z_{33}+2z_{12}}$$
.

On the other hand, if the two terms of $((\mathcal{T}_{M}^{*}(p))^{2})_{12}$ do not collapse into each other, that means that $z_{11} + z_{12} \neq z_{13} + z_{14}$. From Eq. (26), this implies that in fact, $z_{11} + z_{12} \neq z_{13} + z_{14} \neq z_{33} + z_{34}$ are pairwise distinct. So, it must either be the case that $z_{11} + z_{12} > z_{13} + z_{14} > z_{33} + z_{34}$, or it must be the case that $z_{33} + z_{34} > z_{13} + z_{14} > z_{11} + z_{12}$. Depending on which of the two cases occur, the leading degree term of the LHS of Eq. (27) will be, respectively, either

 $4p^{3z_{11}+2z_{33}+z_{12}+z_{13}+z_{14}}$ or $4p^{2z_{11}+3z_{33}+z_{34}+z_{13}+z_{14}}$.

- Step 2. Case 1. or Case 2. hold in Eq. (26). Case 0. does not. The leading degree term of the RHS of Eq. (27) is 4p<sup>4z₁₁+2z₁₃+2z₁₄. Furthermore, we have
 </sup>
 - (A) $z_{11} + z_{14} = z_{12} + z_{13} > z_{13} + z_{34}$, or
 - (B) $z_{11} + z_{14} = z_{13} + z_{34} > z_{12} + z_{13}$.

and the leading degree term of $((\mathcal{T}^*_M(p))^2)_{14}$ is obtained by combining two terms $p^{z_{11}+z_{14}} + p^{z_{12}+z_{13}}$ in case (A), or $p^{z_{11}+z_{14}} + p^{z_{13}+z_{34}}$ in case (B).

We will now analyze the leading degree term of the RHS of Eq. (27). Let us first focus on the terms of

$$((\mathcal{T}_M^*(p))^2)_{13} = p^{z_{11}+z_{13}} + p^{z_{12}+z_{14}} + p^{z_{13}+z_{33}} + p^{z_{14}+z_{34}}.$$

We note that $z_{11}+z_{13} > z_{12}+z_{14}$, since $z_{11} > z_{12}$, and $z_{13} \ge z_{14}$. Similarly, $z_{11}+z_{13} > z_{13}+z_{33}$, since $z_{11} > z_{33}$. Finally, $z_{11} + z_{13} > z_{14} + z_{34}$. So, the leading degree term of $((\mathcal{T}_M^*(p))^2)_{13}$ is $p^{z_{11}+z_{13}}$ with coefficient 1. Now, we focus on the terms of

$$((\mathcal{T}_M^*(p))^2)_{14} = p^{z_{11}+z_{14}} + p^{z_{12}+z_{13}} + p^{z_{13}+z_{34}} + p^{z_{14}+z_{33}}.$$

Since $z_{11} > z_{33}$, it follows that $z_{11} + z_{14} > z_{14} + z_{33}$. So, the leading term coefficient of $((\mathcal{T}_M^*(p))^2)_{14}$ cannot be 4. Since the square of this coefficient must be 4 or 16 to match that of the LHS of Eq. (27), we see that this coefficient must be 2, and the leading term coefficient of the LHS of Eq. (27) must be 4 (not 16). In particular, this also rules out the possibility that $z_{11} + z_{12} = z_{13} + z_{14} = z_{33} + z_{34}$, i.e., Case 0. in Eq. (26) does not hold, and Case 1. or Case 2. hold in Eq. (26).

Now, in $((\mathcal{T}_M^*(p))^2)_{14}$, we have seen that the leading degree term cannot be $p^{z_{14}+z_{33}}$. Similarly, it is not possible that $z_{12} + z_{13} = z_{13} + z_{34}$, since $z_{12} \neq z_{34}$. So, the leading degree term of $((\mathcal{T}_M^*(p))^2)_{14}$ must either be $p^{z_{11}+z_{14}} + p^{z_{12}+z_{13}}$, or $p^{z_{11}+z_{14}} + p^{z_{13}+z_{34}}$. Thus, either (A) or (B) hold. In either of these cases, we find that the leading degree term of the RHS of Eq. (27) is

$$4p^{4z_{11}+2z_{13}+2z_{14}}$$

• Step 3. Case 1. does not hold in Eq. (26), which implies that only Case 2 is viable. The leading degree term of the LHS of Eq. (27) is $4p^{2z_{11}+3z_{33}+z_{34}+z_{13}+z_{14}}$.

By Eq. (26), $z_{13} + z_{14}$ is the average of $z_{11} + z_{12}$ and $z_{33} + z_{34}$. Assume Case 1. holds, i.e., $z_{11} + z_{12} > z_{13} + z_{14} > z_{33} + z_{34}$. In this case, by Step 1. the leading degree term of the first term of $\xi((\mathcal{T}_M^*(p))^2)$ is $4p^{3z_{11}+2z_{33}+z_{12}+z_{13}+z_{14}}$. Since it has to equal the leading degree term of the second term, this means that $3z_{11} + 2z_{33} + z_{12} + z_{13} + z_{14} = 4z_{11} + 2z_{13} + 2z_{14}$. Simplifying, we see that

$$2z_{33} + z_{12} = z_{11} + z_{13} + z_{14}.$$
(28)

But also, for

$$((\mathcal{T}_M^*(p))^2)_{14} = p^{z_{11}+z_{14}} + p^{z_{12}+z_{13}} + p^{z_{13}+z_{34}} + p^{z_{14}+z_{33}}$$

in case (A), $z_{11} + z_{14} = z_{12} + z_{13}$, the leading degree term of $((\mathcal{T}_M^*(p))^2)_{14}$ is $p^{z_{11}+z_{14}} + p^{z_{12}+z_{13}}$. Together with Eq. (28), this implies that $2z_{13} = 2z_{33}$, which is not possible since $z_{33} > z_{13}$. The other possibility is case (B), $z_{11} + z_{14} = z_{13} + z_{34}$, the leading degree term of $((\mathcal{T}_M^*(p))^2)_{14}$ is $p^{z_{11}+z_{14}} + p^{z_{13}+z_{34}}$. Eq. (28) implies that $2z_{33} + z_{12} = 2z_{13} + z_{34}$. Since $p^{z_{13}+z_{34}}$ has the highest degree in $((\mathcal{T}_M^*(p))^2)_{14}$, that also implies that (note that $z_{11} = z_{13} - z_{14} + z_{34}$ from Case (B))

$$z_{13} + z_{34} = z_{13} - z_{14} + z_{34} + z_{14} = z_{11} + z_{14} > z_{14} + z_{33}.$$

So, $2z_{33} + z_{12} = 2z_{13} + z_{34} > z_{13} + (z_{14} + z_{33})$, which means that $z_{33} + z_{12} > z_{13} + z_{14}$. Since we have assumed that we are in Case 1., $z_{11} + z_{12} > z_{13} + z_{14} > z_{33} + z_{34}$, we see that $z_{33} + z_{12} > z_{13} + z_{14} > z_{33} + z_{34}$. So, $z_{12} > z_{34}$. However, since $p^{z_{13}+z_{34}}$ has the highest degree in $((\mathcal{T}_M^*(p))^2)_{14}$, we also know that $z_{13} + z_{34} > z_{12} + z_{13}$. But this implies that $z_{34} > z_{12}$, which is a contradiction. So, it is not possible that $z_{11} + z_{12} > z_{13} + z_{14} > z_{33} + z_{34}$. We conclude that in fact, Case 1. is impossible, only Case 2. remains, namely

$$z_{33} + z_{34} > z_{13} + z_{14} > z_{11} + z_{12}, (29)$$

and the leading degree term of the LHS of Eq. (27) is $4p^{2z_{11}+3z_{33}+z_{34}+z_{13}+z_{14}}$.

• Step 4. Case (A) is impossible, which implies that only Case (B) is viable.

Since the leading degree terms of the LHS and the RHS of Eq. (27) are equal, this means that $2z_{11} + 3z_{33} + z_{34} + z_{13} + z_{14} = 4z_{11} + 2z_{13} + 2z_{14}$. From Eq. (26), we know that $2z_{13} + 2z_{14} = z_{11} + z_{33} + z_{12} + z_{34}$. Substituting into the equation above, we get that $2z_{11} + 3z_{33} + z_{34} + z_{13} + z_{14} = 5z_{11} + z_{33} + z_{34} + z_{12}$. Simplifying, we get

$$2z_{33} + z_{13} + (z_{14} + z_{11}) = 4z_{11} + z_{12}.$$
(30)

Assume Case (A), $z_{11} + z_{14} = z_{12} + z_{13}$. Together with Eq. (30), this implies that

$$2z_{33} + 2z_{13} = 4z_{11},$$

which is not possible since $z_{11} > z_{33}, z_{13}$. So, we conclude that Case (A) is impossible, and only Case (B) is viable, i.e., the leading degree term of $((\mathcal{T}_M^*(p))^2)_{14}$ is $p^{z_{11}+z_{14}}+p^{z_{13}+z_{34}}$, and we have

$$z_{11} + z_{14} = z_{13} + z_{34} > z_{12} + z_{13}.$$
(31)

The only case remaining is Case 2. in combination of Case (B).

• Step 5. Case 2. in combination of Case (B) is impossible.

We will now consider the least degree term of the LHS and RHS of Eq. (27). From Case 2. Eq. (29), we note that the least degree term of

$$((\mathcal{T}_M^*(p))^2)_{12} = 2p^{z_{11}+z_{12}} + 2p^{z_{13}+z_{14}}$$

is $2p^{z_{11}+z_{12}}$, and the least degree term of

$$((\mathcal{T}_M^*(p))^2)_{34} = 2p^{z_{13}+z_{14}} + 2p^{z_{33}+34}$$

is $2p^{z_{13}+z_{14}}$. Since $z_{11}+z_{12} < z_{33}+z_{34}$, and $z_{11} > z_{33}$, we conclude that $z_{12} < z_{34}$. Similarly, since $z_{11}+z_{12} < z_{13}+z_{14}$, we also conclude that $z_{12} < z_{13}, z_{14}$. So, the least degree term of

$$((\mathcal{T}_M^*(p))^2)_{11} = p^{2z_{11}} + p^{2z_{12}} + p^{2z_{13}} + p^{2z_{14}}$$

is $p^{2z_{12}}$. From case (B) Eq. (31), we see that $z_{11} + z_{14} = z_{13} + z_{34}$. Since $z_{11} > z_{34}, z_{13}$, we see that $z_{14} < z_{13}$, and $z_{14} < z_{34}$. So, the least degree term of

$$((\mathcal{T}_M^*(p))^2)_{33} = p^{2z_{13}} + p^{2z_{14}} + p^{2z_{33}} + p^{2z_{34}}$$

is $p^{2z_{14}}$. Summing up, the least degree term of the LHS of Eq. (27) is precisely

$$4p^{z_{11}+3z_{12}+z_{13}+3z_{14}}$$

As for the RHS of Eq. (27), we note that since $z_{14} < z_{13}$, and $z_{12} < z_{11}, z_{33}, z_{34}$, the least degree term of

$$((\mathcal{T}_M^*(p))^2)_{13} = p^{z_{11}+z_{13}} + p^{z_{12}+z_{14}} + p^{z_{13}+z_{33}} + p^{z_{14}+z_{34}}$$

is $p^{z_{12}+z_{14}}$. This means that the least degree term of $((\mathcal{T}_M^*(p))^2)_{14}$ must have a coefficient of exactly 2 for the coefficients of the LHS and RHS of Eq. (27) to be equal. But we are in case (B), and so $p^{z_{11}+z_{14}} + p^{z_{13}+z_{34}}$ is the leading degree term of

$$((\mathcal{T}_M^*(p))^2)_{14} = p^{z_{11}+z_{14}} + p^{z_{12}+z_{13}} + p^{z_{13}+z_{34}} + p^{z_{14}+z_{33}}$$

So, the least degree term must be the combined term from $p^{z_{12}+z_{13}} + p^{z_{14}+z_{33}}$. In this case, the least degree term of the RHS of Eq. (27) is

$$4p^{2z_{12}+2z_{14}+2z_{12}+2z_{13}}.$$

Since this must be equal to the least degree term of the LHS of Eq. (27), we find that $z_{11} + 3z_{12} + z_{13} + 3z_{14} = 4z_{12} + 2z_{14} + 2z_{13}$. Simplifying, we get

$$z_{11} + z_{14} = z_{12} + z_{13},$$

which contradicts case (B) Eq. (31). So, we see that in fact, Case 2. in combination of case (B) is impossible. This implies that our original assumption that $\xi(((\mathcal{T}_M^*(p))^2)) = 0$ for all $p \in \mathbb{R}$ must be false.

Now, if we let $\xi_2 : N \mapsto \xi(N^2)$, we see that $\xi_2(\mathcal{T}_M^*(p))$ is not the zero function. From the construction of \mathcal{T}_M^* , this implies that $\xi_2(\mathcal{T}_M(\mathbf{p}))$ is not the zero function either. So, we can use Lemma 28 to find $M' \in \mathfrak{R}(M, \mathcal{F} \cup \{\rho_{\texttt{tensor}}, \zeta, \xi_2\}) \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{>0})$. Since $F(M') \neq 0$ for all $F \in \mathcal{F} \cup \{\rho_{\texttt{tensor}}, \zeta\}$, and $\xi((M')^2) \neq 0$, we can use Corollary 33 to find some $N \in \mathfrak{R}(M', \mathcal{F} \cup \{\rho_{\texttt{tensor}}, \zeta, \xi\}) \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{>0})$. Since $\zeta(N) \neq 0$, if N is not of Form (III), then it must be either diagonal distinct, or isomorphic to a matrix of Form (I) or Form (II), or satisfies $N_{11} = N_{22} \neq N_{33} = N_{44}$, but is not of Form (III). In any case, since $\rho_{\texttt{tensor}}(N) \neq 0$, we see that $\operatorname{Pl-GH}(N) \leq \operatorname{Pl-GH}(M)$ is #P-hard, due to either Theorem 68, or Lemma 73 and Lemma 82, or Lemma 70. On the other hand, if N is of Form (III), it is the required matrix (satisfying $\xi(N) \neq 0$).

We will now need a few more technical lemmas that show that since $\rho_{tensor}(M) \neq 0$, M may be assumed to have some more structure.

Lemma 89. Let $M \in Sym_4^{pd}(\mathbb{R}_{>0})$ be of Form (III) such that $\rho_{tensor}(M) \neq 0$. Let \mathcal{F} be a countable set of $Sym_4(\mathbb{R})$ -polynomials such that $F(M) \neq 0$ for all $F \in \mathcal{F}$. Then, either PI-GH(M) is #P-hard, or there exists some $N \in \mathfrak{R}(M, \mathcal{F} \cup \mathcal{F}' \cup \{\rho_{tensor}\}) \cap Sym_4^{pd}(\mathbb{R}_{>0})$ of Form (III), where $\mathcal{F}' = \{\xi_{c_1,c_2} : c_1, c_2 \in (\mathbb{Z}_{>0})\}$ for the $Sym_4(\mathbb{R})$ -polynomials

$$\xi_{c_1,c_2}: N \mapsto (N_{13}^{c_1} N_{12}^{c_2} - N_{14}^{c_1} N_{11}^{c_2})^2 + (N_{33}^{c_1} N_{14}^{c_2} - N_{34}^{c_1} N_{13}^{c_2})^2.$$

Proof. We let $\zeta : N \mapsto (N_{11} - N_{33})(N_{11} - N_{44})(N_{22} - N_{33})(N_{22} - N_{44})$. We note that $\zeta(M) \neq 0$. Without loss of generality, we may first replace M with the matrix $M' \in \mathfrak{R}(M, \mathcal{F} \cup \{\rho_{\mathsf{tensor}}, \zeta\}) \cap$ $\operatorname{Sym}_{4}^{\mathsf{pd}}(\mathbb{R}_{>0})$ whose existence is guaranteed by Lemma 86. So, we may assume that $M_{ii} > M_{jk}$ for all $i \in [4], j \neq k \in [4]$. Now, we let the entries of M be generated by some $\{g_t\}_{t \in [d]}$. We may replace M with some cM, as guaranteed by Lemma 4 such that $e_{ijt} \geq 0$ for all $i, j \in [4], t \in [d]$. We also note that $e_{ij0} = 0$ for all $i, j \in [4]$. We will now define the function $\widehat{\mathcal{T}_M} : \mathbb{R}^{d+1} \to \operatorname{Sym}_4(\mathbb{R})$ just as we did in Eq. (24). Then, we similarly define $\mathcal{T}_M^* : \mathbb{R} \to \operatorname{Sym}_4(\mathbb{R})$ as in Eq. (25), such that

$$\mathcal{T}_M^*(p)_{ij} = p^{z_{ij}}$$

for some integers z_{ij} . We may assume that $z_{11} \neq z_{33}$, and that $z_{ii} > z_{jk}$ for all $i \in [4]$, and $j \neq k \in [4]$.

We will now consider $\xi_{c_1,c_2}((\mathcal{T}^*_M(p))^2)$. First we want to show that if $\xi_{c_1,c_2}((\mathcal{T}^*_M(p))^2) = 0$, for all $p \in \mathbb{R}$, then $c_1 = c_2$. We note that

$$((\mathcal{T}_{M}^{*}(p))_{13}^{2})^{c_{1}}((\mathcal{T}_{M}^{*}(p))_{12}^{2})^{c_{2}} = (p^{z_{11}+z_{13}}+p^{z_{12}+z_{14}}+p^{z_{13}+z_{33}}+p^{z_{14}+z_{34}})^{c_{1}} \cdot (2p^{z_{11}+z_{12}}+2p^{z_{13}+z_{14}})^{c_{2}}, \text{ and}$$

$$((\mathcal{T}_{M}^{*}(p))_{14}^{2})^{c_{1}}((\mathcal{T}_{M}^{*}(p))_{11}^{2})^{c_{2}} = (p^{z_{11}+z_{14}} + p^{z_{12}+z_{13}} + p^{z_{13}+z_{34}} + p^{z_{14}+z_{33}})^{c_{1}} \cdot (p^{2z_{11}} + p^{2z_{12}} + p^{2z_{13}} + p^{2z_{14}})^{c_{2}}.$$

The identity $\xi_{c_1,c_2}((\mathcal{T}^*_M(p))^2) = 0$ for all $p \in \mathbb{R}$ implies that both terms in ξ_{c_1,c_2} are 0. In particular, $((\mathcal{T}^*_M(p))^2_{13})^{c_1}((\mathcal{T}^*_M(p))^2_{12})^{c_2} = ((\mathcal{T}^*_M(p))^2_{14})^{c_1}((\mathcal{T}^*_M(p))^2_{11})^{c_2}$, for all $p \in \mathbb{R}$, we see that the coefficient of the leading term of the LHS here is a (possibly non-trivial) multiple of 2^{c_2} . As for the RHS, by construction, the leading term of

$$(\mathcal{T}_M^*(p))_{11}^2 = p^{2z_{11}} + p^{2z_{12}} + p^{2z_{13}} + p^{2z_{14}}$$

will be $p^{2z_{11}}$. So, the leading term of $((\mathcal{T}_M^*(p))_{11}^2)^{c_2}$ will be $p^{2c_2z_{11}}$, with a coefficient of 1. So, this means that some terms within $((\mathcal{T}_M^*(p))_{14}^2)^{c_1}$ must be equal to each other. In fact, since the leading term coefficient of this term must be a multiple of 2^{c_2} , it must either be the case that the leading term of $(\mathcal{T}_M^*(p))_{14}^2$ must have either 2 or all 4 of the terms have the same degree. If all four terms have the same degree, that would imply that $z_{11} + z_{14} = z_{33} + z_{14}$, which is not true since $z_{11} \neq z_{33}$. So, this means that the leading term coefficient of the RHS is precisely 2^{c_1} . Since this is a multiple of 2^{c_1} , this implies that $c_1 \geq c_2$.

On the other hand, we can apply the same argument on the second term of the identity $\xi_{c_1,c_2}((\mathcal{T}_M^*(p))^2) = 0$, which is $((\mathcal{T}_M^*(p))^2_{33})^{c_1}((\mathcal{T}_M^*(p))^2_{14})^{c_2} = ((\mathcal{T}_M^*(p))^2_{34})^{c_1}((\mathcal{T}_M^*(p))^2_{13})^{c_2}$. Then we find that $c_2 \ge c_1$, which implies that in fact, $c_1 = c_2$.

Now, if $c_1 = c_2$, let us consider the matrix M itself. If $(M_{13}M_{12})^{c_1} = (M_{11}M_{14})^{c_1}$, and $(M_{33}M_{14})^{c_1} = (M_{13}M_{34})^{c_1}$, since $M \in \text{Sym}_4(\mathbb{R}_{>0})$, that implies that $M_{11}M_{14} = M_{12}M_{13}$, and $M_{14}M_{33} = M_{13}M_{34}$. We also note that $M_{14} = M_{23}$, and $M_{11}M_{44} = M_{22}M_{33}$, since M is of Form (III). So, this implies that $\rho_{\text{tensor}}(M) = 0$, which contradicts our assumption about M.

So, we see that for all ξ_{c_1,c_2} , either $\xi_{c_1,c_2}(M) \neq 0$, or there exists some $p \in \mathbb{R}$ such that $\xi_{c_1,c_2}((\mathcal{T}_M^*(p))^2) \neq 0$. We will now let $\mathcal{F}'' = \{F'': N \mapsto \xi_{c_1,c_2}(N^2) \mid c_1 \neq c_2 \in \mathbb{Z}_{>0}\}$, and $\mathcal{F}''' = \{\xi_{c,c}: c \in \mathbb{Z}_{>0}\}$. We can now use Lemma 28 to find $M'' \in \mathfrak{R}(M, \mathcal{F} \cup \mathcal{F}'' \cup \mathcal{F}''' \cup \{\rho_{\text{tensor}}, \zeta\}) \cap \text{Sym}_4^{\text{pd}}(\mathbb{R}_{>0})$. Since $F(M'') \neq 0$ for all $F \in \mathcal{F} \cup \mathcal{F}''' \cup \{\rho_{\text{tensor}}, \zeta\}$, and $\xi_{c_1,c_2}(M^2) \neq 0$ for all $c_1 \neq c_2 \in \mathbb{Z}_{>0}$, we can then use Corollary 33 to find some $N \in \mathfrak{R}(M, \mathcal{F} \cup \mathcal{F}' \cup \{\rho_{\text{tensor}}, \zeta\}) \cap \text{Sym}_4^{\text{pd}}(\mathbb{R}_{>0})$. Since $\zeta(N) \neq 0$, if N is not of Form (III), then it must either be diagonal distinct, or isomorphic to a matrix of Form (I) or Form (II), or satisfies $N_{11} = N_{22} \neq N_{33} = N_{44}$, but is not of Form (III). In any case, since $\rho_{\text{tensor}}(N) \neq 0$, we see that $\text{Pl-GH}(N) \leq \text{Pl-GH}(M)$ is #P-hard, due to either Theorem 68, or Lemma 73 and Lemma 82, or Lemma 70. On the other hand, if N is of Form (III), it is the required matrix.

Lemma 90. Let $M \in Sym_4^{pd}(\mathbb{R}_{>0})$ be of Form (III) such that $\rho_{tensor}(M) \neq 0$. Let \mathcal{F} be a countable set of $Sym_4(\mathbb{R})$ -polynomials such that $F(M) \neq 0$ for all $F \in \mathcal{F}$. Then, either Pl-GH(M) is #P-hard, or there exists some $N \in \mathfrak{R}(M, \mathcal{F} \cup \{\rho_{tensor}\}) \cap Sym_4^{pd}(\mathbb{R}_{>0})$ of Form (III), such that

$$\left(\log\left(\frac{N_{13}}{N_{14}}\right)\right)^2 \neq \log\left(\frac{N_{11}}{N_{12}}\right) \cdot \log\left(\frac{N_{33}}{N_{34}}\right),$$

and $N_{ii} > N_{jk}$ for all $i \in [4]$, $j \neq k \in [4]$, and $N_{11}N_{33}N_{12}N_{34} \neq (N_{13}N_{14})^2$.

Proof. We will let $\zeta : N \mapsto (N_{11} - N_{33})(N_{11} - N_{44})(N_{22} - N_{33})(N_{22} - N_{44})$, and $\zeta' : N \mapsto N_{11}N_{33}N_{12}N_{34} - (N_{13}N_{14})^2$. We note that $\zeta(M) \neq 0$. We will also let $\mathcal{F}' = \{\xi_{c_1,c_2} : c_1, c_2 \in (\mathbb{Z}_{>0})\}$ as defined in Lemma 89, such that .

$$\xi_{c_1,c_2}: N \mapsto (N_{13}^{c_1} N_{12}^{c_2} - N_{14}^{c_1} N_{11}^{c_2})^2 + (N_{33}^{c_1} N_{14}^{c_2} - N_{34}^{c_1} N_{13}^{c_2})^2.$$

We can first find $M' \in \mathfrak{R}(M, \mathcal{F} \cup \mathcal{F}' \cup \{\rho_{\texttt{tensor}}, \zeta\}) \cap \operatorname{Sym}_q^{pd}(\mathbb{R}_{>0})$ using Lemma 89. We can now use Lemma 88 to find $M'' \in \mathfrak{R}(M, \mathcal{F} \cup \mathcal{F}' \cup \{\rho_{\texttt{tensor}}, \zeta, \zeta'\}) \cap \operatorname{Sym}_q^{pd}(\mathbb{R}_{>0})$. Finally, we can use Lemma 86 to find $M''' \in \mathfrak{R}(M, \mathcal{F} \cup \mathcal{F}' \cup \{\rho_{\texttt{tensor}}, \zeta, \zeta'\}) \cap \operatorname{Sym}_q^{pd}(\mathbb{R}_{>0})$ such that $(M''')_{ii} > (M''')_{jk}$ for all $i \in [4]$, $j \neq k \in [4]$. Since $\zeta(M''') \neq 0$, if M''' is not of Form (III), then it must either be diagonal distinct, or isomorphic to a matrix of Form (I) or Form (II), or satisfies $(M''')_{11} = (M''')_{22} \neq (M''')_{33} = (M''')_{44}$, but is not of Form (III). In any case, since $\rho_{\texttt{tensor}}(M''') \neq 0$, we see that $\texttt{Pl-GH}(M''') \leq \texttt{Pl-GH}(M)$ is #P-hard, due to either Theorem 68, or Lemma 73 and Lemma 82, or Lemma 70. We may now assume that M''' is of Form (III).

For convenience, we can rename this M''' as M. We also note that if $M_{13} = M_{14}$, we are already done, since $M_{11}/M_{12} > 1$, $M_{33}/M_{34} > 1$, which implies that

$$\log\left(\frac{M_{11}}{M_{12}}\right) \cdot \log\left(\frac{M_{33}}{M_{34}}\right) > 0 = \left(\log\left(\frac{M_{13}}{M_{14}}\right)\right)^2.$$

So, we may assume that $M_{13} \neq M_{14}$. Now, we can assume without loss of generality that $M_{11} > M_{33}$, and that $M_{13} > M_{14}$ by permuting the rows and columns of M (first by a possible switch $\{1, 2\} \leftrightarrow \{3, 4\}$ and then a possible flip $3 \leftrightarrow 4$).

Now, we may let the entries of M be generated by some $\{g_t\}_{t\in[d]}$, such that $e_{ijt} \ge 0$ for all $i, j \in [4], t \in [d]$. So, for all $\mathbf{p} \in (\mathbb{R}_{>0})^d$,

$$\left(\log\left(\frac{\mathcal{T}_M(\mathbf{p})_{13}}{\mathcal{T}_M(\mathbf{p})_{14}}\right)\right)^2 = \left(\sum_{t \in [d]} (e_{13t} - e_{14t})\log(p_t)\right)^2, \text{ and}$$

$$\left(\log\left(\frac{\mathcal{T}_M(\mathbf{p})_{33}}{\mathcal{T}_M(\mathbf{p})_{34}}\right)\right) \left(\log\left(\frac{\mathcal{T}_M(\mathbf{p})_{11}}{\mathcal{T}_M(\mathbf{p})_{12}}\right)\right) = \left(\sum_{t \in [d]} (e_{33t} - e_{34t})\log(p_t)\right) \cdot \left(\sum_{t \in [d]} (e_{11t} - e_{12t})\log(p_t)\right).$$

We note that none of these three functions are the constant zero function. For example, since $M_{11} > M_{12}$, we have $(e_{111}, \ldots, e_{11d}) \neq (e_{121}, \ldots, e_{12d})$ and thus $\sum_{t \in [d]} (e_{11t} - e_{12t}) \log(p_t)$ is not constant zero. The other two being not constant zero follow similarly from $M_{13} > M_{14}$ and $M_{33} > M_{44}$. We will now make use of the following claim, which we shall prove shortly.

Claim 91. Let $\mathbf{0} \neq (a_1, \ldots, a_n), (b_1, \ldots, b_n), (c_1, \ldots, c_n) \in \mathbb{Z}^n$ such that

$$\left(\sum_{i\in[n]}a_ix_i\right)^2 = \left(\sum_{i\in[n]}b_ix_i\right)\left(\sum_{i\in[n]}c_ix_i\right),$$

for all $(x_1, \ldots, x_n) \in \mathbb{R}^n$, then there exists a rational constant κ such that

$$(c_1,\ldots,c_n)=\kappa\cdot(a_1,\ldots,a_n)=\kappa^2\cdot(b_1,\ldots,b_n).$$

We will now define the function $\Omega : (\mathbb{R}_{>0})^d \to \mathbb{R}$ such that

$$\Omega(\mathbf{p}) = \left(\log\left(\frac{\mathcal{T}_M(\mathbf{p})_{13}}{\mathcal{T}_M(\mathbf{p})_{14}}\right)\right)^2 - \left(\log\left(\frac{\mathcal{T}_M(\mathbf{p})_{33}}{\mathcal{T}_M(\mathbf{p})_{34}}\right)\right) \cdot \left(\log\left(\frac{\mathcal{T}_M(\mathbf{p})_{11}}{\mathcal{T}_M(\mathbf{p})_{12}}\right)\right)$$

Claim 91 tells us that if $\Omega(\mathbf{p}) = 0$ for all $\mathbf{p} \in (\mathbb{R}_{>0})^d$, then there exists some rational $\kappa \in \mathbb{Q}$ such that

$$(e_{111} - e_{121}, \dots, e_{11d} - e_{12d}) = \kappa \cdot (e_{131} - e_{141}, \dots, e_{13d} - e_{14d}) = \kappa^2 \cdot (e_{331} - e_{341}, \dots, e_{33d} - e_{34d}).$$

This implies that

$$\frac{(\log (M_{11}/M_{12}))}{(\log (M_{13}/M_{14}))} = \frac{(e_{111} - e_{121})\log(g_1) + \dots + (e_{11d} - e_{12d})\log(g_d)}{(e_{131} - e_{141})\log(g_1) + \dots + (e_{13d} - e_{14d})\log(g_d)} = \kappa = \frac{c_1}{c_2}, \text{ and}$$
$$\frac{(\log (M_{13}/M_{14}))}{(\log (M_{33}/M_{34}))} = \frac{(e_{131} - e_{141})\log(g_1) + \dots + (e_{13d} - e_{14d})\log(g_d)}{(e_{331} - e_{341})\log(g_1) + \dots + (e_{33d} - e_{34d})\log(g_d)} = \kappa = \frac{c_1}{c_2},$$

for some positive integers c_1, c_2 . But this means that

$$\left(\frac{M_{11}}{M_{12}}\right)^{c_2} = \left(\frac{M_{13}}{M_{14}}\right)^{c_1}$$
, and $\left(\frac{M_{13}}{M_{14}}\right)^{c_2} = \left(\frac{M_{33}}{M_{34}}\right)^{c_1}$

In particular, we see that

$$((M)_{13})^{c_1}((M)_{12})^{c_2} = ((M)_{11})^{c_2}((M)_{14})^{c_1}$$
, and $((M)_{13})^{c_2}((M)_{34})^{c_1} = ((M)_{33})^{c_1}((M)_{14})^{c_2}$.

This contradicts our assumption that $\xi_{c_1,c_2}(M) \neq 0$. So, our assumption that $\Omega(\mathbf{p}) = 0$ for all $\mathbf{p} \in (\mathbb{R}_{>0})^d$ must be false. Let $\mathbf{p}' = (p'_1, \ldots, p'_d) \in (\mathbb{R}_{>0})^d$ such that $\Omega(\mathbf{p}') \neq 0$. We can now define $\mathbf{p} : \mathbb{R} \to \mathbb{R}^d$, such that

$$\mathbf{p}(t) = t \cdot \mathbf{p}' + (1-t) \cdot (g_1, \dots, g_d).$$

We note that $\mathbf{p}(0) = (g_1, \ldots, g_d) \in (\mathbb{R}_{>0})^d$, and $\mathbf{p}(1) = \mathbf{p}' \in (\mathbb{R}_{>0})^d$. Therefore, for all $t \in [0, 1]$, and for all $i \in [d]$, $\mathbf{p}(t)_i = t \cdot p'_i + (1 - t)g_i > 0$. Moreover, since $\mathbf{p}(t)$ is continuous as a function of t, we see that there exists some $\delta_1, \delta_2 > 0$ such that $\mathbf{p}(t) \in (\mathbb{R}_{>0})^d$ for all $t \in (-\delta_1, \delta_1)$, and $t \in (1 - \delta_2, 1 + \delta_2)$. We will now let $U = (-\delta_1, 1 + \delta_2) \subset \mathbb{R}$ be an open set, such that $\mathbf{p}(t) \in (\mathbb{R}_{>0})^d$ for all $t \in U$. We now define $\omega : U \to \mathbb{R}$ such that

$$\omega(t) = \Omega(\mathbf{p}(t)).$$

Clearly, ω is a real valued analytic function on U. Moreover, by our choice of \mathbf{p}' , we know that $\omega(1) = \Omega(\mathbf{p}') \neq 0$. So, ω is a non-zero real analytic function on the open set U. So, we know ([KP02], Corollary 1.2.7) that the set of zeros of ω within [0, 1] cannot have an accumulation point. In particular, there can only be finitely many zeros of ω within [0, 1]. We can denote the set of zeros as $\emptyset_{\omega} = \{t \in [0, 1] : \omega(t) = 0\}$.

Since $\mathbf{p}(0) = (g_1, \ldots, g_d)$, we see that $\mathcal{T}_M(\mathbf{p}(0)) = M$. So, there exists some $0 < \delta < 1$ such that $\mathcal{T}_M(\mathbf{p}(t))_{ii} > \mathcal{T}_M(\mathbf{p}(t))_{jk} > 0$, for all $i \in [4], j \neq k \in [4]$, and $t \in (0, \delta)$. We also note that the eigenvalues of $\mathcal{T}_M(\mathbf{p}(t))$ are continuous as functions of t. Since $\mathcal{T}_M(\mathbf{p}(0)) = M$ is positive definite, this implies that there exists some $0 < \delta' < 1$ such that $\mathcal{T}_M(\mathbf{p}(t))$ is positive definite for all $t \in (0, \delta')$. We will now let $\mathcal{F}'' = \mathcal{F} \cup \mathcal{F}' \cup \{\rho_{\mathsf{tensor}}, \zeta, \zeta'\}$. We note that \mathcal{F}'' is a countable set of $\operatorname{Sym}_4(\mathbb{R})$ -polynomials. So, $F(\mathcal{T}_M(\mathbf{p}(t)))$ is a polynomial in t for all $F \in \mathcal{F}''$. Since $F(M) \neq 0$ for all $F \in \mathcal{F}''$. We let $\emptyset_F = \{t \in (0, 1) : F(\mathcal{T}_M(\mathbf{p}(t))) = 0\}$ for all $F \in \mathcal{F}''$. Each of these is a finite set, and therefore, $\bigcup_{F \in \mathcal{F}''} \emptyset_F$ is a countable set.

We can therefore find $\mathbf{p}^* = \mathbf{p}(t^*)$ for some $t^* \in ((0, \delta) \cap (0, \delta')) \setminus (\bigcup_{F \in \mathcal{F}''} \emptyset_F \cup \emptyset_{\omega})$. By our choice of t^* , we see that $F(\mathcal{T}_M(\mathbf{p}^*)) \neq 0$ for all $F \in \mathcal{F} \cup \mathcal{F}' \cup \{\rho_{\texttt{tensor}}, \zeta, \zeta'\}$, and $\mathcal{T}_M(\mathbf{p}^*) \in \text{Sym}_4^{pd}(\mathbb{R}_{>0})$. Moreover, we also see that $\Omega(\mathbf{p}^*) \neq 0$. So, $N = \mathcal{T}_M(\mathbf{p}^*) \in \mathfrak{R}(M, \mathcal{F} \cup \mathcal{F}' \cup \{\rho_{\texttt{tensor}}, \zeta, \zeta'\}) \cap$ $\text{Sym}_4^{pd}(\mathbb{R}_{>0})$ such that $\Omega(N) \neq 0$. Since $\zeta(N) \neq 0$, if N is not of Form (III), then it must either be diagonal distinct, or isomorphic to a matrix of Form (I) or Form (II), or satisfies $N_{11} = N_{22} \neq N_{33} =$ N_{44} , but is not of Form (III). In any case, since $\rho_{\texttt{tensor}}(N) \neq 0$, we see that $\texttt{P1-GH}(N) \leq \texttt{P1-GH}(M)$ is #P-hard, due to either Theorem 68, or Lemma 73 and Lemma 82, or Lemma 70. On the other hand, if N is of Form (III), it is the required matrix.

We will now prove Claim 91.

Proof of Claim 91. The zero sets defined by the LHS and the RHS are respectively the hyperplane $\sum_{i \in [n]} a_i x_i = 0$, and the union of the hyperplanes $\sum_{i \in [n]} b_i x_i = 0$ and $\sum_{i \in [n]} c_i x_i = 0$. Thus they

must all coincide. Their (non-zero) normal vectors must be proportional. Therefore there exist constants $\kappa, \kappa' \in \mathbb{Q}_{\neq 0}$ such that $(c_1, \ldots, c_n) = \kappa \cdot (a_1, \ldots, a_n)$ and $(b_1, \ldots, b_n) = \kappa' \cdot (a_1, \ldots, a_n)$. Then the given equality in the lemma statement implies that $\kappa' = \kappa^{-1}$.

Lemma 92. Let $M \in Sym_4^{pd}(\mathbb{R}_{>0})$ be of Form (III) such that $\rho_{tensor}(M) \neq 0$. Let \mathcal{F} be a countable set of $Sym_4(\mathbb{R})$ -polynomials such that $F(M) \neq 0$ for all $F \in \mathcal{F}$. Let $\mathbf{0} \neq \mathbf{x} \in \chi_4$ of support size greater than 2. Then, either Pl-GH(M) is #P-hard, or there exists some $N \in \mathfrak{R}(M, \mathcal{F} \cup \{\Psi_{\mathbf{x}}, \rho_{tensor}\}) \cap Sym_4^{pd}(\mathbb{R}_{>0})$ of Form (III).

Proof. We can replace M with the matrix M' that is obtained from Lemma 90. We may also assume without loss of generality that $M_{11} > M_{33}$, and that $M_{13} \ge M_{14}$ by permuting the rows and columns of M (first by a possible switch $\{1, 2\} \leftrightarrow \{3, 4\}$ and then a possible flip $3 \leftrightarrow 4$). We will now let the entries of M be generated by some $\{g_t\}_{t\in[d]}$. Using Lemma 4, we may replace M with some cM such that $e_{ijt} \ge 0$ for all $i, j \in [4], t \in [d]$. Since $M \in \text{Sym}_4(\mathbb{R}_{>0})$, we note that $e_{ij0} = 0$ for all $i, j \in [4]$.

We will now define the function $\widehat{\mathcal{T}_M}: \mathbb{R}^{d+1} \to \operatorname{Sym}_4(\mathbb{R})$, just as in Eq. (24), such that

$$\widehat{\mathcal{T}_M}(p, z_1, \dots, z_d)_{ij} = \mathcal{T}_M(p^{z_1}, \dots, p^{z_d})$$

Following the same argument as in the paragraph after Eq. (24), we can pick some rational z_1^*, \ldots, z_d^* such that $\widehat{\mathcal{T}}_M(e, z_1^*, \ldots, z_d^*)$ satisfies the properties of M, that

$$M_{11}M_{33}M_{12}M_{34} \neq (M_{13}M_{14})^2, \quad \left(\log\left(\frac{M_{13}}{M_{14}}\right)\right)^2 \neq \left(\log\left(\frac{M_{11}}{M_{12}}\right)\right) \cdot \left(\log\left(\frac{M_{33}}{M_{34}}\right)\right),$$

 $M_{11} > M_{33}, M_{13} \ge M_{14}$, and $M_{ii} > M_{jk}$ for all $i \in [4], j \ne k \in [4]$, by permuting the rows and columns of M (first by a possible switch $\{1, 2\} \leftrightarrow \{3, 4\}$ and then a possible flip $3 \leftrightarrow 4$). We then let $Z^* \in \mathbb{Z}_{>0}$ such that $Z^* z_t^* \in \mathbb{Z}_{>0}$ for all $t \in [d]$. We can then define $\mathcal{T}_M^* : \mathbb{R} \to \text{Sym}_4(\mathbb{R})$, as in Eq. (25) such that

$$\mathcal{T}_M^*(p)_{ij} = \widehat{\mathcal{T}_M}(p, Z^* \cdot z_i^*, \dots, Z^* \cdot z_d^*)_{ij} = p^{z_{ij}},$$

for some $z_{ij} \in \mathbb{Z}_{\geq 0}$ for all $i, j \in [4]$. We may assume from this construction that $z_{ii} > z_{jk}$ for all $i \in [4], j \neq k \in [4], z_{11} > z_{33}, z_{13} \geq z_{14}, z_{11} + z_{33} + z_{12} + z_{34} \neq 2(z_{13} + z_{14})$, and that $(z_{13} - z_{14})^2 \neq (z_{11} - z_{12})(z_{33} - z_{34})$.

We see that

$$\mathcal{T}_{M}^{*}(p) = \begin{pmatrix} p^{z_{11}} & p^{z_{12}} & p^{z_{13}} & p^{z_{14}} \\ p^{z_{12}} & p^{z_{11}} & p^{z_{14}} & p^{z_{13}} \\ p^{z_{13}} & p^{z_{14}} & p^{z_{33}} & p^{z_{34}} \\ p^{z_{14}} & p^{z_{13}} & p^{z_{34}} & p^{z_{33}} \end{pmatrix}$$

It can be verified that the eigenvalues of $\mathcal{T}_M^*(p)$ are:

$$\lambda_{1}(\mathcal{T}_{M}^{*}(p)) = \frac{1}{2} (\mu_{1}(p) + \nu_{1}(p)),$$

$$\lambda_{2}(\mathcal{T}_{M}^{*}(p)) = \frac{1}{2} (\mu_{1}(p) - \nu_{1}(p)),$$

$$\lambda_{3}(\mathcal{T}_{M}^{*}(p)) = \frac{1}{2} (\mu_{2}(p) + \nu_{2}(p)),$$

$$\lambda_{4}(\mathcal{T}_{M}^{*}(p)) = \frac{1}{2} (\mu_{2}(p) - \nu_{2}(p)),$$
(32)

where

$$\mu_1(p) = p^{z_{11}} + p^{z_{33}} + p^{z_{12}} + p^{z_{34}},$$

$$\mu_2(p) = p^{z_{11}} + p^{z_{33}} - p^{z_{12}} - p^{z_{34}},$$

$$\nu_1(p) = \sqrt{(p^{z_{11}} - p^{z_{33}} + p^{z_{12}} - p^{z_{34}})^2 + 4(p^{z_{13}} + p^{z_{14}})^2}, \text{ and}$$

$$\nu_2(p) = \sqrt{(p^{z_{11}} - p^{z_{33}} - p^{z_{12}} + p^{z_{34}})^2 + 4(p^{z_{13}} - p^{z_{14}})^2}.$$

We will now make use of the following claim, which we shall prove shortly.

Claim 93.

$$\lim_{p \to \infty} \frac{\lambda_1(\mathcal{T}_M^*(p))}{p^{z_{11}}} = \lim_{p \to \infty} \frac{\lambda_3(\mathcal{T}_M^*(p))}{p^{z_{11}}} = 1,$$
$$\lim_{p \to \infty} \frac{\lambda_2(\mathcal{T}_M^*(p))}{p^{z_{33}}} = \lim_{p \to \infty} \frac{\lambda_4(\mathcal{T}_M^*(p))}{p^{z_{33}}} = 1.$$

We can see that for large enough values of p, $\lambda_i(\mathcal{T}_M^*(p)) > 0$ for all $i \in [4]$. Define the function $\varphi_{\mathbf{x}} : (\mathbb{R}_{\neq 0})^4 \to \mathbb{R}$ as $\varphi_{\mathbf{x}}(\alpha_1, \ldots, \alpha_4) = \prod_{i \in [4]} (\alpha_i)^{x_i}$. For large enough values of p, Claim 93 implies that $\varphi_{\mathbf{x}}(\lambda_1(\mathcal{T}_M^*(p)), \ldots, \lambda_4(\mathcal{T}_M^*(p)))$ is well-defined. Moreover, we note that

$$\phi_{\mathbf{x}}(\lambda_1(\mathcal{T}_M^*(p)),\ldots,\lambda_4(\mathcal{T}_M^*(p))) = 0 \iff \varphi_{\mathbf{x}}(\lambda_1(\mathcal{T}_M^*(p)),\ldots,\lambda_4(\mathcal{T}_M^*(p))) = 1.$$

We can analyze $\phi_{\mathbf{x}}(\lambda_1(\mathcal{T}_M^*(p)), \ldots, \lambda_4(\mathcal{T}_M^*(p)))$ by studying $\varphi_{\mathbf{x}}(\lambda_1(\mathcal{T}_M^*(p)), \ldots, \lambda_4(\mathcal{T}_M^*(p)))$. We note that

$$\varphi_{\mathbf{x}}(\lambda_{1}(\mathcal{T}_{M}^{*}(p)),\ldots,\lambda_{4}(\mathcal{T}_{M}^{*}(p))) = (p^{z_{11}})^{x_{1}+x_{3}}(p^{z_{33}})^{x_{2}+x_{4}} \left(\frac{\lambda_{1}(\mathcal{T}_{M}^{*}(p))}{p^{z_{11}}}\right)^{x_{1}} \cdot \left(\frac{\lambda_{2}(\mathcal{T}_{M}^{*}(p))}{p^{z_{33}}}\right)^{x_{2}} \cdot \left(\frac{\lambda_{3}(\mathcal{T}_{M}^{*}(p))}{p^{z_{11}}}\right)^{x_{3}} \cdot \left(\frac{\lambda_{4}(\mathcal{T}_{M}^{*}(p))}{p^{z_{33}}}\right)^{x_{4}}.$$

So,

$$\lim_{p \to \infty} \varphi_{\mathbf{x}}(\lambda_1(\mathcal{T}_M^*(p)), \dots, \lambda_4(\mathcal{T}_M^*(p))) = \lim_{p \to \infty} p^{(z_{11}-z_{33})(x_1+x_3)}$$

Since $z_{11} - z_{33} > 0$ by our assumption, we see that $\lim_{p \to \infty} \varphi_{\mathbf{x}}(\lambda_1(\mathcal{T}_M^*(p)), \dots, \lambda_4(\mathcal{T}_M^*(p))) \neq 1$ if $x_1 + x_3 \neq 0$. In other words, for large enough $p, \phi_{\mathbf{x}}(\lambda_1(\mathcal{T}_M^*(p)), \dots, \lambda_4(\mathcal{T}_M^*(p))) \neq 0$, if $x_1 + x_3 \neq 0$. So, when $x_1 + x_3 \neq 0$, we see from the construction of \mathcal{T}_M^* that there exists some $\mathbf{p}^* = ((p^*)^{Z^* \cdot z_1^*}, \dots, (p^*)^{Z^* \cdot z_d^*})$ such that $\phi_{\mathbf{x}}(\mathcal{T}_M(\mathbf{p}^*)) \neq 0$.

We will now assume that $x_1 + x_3 = 0$. This of course also implies that $x_2 + x_4 = 0$, since $\mathbf{x} \in \chi_4$. If $x_1 = 0$, or $x_2 = 0$, then the support size of \mathbf{x} would be ≤ 2 . We can assume that $x_1 > 0$, and that $x_2 \neq 0$. So, we see that $\phi_{\mathbf{x}}(\lambda_1(\mathcal{T}_M^*(p)), \ldots, \lambda_4(\mathcal{T}_M^*(p))) = 0$ implies that

$$\left(\frac{\lambda_1(\mathcal{T}_M^*(p))}{\lambda_3(\mathcal{T}_M^*(p))}\right)^{x_1} = \left(\frac{\lambda_2(\mathcal{T}_M^*(p))}{\lambda_4(\mathcal{T}_M^*(p))}\right)^{x_2},$$

for all $p \in \mathbb{R}$ where $\lambda_i(\mathcal{T}^*_M(p)) \neq 0$ for all $i \in [4]$.

We will now make use of the following claim, which we shall prove later.

Claim 94. For small $\delta > 0$,

$$\begin{split} \lambda_1(\mathcal{T}_M^*(e^{\delta})) &= 4 + \frac{\delta}{2} \left(z_{11} + z_{33} + z_{12} + z_{34} + 2z_{13} + 2z_{14} \right) + O(\delta^2), \\ \lambda_2(\mathcal{T}_M^*(e^{\delta})) &= \frac{\delta}{2} \left(z_{11} + z_{33} + z_{12} + z_{34} - 2z_{13} - 2z_{14} \right) + O(\delta^2), \\ \lambda_3(\mathcal{T}_M^*(e^{\delta})) &= \frac{\delta}{2} \left(\left(z_{11} + z_{33} - z_{12} - z_{34} \right) + \sqrt{(z_{11} - z_{33} - z_{12} + z_{34})^2 + 4(z_{13} - z_{14})^2} \right) + O(\delta^2), \\ \lambda_4(\mathcal{T}_M^*(e^{\delta})) &= \frac{\delta}{2} \left(\left(z_{11} + z_{33} - z_{12} - z_{34} \right) - \sqrt{(z_{11} - z_{33} - z_{12} + z_{34})^2 + 4(z_{13} - z_{14})^2} \right) + O(\delta^2). \end{split}$$

We can see that,

$$\lim_{\delta \to 0} \lambda_1(\mathcal{T}^*_M(e^{\delta})) = 4.$$

Also $\lim_{\delta \to 0} \lambda_i(\mathcal{T}_M^*(e^{\delta})) = 0$ for i = 2, 3, 4. From our choice of z_{ij} , we know that $z_{11}, z_{33} > z_{12}, z_{34}$. So, we know that $z_{11} + z_{33} - z_{12} - z_{34} > 0$. This implies that

$$\lim_{\delta \to 0} \frac{\lambda_3(\mathcal{T}_M^*(e^{\delta}))}{\delta} = (z_{11} + z_{33} - z_{12} - z_{34}) + \sqrt{(z_{11} - z_{33} - z_{12} + z_{34})^2 + 4(z_{13} - z_{14})^2} > 0.$$

So,

$$0 \neq \lim_{\delta \to 0} \left(\delta^{x_1} \cdot \left(\frac{\lambda_1(\mathcal{T}_M^*(e^{\delta})))}{\lambda_3(\mathcal{T}_M^*(e^{\delta}))} \right)^{x_1} \right) = \lim_{\delta \to 0} \left(\delta^{x_1} \cdot \left(\frac{\lambda_2(\mathcal{T}_M^*(e^{\delta})))}{\lambda_4(\mathcal{T}_M^*(e^{\delta}))} \right)^{x_2} \right).$$
(33)

So, if $x_2 > 0$, then in the expression in Claim 94, if the coefficient of δ in $\lambda_4(\mathcal{T}_M^*(e^{\delta}))$ is non-zero, then the ratio λ_2/λ_4 in Eq. (33) will stay bounded as $\delta \to 0$, leading to a contradiction to Eq. (33) where the limit is non-zero. Hence the coefficient of δ in $\lambda_4(\mathcal{T}_M^*(e^{\delta}))$ must be 0. Similarly, if $x_2 < 0$, the coefficient of δ in $\lambda_2(\mathcal{T}_M^*(e^{\delta}))$ must be 0. Let us first assume that $x_2 > 0$. In that case, we find that

$$z_{11} + z_{33} - z_{12} - z_{34} = \sqrt{(z_{11} - z_{33} - z_{12} + z_{34})^2 + 4(z_{13} - z_{14})^2}$$

On squaring both sides and simplifying, we find that

$$(z_{11}-z_{12})^2 + (z_{33}-z_{34})^2 + 2(z_{11}-z_{12})(z_{33}-z_{34}) = (z_{11}-z_{12})^2 + (z_{33}-z_{34})^2 - 2(z_{11}-z_{12})(z_{33}-z_{34}) + 4(z_{13}-z_{14})^2.$$

This implies that $(z_{11}-z_{12})(z_{33}-z_{34}) = (z_{13}-z_{14})^2$. But by our choice of z_{ij} , we ensured that $(z_{11}-z_{12})(z_{33}-z_{34}) \neq (z_{13}-z_{14})^2$. So, we see that it is not possible that $\phi_{\mathbf{x}}(\lambda_1(\mathcal{T}_M^*(e^{\delta})),\ldots,\lambda_4(\mathcal{T}_M^*(e^{\delta}))) = 0$ for all $\delta \in \mathbb{R}$, if $x_2 > 0$.

On the other hand, if $x_2 < 0$, then from Claim 94, we see that the coefficient of δ in $\lambda_2(\mathcal{T}_M^*(e^{\delta}))$ must be 0. This implies that $z_{11} + z_{33} + z_{12} + z_{34} = 2z_{13} + 2z_{14}$. But once again, from our choice of z_{ij} , we ensured that $z_{11} + z_{33} + z_{12} + z_{34} \neq 2z_{13} + 2z_{14}$. So, we see that it is not possible that $\phi_{\mathbf{x}}(\lambda_1(\mathcal{T}_M^*(e^{\delta})), \ldots, \lambda_4(\mathcal{T}_M^*(e^{\delta}))) = 0$ for all $\delta \in \mathbb{R}$, if $x_2 < 0$ either.

So, we see that for all $\mathbf{x} \in \chi_4$ of support size greater than 2, $\phi_{\mathbf{x}}(\lambda_1(\mathcal{T}_M^*(e^{\delta})), \ldots, \lambda_4(\mathcal{T}_M^*(e^{\delta}))) \neq 0$ for some δ that is small enough. Moreover, this is true for all $\mathbf{y} \in \chi_q$ such that $y_{\sigma(i)} = x_i$, for some $\sigma \in S_4$. This means that there exists some $p^* \in \mathbb{R}$ such that

$$\Psi_{\mathbf{x}}(\mathcal{T}_M^*(p^*)) = \Phi_{\mathbf{x}}(\lambda_1(\mathcal{T}_M^*(p^*)), \dots, \lambda_4(\mathcal{T}_M^*(p^*))) \neq 0.$$

So, Lemma 28 allows us to find the required $N \in \mathfrak{R}(M, \mathcal{F} \cup \{\Psi_{\mathbf{x}}, \rho_{\mathtt{tensor}}\}) \cap \mathrm{Sym}_{4}^{\mathtt{pd}}(\mathbb{R}_{>0})$.

We will now prove Claim 93, and Claim 94

Proof of Claim 93. We note that

$$\lim_{p \to \infty} \frac{\mu_1(p)}{p^{z_{11}}} = \lim_{p \to \infty} \frac{p^{z_{11}} + p^{z_{33}} + p^{z_{12}} + p^{z_{34}}}{p^{z_{11}}} = 1$$

Similarly,

$$\lim_{p \to \infty} \frac{\mu_2(p)}{p^{z_{11}}} = \lim_{p \to \infty} \frac{p^{z_{11}} + p^{z_{33}} - p^{z_{12}} - p^{z_{34}}}{p^{z_{11}}} = 1.$$

We can also see that

$$\lim_{p \to \infty} \frac{\nu_1(p)}{p^{z_{11}}} = \lim_{p \to \infty} \frac{\sqrt{(p^{z_{11}} - p^{z_{33}} + p^{z_{12}} - p^{z_{34}})^2 + 4(p^{z_{13}} + p^{z_{14}})^2}}{p^{z_{11}}} = 1, \text{ and}$$
$$\lim_{p \to \infty} \frac{\nu_2(p)}{p^{z_{11}}} = \sqrt{(p^{z_{11}} - p^{z_{33}} - p^{z_{12}} + p^{z_{34}})^2 + 4(p^{z_{13}} - p^{z_{14}})^2} = 1.$$

This immediately proves that

$$\lim_{p \to \infty} \frac{\lambda_1(\mathcal{T}_M^*(p))}{p^{z_{11}}} = \lim_{p \to \infty} \frac{\lambda_3(\mathcal{T}_M^*(p))}{p^{z_{11}}} = 1.$$

We now note that

$$\lambda_2(\mathcal{T}_M^*(p)) = \frac{1}{2} \left(\frac{(\mu_1(p))^2 - (\nu_1(p))^2}{\mu_1(p) + \nu_1(p)} \right), \quad \text{and} \quad \lambda_4(\mathcal{T}_M^*(p)) = \frac{1}{2} \left(\frac{(\mu_2(p))^2 - (\nu_2(p))^2}{\mu_2(p) + \nu_2(p)} \right).$$

Now, we can see that

$$(\mu_1(p))^2 - (\nu_1(p))^2 = 4(p^{z_{11}} + p^{z_{12}})(p^{z_{33}} + p^{z_{34}}) - 4(p^{z_{13}} + p^{z_{14}})^2, \text{ and}$$
$$(\mu_2(p))^2 - (\nu_2(p))^2 = 4(p^{z_{11}} - p^{z_{12}})(p^{z_{33}} - p^{z_{34}}) - 4(p^{z_{13}} - p^{z_{14}})^2.$$

Since we already know the limiting behaviors of $\mu_1(p) + \nu(p)$, and $\mu_2(p) + \nu_2(p)$, we can see that

$$\lim_{p \to \infty} \frac{\lambda_2(\mathcal{T}_M^*(p))}{p^{z_{33}}} = \lim_{p \to \infty} \frac{\lambda_4(\mathcal{T}_M^*(p))}{p^{z_{33}}} = 1.$$

Proof of Claim 94. Following the Taylor series expansions of $\mu_1(e^{\delta}), \mu_2(e^{\delta}), (\nu_1(e^{\delta}))^2$, and $(\nu_2(e^{\delta}))^2$, we see that

$$\mu_1(e^{\delta}) = 4 + \delta(z_{11} + z_{33} + z_{12} + z_{34}) + O(\delta^2),$$

$$\mu_2(e^{\delta}) = \delta(z_{11} + z_{33} - z_{12} - z_{34}) + O(\delta^2),$$

$$(\nu_1(e^{\delta}))^2 = 16 + 16\delta(z_{13} + z_{14}) + O(\delta^2), \text{ and}$$

$$(\nu_2(e^{\delta}))^2 = \delta^2((z_{11} - z_{33} - z_{12} + z_{34})^2 + 4(z_{13} - z_{14})^2) + O(\delta^3).$$

Then, we can use the square root expansion of $\sqrt{1+x}$ to see that

$$\nu_1(e^{\delta}) = 4 + 2\delta(z_{13} + z_{14}) + O(\delta^2), \text{ and}$$

 $\nu_2(e^{\delta}) = \delta\sqrt{(z_{11} - z_{33} - z_{12} + z_{34})^2 + 4(z_{13} - z_{14})^2} + O(\delta^2).$

Putting together these expressions finishes the proof of this claim.

We can now prove that Pl-GH(M), when M is of Form (III) that is not isomorphic to a tensor product, must also be #P-hard.

Lemma 95. Let $M \in Sym_4^{pd}(\mathbb{R}_{>0})$ be of Form (III) such that $\rho_{tensor}(M) \neq 0$. Then, Pl-GH(M) is #*P*-hard.

Proof. The proof is very similar to the proof of Lemma 85. The only difference is that instead of Lemma 83, we will have to use Lemma 92 to find the matrices N_i . The rest of the proof is identical.

11.4 Form (VI)

Finally, there is only one more form of matrix for us to deal with. We will now prove that Pl-GH(M) is #P-hard, when M is of Form (VI) that is not isomorphic to a tensor product.

$$\begin{pmatrix} a & x & y & z \\ x & a & z & y \\ y & z & a & x \\ z & y & x & a \end{pmatrix}$$

Figure 10: Form (VI)

Lemma 96. Let $M \in Sym_4^{pd}(\mathbb{R}_{>0})$ be of Form (VI) such that $\rho_{tensor}(M) \neq 0$. Let \mathcal{F} be a countable set of $Sym_4(\mathbb{R})$ -polynomials such that $F(M) \neq 0$ for all $F \in \mathcal{F}$. Let $\mathbf{0} \neq \mathbf{x} \in \chi_4$ of support size greater than 2. Then, either Pl-GH(M) is #P-hard, or there exists some $N \in \mathfrak{R}(M, \mathcal{F} \cup \{\Psi_{\mathbf{x}}, \rho_{tensor}\}) \cap Sym_4^{pd}(\mathbb{R}_{>0})$ of Form (VI).

Proof. We can use Lemma 86 to obtain some $M' \in \mathfrak{R}(M, \mathcal{F} \cup \{\rho_{\texttt{tensor}}\}) \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{>0})$, such that $M'_{ii} > M'_{jk}$ for all $i \in [4], j \neq k \in [4]$. If M' is diagonal distinct, then Theorem 68 implies that $\mathsf{Pl-GH}(M') \leq \mathsf{Pl-GH}(M)$ is $\#\mathsf{P}$ -hard. Let us assume that M' is not diagonal distinct. If M' is not of Form (VI), then Theorem 75 implies that either $\mathsf{Pl-GH}(M') \leq \mathsf{Pl-GH}(M)$ is $\#\mathsf{P}$ -hard, or there exists some $N \in \mathfrak{R}(M', \mathcal{F} \cup \{\rho_{\texttt{tensor}}\}) \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{>0})$ such that N is of Form (I), (III), or (IV), and the diagonal entries of N are not all identical. But then, we note that since $\rho_{\texttt{tensor}}(N) \neq 0$, Lemma 82, Lemma 85, and Lemma 95 imply that $\mathsf{Pl-GH}(N) \leq \mathsf{Pl-GH}(M)$ is $\#\mathsf{P}$ -hard. So, we may now assume that M' is of Form (VI). For convenience let us rename the matrix M' as M. If $M_{11}M_{14} = M_{12}M_{13}$, let $\kappa = M_{13}/M_{11} = M_{14}/M_{12}$, then

$$M = \begin{pmatrix} M_{11} & M_{12} \\ M_{12} & M_{11} \end{pmatrix} \otimes \begin{pmatrix} 1 & \kappa \\ \kappa & 1 \end{pmatrix} = \begin{pmatrix} a & x \\ x & a \end{pmatrix} \otimes \begin{pmatrix} 1 & \kappa \\ \kappa & 1 \end{pmatrix}.$$

Similarly, if $M_{11}M_{12} = M_{13}M_{14}$, or $M_{11}M_{13} = M_{12}M_{14}$, then M is isomorphic to a tensor product under the flips $2 \leftrightarrow 4$, or $3 \leftrightarrow 4$, respectively. So, we have $M_{11}M_{12} \neq M_{13}M_{14}$, $M_{11}M_{13} \neq M_{12}M_{14}$, and $M_{11}M_{14} \neq M_{12}M_{13}$, since $\rho_{\texttt{tensor}}(M) \neq 0$.

We will now let the entries of M be generated by some $\{g_t\}_{t\in[d]}$. Using Lemma 4, we may replace M with some cM such that $e_{ijt} \ge 0$ for all $i, j \in [4], t \in [d]$. Since $M \in \text{Sym}_4(\mathbb{R}_{>0})$, we note that $e_{ij0} = 0$ for all $i, j \in [4]$.

We will now define the function $\widehat{\mathcal{T}_M} : \mathbb{R}^{d+1} \to \operatorname{Sym}_q^{\mathbb{R}}$, just as in Eq. (24), such that

$$\widehat{\mathcal{T}_M}(p, z_1, \dots, z_d)_{ij} = \mathcal{T}_M(p^{z_1}, \dots, p^{z_d})$$

We can pick some rational z_1^*, \ldots, z_d^* such that $\widehat{\mathcal{T}}_M(p, z_1^*, \ldots, z_d^*)$ satisfies the properties of M, that

 $M_{11}M_{12} \neq M_{13}M_{14}, \ M_{11}M_{13} \neq M_{12}M_{14}, \ M_{11}M_{14} = M_{12}M_{13}, \ \text{and} \ M_{ii} > M_{jk}$

for all $i \in [4]$, $j \neq k \in [4]$. We then let $Z^* \in \mathbb{Z}_{>0}$ such that $Z^* z_t^* \in \mathbb{Z}_{>0}$ for all $t \in [d]$. We can then define $\mathcal{T}_M^* : \mathbb{R} \to \operatorname{Sym}_4(\mathbb{R})$, as in Eq. (25) such that

$$\mathcal{T}_M^*(p)_{ij} = \widehat{\mathcal{T}_M}(p, Z^* \cdot z_i^*, \dots, Z^* \cdot z_d^*)_{ij} = p^{z_{ij}},$$

for some $z_{ij} \in \mathbb{Z}_{\geq 0}$ for all $i, j \in [4]$. We may assume from this construction that $z_{ii} > z_{jk}$ for all $i \in [4]$, and $j \neq k \in [4]$, $z_{11} + z_{12} \neq z_{13} + z_{14}$, $z_{11} + z_{13} \neq z_{12} + z_{14}$, and $z_{11} + z_{14} \neq z_{12} + z_{13}$.

We see that

$$\mathcal{T}_{M}^{*}(p) = \begin{pmatrix} p^{z_{11}} & p^{z_{12}} & p^{z_{13}} & p^{z_{14}} \\ p^{z_{12}} & p^{z_{11}} & p^{z_{14}} & p^{z_{13}} \\ p^{z_{13}} & p^{z_{14}} & p^{z_{11}} & p^{z_{12}} \\ p^{z_{14}} & p^{z_{13}} & p^{z_{12}} & p^{z_{11}} \end{pmatrix}$$

It can be verified that the eigenvalues of $\mathcal{T}_M^*(p)$ are:

$$\lambda_{1}(\mathcal{T}_{M}^{*}(p)) = p^{z_{11}} + p^{z_{12}} + p^{z_{13}} + p^{z_{14}},$$

$$\lambda_{2}(\mathcal{T}_{M}^{*}(p)) = p^{z_{11}} + p^{z_{12}} - p^{z_{13}} - p^{z_{14}},$$

$$\lambda_{3}(\mathcal{T}_{M}^{*}(p)) = p^{z_{11}} - p^{z_{12}} + p^{z_{13}} - p^{z_{14}},$$

$$\lambda_{4}(\mathcal{T}_{M}^{*}(p)) = p^{z_{11}} - p^{z_{12}} - p^{z_{13}} + p^{z_{14}}.$$
(34)

We will now define the function $\varphi_{\mathbf{x}} : (\mathbb{R}_{\neq 0})^4 \to \mathbb{R}$ such that $\varphi_{\mathbf{x}}(\alpha_1, \ldots, \alpha_4) = \prod_{i \in [4]} (\alpha_i)^{x_i}$. We see that

$$\varphi_{\mathbf{x}}(\lambda_1(\mathcal{T}_M^*(p),\ldots,\lambda_4(\mathcal{T}_M^*(p))=1\iff \phi_{\mathbf{x}}(\lambda_1(\mathcal{T}_M^*(p),\ldots,\lambda_4(\mathcal{T}_M^*(p))=0.$$

We will understand the behavior of $\phi_{\mathbf{x}}$ by studying $\varphi_{\mathbf{x}}.$

From the Taylor series expansions of $\lambda_i(\mathcal{T}^*_M(p))$ for each $i \in [4]$, we can see that for small $\delta > 0$,

$$\begin{split} \lambda_1(\mathcal{T}_M^*(e^{\delta})) &= 4 + \delta \left(z_{11} + z_{12} + z_{13} + z_{14} \right) + O(\delta^2), \\ \lambda_2(\mathcal{T}_M^*(e^{\delta})) &= \delta \left(z_{11} + z_{12} - z_{13} - z_{14} \right) + O(\delta^2), \\ \lambda_3(\mathcal{T}_M^*(e^{\delta})) &= \delta \left(z_{11} - z_{12} + z_{13} - z_{14} \right) + O(\delta^2), \\ \lambda_4(\mathcal{T}_M^*(e^{\delta})) &= \delta \left(z_{11} - z_{12} - z_{13} + z_{14} \right) + O(\delta^2). \end{split}$$

From our choice of z_{ij} , we can see that the coefficients of δ in each of the eigenvalues is non-zero. So, the following limits all exist and are non-zero constants,

$$\lim_{\delta \to 0} \frac{\lambda_i(\mathcal{T}_M^*(e^{\delta}))}{\delta} \neq 0,$$

for all $i \in \{2, 3, 4\}$. For the given $\mathbf{x} \in \chi_4$,

$$\varphi_{\mathbf{x}}\left(\lambda_{1}(\mathcal{T}_{M}^{*}(e^{\delta})),\ldots,\lambda_{4}(\mathcal{T}_{M}^{*}(e^{\delta}))\right) = \delta^{x_{2}+x_{3}+x_{4}} \cdot \left(\lambda_{1}(\mathcal{T}_{M}^{*}(e^{\delta}))\right)^{x_{1}} \cdot \left(\frac{\lambda_{2}(\mathcal{T}_{M}^{*}(e^{\delta}))}{\delta}\right)^{x_{2}} \cdot \left(\frac{\lambda_{3}(\mathcal{T}_{M}^{*}(e^{\delta}))}{\delta}\right)^{x_{3}} \cdot \left(\frac{\lambda_{4}(\mathcal{T}_{M}^{*}(e^{\delta}))}{\delta}\right)^{x_{4}}.$$

So, we see that

$$\lim_{\delta \to 0} \varphi_{\mathbf{x}} \left(\lambda_1(\mathcal{T}_M^*(e^{\delta})), \dots, \lambda_4(\mathcal{T}_M^*(e^{\delta})) \right) = 4^{x_1} \lim_{\delta \to 0} \left[(\delta^{x_1 + x_2 + x_3}) \prod_{i=2,3,4} \left(\frac{\lambda_i(\mathcal{T}_M^*(e^{\delta}))}{\delta} \right)^{x_i} \right]$$

This implies that unless $x_2 + x_3 + x_4 = (-x_1) = 0$,

$$\lim_{\delta \to 0} \varphi_{\mathbf{x}} \left(\lambda_1(\mathcal{T}_M^*(e^{\delta})), \dots, \lambda_4(\mathcal{T}_M^*(e^{\delta})) \right) \neq 1,$$

which in turn implies that

$$\lim_{\delta \to 0} \phi_{\mathbf{x}}(\lambda_1(\mathcal{T}_M^*(e^{\delta})), \dots, \lambda_4(\mathcal{T}_M^*(e^{\delta}))) \neq 0.$$

So, we may assume that $x_1 = 0$. Since **x** has a support size > 2, and $x_2 + x_3 + x_4 = 0$, we now may assume that none of $x_2, x_3, x_4 = 0$.

Suppose $z_{12} = z_{13}$, then from Eq. (34), we see that

$$\lambda_2(\mathcal{T}_M^*(p)) = p^{z_{11}} - p^{z_{14}} = \lambda_3(\mathcal{T}_M^*(p)).$$

So, $\varphi_{\mathbf{x}} (\lambda_1(\mathcal{T}_M^*(p)), \dots, \lambda_4(\mathcal{T}_M^*(p))) = 1$ for all $p \in \mathbb{R}$ implies that $\lambda_2(\mathcal{T}_M^*(p))^{x_2+x_3} = \lambda_4(\mathcal{T}_M^*(p))^{x_2+x_3}$ for all $p \in \mathbb{R}$. If $x_2 + x_3 = 0$, then $x_4 = -(x_2 + x_3) = 0$ as well, and \mathbf{x} would have support size ≤ 2 , so we may assume otherwise. Without loss generality, we may assume that $x_2 + x_3 > 0$. So, we see that $\lambda_2(\mathcal{T}_M^*(p)) = \lambda_4(\mathcal{T}_M^*(p))$ for all $p \in \mathbb{R}$. By Eq. (34), this implies that

$$p^{z_{11}} - p^{z_{14}} = p^{z_{11}} - 2p^{z_{12}} + p^{z_{14}}$$

Simplifying this, we find that $z_{12} = z_{13} = z_{14}$ as well.

We also know that $z_{11} \neq z_{12}$. So, we see that $\mathcal{T}^*_M(p)$ is of the form

$$\mathcal{T}_{M^*}(p) = \begin{pmatrix} p^{z_{11}} & p^{z_{12}} & p^{z_{12}} & p^{z_{12}} \\ p^{z_{12}} & p^{z_{11}} & p^{z_{12}} & p^{z_{12}} \\ p^{z_{12}} & p^{z_{12}} & p^{z_{11}} & p^{z_{12}} \\ p^{z_{12}} & p^{z_{12}} & p^{z_{12}} & p^{z_{11}} \end{pmatrix}.$$

But then, it is already known from Corollary 8 that when p > 1, $Pl-GH(\mathcal{T}_{M^*}(p)) \leq Pl-GH(M)$ is #P-hard.

From the above proof we may now assume we are in the case where $z_{12} \neq z_{13}$. By the same reasoning, we may also assume that in fact, $z_{12} \neq z_{13} \neq z_{14}$ are pairwise distinct. Without loss of generality, we may assume that $x_2 > 0$. We can also assume by symmetry, that $x_3 \geq x_4$. So, the two cases we have to consider are: $x_3 > 0$ (which would imply that $x_4 = -(x_2 + x_3) < 0$), and $x_3 < 0$ (which would also imply that $x_4 \leq x_3 < 0$). Let us first consider the case where $x_3 > 0$. We see that if $\phi_{\mathbf{x}}(\lambda_1(\mathcal{T}_M^*(p)), \ldots, \lambda_4(\mathcal{T}_M^*(p))) = 0$, then

$$(p^{z_{11}} + p^{z_{12}} - p^{z_{13}} - p^{z_{14}})^{x_2} (p^{z_{11}} - p^{z_{12}} + p^{z_{13}} - p^{z_{14}})^{x_3} = (p^{z_{11}} - p^{z_{12}} - p^{z_{13}} + p^{z_{14}})^{x_2 + x_3}.$$

The highest degree term of both the LHS and the RHS are both equal to $p^{z_{11}(x_2+x_3)}$. The second highest degree term will depend on which of z_{12}, z_{13}, z_{14} is the largest. If $z_{12} > z_{13}, z_{14}$, then the second highest degree term of the LHS is $(x_2 - x_3)p^{z_{11}(x_2+x_3-1)+z_{12}}$, while the second highest degree term of the RHS is $(-x_2 - x_3)p^{z_{11}(x_2+x_3-1)+z_{12}}$. These terms are clearly not equal to each other. On the other hand, if $z_{13} > z_{12}, z_{14}$, then the second highest degree term of the LHS is $(-x_2 + x_3)p^{z_{11}(x_2+x_3-1)+z_{13}}$, while the second highest degree term of the RHS is $(-x_2 - x_3)p^{z_{11}(x_2+x_3-1)+z_{13}}$, which are also not equal to each other. Finally, if $z_{14} > z_{12}, z_{13}$, the second highest degree term of the RHS is $(x_2 + x_3)p^{z_{11}(x_2+x_3-1)+z_{14}}$, while the second highest degree term of the RHS is $(x_2 + x_3)p^{z_{11}(x_2+x_3-1)+z_{14}}$, while the second highest degree term of the RHS is $(x_2 + x_3)p^{z_{11}(x_2+x_3-1)+z_{14}}$, while the second highest degree term of the RHS is $(x_2 + x_3)p^{z_{11}(x_2+x_3-1)+z_{14}}$, while the second highest degree term of the RHS is $(x_2 + x_3)p^{z_{11}(x_2+x_3-1)+z_{14}}$, while the second highest degree term of the RHS is $(x_2 + x_3)p^{z_{11}(x_2+x_3-1)+z_{14}}$, while the second highest degree term of the RHS is $(x_2 + x_3)p^{z_{11}(x_2+x_3-1)+z_{14}}$, while the second highest degree term of the RHS is $(x_2 + x_3)p^{z_{11}(x_2+x_3-1)+z_{14}}$, while the second highest degree term of the RHS is not possible that $\phi_{\mathbf{x}}(\lambda_1(\mathcal{T}_M^*(p)), \ldots, \lambda_4(\mathcal{T}_M^*(p))) = 0$.

We will now consider the case where $x_3 < 0$. In this case, if $\phi_{\mathbf{x}}(\lambda_1(\mathcal{T}_M^*(p)), \ldots, \lambda_4(\mathcal{T}_M^*(p))) = 0$, then

$$(p^{z_{11}} + p^{z_{12}} - p^{z_{13}} - p^{z_{14}})^{-x_3 - x_4} = (p^{z_{11}} - p^{z_{12}} + p^{z_{13}} - p^{z_{14}})^{-x_3} (p^{z_{11}} - p^{z_{12}} - p^{z_{13}} + p^{z_{14}})^{-x_4}.$$

The highest degree term of both the LHS and the RHS are both equal to $p^{z_{11}(-x_3-x_4)}$. The second highest degree term will again depend on which of z_{12}, z_{13}, z_{14} is the largest. If $z_{12} > z_{13}, z_{14}$, then the second highest degree term of the LHS is $(-x_3 - x_4)p^{z_{11}(-x_3-x_4-1)+z_{12}}$, while the second

highest degree term of the RHS is $(x_3 + x_4)p^{z_{11}(-x_3-x_4-1)+z_{12}}$. These terms are clearly not equal to each other. On the other hand, if $z_{13} > z_{12}, z_{14}$, then the second highest degree term of the LHS is $(x_3 + x_4)p^{z_{11}(-x_3-x_4-1)+z_{13}}$, while the second highest degree term of the RHS is $(-x_3 + x_4)p^{z_{11}(-x_3-x_4-1)+z_{13}}$, which are also not equal to each other. Finally, if $z_{14} > z_{12}, z_{13}$, the second highest degree term of the LHS is $(x_3 + x_4)p^{z_{11}(-x_3-x_4-1)+z_{14}}$, while the second highest degree term of the RHS is $(x_3 - x_4)p^{z_{11}(-x_3-x_4-1)+z_{14}}$, which are not equal to each other either. This proves that it is not possible that $\phi_{\mathbf{x}}(\lambda_1(\mathcal{T}_M^*(p)), \ldots, \lambda_4(\mathcal{T}_M^*(p))) = 0$, even in this case.

So, we see that for all $\mathbf{x} \in \chi_4$ of support size greater than 2, $\phi_{\mathbf{x}}(\lambda_1(\mathcal{T}_M^*(p), \ldots, \lambda_4(\mathcal{T}_M^*(p))))$ is a non-zero polynomial. Moreover, this is true for all $\mathbf{y} \in \chi_q$ such that $y_{\sigma(i)} = x_i$, for some $\sigma \in S_4$. In particular, this means that there exists some $p^* \in \mathbb{R}$ such that

$$\Psi_{\mathbf{x}}(\mathcal{T}_M^*(p^*)) = \Phi_{\mathbf{x}}(\lambda_1(\mathcal{T}_M^*(p^*)), \dots, \lambda_4(\mathcal{T}_M^*(p^*))) \neq 0.$$

So, Lemma 28 allows us to find some $N = \mathcal{T}_M(\mathbf{p}^*) \in \mathfrak{R}(M, \mathcal{F} \cup \{\Psi_{\mathbf{x}}, \rho_{\texttt{tensor}}\}) \cap \text{Sym}_4^{\texttt{pd}}(\mathbb{R}_{>0})$. If N is diagonal distinct, then Theorem 68 implies that $\texttt{Pl-GH}(N) \leq \texttt{Pl-GH}(M)$ is #P-hard. So, we may assume that N is not diagonal distinct.

Let us now assume that N is not of Form (VI). Now, from Theorem 75, we see that either $P1-GH(N) \leq P1-GH(M)$ is #P-had, or there exists some $N' \in \mathfrak{R}(M, \mathcal{F} \cup \{\Psi_{\mathbf{x}}, \rho_{\mathtt{tensor}}\}) \cap \operatorname{Sym}_{4}^{pd}(\mathbb{R}_{>0})$, that is isomorphic to a matrix of Form (I), (III), or (IV), and does not have diagonal entries that are all identical to each other. But in that case, we note that since $\rho_{\mathtt{tensor}}(N') \neq 0$, Lemma 82, Lemma 85, and Lemma 95 imply that $P1-GH(N') \leq P1-GH(M)$ is #P-hard. Otherwise, N is the required matrix of Form (VI), gaining the crucial property that $\Psi_{\mathbf{x}}(N) \neq 0$.

We can now prove that Pl-GH(M), when M is of Form (VI) that is not isomorphic to a tensor product, must also be #P-hard.

Lemma 97. Let $M \in Sym_4^{pd}(\mathbb{R}_{>0})$ be of Form (VI) such that $\rho_{tensor}(M) \neq 0$. Then, Pl-GH(M) is #P-hard.

Proof. The proof is very similar to the proof of Lemma 85. The only difference is that instead of Lemma 83, we will have to use Lemma 96 to find the matrices N_i . The rest of the proof is identical.

We can finally prove the following theorem.

Theorem 98. Let $M \in Sym_4^{pd}(\mathbb{R}_{>0})$ such that $\rho_{tensor}(M) \neq 0$. Then, Pl-GH(M) is #P-hard.

Proof. If M is diagonal distinct, then Theorem 68 implies that P1-GH(M) is #P-hard. Otherwise, from Theorem 75, we know that either P1-GH(M) is #P-hard, or there exists some $M' \in \mathfrak{R}(M, \mathcal{F}_M \cup \{\rho_{\texttt{tensor}}\}) \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{>0})$ that is of Form (I), (III), (IV), or (VI). In any case, Lemma 82. Lemma 95, Lemma 85, or Lemma 97 allow us to prove that $P1-GH(M') \neq P1-GH(M)$ is #P-hard.

12 Dichotomy for 4×4 matrices

We will now first deal with matrices $M \in \text{Sym}_4^{\text{pd}}(\mathbb{R}_{>0})$ such that M is isomorphic to $A \otimes B$ for some $A, B \in \text{Sym}_2(\mathbb{R})$. We can assume without loss of generality, that in fact, $M = A \otimes B$. In this case, we see that given any graph G = (V, E),

$$Z_{A\otimes B}(G) = \sum_{\sigma:V \to [4]} \prod_{(u,v) \in E} (A \otimes B)_{\sigma(u)\sigma(v)} = \sum_{\sigma_1, \sigma_2:V \to [2]} \prod_{(u,v) \in E} \left((A \otimes B)_{\tau(\sigma_1(u), \sigma_2(u))\tau(\sigma_1(v), \sigma_2(v))} \right),$$

where $\tau(i, j) = 2(i-1) + j$ for all $i, j \in [2]$.

But then, we see that

$$Z_{A\otimes B}(G) = \sum_{\sigma_1, \sigma_2: V \to [2]} \prod_{(u,v) \in E} A_{\sigma_1(u)\sigma_1(v)} B_{\sigma_2(u)\sigma_2(v)}$$
$$= \left(\sum_{\sigma_1: V \to [2]} \prod_{(u,v) \in E} A_{\sigma_1(u)\sigma_1(v)} \right) \cdot \left(\sum_{\sigma_2: V \to [2]} \prod_{(u,v) \in E} B_{\sigma_2(u)\sigma_2(v)} \right)$$
$$= (Z_A(G)) \cdot (Z_B(G)) .$$

So, we see that when $M = A \otimes B$, it is in some sense, equivalent to problems of a smaller size. We shall formalize this notion shortly. In the meantime, we can now immediately prove the following lemma.

Lemma 99. Let $M \in Sym_4^{pd}(\mathbb{R}_{>0})$, such that $M = A \otimes B$ for some $A, B \in Sym_2(\mathbb{R})$. If Pl-GH(A) is polynomial time tractable, then $Pl-GH(M) \equiv Pl-GH(B)$.

Proof. Let $M = A \otimes B$. If A or B has any zero entries entries, then M would have zero entries, so we see that A and B cannot have any zero entries. Now, if we assume that A and B have entries a and b of the opposite signs, then ab < 0 would be an entry of M. So, all the entries of A and B must have the same sign. If this sign is –, then we may replace A and B with -A and -B. So, we may assume that $A, B \in \text{Sym}_2(\mathbb{R}_{>0})$. This implies that for any graph G = (V, E), $Z_A(G) > 0$, and $Z_B(G) > 0$. If we now had oracle access to Pl-GH(B), we can use it to compute $Z_M(G) = Z_A(G) \cdot Z_B(G)$ for all planar G = (V, E). Similarly, if we had oracle access to Pl-GH(M), we could use it to compute $Z_B(G) = \frac{Z_M(G)}{Z_A(G)}$ for any planar G = (V, E).

Remark. Let $M \in Sym_4^{pd}(\mathbb{R}_{>0})$, such that $M = A \otimes B$ for some $A, B \in Sym_2(\mathbb{R})$. Then, as we saw in the proof of Lemma 99, we may assume that in fact, $A, B \in Sym_2(\mathbb{R}_{>0})$. Now, we may let (λ_1, λ_2) be the eigenvalues of A, and (μ_1, μ_2) be the eigenvalues of B. This implies that the eigenvalues of $M = A \otimes B$ are:

 $\lambda_1\mu_1, \quad \lambda_1\mu_2, \quad \lambda_2\mu_1, \quad \lambda_2\mu_2.$

We may assume without loss of generality that $\lambda_1 \geq \lambda_2$, and $\mu_1 \geq \mu_2$. Since $A \in Sym_2(\mathbb{R}_{>0})$, we know from the Perron-Frobenius Theorem, that in fact, $\lambda_1 > |\lambda_2|$. Similarly, since $B \in Sym_2(\mathbb{R}_{>0})$, we see that $\mu_1 > |\mu_2|$. Now, if $\lambda_2 \leq 0$, we note that $\lambda_2\mu_1 \leq 0$ would be an eigenvalue of $A \otimes B = M$, which contradicts our assumption that $M \in Sym_4^{pd}(\mathbb{R}_{>0})$. So, in fact, $\lambda_2 > 0$. Similarly, $\mu_2 > 0$ as well. This means that if $M \in Sym_4^{pd}(\mathbb{R}_{>0})$ is such that $M = A \otimes B$ for some $A, B \in Sym_2(\mathbb{R})$, we may assume that $A, B \in Sym_2^{pd}(\mathbb{R}_{>0})$. Similarly, if $M \in Sym_4^F(\mathbb{R}_{>0})$, and $M = A \otimes B$, we may assume $A, B \in Sym_2(\mathbb{R}_{>0})$. If A or B has an eigenvalue 0, so would $M = A \otimes B$. So, if $M \in Sym_4^F(\mathbb{R}_{>0})$ is such that $M = A \otimes B$ for some $A, B \in Sym_2(\mathbb{R})$, we may assume that $A, B \in Sym_2^F(\mathbb{R}_{>0})$.

We will now state the following theorem [GW20]:

Theorem 100. The problem Pl-GH(M) is #P-hard, for $M = \begin{pmatrix} x & y \\ y & z \end{pmatrix} \in Sym_2(\mathbb{R}_{\geq 0})$, unless $xz = y^2$, y = 0, or x = z, in which case Pl-GH(M) is in polynomial time.

This theorem now allows us to prove the following lemma immediately.

Lemma 101. Let $M \in Sym_4^{pd}(\mathbb{R}_{>0})$, such that $M = A \otimes B$ for some $A, B \in Sym_2^{pd}(\mathbb{R}_{>0})$. Then Pl-GH(M) is polynomial time tractable if $A_{11} = A_{22}$, and $B_{11} = B_{22}$. If $A_{11} = A_{22}$, but $B_{11} \neq B_{22}$, or if $B_{11} = B_{22}$, but $A_{11} \neq A_{22}$, then Pl-GH(M) is #P-hard.

Proof. The tractability part follows from Theorem 100. For the #P-hardness part, by symmetry, let us assume that $A_{11} = A_{22}$, but $B_{11} \neq B_{22}$. From Theorem 100, we see that since $A_{11} = A_{22}$, Pl-GH(A) is polynomial time tractable. On the other hand, we apply Theorem 100 to Pl-GH(B), and see that (1) $B_{11}B_{22} \neq (B_{12})^2$ since $B \in \text{Sym}_2^{\text{pd}}(\mathbb{R}_{>0})$ has full rank, (2) $B_{12} \neq 0$ as $B \in \text{Sym}_2(\mathbb{R}_{>0})$, and (3) $B_{11} \neq B_{22}$ as given. Hence, Pl-GH(B) is #P-hard. Now, since Pl-GH(A) is polynomially tractable, Lemma 99 implies that Pl-GH(M) is #P-hard.

To deal with the case where $A_{11} \neq A_{22}$ and $B_{11} \neq B_{22}$ will require just a little bit more work.

Lemma 102. Let $M \in Sym_4^{pd}(\mathbb{R}_{>0})$, such that $M = A \otimes B$ for some $A, B \in Sym_2^{pd}(\mathbb{R}_{>0})$, that satisfy $A_{11} \neq A_{22}$ and $B_{11} \neq B_{22}$. Let \mathcal{F} be a countable set of $Sym_4(\mathbb{R})$ -polynomials such that $F(M) \neq 0$ for all $F \in \mathcal{F}$. Then, either Pl-GH(M) is #P-hard, or there exists some diagonal distinct $N \in \mathfrak{R}(M, \mathcal{F}) \cap Sym_4^{pd}(\mathbb{R}_{>0})$, such that $\rho_{tensor}(N) = 0$.

Proof. We may assume without loss of generality that $A_{11} > A_{22}$, and that $B_{11} > B_{22}$, by permuting the rows and columns of A and B (and correspondingly, M). We can see that $M_{11} = A_{11}B_{11} > M_{ij} > M_{44}$ for any $(i, j) \neq (1, 1), (4, 4)$. So, rows 1 and i are not order identical for any $i \neq 1$. Similarly, rows 4 and i are not order identical for any $i \neq 4$. We have $M_{11} > M_{22} > M_{44}$ and $M_{11} > M_{33} > M_{44}$. If $M_{22} \neq M_{33}$, then M is already diagonal distinct, and we are done. So, we may assume that $M_{22} = M_{33}$. Let us first assume that rows 2 and 3 are not order identical. Then Mis p.o. distinct, and by Corollary 62 there exists some diagonal distinct $N \in \mathfrak{R}(M, \mathcal{F}) \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{>0})$. If $\rho_{\text{tensor}}(N) \neq 0$, then Theorem 98 immediately implies that $\text{Pl-GH}(N) \leq \text{Pl-GH}(M)$ is #P-hard. On the other hand, if $\rho_{\text{tensor}}(N) = 0$ then N is our required matrix.

Finally, we consider the case where rows 2 and 3 are order identical. We see that the multi-set of elements in row 2 of M is $\{A_{11}B_{12}, A_{11}B_{22}, A_{12}B_{12}, A_{12}B_{22}\}$, and the multi-set of elements in row 3 of M is $\{A_{12}B_{11}, A_{12}B_{12}, A_{22}B_{11}, A_{22}B_{12}\}$. We can see that $A_{12}B_{12}$ and $A_{11}B_{22} = A_{22}B_{11}$ (which are $M_{22} = M_{33}$) appear in both row 2 and row 3. So, it must be the case that either $A_{11}B_{12} = A_{22}B_{12}$ and $A_{12}B_{22} = A_{12}B_{11}$, or $A_{11}B_{12} = A_{12}B_{11}$ and $A_{12}B_{22} = A_{22}B_{12}$. Since we

have assumed that $A_{11} > A_{22}$, and that $B_{11} > B_{22}$, we can see the first option here is not possible. So, we see that in fact,

$$\frac{A_{11}}{A_{12}} = \frac{B_{11}}{B_{12}}$$
, and $\frac{A_{22}}{A_{12}} = \frac{B_{22}}{B_{12}}$

But this means that $B = \kappa \cdot A$ for some constant $\kappa > 0$, and in this case, we see that for any graph G = (V, E),

$$Z_M(G) = Z_{A \otimes B}(G) = Z_A(G) \cdot Z_{\kappa \cdot A}(G) = \kappa^{|E|} \cdot (Z_A(G))^2.$$

Since $A \in \text{Sym}_2(\mathbb{R}_{>0})$, we see that for any graph G = (V, E), $Z_A(G) > 0$. So, given oracle access to Pl-GH(M), we can compute $Z_M(G)$ for any planar graph G = (V, E), and use it to compute

$$Z_A(G) = \sqrt{\frac{Z_M(G)}{\kappa^{|E|}}}.$$

So, we see that $\text{Pl-GH}(A) \leq \text{Pl-GH}(M)$. Since $A \in \text{Sym}_2^{\text{pd}}(\mathbb{R}_{>0})$, and $A_{11} > A_{22}$, Theorem 100 implies that $\text{Pl-GH}(A) \leq \text{Pl-GH}(M)$ is #P-hard.

Lemma 103. Let $M \in Sym_4^{pd}(\mathbb{R}_{>0})$ be a diagonal distinct matrix, such that $M = A \otimes B$ for some $A, B \in Sym_2^{pd}(\mathbb{R})$. Then, Pl-GH(M) is #P-hard.

Proof. Since M is diagonal distinct, it must be the case that $A_{11} \neq A_{22}$, and $B_{11} \neq B_{22}$. By permuting the rows and columns of A and B (and correspondingly, the rows and columns of M), we may assume that $A_{11} > A_{22}$, and that $B_{11} > B_{22}$.

Let \mathcal{L}_M be the lattice of the eigenvalues of M, and let \mathcal{B} be a lattice basis of \mathcal{L}_M . If there is any $\mathbf{x} \in \mathcal{B}$ that is confluent, we can use the combination of Lemma 59, Corollary 50, and Theorem 46, to find some diagonal distinct $N_1 \in \mathfrak{R}(M, \mathcal{F}_M \cup \{\Psi_{\mathbf{x}}, \rho_{\mathsf{diag}}\}) \cap \operatorname{Sym}_4^{\mathsf{pd}}(\mathbb{R}_{>0})$.

We know from Lemma 26 that after repeating this step at most 4 times, we will have some diagonal distinct $N \in \text{Sym}_{4}^{\text{pd}}(\mathbb{R}_{>0})$, such that $\text{Pl-GH}(N) \leq \text{Pl-GH}(M)$, and

$$\mathbf{0} \neq \mathbf{x} \in \overline{\mathcal{L}_N} \implies \mathbf{x} \text{ is not confluent.}$$

Once again, we see that if $\rho_{\text{tensor}}(N) \neq 0$, Theorem 98 immediately implies that $\text{Pl-GH}(N) \leq \text{Pl-GH}(M)$ is #P-hard, so let us assume $\rho_{\text{tensor}}(N) = 0$. Now, since any non-zero $\mathbf{x} \in \overline{\mathcal{L}}_N$ is not confluent, Lemma 64 implies that

$$\overline{\mathcal{L}_N} \subseteq \left\{ (x_1, x_2, x_3, x_4) \in \chi_4 \ \Big| \ |x_1| = |x_2| = |x_3| = |x_4| \right\}.$$
(35)

We can now assume (after some permutation of the rows and columns of N) that $N = A' \otimes B'$ for some $A', B' \in \operatorname{Sym}_2^{pd}(\mathbb{R}_{>0})$. Since N is diagonal distinct, we see that A' and B' are diagonal distinct as well. We may let $A' = H_1 D_1 H_1^T$, and $B' = H_2 D_2 H_2^T$, for some orthogonal matrices H_1, H_2 , and diagonal matrices D_1, D_2 . Since $N = A' \otimes B'$, this implies that $N = (H_1 \otimes H_2)(D_1 \otimes D_2)(H_1 \otimes H_2)^T$. We may further assume that the eigenvalues of A' and B' are (λ_1, λ_2) , and (μ_1, μ_2) respectively. This implies that the eigenvalues of $N = A' \otimes B'$ are:

$$\lambda_1\mu_1, \quad \lambda_1\mu_2, \quad \lambda_2\mu_1, \quad \lambda_2\mu_2.$$

We can clearly see that $(\lambda_1\mu_1)(\lambda_2\mu_2) = (\lambda_1\mu_2)(\lambda_2\mu_1)$. So, $(1, -1, -1, 1) \in \mathcal{L}_N$. Let us now consider any $\mathbf{0} \neq \mathbf{x} \in \mathcal{L}_N$. From Eq. (35), we know that it must be the case that $|x_1| = |x_2| = |x_3| = |x_4|$. We may assume without loss of generality, that $x_1 > 0$. If $x_2 = x_1$, then \mathbf{x} being in χ_4 implies that $x_3 = x_4 = -x_1$. Then $x_1(1, 1, -1, -1) \in \mathcal{L}_N$. This implies that $x_1(1, -1, -1, 1) + x_1(1, 1, -1, -1) =$ $x_1(2, 0, -2, 0) \in \mathcal{L}_N \subseteq \overline{\mathcal{L}_N}$ which contradicts Eq. (35). So, we conclude that $x_2 = -x_1$. By a symmetric argument $x_3 = -x_1$. Since $\mathbf{x} \in \chi_4$, this forces $x_4 = x_1$. Hence $\mathbf{x} = x_1(1, -1, -1, 1)$. Therefore, we conclude that if $\mathbf{x} \in \mathcal{L}_N$, then \mathbf{x} must be a multiple of (1, -1, -1, 1). In other words,

$$\mathcal{L}_N = \{ c \cdot (1, -1, -1, 1) \mid c \in \mathbb{Z} \}.$$

We now define $N' = A' \otimes I$, where I is the 2 by 2 identity matrix. Recall that $N = A' \otimes B' = (H_1 \otimes H_2)(D_1 \otimes D_2)(H_1 \otimes H_2)^T$, and we can now see that $N' = A' \otimes I = (H_1 \otimes H_2)(D_1 \otimes I)(H_1 \otimes H_2)^T$. So, if we let $H = H_1 \otimes H_2$, $D = D_1 \otimes D_2$, and $D' = D_1 \otimes I$, we see that $N = HDH^T$, and $N' = HD'H^T$. Moreover, by construction, we note that the eigenvalues of N' are: $(\lambda_1, \lambda_2, \lambda_1, \lambda_2)$. So, $(1, -1, -1, 1) \in \mathcal{L}_{N'}$, which implies that

$$\mathcal{L}_N \subseteq \mathcal{L}_{N'}.$$

But then, Lemma 13 implies that $Pl-GH(N') \leq Pl-GH(N)$. But by construction of N', we see that for any graph G = (V, E),

$$Z_{N'}(G) = Z_{A' \otimes I}(G) = Z_{A'}(G) \cdot Z_I(G).$$

We note that for any connected graph G = (V, E), $Z_I(G) = 2$, as I is the 2 by 2 identity matrix. This means that if we had oracle access to Pl-GH(N'), we could use it to compute $Z_{A'}(G)$ for any connected (and thus all) planar graphs G = (V, E). So, we see that $\text{Pl-GH}(A') \leq \text{Pl-GH}(N')$. Since $(A')_{11} \neq (A')_{22}$ by construction, Theorem 100 implies that Pl-GH(A') is #P-hard. This proves that $\text{Pl-GH}(A') \leq \text{Pl-GH}(N') \leq \text{Pl-GH}(N) \leq \text{Pl-GH}(N) \leq \text{Pl-GH}(N) = \frac{1}{2}$.

Remark. Let $M = A \otimes B$ for any $M \in Sym_4(\mathbb{R}_{>0})$. In the proof above, we made use of the argument that if the matrices N we obtained using the theorems and lemmas from the previous section satisfy the equation $\rho_{\text{tensor}}(N) \neq 0$, then we can use Theorem 98 to prove the #P-hardness of $\text{Pl-GH}(N) \leq$ Pl-GH(M). There is an alternative route in this proof. This alternative route starts by observing that, in fact, if $\rho_{\text{tensor}}(M) = 0$, then any matrix N that we obtain using the techniques from the rest of this paper will also satisfy the equation $\rho_{\text{tensor}}(N) = 0$. This is because of the following set of properties: $T_n(M) = T_n(A) \otimes T_n(B)$, $S_n(M) = S_n(A) \otimes S_n(B)$, and $R_n(M) = R_n(A) \otimes R_n(B)$, for all $n \geq 1$. Moreover, if we let the entries of M be generated by some $\{g_t\}_{t \in [d]}$, this implies that

$$\rho_{\texttt{tensor}}(M) = 0 \implies \rho_{\texttt{tensor}}(\mathcal{T}_M(\mathbf{p})) = 0$$

for all $\mathbf{p} \in \mathbb{R}^d$. Similarly,

$$\rho_{\texttt{tensor}}(M) = 0 \implies \rho_{\texttt{tensor}}(\mathcal{S}_M(\theta))) = 0, \text{ and } \rho_{\texttt{tensor}}(\mathcal{R}_M(\theta)) = 0,$$

for all $\theta \in \mathbb{R}$.

Lemma 102 and Lemma 103 form the last piece of the puzzle for matricces in $\text{Sym}_4^{pd}(\mathbb{R}_{>0})$, and now we can prove the following theorem.

Theorem 104. Let $M \in Sym_4^{pd}(\mathbb{R}_{>0})$. Then Pl-GH(M) is #P-hard, unless M is isomorphic to $A \otimes B$ for some $A, B \in Sym_2^{pd}(\mathbb{R}_{>0})$ such that $A_{11} = A_{22}$, and $B_{11} = B_{22}$, in which case, Pl-GH(M) is polynomial time tractable.

Proof. We know from Theorem 98 that Pl-GH(M) is #P-hard, unless $\rho_{tensor}(M) = 0$. But, if $\rho_{tensor}(M) = 0$ then (by the defining property of ρ_{tensor}) M is isomorphic to $A \otimes B$ for some $A, B \in Sym_2^{pd}(\mathbb{R}_{>0})$. By permuting the rows and columns of M, we may assume that in fact, $M = A \otimes B$.

We will now consider the diagonal entries of A and B. If $A_{11} = A_{22}$, but $B_{11} \neq B_{22}$, or if $B_{11} = B_{22}$, but $A_{11} \neq A_{22}$, Lemma 101 implies that Pl-GH(M) is #P-hard. If $A_{11} \neq A_{22}$, and $B_{11} \neq B_{22}$, Lemma 102 implies that either Pl-GH(M) is #P-hard, or there exists some diagonal distinct $N \in \mathfrak{R}(M, \mathcal{F}_M) \cap \operatorname{Sym}_4^{pd}(\mathbb{R}_{>0})$ such that $\rho_{\text{tensor}}(N) = 0$. But from Lemma 103, we see that Pl-GH(N) \leq Pl-GH(M) is #P-hard. So, we see that even if M is isomorphic to some $A \otimes B$, unless $A_{11} = A_{22}$ and $B_{11} = B_{22}$, Pl-GH(M) is #P-hard. Moreover, if $A_{11} = A_{22}$, and $B_{11} = B_{22}$, we see from Lemma 101 that Pl-GH(M) is polynomially tractable.

We can extend this dichotomy to all full rank matrices with some additional effort. When we consider $M \in \operatorname{Sym}_{4}^{\mathsf{F}}(\mathbb{R}_{>0})$, we come across a family of matrices for which the gadgets we have constructed so far are insufficient for proving the #P-hardness of Pl-GH(M). So, we will need to introduce a new edge gadget. Given any graph G = (V, E), we will construct the graph BG by replacing each edge of G with the gadget in Fig. 3a. Clearly, this gadget preserves planarity. Moreover, we note that for any $M \in \operatorname{Sym}_{q}(\mathbb{R})$,

$$Z_M(BG) = \sum_{\sigma: V \to [q]} \prod_{(u,v) \in E} \sum_{a,b \in [q]} M_{\sigma(u)a} M_{\sigma(u)b} M_{\sigma(v)a} M_{\sigma(v)b} M_{ab} = \sum_{\sigma: V \to [q]} \prod_{(u,v) \in E} (BM)_{\sigma(u)\sigma(v)},$$

where $BM \in \text{Sym}_q(\mathbb{R})$ such that for all $i, j \in [q]$,

$$(BM)_{ij} = \sum_{a,b\in[q]} M_{ia}M_{ib}M_{ja}M_{jb}M_{ab}$$

Therefore, we see that $Pl-GH(BM) \leq Pl-GH(M)$ for all $M \in \text{Sym}_q(\mathbb{R})$. We will first use this gadget to prove the #P-hardness of one special family of matrices in $\text{Sym}_4^F(\mathbb{R}_{>0})$.

Lemma 105. Let $M \in Sym_4^F(\mathbb{R}_{>0})$ such that

$$M = \begin{pmatrix} p^2 & pq & pq & q^2 \\ pq & q^2 & p^2 & pq \\ pq & p^2 & q^2 & pq \\ q^2 & pq & pq & p^2 \end{pmatrix}$$

where $p \neq q \in \mathbb{R}_{>0}$. Then, Pl-GH(M) is #P-hard.



(a) The edge gadget B. (b) An example graph G, and the graph BG.

Figure 3: The edge gadget B

Proof. We will consider $B(T_n(M))$ for $n \ge 1$. From the definitions, we can verify that

$$\begin{split} &(B(T_nM))_{11} = (B(T_nM))_{44} = p^{10n} + 4p^{7n}q^{3n} + 2p^{6n}q^{4n} + 4p^{4n}q^{6n} + 4p^{3n}q^{7n} + p^{2n}q^{8n}, \\ &(B(T_nM))_{22} = (B(T_nM))_{33} = p^{8n}q^{2n} + 4p^{7n}q^{3n} + 4p^{6n}q^{4n} + 2p^{4n}q^{6n} + 4p^{3n}q^{7n} + q^{10n}, \\ &(B(T_nM))_{12} = (B(T_nM))_{13} = p^{8n}q^{2n} + 2p^{7n}q^{3n} + 3p^{6n}q^{4n} + 4p^{5n}q^{5n} + 3p^{4n}q^{6n} + 2p^{3n}q^{7n} + p^{2n}q^{8n}, \\ &(B(T_nM))_{24} = (B(T_nM))_{34} = p^{8n}q^{2n} + 2p^{7n}q^{3n} + 3p^{6n}q^{4n} + 4p^{5n}q^{5n} + 3p^{4n}q^{6n} + 2p^{3n}q^{7n} + p^{2n}q^{8n}, \\ &(B(T_nM))_{24} = (B(T_nM))_{34} = p^{8n}q^{2n} + 2p^{7n}q^{3n} + 3p^{6n}q^{4n} + 4p^{5n}q^{5n} + 3p^{4n}q^{6n} + 2p^{3n}q^{7n} + p^{2n}q^{8n}, \\ &(B(T_nM))_{14} = (B(T_nM))_{23} = 4p^{6n}q^{4n} + 8p^{5n}q^{5n} + 4p^{4n}q^{6n}. \end{split}$$

So, it is seen that

$$\begin{split} \det(B(T_nM)) &= p^{36n}q^{4n} + 8p^{35n}q^{5n} + 20p^{34n}q^{6n} + 8p^{33n}q^{7n} - 32p^{32n}q^{8n} \\ &\quad -40p^{31n}q^{9n} - 100p^{30n}q^{10n} - 296p^{29n}q^{11n} - 84p^{28n}q^{12n} \\ &\quad +840p^{27n}q^{13n} + 1204p^{26n}q^{14n} + 72p^{25n}q^{15n} - 1440p^{24n}q^{16n} \\ &\quad -2088p^{23n}q^{17n} - 1124p^{22n}q^{18n} + 1496p^{21n}q^{19n} + 3110p^{20n}q^{20n} \\ &\quad + 1496p^{19n}q^{21n} - 1124p^{18n}q^{22n} - 2088p^{17n}q^{23n} - 1440p^{16n}q^{24n} \\ &\quad + 72p^{15n}q^{25n} + 1204p^{14n}q^{26n} + 840p^{13n}q^{27n} - 84p^{12n}q^{28n} \\ &\quad - 296p^{11n}q^{29n} - 100p^{10n}q^{30n} - 40p^{9n}q^{31n} - 32p^{8n}q^{32n} \\ &\quad + 8p^{7n}q^{33n} + 20p^{6n}q^{34n} + 8p^{5n}q^{35n} + p^{4n}q^{36n}. \end{split}$$

We will define the polynomial $f : \mathbb{R} \to \mathbb{R}$ such that

$$\begin{split} f(x) &= x^{36} + 8x^{35} + 20x^{34} + 8x^{33} - 32x^{32} - 40x^{31} - 100x^{30} - 296x^{29} - 84x^{28} \\ &+ 840x^{27} + 1204x^{26} + 72x^{25} - 1440x^{24} - 2088x^{23} + 1124x^{22} + 1496x^{21} + 3110x^{20} \\ &+ 1496x^{19} + 1124x^{18} - 2088x^{17} - 1440x^{16} + 72x^{15} + 1204x^{14} + 840x^{13} \\ &- 84x^{12} - 296x^{11} - 100x^{10} - 40x^9 - 32x^8 + 8x^7 + 20x^6 + 8x^5 + x^4. \end{split}$$

By construction, $\det(B(T_nM)) = q^{40n} f((p/q)^n)$. Since f is a non-zero polynomial, we know that it has finitely many real roots. Since $p \neq q \in \mathbb{R}_{>0}$, we know that if p > q, then $(p/q)^{n+1} > (p/q)^n > 1$

for all $n \ge 1$, and if p < q, then $0 < (p/q)^{n+1} < (p/q)^n$ for all $n \ge 1$. In particular, this implies that for large enough $n \ge 1$, $\det(B(T_n M)) = q^{40n} f((p/q)^n) \ne 0$.

Therefore, we see that for large enough $n \geq 1$, $B(T_nM) \in \operatorname{Sym}_4^F(\mathbb{R}_{>0})$. We will now let $N_n = (B(T_nM))^2$ for all $n \geq 1$. We see that $N_n \in \operatorname{Sym}_4^{pd}(\mathbb{R}_{>0})$ for large enough n. We will now define the $\operatorname{Sym}_4(\mathbb{R})$ -polynomials ξ_1, ξ_2, ξ_3 such that $\xi_1 : N \mapsto (N_{11}N_{14} - N_{12}N_{13}), \xi_2 : N \mapsto (N_{11}N_{13} - N_{12}N_{14}),$ and $\xi_3 : N \mapsto (N_{11}N_{12} - N_{13}N_{14})$. From the construction of ρ_{tensor} in Lemma 65, we can see that if $\xi_1(N) \neq 0$, then $\rho_{\text{tensor}}(N) \neq 0$. We note that the matrix N^{σ} is defined such that $(N^{\sigma})_{ij} = N_{\sigma(i)\sigma(j)}$ for any $\sigma \in S_4$. In fact, given any $\sigma \in S_4$ such that $\{\sigma(1), \sigma(4)\} = \{1, 4\}, \text{ or } \{2, 3\},$ we see that $\xi_1(N) \neq 0$ implies that $\rho_{\text{tensor}}(N^{\sigma}) \neq 0$. Similarly, given any $\sigma \in S_4$ such that $\{\sigma(1), \sigma(3)\} = \{1, 4\},$ or $\{2, 3\}$, we see that $\xi_2(N) \neq 0$ implies that $\rho_{\text{tensor}}(N^{\sigma}) \neq 0$, and given any $\sigma \in S_4$ such that $\{\sigma(1), \sigma(2)\} = \{1, 4\}, \text{ or } \{2, 3\},$ we see that $\xi_3(N) \neq 0$ implies that $\rho_{\text{tensor}}(N^{\sigma}) \neq 0$. Summing up, we find that if $\xi_i(N) \neq 0$ for all $i \in [3]$, then $\rho_{\text{tensor}}(N) \neq 0$.

We will now show that for large enough $n \ge 1$, $\xi_i(N_n) \ne 0$ for all $i \in [3]$. To do this, we will compute $\xi_i(N_n)$ for all $n \ge 1$. It can be verified that

$$\begin{split} \xi_1(N_n) &= \xi_2(N_n) = p^{38n}q^{2n} + 2p^{37n}q^{3n} + 4p^{36n}q^{4n} + 22p^{35n}q^{5n} + 48p^{34n}q^{6n} \\ &+ 82p^{33n}q^{7n} + 204p^{32n}q^{8n} + 358p^{31n}q^{9n} + 396p^{30n}q^{10n} \\ &+ 394p^{29n}q^{11n} + 164p^{28n}q^{12n} - 722p^{27n}q^{13n} - 1968p^{26n}q^{14n} \\ &- 2886p^{25n}q^{15n} - 2740p^{24n}q^{16n} - 610p^{23n}q^{17n} + 2918p^{22n}q^{18n} \\ &+ 5382p^{21n}q^{19n} + 5100p^{20n}q^{20n} + 2242p^{19n}q^{21n} - 1648p^{18n}q^{22n} \\ &- 4042p^{17n}q^{23n} - 3644p^{16n}q^{24n} - 1678p^{15n}q^{25n} + 140p^{14n}q^{26n} \\ &+ 1006p^{13n}q^{27n} + 876p^{12n}q^{28n} + 378p^{11n}q^{29n} + 112p^{10n}q^{30n} \\ &+ 62p^{9n}q^{31n} + 36p^{8n}q^{32n} + 10p^{7n}q^{33n} + p^{6n}q^{34n}, \text{ and} \\ \end{split}$$

We can now define the polynomials $f_1, f_3 : \mathbb{R} \to \mathbb{R}$ such that

$$\begin{split} f_1(x) &= x^{38} + 2x^{37} + 4x^{36} + 22x^{35} + 48x^{34} + 82x^{33} + 204x^{32} + 358x^{31} \\ &\quad + 396x^{30} + 394x^{29} + 164x^{28} - 722x^{27} - 1968x^{26} - 2886x^{25} \\ &\quad - 2740x^{24} - 610x^{23} + 2918x^{22} + 5382x^{21} + 5100x^{20} + 2242x^{19} \\ &\quad - 1648x^{18} - 4042x^{17} - 3644x^{16} - 1678x^{15} + 140x^{14} + 1006x^{13} \\ &\quad + 876x^{12} + 378x^{11} + 112x^{10} + 62x^9 + 36x^8 + 10x^7 + x^6, \end{split}$$

 $-44p^{7n}q^{33n} - 12p^{6n}q^{34n} - 4p^{5n}q^{35n} - p^{4n}q^{36n}$

$$\begin{split} f_3(x) &= 9x^{36} + 20x^{35} + 16x^{34} + 108x^{33} + 250x^{32} + 188x^{31} + 484x^{30} \\ &+ 1236x^{29} + 966x^{28} + 724x^{27} + 1736x^{26} - 244x^{25} \\ &- 4638x^{24} - 4196x^{23} - 2388x^{22} - 5356x^{21} - 4616x^{20} \\ &+ 4060x^{19} + 9088x^{18} + 5604x^{17} + 1454x^{16} + 468x^{15} \\ &- 4x^{14} - 996x^{13} - 1478x^{12} - 1220x^{11} - 728x^{10} \\ &- 348x^9 - 138x^8 - 44x^7 - 12x^6 - 4x^5 - x^4. \end{split}$$

By construction, we see that $\xi_1(N_n) = \xi_2(N_n) = q^{40n} f_1((p/q)^n)$, and $\xi_3(N_n) = q^{40n} f_3((p/q)^n)$. Since f_1 and f_3 are non-zero polynomials, we know that they have finitely many real roots. We are given $p \neq q \in \mathbb{R}_{>0}$. So, $(p/q)^{n+1} > (p/q)^n$ for all $n \geq 1$, or $(p/q)^{n+1} < (p/q)^n$ for all $n \geq 1$. In either case, we see that for large enough $n \geq 1$, $\xi_1(N_n) = \xi_2(N_n) \neq 0$, and $\xi_3(N_n) \neq 0$.

So, there exists some $n^* \geq 1$ such that $N_{n^*} \in \text{Sym}_4^{\text{pd}}(\mathbb{R}_{>0})$, and $\rho_{\text{tensor}}(N_{n^*}) \neq 0$. From Theorem 98, we know that $\text{Pl-GH}(N_{n^*}) \leq \text{Pl-GH}(B(T_{n^*}M)) \leq \text{Pl-GH}(T_{n^*M}) \leq \text{Pl-GH}(M)$ is #P-hard.

With the special case taken care of, we return to the more general case where M is "close" to being a tensor product. We will need a few more lemmas.

Lemma 106. Let $M \in Sym_4^{\mathsf{F}}(\mathbb{R}_{>0})$, such that $\rho_{\texttt{tensor}}(M) \neq 0$, but $\varrho_{\texttt{tensor}}(M^2) = 0$. Then, either P1-GH(M) is #P-hard, or $M = KDK^{\mathsf{T}}$, where

$$K = \frac{1}{2} \begin{pmatrix} 1 & u & v & 1 \\ 1 & -v & u & -1 \\ 1 & v & -u & -1 \\ 1 & -u & -v & 1 \end{pmatrix}$$

for some $u, v \in \mathbb{R}$ with $u^2 + v^2 = 2$, and $D = \kappa \cdot \operatorname{diag}(1, x, y, z)$ for some $\kappa \in \mathbb{R}_{>0}$, and $x, y, z \in \mathbb{R}$ such that 0 < |x|, |y|, |z| < 1. Moreover,

- 1. If $|x| \neq |y|$, then u = v = 1, and z = -xy.
- 2. If x = y, then $z = -x^2$.
- 3. If x = -y, then either $z = x^2$, or $z = -x^2$ and $|u| \neq |v|$.

Remark. Given $u^2 + v^2 = 2$, we have uv = 1 iff u = v. In this case, K is a tensor product. In particular, when u = v = 1, $K = H^{\otimes 2}$, where H is the 2 by 2 Hadamard matrix.

Proof. Since $M^2 \in \text{Sym}_4^{\text{pd}}(\mathbb{R}_{>0})$, such that $\rho_{\text{tensor}}(M^2) = 0$, we may let $M^2 = A \otimes B$ for some $A, B \in \text{Sym}_2^{\text{pd}}(\mathbb{R}_{>0})$. From Theorem 104, we know that $\text{Pl-GH}(M^2) \leq \text{Pl-GH}(M)$ is #P-hard unless $A_{11} = A_{22}, B_{11} = B_{22}$. So, let us now assume that $A_{11} = A_{22}, B_{11} = B_{22}$. We note that in this case, $A = HD_1H^T$, and $B = HD_2H^T$, where

$$H = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1\\ 1 & -1 \end{pmatrix}, \quad D_1 = \begin{pmatrix} A_{11} + A_{12} & 0\\ 0 & A_{11} - A_{12} \end{pmatrix}, \quad \text{and} \quad D_2 = \begin{pmatrix} B_{11} + B_{12} & 0\\ 0 & B_{11} - B_{12} \end{pmatrix}.$$

So, we see that $M^2 = (H \otimes H)(D_1 \otimes D_2)(H \otimes H)^T$. We may now let $\lambda_1 = A_{11} + A_{12} > A_{11} - A_{12} = \lambda_2$ be the eigenvalues of A, and $\mu_1 = B_{11} + B_{12} > B_{11} - B_{12} = \mu_2$ be the eigenvalues of B. So, the eigenvalues of M^2 are $\lambda_1 \mu_1$, $\lambda_1 \mu_2$, $\lambda_2 \mu_1$, and $\lambda_2 \mu_2$.

We will now consider the eigenvalues of M. We know that they must be such that their squares are the eigenvalues of M^2 . Moreover, since $M \in \text{Sym}_4(\mathbb{R}_{>0})$, we know from the Perron-Frobenius Theorem, that M has a unique positive eigenvalue with the largest absolute value. Since $\lambda_1\mu_1$ is the unique eigenvalue of M^2 with the largest absolute value, this implies that $\kappa = \sqrt{\lambda_1\mu_1}$ is the unique positive eigenvalue of M with the largest absolute value. We may now define $x, y, z \in \mathbb{R}_{\neq 0}$ to be such that $\kappa x, \kappa y$, and κz are the eigenvalues of M that satisfy the equations: $(\kappa x)^2 = \lambda_1\mu_2, (\kappa y)^2 = \lambda_2\mu_1$, and $(\kappa z)^2 = \lambda_2\mu_2$. Note that since κ is positive and is the unique eigenvalue with the largest absolute eigenvalue, it follows that |x|, |y|, |z| < 1. Moreover, since $(\lambda_1\mu_1)(\lambda_2\mu_2) = (\lambda_1\mu_2)(\lambda_2\mu_1)$, we see that $z^2 = (xy)^2$. So, either z = xy, or z = -xy. In either case, since |x|, |y| < 1, it follows that $\kappa > \kappa |x|, \kappa |y| > \kappa |z|$. So, 1 > |x|, |y| > |z|.

We will first consider the case where $|x| \neq |y|$. In that case, we see that the eigenvalues of M^2 are all distinct. Then their corresponding eigenspaces are all one dimensional, thus the corresponding unit eigenvectors are unique up to a \pm sign. This means that $H \otimes H$ is the unique orthogonal matrix (up to a ± 1 multiplier per each column) such that $M^2 = (H \otimes H)(D_1 \otimes D_2)(H \otimes H)^T$. Since $M \cdot M^2 = M^3 = M^2 \cdot M$, we know that M and M^2 must be simultaneously diagonalizable. But this then implies that M must also be diagonalized by $H \otimes H$, which precisely takes the form of the matrix K in the statement of this lemma (with u = v = 1). Moreover, if z = xy, that implies that $M = (H \otimes H)(D'_1 \otimes D'_2)(H \otimes H)^T = (HD'_1H^T) \otimes (HD'_2H^T)$, where $D'_1 = \sqrt{\kappa} \cdot \text{diag}(1, y)$, and $D'_2 = \sqrt{\kappa} \cdot \text{diag}(1, x)$. But this implies that $\rho_{\text{tensor}}(M) = 0$, which contradicts our assumption about M. So, our assumption that z = xy must be false. This means that z = -xy, in which case, $M = KDK^T$ with u = v = 1, and $D = \kappa \cdot \text{diag}(1, x, y, -xy)$. So, the statement of the lemma is proved.

Now, let us consider the case where |x| = |y|. We will characterize the set of all matrices H'such that $M^2 = (H')(D_1 \otimes D_2)(H')^{\mathsf{T}}$. We note that the eigenvalues $(\lambda_1\mu_1)$ and $(\lambda_2\mu_2)$ of M^2 have multiplicity 1. So, their corresponding eigenspaces are one dimensional. So, $1/2(1, 1, 1, 1)^{\mathsf{T}}$ and $1/2(1, -1, -1, 1)^{\mathsf{T}}$ must be column 1 and column 4 of H' respectively (upto a factor of ± 1). But since |x| = |y|, it follows that the duplicate eigenvalue $(\lambda_1\mu_2) = \kappa^2 x^2 = \kappa^2 y^2 = (\lambda_2\mu_1)$ has multiplicity 2. So, the corresponding eigenspace has dimension 2. So, all unit vectors that are orthogonal to both $1/2(1, 1, 1, 1)^{\mathsf{T}}$, and $1/2(1, -1, -1, 1)^{\mathsf{T}}$ are unit eigenvectors of M^2 with the eigenvalue $(\lambda_1\mu_2) = (\lambda_2\mu_1)$. Therefore, any unit vectors $\mathbf{v}_1, \mathbf{v}_2$ that are orthogonal to each other, and also to both $(1, 1, 1, 1)^{\mathsf{T}}$ and $(1, -1, -1, 1)^{\mathsf{T}}$ could be columns of H'. We may now assume that $\mathbf{v}_1 = (v_{11}, v_{12}, v_{13}, v_{14})^{\mathsf{T}}$ and $\mathbf{v}_2 = (v_{21}, v_{22}, v_{23}, v_{24})^{\mathsf{T}}$. Since \mathbf{v}_1 and \mathbf{v}_2 are both orthogonal to $(1, 1, 1, 1)^{\mathsf{T}}$ and $(1, -1, -1, 1)^{\mathsf{T}}$, it must be the case that

$$v_{11} + v_{12} + v_{13} + v_{14} = 0, \quad v_{11} - v_{12} - v_{13} + v_{14} = 0 \implies (v_{11} + v_{14}) = (v_{12} + v_{13}) = 0,$$

$$v_{21} + v_{22} + v_{23} + v_{24} = 0, \quad v_{21} - v_{22} - v_{23} + v_{24} = 0 \implies (v_{21} + v_{24}) = (v_{22} + v_{23}) = 0.$$

We may let $u = v_{11}$, and $v = v_{13}$. This implies that $\mathbf{v}_1 = (u, -v, v, -u)^T$ is one of the eigenvectors of M^2 with the eigenvalue $(\lambda_1 \mu_2)$. Since eigenvectors are preserved under scalar multiplication, we

may assume that $u^2 + v^2 = 2$, and then $\mathbf{v}_1 = \frac{1}{2}(u, -v, v, -u)^T$ is a unit vector. Since \mathbf{v}_1 and \mathbf{v}_2 are also orthogonal to each other, this implies that $u(v_{21} - v_{24}) + v(v_{22} - v_{23}) = 0$. Since $v_{24} = -v_{21}$, and $v_{23} = -v_{22}$, this means that $u(v_{21}) - v(v_{22}) = 0$. It follows that we have $\mathbf{v}_2 = \frac{1}{2}(v, u, -u, -v)^T$, as the other unit eigenvector of $(\lambda_2 \mu_1) = (\lambda_1 \mu_2)$.

Summing up, if |x| = |y|, the eigenvectors of M^2 are $\frac{1}{2}(1, 1, 1, 1)^T$, $\frac{1}{2}(u, -v, v, u)^T$, $\frac{1}{2}(v, u, -u, -v)^T$, and $\frac{1}{2}(1, -1, -1, 1)^T$, for all $u, v \in \mathbb{R}$ such that $u^2 + v^2 = 2$. In other words, if $M^2 = (H')(D_1 \otimes D_2)(H')^T$, we see that H' must be of the form of the matrix K in the statement of the lemma. Since $M \cdot M^2 = M^3 = M^2 \cdot M$, we know that M and M^2 must be simultaneously diagonalizable. So, this implies that when |x| = |y|, $M = KDK^T$ for some orthogonal matrix K of the stated form in the statement of this lemma, and $D = \kappa \cdot \text{diag}(1, x, y, z)$, where $z = \pm xy$.

If x = y, then this means that the multiplicity of the eigenvalues $\kappa x = \kappa y$ is once again 2. This implies that any nonzero linear combination of \mathbf{v}_1 and \mathbf{v}_2 is also an eigenvector of M. In particular, that implies that $\left(\frac{u+v}{2}\right)\mathbf{v}_1 + \left(\frac{v-u}{2}\right)\mathbf{v}_2 = \frac{1}{2}(1,-1,1,-1)^T$, and $\left(\frac{u-v}{2}\right)\mathbf{v}_1 + \left(\frac{u+v}{2}\right)\mathbf{v}_2 = \frac{1}{2}(1,1,-1,-1)^T$ are eigenvectors of M as well, with the eigenvalue $\kappa x = \kappa y$. This means that $M = (H \otimes H)D(H \otimes H)^T$. If z = xy, that would again imply that $\rho_{\text{tensor}}(M) = 0$. So, we conclude that $z = -xy = -x^2$, in which case, M is of the form in the statement of the lemma.

Finally, we consider the case where x = -y. We have $z = \pm x^2$. If $z = x^2$, then M is already of the form in the statement of the lemma and we are done. We will now assume that $z = -x^2$. For a contradiction assume |u| = |v|. As $u^2 + v^2 = 2$, we have |u| = |v| = 1. We may assume that u = 1, since unit eigenvectors are preserved when multiplied by ± 1 . If v = 1 as well, we find that $K = H \otimes H$. So, $M = (HD'_1H^T) \otimes (HD'_2H^T)$, where $D'_1 = \sqrt{\kappa} \cdot \text{diag}(1, -x)$, and $D'_2 = \sqrt{\kappa} \cdot \text{diag}(1, x)$, which implies once again that $\rho_{\text{tensor}}(M) = 0$, contradicting our assumption about M. Finally, if v = -1, we will consider the matrix $M' \in \text{Sym}_4^F(\mathbb{R}_{>0})$ such that $(M')_{ij} = M_{\sigma(i)\sigma(j)}$, where $\sigma \in S_4$ is such that $\sigma(1) = 1, \sigma(2) = 3, \sigma(3) = 2$, and $\sigma(4) = 4$. We note that M and M' are isomorphic to each other. Since $M = KDK^T$, we see that $M' = (K')D'(K')^T$, where K' is obtained by applying σ to the rows of K, and multiplying the 3rd column by -1, and D' is obtained by multiplying both the 3rd row and column of D by -1. But then, we see that $K' = H \otimes H$, and $D' = D = \kappa \cdot \text{diag}(1, x, -x, -x^2)$. Hence, $M' = (HD'_1H^T) \otimes (HD'_2H^T)$, where $D'_1 = \sqrt{\kappa} \cdot \text{diag}(1, -x)$, and $D'_2 = \sqrt{\kappa} \cdot \text{diag}(1, x)$. This implies that $\rho_{\text{tensor}}(M') = 0$, which contradicts our assumption that $\rho_{\text{tensor}}(M) \neq 0$. So, we conclude that when x = -y and $z = -x^2$ we have $|u| \neq |v|$.

In all cases, we have proved that either Pl-GH(M) is #P-hard, or M, the diagonalizing K, and the diagonal matrix D, take the form in the statement of the lemma.

Lemma 107. Let $M \in Sym_4^{pd}(\mathbb{R}_{>0})$, such that $M = A \otimes B$ for some $A, B \in Sym_2^{pd}(\mathbb{R}_{>0})$. For any $\sigma \in S_4$, if $\{\sigma(1), \sigma(4)\} = \{1, 2\}$, or $\{3, 4\}$, or $\{1, 3\}$, or $\{2, 4\}$, then $\varrho_{tensor}(M^{\sigma}) \neq 0$, where $(M^{\sigma})_{ij} = M_{\sigma(i)\sigma(j)}$ for all $i, j \in [4]$.

Proof. Since $M = A \otimes B$, we have $\rho_{\texttt{tensor}}(M) = 0$, where

$$M = \begin{pmatrix} A_{11}B_{11} & A_{11}B_{12} & A_{12}B_{11} & A_{12}B_{12} \\ A_{11}B_{12} & A_{11}B_{22} & A_{12}B_{12} & A_{12}B_{22} \\ A_{12}B_{11} & A_{12}B_{12} & A_{22}B_{11} & A_{22}B_{12} \\ A_{12}B_{12} & A_{12}B_{22} & A_{22}B_{12} & A_{22}B_{22} \end{pmatrix}$$

We note from the definition of ρ_{tensor} in Lemma 65 that $\rho_{\text{tensor}}(N) = 0$ implies that $N_{14} = N_{23}$, and $N_{i1}N_{i4} = N_{i2}N_{i3}$ for all $i \in [4]$.

We will first consider $\sigma \in S_4$ such that $\{\sigma(1), \sigma(4)\} = \{1, 2\}$, or $\{3, 4\}$. So, $\rho_{\text{tensor}}(M^{\sigma}) = 0$ implies that $M_{\sigma(1)\sigma(4)} = M_{\sigma(2)\sigma(3)}$. This implies that $M_{12} = (A_{11}B_{12}) = (A_{22}B_{12}) = M_{34}$, which is only possible if $A_{11} = A_{22}$, since $B_{12} > 0$. We also note that $\rho_{\text{tensor}}(M^{\sigma}) = 0$ implies that $M_{1\sigma(1)}M_{1\sigma(4)} = M_{1\sigma(2)}M_{1\sigma(3)}$. This implies that $M_{11}M_{12} = M_{13}M_{14}$. In other words, it must be the case that $(A_{11}B_{11})(A_{11}B_{12}) = (A_{12}B_{11})(A_{12}B_{12})$. Since $A_{11}, A_{12}, B_{11}, B_{12} > 0$, this is only possible if $A_{11} = A_{12}$. But if $A_{11} = A_{12} = A_{22}$, that means A is a rank 1 matrix, which contradicts the fact that $A \in \text{Sym}_2^{\text{pd}}(\mathbb{R}_{>0})$. So, we conclude that if $\{\sigma(1), \sigma(4)\} = \{1, 2\}$, or if $\{\sigma(1), \sigma(4)\} = \{3, 4\}$ then $\rho_{\text{tensor}}(M^{\sigma}) \neq 0$.

For $\sigma \in S_4$ such that $\{\sigma(1), \sigma(4)\} = \{1, 3\}$, or $\{2, 4\}$, let $\tau \in S_4$ be the transposition (23). Then $M^{\tau} = (A \otimes B)^{\tau} = B \otimes A$. Now $M_{\sigma(1)\sigma(4)} = M_{\sigma(2)\sigma(3)}$ becomes $M_{13} = M_{24}$, i.e., $(M^{\tau})_{12} = (M^{\tau})_{34}$. Similarly, $M_{1\sigma(1)}M_{1\sigma(4)} = M_{1\sigma(2)}M_{1\sigma(3)}$ is $M_{11}M_{13} = M_{12}M_{14}$, i.e., $(M^{\tau})_{11}(M^{\tau})_{12} = (M^{\tau})_{13}(M^{\tau})_{14}$. Since M^{τ} is also a tensor product, only switching the roles of $A, B \in \text{Sym}_2^{\text{pd}}(\mathbb{R}_{>0})$ in M, $\rho_{\text{tensor}}(M^{\sigma}) = 0$ leads to the same contradiction as in the previous paragraph (with M^{τ} in place of M). So, $\rho_{\text{tensor}}(M^{\sigma}) \neq 0$ when $\{\sigma(1), \sigma(4)\} = \{1, 3\}$, or $\{2, 4\}$.

Lemma 108. Let $M \in Sym_4^{\mathsf{F}}(\mathbb{R}_{>0})$ be a matrix of the form $M = KDK^{\mathsf{T}}$, where the matrix K and D are as in Lemma 106, such that $|x| \neq |y|$, u = v = 1, and z = -xy. Then, Pl-GH(M) is #P-hard.

Proof. We will first define the Sym₄(\mathbb{R})-polynomial ξ such that $\xi : N \mapsto (N_{11}N_{14} - N_{12}N_{13})$. We note from our choice of M, that $M = KDK^{\mathsf{T}}$, where

and $D = \kappa \cdot \operatorname{diag}(1, x, y, -xy)$ for some $\kappa \in \mathbb{R}_{>0}$, and $x, y \in \mathbb{R}$ such that 0 < |x|, |y| < 1, and $|x| \neq |y|$. We note that for any odd $n \ge 1$, $M^n = KD^nK^{\mathsf{T}}$, where $D^n = (\kappa)^n \cdot \operatorname{diag}(1, x^n, y^n, -(xy)^n)$. We note that

$$(M^{n})_{11} = (M^{n})_{22} = (M^{n})_{33} = (M^{n})_{44} = \kappa^{n} \cdot \left(\frac{1+x^{n}+y^{n}-(xy)^{n}}{4}\right),$$
$$(M^{n})_{12} = (M^{n})_{34} = \kappa^{n} \cdot \left(\frac{1-x^{n}+y^{n}+(xy)^{n}}{4}\right),$$
$$(M^{n})_{13} = (M^{n})_{24} = \kappa^{n} \cdot \left(\frac{1+x^{n}-y^{n}+(xy)^{n}}{4}\right),$$
$$(M^{n})_{14} = (M^{n})_{23} = \kappa^{n} \cdot \left(\frac{1-x^{n}-y^{n}-(xy)^{n}}{4}\right).$$

We will now let $N_n = (T_2((4/\kappa^n)M^n))^2$ for any odd $n \ge 1$. We note that $(N_n)_{11} = (1+x^n+y^n-(xy)^n)^4 + (1-x^n+y^n+(xy)^n)^4 + (1+x^n-y^n+(xy)^n)^4 + (1-x^n-y^n-(xy)^n)^4,$

$$(N_n)_{12} = 2((1+x^n+y^n-(xy)^n)(1-x^n+y^n+(xy)^n))^2 + 2((1+x^n-y^n+(xy)^n)(1-x^n-y^n-(xy)^n))^2,$$

$$(N_n)_{13} = 2((1+x^n+y^n-(xy)^n)(1+x^n-y^n+(xy)^n))^2 + 2((1-x^n+y^n+(xy)^n)(1-x^n-y^n-(xy)^n))^2,$$

$$(N_n)_{14} = 2((1+x^n+y^n-(xy)^n)(1-x^n-y^n-(xy)^n))^2 + 2((1-x^n+y^n+(xy)^n)(1+x^n-y^n+(xy)^n))^2.$$

After simplifying these equations, and computing $\xi(N_n)$, we find that

After simplifying these equations, and computing $\xi(N_n)$, we find that

$$\xi(N_n) = -1024x^{2n}y^{2n} \left(1 - 2y^{2n} - 2x^{2n} + y^{4n} + x^{4n} + 4x^{2n}y^{2n} - 2x^{2n}y^{4n} - 2x^{4n}y^{2n} + x^{4n}y^{4n}\right).$$

Since |x|, |y| < 1, we know that for large enough n,

$$1 > |2y^{2n}| + |2x^{2n}| + |y^{4n}| + |x^{4n}| + |4x^{2n}y^{2n}| + |x^{2n}y^{4n}| + |2x^{4n}y^{2n}| + |x^{4n}y^{4n}|$$

Moreover, we know that $|x|, |y| \neq 0$. So, we see that there exists some odd n^* , such that $\xi(N_{n^*}) \neq 0$. Since $\xi(N_{n^*}) = (4/\kappa^{n^*})^4 \xi((T_2(M^{n^*}))^2)$, this implies that $\xi((T_2(M^{n^*}))^2) \neq 0$.

We will now let $M' = M^{n^*}$. Since $M \in \text{Sym}_4^F(\mathbb{R}_{>0})$, it follows that $M' \in \text{Sym}_4^F(\mathbb{R}_{>0})$ as well. From the construction of ρ_{tensor} in Lemma 65, we see that $\xi(N) \neq 0$ implies that $\rho_{\text{tensor}}(N^{\sigma}) \neq 0$ for all $\sigma \in S_4$ such that $\{\sigma(1), \sigma(4)\} = \{1, 4\}$, or $\{2, 3\}$. We will now define

$$T_1 = \left\{ \sigma \in S_4 : \{ \sigma(1), \sigma(4) \} \in \{ \{1, 4\}, \{2, 3\} \} \right\},\$$

and define the Sym₄(\mathbb{R})-polynomial ζ_1 such that $\zeta_1 : N \mapsto \prod_{\sigma \in T_1} \rho_{\texttt{tensor}}((N^2)^{\sigma})$. By construction, we see that $\zeta_1(T_2(M')) \neq 0$. We also note that $(M')^2 = M^{2n^*} = (M^2)^{n^*}$. By our choice of M, we know that $\rho_{\texttt{tensor}}(M^2) = 0$. So, there exist $A, B \in \text{Sym}_2^{pd}(\mathbb{R}_{>0})$ such that $M^2 = A \otimes B$. But this means that $M^{2n^*} = A^{n^*} \otimes B^{n^*}$, which means that $\rho_{\texttt{tensor}}((M')^2) = 0$ as well. So, from Lemma 107, we note that for any $\sigma \in S_4$, such that $\{\sigma(1), \sigma(4)\} = \{1, 2\}$, or $\{3, 4\}$, or $\{1, 3\}$, or $\{2, 4\}, \rho_{\texttt{tensor}}(((M')^2)^{\sigma}) \neq 0$. We will now let

$$T_2 = S_4 \setminus T_1 = \left\{ \sigma \in S_4 : \{\sigma(1), \sigma(4)\} \in \{\{1, 2\}, \{3, 4\}, \{1, 3\}, \{2, 4\}\} \right\},\$$

and define the Sym₄(\mathbb{R})-polynomial ζ_2 such that $\zeta_2 : N \mapsto \prod_{\sigma \in T_2} \rho_{\texttt{tensor}}((N^2)^{\sigma})$. We have just seen that $\zeta_2(M') \neq 0$.

Since $M' \in \operatorname{Sym}_4^F(\mathbb{R}_{>0})$, Lemma 28 lets us find $M'' \in \mathfrak{R}(M', \{\zeta_1, \zeta_2\}) \cap \operatorname{Sym}_4^F(\mathbb{R}_{>0})$. Since $\zeta_1(M'') \neq 0$, and $\zeta_2(M'') \neq 0$, it follows that $\rho_{\operatorname{tensor}}((M'')^2) \neq 0$. Since $(M'')^2 \in \operatorname{Sym}_4^{\operatorname{pd}}(\mathbb{R}_{>0})$, Theorem 98 implies that $\operatorname{Pl-GH}((M'')^2) \leq \operatorname{Pl-GH}(M'') \leq \operatorname{Pl-GH}(M')$ is $\#\operatorname{P-hard}$. Since $M' = M^{n^*}$ for some integer $n^* \geq 1$, we also see that $\operatorname{Pl-GH}(M') \leq \operatorname{Pl-GH}(M)$. This proves that $\operatorname{Pl-GH}(M)$ is $\#\operatorname{P-hard}$.

Remark. We should note that in Lemma 108, we do not require that $\rho_{tensor}(M) \neq 0$. We only require that M be of the form $M = KDK^{T}$, where K and D are as in Lemma 106 (with the stipulation on x, y, z, u, v in the statement of Lemma 108). In this case, it trivially follows that $\rho_{tensor}(M^2) = 0$, which is the only property of M that is used in the proof. This will also be true for Lemma 109, Lemma 110, and Lemma 111 below.

Lemma 109. Let $M \in Sym_4^{\mathsf{F}}(\mathbb{R}_{>0})$ be a matrix of the form $M = KDK^{\mathsf{T}}$, where the matrix K and D are as in Lemma 106, such that x = y, and $z = -x^2$. Then, Pl-GH(M) is #P-hard.

Proof. We will define ξ such that $\xi : N \mapsto (N_{11}N_{14} - N_{12}N_{13})$, as in Lemma 108. We note from Lemma 106, that $M = KDK^{\mathsf{T}}$, where

$$K = \frac{1}{2} \begin{pmatrix} 1 & u & v & 1 \\ 1 & -v & u & -1 \\ 1 & v & -u & -1 \\ 1 & -u & -v & 1 \end{pmatrix}$$

for some $u, v \in \mathbb{R}$ with $u^2 + v^2 = 2$, and $D = \kappa \cdot \operatorname{diag}(1, x, x, -x^2)$ for some $\kappa \in \mathbb{R}_{>0}$, and $x \in \mathbb{R}$ such that |x| < 1. For odd $n \ge 1$, we note that $M^n = KD^nK^T$, where $D^n = (\kappa)^n \cdot (1, x^n, x^n, -x^{2n})$. So, we note that (since $u^2 + v^2 = 2$),

$$\begin{split} (M^{n})_{11} &= (M^{n})_{22} = (M^{n})_{33} = (M^{n})_{44} = \kappa^{n} \cdot \left(\frac{1 + (u^{2} + v^{2})x^{n} - x^{2n}}{4}\right) = \kappa^{n} \cdot \left(\frac{1 + 2x^{n} - x^{2n}}{4}\right), \\ (M^{n})_{12} &= (M^{n})_{34} = (M^{n})_{13} = (M^{n})_{24} = \kappa^{n} \cdot \left(\frac{1 + (uv)(-x + x) + x^{2n}}{4}\right) = \kappa^{n} \cdot \left(\frac{1 + x^{2n}}{4}\right), \\ (M^{n})_{14} &= (M^{n})_{23} = \kappa^{n} \cdot \left(\frac{1 - (u^{2} + v^{2})x - x^{2n}}{4}\right) = \kappa^{n} \cdot \left(\frac{1 - 2x^{n} - x^{2n}}{4}\right). \end{split}$$

We will let $N_n = (T_2((4/\kappa^n)M^n))^2$ for any odd $n \ge 1$. We note that

$$(N_n)_{11} = (1 + 2x^n - x^{2n})^4 + 2(1 + x^{2n})^4 + (1 - 2x^n - x^{2n})^4,$$

$$(N_n)_{12} = (N_n)_{13} = 2((1 + 2x^n - x^{2n})(1 + x^{2n}))^2 + 2((1 - 2x^n - x^{2n})(1 + x^{2n}))^2,$$

$$(N_n)_{14} = 2((1 + 2x^n - x^{2n})(1 - 2x^n - x^{2n}))^2 + 2(1 + x^{2n})^4.$$

After simplifying and computing $\xi(N_n)$, we find that

$$\xi(N_n) = (-1024)x^{4n} (1 - x^{2n})^4.$$

Since 0 < |x| < 1, we see that there exists some odd $n^* \ge 1$ such that $\xi(N_{n^*}) \neq 0$. Since $\xi(N_{n^*}) = (4/\kappa^{n^*})^4 \xi((T_2(M^{n^*}))^2)$, this implies that $\xi((T_2(M^{n^*}))^2) \neq 0$.

We will now let $M' = M^{n^*}$. Now, by the same argument as in the proof of Lemma 108, we can now find some $M'' \in \operatorname{Sym}_4^{\mathsf{F}}(\mathbb{R}_{>0})$, such that $\operatorname{Pl-GH}(M'') \leq \operatorname{Pl-GH}(M')$, and $\rho_{\operatorname{tensor}}((M'')^2) \neq 0$. So, Theorem 98 once again lets us prove that $\operatorname{Pl-GH}((M'')^2) \leq \operatorname{Pl-GH}(M') \leq \operatorname{Pl-GH}(M)$ is #P-hard. \Box

Lemma 110. Let $M \in Sym_4^{\mathsf{F}}(\mathbb{R}_{>0})$ be a matrix of the form $M = KDK^{\mathsf{T}}$, where the matrix K and D are as in Lemma 106, such that x = -y, $z = -x^2$, and $|u| \neq |v|$. Then, Pl-GH(M) is #P-hard.

Proof. We will once again define ξ such that $\xi : N \mapsto (N_{11}N_{14} - N_{12}N_{13})$. We note from Lemma 106, that $M = KDK^{\mathsf{T}}$, where

$$K = \frac{1}{2} \begin{pmatrix} 1 & u & v & 1 \\ 1 & -v & u & -1 \\ 1 & v & -u & -1 \\ 1 & -u & -v & 1 \end{pmatrix}$$

for some $u, v \in \mathbb{R}$ with $u^2 + v^2 = 2$, $|u| \neq |v|$, and $D = \kappa \cdot \operatorname{diag}(1, x, -x, -x^2)$ for some $\kappa \in \mathbb{R}_{>0}$, and $x, y, z \in \mathbb{R}$ such that |x|, |y|, |z| < 1.

We note that for odd $n \ge 1$, $M^n = KD^nK^T$, where $D^n = (\kappa)^n \cdot \text{diag}(1, x^n - x^n, -x^{2n})$. So, we note that

$$(M^{n})_{11} = (M^{n})_{22} = (M^{n})_{33} = (M^{n})_{44} = \kappa^{n} \cdot \left(\frac{1 + (u^{2} - v^{2})x^{n} - x^{2n}}{4}\right),$$
$$(M^{n})_{12} = (M^{n})_{34} = \kappa^{n} \cdot \left(\frac{1 - 2uvx^{n} + x^{2n}}{4}\right),$$
$$(M^{n})_{13} = (M^{n})_{24} = \kappa^{n} \cdot \left(\frac{1 + 2uvx^{n} + x^{2n}}{4}\right),$$
$$(M^{n})_{14} = (M^{n})_{23} = \kappa^{n} \cdot \left(\frac{1 + (v^{2} - u^{2})x^{n} - x^{2n}}{4}\right).$$

We will now let $N_n = (T_2((4/\kappa^n)M^n))^2$ for any odd $n \ge 1$. We note that

$$\begin{split} (N_n)_{11} &= (1 + (u^2 - v^2)x^n - x^{2n})^4 + (1 - 2uvx^n + x^{2n})^4 + (1 + 2uvx^n + x^{2n})^4 + (1 + (v^2 - u^2)x^n - x^{2n})^4, \\ (N_n)_{12} &= 2((1 + (u^2 - v^2)x^n - x^{2n})(1 - 2uvx^n + x^{2n}))^2 + 2((1 + (v^2 - u^2)x^n - x^{2n})(1 + 2uvx^n + x^{2n}))^2, \\ (N_n)_{13} &= 2((1 + (u^2 - v^2)x^n - x^{2n})(1 + 2uvx^n + x^{2n}))^2 + 2((1 + (v^2 - u^2)x^n - x^{2n})(1 - 2uvx^n + x^{2n}))^2, \\ (N_n)_{14} &= 2((1 + (u^2 - v^2)x^n - x^{2n})(1 + (v^2 - u^2)x^n - x^{2n}))^2 + 2((1 - 2uvx^n + x^{2n})(1 + 2uvx^n + x^{2n}))^2. \\ \text{If we compute } \xi(N_n), \text{ it can be verified that} \end{split}$$

$$\xi(N_n) = f_4(u, v)x^{4n} + f_6(u, v)x^{6n} + f_8(u, v)x^{8n} + f_{10}(u, v)x^{10n} + f_{12}(u, v)x^{12n},$$

where

$$f_4(u,v) = f_{12}(u,v) = -48u^8 - 48v^8 + 576u^6v^2 + 576u^2v^6 - 128u^4 - 128v^4 + 768u^2v^2 - 1824u^4v^4 + 256,$$

$$f_{6}(u,v) = f_{10}(u,v) = 16u^{12} - 160u^{10}v^{2} + 240u^{8}v^{4} + 832u^{6}v^{6} + 240u^{4}v^{8} - 160u^{2}v^{10} + 16v^{12} + 128u^{8} - 512u^{6}v^{2} - 1280u^{4}v^{4} - 512u^{2}v^{6} + 128v^{8} + 256u^{4} + 512u^{2}v^{2} + 256v^{4},$$

$$\begin{split} f_8(u,v) &= 4u^{16} - 32u^{14}v^2 - 16u^{12}v^4 + 288u^{10}v^6 + 536u^8v^8 + 288u^6v^{10} - 16u^4v^{12} - 32u^2v^{14} + 4v^{16}v^6 \\ &\quad - 32u^{12} + 64u^{10}v^2 + 544u^8v^4 + 896u^6v^6 + 544u^4v^8 + 64u^2v^{10} - 32v^{12} - 288u^8 \\ &\quad + 384u^6v^2 - 4800u^4v^4 + 384u^2v^6 - 288v^8 - 256u^4 + 1536u^2v^2 - 256v^4 + 512. \end{split}$$

While these equations appear a bit intimidating, we will focus on $f_4(u, v)$. We note that $u^2 + v^2 = 2$. So, $f_4(u, v) = 0$ if and only if

$$-16\left(3u^8 + 3(2-u^2)^4 - 36u^6(2-u^2) - 36u^2(2-u^2)^3 + 8u^4 + 8(2-u^2)^2 - 48u^2(2-u^2) + 114u^2(2-u^2)^2 - 16\right) = 0.$$
After simplifying, we find that $f_4(u, v) = 0$ if and only if

$$-1024(3u^8 - 12u^6 + 16u^4 - 8u^2 + 1) = 0.$$

This is a degree 4 polynomial in u^2 , and it can be verified that

$$3u^8 - 12u^6 + 16u^4 - 8u^2 + 1 = 3(u^2 - 1)^2 \left(u^2 - 1 - \frac{\sqrt{6}}{3}\right) \left(u^2 - 1 + \frac{\sqrt{6}}{3}\right).$$

Since we know that $|u| \neq |v|$, we can see that it is not possible that $u^2 = 1$. So, $f_4(u, v) = 0$ if and only if $u^2 = (1 - \sqrt{6}/3)$ and $v^2 = (1 + \sqrt{6}/3)$, or if $u^2 = (1 + \sqrt{6}/3)$ and $v^2 = (1 - \sqrt{6}/3)$. But in either case, for these values of u^2 and v^2 , it is seen that $f_6(u, v) = (16384/9) \neq 0$. Summing up, we find that if $f_4(u, v) = 0$, then $f_6(u, v) \neq 0$. If $f_4(u, v) \neq 0$, then we note that since |x| < 1, for large enough $n \ge 1$, $|f_4(u, v)| > |f_6(u, v)|x^{2n} + |f_8(u, v)|x^{4n} + |f_{10}(u, v)|x^{6n} + |f_{12}(u, v)|x^{8n}$. So, there exists some odd $n^* \ge 1$ such that $\xi(N_{n^*}) \neq 0$. On the other hand, if $f_4(u, v) = f_{12}(u, v) = 0$, we note that for large enough $n \ge 1$, $|f_6(u, v)| > |f_8(u, v)|x^{2n} + |f_{10}(u, v)|x^{4n}$. So, once again there exists some odd $n^* \ge 1$ such that $\xi(N_{n^*}) \neq 0$. Since $\xi(N_{n^*}) = (4/\kappa^{n^*})^4 \xi((T_2(M^{n^*}))^2)$, this implies that $\xi((T_2(M^{n^*}))^2) \neq 0$.

We will now let $M' = M^{n^*}$. Now, by the same argument as in the proof of Lemma 108, we can now find some $M'' \in \text{Sym}_4^F(\mathbb{R}_{>0})$, such that $\text{Pl-GH}(M'') \leq \text{Pl-GH}(M')$, and $\rho_{\text{tensor}}((M'')^2) \neq 0$. So, Theorem 98 once again lets us prove that $\text{Pl-GH}((M'')^2) \leq \text{Pl-GH}(M') \leq \text{Pl-GH}(M)$ is #P-hard. \square

Lemma 111. Let $M \in Sym_4^{\mathsf{F}}(\mathbb{R}_{>0})$ be a matrix of the form $M = KDK^{\mathsf{T}}$, where the matrix K and D are as in Lemma 106, such that x = -y, and $z = x^2$. Then, $\mathsf{Pl-GH}(M)$ is #P-hard.

Proof. We note from Lemma 106, that $M = KDK^{T}$, where

$$K = \frac{1}{2} \begin{pmatrix} 1 & u & v & 1 \\ 1 & -v & u & -1 \\ 1 & v & -u & -1 \\ 1 & -u & -v & 1 \end{pmatrix}$$

for some $u, v \in \mathbb{R}$ with $u^2 + v^2 = 2$, and $D = \kappa \cdot \operatorname{diag}(1, x, -x, x^2)$ for some $\kappa \in \mathbb{R}_{>0}$, and $x \in \mathbb{R}$ such that |x| < 1.

We will first consider the case where u = 0. Since $u^2 + v^2 = 2$, this implies that $|v| = \sqrt{2}$. Since eigenvectors are equivalent up to scaling, we may assume that in fact, $v = \sqrt{2}$. In this case, we find that since $M = KDK^{T}$,

$$M = \frac{1}{2} \begin{pmatrix} (1-2x+x^2) & (1-x^2) & (1-x^2) & (1+2x+x^2) \\ (1-x^2) & (1+2x+x^2) & (1-2x+x^2) & (1-x^2) \\ (1-x^2) & (1-2x+x^2) & (1+2x+x^2) & (1-x^2) \\ (1+2x+x^2) & (1-x^2) & (1-x^2) & (1-2x+x^2) \end{pmatrix}$$
$$= \frac{1}{2} \begin{pmatrix} (1-x)^2 & (1-x)(1+x) & (1-x)(1+x) & (1+x)^2 \\ (1-x)(1+x) & (1-x)^2 & (1-x)(1+x) & (1-x)(1+x) \\ (1-x)(1+x) & (1-x)^2 & (1-x)(1+x) & (1-x)(1+x) \\ (1+x)^2 & (1-x)(1+x) & (1-x)(1+x) & (1-x)^2 \end{pmatrix}$$

Since $x \neq 0$, we see that this matrix M has the exact form as in Lemma 105 with $p = (1/\sqrt{2})(1-x)$, and $q = (1/\sqrt{2})(1+x)$. So, we already know that Pl-GH(M) is #P-hard. Similarly, when v = 0, it is seen that M has the exact form as in Lemma 105 with $p = (1/\sqrt{2})(1+x)$, and $q = (1/\sqrt{2})(1-x)$, and therefore, Pl-GH(M) is #P-hard.

We can therefore now assume that $u \neq 0$ and $|u| \neq \sqrt{2}$. We will again define ξ such that $\xi : N \mapsto (N_{11}N_{14} - N_{12}N_{13})$. We note that for odd $n \geq 1$, $M^n = KD^nK^T$, where $D^n = (\kappa)^n \cdot \text{diag}(1, x^n - x^n, x^{2n})$. So, we note that

$$(M^{n})_{11} = (M^{n})_{22} = (M^{n})_{33} = (M^{n})_{44} = \kappa^{n} \cdot \left(\frac{1 + (u^{2} - v^{2})x^{n} + x^{2n}}{4}\right),$$
$$(M^{n})_{12} = (M^{n})_{34} = \kappa^{n} \cdot \left(\frac{1 - 2uvx^{n} - x^{2n}}{4}\right),$$
$$(M^{n})_{13} = (M^{n})_{24} = \kappa^{n} \cdot \left(\frac{1 + 2uvx^{n} - x^{2n}}{4}\right),$$
$$(M^{n})_{14} = (M^{n})_{23} = \kappa^{n} \cdot \left(\frac{1 + (v^{2} - u^{2})x^{n} + x^{2n}}{4}\right).$$

We will now let $N_n = (T_2((4/\kappa^n)M^n))^2$ for any odd $n \ge 1$. We note that

 $(N_n)_{11} = (1 + (u^2 - v^2)x^n + x^{2n})^4 + (1 - 2uvx^n - x^{2n})^4 + (1 + 2uvx^n - x^{2n})^4 + (1 + (v^2 - u^2)x^n + x^{2n})^4,$ $(N_n)_{12} = 2((1 + (u^2 - v^2)x^n + x^{2n})(1 - 2uvx^n - x^{2n}))^2 + 2((1 + (v^2 - u^2)x^n + x^{2n})(1 + 2uvx^n - x^{2n}))^2,$ $(N_n)_{13} = 2((1 + (u^2 - v^2)x^n + x^{2n})(1 + 2uvx^n - x^{2n}))^2 + 2((1 + (v^2 - u^2)x^n + x^{2n})(1 - 2uvx^n - x^{2n}))^2,$ $(N_n)_{14} = 2((1 + (u^2 - v^2)x^n + x^{2n})(1 + (v^2 - u^2)x^n + x^{2n}))^2 + 2((1 - 2uvx^n - x^{2n})(1 + 2uvx^n - x^{2n}))^2.$ If we compute $\xi(N_n)$, it can be verified that

$$\xi(N_n) = f_4(u,v)x^{4n} + f_6(u,v)x^{6n} + f_8(u,v)x^{8n} + f_{10}(u,v)x^{10n} + f_{12}(u,v)x^{12n},$$

where

$$f_4(u,v) = f_{12}(u,v) = -48u^8 - 48v^8 + 576u^6v^2 + 576u^2v^6 + 128u^4 + 128v^4 - 768u^2v^2 - 1824u^4v^4 + 256,$$

$$f_{6}(u,v) = f_{10}(u,v) = 16u^{12} - 160u^{10}v^{2} + 240u^{8}v^{4} + 832u^{6}v^{6} + 240u^{4}v^{8} - 160u^{2}v^{10} + 16v^{12} - 128u^{8} + 512u^{6}v^{2} + 1280u^{4}v^{4} + 512u^{2}v^{6} - 128v^{8} + 256u^{4} + 512u^{2}v^{2} + 256v^{4},$$

$$\begin{split} f_8(u,v) &= 4u^{16} - 32u^{14}v^2 - 16u^{12}v^4 + 288u^{10}v^6 + 536u^8v^8 + 288u^6v^{10} - 16u^4v^{12} - 32u^2v^{14} + 4v^{16}v^6 + 32u^{12} - 64u^{10}v^2 - 544u^8v^4 - 896u^6v^6 - 544u^4v^8 - 64u^2v^{10} + 32v^{12} - 288u^8 + 384u^6v^2 - 4800u^4v^4 + 384u^2v^6 - 288v^8 + 256u^4 - 1536u^2v^2 + 256v^4 + 512. \end{split}$$

We will focus on $f_4(u, v)$. We note that $u^2 + v^2 = 2$. So, $f_4(u, v) = 0$ if and only if

$$-16\left(3u^8 + 3(2-u^2)^4 - 36u^6(2-u^2) - 36u^2(2-u^2)^3 - 8u^4 - 8(2-u^2)^2 + 48u^2(2-u^2) + 114u^2(2-u^2)^2 - 16\right) = 0.$$

After simplifying, we find that $f_4(u, v) = 0$ if and only if

$$-1024(3u^8 - 12u^6 + 14u^4 - 4u^2) = 0.$$

It can be verified that

$$3u^8 - 12u^6 + 14u^4 - 4u^2 = 3u^2(u^2 - 2)\left(u^2 - 1 - \frac{1}{\sqrt{3}}\right)\left(u^2 - 1 + \frac{1}{\sqrt{3}}\right).$$

Since we have already assumed that that $u \neq 0$ and $|u| \neq \sqrt{2}$, we see that $f_4(u, v) = 0$ if and only if $u^2 = (1 - 1/\sqrt{3})$ and $v^2 = (1 + 1/\sqrt{3})$, or if $u^2 = (1 + 1/\sqrt{3})$ and $v^2 = (1 - 1/\sqrt{3})$. In either case, it is seen that $f_6(u, v) = (16384/9) \neq 0$. So, we find that if $f_4(u, v) = 0$, then $f_6(u, v) \neq 0$. Following the same argument as in the proof Lemma 110 lets us find some odd $n^* \geq 1$ such that $\xi(N_{n^*}) \neq 0$. Since $\xi(N_{n^*}) = (4/\kappa^{n^*})^4 \xi((T_2(M^{n^*}))^2)$, this implies that $\xi((T_2(M^{n^*}))^2) \neq 0$.

We will now let $M' = M^{n^*}$. Now, by the same argument as in the proof of Lemma 108, we can now find some $M'' \in \text{Sym}_4^F(\mathbb{R}_{>0})$, such that $\text{Pl-GH}(M'') \leq \text{Pl-GH}(M')$, and $\rho_{\text{tensor}}((M'')^2) \neq 0$. So, Theorem 98 once again lets us prove that $\text{Pl-GH}((M'')^2) \leq \text{Pl-GH}(M') \leq \text{Pl-GH}(M)$ is #P-hard. \square

We are finally ready to prove the following theorem.

Theorem 112. Let $M \in Sym_4^{\mathsf{F}}(\mathbb{R}_{>0})$. Then $\mathsf{Pl-GH}(M)$ is #P-hard, unless M is isomorphic to $A \otimes B$ for some $A, B \in Sym_2^{\mathsf{F}}(\mathbb{R}_{>0})$ such that $A_{11} = A_{22}$, and $B_{11} = B_{22}$, in which case, $\mathsf{Pl-GH}(M)$ is polynomial time tractable.

Proof. We will let $N = M^2$. Since $M \in \text{Sym}_4^F(\mathbb{R}_{>0})$, we see that $N \in \text{Sym}_4^{pd}(\mathbb{R}_{>0})$. We know from Theorem 104 that unless N is isomorphic to $A' \otimes B'$ for some $A', B' \in \text{Sym}_2^{pd}(\mathbb{R}_{>0})$ that satisfy the conditions that $(A')_{11} = (A')_{22}$, and $(B')_{11} = (B')_{22}$, $\text{Pl-GH}(N) \leq \text{Pl-GH}(M)$ is #P-hard.

We will now consider M such that $\rho_{tensor}(M) \neq 0$, but $\rho_{tensor}(N) = 0$. We may assume that after permutation of rows and columns of M (and correspondingly N), that $N = A' \otimes B'$ for some A', and B' as above. In that case, we see that $\rho_{tensor}(M) \neq 0$, but $\rho_{tensor}(N) = 0$. So, from Lemma 106, Lemma 108, Lemma 109, Lemma 110, and Lemma 111, we see that Pl-GH(M) is #P-hard.

Finally, we consider the case where $\rho_{\text{tensor}}(M) = 0$. We may assume that after permutation of rows and columns of M (and correspondingly N), that $N = A' \otimes B'$ for some A', and B' as above. Since $(A')_{11} = (A')_{22}$, we note that $A' = HD'_1H^{\mathsf{T}}$, where

$$H = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1\\ 1 & -1 \end{pmatrix}, \text{ and } D'_1 = \begin{pmatrix} A'_{11} + A'_{12} & 0\\ 0 & A'_{11} - A'_{12} \end{pmatrix}$$

Similarly, since $(B')_{11} = (B')_{22}$, we can verify that $B' = HD'_2H^{\mathsf{T}}$, where H is the same matrix as above, and

$$D_2' = \begin{pmatrix} B_{11}' + B_{12}' & 0\\ 0 & B_{11}' - B_{12}' \end{pmatrix}$$

We may let $\lambda'_1 = A'_{11} + A'_{12}$, and $\lambda'_2 = A'_{11} - A'_{12}$. Similarly, we may let $\mu'_1 = B'_{11} + B'_{12}$, and $\mu'_2 = B'_{11} - B'_{12}$. We note that since $A'_{12}, B'_{12} > 0$, it follows that $\lambda'_1 > \lambda'_2$, and $\mu'_1 > \mu'_2$. Moreover,

since $A', B' \in \text{Sym}_2^{\text{pd}}(\mathbb{R}_{>0})$, we also see that $\lambda'_2, \mu'_2 > 0$. Now, it follows that $\lambda'_1\mu'_1 > \lambda'_1\mu'_2 > \lambda'_2\mu'_2$, and that $\lambda'_1\mu'_1 > \lambda'_2\mu'_1 > \lambda'_2\mu'_2$. Now, if we let $H' = H \otimes H$, and $D' = D'_1 \otimes D'_2$, we see that $N = A' \otimes B' = (H')D'(H')^{\mathsf{T}}$, where

We will now consider the eigenvalues of the matrix M. Since $M^2 = N$, we know that the squares of the eigenvalues of M must precisely be the eigenvalues of N. Moreover, since $M \in \text{Sym}_4(\mathbb{R}_{>0})$, from the Perron-Frobenius Theorem, we know that M has a unique positive eigenvalue with the largest absolute value. Since $(\lambda'_1\mu'_1)$ is the unique eigenvalue of N with the largest absolute value, it follows that $\kappa_1 = \sqrt{\lambda'_1\mu'_1}$ is one of the eigenvalues of M. If we now let $\lambda_1 = \sqrt{\lambda'_1}$, and $\mu_1 = \sqrt{\mu'_1}$, this means that $\kappa_1 = \lambda_1\mu_1$ is an eigenvalue of M. We also know that there exists some eigenvalue κ_2 of M such that $(\kappa_2)^2 = \lambda'_2\mu'_1$. We will let $\lambda_2 = (\kappa_2/\mu_1) = \pm \sqrt{\lambda'_2}$. Therefore, $\lambda_2\mu_1$ is an eigenvalue of M. Similarly, we know that there exists an eigenvalue κ_3 of M such that $(\kappa_3)^2 = \lambda'_1\mu'_2$. We can then let $\mu_2 = (\kappa_3/\lambda_1) = \pm \sqrt{\mu'_2}$, such that $\lambda_1\mu_2$ is an eigenvalue of M. Finally, we know that there exists some eigenvalue κ_4 of M such that $(\kappa_4)^2 = \lambda'_2\mu'_2$. Therefore, we see that $\kappa_4 = \pm \sqrt{\lambda'_2\mu'_2} = \pm \lambda_2\mu_2$. As $\rho_{\text{tensor}}(M) = 0$, M is isomorphic to a tensor product. Then its 4 eigenvalues must satisfy an equation of the form $\kappa_i\kappa_j = \kappa_k\kappa_\ell$, for some i, j, k, ℓ with $\{i, j, k, \ell\} = \{1, 2, 3, 4\}$. Since κ_1 and κ_4 have the maximum and minimum absolute value respectively among the 4 eigenvalues, we must have $\kappa_1\kappa_4 = \kappa_2\kappa_3$. Since $\kappa_1 = \lambda_1\mu_1$, $\kappa_2 = \lambda_2\mu_1$, $\kappa_3 = \lambda_1\mu_2$, we must have $\kappa_4 = \lambda_2\mu_2$.

We now note that since $M^2 = N$, $N \cdot M = M^3 = M \cdot N$. Since these matrices commute, we see that they can both be diagonalized by some orthogonal matrix K. If we let $D = \text{diag}(\lambda_1\mu_1, \lambda_1\mu_2, \lambda_2\mu_1, \lambda_2\mu_2) = D_1 \otimes D_2$, where $D_1 = \text{diag}(\lambda_1, \lambda_2)$, and $D_2 = \text{diag}(\mu_1, \mu_2)$, We see that there exists some K such that $N = KD'K^T$, and $M = KDK^T$.

If $\lambda'_1 \mu'_2 \neq \lambda'_2 \mu'_1$, then N has distinct eigenvalues, which implies that $H' = H \otimes H$ is the only matrix (upto scaling each column by -1) which can diagonalize N. This would then imply that H' can also diagonalize M. This means that we can take $K = H' = (H \otimes H)$. In other words, $M = (HD_1H^T) \otimes (HD_2H^T)$.

If $\lambda'_1\mu'_2 = \lambda'_2\mu'_1$, then N can be diagonalized by any K such that

$$K = \frac{1}{2} \begin{pmatrix} 1 & u & v & 1 \\ 1 & -v & u & -1 \\ 1 & v & -u & -1 \\ 1 & -u & -v & 1 \end{pmatrix}$$

for $u, v \in \mathbb{R}$ such that $u^2 + v^2 = 2$. Moreover, $\lambda'_1 \mu'_2 = \lambda'_2 \mu'_1$ implies that $\lambda_1 \mu_2 = \pm \lambda_2 \mu_1$. If in fact, $\lambda_1 \mu_2 = \lambda_2 \mu_1$, we can replace the two middle columns of K with any two orthogonal vectors that lie in their span. So, we see that M can also be diagonalized by $H' = H \otimes H$. So, once again, we find that $M = (HD_1H^T) \otimes (HD_2H^T)$. Finally, if $\lambda_1 \mu_2 = -\lambda_2 \mu_1$, we see that $M = KDK^T$, such that $D = (\lambda_1 \mu_1) \cdot \text{diag}(1, x, -x, -x^2)$, where $x = (\mu_2/\mu_1) = -(\lambda_2/\lambda_1)$. So, from Lemma 110 we see that unless |u| = |v| = 1, P1-GH(M) is #P-hard. Now assume |u| = |v| = 1. Since columns of K can be scaled by -1, we may assume that u = 1. If v = 1, then we see that once again, M is diagonalized by $K = H' = H \otimes H$. So, $M = (HD_1H^T) \otimes (HD_2H^T)$. If v = -1, then we claim that $M = (HD_2H^T) \otimes (HD_1H^T)$. Indeed, if we let

then $M = KDK^{\mathsf{T}} = K(D_1 \otimes D_2)K^{\mathsf{T}} = K'(D_2 \otimes D_1)K^{\mathsf{T}} = (HD_2H^{\mathsf{T}}) \otimes (HD_1H^{\mathsf{T}}).$

So, in any case, we see that $M = A \otimes B$ for some $A, B \in \text{Sym}_2^F(\mathbb{R}_{>0})$. Moreover, since $(A')_{11} = (A')_{22}$, and $(B')_{11} = (B')_{22}$, we see that $N_{11} = N_{22} = N_{33} = N_{44}$. Now, $N_{11} = N_{22}$ implies that

$$((A_{11})^2 + (A_{12})^2) \cdot ((B_{11})^2 + (B_{12})^2) = (M_{11})^2 + (M_{12})^2 + (M_{13})^2 + (M_{14})^2$$

= $(M_{12})^2 + (M_{22})^2 + (M_{23})^2 + (M_{24})^2 = ((A_{11})^2 + (A_{12})^2) \cdot ((B_{12})^2 + (B_{22})^2),$

which implies that $(B_{11})^2 = (B_{22})^2$. Since $B \in \text{Sym}_2(\mathbb{R}_{>0})$, in fact, we see that $B_{11} = B_{22}$. Similarly, $N_{11} = N_{33}$ would imply that $A_{11} = A_{22}$ as well. So, if $(A')_{11} = (A')_{22}$, and $(B')_{11} = (B')_{22}$, we see that $A_{11} = A_{22}$, and $B_{11} = B_{22}$ as well, in which case, Lemma 101 implies that P1-GH(M) is polynomial time tractable.

So, we see that Pl-GH(M) is #P-hard, unless M is isomorphic to $A \otimes B$ for some $A, B \in Sym_2^F(\mathbb{R}_{>0})$ such that $A_{11} = A_{22}$, and $B_{11} = B_{22}$, in which case, Pl-GH(M) is polynomial time tractable.

We can extend this dichotomy to all non-negative real valued full rank matrices with minimal effort.

Definition 113. Let $M \in Sym_q(\mathbb{R}_{\geq 0})$. We say that M is domain separable if it is isomorphic to some $A \oplus B = \begin{pmatrix} A & \mathbf{0} \\ \mathbf{0} & B \end{pmatrix}$ for some non-empty matrices $A \in Sym_{q_1}(\mathbb{R}_{\geq 0})$, and $B \in Sym_{q_2}(\mathbb{R}_{\geq 0})$ where $q_1 + q_2 = q$.

It is known from [CCL13] (Lemma 4.6, p. 940, the proof of which uses Lemma 4.1, p. 937, called the first pinning lemma) that GH(M) is #P-hard iff at least one of GH(A) or GH(B) is #P-hard, and that GH(M) is polynomial time tractable iff both GH(A) and GH(B) are polynomial time tractable. Furthermore, the proof of Lemma 4.1 in [CCL13] uses only planar gadgets (and in that proof we can place each identifying vertex, called w and w^* 's in the paper, to be on the outer face). Thus, this proof works for planar graphs, i.e., Pl-GH(M) is #P-hard iff at least one of Pl-GH(A) or Pl-GH(B)is #P-hard, and that Pl-GH(M) is polynomial time tractable iff both Pl-GH(A) and Pl-GH(B) are polynomial time tractable. Now, we already have a dichotomy from [GW20; CM23] for Pl-GH(M)when $M \in Sym_{q}(\mathbb{R}_{\geq 0})$ for q < 4. So, that allows us to handle domain separable matrices with ease.

Lemma 114. Let $M \in Sym_4^{\mathsf{F}}(\mathbb{R}_{\geq 0})$ such that $M = A \oplus B$ is domain separable. Then $\mathsf{Pl-GH}(M)$ is #P-hard unless $\mathsf{Pl-GH}(A)$ and $\mathsf{Pl-GH}(B)$ are both polynomial time tractable, in which case, $\mathsf{Pl-GH}(M)$ is also polynomial time tractable.

Definition 115. Let $M \in Sym_q(\mathbb{R}_{\geq 0})$. We say that M is bipartite if there exists some $A \in \mathbb{R}^{q_1 \times q_2}$ for some $q_1 + q_2 = q$, such that $M = \begin{pmatrix} 0 & A \\ A^T & 0 \end{pmatrix}$.

Lemma 116. Let $M \in Sym_4^{\mathsf{F}}(\mathbb{R}_{\geq 0})$ be bipartite. Then $\mathsf{Pl-GH}(M)$ is #P-hard unless M is isomorphic to $A \otimes B$ for some $A, B \in Sym_2^{\mathsf{F}}(\mathbb{R}_{\geq 0})$ such that $A_{11} = A_{22}$, and $B_{11} = B_{22}$, in which case, $\mathsf{Pl-GH}(M)$ is also polynomial time tractable.

Proof. We may assume that $M = \begin{pmatrix} \mathbf{0} & A \\ A^{\mathsf{T}} & \mathbf{0} \end{pmatrix}$, for some $A \in (\mathbb{R}_{\geq 0})^{q_1 \times q_2}$, where $q_1 + q_2 = 4$. If $q_1 = 1$ (or $q_2 = 1$), we can see that M has rank at most 2. So, it is not possible that $M \in \operatorname{Sym}_4^{\mathsf{F}}(\mathbb{R}_{\geq 0})$.

So, we may assume that $q_1 = q_2 = 2$. Then, M must be (upto some isomorphism) of the form

$$M = \begin{pmatrix} 0 & 0 & M_{13} & M_{14} \\ 0 & 0 & M_{23} & M_{24} \\ M_{13} & M_{23} & 0 & 0 \\ M_{14} & M_{24} & 0 & 0 \end{pmatrix}$$

But then, we see that $(T_n M)^2 = A_n \oplus A_n$, where

$$A_{n} = \begin{pmatrix} (M_{13})^{2n} + (M_{14})^{2n} & (M_{13}M_{23})^{n} + (M_{14}M_{24})^{n} \\ (M_{13}M_{23})^{n} + (M_{14}M_{24})^{n} & (M_{23})^{2n} + (M_{24})^{2n} \end{pmatrix}$$

If there exists some $n \ge 1$ such that $(A_n)_{11} \ne (A_n)_{22}$, then from Theorem 100 and Lemma 114, it follows that $\text{Pl-GH}((T_n M)^2) \le \text{Pl-GH}(M)$ is #P-hard.

If $(A_n)_{11} = (A_n)_{22}$ for all $n \ge 1$, that implies that $\{M_{13}, M_{14}\} = \{M_{23}, M_{24}\}$ as multi-sets. But if $M_{13} = M_{23}$, and $M_{14} = M_{24}$, then M can once again, not be full rank. This implies that $M_{13} = M_{24}$, and $M_{14} = M_{23}$. But in this case, we see that $M = B \otimes A$, where $B = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$. In this case, we see from Theorem 100 that Pl-GH(A) and Pl-GH(B) are both polynomial time tractable. Since $Z_M(G) = Z_A(G) \cdot Z_B(G)$ for all graphs G = (V, E), this implies that Pl-GH(M) is also polynomial time tractable.

This lets us prove the following dichotomy theorem.

Theorem 117. Let $M \in Sym_4^F(\mathbb{R}_{\geq 0})$. Then Pl-GH(M) is #P-hard, unless one of the following conditions is true, in which case, Pl-GH(M) is polynomial tractable.

- (1) (direct sum) $M \cong A \oplus B$ for polynomially tractable Pl-GH(A), Pl-GH(B),
- (2) (tensor product) $M \cong A \otimes B$ for polynomially tractable Pl-GH(A), Pl-GH(B).

Proof. If M is domain separable, i.e., $M \cong A \oplus B$ for any A, B, we know from Lemma 114 that Pl-GH(M) is #P-hard, unless Pl-GH(A), and Pl-GH(B) are both polynomially tractable, in which case, Pl-GH(M) is also polynomially tractable. If M is bipartite, we know from Lemma 116 that Pl-GH(M) is #P-hard, unless $M \cong A \otimes B$, where Pl-GH(A), and Pl-GH(B) are both polynomially tractable, in which case, Pl-GH(M) is also polynomially tractable.

Now, we consider any M that is neither domain separable, nor bipartite. The underlying graph of M where an edge (i, j) exists iff $M_{ij} > 0$ is connected and non-bipartite. In this case, since $M \in \operatorname{Sym}_4(\mathbb{R}_{\geq 0})$, there exists some $n^* \geq 1$ such that $M^n \in \operatorname{Sym}_4^F(\mathbb{R}_{>0})$ for all $n \geq n^*$. We let $n \geq n^*$ be some odd integer. From Theorem 112, we know that $\operatorname{Pl-GH}(M^n) \leq \operatorname{Pl-GH}(M)$ is $\#\operatorname{P-hard}$, unless M^n is isomorphic to $A' \otimes B'$, for some polynomial time tractable $\operatorname{Pl-GH}(A')$, $\operatorname{Pl-GH}(B')$, where $A', B' \in \operatorname{Sym}_2^F(\mathbb{R}_{>0})$, such that $(A')_{11} = (A')_{22}$, and $(B')_{11} = (B')_{22}$.

Without loss of generality, we will now consider the case where $M^n = A' \otimes B'$ for some $A', B' \in \text{Sym}_2^{\mathsf{F}}(\mathbb{R}_{>0})$. Since $(A')_{11} = (A')_{22}$, and $(B')_{11} = (B')_{22}$, we know that $A' = HD'_1H^{\mathsf{T}}$, and $B' = HD'_2H^{\mathsf{T}}$, where

$$H = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1\\ 1 & -1 \end{pmatrix}, D'_1 = \begin{pmatrix} A'_{11} + A'_{12} & 0\\ 0 & A'_{11} - A'_{12} \end{pmatrix}, D'_2 = \begin{pmatrix} B'_{11} + B'_{12} & 0\\ 0 & B'_{11} - B'_{12} \end{pmatrix}$$

So, if we let $\lambda'_1 = A'_{11} + A'_{12}$, $\lambda'_2 = A'_{11} - A'_{12}$, $\mu'_1 = B'_{11} + B'_{12}$, and $\mu'_2 = B'_{11} - B'_{12}$, we see that $M^n = A' \otimes B' = (H \otimes H)D'(H \otimes H)^T$, where $D' = \operatorname{diag}(\lambda'_1\mu'_1, \lambda'_1\mu'_2, \lambda'_2\mu'_1, \lambda'_2\mu'_2)$. We also note that since $A', B' \in \operatorname{Sym}_2(\mathbb{R}_{>0})$, $\lambda'_1 > |\lambda'_2|$, and $\mu'_1 > |\mu'_2|$. So, $|\lambda'_1\mu'_1| > |\lambda'_2\mu'_1| > |\lambda'_2\mu'_2|$, and $|\lambda'_1\mu'_1| > |\lambda'_1\mu'_2| > |\lambda'_2\mu'_2|$. If $\lambda'_2\mu'_1 \neq \lambda'_1\mu'_2$, then M^n would have four distinct eigenvalues. Otherwise, it will have three distinct eigenvalues, with one of them being repeated.

Since $A' \otimes B' = M^n$, we may assume that the eigenvalues of M are: $\nu_1, \nu_2, \nu_3, \nu_4$, such that $(\nu_1)^n = \lambda'_1 \mu'_1, (\nu_2)^n = \lambda'_1 \mu'_2, (\nu_3)^n = \lambda'_2 \mu'_1, (\nu_4)^n = \lambda'_2 \mu'_2$. Since n is an odd integer, and $\nu_i, \lambda'_i, \mu'_i$ are all real, this implies that $\nu_1 = (\lambda'_1 \mu'_1)^{\frac{1}{n}}, \nu_2 = (\lambda'_1 \mu'_2)^{\frac{1}{n}}, \nu_3 = (\lambda'_2 \mu'_1)^{\frac{1}{n}}, \nu_4 = (\lambda'_2 \mu'_2)^{\frac{1}{n}}$. Now, if we let $\lambda_i = (\lambda'_i)^{\frac{1}{n}}$, and $\mu_i = (\mu'_i)^{\frac{1}{n}}$ for $i \in [2]$, we find that $\nu_1 = \lambda_1 \mu_1, \nu_2 = \lambda_1 \mu_2, \nu_3 = \lambda_2 \mu_1, \nu_4 = \lambda_2 \mu_2$.

Since $M^n \cdot M = M \cdot M^n$, we know that M and M^n can be simultaneously diagonalized by the same orthogonal matrix. Now, if $\lambda'_2 \mu'_1 \neq \lambda'_1 \mu'_2$, we know that the diagonalizing orthogonal matrix for M^n is essentially unique, whose columns are the unit column eigenvectors of M^n corresponding to the respective eigenvalues. This is the matrix $(H \otimes H)K$, where $K = \begin{pmatrix} \pm 1 & 0 & 0 & 0 \\ 0 & \pm 1 & 0 & 0 & 0 \\ 0 & 0 & \pm 1 & 0 \\ 0 & 0 & \pm 1 & 0 \end{pmatrix}$. Note that for any diagonal matrix D', we have KD' = D'K. This implies that $M = (HD_1H^T) \otimes (HD_2H^T)$, where $D_1 = \operatorname{diag}(\lambda_1, \lambda_2)$, and $D_2 = \operatorname{diag}(\mu_1, \mu_2)$. On the other hand, if $\lambda'_2 \mu'_1 = \lambda'_1 \mu'_2$, we see that $(\lambda'_2 \mu'_1)^{\frac{1}{n}} = (\lambda'_1 \mu'_2)^{\frac{1}{n}}$. So, $\nu_3 = \lambda_2 \mu_1 = \lambda_1 \mu_2 = \nu_2$ as well. Therefore, we have the following orthogonal decomposition of $\mathbb{R}^4 = V_1 \oplus V_{23} \oplus V_4$ as a direct sum: V_1 (respectively, V_4) is a one-dimensional eigenspace corresponding to the eigenvalue $\nu'_1 = \lambda_1 \mu_1$ (respectively, $\nu_4 = \lambda_2 \mu_2$) of M; V_{23} is a two-dimensional eigenspace where both M^n and M act as scalar matrices. Thus any orthogonal matrix that diagonalizes M^n also diagonalizes M. So, once again, we see that the matrix M can be diagonalized by $H \otimes H$. So, in either case, we find that $M = (HD_1H^T) \otimes (HD_2H^T)$, where $D_1 = \operatorname{diag}(\lambda_1, \lambda_2)$, and $D_2 = \operatorname{diag}(\mu_1, \mu_2)$.

If we let $A = HD_1H^T$, and $B = HD_2H^T$, we see that $M = A \otimes B$. Moreover, we find that $A_{11} = \frac{1}{2}(\lambda_1 + \lambda_2) = A_{22}$, and $B_{11} = \frac{1}{2}(\mu_1 + \mu_2) = B_{22}$. So, we see that when $M^n = A' \otimes B'$, where $(A')_{11} = (A')_{22}$, and $(B')_{11} = (B')_{22}$, then $M = A \otimes B$ for some A, B such that $A_{11} = A_{22}$, and $B_{11} = B_{22}$. Moreover, in this case, it is easily seen that P1-GH(M) is polynomially tractable.

References

- [Ats+19] Albert Atserias, Laura Mančinska, David E. Roberson, Robert Šámal, Simone Severini, and Antonios Varvitsiotis. "Quantum and non-signalling graph isomorphisms". In: Journal of Combinatorial Theory, Series B 136 (2019), pp. 289–328.
- [Bac17] Miriam Backens. "A New Holant Dichotomy Inspired by Quantum Computation". In: 44th International Colloquium on Automata, Languages, and Programming, ICALP 2017, July 10-14, 2017, Warsaw, Poland. Vol. 80. LIPIcs. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2017, 16:1–16:14.
- [Bac18] Miriam Backens. "A Complete Dichotomy for Complex-Valued Holant^c". In: 45th International Colloquium on Automata, Languages, and Programming, ICALP 2018, July 9-13, 2018, Prague, Czech Republic. Vol. 107. LIPIcs. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2018, 12:1–12:14.
- [BG05] Andrei Bulatov and Martin Grohe. "The complexity of partition functions". In: *Theo*retical Computer Science 348.2-3 (2005), pp. 148–186.
- [Bul+12] Andrei Bulatov, Martin Dyer, Leslie Ann Goldberg, Markus Jalsenius, Mark Jerrum, and David Richerby. "The complexity of weighted and unweighted # CSP". In: Journal of Computer and System Sciences 78.2 (2012), pp. 681–688.
- [Bul13] Andrei A. Bulatov. "The complexity of the counting constraint satisfaction problem". In: Journal of the ACM (JACM) 60.5 (2013), pp. 1–41.
- [CC17] Jin-Yi Cai and Xi Chen. "Complexity of counting CSP with complex weights". In: Proceedings of the forty-fourth annual ACM symposium on Theory of computing (STOC) (2012), pp. 909–920. Vol. 64(3). Journal of the ACM. 2017, pp. 1–39.
- [CCL13] Jin-Yi Cai, Xi Chen, and Pinyan Lu. "Graph homomorphisms with complex values: A dichotomy theorem". In: *SIAM Journal on Computing* 42.3 (2013), pp. 924–1029.
- [CCL16] Jin-Yi Cai, Xi Chen, and Pinyan Lu. "Nonnegative weighted # CSP: an effective complexity dichotomy". In: *SIAM Journal on Computing* 45.6 (2016), pp. 2177–2198.
- [CF19] Jin-Yi Cai and Zhiguo Fu. "Holographic algorithm with Matchgates is universal for planar # CSP over boolean domain". In: SIAM Journal on Computing 51.2 (2019), STOC17–50.
- [CGW16] Jin-Yi Cai, Heng Guo, and Tyson Williams. "A complete dichotomy rises from the capture of vanishing signatures". In: SIAM Journal on Computing 45.5 (2016), pp. 1671– 1728.
- [CLX09] Jin-Yi Cai, Pinyan Lu, and Mingji Xia. "Holant problems and counting CSP". In: Proceedings of the forty-first annual ACM symposium on Theory of computing. 2009, pp. 715–724.

- [CM23] Jin-Yi Cai and Ashwin Maran. "The complexity of counting planar graph homomorphisms of domain size 3". In: Proceedings of the 55th Annual ACM Symposium on Theory of Computing. 2023, pp. 1285–1297.
- [DG00] Martin Dyer and Catherine Greenhill. "The complexity of counting graph homomorphisms (extended abstract)". In: Proceedings of the Eleventh Annual ACM-SIAM Symposium on Discrete Algorithms, January 9-11, 2000, San Francisco, CA, USA. Ed. by David B. Shmoys. ACM/SIAM, 2000, pp. 246–255.
- [DR10] Martin E. Dyer and David M. Richerby. "On the complexity of # CSP". In: *Proceedings* of the forty-second ACM symposium on Theory of computing. 2010, pp. 725–734.
- [DR11] Martin E. Dyer and David M. Richerby. "The # CSP Dichotomy is Decidable". In: 28th International Symposium on Theoretical Aspects of Computer Science (STACS 2011).
 Ed. by Thomas Schwentick and Christoph Dürr. Vol. 9. Dagstuhl, Germany: Schloss Dagstuhl-Leibniz-Zentrum fuer Informatik, 2011, pp. 261–272. DOI: 10.4230/LIPIcs.STACS.2011.261.
- [DR13] Martin E. Dyer and David M. Richerby. "An effective dichotomy for the counting constraint satisfaction problem". In: *SIAM Journal on Computing* 42.3 (2013), pp. 1245– 1274.
- [FY14] Zhiguo Fu and Fengqin Yang. "Holographic algorithms on bases of rank 2". In: Information Processing Letters 114.11 (2014), pp. 585–590.
- [FYY19] Zhiguo Fu, Fengqin Yang, and Minghao Yin. "On blockwise symmetric matchgate signatures and higher domain # CSP". In: Information and Computation 264 (2019), pp. 1– 11.
- [Ger31] Semyon Aranovich Gershgorin. "Uber die abgrenzung der eigenwerte einer matrix". In: Известия Российской академии наук. Серия математическая 6 (1931), pp. 749– 754.
- [Gol+10] Leslie Ann Goldberg, Martin Grohe, Mark Jerrum, and Marc Thurley. "A complexity dichotomy for partition functions with mixed signs". In: SIAM Journal on Computing 39.7 (2010), pp. 3336–3402.
- [GW20] Heng Guo and Tyson Williams. "The complexity of planar Boolean #CSP with complex weights". In: Journal of Computer and System Sciences 107 (2020), pp. 1–27.
- [HM87] Gary Harris and Clyde Martin. "Shorter notes: The roots of a polynomial vary continuously as a function of the coefficients". In: *Proceedings of the American Mathematical Society* (1987), pp. 390–392.
- [Jac85] N. Jacobson. Basic Algebra. Basic Algebra v. 2. W.H. Freeman, 1985. ISBN: 9780716719335. URL: https://books.google.com/books?id=oNmDSAAACAAJ.
- [Kas61] Pieter W. Kasteleyn. "The statistics of dimers on a lattice: I. The number of dimer arrangements on a quadratic lattice". In: *Physica* 27.12 (1961), pp. 1209–1225.
- [Kas63] Pieter W. Kasteleyn. "Dimer statistics and phase transitions". In: Journal of Mathematical Physics 4.2 (1963), pp. 287–293.

- [Kas67] Pieter Kasteleyn. "Graph theory and crystal physics". In: *Graph theory and theoretical physics* (1967), pp. 43–110.
- [KP02] Steven G. Krantz and Harold R. Parks. A Primer of Real Analytic Functions (Second ed.) Birkhauser, 2002.
- [LMR20] Martino Lupini, Laura Mančinska, and David E. Roberson. "Nonlocal games and quantum permutation groups". In: *Journal of Functional Analysis* 279.5 (2020), p. 108592.
- [Lov12] László Lovász. Large networks and graph limits. Vol. 60. American Mathematical Soc., 2012.
- [MR20] Laura Mančinska and David E. Roberson. "Quantum isomorphism is equivalent to equality of homomorphism counts from planar graphs". In: 2020 IEEE 61st Annual Symposium on Foundations of Computer Science (FOCS). IEEE. 2020, pp. 661–672.
- [TF61] Harold NV. Temperley and Michael E. Fisher. "Dimer problem in statistical mechanicsan exact result". In: *Philosophical Magazine* 6.68 (1961), pp. 1061–1063.
- [Val01] Leslie G. Valiant. "Quantum computers that can be simulated classically in polynomial time". In: Proceedings of the thirty-third annual ACM symposium on Theory of computing. 2001, pp. 114–123.
- [Val08] Leslie G Valiant. "Holographic algorithms". In: SIAM Journal on Computing 37.5 (2008), pp. 1565–1594.
- [Val79] Leslie G. Valiant. "The complexity of computing the permanent". In: Theoretical computer science 8.2 (1979), pp. 189–201.
- [Ver05] Dirk Vertigan. "The computational complexity of Tutte invariants for planar graphs". In: SIAM Journal on Computing 35.3 (2005), pp. 690–712.
- [YF22] Peng Yang and Zhiguo Fu. "Local holographic transformations: tractability and hardness". In: Frontiers of Computer Science 17.2 (2022), pp. 1–11.