## Local unitarty equivalence and entanglement by Bargmann invariants

Lin Zhang<sup>1</sup>\*, Bing Xie<sup>1</sup>, Yuanhong Tao<sup>2</sup>

<sup>1</sup>School of Science, Hangzhou Dianzi University, Hangzhou 310018, People's Republic of China <sup>2</sup>College of Science, Zhejiang University of Science and Technology, Hangzhou, Zhejiang 310023, PR China

15Dec2024

#### Abstract

The study of quantum states frequently examines the connection between non-local effects in quantum mechanics and quantities that remain unchanged under local unitary transformations. Among these concepts, local unitary equivalence, defined through local unitary transformations, holds significant importance in quantum state classification and resource theory. This paper focuses on the fundamental issue of local unitary equivalence for multipartite quantum states within quantum information theory, with the aim of identifying the comprehensive set of invariants that define their local unitary orbits. These invariants are crucial for deriving polynomial invariants and describing physical properties that remain invariant under local unitary transformations. Specifically, the research delves into the characterization of local unitary equivalence and the detection of entanglement using local unitary Bargmann invariants, utilizing the generalized Schur-Weyl duality to analyze tensor product symmetries. Taking the two-qubit system as an example, our study demonstrates the measurability of the invariants that determine local unitary equivalence and establishes a relationship between Makhlin<sub>i</sub> s fundamental invariants (a complete set of 18 local unitary invariants for two-qubit states) and local unitary Bargmann invariants. These Bargmann invariants are related to the trace of density operator products with marginal states and can be measured through a cycle test, an extension of the SWAP test. These findings offer a practical criterion for en- tanglement detection based on local unitary Bargmann invariants, contributing to the advancement of quantum information theory and its applications.

<sup>\*</sup>E-mail: godyalin@163.com

## Contents

1	Intro	oduction	3
2	Barg	mann invariants and local equivalence of quantum states	5
3	Enta	inglement criterion via LU Bargmann invariants	7
4	Disc	cussion and conclusion	7
Appendices			11
A	Proc	of of Proposition 2.1	11
	A.1	Invariant theory	11
	A.2	The generalized Schur-Weyl duality	12
	A.3	Proof of Proposition 2.1	13
B	Proc	of of Theorem 2.2	14
	<b>B</b> .1	Product formula for two-qubit observables	15
		B.1.1 Product formula	18
		B.1.2 Auxiliary results	24
		B.1.3 Recurrence relation for the matrix power	33
	B.2	Some results about products involved two-qubit states	35
	B.3	Revisiting local unitary invariants	40
	B.4	Proof of Theorem 2.2	48
C	Proc	of of Theorem 3.1	48
	<b>C</b> .1	Entanglement criterion by Makhlin's invariants	48
	C.2	Proof of Theorem 3.1	52

## 1 Introduction

In the rapidly evolving field of quantum information science, understanding and manipulating quantum states is paramount. Among the myriad phenomena that quantum mechanics offers, local unitary equivalence and entanglement stand out as fundamental yet intricate concepts. Local unitary equivalence, which posits that certain quantum states are indistinguishable under local operations and classical communication (LOCC), lies at the heart of quantum state classification and resource theories. Entanglement, on the other hand, serves as a cornerstone for quantum computing [1], quantum cryptography [2], and various quantum communication protocols [3, 4, 5], underscoring its pivotal role in harnessing the power of quantum mechanics.

The local unitary equivalence [6, 7, 8, 9, 10], defined through local unitary transformations, holds significant importance in quantum information science because the importance of local unitary transformations lies in their crucial roles in quantum state manipulation, quantum algorithm design, and quantum state classification. The elaboration on these key roles can be described in detail:

- Local unitary transformations are indispensable tools in quantum state manipulation [11]. They enable precise control over the local properties of quantum states without altering the overall properties of the quantum system. Specifically, local unitary transformations can act on certain or some subsystems within a quantum system, thereby changing the quantum states of these subsystems while maintaining the inner product and norm of the overall system's quantum state. This characteristic makes local unitary transformations widely applicable in processes such as quantum state preparation, manipulation, and measurement. For instance, in quantum computing, we can utilize local unitary transformations to realize interactions and information transfer between qubits, thereby constructing complex quantum algorithms and quantum networks.
- Local unitary transformations play a pivotal role in quantum algorithm design [12], a core issue in the field of quantum computing. On the one hand, local unitary transformations can serve as basic operational units for quantum algorithms, with complex quantum algorithms being constructed by combining different local unitary transformations. On the other hand, local unitary transformations can be used to optimize the performance of quantum algorithms. For example, in quantum search algorithms, we can adjust the structure of the search space through local unitary transformations, thereby improving search efficiency. Additionally, local unitary transformation, providing more possibilities and flexibility for quantum algorithm design.

• Local unitary transformations play a key role in quantum state classification [13], an important problem in quantum information science. Firstly, local unitary transformations can be used to determine whether two quantum states are equivalent. In quantum state classification, we need to ascertain which quantum states are equivalent under local unitary transformations, meaning they can be transformed into each other through local unitary transformations. This equivalence judgment is significant for understanding the properties of quantum states and constructing quantum algorithms. Secondly, local unitary transformations can be used to study the entanglement properties of quantum states. Entanglement is an important characteristic of quantum states, describing non-classical correlations between quantum states. Local unitary transformations can preserve the entanglement properties of quantum states, providing a powerful tool for studying entanglement phenomena.

The characterization of local unitary equivalence and the detection of entanglement are crucial for advancing our understanding and applications of quantum systems. Despite significant progress, these tasks remain challenging due to the complex nature of quantum states and the high-dimensional spaces they inhabit. The complex interaction between local unitary transformations and global quantum characteristics requires a refined method to distinguish equivalent states and efficiently recognize entangled states.

This paper delves into the problem of characterizing local unitary equivalence of multipartite states and detecting entanglement in quantum systems by using locally unitary Bargmann invariants, which are involved in many protocols such as quantum fingerprinting [14], etc. The Bargmann invariants are a gauge-invariant quantity that is related to the geometric phase acquired by a quantum system undergoing a cyclic evolution [15], and are also connected to Wigner rotations and null phase curves [16], besides they are used in Kirkwood-Dirac quasi-probability representation [17, 18, 19] and used as witness of quantum imaginarity [20], as well. Note that Bargmann invariants are also known as multivariate traces [21], which can be estimated by a constant quantum depth circuit [22] while respecting near-term quantum architecture constraints. With the constant quantum depth circuit, theoretical speaking, we can use it to test the LU equivalence operationally by experiment. We aim to provide a comprehensive framework that leverages theoretical insights and practical algorithms to tackle these challenges. By exploring the mathematical structure of quantum states and the properties of local unitary transformations, we seek to elucidate the conditions under which two quantum states are locally equivalent. Furthermore, we propose and analyze the proposed entanglement detection methods by Bargmann invariants.

Our work is structured as follows: We first review the foundational concepts of local unitary equivalence, setting the stage for our subsequent discussions. We then delve into the theoretical underpinnings of local unitary transformations and their implications for quantum state classifi-

cation. Next, we present a criterion of entanglement detection by Bargmann invariants which are locally unitary invariant, discussing their strengths and limitations. We conclude by proposing novel methods and algorithms that address the current challenges in characterizing local unitary equivalence and detecting entanglement, and we outline potential directions for future research. By contributing to the ongoing discourse on these foundational quantum phenomena, we hope to pave the way for new advancements in quantum information science and technology.

## 2 Bargmann invariants and local equivalence of quantum states

Before proceeding, let us fix notations used in this paper. Given two tuples of N states  $\Psi = (\rho_1, \ldots, \rho_N)$  and  $\Psi' = (\rho'_1, \ldots, \rho'_N)$  acting on Hilbert space  $\mathbb{C}^d$ , if there exists a unitary  $\mathbf{U} \in U(d)$ , the unitary group acting on  $\mathbb{C}^d$ , such that  $\rho'_k = \mathbf{U}\rho_k\mathbf{U}^\dagger$  for each  $k = 1, 2, \ldots, N$ , we say  $\Psi$  and  $\Psi'$  are *unitarily equivalent*. If there exists a set of invariant properties allows us to decide whether two tuples of states are unitarily equivalent, this set is said to be complete.

Consider a tuple of *N* pure/mixed quantum states  $\Psi = (\rho_1, ..., \rho_N)$ , where states  $\rho_k$ 's act on the same underlying Hilbert space. The *Bargmann invariant* (aka multivariate traces [21, 22]) of this tuple of states is defined as

$$\Delta_{12\cdots N}(\Psi) = \operatorname{Tr}\left(\rho_1 \rho_2 \cdots \rho_N\right). \tag{2.1}$$

Bargmann invariants can be used to describe the unitarily equivalence between tuples of states. In fact, we have already known the following result [23]: Let  $\Psi = (\rho_1, \ldots, \rho_N)$  and  $\Psi' = (\rho'_1, \ldots, \rho'_N)$  be two tuples of mixed states on  $\mathbb{C}^d$ . Then both  $\Psi$  and  $\Psi'$  are unitarily equivalent if and only if, for every  $m \in \mathbb{N}$  and for every sequence  $i_1, i_2, \ldots, i_m$  of numbers from  $\{1, \ldots, N\}$ , the corresponding Bargmann invariants of degree m agree

$$\operatorname{Tr}\left(\rho_{i_{1}}\rho_{i_{2}}\cdots\rho_{i_{m}}\right)=\operatorname{Tr}\left(\rho_{i_{1}}^{\prime}\rho_{i_{2}}^{\prime}\cdots\rho_{i_{m}}^{\prime}\right).$$
(2.2)

Recently, quantum circuits such as cycle test was introduced, which enable the *direct measurement* of complete sets of Bargmann invariants for a tuple of quantum states [21]. Motivated by this result, we will investigate the locally unitary equivalent of tuples of multipartite states using locally unitary Bargmann invariants.

The same paradigm in the last section motivated the usage of invariant polynomials in the context of classification of entanglement classes subject to local unitary transformation. Let  $V = \text{Herm}(\mathbb{C}^{d_1} \otimes \cdots \otimes \mathbb{C}^{d_N})$ , the Hermitian matrices acting on the tensor space, and denote the local unitary group by  $U(d) = U(d_1) \otimes \cdots \otimes U(d_N)$ . Define U(d) acts on V by conjugation, i.e., for any  $g \in U(d)$  and  $X \in V$ , we get the LU action of U(d) on V via  $\tau_g X = gXg^{\dagger}$ . In fact, given two tuples of multipartite states on  $\mathbb{C}^{d_1} \otimes \cdots \otimes \mathbb{C}^{d_N}$ ,  $\Psi = (\rho_1, \dots, \rho_N)$  and  $\Psi' = (\rho'_1, \dots, \rho'_N)$ , they are locally

unitarily (LU) equivalent in the sense that  $\rho'_i = g\rho_i g^{\dagger}$  for all i = 1, ..., N and some  $g \in U(d)$ . That is, there exist a collection of unitary operators  $U_i \in U(d_i)$  such that  $g = U_1 \otimes \cdots \otimes U_N$ , where i = 1, ..., N, and

$$\rho_i' = (\boldsymbol{U}_1 \otimes \cdots \otimes \boldsymbol{U}_N) \rho_i (\boldsymbol{U}_1 \otimes \cdots \otimes \boldsymbol{U}_N)^{\dagger}$$
(2.3)

for each i = 1, ..., N. Clearly, when N = 1, this problem is reduced to a well-known locally unitary equivalence of two multipartite states. In what follows, we characterize locally unitary equivalence between two multipartite states using measurable quantities which can be expressed as a linear combination of locally unitary Bargmann invariants.

**Proposition 2.1.** For any two N-partite states  $\rho$  and  $\sigma$  acting on  $\mathbb{C}^{d_1} \otimes \cdots \otimes \mathbb{C}^{d_N}$ , they are LU equivalent, *i.e.*,  $\sigma = g\rho g^{\dagger}$  for some  $g \in U(d)$ , if and only if, for arbitrary positive integer n, it holds that

$$\operatorname{Tr}\left(\sigma^{\otimes n}\mathbf{P}_{d,n}(\boldsymbol{\pi}) = \operatorname{Tr}\left(\rho^{\otimes n}\mathbf{P}_{d,n}(\boldsymbol{\pi})\right),\tag{2.4}$$

where the meaning of  $\mathbf{P}_{d,n}(\pi)$  will be explained immediately for all  $\pi := (\pi_1, \ldots, \pi_N) \in S_n^N$ , the Cartesian product of N copies of the permutation group of n distinct elements.

It is easily seen that these quantities involved in Eq. (2.4) can be shown to be a linear combination of locally unitary Bargmann invariants  $\text{Tr}(\rho_{i_1} \cdots \rho_{i_n})$ , where each  $\rho_{i_k}$ 's is taken from the sequence of states  $\{\rho_{\Lambda} : \Lambda \subseteq \{1, \ldots, N\}\}$ , where  $\rho_{\Lambda} = \text{Tr}_{\bar{\Lambda}}(\rho)$ , where  $\bar{\Lambda} := \{1, \ldots, N\}\setminus\Lambda$ . Due to the measurability of Bargmann invariants, the above result in fact leads an operational test for LU equivalence.

In order to derive a further result, let us focus on the special bipartite case. Given two bipartite states  $\rho_{AB}$  and  $\rho'_{AB}$  on  $\mathbb{C}^m \otimes \mathbb{C}^n (m = n = 2)$ , let  $\Psi = (\rho_{AB}, \rho_A \otimes \mathbb{1}_B, \mathbb{1}_A \otimes \rho_B)$ , where  $\mathbb{1}_X (X = A, B)$  is the identity operator, and  $\Psi' = (\rho'_{AB}, \rho'_A \otimes \mathbb{1}_B, \mathbb{1}_A \otimes \rho'_B)$ , we will study the locally unitary equivalence of two tuples  $\Psi$  and  $\Psi'$ . It is easily seen that  $\rho_{AB}$  is LU equivalent to  $\rho'_{AB}$  if and only if  $\Psi$  is LU equivalent to  $\Psi'$ .

Although a complete set of LU invariants of two-qubit states is given already by Makhlin in 2002, we would like here to work out a complete set of LU invariants of two-qubit states in terms of Bargmann invariants which are measurable quantities, interested by experimenter.

**Theorem 2.2.** Given any two-qubit state  $\rho_{AB} \in D(\mathbb{C}^2 \otimes \mathbb{C}^2)$ . Denote  $X_0 = \rho_{AB}, X_1 = \rho_A \otimes \mathbb{1}_B$ , and  $X_2 = \mathbb{1}_A \otimes \rho_B$ . All LU invariant polynomials of two-qubit states can be generated by the following 18

locally unitary Bargmann invariants  $B_k(k = 1, ..., 18)$ , where the meanings of  $B_k$ 's are given below:

$$B_{1} := \operatorname{Tr} (X_{0}X_{1}), B_{2} := \operatorname{Tr} (X_{0}X_{2}), B_{3} := \operatorname{Tr} (X_{0}X_{1}X_{2}), B_{4} := \operatorname{Tr} (X_{0}^{2}),$$

$$B_{5} := \operatorname{Tr} (X_{0}^{2}X_{1}X_{2}), B_{6} := \operatorname{Tr} (X_{0}^{3}), B_{7} := \operatorname{Tr} (X_{0}^{3}X_{1}), B_{8} := \operatorname{Tr} (X_{0}^{3}X_{2}),$$

$$B_{9} := \operatorname{Tr} (X_{0}^{3}X_{1}X_{2}), B_{10} := \operatorname{Tr} (X_{0}^{4}), B_{11} := \operatorname{Tr} (X_{0}^{2}X_{1}X_{0}^{2}X_{1}), B_{12} := \operatorname{Tr} (X_{0}^{2}X_{2}X_{0}^{2}X_{2}),$$

$$B_{13} := \operatorname{Tr} (X_{0}X_{1}X_{2}X_{0}^{2}X_{1}), B_{14} := \operatorname{Tr} (X_{0}X_{1}X_{2}X_{0}^{2}X_{2}), B_{15} := \operatorname{Tr} (X_{0}X_{1}X_{2}X_{0}^{3}X_{1}),$$

$$B_{16} := \operatorname{Tr} (X_{0}X_{1}X_{2}X_{0}^{3}X_{2}), B_{17} := \operatorname{Tr} (X_{0}X_{1}X_{0}^{2}X_{1}X_{0}^{3}X_{1}), B_{18} := \operatorname{Tr} (X_{0}X_{2}X_{0}^{2}X_{2}X_{0}^{3}X_{2}).$$
(2.5)

Moreover, the specific expressions for all Makhlin invariants  $L_k$ 's are analytically expressed by using Bargmann invariants  $B_k$ 's given in Appendix.

The proof of this theorem can be finished by finding analytical relations between Makhlin invariants  $L_k$ 's and Bargmann invariants  $B_k$ 's. In other words, as generators of a complete set of LU invariants,  $B_k$ 's are more important than  $L_k$ 's because  $B_k$ 's are measurable by the recent proposed quantum circuit, the named cycle test. Therefore, we can determine whether two *unknown* two-qubit states are LU equivalent if and only if they have the same values on the 18 Bargmann generators by measurement.

## 3 Entanglement criterion via LU Bargmann invariants

From the connection between Makhlin invariants and Bargmann invariants, we will get a physical and operational criterion in entanglement detection. In fact, we get the following:

**Theorem 3.1.** A two-qubit state  $\rho_{AB}$  is entangled if and only if the following inequality holds true:

$$6(B_1 + B_2 - B_1B_2 - B_4 - B_{10}) + 12(B_5 - B_3) + 3B_4^2 + 4B_6 < 1,$$
(3.1)

where the meanings of  $B_k$ 's here are taken from Eq. (2.5). Explicitly, Eq. (3.1) can be equivalently rewritten as

$$6 \left[ \operatorname{Tr} \left( \rho_{A}^{2} \right) + \operatorname{Tr} \left( \rho_{B}^{2} \right) - \operatorname{Tr} \left( \rho_{A}^{2} \right) \operatorname{Tr} \left( \rho_{B}^{2} \right) - \operatorname{Tr} \left( \rho_{AB}^{2} \right) - \operatorname{Tr} \left( \rho_{AB}^{4} \right) \right] + 12 \left[ \operatorname{Tr} \left( \rho_{AB}^{2} (\rho_{A} \otimes \rho_{B}) \right) - \operatorname{Tr} \left( \rho_{AB} (\rho_{A} \otimes \rho_{B}) \right) \right] + 3 \left[ \operatorname{Tr} \left( \rho_{AB}^{2} \right) \right]^{2} + 4 \operatorname{Tr} \left( \rho_{AB}^{3} \right) < 1.$$
(3.2)

## 4 Discussion and conclusion

In [25], the authors proposed a test for entanglement of two-qubit states. But, it is hard to determine the separability of an *unknown state* in practice because one has to check their inequality for all sets of local testing observables being complementary. We see from Eq. (3.1) or Eq. (3.2) that in order to determine entanglement in an *unknown* two-qubit state, it suffices to measure only 7 locally unitarily Bargmann invariants for such two-qubit state.

In this work, we have explored the local unitary equivalence of multipartite states using Bargmann invariants. We have identified a complete set of 18 Bargmann generators that constitute local unitary (LU) invariants for two-qubit states. Building on this foundation, we propose a method to characterize entanglement in unknown two-qubit states by measuring a subset of seven out of these 18 Bargmann generators. Our approach can be extended to higher-dimensional state spaces. Our findings also inspire novel experimental designs to test entanglement in unknown quantum states. In future research, we plan to investigate the relationships between the moments of the probability distribution of random measurements [26] and Bargmann invariants.

#### Acknowledgments

This research is supported by Zhejiang Provincial Natural Science Foundation of China under Grant No. LZ23A010005 and by NSFC under Grant No.11971140.

## References

- R. Jozsa and N. Linden, On the role of entanglement in quantum-computational speed-up, Proc. R. Soc. Lond. A 459, 2011(2003).
- [2] C. Portmann and R. Renner, Security in quantum cryptography, Rev. Math. Phys. 94, 025008(2022).
- [3] C.H. Bennett and S.J. Wiesner, Communication via one- and two-particle operators on Einstein-Podolsky-Rosen states, Phys. Rev. Lett. 69, 2881(1992).
- [4] C.H. Bennett, G. Brassard, C. Crépeau, R. Jozsa, A. Peres, and W. Wootters, Teleporting an unknown quantum state via dual classical and EPR channels, Phys. Rev. Lett. 70, 1895(1993).
- [5] D. Bouwmeester, J.W. Pan, K. Mattle, M. Eibl, H. Weinfurter, and A. Zeilinger, Experimental quantum teleportation, Nature **390**, 575 (1997).
- [6] Y. Makhlin, Nonlocal properties of two-qubit gates and mixed states, and the optimization of quantum computations, Quant Inf Process **1**, 243-252 (2002).
- [7] B. Kraus, Local unitary equivalence of multipartite pure states, Phys. Rev. Lett. **104**, 020504 (2010).
- [8] C. Zhou, T. Zhang, S-M. Fei, N. Jing, and X. Li-Jost, Local unitary equivalence of arbitrary dimensional bipartite quantum states, Phys. Rev. A **86**, 010303(R) (2012).

- [9] N. Jing, S-M. Fei, M. Li, X. Li-Jost, and T. Zhang, Local unitary invariants of generic multiqubit states, Phys. Rev. A **92**, 022306 (2015).
- [10] A.M. Martins, Necessary and Sufficient Conditions for Local Unitary Equivalence of Multiqubit States, Phys. Rev. A 91, 042308 (2015)
- [11] G. Gour, N.R. Wallach, Necessary and sufficient conditions for local manipulation of multipartite pure quantum states, New. J. Phys. **13**, 073013 (2011).
- [12] J.J. Vartiainen, Unitary Transformations for Quantum Computing, PhD Thesis (2005).
- [13] G. Gour, N.R. Wallach, Classification of multipartite entanglement of all finite dimensionality, Phys. Rev. Lett. 111, 060502 (2013).
- [14] H. Buhrman, R. Cleve, J. Watrous, and R. de Wolf, Quantum fingerprinting, Phys. Rev. Lett. 87, 167902(2001).
- [15] R. Simon and N. Mukunda, Bargmann invariant and the geometry of the Güoy effect, Phys. Rev. Lett. 70, 880(1993).
- [16] N. Mukunda, P.K. Aravind, and R. Simon, Wigner rotations, Bargmann invariants and geometric phases, J. Phys. A : Math. Gen. 36, 2347 (2003).
- [17] J.G. Kirkwood, Quantum statistics of almost classical assemblies, 44, 31(1933).
- [18] P.A.M. Dirac, On the analogy between classical and quantum mechanics, Rev. Math. Phys. 17, 195(1945).
- [19] C. Bamber and J.S. Lundeen, Observing Dirac's classical phase space analog to the quantum state, Phys. Rev. Lett. **112**, 31(2014).
- [20] C. Fernandes, R. Wagner, L. Novo, and E.F. Galvão, Unitary-Invariant Witnesses of Quantum Imaginarity, Phys. Rev. Lett. 133, 190201 (2024).
- [21] M. Oszmaniec, D.J. Brod and E.F. Galvão, Measuring relational information between quantum states, and applications, New. J. Phys. 26, 013053 (2024).
- [22] Y. Quek, E. Kaur, and M.M. Wilde, Multivariate trace estimation in constant quantum depth, Quantum 8, 1220 (2024).
- [23] C. Procesi, The invariant theory of  $n \times n$  matrices, Adv.Math. **19**, 306 (1976).
- [24] P. Vrana, Group representations in entanglement theory, PhD Thesis (2012).

- [25] S. Yu, J-W. Pan, Z-B. Chen, and Y-D. Zhang, Comprehensive test of entanglement for twolevel systems via the indeterminacy relationship, Phys. Rev. Lett. 91, 217903 (2003).
- [26] N. Wyderka, A. Ketterer, S. Imai, J.L. Bönsel, D.E. Jones, B.T. Kirby, X-D. Yu, and O. Gühne, Complete characterization of quantum correlations by randomized measurements, Phys. Rev. Lett. 131, 090201 (2023).
- [27] L. Zhang, Matrix integrals over unitary groups: An application of Schur-Weyl duality, arXiv:1408.3782v6
- [28] R.A. Horn and C.R. Johnson, Matrix Analysis, Cambridge University Press (2nd Ed), (2013).

# Appendices

In what follows, we proceed to prove each of them. When deriving these main results, we present some essential tools that facilitate the obtainment of additional findings. For instance, we establish a rigorous relationship concerning the conversion between Makhlin's fundamental invariants and LU Bargmann's invariants. With these preparations, we can calculate arbitrary locally unitary Bargmann invariants Tr ( $\rho_{i_1} \cdots \rho_{i_N}$ ), where each  $\rho_{i_k}$  is from the set { $\rho_{AB}$ ,  $\rho_A \otimes \mathbb{1}_B$ ,  $\mathbb{1}_A \otimes \rho_B$ } for any two-qubit state  $\rho_{AB}$ , up to ignoring dimensional factors.

## A Proof of Proposition 2.1

For the proof of Proposition 2.1 in the main text, we used a lot of tools which cannot be explained in detail within the confines of that proposition's discussion. Now in this section, we will present a more comprehensive and detailed exploration of these tools, providing the necessary background, definitions, and explanations to fully understand their application in the proof. This deeper dive will not only clarify the intricacies of the proof but also enhance the reader's grasp of the underlying mathematical concepts and techniques. By doing so, we aim to make the proof of Proposition 2.1 more accessible and insightful for a broader audience.

#### A.1 Invariant theory

Let *K* be a compact group and let

$$\Pi: K \ni g \mapsto \Pi_g \in \mathsf{GL}(V) \tag{A.1}$$

be a representation of *K* in a finite dimensional real vector space *V*. Since *K* is compact, we can assume that  $\Pi_g$  is an orthogonal transformation. That is,

$$\Pi: K \ni g \mapsto \Pi_g \in \mathsf{O}(V). \tag{A.2}$$

The space of all real polynomials on *V* is denoted by  $\mathbb{R}[V]$ . We will denote by  $\mathbb{R}[V]_n$  the space of real homogeneous polynomials on *V* of degree *n*. Homogeneous polynomials of degree *n* are mappings of the form:

$$p(\boldsymbol{v}) = \left\langle \tilde{\boldsymbol{p}}, \boldsymbol{v}^{\otimes n} \right\rangle \tag{A.3}$$

where  $\langle \cdot, \cdot \rangle$  is the *K*-invariant inner product in  $V^{\otimes n}$  (induced by the inner product on *V*), and  $\tilde{p} \in V^{\otimes n}$  is a tensor encoding the polynomial *p*.

Invariant homogeneous polynomials of degree n are polynomials that must satisfy

$$p(\Pi_{g^{-1}}\boldsymbol{v}) = p(\boldsymbol{v}) \tag{A.4}$$

for every  $v \in V$  and  $g \in K$ . This is equivalent to

$$\langle \tilde{\boldsymbol{p}}, \boldsymbol{v}^{\otimes n} \rangle = \langle \tilde{\boldsymbol{p}}, (\Pi_{g^{-1}} \boldsymbol{v})^{\otimes n} \rangle = \langle \tilde{\boldsymbol{p}}, \Pi_{g^{-1}}^{\otimes n} \boldsymbol{v}^{\otimes n} \rangle = \langle \Pi_{g}^{\otimes n} \tilde{\boldsymbol{p}}, \boldsymbol{v}^{\otimes n} \rangle,$$
 (A.5)

which implies

$$\Pi_g^{\otimes k} \tilde{p} = \tilde{p} \tag{A.6}$$

for every  $g \in K$ .

Denote the set of all *K*-invariant polynomials by  $\mathbb{R}[V]^K$ . It is well known result in invariant theory that in the case of compact groups we can use invariant polynomials in  $\mathbb{R}[V]^K$  to decide about equivalence of elements of *V* under the action of *K*.

**Proposition A.1** ([24]). For  $u, v \in V$ , we have  $v = \prod_g u$ , for some  $g \in K$  if and only if for every invariant polynomial  $p \in \mathbb{R}[V]^K$ , we have p(v) = p(u).

Because every polynomial can be decomposed into the direct sum of homogeneous polynomials, this implies  $\mathbb{R}[V]^K = \bigoplus_{n=1}^{\infty} \mathbb{R}[V]_n^K$ . Then the above Proposition A.1 can be restated as

**Proposition A.2.** For  $u, v \in V$ , we have  $v = \prod_g u$ , for some  $g \in K$  if and only if for every K-invariant homogeneous polynomial  $p_n$  of degree n, we have  $p_n(v) = p_n(u)$ , where n = 1, 2, ...

### A.2 The generalized Schur-Weyl duality

Consider a system of *n* qudits, acting on  $(\mathbb{C}^d)^{\otimes n}$  each with a standard local computational basis  $\{|i\rangle, i = 1, ..., d\}$ . The Schur-Weyl duality relates transforms on the system performed by local *d*-dimensional unitary operations to those performed by permutation of the qudits. Recall that the symmetric group  $S_n$  is the group of all permutations of *n* objects. This group is naturally represented in our system by

$$\mathbf{P}_{d,n}(\pi)|i_{1}\cdots i_{n}\rangle := |i_{\pi^{-1}(1)}\cdots i_{\pi^{-1}(n)}\rangle, \tag{A.7}$$

where  $\pi \in S_n$  is a permutation and  $|i_1 \cdots i_n\rangle$  is shorthand for  $|i_1\rangle \otimes \cdots \otimes |i_n\rangle$ . Let U(d) denote the group of  $d \times d$  unitary operators. This group is naturally represented in our system by

$$\mathbf{Q}_{d,n}(\mathbf{U})|i_1\cdots i_n\rangle := \mathbf{U}|i_1\rangle \otimes \cdots \otimes \mathbf{U}|i_n\rangle, \tag{A.8}$$

where  $\boldsymbol{U} \in U(d)$ . In fact,  $\boldsymbol{Q}_{d,n}(\boldsymbol{U}) = \boldsymbol{U}^{\otimes n}$ , which is called the *collective action* of  $\boldsymbol{U} \in U(d)$ . Thus we have the following famous result:

**Theorem A.3** (Schur, [27]). Let  $\mathcal{A} = \operatorname{span}_{\mathbb{C}} \{ \mathbf{P}_{d,n}(\pi) : \pi \in S_k \}$  and  $\mathcal{B} = \operatorname{span}_{\mathbb{C}} \{ \mathbf{Q}_{d,n}(\mathbf{U}) : \mathbf{U} \in U(d) \}$ . *Then:* 

$$\mathcal{A}' = \mathcal{B} \quad and \quad \mathcal{A} = \mathcal{B}'.$$
 (A.9)

When treated as matrix algebras, such pairs (A, B) are known as *dual reductive pairs* since the collective action of the unitary group on the tensor space and the permutation action of tensor factors are mutual commutants.

In fact, the above dual theorem by Schur can be generalized. Consider the local unitary group  $U(d) = U(d_1) \otimes \cdots \otimes U(d_N)$ , where  $d = (d_1, \ldots, d_N)$  are positive integer dimensions, which is a subgroup of  $GL(d) = GL(d_1, \mathbb{C}) \otimes \cdots \otimes GL(d_N, \mathbb{C})$ . Let  $V_i$  be a  $d_i$ -dimensional complex Hilbert space and  $V = V_1 \otimes \cdots \otimes V_n$ . Then U(d) acts on the vector space  $End(V) = \bigotimes_{i=1}^N End(V_i)$ , where  $End(V_i)$  is the set of all endomorphisms from V to itself, by

$$M \mapsto gMg^{\dagger} \quad (g = U_1 \otimes \cdots \otimes U_N \in U(d), M \in End(V))$$
 (A.10)

which is obtained by linear extension of the action:  $\bigotimes_{i=1}^{N} X_i \mapsto \bigotimes_{i=1}^{N} U_i X_i U_i^{\dagger}$ , where  $X_i \in \text{End}(V_i)$ and  $U_i \in U(d_i)$ .

Consider the representation of U(d) on  $End(V^{\otimes n})$ , defined by

$$\mathbf{Q}_{d,n}(\boldsymbol{U}_1,\ldots,\boldsymbol{U}_N) \stackrel{\text{def}}{=} \mathbf{Q}_{d_1,n}(\boldsymbol{U}_1) \otimes \cdots \otimes \mathbf{Q}_{d_N,n}(\boldsymbol{U}_N), \tag{A.11}$$

where  $\mathbf{Q}_{d_i,n}(\mathbf{U}_i) = \mathbf{U}_i^{\otimes n}$  for  $\mathbf{U}_i \in U(d_i)$ . Denote the *N*-fold Cartesian product  $S_n^N := S_n \times \cdots \times S_n$  of the symmetric group  $S_n$  of order *n*. The action of  $S_n^N$  on  $End(V^{\otimes n})$  is defined by

$$\mathbf{P}_{d,n}(\pi_1,\ldots,\pi_N) \stackrel{\text{def}}{=} \mathbf{P}_{d_1,n}(\pi_1) \otimes \cdots \otimes \mathbf{P}_{d_N,n}(\pi_N), \tag{A.12}$$

where  $\mathbf{P}_{d_i,n}(\pi_i) \in \text{End}(V_i^{\otimes n})$  for  $\pi_i \in S_n$  with its definition taken from Eq. (A.7).

Theorem A.4 (The generalized Schur-Weyl duality, [?, ?]). Let

$$\widetilde{\mathcal{A}} := \operatorname{span}_{\mathbb{C}} \left\{ \mathbf{P}_{d,n}(\pi_1, \dots, \pi_N) : (\pi_1, \dots, \pi_N) \in S_n^N \right\},$$
(A.13)

$$\widetilde{\mathcal{B}} := \operatorname{span}_{\mathbb{C}} \left\{ \mathbf{Q}_{d,n}(\mathbf{U}_{1},\ldots,\mathbf{U}_{N}) : (\mathbf{U}_{1},\ldots,\mathbf{U}_{N}) \in \mathsf{U}(d) \right\}.$$
(A.14)

Then it holds that

$$\widetilde{\mathcal{A}}' = \widetilde{\mathcal{B}} \quad and \quad \widetilde{\mathcal{B}}' = \widetilde{\mathcal{A}}.$$
 (A.15)

#### A.3 Proof of Proposition 2.1

Let  $V = \text{Herm}(\mathbb{C}^{d_1} \otimes \cdots \otimes \mathbb{C}^{d_N})$ , the Hermitian matrices acting on the tensor space, and denote the local unitary group by  $U(d) = U(d_1) \otimes \cdots \otimes U(d_N)$ . Define U(d) acts on V by conjugation, i.e., for any  $g \in U(d)$  and  $X \in V$ , we get the LU action of U(d) on V via  $\tau_g X = gXg^{\dagger}$ . In fact, given two tuples of multipartite states on  $\mathbb{C}^{d_1} \otimes \cdots \otimes \mathbb{C}^{d_N}$ ,  $\Psi = (\rho_1, \dots, \rho_N)$  and  $\Psi' = (\rho'_1, \dots, \rho'_N)$ , they are locally unitarily (LU) equivalent in the sense that  $\rho'_i = g\rho_i g^{\dagger}$  for all  $i = 1, \dots, N$  and some  $g \in U(d)$ . That is, there exist a collection of unitary operators  $U_i \in U(d_i)$  such that  $g = U_1 \otimes \cdots \otimes U_N$ , where i = 1, ..., N, and

$$\rho'_i = (\boldsymbol{U}_1 \otimes \cdots \otimes \boldsymbol{U}_N) \rho_i (\boldsymbol{U}_1 \otimes \cdots \otimes \boldsymbol{U}_N)^{\dagger}$$
(A.16)

for each  $i = 1, \ldots, N$ .

Proof of Proposition 2.1. Clearly  $\rho \in V$ , then  $g = \mathbf{U}_1 \otimes \cdots \otimes \mathbf{U}_N \in U(d)$  acts on  $\rho$  via  $\tau_g \rho = g\rho g^{\dagger}$ . The space of all real polynomials on V is denoted by  $\mathbb{R}[V]$ . We will denote by  $\mathbb{R}[V]_n$  the space of real homogenous polynomials on V of degree n. We have already known that each homogeneous polynomials of degree n are mappings of the form  $p(v) = \langle \tilde{p}, v^{\otimes n} \rangle$ , where  $v \in V$  and  $\tilde{p} \in V^{\otimes n}$ . Thus  $\tilde{p}$  defines an U(d)-invariant polynomial  $p \in \mathbb{R}[V]_n^{U(d)}$  if and only if  $\tau_g^{\otimes n} \tilde{p} = \tilde{p}$  for all  $g \in U(d)$ . Now for both  $\rho$  and  $\sigma$  satisfying  $\sigma = g\rho g^{\dagger}$  for some  $g \in U(d)$  if and only if  $p(\rho) = p(\sigma)$  for  $\forall p \in \mathbb{R}[V]^{U(d)}$  [24]. By virtue of the above this is equivalent to demanding that for all  $g \in U(d)$ 

$$\tilde{p} = \tau_g^{\otimes n} \tilde{p} = \mathbf{Q}_{d,n}(\bar{g}) \tilde{p} \mathbf{Q}_{d,n}(\bar{g}),$$

where  $\mathbf{Q}_{d,n}(\bar{g}) = \mathbf{U}_1^{\otimes n} \otimes \cdots \otimes \mathbf{U}_N^{\otimes n}$  for  $\bar{g} := (\mathbf{U}_1, \dots, \mathbf{U}_N)$ . This amounts to requiring  $[\mathbf{Q}_{d,n}(\bar{g}), \tilde{p}] = 0$  implying that  $\tilde{p} \in \tilde{\mathcal{B}}' = \tilde{\mathcal{A}}$  by the generalized Schur-Weyl duality. Thus  $\tilde{p}$  can be expanded into a linear combination of  $\mathbf{P}_{d,n}(\pi)$ 's for  $\pi = (\pi_1, \dots, \pi_N) \in S_n^N$ , where  $\mathbf{P}_{d,n}(\pi) := \mathbf{P}_{d_1,n}(\pi_1) \otimes \cdots \otimes \mathbf{P}_{d_N,n}(\pi_N)$  for each permutation  $\pi_i \in S_n(i = 1, \dots, n)$ . Here, for  $d = d_1, \dots, d_N$ ,  $\mathbf{P}_{d,n}$  acting on  $(\mathbb{C}^d)^{\otimes n}$  via the action on computational basis vectors is defined by  $\mathbf{P}_{d,n}|i_1\cdots i_n\rangle := |i_{\pi^{-1}(1)}\cdots i_{\pi^{-1}(n)}\rangle$ . By the generalized Schur-Weyl duality, it is easily seen that  $\sigma = g\rho g^{\dagger}$  if and only if  $\operatorname{Tr}(\sigma^{\otimes n}\mathbf{P}_{d,n}(\pi)) = \operatorname{Tr}(\rho^{\otimes n}\mathbf{P}_{d,n}(\pi))$ , where  $n = 1, 2, \dots$  and  $\pi \in S_n^N$ . This completes the proof.

## **B Proof of Theorem 2.2**

In this section, we first establish an intriguing formula (Lemma B.5) concerning operator products. Subsequently, we reformulate the 18 Makhlin invariants  $I_k$ 's using 18 LU invariant generators, denoted as  $L_k$ 's (Proposition B.22). With these foundational steps completed, we can express all 18 Bargmann generators  $B_k$ 's as polynomials in terms of the 18 LU invariant generators  $L_k$ 's (see Lemma B.23). Building on this, we derive expressions for the  $L_k$ 's in terms of the  $B_k$ 's. Through the interrelationships between the  $L_k$ 's and  $B_k$ 's, we deduce that the set of 18 Bargmann invariants  $B_k$ 's constitutes a complete set that determines the local unitary equivalence of two-qubit states.

#### **B.1** Product formula for two-qubit observables

Let us fix some notations used in this section. Firstly, we recall the notion of the *cross product* in the real Euclidean space  $\mathbb{R}^3$ . We will make the convention by assuming that the cross product of two row(column) vectors will be a row(column) vector according to the definition of the cross product. For instance, for two column vectors  $\mathbf{x} = (x_1, x_2, x_3)^{\mathsf{T}}$  and  $\mathbf{y} = (y_1, y_2, y_3)^{\mathsf{T}}$  in  $\mathbb{R}^3$ , where  $\mathsf{T}$  means the transpose, their cross product  $\mathbf{x} \times \mathbf{y}$  is identified with

$$\mathbf{x} imes \mathbf{y} = \left( \left| \begin{array}{ccc} x_2 & x_3 \\ y_2 & y_3 \end{array} \right|, - \left| \begin{array}{ccc} x_1 & x_3 \\ y_1 & y_3 \end{array} \right|, \left| \begin{array}{ccc} x_1 & x_2 \\ y_1 & y_2 \end{array} \right| 
ight)^{\mathsf{T}}.$$

Moreover the cross product  $x^{\mathsf{T}} \times y^{\mathsf{T}}$  is identified with

$$\mathbf{x}^{\mathsf{T}} \times \mathbf{y}^{\mathsf{T}} = \left( \left| \begin{array}{ccc} x_2 & x_3 \\ y_2 & y_3 \end{array} \right|, - \left| \begin{array}{ccc} x_1 & x_3 \\ y_1 & y_3 \end{array} \right|, \left| \begin{array}{ccc} x_1 & x_2 \\ y_1 & y_2 \end{array} \right| \right).$$

According to this convention, we find that  $(x \times y)^{\mathsf{T}} = x^{\mathsf{T}} \times y^{\mathsf{T}}$ .

In what follows, we will use exchangeably the notation of column (row) vector  $x(x^{T})$  and the Dirac notation ket (bra)  $|x\rangle(\langle x|)$ . The inner products between two *real* 3-dimensional column vectors x and y and two *real* 3 × 3 matrices M and N, are defined by, respectively,

$$\langle x, y \rangle := x^{\mathsf{T}}y$$
 and  $\langle M, N \rangle := \operatorname{Tr}(M^{\mathsf{T}}N)$ 

where Tr stands for the usual matrix trace. We often write  $\langle x, My \rangle$  as  $\langle x|M|y \rangle$ . Denote  $|x| := \sqrt{\langle x, x \rangle}$  and  $||M|| := \sqrt{\langle M, M \rangle}$ .

We also use the notion of the cofactors [28] of entries in a matrix is defined as follows.

**Definition B.1.** For any (real or complex) square matrix  $M = (m_{ij})_{n \times n}$ , the so-called *cofactor* of entry  $m_{ij}$  is defined as the factor  $(-1)^{i+j}$  times the determinant of the  $(n-1) \times (n-1)$  matrix (denoted by  $M[\hat{i}|\hat{j}]$ ) obtained by deleting the *i*-th row and *j*-th column of M. That is, the cofactor of  $m_{ij}$  is

$$\widehat{m}_{ij} \stackrel{\text{def}}{=} (-1)^{i+j} \det\left(\boldsymbol{M}[\hat{i}|\hat{j}]\right). \tag{B.1}$$

Denote by  $\widehat{M} := (\widehat{m}_{ij})_{n \times n}$ , which is called the *cofactor matrix*. Then  $M^* \stackrel{\text{def}}{=} \widehat{M}^{\mathsf{T}}$  is called the *adjugate matrix* of M.

In Linear Algebra, for any two square matrices *M* and *N* of order *n*, it is well-known that

$$\widehat{M}^{\mathsf{T}} = (\widehat{M})^{\mathsf{T}}$$
 and  $\widehat{M}\widehat{N} = \widehat{M}\widehat{N}.$  (B.2)

Let the characteristic polynomial of the  $n \times n$  matrix M be  $f_n(\lambda)$ . Then

$$f_n(x) = \sum_{k=0}^n (-1)^k e_k(\mathbf{M}) x^{n-k},$$
(B.3)

where

$$\begin{cases} e_0(\boldsymbol{M}) &\equiv 1\\ e_1(\boldsymbol{M}) &= \operatorname{Tr}(\boldsymbol{M})\\ &\vdots \\ e_{n-1}(\boldsymbol{M}) &= \operatorname{Tr}\left(\widehat{\boldsymbol{M}}\right)\\ e_n(\boldsymbol{M}) &= \det(\boldsymbol{M}). \end{cases}$$
(B.4)

We can use Hamilton-Cayley theorem in Linear Algebra, together with the continuity argument, to give a formula towards the computation of adjugate matrix, which can be described as follows:

**Proposition B.2.** For any  $n \times n$  matrix M, its adjugate matrix can be determined by

$$M^* = \sum_{k=0}^{n-1} e_k(M) (-M)^{n-1-k}.$$
(B.5)

Proof. Indeed, This indicates by Hamilton-Cayley Theorem that

$$M^n - e_1(M)M^{n-1} + \dots + (-1)^{n-1}e_{n-1}(M)M + (-1)^n \det(M)\mathbb{1}_n = 0.$$

Thus

$$\left(M^{n-1} - e_1(M)M^{n-2} + (-1)^{n-1}e_{n-1}(M)\mathbb{1}_n\right)M = (-1)^{n-1}\det(M)\mathbb{1}_n = (-1)^{n-1}M^*M.$$

Then

$$M^* = (-M)^{n-1} + e_1(M)(-M)^{n-2} + \dots + e_k(M)(-M)^{n-1-k} + e_{n-1}(M)\mathbb{1}_n$$
  
=  $\sum_{k=0}^{n-1} e_k(M)(-M)^{n-1-k}$ 

holds true if M is invertible. By the continuity argument, this holds true for all square matrix M.

**Corollary B.3.** *For any square matrix*  $M \in \mathbb{R}^{3 \times 3}$ *, it holds that* 

(i) 
$$\operatorname{Tr}\left(\widehat{M}\right) = \frac{\operatorname{Tr}(M)^2 - \operatorname{Tr}(M^2)}{2};$$
  
(ii)  $\widehat{M}^{\mathsf{T}}\widehat{M} = (M^{\mathsf{T}}M)^2 - \langle M, M \rangle M^{\mathsf{T}}M + \langle \widehat{M}, \widehat{M} \rangle \mathbb{1}_3;$   
(iii)  $\langle \widehat{M}, \widehat{M} \rangle = \frac{1}{2} \left( \langle M, M \rangle^2 - \langle M^{\mathsf{T}}M, M^{\mathsf{T}}M \rangle \right);$ 

(iv)  $\widehat{M} = M^4 - c_2(M)M^2 + c_1(M)M + c_0(M)\mathbb{1}_3$ , where three coefficients  $c_k(M)(k = 0, 1, 2)$  are identified with

$$\begin{cases} c_0(M) &= \frac{-\operatorname{Tr}(M)^4 + 2\operatorname{Tr}(M)^2 \operatorname{Tr}(M^2) + \operatorname{Tr}(M^2)^2 - 2\operatorname{Tr}(M^4)}{8}, \\ c_1(M) &= \frac{\operatorname{Tr}(M) \left( \operatorname{Tr}(M)^2 - \operatorname{Tr}(M^2) \right)}{2}, \\ c_2(M) &= \frac{\operatorname{Tr}(M)^2 + \operatorname{Tr}(M^2)}{2}. \end{cases}$$
(B.6)

*Proof.* The proof is conceptually simple. We can also use MATHEMATICA to do this. In what follows, we give analytical reasoning. By Proposition B.2, we see that

$$M^* = \widehat{M}^{\mathsf{T}} = M^2 - \operatorname{Tr}(M) M + \operatorname{Tr}\left(\widehat{M}\right) \mathbb{1}_3.$$
(B.7)

(i) By taking the traces on both sides, we get that

$$\operatorname{Tr}\left(\widehat{M}\right) = rac{\operatorname{Tr}\left(M\right)^2 - \operatorname{Tr}\left(M^2\right)}{2}.$$

(ii) Now we use  $M^{\mathsf{T}}M$  to replace M in Eq. (B.7), then

$$\widehat{\boldsymbol{M}}^{\mathsf{T}}\widehat{\boldsymbol{M}} = \widehat{\boldsymbol{M}^{\mathsf{T}}\boldsymbol{M}} = (\boldsymbol{M}^{\mathsf{T}}\boldsymbol{M})^2 - \operatorname{Tr}(\boldsymbol{M}^{\mathsf{T}}\boldsymbol{M})\,\boldsymbol{M}^{\mathsf{T}}\boldsymbol{M} + \operatorname{Tr}\left(\widehat{\boldsymbol{M}^{\mathsf{T}}\boldsymbol{M}}\right)\mathbb{1}_3$$
$$= (\boldsymbol{M}^{\mathsf{T}}\boldsymbol{M})^2 - \langle \boldsymbol{M}, \boldsymbol{M} \rangle \boldsymbol{M}^{\mathsf{T}}\boldsymbol{M} + \langle \widehat{\boldsymbol{M}}, \widehat{\boldsymbol{M}} \rangle \mathbb{1}_3.$$

(iii) By taking the traces on both sides of the identity in (ii), after simplifying it, we get the desired result.

(iv) Apparently,

$$\widehat{\widehat{M}} = \left(\widehat{M}^2 - \operatorname{Tr}\left(\widehat{M}\right)\widehat{M} + \operatorname{Tr}\left(\widehat{\widehat{M}}\right)\mathbb{1}_3\right)^{\mathsf{T}} \\ = \left(\widehat{M}^2\right)^{\mathsf{T}} - \operatorname{Tr}\left(\widehat{M}\right)\widehat{M}^{\mathsf{T}} + \operatorname{Tr}\left(\widehat{\widehat{M}}\right)\mathbb{1}_3,$$

where

$$\left(\widehat{M}^2\right)^{\mathsf{T}} = \widehat{M}^2^{\mathsf{T}} = M^4 - \operatorname{Tr}\left(M^2\right)M^2 + \operatorname{Tr}\left(\widehat{M}^2\right)\mathbb{1}_3.$$

Thus substituting this into the expression of  $\widehat{\widehat{M}}$ , we get that

$$\begin{aligned} \widehat{\widehat{M}} &= \left( M^4 - \operatorname{Tr} \left( M^2 \right) M^2 + \operatorname{Tr} \left( \widehat{M^2} \right) \mathbb{1}_3 \right) \\ &- \operatorname{Tr} \left( \widehat{M} \right) \left( M^2 - \operatorname{Tr} \left( M \right) M + \operatorname{Tr} \left( \widehat{M} \right) \mathbb{1}_3 \right) + \operatorname{Tr} \left( \widehat{\widehat{M}} \right) \mathbb{1}_3 \\ &= M^4 - \left[ \operatorname{Tr} \left( M^2 \right) + \operatorname{Tr} \left( \widehat{M} \right) \right] M^2 + \operatorname{Tr} \left( M \right) \operatorname{Tr} \left( \widehat{M} \right) M \\ &+ \left[ \operatorname{Tr} \left( \widehat{M^2} \right) - \operatorname{Tr} \left( \widehat{M} \right)^2 + \operatorname{Tr} \left( \widehat{\widehat{M}} \right) \right] \mathbb{1}_3. \end{aligned}$$

Using many times the result obtained in (i), finally we obtain the desired identity.

#### **B.1.1** Product formula

As conventions, three Pauli matrices are given below:

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$
(B.8)

For any two-qubit *observable X*, we can decompose it as

$$X = t \mathbb{1}_4 + \mathbf{r} \cdot \boldsymbol{\sigma} \otimes \mathbb{1}_2 + \mathbb{1}_2 \otimes \mathbf{s} \cdot \boldsymbol{\sigma} + \sum_{i,j=1}^3 t_{ij} \sigma_i \otimes \sigma_j,$$
(B.9)

where  $t \in \mathbb{R}$ ,  $\mathbf{r} := (r_1, r_2, r_3)^{\mathsf{T}}$ ,  $\mathbf{s} := (s_1, s_2, s_3)^{\mathsf{T}} \in \mathbb{R}^3$ , and  $\mathbf{T} := (t_{ij})_{3\times 3} \in \mathbb{R}^{3\times 3}$ . Here  $\mathbf{r} \cdot \mathbf{\sigma} := \sum_{i=1}^3 r_i \sigma_i$ . By mimicking this notation, we introduce the following notation:  $\mathbf{F}_k = (\varepsilon_{ijk})_{3\times 3}$ , where  $\varepsilon_{ijk} := \operatorname{sign}[(j-i)(k-i)(k-j)]$  for  $i, j, k \in [3] := \{1, 2, 3\}$ . Indeed,

$$F_{1} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix}, \quad F_{2} = \begin{pmatrix} 0 & 0 & -1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \quad F_{3} = \begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$
(B.10)

Denote  $x \cdot \mathcal{F} := \sum_{k=1}^{3} x_k F_k$ , where  $\mathcal{F} := (F_1, F_2, F_3)$ . It is easily seen that the cross product can be realized as

$$\mathbf{x} \times \mathbf{y} = \left( \langle \mathbf{x} | F_1 | \mathbf{y} \rangle, \langle \mathbf{x} | F_2 | \mathbf{y} \rangle, \langle \mathbf{x} | F_3 | \mathbf{y} \rangle \right)^{\mathsf{T}}.$$
(B.11)

For convenience, we parameterize X in the notation (t, r, s, T) for X, denoted by  $X \approx (t, r, s, T)$ , and (t', r', s', T') for X', denoted by  $X' \approx (t', r', s', T')$ , respectively. Consider the product  $\tilde{X} := XX'$  with parameters  $(\tilde{t}, \tilde{r}, \tilde{s}, \tilde{T})$ .

In order to describe our product formula for  $\tilde{X}$ , we introduce the following notations: Denote

$$\Omega(M,N) := \begin{pmatrix} e_2^{\mathsf{T}}M \times e_3^{\mathsf{T}}N + e_2^{\mathsf{T}}N \times e_3^{\mathsf{T}}M \\ e_3^{\mathsf{T}}M \times e_1^{\mathsf{T}}N + e_3^{\mathsf{T}}N \times e_1^{\mathsf{T}}M \\ e_1^{\mathsf{T}}M \times e_2^{\mathsf{T}}N + e_1^{\mathsf{T}}N \times e_2^{\mathsf{T}}M \end{pmatrix},$$
(B.12)

where  $M, N \in \mathbb{R}^{3\times3}$  and  $\{e_1, e_2, e_3\}$  is the computational basis of  $\mathbb{R}^3$ , defined by  $e_1 = (1, 0, 0)^{\mathsf{T}}, e_2 = (0, 1, 0)^{\mathsf{T}}$ , and  $e_3 = (0, 0, 1)^{\mathsf{T}}$ . Clearly  $\Omega$  is symmetric bilinear mapping in the sense that  $\Omega(M, N) = \Omega(N, M)$ . Let

$$\Psi(\mathbf{x}, \mathbf{M}, \mathbf{y}) := (\mathbf{x} \cdot \mathcal{F})^{\mathsf{T}} \mathbf{M} + \mathbf{M}(\mathbf{y} \cdot \mathcal{F}), \tag{B.13}$$

where  $x, y \in \mathbb{R}^3$  and  $M \in \mathbb{R}^{3 \times 3}$ .

**Proposition B.4.** For the matrix  $\Omega(M, N)$ , its entries can be identified as

$$\Omega(\boldsymbol{M},\boldsymbol{N})_{p,q} = -\langle \boldsymbol{F}_p \boldsymbol{M} \boldsymbol{F}_q, \boldsymbol{N} \rangle \quad (\forall p,q \in \{1,2,3\}).$$
(B.14)

Moreover, it holds that

$$\Omega(\boldsymbol{M},\boldsymbol{N}) = \frac{1}{2} \sum_{i,j=1}^{3} |\boldsymbol{e}_i \times \boldsymbol{e}_j\rangle \langle \boldsymbol{e}_i^{\mathsf{T}} \boldsymbol{M} \times \boldsymbol{e}_j^{\mathsf{T}} \boldsymbol{N} + \boldsymbol{e}_i^{\mathsf{T}} \boldsymbol{N} \times \boldsymbol{e}_j^{\mathsf{T}} \boldsymbol{M}|.$$
(B.15)

*Proof.* For the first row of  $\Omega(M, N)$ , we find that

$$e_{2}^{\mathsf{T}}M \times e_{3}^{\mathsf{T}}N + e_{2}^{\mathsf{T}}N \times e_{3}^{\mathsf{T}}M = \left( \langle e_{2} | MF_{1}N^{\mathsf{T}} + NF_{1}M^{\mathsf{T}} | e_{3} \rangle, \langle e_{2} | MF_{2}N^{\mathsf{T}} + NF_{2}M^{\mathsf{T}} | e_{3} \rangle, \langle e_{2} | MF_{3}N^{\mathsf{T}} + NF_{3}M^{\mathsf{T}} | e_{3} \rangle \right).$$

Next, we determine such three components as follows. In fact,  $MF_jN^{\mathsf{T}} + NF_jM^{\mathsf{T}}$  is skew-Hermitian, and thus it can be decomposed as

$$MF_{j}N^{\mathsf{T}} + NF_{j}M^{\mathsf{T}} = c_{1}^{(j)}F_{1} + c_{2}^{(j)}F_{2} + c_{3}^{(j)}F_{3}.$$

This implies that

$$\operatorname{Tr}\left(F_{i}(\boldsymbol{M}\boldsymbol{F}_{j}\boldsymbol{N}^{\mathsf{T}}+\boldsymbol{N}\boldsymbol{F}_{j}\boldsymbol{M}^{\mathsf{T}})\right)=c_{1}^{(j)}\operatorname{Tr}\left(F_{i}F_{1}\right)+c_{2}^{(j)}\operatorname{Tr}\left(F_{i}F_{2}\right)+c_{3}^{(j)}\operatorname{Tr}\left(F_{i}F_{3}\right).$$

That is,

$$\operatorname{Tr}\left(F_{i}NF_{j}M^{\mathsf{T}}\right) = -c_{1}^{(j)}\delta_{1i} - c_{2}^{(j)}\delta_{2i} - c_{3}^{(j)}\delta_{3i} \Longrightarrow c_{i}^{(j)} = -\operatorname{Tr}\left((F_{i}MF_{j})^{\mathsf{T}}N\right) = -\langle F_{i}MF_{j},N\rangle.$$

From this observation, we get that  $\langle e_2 | MF_j N^{\mathsf{T}} + NF_j M^{\mathsf{T}} | e_3 \rangle = -\operatorname{Tr} (F_1 MF_j N^{\mathsf{T}})$ , which implies that

$$\boldsymbol{e}_{2}^{\mathsf{T}}\boldsymbol{M}\times\boldsymbol{e}_{3}^{\mathsf{T}}\boldsymbol{N}+\boldsymbol{e}_{2}^{\mathsf{T}}\boldsymbol{N}\times\boldsymbol{e}_{3}^{\mathsf{T}}\boldsymbol{M}=-\left(\left\langle \boldsymbol{F}_{1}\boldsymbol{M}\boldsymbol{F}_{1},\boldsymbol{N}\right\rangle ,\left\langle \boldsymbol{F}_{1}\boldsymbol{M}\boldsymbol{F}_{2},\boldsymbol{N}\right\rangle ,\left\langle \boldsymbol{F}_{1}\boldsymbol{M}\boldsymbol{F}_{3},\boldsymbol{N}\right\rangle \right)$$

Similar procedures for 2nd and 3rd rows are performed, respectively, and thus we get the desired result:  $\Omega(M, N)_{p,q} = -\langle F_p M F_q, N \rangle$ . The second item can be checked as follows: Clearly i = j,  $|e_i \times e_j\rangle\langle e_i^{\mathsf{T}}M \times e_j^{\mathsf{T}}N + e_i^{\mathsf{T}}N \times e_j^{\mathsf{T}}M| = 0$  due to the fact that  $e_i \times e_j = 0$  if i = j. Besides, for  $i \neq j$ ,

$$|e_i \times e_j\rangle\langle e_i^{\mathsf{T}} M \times e_j^{\mathsf{T}} N + e_i^{\mathsf{T}} N \times e_j^{\mathsf{T}} M| = |e_j \times e_i\rangle\langle e_j^{\mathsf{T}} M \times e_i^{\mathsf{T}} N + e_j^{\mathsf{T}} N \times e_i^{\mathsf{T}} M|.$$

It suffices to consider (i, j) = (1, 2), (1, 3), (2, 3). Note that  $e_1 \times e_2 = e_3, e_2 \times e_3 = e_1$ , and  $e_3 \times e_1 = e_2$ . Thus we get that

$$\begin{aligned} \frac{1}{2} \sum_{i,j=1}^{3} |e_i \times e_j\rangle \langle e_i^{\mathsf{T}} M \times e_j^{\mathsf{T}} N + e_i^{\mathsf{T}} N \times e_j^{\mathsf{T}} M| &= \sum_{1 \leq i < j \leq 3} |e_i \times e_j\rangle \langle e_i^{\mathsf{T}} M \times e_j^{\mathsf{T}} N + e_i^{\mathsf{T}} N \times e_j^{\mathsf{T}} M| \\ &= |e_1 \times e_2\rangle \langle e_1^{\mathsf{T}} M \times e_2^{\mathsf{T}} N + e_1^{\mathsf{T}} N \times e_2^{\mathsf{T}} M| + |e_1 \times e_3\rangle \langle e_1^{\mathsf{T}} M \times e_3^{\mathsf{T}} N + e_1^{\mathsf{T}} N \times e_3^{\mathsf{T}} M| \\ &+ |e_2 \times e_3\rangle \langle e_2^{\mathsf{T}} M \times e_3^{\mathsf{T}} N + e_2^{\mathsf{T}} N \times e_3^{\mathsf{T}} M| \\ &= |e_3\rangle \langle e_1^{\mathsf{T}} M \times e_2^{\mathsf{T}} N + e_1^{\mathsf{T}} N \times e_2^{\mathsf{T}} M| + |e_2\rangle \langle e_3^{\mathsf{T}} M \times e_1^{\mathsf{T}} N + e_3^{\mathsf{T}} N \times e_1^{\mathsf{T}} M| \\ &+ |e_1\rangle \langle e_2^{\mathsf{T}} M \times e_3^{\mathsf{T}} N + e_2^{\mathsf{T}} N \times e_3^{\mathsf{T}} M|, \end{aligned}$$

which implies the desired result when witting is as a matrix form.

We have the following formula for the product  $\tilde{X} = XX'$  of X and X'.

**Lemma B.5** (Product formula of two-qubit observables). If  $X \approx (t, r, s, T)$  and  $X' \approx (t', r', s', T')$ , then  $\tilde{X} \approx (\tilde{t}, \tilde{r}, \tilde{s}, \tilde{T})$  is given by the following formulae:

$$\begin{cases} \tilde{t} = tt' + \langle \boldsymbol{r}, \boldsymbol{r}' \rangle + \langle \boldsymbol{s}, \boldsymbol{s}' \rangle + \langle \boldsymbol{T}, \boldsymbol{T}' \rangle, \\ \tilde{r} = t'\boldsymbol{r} + t\boldsymbol{r}' + \boldsymbol{T}'\boldsymbol{s} + \boldsymbol{T}\boldsymbol{s}' + i\left(\boldsymbol{r} \times \boldsymbol{r}' + \sum_{i=1}^{3} \boldsymbol{T}\boldsymbol{e}_{i} \times \boldsymbol{T}'\boldsymbol{e}_{i}\right), \\ \tilde{s} = t'\boldsymbol{s} + t\boldsymbol{s}' + {\boldsymbol{T}'}^{\mathsf{T}}\boldsymbol{r} + {\boldsymbol{T}}^{\mathsf{T}}\boldsymbol{r}' + i\left(\boldsymbol{s} \times \boldsymbol{s}' + \sum_{i=1}^{3} {\boldsymbol{T}}^{\mathsf{T}}\boldsymbol{e}_{i} \times {\boldsymbol{T}'}^{\mathsf{T}}\boldsymbol{e}_{i}\right), \\ \tilde{\boldsymbol{T}} = t'\boldsymbol{T} + t\boldsymbol{T}' + |\boldsymbol{r}\rangle\langle\boldsymbol{s}'| + |\boldsymbol{r}'\rangle\langle\boldsymbol{s}| - \Omega(\boldsymbol{T}, \boldsymbol{T}') + i\left(\Psi(\boldsymbol{r}, \boldsymbol{T}', \boldsymbol{s}) - \Psi(\boldsymbol{r}', \boldsymbol{T}, \boldsymbol{s}')\right). \end{cases}$$
(B.16)

Moreover,  $\operatorname{Tr}(XX') = 4(tt' + \langle \mathbf{r}, \mathbf{r}' \rangle + \langle \mathbf{s}, \mathbf{s}' \rangle + \langle \mathbf{T}, \mathbf{T}' \rangle).$ 

Proof. The proof is conceptually, but needs tedious algebraic computations. Indeed,

$$\begin{split} \tilde{t} &= \frac{1}{4} \operatorname{Tr} \left( \tilde{X} \right) = \frac{1}{4} \operatorname{Tr} \left( X X' \right), \\ \tilde{r}_i &= \frac{1}{4} \operatorname{Tr} \left( X X' (\sigma_i \otimes \mathbb{1}_2) \right), \\ \tilde{s}_j &= \frac{1}{4} \operatorname{Tr} \left( X X' (\mathbb{1}_2 \otimes \sigma_j) \right), \\ \tilde{t}_{ij} &= \frac{1}{4} \operatorname{Tr} \left( X X' (\sigma_i \otimes \sigma_j) \right). \end{split}$$

The next step is to check the correctness of the desired formula. This can be done by using the symbolic computation of the mathematical software MATHEMATICA. Assume that  $X \approx (t, r, s, T)$  and  $X' \approx (t', r', s', T')$ . Then

$$\begin{split} \mathbf{X}\mathbf{X}' &= \left(t\mathbb{1}_4 + \mathbf{r}\cdot\boldsymbol{\sigma}\otimes\mathbb{1}_2 + \mathbb{1}_2\otimes\mathbf{s}\cdot\boldsymbol{\sigma} + \sum_{i,j=1}^3 t_{ij}\sigma_i\otimes\sigma_j\right) \\ &\times \left(t'\mathbb{1}_4 + \mathbf{r}'\cdot\boldsymbol{\sigma}\otimes\mathbb{1}_2 + \mathbb{1}_2\otimes\mathbf{s}'\cdot\boldsymbol{\sigma} + \sum_{i,j=1}^3 t'_{ij}\sigma_i\otimes\sigma_j\right) \\ &= \left(tt'\mathbb{1}_4 + t\mathbf{r}'\cdot\boldsymbol{\sigma}\otimes\mathbb{1}_2 + \mathbb{1}_2\otimes t\mathbf{s}'\cdot\boldsymbol{\sigma} + \sum_{i,j=1}^3 tt'_{ij}\sigma_i\otimes\sigma_j\right) \\ &+ \left(t'\mathbf{r}\cdot\boldsymbol{\sigma}\otimes\mathbb{1}_2 + (\mathbf{r}\cdot\boldsymbol{\sigma})(\mathbf{r}'\cdot\boldsymbol{\sigma})\otimes\mathbb{1}_2 + \mathbf{r}\cdot\boldsymbol{\sigma}\otimes\mathbf{s}'\cdot\boldsymbol{\sigma} + \sum_{i,j=1}^3 t'_{ij}\sigma_i\otimes(\mathbf{s}\cdot\boldsymbol{\sigma})\sigma_j\right) \\ &+ \left(\mathbb{1}_2\otimes t'\mathbf{s}\cdot\boldsymbol{\sigma} + \mathbf{r}'\cdot\boldsymbol{\sigma}\otimes\mathbf{s}\cdot\boldsymbol{\sigma} + \mathbb{1}_2\otimes(\mathbf{s}\cdot\boldsymbol{\sigma})(\mathbf{s}'\cdot\boldsymbol{\sigma}) + \sum_{i,j=1}^3 t'_{ij}\sigma_i\otimes(\mathbf{s}\cdot\boldsymbol{\sigma})\sigma_j\right) \\ &+ \left(t'\sum_{i,j=1}^3 t_{ij}\sigma_i\otimes\sigma_j + \sum_{i,j=1}^3 t_{ij}\sigma_i(\mathbf{r}'\cdot\boldsymbol{\sigma})\otimes\sigma_j + \sum_{i,j=1}^3 t_{ij}\sigma_i\otimes\sigma_j(\mathbf{s}'\cdot\boldsymbol{\sigma})\right) \\ &+ \left(\sum_{i,j=1}^3 t_{ij}\sigma_i\otimes\sigma_j\right) \left(\sum_{i,j=1}^3 t'_{ij}\sigma_i\otimes\sigma_j\right). \end{split}$$

Furthermore

$$\begin{split} XX' &= tt'\mathbb{1}_4 + (tr' + t'r) \cdot \sigma \otimes \mathbb{1}_2 + \mathbb{1}_2 \otimes (ts' + t's) \cdot \sigma + \sum_{i,j=1}^3 (tt'_{ij} + t't_{ij})\sigma_i \otimes \sigma_j \\ &+ \left( (r \cdot \sigma)(r' \cdot \sigma) \otimes \mathbb{1}_2 + r \cdot \sigma \otimes s' \cdot \sigma + \sum_{i,j=1}^3 t'_{ij}(r \cdot \sigma)\sigma_i \otimes \sigma_j + \sum_{i,j=1}^3 t_{ij}\sigma_i(r' \cdot \sigma) \otimes \sigma_j \right) \\ &+ \left( r' \cdot \sigma \otimes s \cdot \sigma + \mathbb{1}_2 \otimes (s \cdot \sigma)(s' \cdot \sigma) + \sum_{i,j=1}^3 t'_{ij}\sigma_i \otimes (s \cdot \sigma)\sigma_j + \sum_{i,j=1}^3 t_{ij}\sigma_i \otimes \sigma_j(s' \cdot \sigma) \right) \\ &+ \left( \sum_{i,j=1}^3 t_{ij}\sigma_i \otimes \sigma_j \right) \left( \sum_{i,j=1}^3 t'_{ij}\sigma_i \otimes \sigma_j \right). \end{split}$$

Note that  $(\mathbf{r} \cdot \boldsymbol{\sigma})(\mathbf{r}' \cdot \boldsymbol{\sigma}) \otimes \mathbb{1}_2 = \langle \mathbf{r}, \mathbf{r}' \rangle \mathbb{1}_4 + i(\mathbf{r} \times \mathbf{r}') \cdot \boldsymbol{\sigma} \otimes \mathbb{1}_2$  and  $\mathbb{1}_2 \otimes (\mathbf{s} \cdot \boldsymbol{\sigma})(\mathbf{s}' \cdot \boldsymbol{\sigma}) = \langle \mathbf{s}, \mathbf{s}' \rangle \mathbb{1}_4 + \mathbb{1}_2 \otimes i(\mathbf{s} \times \mathbf{s}') \cdot \boldsymbol{\sigma}$ . Then we see that

$$\begin{split} \mathbf{X}\mathbf{X}' &= (tt' + \langle \mathbf{r}, \mathbf{r}' \rangle + \langle \mathbf{s}, \mathbf{s}' \rangle) \mathbb{1}_4 + (t\mathbf{r}' + t'\mathbf{r} + \mathbf{i}\mathbf{r} \times \mathbf{r}') \cdot \boldsymbol{\sigma} \otimes \mathbb{1}_2 \\ &+ \mathbb{1}_2 \otimes (t\mathbf{s}' + t'\mathbf{s} + \mathbf{i}\mathbf{s} \times \mathbf{s}') \cdot \boldsymbol{\sigma} + \sum_{i,j=1}^3 (tT' + t'T)_{ij}\sigma_i \otimes \sigma_j \\ &+ \left(\mathbf{r} \cdot \boldsymbol{\sigma} \otimes \mathbf{s}' \cdot \boldsymbol{\sigma} + \sum_{i,j=1}^3 t'_{ij}(\mathbf{r} \cdot \boldsymbol{\sigma})\sigma_i \otimes \sigma_j + \sum_{i,j=1}^3 t_{ij}\sigma_i(\mathbf{r}' \cdot \boldsymbol{\sigma}) \otimes \sigma_j\right) \\ &+ \left(\mathbf{r}' \cdot \boldsymbol{\sigma} \otimes \mathbf{s} \cdot \boldsymbol{\sigma} + \sum_{i,j=1}^3 t'_{ij}\sigma_i \otimes (\mathbf{s} \cdot \boldsymbol{\sigma})\sigma_j + \sum_{i,j=1}^3 t_{ij}\sigma_i \otimes \sigma_j(\mathbf{s}' \cdot \boldsymbol{\sigma})\right) \\ &+ \left(\sum_{i,j=1}^3 t_{ij}\sigma_i \otimes \sigma_j\right) \left(\sum_{i,j=1}^3 t'_{ij}\sigma_i \otimes \sigma_j\right). \end{split}$$

Now we use the fact that  $\sigma_i \sigma_j = i \sum_{k=1}^3 \varepsilon_{ijk} \sigma_k + \delta_{ij} \mathbb{1}_2$  and get that

$$\mathbf{r} \cdot \boldsymbol{\sigma} \otimes \mathbf{s}' \cdot \boldsymbol{\sigma} = \sum_{i,j=1}^{3} (|\mathbf{r}\rangle \langle \mathbf{s}'|)_{ij} \sigma_i \otimes \sigma_j,$$
  
$$\mathbf{r}' \cdot \boldsymbol{\sigma} \otimes \mathbf{s} \cdot \boldsymbol{\sigma} = \sum_{i,j=1}^{3} (|\mathbf{r}'\rangle \langle \mathbf{s}|)_{ij} \sigma_i \otimes \sigma_j.$$

We also have

$$\begin{split} &\sum_{i,j=1}^{3} t'_{ij}(\mathbf{r}\cdot\boldsymbol{\sigma})\sigma_{i}\otimes\sigma_{j} = \sum_{k=1}^{3}\sum_{i,j=1}^{3} t'_{ij}r_{k}\sigma_{k}\sigma_{i}\otimes\sigma_{j} = \sum_{i,j,k=1}^{3} t'_{ij}r_{k}\left(\mathrm{i}\sum_{j'=1}^{3}\varepsilon_{kij'}\sigma_{j'}+\delta_{ki}\mathbb{1}_{2}\right)\otimes\sigma_{j} \\ &=\mathrm{i}\sum_{j',j=1}^{3}\left(\sum_{k,i=1}^{3} t'_{ij}\varepsilon_{kij'}r_{k}\right)\sigma_{j'}\otimes\sigma_{j}+\mathbb{1}_{2}\otimes\sum_{i,j,k=1}^{3}\left(t'_{ij}\delta_{ki}r_{k}\right)\sigma_{j} \\ &=\mathrm{i}\sum_{j',j=1}^{3}\left(\left(\mathbf{r}\cdot\mathcal{F}\right)^{\mathsf{T}}\mathbf{T}'\right)_{j'j}\sigma_{j'}\otimes\sigma_{j}+\mathbb{1}_{2}\otimes\left(\mathbf{T}'^{\mathsf{T}}\mathbf{r}\right)\cdot\boldsymbol{\sigma}=\mathrm{i}\sum_{i,j=1}^{3}\left(\left(\mathbf{r}\cdot\mathcal{F}\right)^{\mathsf{T}}\mathbf{T}'\right)_{ij}\sigma_{i}\otimes\sigma_{j}+\mathbb{1}_{2}\otimes\left(\mathbf{T}'^{\mathsf{T}}\mathbf{r}\right)\cdot\boldsymbol{\sigma} \end{split}$$

and

$$\begin{split} &\sum_{i,j=1}^{3} t_{ij} \sigma_i (\mathbf{r}' \cdot \boldsymbol{\sigma}) \otimes \sigma_j = \sum_{k=1}^{3} \sum_{i,j=1}^{3} t_{ij} r'_k \sigma_i \sigma_k \otimes \sigma_j = \sum_{i,j,k=1}^{3} t_{ij} r'_k \left( i \sum_{j'=1}^{3} \varepsilon_{ikj'} \sigma_{j'} + \delta_{ik} \mathbb{1}_2 \right) \otimes \sigma_j \\ &= -i \sum_{j',j=1}^{3} \left( \sum_{i,k=1}^{3} t_{ij} \varepsilon_{ij'k} r'_k \right) \sigma_{j'} \otimes \sigma_j + \mathbb{1}_2 \otimes \sum_{i,j,k=1}^{3} \left( t_{ij} \delta_{ik} r'_k \right) \sigma_j \\ &= -i \sum_{j',j=1}^{3} \left( (\mathbf{r}' \cdot \mathcal{F})^{\mathsf{T}} \mathbf{T} \right)_{j'j} \sigma_{j'} \otimes \sigma_j + \mathbb{1}_2 \otimes (\mathbf{T}^{\mathsf{T}} \mathbf{r}) \cdot \boldsymbol{\sigma} = -i \sum_{i,j=1}^{3} \left( (\mathbf{r}' \cdot \mathcal{F})^{\mathsf{T}} \mathbf{T} \right)_{ij} \sigma_i \otimes \sigma_j + \mathbb{1}_2 \otimes (\mathbf{T}^{\mathsf{T}} \mathbf{r}') \cdot \boldsymbol{\sigma}. \end{split}$$

Similarly, we get that

$$\sum_{i,j=1}^{3} t'_{ij} \sigma_i \otimes (s \cdot \sigma) \sigma_j = \mathrm{i} \sum_{i,j=1}^{3} \left( T'(s \cdot \mathcal{F}) \right)_{ij} \sigma_i \otimes \sigma_j + (T's) \cdot \sigma \otimes \mathbb{1}_2,$$
$$\sum_{i,j=1}^{3} t_{ij} \sigma_i \otimes \sigma_j (s' \cdot \sigma) = -\mathrm{i} \sum_{i,j=1}^{3} \left( T(s' \cdot \mathcal{F}) \right)_{ij} \sigma_i \otimes \sigma_j + (Ts') \cdot \sigma \otimes \mathbb{1}_2.$$

At last,

$$\begin{split} &\left(\sum_{i,j=1}^{3} t_{ij}\sigma_{i}\otimes\sigma_{j}\right)\left(\sum_{i,j=1}^{3} t_{ij}'\sigma_{i}\otimes\sigma_{j}\right)=\sum_{i,j,k,l=1}^{3} t_{ij}t_{kl}'\sigma_{i}\sigma_{k}\otimes\sigma_{j}\sigma_{l}\\ &=\sum_{i,j,k,l=1}^{3} t_{ij}t_{kl}'\left(\mathrm{i}\sum_{p=1}^{3} \varepsilon_{ikp}\sigma_{p}+\delta_{ik}\mathbb{1}_{2}\right)\otimes\left(\mathrm{i}\sum_{q=1}^{3} \varepsilon_{jlq}\sigma_{q}+\delta_{jl}\mathbb{1}_{2}\right)\\ &=-\sum_{i,j,k,l,p,q=1}^{3} t_{ij}t_{kl}'\varepsilon_{ikp}\varepsilon_{jlq}\sigma_{p}\otimes\sigma_{q}+\mathrm{i}\sum_{i,j,k,l,p=1}^{3} t_{ij}t_{kl}'\varepsilon_{ikp}\delta_{jl}\sigma_{p}\otimes\mathbb{1}_{2}+\mathrm{i}\sum_{i,j,k,l,q=1}^{3} t_{ij}t_{kl}'\varepsilon_{jlq}\delta_{ik}\mathbb{1}_{2}\otimes\sigma_{q}\\ &+\sum_{i,j,k,l=1}^{3} t_{ij}t_{kl}'\delta_{ik}\delta_{jl}\mathbb{1}_{2}\otimes\mathbb{1}_{2}\\ &=\sum_{p,q=1}^{3}\left(\sum_{i,j,k,l=1}^{3} t_{ij}t_{kl}'\varepsilon_{ikp}\varepsilon_{ljq}\right)\sigma_{p}\otimes\sigma_{q}+\mathrm{i}\sum_{p=1}^{3}\left(\sum_{i,j,k,l=1}^{3} t_{ij}t_{kl}'\varepsilon_{ikp}\delta_{jl}\right)\sigma_{p}\otimes\mathbb{1}_{2}\\ &+\mathrm{i}\mathbb{1}_{2}\otimes\sum_{q=1}^{3}\left(\sum_{i,j,k,l=1}^{3} t_{ij}t_{kl}'\varepsilon_{jlq}\delta_{ik}\right)\sigma_{q}+\langle T,T'\rangle\mathbb{1}_{4}\\ &=-\sum_{p,q=1}^{3}\Omega(T,T')_{p,q}\sigma_{p}\otimes\sigma_{q}+\mathrm{i}\left(\sum_{i=1}^{3} Te_{i}\times T'e_{i}\right)\cdot\sigma\otimes\mathbb{1}_{2}+\mathrm{i}\mathbb{1}_{2}\otimes\left(\sum_{i=1}^{3} T^{\mathsf{T}}e_{i}\times T'^{\mathsf{T}}e_{i}\right)\cdot\sigma\\ &+\langle T,T'\rangle\mathbb{1}_{4},\end{split}$$

where we used the facts that

(1) 
$$\Omega(\mathbf{T}, \mathbf{T}')_{p,q} = -\langle \mathbf{F}_p \mathbf{T} \mathbf{F}_q, \mathbf{T}' \rangle;$$
  
(2)  $(\sum_{j=1}^{3} \mathbf{T} \mathbf{e}_j \times \mathbf{T}' \mathbf{e}_j) \cdot \boldsymbol{\sigma} = \sum_{j=1}^{3} \left( \sum_{i,k,p=1}^{3} (\mathbf{T} \mathbf{e}_j)_i (\mathbf{T}' \mathbf{e}_j)_k \varepsilon_{ikp} \sigma_p \right) = \sum_{j=1}^{3} \left( \sum_{i,k,p=1}^{3} (\mathbf{T} \mathbf{e}_j)_i (\mathbf{T}' \mathbf{e}_l)_k \varepsilon_{ikp} \delta_{jl} \sigma_p \right) = \sum_{i,j,k,l,p=1}^{3} t_{ij} t'_{kl} \varepsilon_{ikp} \delta_{jl} \sigma_p;$ 

(3) 
$$\left(\sum_{i=1}^{3} T^{\mathsf{T}} \boldsymbol{e}_{i} \times T'^{\mathsf{T}} \boldsymbol{e}_{i}\right) \cdot \boldsymbol{\sigma} = \sum_{i,j,k,l,q=1}^{3} t_{ij} t'_{kl} \varepsilon_{jlq} \delta_{ik} \sigma_{q}.$$

We are done.

The advantage of this product formula for two-qubit observables lies in its independence from the components of vectors (or matrix entries).

**Corollary B.6.** The commutator [X, X'] := XX' - X'X is identified as

$$[\mathbf{X}, \mathbf{X}'] = 2\mathbf{i} \Big[ \Big( \mathbf{r} \times \mathbf{r}' + \sum_{i=1}^{3} T \mathbf{e}_i \times T' \mathbf{e}_i \Big) \cdot \sigma \otimes \mathbb{1}_2 + \mathbb{1}_2 \otimes \Big( \mathbf{s} \times \mathbf{s}' + \sum_{i=1}^{3} T^{\mathsf{T}} \mathbf{e}_i \times T'^{\mathsf{T}} \mathbf{e}_i \Big) \cdot \sigma \\ + \sum_{i,j=1}^{3} \Big( \Psi(\mathbf{r}, \mathbf{T}', \mathbf{s}) - \Psi(\mathbf{r}', \mathbf{T}, \mathbf{s}') \Big)_{ij} \sigma_i \otimes \sigma_j \Big].$$
(B.17)

*Moreover*  $[\mathbf{X}, \mathbf{X}'] = 0$  *if and only if* 

$$\begin{cases} \mathbf{r} \times \mathbf{r}' + \sum_{i=1}^{3} T \mathbf{e}_i \times T' \mathbf{e}_i &= \mathbf{0}, \\ \mathbf{s} \times \mathbf{s}' + \sum_{i=1}^{3} T^{\mathsf{T}} \mathbf{e}_i \times T'^{\mathsf{T}} \mathbf{e}_i &= \mathbf{0}, \\ \Psi(\mathbf{r}, T', \mathbf{s}) &= \Psi(\mathbf{r}', T, \mathbf{s}'). \end{cases}$$
(B.18)

Proposition B.7. It holds that

$$\sum_{i=1}^{3} A \boldsymbol{e}_i \times \boldsymbol{B} \boldsymbol{e}_i = \sum_{i=1}^{3} (A \boldsymbol{B}^{\mathsf{T}} \boldsymbol{e}_i) \times \boldsymbol{e}_i = \sum_{i=1}^{3} \boldsymbol{e}_i \times (\boldsymbol{B} \boldsymbol{A}^{\mathsf{T}} \boldsymbol{e}_i),$$
(B.19)

where  $A, B \in \mathbb{R}^{3 \times 3}$ .

Proof. Indeed,

$$\sum_{i=1}^{3} Ae_i \times Be_i = \sum_{i=1}^{3} Ae_i \times \sum_{j=1}^{3} |e_j\rangle \langle e_j | Be_i = \sum_{j=1}^{3} \sum_{i=1}^{3} Ae_i \langle e_j | B | e_i \rangle \times e_j$$
$$= \sum_{j=1}^{3} \sum_{i=1}^{3} A |e_i\rangle \langle e_i | B^{\mathsf{T}} | e_j \rangle \times e_j = \sum_{j=1}^{3} A \sum_{i=1}^{3} |e_i\rangle \langle e_i | B^{\mathsf{T}} | e_j \rangle \times e_j$$
$$= \sum_{j=1}^{3} AB^{\mathsf{T}} e_j \times e_j,$$

completing the proof.

**Corollary B.8.** *For*  $A, B \in \mathbb{R}^{3 \times 3}$ *, we have* 

$$\sum_{k=1}^{3} A\boldsymbol{e}_{k} \times \boldsymbol{B}\boldsymbol{e}_{k} = -\frac{1}{2} \operatorname{Tr} \left( \mathcal{F} (\boldsymbol{A}\boldsymbol{B}^{\mathsf{T}} - \boldsymbol{B}\boldsymbol{A}^{\mathsf{T}}) \right) = -\frac{1}{2} \sum_{k=1}^{3} \operatorname{Tr} \left( \boldsymbol{F}_{k} (\boldsymbol{A}\boldsymbol{B}^{\mathsf{T}} - \boldsymbol{B}\boldsymbol{A}^{\mathsf{T}}) \right) \boldsymbol{e}_{k}, \tag{B.20}$$

where

$$\operatorname{Tr}\left(\mathcal{F}(AB^{\mathsf{T}}-BA^{\mathsf{T}})\right) := \left(\operatorname{Tr}\left(F_{1}(AB^{\mathsf{T}}-BA^{\mathsf{T}})\right), \operatorname{Tr}\left(F_{2}(AB^{\mathsf{T}}-BA^{\mathsf{T}})\right), \operatorname{Tr}\left(F_{3}(AB^{\mathsf{T}}-BA^{\mathsf{T}})\right)\right).$$

*Proof.* Indeed, by Proposition B.7, we get that

$$\sum_{k=1}^{3} A \boldsymbol{e}_{k} \times B \boldsymbol{e}_{k} = \sum_{i=1}^{3} (A B^{\mathsf{T}} \boldsymbol{e}_{i}) \times \boldsymbol{e}_{i} = \sum_{i=1}^{3} \boldsymbol{e}_{i} \times (B A^{\mathsf{T}} \boldsymbol{e}_{i}) = -\sum_{i=1}^{3} (B A^{\mathsf{T}} \boldsymbol{e}_{i}) \times \boldsymbol{e}_{i},$$

implying that

$$\begin{split} &\sum_{k=1}^{3} A\boldsymbol{e}_{k} \times B\boldsymbol{e}_{k} = -\frac{1}{2} \sum_{i=1}^{3} \boldsymbol{e}_{i} \times \left[ (AB^{\mathsf{T}} - BA^{\mathsf{T}}) \boldsymbol{e}_{i} \right] \\ &= -\frac{1}{2} \left( \sum_{i=1}^{3} \left\langle \boldsymbol{e}_{i} \middle| F_{1} (AB^{\mathsf{T}} - BA^{\mathsf{T}}) \middle| \boldsymbol{e}_{i} \right\rangle, \sum_{i=1}^{3} \left\langle \boldsymbol{e}_{i} \middle| F_{2} (AB^{\mathsf{T}} - BA^{\mathsf{T}}) \middle| \boldsymbol{e}_{i} \right\rangle, \sum_{i=1}^{3} \left\langle \boldsymbol{e}_{i} \middle| F_{3} (AB^{\mathsf{T}} - BA^{\mathsf{T}}) \middle| \boldsymbol{e}_{i} \right\rangle \\ &= -\frac{1}{2} \left( \operatorname{Tr} \left( F_{1} (AB^{\mathsf{T}} - BA^{\mathsf{T}}) \right), \operatorname{Tr} \left( F_{2} (AB^{\mathsf{T}} - BA^{\mathsf{T}}) \right), \operatorname{Tr} \left( F_{3} (AB^{\mathsf{T}} - BA^{\mathsf{T}}) \right) \right). \end{split}$$

This can be written down in a simplified notation:

$$\operatorname{Tr}\left(\mathcal{F}(AB^{\mathsf{T}} - BA^{\mathsf{T}})\right) := \left(\operatorname{Tr}\left(F_{1}(AB^{\mathsf{T}} - BA^{\mathsf{T}})\right), \operatorname{Tr}\left(F_{2}(AB^{\mathsf{T}} - BA^{\mathsf{T}})\right), \operatorname{Tr}\left(F_{3}(AB^{\mathsf{T}} - BA^{\mathsf{T}})\right)\right).$$
  
e are done.

We

#### **B.1.2** Auxiliary results

To establish a rigorous relationship between Makhlin's invariants and the Bargmann invariants under local unitary (LU) transformations, we need to perform detailed calculations. Throughout this process, numerous intriguing insights and findings will emerge, which can be immediately utilized for simplifications and reductions.

**Lemma B.9.** For two given vectors  $\mathbf{x} = (x_1, x_2, x_3)^{\mathsf{T}}, \mathbf{y} = (y_1, y_2, y_3)^{\mathsf{T}} \in \mathbb{R}^3$ , it holds that

(i)  $(\mathbf{x} \cdot \mathcal{F})^{\mathsf{T}} = -\mathbf{x} \cdot \mathcal{F};$ 

(*ii*) 
$$(\mathbf{x} \cdot \mathcal{F})^{\mathsf{T}} \mathbf{y} = \mathbf{x} \times \mathbf{y};$$

(iii) 
$$\mathbf{x} \cdot \mathcal{F} = \sum_{j=1}^{3} |\mathbf{e}_{j} \times \mathbf{x}\rangle \langle \mathbf{e}_{j}| = \sum_{j=1}^{3} |\mathbf{e}_{j}\rangle \langle \mathbf{x} \times \mathbf{e}_{j}|.$$
  
(iv)  $(\mathbf{x} \cdot \mathcal{F})^{\mathsf{T}}(\mathbf{y} \cdot \mathcal{F}) = \sum_{j=1}^{3} \mathbf{F}_{j} |\mathbf{x}\rangle \langle \mathbf{y}| \mathbf{F}_{j}^{\mathsf{T}} = \langle \mathbf{y}, \mathbf{x}\rangle \mathbb{1}_{3} - |\mathbf{y}\rangle \langle \mathbf{x}| \text{ and thus } \langle \mathbf{x} \cdot \mathcal{F}, \mathbf{y} \cdot \mathcal{F} \rangle = 2 \langle \mathbf{x}, \mathbf{y} \rangle;$   
(v)  $(\mathbf{x} \times \mathbf{y}) \cdot \mathcal{F} = |\mathbf{x}\rangle \langle \mathbf{y}| - |\mathbf{y}\rangle \langle \mathbf{x}|.$ 

Proof. For the 1st item, it is trivial result. For the 2nd item, in fact, we can check this identity directly as follows:

$$(\mathbf{x} \cdot \mathcal{F})^{\mathsf{T}} \mathbf{y} = - \begin{pmatrix} 0 & x_3 & -x_2 \\ -x_3 & 0 & x_1 \\ x_2 & -x_1 & 0 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} = - \begin{pmatrix} x_3 y_2 - x_2 y_3 \\ x_1 y_3 - x_3 y_1 \\ x_2 y_1 - x_1 y_2 \end{pmatrix}$$
$$= \begin{pmatrix} \left| \begin{array}{c} x_2 & x_3 \\ y_2 & y_3 \end{array}\right|, - \left| \begin{array}{c} x_1 & x_3 \\ y_1 & y_3 \end{array}\right|, \left| \begin{array}{c} x_1 & x_2 \\ y_1 & y_2 \end{array}\right| \end{pmatrix}^{\mathsf{T}} \\ = \mathbf{x} \times \mathbf{y}.$$

The 3rd item can be also calculated immediately. Indeed, note that  $(\mathbf{x} \cdot \mathcal{F})^{\mathsf{T}} \mathbf{e}_j = \mathbf{x} \times \mathbf{e}_j$  or in Dirac notation,

$$(\boldsymbol{x}\cdot\boldsymbol{\mathcal{F}})^{\mathsf{T}}|\boldsymbol{e}_{j}\rangle = |\boldsymbol{x}\times\boldsymbol{e}_{j}\rangle, \quad j = 1, 2, 3,$$

we get that

$$\begin{aligned} \mathbf{x} \cdot \mathcal{F} &= -(\mathbf{x} \cdot \mathcal{F})^{\mathsf{T}} \sum_{j=1}^{3} |\mathbf{e}_{j}\rangle \langle \mathbf{e}_{j}| = -\sum_{j=1}^{3} (\mathbf{x} \cdot \mathcal{F})^{\mathsf{T}} |\mathbf{e}_{j}\rangle \langle \mathbf{e}_{j}| \\ &= -\sum_{j=1}^{3} |\mathbf{x} \times \mathbf{e}_{j}\rangle \langle \mathbf{e}_{j}| = \sum_{j=1}^{3} |\mathbf{e}_{j} \times \mathbf{x}\rangle \langle \mathbf{e}_{j}|. \end{aligned}$$

Analogously,

$$x\cdot \mathcal{F} \;\;=\;\; \sum_{j=1}^3 |e_j
angle\langle e_j|(x\cdot \mathcal{F}) = \sum_{j=1}^3 |e_j
angle\langle x imes e_j|.$$

For the 4th item, Furthermore,

$$\begin{aligned} (\boldsymbol{x} \cdot \boldsymbol{\mathcal{F}})^{\mathsf{T}}(\boldsymbol{y} \cdot \boldsymbol{\mathcal{F}}) &= (\boldsymbol{x} \cdot \boldsymbol{\mathcal{F}})^{\mathsf{T}} \sum_{j=1}^{3} |\boldsymbol{e}_{j}\rangle \langle \boldsymbol{e}_{j}| (\boldsymbol{y} \cdot \boldsymbol{\mathcal{F}}) = \sum_{j=1}^{3} (\boldsymbol{x} \cdot \boldsymbol{\mathcal{F}})^{\mathsf{T}} |\boldsymbol{e}_{j}\rangle \langle \boldsymbol{e}_{j}| (\boldsymbol{y} \cdot \boldsymbol{\mathcal{F}}) \\ &= \sum_{j=1}^{3} |\boldsymbol{x} \times \boldsymbol{e}_{j}\rangle \langle \boldsymbol{y} \times \boldsymbol{e}_{j}| = \sum_{j=1}^{3} |\boldsymbol{e}_{j} \times \boldsymbol{x}\rangle \langle \boldsymbol{e}_{j} \times \boldsymbol{y}| = \sum_{j=1}^{3} \boldsymbol{F}_{j} |\boldsymbol{x}\rangle \langle \boldsymbol{y}| \boldsymbol{F}_{j}^{\mathsf{T}} \end{aligned}$$

Note that  $\langle F_i, F_j \rangle = 2\delta_{ij}$ . We get that  $\langle x \cdot \mathcal{F}, y \cdot \mathcal{F} \rangle = 2\langle x, y \rangle$ . For the last item, we see that

$$\begin{aligned} (\boldsymbol{x} \times \boldsymbol{y}) \cdot \mathcal{F} &= \sum_{k=1}^{3} \langle \boldsymbol{x} \big| \boldsymbol{F}_{k} \big| \boldsymbol{y} \rangle \boldsymbol{F}_{k} = \sum_{k=1}^{3} \operatorname{Tr} \left( \boldsymbol{F}_{k} \big| \boldsymbol{y} \rangle \langle \boldsymbol{x} \big| \right) \boldsymbol{F}_{k} = -\sum_{k=1}^{3} \operatorname{Tr} \left( \boldsymbol{F}_{k} \big| \boldsymbol{x} \rangle \langle \boldsymbol{y} \big| \right) \boldsymbol{F}_{k} \\ &= -\sum_{k=1}^{3} \frac{1}{2} \operatorname{Tr} \left( \boldsymbol{F}_{k} (|\boldsymbol{x}\rangle \langle \boldsymbol{y} | - |\boldsymbol{y}\rangle \langle \boldsymbol{x} |) \right) \boldsymbol{F}_{k} = |\boldsymbol{x}\rangle \langle \boldsymbol{y} | - |\boldsymbol{y}\rangle \langle \boldsymbol{x} |. \end{aligned}$$

This completes the proof.

In fact, the 2nd item in Lemma B.9 can be viewed as the implementation of cross product by matrix multiplication. This observation is simple but very important throughout this paper.

Another important fact is paramount in the following development. In fact,

**Lemma B.10.** For arbitrary two matrices  $M, N \in \mathbb{R}^{3 \times 3}$  and any two vectors  $x, y \in \mathbb{R}^3$ , it holds that

(i) 
$$\Omega(M, M) = 2\widehat{M}$$
.

(ii)  $M(x \cdot \mathcal{F})N^{\mathsf{T}} + N(x \cdot \mathcal{F})M^{\mathsf{T}} = (\Omega(M, N)x) \cdot \mathcal{F}$ . In particular, for M = N, we get that

$$M(x \cdot \mathcal{F})M^{\mathsf{T}} = \frac{1}{2}(\Omega(M, M)x) \cdot \mathcal{F} = (\widehat{M}x) \cdot \mathcal{F}.$$
(B.21)

(iii) 
$$M^{\mathsf{T}}[(Mx) \cdot \mathcal{F}]M = \det(M)(x \cdot \mathcal{F}) \text{ and } M[(M^{\mathsf{T}}x) \cdot \mathcal{F}]M^{\mathsf{T}} = \det(M)(x \cdot \mathcal{F}).$$
  
(iv)  $\langle (x \cdot \mathcal{F})M, M(y \cdot \mathcal{F}) \rangle = 2\langle x | \widehat{M} | y \rangle.$ 

*Proof.* For the 1st item, the proof is can be obtained immediately by direct computation. Indeed, using Proposition B.4, for M = N, we get that

$$\Omega(M, M)_{p,q} = -\langle F_p M F_q, M \rangle = \langle F_p M, M F_q \rangle$$
  
=  $\langle (e_p \cdot F) M, M(e_q \cdot F) \rangle = 2 \langle e_p | \widehat{M} | e_q \rangle,$ 

implying that  $\Omega(M, M) = 2\widehat{M}$ . For the 2nd item, it is easily seen that

$$\left(M(x \cdot \mathcal{F})N^{\mathsf{T}} + N(x \cdot \mathcal{F})M^{\mathsf{T}}\right)^{\mathsf{T}} = -\left(M(x \cdot \mathcal{F})N^{\mathsf{T}} + N(x \cdot \mathcal{F})M^{\mathsf{T}}\right).$$

Thus it can be decomposed as

$$M(\mathbf{x} \cdot \mathcal{F})N^{\mathsf{T}} + N(\mathbf{x} \cdot \mathcal{F})M^{\mathsf{T}} = \sum_{k=1}^{3} c_{k}(M, N)F_{k},$$

where the coefficients  $c_k$  can be identified with

$$c_k = -\frac{1}{2} \operatorname{Tr} \left( \boldsymbol{M}(\boldsymbol{x} \cdot \boldsymbol{\mathcal{F}}) \boldsymbol{N}^{\mathsf{T}} \boldsymbol{F}_k \right) - \frac{1}{2} \operatorname{Tr} \left( \boldsymbol{N}(\boldsymbol{x} \cdot \boldsymbol{\mathcal{F}}) \boldsymbol{M}^{\mathsf{T}} \boldsymbol{F}_k \right) \\ = \langle \boldsymbol{e}_k, \Omega(\boldsymbol{M}, \boldsymbol{N}) \boldsymbol{x} \rangle,$$

implying that  $M(x \cdot \mathcal{F})N^{\mathsf{T}} + N(x \cdot \mathcal{F})M^{\mathsf{T}} = (\Omega(M, N)x) \cdot \mathcal{F}$ . In particular, for M = N, the desired identity follows immediately from  $\Omega(M, M) = 2\widehat{M}$ . For the 3rd item, we see from the obtained result in (ii) that

$$M^{\mathsf{T}}[(Mx)\cdot\mathcal{F}]M = (\widehat{M}^{\mathsf{T}}Mx)\cdot\mathcal{F} = \det(M)(x\cdot\mathcal{F}).$$

For the 4th item,

$$\begin{split} \left\langle (x \cdot \mathcal{F})M, M(y \cdot \mathcal{F}) \right\rangle &= \operatorname{Tr} \left( M^{\mathsf{T}}(x \cdot \mathcal{F})^{\mathsf{T}}M(y \cdot \mathcal{F}) \right) = \operatorname{Tr} \left( (x \cdot \mathcal{F})^{\mathsf{T}}M(y \cdot \mathcal{F})M^{\mathsf{T}} \right) \\ &= \operatorname{Tr} \left( (x \cdot \mathcal{F})^{\mathsf{T}}[(\widehat{M}y) \cdot \mathcal{F}] \right) = \left\langle x \cdot \mathcal{F}, (\widehat{M}y) \cdot \mathcal{F} \right\rangle \\ &= 2 \langle x | \widehat{M} | y \rangle. \end{split}$$

In the first equality, we used the definition of Hilbert-Schmidt inner product. For the 2nd equality, we used the cyclicity of trace. In the 3rd equality, we used the obtained result in (ii). In the last equality, we used the fact obtained in (iii) of Lemma B.9.  $\Box$ 

**Corollary B.11.** For an arbitrary invertible matrix  $L \in \mathbb{R}^{3 \times 3}$  and any two vectors  $u, v \in \mathbb{R}^3$ , we have that

(i)  $L(\mathbf{u} \times \mathbf{v}) = \widehat{L}\mathbf{u} \times (L^{\mathsf{T}})^{-1}\mathbf{v}$ , in fact, we see that

$$L(\boldsymbol{u} \times \boldsymbol{v}) = \widehat{L}\boldsymbol{u} \times (L^{\mathsf{T}})^{-1}\boldsymbol{v}$$
(B.22)

$$= \det(L)\left((L^{\mathsf{T}})^{-1}\boldsymbol{u}\times(L^{\mathsf{T}})^{-1}\boldsymbol{v}\right)$$
(B.23)

$$= \frac{1}{\det(L)} \left( \widehat{L}u \times \widehat{L}v \right). \tag{B.24}$$

In particular, for  $R \in SO(3)$ , the special orthogonal group of order 3, we recover the well-known formula:

$$\boldsymbol{R}(\boldsymbol{u} \times \boldsymbol{v}) = \boldsymbol{R}\boldsymbol{u} \times \boldsymbol{R}\boldsymbol{v}. \tag{B.25}$$

(ii)  $Lu \times Lv = \det(L)(L^{\mathsf{T}})^{-1}(u \times v) = \widehat{L}(u \times v).$ 

*Proof.* It suffices to show that  $L(u \times v) = \hat{L}u \times (L^{T})^{-1}v$ . Indeed, via the fact that  $u \times v = (u \cdot F)^{T}v$ , using the result in (ii) of Lemma B.10

$$L(\boldsymbol{u} \times \boldsymbol{v}) = L(\boldsymbol{u} \cdot \boldsymbol{F})^{\mathsf{T}} \boldsymbol{v} = L(\boldsymbol{u} \cdot \boldsymbol{\mathcal{F}})^{\mathsf{T}} L^{\mathsf{T}} (L^{\mathsf{T}})^{-1} \boldsymbol{v} = [L(\boldsymbol{u} \cdot \boldsymbol{\mathcal{F}}) L^{\mathsf{T}}]^{\mathsf{T}} (L^{\mathsf{T}})^{-1} \boldsymbol{v}$$
$$= [(\widehat{L}\boldsymbol{u}) \cdot \boldsymbol{\mathcal{F}}]^{\mathsf{T}} (L^{\mathsf{T}})^{-1} \boldsymbol{v} = \widehat{L}\boldsymbol{u} \times (L^{\mathsf{T}})^{-1} \boldsymbol{v}.$$

Due to the fact that  $\widehat{L}L^{\mathsf{T}} = \det(L)\mathbb{1}_3$  and *L* is invertible, we get the other two forms of this formula. In particular, for  $L = R \in \mathsf{SO}(3)$ , then  $\det(R) = 1$  and  $\widehat{R} = R$ , which leads to the desired identity.

The above results are obtained under the invertibility condition. In fact, we can remove such condition, that is, the following identities holds for any matrix  $L \in \mathbb{R}^{3 \times 3}$ :

$$\det(L)L(u \times v) = \widehat{L}u \times \widehat{L}v, \qquad (B.26)$$

$$Lu \times Lv = \widehat{L}(u \times v). \tag{B.27}$$

**Corollary B.12.** For any two matrices  $M, N \in \mathbb{R}^{3 \times 3}$  and any two vectors  $u, v \in \mathbb{R}^3$ , it holds that

$$Mu \times Nv + Nu \times Mv = \Omega(M, N)(u \times v).$$
(B.28)

In particular, for M = N, we get that  $Mu \times Mv = \frac{1}{2}\Omega(M, M)(u \times v) = \widehat{M}(u \times v)$ .

Proof. In fact,

$$\begin{aligned} Mu \times Nv + Nu \times Mv &= \left( \langle Mu | F_1 | Nv \rangle, \langle Mu | F_2 | Nv \rangle, \langle Mu | F_3 | Nv \rangle \right)^{\mathsf{T}} \\ &+ \left( \langle Nu | F_1 | Mv \rangle, \langle Nu | F_2 | Mv \rangle, \langle Nu | F_3 | Mv \rangle \right)^{\mathsf{T}}, \end{aligned}$$

which is equal to

$$\left(\left\langle u \middle| M^{\mathsf{T}} F_1 N + N^{\mathsf{T}} F_1 M \middle| v \right\rangle, \left\langle u \middle| M^{\mathsf{T}} F_2 N + N^{\mathsf{T}} F_2 M \middle| v \right\rangle, \left\langle u \middle| M^{\mathsf{T}} F_3 N + N^{\mathsf{T}} F_3 M \middle| v \right\rangle\right)^{\mathsf{T}}.$$

Now we can easily check that

$$\boldsymbol{M}^{\mathsf{T}}\boldsymbol{F}_{i}\boldsymbol{N}+\boldsymbol{N}^{\mathsf{T}}\boldsymbol{F}_{i}\boldsymbol{M}=-\sum_{j=1}^{3}\left\langle \boldsymbol{F}_{i}\boldsymbol{M}\boldsymbol{F}_{j},\boldsymbol{N}
ight
angle \boldsymbol{F}_{j},$$

implying that

$$\langle \boldsymbol{u} | \boldsymbol{M}^{\mathsf{T}} \boldsymbol{F}_{i} \boldsymbol{N} + \boldsymbol{N}^{\mathsf{T}} \boldsymbol{F}_{i} \boldsymbol{M} | \boldsymbol{v} \rangle = -\sum_{j=1}^{3} \langle \boldsymbol{F}_{i} \boldsymbol{M} \boldsymbol{F}_{j}, \boldsymbol{N} \rangle \langle \boldsymbol{u} | \boldsymbol{F}_{j} | \boldsymbol{v} \rangle = \sum_{j=1}^{3} \Omega(\boldsymbol{M}, \boldsymbol{N})_{i,j} \langle \boldsymbol{u} | \boldsymbol{F}_{j} | \boldsymbol{v} \rangle$$
$$= [\Omega(\boldsymbol{M}, \boldsymbol{N})(\boldsymbol{u} \times \boldsymbol{v})]_{i}.$$

That is, the desired result is true.

**Corollary B.13.** *For any two matrices*  $M, N \in \mathbb{R}^{3 \times 3}$ *, it holds that* 

$$\Omega(M,N) = \widehat{M+N} - \left(\widehat{M} + \widehat{N}\right).$$
(B.29)

Proof. Indeed, using Corollary B.12, we get that

$$(M+N)u \times (M+N)v = \widehat{M+N}(u \times v),$$
$$Mu \times Mv = \widehat{M}(u \times v),$$
$$Nu \times Nv = \widehat{N}(u \times v),$$

and thus

$$\begin{aligned} Mu \times Nv + Nu \times Mv &= (M+N)u \times (M+N)v - Mu \times Mv - Nu \times Nv \\ &= \widehat{M+N}(u \times v) - \widehat{M}(u \times v) - \widehat{N}(u \times v) \\ &= \left(\widehat{M+N} - \widehat{M} - \widehat{N}\right)(u \times v) = \Omega(M,N)(u \times v), \end{aligned}$$

implying that  $\Omega(M, N) = \widehat{M+N} - \widehat{M} - \widehat{N}$ .

**Corollary B.14.** For any matrix  $M \in \mathbb{R}^{3 \times 3}$  and any vectors  $x, y \in \mathbb{R}^3$ , it holds that

(i)  $[(My) \cdot \mathcal{F}]M = \widehat{M}(y \cdot \mathcal{F}).$ 

(*ii*) 
$$M[(M^{\mathsf{T}}x) \cdot \mathcal{F}] = (x \cdot \mathcal{F})\widehat{M}.$$

*Proof.* We can prove these results in two steps:

• Assume that *M* is invertible. Then by the result in (iii) of Lemma B.10, we get that

$$M^{\mathsf{T}}[(My) \cdot \mathcal{F}]M = \det(M)(y \cdot \mathcal{F}) = M^{\mathsf{T}}M(y \cdot \mathcal{F}).$$

Because *M* is invertible, i.e.,  $M^{\mathsf{T}}$  is also invertible, we get that

$$[(My)\cdot\mathcal{F}]M=\widehat{M}(y\cdot\mathcal{F}).$$

Analogously, we also have that  $M[(M^{\mathsf{T}}x) \cdot \mathcal{F}] = (x \cdot \mathcal{F})\widehat{M}$ .

Now if *M* is not invertible, then we can take a net by using SVD such that *M* can be approximated in any precision by such net. Indeed, via SVD, there there exist two orthogonal matrices *P* and *Q* in O(3) such that *M* = *P*Σ*Q*<sup>T</sup>, where Σ = diag(*m*<sub>1</sub>, *m*<sub>2</sub>, *m*<sub>3</sub>) consists of singular values of *M*. Let such net {*M*<sub>ε</sub> : *ε* > 0} be given by *M*<sub>ε</sub> = *P*(Σ + *ε*1<sub>3</sub>)*Q*<sup>T</sup> for small enough *ε* > 0. Now lim<sub>ε→0<sup>+</sup></sub> *M*<sub>ε</sub> = *M* and

$$[(M_{\epsilon}y)\cdot \mathcal{F}]M_{\epsilon}=\widehat{M}_{\epsilon}(y\cdot \mathcal{F}).$$

The proof can be finished by taking the limit for  $\epsilon \rightarrow 0^+$  on both sides of the above expression due to the continuity argument and the fact that

$$\lim_{\epsilon \to 0^+} \widehat{M}_{\epsilon} = \widehat{M}.$$
(B.30)

To this end, using the result (i) in Lemma B.3, we see that

$$\lim_{\epsilon \to 0^+} \operatorname{Tr}\left(\widehat{M}_{\epsilon}\right) = \frac{1}{2} \left( \lim_{\epsilon \to 0^+} \operatorname{Tr}\left(M_{\epsilon}\right)^2 - \lim_{\epsilon \to 0^+} \operatorname{Tr}\left(M_{\epsilon}^2\right) \right) \\ = \frac{1}{2} \left( \lim_{\epsilon \to 0^+} \operatorname{Tr}\left(M\right)^2 - \lim_{\epsilon \to 0^+} \operatorname{Tr}\left(M^2\right) \right) = \operatorname{Tr}\left(\widehat{M}\right).$$

By Proposition B.2, we get

$$\begin{split} \lim_{\epsilon \to 0^+} \widehat{M}_{\epsilon} &= \lim_{\epsilon \to 0^+} \left( M_{\epsilon}^2 - \operatorname{Tr} \left( M_{\epsilon} \right) M_{\epsilon} + \operatorname{Tr} \left( \widehat{M}_{\epsilon} \right) \mathbb{1}_3 \right)^{\mathsf{T}} \\ &= \left( \lim_{\epsilon \to 0^+} M_{\epsilon}^2 - \lim_{\epsilon \to 0^+} \operatorname{Tr} \left( M_{\epsilon} \right) \lim_{\epsilon \to 0^+} M_{\epsilon} + \lim_{\epsilon \to 0^+} \operatorname{Tr} \left( \widehat{M}_{\epsilon} \right) \mathbb{1}_3 \right)^{\mathsf{T}} \\ &= \left( M^2 - \operatorname{Tr} \left( M \right) M + \operatorname{Tr} \left( \widehat{M} \right) \mathbb{1}_3 \right)^{\mathsf{T}} = \widehat{M}. \end{split}$$

The proof is complete.

Next we summarize important properties concerning  $\Omega$ .

**Lemma B.15.** For  $\Omega$ , defined in Eq. (B.12), it holds that

- (i)  $\Omega(T, |a\rangle\langle b|) = (a \cdot \mathcal{F})T(b \cdot \mathcal{F})^{\mathsf{T}}.$
- (ii)  $\Omega(M, \widehat{T}) = \operatorname{Tr}(M^{\mathsf{T}}T) T TM^{\mathsf{T}}T$ , in particular,  $\Omega(T, \widehat{T}) = ||T||^2 T TT^{\mathsf{T}}T$ .
- (iii)  $\Omega(T, AT) = \operatorname{Tr}(A) \widehat{T} A^{\mathsf{T}} \widehat{T}$ ; in particular,  $\Omega(T, TT^{\mathsf{T}}T) = \|T\|^2 \widehat{T} \det(T)T$ .

(*iv*) 
$$\Omega(T, TB) = \operatorname{Tr}(B) \widehat{T} - \widehat{T}B^{\mathsf{T}}$$
.

(v) 
$$\Omega(T, (\mathbf{r} \cdot \mathcal{F})T(\mathbf{s} \cdot \mathcal{F})^{\mathsf{T}}) = \langle \mathbf{r} | T | \mathbf{s} \rangle T + ||T||^2 |\mathbf{r}\rangle \langle \mathbf{s}| - (|\mathbf{r}\rangle \langle \mathbf{s}|T^{\mathsf{T}}T + TT^{\mathsf{T}}|\mathbf{r}\rangle \langle \mathbf{s}|).$$

(vi)  $\Omega(T, x \cdot \mathcal{F}) = Tx \cdot \mathcal{F} + |n\rangle \langle x|$ , where  $n = \sum_{i=1}^{3} Te_i \times e_i$  is determined from  $T - T^{\mathsf{T}} = n \cdot \mathcal{F}$ .

(vii)  $\Omega(\mathbf{x} \cdot \mathcal{F}, \mathbf{y} \cdot \mathcal{F}) = |\mathbf{x}\rangle \langle \mathbf{y}| + |\mathbf{y}\rangle \langle \mathbf{x}|$ . In particular,  $\Omega(\mathbf{x} \cdot \mathcal{F}, \mathbf{x} \cdot \mathcal{F}) = 2|\mathbf{x}\rangle \langle \mathbf{x}|$ . *Proof.* (i) Indeed, for  $T' = |\mathbf{a}\rangle \langle \mathbf{b}|$ , we see that

$$\Omega(T, T') = \begin{pmatrix} a_{3}e_{2}^{\mathsf{T}}T \times b^{\mathsf{T}} \\ a_{1}e_{3}^{\mathsf{T}}T \times b^{\mathsf{T}} \\ a_{2}e_{1}^{\mathsf{T}}T \times b^{\mathsf{T}} \end{pmatrix} - \begin{pmatrix} a_{2}e_{3}^{\mathsf{T}}T \times b^{\mathsf{T}} \\ a_{3}e_{1}^{\mathsf{T}}T \times b^{\mathsf{T}} \\ a_{1}e_{2}^{\mathsf{T}}T \times b^{\mathsf{T}} \end{pmatrix}$$
$$= \operatorname{diag}(a_{3}, a_{1}, a_{2}) \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix} \begin{pmatrix} e_{1}^{\mathsf{T}}T \times b^{\mathsf{T}} \\ e_{2}^{\mathsf{T}}T \times b^{\mathsf{T}} \\ e_{3}^{\mathsf{T}}T \times b^{\mathsf{T}} \end{pmatrix}$$
$$-\operatorname{diag}(a_{2}, a_{3}, a_{1}) \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} e_{1}^{\mathsf{T}}T \times b^{\mathsf{T}} \\ e_{3}^{\mathsf{T}}T \times b^{\mathsf{T}} \\ e_{3}^{\mathsf{T}}T \times b^{\mathsf{T}} \end{pmatrix},$$

which is just equal to  $(a \cdot \mathcal{F})T(b \cdot \mathcal{F})^{\mathsf{T}}$ , where we used the facts that

$$\operatorname{diag}(a_{3}, a_{1}, a_{2}) \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix} - \operatorname{diag}(a_{2}, a_{3}, a_{1}) \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} = \boldsymbol{a} \cdot \mathcal{F}$$
$$\begin{pmatrix} \boldsymbol{e}_{1}^{\mathsf{T}} \mathbf{T} \times \boldsymbol{b}^{\mathsf{T}} \\ \boldsymbol{e}_{2}^{\mathsf{T}} \mathbf{T} \times \boldsymbol{b}^{\mathsf{T}} \\ \boldsymbol{e}_{3}^{\mathsf{T}} \mathbf{T} \times \boldsymbol{b}^{\mathsf{T}} \end{pmatrix} = \boldsymbol{T} (\boldsymbol{b} \cdot \mathcal{F})^{\mathsf{T}}.$$

 $\left(e_{3}^{\mathsf{T}}T \times b^{\mathsf{T}}\right)$ (ii) The correctness of this result can be directly checked by MATHEMATICA. In what follows, we infer it by analytical method. In fact, using the result obtained in (i) previously,

$$\begin{aligned} \Omega(|a\rangle\langle b|, \widehat{T}) &= & \Omega(\widehat{T}, |a\rangle\langle b|) = (a \cdot \mathcal{F})\widehat{T}(b \cdot \mathcal{F})^{\mathsf{T}} \\ &= & (a \cdot \mathcal{F})(Tb \cdot \mathcal{F})^{\mathsf{T}}T = (\langle a|T|b\rangle \mathbb{1}_3 - T|b\rangle\langle a|)T \\ &= & \operatorname{Tr}(|b\rangle\langle a|T)T - T|b\rangle\langle a|T. \end{aligned}$$

Here in the 3rd equality, we used the first property in Corollary B.14; and in the 4th equality, we used the third property in Lemma B.9. Now using SVD of M:  $M = \sum_{j=1}^{3} s_j |a_j\rangle \langle b_j|$ , we can finish the proof:

$$\Omega(\boldsymbol{M}, \boldsymbol{\widehat{T}}) = \operatorname{Tr}(\boldsymbol{M}^{\mathsf{T}}\boldsymbol{T}) \boldsymbol{T} - \boldsymbol{T}\boldsymbol{M}^{\mathsf{T}}\boldsymbol{T}.$$

Indeed, by the bi-linearity of  $\Omega(\cdot,\cdot),$ 

and

$$\Omega(T, \widehat{T}) = \sum_{j=1}^{3} s_{j} \Omega(|a_{j}\rangle \langle b_{j}|, \widehat{T}) = \sum_{j=1}^{3} s_{j} \left( \operatorname{Tr} \left( |b_{j}\rangle \langle a_{j}|T \right) T - T |b_{j}\rangle \langle a_{j}|T \right) \right)$$
$$= \operatorname{Tr} \left( \sum_{j=1}^{3} s_{j} |b_{j}\rangle \langle a_{j}|T \right) T - T \sum_{j=1}^{3} s_{j} |b_{j}\rangle \langle a_{j}|T \right)$$
$$= \operatorname{Tr} \left( T^{\mathsf{T}}T \right) T - T T^{\mathsf{T}}T = ||T||^{2} T - T T^{\mathsf{T}}T.$$

(iii) We prove a stronger result: If T' = AT for any  $3 \times 3$  real matrix A, where  $e_k^T A = (a_{k1}, a_{k2}, a_{k3})(k \in [3])$ ,

$$\begin{split} \Omega(T, AT) &= \begin{pmatrix} e_2^{\mathsf{T}}T \times (a_{31}e_1^{\mathsf{T}}T + a_{33}e_3^{\mathsf{T}}T) + (a_{21}e_1^{\mathsf{T}}T + a_{22}e_2^{\mathsf{T}}T) \times e_3^{\mathsf{T}}T \\ e_3^{\mathsf{T}}T \times (a_{11}e_1^{\mathsf{T}}T + a_{12}e_2^{\mathsf{T}}T) + (a_{32}e_2^{\mathsf{T}}T + a_{33}e_3^{\mathsf{T}}T) \times e_1^{\mathsf{T}}T \\ e_1^{\mathsf{T}}T \times (a_{22}e_2^{\mathsf{T}}T + a_{23}e_3^{\mathsf{T}}T) + (a_{11}e_1^{\mathsf{T}}T + a_{13}e_3^{\mathsf{T}}T) \times e_2^{\mathsf{T}}T \end{pmatrix} \\ &= \begin{pmatrix} (a_{22} + a_{33})e_2^{\mathsf{T}}T \times e_3^{\mathsf{T}}T \\ (a_{11} + a_{33})e_3^{\mathsf{T}}T \times e_1^{\mathsf{T}}T \\ (a_{11} + a_{22})e_1^{\mathsf{T}}T \times e_2^{\mathsf{T}}T \end{pmatrix} + \begin{pmatrix} a_{31}e_2^{\mathsf{T}}T \times e_1^{\mathsf{T}}T \\ a_{23}e_1^{\mathsf{T}}T \times e_3^{\mathsf{T}}T \\ a_{23}e_1^{\mathsf{T}}T \times e_3^{\mathsf{T}}T \end{pmatrix} + \begin{pmatrix} a_{21}e_1^{\mathsf{T}}T \times e_3^{\mathsf{T}}T \\ a_{12}e_2^{\mathsf{T}}T \times e_3^{\mathsf{T}}T \\ a_{12}e_2^{\mathsf{T}}T \times e_3^{\mathsf{T}}T \end{pmatrix} \\ &= [\operatorname{Tr}(A) - \operatorname{diag}(a_{11}, a_{22}, a_{33})]\widehat{T} - \begin{pmatrix} a_{31}e_1^{\mathsf{T}}T \times e_3^{\mathsf{T}}T \\ a_{22}e_3^{\mathsf{T}}T \times e_1^{\mathsf{T}}T \\ a_{23}e_3^{\mathsf{T}}T \times e_1^{\mathsf{T}}T \end{pmatrix} - \begin{pmatrix} a_{21}e_3^{\mathsf{T}}T \times e_2^{\mathsf{T}}T \\ a_{32}e_1^{\mathsf{T}}T \times e_3^{\mathsf{T}}T \\ a_{12}e_2^{\mathsf{T}}T \times e_3^{\mathsf{T}}T \\ a_{23}e_3^{\mathsf{T}}T \times e_1^{\mathsf{T}}T \end{pmatrix} \\ &= \operatorname{Tr}(A)\widehat{T} - \operatorname{diag}(a_{11}, a_{22}, a_{33})\widehat{T} - \begin{pmatrix} 0 & a_{21} & a_{31} \\ a_{12} & 0 & a_{32} \\ a_{13} & a_{23} & 0 \end{pmatrix} \widehat{T} = \operatorname{Tr}(A)\widehat{T} - A^{\mathsf{T}}\widehat{T}. \end{split}$$

That is, for T' = AT,

$$\Omega(T, AT) = \operatorname{Tr}(A)\,\widehat{T} - A^{\mathsf{T}}\widehat{T}.\tag{B.31}$$

Letting in the above  $A = TT^{\mathsf{T}}$ , we get that

$$\operatorname{Tr}(A)\,\widehat{T} - A^{\mathsf{T}}\widehat{T} = \operatorname{Tr}(TT^{\mathsf{T}})\,\widehat{T} - TT^{\mathsf{T}}\widehat{T} = \langle T,T\rangle\widehat{T} - \det(T)T.$$

The another approach to this result can be described as follows. Indeed, *A* can be decomposed as  $A = \sum_{i=1}^{3} s_i |x_i\rangle \langle y_i|$  by SVD. Then

$$\begin{split} \Omega(\boldsymbol{T},\boldsymbol{A}\boldsymbol{T}) &= \sum_{i=1}^{3} s_{i} \Omega(\boldsymbol{T},|\boldsymbol{x}_{i}\rangle\langle\boldsymbol{y}_{i}|\boldsymbol{T}) = \sum_{j=1}^{3} s_{i} (\boldsymbol{x}_{i} \cdot \boldsymbol{\mathcal{F}}) \boldsymbol{T} ((\boldsymbol{T}^{\mathsf{T}}\boldsymbol{y}_{j}) \cdot \boldsymbol{\mathcal{F}})^{\mathsf{T}} \\ &= \sum_{i=1}^{3} s_{i} (\boldsymbol{x}_{i} \cdot \boldsymbol{\mathcal{F}}) (\boldsymbol{y}_{i} \cdot \boldsymbol{\mathcal{F}})^{\mathsf{T}} \widehat{\boldsymbol{T}} = \sum_{i=1}^{3} s_{i} \left( \langle \boldsymbol{y}_{i}, \boldsymbol{x}_{i} \rangle \mathbb{1}_{3} - |\boldsymbol{y}_{i}\rangle\langle\boldsymbol{x}_{i}| \right) \widehat{\boldsymbol{T}} \\ &= \operatorname{Tr}(\boldsymbol{A}) \widehat{\boldsymbol{T}} - \boldsymbol{A}^{\mathsf{T}} \widehat{\boldsymbol{T}}. \end{split}$$

(iv) By SVD of B,  $B = \sum_{i=1}^{3} s_{i} |x_{i}\rangle \langle y_{j}|$ . Now

$$\Omega(T,TB) = \sum_{j=1}^{3} s_j \Omega(T,T|x_j\rangle\langle y_j|) = \sum_{j=1}^{3} s_j \Omega(T,|Tx_j\rangle\langle y_j|).$$

Using (i), we get that

$$\begin{aligned} \Omega(\boldsymbol{T},\boldsymbol{T}\boldsymbol{B}) &= \sum_{j=1}^{3} s_{j} \Omega(\boldsymbol{T},|\boldsymbol{T}\boldsymbol{x}_{j}\rangle\langle\boldsymbol{y}_{j}|) = \sum_{j=1}^{3} s_{j} ((\boldsymbol{T}\boldsymbol{x}_{j})\cdot\boldsymbol{\mathcal{F}})\boldsymbol{T}(\boldsymbol{y}_{j}\cdot\boldsymbol{\mathcal{F}})^{\mathsf{T}} \\ &= \sum_{j=1}^{3} s_{j} \widehat{\boldsymbol{T}}(\boldsymbol{x}_{j}\cdot\boldsymbol{\mathcal{F}})(\boldsymbol{y}_{j}\cdot\boldsymbol{\mathcal{F}})^{\mathsf{T}} = \widehat{\boldsymbol{T}} \sum_{j=1}^{3} s_{j} (\boldsymbol{x}_{j}\cdot\boldsymbol{\mathcal{F}})(\boldsymbol{y}_{j}\cdot\boldsymbol{\mathcal{F}})^{\mathsf{T}} \\ &= \widehat{\boldsymbol{T}} \sum_{j=1}^{3} s_{j} \left( \langle \boldsymbol{y}_{j},\boldsymbol{x}_{j}\rangle \mathbb{1}_{3} - |\boldsymbol{y}_{j}\rangle\langle\boldsymbol{x}_{j}| \right) = \mathrm{Tr}\left(\boldsymbol{B}\right) \widehat{\boldsymbol{T}} - \widehat{\boldsymbol{T}}\boldsymbol{B}^{\mathsf{T}}. \end{aligned}$$

(v) Note that

$$(\mathbf{r}\cdot\mathcal{F})\mathbf{T}(\mathbf{s}\cdot\mathcal{F})^{\mathsf{T}} = \begin{pmatrix} \mathbf{e}_{1}^{\mathsf{T}}(\mathbf{r}\cdot\mathcal{F})\mathbf{T}\times\mathbf{s}^{\mathsf{T}} \\ \mathbf{e}_{2}^{\mathsf{T}}(\mathbf{r}\cdot\mathcal{F})\mathbf{T}\times\mathbf{s}^{\mathsf{T}} \\ \mathbf{e}_{3}^{\mathsf{T}}(\mathbf{r}\cdot\mathcal{F})\mathbf{T}\times\mathbf{s}^{\mathsf{T}} \end{pmatrix}$$

implies that

$$e_k^{\mathsf{T}}(\boldsymbol{r}\cdot\mathcal{F})T(\boldsymbol{s}\cdot\mathcal{F})^{\mathsf{T}}=e_k^{\mathsf{T}}(\boldsymbol{r}\cdot\mathcal{F})T\times\boldsymbol{s}^{\mathsf{T}}$$

Using the facts that

$$(u \times v) \times w = \langle w, u \rangle v - \langle w, v \rangle u,$$
  
 $u \times (v \times w) = \langle u, w \rangle v - \langle u, v \rangle w,$ 

we get that

$$\begin{split} \Omega(T, (r \cdot \mathcal{F})T(s \cdot \mathcal{F})^{\mathsf{T}}) &= \begin{pmatrix} (e_2^{\mathsf{T}}(r \cdot \mathcal{F})T \times s^{\mathsf{T}}) \times e_3^{\mathsf{T}}T + e_2^{\mathsf{T}}T \times (e_3^{\mathsf{T}}(r \cdot \mathcal{F})T \times s^{\mathsf{T}}) \\ (e_3^{\mathsf{T}}(r \cdot \mathcal{F})T \times s^{\mathsf{T}}) \times e_1^{\mathsf{T}}T + e_3^{\mathsf{T}}T \times (e_1^{\mathsf{T}}(r \cdot \mathcal{F})T \times s^{\mathsf{T}}) \\ (e_1^{\mathsf{T}}(r \cdot \mathcal{F})T \times s^{\mathsf{T}}) \times e_2^{\mathsf{T}}T + e_3^{\mathsf{T}}T \times (e_2^{\mathsf{T}}(r \cdot \mathcal{F})T \times s^{\mathsf{T}}) \\ (e_1^{\mathsf{T}}(r \cdot \mathcal{F})T \times s^{\mathsf{T}}) \times e_2^{\mathsf{T}}T + e_1^{\mathsf{T}}T \times (e_2^{\mathsf{T}}(r \cdot \mathcal{F})T \times s^{\mathsf{T}}) \\ (Ts)_1 e_3^{\mathsf{T}}(r \cdot \mathcal{F})T - (Ts)_2 e_3^{\mathsf{T}}(r \cdot \mathcal{F})T \\ (Ts)_2 e_1^{\mathsf{T}}(r \cdot \mathcal{F})T - (Ts)_1 e_2^{\mathsf{T}}(r \cdot \mathcal{F})T \end{pmatrix} + \begin{pmatrix} \langle e_2 | \{r \cdot \mathcal{F}, TT^{\mathsf{T}}\} | e_3 \rangle \\ \langle e_3 | \{r \cdot \mathcal{F}, TT^{\mathsf{T}}\} | e_1 \rangle \\ \langle e_1 | \{r \cdot \mathcal{F}, TT^{\mathsf{T}}\} | e_2 \end{pmatrix} \end{pmatrix} s^{\mathsf{T}} \\ &= [(Ts) \cdot \mathcal{F}]^{\mathsf{T}}(r \cdot \mathcal{F})T + (\langle T, T \rangle \mathbb{1}_3 - TT^{\mathsf{T}}) | r \rangle \langle s| \\ &= \langle r | T | s \rangle T - |r \rangle \langle s | T^{\mathsf{T}}T + (||T||^2 \mathbb{1}_3 - TT^{\mathsf{T}}) | r \rangle \langle s|. \end{split}$$

Here  $\{A, B\} := AB + BA$ . Other items can be checked by direct calculation. This completes the proof.

**Lemma B.16.** For  $\Omega$ , defined in Eq. (B.12), it holds that

- (i)  $\Omega(A\widehat{T}, B) = \Omega(A, BT^{\mathsf{T}})T.$
- (*ii*)  $\Omega(\widehat{T}A, B) = T\Omega(A, T^{\mathsf{T}}B).$

*Proof.* For the 1st item, note that

$$\Omega(A\widehat{T}, |a\rangle\langle b|) = (a \cdot \mathcal{F})A\widehat{T}(b \cdot \mathcal{F})^{\mathsf{T}} = (a \cdot \mathcal{F})A(Tb \cdot \mathcal{F})^{\mathsf{T}}T$$
$$= \Omega(A, |a\rangle\langle b|T^{\mathsf{T}})T,$$

implying that

$$\Omega(A\widehat{T},B)=\Omega(A,BT^{\mathsf{T}})T.$$

For the 2nd item, we see that

$$\Omega(\widehat{T}A, |a\rangle\langle b|) = (a \cdot \mathcal{F})\widehat{T}A(b \cdot \mathcal{F})^{\mathsf{T}} = T(T^{\mathsf{T}}a \cdot \mathcal{F})A(b \cdot \mathcal{F})^{\mathsf{T}}$$
$$= T\Omega(A, T^{\mathsf{T}}|a\rangle\langle b|),$$

implying that  $\Omega(\widehat{T}A, B) = T\Omega(A, T^{\mathsf{T}}B)$ .

### **B.1.3** Recurrence relation for the matrix power

Let 
$$X^{1} \approx (t^{(1)}, r^{(1)}, s^{(1)}, T^{(1)}) = (t, r, s, T)$$
 and  $X^{k} \approx (t^{(k)}, r^{(k)}, s^{(k)}, T^{(k)})$ , i.e.,  
 $X^{k} = t^{(k)} \mathbb{1}_{4} + r^{(k)} \cdot \sigma \otimes \mathbb{1}_{2} + \mathbb{1}_{2} \otimes s^{(k)} \cdot \sigma + \sum_{i,j=1}^{3} t^{(k)}_{ij} \sigma_{i} \otimes \sigma_{j} \quad (k \ge 1),$ 
(B.32)

where  $T^{(k)} := \left(t_{ij}^{(k)}\right)_{3\times 3}$ . By Lemma B.5, we get that

**Corollary B.17.** The recurrence relations of coefficients between  $X^{k+1} = X^k X \approx (t^{(k+1)}, r^{(k+1)}, s^{(k+1)}, T^{(k+1)})$ and  $X^k \approx (t^{(k)}, r^{(k)}, s^{(k)}, T^{(k)})$  can be identified as:

$$\begin{cases} t^{(k+1)} = t^{(k)}t + \langle \mathbf{r}^{(k)}, \mathbf{r} \rangle + \langle \mathbf{s}^{(k)}, \mathbf{s} \rangle + \langle \mathbf{T}^{(k)}, \mathbf{T} \rangle \\ \mathbf{r}^{(k+1)} = t\mathbf{r}^{(k)} + t^{(k)}\mathbf{r} + \mathbf{T}\mathbf{s}^{(k)} + \mathbf{T}^{(k)}\mathbf{s} \\ \mathbf{s}^{(k+1)} = t\mathbf{s}^{(k)} + t^{(k)}\mathbf{s} + \mathbf{T}^{\mathsf{T}}\mathbf{r}^{(k)} + \mathbf{T}^{(k)^{\mathsf{T}}}\mathbf{r} \\ \mathbf{T}^{(k+1)} = |\mathbf{r}^{(k)}\rangle\langle \mathbf{s}| + |\mathbf{r}\rangle\langle \mathbf{s}^{(k)}| + t\mathbf{T}^{(k)} + t^{(k)}\mathbf{T} - \Omega(\mathbf{T}^{(k)}, \mathbf{T}) \end{cases}$$
(B.33)

where  $k \ge 1$ .

*Proof.* Using Corollary B.6, we see that  $[X^k, X] = 0$  if and only if

$$r^{(k)} \times r + \sum_{i=1}^{3} T^{(k)} e_i \times T e_i = 0,$$
  

$$s^{(k)} \times s + \sum_{i=1}^{3} T^{(k)^{\mathsf{T}}} e_i \times T^{\mathsf{T}} e_i = 0,$$
  

$$\Psi(r^{(k)}, T, s^{(k)}) = \Psi(r, T^{(k)}, s)$$

The recurrence relation is obtained immediately.

Using the previous results, we can list here the coefficients of  $X^k (1 \le k \le 4)$  below:

• For k = 2,  $X^2 \approx (t^{(2)}, r^{(2)}, s^{(2)}, T^{(2)})$  can be identified as

$$\begin{cases} t^{(2)} = t^2 + |\mathbf{r}|^2 + |\mathbf{s}|^2 + ||\mathbf{T}||^2, \\ \mathbf{r}^{(2)} = 2t \cdot \mathbf{r} + 2\mathbf{T}\mathbf{s}, \\ \mathbf{s}^{(2)} = 2t \cdot \mathbf{s} + 2\mathbf{T}^{\mathsf{T}}\mathbf{r}, \\ \mathbf{T}^{(2)} = 2t \cdot \mathbf{T} + 2|\mathbf{r}\rangle\langle \mathbf{s}| - 2\widehat{\mathbf{T}}. \end{cases}$$

• For k = 3,  $X^3 \approx (t^{(3)}, r^{(3)}, s^{(3)}, T^{(3)})$  can be identified as

$$\begin{cases} t^{(3)} &= t^3 + 3t(|\mathbf{r}|^2 + |\mathbf{s}|^2 + ||\mathbf{T}||^2) + 6(\langle \mathbf{r}|\mathbf{T}|\mathbf{s}\rangle - \det(\mathbf{T})), \\ \mathbf{r}^{(3)} &= \left(3t^2 + |\mathbf{r}|^2 + 3|\mathbf{s}|^2 + ||\mathbf{T}||^2\right)\mathbf{r} + 2\mathbf{T}\mathbf{T}^{\mathsf{T}}\mathbf{r} + 6t\mathbf{T}\mathbf{s} - 2\widehat{\mathbf{T}}\mathbf{s}, \\ \mathbf{s}^{(3)} &= \left(3t^2 + 3|\mathbf{r}|^2 + |\mathbf{s}|^2 + ||\mathbf{T}||^2\right)\mathbf{s} + 2\mathbf{T}^{\mathsf{T}}\mathbf{T}\mathbf{s} + 6t\mathbf{T}^{\mathsf{T}}\mathbf{r} - 2\widehat{\mathbf{T}}^{\mathsf{T}}\mathbf{r}, \\ \mathbf{T}^{(3)} &= \left(3t^2 + |\mathbf{r}|^2 + |\mathbf{s}|^2 + 3||\mathbf{T}||^2\right)\mathbf{T} + 6t(|\mathbf{r}\rangle\langle \mathbf{s}| - \widehat{\mathbf{T}}) \\ &+ 2\left(|\mathbf{r}\rangle\langle \mathbf{r}|\mathbf{T} + \mathbf{T}|\mathbf{s}\rangle\langle \mathbf{s}| - \mathbf{T}\mathbf{T}^{\mathsf{T}}\mathbf{T} - \Omega(\mathbf{T}, |\mathbf{r}\rangle\langle \mathbf{s}|)\right). \end{cases}$$

• For k = 4,  $X^4 \approx \left(t^{(4)}, r^{(4)}, s^{(4)}, T^{(4)}\right)$  can be identified as

$$\begin{cases} t^{(4)} &= t^4 + |\mathbf{r}|^4 + |\mathbf{s}|^4 + ||\mathbf{T}||^4 + 6|\mathbf{r}|^2|\mathbf{s}|^2 + 6t^2(|\mathbf{r}|^2 + |\mathbf{s}|^2 + ||\mathbf{T}||^2) \\ &+ 2(|\mathbf{r}|^2 + |\mathbf{s}|^2) ||\mathbf{T}||^2 + 4(\langle \mathbf{r}|\mathbf{T}\mathbf{T}^{\mathsf{T}}|\mathbf{r} \rangle + \langle \mathbf{s}|\mathbf{T}^{\mathsf{T}}\mathbf{T}|\mathbf{s} \rangle + \langle \hat{\mathbf{T}}, \hat{\mathbf{T}} \rangle) \\ &+ 24t(\langle \mathbf{r}|\mathbf{T}|\mathbf{s} \rangle - \det(\mathbf{T})) - 8\langle \mathbf{r}|\hat{\mathbf{T}}|\mathbf{s} \rangle, \\ \mathbf{r}^{(4)} &= 4\Big[ \left( t(t^2 + |\mathbf{r}|^2 + 3|\mathbf{s}|^2 + ||\mathbf{T}||^2) + 2\langle \mathbf{r}|\mathbf{T}|\mathbf{s} \rangle - 2\det(\mathbf{T}) \right) \mathbf{r} \\ &+ (3t^2 + |\mathbf{r}|^2 + |\mathbf{s}|^2 + ||\mathbf{T}||^2) \mathbf{T}\mathbf{s} + 2t\mathbf{T}\mathbf{T}^{\mathsf{T}}\mathbf{r} - 2t\hat{\mathbf{T}}\mathbf{s} \Big], \\ \mathbf{s}^{(4)} &= 4\Big[ \left( t(t^2 + 3|\mathbf{r}|^2 + |\mathbf{s}|^2 + ||\mathbf{T}||^2) + 2\langle \mathbf{r}|\mathbf{T}|\mathbf{s} \rangle - 2\det(\mathbf{T}) \right) \mathbf{s} \\ &+ (3t^2 + |\mathbf{r}|^2 + |\mathbf{s}|^2 + ||\mathbf{T}||^2) \mathbf{T}^{\mathsf{T}}\mathbf{r} + 2t\mathbf{T}^{\mathsf{T}}\mathbf{T}\mathbf{s} - 2t\hat{\mathbf{T}}^{\mathsf{T}}\mathbf{r} \Big], \\ \mathbf{T}^{(4)} &= 4\Big[ \left( t(t^2 + |\mathbf{r}|^2 + |\mathbf{s}|^2 + 3||\mathbf{T}||^2) + 2\langle \mathbf{r}|\mathbf{T}|\mathbf{s} \rangle - 2\det(\mathbf{T}) \right) \mathbf{T} \\ &+ (3t^2 + |\mathbf{r}|^2 + |\mathbf{s}|^2 + ||\mathbf{T}||^2)(|\mathbf{r}\rangle\langle \mathbf{s}| - \hat{\mathbf{T}}) \\ &+ 2t(|\mathbf{r}\rangle\langle \mathbf{r}|\mathbf{T} + \mathbf{T}|\mathbf{s}\rangle\langle \mathbf{s}| - \mathbf{T}^{\mathsf{T}}\mathbf{T} - \Omega(\mathbf{T}, |\mathbf{r}\rangle\langle \mathbf{s}|)) \Big] \end{cases}$$

For instance, we give the details in calculating  $T^{(4)}$ :

$$T^{(4)} = |\mathbf{r}^{(3)}\rangle\langle \mathbf{s}| + |\mathbf{r}\rangle\langle \mathbf{s}^{(3)}| + tT^{(3)} + t^{(3)}T - \Omega(T^{(3)}, T).$$

In what follows, we calculate it term by term:

(i) 
$$|\mathbf{r}^{(3)}\rangle\langle \mathbf{s}| = (3t^2 + |\mathbf{r}|^2 + 3|\mathbf{s}|^2 + ||\mathbf{T}||^2)|\mathbf{r}\rangle\langle \mathbf{s}| + 2\mathbf{T}\mathbf{T}^{\mathsf{T}}|\mathbf{r}\rangle\langle \mathbf{s}| + 2(3t\mathbf{T} - \widehat{\mathbf{T}})|\mathbf{s}\rangle\langle \mathbf{s}|$$

- (ii)  $|\mathbf{r}\rangle\langle s^{(3)}| = (3t^2 + 3|\mathbf{r}|^2 + |\mathbf{s}|^2 + ||\mathbf{T}||^2)|\mathbf{r}\rangle\langle s| + 2|\mathbf{r}\rangle\langle s|\mathbf{T}^{\mathsf{T}}\mathbf{T} + 2|\mathbf{r}\rangle\langle \mathbf{r}|(3t\mathbf{T} \widehat{\mathbf{T}})$
- (iii)  $tT^{(3)} = 2t(|\mathbf{r}\rangle\langle\mathbf{r}|\mathbf{T}+\mathbf{T}|\mathbf{s}\rangle\langle\mathbf{s}|) + t(3t^2 + |\mathbf{r}|^2 + |\mathbf{s}|^2 + 3 ||\mathbf{T}||^2)\mathbf{T} + 6t^2(|\mathbf{r}\rangle\langle\mathbf{s}| \widehat{\mathbf{T}}) 2t[(\mathbf{r}\cdot\mathcal{F})\mathbf{T}(\mathbf{s}\cdot\mathcal{F})^{\mathsf{T}} + \mathbf{T}\mathbf{T}^{\mathsf{T}}\mathbf{T}]$
- (iv)  $t^{(3)}T = \left[t^3 + 3t(|\mathbf{r}|^2 + |\mathbf{s}|^2 + ||\mathbf{T}||^2) + 6(\langle \mathbf{r}|\mathbf{T}|\mathbf{s}\rangle \det(\mathbf{T}))\right]T$
- (v) Now we calculate  $\Omega(T^{(3)}, T)$ . Indeed,

$$\begin{split} \Omega(T^{(3)},T) &= 2[\Omega(|\mathbf{r}\rangle\langle\mathbf{r}|T,T) + \Omega(T|s\rangle\langle s|,T)] + (3t^2 + |\mathbf{r}|^2 + |s|^2 + 3 \|T\|^2)\Omega(T,T) \\ &\quad 6t[\Omega(|\mathbf{r}\rangle\langle s|,T) - \Omega(\widehat{T},T)] - 2[\Omega((\mathbf{r}\cdot\mathcal{F})T(s\cdot\mathcal{F})^{\mathsf{T}},T) + \Omega(TT^{\mathsf{T}}T,T)] \\ &= 2[(\mathbf{r}\cdot\mathcal{F})T((T^{\mathsf{T}}\mathbf{r})\cdot\mathcal{F})^{\mathsf{T}} + ((Ts)\cdot\mathcal{F})T(s\cdot\mathcal{F})^{\mathsf{T}}] + 2(3t^2 + |\mathbf{r}|^2 + |s|^2 + 3 \|T\|^2)\widehat{T} \\ &\quad + 6t[(\mathbf{r}\cdot\mathcal{F})T(s\cdot\mathcal{F})^{\mathsf{T}} - \langle T,T\rangle T + TT^{\mathsf{T}}T] \\ &\quad - 2[\langle\mathbf{r}|T|s\rangle T - |\mathbf{r}\rangle\langle s|T^{\mathsf{T}}T + (\|T\|^2\mathbb{1}_3 - TT^{\mathsf{T}})|\mathbf{r}\rangle\langle s|] - 2[\langle T,T\rangle\widehat{T} - \det(T)T]. \end{split}$$

Thus

$$T^{(4)} = 4 \Big[ \Big( t^3 + t(|\mathbf{r}|^2 + |\mathbf{s}|^2 + 3 ||\mathbf{T}||^2) + 2 \langle \mathbf{r} | \mathbf{T} | \mathbf{s} \rangle - 2 \det(\mathbf{T}) \Big) \mathbf{T} \\ + (3t^2 + |\mathbf{r}|^2 + |\mathbf{s}|^2 + ||\mathbf{T}||^2) (|\mathbf{r}\rangle\langle \mathbf{s}| - \widehat{\mathbf{T}}) \\ + 2t (|\mathbf{r}\rangle\langle \mathbf{r} | \mathbf{T} + \mathbf{T} | \mathbf{s} \rangle\langle \mathbf{s} | - \mathbf{T} \mathbf{T}^{\mathsf{T}} \mathbf{T} - (\mathbf{r} \cdot \mathcal{F}) \mathbf{T} (\mathbf{s} \cdot \mathcal{F})^{\mathsf{T}}) \Big] \\ = 4 \Big[ \Big( t^3 + t(|\mathbf{r}|^2 + |\mathbf{s}|^2 + 3 ||\mathbf{T}||^2) + 2 \langle \mathbf{r} | \mathbf{T} | \mathbf{s} \rangle - 2 \det(\mathbf{T}) \Big) \mathbf{T} \\ + (3t^2 + |\mathbf{r}|^2 + |\mathbf{s}|^2 + ||\mathbf{T}||^2) (|\mathbf{r}\rangle\langle \mathbf{s}| - \widehat{\mathbf{T}}) \\ + 2t (|\mathbf{r}\rangle\langle \mathbf{r} | \mathbf{T} + \mathbf{T} | \mathbf{s} \rangle\langle \mathbf{s} | - \mathbf{T} \mathbf{T}^{\mathsf{T}} \mathbf{T} - \Omega(\mathbf{T}, |\mathbf{r}\rangle\langle \mathbf{s} |)) \Big].$$

## B.2 Some results about products involved two-qubit states

We have already known that

$$\boldsymbol{X}^{k} \approx \left(t^{(k)}, \boldsymbol{r}^{(k)}, \boldsymbol{s}^{(k)}, \boldsymbol{T}^{(k)}\right), \qquad (B.34)$$

$$\rho_A \otimes \mathbb{1}_B \approx \left(\frac{1}{2}, \frac{a}{2}, \mathbf{0}, \mathbf{0}\right),$$
(B.35)

$$\mathbb{1}_A \otimes \rho_B \approx \left(\frac{1}{2}, \mathbf{0}, \frac{b}{2}, \mathbf{0}\right), \qquad (B.36)$$

$$\rho_A \otimes \rho_B \approx \left(\frac{1}{4}, \frac{a}{4}, \frac{b}{4}, \frac{|a\rangle\langle b|}{4}\right)$$
(B.37)

**Proposition B.18.** Let  $\rho_{AB}^k \approx \frac{1}{4^k} \left( c^{(k)}, \boldsymbol{x}^{(k)}, \boldsymbol{y}^{(k)}, \boldsymbol{Z}^{(k)} \right)$ , where k = 2, 3, 4. We have the following results:

(*i*) For k = 2,

$$\begin{cases} c^{(2)} = 1 + |a|^{2} + |b|^{2} + ||C||^{2}, \\ x^{(2)} = 2a + 2Cb, \\ y^{(2)} = 2b + 2C^{\mathsf{T}}a, \\ Z^{(2)} = 2\left(C + |a\rangle\langle b| - \widehat{C}\right). \end{cases}$$
(B.38)

(*ii*) For k = 3,

$$\begin{cases} c^{(3)} = 1 + 3\left(|a|^{2} + |b|^{2} + ||C||^{2}\right) + 6\left(\langle a|C|b\rangle - \det(C)\right), \\ x^{(3)} = \left(3 + |a|^{2} + 3|b|^{2} + ||C||^{2}\right)a + 2CC^{\mathsf{T}}a + 6Cb - 2\widehat{C}b, \\ y^{(3)} = \left(3 + 3|a|^{2} + |b|^{2} + ||C||^{2}\right)b + 2C^{\mathsf{T}}Cb + 6C^{\mathsf{T}}a - 2\widehat{C}^{\mathsf{T}}a, \\ Z^{(3)} = \left(3 + |a|^{2} + |b|^{2} + 3||C||^{2}\right)C + 6\left(|a\rangle\langle b| - \widehat{C}\right) \\ + 2\left(|a\rangle\langle a|C + C|b\rangle\langle b| - CC^{\mathsf{T}}C - \Omega(C, |a\rangle\langle b|)\right). \end{cases}$$
(B.39)

(*iii*) For k = 4,

$$\begin{cases} c^{(4)} = 1 + 6\left(|a|^{2} + |b|^{2} + |a|^{2} |b|^{2}\right) + |a|^{4} + |b|^{4} + ||C||^{4} + 24\langle a|C|b\rangle \\ + 2||C||^{2}\left(3 + |a|^{2} + |b|^{2}\right) + 4\langle a|CC^{\mathsf{T}}|a\rangle + 4\langle b|C^{\mathsf{T}}C|b\rangle + 4\left|\widehat{C}\right|^{2} \\ - 8\langle a|\widehat{C}|b\rangle - 24\det(C), \\ x^{(4)} = 4\left(1 + |a|^{2} + 3|b|^{2} + ||C||^{2} + 2\langle a|C|b\rangle - 2\det(C)\right)a + 8CC^{\mathsf{T}}a \\ + 4\left(3 + |a|^{2} + |b|^{2} + ||C||^{2}\right)Cb - 8\widehat{C}b, \\ y^{(4)} = 4\left(1 + 3|a|^{2} + |b|^{2} + ||C||^{2}\right)C^{\mathsf{T}}a - 8\widehat{C}^{\mathsf{T}}a, \\ Z^{(4)} = 4\left(1 + |a|^{2} + |b|^{2} + ||C||^{2}\right)C^{\mathsf{T}}a - 8\widehat{C}^{\mathsf{T}}a, \\ Z^{(4)} = 4\left(1 + |a|^{2} + |b|^{2} + 3||C||^{2} + 2\langle a|C|b\rangle - 2\det(C)\right)C \\ + 4\left(3 + |a|^{2} + |b|^{2} + ||C||^{2}\right)(|a\rangle\langle b| - \widehat{C}) \\ + 8\left(|a\rangle\langle a|C + C|b\rangle\langle b| - CC^{\mathsf{T}}C - \Omega(C, |a\rangle\langle b|)\right). \end{cases}$$
(B.40)

*Proof.* The proof follows immediately when we let

$$(t, \mathbf{r}, \mathbf{s}, \mathbf{T}) = \frac{1}{4} (1, \mathbf{a}, \mathbf{b}, \mathbf{C})$$

in Corollary B.17.

Proposition B.19. Let

$$\begin{split} \boldsymbol{X}^{k}(\rho_{A}\otimes\mathbb{1}_{B}) &\approx \frac{1}{2}(\tilde{c}_{A}^{(k)}, \tilde{\boldsymbol{x}}_{A}^{(k)}, \tilde{\boldsymbol{y}}_{A}^{(k)}, \tilde{\boldsymbol{Z}}_{A}^{(k)}), \\ \boldsymbol{X}^{k}(\mathbb{1}_{A}\otimes\rho_{B}) &\approx \frac{1}{2}(\tilde{c}_{B}^{(k)}, \tilde{\boldsymbol{x}}_{B}^{(k)}, \tilde{\boldsymbol{y}}_{B}^{(k)}, \tilde{\boldsymbol{Z}}_{B}^{(k)}), \\ \boldsymbol{X}^{k}(\rho_{A}\otimes\rho_{B}) &\approx \frac{1}{4}(\tilde{c}_{AB}^{(k)}, \tilde{\boldsymbol{x}}_{AB}^{(k)}, \tilde{\boldsymbol{y}}_{AB}^{(k)}, \tilde{\boldsymbol{Z}}_{AB}^{(k)}). \end{split}$$

We have the following results:

(i) 
$$\mathbf{X}^{k}(\rho_{A} \otimes \mathbb{1}_{B}) \approx \frac{1}{2} (\tilde{c}_{A}^{(k)}, \tilde{\mathbf{x}}_{A}^{(k)}, \tilde{\mathbf{y}}_{A}^{(k)}, \tilde{\mathbf{Z}}_{A}^{(k)})$$
 is determined by  

$$\begin{cases} \tilde{c}_{A}^{(k)} = t^{(k)} + \langle \mathbf{r}^{(k)}, \mathbf{a} \rangle, \\ \tilde{\mathbf{x}}_{A}^{(k)} = \mathbf{r}^{(k)} + t^{(k)}\mathbf{a} + \mathrm{i}\mathbf{r}^{(k)} \times \mathbf{a}, \\ \tilde{\mathbf{y}}_{A}^{(k)} = \mathbf{s}^{(k)} + \mathbf{T}^{(k)^{\mathsf{T}}}\mathbf{a}, \\ \tilde{\mathbf{Z}}_{A}^{(k)} = |\mathbf{a}\rangle\langle \mathbf{s}^{(k)}| + \mathbf{T}^{(k)} - \mathrm{i}(\mathbf{a}\cdot\mathcal{F})^{\mathsf{T}}\mathbf{T}^{(k)}, \end{cases}$$
(B.41)

(ii)  $\mathbf{X}^{k}(\mathbb{1}_{A} \otimes \rho_{B}) \approx \frac{1}{2}(\tilde{c}_{B}^{(k)}, \tilde{\mathbf{x}}_{B}^{(k)}, \tilde{\mathbf{y}}_{B}^{(k)}, \tilde{\mathbf{Z}}_{B}^{(k)})$  is determined by  $\begin{cases} \tilde{c}_{B}^{(k)} = t^{(k)} + \langle \mathbf{s}^{(k)}, \mathbf{b} \rangle, \\ \tilde{\mathbf{x}}_{B}^{(k)} = \mathbf{r}^{(k)} + \mathbf{T}^{(k)}\mathbf{b}, \\ \tilde{\mathbf{y}}_{R}^{(k)} = \mathbf{s}^{(k)} + t^{(k)}\mathbf{b} + \mathbf{i}\mathbf{s}^{(k)} \end{cases}$ 

$$\begin{cases} \tilde{c}_B^{(k)} &= t^{(k)} + \langle \boldsymbol{s}^{(k)}, \boldsymbol{b} \rangle, \\ \tilde{\boldsymbol{x}}_B^{(k)} &= \boldsymbol{r}^{(k)} + \boldsymbol{T}^{(k)} \boldsymbol{b}, \\ \tilde{\boldsymbol{y}}_B^{(k)} &= \boldsymbol{s}^{(k)} + t^{(k)} \boldsymbol{b} + i \boldsymbol{s}^{(k)} \times \boldsymbol{b}, \\ \tilde{\boldsymbol{Z}}_B^{(k)} &= |\boldsymbol{r}^{(k)} \rangle \langle \boldsymbol{b} | + \boldsymbol{T}^{(k)} - i \boldsymbol{T}^{(k)} (\boldsymbol{b} \cdot \boldsymbol{\mathcal{F}}), \end{cases}$$
(B.42)

(iii)  $\mathbf{X}^{k}(\rho_{A} \otimes \rho_{B}) \approx \frac{1}{4}(\tilde{c}_{AB}^{(k)}, \tilde{\mathbf{x}}_{AB}^{(k)}, \tilde{\mathbf{y}}_{AB}^{(k)}, \tilde{\mathbf{Z}}_{AB}^{(k)})$  is determined by

$$\begin{cases} \tilde{c}_{AB}^{(k)} = t^{(k)} + \langle \mathbf{r}^{(k)}, \mathbf{a} \rangle + \langle \mathbf{s}^{(k)}, \mathbf{b} \rangle + \langle \mathbf{a} | \mathbf{T}^{(k)} | \mathbf{b} \rangle, \\ \tilde{\mathbf{x}}_{AB}^{(k)} = \mathbf{r}^{(k)} + (t^{(k)} + \langle \mathbf{s}^{(k)}, \mathbf{b} \rangle) \mathbf{a} + \mathbf{T}^{(k)} \mathbf{b} + \mathbf{i} (\mathbf{r}^{(k)} \times \mathbf{a} + \mathbf{T}^{(k)} \mathbf{b} \times \mathbf{a}), \\ \tilde{\mathbf{y}}_{AB}^{(k)} = \mathbf{s}^{(k)} + (t^{(k)} + \langle \mathbf{r}^{(k)}, \mathbf{a} \rangle) \mathbf{b} + \mathbf{T}^{(k)^{\mathsf{T}}} \mathbf{a} + \mathbf{i} (\mathbf{s}^{(k)} \times \mathbf{b} + \mathbf{T}^{(k)^{\mathsf{T}}} \mathbf{a} \times \mathbf{b}), \\ \tilde{\mathbf{Z}}_{AB}^{(k)} = |\mathbf{r}^{(k)}\rangle \langle \mathbf{b} | + |\mathbf{a}\rangle \langle \mathbf{s}^{(k)} | + t^{(k)} | \mathbf{a}\rangle \langle \mathbf{b} | + \mathbf{T}^{(k)} - \Omega(\mathbf{T}^{(k)}, |\mathbf{a}\rangle \langle \mathbf{b} |) \\ + \mathbf{i} \left( \Psi(\mathbf{r}^{(k)}, |\mathbf{a}\rangle \langle \mathbf{b} |, \mathbf{s}^{(k)}) - \Psi(\mathbf{a}, \mathbf{T}^{(k)}, \mathbf{b}) \right). \end{cases}$$
(B.43)

Proposition B.20. Let

$$\begin{array}{lll}
\rho_{AB}^{k}(\rho_{A}\otimes\mathbb{1}_{B}) &\approx & \frac{1}{2\cdot4^{k}}(c_{A}^{(k)},\boldsymbol{x}_{A}^{(k)},\boldsymbol{y}_{A}^{(k)},\boldsymbol{Z}_{A}^{(k)}), \\
\rho_{AB}^{k}(\mathbb{1}_{A}\otimes\rho_{B}) &\approx & \frac{1}{2\cdot4^{k}}(c_{B}^{(k)},\boldsymbol{x}_{B}^{(k)},\boldsymbol{y}_{B}^{(k)},\boldsymbol{Z}_{B}^{(k)}), \\
\rho_{AB}^{k}(\rho_{A}\otimes\rho_{B}) &\approx & \frac{1}{4^{k+1}}(c_{AB}^{(k)},\boldsymbol{x}_{AB}^{(k)},\boldsymbol{y}_{AB}^{(k)},\boldsymbol{Z}_{AB}^{(k)}).
\end{array}$$

Then we get the following statements:

(i) For k = 1, it holds that

$$\begin{cases} c_A^{(1)} &= 1 + |a|^2, \\ x_A^{(1)} &= 2a, \\ y_A^{(1)} &= b + C^{\mathsf{T}}a, \\ Z_A^{(1)} &= C + |a\rangle\langle b| - \mathrm{i}(a \cdot \mathcal{F})^{\mathsf{T}}C, \end{cases} \begin{cases} c_B^{(1)} &= 1 + |b|^2, \\ x_B^{(1)} &= 1 + |b|^2, \\ x_B^{(1)} &= a + Cb, \\ y_B^{(1)} &= 2b, \\ Z_B^{(1)} &= C + |a\rangle\langle b| - \mathrm{i}C(b \cdot \mathcal{F}). \end{cases}$$

and

$$\begin{cases} c_{AB}^{(1)} &= 1 + |a|^2 + |b|^2 + \langle a|C|b \rangle \\ x_{AB}^{(1)} &= (2 + |b|^2)a + Cb + iCb \times a \\ y_{AB}^{(1)} &= (2 + |a|^2)b + C^{\mathsf{T}}a + iC^{\mathsf{T}}a \times b \\ Z_{AB}^{(1)} &= C + 3|a \rangle \langle b| - \Omega(C, |a \rangle \langle b|) - i\Psi(a, C, b). \end{cases}$$

(*ii*) For k = 2, it holds that

$$\begin{cases} c_A^{(2)} &= 1+3 |a|^2 + |b|^2 + ||C||^2 + 2\langle a|C|b \rangle, \\ x_A^{(2)} &= (3+|a|^2+|b|^2+||C||^2)a + 2Cb + 2iCb \times a, \\ y_A^{(2)} &= 2(1+|a|^2)b + 4C^{\mathsf{T}}a - 2\widehat{C}^{\mathsf{T}}a, \\ Z_A^{(2)} &= 2(C-\widehat{C}) + 2|a\rangle\langle a|C+4|a\rangle\langle b| - 2i(a \cdot \mathcal{F})^{\mathsf{T}}(C-\widehat{C}), \end{cases}$$

$$\begin{cases} c_B^{(2)} &= 1 + |a|^2 + 3 |b|^2 + ||C||^2 + 2\langle a |C|b \rangle, \\ x_B^{(2)} &= 2(1 + |b|^2)a + 4Cb - 2\widehat{C}b, \\ y_B^{(2)} &= (3 + |a|^2 + |b|^2 + ||C||^2)b + 2C^{\mathsf{T}}a + 2\mathrm{i}C^{\mathsf{T}}a \times b, \\ Z_B^{(2)} &= 2(C - \widehat{C}) + 2C|b \rangle \langle b| + 4|a \rangle \langle b| - 2\mathrm{i}(C - \widehat{C})(b \cdot \mathcal{F}). \end{cases}$$

and

$$\begin{cases} c_{AB}^{(2)} &= 1+3 |a|^2 + 3 |b|^2 + 2 |a|^2 |b|^2 + ||C||^2 + 6\langle a|C|b \rangle - 2\langle a|\widehat{C}|b \rangle, \\ x_{AB}^{(2)} &= (3+|a|^2+5|b|^2+||C||^2+2\langle a|C|b \rangle)a + 2(2Cb-\widehat{C}b) + 2i(2Cb-\widehat{C}b) \times a, \\ y_{AB}^{(2)} &= (3+5|a|^2+|b|^2+||C||^2+2\langle a|C|b \rangle)b + 2(2C^{\mathsf{T}}a-\widehat{C}^{\mathsf{T}}a) + 2i(2C^{\mathsf{T}}a-\widehat{C}^{\mathsf{T}}a) \times b, \\ Z_{AB}^{(2)} &= 2(C-\widehat{C}) + 2(|a\rangle\langle a|C+C|b\rangle\langle b|) + (7+|a|^2+|b|^2+||C||^2)|a\rangle\langle b| \\ &- 2\Omega(C-\widehat{C},|a\rangle\langle b|) + 2i(\Psi(Cb,|a\rangle\langle b|,C^{\mathsf{T}}a) - \Psi(a,C-\widehat{C},b)). \end{cases}$$

(iii) For k = 3, it holds that

$$\begin{cases} c_A^{(3)} &= 1 + 6 |a|^2 + |a|^4 + 3 |b|^2 (1 + |a|^2) + (3 + |a|^2) ||C||^2 + 12 \langle a|C|b \rangle + 2 \langle a|CC^{\mathsf{T}}|a \rangle \\ &- 6 \det(C) - 2 \langle a|\widehat{C}|b \rangle, \\ x_A^{(3)} &= (4 + 4 |a|^2 + 6 |b|^2 + 4 ||C||^2 + 6 \langle a|C|b \rangle - 6 \det(C))a + 2(CC^{\mathsf{T}}a + 3Cb - \widehat{C}b) \\ &+ 2i(CC^{\mathsf{T}}a + 3Cb - \widehat{C}b) \times a, \\ y_A^{(3)} &= (3 + 9 |a|^2 + |b|^2 + ||C||^2 + 2 \langle a|C|b \rangle)b + 2C^{\mathsf{T}}Cb + (9 + 3 |a|^2 + |b|^2 + 3 ||C||^2)C^{\mathsf{T}}a \\ &- 2C^{\mathsf{T}}CC^{\mathsf{T}}a - 8\widehat{C}^{\mathsf{T}}a, \\ Z_A^{(3)} &= (3 + |a|^2 + |b|^2 + 3 ||C||^2)C - 6\widehat{C} + 2|a\rangle\langle a|(4C - \widehat{C}) + 2C|b\rangle\langle b| + 2|a\rangle\langle b|C^{\mathsf{T}}C \\ &+ (9 + 3 |a|^2 + |b|^2 + ||C||^2)|a\rangle\langle b| - 2CC^{\mathsf{T}}C - 2\Omega(C, |a\rangle\langle b|) \\ &- i(a \cdot \mathcal{F})^{\mathsf{T}} \left[ (3 + |a|^2 + |b|^2 + 3 ||C||^2)C - 6\widehat{C} + 2(C|b\rangle\langle b| - CC^{\mathsf{T}}C - \Omega(C, |a\rangle\langle b|)) \right], \end{cases}$$

$$\begin{cases} c_B^{(3)} &= 1 + 6 |b|^2 + |b|^4 + 3 |a|^2 (1 + |b|^2) + (3 + |b|^2) ||C||^2 + 12 \langle a|C|b \rangle + 2 \langle b|C^{\mathsf{T}}C|b \rangle \\ &- 6 \det(C) - 2 \langle a|\widehat{C}|b \rangle, \\ x_B^{(3)} &= (3 + |a|^2 + 9 |b|^2 + ||C||^2 + 2 \langle a|C|b \rangle) a + 2CC^{\mathsf{T}}a + (9 + |a|^2 + 3 |b|^2 + 3 ||C||^2) Cb \\ &- 2CC^{\mathsf{T}}Cb - 8\widehat{C}b, \\ y_B^{(3)} &= (4 + 6 |a|^2 + 4 |b|^2 + 4 ||C||^2 + 6 \langle a|C|b \rangle - 6 \det(C))b + 2(C^{\mathsf{T}}Cb + 3C^{\mathsf{T}}a - \widehat{C}^{\mathsf{T}}a) \\ &+ 2i(C^{\mathsf{T}}Cb + 3C^{\mathsf{T}}a - \widehat{C}^{\mathsf{T}}a) \times b, \\ Z_B^{(3)} &= (3 + |a|^2 + |b|^2 + 3 ||C||^2)C - 6\widehat{C} + 2(4C - \widehat{C})|b\rangle\langle b| + 2|a\rangle\langle a|C + 2CC^{\mathsf{T}}|a\rangle\langle b| \\ &+ (9 + |a|^2 + 3 |b|^2 + ||C||^2)|a\rangle\langle b| - 2CC^{\mathsf{T}}C - 2\Omega(C, |a\rangle\langle b|) \\ &- i \left[ (3 + |a|^2 + |b|^2 + 3 ||C||^2)C - 6\widehat{C} + 2(|a\rangle\langle a|C - CC^{\mathsf{T}}C - \Omega(C, |a\rangle\langle b|)) \right] (b \cdot \mathcal{F}), \end{cases}$$

and

$$\begin{cases} c_{AB}^{(3)} &= 1 + 6(|a|^{2} + |b|^{2}) + 12|a|^{2}|b|^{2} + |a|^{4} + |b|^{4} + (3 + |a|^{2} + |b|^{2}) ||C||^{2} \\ &+ 3(7 + |a|^{2} + |b|^{2} + ||C||^{2})\langle a|C|b\rangle - 10\langle a|\widehat{C}|b\rangle - 6\det(C) \\ &+ 2(\langle a|CC^{\mathsf{T}}|a\rangle + \langle b|C^{\mathsf{T}}C|b\rangle - \langle a|CC^{\mathsf{T}}C|b\rangle), \\ x_{AB}^{(3)} &= \left(4 + 4|a|^{2} + 15|b|^{2} + |b|^{4} + 4||C||^{2} + (3|a|^{2} + ||C||^{2})|b|^{2} + 14\langle a|C|b\rangle + 2\langle b|C^{\mathsf{T}}C|b\rangle \\ &- 2\langle a|\widehat{C}|b\rangle - 6\det(C)\right)a + 2CC^{\mathsf{T}}a + (9 + |a|^{2} + 3|b|^{2} + 3||C||^{2})Cb - 8\widehat{C}b - 2CC^{\mathsf{T}}Cb \\ &+ i\left(2CC^{\mathsf{T}}a + (9 + |a|^{2} + 3|b|^{2} + 3||C||^{2})Cb - 8\widehat{C}b - 2CC^{\mathsf{T}}Cb\right) \times a, \\ y_{AB}^{(3)} &= \left(4 + 15|a|^{2} + |a|^{4} + 4|b|^{2} + 4||C||^{2} + (3|b|^{2} + ||C||^{2})|a|^{2} + 14\langle a|C|b\rangle + 2\langle a|CC^{\mathsf{T}}|a\rangle \\ &- 2\langle a|\widehat{C}|b\rangle - 6\det(C)\right)b + 2C^{\mathsf{T}}Cb + (9 + 3|a|^{2} + |b|^{2} + 3||C||^{2})C^{\mathsf{T}}a - 8\widehat{C}^{\mathsf{T}}a - 2C^{\mathsf{T}}CC^{\mathsf{T}}a \\ &+ i\left(2C^{\mathsf{T}}b + (9 + 3|a|^{2} + |b|^{2} + 3||C||^{2})C^{\mathsf{T}}a - 8\widehat{C}^{\mathsf{T}}a - 2C^{\mathsf{T}}CC^{\mathsf{T}}a \\ &+ i\left(2C^{\mathsf{T}}Cb + (9 + 3|a|^{2} + |b|^{2} + 3||C||^{2})C^{\mathsf{T}}a - 8\widehat{C}^{\mathsf{T}}a - 2C^{\mathsf{T}}CC^{\mathsf{T}}a \\ &+ i\left(2C^{\mathsf{T}}b + (9 + 3|a|^{2} + |b|^{2} + 3||C||^{2})C^{\mathsf{T}}a - 8\widehat{C}^{\mathsf{T}}a - 2C^{\mathsf{T}}CC^{\mathsf{T}}a \\ &+ i\left(2C^{\mathsf{T}}b + (9 + 3|a|^{2} + |b|^{2} + 3||C||^{2})C^{\mathsf{T}}a - 8\widehat{C}^{\mathsf{T}}a - 2C^{\mathsf{T}}CC^{\mathsf{T}}a \\ &+ i\left(2C^{\mathsf{T}}b + (9 + 3|a|^{2} + |b|^{2} + 3||C||^{2})C^{\mathsf{T}}a - 8\widehat{C}^{\mathsf{T}}a - 2C^{\mathsf{T}}CC^{\mathsf{T}}a \\ &+ i\left(2C^{\mathsf{T}}b + (9 + 3|a|^{2} + |b|^{2} + 3||C||^{2})C^{\mathsf{T}}a - 8\widehat{C}^{\mathsf{T}}a - 2C^{\mathsf{T}}CC^{\mathsf{T}}a \\ &+ \left(13 + 7(|a|^{2} + |b|^{2} + 3||C||^{2})C^{\mathsf{T}} - 2C^{\mathsf{T}}a - 6\widehat{C}^{\mathsf{T}}a - 2C^{\mathsf{T}}b\rangle \langle b| \\ &+ \left(13 + 7(|a|^{2} + |b|^{2} + 3||C||^{2})C - 2CC^{\mathsf{T}}C - 6\widehat{C}, |a\rangle \langle b| \right) \\ &- \Omega((5 + |a|^{2} + |b|^{2} + 3||C||^{2})C - 2CC^{\mathsf{T}}Cb + 6C^{\mathsf{T}}a - 2\widehat{C}^{\mathsf{T}}a) \\ &- i\left((3 + |a|^{2} + |b|^{2} + 3||C||^{2})\Psi(a, C, b) - 6\Psi(a, \widehat{C}, b) - 2\Psi(a, CC^{\mathsf{T}}C, b)\right) \\ &+ 2i\left(|b|^{2}(a \cdot \mathcal{F})C + |a|^{2}C(b \cdot \mathcal{F})^{\mathsf{T}}\right).$$

#### **B.3** Revisiting local unitary invariants

For any two-qubit state  $\rho_{AB}$ , decomposed as

$$\rho_{AB} = \frac{1}{4} \left( \mathbb{1} \otimes \mathbb{1} + \boldsymbol{a} \cdot \boldsymbol{\sigma} \otimes \mathbb{1} + \mathbb{1} \otimes \boldsymbol{b} \cdot \boldsymbol{\sigma} + \sum_{i,j=1}^{3} c_{ij} \sigma_i \otimes \sigma_j \right),$$
(B.44)

where  $\mathbf{a} = (a_1, a_2, a_3)^{\mathsf{T}}$  and  $\mathbf{b} = (b_1, b_2, b_3)^{\mathsf{T}}$  are in  $\mathbb{R}^3$ , and  $\mathbf{C} = (c_{ij})_{3\times 3} \in \mathbb{R}^{3\times 3}$ . Its two reduced states are given by, respectively  $\rho_A = \frac{1}{2}(\mathbb{1}_2 + \mathbf{a} \cdot \boldsymbol{\sigma})$  and  $\rho_B = \frac{1}{2}(\mathbb{1}_2 + \mathbf{b} \cdot \boldsymbol{\sigma})$ . In 2002, Makhlin had published the following well-known result<sup>1</sup>:

**Proposition B.21** ([6]). *For any mixed two-qubit states*  $\rho_{AB}$ ,  $\rho'_{AB} \in D(\mathbb{C}^2 \otimes \mathbb{C}^2)$ , *both are LU equivalent* 

<sup>&</sup>lt;sup>1</sup>Here we reformulate those 18 LU invariants for our convenience. They are also termed Makhlin's invariants.

if and only if the following 18-tuple  $(I_1, \ldots, I_{18})$  are the same for both  $\rho_{AB}$  and  $\rho'_{AB}$ , where

$$I_{1} = \det(C), I_{2} = \langle C, C \rangle, I_{3} = \langle C^{\mathsf{T}}C, C^{\mathsf{T}}C \rangle,$$

$$I_{4} = \langle a, a \rangle, I_{5} = \langle a | CC^{\mathsf{T}} | a \rangle, I_{6} = \langle a | (CC^{\mathsf{T}})^{2} | a \rangle,$$

$$I_{7} = \langle b, b \rangle, I_{8} = \langle b | C^{\mathsf{T}}C | b \rangle, I_{9} = \langle b | (C^{\mathsf{T}}C)^{2} | b \rangle,$$

$$I_{10} = a \cdot (CC^{\mathsf{T}}a \times (CC^{\mathsf{T}})^{2}a), I_{11} = b \cdot (C^{\mathsf{T}}Cb \times (C^{\mathsf{T}}C)^{2}b),$$

$$I_{12} = \langle a | C | b \rangle, I_{13} = \langle a | CC^{\mathsf{T}}C | b \rangle, I_{14} = \langle (a \cdot \mathcal{F})C, C(b \cdot \mathcal{F}) \rangle,$$

$$I_{15} = a \cdot (CC^{\mathsf{T}}a \times Cb), I_{16} = C^{\mathsf{T}}a \cdot (b \times C^{\mathsf{T}}Cb),$$

$$I_{17} = C^{\mathsf{T}}a \cdot (C^{\mathsf{T}}CC^{\mathsf{T}}a \times b), I_{18} = a \cdot (Cb \times CC^{\mathsf{T}}Cb).$$

Here we deliberately omit the constant factor in Makhlin's invariants. For our purposes, we will give another 18-tuple of invariants in replacement of Makhlin's invariants.

**Proposition B.22** ([6]). For any mixed two-qubit states  $\rho_{AB}$ ,  $\rho'_{AB} \in D(\mathbb{C}^2 \otimes \mathbb{C}^2)$ , both are LU equivalent if and only if the following 18-tuple  $(L_1, \ldots, L_{18})$  are the same for both  $\rho_{AB}$  and  $\rho'_{AB}$ , where

$$L_{1} = \det(C), L_{2} = \langle C, C \rangle, L_{3} = \langle \widehat{C}, \widehat{C} \rangle,$$

$$L_{4} = \langle a, a \rangle, L_{5} = \langle a | CC^{\mathsf{T}} | a \rangle, L_{6} = \langle a | \widehat{CC^{\mathsf{T}}} | a \rangle,$$

$$L_{7} = \langle b, b \rangle, L_{8} = \langle b | C^{\mathsf{T}} C | b \rangle, L_{9} = \langle b | \widehat{C^{\mathsf{T}} C} | b \rangle,$$

$$L_{10} = a \cdot (CC^{\mathsf{T}} a \times \widehat{CC^{\mathsf{T}}} a), L_{11} = b \cdot (C^{\mathsf{T}} Cb \times \widehat{C^{\mathsf{T}}} Cb),$$

$$L_{12} = \langle a | C | b \rangle, L_{13} = \langle a | CC^{\mathsf{T}} C | b \rangle, L_{14} = \langle a | \widehat{C} | b \rangle,$$

$$L_{15} = b \cdot (C^{\mathsf{T}} a \times \widehat{C^{\mathsf{T}}} a), L_{16} = a \cdot (Cb \times \widehat{C}b),$$

$$L_{17} = \widehat{C} b \cdot (a \times CC^{\mathsf{T}} a), L_{18} = \widehat{C}^{\mathsf{T}} a \cdot (b \times C^{\mathsf{T}} Cb).$$

*Proof.* Note that we can find out the following relations

- (1)  $I_k = L_k$ , where  $k \in \{1, 2, 4, 5, 7, 8, 10, 11, 12, 13, 14, 17, 18\}$
- (2)  $I_3 = L_2^2 2L_3$
- (3)  $I_6 = L_6 + L_2 L_5 L_3 L_4$
- $(4) I_9 = L_9 + L_2 L_8 L_3 L_7$
- (5)  $I_k = -L_k$ , where  $k \in \{15, 16\}$

Indeed, the first one is trivial. For the 2nd item, note that  $2\langle \hat{C}, \hat{C} \rangle = \langle C, C \rangle^2 - \langle C^{\mathsf{T}}C, C^{\mathsf{T}}C \rangle$ . This implies that the desired result. For the 3rd item,

$$\widehat{CC}^{\mathsf{T}} = (CC^{\mathsf{T}})^2 - \langle C, C \rangle CC^{\mathsf{T}} + \langle \widehat{C}, \widehat{C} \rangle \mathbb{1}_3$$

implying that

$$\langle a | \widehat{CC^{\mathsf{T}}} | a \rangle = \langle a | (CC^{\mathsf{T}})^2 | a \rangle - \langle C, C \rangle \langle a | CC^{\mathsf{T}} | a \rangle + \langle \widehat{C}, \widehat{C} \rangle \langle a | a \rangle.$$

That is,

$$L_6 = I_6 - L_2 L_5 + L_3 L_4. \tag{B.45}$$

For the 4th item,

$$\widehat{C^{\mathsf{T}}C} = (C^{\mathsf{T}}C)^2 - \langle C, C \rangle C^{\mathsf{T}}C + \langle \widehat{C}, \widehat{C} \rangle \mathbb{1}_3$$

implying that

$$\langle \boldsymbol{b} | \widehat{\boldsymbol{C}^{\mathsf{T}} \boldsymbol{C}} | \boldsymbol{b} \rangle = \langle \boldsymbol{b} | (\boldsymbol{C}^{\mathsf{T}} \boldsymbol{C})^2 | \boldsymbol{b} \rangle - \langle \boldsymbol{C}, \boldsymbol{C} \rangle \langle \boldsymbol{b} | \boldsymbol{C}^{\mathsf{T}} \boldsymbol{C} | \boldsymbol{b} \rangle + \langle \widehat{\boldsymbol{C}}, \widehat{\boldsymbol{C}} \rangle \langle \boldsymbol{b} | \boldsymbol{b} \rangle.$$
(B.46)

That is,

$$L_9 = I_9 - L_2 L_8 + L_3 L_7. \tag{B.47}$$

For the equality of  $I_{10/11} = L_{10/11}$ 

$$(CC^{\mathsf{T}})^2 a = \widehat{CC^{\mathsf{T}}} a + \langle C, C \rangle CC^{\mathsf{T}} a - \langle \widehat{C}, \widehat{C} \rangle a.$$

Then

$$CC^{\mathsf{T}}a \times (CC^{\mathsf{T}})^2 a = CC^{\mathsf{T}}a \times \widehat{CC^{\mathsf{T}}}a - \langle \widehat{C}, \widehat{C} \rangle CC^{\mathsf{T}}a \times a,$$

implying that

$$I_{10} = \boldsymbol{a} \cdot (\boldsymbol{C}\boldsymbol{C}^{\mathsf{T}}\boldsymbol{a} \times (\boldsymbol{C}\boldsymbol{C}^{\mathsf{T}})^{2}\boldsymbol{a}) = \boldsymbol{a} \cdot \left(\boldsymbol{C}\boldsymbol{C}^{\mathsf{T}}\boldsymbol{a} \times \widehat{\boldsymbol{C}\boldsymbol{C}^{\mathsf{T}}}\boldsymbol{a}\right) = L_{10}.$$

$$(C^{\mathsf{T}}C)^{2}b = \widehat{C^{\mathsf{T}}C}b + \langle C, C \rangle C^{\mathsf{T}}Cb - \langle \widehat{C}, \widehat{C} \rangle b.$$

Then

$$C^{\mathsf{T}}Cb \times (C^{\mathsf{T}}C)^{2}b = C^{\mathsf{T}}Cb \times \widehat{C^{\mathsf{T}}Cb} - \langle \widehat{C}, \widehat{C} \rangle C^{\mathsf{T}}Cb \times b,$$

implying that

$$I_{11} = \boldsymbol{b} \cdot (\boldsymbol{C}^{\mathsf{T}} \boldsymbol{C} \boldsymbol{b} \times (\boldsymbol{C}^{\mathsf{T}} \boldsymbol{C})^{2} \boldsymbol{b}) = \boldsymbol{b} \cdot \left(\boldsymbol{C}^{\mathsf{T}} \boldsymbol{C} \boldsymbol{b} \times \widehat{\boldsymbol{C}^{\mathsf{T}} \boldsymbol{C} \boldsymbol{b}}\right) = L_{11}.$$

For the 5th item,

$$I_{15} = a \cdot (CC^{\mathsf{T}}a \times Cb) = \langle a, CC^{\mathsf{T}}a \times Cb \rangle = \langle a, \widehat{C}(C^{\mathsf{T}}a \times b) \rangle$$
$$= \langle \widehat{C}^{\mathsf{T}}a, C^{\mathsf{T}}a \times b \rangle = b \cdot (\widehat{C}^{\mathsf{T}}a \times C^{\mathsf{T}}a) = -b \cdot (C^{\mathsf{T}}a \times \widehat{C}^{\mathsf{T}}a) = -L_{15}$$

Similarly, we get that  $I_{16} = -L_{16}$ . Indeed,

$$I_{16} = C^{\mathsf{T}} a \cdot (b \times C^{\mathsf{T}} C b) = b \cdot (C^{\mathsf{T}} C b \times C^{\mathsf{T}} a) = \langle b, C^{\mathsf{T}} C b \times C^{\mathsf{T}} a \rangle$$
$$= \langle b, \widehat{C}^{\mathsf{T}} (C b \times a) \rangle = \langle \widehat{C} b, (C b \times a) \rangle = a \cdot (\widehat{C} b \times C b)$$
$$= -a \cdot (C b \times \widehat{C} b) = -L_{16}.$$

We also note that

$$I_{17} = b \cdot (C^{\mathsf{T}} a \times C^{\mathsf{T}} C C^{\mathsf{T}} a) = \langle b, \widehat{C}^{\mathsf{T}} (a \times C C^{\mathsf{T}} a) \rangle = \langle \widehat{C} b, a \times C C^{\mathsf{T}} a \rangle = L_{17}$$

and

$$I_{18} = a \cdot (Cb \times CC^{\mathsf{T}}Cb) = \left\langle a, \widehat{C}(b \times C^{\mathsf{T}}Cb) \right\rangle = \left\langle \widehat{C}^{\mathsf{T}}a, b \times C^{\mathsf{T}}Cb \right\rangle = L_{18}.$$

From the above discussion, we can see that the invariant ring generated by 18 Makhlin's invariants  $I_k$  (k = 1, ..., 18) can also be generated by our proposed 18 invariants  $L_k$  (k = 1, ..., 18).

Based on this observation, we can infer the following results:

**Lemma B.23.** For any two-qubit state  $\rho_{AB}$  decomposed as in Eq. (B.44) above, let  $X_0 = \rho_{AB}, X_1 = \rho_A \otimes \mathbb{1}_2$ , and  $X_2 = \mathbb{1}_2 \otimes \rho_B$ , it holds that

(1)  $B_1 = \text{Tr} (X_0 X_1) = \frac{1+L_4}{2}$ . (2)  $B_2 = \text{Tr} (X_0 X_2) = \frac{1+L_7}{2}$ . (3)  $B_3 = \text{Tr} (X_0 X_1 X_2) = \frac{1+L_4+L_7+L_{12}}{4}$ . (4)  $B_4 = \text{Tr} (X_0^2) = \frac{1+L_2+L_4+L_7}{4}$ . (5)  $B_5 = \text{Tr} (X_0^2 X_1 X_2) = \frac{1+L_2+3L_4+3L_7+2L_4L_7+6L_{12}-2L_{14}}{16}$ . (6)  $B_6 = \text{Tr} (X_0^3) = \frac{1-6L_1+3L_2+3L_4+3L_7+6L_{12}}{16}$ . (7)  $B_7 = \text{Tr} (X_0^3 X_1) = \frac{1-6L_1+L_2(3+L_4)+6L_4+L_4^2+2L_5+3(1+L_4)L_7+12L_{12}-2L_{14}}{32}$ . (8)  $B_8 = \text{Tr} (X_0^3 X_2) = \frac{1-6L_1+L_2(3+L_4)+6L_7+L_7^2+2L_8+3(1+L_7)L_4+12L_{12}-2L_{14}}{32}$ . (9)  $B_9 = \text{Tr} (X_0^3 X_1 X_2)$  is given by

$$B_{9} = \frac{1}{64} \Big[ 1 + 6(L_{4} + L_{7}) + 12L_{4}L_{7} + L_{4}^{2} + L_{7}^{2} + (3 + L_{4} + L_{7})L_{2} \\ + 3(7 + L_{2} + L_{4} + L_{7})L_{12} + 2(L_{5} + L_{8}) - 6L_{1} - 2L_{13} - 10L_{14} \Big].$$

(10)  $B_{10} = \text{Tr} \left( \mathbf{X}_{0}^{4} \right)$  is given by

$$B_{10} = \frac{1}{64} \Big[ 1 + 6 \left( L_4 + L_7 + L_4 L_7 \right) + L_4^2 + L_7^2 + \left( 6 + L_2 + 2L_4 + 2L_7 \right) L_2 + 24L_{12} + 4 \left( L_3 + L_5 + L_8 - 2L_{14} - 6L_1 \right) \Big].$$

(11)  $B_{11} = \text{Tr} \left( X_0^2 X_1 X_0^2 X_1 \right)$  is given by

$$B_{11} = \frac{1}{256} \Big[ 8L_{12}^2 + 8L_{12}(6 + 6L_4 + L_7 + L_2) + 4(7 + L_4)L_5 - 8(3 + L_4)L_{14} + 8L_6 + 4(1 - L_4)L_8 \\ - 8(3 + L_4)L_1 + 4(1 - L_4)L_3 + (1 + L_4)L_2^2 + 2(1 + L_4)(3 + L_4 + L_7)L_2 \\ + (1 + 15L_4 + 15L_4^2 + L_4^3 + 6L_7 + 36L_4L_7 + 6L_4^2L_7 + L_7^2 + L_4L_7^2) \Big].$$

(12) 
$$B_{12} = \text{Tr} \left( X_0^2 X_2 X_0^2 X_2 \right)$$
 is given by

$$B_{12} = \frac{1}{256} \Big[ 8L_{12}^2 + 8L_{12}(6 + 6L_7 + L_4 + L_2) + 4(7 + L_7)L_8 - 8(3 + L_7)L_{14} + 8L_9 + 4(1 - L_7)L_5 - 8(3 + L_7)L_1 + 4(1 - L_7)L_3 + (1 + L_7)L_2^2 + 2(1 + L_7)(3 + L_4 + L_7)L_2 + (1 + 15L_7 + 15L_7^2 + L_7^3 + 6L_4 + 36L_4L_7 + 6L_7^2L_4 + L_4^2 + L_7L_4^2) \Big].$$

(13)  $B_{13} = \text{Tr} \left( X_0 X_1 X_2 X_0^2 X_1 \right)$  is given by

$$B_{13} = \frac{1}{128} \Big[ 4L_{12}^2 + L_{12}(30 + 6L_7 + 18L_4 + 2L_2) + (3 + L_4 + L_7 + L_4L_7)L_2 + 2(1 - L_4)L_8 + 8L_5 - 2(5 + L_4)L_{14} - 2(3 - L_4)L_1 + 4iL_{15} + (1 + 6L_7 + L_7^2 + 10L_4 + 27L_4L_7 + L_7^2L_4 + 5L_4^2 + 3L_7L_4^2) \Big].$$

(14)  $B_{14} = \text{Tr} \left( X_0 X_1 X_2 X_0^2 X_2 \right)$  is given by

$$B_{14} = \frac{1}{128} \Big[ 4L_{12}^2 + L_{12}(30 + 6L_4 + 18L_7 + 2L_2) + (3 + L_4 + L_7 + L_4L_7)L_2 + 2(1 - L_7)L_5 + 8L_8 - 2(5 + L_7)L_{14} - 2(3 - L_7)L_1 + 4iL_{16} + (1 + 6L_4 + L_4^2 + 10L_7 + 27L_4L_7 + L_4^2L_7 + 5L_7^2 + 3L_4L_7^2) \Big].$$

(15)  $B_{15} = \text{Tr} \left( X_0 X_1 X_2 X_0^3 X_1 \right)$  is given by

$$B_{15} = \frac{1}{512} \Big[ 1 + L_4^3 + 26L_7L_4^2 + 15L_4^2 + 13L_7^2L_4 + 76L_4L_7 + 15L_4 + 5L_7^2 + 4L_3 + 10L_7 - L_2^2(L_4 - 1) \\ + 26L_5 + 12L_8 + 6(L_4 + L_7)L_5 - 4L_6 + 68L_{12} + 88L_4L_{12} + 4L_4^2L_{12} + 4L_5L_{12} + 44L_7L_{12} \\ + 12L_4L_7L_{12} + 28L_{12}^2 - 4L_1(6 + 2L_7 + L_4(L_7 + 4) + 3L_{12}) \\ + 2L_2(3 + L_5 + 3L_7 + 12L_{12} + L_4(4L_7 + 2L_{12} + 5)) \\ - 4L_4L_{13} - 4L_{13} - 12L_4L_{14} - 4L_7L_{14} - 4L_{12}L_{14} - 44L_{14} + i(16L_{15} - 4L_{16} + 4L_{17}) \Big].$$

## (16) $B_{16} = \text{Tr} \left( X_0 X_1 X_2 X_0^3 X_2 \right)$ is given by

$$B_{16} = \frac{1}{512} \Big[ 1 + L_7^3 + 26L_4L_7^2 + 15L_7^2 + 13L_4^2L_7 + 76L_4L_7 + 15L_7 + 5L_4^2 + 4L_3 + 10L_4 - L_2^2(L_7 - 1) \\ + 26L_8 + 12L_5 + 6(L_4 + L_7)L_8 - 4L_9 + 68L_{12} + 88L_7L_{12} + 4L_7^2L_{12} + 4L_8L_{12} + 44L_4L_{12} \\ + 12L_4L_7L_{12} + 28L_{12}^2 - 4L_1(6 + 2L_4 + L_7(L_4 + 4) + 3L_{12}) \\ + 2L_2(3 + L_8 + 3L_4 + 12L_{12} + L_7(4L_4 + 2L_{12} + 5)) \\ - 4L_7L_{13} - 4L_{13} - 12L_7L_{14} - 4L_4L_{14} - 4L_{12}L_{14} - 44L_{14} + i(16L_{16} - 4L_{15} + 4L_{18}) \Big].$$

(17)  $B_{17} = \text{Tr} \left( X_0 X_1 X_0^2 X_1 X_0^3 X_1 \right)$  is given by

$$\begin{split} B_{17} &= \frac{1}{8192} \Big[ 1 + L_7^3 + 15L_7^2 + 12L_3L_7 + 15L_7 + 48L_1^2 + 60L_3 - L_2^3(L_4 - 1) + 36L_4 + 48L_3L_4 + 315L_4L_7 \\ &\quad + 3L_4L_7^3 + 150L_4L_7^2 + 75L_4^2L_7^2 + 126L_4^2 - 12L_3L_4^2 + 525L_4^2L_7 + 9L_4^4 + 84L_4^3 + 105L_4^3L_7 \\ &\quad + 60L_8 + 48L_4L_8 + 12L_7L_8 - 4L_4L_7L_8 - 4L_9L_4 - 12L_8L_4^2 + 4L_5L_7^2 + 224L_5 + 132L_5L_7 \\ &\quad + 448L_4L_5 - 4L_5L_8 + 108L_4L_7L_5 + 96L_4^2L_5 + 32L_5^2 + 8L_4L_6 - 12L_7L_6 + 24L_6 \\ &\quad + 24L_3L_{12} + 300L_7L_{12} + 210L_{12} + 18L_{12}L_7^2 + 1050L_4L_{12} + 24L_8L_{12} + 600L_4L_7L_{12} - 8L_3L_4L_{12} \\ &\quad + 30L_4^3L_{12} + 60L_4^2L_7L_{12} + 630L_4^2L_{12} + 6L_4L_7^2L_{12} - 8L_4L_8L_{12} + 336L_5L_{12} + 48L_4L_5L_{12} \\ &\quad - 16L_6L_{12} + 8L_5L_7L_{12} + 16L_{12}^3 + 552L_{12}^2 + 312L_4L_{12}^2 + 40L_7L_{12}^2 + 8L_{12}L_{13} \\ &\quad + L_2^2(4L_5 + (L_4 + 3)L_7 - L_4(L_4 + 2(L_{12} - 9)) + 18L_{12} + 15) \\ &\quad + L_2((5L_4 + 3)L_7^2 + 2(21L_4^2 + 84L_4 + 4L_5 + 2(L_4 + 9)L_{12} + 15)L_7 + 36L_{12}^2 - 4L_3(L_4 - 3) \\ &\quad + 91L_4 + 132L_5 + L_4(L_4(9L_4 + 77) + 44L_5 - 4L_8) - 8L_6 + 12L_8 + 4(3L_4(L_4 + 22) + 2L_5)L_{12} \\ &\quad + 300L_{12} - 96L_{14} + 15) - 400L_4L_{14} - 48L_4L_7L_4 - 96L_7L_4 - 240L_{14} - 64L_4^2L_{14} - 48L_5L_{14} \\ &\quad + 20L_{14}^2 - 32L_4L_{12}L_{14} - 256L_{12}L_{14} + 8L_1(L_2((L_4 - 6)L_4 - 15) - 10L_5 + ((L_4 - 16)L_4 - 15)L_7 \\ &\quad - 48L_{12} + 6L_4 - L_4(L_4(L_4 + 27) + 12L_{12} - 2L_4 + 61) - 15) + 16i(L_4L_{18} - L_{12}L_{15} - L_{10}) \Big]. \end{split}$$

(18) 
$$B_{18} = \text{Tr} \left( X_0 X_2 X_0^2 X_2 X_0^3 X_2 \right)$$
 is given by

$$\begin{split} B_{18} &= \frac{1}{8192} \Big[ 1 + L_4^3 + 15L_4^2 + 12L_3L_4 + 15L_4 + 48L_1^2 + 60L_3 - L_2^3(L_7 - 1) + 36L_7 + 48L_3L_7 + 315L_4L_7 \\ &\quad + 3L_7L_4^3 + 150L_7L_4^2 + 75L_4^2L_7^2 + 126L_7^2 - 12L_3L_7^2 + 525L_7^2L_4 + 9L_7^4 + 84L_7^3 + 105L_7^3L_4 \\ &\quad + 60L_5 + 48L_7L_5 + 12L_4L_5 - 4L_4L_7L_5 - 4L_6L_7 - 12L_5L_7^2 + 4L_8L_4^2 + 224L_8 + 132L_8L_4 \\ &\quad + 448L_7L_8 - 4L_5L_8 + 108L_4L_7L_8 + 96L_7^2L_8 + 32L_8^2 + 8L_7L_9 - 12L_4L_9 + 24L_9 \\ &\quad + 24L_3L_{12} + 300L_4L_{12} + 210L_{12} + 18L_{12}L_4^2 + 1050L_7L_{12} + 24L_5L_{12} + 600L_4L_7L_{12} - 8L_3L_7L_{12} \\ &\quad + 30L_7^3L_{12} + 60L_7^2L_4L_{12} + 630L_7^2L_{12} + 6L_7L_4^2L_{12} - 8L_7L_5L_{12} + 336L_8L_{12} + 48L_7L_8L_{12} \\ &\quad - 16L_9L_{12} + 8L_4L_8L_{12} + 16L_{12}^3 + 552L_{12}^2 + 312L_7L_{12}^2 + 40L_4L_{12}^2 + 8L_{12}L_{13} \\ &\quad + L_2^2(4L_8 + (L_7 + 3)L_4 - L_7(L_7 + 2(L_{12} - 9)) + 18L_{12} + 15) \\ &\quad + L_2((5L_7 + 3)L_4^2 + 2(21L_7^2 + 84L_7 + 4L_8 + 2(L_7 + 9)L_{12} + 15)L_4 + 36L_{12}^2 - 4L_3(L_7 - 3) \\ &\quad + 91L_7 + 132L_8 + L_7(L_7(9L_7 + 77) + 44L_8 - 4L_5) - 8L_9 + 12L_5 + 4(3L_7(L_7 + 22) + 2L_8)L_{12} \\ &\quad + 300L_{12} - 96L_{14} + 15) - 400L_7L_{14} - 48L_4L_7L_{14} - 96L_4L_{14} - 240L_{14} - 64L_7^2L_{14} - 48L_8L_{14} \\ &\quad + 20L_{14}^2 - 32L_7L_{12}L_{14} - 256L_{12}L_{14} + 8L_1(L_2((L_7 - 6)L_7 - 15) - 10L_8 + ((L_7 - 16)L_7 - 15)L_4 \\ &\quad - 48L_{12} + 6L_{14} - L_7(L_7(L_7 + 27) + 12L_{12} - 2L_{14} + 61) - 15) + 16i(L_7L_{17} - L_{12}L_{16} - L_{11})\Big]. \end{split}$$

Those Makhlin invariants  $L_k$ 's can be also expressed by using Bargmann invariants  $B_k$ 's below:

- $L_1 = \frac{2}{3}(1 3B_1 3B_2 + 6B_3 + 3B_4 4B_6).$
- $L_2 = 1 2B_1 2B_2 + 4B_4$ .
- $L_3 = 4(1 + B_1B_2 3B_1 3B_2 + 6B_3 + B_4 B_4^2 + B_1B_4 + B_2B_4 4B_7 4B_8 + 4B_{10}).$
- $L_4 = 2B_1 1$ .
- $L_5 = 2B_2 + 4B_4 4B_1B_4 8B_5 8B_6 + 16B_7 1$ .
- $L_6 = \frac{4}{3} \Big( 1 + 4B_1 9B_1^2 18B_1B_2 + 6B_1^2B_2 3B_2^2 + 24B_1B_3 + 12B_2B_3 12B_3^2 12B_4 + 6B_2B_4 + 18B_1B_4 + 6B_1^2B_4 12B_3B_4 + 3B_4^2 6B_1B_4^2 + 12B_5 12B_1B_5 + 20B_6 4B_1B_6 24B_7 24B_1B_7 12B_{10} + 12B_1B_{10} + 24B_{11} \Big).$
- $L_7 = 2B_2 1$ .
- $L_8 = 2B_1 4B_2B_4 + 4B_4 8B_5 8B_6 + 16B_8 1$ .
- $L_9 = \frac{4}{3} \Big( 1 + 4B_2 9B_2^2 18B_1B_2 + 6B_1B_2^2 3B_1^2 + 24B_2B_3 + 12B_1B_3 12B_3^2 12B_4 + 6B_1B_4 + 18B_2B_4 + 6B_2^2B_4 12B_3B_4 + 3B_4^2 6B_2B_4^2 + 12B_5 12B_2B_5 + 20B_6 4B_2B_6 24B_8 24B_2B_8 12B_{10} + 12B_2B_{10} + 24B_{12} \Big).$
- $L_{10} = \frac{2}{3}i(27 97B_1 + 114B_1^2 46B_1^3 81B_2 + 178B_1B_2 64B_1^2B_2 + 78B_2^2 + 108B_1B_2^2 + 18B_2^3 + 172B_3 368B_1B_3 + 168B_1^2B_3 384B_2B_3 288B_1B_2B_3 144B_2^2B_3 + 456B_3^2 288B_3^3 + 120B_1B_3^2 + 360B_2B_3^2 18B_4 + 54B_1B_4 + 54B_2B_4 137B_1^2B_4 + 48B_1^3B_4 390B_1B_2B_4 141B_2^2B_4 + 72B_1B_2^2B_4 108B_3B_4 + 660B_1B_3B_4 + 48B_1^2B_3B_4 + 540B_2B_3B_4 192B_1B_2B_3B_4 480B_3^2B_4 129B_4^2 + 261B_1B_4^2 + 72B_1^2B_4^2 + 81B_2B_4^2 48B_1B_2B_4^2 144B_3B_4^2 12B_5 68B_1B_5 + 96B_1^2B_5 + 36B_2B_5 144B_1B_2B_5 + 144B_3B_5 96B_1B_3B_5 + 60B_4B_5 + 60B_1B_4B_5 36B_2B_4B_5 + 96B_3B_4B_5 + 48B_4^2B_5 + 88B_6 92B_1B_6 32B_1^3B_6 228B_2B_6 40B_1B_2B_6 + 64B_1^2B_2B_6 + 488B_3B_6 + 32B_1B_3B_6 + 336B_4B_6 + 32B_1^2B_4B_6 36B_2B_4B_6 + 96B_3B_4B_6 + 48B_4^2B_6 64B_1B_5B_6 40B_6^2 16B_7 + 132B_1B_7 + 96B_1^2B_7 + 228B_2B_7 + 384B_1B_2B_7 624B_3B_7 768B_1B_3B_7 552B_4B_7 96B_1B_4B_7 + 72B_2B_4B_7 192B_3B_4B_7 96B_4^2B_7 + 144B_5B_7 + 400B_6B_7 768B_7^2 + 36B_1B_8 96B_1^2B_8 + 84B_2B_8 48B_1B_2B_8 240B_3B_8 + 288B_1B_3B_8 24B_4B_8 + 72B_1B_4B_8 48B_5B_8 48B_6B_8 + 96B_7B_8 + 24B_9 48B_1B_9 144B_2B_9 + 192B_1B_2B_9 + 96B_3B_4B_1 12B_{11} 192B_1B_{11} 72B_1B_{12} 72B_2B_{11} + 192B_3B_{11} + 96B_4B_{11} + 36B_{12} + 96B_{13} 192B_1B_{13} 96B_2B_{13} + 192B_3B_{13} 192B_{14} + 384B_1B_{14} 384B_1B_{16} + 192B_{16} + 768B_{17}).$
- $L_{11} = \frac{2}{3}i\left(27 97B_2 + 114B_2^2 46B_2^3 81B_1 + 178B_1B_2 64B_2^2B_1 + 78B_1^2 + 108B_2B_1^2 + 18B_1^3 + 172B_3 368B_2B_3 + 168B_2^2B_3 384B_1B_3 288B_1B_2B_3 144B_1^2B_3 + 456B_3^2 288B_3^3 + 120B_2B_3^2 + 360B_1B_3^2 18B_4 + 54B_1B_4 + 54B_2B_4 137B_2^2B_4 + 48B_2^3B_4 390B_1B_2B_4 141B_1^2B_4 + 72B_1^2B_2B_4 128B_1B_2B_4 141B_1^2B_4 + 72B_1^2B_2B_4 128B_1B_2B_4 141B_1^2B_4 + 72B_1^2B_2B_4 128B_1B_2B_4 141B_1^2B_4 + 72B_1^2B_2B_4 128B_1B_2B_4 128B_1B_2B_2B_4 128B_1B_2B_4 128B_1B_2B_2B_4 128B_1B_2B_4 128B_1B_2B_4 128B_1B_2B_2B_4 128B_1B_2B_2B_4 128B_1B_2B_2B_4 128B_1B_2B_2B_4 128B_1B_2B_2 128B_1B_2B_2 128B_1B_2B_2 128B_1B_2B_2 128B_1B_2B_2 128B_1B_2 12$

$$\begin{split} &108B_{3}B_{4}+660B_{2}B_{3}B_{4}+48B_{2}^{2}B_{3}B_{4}+540B_{1}B_{3}B_{4}-192B_{1}B_{2}B_{3}B_{4}-480B_{3}^{2}B_{4}-129B_{4}^{2}+261B_{2}B_{4}^{2}+\\ &72B_{2}^{2}B_{4}^{2}+81B_{1}B_{4}^{2}-48B_{1}B_{2}B_{4}^{2}-144B_{3}B_{4}^{2}-12B_{5}-68B_{2}B_{5}+96B_{2}^{2}B_{5}+36B_{1}B_{5}-144B_{1}B_{2}B_{5}+\\ &144B_{3}B_{5}-96B_{2}B_{3}B_{5}+60B_{4}B_{5}+60B_{2}B_{4}B_{5}-36B_{1}B_{4}B_{5}+96B_{3}B_{4}B_{5}+48B_{4}^{2}B_{5}+88B_{6}-92B_{2}B_{6}-\\ &32B_{2}^{3}B_{6}-228B_{1}B_{6}-40B_{1}B_{2}B_{6}+64B_{2}^{2}B_{1}B_{6}+488B_{3}B_{6}+32B_{2}B_{3}B_{6}+336B_{4}B_{6}+32B_{2}^{2}B_{4}B_{6}-\\ &36B_{1}B_{4}B_{6}+96B_{3}B_{4}B_{6}+48B_{4}^{2}B_{6}+48B_{5}B_{6}-64B_{2}B_{5}B_{6}-40B_{6}^{2}-16B_{8}+132B_{2}B_{8}+96B_{2}^{2}B_{8}+\\ &228B_{1}B_{8}+384B_{1}B_{2}B_{8}-624B_{3}B_{8}-768B_{2}B_{3}B_{8}-552B_{4}B_{8}-96B_{2}B_{4}B_{8}+72B_{1}B_{4}B_{8}-192B_{3}B_{4}B_{8}-\\ &96B_{4}^{2}B_{8}+144B_{5}B_{8}+400B_{6}B_{8}-768B_{8}^{2}+36B_{2}B_{7}-96B_{2}^{2}B_{7}+84B_{1}B_{7}-48B_{1}B_{2}B_{7}-240B_{3}B_{7}+\\ &288B_{2}B_{3}B_{7}-24B_{4}B_{7}+72B_{2}B_{4}B_{7}-48B_{5}B_{7}-48B_{6}B_{7}+96B_{7}B_{8}+24B_{9}-48B_{2}B_{9}-144B_{1}B_{9}+\\ &192B_{1}B_{2}B_{9}+96B_{3}B_{9}-90B_{10}+138B_{2}B_{10}-96B_{2}^{2}B_{10}+162B_{1}B_{10}-144B_{1}B_{2}B_{10}-288B_{3}B_{10}+\\ &192B_{2}B_{3}B_{10}-144B_{4}B_{10}+96B_{2}B_{4}B_{10}-12B_{12}-192B_{2}B_{12}-72B_{1}B_{12}-72B_{2}B_{11}+192B_{3}B_{12}+\\ &96B_{4}B_{12}+36B_{11}+96B_{14}-192B_{2}B_{14}-96B_{1}B_{14}+192B_{3}B_{14}-192B_{13}+384B_{2}B_{13}-384B_{2}B_{15}+\\ &192B_{15}+768B_{18} \Big). \end{split}$$

- $L_{12} = 1 2B_1 2B_2 + 4B_3$ .
- $L_{13} = 12(B_1 + B_2) 12(B_1 + B_2)B_4 36B_3 + 24B_3B_4 + 24B_5 8B_6 + 16(B_7 + B_8) 32B_9 3.$
- $L_{14} = 2(1 3B_1 3B_2 + 2B_1B_2 + 6B_3 + B_4 4B_5).$
- $L_{15} = \frac{4}{3}i\Big(-1+5B_1-6B_1^2+3B_2+3B_1B_2-12B_3+6B_1B_3-6B_2B_3+12B_3^2+6B_4-12B_1B_4-6B_2B_4+6B_1B_2B_4+6B_3B_4-6B_5+12B_1B_5-14B_6+4B_1B_6+24B_7+12B_8-12B_1B_8-24B_{13}\Big).$
- $L_{16} = \frac{4}{3}i\Big(-1+5B_2-6B_2^2+3B_1+3B_1B_2-12B_3+6B_2B_3-6B_1B_3+12B_3^2+6B_4-12B_2B_4-6B_1B_4+6B_1B_2B_4+6B_3B_4-6B_5+12B_2B_5-14B_6+4B_2B_6+24B_8+12B_7-12B_2B_7-24B_{14}\Big).$
- $L_{17} = \frac{4}{3}i\Big(-9 + 15B_1 + 6B_1^2 + 19B_2 11B_1B_2 24B_3 6B_1B_3 + 6B_2B_3 12B_3^2 + 18B_4 24B_1B_4 + 6B_1^2B_4 30B_2B_4 + 12B_1B_2B_4 + 42B_3B_4 24B_1B_3B_4 + 6B_4^2 6B_1B_4^2 + 6B_5 48B_1B_5 + 12B_2B_5 12B_4B_5 18B_6 4B_2B_6 + 8B_1B_2B_6 12B_4B_6 12B_7 12B_2B_7 + 48B_3B_7 + 24B_4B_7 + 24B_1B_8 + 48B_1B_9 + 24B_{10} 12B_1B_{10} 24B_{11} + 96B_{13} 24B_{14} 96B_{15}\Big).$
- $L_{18} = \frac{4}{3}i\Big(-9 + 15B_2 + 6B_2^2 + 19B_1 11B_1B_2 24B_3 6B_2B_3 + 6B_1B_3 12B_3^2 + 18B_4 24B_2B_4 + 6B_2^2B_4 30B_1B_4 + 12B_1B_2B_4 + 42B_3B_4 24B_2B_3B_4 + 6B_4^2 6B_2B_4^2 + 6B_5 48B_2B_5 + 12B_1B_5 12B_4B_5 18B_6 4B_1B_6 + 8B_1B_2B_6 12B_4B_6 12B_8 12B_1B_8 + 48B_3B_8 + 24B_4B_8 + 24B_2B_7 + 48B_2B_9 + 24B_{10} 12B_2B_{10} 24B_{12} + 96B_{14} 24B_{13} 96B_{16}\Big).$

*Proof.* The correctness of all of these results can be checked by the mathematical software MATH-EMATICA. We remark here that getting these results is more difficult than checking the above formulas. All materials preceding this lemma serve as preparations for simplifying the calculations in the proof of this lemma. In fact, we expand  $B_k$ 's by using the Bloch decomposition of  $\rho_{AB}$ . Through tedious algebraic computations and simplifications, utilizing the results from Subsections B.1, B.2, and B.3, we obtain the desired results.

#### **B.4 Proof of Theorem 2.2**

With the above preparations, now we can present the proof of Theorem 2.2.

*Proof of Theorem* 2.2. We have already known that the generators of the invariant ring generated by LU invariant polynomials for two-qubit states is given by a complete set of 18 Makhlin's fundamental invariants  $I_k$ 's, where  $I_k$ 's can be generated by  $L_k$ 's in Proposition B.22. From Lemma B.23, we see that  $L_k$ 's can be generated by  $B_k$ 's. Therefore,  $B_k$ 's can generate the invariant ring.

## C Proof of Theorem 3.1

#### C.1 Entanglement criterion by Makhlin's invariants

Let the partial trace with respect to either one subsystem of  $\rho_{AB}$  be given by  $\rho_{AB}^{\Gamma} = \rho_{AB}^{T_A}$  or  $\rho_{AB}^{T_B}$ . We have the following result:

**Lemma C.1.** All eigenvalues of the operator  $X := 4\rho_{AB} - \mathbb{1}_2 \otimes \mathbb{1}_2$  are determined by its characteristic polynomial equation  $x^4 + px^2 + qx + r = 0$ , where

$$\begin{cases} p = -2(L_2 + L_4 + L_7), \\ q = -8(L_{12} - L_1), \\ r = L_2^2 + 2(L_4 + L_7)L_2 + (L_4 - L_7)^2 - 4(L_3 + L_5 + L_8) + 8L_{14}. \end{cases}$$
(C.1)

*Here the meaning of*  $L_k$ 's can be found in Proposition B.22.

*Proof.* The proof is obtained by direct and tedious computations. It is omitted here.

We remark here that the correctness of the above result can also be checked by employing symbolic computation function of MATHEMATICA. Apparently, getting this result is more difficult than checking the correctness of it. Based on the above result presented in Lemma C.1, we can derive the following characterization of entanglement in two-qubit system. Basically, it is another equivalent reformulation of Positive Partial-Tranpose criteria for two-qubit system. More importantly, our reformulation can be viewed as the first criterion using locally unitary invariants.

For any two-qubit state  $\rho_{AB}$ , parameterized as in Eq. (B.44), note that

$$\operatorname{Tr}(\rho_A^2) = \frac{1+L_4}{2}, \quad \operatorname{Tr}(\rho_B^2) = \frac{1+L_7}{2}, \quad \operatorname{Tr}(\rho_{AB}^2) = \frac{1+L_2+L_4+L_7}{4}, \quad (C.2)$$

from the facts that  $\operatorname{Tr}(\rho_A^2)$ ,  $\operatorname{Tr}(\rho_B^2) \in [\frac{1}{2}, 1]$  and  $\operatorname{Tr}(\rho_{AB}^2) \in [\frac{1}{4}, 1]$ , we get that

$$\begin{cases} 0 \leqslant L_4 \leqslant 1, \\ 0 \leqslant L_7 \leqslant 1, \\ 0 \leqslant L_2 + L_4 + L_7 \leqslant 3. \end{cases}$$
(C.3)

It follows from Lemma C.1, we get the characteristic polynomial equation is given by

$$\lambda^4 - \lambda^3 + \frac{p+6}{16}\lambda^2 - \frac{2p-q+4}{64}\lambda + \frac{p-q+r+1}{256} = 0.$$
 (C.4)

Recall a result in [?]: Consider an algebraic equation of degree  $N \ge 1$ ,

$$\prod_{k=1}^{N} (x - x_k) = \sum_{\ell=0}^{N} (-1)^{\ell} e_{\ell} x^{N-\ell} = 0 \quad (e_0 = 1),$$
(C.5)

which has only real roots  $x_k \in \mathbb{R}(k = 1, ..., N)$ . The necessary and sufficient condition that all the roots  $x_k$ 's to be non-negative is that all the coefficients  $e_\ell$ 's are non-negative. That is,

$$(\forall k \in [N] : x_k \ge 0) \iff (\forall \ell \in [N] : e_\ell \ge 0, e_0 \equiv 1).$$
(C.6)

From the above result, we can present a following result about the positivity of Hermitian matrix *X*:

**Proposition C.2.** For a Hermitian complex matrix  $\mathbf{X} \in \mathbb{C}^{N \times N}$ , denote  $p_k(\mathbf{X}) := \text{Tr}(\mathbf{X}^k)$ , then its characteristic polynomial is given by

$$\det(xI_N - X) = \sum_{k=0}^{N} (-1)^k e_k(X) x^{N-k}$$

where

$$e_{k}(\mathbf{X}) = \frac{1}{k!} \begin{vmatrix} p_{1}(\mathbf{X}) & 1 & 0 & \cdots & 0 \\ p_{2}(\mathbf{X}) & p_{1}(\mathbf{X}) & 2 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ p_{k-1}(\mathbf{X}) & p_{k-2}(\mathbf{X}) & p_{k-3}(\mathbf{X}) & \cdots & k-1 \\ p_{k}(\mathbf{X}) & p_{k-1}(\mathbf{X}) & p_{k-2}(\mathbf{X}) & \cdots & p_{1}(\mathbf{X}) \end{vmatrix} \quad (k \ge 1).$$

Then we have

$$\mathbf{X} \ge \mathbf{0} \iff \begin{vmatrix} p_1(\mathbf{X}) & 1 & 0 & \cdots & 0 \\ p_2(\mathbf{X}) & p_1(\mathbf{X}) & 2 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ p_{k-1}(\mathbf{X}) & p_{k-2}(\mathbf{X}) & p_{k-3}(\mathbf{X}) & \cdots & k-1 \\ p_k(\mathbf{X}) & p_{k-1}(\mathbf{X}) & p_{k-2}(\mathbf{X}) & \cdots & p_1(\mathbf{X}) \end{vmatrix} \ge 0 \quad (k = 1, 2, \dots, N).$$

*Proof.* Since **X** is Hermitian matrix, it follows that its characteristic polynomial  $det(x\mathbb{1}_N - \mathbf{X}) = \sum_{k=0}^{N} (-1)^k e_k(\mathbf{X}) x^{N-k}$  has only real roots. These real roots are non-negative if and only if  $\mathbf{X} \ge \mathbf{0}$ . Therefore  $\mathbf{X} \ge \mathbf{0}$  if and only if  $e_k(\mathbf{X}) \ge 0$ , where k = 1, ..., N

From the above result, the non-negativeness of  $\rho_{AB}$  is guaranteed by the following inequalities [?]:

$$\begin{cases} p+6 & \ge 0\\ 2p-q+4 & \ge 0 \iff \\ p-q+r+1 & \ge 0 \end{cases} \begin{pmatrix} p & \ge -6\\ q & \le 2p+4\\ r & \ge q-p-1. \end{cases}$$
(C.7)

Based on both Eq. (C.3) and Eq. (C.7), we can summarize the above discussion into the following result:

**Proposition C.3.** For any Hermitian matrix  $\rho_{AB}$  of fixed trace one, parameterized as

$$\rho_{AB} = \frac{1}{4} \left( \mathbb{1}_2 \otimes \mathbb{1}_2 + \boldsymbol{a} \cdot \boldsymbol{\sigma} \otimes \mathbb{1}_2 + \mathbb{1}_2 \otimes \boldsymbol{b} \cdot \boldsymbol{\sigma} + \sum_{i,j=1}^3 c_{ij} \sigma_i \otimes \sigma_j \right),$$
(C.8)

where  $\mathbf{a} = (a_1, a_2, a_3)^{\mathsf{T}}$  and  $\mathbf{b} = (b_1, b_2, b_3)^{\mathsf{T}}$  are in  $\mathbb{R}^3$ , and  $\mathbf{C} = (c_{ij})_{3\times 3} \in \mathbb{R}^{3\times 3}$ , the necessary and sufficient condition for the non-negativeness  $\rho_{AB} \ge \mathbf{0}$  if and only if the following inequalities concerning the 3-tuple  $(\mathbf{a}, \mathbf{b}, \mathbf{C})$  are true:

$$\begin{cases} 0 \leq L_4 \leq 1, \\ 0 \leq L_7 \leq 1, \\ 0 \leq L_2 + L_4 + L_7 \leq 3, \\ L_2 + L_4 + L_7 \leq 1 + 2(L_{12} - L_1), \\ (L_2 + L_4 + L_7 - 1)^2 - 4(L_3 + L_4L_7 + L_5 + L_8) + 8(L_{12} + L_{14} - L_1) \geq 0. \end{cases}$$
(C.9)

The above constraints about the 3-tuple (a, b, C) can be equivalently to reformulated via locally unitary Bargmann invariants:

$$\begin{cases} 1 + 2 \operatorname{Tr} (\rho_{AB}^{3}) & \geqslant 3 \operatorname{Tr} (\rho_{AB}^{2}) ,\\ 1 + 3 [\operatorname{Tr} (\rho_{AB}^{2})]^{2} + 8 \operatorname{Tr} (\rho_{AB}^{3}) & \geqslant 6 \operatorname{Tr} (\rho_{AB}^{4}) + 6 \operatorname{Tr} (\rho_{AB}^{2}) . \end{cases}$$
(C.10)

**Lemma C.4** (Detection of entanglement via locally unitary invariants). For any given two-qubit state  $\rho_{AB}$ , parameterized as in (B.44), which is entangled if and only if 9 invariants of 18 Makhlin invariants are satisfying the following inequality:

$$1 + (|a|^{2} - |b|^{2})^{2} + 2(|a|^{2} + |b|^{2})\langle C, C \rangle + 2\langle C^{\mathsf{T}}C, C^{\mathsf{T}}C \rangle + 8(\langle a|C|b \rangle + \det(C))$$
  
$$< \langle C, C \rangle^{2} + 2(|a|^{2} + |b|^{2} + \langle C, C \rangle) + 4(\langle a|CC^{\mathsf{T}}|a \rangle + \langle b|C^{\mathsf{T}}C|b \rangle) + 8\langle a|\widehat{C}|b \rangle.$$
(C.11)

*Proof.* All eigenvalues of the operator  $\Upsilon := 4\rho_{AB}^{\Gamma} - \mathbb{1}_2 \otimes \mathbb{1}_2$  are determined by its characteristic polynomial equation  $y^4 + \tilde{p}y^2 + \tilde{q}y + \tilde{r} = 0$ , where

$$\begin{split} \tilde{p} &= -2\left(|a|^2 + |b|^2 + \langle C, C \rangle\right), \quad \tilde{q} = -8\left(\langle a|C|b \rangle + \det(C)\right), \\ \tilde{r} &= \left(|a|^2 - |b|^2\right)^2 + 2(|a|^2 + |b|^2)\langle C, C \rangle + 2\langle C^{\mathsf{T}}C, C^{\mathsf{T}}C \rangle - \langle C, C \rangle^2 \\ &- 4\left(\langle a|CC^{\mathsf{T}}|a \rangle + \langle b|C^{\mathsf{T}}C|b \rangle\right) - 8\langle a|\widehat{C}|b \rangle. \end{split}$$

Note that  $\det(\rho_{AB}^{\Gamma}) = \frac{\tilde{p} - \tilde{q} + \tilde{r} + 1}{256}$ . Thus  $\rho_{AB}$  is entangled if and only if  $\det(\rho_{AB}^{\Gamma}) < 0$ . Therefore we get the desired inequality.

**Example C.5** (The family of two-qubit Werner states). Two-qubit Wener state of single parameter is defined by  $\rho_w = w |\psi^-\rangle \langle \psi^-| + (1-w) \frac{\mathbb{1}_4}{4}$ , where  $|\psi^-\rangle = \frac{|01\rangle - |10\rangle}{\sqrt{2}}$  and  $w \in [0,1]$ , which can be rewritten as

$$\rho_w = \frac{1}{4} \left( \mathbb{1}_2 \otimes \mathbb{1}_2 - w \sum_{k=1}^3 \sigma_k \otimes \sigma_k \right).$$

In such a case, a = b = 0 and  $C = -w\mathbb{1}_3$ . Then two-qubit Werner state  $\rho_w$  is entangled if and only if Eq. (C.11) becomes

$$1 + 2\langle C^{\mathsf{T}}C, C^{\mathsf{T}}C \rangle + 8 \det(C) < \langle C, C \rangle^{2} + 2\langle C, C \rangle$$

$$\iff 1 + 6w^{4} - 8w^{3} < 9w^{4} + 6w^{2} \iff \frac{1}{3} < w \leq 1.$$
(C.12)

**Example C.6** (The family of two-qubit Bell-diagonal states). Two-qubit Bell-diagonal state of three parameters is defined by

$$\rho_{\text{Bell}} = \frac{1}{4} \left( \mathbb{1}_2 \otimes \mathbb{1}_2 + \sum_{k=1}^3 t_k \sigma_k \otimes \sigma_k \right),$$

where  $t = (t_1, t_2, t_3) \in D$  (specified later). The set *D* is a bounded and closed region:  $D \subset [-1, 1]^3$ . The above mentioned *D* is determined by

$$\begin{cases} 1 - t_1 - t_2 - t_3 \ge 0, \\ 1 - t_1 + t_2 + t_3 \ge 0, \\ 1 + t_1 - t_2 + t_3 \ge 0, \\ 1 + t_1 + t_2 - t_3 \ge 0. \end{cases}$$

In this case, a = b = 0 and  $C = \text{diag}(t_1, t_2, t_3)$ . Now two-qubit Bell-diagonal state  $\rho_{\text{Bell}}$  is entangled if and only if Eq. (C.11) becomes

$$1 + 2\langle C^{\mathsf{T}}C, C^{\mathsf{T}}C \rangle + 8\det(C) < \langle C, C \rangle^{2} + 2\langle C, C \rangle$$
$$\iff 1 + 2\sum_{j=1}^{3} t_{j}^{4} + 8t_{1}t_{2}t_{3} < \left(\sum_{j=1}^{3} t_{j}^{2}\right)^{2} + 2\sum_{j=1}^{3} t_{j}^{2}.$$

Note that

$$\left(\sum_{j=1}^{3} t_{j}^{2}\right)^{2} + 2\sum_{j=1}^{3} t_{j}^{2} - 2\sum_{j=1}^{3} t_{j}^{4} - 8t_{1}t_{2}t_{3} - 1$$
  
=  $-(t_{1} - t_{2} - t_{3} + 1)(t_{1} + t_{2} - t_{3} - 1)(t_{1} - t_{2} + t_{3} - 1)(t_{1} + t_{2} + t_{3} + 1) > 0,$ 

which is equivalent to  $|t_1| + |t_2| + |t_3| > 1$ .

## C.2 Proof of Theorem 3.1

*Proof of Theorem 3.1.* Note we obtained that a complete set of LU Bargmann invariants  $\{B_k : k = 1, ..., 18\}$  for the invariant ring of LU invariant polynomials for two-qubit states. Using the 18 Bargmann generators, we can test the LU equivalence of two-qubit states by experiment via measuring Bargmann invariants. Besides, we can use 7 Bargmann invariants to test entanglement of two-qubit states: By using Lemma B.23, Eq. (C.11) can be equivalently transformed into the following form:

$$6(B_1 + B_2 - B_1B_2 - B_4 - B_{10}) + 12(B_5 - B_3) + 3B_4^2 + 4B_6 < 1.$$

This completes the proof.