

SUBSPACE CORRECTION METHODS FOR SEMICOERCIVE AND NEARLY SEMICOERCIVE CONVEX OPTIMIZATION WITH APPLICATIONS TO NONLINEAR PDES*

YOUNG-JU LEE[†] AND JONGHO PARK[‡]

Abstract. We present new convergence analyses for subspace correction methods for semicoercive and nearly semicoercive convex optimization problems, generalizing the theory of singular and nearly singular linear problems to the nonlinear domain. Our results demonstrate that the elegant theoretical framework developed for singular and nearly singular linear problems can be extended to semicoercive and nearly semicoercive convex optimization problems. For semicoercive problems, we show that the convergence rate can be estimated in terms of a seminorm stable decomposition over the subspaces and the kernel of the problem, aligning with the theory for singular linear problems. For nearly semicoercive problems, we establish a parameter-independent convergence rate, assuming the kernel of the semicoercive part can be decomposed into a sum of local kernels, which aligns with the theory for nearly singular problems. To demonstrate the applicability of our results, we provide convergence analyses of two-level additive Schwarz methods for solving a nonlinear Neumann boundary value problem and its perturbation within the proposed abstract framework.

Key words. Semicoercive problems, Nearly semicoercive problems, Subspace correction methods, Convex optimization, Nonlinear PDEs

AMS subject classifications. 65J20, 65N20, 65N55, 90C22, 90C25

1. Introduction. Many important linear problems arise in science and engineering as either singular or nearly singular. These problems can be characterized as systems, which have a nontrivial null space or near null space and they appear in various applications, which include finite element discretizations of the Poisson equation with pure Neumann boundary conditions, and/or variational problems of $H(\text{div})$ and $H(\text{curl})$ [5, 9]. An important class of nearly singular problems can also be found at nearly incompressible linear elastic equations [31]. Nearly singular problems also occur when solving indefinite systems arising from mixed finite element discretizations of the Navier–Stokes equations [25], as well as in more complex systems such as non-Newtonian fluids [35] and fluid–structure interaction problems [53]. In particular, the nearly incompressible linear elasticity problem arises as a subproblem to be solved [31], when the augmented Lagrangian Uzawa method is employed [33].

Due to the significance of singular and nearly singular linear problems in scientific computing, as highlighted by the numerous examples discussed above, there has been extensive research on numerical methods for solving these problems. The theory of basic iterative methods for singular problems was first introduced in [28], with more refined results presented in later works such as [14, 32, 50]. In addition, subspace correction methods [52, 54], which offer a general framework for a variety of iterative methods, ranging from basic methods to advanced ones like domain decomposition and multigrid methods, were rigorously analyzed for singular linear problems in [34, 50] and for nearly singular linear problems in [33, 51]. Building on the theory of subspace correction methods, several applications have been developed for the specific examples

*Submitted to arXiv.

Funding: Young-Ju Lee’s work was supported by NSF-DMS 2208499.

[†]Department of Mathematics, Texas State University, San Marcos, TX 78666, USA (yjlee@txstate.edu).

[‡]Applied Mathematics and Computational Sciences Program, Computer, Electrical and Mathematical Science and Engineering Division, King Abdullah University of Science and Technology (KAUST), Thuwal 23955, Saudi Arabia (jongho.park@kaust.edu.sa).

discussed above; see, for instance, [25, 31, 53].

A natural generalization of the concept of singularity in linear problems to convex optimization problems is semicoercivity [2]. Intuitively, a convex functional is said to be semicoercive if it is flat along some subspace and increases to infinity in the other directions; the rigorous definition of semicoercivity can be found in [2], as well as in section 2. In the special case of quadratic functionals, semicoercivity is equivalent to the singularity of the corresponding linear problem. As a result, semicoercive problems, like singular ones, arise frequently in various nonlinear applications. Consequently, there has been some research on efficient numerical solvers for particular semicoercive problems, such as [21, 23].

The goal of this paper is to extend the well-established theory of subspace correction methods for singular and nearly singular problems [33, 34, 50, 51] to semicoercive and nearly semicoercive convex optimization problems. This extension is motivated by recent developments in the convergence theory of subspace correction methods for convex optimization. While early results on the convex theory can be found in [6, 7, 15, 47, 48], more refined and general convergence results have been presented recently in [39, 41, 45]. Moreover, this convergence theory has been applied to several interesting nonlinear partial differential equations (PDEs) and variational inequalities in [30, 43, 44]. However, there has been no work addressing the theory of subspace correction methods for semicoercive or nearly semicoercive problems; all existing studies require coercivity or even stronger assumptions, such as uniform convexity.

In this paper, we present new convergence analyses for subspace correction methods for semicoercive and nearly semicoercive convex optimization problems in Banach spaces. Roughly speaking, the main results of this paper combine the theories for singular and nearly singular problems [33, 34, 51] with those for convex optimization problems [39, 45] for subspace correction methods. For semicoercive problems, we prove that the convergence rate of a subspace correction method can be estimated in terms of a seminorm stable decomposition over the subspaces and the kernel of the problem. This aligns with the theory for singular linear problems established in [34, 50]. The analysis is analogous to that for coercive problems, except that special care must be taken with the kernel of the problem. For nearly semicoercive problems, in the same spirit as in [33], we obtain a parameter-independent convergence rate estimate for subspace correction methods, under the assumption that the kernel of the semicoercive part, i.e., the near null space, can be decomposed into a sum of local near null spaces. We note that the analysis of nearly singular linear problems in [33] relies heavily on orthogonality in Hilbert spaces. However, such orthogonality does not directly apply in our setting, as we are dealing with nonlinear problems posed in Banach spaces. Therefore, we carefully extend the theory of nearly singular linear problems to nearly semicoercive convex optimization problems by employing nonlinear orthogonal decompositions in Banach spaces, which were introduced in [3, 4]. As examples of the applications of the convergence theory proposed in this paper, we provide convergence analyses of two-level additive Schwarz methods for solving a Neumann boundary value problem involving the s -Laplacian [30] and its perturbation.

The rest of this paper is organized as follows. In section 2, we review the notion of semicoercivity and provide a characterization in terms of seminorms. In section 3, we review subspace correction methods for convex optimization. In section 4, we present a convergence analysis of subspace correction methods for semicoercive convex optimization. In section 5, we present a convergence analysis for subspace correction methods for nearly semicoercive convex optimization. In section 6, we present several

applications of the proposed convergence theory to domain decomposition methods for nonlinear PDEs. In [section 7](#), we conclude the paper with remarks.

2. Semicoercive functionals. In this section, we present the notion of semicoercive functionals [\[2\]](#) and their characterization in terms of seminorms. Additionally, we explore the relation between semicoercive convex functionals and semidefinite linear problems [\[32, 50\]](#).

We first recall the definition of semicoercive functionals introduced in [\[2\]](#). Let V be a reflexive Banach space equipped with a norm $\|\cdot\|$. A proper functional $F: V \rightarrow \overline{\mathbb{R}}$ is said to be *semicoercive* if there exists a closed subspace \mathcal{N} of V such that

$$(2.1) \quad F(v) = F(v + \phi) \quad \forall v \in V, \phi \in \mathcal{N},$$

and the quotient functional $\bar{F}: V/\mathcal{N} \rightarrow \overline{\mathbb{R}}$ defined by

$$\bar{F}(v + \mathcal{N}) = F(v), \quad v \in V,$$

is coercive in the sense that

$$(2.2) \quad \bar{F}(v + \mathcal{N}) \rightarrow \infty \quad \text{as } \|v + \mathcal{N}\|_{V/\mathcal{N}} \rightarrow \infty,$$

where $\|\cdot\|_{V/\mathcal{N}}$ denotes the quotient norm given by

$$\|v + \mathcal{N}\|_{V/\mathcal{N}} = \inf_{\phi \in \mathcal{N}} \|v + \phi\|, \quad v \in V.$$

If such a subspace \mathcal{N} exists, then it can be easily shown that \mathcal{N} is unique. We refer to this subspace as the *kernel* of F , denoted by $\mathcal{N} = \ker F$.

As described above, semicoercivity is defined in terms of a quotient space. However, when developing the convergence theory for subspace correction methods and its applications, it is more convenient to work with seminorms rather than quotient spaces. In [Lemma 2.1](#), we present a condition under which a seminorm can be characterized as a quotient norm. Given a seminorm $|\cdot|$ on V , we define the *kernel* $\ker |\cdot|$ of $|\cdot|$ as

$$\ker |\cdot| = \{v \in V : |v| = 0\}.$$

LEMMA 2.1. *Let V be a Banach space. A seminorm $|\cdot|$ on V is equivalent to the quotient norm $\|\cdot + \mathcal{N}\|_{V/\mathcal{N}}$ for some closed subspace \mathcal{N} of V if and only if it satisfies the following:*

- (i) $|\cdot|$ is continuous on V .
- (ii) There exists a positive constant C such that

$$(2.3) \quad \inf_{\phi \in \ker |\cdot|} \|v + \phi\| \leq C|v| \quad \forall v \in V.$$

Proof. Suppose that we have a seminorm $|\cdot|$ on V that is equivalent to the quotient norm $\|\cdot + \mathcal{N}\|_{V/\mathcal{N}}$ for some closed subspace \mathcal{N} of V . It follows directly that $\mathcal{N} = \ker F$, making both (i) and (ii) straightforward.

Conversely, let $|\cdot|$ be a seminorm on V that satisfies (i) and (ii). We set $\mathcal{N} = \ker |\cdot|$. Due to the continuity of $|\cdot|$, we have

$$(2.4) \quad |v| = |v + \phi| \leq C'\|v + \phi\|$$

for any $\phi \in \mathcal{N}$, where C' is a positive constant. By combining [\(2.3\)](#) and [\(2.4\)](#), we deduce that $|\cdot|$ is equivalent to $\|\cdot + \mathcal{N}\|_{V/\mathcal{N}}$. \square

Remark 2.2. The condition (2.3) is satisfied by many seminorms commonly encountered in PDEs. For instance, the Bramble–Hilbert lemma [10] ensures that if $V = W^{m,p}(\Omega)$, where $m \in \mathbb{Z}_{>0}$, $p \in [1, \infty]$, and the domain $\Omega \subset \mathbb{R}^d$ satisfies certain geometric conditions [22, 24], then (2.3) holds for the Sobolev seminorm $|\cdot|_{W^{m,p}(\Omega)}$.

Thanks to Lemma 2.1, we are able to characterize semicoercivity in terms of seminorms; see Proposition 2.3.

PROPOSITION 2.3. *Let V be a reflexive Banach space. A proper functional $F: V \rightarrow \overline{\mathbb{R}}$ is semicoercive if and only if there exists a continuous seminorm $|\cdot|$ on V that satisfies (2.3) and the following:*

- (i) $F(v) = F(v + \phi)$ for any $v \in V$ and $\phi \in \ker |\cdot|$.
- (ii) $F(v) \rightarrow \infty$ as $|v| \rightarrow \infty$.

Proof. If we have a semicoercive functional F , then the seminorm $|\cdot|$ defined as

$$|v| = \inf_{\phi \in \ker F} \|v + \phi\|, \quad v \in V,$$

is continuous and satisfies (2.3), (i) and (ii). Conversely, if we have a continuous seminorm $|\cdot|$ on V that satisfies (2.3), (i), and (ii), then we can readily deduce, using Lemma 2.1, that F is semicoercive with the kernel $\ker F = \ker |\cdot|$. \square

We conclude this section by demonstrating that minimizing semicoercive and convex energy functionals generalizes solving semidefinite linear problems. Let H be a Hilbert space equipped with an inner product (\cdot, \cdot) . We consider the following semidefinite linear problem:

$$(2.5) \quad Au = f,$$

where $A: H \rightarrow H$ is a continuous, symmetric and positive semidefinite linear operator, and $f \in \text{ran } A$. It is straightforward to verify that $u \in H$ solves (2.5) if and only if it minimizes the following quadratic energy functional:

$$(2.6) \quad F(v) = \frac{1}{2}(Av, v) - (f, v), \quad v \in H.$$

That is, the semidefinite linear problem (2.5) is equivalent to the minimization problem given by (2.6). Clearly, the energy functional F in (2.6) is convex and semicoercive with the kernel $\ker F = \ker A$. Hence, we conclude that the semidefinite linear problem (2.5) is a special case of semicoercive convex optimization.

3. Subspace correction methods for convex optimization. In this section, we briefly summarize subspace correction methods for convex optimization, which have been extensively studied in the literature, e.g., [39, 45, 47, 48]. For simplicity, we focus on the case of exact local problems only; the case of inexact local problems [39, 45] will be considered in Appendix A.

We consider the following abstract convex optimization problem on a reflexive Banach space V :

$$(3.1) \quad \min_{v \in V} F(v),$$

where $F: V \rightarrow \overline{\mathbb{R}}$ is a Gâteaux differentiable and convex functional. We can readily verify that the problem (3.1) admits a solution (may not be unique) $u \in V$ if F is semicoercive.

We assume that the solution space V of (3.1) admits a space decomposition

$$(3.2) \quad V = \sum_{j=1}^N V_j,$$

where each V_j , $j \in [N] = \{1, 2, \dots, N\}$, is a closed subspace of V . One important property of the space decomposition (3.2) is the stable decomposition property. Namely, we have

$$(3.3) \quad \sup_{\|w\|=1} \inf_{\sum_{j=1}^N w_j = w} \left(\sum_{j=1}^N \|w_j\|^q \right)^{\frac{1}{q}} < \infty,$$

for any $q \in [1, \infty)$, where w and w_j are taken from V and V_j , respectively. This property follows directly from the open mapping theorem (see [54, Equation (2.15)] and [45, Equation (2)]).

Meanwhile, the convexity of the energy functional F in (3.1) implies the following inequality, known as the strengthened convexity condition [39, Assumption 4.2], holds for some $\tau > 0$:

$$(3.4) \quad (1 - \tau N)F(v) + \tau \sum_{j=1}^N F(v + w_j) \geq F \left(v + \tau \sum_{j=1}^N w_j \right), \quad v \in V, \quad w_j \in V_j.$$

A positive constant τ_0 is defined as the maximum τ that satisfies the strengthened convexity condition, i.e.,

$$(3.5) \quad \tau_0 = \max \{ \tau > 0 : \text{The inequality (3.4) holds} \}.$$

Then it is clear that (3.4) holds for every $\tau > 0$ less than or equal to τ_0 . While we have a trivial estimate $\tau_0 \geq 1/N$, in many applications, better estimates for τ_0 , which are often independent of N , can be obtained using a coloring argument [39, 48]. One may refer to [39, Section 4.1] for a discussion on the relation between (3.4) and strengthened Cauchy–Schwarz inequalities, which plays a crucial role in the analysis of multilevel methods for linear problems [49, 52].

Subspace correction methods involve local problems defined in the subspaces $\{V_j\}_{j=1}^N$. Given $v \in V$, the optimal residual in the subspace V_j , which the energy functional F , is obtained by solving the minimization problem

$$(3.6) \quad \min_{w_j \in V_j} \{ F_j(w_j; v) := F(v + w_j) \}.$$

The parallel subspace correction method, also known as the additive Schwarz method in the literature on domain decomposition methods, for solving (3.1) with the local problem (3.6) is presented in Algorithm 3.1. Note that the upper bound τ_0 for the step size τ was given in (3.5).

Remark 3.1. Incorporating acceleration schemes designed for first-order methods for convex optimization (see, e.g., [38]) into Algorithm 3.1 yields accelerated variants of the parallel subspace correction method [40, 41]. These accelerated methods typically exhibit faster convergence compared to the unaccelerated version, with only a marginal increase in computational cost per iteration. However, a detailed discussion of these accelerated methods is beyond the scope of this paper.

Algorithm 3.1 Parallel subspace correction method

Given the step size $\tau \in (0, \tau_0]$:
 Choose $u^{(0)} \in V$.
for $n = 0, 1, 2, \dots$ **do**
 for $j \in [N]$ **in parallel do**
 $w_j^{(n+1)} \in \arg \min_{w_j \in V_j} F_j(w_j; u^{(n)})$
 end for
 $u^{(n+1)} = u^{(n)} + \tau \sum_{j=1}^N w_j^{(n+1)}$
end for

Algorithm 3.2 Successive subspace correction method

Choose $u^{(0)} \in V$.
for $n = 0, 1, 2, \dots$ **do**
 for $j = 1, 2, \dots, N$ **do**
 $w_j^{(n+1)} \in \arg \min_{w_j \in V_j} F(w_j; u^{(n+\frac{j-1}{N})})$
 $u^{(n+\frac{j}{N})} = u^{(n+\frac{j-1}{N})} + w_j^{(n+1)}$
 end for
end for

Another type of subspace correction method, known as the successive subspace correction method, is presented in [Algorithm 3.2](#). Also referred to as the multiplicative Schwarz method, this method involves solving the local problems sequentially.

In this paper, we focus on the parallel subspace correction method. We note that the relation between the successive and parallel subspace correction methods for linear problems has been investigated in [\[11, 27, 49\]](#), where it was shown that the convergence rate of the successive method achieves a bound in terms of the parallel method. Moreover, the sharp convergence estimate for the successive subspace correction method, so called the Xu–Zikatanov identity, is well-established in the literature [\[12, 18, 34, 54\]](#). However, tight convergence results for the successive subspace correction method for convex optimization remain open. Some existing estimates for the successive subspace correction method for convex optimization can be found in [\[6, 7, 15, 47, 48\]](#).

4. Convergence analysis for semicoercive problems. In this section, we present a new convergence theory for subspace correction methods for semicoercive convex optimization. Throughout this section, we assume that the energy functional F in the problem [\(3.1\)](#) is semicoercive with respect to a seminorm $|\cdot|$ in the sense of [Proposition 2.3](#), and denote $\mathcal{N} = \ker F = \ker |\cdot|$. To this end, we derive convergence rate estimates for the parallel subspace correction in terms of a seminorm stable decomposition over the subspaces $\{V_j\}$ and the kernel \mathcal{N} . These results align with the sharp theory of singular linear problems established in [\[34, 50\]](#).

Remark 4.1. Even if the energy functional F is semicoercive with a nontrivial kernel, the kernel of each local problem in subspace correction methods may still be trivial, i.e., $V_j \cap \ker F = \{0\}$.

4.1. Descent property. We first show that the parallel subspace correction method for solving the semicoercive problem (3.1) achieves a certain descent property on the energy. This descent property will play a central role in analyzing the convergence rate.

Let V^* be the topological dual space of the reflexive Banach space V . Given $v \in V$ and $w_j \in V_j$, we denote the Bregman distance associated with F between $v + w_j$ and v by $d_j(w_j; v)$, i.e.,

$$(4.1) \quad d_j(w_j; v) = F_j(w_j; v) - F(v) - \langle F'(v), w_j \rangle, \quad v \in V, \quad w_j \in V_j,$$

where $F'(v) \in V^*$ is the Gâteaux derivative of F at v , and $\langle \cdot, \cdot \rangle$ represents the duality pairing on V .

In Lemma 4.2, we present a generalized additive Schwarz lemma [30, 39, 45] for semicoercive problems, which states that the parallel subspace correction method can be viewed as a gradient descent method endowed with a specific nonlinear metric-like functional.

LEMMA 4.2 (generalized additive Schwarz lemma). *Let V be a reflexive Banach space, and let $F: V \rightarrow \mathbb{R}$ be a Gâteaux differentiable, convex, and semicoercive functional with the kernel \mathcal{N} . For $v \in V$, we have*

$$(4.2) \quad \hat{w} := \sum_{j=1}^N \hat{w}_j \in \arg \min_{v \in V} \left\{ \langle F'(v), w \rangle + \inf_{\phi \in \mathcal{N}} \inf_{\sum_{j=1}^N w_j = w + \phi} \sum_{j=1}^N d_j(w_j; v) \right\},$$

where $\hat{w}_j \in V_j$, $j \in [N]$, is given by

$$(4.3) \quad \hat{w}_j \in \arg \min_{w_j \in V_j} F(w_j; v) = \arg \min_{w_j \in V_j} \{ \langle F'(v), w_j \rangle + d_j(w_j; v) \}.$$

Moreover, we have

$$(4.4) \quad \inf_{\phi \in \mathcal{N}} \inf_{\sum_{j=1}^N w_j = \hat{w} + \phi} \sum_{j=1}^N d_j(w_j; v) = \sum_{j=1}^N d_j(\hat{w}_j; v).$$

Proof. Throughout the proof, we write

$$d(w; v) = \inf_{\phi \in \mathcal{N}} \inf_{\sum_{j=1}^N w_j = w + \phi} \sum_{j=1}^N d_j(w_j; v), \quad w \in V.$$

We take any $w \in V$. For any $\phi \in \mathcal{N}$ and $w_j \in V_j$, $j \in [N]$, such that $w = \sum_{j=1}^N w_j + \phi$, we get

$$(4.5) \quad \begin{aligned} \langle F'(v), \hat{w} \rangle + d(\hat{w}; v) &\leq \sum_{j=1}^N (\langle F'(v), \hat{w}_j \rangle + d_j(\hat{w}_j; v)) \\ &\stackrel{(4.3)}{\leq} \sum_{j=1}^N (\langle F'(v), w_j \rangle + d_j(w_j; v)) \\ &= \langle F'(v), w \rangle + \sum_{j=1}^N d_j(w_j; v). \end{aligned}$$

where the first inequality holds because $\hat{w} = 0 + \sum_{j=1}^N \hat{w}_j \in \mathcal{N} + \sum_{j=1}^N V_j$, and the last equality follows from $\langle F'(v), \phi \rangle = 0$. By minimizing the last line of (4.5) over all $(w_j)_{j=1}^N$ and ϕ , we obtain

$$(4.6) \quad \langle F'(v), \hat{w} \rangle + d(\hat{w}; v) \leq \sum_{j=1}^N (\langle F'(v), \hat{w}_j \rangle + d_j(\hat{w}_j; v)) \leq \langle F'(v), w \rangle + d(w; v),$$

which implies (4.2). Then, setting $w = \hat{w}$ in (4.6) yields (4.4). \square

Using Lemma 4.2, we can establish a descent property of the parallel subspace correction method for semicoercive problems, as presented in Lemma 4.3. We note that a corresponding result for coercive problems can be found in [45, Theorem 1].

LEMMA 4.3. *Let V be a reflexive Banach space, and let $F: V \rightarrow \mathbb{R}$ be a Gâteaux differentiable, convex, and semicoercive functional with the kernel \mathcal{N} . In Algorithm 3.1, we have*

$$F(u^{(n+1)}) \leq F(u^{(n)}) + \tau \min_{w \in V} \left\{ \langle F'(u^{(n)}), w \rangle + \inf_{\phi \in \mathcal{N}} \inf_{\sum_{j=1}^N w_j = w + \phi} \sum_{j=1}^N d_j(w_j; u^{(n)}) \right\}, \quad n \geq 0.$$

Proof. Take any $n \geq 0$. Since the strengthened convexity condition (3.4) implies

$$(4.7) \quad F(u^{(n+1)}) \leq (1 - \tau N)F(u^{(n)}) + \tau \sum_{j=1}^N F(u^{(n)} + w_j^{(n+1)}),$$

it suffices to estimate the term $\sum_{j=1}^N F(u^{(n)} + w_j^{(n+1)})$. It follows that

$$(4.8) \quad \begin{aligned} \sum_{j=1}^N F(u^{(n)} + w_j^{(n+1)}) &= NF(u^{(n)}) + \sum_{j=1}^N (\langle F'(u^{(n)}), w_j^{(n+1)} \rangle + d_j(w_j^{(n+1)}; u^{(n)})) \\ &= NF(u^{(n)}) + \min_{w \in V} \left\{ \langle F'(u^{(n)}), w \rangle + \inf_{\phi \in \mathcal{N}} \inf_{\sum_{j=1}^N w_j = w + \phi} \sum_{j=1}^N d_j(w_j; u^{(n)}) \right\}, \end{aligned}$$

where the last inequality is due to Lemma 4.2. Combining (4.7) and (4.8) yields the desired result. \square

A straightforward consequence of Lemma 4.3 is that the energy sequence generated by the parallel subspace correction method is decreasing; see Corollary 4.4.

COROLLARY 4.4. *Let V be a reflexive Banach space, and let $F: V \rightarrow \mathbb{R}$ be a Gâteaux differentiable, convex, and semicoercive functional. In Algorithm 3.1, the energy sequence $\{F(u^{(n)})\}$ is decreasing.*

4.2. Convergence rate analysis. Now, we derive convergence rate estimates for the parallel subspace correction method for solving the semicoercive problem (3.1) based on the descent property presented in Lemma 4.3.

A key ingredient in the convergence rate analysis is a stable decomposition property of the seminorm $|\cdot|$, similar to (3.3). To establish this property, we assume

that the seminorm satisfies the conditions stated in [Lemma 2.1](#), as well as the local condition described in [Assumption 4.5](#). It is worth noting that a similar assumption was made in the theory of singular linear problems; see [[34](#), Equation (A1)].

Assumption 4.5. For each $j \in [N]$, there exists a positive constant C_j such that

$$\inf_{\phi_j \in V_j \cap \ker |\cdot|} \|v_j + \phi_j\| \leq C_j |v_j| \quad \forall v_j \in V_j.$$

Remark 4.6. As discussed in [[34](#), Example A.1], [Assumption 4.5](#) does not follow from (2.3) in general, and needs to be added as an additional assumption. In the same spirit as [Remark 2.2](#), this condition is satisfied by many seminorms commonly encountered in PDEs.

In [Lemma 4.7](#), we show that, under [Assumption 4.5](#), the stable decomposition property holds with respect to the seminorm $|\cdot|$ as well if the kernel of the seminorm is included as an additional subspace.

LEMMA 4.7. *Let $|\cdot|$ be a continuous seminorm on a Banach space V that satisfies (2.3). In addition, suppose that [Assumption 4.5](#) holds. Then we have*

$$\sup_{|w|=1} \inf_{\phi \in \ker |\cdot|} \inf_{\sum_{j=1}^N w_j = w + \phi} \left(\sum_{j=1}^N \|w_j\|^q \right)^{\frac{1}{q}} < \infty,$$

for any $q \in [1, \infty)$, where w and w_j are taken from V and V_j , respectively.

Proof. Given $w \in V$ with $|w| = 1$, let $((\tilde{w}_j)_{j=1}^N, \tilde{\phi})$ be a minimizer of $\sum_{j=1}^N |w_j|^q$ over $w_j \in V_j$, $j \in [N]$, and $\phi \in \mathcal{N} := \ker |\cdot|$ such that $\sum_{j=1}^N w_j = w + \phi$. Since $|\cdot|$ is invariant under addition by an element of \mathcal{N} , we may assume that

$$\|\tilde{w}_j\| = \inf_{\phi_j \in V_j \cap \mathcal{N}} \|\tilde{w}_j + \phi_j\|.$$

It follows that

$$\begin{aligned} \inf_{\phi \in \mathcal{N}} \inf_{\sum_{j=1}^N w_j = w + \phi} \left(\sum_{j=1}^N \|w_j\|^q \right)^{\frac{1}{q}} &\leq \left(\sum_{j=1}^N \|\tilde{w}_j\|^q \right)^{\frac{1}{q}} \\ &\leq C \left(\sum_{j=1}^N |\tilde{w}_j|^q \right)^{\frac{1}{q}} = C \inf_{\phi \in \mathcal{N}} \inf_{\sum_{j=1}^N w_j = w + \phi} \left(\sum_{j=1}^N |w_j|^q \right)^{\frac{1}{q}} \end{aligned}$$

for some positive constant C , where the second inequality is due to [Assumption 4.5](#). Therefore, it suffices to prove

$$(4.9) \quad \sup_{|w|=1} \inf_{\phi \in \ker |\cdot|} \inf_{\sum_{j=1}^N w_j = w + \phi} \left(\sum_{j=1}^N |w_j|^q \right)^{\frac{1}{q}} < \infty.$$

With an abuse of notation, we write

$$V_j / \mathcal{N} = \{v_j + \mathcal{N} : v_j \in V_j\}.$$

Since $V/\mathcal{N} = \sum_{j=1}^N V_j/\mathcal{N}$ and both V/\mathcal{N} and each V_j/\mathcal{N} are Banach spaces, the open mapping theorem implies that

$$\sup_{\|w+\mathcal{N}\|_{V/\mathcal{N}}=1} \inf_{\sum_{j=1}^N (w_j+\mathcal{N})=w+\mathcal{N}} \left(\sum_{j=1}^N \|w_j+\mathcal{N}\|_{V_j/\mathcal{N}}^q \right)^{\frac{1}{q}} < \infty,$$

where w and w_j are taken from V and V_j , respectively (cf. (3.3)). From Lemma 2.1, we know that $|\cdot|$ is equivalent to $\|\cdot+\mathcal{N}\|_{V/\mathcal{N}}$. Moreover, $\sum_{j=1}^N (w_j+\mathcal{N})=w+\mathcal{N}$ is equivalent to $\sum_{j=1}^N w_j = w + \phi$ for some $\phi \in \mathcal{N}$. Accordingly, we obtain (4.9), which completes the proof. \square

Meanwhile, from Lemma 4.3, we observe that the convergence behavior of the parallel subspace correction method depends on the infimum of the sum of the local functionals $d_j(w_j; v)$, $j \in [N]$. To establish an upper bound for this infimum, we require a smoothness assumption on each $d_j(w_j; v)$, which ensures that $d_j(w_j; v)$ can be bounded above by a power of the norm $\|w_j\|^q$; see Assumption 4.8. We note that, throughout this paper, we adopt the convention $0/0 = 0$ for arguments of sup and $0/0 = \infty$ for arguments of inf.

Assumption 4.8 (local smoothness). For some $q > 1$, each $d_j(w_j; v)$, $j \in [N]$, satisfies the following: for any $|\cdot|$ -bounded convex subset $K \subset V$ and $\|\cdot\|$ -bounded convex subset $K_j \subset V_j$ satisfying $0 \in K_j$, we have

$$\sup_{v \in K, w_j \in K_j} \frac{d_j(w_j; v)}{\|w_j\|^q} < \infty.$$

In the case of exact local problems (3.6), an easy-to-check sufficient condition for Assumption 4.8 is the weak smoothness [42] of the energy functional F , which is valid in many applications involving nonlinear PDEs (see, e.g., [19, 30, 48]). We summarize this result in Proposition 4.9. In what follows, given $v, w \in V$, we denote the Bregman distance associated with F between $v+w$ and v by $d_F(w; v)$, i.e.,

$$(4.10) \quad d_F(w; v) = F(v+w) - F(v) - \langle F'(v), w \rangle, \quad v, w \in V.$$

PROPOSITION 4.9. *Let V be a reflexive Banach space, and let $F: V \rightarrow \mathbb{R}$ be a Gâteaux differentiable, convex, and semicoercive functional with respect to a seminorm $|\cdot|$ in the sense of Proposition 2.3. Suppose that each $d_j(w_j; v)$, $j \in [N]$, is given by (3.6) and (4.1). For some $q > 1$, if F is locally q -weakly smooth, i.e., if*

$$(4.11) \quad \sup_{v, v+w \in K} \frac{d_F(w; v)}{\|w\|^q} < \infty,$$

for any $\|\cdot\|$ -bounded convex subset K of V , then Assumption 4.8 holds.

Proof. Throughout this proof, let C denote a general positive constant. First, we prove that the local smoothness with respect to the norm $\|\cdot\|$ implies the local smoothness with respect to the seminorm $|\cdot|$ as well, i.e.,

$$(4.12) \quad \sup_{v, v+w \in K} \frac{d_F(w; v)}{|w|^q} < \infty,$$

for any $|\cdot|$ -bounded convex subset K of V . Suppose that (4.11) holds, and take any $|\cdot|$ -bounded convex subset K of V . We define the set \widehat{K} as the convex hull of the

following set:

$$\left\{ v + \phi : v \in K, \phi = \arg \min_{\psi \in \ker |\cdot|} \|v + \psi\| \right\}.$$

Then (2.3) implies that \widehat{K} is $\|\cdot\|$ -bounded. Hence, for any $v, v + w \in K$, we have

$$d_F(w; v) = d_F(\hat{w}; \hat{v}) \stackrel{(4.11)}{\leq} C \|\hat{w}\|^q \stackrel{(2.3)}{\leq} C |\hat{w}|^q = C |w|^q,$$

where $\hat{v} \in \widehat{K}$ is defined as

$$\hat{v} = v + \phi, \quad \phi = \arg \min_{\psi \in \ker |\cdot|} \|v + \psi\|,$$

and $\hat{w} \in \widehat{K}$ is defined similarly. This implies that (4.12) holds.

Now, it is enough to show that (4.12) implies Assumption 4.8. Suppose that (4.12) holds. Take any $j \in [N]$, $|\cdot|$ -bounded convex subset K of V , and $\|\cdot\|$ -bounded convex subset K_j of V_j such that $0 \in K_j$. It is straightforward to verify that $K + K_j$ is $|\cdot|$ -bounded. Moreover, if $v \in K$ and $w_j \in K_j$, then we have $v, v + w_j \in K + K_j$. Applying (4.12) with the set $K + K_j$ yields the desired result. \square

In Lemma 4.10, we combine Lemma 4.7 and Assumption 4.8 to show that the infimum of the sum of the local functionals $d_j(w_j; v)$ appeared in Lemma 4.3 can be bounded above a power of the seminorm $|w|^q$.

LEMMA 4.10 (stable decomposition). *Let V be a reflexive Banach space, and let $F: V \rightarrow \mathbb{R}$ be a Gâteaux differentiable, convex, and semicoercive functional with respect to a seminorm $|\cdot|$ in the sense of Proposition 2.3. Suppose that Assumptions 4.5 and 4.8 hold. For any $|\cdot|$ -bounded convex subset $K \subset V$, the following holds:*

$$(4.13) \quad C_K := q \sup_{v, v+w \in K} \inf_{\phi \in \ker |\cdot|} \inf_{\sum_{j=1}^N w_j = w + \phi} \frac{\sum_{j=1}^N d_j(w_j; v)}{|w|^q} < \infty.$$

Proof. We take any $|\cdot|$ -bounded convex subset K of V , and write

$$M_K := \sup_{v \in K} |v| < \infty.$$

Choose any $v, v + w \in K$. We first observe that

$$|w| \leq |v| + |v + w| \leq 2M_K.$$

Let $((\tilde{w}_j)_{j=1}^N, \tilde{\phi})$ be a minimizer of $\sum_{j=1}^N \|w_j\|^q$ over $w_j \in V_j$, $j \in [N]$, and $\phi \in \mathcal{N} := \ker |\cdot|$ such that $\sum_{j=1}^N w_j = w + \phi$. By Lemma 4.7, each $\|\tilde{w}_j\|$ is bounded by a constant depending on K only. Namely, we have

$$\|\tilde{w}_j\| \leq \left(\sum_{j=1}^N \|\tilde{w}_j\|^q \right)^{\frac{1}{q}} \leq C |w| \leq 2CM_K,$$

for some $C > 0$. If we set $K_j = \{w_j \in V_j : \|w_j\| \leq 2CM_K\}$ for each $j \in [N]$ in

Assumption 4.8, then we obtain

$$L_{j,K} := \sup_{v \in K, w_j \in K_j} \frac{d_j(w_j; v)}{\|w_j\|^q} < \infty.$$

Note that the constant $L_{j,K}$ depends on j , q , and K only. It follows that

$$\inf_{\phi \in \mathcal{N}} \inf_{\sum_{j=1}^N w_j = w + \phi} d_j(w_j; v) \leq \sum_{j=1}^N d_j(\tilde{w}_j; v) \leq \sum_{j=1}^N L_{j,K} \|\tilde{w}_j\|^q \leq C^q \left(\max_{j \in [N]} L_{j,K} \right) |w|^q.$$

As the constant in the rightmost-hand side of the above equation depends on q and K only, the proof is complete. \square

Given the initial iterate $u^{(0)} \in V$ of Algorithm 3.1, we define

$$(4.14a) \quad K_0 = \{v \in V : F(v) \leq F(u^{(0)})\},$$

$$(4.14b) \quad R_0 = \sup_{v \in K_0} |v - u|.$$

The convexity and semicoercivity of F implies that K_0 is $|\cdot|$ -bounded and convex, and that $R_0 < \infty$. Moreover, Corollary 4.4 implies that the sequence $\{u^{(n)}\}$ generated by Algorithm 3.1 is contained in K_0 . Consequently, thanks to Lemma 4.10, we have

$$\inf_{\phi \in \mathcal{N}} \inf_{\sum_{j=1}^N w_j = w + \phi} \sum_{j=1}^N d_j(w_j; u^{(n)}) \leq \frac{C_{K_0}}{q} |w|^q,$$

for any $w \in V$ such that $u^{(n)} + w \in K$ and $n \geq 0$, where C_{K_0} was given in (4.13). Using a similar argument as in [39, 41], we are able to derive the following convergence theorem for the parallel subspace correction method for solving (3.1). A proof of Theorem 4.11 can be found in Appendix B.

THEOREM 4.11. *Let V be a reflexive Banach space, and let $F: V \rightarrow \mathbb{R}$ be a Gâteaux differentiable, convex, and semicoercive functional with respect to a seminorm $|\cdot|$ in the sense of Proposition 2.3. Suppose that Assumption 4.8 holds. In Algorithm 3.1, let $\zeta_n = F(u^{(n)}) - F(u)$ for $n \geq 0$. If $\zeta_0 > C_{K_0} R_0^q$, then we have*

$$\zeta_1 \leq \left(1 - \tau \left(1 - \frac{1}{q} \right) \right) \zeta_0,$$

where C_{K_0} and R_0 were given in (4.13) and (4.14). Otherwise, we have

$$\zeta_n \leq \frac{C}{(n + (C/\zeta_0)^{1/\beta})^\beta}, \quad n \geq 0,$$

where

$$\beta = q - 1, \quad C = \left(\frac{q}{\tau} \right)^{q-1} C_{K_0} R_0^q.$$

In the same spirit as the convergence theory for coercive problems [39, 41, 45], we are able to obtain an improved convergence rate of the parallel subspace correction method if we additionally assume that the energy functional F is sharp [46] around a minimizer. We formally summarize this sharpness assumption in Assumption 4.12.

Assumption 4.12 (sharpness). For some $p > 1$, the function F satisfies the following: for any $|\cdot|$ -bounded convex subset $K \subset V$ satisfying $u \in K$, we have

$$(4.15) \quad \mu_K := p \inf_{v \in K} \frac{F(v) - F(u)}{|v - u|^p} > 0.$$

In [Theorem 4.13](#), we provide convergence rate estimates for the parallel subspace correction method under the additional assumption described in [Assumption 4.12](#). A proof of [Theorem 4.13](#) is given in [Appendix B](#).

THEOREM 4.13. *Let V be a reflexive Banach space, and let $F: V \rightarrow \mathbb{R}$ be a Gâteaux differentiable, convex, and semicoercive functional with respect to a seminorm $|\cdot|$ in the sense of [Proposition 2.3](#). Suppose that [Assumptions 4.8](#) and [4.12](#) hold. In [Algorithm 3.1](#), let $\zeta_n = F(u^{(n)}) - F(u)$ for $n \geq 0$. Then we have the following:*

(a) *In the case $p = q$, we have*

$$\zeta_n \leq \left(1 - \tau \left(1 - \frac{1}{q} \right) \min \left\{ 1, \frac{\mu_{K_0}}{qC_{K_0}} \right\}^{\frac{1}{q-1}} \right)^n \zeta_0, \quad n \geq 0,$$

where C_{K_0} and μ_{K_0} were given in [\(4.13\)](#), [\(4.14a\)](#), and [\(4.15\)](#).

(b) *In the case $p > q$, if $\zeta_0 > \left(\frac{p}{\mu_{K_0}} \right)^{\frac{q}{p-q}} C_{K_0}^{\frac{p}{p-q}}$, then we have*

$$\zeta_1 \leq \left(1 - \tau \left(1 - \frac{1}{q} \right) \right) \zeta_0.$$

Otherwise, we have

$$\zeta_n \leq \frac{C}{(n + (C/\zeta_0)^{1/\beta})^\beta}, \quad n \geq 0,$$

where

$$\beta = \frac{p(q-1)}{p-q}, \quad C = \left(\frac{pq}{(p-q)\tau} \right)^{\frac{p(q-1)}{p-q}} \left(\frac{p}{\mu_{K_0}} \right)^{\frac{q}{p-q}} C_{K_0}^{\frac{p}{p-q}}.$$

Remark 4.14. By setting $\mathcal{N} = \{0\}$ in [Theorems 4.11](#) and [4.13](#), we recover the convergence results for the parallel subspace correction method for coercive problems [\[39, 45\]](#).

5. Convergence analysis for nearly semicoercive problems. In [\[33, 51\]](#), it was proven that subspace correction methods for nearly singular linear systems on Hilbert spaces achieve convergence rate estimates that are independent of the nearly singular behavior of the systems, assuming appropriate conditions on the space decompositions. Namely, if the near null space can be decomposed into a sum of local near null spaces, parameter-independent estimates for the convergence rates can be established [\[33, Theorems 4.2 and 4.3\]](#). In this section, we extend these results to nearly semicoercive convex optimization problems in Banach spaces, thereby enabling applications to a broader class of nonlinear problems.

5.1. Orthogonal decompositions of Banach spaces. A key tool used in the convergence analysis of nearly singular linear systems on Hilbert spaces in [\[33, 51\]](#) is the fact that the kernel of the singular part admits an orthogonal complement; see [\[33,](#)

Lemma 4.5]. Unfortunately, in general Banach spaces, not every closed subspace has a complement [36], and moreover, there is no inherent orthogonality structure. Therefore, to analyze nearly coercive problems in Banach spaces, we must employ alternative tools.

For this purpose, we present decomposition results for Banach spaces based on a generalized notion of orthogonality, introduced in [3, 4]. To achieve this, we require a stronger assumption on Banach spaces than just reflexivity. Specifically, let V be a uniformly convex and uniformly smooth Banach space, which ensures that V is reflexive [20, Theorems II.2.9 and II.2.15]. It is also worth noting that many Sobolev spaces associated with nonlinear PDEs are uniformly convex and uniformly smooth [1].

The normalized duality mapping $J: V \rightrightarrows V^*$ on V is defined as

$$J(v) = \{v^* \in V^* : \langle v^*, v \rangle = \|v^*\| \|v\|, \|v^*\| = \|v\|\}, \quad v \in V.$$

Note that the uniform convexity and uniform smoothness of V ensure that J is bijective [20, Proposition II.3.6], allowing us to identify J as a mapping $J: V \rightarrow V^*$. Similarly, we define $J^*: V^* \rightrightarrows V$ as the normalized duality mapping on V^* :

$$J^*(v^*) = \{v \in V : \langle v^*, v \rangle = \|v^*\| \|v\|, \|v\| = \|v^*\|\}, \quad v^* \in V^*.$$

Then, we have $J^* = J^{-1}$.

Given a closed subspace \mathcal{M} of V , we denote its polar set as \mathcal{M}° :

$$\mathcal{M}^\circ = \{v^* \in V^* : \langle v^*, v \rangle = 0 \text{ for all } v \in \mathcal{M}\}.$$

The following proposition provides an orthogonal decomposition of V into \mathcal{M} and its complement, which is a nonlinear manifold in general [4], involving \mathcal{M}° and the normalized duality mapping J^* . While this result appeared previously in [4, Theorem 2.13], we include the proof of this result for the sake of completeness.

PROPOSITION 5.1. *Let V be a uniformly convex and uniformly smooth Banach space, and let \mathcal{M} be a closed subspace of V . Then, V admits a decomposition $V = \mathcal{M} + J^*\mathcal{M}^\circ$, which satisfies the following:*

- (i) *Each element $v \in V$ has a unique decomposition $v = \phi + \xi$ with $\phi \in \mathcal{M}$ and $\xi \in J^*\mathcal{M}^\circ$. Moreover, we have*

$$(5.1) \quad \phi = \arg \min_{w \in \mathcal{M}} \|v - w\|.$$

- (ii) *$\langle J\xi, \phi \rangle = 0$ for any $\phi \in \mathcal{M}$ and $\xi \in J^*\mathcal{M}^\circ$.*

- (iii) *$\mathcal{M} \cap J^*\mathcal{M}^\circ = \{0\}$.*

Proof. The validity of (ii) follows directly from the definition of the polar set \mathcal{M}° . To prove (iii), let $v \in \mathcal{M} \cap J^*\mathcal{M}^\circ$. Since $v \in \mathcal{M}$ and $Jv = (J^*)^{-1}v \in \mathcal{M}^\circ$, we have $\langle Jv, v \rangle = 0$. This implies $v = 0$ because of the strict monotonicity of J [20, Theorem II.1.8].

Finally, we prove (i) using an argument similar as in [3]. Take any $v \in V$, and we define $\phi \in \mathcal{M}$ as in (5.1). The optimality condition of ϕ reads as

$$\langle J(v - \phi), w \rangle = 0 \quad \forall w \in \mathcal{M},$$

which is equivalent to $J(v - \phi) \in \mathcal{M}^\circ$. Thus, we have $\xi := v - \phi \in J^*\mathcal{M}^\circ$, leading to the desired decomposition $v = \phi + \xi \in \mathcal{M} + J^*\mathcal{M}^\circ$. The uniqueness of this decomposition follows directly from (iii). \square

As a direct consequence of [Proposition 5.1](#), we obtain the following corollary, which will play an important role in the convergence analysis of nearly semicoercive problems.

COROLLARY 5.2. *Let V be a uniformly convex and uniformly smooth Banach space, and let \mathcal{M} be a closed subspace of V . Then we have the following:*

(a) *For any $q \geq 1$, there exists a positive constant C_q , depending only on q , such that*

$$(5.2) \quad \|\phi\|^q + \|\xi\|^q \leq C_q \|\phi + \xi\|^q, \quad \phi \in \mathcal{M}, \xi \in J^* \mathcal{M}^\circ.$$

(b) *For any $\xi \in J^* \mathcal{M}^\circ$, we have*

$$\|\xi\| = \min_{w \in \mathcal{M}} \|\xi + w\|.$$

Proof. Take any $\phi \in \mathcal{M}$ and $\xi \in J^* \mathcal{M}^\circ$. By [Proposition 5.1](#), we have $\phi = \arg \min_{w \in \mathcal{M}} \|\phi + \xi - w\|$, which implies $\|\xi\| \leq \|\phi + \xi\|$. Since ϕ is arbitrary, we deduce that (b) holds. To show (a), we observe that $\|\xi\| \leq \|\phi + \xi\|$ and that $\|\phi\| \leq \|\phi + \xi\| + \|\xi\| \leq 2\|\phi + \xi\|$. Hence, we get $\|\phi\|^q + \|\xi\|^q \leq (2^q + 1)\|\phi + \xi\|^q$, which completes the proof. \square

Remark 5.3. If V is a Hilbert space, then [Proposition 5.1](#) holds naturally when we identify the topological dual space V^* with V , and the duality pairing with the inner product. Moreover, in this setting, we have the Pythagorean property, which is a special case of [Corollary 5.2](#); see [[13](#), Corollary 5.4].

5.2. Parameter-independent estimates. Now, we consider the model nearly semicoercive convex optimization of the form

$$(5.3) \quad \min_{v \in V} \{F(v) := F_0(v) + \epsilon F_1(v)\},$$

where V is a uniformly convex and uniformly smooth Banach space, $F_0: V \rightarrow \mathbb{R}$ and $F_1: V \rightarrow \mathbb{R}$ are Gâteaux differentiable and convex functionals, and $\epsilon > 0$. We further assume that F_0 is semicoercive with respect to a seminorm $|\cdot|$ in the sense of [Proposition 2.3](#), and that F_1 is coercive.

In subspace correction methods (see [Algorithms 3.1](#) and [3.2](#)) for solving (5.3) under the space decomposition (3.2), the local energy functionals F_j , $j \in [N]$, must be specified. We assume that each F_j is given by

$$F_j(w_j; v) = F_{0,j}(w_j; v) + \epsilon F_{1,j}(w_j; v), \quad v \in V, w_j \in V_j,$$

where $F_{0,j}$ and $F_{1,j}$ are specified in the case of exact local problems as follows:

$$(5.4) \quad \begin{aligned} F_{0,j}(w_k; v) &= F_0(v + w_j), \\ F_{1,j}(w_k; v) &= F_1(v + w_j), \end{aligned} \quad v \in V, w_j \in V_j.$$

The general case of inexact local problems [[39](#), [45](#)] will be considered in [Appendix A](#). Similar as in (4.1), we define

$$\begin{aligned} d_{0,j}(w_j; v) &= F_{0,j}(w_j; v) - F(v) - \langle F'_0(v), w_j \rangle, \\ d_{1,j}(w_j; v) &= F_{1,j}(w_j; v) - F(v) - \langle F'_1(v), w_j \rangle, \end{aligned} \quad v \in V, w_j \in V_j.$$

Then we have

$$d_j(w_j; v) = d_{0,j}(w_j; v) + \epsilon d_{1,j}(w_j; v), \quad v \in V, w_j \in V_j,$$

where $d_j(w_j; v)$ was given in (4.1).

To guarantee the convergence of the parallel subspace correction method for solving (5.3), we require a smoothness assumption on each $d_{0,j}(w_j; v)$ and $d_{1,j}(w_j; v)$, analogous to Assumption 4.8. This assumption is formally stated in Assumption 5.4.

Assumption 5.4 (local smoothness). For some $q > 1$, each $d_{0,j}(w_j; v)$ and $d_{1,j}(w_j; v)$, $j \in [N]$, satisfy the following: for any $\|\cdot\|$ -bounded convex subset $K \subset V$ and $\|\cdot\|$ -bounded convex subset $K_j \subset V_j$ satisfying $0 \in K_j$, we have

$$\sup_{v \in K, w_j \in K_j} \frac{d_{0,j}(w_j; v)}{\|w_j\|^q} < \infty \quad \text{and} \quad \sup_{v \in K, w_j \in K_j} \frac{d_{1,j}(w_j; v)}{\|w_j\|^q} < \infty.$$

For $q > 1$, we define an ϵ -dependent norm $\|\cdot\|_{\epsilon,q}$ on V as

$$\|v\|_{\epsilon,q} := (|v|^q + \epsilon\|v\|^q)^{\frac{1}{q}}, \quad v \in V.$$

It is straightforward to verify that the norm $\|\cdot\|_{\epsilon,q}$ is equivalent to the original norm $\|\cdot\|$.

Under the smoothness assumption stated in Assumption 5.4, Theorems 4.11 and 4.13 (see also [39, 41, 45]) indicate that a key factor determining the convergence rate of the parallel subspace correction method (Algorithm 3.1) for solving (5.3) is the following constant:

$$(5.5) \quad C_{K_0} := q \sup_{v, v+w \in K_0} \inf_{\sum_{j=1}^N w_j = w} \frac{\sum_{j=1}^N d_j(w_j; v)}{\|w\|_{\epsilon,q}^q},$$

where the set K_0 was given in (4.14a), and each w_j belongs to V_j .

In order to derive an ϵ -independent upper bound for C_{K_0} , we need to impose additional assumptions on the space decomposition and the local problems. The first of these assumptions, summarized in Assumption 5.5, requires that the kernel of the semicoercive functional F_0 in (5.3) can be decomposed into a sum of local kernels (cf. [33, equation (A1)]).

Assumption 5.5 (kernel decomposition). The kernel $\mathcal{N} = \ker F_0$ of the semicoercive functional F_0 in (5.3) admits a decomposition $\mathcal{N} = \sum_{j=1}^N (V_j \cap \mathcal{N})$.

The second assumption, summarized in Assumption 5.6, is that the functional $d_{1,j}(\cdot; v)$ satisfies a triangle inequality-like property.

Assumption 5.6 (triangle inequality-like property). For any bounded convex subset $K \subset V$, there exists a positive constant $C_{K,\text{tri}}$ such that

$$(5.6) \quad d_{1,j}(v_j + w_j; v) \leq C_{K,\text{tri}} (d_{1,j}(v_j; v) + d_{1,j}(w_j; v)), \quad v \in K, v_j, w_j \in V_j, j \in [N].$$

In the case of exact local problems (5.4), a straightforward sufficient condition for Assumption 5.6 is that the Bregman distance d_{F_1} associated with the functional F_1 (cf. (4.10)) satisfies the following triangle-inequality-like property:

$$d_{F_1}(w_1 + w_2; v) \leq C_{K,\text{tri}} (d_{F_1}(w_1; v) + d_{F_1}(w_2; v)), \quad v \in K, w_1, w_2 \in V,$$

for some positive constant $C_{K,\text{tri}}$. This property is indeed satisfied by a broad class of convex functionals, making it a practical and verifiable condition. We provide a detailed discussion of this property in Appendix C.

Now, we are ready to present the main result of this section, [Theorem 5.7](#), which provides an ϵ -independent convergence rate estimate of the parallel subspace correction method for solving (5.3). We note that [Theorem 5.7](#) generalizes the existing result [51, Theorem 3.1] for nearly singular linear problems to the context of nearly semicoercive convex optimization problems.

THEOREM 5.7. *Let V be a uniformly convex and uniformly smooth Banach space. Suppose that [Assumptions 5.4](#) to [5.6](#) hold. Then the constant C_{K_0} given in (5.5) has an upper bound independent of ϵ . More precisely, we have*

$$C_{K_0} \leq qC_q \sup_{\substack{v \in K_0, \phi \in \mathcal{N}, \xi \in J^*\mathcal{N}^\circ, \\ v + \phi + \xi \in K_0}} \left[C_q C_{K_0, \text{tri}} \inf_{\sum_{j=1}^N \phi_j = \phi} \frac{\sum_{j=1}^N d_{1,j}(\phi_j; v)}{\|\phi\|^q} \right. \\ \left. + \inf_{\sum_{j=1}^N \xi_j = \xi} \left(\frac{\sum_{j=1}^N d_{0,j}(\xi_j; v)}{\|\xi\|^q} + C_q C_{K_0, \text{tri}} \frac{\sum_{j=1}^N d_{1,j}(\xi_j; v)}{\|\xi\|^q} \right) \right] < \infty,$$

where C_q and $C_{K_0, \text{tri}}$ were given in (5.2) and (5.6), respectively, and each ϕ_j and ξ_j are taken from $V_j \cap \mathcal{N}$ and V_j , respectively.

Proof. Invoking [Proposition 5.1](#), we have

$$(5.7) \quad C_{K_0} = q \sup_{\substack{v \in K_0, \phi \in \mathcal{N}, \xi \in J^*\mathcal{N}^\circ, \\ v + \phi + \xi \in K_0}} \inf_{\sum_{j=1}^N \phi_j = \phi, \sum_{j=1}^N \xi_j = \xi} \frac{\sum_{j=1}^N d_j(\phi_j + \xi_j; v)}{\|\phi + \xi\|_{\epsilon, q}^q}.$$

To estimate the right-hand side of (5.7), we choose any $v \in K_0$, $\phi \in \mathcal{N}$, and $\xi \in J^*\mathcal{N}^\circ$ such that $v + \phi + \xi \in K_0$. For simplicity, we assume that $\phi \neq 0$ and $\xi \neq 0$; the cases where either $\phi = 0$ or $\xi = 0$ are straightforward. Thanks to (3.2) and [Assumption 5.5](#), we can decompose ϕ and ξ as $\phi = \sum_{j=1}^N \phi_j$ and $\xi = \sum_{j=1}^N \xi_j$, where each ϕ_j belongs to $V_j \cap \mathcal{N}$ and each ξ_j belongs to V_j .

We first deduce an upper bound for $\sum_{j=1}^N d_j(\phi_j + \xi_j; v)$ (cf. [33, Lemma 4.5]):

$$(5.8) \quad \sum_{j=1}^N d_j(\phi_j + \xi_j; v) = \sum_{j=1}^N d_{0,j}(\xi_j; v) + \epsilon \sum_{j=1}^N d_{1,j}(\phi_j + \xi_j; v) \\ \leq \sum_{j=1}^N d_{0,j}(\xi_j; v) + C_{K_0, \text{tri}} \epsilon \sum_{j=1}^N (d_{1,j}(\phi_j; v) + d_{1,j}(\xi_j; v)),$$

where the inequality is due to [Assumption 5.6](#). Next, by invoking [Corollary 5.2\(a\)](#), we obtain a lower bound for $\|\phi + \xi\|_{\epsilon, q}^q$ as follows:

$$(5.9) \quad \|\phi + \xi\|_{\epsilon, q}^q = |\xi|^q + \epsilon \|\phi + \xi\|^q \geq |\xi|^q + C_q^{-1} \epsilon (\|\phi\|^q + \|\xi\|^q).$$

Combining (5.8) and (5.9) yields

$$(5.10) \quad \frac{\sum_{j=1}^N d_j(\phi_j + \xi_j; v)}{\|\phi + \xi\|_{\epsilon, q}^q} \leq \frac{\sum_{j=1}^N d_{0,j}(\xi_j; v)}{|\xi|^q} + C_q C_{K_0, \text{tri}} \sum_{j=1}^N \left(\frac{d_{1,j}(\phi_j; v)}{\|\phi\|^q} + \frac{d_{1,j}(\xi_j; w)}{\|\xi\|^q} \right).$$

Since the decompositions $\phi = \sum_{j=1}^N \phi_j$ and $\xi = \sum_{j=1}^N \xi_j$ were arbitrarily chosen, by invoking (5.7) and (5.10), we obtain

$$(5.11) \quad C_{K_0} \leq q \sup_{\substack{v \in K_0, \phi \in \mathcal{N}, \xi \in J^* \mathcal{N}^\circ, \\ v + \phi + \xi \in K_0}} \left[C_q C_{K_0, \text{tri}} \inf_{\sum_{j=1}^N \phi_j = \phi} \frac{\sum_{j=1}^N d_{1,j}(\phi_j; v)}{\|\phi\|^q} \right. \\ \left. + \inf_{\sum_{j=1}^N \xi_j = \xi} \left(\frac{\sum_{j=1}^N d_{0,j}(\xi_j; v)}{|\xi|^q} + C_q C_{K_0, \text{tri}} \frac{\sum_{j=1}^N d_{1,j}(\xi_j; v)}{\|\xi\|^q} \right) \right].$$

It remains to show that the right-hand side of (5.11) is finite. We write

$$M_{K_0} := \sup_{v \in K_0} \|v\| < \infty.$$

Then, for any $v \in K_0$, $\phi \in \mathcal{N}$, and $\xi \in J^* \mathcal{N}^\circ$ such that $v + \phi + \xi \in K_0$, we have

$$\|\xi\| \stackrel{(i)}{\leq} \|\phi + \xi\| \leq \|v\| + \|v + \phi + \xi\| \leq 2M_{K_0}$$

and

$$\|\phi\| \leq \|v + \phi + \xi\| + \|v\| + \|\xi\| \leq 3M_{K_0},$$

where (i) is due to Corollary 5.2(b). In addition, thanks to (2.3) and Corollary 5.2(b), we have

$$|\xi|^q \geq C \inf_{w \in \mathcal{N}} \|\xi + w\|^q = C \|\xi\|^q.$$

for some positive constant C . With these bounds, the finiteness of the right-hand side of (5.11) follows directly from a similar argument as in the proof of Lemma 4.10. \square

As presented in Theorem 4.13, an enhanced convergence rate estimate for the parallel subspace correction method can be achieved if we have an additional assumption that the energy functional is sharp. In the following, we present a relevant result for nearly semicoercive problems. First, we present in Assumption 5.8, which introduces an appropriate sharpness assumption for the nearly semicoercive problem (5.3), where d_{F_0} and d_{F_1} are defined in the same manner as in (4.10). It is readily observed that Assumption 5.8 provides a sufficient condition to guarantee that the energy functional F in (5.3) satisfies Assumption 4.12.

Assumption 5.8 (uniform convexity). For some $p > 1$, we have the following:

(a) For any $|\cdot|$ -bounded convex subset K of V , we have

$$\mu_{0,K} := p \inf_{v, v+w \in K} \frac{d_{F_0}(w; v)}{|w|^p} > 0.$$

(b) For any $\|\cdot\|$ -bounded convex subset K of V , we have

$$\mu_{1,K} := p \inf_{v, v+w \in K} \frac{d_{F_1}(w; v)}{\|w\|^p} > 0.$$

Under Assumptions 5.4 and 5.8, Theorem 4.13 indicates that the constant μ_{K_0} given below is a critical factor in determining the convergence rate of the subspace

correction method, alongside C_{K_0} :

$$(5.12) \quad \mu_{K_0} := p \inf_{v \in K_0} \frac{F(v) - F(u)}{\|v - u\|_{\epsilon, q}^p},$$

where the set K_0 was given in (4.14a). The following theorem states that μ_{K_0} has a lower bound independent of ϵ if ϵ is small enough.

THEOREM 5.9. *Let V be a uniformly convex and uniformly smooth Banach space. Suppose that Assumptions 5.4 and 5.8 hold with $p \geq q$. If $\epsilon \in (0, 1)$, then the constant μ_{K_0} given in (5.12) has a lower bound independent of ϵ .*

Proof. Take any $v \in K_0 \setminus \{0\}$, and let $w = v - u$. We derive an upper bound for $\|w\|_{\epsilon, q}^p$ as follows:

$$(5.13) \quad \|w\|_{\epsilon, q}^p = (|w|^q + \epsilon \|w\|^q)^{\frac{p}{q}} \leq 2^{\frac{p}{q}-1} (|w|^p + \epsilon^{\frac{p}{q}} \|w\|^p) \leq 2^{\frac{p}{q}-1} (|w|^p + \epsilon \|w\|^p),$$

where the first inequality follows from the elementary inequality

$$(a + b)^s \leq 2^{s-1} (a^s + b^s), \quad a, b \geq 0, \quad s \geq 1,$$

and the second inequality is because of $\epsilon \in (0, 1)$. Note that the set K_0 is both $|\cdot|$ -bounded and $\|\cdot\|$ -bounded. Hence, it follows that

$$\mu_{K_0} \stackrel{(5.13)}{\geq} \frac{d_{F_0}(w; u) + \epsilon d_{F_1}(w; u)}{2^{\frac{p}{q}-1} (|w|^p + \epsilon \|w\|^p)} \stackrel{(i)}{\geq} \frac{\mu_{0, K_0} |w|^p + \epsilon \mu_{1, K_0} \|w\|^p}{p 2^{\frac{p}{q}-1} (|w|^p + \epsilon \|w\|^p)} \geq \frac{\min\{\mu_{0, K_0}, \mu_{1, K_0}\}}{p 2^{\frac{p}{q}-1}},$$

which completes the proof, where (i) is due to Assumption 5.8. \square

6. Applications. In this section, we present several applications of the proposed convergence theory for subspace correction methods. We analyze the convergence of two-level domain decomposition methods for solving some nonlinear PDEs with associated energy functionals that are either semicoercive or nearly semicoercive.

Let Ω be a bounded polyhedral domain in \mathbb{R}^d , and let \mathcal{T}_h be a quasi-uniform triangulation of Ω with h the characteristic element diameter. We denote by $S_h(\Omega)$ the continuous and piecewise linear finite element space defined on \mathcal{T}_h :

$$S_h(\Omega) = \{v \in C(\Omega) : v|_T \in \mathbb{P}_1(T) \text{ for all } T \in \mathcal{T}_h\}.$$

Assume that we also have a quasi-uniform triangulation \mathcal{T}_H , with \mathcal{T}_h a refinement of \mathcal{T}_H . We define the coarse finite element space $S_H(\Omega)$ similarly to $S_h(\Omega)$:

$$S_H(\Omega) = \{v \in C(\Omega) : v|_T \in \mathbb{P}_1(T) \text{ for all } T \in \mathcal{T}_H\}.$$

Since $S_H(\Omega) \subset S_h(\Omega)$, the natural embedding operator $I_0: S_H(\Omega) \rightarrow S_h(\Omega)$ is well-defined.

Let $\{\Omega_j\}_{j=1}^N$ be a quasi-uniform overlapping domain decomposition of Ω , where each subdomain Ω_j is a union of \mathcal{T}_h -elements and has diameter of order H . The overlap width among the subdomains is measured by a parameter δ . For each $j \in [N]$, we define the local finite element space $S_h(\Omega_j)$ as follows:

$$S_h(\Omega_j) = \{v \in C(\Omega_j) : v|_T \in \mathbb{P}_1(T) \text{ for all } T \in \mathcal{T}_h|_{\Omega_j}, v = 0 \text{ on } \partial\Omega_j \setminus \partial\Omega\}.$$

The operator $I_j: S_h(\Omega_j) \rightarrow S_h(\Omega)$ is then defined as the extension-by-zero operator.

In what follows, the notation $A \lesssim B$ means that there exists a constant $c > 0$ independent of h, H , and δ , such that $A \leq cB$.

6.1. A semicoercive problem. We consider the following s -Laplacian equation with the homogeneous Neumann boundary condition:

$$\begin{aligned} -\nabla \cdot (|\nabla u|^{s-2} \nabla u) &= f \quad \text{in } \Omega, \\ \frac{\partial u}{\partial \nu} &= 0 \quad \text{on } \partial\Omega, \end{aligned}$$

where $s > 1$, $f \in (W^{1,s}(\Omega))^*$, and ν is the unit outer normal to $\partial\Omega$. For this problem to be solvable, it is required that f satisfies the compatibility condition $\langle f, 1 \rangle = 0$. This equation is well-known to have the following variational formulation:

$$\min_{v \in W^{1,s}(\Omega)} \left\{ \frac{1}{s} \int_{\Omega} |\nabla v|^s dx - \langle f, v \rangle \right\}.$$

To solve this variational problem numerically, we consider the following finite element discretization defined on $S_h(\Omega)$:

$$(6.1) \quad \min_{v \in S_h(\Omega)} \left\{ \frac{1}{s} \int_{\Omega} |\nabla v|^s dx - \langle f, v \rangle \right\}.$$

Error estimates for (6.1) can be found in, e.g., [8]. We observe that (6.1) is a specific instance of (3.1). Namely, we obtain (6.1) by setting

$$V = S_h(\Omega), \quad F(v) = \frac{1}{s} \int_{\Omega} |\nabla v|^s dx - \langle f, v \rangle.$$

in (3.1). It is straightforward to verify that F is semicoercive with respect to the $W^{1,s}(\Omega)$ -seminorm, whose kernel is $\text{span}\{1\}$.

In the following, we analyze a two-level additive Schwarz method for solving the Neumann boundary value problem (6.1). We note that the case of the Dirichlet boundary condition was considered in several existing works, e.g., [30, 48]. We define the subspaces $\{V_j\}_{j=0}^N$ of $V = S_h(\Omega)$ as follows:

$$(6.2) \quad V_0 = I_0 S_H(\Omega), \quad V_j = I_j S_h(\Omega_j), \quad j \in [N],$$

so that we have the two-level space decomposition

$$V = V_0 + \sum_{j=1}^N V_j.$$

If we employ this two-level space decomposition and the exact local problems (3.6) in the parallel subspace correction method presented in Algorithm 3.1, then we obtain the two-level additive Schwarz method.

Thanks to Theorem 4.13, it suffices to verify Assumptions 4.8 and 4.12 and estimate the constants τ_0 , μ_{K_0} , and C_{K_0} given in (3.5), (4.14a), and (4.13), respectively, to estimate the convergence rate of Algorithm 3.1. The strengthened convexity parameter τ_0 has a lower bound $\tau_0 \geq \frac{1}{5}$, due to a usual coloring argument [39, Section 5.1]. Proceeding as in [39, Section 6.1], we can verify that Assumptions 4.8 and 4.12 hold with $p = \max\{s, 2\}$, $q = \min\{s, 2\}$, and $\mu_{K_0} \gtrsim 1$ (cf. [39, Equations (6.6) and (6.7)]). Finally, using a $W^{1,s}$ -stability estimate for L^2 -projection and the Poincaré inequality (see the discussion for [48, Lemma 4.1] for details), we deduce that the stable

decomposition parameter C_{K_0} satisfies

$$C_{K_0} \lesssim 1 + \left(\frac{H}{\delta} \right)^{\frac{q(s-1)}{s}}.$$

In conclusion, by [Theorem 4.13](#), the two-level additive Schwarz method for solving [\(6.1\)](#) enjoys the following estimate:

$$F(u^{(n)}) - F(u) \lesssim \frac{1 + (H/\delta)^{\frac{pq(s-1)}{(p-q)s}}}{n^{\frac{p(q-1)}{p-q}}}, \quad n \geq 0.$$

6.2. A nearly semicoercive problem. As a next example, we consider an s -Laplacian equation with a perturbation term, given by

$$\begin{aligned} -\nabla \cdot (|\nabla u|^{s-2} \nabla u) + \epsilon u &= f && \text{in } \Omega, \\ \frac{\partial u}{\partial \nu} &= 0 && \text{on } \partial\Omega, \end{aligned}$$

where $s > 1$, $f \in (W^{1,s}(\Omega))^*$, and $\epsilon > 0$. The above equation admits the following variational formulation:

$$\min_{v \in W^{1,s}(\Omega)} \left\{ \frac{1}{s} \int_{\Omega} |\nabla v|^s dx + \frac{\epsilon}{2} \int_{\Omega} v^2 dx - \langle f, v \rangle \right\}.$$

The finite element discretization of this variational formulation defined on $S_h(\Omega)$ is given by

$$(6.3) \quad \min_{v \in S_h(\Omega)} \left\{ \frac{1}{s} \int_{\Omega} |\nabla v|^s dx + \frac{\epsilon}{2} \int_{\Omega} v^2 dx - \langle f, v \rangle \right\}.$$

We observe that, if we set

$$V = S_h(\Omega), \quad F_0(v) = \frac{1}{s} \int_{\Omega} |\nabla v|^s dx - \langle f, v \rangle, \quad F_1(v) = \frac{1}{2} \int_{\Omega} v^2 dx$$

in the abstract nearly semicoercive problem [\(5.3\)](#), then we obtain [\(6.3\)](#).

In the following, we prove that the convergence rate of the two-level additive Schwarz method, specifically [Algorithm 3.1](#) equipped with the two-level space decomposition [\(6.2\)](#) and the exact local problems, for solving [\(6.3\)](#), achieves an ϵ -independent estimate when the perturbation parameter ϵ is sufficiently small. In the framework introduced in [section 5](#), we set the seminorm $|\cdot|$ and norm $\|\cdot\|$ as $|\cdot| = |\cdot|_{W^{1,s}(\Omega)}$ and $\|\cdot\| = \|\cdot\|_{L^2(\Omega)}$, respectively. Using the same argument as in the previous example, we verify that [Assumptions 5.4](#) and [5.8](#) hold with $p = \max\{s, 2\}$ and $q = \min\{s, 2\}$. It is clear that [Assumption 5.6](#) holds with $C_{K,\text{tri}} = 2$ (cf. [Example C.1](#)). Moreover, [Assumption 5.5](#) is satisfied since $\text{span}\{1\} \subset V_0$. Therefore, by [Theorems 5.7](#) and [5.9](#), we conclude that the convergence rate of the two-level additive Schwarz method for solving [\(6.3\)](#) achieves an ϵ -independent estimate.

7. Concluding remarks. In this paper, we presented a convergence analysis of subspace correction methods for semicoercive and nearly semicoercive convex optimization problems, generalizing the theory of singular [\[34, 50\]](#) and nearly singular [\[33, 51\]](#) linear problems. The central message is that the elegant theoretical results

developed for linear problems can be directly extended to convex optimization problems. Given the wide applicability of the linear theory to various PDEs [25, 31, 53], we anticipate that the convex theory introduced in this paper will similarly have broad applications to nonlinear PDEs.

We conclude this paper by discussing potential avenues for future work related to the abstract theory. While this paper focused on smooth convex optimization problems, a natural extension would be to explore constrained or nonsmooth convex optimization problems [6, 7, 15, 39]. However, we anticipate that this direction may not yield favorable results. Specifically, in [17, 29], it was shown that domain decomposition methods for dual total variation minimization, a semicoercive convex problem with pointwise constraints, only achieve sublinear convergence rates, despite its semicoercive structure.

On the other hand, a recent work [44] demonstrated that domain decomposition methods for certain semilinear elliptic problems achieve convergence rates that are independent of the nonlinearity of the problems. We expect that there may be a connection between the result in [44] and the nearly semicoercive theory presented in this paper, though further investigation is needed.

Appendix A. Inexact local problems. Subspace correction methods often incorporate inexact local problems. Namely, each local energy functional F_j , $j \in [N]$, in Algorithms 3.1 and 3.2 may not be defined exactly as in (3.6), but rather as an approximation. From this perspective, the convergence analyses of subspace correction methods for convex optimization presented in [39, 41, 45] were conducted allowing inexact local problems. In this appendix, we demonstrate how the convergence analysis in this paper can be extended to accommodate inexact local problems.

A.1. Semicoercive problems. We first consider subspace correction methods for semicoercive problems discussed in section 4. That is, in the model problem (3.1), we assume that F is semicoercive with respect to a seminorm $|\cdot|$ in the sense of Proposition 2.3, and denote $\mathcal{N} = \ker F = \ker |\cdot|$. We assume that each local energy functional F_j , $j \in [N]$, is not necessarily given by the exact one (3.6), but rather by any functional satisfying the smoothness condition stated in Assumption 4.8, where d_j is still defined as in (4.1). Additionally, we require the further assumptions on F_j summarized in Assumption A.1 (cf. [45, Assumption 1]).

Assumption A.1 (local problems). For any $j \in [N]$ and $v \in V$, the local energy functional $F_j(\cdot; v): V_j \rightarrow \mathbb{R}$ satisfies the following:

- (a) (convexity) The functional $F_j(\cdot; v): V_j \rightarrow \mathbb{R}$ is Gâteaux differentiable, semicoercive, and convex.
- (b) (consistency) We have

$$F_j(0; v) = F(v),$$

and

$$\langle F'_j(0; v), w_j \rangle = \langle F'(v), w_j \rangle \quad \forall w_j \in V_j.$$

- (c) (stability) For some $\omega \in (0, 1] \cup (1, \rho)$, we have

$$(A.1) \quad d_F(w_j; v) \leq \omega d_j(w_j; v) \quad \forall w_j \in V_j,$$

where the constant ρ is defined as

$$(A.2) \quad \rho = \min_{j \in [N]} \inf_{d_j(w_j; v) \neq 0} \frac{\langle d'_j(w_j; v), w_j \rangle}{d_j(w_j; v)}.$$

If the local energy functional F_j is given by (3.6), then it clearly satisfies [Assumption A.1](#). One can verify without difficulty that [Assumption A.1\(a, b\)](#) implies that the constant ρ defined in (A.2) satisfies $\rho \geq 1$. For completeness, we provide an example below, as also given in [45, Example 2].

Example A.2. Suppose that the local energy functional F_j is given by

$$F_j(w_j; v) = F(v) + \langle F'(v), w_j \rangle + \frac{M}{s} \|w_j\|^s, \quad v \in V, \quad w_j \in V_j,$$

for some $s > 1$ and $M > 0$. It is clear that [Assumption A.1\(a, b\)](#) holds. Moreover, for any $v \in V$ and $w_j \in V_j \setminus \{0\}$, we have (see [55])

$$\frac{\langle d'_j(w_j; v), w_j \rangle}{d_j(w_j; v)} = s.$$

This implies $\rho = s$.

The additional assumptions for local problems presented in [Assumption A.1](#) are motivated by their role in ensuring a sufficient decrease property for the local problems. More precisely, these assumptions guarantee that solving a local problem satisfying leads to a reduction in the energy F ; see [Lemma A.3](#).

LEMMA A.3. *Let V be a reflexive Banach space, and let $F: V \rightarrow \mathbb{R}$ be a Gâteaux differentiable, convex, and semicoercive convex functional with the kernel \mathcal{N} . For $j \in [N]$ and $v \in V$, let*

$$\hat{w}_j \in \arg \min_{w_j \in V_j} F_j(w_j; v).$$

Under [Assumption A.1](#), we have

$$F(v) - F(v + \hat{w}_j) \geq \left(1 - \frac{\omega}{\rho}\right) \langle d'_j(\hat{w}_j; v), \hat{w}_j \rangle \geq 0.$$

Proof. By [Assumption A.1\(b\)](#), we have

$$\hat{w}_j \in \arg \min_{w_j \in V_j} \{\langle F'(v), w_j \rangle + d_j(w_j; v)\},$$

in which the optimality condition reads as

$$(A.3) \quad \langle F'(v), w_j \rangle + \langle d'_j(\hat{w}_j; v), w_j \rangle = 0 \quad \forall w_j \in V_j.$$

It follows that

$$\begin{aligned} F(v + \hat{w}_j) &\stackrel{(A.1)}{\leq} F(v) + \langle F'(v), \hat{w}_j \rangle + \omega d_j(\hat{w}_j; v) \\ &\stackrel{(A.2)}{\leq} F(v) + \langle F'(v), \hat{w}_j \rangle + \frac{\omega}{\rho} \langle d'_j(\hat{w}_j; v), \hat{w}_j \rangle \\ &\stackrel{(A.3)}{=} F(v) - \left(1 - \frac{\omega}{\rho}\right) \langle d'_j(\hat{w}_j; v), \hat{w}_j \rangle. \end{aligned}$$

Finally, since $\rho \geq 1$ and $\omega \leq \rho$, we have

$$\left(1 - \frac{\omega}{\rho}\right) \langle d'_j(\hat{w}_j; v), \hat{w}_j \rangle \geq 0,$$

which completes the proof. \square

Using [Lemma A.3](#), we are able to derive a descent property of the parallel subspace correction method with inexact local problems, as summarized in [Lemma A.4](#). This result generalizes [Lemma 4.3](#).

LEMMA A.4. *Let V be a reflexive Banach space, and let $F: V \rightarrow \mathbb{R}$ be a Gâteaux differentiable, convex, and semicoercive convex functional with the kernel \mathcal{N} . In [Algorithm 3.1](#), suppose that [Assumption A.1](#) holds. Then we have*

$$F(u^{(n+1)}) \leq F(u^{(n)}) + \tau\theta \min_{w \in V} \left\{ \langle F'(u^{(n)}), w \rangle + \inf_{\phi \in \mathcal{N}} \inf_{\sum_{j=1}^N w_j = w + \phi} \sum_{j=1}^N d_j(w_j; u^{(n)}) \right\}, \quad n \geq 0,$$

where the constant θ is given by

$$\theta = \begin{cases} 1, & \text{if } \omega \in [0, 1], \\ \frac{\rho - \omega}{\rho - 1}, & \text{if } \omega \in (1, \rho). \end{cases}$$

Proof. Take any $n \geq 0$. By [\(4.7\)](#), it suffices to estimate $\sum_{j=1}^N F(u^{(n)} + w_j^{(n+1)})$. As the case $\omega \in (0, 1]$ can be proven with the same argument as in [Lemma 4.3](#), we focus only on the case $\omega \in (1, \rho)$. It follows that

$$\begin{aligned} & \text{(A.4)} \\ & \sum_{j=1}^N F(u^{(n)} + w_j^{(n+1)}) \stackrel{\text{(A.1)}}{\leq} NF(u^{(n)}) + \sum_{j=1}^N \left(\langle F'(u^{(n)}), w_j^{(n+1)} \rangle + \omega d_j(w_j^{(n+1)}; u^{(n)}) \right) \\ & = NF(u^{(n)}) + \omega \min_{w \in V} \left\{ \langle F'(u^{(n)}), w \rangle + \inf_{\phi \in \mathcal{N}} \inf_{\sum_{j=1}^N w_j = w + \phi} \sum_{j=1}^N d_j(w_j; u^{(n)}) \right\} \\ & \quad - (\omega - 1) \sum_{j=1}^N \langle F'(u^{(n)}), w_j^{(n+1)} \rangle. \end{aligned}$$

Note that [Lemma A.3](#) implies

$$\begin{aligned} & \text{(A.5)} \quad -\langle F'(u^{(n)}), w_j^{(n+1)} \rangle \stackrel{\text{(A.3)}}{=} \langle d'_j(w_j^{(n+1)}; u^{(n)}), w_j^{(n+1)} \rangle \\ & \leq \frac{\rho}{\rho - \omega} \left(F(u^{(n)}) - F(u^{(n)} + w_j^{(n+1)}) \right). \end{aligned}$$

By [\(A.4\)](#) and [\(A.5\)](#), we get

$$\begin{aligned} & \text{(A.6)} \quad \sum_{j=1}^N F(u^{(n)} + w_j^{(n+1)}) \\ & \leq NF(u^{(n)}) + \frac{\rho - \omega}{\rho - 1} \min_{w \in V} \left\{ \langle F'(u^{(n)}), w \rangle + \inf_{\phi \in \mathcal{N}} \inf_{\sum_{j=1}^N w_j = w + \phi} \sum_{j=1}^N d_j(w_j; u^{(n)}) \right\}. \end{aligned}$$

Finally, combining [\(4.7\)](#) and [\(A.6\)](#) completes the proof. \square

The only distinction in the descent property for the case of inexact local problems, as described in [Lemma A.4](#), compared to the case of exact local problems in [Lemma 4.3](#), is that τ in [Lemma 4.3](#) is replaced with $\tau\theta$ in [Lemma A.4](#). Consequently, by following the same arguments as those in [section 4](#), we can establish convergence theorems analogous to [Theorems 4.11](#) and [4.13](#) for inexact local problems. As the statements of these results are identical to those of [Theorems 4.11](#) and [4.13](#), except for replacing τ with $\tau\theta$, we omit them here for brevity.

A.2. Nearly semicoercive problems. Next, we consider subspace correction methods for the nearly semicoercive problems discussed in [section 5](#). As outlined in [section 5](#), the analysis for nearly semicoercive problems relies on the assumptions stated in [Assumptions 5.4](#) to [5.6](#) and [5.8](#). Among these, the assumptions specifically related to the local problems are [Assumptions 5.4](#) and [5.6](#). Since the analysis of nearly semicoercive problems builds upon the coercive theory (using the particular norm $|\cdot|_{\epsilon,q}$), we can derive analogous results to [Theorems 5.7](#) and [5.9](#), provided that additional assumptions on the local problems—such as convexity, consistency, and stability, as described in [Assumption A.1](#)—are satisfied. For the sake of brevity, we omit the detailed derivations.

Appendix B. Proofs of the convergence theorems. In this appendix, we provide proofs of the convergence theorems of the parallel subspace correction method for semicoercive problems discussed in this paper, namely, [Theorems 4.11](#) and [4.13](#). The proofs presented in this section use similar arguments as in [\[39, 41\]](#).

We begin by presenting several elementary lemmas. We note that [Lemma B.1](#) also appeared in [\[41, Lemma 3.8\]](#).

LEMMA B.1. *Let $a, b > 0$, $q > 1$, and $T > 0$. The minimum of the function $g(t) = \frac{a}{q}t^q - bt$, $t \in [0, T]$, is given as follows:*

$$\min_{t \in [0, T]} g(t) = \begin{cases} \frac{a}{q}T^q - bT < -bT \left(1 - \frac{1}{q}\right) & \text{if } aT^{q-1} - b < 0, \\ -b \left(1 - \frac{1}{q}\right) \left(\frac{b}{a}\right)^{\frac{1}{q-1}} & \text{if } aT^{q-1} - b \geq 0. \end{cases}$$

The following lemma, also introduced in [\[26, Lemma 1.1\]](#), can be proven easily by invoking [\[41, Lemma 3.7\]](#).

LEMMA B.2. *Let $\{a_n\}$ be a sequence of positive real numbers that satisfies*

$$a_n - a_{n+1} \geq Ca_n^\gamma, \quad n \geq 0,$$

for some $C > 0$ and $\gamma > 1$. Then with $\beta = \frac{1}{\gamma-1}$, we have

$$a_n \leq \left(\frac{\beta}{Cn + \beta a_0^{-1/\beta}} \right)^\beta, \quad n \geq 0.$$

Thanks to [Lemma 4.3](#) (see [Lemma A.4](#) for the case of inexact local problems), for any $n \geq 0$, it suffices to estimate

$$\min_{w \in V} \left\{ \langle F'(u^{(n)}), w \rangle + \inf_{\phi \in \mathcal{N}} \inf_{\sum_{j=1}^N w_j = w + \phi} \sum_{j=1}^N d_j(w_j; u^{(n)}) \right\}.$$

It follows that

$$\begin{aligned}
& \min_{w \in V} \left\{ \langle F'(u^{(n)}), w \rangle + \inf_{\phi \in \mathcal{N}} \inf_{\sum_{j=1}^N w_j = w + \phi} \sum_{j=1}^N d_j(w_j; u^{(n)}) \right\} \\
& \leq \min_{u^{(n)} + w \in K_0} \left\{ \langle F'(u^{(n)}), w \rangle + \inf_{\phi \in \mathcal{N}} \inf_{\sum_{j=1}^N w_j = w + \phi} \sum_{j=1}^N d_j(w_j; u^{(n)}) \right\} \\
\text{(B.1)} \quad & \stackrel{\text{(i)}}{\leq} \min_{u^{(n)} + w \in K_0} \left\{ \langle F'(u^{(n)}), w \rangle + \frac{C_{K_0}}{q} |w|^q \right\} \\
& \stackrel{\text{(ii)}}{\leq} \min_{\alpha \in [0,1]} \left\{ \alpha \langle F'(u^{(n)}), u - u^{(n)} \rangle + \frac{\alpha^q C_{K_0}}{q} |u - u^{(n)}|^q \right\} \\
& \stackrel{\text{(iii)}}{\leq} \min_{\alpha \in [0,1]} \left\{ -\alpha \zeta_n + \frac{\alpha^q C_{K_0}}{q} |u - u^{(n)}|^q \right\},
\end{aligned}$$

where (i) follows from [Lemma 4.10](#), (ii) uses the substitution $w = \alpha(u - u^{(n)})$ for $\alpha \in [0, 1]$, (iii) is due to the convexity of F , and $\zeta_n = F(u^{(n)}) - F(u)$. Both [Theorems 4.11](#) and [4.13](#) can be proven by using (B.1), as presented in the remainder of this appendix.

B.1. Proof of [Theorem 4.11](#). We proceed to estimate the last line of (B.1) as follows:

$$\begin{aligned}
\text{(B.2)} \quad & \min_{\alpha \in [0,1]} \left\{ -\alpha \zeta_n + \frac{\alpha^q C_{K_0}}{q} |u - u^{(n)}|^q \right\} \stackrel{\text{(4.14b)}}{\leq} \min_{\alpha \in [0,1]} \left\{ -\alpha \zeta_n + \frac{\alpha^q C_{K_0} R_0^q}{q} \right\} \\
& \leq \begin{cases} -\left(1 - \frac{1}{q}\right) \zeta_n & \text{if } \zeta_n > C_{K_0} R_0^q, \\ -\left(1 - \frac{1}{q}\right) \frac{\zeta_n^{\frac{q}{q-1}}}{(C_{K_0} R_0^q)^{\frac{1}{q-1}}} & \text{if } \zeta_n \leq C_{K_0} R_0^q, \end{cases}
\end{aligned}$$

where the last inequality is due to [Lemma B.1](#). Combining [Lemma 4.3](#), (B.1), and (B.2), we obtain

$$\zeta_{n+1} \leq \begin{cases} \left(1 - \tau \left(1 - \frac{1}{q}\right)\right) \zeta_n & \text{if } \zeta_n > C_{K_0} R_0^q, \\ \zeta_n - \tau \left(1 - \frac{1}{q}\right) \frac{\zeta_n^{\frac{q}{q-1}}}{(C_{K_0} R_0^q)^{\frac{1}{q-1}}} & \text{if } \zeta_n \leq C_{K_0} R_0^q. \end{cases}$$

Note that, by [Corollary 4.4](#), the condition $\zeta_0 \leq C_{K_0} R_0^q$ ensures $\zeta_n \leq C_{K_0} R_0^q$. Finally, invoking [Lemma B.2](#) completes the proof of [Theorem 4.11](#).

B.2. Proof of [Theorem 4.13](#). In the case of [Theorem 4.13](#), an alternative upper bound for the last line of (B.1) can be derived by invoking [Assumption 4.12](#):

$$\text{(B.3)} \quad \min_{\alpha \in [0,1]} \left\{ -\alpha \zeta_n + \frac{\alpha^q C_{K_0}}{q} |u - u^{(n)}|^q \right\} \leq \min_{\alpha \in [0,1]} \left\{ -\alpha \zeta_n + \frac{\alpha^q p^{\frac{q}{p}} C_{K_0}}{q \mu_{K_0}^{\frac{q}{p}}} \zeta_n^{\frac{q}{p}} \right\}.$$

We first consider the case $p = q$. It follows by [Lemma B.1](#) that

$$(B.4) \quad \min_{\alpha \in [0,1]} \left\{ -\alpha \zeta_n + \frac{\alpha^q p^{\frac{q}{p}} C_{K_0}}{q \mu_{K_0}^{\frac{q}{p}}} \zeta_n^{\frac{q}{p}} \right\} = \min_{\alpha \in [0,1]} \left\{ -\alpha \zeta_n + \frac{\alpha^q q C_{K_0}}{q \mu_{K_0}} \zeta_n \right\} \\ \leq \zeta_n \left(1 - \frac{1}{q} \right) \min \left\{ 1, \frac{\mu_{K_0}}{q C_{K_0}} \right\}^{\frac{1}{q-1}}.$$

By combining [Lemma 4.3](#), [\(B.1\)](#), [\(B.3\)](#), and [\(B.4\)](#), we obtain the desired result.

Next, we consider the case $p > q$. By [Lemma B.1](#), we have

$$(B.5) \quad \min_{\alpha \in [0,1]} \left\{ -\alpha \zeta_n + \frac{\alpha^q p^{\frac{q}{p}} C_{K_0}}{q \mu_{K_0}^{\frac{q}{p}}} \zeta_n^{\frac{q}{p}} \right\} \\ \leq \begin{cases} -\left(1 - \frac{1}{q}\right) \zeta_n & \text{if } \zeta_n > \left(\frac{p}{\mu_{K_0}}\right)^{\frac{q}{p-q}} C_{K_0}^{\frac{p}{p-q}}, \\ -\left(1 - \frac{1}{q}\right) \left(\frac{\mu_{K_0}}{p}\right)^{\frac{q}{p(q-1)}} \frac{\zeta_n^{\frac{q(p-1)}{p(q-1)}}}{C_{K_0}^{\frac{q-1}{q}}} & \text{if } \zeta_n \leq \left(\frac{p}{\mu_{K_0}}\right)^{\frac{q}{p-q}} C_{K_0}^{\frac{p}{p-q}}, \end{cases}$$

Combining [Lemma 4.3](#), [\(B.1\)](#), [\(B.3\)](#), and [\(B.5\)](#), we get

$$\zeta_{n+1} \leq \begin{cases} \left(1 - \tau \left(1 - \frac{1}{q}\right)\right) \zeta_n & \text{if } \zeta_n > \left(\frac{p}{\mu_{K_0}}\right)^{\frac{q}{p-q}} C_{K_0}^{\frac{p}{p-q}}, \\ \zeta_n - \tau \left(1 - \frac{1}{q}\right) \left(\frac{\mu_{K_0}}{p}\right)^{\frac{q}{p(q-1)}} \frac{\zeta_n^{\frac{q(p-1)}{p(q-1)}}}{C_{K_0}^{\frac{q-1}{q}}} & \text{if } \zeta_n \leq \left(\frac{p}{\mu_{K_0}}\right)^{\frac{q}{p-q}} C_{K_0}^{\frac{p}{p-q}}. \end{cases}$$

Invoking [Lemma B.2](#) completes the proof of [Theorem 4.13](#).

Appendix C. Triangle inequality-like properties of convex functionals.

As presented in [Assumption 5.6](#), the convergence analysis of nearly semicoercive problems introduced in this paper requires an assumption that each local energy functional satisfies a certain triangle inequality-like property. To describe this triangle inequality-like property in detail, let $F: V \rightarrow \mathbb{R}$ be a Gâteaux differentiable and convex functional defined on a Banach space V . The property states that, for any bounded and convex subset K of V , there exists a positive constant $C_{K,\text{tri}}$ such that

$$(C.1) \quad d_F(w_1 + w_2; v) \leq C_{K,\text{tri}} (d_F(w_1; v) + d_F(w_2; v)), \quad v \in K, w_1, w_2 \in V,$$

where d_F denotes the Bregman distance associated with F given in [\(4.10\)](#). In this appendix, we provide several examples of convex functionals that satisfy the triangle inequality-like property [\(C.1\)](#).

Example C.1 (quadratic functionals on Hilbert spaces). Let H be a Hilbert space equipped with an inner product (\cdot, \cdot) . We consider the quadratic functional previously given in [\(2.6\)](#):

$$F(v) = \frac{1}{2}(Av, v) - (f, v), \quad v \in H,$$

where $A: H \rightarrow H$ is a continuous, symmetric and positive semidefinite linear operator, and $f \in H$. The Bregman distance d_F is given by

$$d_F(w; v) = \frac{1}{2}(Aw, w), \quad v, w \in H.$$

Thanks to the Cauchy–Schwarz inequality, we can readily deduce that (C.1) holds with $C_{K,\text{tri}} = 2$.

Example C.2 (smooth and strongly convex functionals). Let F be an L -smooth and μ -strongly convex functional defined on a Banach space V , i.e.,

$$\frac{\mu}{2}\|w\|^2 \leq d_F(w; v) \leq \frac{L}{2}\|w\|^2, \quad v, w \in V.$$

for some $L, \mu > 0$. Examples of such smooth and strongly convex functionals can be found in the literature, e.g., [16, 19, 47]. For any $v, w_1, w_2 \in V$, it follows that

$$d_F(w_1 + w_2; v) \leq \frac{L}{2}\|w_1 + w_2\|^2 \leq L(\|w_1\|^2 + \|w_2\|^2) \leq \frac{2L}{\mu}(d_F(w_1; v) + d_F(w_2; v)),$$

which proves (C.1).

Example C.3 (s -Laplacian energy). Let Ω be a bounded polyhedral domain in \mathbb{R}^d . We consider the convex functional F , defined on $V = W^{1,s}(\Omega)$ with $s > 1$, which arises in variational formulations of s -Laplacian problems, as discussed in section 6:

$$F(v) = \frac{1}{s} \int_{\Omega} |\nabla v|^s dx, \quad v \in V.$$

In [30, Lemma 3.3], it was proven that the Bregman distance $d_F(w; v)$ is equivalent, up to a multiplicative constant, to the squared quasi-norm $\|w\|_{(\nabla v)}^2$ introduced [8, 37], which is given by

$$\|w\|_{(\nabla v)}^2 = \int_{\Omega} (|\nabla w| + |\nabla v|)^{s-2} |\nabla w|^2 dx, \quad v, w \in V.$$

By leveraging this equivalence relation and the triangle-like property of the quasi-norm established in [37, Lemma 5.4], we can deduce that the functional F satisfies the triangle-like property (C.1).

REFERENCES

- [1] R. A. ADAMS, *Sobolev Spaces*, Academic Press [Harcourt Brace Jovanovich, Publishers], New York-London, 1975.
- [2] S. ADLY, E. ERNST, AND M. THÉRA, *A characterization of convex and semicoercive functionals*, J. Convex Anal., 8 (2001), pp. 127–148.
- [3] Y. I. ALBER, *Decomposition theorems in Banach spaces*, in Operator theory and its applications (Winnipeg, MB, 1998), vol. 25 of Fields Inst. Commun., Amer. Math. Soc., Providence, RI, 2000, pp. 77–93.
- [4] Y. I. ALBER, *James orthogonality and orthogonal decompositions of Banach spaces*, J. Math. Anal. Appl., 312 (2005), pp. 330–342.
- [5] D. N. ARNOLD, R. S. FALK, AND R. WINTHER, *Multigrid in $H(\text{div})$ and $H(\text{curl})$* , Numer. Math., 85 (2000), pp. 197–217.
- [6] L. BADEA AND R. KRAUSE, *One-and two-level Schwarz methods for variational inequalities of the second kind and their application to frictional contact*, Numer. Math., 120 (2012), pp. 573–599.
- [7] L. BADEA, X.-C. TAI, AND J. WANG, *Convergence rate analysis of a multiplicative Schwarz method for variational inequalities*, SIAM J. Numer. Anal., 41 (2003), pp. 1052–1073.
- [8] J. W. BARRETT AND W. B. LIU, *Finite element approximation of the p -Laplacian*, Math. Comp., 61 (1993), pp. 523–537.
- [9] P. BOCHEV AND R. B. LEHOUCQ, *On the finite element solution of the pure Neumann problem*, SIAM Rev., 47 (2005), pp. 50–66.

- [10] J. H. BRAMBLE AND S. R. HILBERT, *Estimation of linear functionals on Sobolev spaces with application to Fourier transforms and spline interpolation*, SIAM J. Numer. Anal., 7 (1970), pp. 112–124.
- [11] J. H. BRAMBLE, J. E. PASCIAK, J. P. WANG, AND J. XU, *Convergence estimates for product iterative methods with applications to domain decomposition*, Math. Comp., 57 (1991), pp. 1–21.
- [12] S. C. BRENNER, *An additive analysis of multiplicative Schwarz methods*, Numer. Math., 123 (2013), pp. 1–19.
- [13] H. BREZIS, *Functional Analysis, Sobolev Spaces and Partial Differential Equations*, Springer, New York, 2011.
- [14] Z.-H. CAO, *On the convergence of general stationary linear iterative methods for singular linear systems*, SIAM J. Matrix Anal. Appl., 29 (2008), pp. 1382–1388.
- [15] C. CARSTENSEN, *Domain decomposition for a non-smooth convex minimization problem and its application to plasticity*, Numer. Linear Algebra Appl., 4 (1997), pp. 177–190.
- [16] A. CHAMBOLLE AND T. POCK, *An introduction to continuous optimization for imaging*, Acta Numer., 25 (2016), pp. 161–319.
- [17] H. CHANG, X.-C. TAI, L.-L. WANG, AND D. YANG, *Convergence rate of overlapping domain decomposition methods for the Rudin–Osher–Fatemi model based on a dual formulation*, SIAM J. Imaging Sci., 8 (2015), pp. 564–591.
- [18] L. CHEN, *Deriving the X–Z identity from auxiliary space method*, in Domain Decomposition Methods in Science and Engineering XIX, vol. 78 of Lect. Notes Comput. Sci. Eng., Springer, Heidelberg, 2011, pp. 309–316.
- [19] L. CHEN, X. HU, AND S. WISE, *Convergence analysis of the fast subspace descent method for convex optimization problems*, Math. Comp., 89 (2020), pp. 2249–2282.
- [20] I. CIORANESCU, *Geometry of Banach spaces, duality mappings and nonlinear problems*, vol. 62, Kluwer Academic Publishers Group, Dordrecht, 1990.
- [21] G. G. DE DIEGO, P. E. FARRELL, AND I. J. HEWITT, *On the finite element approximation of a semicoercive Stokes variational inequality arising in glaciology*, SIAM J. Numer. Anal., 61 (2023), pp. 1–25.
- [22] S. DEKEL AND D. LEVIATAN, *The Bramble–Hilbert lemma for convex domains*, SIAM J. Math. Anal., 35 (2004), pp. 1203–1212.
- [23] Z. E. DOSTÁL, D. HORÁK, AND D. STEFANICA, *A scalable FETI-DP algorithm for a semicoercive variational inequality*, Comput. Methods Appl. Mech. Engrg., 196 (2007), pp. 1369–1379.
- [24] T. DUPONT AND R. SCOTT, *Polynomial approximation of functions in Sobolev spaces*, Math. Comp., 34 (1980), pp. 441–463.
- [25] P. E. FARRELL, L. MITCHELL, AND F. WECHSUNG, *An augmented Lagrangian preconditioner for the 3D stationary incompressible Navier–Stokes equations at high Reynolds number*, SIAM J. Sci. Comput., 41 (2019), pp. A3073–A3096.
- [26] G. N. GRAPIGLIA AND Y. NESTEROV, *Regularized Newton methods for minimizing functions with Hölder continuous Hessians*, SIAM J. Optim., 27 (2017), pp. 478–506.
- [27] M. GRIEBEL AND P. OSWALD, *On the abstract theory of additive and multiplicative Schwarz algorithms*, Numer. Math., 70 (1995), pp. 163–180.
- [28] H. B. KELLER, *On the solution of singular and semidefinite linear systems by iteration*, J. Soc. Indust. Appl. Math. Ser. B Numer. Anal., 2 (1965), pp. 281–290.
- [29] C.-O. LEE AND J. PARK, *Fast nonoverlapping block Jacobi method for the dual Rudin–Osher–Fatemi model*, SIAM J. Imaging Sci., 12 (2019), pp. 2009–2034.
- [30] Y.-J. LEE AND J. PARK, *On the linear convergence of additive Schwarz methods for the p-Laplacian*, IMA J. Numer. Anal., (2024), <https://doi.org/10.1093/imanum/drae068>.
- [31] Y.-J. LEE, J. WU, AND J. CHEN, *Robust multigrid method for the planar linear elasticity problems*, Numer. Math., 113 (2009), pp. 473–496.
- [32] Y.-J. LEE, J. WU, J. XU, AND L. ZIKATANOV, *On the convergence of iterative methods for semidefinite linear systems*, SIAM J. Matrix Anal. Appl., 28 (2006), pp. 634–641.
- [33] Y.-J. LEE, J. WU, J. XU, AND L. ZIKATANOV, *Robust subspace correction methods for nearly singular systems*, Math. Models Methods Appl. Sci., 17 (2007), pp. 1937–1963.
- [34] Y.-J. LEE, J. WU, J. XU, AND L. ZIKATANOV, *A sharp convergence estimate for the method of subspace corrections for singular systems of equations*, Math. Comp., 77 (2008), pp. 831–850.
- [35] Y.-J. LEE AND J. XU, *New formulations, positivity preserving discretizations and stability analysis for non-Newtonian flow models*, Comput. Methods Appl. Mech. Engrg., 195 (2006), pp. 1180–1206.
- [36] J. LINDENSTRAUSS AND L. TZAFRIRI, *On the complemented subspaces problem*, Israel J. Math.,

- 9 (1971), pp. 263–269.
- [37] W. LIU AND N. YAN, *Quasi-norm local error estimators for p -Laplacian*, SIAM J. Numer. Anal., 39 (2001), pp. 100–127.
 - [38] Y. NESTEROV, *Gradient methods for minimizing composite functions*, Math. Program., 140 (2013), pp. 125–161.
 - [39] J. PARK, *Additive Schwarz methods for convex optimization as gradient methods*, SIAM J. Numer. Anal., 58 (2020), pp. 1495–1530.
 - [40] J. PARK, *Accelerated additive Schwarz methods for convex optimization with adaptive restart*, J. Sci. Comput., 89 (2021), p. Paper No. 58.
 - [41] J. PARK, *Additive Schwarz methods for convex optimization with backtracking*, Comput. Math. Appl., 113 (2022), pp. 332–344.
 - [42] J. PARK, *Fast gradient methods for uniformly convex and weakly smooth problems*, Adv. Comput. Math., 48 (2022), p. Paper No. 34.
 - [43] J. PARK, *Additive Schwarz methods for fourth-order variational inequalities*, J. Sci. Comput., 101 (2024), p. Paper No. 74.
 - [44] J. PARK, *Additive Schwarz methods for semilinear elliptic problems with convex energy functionals: Convergence rate independent of nonlinearity*, SIAM J. Sci. Comput., 46 (2024), pp. A1373–A1396.
 - [45] J. PARK AND J. XU, *Theory of parallel subspace correction methods for smooth convex optimization*. Submitted to the DD28 Proceedings.
 - [46] V. ROULET AND A. D’ASPROMONT, *Sharpness, restart, and acceleration*, SIAM J. Optim., 30 (2020), pp. 262–289.
 - [47] X.-C. TAI AND M. ESPEDAL, *Rate of convergence of some space decomposition methods for linear and nonlinear problems*, SIAM J. Numer. Anal., 35 (1998), pp. 1558–1570.
 - [48] X.-C. TAI AND J. XU, *Global and uniform convergence of subspace correction methods for some convex optimization problems*, Math. Comp., 71 (2002), pp. 105–124.
 - [49] A. TOSELLI AND O. WIDLUND, *Domain Decomposition Methods—Algorithms and Theory*, Springer, Berlin, 2005.
 - [50] J. WU, Y.-J. LEE, J. XU, AND L. ZIKATANOV, *Convergence analysis on iterative methods for semidefinite systems*, J. Comput. Math., 26 (2008), pp. 797–815.
 - [51] J. WU AND H. ZHENG, *Parallel subspace correction methods for nearly singular systems*, J. Comput. Appl. Math., 271 (2014), pp. 180–194.
 - [52] J. XU, *Iterative methods by space decomposition and subspace correction*, SIAM Rev., 34 (1992), pp. 581–613.
 - [53] J. XU AND K. YANG, *Well-posedness and robust preconditioners for discretized fluid–structure interaction systems*, Comput. Methods Appl. Mech. Engrg., 292 (2015), pp. 69–91.
 - [54] J. XU AND L. ZIKATANOV, *The method of alternating projections and the method of subspace corrections in Hilbert space*, J. Amer. Math. Soc., 15 (2002), pp. 573–597.
 - [55] Z.-B. XU AND G. F. ROACH, *Characteristic inequalities of uniformly convex and uniformly smooth Banach spaces*, J. Math. Anal. Appl., 157 (1991), pp. 189–210.