# How Green Can AI Be? A Study of Trends in Machine Learning Environmental Impacts

Clément Morand<sup>1</sup>, Anne-Laure Ligozat<sup>1,2</sup>, Aurélie Névéol<sup>1</sup>

<sup>1</sup>Université Paris-Saclay, CNRS, LISN,<sup>2</sup>ENSIIE name.surname@lisn.upsaclay.fr

#### Abstract

The compute requirements associated with training *Artificial Intelligence* (AI) models have increased exponentially over time. Optimisation strategies aim to reduce the energy consumption and environmental impacts associated with AI, possibly shifting impacts from the use phase to the manufacturing phase in the life-cycle of hardware. This paper investigates the evolution of individual graphics cards production impacts and of the environmental impacts associated with training Machine Learning (ML) models over time.

We collect information on graphics cards used to train ML models and released between 2013 and 2023. We assess the environmental impacts associated with the production of each card to visualize the trends on the same period. Then, using information on notable AI systems from the Epoch AI dataset we assess the environmental impacts associated with training each system.

The environmental impacts of graphics cards production have increased continuously. The energy consumption and environmental impacts associated with training models have increased exponentially, even when considering reduction strategies such as location shifting to places with less carbon intensive electricity mixes.

These results suggest that current impact reduction strategies cannot curb the growth in the environmental impacts of AI. This is consistent with rebound effect, where the efficiency increases fuel the creation of even larger models thereby cancelling the potential impact reduction. Furthermore, these results highlight the importance of considering the impacts of hardware over the entire life-cycle rather than the sole usage phase in order to avoid impact shifting.

The environmental impact of AI cannot be reduced without reducing AI activities as well as increasing efficiency.

#### 1 Introduction

Environmental impacts such as the carbon footprint, water usage, metallic resource depletion or toxicity to human and non human life caused by ML raise increasing concern. Strategies that optimize the energy consumption of models training have been developed to mitigate the environmental impacts of the sector. *Technical optimisations* adjust model architecture to offer the same task performance in smaller models. *Shift optimisations* displaces computation towards less carbon-intensive mixes performing compute when more renewable energy is available or purchasing of renewable energy. Current strategies also rely on frequent hardware updates to benefit from more energy efficient recent hardware.

Major AI companies have claimed that these strategies would mitigate and eventually reduce the carbon footprint of ML training (Patterson et al. 2022). However, they fail to account for two important factors. First, frequent hardware upgrades cause impact shifting (i.e., reducing the environmental impacts from one life cycle phase or impact category at the detriment of other life cycle phases and/or impact categories), which limits the overall benefit. Second, optimisation often accompany rebound effect (i.e., optimization can lead to a less than expected decrease, or even increase the total environmental impacts of the sector).

Herein, we consider impact shifting, rebound effects and other digital trends to question "How green can AI be?" Specifically, we investigate how the impacts of individual graphics cards have evolved in the past 10 years and how the environmental impacts associated with training ML models have evolved over this period. We focus on the past decade (2013-2023) and conduct a detailed study of the characteristics and environmental impacts of the production of NVIDIA workstation graphics cards. We link information on hardware to ML models trained using them. Finally, we question the global effect of the impact mitigating strategies.

The main contributions of this paper are as follows:

- An assessment of the environmental impact of graphic cards released between 2013 and 2023
- A detailed analysis of the environmental impact of ML models, including impact shifting and rebound effect
- Evidence that the impact of AI is increasing in spite of the promises of current reduction strategies. The material used in our analysis are shared to foster reroducibility.

#### 2 State of the art

After the high level of carbon emissions associated with training Natural Language processing models was reported (Strubell, Ganesh, and McCallum 2019), researchers stated the need for a "Green AI" (Schwartz et al. 2020), soon structured as an entire research field (Verdecchia, Sallou, and Cruz 2023). Green AI research includes the development of carbon footprint reporting tools, such as *Green Algorithms* (Lannelongue, Grealey, and Inouye 2021) or *Car-* *bonTracker* (Anthony, Kanding, and Selvan 2020). Recent reviews offer a comparison of tools focusing on measuring the impact of the use phase of computation (Bouza, Bugeau, and Lannelongue 2023; Jay et al. 2023; Bannour et al. 2021). Subsequently, *MLCA* was developed to also account for other phases of the life cycle of the hardware used and for abiotic depletion potential and primary energy demand in addition to the carbon footprint (Morand, Névéol, and Ligozat 2024).

These tools have been used for individual reporting of the carbon footprint associated with training ML models, e.g., (Luccioni, Viguier, and Ligozat 2023). Optimisation techniques have been developed and used to build less carbon intensive models (Wu et al. 2022; Patterson et al. 2022). However, we do not know of studies of the environmental impacts at the scale of the ML sector. Thompson et al. (2023) and Sevilla et al. (2022) have shown that compute requirements for ML models training over time follow an exponential growth. Desislavov, Martínez-Plumed, and Hernández-Orallo (2023) studied the energy demand for ML models inferences over time, also showing a tendency towards a growth in energy demand per inference over time. A position paper by de Vries (2023) discussed the potential growth of the ML sector with the recent surge in demand for large language models freely accessible as chatbots.

The energy efficiency of graphics cards has increased exponentially which leads some to think that frequently replacing hardware will shrink the carbon footprint of training models (Patterson et al. 2022). However, the computation requirements to train models have also increased exponentially (Sevilla et al. 2022; Thompson et al. 2023). This seems to indicate the presence of a *producer rebound* effect where the efficiency increases fuel the creation of even larger models thereby canceling the potential impact reduction (Coroama and Mattern 2019). Bol, Pirson, and Dekimpe (2021) and Gossart (2015) showed that optimisations are often absorbed by the growth of the *Information and Communication Technologies* (ICT) sector. Does this observation also hold for the case of the AI sector? Masanet et al. (2020) has shown that despite an exponential growth in computation demand between 2012 and 2018, the total energy consumption of datacenters had only increased by around 5% indicates that "A [...] source of higher electricity consumption is coming from energy-intensive data centres, artificial intelligence (AI) and cryptocurrencies, which could double by 2026."

Overall, previous work led to the development of tools for measuring the impact of specific AI algorithms. Studies at scale have addressed parts of the AI sector only such as datacenters, or have omitted rebound effect and hardware production impacts. Herein, we present a comprehensive study of the impact of ML hardware and models over a decade, including rebound effect and hardware production.

## 3 Methodology

Details on the sources, processes and methodological choices are available in the accompanying code and data.

#### 3.1 Graphics cards production impacts

To better understand the extent of the impact shifting, we study the evolution of graphics cards characteristics over time and the associated evolution in the environmental impacts of producing graphics cards over time. Life cycle assessments of ICT equipment have shown the importance of *Integrated Circuits (IC)* in the environmental impacts of ICT equipment (Clément, Jacquemotte, and Hilty 2020). ICs come in two forms in graphics cards: GPU (logic type ICs) and memory (memory type ICs). The surface of the GPU is indicated by the GPU *die area*. Contributors to the impact of producing ICs include the surface of the IC (i.e., the die area of the GPU and the surface of memory type IC for the memory chips), as well as how finely the circuits are printed on the semiconducting material. Thus, if the die area increases we can expect an increase in the environmental impacts of the device. Furthermore, (Pirson et al. 2023) have shown that, with finer technological nodes, the environmental impacts per produced  $\text{cm}^2$  of die increase. Thus, as the quantity of memory increases (probable increase in memory type IC surface), the GPU die size increases and the technological node gets finer (latest GPUs processed at 5nm), we can estimate that the environmental impacts of graphics cards production increases.

We study the evolution of the characteristics of graphics cards over time to check these hypotheses. We focus on NVIDIA workstation graphics cards as NVIDIA is the lead provider of graphics cards for servers. Additional cards frequently used to train ML models are also included in our study: Google's TPU, Huawei or Cerebras or NVIDIA non workstation cards such as the Geforce GTX 1080 Ti.

NVIDIA workstation graphics cards A dataset gathering information on 167 NVIDIA workstation graphics cards models released between 2013 and 2023 included is curated. The main information gathered for each model includes: Release date, die area, technological node, memory type, memory size, *Thermal Design Power* (TDP) and compute power (Single, Double, Tensor and Half floating precision). The dataset is based on data retrieved from the TechPowerUp GPU database<sup>1</sup>. Another dataset of 76 graphics cards models based on a wikipedia page listing NVIDIA graphics cards<sup>2</sup> was retrieved. We merged the two datasets to cross-validate the specifications of the different cards. This validated information on 74 out of the 167 models (44%) In cases of divergent information, NVIDIA's published datasheets are taken as reference. Our final dataset is available with the accompanying code.

Other graphics cards To assess the environmental impacts of ML models, we gather information on other graphics card models used for training. We relied on different sources: the TechPowerUp database provided details for NVIDIA non-workstation cards (11 card models), the Google Cloud Platform documentation and publications by

<sup>1</sup> https://www.techpowerup.com/gpu-specs/ on 12/12/2023

<sup>&</sup>lt;sup>2</sup>https://en.wikipedia.org/wiki/List\_of\_Nvidia\_graphics processing units on April 12, 2024

Google on their hardware, and website of the manufacturer/press releases/benchmarks for the other cards (Cerebras CS-2, Huawei Ascend 910 and AMD Instinct MI250X).

### 3.2 ML models training impacts

The studies on ML models training have been conducted using the Epoch AI Notable systems database (Epoch AI 2022) retrieved on July 19, 2024. This database is the most comprehensive one on ML systems to our knowledge. It gathers extensive information on a large variety of *notable* ML systems (including but not limited to: name, publication date, number of *floating point operations* (FLOP) during training, origin country of model producers, training duration, training hardware and hardware quantity). Notable systems are "Models that have advanced the state of the art, had a large influence in the field's history, or had a large impact within the world"<sup>3</sup>. To estimate the environmental impacts of training a ML model, information on the training duration, training hardware, hardware usage during training and source of electricity is needed (Morand, Névéol, and Ligozat 2024).

Ambiguous card names Multiple card names are documented for 5 ML models (one per training process in multiple steps carried on different hardware). These models are excluded from analyses as it is unclear which hardware is used for which share of the processes. For some other models the hardware name itself is ambiguous. For instance, it might be indicated as "A100", which could refer to several cards. Each of these graphics cards can differ in some characteristics (e.g., estimated environmental impact, energy consumption or compute power). In such cases, the most plausible candidate is chosen as the reference value, and the other options are used to compute a value interval.

Training duration The number of GPU-hours required for training a model can be estimated in two ways. For models specifying duration and hardware quantity, these two values can be multiplied to obtain an estimate referred to as  $GPU-h_1$ . It is the most reliable as it uses information retrieved directly from the papers presenting the models. However, Table 1 shows the low coverage of this method in the database. For models were the compute required for training (FLOP) and hardware used are specified, another option is to divide the number of FLOP by the compute power of the card used. The compute power values that are used are the maximum of the peak performance in single, half or tensor floating point precision. This second estimation, referred to as  $GPU-h<sub>2</sub>$ , should lead to an under-estimation of the number of GPU-hours, as the hardware does not always operate at peak performance. For each of the 106 models where both estimates are available, we compare them to obtain an estimate for the ratio of hardware performance and to validate the estimates from  $GPU-h<sub>2</sub>$ . this process revealed anomalies, especially on fine-tuned models where  $GPU-h<sub>2</sub>$  corresponds to training the base model and  $GPU-h_1$  correspond to the fine-tuning process. Anomalies also occur when incorrect compute power values are used to estimate the number

of FLOP in the EpochAI data-base. These anomalies should not pose problems as we use  $GPU-h_1$  for these models.

We build a linear model to predict  $GPU-h_1$  using  $GPU-h<sub>2</sub>$ . This model is computed on 87 observation excluding the anomalies. To ensure a linear relation, we estimate  $log(GPU-h_1) \sim log(GPU-h_2)$  The linear regression analysis revealed a statistically significant model  $(F(1,85) = 4161,$  $p < 2.2e-16$ , with an adjusted R<sup>2</sup> of 0.98, meaning that 98 percent of the variance in the observations is explained by our model. The model equation is  $log(GPU-h_1) = 1.31 +$  $1.00 \log(\text{GPU-h}_2)$  with a standard error of 0.16 for the intercept and 0.02 for the regression coefficient. This indicates that an increase of 1 for the  $log(GPU-h_2)$  value leads to an average increase of 1.00 units in  $log(GPU-h_1)$ . This positive relationship between  $log(GPU-h_2)$  and  $log(GPU-h_1)$ was found to be statistically significant (t(85) = 64.50, p  $< 2e - 16$ ), affirming the predictive power of log(GPU-h<sub>2</sub>) on  $log(GPU-h_1)$ . In addition to the regression analysis, a scatterplot with the fitted regression line were examined to ensure model assumptions were met. The residuals were normally distributed (Shapiro-Wilk W = .98, p = .24), homoscedasticity was confirmed (studentized Breusch-Pagan test =  $.86$ ,  $p = .35$ ), and the residuals appeared to be independent (Durbin-Watson  $D = 1.67$ ,  $p = .055$ ). This model correspond to using a constant performance ratio of  $\simeq 27\%$ The final estimation for the number of GPU hours is thus as follows: we use  $GPU-h_1$  for the 112 models where it is available and use  $GPU-h_2$  for another 93 models.

Server characteristics For NVIDIA workstations, we supposed that servers contain 4 graphics cards and 2 CPUs. For NVIDIA non-workstation cards, we supposed servers with 2 graphics cards for 2 CPUs. Without information on memory usage per model training, we did not account for the presence of memory in the servers, which largely underestimates the production impacts of the servers. For non-NVIDIA hardware, we searched documentation (e.g., Google Cloud Platform documentation, publications by Google) to obtain information on the number of chips and processors per server. For instance, for TPUv3, a server with two CPUs manages every four TPU chips.

We use values consistent with hyper-scale datacenters for the hardware lifespan, average utilization over its life-cycle and infrastructure energy consumption. We use a lifespan of 3 years for the hardware (Ostrouchov et al. 2020), and, using information from Meta, we choose a close to optimal PUE of 1.1 and a hardware utilization of 50 Using a constant PUE of 1.1 overlooks the significant reduction in PUE of datacenters used to train ML models between 2012 and 2018 (Masanet et al. 2020), leading to an under-estimation of the infrastructure consumption for models released before the transition towards hyper-scaler clouds.

Hardware consumption For the hardware usage during training, a value of 100% was considered both for GPUs and CPUs. This leads to an overestimate of the energy consumption as processors do not function at max power during the entirety of training. This overestimation should be at most twice the actual consumption value. As we do not account for the memory quantity because of lack of data, we also

<sup>3</sup> https://epochai.org/data/notable-ai-models-documentation

		<b>Systems</b>	<b>FLOP</b>	hardware	both	training duration	hardware quantity	both	+ hardware
Number		825	386	244	206	142	166	112	107
Coverage $(\%)$		100	46.8	29.6	24.9	17.2	20.1	13.6	13.0
Confidence	Confident	170	116	109	97	77	72	59	58
	Likely	96	65	54	46	36	30	20	18
	Speculative	51	36	13	13	20	10		
	Unknown	508	169	63	50	33	30	24	24

Table 1: Description of the Epoch AI dataset. The values show the number of entries for each information type. Confidence scores are reported by database authors for Training compute, Parameters, and Training dataset size.

do not account for its associated energy consumption, leading to an under-estimation of the total energy consumption. This under-estimation should not be too large as processors consumption are generally the dominating source of energy consumption when training ML models (Jay et al. 2023).

Environmental impact of the energy usage The energy mix used to evaluate the impacts of each model corresponds to that of the countries of the ML systems producers. If multiple countries participate in the creation of the system, the energy mixes of each implicated country are considered to create a value interval, and the mix of the country indicated in first position is considered as the reference value.

Impact of strategies for reducing the environmental impacts of the energy usage Different scenarios of reduction of the carbon intensity of the energy mixes over time are considered. These scenarios assess the impact of shifting compute locations and investing in decarbonizing electricity for datacenters. The two techniques of compute location shifting and 'greening' the used electricity mixes are important in the strategies of decarbonation of the tech companies (Patterson et al. 2022). Each scenario comes in the form of a continuous reduction of X 2019 is chosen as a starting year as it is the year of publication of the paper by (Strubell, Ganesh, and McCallum 2019) which first raised concerns on the carbon footprint of training ML models. Scenarios of a reduction up to 25% per year are explored. The carbon intensity of the mix used for training a model is thus multiplied by  $(1 - \text{ratio})^n$  where n stand for the number of years since 2019 at release date of the system.

### 3.3 Metrics used in this study

We assess the energy consumption, carbon footprint and metallic resource usage associated with hardware production and training ML systems. The carbon footprint is assessed according to *Global Warming Potential* (GWP)<sup>4</sup>, measured in  $kgCO<sub>2</sub>$  eq for the emissions of greenhouse gases such as  $CO<sub>2</sub>$  (Forster et al. 2023). Metallic resource usage is assessed through *Abiotic Depletion Potential* (ADPe) measured in *kilograms antimony equivalent* (kgSb eq) (van Oers, Guinée, and Heijungs 2020). Metrics are computed using the MLCA tool (Morand, Névéol, and Ligozat 2024).

#### 3.4 Software used in the analysis

Data manipulation and statistical analyses have been performed using emacs Org mode 9.1.9, Python 3.8.10 using the pandas version 2.0.3 library and R version 3.6.3 (2020- 02-29) with the ggplot2\_3.4.3, dplyr\_1.1.3, lmtest\_0.9-40, stringr $1.5.0$  and zoo $1.8-12$  libraries. Platform:  $x86.64$ -pclinux-gnu (64-bit) Running under: Ubuntu 20.04.6 LTS

### 4 Impact shifting and hardware production

Computing facilities have been getting increasingly energy efficient and can perform much more computation in a fixed duration using less energy. This efficiency is partly obtained by regular hardware updates in data centers, thus incurring additional equipment production and end-of-life environmental costs. Frequently changing hardware relies on *impact shifting*, i.e., reducing one type of environmental impact in one specific phase of the life-cycle of a product at the detriment of other categories of environmental impacts or other life cycle phases. Furthermore, graphics cards are getting increasingly more technologically advanced, leading to possible increases in their production impacts.

# 4.1 Evolution of graphics cards characteristics

Figure 1 shows the evolution of the characteristics of NVIDIA workstation graphics cards from 2013 to 2023. Cards are being produced with decreasing technological nodes. Die area has increased linearly while memory size has grown exponentially (around 30% Cummulative Annual Growth Rate). Exponential growth in the memory size does not necessarily mean a proportional increase in memory IC area as Moore's Law has allowed to exponentially miniaturize ICs.

Figure 2 shows the evolution of the energy consumption of NVIDIA workstation graphics cards from 2013 to 2023, estimated by their *Termal Design Power (TDP)*. Even if the energy consumption per operation has decreased over time, the total energy consumption of a card has slightly increased over time. This observation indicates a form of rebound effect, where the energy efficiency improvements on the cards have allowed to increase to number of operations realised on a card at a fixed energy consumption.

### 4.2 Impacts of graphics cards manufacturing

Figure 3 shows the evolution in the estimated environmental impacts of graphics cards production, in terms of (GWP)

<sup>4</sup> using a time horizon of 100 years (GWP100)



Figure 1: Evolution of the characteristics of NVIDIA workstation graphics cards from 2013 to 2023



Figure 2: Evolution of the energy consumption of NVIDIA workstation graphics cards from 2013 to 2023

and (ADP). The overall trend suggests that the production impacts have increased over time for both impact categories, as was expected from the observations on the graphics cards characteristics. Overall, all the important characteristics of graphics cards produced by NVIDIA for workstation have increased, leading to an increase in the environmental impacts incurred by the production of these cards over time.

### 4.3 Evolution of the hardware requirements

Figures 4a and 4b show the production impacts of graphics cards used to train ML systems in the Epoch AI data-set. The models of graphics cards used to train ML systems have followed a similar trend to that of the produced workstation cards, confirming that the production impacts of graphics cards used to train ML models have increased over time.

Figure 5 shows the evolution of the number of graphics cards used to train models in the Epoch AI database. Hardware quantity has increased exponentially. Combining this observation with the observation that the graphics cards used are more impacting at production indicate that the production of the hardware used to train ML models cause significant and growing environmental impacts that need to be addressed as well as the energy consumption of the hardware.

# 5 Optimization Strategies in a growing sector

### 5.1 Trends in ML models training

Energy consumption of models training Figure 6 shows that energy consumption has increased exponentially even

when low-energy demanding models are used. There are too few low energy intensive models to draw reliable conclusions. Nonetheless, data suggests an increase from 2012 to 2019, and a decrease from 2019 to 2023, bringing energy consumption to the level of 2012. The advent of green AI research from 2019 could be a contributing factor. The less energy intensive models could also be fine-tuned versions of pre-existing models that build on the larger models to attain good performance with a small overhead.

Environmental impacts of model training Figure 7 presents the estimated training impacts in terms of GWP (Figure 7a) and ADP (Figure 7b). Both environmental indicators have increased exponentially between 2012 and 2024.

Table 2 summarizes the distribution of the shares of embodied impacts, i.e., the share of the total impacts that can be attributed to hardware production. The estimations for the share of GWP are lower than expected (embodied impacts typically amount to a third of the total footprint of model training (Gupta et al. 2022; Wu et al. 2024)). This underestimation comes from the absence of accounting for the memory used to train models. ADP, however, comes close to exclusively from hardware production. Solely reducing the energy impacts from ML models training will not be sufficient to solve all of the environmental impacts of AI.

		$\vert$ Min $\vert$ Q1 $\vert$ Q2 $\vert$ Mean $\vert$ Q3 $\vert$ Max	
ADP $(\%)$   50   100   100   99.2   100   100			
GWP $(\%)$   5.8   11   15   15   16   47			

Table 2: Summary of the share of embodied impacts on the total impacts associated with training models Q1, Q2 and Q3 respectively refer to the first, second and third quartiles.

# 5.2 Is location shifting the solution?

We compare the impact of models trained with real electricity mixes with a simulated annual reduction of 25% of the carbon intensity of the electricity mixes. Models with a carbon footprint inferior to 50 kgCO<sub>2</sub> eq are excluded (16) models in real electricity mixes, vs. 21 in the simulation). Figure 8 shows that the carbon footprint of models released from 2019 is increasing, suggesting that the reduction strategy fails to curb the exponential growth of models impact.

It can be noted that the carbon intensity of electricity mixes is bound to around  $15\text{-}20gCO_2$  eq/kWh, based on the current world lower intensity mixes. Thus, even if carbon intensity reductions were quick enough to counterbalance the increase in energy consumption, the benefits would be short lived, under a steady trend in energy consumption. In addition, the production phase of hardware also has an increasing impact which is not curbed by reducing carbon intensity. Finally, the geographic changes intended to reduce carbon intensity could incentivizes higher impact for datacenter facilities (Velkova 2019).

### 6 Discussion

Rebound effect is prevalent in the AI sector Our observations on the trends in ML models training energy consump-



Figure 3: Evolution of the production impacts (left in GWP, right in ADP) of NVIDIA workstation graphics cards.



Figure 4: Evolution of the production impacts (left in GWP, right in ADP) of the graphics cards used for training ML systems in the Epoch AI data-set. Value intervals correspond to cases of ambiguous card names. The blue line represents the trend in the reference values while the red line represents the trend on the minimal values.



Figure 5: Evolution of the number of graphics cards used for training ML models in the Epoch AI dataset. Trend was computed using the *Weighted Least Square* (WLS) method to account for heteroscedasticity

tion, showing the prevalence of the rebound effect are consistent with diverse macro observations: Wu et al. (2022) identify a rebound effect at Facebook while Patterson et al. (2022) show that a constant share of Google's energy consumption is attributable to AI when the total consumption increased dramatically. The conclusions of our study should also apply to inferences. de Vries (2023) discusses rising inferences and Desislavov, Martínez-Plumed, and Hernández-Orallo (2023) suggest that inferences are also getting more



Figure 6: Evolution of the energy consumption of training ML models over time. Value intervals account for ambiguous card names. Trend has been computed using WLS to account for heteroscedasticity.

computationally and energy expensive over time.

Greener energy cannot void carbon impact. As shown in Figure 8, reduction of the carbon intensity of the electricity used seems insufficient to curb the exponential growth of the carbon footprint of training ML models. Furthermore, the electricity consumption of data-centers destabilizes local electricity grids (Ortar et al. 2023), potentially causing the prolongation of fossil fuel power plants (Chu 2024; Ambrose 2024). Matching the carbon footprint of AI



Figure 7: Environmental impacts in GWP (a) and ADP (b) of training ML models over time. Value intervals account for ambiguous card names and model producers from multiple countries. Trends computed using WLS to account for heteroscedasticity.



Figure 8: Estimated carbon footprint of training ML models released after 2019, with or without reduction of the carbon intensity of the used electricity.

through carbon offsetting also has limited potential (Greenfield 2023; Lohmann 2009; Bol, Pirson, and Dekimpe 2021). Impacts go beyond carbon footprint Our study focused on the carbon footprint and metallic resource depletion incurred by training models. Both metrics have increased over time, but the carbon footprint comes mostly from the energy consumption while metallic resource depletion comes from hardware production, showing the importance of multicriteria environmental assessments. AI also causes multiple other environmental impacts that are harder to quantify, such as water usage (Mytton 2021), that creates a competition for resources, especially in urban areas (Roussilhe et al. 2024).

The mining process and the disposal of hardware at its end of life also cause the destruction of ecosystems (Comber and Eriksson 2023) and create multiple pollutants that are toxic to human and non-human life (World Health Organization 2021). AI also has significant social consequences and poses numerous ethical challenges (Bender et al. 2021; Jiang et al. 2023; Dauvergne 2021; Keyes 2018; Shi et al. 2023).

# 7 Limitations

We acknowledge limitations in our study, which stem from the material and methods available to conduct the analysis. Nonetheless, this remains, to our knowledge, the most comprehensive study on the global impact of the AI sector. Limitations of the material and methods. The production impact assessment in MLCA does not account for technological node, and supposes a fixed memory density (while it has increase over time), leading to production impacts estimate being inexact. MLCA currently does not account for the end of life of the hardware nor for the life cycle of datacenter buildings or cooling infrastructure. The Epoch AI database is incomplete (models and model information are missing), which leads to some uncertainty on the observed trends, as the excluded models could have an impact.

Modeling hypotheses are needed for analysis. Hypotheses sometimes lead to under or over-estimation of some parameters such as GPU consumption or training duration. For example, the impact of the BLOOM model is overestimated in our study. It listed in Epoch AI as an international collaboration, so the world average electricity mix is used in our analysis although it was trained in France, which has a lower carbon intensity than world average. Lacking information, we supposed that all models were trained in hyper-scale data-centers, leading to using a quasi optimal PUE for all models, masking the increase in PUE over the last decade.

This study is limited to model training. It does not account for inferences (which are becoming increasingly prevalent over the life cycle of models), model retraining or multiple training iterations during model development.

### 8 Conclusion and Future work

In this paper, we curated a graphics cards dataset and showed that the production impacts of the hardware used for training ML models have continuously increased over time. Using the Epoch AI dataset, we showed that the energy consumption and environmental impacts associated with training models have also increased, even when considering reduction strategies such as location shifting. We demonstrate the limits of impact shifting strategies. Our results suggest that current impact reduction strategies alone cannot curb the growth in the environmental impacts of AI. Impact reduction must be combined with a broader reflection on the place and role of AI in a sustainable society.

# References

Ambrose, J. 2024. Ireland's datacentres overtake electricity use of all urban homes combined. The Guardian, https://www.theguardian.com/world/article/2024/jul/23/ ireland-datacentres-overtake-electricity-use-of-all-homescombined-figures-show.

Anthony, L. F. W.; Kanding, B.; and Selvan, R. 2020. Carbontracker: Tracking and Predicting the Carbon Footprint of Training Deep Learning Models. arXiv:2007.03051.

Bannour, N.; Ghannay, S.; Névéol, A.; and Ligozat, A.-L. 2021. Evaluating the carbon footprint of NLP methods: a survey and analysis of existing tools. In Moosavi, N. S.; Gurevych, I.; Fan, A.; Wolf, T.; Hou, Y.; Marasović, A.; and Ravi, S., eds., *Proceedings of the Second Workshop on Simple and Efficient Natural Language Processing*, 11–21. Virtual: Association for Computational Linguistics.

Bender, E. M.; Gebru, T.; McMillan-Major, A.; and Shmitchell, S. 2021. On the Dangers of Stochastic Parrots: Can Language Models Be Too Big? In *Proceedings of the 2021 ACM Conference on Fairness, Accountability, and Transparency*, FAccT '21, 610–623. New York, NY, USA: Association for Computing Machinery. ISBN 9781450383097.

Bol, D.; Pirson, T.; and Dekimpe, R. 2021. Moore's Law and ICT Innovation in the Anthropocene. In *2021 Design, Automation & Test in Europe Conference & Exhibition (DATE)*, 19–24.

Bouza, L.; Bugeau, A.; and Lannelongue, L. 2023. How to estimate carbon footprint when training deep learning models? A guide and review. *Environmental Research Communications*, 5(11): 115014.

Chu, A. 2024. US slows plans to retire coalfired plants as power demand from AI surges. Financial Times, https://www.ft.com/content/ddaac44b-e245- 4c8a-bf68-c773cc8f4e63.

Clément, L.-P. P.-V.; Jacquemotte, Q. E.; and Hilty, L. M. 2020. Sources of variation in life cycle assessments of smartphones and tablet computers. *Environmental Impact Assessment Review*, 84: 106416.

Comber, R.; and Eriksson, E. 2023. Computing as Ecocide. In *Ninth Computing within Limits 2023*. LIMITS. Https://limits.pubpub.org/pub/a8h46wqy.

Coroama, V. C.; and Mattern, F. 2019. Digital Rebound - Why Digitalization Will not Redeem us our Environmental Sins. In Wolff, A., ed., *Proceedings of the 6th International Conference on ICT for Sustainability, ICT4S 2019, Lappeenranta, Finland, June 10-14, 2019*, volume 2382 of *CEUR Workshop Proceedings*. CEUR-WS.org.

Dauvergne, P. 2021. The globalization of artificial intelligence: consequences for the politics of environmentalism. *Globalizations*, 18(2): 285–299.

de Vries, A. 2023. The growing energy footprint of artificial intelligence. *Joule*, 7(10): 2191–2194.

Desislavov, R.; Martínez-Plumed, F.; and Hernández-Orallo, J. 2023. Trends in AI inference energy consumption: Beyond the performance-vs-parameter laws of deep learning. *Sustainable Computing: Informatics and Systems*, 38: 100857.

Epoch AI. 2022. Parameter, Compute and Data Trends in Machine Learning. Accessed: 2024-07-19.

Forster, P.; Storelvmo, T.; Armour, K.; Collins, W.; Dufresne, J.-L.; Frame, D.; Lunt, D.; Mauritsen, T.; Palmer, M.; Watanabe, M.; Wild, M.; and Zhang, H. 2023. The Earth's Energy Budget, Climate Feedbacks and Climate Sensitivity. In Masson-Delmotte, V.; Zhai, P.; Pirani, A.; Connors, S.; Péan, C.; Berger, S.; Caud, N.; Chen, Y.; Goldfarb, L.; Gomis, M.; Huang, M.; Leitzell, K.; Lonnoy, E.; Matthews, J.; Maycock, T.; Waterfield, T.; Yelekçi, O.; Yu, R.; and Zhou, B., eds., *Climate Change 2021 – The Physical Science Basis: Working Group I Contribution to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, 923–1054. Cambridge University Press.

Gossart, C. 2015. Rebound Effects and ICT: A Review of the Literature. In Hilty, L. M.; and Aebischer, B., eds., *ICT Innovations for Sustainability*, 435–448. Cham: Springer International Publishing. ISBN 978-3-319-09228-7.

Greenfield, P. 2023. Revealed: more than 90% of rainforest carbon offsets by biggest certifier are worthless, analysis shows. The Guardian, https://www.theguardian.com/ environment/2023/jan/18/revealed-forest-carbon-offsetsbiggest-provider-worthless-verra-aoe.

Gupta, U.; Kim, Y. G.; Lee, S.; Tse, J.; Lee, H.-H. S.; Wei, G.-Y.; Brooks, D.; and Wu, C.-J. 2022. Chasing Carbon: The Elusive Environmental Footprint of Computing.

Jay, M.; Ostapenco, V.; Lefevre, L.; Trystram, D.; Orgerie, A.-C.; and Fichel, B. 2023. An experimental comparison of software-based power meters: focus on CPU and GPU. In *2023 IEEE/ACM 23rd International Symposium on Cluster, Cloud and Internet Computing (CCGrid)*, 106–118.

Jiang, H. H.; Brown, L.; Cheng, J.; Khan, M.; Gupta, A.; Workman, D.; Hanna, A.; Flowers, J.; and Gebru, T. 2023. AI Art and Its Impact on Artists. In *Proceedings of the 2023 AAAI/ACM Conference on AI, Ethics, and Society*, AIES '23, 363–374. New York, NY, USA: Association for Computing Machinery. ISBN 9798400702310.

Keyes, O. 2018. The Misgendering Machines: Trans/HCI Implications of Automatic Gender Recognition. *Proc. ACM Hum. Comput. Interact.*, 2(CSCW): 88:1–88:22.

Lannelongue, L.; Grealey, J.; and Inouye, M. 2021. Green Algorithms: Quantifying the Carbon Footprint of Computation. *Advanced Science*, 8(12): 2100707.

Lohmann, L. 2009. Toward a different debate in environmental accounting: The cases of carbon and cost–benefit. *Accounting, Organizations and Society*, 34(3): 499–534.

Luccioni, A. S.; Viguier, S.; and Ligozat, A.-L. 2023. Estimating the Carbon Footprint of BLOOM, a 176B Parameter Language Model. *Journal of Machine Learning Research*, 24(253): 1–15.

Masanet, E.; Shehabi, A.; Lei, N.; Smith, S.; and Koomey, J. 2020. Recalibrating global data center energy-use estimates. *Science*, 367(6481): 984–986.

Morand, C.; Névéol, A.; and Ligozat, A.-L. 2024. MLCA: a tool for Machine Learning Life Cycle Assessment. In *2024 International Conference on ICT for Sustainability (ICT4S)*. Stockholm, Sweden.

Mytton, D. 2021. Data center water consumption. *npj Clean Water*, 4.

Ortar, N.; Taylor, A. R. E.; Velkova, J.; Brodie, P.; Johnson, A.; Marquet, C.; Pollio, A.; and Cirolia, L. 2023. *4 Powering 'smart' futures: data centres and the energy politics of digitalisation*, 125–168. Berlin, Boston: De Gruyter. ISBN 9783110745641.

Ostrouchov, G.; Maxwell, D.; Ashraf, R. A.; Engelmann, C.; Shankar, M.; and Rogers, J. H. 2020. GPU Lifetimes on Titan Supercomputer: Survival Analysis and Reliability. In *SC20: International Conference for High Performance Computing, Networking, Storage and Analysis*, 1–14.

Patterson, D.; Gonzalez, J.; Hölzle, U.; Le, Q.; Liang, C.; Munguia, L.-M.; Rothchild, D.; So, D. R.; Texier, M.; and Dean, J. 2022. The Carbon Footprint of Machine Learning Training Will Plateau, Then Shrink. *Computer*, 55(7): 18– 28.

Pirson, T.; Delhaye, T. P.; Pip, A. G.; Le Brun, G.; Raskin, J.-P.; and Bol, D. 2023. The Environmental Footprint of IC Production: Review, Analysis, and Lessons From Historical Trends. *IEEE Transactions on Semiconductor Manufacturing*, 36(1): 56–67.

Roussilhe, G.; Pirson, T.; Xhonneux, M.; and Bol, D. 2024. From silicon shield to carbon lock-in? The environmental footprint of electronic components manufacturing in Taiwan (2015–2020). *Journal of Industrial Ecology*, n/a(n/a): 15.

Schwartz, R.; Dodge, J.; Smith, N. A.; and Etzioni, O. 2020. Green AI. *Commun. ACM*, 63(12): 54–63.

Sevilla, J.; Heim, L.; Ho, A.; Besiroglu, T.; Hobbhahn, M.; and Villalobos, P. 2022. Compute Trends Across Three Eras of Machine Learning. In *2022 International Joint Conference on Neural Networks (IJCNN)*, 1–8.

Shi, M.; Currier, K.; Liu, Z.; Janowicz, K.; Wiedemann, N.; Verstegen, J.; McKenzie, G.; Graser, A.; Zhu, R.; and Mai, G. 2023. Thinking Geographically about AI Sustainability. *AGILE: GIScience Series*, 4: 42.

Strubell, E.; Ganesh, A.; and McCallum, A. 2019. Energy and Policy Considerations for Deep Learning in NLP. In Korhonen, A.; Traum, D. R.; and Màrquez, L., eds., *Proceedings of the 57th Conference of the Association for Computational Linguistics, ACL 2019, Florence, Italy, July 28- August 2, 2019, Volume 1: Long Papers*, 3645–3650. Association for Computational Linguistics.

Thompson, N.; Greenewald, K.; Lee, K.; and Manso, G. F. 2023. The Computational Limits of Deep Learning. In *Ninth Computing within Limits 2023*. LIMITS. Https://limits.pubpub.org/pub/wm1lwjce.

van Oers, L.; Guinée, J. B.; and Heijungs, R. 2020. Abiotic resource depletion potentials (ADPs) for elements revisited—updating ultimate reserve estimates and introducing time series for production data. *The International Journal of Life Cycle Assessment*, 25: 294–308.

Velkova, J. 2019. Data Centres as Impermanent Infrastructures. *Culture Machine*, 18.

Verdecchia, R.; Sallou, J.; and Cruz, L. 2023. A systematic review of Green AI. *WIREs Data Mining and Knowledge Discovery*, 13(4): e1507.

World Health Organization. 2021. *Children and digital dumpsites: e-waste exposure and child health*. World Health Organization.

Wu, C.; Raghavendra, R.; Gupta, U.; Acun, B.; Ardalani, N.; Maeng, K.; Chang, G.; Behram, F. A.; Huang, J.; Bai, C.; Gschwind, M.; Gupta, A.; Ott, M.; Melnikov, A.; Candido, S.; Brooks, D.; Chauhan, G.; Lee, B.; Lee, H. S.; Akyildiz, B.; Balandat, M.; Spisak, J.; Jain, R.; Rabbat, M.; and Hazelwood, K. M. 2022. Sustainable AI: Environmental Implications, Challenges and Opportunities. In Marculescu, D.; Chi, Y.; and Wu, C., eds., *Proceedings of Machine Learning and Systems 2022, MLSys 2022, Santa Clara, CA, USA, August 29 - September 1, 2022*. mlsys.org.

Wu, C.-J.; Acun, B.; Raghavendra, R.; and Hazelwood, K. 2024. Beyond Efficiency: Scaling AI Sustainably. arXiv:2406.05303.