DOI: xxx/xxxx

RESEARCH ARTICLE

Biomarker combination based on the Youden index with and without gold standard

Ao Sun¹ | Yanting Li² | Xiao-Hua Zhou*³

- ¹Center of Data Science, Peking University, Beijing, China
- ²Rheumatology Department, Guang'anmen Hospital, Beijing, China
- ³Department of Biostatistics and Beijing International Center for Mathematical Research, Peking University, Beijing, China

Correspondence

Xiao-Hua Zhou, Department of Biostatistics and Beijing International Center for Mathematical Research, Peking University, Beijing, China. Email: azhou@math.pku.edu.cn

Abstract

In clinical practice, multiple biomarkers are often measured on the same subject for disease diagnosis, and combining them can improve diagnostic accuracy. Existing studies typically combine multiple biomarkers by maximizing the Area Under the ROC Curve (AUC), assuming a gold standard exists or that biomarkers follow a multivariate normal distribution. However, practical diagnostic settings require both optimal combination coefficients and an effective cutoff value, and the reference test may be imperfect. In this paper, we propose a two-stage method for identifying the optimal linear combination and cutoff value based on the Youden index. First, it maximizes an approximation of the empirical AUC to estimate the optimal linear coefficients for combining multiple biomarkers. Then, it maximizes the empirical Youden index to determine the optimal cutoff point for disease classification. Under the semiparametric single index model and regularity conditions, the estimators for the linear coefficients, cutoff point, and Youden index are consistent. This method is also applicable when the reference standard is imperfect. We demonstrate the performance of our method through simulations and apply it to construct a diagnostic scale for Chinese medicine.

KEYWORDS:

Youden index; linear combination; imperfect reference test; Chinese medicine scales

1 | INTRODUCTION

In fields like meteorology, economics, and computer science, leveraging multifaceted information for classification and prediction is crucial and has long been studied. Similarly, in medical practice, disease classification and prediction using clinical and laboratory data are key areas of research. Diagnostic studies often involve multiple tests on individuals, and no single test or biomarker is sufficient for precise diagnosis or prognosis. Therefore, integrating various sources of information is essential to enhance diagnostic accuracy ¹.

Various methods exist for combining multiple biomarkers, with the simplest and most popular being linear combination. Pepe et al.² employed likelihood maximization to estimate linear combination coefficients. Another common objective for linear combination is maximizing the area under the ROC curve (AUC), the most widely used index for summarizing the ROC curve, which evaluates the diagnostic accuracy of continuous or ordinal biomarkers³. Su and Liu⁴, Reiser and Faraggi⁵, Liu et al.⁶ optimized the AUC under the assumption that biomarkers follow a mixture of multivariate normal distributions. Pepe et al.² relaxed the assumption of a parametric distribution, advocating for using the empirical AUC as the objective function and

performing a grid search for the optimal linear combinations in a p-dimensional space. However, grid search becomes computationally challenging with more than three biomarkers, leading to the development of more efficient combination methods such as the min-max approach⁷, stepwise method⁸, and approximation methods^{9,10}.

While the AUC offers a comprehensive overview of diagnostic accuracy across all possible thresholds, a specific cutoff point is essential for diagnostic purposes. Surprisingly, very few research articles have explored combining biomarkers using alternative summary statistics of the ROC curve as objective functions beyond AUC. Some studies have numerically searched for optimal linear combinations that maximize sensitivity at a fixed specificity ^{11,12}, while several methods have identified the best linear combination by maximizing partial AUC ^{13,14,15}.

The Youden index is another widely used summary index of the ROC curve ¹⁶. It represents the maximum of the sum of sensitivity and specificity minus one, ranging from 0 to 1, where 1 indicates complete separation of biomarker distributions in healthy and diseased populations, and 0 indicates complete overlap. In medical and biological sciences, it is extensively used for selecting diagnostic thresholds due to its seamless integration with the ROC framework ¹⁷ and its direct measurement of the maximum overall correct classification rate achievable by a biomarker. Combining biomarkers using the Youden index yields the highest possible overall correct classification rate at the diagnostic threshold among all feasible linear combinations, resulting in optimal diagnostic accuracy.

Existing methods for maximizing the Youden index often overlook the properties of the estimator and the inference process, focusing primarily on scenarios with a gold standard ^{18,19,20}. In practice, labels for diseased and healthy individuals are often based on imperfect reference tests, leading to potential misclassifications. Ignoring these misclassifications during diagnostic test development can severely bias accuracy parameter estimates ²¹. To address the absence of a gold standard, latent-class models with two latent classes have been proposed. These models linearly combine continuous biomarkers to maximize AUC under the assumption of a multivariate normal distribution of biomarker values ^{22,23,24}. However, in real-world datasets, the assumption of a multivariate normal distribution is often too restrictive to be appropriate.

In this article, we introduce a two-stage method for estimating optimal combination coefficients and threshold, along with the corresponding Youden index. We establish the uniform consistency of the proposed estimators and propose a procedure to construct the confidence interval for the Youden index. Additionally, we demonstrate that our method achieves optimal combination coefficients and thresholds even when the reference standard is imperfect. To our knowledge, this is the first paper to address linear combination of biomarkers without parametric assumptions in the presence of an imperfect reference standard.

The remainder of the paper is organized as follows. Sections 2.1 and 2.2 introduce notations and provide a brief review of existing methods. Section 3 presents the newly proposed two-stage combination method. Section 4 discusses the linear combination approach when the reference test is imperfect. Numerical studies, aimed at evaluating the performance of the method and applying it to a real data example, are presented in Section 5. Finally, Section 6 offers a discussion.

2 | PRELIMINARIES

2.1 | Notations

Consider a scenario involving n patients, categorized into two groups: diseased and healthy. Let D denote the presence (D=1) or absence (D=0) of the disease. There are n_1 diseased subjects and n_0 healthy subjects. Each patient is measured across p biomarkers, denoted as T_1, \dots, T_p . For the diseased group, let T^1 represent the p-dimensional vector of biomarker measurements, where $T_i^1 = (T_{i1}^1, T_{i2}^1, \dots, T_{ip}^1)$ denotes the observed biomarker values for the ith diseased subject ($i=1,\dots,n_1$). Here, T_{ik}^1 is the value of the kth biomarker for the ith individual. Similarly, for the healthy group, $T_j^0 = (T_{j1}^0, T_{j2}^0, \dots, T_{jp}^0)$ represents the biomarker values for the jth healthy subject ($j=1,\dots,n_0$), and T_{jk}^0 is the measurement of the kth biomarker for the jth healthy individual.

We consider linear combination scores of the form $\omega_0'T = T_1 + \cdots + \omega_{0p}T_p$, where $\omega_0 = (1, \omega_{02}, \cdots, \omega_{0p})$ is the vector of combination coefficients. This linear score omits an intercept, and the coefficient for T_1 is set to 1 without loss of generality. If $\omega_{01} > 0$, a linear predictor of the form $\gamma + \omega_{01} \cdot \omega_0'T$ exceeding a threshold is equivalent to using $\omega_0'T$. Additionally, we can redefine T_1 as $-T_1$ to ensure $\omega_{01} > 0$, covering all possible cases. Since the receiver operating characteristic (ROC) curves for $\omega_0'T$ and $\gamma + \omega_{01} \cdot \omega_0'T$ are identical, it suffices to focus on the linear combination $\omega_0'T$. If the relationship between T and D can be described using a semiparametric single index model, the probability of disease can be expressed as: $\mathbb{P}(D=1|T) = H(\omega_0'T)$, where H is an unknown increasing link function. According to Neyman and Pearson 25, the combination score $\omega_0'T$ yields the optimal ROC curve, meaning no other combination score based on T can outperform it at any accuracy point (FPR, TPR) on the

ROC curve. Therefore, for a fixed false positive rate (FPR), the true positive rate (TPR) for the rule $\omega'_0 T > c$ is higher than that of any other score with the same FPR. The Youden index, which quantifies the maximum vertical distance between the ROC curve and the diagonal $y = x^{26}$, is defined as:

$$J_{\omega_0} = \max_{c} \left\{ Se_{\omega_0}(c) + Sp_{\omega_0}(c) - 1 \right\} = \max_{c} \left\{ F_{0\omega_0}(c) - F_{1\omega_0}(c) \right\},\tag{1}$$

where $\mathrm{Se}_{\omega_0}(c)=1-F_{1\omega_0}(c)$ is the sensitivity, and $\mathrm{Sp}_{\omega_0}(c)=F_{0\omega_0}(c)$ is the specificity at threshold c. Here, $F_{1\omega_0}(\cdot)$ and $F_{0\omega_0}(\cdot)$ denote the cumulative distribution functions of $\omega_0'T^1$ and $\omega_0'T^0$, respectively, while $f_{0\omega_0}(\cdot)$ and $f_{1\omega_0}(\cdot)$ are their corresponding densities. Given the optimality of the ROC curve, the score $\omega_0'T$ achieves the maximum Youden index among all possible combination scores.

We typically assume that the supports of $\omega_0'T^1$ and $\omega_0'T^0$ overlap. Moreover, we assume the existence of a threshold c_0 such that $f_{1\omega_0}(c_0)=f_{0\omega_0}(c_0)$, with $f_{0\omega_0}(t)< f_{1\omega_0}(t)$ for $t< c_0$ and $f_{0\omega_0}(t)> f_{1\omega_0}(t)$ for $t>c_0$. This condition holds, for instance, when the likelihood ratio is monotonic. It implies that $\omega_0'T^0$ is stochastically smaller than $\omega_0'T^1$, i.e. $F_{1\omega_0}(t)\geq F_{0\omega_0}(t)$ for all t. Under this assumption, the sum $Se_{\omega_0}(c)+Sp_{\omega_0}(c)$ is maximized by selecting $c=c_0$. The Youden index for the linear combination score $\omega_0'T$, denoted as J_{ω_0} , then can be given by:

$$J_{\omega_0} = F_{0\omega_0}(c_0) - F_{1\omega_0}(c_0). \tag{2}$$

Our primary objective is to estimate the optimal combination coefficients ω_0 that maximize the Youden index, the corresponding cutoff point c_0 , and the maximum Youden index J_{ω_0} .

2.2 | Existing methods

Existing methods for finding optimal linear combination of multiple biomarkers to maximize the Youden index can be categorized into empirical searching methods and derivation-based numerical searching methods. These methods lack well-defined estimator properties and do not address scenarios where the reference test is imperfect.

2.2.1 | Empirical searching methods

Given ω , denote the optimal cutoff point as c_{ω} . The empirical estimate of the Youden index J_{ω} is:

$$\widehat{J}_{\omega} = \frac{\sum_{j=1}^{n_0} \mathbb{I}\left(\omega' T_j^0 \le c_{\omega}\right)}{n_0} - \frac{\sum_{i=1}^{n_1} \mathbb{I}\left(\omega' T_i^1 \le c_{\omega}\right)}{n_1}.$$
(3)

Optimizing this non-continuous objective function via grid search for multiple biomarkers has a computational cost of $\mathcal{O}(n^p)$, where n is the sample size and p is the number of biomarkers.

Two empirical searching methods, the min-max method and the stepwise method, have been proposed to address this computational challenge 22 . The min-max method simplifies the optimization by only considering the minimum and maximum values of the biomarker measurements, reducing the *p*-dimensional grid search to a single coefficient ω . However, a key limitation of the min-max approach is its reliance solely on the minimum and maximum values of the biomarkers, which may not result in the optimal linear combination.

The stepwise approach reduces computational complexity by initially calculating the empirical Youden index for each biomarker and ranking them. It starts by combining the two biomarkers with the highest Youden indices, using grid search to optimize the coefficients, then sequentially adds additional biomarkers. While effective, this stepwise method is not guaranteed to find the global optimum and can be cumbersome when dealing with binary biomarkers.

2.2.2 | Derivation-based numerical searching methods

There are two derivation-based methods for numerically searching for the optimal linear combination maximizing the Youden index ²². One method is parametric, assuming biomarkers follow a multivariate normal distribution in both diseased and healthy states. Under this assumption, explicit solutions for the optimal cutoff value and Youden index can be derived. However, this assumption may not hold in real-world scenarios.

The other method is non-parametric, using an approximation for the indicator function in objective function (3) and simultaneously maximizing it with respect to ω and c. The estimator $(\hat{\omega}^{SIM}, \hat{c}^{SIM})$ is defined:

$$(\widehat{\omega}^{\text{SIM}}, \widehat{c}^{\text{SIM}}) = \arg\max_{\omega, c} \left\{ \frac{1}{n_0} \sum_{j=1}^{n_0} \Phi\left(\frac{c - \omega' T_j^0}{h}\right) - \frac{1}{n_1} \sum_{i=1}^{n_1} \Phi\left(\frac{c - \omega' T_i^1}{h}\right) \right\}.$$

However, the coefficients ω and the cutoff value c_{ω} are highly correlated, making simultaneous optimization with quasi-Newton algorithm prone to converging to local minima 27 .

3 | THE PROPOSED METHOD

In this section, we propose a two-stage method to estimate the optimal linear coefficients and cutoff point, establish the consistency of the corresponding estimators, and introduce a new confidence interval for the optimal Youden index.

3.1 | Estimation

Given that under the single index model, the optimal combination score with the largest Youden index also has the largest AUC, we propose a two-stage approach to estimate the coefficients ω_0 and the cutoff point c_0 . In the first stage, we aim to maximize the empirical AUC to estimate the optimal linear combination coefficients ω_0 . The empirical AUC of a linear risk score, $\omega' T$, is defined as:

$$\widehat{AUC}(\omega) = \frac{\sum_{i=1}^{n_1} \sum_{j=1}^{n_0} \left\{ \mathbb{I}\left[\omega' T_i^1 > \omega' T_j^0\right] + \frac{1}{2} \mathbb{I}\left[\omega' T_i^1 = \omega' T_j^0\right] \right\}}{n_1 n_0},\tag{4}$$

where $\mathbb{I}(\cdot)$ is the indicator function. This objective function is discontinuous due to the indicator function, necessitating brute force search or specialized algorithms. The computational cost grows exponentially with n^p , making optimization based on $\widehat{AUC}(\omega)$ impractical for large numbers of biomarkers.

To address this, we approximate the indicator function with a differentiable function. According to Lin et al. ²⁸, using the standard normal distribution function for this approximation is more accurate and stable than using the sigmoid function. Therefore, we use $\Phi\left(\left(\omega'T_i^1-\omega'T_j^0\right)/h_n\right)$ as a smooth approximation to $\mathbb{E}\left[\omega'T_i^1>\omega'T_j^0\right]+\frac{1}{2}\mathbb{E}\left[\omega'T_i^1=\omega'T_j^0\right]$, where $\Phi(\cdot)$ is the cumulative distribution function of the standard normal variable and h_n is a bandwidth chosen to converge to zero, specifically $h_n=(n_1n_0)^{-0.1}$, as recommended by Vexler et al. ²⁹ The resulting estimator, denoted as $\widehat{\omega}$, is defined as:

$$\widehat{\omega} = \arg\max_{\omega} \left\{ \frac{1}{n_1 n_0} \sum_{i=1}^{n_1} \sum_{j=1}^{n_0} \Phi\left(\frac{\omega' T_i^1 - \omega' T_i^0}{h}\right) \right\}. \tag{5}$$

The estimator is identifiable up to a scale constant if at least one component of T is continuous.

After obtaining the estimator of coefficients, the optimal cutoff point is estimated by:

$$\widehat{c}^{\text{TS}} = \operatorname{median} \left\{ c : \max_{c} \frac{\sum_{j=1}^{n_0} \mathbb{I}\left(\widehat{\omega}' T_j^0 \le c\right)}{n_0} - \frac{\sum_{i=1}^{n_1} \mathbb{I}\left(\widehat{\omega}' T_i^1 \le c\right)}{n_1} \right\}.$$
 (6)

Alternatively, the maximum or minimum value instead of the median can be used in (6).

The estimated Youden index, denoted as \hat{J}^{TS} , is given by:

$$\widehat{J}^{\text{TS}} = \frac{\sum_{j=1}^{n_0} \mathbb{I}\left(\widehat{\omega}' T_j^0 \le \widehat{c}^{\text{TS}}\right)}{n_0} - \frac{\sum_{i=1}^{n_1} \mathbb{I}\left(\widehat{\omega}' T_i^1 \le \widehat{c}^{\text{TS}}\right)}{n_1}.$$
 (7)

3.2 | Asymptotic properties

The asymptotic properties of $\widehat{\omega}$ have been proven by Ma and Huang⁹. For completeness, we briefly review these properties. Let $\theta_0 = (\omega_{02}, \cdots, \omega_{0p})$, and $\widehat{\theta}_n = (\widehat{\omega}_2, \cdots, \widehat{\omega}_p)$. For the consistency of $\widehat{\theta}_n$, we assume the following conditions: (A1) The true parameter value θ_0 is an interior point of θ , which is a compact subset of \mathbb{R}^{p-1} . (A2) Let S_T denote the support of the biomarker

vector T, where S_T is not contained in any proper linear subspace of \mathbb{R}^p and the first component of T has an everywhere positive density, conditional on the other components.

Lemma 1. Suppose assumptions (A1) and (A2) hold and $h_n \to 0$ as $n \to \infty$. Then $\hat{\omega} \to_P \omega_0$ as $n \to \infty$.

For the consistency of \hat{c}^{TS} and \hat{J}^{TS} , we assume: (B) For any $\delta > 0$, there exists $\epsilon > 0$, such that $\sup_{|x-c_0|>\delta}\left[F_{0\omega_0}(x)-F_{1\omega_0}(x)\right] < F_{0\omega_0}(c_0)-F_{1\omega_0}(c_0)-\epsilon$. This condition is slightly weaker than the assumption in Section 2.1 and ensures that the maximum of $F_{0\omega_0}(x)-F_{1\omega_0}(x)$ occurs at c_0 and is separated from other values by at least ϵ .

Theorem 1. Suppose assumptions (A1), (A2) and (B) hold and $h_n \to 0$ as $n \to \infty$. Then $\hat{c}^{TS} \to_P c_0$ and $\hat{J}^{TS} \to_P J_{\infty}$ as $n \to \infty$.

Theorem 1 provides asymptotic consistency for estimators of the cutoff point and Youden index, with the proof presented in Appendix A.

3.3 | Inference for Youden index

According to Shan³⁰, the coverage property of Wald-type confidence intervals for the Youden index is generally unsatisfactory and we use the square-and-add limits based on the Wilson score method to construct the non-parametric confidence interval for the Youden index J_{ω_0} .

For simplicity, denote $p_1 = P(T^1 \le c_0)$ and $p_0 = P(T^0 \le c_0)$. The Wilson confidence intervals for p_1 , denoted as (l_1, u_1) , are the roots of the following equality:

$$(p_1 - \hat{p}_1)^2 = z_{1-\alpha/2}^2 \frac{p_1(1-p_1)}{n_1}.$$

From this, we can calculate:

$$l_1 = \frac{1}{1 + z_{1-\alpha/2}^2/n_1} \left[\hat{p}_1 + \frac{z_{1-\alpha/2}^2}{2n_1} - z_{1-\alpha/2} \sqrt{\frac{\hat{p}_1(1-\hat{p}_1)}{n_1} + \frac{z_{1-\alpha/2}^2}{4n_1^2}} \right],$$

and

$$u_1 = \frac{1}{1 + z_{1-\alpha/2}^2/n_1} \left[\hat{p}_1 + \frac{z_{1-\alpha/2}^2}{2n_1} + z_{1-\alpha/2} \sqrt{\frac{\hat{p}_1(1-\hat{p}_1)}{n_1} + \frac{z_{1-\alpha/2}^2}{4n_1^2}} \right].$$

The confidence intervals for $\text{Var}(\widehat{p}_1)$ can be estimated by $\widehat{\text{Var}}_l(\widehat{p}_1) = l_1(1 - l_1)/n_1$, $\widehat{\text{Var}}_u(\widehat{p}_1) = u_1(1 - u_1)/n_1$.

The Wilson confidence interval for p_0 can be calculated similarly and denoted as (l_0, u_0) . The confidence intervals for $Var(\widehat{p}_0)$ can be estimated as $\widehat{Var}_l(\widehat{p}_0) = l_0(1 - l_0)/n_0$ and $\widehat{Var}_u(\widehat{p}_0) = u_0(1 - u_0)/n_0$.

Assuming independence of test results between the healthy and diseased groups, after obtaining $\widehat{\omega}$ and \widehat{c}^{TS} , the variance of \widehat{J}^{TS} can be consistently estimated as $\widehat{\text{Var}}(\widehat{J}^{TS}) = \widehat{\text{Var}}(\widehat{p}_0 - \widehat{p}_1) = \widehat{\text{Var}}(\widehat{p}_0) + \widehat{\text{Var}}(\widehat{p}_1)$.

We use the AC estimators for p_1 and p_0 , as proposed by Agresti and Choll³¹, denoted as \hat{p}_1 and \hat{p}_0 , which are as follows:

$$\widehat{p}_0 = \frac{\sum_{j=1}^{n_0} \mathbb{I}\left(\widehat{\omega}' T_j^0 \leq \widehat{c}\right) + \frac{1}{2} z_{1-\alpha/2}^2}{n_0 + z_{1-\alpha/2}^2}, \ \widehat{p}_1 = \frac{\sum_{i=1}^{n_1} \mathbb{I}\left(\widehat{\omega}' T_i^1 \leq \widehat{c}\right) + \frac{1}{2} z_{1-\alpha/2}^2}{n_1 + z_{1-\alpha/2}^2}.$$

And the adjusted estimator for the Youden index is defined as:

$$\widehat{J}^{\text{AC}} = \frac{\sum_{j=1}^{n_0} \mathbb{I}\left(\widehat{\omega}' T_j^0 \leq \widehat{c}\right) + \frac{1}{2} z_{1-\alpha/2}^2}{n_0 + z_{1-\alpha/2}^2} - \frac{\sum_{i=1}^{n_1} \mathbb{I}\left(\widehat{\omega}' T_i^1 \leq \widehat{c}\right) + \frac{1}{2} z_{1-\alpha/2}^2}{n_1 + z_{1-\alpha/2}^2},$$

According to Zhou and Qin³², the adjusted estimator \hat{J}^{AC} has smaller bias than the empirical estimator \hat{J}^{TS} when $J \leq 0.8$ and when sample size are small. Therefore, we propose calculating the confidence interval of J_0 as:

$$(J_L,J_U) = \left(\widehat{J}^{\mathrm{AC}} - z_{1-\alpha/2} \sqrt{\widehat{\mathrm{Var}}_l(\widehat{p}_0) + \widehat{\mathrm{Var}}_u(\widehat{p}_1)}, \widehat{J}^{\mathrm{AC}} + z_{1-\alpha/2} \sqrt{\widehat{\mathrm{Var}}_u(\widehat{p}_0) + \widehat{\mathrm{Var}}_l(\widehat{p}_1)}\right).$$

This confidence interval does not require bootstrap sampling, making it computationally efficient.

4 | WHEN THE REFERENCE TEST IS IMPERFECT

In numerous scenarios, while the true disease status, or gold standard, have a well-defined clinical definition, practical constraints can render it unobservable or latent. Instead, we observe an imperfect reference test that is prone to errors ^{33,34}. Existing methods for combining biomarkers, when only imperfect gold standards are available, assume a multivariate normal distribution for the joint distribution of biomarkers and focus on maximizing the AUC rather than the Youden index ^{22,23,24}. We will identify the optimal linear combination and cutoff point that maximizes the Youden index without making parametric assumptions.

Denote the imperfect reference test as R. The positive predictive value (ppv) of R is defined as ppv = $\mathbb{P}(D=1|R=1)$ and the negative predictive value (npv) is defined as npv = $\mathbb{P}(D=0|R=0)$. We make two general assumptions:

Assumption 1. ppv + npv > 1.

Assumption 2. Given the true disease status, biomarkers and the imperfect reference test are independent.

Assumption 1 indicates that the reference test is more accurate than random guessing, while Assumption 2 suggests that biomarkers and the reference standard are improbable to misdiagnose the same patients.

When considering the imperfect test R as the reference, for a linear combination $\omega'T$, we denote the corresponding proxy AUC as \widetilde{AUC}_{ω} , which is defined as:

$$\widetilde{AUC}(\omega) = P(\omega'T_i > \omega'T_i | R_i = 1, R_i = 0), \tag{8}$$

and its relationship with AUC(ω), which is the true AUC for score $\omega'T$, can be expressed as:

$$\widetilde{AUC}(\omega) = (ppv + npv - 1)AUC(\omega) - \frac{1}{2}(ppv + npv) + 1.$$
(9)

Similarly, we denote the Youden index when R is regarded as the reference as \widetilde{J}_{ω} , defined as:

$$\widetilde{J}_{\omega} = \max_{\alpha} \left\{ P(\omega'T > c | R = 1) + P(\omega'T < c | R = 0) - 1 \right\},\tag{10}$$

and its relationship with J_{ω} , which is the true Youden index, can be expressed as:

$$\widetilde{J}_{\omega} = (ppv + npv - 1)J_{\omega}. \tag{11}$$

Under assumption 1, $\widetilde{AUC}(\omega)$ and \widetilde{J}_{ω} increase monotonically with $AUC(\omega)$ and J_{ω} , respectively. Consequently, the combination coefficients ω_0 that maximize $\widetilde{AUC}(\omega)$ also maximize $\widetilde{AUC}(\omega)$. Similarly, the cutoff point c_0 that achieves J_{ω_0} also achieves the maximum \widetilde{J}_{ω_0} . The derivation of (9) and (11) can be found in Appendix B.

4.1 | Estimation

Assume there are \tilde{n}_1 subjects with diseased imperfect reference result and \tilde{n}_0 subjects with healthy imperfect reference result. Denote the biomarker measurements, whose result of imperfect reference standard is diseased, as \tilde{T}^1 , and \tilde{T}^0 as the measurements with healthy reference result. Denote the proxy AUC estimator as $\check{\omega}$, which is calculated as:

$$\breve{\omega} = \arg\max_{\omega} \left\{ \frac{1}{\tilde{n}_1 \tilde{n}_0} \sum_{i=1}^{\tilde{n}_1} \sum_{j=1}^{\tilde{n}_0} \Phi\left(\frac{\omega' \tilde{T}_i^1 - \omega' \tilde{T}_i^0}{h}\right) \right\}.$$
(12)

This estimator is also only identifiable up to a scale constant if at least one component of T is continuous.

After obtaining the estimator of coefficients, the cutoff point is estimated by

$$\check{c} = \operatorname{median} \left\{ c : \max_{c} \frac{\sum_{j=1}^{\tilde{n}_{0}} \mathbb{I}\left(\check{\omega}'\widetilde{T}_{j}^{0} \leq c\right)}{\tilde{n}_{0}} - \frac{\sum_{i=1}^{\tilde{n}_{1}} \mathbb{I}\left(\check{\omega}'\widetilde{T}_{i}^{1} \leq c\right)}{\tilde{n}_{1}} \right\}.$$
(13)

Alternatively, we can use the maximum or minimum value instead of the median in (13).

The estimated proxy Youden index, denoted as J, is given by

$$\check{J} = \frac{\sum_{j=1}^{\tilde{n}_0} \mathbb{I}\left(\check{\omega}'\widetilde{T}_j^0 \le \check{c}\right)}{\tilde{n}_0} - \frac{\sum_{i=1}^{\tilde{n}_1} \mathbb{I}\left(\check{\omega}'\widetilde{T}_i^1 \le \check{c}\right)}{\tilde{n}_1}.$$
(14)

4.2 | Asymtotic properties

For identifiability, we also assume the first component of ω_0 is equal to 1. The asymptotic properties of $\check{\omega}$ can be seen a simple extension to Lemma 1.

Theorem 2. Suppose assumptions (A1) and (A2) hold and $h_n \to 0$ as $n \to \infty$. Then $\check{\omega} \to_P \omega_0$ as $n \to \infty$.

The consistency of \check{c} and \check{J} can also be seen as a simple extension to Theorem 1. We assume that: (C) For any $\delta>0$, there exists $\epsilon>0$, such that $\sup_{|x-c_0|>\delta}\left[\widetilde{F}_{0\omega_0}(x)-\widetilde{F}_{1\omega_0}(x)\right]<\widetilde{F}_{0\omega_0}(c_0)-\widetilde{F}_{1\omega_0}(c_0)-\epsilon$, where $\widetilde{F}_{0\omega_0}(x)$ is defined as $\mathbb{P}(\omega'T\leq x|R=0)$ and $\widetilde{F}_{1\omega_0}(x)$ is defined as $\mathbb{P}(\omega'T\leq x|R=1)$.

Theorem 3. Suppose assumption (A1), (A2) and (C) hold, and $h_n \to 0$ as $n \to \infty$. Then $\check{c} \to_P c_0$ and $(ppv + npv - 1)\check{J} \to_P J_{\omega_0}$ as $n \to \infty$.

In conclusion, when only an imperfect reference test is available and the two general assumptions hold, we can effectively treat the imperfect reference test as the gold standard. We can then utilize the proposed two-stage method to determine the optimal coefficients and cutoff value.

5 | NUMERICAL STUDIES

This section comprises four parts: Section 5.1 presents the coverage probabilities and interval lengths of the proposed confidence interval. Section 5.2 compares our proposed two-stage method with the simultaneous optimization approach under both perfect and imperfect reference tests. Section 5.3 evaluates the performance of the two-stage method relative to the simultaneous optimization method across scenarios where the single-index model holds and where it does not. Lastly, Section 5.4 demonstrates the application of our proposed method to real-world data.

5.1 | Coverage probabilities and lengths of confidence intervals

We assess the coverage probabilities and average lengths of the proposed confidence interval and demonstrate its effectiveness by generating 1000 random samples with a total size of n. Specially, we drew samples of size $\pi \times n$ from the distribution function for test results of diseased patients and another independent sample of size $(1 - \pi) \times n$ from the distribution function for healthy subjects. The sample size n were set to 100, 200 and 400, with prevalence π set to 0.25, 0.5 and 0.75, respectively. Through simulation, we compare the coverage rate of our confidence interval with that of the NP confidence interval proposed by Shan³⁰, which uses the same variance estimators but empirical Youden index estimator.

In our study, multivariate normal distributions were chosen as the underlying distributions. Specifically, we selected $T^0 \sim MVN\left(0,I_5\right)$ for healthy subjects and $T^1 \sim MVN\left(\mu_1,I_5\right)$ for diseased subjects. The optimal linear coefficient is $\omega_0 = \mu_1$, and μ_1 was chosen to achieve predefined Youden index values of 0.45, 0.60, 0.70, and 0.85. The results are summarized in Table 1.

From the results, it is evident that as the sample size increases, the length of the confidence interval decreases. In balanced data scenarios ($\pi = 0.50$), the confidence interval is narrower compared to unbalanced data scenarios ($\pi = 0.25$ or 0.75). Additionally, the larger the true Youden index value, the narrower the confidence interval. Compared to the previously used NP confidence interval, our proposed confidence interval generally has higher coverage rates, consistently achieving the nominal 95% level.

5.2 | Investigation of performance with perfect and imperfect reference tests

We conducted simulation studies to evaluate the performance of our newly proposed two-stage method under perfect and imperfect reference tests when the single index model is satisfied. The Youden index was predefined as 0.45 and 0.70, with sample sizes of 200, 400, and 800 and prevalences of 0.25, 0.5 and 0.75. These were randomly split into training and testing sets with equal sample size. The sensitivity and specificity of the reference test were set to 1, 0.95, 0.90, 0.85, respectively. For each scenario, 1000 random samples were generated.

The results are displayed in Table 2 and indicate that as the sensitivity and specificity of the reference test decrease, performance deteriorates in both training and testing sets. As the sample size increases, the Youden index values approach the

true value J_0 in both the training and testing set. And in all scenarios, our method outperforms the simultaneous optimization method, with the advantage becoming more pronounced as the reference test's accuracy decreases.

5.3 | Comparison of the proposed method and existing method

We also conducted simulation studies to compare the performance of our newly proposed two-stage method with the existing simultaneous optimization method under scenarios where the single index model is satisfied and not satisfied. We present results for two scenarios: multivariate normal distribution with equal covariance matrix and unequal covariance matrices. When the covariance matrices differ between diseased and healthy populations, the single index model is not satisfied.

In the scenario with equal covariance matrices, diseased and healthy samples were drawn from multivariate normal distributions with mean vectors $\mu_0 = (0,0,0,0,0)'$ and $\mu_1 = (0.4,0.7,1.0,1.3,1.6)'$. Both groups shared the same covariance matrix, $\Sigma_1 = \Sigma_0 = (1-\gamma)I_5 + \gamma J_5$, where I_5 is the identity matrix, J_5 is a matrix of all ones, and γ was set to 0.3, 0.5, and 0.7. In the scenario with unequal covariance matrices, diseased and healthy samples were drawn from multivariate normal distributions with the same mean vectors but different covariance matrices: $\Sigma_1 = 0.3I_5 + 0.7J_5$ and $\Sigma_0 = 0.7I_5 + 0.3J_5$.

For both settings, prevalences were set to 0.25, 0.5 and 0.75. We generated 1000 random samples for each simulation setting, with sample size of n = 200, n = 400, and n = 800. Each dataset was randomly split into equal-sized training and testing sets. The coefficients, mean and variance of the estimated Youden index in the training and testing sets are presented in Tables 3 and 4.

The simulation results demonstrate that our newly proposed method performs well in both scenarios, whether the single index model holds (Table 3) or not (Table 4). Our two-stage method consistently achieves higher Youden index values compared to the simultaneous optimization method. Furthermore, our method exhibits lower variance, indicating greater stability in performance.

We conducted additional simulations with all biomarkers being binary, where the cutoff value and linear coefficients may not be unique, to assess whether our method can achieve a higher Youden index and thereby evaluate its robustness. Five biomarkers were generated following a Bernoulli distribution with parameters p = (0.7, 0.6, 0.5, 0.4, 0.3) respectively. The risk score was modeled using logistic regression, where coefficients were chose to achieve prevalences of 0.25, 0.5 and 0.75. For each scenario, sample sizes of 200, 400 and 800 were used, with samples equally divided into training and testing sets. The results are presented in Table 5. The findings indicate that even with all biomarkers being binary, our method demonstrates superior performance with lower variance compared to the simultaneous optimization method. As the sample size increases, the Youden index in the testing sets improves while variance decreases.

5.4 | Data example

Rheumatoid arthritis (RA) is a chronic autoimmune disease that primarily affects joints and surrounding tissues, causing inflammation, joint damage, disability, and work loss³⁵. RA patients face increased mortality risks from cardiovascular diseases, respiratory conditions, and infections compared to the general population³⁶. Pharmacological treatments, including various forms of disease-modifying antirheumatic drugs (DMARDs) and glucocorticoids, aim to achieve sustained remission or low disease activity³⁷. However, some patients experience poor responses or adverse effects, necessitating exploration of alternative therapies³⁸.

Nowadays, traditional Chinese medicine (TCM) is increasingly recognized worldwide, with evidence suggesting that integrative TCM and Western medicine can improve treatment outcomes compared to Western medicine alone ³⁹. In TCM, syndrome differentiation (ZHENG) is central to treatment, functioning as a diagnostic framework guided by TCM theory. However, syndrome differentiation relies heavily on practitioners' subjective judgment, influenced by clinical experience and training. This variability often results in diagnostic inconsistencies, with reported inter-practitioner agreement as low as 30% ⁴⁰. To address this, standardized diagnostic scales for various conditions, including coronary heart disease, heart failure, and ulcerative colitis, have been developed ^{41,42,43}.

RA has been a key focus of traditional Chinese medicine (TCM) for over 2000 years, rooted in classical texts like the Inner Canon of Huangdi (Huangdi Neijing). Damp-Heat Impeding Syndrome (DHIS) is a common TCM pattern in RA, affecting 43.86% of cases 44. Network meta-analyses suggest that herbs targeting DHIS can reduce inflammation, alleviate symptoms, and improve drug tolerance 45. Diagnostic scales for DHIS have been developed, but they primarily rely on the Delphi method, which collects expert opinions through multiple rounds of anonymous surveys 46. This method depends solely on expert consensus

without incorporating data-driven analysis, limiting its reliability. The latest DHIS diagnostic scale yields a Youden index of just 0.28, with a specificity of 1 but a sensitivity of only 0.28, indicating overly conservative performance.⁴⁷

In this study, we construct a diagnostic scale for Damp-Heat Impeding Syndrome (DHIS) using our newly proposed method. We utilize data from the China Rheumatoid Arthritis Registry of Patients with Chinese Medicine (CERTAIN), the first prospective, multicenter registry for RA patients receiving TCM treatment 48. CERTAIN includes over 14,000 RA patients from 145 centers across 30 provinces in China. Our analysis focuses on 4,310 patients with clinician-confirmed diagnoses and complete records of 27 symptoms relevant to DHIS, along with two supporting laboratory indicators. To ensure data accuracy, clinical symptoms—including tongue and pulse manifestations—were initially recorded by a junior physician and subsequently reviewed and corrected by a senior physician. Syndrome diagnoses were independently made by an associate chief physician or higher. Therefore, it is reasonable to assume that the recorded symptoms and syndrome diagnoses may introduce independent errors in classification 49.

We split the dataset into a training set (70%) and a testing set (30%) to compare the performance of our proposed two-stage method with the existing simultaneous optimization method for constructing TCM scales to diagnose dampness-heat obstruction syndrome. The results, summarized in Table 6, include the optimal Youden index, sensitivity, and specificity for each method. The two-stage method achieves a Youden index of 0.69 on the training set—20% higher than the 0.58 achieved by the simultaneous optimization method. Both methods yield similar Youden indices on the testing set, indicating the stability of the two-stage approach and the absence of overfitting.

6 | DISCUSSION

In this article, we introduce a novel method for linearly combining multiple biomarkers to maximize the Youden index under both perfect and imperfect reference standard scenarios. Our approach operates in two stages: first optimizing an empirical AUC approximation to determine optimal linear coefficients, and then maximizing the empirical Youden index to find the optimal disease classification cutoff. Under the semiparametric single index model and certain regularity conditions, our method provides consistent estimators for the linear coefficients, cutoff point, and Youden index. Furthermore, we propose a confidence interval for the Youden index using the AC estimator and Wilson score method, demonstrating robust coverage rates and computational efficiency.

Additionally, we used simulations to compare the performance of our newly proposed two-stage method with the existing simultaneous optimization method. Simulations were conducted under scenarios where the single-index model holds and where it does not. Furthermore, we evaluated the robustness of our method by performing simulations with all biomarkers set as binary. Such binary biomarker settings are common in diagnostic and prognostic assays, particularly in traditional Chinese medicine. Through simulations and a real data example, we demonstrated that our proposed method outperforms the existing approach in nearly all scenarios considered. Notably, when all biomarkers are binary, our method also performs well, with the estimated Youden index exhibiting strong performance on both training and testing datasets. Extensive simulation studies further validate the performance of our method and assess its effectiveness across various conditions.

However, we observe that the performance of the Youden index in the testing set consistently lags behind that in the training set for both methods. Improving generalization remains an open research challenge. Additionally, the assumption of a single index model may not universally hold. In clinical settings, it may not be practical to combine all biomarkers available, especially when some provide little information. Including such biomarkers could even degrade the performance of the combination score's Youden index. Hence, our future research will focus on relaxing the single index model assumption and devising methods to select an optimal subset of biomarkers for efficient and effective linear combination in practical applications.

References

1. Schindler SE, Bateman RJ. Combining blood-based biomarkers to predict risk for Alzheimer's disease dementia. *Nature Aging* 2021; 1(1): 26–28.

- 2. Pepe MS, Cai T, Longton G. Combining predictors for classification using the area under the receiver operating characteristic curve. *Biometrics* 2006; 62(1): 221–229.
- 3. Zhou XH, Obuchowski NA, McClish DK. Statistical methods in diagnostic medicine. 712. John Wiley & Sons. 2011.
- 4. Su JQ, Liu JS. Linear combinations of multiple diagnostic markers. *Journal of the American Statistical Association* 1993; 88(424): 1350–1355.
- 5. Reiser B, Faraggi D. Confidence intervals for the generalized ROC criterion. *Biometrics* 1997: 644–652.
- 6. Liu A, Schisterman EF, Zhu Y. On linear combinations of biomarkers to improve diagnostic accuracy. *Statistics in medicine* 2005; 24(1): 37–47.
- 7. Liu C, Liu A, Halabi S. A min-max combination of biomarkers to improve diagnostic accuracy. *Statistics in medicine* 2011; 30(16): 2005–2014.
- 8. Kang L, Xiong C, Crane P, Tian L. Linear combinations of biomarkers to improve diagnostic accuracy with three ordinal diagnostic categories. *Statistics in medicine* 2013; 32(4): 631–643.
- 9. Ma S, Huang J. Combining multiple markers for classification using ROC. *Biometrics* 2007; 63(3): 751–757.
- 10. Huang X, Oin G, Fang Y. Optimal combinations of diagnostic tests based on AUC. Biometrics 2011; 67(2): 568–576.
- 11. Gao F, Xiong C, Yan Y, Yu K, Zhang Z. Estimating optimum linear combination of multiple correlated diagnostic tests at a fixed specificity with receiver operating characteristic curves. *Journal of Data Science* 2008; 6(1): 105–123.
- 12. Meisner A, Carone M, Pepe MS, Kerr KF. Combining biomarkers by maximizing the true positive rate for a fixed false positive rate. *Biometrical Journal* 2021; 63(6): 1223–1240.
- 13. Ma H, Halabi S, Liu A, others . On the use of min-max combination of biomarkers to maximize the partial area under the ROC curve. *Journal of probability and statistics* 2019; 2019.
- 14. Zhang Z, Lu Y, Tian L. On feature ensemble optimizing the sensitivity and partial ROC curve. *Statistica Sinica* 2019; 29(3): 1395–1418.
- 15. Yan Q, Bantis LE, Stanford JL, Feng Z. Combining multiple biomarkers linearly to maximize the partial area under the ROC curve. *Statistics in medicine* 2018; 37(4): 627–642.
- 16. Youden WJ. Index for rating diagnostic tests. Cancer 1950; 3(1): 32–35.
- 17. Perkins NJ, Schisterman EF. The inconsistency of "optimal" cutpoints obtained using two criteria based on the receiver operating characteristic curve. *American journal of epidemiology* 2006; 163(7): 670–675.
- 18. Yin J, Tian L. Optimal linear combinations of multiple diagnostic biomarkers based on Youden index. *Statistics in medicine* 2014; 33(8): 1426–1440.
- 19. Aznar-Gimeno R, Esteban LM, Hoyo-Alonso dR, Borque-Fernando Á, Sanz G. A Stepwise Algorithm for Linearly Combining Biomarkers under Youden Index Maximization. *Mathematics* 2022; 10(8): 1221.
- 20. Aznar-Gimeno R, Esteban LM, Sanz G, Hoyo-Alonso dR. Comparing the Min–Max–Median/IQR Approach with the Min–Max Approach, Logistic Regression and XGBoost, Maximising the Youden Index. *Symmetry* 2023; 15(3): 756.
- 21. Lu Y, Dendukuri N, Schiller I, Joseph L. A Bayesian approach to simultaneously adjusting for verification and reference standard bias in diagnostic test studies. *Statistics in medicine* 2010; 29(24): 2532–2543.

22. Yu B, Zhou C, Bandinelli S. Combining multiple continuous tests for the diagnosis of kidney impairment in the absence of a gold standard. *Statistics in medicine* 2011; 30(14): 1712–1721.

- 23. García Barrado L, Coart E, Burzykowski T, Initiative ADN. Development of a diagnostic test based on multiple continuous biomarkers with an imperfect reference test. *Statistics in medicine* 2016; 35(4): 595–608.
- 24. García Barrado L, Coart E, Burzykowski T. Estimation of diagnostic accuracy of a combination of continuous biomarkers allowing for conditional dependence between the biomarkers and the imperfect reference-test. *Biometrics* 2017; 73(2): 646–655.
- Neyman J, Pearson ES. IX. On the problem of the most efficient tests of statistical hypotheses. *Philosophical Transactions of the Royal Society of London. Series A, Containing Papers of a Mathematical or Physical Character* 1933; 231(694-706): 289–337.
- 26. Schisterman EF, Perkins NJ, Liu A, Bondell H. Optimal cut-point and its corresponding Youden Index to discriminate individuals using pooled blood samples. *Epidemiology* 2005; 16(1): 73–81.
- 27. Wright SJ. Numerical optimization. 2006.
- 28. Lin H, Zhou L, Peng H, Zhou XH. Selection and combination of biomarkers using ROC method for disease classification and prediction. *Canadian Journal of Statistics* 2011; 39(2): 324–343.
- 29. Vexler A, Liu A, Schisterman EF, Wu C. Note on distribution-free estimation of maximum linear separation of two multivariate distributions. *Nonparametric Statistics* 2006; 18(2): 145–158.
- 30. Shan G. Improved confidence intervals for the Youden index. *PloS one* 2015; 10(7): e0127272.
- 31. Agresti A, Coull BA. Approximate is better than "exact" for interval estimation of binomial proportions. *The American Statistician* 1998; 52(2): 119–126.
- 32. Zhou H, Qin G. New nonparametric confidence intervals for the Youden index. *Journal of biopharmaceutical statistics* 2012; 22(6): 1244–1257.
- 33. Dawid AP, Skene AM. Maximum likelihood estimation of observer error-rates using the EM algorithm. *Journal of the Royal Statistical Society: Series C (Applied Statistics)* 1979; 28(1): 20–28.
- 34. Walter SD, Irwig LM. Estimation of test error rates, disease prevalence and relative risk from misclassified data: a review. *Journal of clinical epidemiology* 1988; 41(9): 923–937.
- Gravallese EM, Firestein GS. Rheumatoid arthritis—common origins, divergent mechanisms. New England Journal of Medicine 2023; 388(6): 529–542.
- 36. Turk MA, Liu Y, Pope JE. Non-pharmacological interventions in the treatment of rheumatoid arthritis: a systematic review and meta-analysis. *Autoimmunity reviews* 2023; 22(6): 103323.
- 37. Smolen JS, Landewé RB, Bergstra SA, et al. EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2022 update. *Annals of the rheumatic diseases* 2023; 82(1): 3–18.
- 38. Nagy G, Roodenrijs NM, Welsing PM, et al. EULAR definition of difficult-to-treat rheumatoid arthritis. *Annals of the rheumatic diseases* 2021; 80(1): 31–35.
- Xing Q, Fu L, Yu Z, Zhou X. Efficacy and safety of integrated traditional Chinese medicine and Western medicine on the treatment of rheumatoid arthritis: a meta-analysis. *Evidence-Based Complementary and Alternative Medicine* 2020; 2020(1): 4348709.
- 40. Zhang GG, Singh B, Lee W, Handwerger B, Lao L, Berman B. Improvement of agreement in TCM diagnosis among TCM practitioners for persons with the conventional diagnosis of rheumatoid arthritis: effect of training. *The Journal of Alternative and Complementary Medicine* 2008; 14(4): 381–386.

41. Fang G, Zhang Ll, Ren Q, et al. Development of a Diagnostic Questionnaire for Damp Phlegm Pattern and Blood Stasis Pattern in Coronary Heart Disease Patients (CHD-DPBSPQ). *Evidence-Based Complementary and Alternative Medicine* 2019; 2019(1): 6856085.

- 42. Leung AYL, Zhang J, Chan CY, et al. Validation of evidence-based questionnaire for TCM syndrome differentiation of heart failure and evaluation of expert consensus. *Chinese Medicine* 2023; 18(1): 70.
- 43. Chen XI, Wen Y, Wu Zc, et al. Development of a Traditional Chinese Medicine Syndrome-Specific Scale for Ulcerative Colitis: The Large Intestine Dampness-Heat Syndrome Questionnaire. *Evidence-Based Complementary and Alternative Medicine* 2018; 2018(1): 4039019.
- 44. Wang J, Gong X, Tang X, Liu H, Liu J, He D. Multi-center cross-sectional survey of characteristics of Chinese medicine syndromes in 1602 rheumatoid arthritis patients. *Journal of Traditional Chinese Medicine* 2018; 59(11): 963-967.
- 45. Li S, Liu D, Chen Z, et al. Comparative efficacy and safety of four classical prescriptions for clearing damp-heat recommended by clinical guidelines in treating rheumatoid arthritis: a network meta-analysis. *Annals of Palliative Medicine* 2021; 10(7): 7298328–7297328.
- 46. Graham B, Regehr G, Wright JG. Delphi as a method to establish consensus for diagnostic criteria. *Journal of clinical epidemiology* 2003; 56(12): 1150–1156.
- 47. Jiang Q, Wang H, Gong X, Luo C. Guidelines of diagnosis and treatment of rheumatoid arthritis disease and syndrome combination. *J Tradit Chin Med* 2018; 59(20): 1794–800.
- 48. Gong X, Liu Wx, Li D, et al. China rheumatoid arthritis registry of patients with Chinese medicine (CERTAIN): Rationale, design, and baseline characteristics of the first 11,764 enrollees. *Phytomedicine* 2022; 104: 154236.
- 49. Wang XN, Zhou V, Liu Q, Gao Y, Zhou XH. Evaluation of the accuracy of diagnostic scales for a syndrome in Chinese medicine in the absence of a gold standard. *Chinese Medicine* 2016; 11: 1–7.

APPENDIX

A PROOF OF THEOREM 1

In our article, the true optimal cutoff value c_0 is defined as:

$$c_0 = \arg \max_{c} \left\{ F_{0\omega_0}(c) - F_{1\omega_0}(c) \right\},$$
 (A1)

and the optimal Youden index is defined as:

$$J_{\omega_0} = \max \left\{ Se_{\omega_0}(c) + Sp_{\omega_0}(c) - 1 \right\} = F_{0\omega_0}(c_0) - F_{1\omega_0}(c_0). \tag{A2}$$

Define the empirical distribution functions as follows:

$$\widehat{F}_{0\widehat{\omega}}(c) = \frac{1}{n_0} \sum_{j=1}^{n_0} \mathbb{I}\left(\widehat{\omega}' T_j^0 \leq c\right),$$

$$\widehat{F}_{0\omega_0}(c) = \frac{1}{n_0} \sum_{j=1}^{n_0} \mathbb{I}\left(\omega_0' T_j^0 \leq c\right),$$

$$F_{0\omega_0}(c) = \mathbb{P}\left(\omega_0' T^0 \le c\right).$$

Similarly, define $\hat{F}_{1\hat{\omega}}(c)$, $\hat{F}_{1\omega_0}(c)$, and $F_{1\omega_0}(c)$ analogously. Our estimator is defined as:

$$\widehat{c}^{\mathrm{TS}} = \operatorname{median} \left\{ c : \max_{c} \widehat{F}_{0\widehat{\omega}}(c) - \widehat{F}_{1\widehat{\omega}}(c) \right\}, \tag{A3}$$

$$\hat{J}^{TS} = \hat{F}_{0\hat{\omega}}(\hat{c}^{TS}) - \hat{F}_{1\hat{\omega}}(\hat{c}^{TS}). \tag{A4}$$

Alternatively, the maximum or minimum value instead of the median can be used in (A3).

First, we aim to prove that for any given c, $\widehat{F}_{0\widehat{\omega}}(c) \stackrel{P}{\to} F_{0\omega_0}(c)$ and $\widehat{F}_{1\widehat{\omega}}(c) \stackrel{P}{\to} F_{1\omega_0}(c)$ as $n \to \infty$. Using the triangle inequality:

$$\left| \hat{F}_{0\hat{\omega}}(c) - F_{0\omega_0}(c) \right| \le \left| \hat{F}_{0\hat{\omega}}(c) - \hat{F}_{0\omega_0}(c) \right| + \left| \hat{F}_{0\omega_0}(c) - F_{0\omega_0}(c) \right|. \tag{A5}$$

According to Glivenko-Cantelli theorem,

$$\sup_{c} \left| \widehat{F}_{0\omega_0}(c) - F_{0\omega_0}(c) \right| \stackrel{a.s.}{\to} 0.$$

Next we will prove that the first term on the right-hand side of inequality (A5) also converges to 0 in probability.

Since $\widehat{\omega} \stackrel{P}{\to} \omega_0$, for any $\epsilon > 0$, for any $\delta > 0$, there exists N, such that when n > N,

$$\mathbb{P}\left(\|\widehat{\omega} - \omega_0\| < \epsilon\right) > 1 - \delta.$$

Since ω is a finite-dimensional vector, we have

$$|\widehat{\omega}'T - \omega_0'T| \le \|\widehat{\omega} - \omega_0\| \|T\|.$$

For any $\epsilon > 0$, define:

$$A_{\epsilon} = \left\{ T : |\widehat{\omega}'T - \omega_0'T| < \epsilon \right\},\,$$

and we have $\mathbb{P}(A_{\epsilon}) \to 1$. That is, for any given c, for any $\delta > 0$, $\epsilon > 0$, there exists N, such that when n > N,

$$\mathbb{P}\left(\left|\mathbb{I}(\widehat{\omega}'T\leq c)-\mathbb{I}(\omega_0'T\leq c)\right|>\epsilon\right)<\delta. \tag{A6}$$

Then we have $\left| \widehat{F}_{0\hat{\omega}}(c) - \widehat{F}_{0\omega_0}(c) \right| \stackrel{P}{\to} 0$ as $n \to \infty$.

Similarly, we can prove that $\left| \hat{F}_{1\hat{\omega}}(c) - \hat{F}_{1\omega_0}(c) \right| \stackrel{P}{\to} 0$ as $n \to \infty$.

Using condition (B), let $e' = \frac{\epsilon}{5}$. From the convergence of $\widehat{F}_{0\widehat{\omega}}$ and $\widehat{F}_{1\widehat{\omega}}$, for large enough n_0 and n_1 , we have

$$F_{0\omega_0}(c) - F_{1\omega_0}(c) - 2\epsilon' < \widehat{F}_{0\widehat{\omega}}(c) - \widehat{F}_{1\widehat{\omega}}(c) < F_{0\omega_0}(c) - F_{1\omega_0}(c) + 2\epsilon' \quad \text{for all } c.$$

Hence

$$\begin{split} \sup_{|c-c_0|>\delta} \left[\widehat{F}_{0\widehat{\omega}}(c) - \widehat{F}_{1\widehat{\omega}}(c) \right] &\leq 2\epsilon' + \sup_{|c-c_0|>\delta} \left(F_{0\omega_0}(c) - F_{1\omega_0}(c) \right) \\ &< F_{0\omega_0}(c_0) - F_{1\omega_0}(c_0) - 3\epsilon' \quad \text{by condition } (B) \end{split}$$

Because

$$\widehat{F}_{0\widehat{\omega}}(c_0) - \widehat{F}_{1\widehat{\omega}}(c_0) \overset{P}{\to} F_{0\omega_0}(c_0) - F_{1\omega_0}(c_0),$$

for large enough n_0 and n_1 , we have:

$$\widehat{F}_{0\widehat{\omega}}(c_0) - \widehat{F}_{1\widehat{\omega}}(c_0) > F_{0\omega_0}(c_0) - F_{1\omega_0}(c_0) - \epsilon'.$$

Thus,

$$\sup_{|c-c_0|>\delta} \left[\widehat{F}_{0\widehat{\omega}}(c) - \widehat{F}_{1\widehat{\omega}}(c) \right] < \widehat{F}_{0\widehat{\omega}}(c_0) - \widehat{F}_{1\widehat{\omega}}(c_0).$$

Therefore,

$$\sup_{|c-c_0|<\delta} \left[\widehat{F}_{0\widehat{\omega}}(c) - \widehat{F}_{1\widehat{\omega}}(c) \right] = \sup_{c} \left(\widehat{F}_{0\widehat{\omega}}(c) - \widehat{F}_{1\widehat{\omega}}(c) \right)$$

This implies that $\hat{c}^{TS} \stackrel{P}{\rightarrow} c_0$.

Note on median use: In the definition of \hat{c}^{TS} , the median is used to handle potential ties and to provide a robust estimate of the cutoff point. However, in the proof of consistency, the focus is on showing that the entire distribution functions $\hat{F}_{0\hat{\omega}}$ and $\hat{F}_{1\hat{\omega}}$ converge to their true counterparts. The convergence of the distribution functions ensures that the median (or any other robust central tendency measure such as maximum or minimum) of the cutoff points will converge to the true cutoff c_0 . Therefore, the specific use of the median does not affect the overall consistency proof.

Next, we will prove the consistency of \hat{J}^{TS} . Since the convergence of $\hat{F}_{0\hat{\omega}}(c)$ and $\hat{F}_{1\hat{\omega}}(c)$ to $F_{0\omega_0}(c)$ and $F_{1\omega_0}(c)$ hold uniformly over c, we can write:

$$\sup_{c} \left| \widehat{F}_{0\widehat{\omega}}(c) - F_{0\omega_0}(c) \right| \stackrel{P}{\to} 0,$$

$$\sup_{c} \left| \widehat{F}_{1\widehat{\omega}}(c) - F_{1\omega_0}(c) \right| \stackrel{P}{\to} 0.$$

Using the triangle inequality, we have:

$$\left|\left(\widehat{F}_{0\widehat{\omega}}(c)-\widehat{F}_{1\widehat{\omega}}(c)\right)-\left(F_{0\omega_0}(c)-F_{1\omega_0}(c)\right)\right|\leq \left|\left(\widehat{F}_{0\widehat{\omega}}(c)-\widehat{F}_{1\widehat{\omega}}(c)\right)\right|+\left|\left(F_{0\omega_0}(c)-F_{1\omega_0}(c)\right)\right|.$$

Since both the terms on the right-hand side converge uniformly to 0, we get:

$$\sup_{c} \left| \left(\widehat{F}_{0\widehat{\omega}}(c) - \widehat{F}_{1\widehat{\omega}}(c) \right) - \left(F_{0\omega_0}(c) - F_{1\omega_0}(c) \right) \right| \stackrel{P}{\to} 0.$$

Since c_0 maximizes $F_{0\omega_0}(c) - F_{1\omega_0}(c)$,

$$\left(\widehat{F}_{0\widehat{\omega}}(c_0) - \widehat{F}_{1\widehat{\omega}}(c_0)\right) \overset{P}{\to} \left(F_{0\omega_0}(c_0) - F_{1\omega_0}(c_0)\right).$$

Hence,

$$\sup_{c} \left(\widehat{F}_{0\widehat{\omega}}(c) - \widehat{F}_{1\widehat{\omega}}(c) \right) \stackrel{P}{\to} \sup_{c} \left(F_{0\omega_{0}}(c) - F_{1\omega_{0}}(c) \right).$$

Thus,

$$\left(\widehat{F}_{0\widehat{\omega}}(\widehat{c}^{\mathrm{TS}}) - \widehat{F}_{1\widehat{\omega}}(\widehat{c}^{\mathrm{TS}})\right) \overset{P}{\to} \left(F_{0\omega_0}(c_0) - F_{1\omega_0}(c_0)\right).$$

Therefore, $\widehat{J}^{TS} \stackrel{P}{\rightarrow} J_{\omega_0}$.

B DERIVATION OF RELATIONSHIP BETWEEN THE TRUE INDEX AND THE PROXY INDEX

The relationship of $\widetilde{AUC}(\omega)$ and $AUC(\omega)$ is derived as follows:

$$\begin{split} \widetilde{\text{AUC}}(\omega) &= \mathbb{P}(\omega'T_i > \omega'T_j | R_i = 1, R_j = 0) \\ &= \mathbb{P}(\omega'T_i > \omega'T_j | R_i = 1, R_j = 0, D_i = 1, D_j = 0) \mathbb{P}(D_i = 1, D_j = 0 | R_i = 1, R_j = 0) \\ &+ \mathbb{P}(\omega'T_i > \omega'T_j | R_i = 1, R_j = 0, D_i = 0, D_j = 1) \mathbb{P}(D_i = 0, D_j = 1 | R_i = 1, R_j = 0) \\ &+ \mathbb{P}(\omega'T_i > \omega'T_j | R_i = 1, R_j = 0, D_i = 1, D_j = 1) \mathbb{P}(D_i = 1, D_j = 1 | R_i = 1, R_j = 0) \\ &+ \mathbb{P}(\omega'T_i > \omega'T_j | R_i = 1, R_j = 0, D_i = 0, D_j = 0) \mathbb{P}(D_i = 0, D_j = 0 | R_i = 1, R_j = 0) \\ &= ppv \cdot npv \cdot \text{AUC}(\omega) + (1 - ppv)(1 - npv)(1 - \text{AUC}(\omega)) \\ &+ \frac{1}{2} \cdot ppv \cdot (1 - npv) + \frac{1}{2} \cdot (1 - ppv) \cdot npv \\ &= (ppv + npv - 1) \text{AUC}(\omega) - \frac{1}{2}(ppv + npv) + 1. \end{split}$$

To derive the relationship between \widetilde{J}_{ω} and J_{ω} , we first define $\widetilde{J}_{\omega}(c)$ as $\mathbb{P}(\omega'T > c | R = 1) + \mathbb{P}(\omega'T < c | R = 0) - 1$ for a given c, and define $J_{\omega}(c)$ as $\mathbb{P}(\omega'T > c | D = 1) + \mathbb{P}(\omega'T < c | D = 0) - 1$. The relationship between $\widetilde{J}_{\omega}(c)$ and $J_{\omega}(c)$ is as follows:

$$\begin{split} \widetilde{J}_{\omega}(c) &= \mathbb{P}(\omega'T > c | R = 1) + \mathbb{P}(\omega'T < c | R = 0) - 1 \\ &= \mathbb{P}(\omega'T > c | R = 1) + \mathbb{P}(\omega'T < c | R = 0) - 1 \\ &= \mathbb{P}(\omega'T > c | R = 1) + \mathbb{P}(\omega'T < c | R = 0) - 1 \\ &= \mathbb{P}(\omega'T > c | R = 1, D = 1) \mathbb{P}(D = 1 | R = 1) + \mathbb{P}(\omega'T > c | R = 1, D = 0) \mathbb{P}(D = 0 | R = 1) \\ &+ \mathbb{P}(\omega'T < c | R = 0, D = 1) \mathbb{P}(D = 1 | R = 0) + \mathbb{P}(\omega'T < c | R = 0, D = 0) \mathbb{P}(D = 0 | R = 0) - 1 \\ &= (ppv + npv - 1) \left(\mathbb{P}(\omega'T > c | D = 1) + \mathbb{P}(\omega'T < c | D = 0) - 1 \right) \\ &= (ppv + npv - 1) J_{\omega}(c). \end{split}$$

We can see that $\widetilde{J}_{\omega}(c)$ is an increasing function of $J_{\omega}(c)$. Therefore, we have

$$\arg\max_{c} \widetilde{J}_{\omega}(c) = \arg\max_{c} J_{\omega}(c) \triangleq c_{\omega}$$
 (B9)

and

$$\widetilde{J}_{\omega}(c_{\omega}) = \widetilde{J}_{\omega} = (ppv + npv - 1)J_{\omega} = (ppv + npv - 1)J_{\omega}(c_{\omega}). \tag{B10}$$

TABLE 1 Coverage rates (CR), average lengths (AL), lower limit (LL), upper limit (UL) of our newly proposed 95% confidence intervals for the Youden index, as well as the coverage rates (CR_NP) of the 95% confidence intervals using the NP method based on 1000 replications.

Youden index value	n ₁	n_0	CR_NP	CR	AL	LL	UL
$\overline{J_0 = 0.45}$	50	50	0.854	0.944	0.3225	0.3257	0.6482
	100	100	0.890	0.948	0.2352	0.3586	0.5938
	200	200	0.902	0.930	0.1692	0.3837	0.5530
	75	25	0.851	0.963	0.3643	0.3032	0.6674
	150	50	0.885	0.937	0.2662	0.3446	0.6108
	300	100	0.898	0.936	0.1933	0.3772	0.5705
	25	75	0.848	0.965	0.3616	0.3054	0.6671
	50	150	0.890	0.961	0.2649	0.3431	0.6080
	100	300	0.903	0.951	0.1922	0.3763	0.5685
$J_0 = 0.60$	50	50	0.888	0.989	0.2941	0.4583	0.7524
	100	100	0.896	0.972	0.2119	0.5052	0.7171
	200	200	0.918	0.955	0.1521	0.5350	0.6871
	75	25	0.886	0.981	0.3359	0.4318	0.7677
	150	50	0.905	0.977	0.2425	0.4870	0.7295
	300	100	0.902	0.955	0.1734	0.5273	0.7007
	25	75	0.894	0.984	0.3326	0.4339	0.7665
	50	150	0.885	0.968	0.2404	0.4871	0.7275
	100	300	0.912	0.950	0.1730	0.5250	0.6979
$J_0 = 0.70$	50	50	0.929	0.978	0.2694	0.5486	0.8179
·	100	100	0.906	0.970	0.1920	0.6002	0.7921
	200	200	0.914	0.962	0.1362	0.6369	0.7731
	75	25	0.927	0.992	0.3116	0.5181	0.8297
	150	50	0.920	0.970	0.2214	0.5806	0.8020
	300	100	0.924	0.967	0.1568	0.6240	0.7808
	25	75	0.937	0.984	0.3096	0.5153	0.8249
	50	150	0.914	0.972	0.2182	0.5840	0.8021
	100	300	0.922	0.969	0.1564	0.6230	0.7793
$J_0 = 0.85$	50	50	0.979	0.986	0.2191	0.6905	0.9096
	100	100	0.952	0.978	0.1492	0.7531	0.9022
	200	200	0.955	0.976	0.1037	0.7885	0.8922
	75	25	1.000	0.963	0.2664	0.6453	0.9117
	150	50	0.964	0.969	0.1778	0.7270	0.9048
	300	100	0.947	0.972	0.1208	0.7761	0.8969
	25	75	0.999	0.963	0.2645	0.6464	0.9109
	50	150	0.963	0.968	0.1757	0.7290	0.9048
	100	300	0.954	0.962	0.1201	0.7745	0.8947

Sun ET AL 17

TABLE 2 The performance of our newly-proposed two-stage method (TSM) and existing simultaneous optimization method (SIM) when the reference standard is perfect and imperfect

1				Se=Sp=1		Se=Sp=0	.95	Se=Sp=0	.90	Se=Sp=0.85	
J_0	π	n	Method	Train	Test	Train	Test	Train	Test	Train	Test
0.45	0.50	200	TSM	0.5527	0.3901	0.5280	0.3753	0.5043	0.3602	0.4692	0.3340
			SIM	0.4928	0.2872	0.4656	0.2744	0.4337	0.2639	0.4033	0.2440
		400	TSM	0.5125	0.4172	0.5010	0.4104	0.4881	0.4034	0.4696	0.3884
			SIM	0.4683	0.3233	0.4419	0.3096	0.4226	0.2981	0.3955	0.2841
		800	TSM	0.4842	0.4270	0.4776	0.4249	0.4701	0.4197	0.4608	0.4132
			SIM	0.4416	0.3491	0.4310	0.3457	0.4144	0.3325	0.3959	0.3243
	0.75	200	TSM	0.5695	0.3817	0.5374	0.3612	0.4997	0.3403	0.4564	0.3142
			SIM	0.5087	0.2702	0.4734	0.2554	0.4390	0.2487	0.3993	0.2242
		400	TSM	0.5274	0.4084	0.5093	0.4007	0.4855	0.3817	0.4571	0.3616
			SIM	0.4782	0.3047	0.4530	0.2953	0.4237	0.2813	0.3899	0.2644
		800	TSM	0.4951	0.4234	0.4852	0.4166	0.4696	0.4092	0.4553	0.3988
			SIM	0.4520	0.3367	0.4356	0.3280	0.4135	0.3165	0.3906	0.3024
	0.25	200	TSM	0.5811	0.3754	0.5420	0.3539	0.5019	0.3307	0.4556	0.3047
			SIM	0.5148	0.2702	0.4720	0.2513	0.4319	0.2336	0.3850	0.2115
		400	TSM	0.5235	0.4084	0.5056	0.3984	0.4856	0.3868	0.4620	0.3716
			SIM	0.4720	0.3144	0.4439	0.2993	0.4125	0.2848	0.3814	0.2663
		800	TSM	0.4905	0.4239	0.4805	0.4198	0.4675	0.4094	0.4551	0.3996
			SIM	0.4487	0.3434	0.4273	0.3318	0.4060	0.3191	0.3784	0.2978
0.70	0.50	200	TSM	0.7761	0.6603	0.7543	0.7292	0.6468	0.6295	0.6986	0.6038
			SIM	0.6917	0.5239	0.6593	0.5000	0.6137	0.4744	0.5724	0.4400
		400	TSM	0.7430	0.6792	0.7330	0.6721	0.7179	0.6618	0.7028	0.6513
			SIM	0.6922	0.5843	0.6657	0.5665	0.6398	0.5453	0.6075	0.5228
		800	TSM	0.7259	0.6905	0.7204	0.6861	0.7128	0.6818	0.7044	0.6757
			SIM	0.7046	0.6415	0.6890	0.6293	0.6681	0.6137	0.6470	0.5944
	0.75	200	TSM	0.7855	0.6491	0.7515	0.6294	0.7180	0.6131	0.6721	0.5683
			SIM	0.6909	0.4997	0.6537	0.4651	0.6167	0.4495	0.5712	0.4184
		400	TSM	0.7541	0.6684	0.7343	0.6557	0.7074	0.6393	0.6793	0.6187
			SIM	0.6899	0.5476	0.6600	0.5311	0.6239	0.5011	0.5823	0.4711
		800	TSM	0.7335	0.6855	0.7227	0.6772	0.7091	0.6687	0.6912	0.6539
			SIM	0.7017	0.6155	0.6784	0.5991	0.6518	0.5773	0.6172	0.5482
	0.25	200	TSM	0.7906	0.6409	0.7430	0.6201	0.7127	0.5906	0.6602	0.5530
			SIM	0.6975	0.4974	0.6370	0.4617	0.5852	0.4264	0.5363	0.3921
		400	TSM	0.7558	0.6740	0.7370	0.6609	0.7152	0.6485	0.6912	0.6300
			SIM	0.7041	0.5830	0.6636	0.5523	0.6163	0.5194	0.5783	0.4867
		800	TSM	0.7334	0.6844	0.7239	0.6786	0.7121	0.6680	0.6954	0.6562
			SIM	0.7062	0.6266	0.6753	0.6014	0.6440	0.5750	0.6080	0.5471

TABLE 3 The performance of our newly-proposed two-stage method (TSM) and existing simultaneous optimization method (SIM) with samples from multivariate normal distributions with equal covariance matrices

		Estimated	Youden in	dex		Variance of Youden index				
		Training s	et	Testing se	t	Training s	set	Testing set	t	
Prevalence	Sample size	TSM	SIM	TSM	SIM	TSM	SIM	TSM	SIM	
		$\Sigma_1 = \Sigma_1$	$\Sigma_0 = 0.3 * .$	$I_5 + 0.7 * J_5$	(large cor	relation)				
$\pi = 0.5$	n = 200	0.7696	0.7443	0.6570	0.6141	0.0041	0.0048	0.0064	0.0081	
	n = 400	0.7402	0.7303	0.6744	0.6465	0.0019	0.0024	0.0029	0.0039	
	n = 800	0.7249	0.7194	0.6862	0.6623	0.0011	0.0014	0.0014	0.0019	
$\pi = 0.75$	n = 200	0.7893	0.7561	0.6506	0.5944	0.0052	0.0072	0.0093	0.0124	
	n = 400	0.7513	0.7370	0.6718	0.6364	0.0025	0.0031	0.0044	0.0062	
	n = 800	0.7291	0.7225	0.6819	0.6518	0.0014	0.0017	0.0019	0.0026	
$\pi = 0.25$	n = 200	0.7850	0.7617	0.6467	0.6152	0.0045	0.0052	0.0100	0.0117	
	n = 400	0.7562	0.7442	0.6683	0.6373	0.0027	0.0030	0.0041	0.0050	
	n = 800	0.7330	0.7255	0.6826	0.6552	0.0015	0.0017	0.0019	0.0025	
		$\Sigma_1 = \Sigma_0$	$= 0.5 * I_{s}$	$s_5 + 0.5 * J_5$	(medium co	rrelation)				
$\pi = 0.5$	n = 200	0.7279	0.7003	0.6009	0.5485	0.0043	0.0053	0.0072	0.0096	
	n = 400	0.6939	0.6795	0.6214	0.5819	0.0024	0.0032	0.0035	0.0052	
	n = 800	0.6743	0.6692	0.6353	0.6049	0.0013	0.0019	0.0017	0.0026	
$\pi = 0.75$	n = 200	0.7485	0.7070	0.5951	0.5283	0.0060	0.0070	0.0097	0.0140	
	n = 400	0.7034	0.6860	0.6160	0.5666	0.0030	0.0040	0.0046	0.0074	
	n = 800	0.6813	0.6726	0.6297	0.5916	0.0016	0.0025	0.0024	0.0036	
$\pi = 0.25$	n = 200	0.7434	0.7155	0.5918	0.5432	0.0052	0.0058	0.0106	0.0127	
	n = 400	0.7055	0.6900	0.6163	0.5713	0.0029	0.0040	0.0047	0.0062	
	n = 800	0.6846	0.6785	0.6289	0.5990	0.0017	0.0024	0.0021	0.0033	
		$\Sigma_1 =$	$\Sigma_0 = 0.7 *$	$I_5 + 0.3 * J$	(low corr	elation)				
$\pi = 0.5$	n = 200	0.7342	0.6946	0.6097	0.5340	0.0043	0.0060	0.0071	0.0109	
	n = 400	0.7008	0.6785	0.6332	0.5755	0.0022	0.0041	0.0035	0.0073	
	n = 800	0.6812	0.6714	0.6402	0.6052	0.0012	0.0023	0.0017	0.0036	
$\pi = 0.75$	n = 200	0.7544	0.7045	0.5972	0.5050	0.0054	0.0073	0.0090	0.0145	
	n = 400	0.7094	0.6795	0.6226	0.5560	0.0027	0.0046	0.0044	0.0081	
	n = 800	0.6855	0.6698	0.6368	0.5896	0.0016	0.0033	0.0022	0.0047	
$\pi = 0.25$	n = 200	0.7425	0.7015	0.6043	0.5295	0.0049	0.0071	0.0106	0.0142	
	n = 400	0.7136	0.6914	0.6250	0.5689	0.0029	0.0047	0.0040	0.0079	
	n = 800	0.6899	0.6778	0.6375	0.5980	0.0017	0.0031	0.0020	0.0043	

TABLE 4 The performance of our newly-proposed two-stage method (TSM) and existing simultaneous optimization method (SIM) with samples from multivariate normal distributions with unequal covariance matrices

		Estimated	Youden in	dex		Variance of Youden index				
Prevalence		Training set		Testing set		Training set		Testing set		
	Sample size	TSM	SIM	TSM	SIM	TSM	SIM	TSM	SIM	
$\pi = 0.5$	n = 200	0.7302	0.7042	0.6043	0.5584	0.0045	0.0052	0.0069	0.0089	
	n = 400	0.6968	0.6842	0.6191	0.5855	0.0022	0.0027	0.0034	0.0047	
	n = 800	0.6744	0.6694	0.6341	0.6048	0.0016	0.0011	0.0016	0.0024	
$\pi = 0.75$	n = 200	0.7507	0.7192	0.5971	0.5322	0.0056	0.0058	0.0093	0.0129	
	n = 400	0.7064	0.6934	0.6160	0.5725	0.0029	0.0034	0.0050	0.0073	
	n = 800	0.6828	0.6761	0.6290	0.5944	0.0017	0.0019	0.0021	0.0033	
$\pi = 0.25$	n = 200	0.7390	0.7139	0.5998	0.5532	0.0049	0.0052	0.0101	0.0112	
	n = 400	0.7056	0.6937	0.6210	0.5820	0.0029	0.0035	0.0043	0.0052	
	n = 800	0.6815	0.6758	0.6320	0.6030	0.0015	0.0019	0.0022	0.0028	

TABLE 5 The performance of our newly-proposed two-stage method (TSM) and existing simultaneous optimization method (SIM) with all biomarkers being binary

		Estimated	Estimated Youden index				Variance of Youden index				
		Training set		Testing set		Training set		Testing set			
Prevalence	Sample size	TSM	SIM	TSM	SIM	TSM	SIM	TSM	SIM		
			$\omega = (-2.22)$	5, -0.80, 0.6	5, 2.10, 3.55)					
$\pi = 0.5$	n = 200	0.6475	0.6202	0.5495	0.5241	0.0046	0.0052	0.0086	0.0101		
	n = 400	0.6235	0.5983	0.5688	0.5479	0.0026	0.0028	0.0041	0.0044		
	n = 800	0.6149	0.5921	0.5818	0.5622	0.0014	0.0016	0.0019	0.0022		
			$\omega = (-1.0)$	00, 0.10, 1.20	, 2.30, 3.40)						
$\pi = 0.75$	n = 200	0.6031	0.5597	0.4913	0.4503	0.0072	0.0075	0.0126	0.0150		
	n = 400	0.5781	0.5390	0.5159	0.4806	0.0044	0.0046	0.0072	0.0071		
	n = 800	0.5675	0.5256	0.5318	0.4968	0.0028	0.0030	0.0038	0.0038		
			$\omega = (-2.40)$, -1.30, -0.2	20, 0.90, 2.00	0)					
$\pi = 0.25$	n = 200	0.6178	0.5853	0.4786	0.4501	0.0065	0.0070	0.0125	0.0142		
	n = 400	0.5828	0.5540	0.5011	0.4776	0.0036	0.0041	0.0058	0.0063		
	n = 800	0.5632	0.5371	0.5206	0.4978	0.0019	0.0022	0.0028	0.0032		

TABLE 6 Data for diagnosing dampness-heat obstruction syndrome.

	Estimated You	den index	Sensitivity		Specificity		
Method	Training	Testing	Training	Testing	Training	Testing	
TSM	0.6924	0.6928	0.7235	0.7158	0.9688	0.9769	
SIM	0.5783	0.5728	0.8333	0.8183	0.7450	0.7544	