# Natural Orbital Non-Orthogonal Configuration Interaction

Daniel Graf<sup>\*,†,‡</sup> and Alex J. W. Thom<sup>†</sup>

†Yusuf Hamied Department of Chemistry, University of Cambridge, Cambridge ‡Department of Chemistry, Ludwig-Maximilians-Universität München, Munich

E-mail:<daniel.graf@cup.uni-muenchen.de>

#### Abstract

Non-orthogonal configuration interaction (NOCI) is a generalization of the standard orthogonal configuration interaction (CI) method and offers a highly flexible framework for describing ground and excited electronic states. However, this flexibility also comes with challenges, as there is still no clear or generally accepted approach for constructing a compact and accurate state basis for NOCI. In this work, we take a step toward addressing this challenge by introducing a novel NOCI approach designed with three primary objectives: (1) ensuring the method is systematic, (2) achieving a compact NOCI expansion, and (3) treating all electronic states of interest on equal footing. The development of our approach is presented step by step, with each building block evaluated and validated through applications to simple model systems, demonstrating its effectiveness and potential.

#### Introduction

Excited states lie at the core of many fundamental processes in nature, such as photosynthe-sis,<sup>[1](#page-13-0)[,2](#page-13-1)</sup> human vision,<sup>[3,](#page-14-0)[4](#page-14-1)</sup> photoinduced damage of essential biomolecules,<sup>[5](#page-14-2)[,6](#page-14-3)</sup> and even theories concerning the origin of life<sup>[7](#page-14-4)[,8](#page-14-5)</sup> — a detailed understanding of these processes is therefore highly desirable. While spectroscopic methods provide valuable insights, they face limitations in both spatial and temporal resolution, highlighting the indispensable role of theoretical investigations. However, accurately describing such processes remains a formidable challenge. The primary difficulty arises from the need to treat multiple electronic states on the same footing, requiring a balanced treatment of both dynamical and static electron correlation, while at the same time ensuring a feasible computational cost.

Currently, time-dependent density functional theory<sup>[9](#page-14-6)</sup> (TDDFT) is the most widely used framework for excited-state calculations due to its excellent cost-to-performance ratio. While TDDFT is, in principle, an exact theory,  $10$  it depends on an unknown modification of the Hamiltonian — specifically, the exchange-correlation potential. Consequently, the accuracy of TDDFT is critically determined by the choice of exchange-correlation functional. However, given the vast array of available approximations,  $^{11}$  $^{11}$  $^{11}$  selecting an appropriate functional is often challenging. Additionally, in TDDFT, excitation energies are typically computed in the linear response regime, <sup>[12](#page-15-0)</sup> which results in an incapability to describe double excitations. The most significant limitation of TDDFT, however, is its inability to accurately describe static correlation effects in the electronic ground state.<sup>[13](#page-15-1)</sup> This deficiency leads to an imbalance in the representation of the system's electronic states, undermining its reliability for systems where static correlation plays a major role.

The most widely used method for addressing the static correlation problem in the ground state, thereby enabling a balanced description of different electronic states, is the complete active space self-consistent field (CASSCF) method<sup>[14](#page-15-2)</sup> — the multi-reference analogue of Hartree–Fock theory. While CASSCF is highly effective at capturing static correlation effects, its computational cost increases combinatorially with the active space size and, therefore, one is frequently restricted to small active spaces including only the very most important orbitals. The selection of these active orbitals is far from trivial and necessitates expert knowledge, preventing its use as a black-box method. Often it is not even possible to include all orbitals which should be included due to the enormous cost of the method, resulting in a (possibly) unsatisfactory description of the system at hand.

When considering the calculation of excited states, additional challenges arise regarding the orbital optimization: One approach is to target the excited state explicitly using a state-specific formalism. However, this method inherently biases the calculation towards the selected state and is prone to variational collapse onto the ground state.<sup>[15](#page-15-3)</sup> An alternative is the state-averaging formalism, which optimizes a weighted average of the energies of multiple states, allowing for simultaneous computation of several states. [16](#page-15-4) Unfortunately, the resulting wave functions are not true stationary points on the energy surface, which limits their accuracy compared to states obtained from state-specific procedures.

Motivated by the aforementioned challenges, we here propose a non-orthogonal configuration interaction (NOCI) approach based on natural orbitals for multiple electronic states of interest. The approach was designed with three key objectives in mind: (1) being systematic, (2) achieving a compact CI expansion, and (3) ensuring a balanced treatment of the system's electronic states. We will demonstrate, using simple test cases, that the presented approach forms a promising starting point to achieve these objectives.

# Non-orthogonal Configuration Interaction

Non-orthogonal configuration interaction is a generalized extension of the well-established orthogonal configuration interaction (CI) method. In the NOCI framework, the generalized eigenvalue problem to determine the electronic states is expressed as

<span id="page-2-0"></span>
$$
\sum_{J} (H_{IJ} - ES_{IJ}) C_J = 0.
$$
 (1)

where  $H_{IJ}$  and  $S_{IJ}$  represent Hamiltonian and overlap matrix elements in the basis of Slater determinants;  $C_J$  and  $E$  denote the expansion coefficients and the eigenvalue — the energy — of a specific state, respectively. The matrix elements can either be computed using Löwdin's pairing approach<sup>[17,](#page-15-5)[18](#page-15-6)</sup> in conjunction with the generalized Slater-Condon rules, or by employing the recently introduced generalized Wick's theorem.<sup>[19](#page-15-7)</sup>

NOCI offers significant flexibility in constructing the basis of Slater determinants, but this flexibility also poses a key challenge: defining an accurate and compact basis for the states of interest remains an open problem. In this work, we propose a novel approach utilizing principal natural orbital determinants to address this challenge.

#### Principal Natural Orbital Determinants as NOCI Basis

The natural orbitals (NOs) of a given electronic state are the eigenstates of the respective one-particle reduced density matrix (1-RDM):

$$
\gamma[\Psi]|\phi_p\rangle = n_p |\phi_p\rangle \tag{2}
$$

with the 1-RDM defined in a real-space basis according to

$$
\gamma[\Psi] = N \int \Psi^*(\mathbf{x}_1, \mathbf{x}_2, \dots) \Psi(\mathbf{x}_1', \mathbf{x}_2, \dots) d\mathbf{x}_2 d\mathbf{x}_3 \dots
$$
\n(3)

The corresponding eigenvalues are the natural occupation numbers  $\{n_p\}$  for which:

$$
n_p \in [0, 1]; \qquad \sum_p n_p = N_{\rm el} \tag{4}
$$

where  $N_{\rm el}$  denotes the number of electrons in the system. The NOs form an orthonormal basis and the determinant constructed from the  $N_{el}$  NOs with the highest occupation numbers is referred to as the principal natural orbital determinant. [20](#page-15-8) In the following, we will denote the principal NO determinant for state I (whose wavefunction is  $|\Psi_I\rangle$ ) as  $|\Phi_I\rangle$ . Since  $\gamma[\Phi_I]$  is the best possible idempotent approximation of  $\gamma[\Psi_I]$ , CI expansions based on these determinants are highly compact. [20,](#page-15-8)[21](#page-15-9)

The central idea of this work is to form, for each state of interest, a compact *orthogonal* CI basis based on its principal NO determinant. Since the CI expansions of the different states of interest are non-orthogonal to each other, this approach naturally leads to a NOCI problem.

To evaluate the benefits of including principal NO determinants from multiple states in the CI expansion — thereby transforming the approach into a NOCI one — we conducted tests on three simple systems taken from the QUEST database. [22](#page-16-0) Table [1](#page-5-0) summarizes the energies of the two lowest electronic states of these systems, calculated using CISD, CASSCF, CASCI, and the proposed NOCI approach. The NOCI approach, referred to as natural orbital NOCI (NO-NOCI), is annotated with numbers in parentheses specifying the number of principal NO determinants, followed by the size of the active space. Since we aim for a compact state basis and double excitations should have the largest contribution to the correlation energy, only single and double excitations are included in the presented NOCI approach, denoted as NO-NOCISD. The principal NO determinants for the NO-NOCISD variants were obtained from preceding CASCI(4e, 12o) calculations based on  $PBE^{23}$  $PBE^{23}$  $PBE^{23}$  references. The rational behind this decision was to lift Brillouin's Theorem to include as much information as possible in a CI expansion of limited length.

Inspecting Table [1,](#page-5-0) several observations are particularly noteworthy. Most importantly, when comparing the NOCI approach using only the principal NO determinant of the ground state with the approach including the principal NO determinants of both the ground and first excited states, the latter provides a significantly improved description of the first excited state. Table [2](#page-6-0) further illustrates the impact of the initial guess quality and the size of the single and double excitation space on the performance of the NO-NOCISD(1d) approach. As the quality of the initial guess increases, the NOs become more and more tailored to the state they were obtained for  $-$  in this case, the ground state. However, this results in a progressively poorer description of the first excited state. Expanding the excitation space in the NO-NOCISD(1d) approach significantly enhances the description of the excited state,

<span id="page-5-0"></span>Table 1: Energies of the two lowest lying electronic states of the three test systems formaldehyde, dinitrogen, and water obtained with various electronic structure methods. In all calculations, the def2-TZVP basis was employed. The NOCI methods employ CASCI(4e, 12o) principal NO determinants. The theorectical best estimate (TBE) at the complete basis set (CBS) limit was taken from Ref. [22](#page-16-0)

Method	No. Determinants	$E_0$ [H]	$E_1$ [H]	$\Delta E$ [eV]				
Formaldehyde								
CISD(2e, 6o)	36	$-113.896215$	$-113.753047$	3.90				
$NO-NOCISD(1d, 2e, 6o)$	36	$-113.926962$	$-113.718997$	5.66				
$NO-NOCISD(2d, 2e, 4o)$	32	$-113.929377$	$-113.783074$	3.98				
CASSCF(2e, 6o)	36	$-113.900280$	$-113.811091$	2.43				
CASCI(4e, 12o)	4356	$-113.938892$	$-113.786321$	4.15				
CASSCF(4e, 12o)	4356	$-113.993548$	$-113.848675$	3.94				
TBE(Full) CBS				3.58				
Dinitrogen								
$\overline{\text{CISD}}(2e, 6o)$	36	$-108.975125$	$-108.674581$	8.18				
$NO-NOCISD(1d, 2e, 6o)$	36	$-109.005804$	$-108.705817$	8.16				
$NO-NOCISD(2d, 2e, 4o)$	32	$-109.004870$	$-108.727730$	7.54				
CASSCF(2e, 6o)	36	$-108.995599$	$-108.689937$	8.32				
CASCI(4e, 12o)	4356	$-109.017840$	$-108.730362$	7.82				
CASSCF(4e, 12o)	4356	$-109.056915$	$-108.763409$	7.99				
TBE(Full) CBS				7.74				
Water								
CISD(2e, 6o)	36	$-76.057893$	$-75.766529$	7.93				
$NO-NOCISD(1d, 2e, 6o)$	36	$-76.061898$	$-75.464174$	16.26				
$NO-NOCISD(2d, 2e, 4o)$	32	$-76.061919$	$-75.794976$	7.26				
CASSCF(2e, 6o)	36	$-76.052454$	$-75.815680$	6.44				
CASCI(4e, 12o)	4356	$-76.085993$	$-75.803505$	7.69				
CASSCF(4e, 12o)	4356	$-76.108827$	$-75.858557$	6.81				
TBE(Full) CBS				7.33				

<span id="page-6-0"></span>Table 2: Impact of the initial guess quality and the size of the excitation space on the performance of the NO-NOCISD(1d) approach. All calculations were performed with the def2-TZVP basis set.

Method	$E_0$ [H]	$E_1$ [H]				
CASCI(2e, 40) Initial Guess						
$\overline{NO}$ -NOCISD $(1d, 2e, 6o)$	$-76.057894$	$-75.766529$				
$NO-NOCISD(1d, 2e, 8o)$	$-76.058190$	$-75.768593$				
$NO-NOCISD(1d, 2e, 10o)$	$-76.058768$	$-75.771224$				
$CASCI(4e, 12o)$ Initial Guess						
$\overline{\text{NO-NOCISD}}(1d, 2e, 60)$	$-76.061898$	$-75.464174$				
NO-NOCISD(1d, 2e, 8o)	$-76.062828$	$-75.516464$				
$NO-NOCISD(1d, 2e, 10o)$	$-76.062859$	$-75.685024$				
$CASCI(6e, 14o)$ Initial Guess						
$NO-NOCISD(1d, 2e, 6o)$	$-76.066963$	$-75.360981$				
$NO-NOCISD(1d, 2e, 8o)$	$-76.067515$	$-75.365945$				
$NO-NOCISD(1d, 2e, 10o)$	$-76.067968$	$-75.401444$				

while the improvement for the ground state is comparatively moderate. This nicely demonstrates the ordering of the NOs according to their relevance for describing the electronic state they were obtained for. Moreover, it underscores the necessity of including the principal NO determinants for all states of interest within the NOCI framework to achieve a compact and accurate basis.

When comparing NO-NOCISD(2d, 2e, 4o) with CISD using a similar number of determinants, NO-NOCISD(2d, 2e, 4o) shows significantly higher accuracy for the absolute energies of the two states, although CISD performs well in describing their energy difference. Similarly, when comparing NO-NOCISD(2d, 2e, 4o) with CASSCF employing a comparable number of determinants, the NOCI approach demonstrates overall better performance, particularly in reproducing the energy difference between the states.

It is also noteworthy that NO-NOCISD(2d, 2e, 4o) recovers the energies obtained with  $CASCI(4e, 12o)$  — which was used to generate the principal NO determinants — to a very good extent while utilising only about 1% of the determinants required by CASCI. To be more explicit, the worst agreement is observed for  $E_0$  of water, where NO-NOCISD(2d, 2e, 4o) still recovers 99.97% of the total CASSCI(4e, 12o) energy.

In summary, incorporating the principal NO determinants of multiple states proves highly advantageous. It enables the construction of a compact NOCI basis that achieves comparable accuracy across the states of interest.

### Orthogonal Determinant Selection

Thanks to the orthogonality "within" each state of interest, we can compute the corresponding Hamiltonian elements more efficiently and further leverage established algorithms to identify the most significant determinants.

Adopting the strategy used in heat-bath CI,  $^{24,25}$  $^{24,25}$  $^{24,25}$  $^{24,25}$  we include, for each state of interest, only the singly and doubly excited determinants that are connected to the respective principal NO determinant by a Hamiltonian matrix element with a magnitude exceeding a specified threshold  $\epsilon$ . Since these determinants are othogonal to each other, the respective matrix elements of the Hamiltonian can be evaluated according to

$$
H(r \leftarrow p) = \Gamma_{rp}^{I} \left( h_{rp} + \sum_{q \in \text{occ}} \left( g_{rqpq} - g_{qrpq} \right) \right)
$$
 (5)

$$
H(rs \leftarrow pq) = \Gamma_{rp}^I \Gamma_{sq}^J (g_{rspq} - g_{srpq})
$$
\n
$$
(6)
$$

with

$$
h_{rp} = \int \phi_r^*(\mathbf{x}) \left( -\frac{1}{2} \nabla^2 - \sum_A \frac{Z_A}{|\mathbf{r} - \mathbf{R}_A|} \right) \phi_p(\mathbf{x}) \, \mathrm{d}\mathbf{x} \tag{7}
$$

$$
g_{rspq} = \int \phi_r^*(\mathbf{x}_1) \phi_s^*(\mathbf{x}_2) \frac{1}{|\mathbf{r}_1 - \mathbf{r}_2|} \phi_p(\mathbf{x}_1) \phi_q(\mathbf{x}_2) \, \mathrm{d}\mathbf{x}_1 \, \mathrm{d}\mathbf{x}_2 \tag{8}
$$

and  $\Gamma_{rp}^I = (-1)^n$ , where *n* is the number of occupied spin-orbitals between *p* and *r* in state I. As we are concerned only with the magnitude of the matrix elements and not their sign, double excitations can be handled very efficiently. This is because the magnitude depends solely on the four orbitals whose occupations change during the excitation and not the other orbitals.

To assess the impact of the determinant selection, we conducted tests on the water molecule using the def2-TZVP basis set. A comparison of Fig[.1a](#page-8-0) and Fig[.1b](#page-8-0) reveals a significant improvement in the energies of both the ground and first excited states for small selection thresholds in the case of NO-NOCISD(2d, 2e, 8o). This improvement can be attributed to the non-orthogonal mixing of determinants from the two states, as indicated by the simultaneous energy jumps observed in both states, which are absent in the NO-NOCISD(1d, 2e, 8o) case.

<span id="page-8-0"></span>

Figure 1: Energies of the ground and the first excited state of  $H_2O$  evaluated with two different NO-NOCISD variants in the zeroth iteration. The def2-TZVP basis set was employed. The initial guess was obtained with CASCI(4e, 12o).

These results suggest a critical role of small Hamiltonian matrix elements in enhancing the ability of the NOCI approach to improve state descriptions through non-orthogonal mixing. Incorporating non-orthogonal information into the selection process could further improve the accuracy; however, this would entail significantly higher computational costs. For the sake of efficiency, we will adhere to the orthogonal approach outlined above in this work.

#### Iteratively Recalculating the Principal NO Determinant

To further reduce the size of the CI expansion, we tested an iterative process that involves repeatedly computing the 1-RDMs of the states of interest, forming the corresponding principal NO determinants, and selecting the most important determinants within a predefined active space using the procedure described earlier. The underlying idea is to leverage the non-orthogonal flexibility of the framework to encode information during the orbital optimization process, ultimately resulting in a more compact state basis. A further advantage of this approach is its potential to reduce dependence on the initial guess, which has thus far played a critical role since the principal NO determinants were obtained from the initial guesses for the 1-RDMs of the states of interest. While the influence of the initial guesses naturally decreases with larger active excitation spaces within the NOCI framework, the presented approach prioritizes maintaining a compact CI expansion, making small active spaces preferable.

For our investigations we again used water as test system and focused on two questions: 1) Does the iterative procedure make the CI expansion more compact? 2) Does it affect the accuracy?

1) Fig. [2](#page-10-0) and Fig. [3](#page-10-1) again show the energies of the ground and the first excited states as functions of the determinant selection threshold; this time, however, for iteration 1 and 2, respectively. Notably, the threshold at which the determinants of the two states begin to mix increases with each iteration. Additionally, fewer determinants are needed to achieve comparable accuracy (see inset). These results hence suggest that the NOCI expansion size can indeed be reduced by the employed iterative procedure.

2) Tab. [3](#page-11-0) summarizes the energies of the ground and first excited states as a function of the iteration, comparing a less accurate initial guess (left) with a significantly better initial guess (right). Two key observations can be made: First, for the less accurate initial guess, the iterative procedure leads to notable improvements in accuracy. Second, for the better initial guess, the iterative procedure does not enhance the accuracy but also does not lead to any

<span id="page-10-0"></span>

Figure 2: Energies of the ground and the first excited state of  $H_2O$  evaluated with two different NO-NOCISD variants in the first iteration. The def2-TZVP basis set was employed. The initial guess was obtained with CASCI(4e, 12o).

<span id="page-10-1"></span>

Figure 3: Energies of the ground and the first excited state of  $H_2O$  evaluated with two different NO-NOCISD variants in the second iteration. The def2-TZVP basis set was employed. The initial guess was obtained with CASCI(4e, 12o).

significant deterioration. These findings suggest that the iterative procedure is particularly beneficial when starting from poor initial guesses, while posing low risk to accuracy when the initial guess is already good. Naturally, larger active excitation spaces within the NOCI framework increase the potential for energy improvements during the iterative process and thereby further reduce the dependence on the initial guesses; however, since this approach prioritizes compactness, we intentionally limit the size of the active spaces.

<span id="page-11-0"></span>Table 3: NO-NOCISD(2d, 2e, 8o) energies of the ground and first excited states of  $H_2O$ across different iterations, starting from two initial guesses of different quality. The def2- TZVP basis set was employed.

Iteration			$CASCI(2e, 40)$ Initial Guess $CASCI(4e, 120)$ Initial Guess		
	$E_0$	$E_1$	$E_0$	$E_1$	
$\left( \right)$	$-76.058190$	$-75.768593$	$-76.071585$	$-75.796357$	
1	$-76.060309$	$-75.769467$	$-76.070916$	$-75.796386$	
$\mathcal{D}_{\mathcal{L}}$	$-76.061190$	$-75.769925$	$-76.070839$	$-75.796397$	
3	$-76.061207$	$-75.769931$	$-76.070801$	$-75.796402$	
4	$-76.061205$	$-75.769932$	$-76.070778$	$-75.796410$	

Regarding initial guesses for the proposed approach, it would be valuable to further explore methods other than CASSCI. Particularly interesting are methods that 1) treat different states on an equal footing and 2) are computationally inexpensive. Potential candidates include spin-flip TDDFT,  $26-29$  $26-29$  the constraint-based orbital-optimized excitation (COOX) method,<sup>[30](#page-16-6)</sup> and the maximum overlap method.<sup>[31](#page-17-0)</sup> However, investigations in these directions are left for future work.

# Final Procedure

Based on the findings of the preceding sections, we propose the following procedure for constructing a compact NOCI basis:

- 1. Define the states of interest (number of roots).
- 2. Compute initial guesses for the 1-RDMs of the states of interest.
- 3. Construct the principal NO determinant for each state of interest and include them in the NOCI expansion.
- 4. For each principal NO determinant, generate and include only those singly and doubly excited determinants within a predefined active space that are connected to the respective principal NO determinant by a Hamiltonian matrix element exceeding an adaptive threshold  $\epsilon$  in magnitude.
	- Adjust  $\epsilon$  to account for the increasing magnitude of the Hamiltonian matrix elements during the iterative process and to ensure that the NOCI expansion size is reduced.
- 5. Solve the generalized eigenvalue problem defined in Eq. [1.](#page-2-0)
- 6. Recalculate the 1-RDMs based on the updated NOCI states.
- 7. Repeat steps 3 through 6 until the degradation in the energy values of the states of interest exceeds a predefined threshold.

### Computational Details

The NO-NOCI approach was implemented in an in-house Python program interfaced with the FermiONs + program package.<sup>[32](#page-17-1)[,33](#page-17-2)</sup> All CISD and CASCI calculations were carried out using FermiONs++, while CASSCF calculations were performed with the ORCA quantum chemistry program package. [34–](#page-17-3)[38](#page-17-4) The def2-TZVP basis set [39](#page-17-5) was used consistently for all calculations.

# Conclusion

In this work, we introduced a novel procedure for constructing a compact NOCI basis to efficiently treat ground and excited states. The core idea is to include the principal natural orbital determinant of each state of interest in the NOCI expansion, ensuring all states are treated on equal footing. To maintain compactness, only the most important single and double excitations from these reference NO determinants are added to the expansion. The inclusion of multiple NO determinants naturally leads to a NOCI framework; however, the orthogonality "within" each state allows for efficient calculation of Hamiltonian matrix elements and determinant selection.

We employ an iterative procedure where the 1-RDMs of all states of interest are recalculated at each step to update the principal NO determinants. This approach fully exploits the flexibility of the non-orthogonal framework, enabling a further reduction in the expansion size without significant deterioration of accuracy.

Overall, this work represents a promising step toward a systematic procedure for generating compact and accurate NOCI expansions. However, we emphasize that this study serves as a proof of concept, and further testing and refinement will be needed. We hope that this work inspires further exploration and development in this area.

#### Acknowledgement

D. G. acknowledges funding by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – 498448112. D. G. thanks J. Kussmann (LMU Munich) for providing a development version of the FermiONs++ software package.

#### References

- <span id="page-13-0"></span>(1) Cerullo, G.; Polli, D.; Lanzani, G.; De Silvestri, S.; Hashimoto, H.; Cogdell, R. J. Photosynthetic light harvesting by carotenoids: detection of an intermediate excited state. Science 2002, 298, 2395–2398.
- <span id="page-13-1"></span>(2) Cheng, Y.-C.; Fleming, G. R. Dynamics of light harvesting in photosynthesis. Annu. Rev. Phys. Chem. 2009, 60, 241–262.
- <span id="page-14-0"></span>(3) Herbst, J.; Heyne, K.; Diller, R. Femtosecond infrared spectroscopy of bacteriorhodopsin chromophore isomerization. Science 2002, 297, 822–825.
- <span id="page-14-1"></span>(4) Tapavicza, E.; Tavernelli, I.; Rothlisberger, U. Trajectory surface hopping within linear response time-dependent density-functional theory. Phys. Rev. Lett. 2007, 98, 023001.
- <span id="page-14-2"></span>(5) Schultz, T.; Samoylova, E.; Radloff, W.; Hertel, I. V.; Sobolewski, A. L.; Domcke, W. Efficient deactivation of a model base pair via excited-state hydrogen transfer. Science 2004, 306, 1765–1768.
- <span id="page-14-3"></span>(6) Schreier, W. J.; Schrader, T. E.; Koller, F. O.; Gilch, P.; Crespo-Hernández, C. E.; Swaminathan, V. N.; Carell, T.; Zinth, W.; Kohler, B. Thymine dimerization in DNA is an ultrafast photoreaction. Science 2007, 315, 625–629.
- <span id="page-14-4"></span>(7) Roberts, S. J.; Szabla, R.; Todd, Z. R.; Stairs, S.; Bučar, D.-K.; Šponer, J.; Sasselov, D. D.; Powner, M. W. Selective prebiotic conversion of pyrimidine and purine anhydronucleosides into Watson-Crick base-pairing arabino-furanosyl nucleosides in water. Nat. Commun. 2018, 9, 4073.
- <span id="page-14-5"></span>(8) Xu, J.; Chmela, V.; Green, N. J.; Russell, D. A.; Janicki, M. J.; Góra, R. W.; Szabla, R.; Bond, A. D.; Sutherland, J. D. Selective prebiotic formation of RNA pyrimidine and DNA purine nucleosides. Nature 2020, 582, 60–66.
- <span id="page-14-6"></span>(9) Runge, E.; Gross, E. K. Density-functional theory for time-dependent systems. Phys. Rev. Lett. 1984, 52, 997.
- <span id="page-14-7"></span>(10) Hirata, S.; Head-Gordon, M. Time-dependent density functional theory within the Tamm–Dancoff approximation. Chem. Phys. Lett. 1999, 314, 291–299.
- <span id="page-14-8"></span>(11) Mardirossian, N.; Head-Gordon, M. Thirty years of density functional theory in computational chemistry: an overview and extensive assessment of 200 density functionals. Mol. Phys. 2017, 115, 2315–2372.
- <span id="page-15-0"></span>(12) Casida, M. E. Recent Advances In Density Functional Methods: (Part I); World Scientific, 1995; pp 155–192.
- <span id="page-15-1"></span>(13) Matsika, S. Electronic Structure Methods for the Description of Nonadiabatic Effects and Conical Intersections. Chem. Rev. 2021, 121, 9407–9449.
- <span id="page-15-2"></span>(14) Roos, B. O.; Taylor, P. R.; Sigbahn, P. E. A complete active space SCF method (CASSCF) using a density matrix formulated super-CI approach. Chem. Phys. 1980, 48, 157–173.
- <span id="page-15-3"></span>(15) Tran, L. N.; Shea, J. A.; Neuscamman, E. Tracking excited states in wave function optimization using density matrices and variational principles. J. Chem. Theory Comput. 2019, 15, 4790–4803.
- <span id="page-15-4"></span>(16) Werner, H.; Meyer, W. A quadratically convergent MCSCF method for the simultaneous optimization of several states. J. Chem. Phys. 1981, 74, 5794–5801.
- <span id="page-15-5"></span>(17) Löwdin, P. Band Theory, Valence Bond, and Tight-Binding Calculations. J. Appl. Phys. 1962, 33, 251–280.
- <span id="page-15-6"></span>(18) Amos, A. T.; Hall, G. G.; Jones, H. Single determinant wave functions. Proc. R. Soc. London Ser. A Math. Phys. Sci. 1961, 263, 483–493.
- <span id="page-15-7"></span>(19) Burton, H. G. A. Generalized nonorthogonal matrix elements: Unifying Wick's theorem and the Slater–Condon rules. J. Chem. Phys. 2021, 154, 144109.
- <span id="page-15-8"></span>(20) Yu, J. M.; Tsai, J.; Rajabi, A.; Rappoport, D.; Furche, F. Natural determinant reference functional theory. J. Chem. Phys. 2024, 160, 044102.
- <span id="page-15-9"></span>(21) Coe, J. P.; Paterson, M. J. Investigating Multireference Character and Correlation in Quantum Chemistry. J. Chem. Theory Comput. 2015, 11, 4189–4196.
- <span id="page-16-0"></span>(22) Loos, P.-F.; Scemama, A.; Blondel, A.; Garniron, Y.; Caffarel, M.; Jacquemin, D. A Mountaineering Strategy to Excited States: Highly Accurate Reference Energies and Benchmarks. J. Chem. Theory Comput. 2018, 14, 4360–4379.
- <span id="page-16-1"></span>(23) Perdew, J. P.; Burke, K.; Ernzerhof, M. Generalized Gradient Approximation Made Simple. Phys. Rev. Lett. 1996, 77, 3865–3868.
- <span id="page-16-2"></span>(24) Holmes, A. A.; Changlani, H. J.; Umrigar, C. J. Efficient Heat-Bath Sampling in Fock Space. J. Chem. Theory Comput. 2016, 12, 1561–1571.
- <span id="page-16-3"></span>(25) Holmes, A. A.; Tubman, N. M.; Umrigar, C. J. Heat-Bath Configuration Interaction: An Efficient Selected Configuration Interaction Algorithm Inspired by Heat-Bath Sampling. J. Chem. Theory Comput. 2016, 12, 3674–3680.
- <span id="page-16-4"></span>(26) Shao, Y.; Head-Gordon, M.; Krylov, A. I. The spin–flip approach within time-dependent density functional theory: Theory and applications to diradicals. J. Chem. Phys. 2003, 118, 4807–4818.
- (27) Orms, N.; Krylov, A. I. Singlet–triplet energy gaps and the degree of diradical character in binuclear copper molecular magnets characterized by spin-flip density functional theory. Phys. Chem. Chem. Phys. 2018, 20, 13127–13144.
- (28) Krylov, A. I. Reviews in Computational Chemistry; 2017; pp 151–224.
- <span id="page-16-5"></span>(29) Bernard, Y. A.; Shao, Y.; Krylov, A. I. General formulation of spin-flip time-dependent density functional theory using non-collinear kernels: Theory, implementation, and benchmarks. J. Chem. Phys. 2012, 136, 204103.
- <span id="page-16-6"></span>(30) Kussmann, J.; Lemke, Y.; Weinbrenner, A.; Ochsenfeld, C. A Constraint-Based Orbital-Optimized Excited State Method (COOX). J. Chem. Theory Comput. 2024, 20, 8461– 8473.
- <span id="page-17-0"></span>(31) Gilbert, A. T.; Besley, N. A.; Gill, P. M. Self-consistent field calculations of excited states using the maximum overlap method (MOM). J. Phys. Chem. A 2008, 112, 13164–13171.
- <span id="page-17-1"></span>(32) Kussmann, J.; Ochsenfeld, C. Pre-selective screening for matrix elements in linearscaling exact exchange calculations. J. Chem. Phys. 2013, 138, 134114.
- <span id="page-17-2"></span>(33) Kussmann, J.; Ochsenfeld, C. Preselective Screening for Linear-Scaling Exact Exchange-Gradient Calculations for Graphics Processing Units and General Strong-Scaling Massively Parallel Calculations. J. Chem. Theory Comput. 2015, 11, 918–922.
- <span id="page-17-3"></span>(34) Neese, F.; Wennmohs, F.; Becker, U.; Riplinger, C. The ORCA quantum chemistry program package. J. Chem. Phys. 2020, 152, 224108.
- (35) Neese, F. Software update: The ORCA program system—Version 5.0. WIREs Comput. Mol. Sci. 2022, 12, e1606.
- (36) Kollmar, C.; Sivalingam, K.; Helmich-Paris, B.; Angeli, C.; Neese, F. A perturbationbased super-CI approach for the orbital optimization of a CASSCF wave function. J. Comput. Chem. 2019, 40, 1463–1470.
- (37) Neese, F. The SHARK integral generation and digestion system. J. Comput. Chem. 2023, 44, 381–396.
- <span id="page-17-4"></span>(38) Ugandi, M.; Roemelt, M. A recursive formulation of one-electron coupling coefficients for spin-adapted configuration interaction calculations featuring many unpaired electrons. Int. J. Quantum Chem. 2023, 123, e27045.
- <span id="page-17-5"></span>(39) Weigend, F.; Ahlrichs, R. Balanced basis sets of split valence, triple zeta valence and quadruple zeta valence quality for H to Rn: Design and assessment of accuracy. Phys. Chem. Chem. Phys. 2005, 7, 3297–3305.