### ON THE HOPF SUPERALGEBRA OF SYMMETRIC FUNCTIONS IN SUPERSPACE

MASAMUNE HATTORI, RENTA YAGI, SHINTAROU YANAGIDA

ABSTRACT. We introduce a superspace analogue of combinatorial Hopf algebras (Aguiar-Bergeron-Sottile, 2006), and show that the Hopf superalgebra of quasi-symmetric (resp. symmetric) functions in superspace (Fishel-Lapointe-Pinto, 2019) is a terminal object in the category of all (resp. cocommutative) combinatorial Hopf superalgebras. We also introduce a superspace analogue of chromatic symmetric functions of graphs (Stanley, 1995) using the chromatic Hopf superalgebra of two-colored graphs.

#### CONTENTS

Introduction	1
Hopf superalgebra	3
Symmetric and quasi-symmetric functions in superspace	3
Combinatorial Hopf superalgebra	6
A superspace analogue of chromatic symmetric functions	8
oferences	13
	Hopf superalgebra Symmetric and quasi-symmetric functions in superspace Combinatorial Hopf superalgebra A superspace analogue of chromatic symmetric functions ferences

### 1. INTRODUCTION

**Background.** The ring of symmetric functions [M95, Chap. I], [S99, Chap. 7] is a commonplace of various mathematics, including representation theory, (quantum) integrable systems, and combinatorics. It consists of the symmetric polynomials in infinitely many variables  $x = (x_1, x_2, ...)$ , and the ring structure is given by addition and multiplication of such polynomials. A closely related object is the ring of quasi-symmetric functions [G84], which can be seen as a refinement of symmetric functions and provides many applications in enumeration problems of combinatorics and algebraic geometry.

Both of these rings have the Hopf algebra structures [GR, Chap. 2, Chap. 5]. Among the vast amount of studies on these Hopf algebras, let us recall the work of Aguiar, Bergeron and Sottile [ABS06]. They introduced the notion of a combinatorial Hopf algebra as a graded connected Hopf algebra H over a field  $\mathbf{k}$  equipped with an algebra morphism  $\zeta: H \to \mathbf{k}$  called a *character*, and showed that the Hopf algebra of quasi-symmetric (resp. symmetric) functions is a terminal object in the category of all (resp. cocommutative) combinatorial Hopf algebras. It gives a characterization of the Hopf algebras of (quasi-)symmetric functions, and explains the ubiquity of (quasi-)symmetric functions in representation theory and combinatorics. Let us also refer to [GR, Chap. 7] for more explanation of Aguiar-Bergeron-Sottile theory.

Over the past 20 years, a superspace extension of the theory of symmetric functions has been developed and has received much attention [A+, B+12, DLM03, DLM06, DLM07]. In the superspace setting, one considers the polynomials in infinitely many supervariables  $(x; \theta) = (x_1, x_2, \ldots; \theta_1, \theta_2, \ldots)$ , where  $x_i$ 's are the usual commuting variables but  $\theta_i$ 's are the anti-commuting variables  $(\theta_i \theta_j = -\theta_j \theta_i$  and  $\theta_i^2 = 0)$ . Considering the "diagonal permutation of supervariables" (see § 3.1, (3.1) for the precise definition), one has the notion of symmetric functions in superspace [A+, §2].

Recently, Fishel, Lapointe and Pinto [FLP19] investigated a superspace analogue of the rich connection between symmetric function theory and quasi-symmetric functions, and revealed the Hopf algebra structure

GRADUATE SCHOOL OF MATHEMATICS, NAGOYA UNIVERSITY. FUROCHO, CHIKUSAKU, NAGOYA, JAPAN, 464-8602.

*E-mail address*: m21039e@math.nagoya-u.ac.jp, renta.yagi.e6@math.nagoya-u.ac.jp, yanagida@math.nagoya-u.ac.jp. *Date*: 2024.12.23.

Key words and phrases. Hopf superalgebras, super-characters, (quasi-)symmetric functions in superspace, chromatic symmetric functions in superspace.

of the symmetric functions in superspace  $\Lambda$  and of the quasi-symmetric functions in superspace sQSym. These Hopf superalgebras (see Definition 2.1 for the precise meaning) are natural super-analogue of the Hopf algebras of (quasi-)symmetric functions. We will review these topics in § 3.

Main results. In this paper, we investigate a superspace analogue of Aguiar-Bergeron-Sottile theory, and give a characterization of the Hopf superalgebras of (quasi-)symmetric functions in superspace.

Let us explain the idea of our superspace analogue. Since a combinatorial Hopf algebra H is equipped with a character  $\zeta \colon H \to \mathbf{k}$ , we need a natural superspace analogue of a character. Now, we recall some basic theory of (affine) super algebraic geometry [KV11, §1, §2]. A character  $\zeta$  can be regarded as a morphism from the point Spec  $\mathbf{k}$  to "the non-commutative scheme Spec H". According to [KV11, §2.2], as a superspace analogue of the point Spec  $\mathbf{k}$ , we may consider the " $\mathcal{N} = 1$  supersymmetric particle" Spec  $\mathbf{k}[\varepsilon]$ , where  $\mathbf{k}[\varepsilon] = \mathbf{k} + \mathbf{k}\varepsilon$  is the exterior algebra of one variable  $\varepsilon$  (the commutative superalgebra in one odd variable  $\varepsilon$ ). This leads us to the following definition.

**Definition** (Definition 4.1). Let H be a Hopf superalgebra over  $\mathbf{k}$ .

- (1) An even superalgebra morphism  $\zeta \colon H \to \mathbf{k}[\varepsilon]$  is called a *supercharacter* of H.
- (2) *H* is called a *combinatorial Hopf superalgebra* if it is connected graded (see Definition 2.2) and has a supercharacter  $\zeta : H \to \mathbf{k}[\varepsilon]$ .

We will see in §4 that  $\Lambda$  and sQSym are combinatorial Hopf superalgebras with supercharacter  $\zeta_S$  (4.2) and  $\zeta_Q$  (4.1). The main results of this note are:

**Theorem 1** (Theorem 4.2, Proposition 4.4). The Hopf superalgebra sQSym (resp.  $\Lambda$ ) is a terminal object in the category of all (resp. cocommutative) combinatorial Hopf superalgebras. In other words, we have:

- Let  $A = \bigoplus_{k>0} A^k$  be a combinatorial Hopf superalgebra with supercharacter  $\zeta \colon A \to \mathbf{k}[\varepsilon]$ .
- (1) There is a unique graded even morphism  $\Psi: A \to \mathrm{sQSym}$  of graded Hopf superalgebras such that  $\zeta = \zeta_Q \circ \Psi$ , where  $\zeta_Q$  is the supercharacter (4.1).
- (2) If A is cocommutative, then there is a unique graded even morphism  $\Psi: A \to \Lambda$  of graded Hopf superalgebras such that  $\zeta = \zeta_S \circ \Psi$ , where  $\zeta_S$  is the supercharacter (4.2).

As another example of a combinatorial Hopf superalgebra, we will investigate in the final §5 a superspace analogue of chromatic symmetric functions of graphs [S95]. Extending the non-superspace case [GR, §7.3], we consider the space  $\mathcal{G}$  spanned by the basis elements [G] associated to two-colored graphs G = (V, W, E)(the set W of white vertices, the set  $V \setminus W$  of black vertices and the set E of edges), which is a cocommutative combinatorial Hopf superalgebra (Theorem 5.2). Then, by the above Theorem 1 (2), we have a graded even morphism  $\Psi_{ch}: \mathcal{G} \to \Lambda$  of graded Hopf superalgebras.

**Definition** (Definition 5.3). For a two-colored graph G, we call  $\Psi_{ch}([G])$  the chromatic symmetric function in superspace associated to G.

If G = (V, W, E) satisfies  $W = \emptyset$ , then  $\Psi_{ch}([G])$  is equal to Stanley's chromatic symmetric function of the graph (V, E). We will show in Proposition 5.6 an expansion formula of chromatic symmetric functions in superspace.

**Organization.** Let us explain the organization of this note.

§ 2 and § 3 are preliminary. In § 2, we fix our terminology on Hopf algebras in super setting. In § 3, we review the Hopf superalgebras  $\Lambda$  and sQSym of symmetric and quasi-symmetric functions in superspace, largely following [A+] and [FLP19].

§4 is the main part of this paper. We introduce the notion of a combinatorial Hopf superalgebra in Definition 4.1, and characterize sQSym (resp.  $\Lambda$ ) as a terminal object of the category of all (resp. cocommutative) Hopf superalgebras in Theorem 4.2 (resp. Proposition 4.4). This can be regarded as a superspace extension of Aguiar-Bergeron-Sottile theory.

The final §5 gives a superspace analogue of chromatic symmetric functions. We introduce the chromatic Hopf superalgebra and show it is combinatorial (Theorem 5.2). Then, we introduce in Definition 5.3 the notion of a chromatic symmetric function in superspace using the universality of  $\Lambda$ , and study some examples (Proposition 5.5) and expansion formula (Proposition 5.6).

Global notation. The following list gives an overview of the terminology and notations used in the text.

- The symbol  $\mathbb{N}$  denotes the set  $\{0, 1, 2, ...\}$  of all nonnegative integers.
- We use the standard symbols  $\mathbb{Z}$  and  $\mathbb{Q}$  of integers and rational numbers.
- For a (finite) set S, we denote its cardinality by #S.
- A ring or an algebra means a unital associative one unless otherwise stated.
- A coalgebra means a counital coassociative one unless otherwise stated.
- The symbol  $\partial_x$  denotes the partial differential  $\frac{\partial}{\partial x}$  with respect to x.
- The symbol  $\delta_{i,j}$  denotes the Kronecker delta.
- We follow [DM99] and [BE17, §1] for the terminology of super mathematics. We denote the cyclic group of order 2 by Z/2Z = {0,1}, and a Z/2Z-grading is called *parity*. The parity of an object x is denoted by |x| ∈ Z/2Z.

#### 2. Hopf superalgebra

Let us fix the terminology and notation of what we will call Hopf superalgebras.

Let **k** be a field of characteristic 0, and consider linear spaces and linear maps over **k**. A superspace means a  $\mathbb{Z}/2\mathbb{Z}$ -graded linear space. For a superspace V, the even and odd homogeneous parts are denoted by  $V_{\overline{0}}$  and  $V_{\overline{1}}$ , respectively. A linear map  $f: V \to W$  between superspaces V and W is even (resp. odd) if it preserves (resp. reverses) the parity of elements. A linear map f decomposes as  $f = f_{\overline{0}} + f_{\overline{1}}$  with  $f_{\overline{0}}$  even and  $f_{\overline{1}}$  odd. This makes the set Hom(V, W) of linear maps into a superspace. Then, superspaces and linear maps between them form a supercategory SVec in the sense of [BE17, Definition 1.1].

Svec has the standard structure of a monoidal supercategory [BE17, Definition 1.4] with tensor product  $(V \otimes W)_{\overline{0}} := V_{\overline{0}} \otimes W_{\overline{0}} + V_{\overline{1}} \otimes W_{\overline{1}}$  and  $(V \otimes W)_{\overline{1}} := V_{\overline{0}} \otimes W_{\overline{1}} + V_{\overline{1}} \otimes W_{\overline{0}}$  for superspaces V, W, and  $(f \otimes g)(v \otimes w) := (-1)^{|g||v|} f(v) \otimes g(w)$  for linear maps f, g and  $v \in V, w \in W$ . Here and hereafter we denote by  $|x| \in \mathbb{Z}/2\mathbb{Z}$  the parity of the object x. Note that the composition of tensor products of linear maps has a sign:  $(f \otimes g) \circ (h \otimes i) = (-1)^{|g||h|} (f \circ h) \otimes (g \circ i)$ .

Given a monoidal supercategory C, one has the notions of a *superalgebra object*, of a (*super-*)coalgebra object, and of a (*super-*)bialgebra object in C. Now, let us introduce:

**Definition 2.1.** A Hopf superalgebra H (over  $\mathbf{k}$ ) is a Hopf algebra object in the monoidal supercategory SVec.

The monoidal supercategory SVec is symmetric with braiding  $u \otimes v \mapsto (-1)^{|u||v|} v \otimes u$ . Hence, we have the notions of a *commutative* Hopf superalgebra and of a *cocommutative* Hopf superalgebra.

In preparation for  $\S4$ , let us also introduce:

**Definition 2.2.** Let *H* be a Hopf superalgebra.

(1) H is called graded if it has an additional  $\mathbb{N}$ -grading

$$H = \bigoplus_{k \ge 0} H^k = \bigoplus_{k \ge 0} (H^k_{\overline{0}} \oplus H^k_{\overline{1}}).$$

$$(2.1)$$

- (2) *H* is called *graded connected* if it is graded,  $H = \bigoplus_{k>0} H^k$ , and  $H^0 = \mathbf{k}$ .
- (3) A graded morphism  $H \to H'$  of Hopf superalgebras is a morphism of Hopf superalgebras which preserves the N-grading.

**Remark 2.3.** In this note, we will always denote the  $\mathbb{N}$ -grading of a graded Hopf superalgebra H by the upper index  $H^*$ , and the parity ( $\mathbb{Z}/2\mathbb{Z}$ -grading) by the lower index  $H_{\overline{*}}$  as in (2.1).

For a graded Hopf superalgebra  $H = \bigoplus_{k\geq 0} H^k$ , we denote the graded dual by  $H^o := \bigoplus_{k\geq 0} (H^k)^*$ ,  $(H^k)^* := \operatorname{Hom}(H^k, \mathbf{k})$ . If dim  $H^k < \infty$  for each  $k \geq 0$ , then the graded dual  $H^o$  has a natural Hopf superalgebra structure. (See [GR, §1.6] for the even case. For the existence of the antipode, use [FLP19, 2.1] and [GR, Proposition 1.4.16].)

**Definition 2.4.** We call the Hopf superalgebra structure on  $H^o$  the Hopf dual of H.

### 3. Symmetric and quasi-symmetric functions in superspace

Here we recall from [FLP19, §§3–5] the Hopf superalgebras  $\Lambda$  and sQSym of symmetric and quasisymmetric functions in superspace. Let **k** be a field of characteristic 0. 3.1. Symmetric functions in superspace. Following [A+, §2] and [FLP19, §3], we denote by  $\mathbf{k}[x,\theta]_N$  the polynomial algebra in the supervariables

$$(x;\theta) = (x_1,\ldots,x_N;\theta_1,\ldots,\theta_N)$$

with even x and odd  $\theta$ , and consider the action of the symmetric group  $\mathfrak{S}_N$  on  $\mathbf{k}[x,\theta]_N$  in the diagonal way:

$$\sigma(x_1, \dots, x_N; \theta_1, \dots, \theta_N) \coloneqq (x_{\sigma(1)}, \dots, x_{\sigma(N)}; \theta_{\sigma(1)}, \dots, \theta_{\sigma(N)}) \quad (\sigma \in \mathfrak{S}_N).$$
(3.1)

The invariant ring  $\mathbf{k}[x,\theta]_N^{\mathfrak{S}_N}$  is called the ring of symmetric polynomials in N supervariables. It is doubly graded:

$$\mathbf{k}[x,\theta]_N^{\mathfrak{S}_N} = \bigoplus_{n \ge 0} \bigoplus_{0 \le m \le N} (\mathbf{k}[x,\theta]_N^{\mathfrak{S}_N})_{n,m},$$

where  $(\mathbf{k}[x,\theta]_N^{\mathfrak{S}_N})_{n,m}$  is the space of homogeneous elements of degree n in x and of degree m in  $\theta$ . The degree n with respect to x called the *total degree*, and that with respect to  $\theta$  called the *fermionic degree*.

The space  $\Lambda$  of symmetric functions in superspace is defined by inverse limit [A+, §2.2], [FLP19, §3.1]. That is, for M > N, we consider a linear map  $(\mathbf{k}[x, \theta]_M^{\mathfrak{S}_M})_{n,m} \to (\mathbf{k}[x, \theta]_N^{\mathfrak{S}_N})_{n,m}$  defined by  $x_{N+1} = \cdots = x_M = 0$ ,  $\theta_{N+1} = \cdots = \theta_M = 0$ , while send the other variables identically. Then we have an inverse system  $\{(\mathbf{k}[x, \theta]_N^{\mathfrak{S}_N})_{n,m}\}_N$  of linear spaces for each  $n, m \ge 0$ . Now we define a doubly graded linear space  $\Lambda$  by

$$\mathbf{\Lambda} \coloneqq \bigoplus_{n,m \ge 0} \mathbf{\Lambda}_{n,m}, \quad \mathbf{\Lambda}_{n,m} \coloneqq \varprojlim_N (\mathbf{k}[x,\theta]_N^{\mathfrak{S}_N})_{n,m},$$

and call it the space of symmetric functions in superspace. For each  $n, m \ge 0$ , the subspace  $\Lambda_{n,m}$  consists of symmetric polynomials in infinite supervariables of total degree n and fermionic degree m. The space  $\Lambda$ is an algebra, inheriting the multiplication on  $\mathbf{k}[x,\theta]_N^{\mathfrak{S}_N}$ .

Using this bigrading, we can introduce an N-graded superalgebra structure on  $\Lambda$ . First, define the parity ( $\mathbb{Z}/2\mathbb{Z}$ -grading) by

$$\mathbf{\Lambda} = \mathbf{\Lambda}_{\overline{0}} \oplus \mathbf{\Lambda}_{\overline{1}}, \quad \mathbf{\Lambda}_{\overline{p}} \coloneqq \bigoplus_{n \ge 0, \ m \equiv \overline{p} \bmod 2} \mathbf{\Lambda}_{n,m} \quad (p = 0, 1).$$
(3.2)

Then,  $\Lambda$  is a commutative superalgebra (in the supercategory SVec). Moreover, it has an additional N-grading

$$\mathbf{\Lambda} = \bigoplus_{k \ge 0} \mathbf{\Lambda}^k, \quad \mathbf{\Lambda}^k := \bigoplus_{n+m=k} \mathbf{\Lambda}_{n,m} \quad (k \ge 0).$$
(3.3)

makes it further into a graded commutative superalgebra.

To explain the Hopf superalgebra structure on  $\Lambda$ , we recall from [A+, §2.3] some basic supersymmetric functions. We will freely use the notions and symbols of superpartitions in [A+, §2.1] and [FLP19, §3]. Let  $\Lambda = (\Lambda^a; \Lambda^s) = (\Lambda_1, \ldots, \Lambda_m; \Lambda_{m+1}, \ldots, \Lambda_N)$  be a superpartition of length  $\leq N$ , total degree n and fermionic degree m. Then, the monomial polynomial  $m_{\Lambda} \in (\mathbf{k}[x, \theta]_N^{\mathfrak{S}_N})_{n,m}$  is defined as

$$m_{\Lambda}(x,\theta) \coloneqq (\#\operatorname{Aut}(\Lambda^s))^{-1} \sum_{\sigma \in \mathfrak{S}_N} \sigma(x^{\lambda}\theta_1 \cdots \theta_m),$$

where Aut( $\Lambda^s$ ) denotes the order of the automorphism group of the partition  $\Lambda^s = (\Lambda_{m+1}, \ldots, \Lambda_n)$  (see [A+, (2.12)] for the detail), and the  $\mathfrak{S}_N$ -action is given by (3.1). The family  $\{m_\Lambda \in (\mathbf{k}[x,\theta]_N^{\mathfrak{S}_N})_{n,m}\}_N$  is stable under the inverse limit, and we have the monomial symmetric function in superspace

$$m_{\Lambda} \in \mathbf{\Lambda}_{n,m} \tag{3.4}$$

for each superpartition  $\Lambda$  of total degree n and fermionic degree m. The family  $\{m_{\Lambda} \mid \Lambda: \text{ superpartitions}\}$  is a basis of the space  $\Lambda$ .

Next, the elementary symmetric functions in superspace  $e_r, \tilde{e}_s \in \Lambda$  are defined by

$$e_r \coloneqq m_{(\emptyset;1^r)}, \quad \tilde{e}_s \coloneqq m_{(0;1^s)} \quad (r \ge 1, \, s \ge 0), \tag{3.5}$$

Here we denoted  $1^r \coloneqq 1, \ldots, 1$ . They generate the superalgebra  $\Lambda$  [A+, (2.20)]:

$$\Lambda \cong \mathbf{k}[e_1, e_2, \dots; \widetilde{e}_0, \widetilde{e}_1, \dots].$$
(3.6)

Below we use the additional symbol  $e_0 := 1 \in \Lambda$ .

Now we can explain the comultiplication  $\Delta$  on  $\Lambda$  [FLP19, Proposition 4.1]. On the generator  $e_r$  and  $\tilde{e}_s$ , it is given by

$$\Delta(e_r) = \sum_{k,l \ge 0, \, k+l=r} e_k \otimes e_l, \quad \Delta(\widetilde{e}_s) = \sum_{k,l \ge 0, \, k+l=s} (\widetilde{e}_k \otimes e_l + e_k \otimes \widetilde{e}_l).$$

By [FLP19, Proposition 4.8], this comultiplication is cocommutative (in the supercategory SVec) and graded with respect to the N-grading (3.3), and a superalgebra morphism  $\Lambda \to \Lambda \otimes \Lambda$ . Since  $\Lambda$  is connected, i.e.  $\Lambda^0 = \Lambda_0^0 \oplus \Lambda_1^0 = \mathbf{k}$ , by [FLP19, Theorem 2.1], the bialgebra  $\Lambda$  has a unique antipode, and thus is a Hopf superalgebra. Let us summarize as:

**Proposition 3.1** ([FLP19, §4]). The space  $\Lambda$  of symmetric functions in superspace has a connected graded Hopf superalgebra structure which is commutative and cocommutative. The parity is (3.2) and the  $\mathbb{N}$ -grading is (3.3).

3.2. Quasi-symmetric functions in superspace. The contents of this subsection are based on [FLP19, §5, §6].

Let us recall the notion of a *dotted composition* from [FLP19, Definition 5.1]. It is a finite sequence  $\alpha = (\alpha_1, \ldots, \alpha_l)$  of entries belonging to the set  $\{1, 2, 3, \ldots\} \cup \{\dot{0}, \dot{1}, \dot{2}, \ldots\}$ . For such  $\alpha$ , we define  $\eta(\alpha) = (\eta_1, \ldots, \eta_l) \in \{0, 1\}^l$  with

$$\eta_k \coloneqq \begin{cases} 1 & (\text{if } \alpha_k \text{ is dotted}) \\ 0 & (\text{otherwise}) \end{cases}$$
(3.7)

The total degree of  $\alpha$  is defined to be  $\sum_{k=1}^{l} \alpha_k$ , where the dotted entries  $\alpha_k = \dot{n}$  are replaced by n. The fermionic degree of  $\alpha$  is defined to be the number of dotted entries in  $\alpha$ .

Let  $(x; \theta) = (x_1, x_2, \ldots; \theta_1, \theta_2, \ldots)$  be the set of infinite supervariables as before, and let R be the ring of formal power series of finite degree in  $(x; \theta)$ . An element  $f \in R$  is called a *quasi-symmetric function in* superspace if for every dotted composition  $\alpha = (\alpha_1, \ldots, \alpha_l)$ , all monomials  $\theta_{i_1}^{\eta_1} \cdots \theta_{i_l}^{\eta_l} x_{i_1}^{\alpha_1} \cdots x_{i_l}^{\alpha_l}$  in f with indices  $i_1 < \cdots < i_l$  have the same coefficient, where  $(\eta_1, \ldots, \eta_l) = \eta(\alpha)$  is given by (3.7).

Quasi-symmetric functions in superspace form a linear space, which is denoted by sQSym. It has a bigrading by total degree n and fermionic degree m [FLP19, (5.2)], which is denoted by

$$\mathrm{sQSym} = \bigoplus_{n,m \geq 0} \mathrm{sQSym}_{n,m}$$

Then, sQSym is a commutative superalgebra whose parity is the fermionic degree modulo 2:

$$sQSym = sQSym_{\overline{0}} \oplus sQSym_{\overline{1}}, \quad sQSym_{\overline{p}} \coloneqq \bigoplus_{n \ge 0, \ m \equiv p \ \text{mod} \ 2} sQSym_{n,m} \quad (p = 0, 1).$$
(3.8)

It has an additional  $\mathbb{N}$ -grading:

$$sQSym = \bigoplus_{k \ge 0} sQSym^k, \quad sQSym^k \coloneqq \bigoplus_{n+m=k} sQSym_{n,m} \quad (k \ge 0).$$
 (3.9)

One can see that the graded superspace  $\Lambda$  of symmetric functions in superspace is a subspace of sQSym.

In order to explain the comultiplication  $\Delta$  of sQSym, let us introduce a natural basis [FLP19, §5.1]. For each dotted composition  $\alpha = (\alpha_1, \ldots, \alpha_l)$ , we define the monomial quasi-symmetric function  $M_{\alpha}$  by

$$M_{\alpha} \coloneqq \sum_{i_1 < \dots < i_l} \theta_{i_1}^{\eta_1} \cdots \theta_{i_l}^{\eta_l} x_{i_1}^{\alpha_1} \cdots x_{i_l}^{\alpha_l}.$$
(3.10)

The family  $\{M_{\alpha} \mid \alpha: \text{ superpartitions}\}$  is a basis of sQSym.

Now, the comultiplication  $\Delta$  is given in terms of  $M_{\alpha}$ ,  $\alpha = (\alpha_1, \ldots, \alpha_l)$ , as

$$\Delta(M_{\alpha}) = \sum_{k=0}^{l} M_{(\alpha_1,\dots,\alpha_k)} \otimes M_{(\alpha_{k+1},\dots,\alpha_l)}.$$

Note that  $\Delta$  is not cocommutative (see [FLP19, Example 5.8] for an explicit example). By [FLP19, Proposition 5.9],  $\Delta$  is graded with respect to the N-grading (3.3), and is a superalgebra morphism sQSym  $\rightarrow$  sQSym  $\otimes$  sQSym. Since sQSym is connected, i.e., sQSym<sup>0</sup> = sQSym<sup>0</sup>  $\oplus$  sQSym<sup>0</sup> = k, by [FLP19, Theorem 2.1], the bialgebra sQSym has a unique antipode, and is a Hopf superalgebra. Let us summarize as:

**Proposition 3.2** ([FLP19, §5]). The space sQSym of quasi-symmetric functions in superspace has a connected graded Hopf superalgebra structure which is commutative but not cocommutative. The parity is (3.8) and the N-grading is (3.9).

# 4. Combinatorial Hopf superalgebra

We consider a superspace analogue of a combinatorial Hopf algebra [ABS06] (see also [GR, §7]).

As before, let **k** be a field of characteristic 0. We denote by  $\mathbf{k}[\varepsilon]$  be the commutative superalgebra over **k** generated by an odd element  $\varepsilon$ . Thus, we have  $\varepsilon^2 = 0$  and  $\mathbf{k}[\varepsilon] = \mathbf{k} + \mathbf{k}\varepsilon$  as linear spaces.

# **Definition 4.1.** Let H be a Hopf superalgebra over $\mathbf{k}$ .

- (1) An even superalgebra morphism  $\zeta \colon H \to \mathbf{k}[\varepsilon]$  is called a *supercharacter* of H.
- (2) A combinatorial Hopf superalgebra is a pair  $(H, \zeta)$  consisting of a connected graded Hopf superalgebra H and a supercharacter  $\zeta \colon H \to \mathbf{k}[\varepsilon]$ .
- (3) Let  $(H, \zeta)$  and  $(H', \zeta')$  be combinatorial Hopf superalgebras. A morphism  $(H, \zeta) \to (H', \zeta')$  of combinatorial Hopf superalgebras is a graded even morphism  $\psi \colon H \to H'$  of Hopf superalgebras such that  $\zeta = \zeta' \circ \psi$ . This yields the category of combinatorial Hopf superalgebras.

Note that, for a supercharacter  $\zeta \colon H \to \mathbf{k}[\varepsilon]$ , we have  $\zeta(1) = 1$ ,  $|\zeta(a)| = |a|$  and  $\zeta(ab) = \zeta(a)\zeta(b)$  for  $a, b \in H$ . In particular, for  $a, b \in H_{\overline{1}}$ , we have  $\zeta(a), \zeta(b) \in \mathbf{k}\varepsilon$  and  $\zeta(ab) = 0$ .

For example, the Hopf superalgebra sQSym has a supercharacter

$$\zeta_Q \colon \mathrm{sQSym} \longrightarrow \mathbf{k}[\varepsilon] \tag{4.1}$$

given by the specialization  $x_1 = 1$ ,  $x_2 = x_3 = \cdots = 0$  and  $\theta_1 = \varepsilon$ ,  $\theta_2 = \theta_3 = \cdots = 0$ . The same specialization defines a supercharcter on the Hopf superalgebra  $\Lambda$ :

$$\zeta_S \colon \mathbf{\Lambda} \longrightarrow \mathbf{k}[\varepsilon]. \tag{4.2}$$

Then, by Propositions 3.1 and 3.2, the pairs (sQSym,  $\zeta_Q$ ) and ( $\Lambda, \zeta_S$ ) are combinatorial Hopf superalgebras. Now, we state the main statement of this note, which is a superspace extension of [GR, Theorem 7.1.3].

**Theorem 4.2.** The pair (sQSym,  $\zeta_Q$ ) is a terminal object of the category of combinatorial Hopf superalgebras. In other words, we have:

Let  $A = \bigoplus_{k \ge 0} A^k$  be a combinatorial Hopf superalgebra with supercharacter  $\zeta \colon A \to \mathbf{k}[\varepsilon]$ . Then, there is a unique graded even morphism  $\Psi \colon A \to \mathrm{sQSym}$  of graded Hopf superalgebras such that  $\zeta = \zeta_Q \circ \Psi$ , where  $\zeta_Q$  is the supercharacter (4.1).

For the proof, we need the Hopf superalgebra sNSym of noncommutative symmetric functions in superspace [FLP19, §6]. It is defined to be the Hopf dual of sQSym (c.f. Definition 2.4). Precisely speaking, its underlying **k**-linear space is the bigraded dual of sQSym, i.e.,  $\text{sNSym} \coloneqq \bigoplus_{n,m\geq 0} \text{sNSym}_{n,m}$  with  $\text{sNSym}_{n,m} \coloneqq (\text{sQSym}_{n,m})^*$ . Also, sNSym has a **k**-basis  $\{H_\alpha\}$  labeled by dotted compositions  $\alpha$  which is dual to the family  $\{M_\alpha\}$  of monomial quasi-symmetric functions in superspace (3.10). By [FLP19, Proposition 6.2], sNSym is isomorphic (as an algebra) to the free associative algebra with noncommuting generators  $\{H_1, H_2, \ldots, ; \tilde{H}_0, \tilde{H}_1, \ldots\}$ , where we set

$$H_r \coloneqq H_{(r)} \quad (r \ge 1), \quad H_s \coloneqq H_{(s)} \quad (s \ge 0). \tag{4.3}$$

The Hopf superalgebra sNSym is connected graded for the grading  $\mathrm{sNSym} = \bigoplus_{k \ge 0} \mathrm{sNSym}^k$  with  $\mathrm{sNSym}^k := \bigoplus_{n+m=k} \mathrm{sNSym}_{n,m}$ . Note that we have  $H_r \in \mathrm{sNSym}_{r,0} \subset \mathrm{sNSym}_{\overline{0}}^r$   $(r \ge 1)$  and  $H_s \in \mathrm{sNSym}_{s,1} \subset \mathrm{sNSym}_{\overline{1}}^{s+1}$   $(s \ge 0)$ .

Let us also remark that giving a supercharacter on sQSym is equivalent to giving a family of functionals in each homogeneous space  $sQSym_{\overline{p}}^{k} = \bigoplus_{n+m=k, m \equiv \overline{p} \mod 2} sQSym_{n,m}$ . Under this equivalence, the supercharacter  $\zeta_Q$  corresponds to the functionals  $H_r, \widetilde{H}_s$ . That is, for  $f \in sQSym^k$ , we have

$$\zeta_Q(f) = \begin{cases} (H_k, f) & (|f| = \overline{0}) \\ (\widetilde{H}_{k-1}, f) & (|f| = \overline{1}) \end{cases},$$
(4.4)

where |p| denotes the parity of f.

Proof of Theorem 4.2. We follow the non-super case [GR, Theorem 7.1.3]. First, note that we can decompose the given  $\zeta \colon A \to \mathbf{k}[\varepsilon]$  as  $\zeta = \zeta_{\overline{0}} \oplus \zeta_{\overline{1}}\varepsilon$ , where  $\zeta_{\overline{p}} \colon A_{\overline{p}} \to \mathbf{k}$  (p = 0, 1) are linear,  $\zeta_{\overline{0}}$  is even and  $\zeta_{\overline{1}}$  is odd. Let us define an even linear map

$$\zeta' \colon \left( \bigoplus_{r \ge 1} \mathbf{k} H_r \right) \oplus \left( \bigoplus_{s \ge 0} \mathbf{k} \widetilde{H}_s \right) \longrightarrow A^o = A^o_{\overline{0}} \oplus A^o_{\overline{1}}, \tag{4.5}$$

where  $A^o := \bigoplus_{k \ge 0} (A^k)^*$  is the graded dual of the graded superspace  $A = \bigoplus_{k \ge 0} A^k$ , by

$$\zeta'(H_r)(a) \coloneqq \begin{cases} \zeta_{\overline{0}}(a) & (a \in A_{\overline{0}}^r) \\ 0 & (a \text{ is in the other homogeneous spaces}) \end{cases},$$
(4.6)

$$\zeta'(\widetilde{H}_s)(a) \coloneqq \begin{cases} \zeta_{\overline{1}}(a) & (a \in A_{\overline{1}}^{s+1}), \\ 0 & (a \text{ is in the other homogeneous spaces}) \end{cases}$$
(4.7)

Since  $sQSym^o = sNSym$  is the free associative superalgebra  $\mathbf{k}\langle H_1, H_2, \ldots; \widetilde{H}_0, \widetilde{H}_1, \ldots \rangle$ , the map  $\zeta'$  extends uniquely to an even morphism  $\Psi^* : sQSym^o \to A^o$  of superalgebras. Moreover, the definition of  $\zeta'$  implies that  $\Psi^*$  preserves the  $\mathbb{N}$ -grading. Since each homogeneous subspace  $sQSym^k$  is finite dimensional, there exists a unique graded even coalgebra morphism  $\Psi : A \to sQSym$  which is a graded dual of  $\Psi^*$ .

Then, for  $a \in A_{\overline{0}}^r \ (r \ge 1)$ , we have

$$(\zeta_Q \circ \Psi)(a) = H_r(\Psi(a)) = \Psi^*(H_r)(a) = \zeta_{\overline{0}}(a) = \zeta(a),$$

where the first equality follows from (4.4). Similarly, for  $a \in A_{\overline{1}}^{s+1}$   $(s \ge 0)$ , we have

$$(\zeta_Q \circ \Psi)(a) = \widetilde{H}_s(\Psi(a))\varepsilon = \Psi^*(\widetilde{H}_s)(a)\varepsilon = \zeta_{\overline{1}}(a)\varepsilon = \zeta(a)$$

Thus, we have a unique graded even coalgebra morphism  $\Psi$  such that  $\zeta = \zeta_Q \circ \Psi$ .

It remains to show that  $\Psi$  is an algebra morphism. Consider the following two diagrams:

$$A^{\otimes 2} \xrightarrow{\mu^{\otimes 2}} A \xrightarrow{\Psi} sQSym \qquad A^{\otimes 2} \xrightarrow{\Psi^{\otimes 2}} sQSym^{\otimes 2} \xrightarrow{\mu} sQSym$$

$$\downarrow^{\zeta} \xrightarrow{\zeta_Q} \downarrow^{\zeta_Q} \qquad \downarrow^{\zeta^{\otimes 2}} \xrightarrow{\zeta_Q} \downarrow^{\zeta^{\otimes 2}} \xrightarrow{\zeta_Q} k \oplus k\varepsilon$$

Since both diagrams commute and both  $\Psi \circ (\mu \otimes \mu)$  and  $\mu \circ (\Psi \otimes \Psi)$  are coalgebra morphisms, the uniqueness proven above implies that  $\Psi \circ (\mu \otimes \mu) = \mu \circ (\Psi \otimes \Psi)$ . The proof is now completed.

We have an explicit formula of the image  $\Psi(a)$  of  $a \in A$  in Theorem 4.2, which is a superspace extension of [GR, (7.1.3)].

**Proposition 4.3.** Let  $(A, \zeta)$  and  $\Psi: A \to \operatorname{sQSym}$  be the same as in Theorem 4.2. Then, for a homogeneous element  $a \in A^k$ ,  $k \ge 0$ , we have

$$\Psi(a) = \sum_{\alpha: \text{ dotted compositions of } k} \zeta_{\alpha}(a) M_{\alpha}, \qquad (4.8)$$

where, for a dotted composition  $\alpha = (\alpha_1, \ldots, \alpha_l)$ , the symbol  $M_{\alpha}$  denotes the monomial quasi-symmetric function (3.10), and the map  $\zeta_{\alpha}$  is the composite

$$\zeta_{\alpha} \colon A^{k} \xrightarrow{\Delta^{(l-1)}} A^{\otimes l} \xrightarrow{\pi_{\alpha}} A^{\alpha_{1}}_{\overline{\eta_{1}}} \otimes \cdots \otimes A^{\alpha_{l}}_{\overline{\eta_{l}}} \xrightarrow{\zeta^{\otimes l}} \mathbf{k}.$$

$$(4.9)$$

Here  $\eta(\alpha) = (\eta_1, \ldots, \eta_l)$  is given in (3.7), the dotted superscript  $\alpha_k = \dot{n}$  of  $A_{\overline{\eta_k}}^{\alpha_k}$  are replaced by n, and the symbol  $\pi_{\alpha}$  is the natural projection.

*Proof.* By the proof of Theorem 4.2, we have for  $a \in A^k$  that

$$\Psi(a) = \sum_{\alpha: \text{ dotted composition of } k} (H_{\alpha}, \Psi(a)) M_{\alpha},$$

and for each  $\alpha = (\alpha_1, \ldots, \alpha_l)$ , we have

 $(H_{\alpha}, \Psi(a)) = (\Psi^{*}(H_{\alpha}), a) = (\Psi^{*}(H_{(\alpha_{1})}) \cdots \Psi^{*}(H_{(\alpha_{l})}), a) = (\zeta^{\otimes l} \circ \pi_{\alpha}) (\Delta^{(l-1)}(a)) = \zeta_{\alpha}(a).$ 

Here we used the equality  $H_{\alpha} = H_{(\alpha_1)} \cdots H_{(\alpha_l)}$  [FLP19, (6.4)] in the second equality, and we used (4.3), (4.6) and (4.7) in the third equality.

Now we turn to the Hopf superalgebra  $\Lambda$  of symmetric functions in superspace (§ 3.1). It is a combinatorial Hopf superalgebra with supercharacter  $\zeta_S$  (4.2), and is cocommutative. We have the following characterization.

**Proposition 4.4.** The pair  $(\Lambda, \zeta_S)$  is a terminal object of the category of cocommutative combinatorial Hopf superalgebras. In other words, we have:

Let A be a cocommutative combinatorial Hopf superalgebra with supercharacter  $\zeta \colon A \to \mathbf{k}[\varepsilon]$ . Then, there is a unique graded even morphism  $\Psi \colon A \to \mathbf{\Lambda}$  of Hopf superalgebras such that  $\zeta = \zeta_S \circ \Psi$ , where  $\zeta_S \colon \mathbf{\Lambda} \to \mathbf{k}[\varepsilon]$ is the supercharacter (4.2).

Moreover, for a homogeneous element  $a \in A^k$ ,  $k \ge 0$ , we have

$$\Psi(a) = \sum_{\Lambda: \text{ superpartitions of } k} \zeta_{\Lambda}(a) m_{\Lambda}, \qquad (4.10)$$

where, for a superpartition  $\Lambda = (\Lambda_1, \ldots, \Lambda_m; \Lambda_{m+1}, \ldots, \Lambda_l)$ , the symbol  $m_{\Lambda}$  denotes the monomial symmetric function in superspace (3.4), and the symbol  $\zeta_{\Lambda}$  denotes the composite (4.9) for the dotted composition  $(\dot{\Lambda}_1, \ldots, \dot{\Lambda}_m, \Lambda_{m+1}, \ldots, \Lambda_l)$ .

*Proof.* It is enough to slightly modify the proof of Theorem 4.2. Note first that the graded dual  $A^o$  is a commutative superalgebra since A is now a cocommutative co(-super)algebra. Then, recalling the description  $\mathbf{\Lambda} \cong \mathbf{k}[e_1, e_2, \ldots; \tilde{e}_0, \tilde{e}_1, \ldots]$  (3.6), we modify (4.5) and define an even linear map

$$\begin{split} \zeta'' \colon \left(\bigoplus_{r \ge 1} \mathbf{k} e_r\right) \oplus \left(\bigoplus_{s \ge 0} \mathbf{k} \widetilde{e}_s\right) &\longrightarrow A^o = A_0^o \oplus A_{\overline{1}}^o, \\ \zeta''(e_r)(a) \coloneqq \begin{cases} \zeta_{\overline{0}}(a) & (a \in A_{\overline{0}}^r) \\ 0 & (a \text{ is in the other homogeneous spaces}) \end{cases}, \\ \zeta'(\widetilde{e}_s)(a) \coloneqq \begin{cases} \zeta_{\overline{1}}(a) & (a \in A_{\overline{1}}^{s+1}), \\ 0 & (a \text{ is in the other homogeneous spaces}) \end{cases}. \end{split}$$

Since  $\Lambda$  is the free commutative superalgebra generated by  $e_r$ 's and  $\tilde{e}_s$ 's, the map  $\zeta''$  extends uniquely to an even morphism  $\Psi^* \colon \Lambda \to A^o$  which preserves the  $\mathbb{N}$ -grading. Then, since  $\Lambda \cong \Lambda^o$ , i.e.,  $\Lambda$  is self-dual [FLP19, Proposition 4.8], we have a graded even morphism  $\Psi \colon A \to \Lambda^o \cong \Lambda$ . The remaining parts of the statement can be shown as in Theorem 4.2.

#### 5. A superspace analogue of chromatic symmetric functions

We consider a superspace analogue of the Hopf algebra of Stanley's chromatic symmetric functions [GR, §7.3].

Let us denote a finite graph by (V, E), where V is the vertex set and E is the edge set (both are assumed to be finite). Consider a triple G = (V, W, E) consisting of a finite graph (V, E) and a subset  $W \subset V$ , equipped

with an order of the connected components of (V, E). Such a triple can be regarded as a two-colored graph with white vertices W and black vertices  $V \setminus W$ . We call G connected if (V, E) is connected. We depict such a triple by white vertices W and black vertices  $V \setminus W$ , and inserting a bar | among the ordered connected components. For example, the diagram below shows a triple G = (V, W, E) with #V = 7, #W = 3 and the number of connected components being 2.

$$\bullet - \bullet - \circ | \bullet - \circ - \circ - \bullet$$

We also denote by  $\emptyset$  the empty triple.

Let  $\mathcal{G}'$  be the k-linear space spanned by [G] corresponding to each triple G, and  $\mathcal{C} \subset \mathcal{G}'$  be the subspace spanned by the elements

$$[G_1|G_2] - (-1)^{\#W_1 \cdot \#W_2}[G_2|G_1] \quad (G_i = (V_i, W_i, E_i), \text{connected}).$$

We define  $\mathcal{G}$  to be the quotient space

$$\mathcal{G} \coloneqq \mathcal{G}' / \mathcal{C},$$

and denote the class in  $\mathcal{G}$  of  $[G] \in \mathcal{G}'$  by the same symbol [G]. In particular, the following holds in  $\mathcal{G}$ .

$$[\circ|\circ] = 0 \tag{5.1}$$

The space  $\mathcal{G}$  has a double grading

$$\mathcal{G} = \bigoplus_{n,m \ge 0} \mathcal{G}_{n,m},$$

where  $\mathcal{G}_{n,m}$  is spanned by [G] with G = (V, W, E), #V = n and #W = m. It induces an N-grading:

$$\mathcal{G} = \bigoplus_{k \ge 0} \mathcal{G}^k, \quad \mathcal{G}^k \coloneqq \bigoplus_{n+m=k} \mathcal{G}_{n,m}.$$
(5.2)

Note that we have

$$\mathcal{G}^0 = \mathbf{k} = \mathbf{k}[\emptyset]. \tag{5.3}$$

We make  $\mathcal{G}$  into a superspace by defining the  $\mathbb{Z}/2\mathbb{Z}$ -grading to be  $m \mod 2$ :

$$\mathcal{G} = \mathcal{G}_{\overline{0}} \oplus, \mathcal{G}_{\overline{1}}, \quad \mathcal{G}_{\overline{p}} \coloneqq \bigoplus_{n \ge 0, m \mod 2 = \overline{p}} \mathcal{G}_{n,m} \quad (p = 0, 1)$$

With these definitions, we see that the subspace

$$\bigoplus_{n>0} \mathcal{G}_{n,0} \subset \mathcal{G} \tag{5.4}$$

is equal to the underlying linear space of the chromatic Hopf algebra in [GR, Definition 7.3.1].

The superspace  $\mathcal{G}$  has the following structure of a connected graded Hopf superalgebra, which is a natural super analogue of [GR, Definition 7.3.1]. The multiplication is

$$[G_1] \cdot [G_2] \coloneqq [G_1|G_2].$$

Then  $\mathcal{G}$  is a commutative superalgebra whose unit is  $[\emptyset]$ , and generated by  $[G_{\text{con}}]$  with connected triples  $G_{\text{con}}$ . Moreover, the subalgebra  $\bigoplus_{n>0} \mathcal{G}_{n,0} \subset \mathcal{G}$  is equal to the non-super case [GR, §7.3]. For example,

 $[\circ] \cdot [\circ] = 0$ 

holds by (5.1), but  $[\bullet|\bullet] \neq 0$  in  $\mathcal{G}$ .

To explain the comultiplication, we give some preliminary. For a triple G = (V, W, E) and a subset  $V' \subset V$ , we define a new triple  $G|_{V'} := (V', W', E')$  by  $W' := W \cap V'$ ,  $E' := \{e \in E \mid e = \{v_1, v_2\} \subset V'\}$  and the order of the connected components  $G|_{V'} = G_1|\cdots|G_l$ ,  $G_i = (V_i, W_i, E_i)$  being such that  $\#W_1 \leq \cdots \leq \#W_l$ (such an order is not unique, but the equivalence class in  $\mathcal{G}$  is uniquely determined). Now, for each connected triple  $G_{\text{con}}$ , we define

$$\Delta([G_{\rm con}]) \coloneqq \frac{1}{2} \sum_{(V_1, V_2): V_1 \sqcup V_2 = V} \left( [G_{\rm con}|_{V_1}] \otimes [G_{\rm con}|_{V_2}] + (-1)^{\#W_1 \# W_2} [G_{\rm con}|_{V_2}] \otimes [G_{\rm con}|_{V_1}] \right)$$

For an arbitrary triple G, we decompose  $G = G_1 | \cdots | G_l$  into connected  $G_i$ 's and define

$$\Delta([G]) \coloneqq \Delta([G_1]) \cdots \Delta([G_l])$$

For example, we have

$$\begin{split} \Delta([\bullet - \bullet]) &= [\bullet - \bullet] \otimes 1 + 2[\bullet] \otimes [\bullet] + 1 \otimes [\bullet - \bullet], \\ \Delta([\bullet - \circ]) &= [\bullet - \circ] \otimes 1 + [\bullet] \otimes [\circ] + [\circ] \otimes [\bullet] + 1 \otimes [\bullet - \circ], \\ \Delta([\circ - \circ]) &= [\circ - \circ] \otimes 1 + 1 \otimes [\circ - \circ], \\ \Delta([\circ - \circ - \bullet]) &= [\circ - \circ - \bullet] \otimes 1 + [\bullet] \otimes [\circ - \circ] + [\circ - \circ] \otimes [\bullet] + 1 \otimes [\circ - \circ - \bullet], \\ \Delta([\bullet - \circ - \bullet]) &= [\circ - \circ - \bullet] \otimes 1 + [\bullet] \otimes [\circ - \circ] + [\circ - \circ] \otimes [\bullet] + 1 \otimes [\circ - \circ - \bullet], \\ \Delta([\bullet - \circ - \bullet]) &= [\bullet - \circ - \bullet] \otimes 1 + 1 \otimes [\bullet - \circ - \bullet] + [\circ - \circ] \otimes [\bullet] + 1 \otimes [\circ - \circ - \bullet], \\ \Delta([\bullet - \circ - \bullet]) &= [\bullet - \circ - \bullet] \otimes 1 + 1 \otimes [\bullet - \circ - \bullet] + 3[\bullet - \circ - \bullet] \otimes [\bullet] \\ &+ 3[\bullet - \circ] \otimes [\bullet] + 3[\bullet] \otimes [\bullet - \circ] + [\circ] \otimes [\bullet] + [\bullet] + [\bullet] \otimes [\circ]. \end{split}$$

The map  $\Delta$  is coassociative and a superalgebra morphism, and we call it the comultiplication of  $\mathcal{G}$ . Moreover, it is cocommutative. Indeed, for any  $G = (V, W, E) \in \mathcal{G}$ , we have

$$\begin{aligned} \tau \circ \Delta([G]) &= \frac{1}{2} \sum_{(V_1, V_2): V_1 \sqcup V_2 = V} \left( \tau([G|_{V_1}] \otimes [G|_{V_2}]) + (-1)^{\#W_1 \# W_2} \tau([G|_{V_2}] \otimes [G|_{V_1}]) \right) \\ &= \frac{1}{2} \sum_{(V_1, V_2): V_1 \sqcup V_2 = V} \left( (-1)^{\#W_1 \# W_2} [G|_{V_2}] \otimes [G|_{V_1}] + [G|_{V_1}] \otimes [G|_{V_2}] \right) = \Delta([G]), \end{aligned}$$

where  $\tau$  denotes the switching: for  $G_i = (V_i, W_i, E_i), i = 1, 2,$ 

$$\tau\colon :\mathcal{G}\otimes\mathcal{G}\longrightarrow\mathcal{G}\otimes\mathcal{G},\quad [G_1]\otimes[G_2]\longmapsto(-1)^{\#W_1\cdot\#W_2}[G_2]\otimes[G_1].$$

The restriction of the comultiplication  $\Delta$  to the subspace  $\bigoplus_{n\geq 0} \mathcal{G}_{n,0} \subset \mathcal{G}$  (5.4) is equal to the non-super case [GR, Definition 7.3.1]. Indeed, for a triple  $G = (V, \emptyset, E)$ , we have

$$\Delta([G]) = \frac{1}{2} \sum_{(V_1, V_2): V_1 \sqcup V_2 = V} \left( [G|_{V_1}] \otimes [G|_{V_2}] + [G|_{V_2}] \otimes [G|_{V_1}] \right)$$
$$= \sum_{(V_1, V_2): V_1 \sqcup V_2 = V} [G|_{V_1}] \otimes [G|_{V_2}].$$
(5.5)

From the discussion so far,  $\mathcal{G}$  is a commutative and cocommutative bi(-super)algebra. By (5.2) and (5.3), it is connected graded. Then, by [FLP19, Theorem 2.1] and [GR, Proposition 1.4.16] (using the N-grading),  $\mathcal{G}$  has a antipode. The obtained Hopf superalgebra  $\mathcal{G}$  is connected graded by (5.2) and (5.3). We can also see that the subspace (5.4) is a connected graded Hopf subalgebra of  $\mathcal{G}$ , and coincides with the *chromatic Hopf algebra* in [GR, Definition 7.3.1].

# **Definition 5.1.** The connected graded Hopf superalgebra $\mathcal{G}$ is called the *chromatic Hopf superalgebra*.

Moreover, the Hopf superalgebra  $\mathcal{G}$  has a supercharacter  $\zeta_{ch}$ . To specify it, it is enough to give the value on each connected triple  $G_{con} = (V, W, E)$ . We set:

$$\zeta_{\rm ch}([G_{\rm con}]) \coloneqq \begin{cases} 1 & \text{if } E = \emptyset \text{ and } \#W = 0 \\ \varepsilon & \text{if } E = \emptyset \text{ and } \#W = 1 \\ 0 & \text{otherwise} \end{cases}$$
(5.6)

For an arbitrary triple  $G = G_1 | \cdots | G_l$  with connected  $G_i$ 's, we set  $\zeta_{ch}([G]) \coloneqq \zeta_{ch}([G_1]) \cdots \zeta_{ch}([G_l])$ , and obtain a supercharacter

$$\zeta_{\rm ch} \colon \mathcal{G} \longrightarrow \mathbf{k}[\varepsilon]. \tag{5.7}$$

If we restrict  $\zeta_{ch}$  to the Hopf subalgebra  $\bigoplus_{n\geq 0} \mathcal{G}_{n,0} \subset \mathcal{G}$ , then we recover the *edge-free character* in [GR, Definition 7.3.16, (7.3.4)].

In summary, we have:

**Theorem 5.2.** The chromatic Hopf superalgebra  $\mathcal{G}$  with the supercharacter  $\zeta_{ch}$  in (5.7) is a combinatorial Hopf superalgebra which is both commutative and cocommutative. It contains the chromatic Hopf algebra  $\bigoplus_{n>0} \mathcal{G}_{n,0}$  in [GR, Definition 7.3.1] as a Hopf subalgebra.

Since  $(\mathcal{G}, \zeta_{ch})$  is a cocommutative combinatorial Hopf superalgebra, by Proposition 4.4, there is a graded even morphism

$$\Psi_{\mathrm{ch}} \colon \mathcal{G} \longrightarrow \mathbf{\Lambda}$$

of Hopf superalgebras. Restricted to the subalgebra  $\bigoplus_{n\geq 0} \mathcal{G}_{n,0}$ , it recovers the map to the space of symmetric functions for the non-super case [GR, Definition 7.3.16]. Since the image of this non-super map consists of *Stanley's chromatic symmetric functions of graphs*, we may introduce:

**Definition 5.3.** For each triple G = (V, W, E), we call the image  $\Psi_{ch}([G]) \in \Lambda$  the chromatic symmetric function in superspace associated to G.

Let us give some examples of chromatic symmetric functions in superspace.

$$\begin{split} \Psi_{\rm ch}([\bullet]) &= e_1, \quad \Psi_{\rm ch}([\bullet - \bullet]) = 2e_2, \\ \Psi_{\rm ch}([\circ]) &= \widetilde{e}_0, \quad \Psi_{\rm ch}([\bullet - \circ]) = \widetilde{e}_1, \quad \Psi_{\rm ch}([\circ - \circ - \bullet]) = 0, \\ \Psi_{\rm ch}([\bullet - \circ \overbrace{\bullet}]) &= m_{(\dot{0},3)} + 6m_{(\dot{0},1,1,1)} + 3m_{(\dot{0},2,1)}, \end{split}$$

where we used the monomial symmetric function in superspace (3.4) and the elementary symmetric functions in superspace (3.5). From the definition (5.6), (5.7), we have:

**Lemma 5.4.** If a triple  $G = G_1 | \cdots | G_l$  has a connected component  $G_k = (V_k, W_k, E_k)$  with  $\#W_k > 1$ , then  $\Psi_{ch}([G]) = 0$ .

Let us give another example of a chromatic symmetric function in superspace, which can be regarded as a superspace analogue of [GR, Example 7.3.18]. We will denote the complete graph on l vertices by  $K_l$ .

**Proposition 5.5.** For  $n \ge 0$ , let  $K_{n+1,1} = (V, W, E)$  be a triple such that (V, E) is the complete graph  $K_{n+1}$  and #W = 1. Then, using the elementary symmetric function  $\tilde{e}_n$  (3.5), we have

$$\Psi_{\rm ch}([K_{n+1,1}]) = n!\,\widetilde{e}_n$$

*Proof.* Let us once admit the equality

$$\Delta^{(n)}([K_{n+1,1}]) = n! ([\circ] \otimes [\bullet]^{\otimes n} + [\bullet] \otimes [\circ] \otimes [\bullet]^{\otimes (n-1)} + \dots + [\bullet]^{\otimes n} \otimes [\circ]) + O,$$
(5.8)

where the last term  $O \in \mathcal{G}^{\otimes n+1}$  satisfies  $\zeta_{ch}^{\otimes (n+1)}(O) = 0$ . For a dotted partition  $\alpha$ , we denote by  $\zeta_{C,\alpha}$  the composite (4.9) for the supercharacter  $\zeta_{ch}$ . Then, for each  $i = 1, \ldots, n+1$ , we have

$$\zeta_{C,\alpha}([K_{n+1,1}]) = \begin{cases} n! & (\alpha = (1^{i-1}, \dot{0}, 1^{n+1-i})) \\ 0 & (\text{otherwise}) \end{cases}$$

Hence, by (4.10) in Proposition 4.4, we have

Δ

$$\Psi_{\rm ch}([K_{n+1,1}]) = n! \, m_{(0;1,\dots,1)} = n! \, \tilde{e}_n$$

We will show (5.8) in a more general form. For any  $2 \le m \le n$ , we have

$${}^{(m)}([K_{n+1,1}]) = n(n-1)\cdots(n-m+1)[K_{n-m+1,1}] \otimes [\bullet]^{\otimes m}$$
  
+  $n(n-1)\cdots(n-m+2)[K_{n-m+1}] \otimes \sum_{k=1}^{m} [\bullet]^{\otimes (k-1)} \otimes [\circ] \otimes [\bullet]^{\otimes (m-k)} + O$  (5.9)

where  $K_l = (V_l, \emptyset, E_l)$  is the triple corresponding to the ordinary complete graph on l vertices, and the term O satisfies  $\zeta_{ch}^{\otimes (m+1)}(O) = 0$ . Note that O contains terms of the form  $[G_0] \otimes [G_1] \otimes \cdots \otimes [G_m]$  such that for some  $k = 0, \ldots, m$  we have  $G_k = (V_k, W_k, E_k)$  with  $\#E_k > 0$ . If m = n, then (5.9) implies (5.8).

As a preliminary of proving (5.9), we calculate  $\Delta(K_l)$  using (5.5). We have

$$\Delta([K_l]) = [K_l] \otimes 1 + l[K_{l-1}] \otimes [\bullet] + O$$

where the term O means the same as (5.9). In particular, for l > 1, we have

$$\Delta([K_l]) = l[K_{l-1}] \otimes [\bullet] + O.$$
(5.10)

Similarly, we have

$$\Delta([K_{l+1,1}]) = \frac{1}{2} \sum_{\#V_1=1, \#V_2=l} ([K_{l+1,1}|_{V_1}] \otimes [K_{l+1,1}|_{V_2}] + [K_{l+1,1}|_{V_2}] \otimes [K_{l+1,1}|_{V_1}]) + \frac{1}{2} \sum_{\#V_1=l, \#V_2=1} ([K_{l+1,1}|_{V_1}] \otimes [K_{l+1,1}|_{V_2}] + [K_{l+1,1}|_{V_2}] \otimes [K_{l+1,1}|_{V_1}]) + \frac{1}{2} \sum_{\text{others}} ([K_{l+1,1}|_{V_1}] \otimes [K_{l+1,1}|_{V_2}] + (-1)^{\#W_1 \#W_2} [K_{l+1,1}|_{V_2}] \otimes [K_{l+1,1}|_{V_1}]) = [K_l] \otimes [\circ] + l[K_{l,1}] \otimes [\bullet] + O.$$
(5.11)

Now we prove (5.9) by induction on m. When m = 2, we have

$$(\Delta \otimes \operatorname{id}) \circ \Delta(K_{n+1,1}) = \Delta(K_n) \otimes [\circ] + n\Delta(K_{n,1}) \otimes [\bullet] + (\operatorname{others})$$
  
=  $n[K_{n-1}] \otimes [\bullet] \otimes [\circ] + n[K_{n-1}] \otimes [\circ] \otimes [\bullet] + n(n-1)[K_{n-1,1}] \otimes [\bullet] \otimes [\bullet] + O,$ 

and (5.9) actually holds. Next, we assume (5.9) for  $2 \le m \le n-1$ . Then for m+1 we have

$$\begin{aligned} \Delta^{(m+1)}([K_{n+1,1}]) &= (\Delta \otimes \operatorname{id}^{\otimes m}) \circ \Delta^{(m)}(K_{n+1,1}) \\ &= n(n-1)\cdots(n-m+1)\cdot\Delta([K_{n-m+1,1}]) \otimes [\bullet]^{\otimes m} \\ &+ n(n-1)\cdots(n-m+2)\cdot\Delta([K_{n-m+1}]) \otimes \sum_{k=1}^{m} [\bullet]^{\otimes (k-1)} \otimes [\circ] \otimes [\bullet]^{\otimes (m-k)} + O \\ &= n(n-1)\cdots(n-m+1)\cdot\left([K_{n-m}] \otimes [\circ] + (n-m)[K_{n-m,1}] \otimes [\bullet]\right) \otimes [\bullet]^{\otimes m} \\ &+ n(n-1)\cdots(n-m+2)\cdot(n-m+1)[K_{n-m}] \otimes [\bullet] \otimes \sum_{k=1}^{m} [\bullet]^{\otimes (k-1)} \otimes [\circ] \otimes [\bullet]^{\otimes (m-k)} + O \\ &= n(n-1)\cdots(n-m)[K_{n-m,1}] \otimes [\bullet]^{\otimes (m+1)} \\ &+ n(n-1)\cdots(n-m+1)[K_{n-m}] \otimes \sum_{k=1}^{m+1} [\bullet]^{\otimes (k-1)} \otimes [\circ] \otimes [\bullet]^{\otimes (m+1-k)} + O, \end{aligned}$$

where we used (5.10) and (5.11) in the third equality. Hence we have (5.9) for m + 1, and the proof is completed.

We close this note by giving a superspace extension of the expansion formula of chromatic symmetric functions [GR, Proposition 7.3.17]. For a triple G = (V, W, E), a proper coloring is a map  $f: V \to \{1, 2, ...\}$  such that  $f(v) \neq f(v')$  for any edge  $e = \{v, v'\} \in E$ .

**Proposition 5.6.** For a triple G = (V, W, E), the chromatic symmetric function in superspace  $\Psi_{ch}([G]) \in \Lambda$  is expressed as

$$\Psi_{\rm ch}([G]) = \sum_{f} \mathbf{x}_{f}, \quad \mathbf{x}_{f} \coloneqq \prod_{w \in W} \theta_{f(w)} \prod_{v \in V \setminus W} x_{f(v)}$$
(5.12)

where f runs over all the proper colorings of G.

*Proof.* For a triple G = (V, W, E) with  $[G] \in \mathcal{G}^k$ , the formula (4.8) says

$$\Psi_{\rm ch}([G]) = \sum_{\alpha} \zeta_{{\rm ch},\alpha}([G]) M_{\alpha},$$

where  $\alpha = (\alpha_1, \ldots, \alpha_l)$  runs over dotted compositions of k.

To interpret the coefficient  $\zeta_{ch,\alpha}([G])$ , we note that, for  $l \geq 2$ , the image of  $[G] \in \mathcal{G}$  under the iterated comultiplication  $\Delta^{(l-1)} \colon \mathcal{G} \to \mathcal{G}^{\otimes l}$  is expressed as

$$\Delta^{(l-1)}([G]) = \frac{1}{2^{l-1}} \sum_{\substack{(V_1, \dots, V_l): \\ V_1 \sqcup \dots \sqcup V_l = V}} \sum_{\sigma} \pm [G|_{V_{\sigma(1)}}] \otimes [G|_{V_{\sigma(2)}}] \otimes \dots \otimes [G|_{V_{\sigma(l)}}],$$

where  $\pm$  denotes some sign, and  $\sigma$  runs over some permutations of  $(1, \ldots, l)$ . Note that the number of such  $\sigma$  is  $2^{l-1}$ . The supercharacter  $\zeta_{ch}^{\otimes l}$  sends each summand to 1 or 0 depending on whether each  $E_i := \{\{v, v'\} \in E \mid v, v' \in V_{\sigma(i)}\}$  is empty or not, i.e., whether the assignment of color *i* to the vertices in  $V_{\sigma(i)}$  gives a proper coloring of *G*.

Hence, the coefficient  $\zeta_{ch,\alpha}([G])$  of  $M_{\alpha}$  in  $\Psi_{ch}([G])$  counts the proper colorings f such that  $\#f^{-1}(i) = \alpha_i + \eta_i$  and  $\#f^{-1}(i) \cap W = \eta_i$  for each color  $i \in \{1, 2, ...\}$ , where  $(\eta_1, ..., \eta_l) = \eta(\alpha)$  is given by (3.7). In other words, for a color i such that  $\alpha_i = \dot{s}$   $(s \ge 0)$ , the counted f satisfies  $\#f^{-1}(i) = s + 1$  and  $\#f^{-1}(i) \cap W = 1$ , and for i such that  $\alpha_i = r$   $(r \ge 1)$ , f satisfies  $\#f^{-1}(i) = r$  and  $\#f^{-1}(i) \cap W = 0$ . Since  $M_{\alpha} = \sum_f \mathbf{x}_f$  with f running over such proper colorings, we have the desired (5.12).

Acknowledgements. The authors would like to thank Professor Hiroaki Kanno for providing the impetus for this research. This work was financially supported by JST SPRING, Grant Number JPMJSP2125. M.H. and R.Y. would like to take this opportunity to thank the "THERS Make New Standards Program for the Next Generation Researchers."

### References

- [ABS06] M. Aguiar, N. Bergeron, F. Sottile, Combinatorial Hopf algebras and generalized Dehn-Sommerville relations, Compos. Math., 142 (2006), 1–30.
- [A+] L. Alarie-Vezina, O. Blondeau-Fournier, P. Desrosiers, L. Lapointe, P. Mathieu, Symmetric functions in superspace: a compendium of results and open problems, expository notes (2019), 38pp., arXiv:1903.07777.
- [B+12] O. Blondeau-Fournier, P. Desrosiers, L. Lapointe, P. Mathieu, Macdonald polynomials in superspace as eigenfunctions of commuting operators, J. Comb., 3 (2012), 495–561.
- [BE17] J. Brundan, A. P. Ellis, Monoidal supercategories, Comm. Math. Phys., 351 (2017), 1045–1089.
- [DM99] P. Deligne, J. W. Morgan, Notes on supersymmetry, in: Quantum Fields and Strings: A Course for Mathematicians, vol. 1, Princeton, NJ, 1996/1997; Amer. Math. Soc., Province RI, 1999, 41–97.
- [DLM03] P. Desrosiers, L. Lapointe, P. Mathieu, Jack polynomials in superspace, Comm. Math. Phys., 242 (2003), 331-360.
- [DLM06] P. Desrosiers, L. Lapointe, P. Mathieu, Classical symmetric functions in superspace, J. Algebraic Comb., 24 (2006), 209–238.
- [DLM07] P. Desrosiers, L. Lapointe, P. Mathieu, Orthogonality of Jack polynomials in superspace, Adv. Math., 212 (2007), 361–388.
- [FLP19] S. Fishel, L. Lapointe, M. E. Pinto, Hopf algebra structure of symmetric and quasisymmetric functions in superspace, J. Comn. Th., Ser. A 166 (2019), 144–170.
- [G84] I. Gessel, Multipartite P-partitions and inner products of skew Schur functions, Contemp. Math., 34 (1984) 289–301.
- [GR] D. Grinberg, V. Reiner, Hopf Algebras in Combinatorics, expository notes (2020), 282 pp., arXiv:1409.8356v7.
- [KV11] M. Kapranov, E. Vasserot, Supersymmetry and the formal loop space, Adv. Math. 227 (2011), 1078–1128.
- [M95] I. G. Macdonald, Symmetric functions and Hall polynomials, 2nd ed., Oxford Math. Monogr., Oxford Univ. Press, New York, 1995.
- [S95] R. Stanley, A symmetric function generalization of the chromatic polynomial of a graph, Adv. Math., 111 (1995), 166–194.
- [S99] R. Stanley, Enumerative Combinatorics, vol. 2, Cambridge Stud. Adv. Math., 62. Cambridge Univ. Press, Cambridge, 1999.