

EHRHART FUNCTIONS OF WEIGHTED LATTICE POINTS

JESÚS A. DE LOERA, CARLOS E. VALENCIA, RAFAEL H. VILLARREAL, AND CHENGYANG WANG

ABSTRACT. This paper studies three different ways to assign weights to the lattice points of a convex polytope and discusses the algebraic and combinatorial properties of the resulting weighted Ehrhart functions and their generating functions and associated rings. These will be called q -weighted, r -weighted, and s -weighted Ehrhart functions, respectively. The key questions we investigate are: *When are the weighted Ehrhart series rational functions, and which classical Ehrhart theory properties are preserved? And, when are the abstract formal power series the Hilbert series of Ehrhart rings of some polytope?* We prove generalizations about weighted Ehrhart h^* -coefficients of q -weighted Ehrhart series and show q - and s -weighted Ehrhart reciprocity theorems. Then, we show the q - and r -weighted Ehrhart rings are the (classical) Ehrhart rings of weight lifting polytopes.

Dedicated to Richard P. Stanley, on the occasion of his 80th birthday and gratefully celebrating his contributions in the intersection of commutative algebra and combinatorics.

1. INTRODUCTION

In our setup, we work with a convex rational polytope \mathcal{P} in \mathbb{R}^s and $S = K[t_1^{\pm 1}, \dots, t_s^{\pm 1}]$ is a Laurent polynomial ring over a field K . Its monomials are abbreviated by $t^a := t_1^{a_1} \cdots t_s^{a_s}$, $a = (a_1, \dots, a_s)$ in \mathbb{Z}^s . We think of monomials as lattice points and rely on their lattice geometry for our analysis.

Ehrhart functions record the number of lattice points inside all integral dilations of a convex polytope. They play an important role in algebraic combinatorics, commutative algebra, and convex geometry (see, for instance, [9, 11, 35]). In recent years, there have been several natural suggestions on how to extend Ehrhart functions by adding weights to the lattice points. Traditionally, the lattice points have a constant weight of 1, and the Ehrhart function is just counting. This paper studies three different (but related) ways to assign weights to the lattice points, as well as the properties of the resulting weighted Ehrhart functions, generating functions, and associated rings.

Here are the key definitions for the rest of the paper:

Definition 1.1. We say a function $w : \mathbb{Z}^s \rightarrow \mathbb{Z}$ is a *weight* of degree e if for all vectors $v \in \mathbb{Z}^s$ and any scalar k , $w(kv) = h(k)w(v)$ where $h(k)$ is a univariate polynomial of degree e . If $\mathbb{Z} \subset K$ and w is extended to $w : K^s \rightarrow K$, we also call w a weight.

Examples of weight functions appear everywhere; here are a few we consider: w is a constant function, w is a homogeneous linear form, $w = w(t_1, \dots, t_s)$ is a homogeneous polynomial, and w is a quasi-polynomial.

2020 *Mathematics Subject Classification.* Primary 52B20; Secondary 13F20, 05A15, 90C10.

Key words and phrases. Lattice polytopes, weights on lattice points, Ehrhart functions and reciprocity laws, Stanley's positivity and monotonicity theorems, Ehrhart rings, h^* -vectors, semigroup rings.

The first and fourth authors were supported by NSF grant DMS-2434665, and the second and third authors were supported by SNII, México.

With weight functions in hand, we can put weights on the lattice points (we count them with a weight) or, equivalently, to the monomials of the Ehrhart ring. We remark we were not the first to introduce or study weights. Some prior history follows after the definitions.

In this paper the *cone over* \mathcal{P} , denoted $\text{cone}(\mathcal{P})$ or $C_{\mathcal{P}}$, is the polyhedral cone in \mathbb{R}^{s+1} generated by

$$(1.1) \quad B_{\mathcal{P}} := \{(x, 1) : x \in \mathcal{P} \cap \mathbb{Z}^s\} \subset \mathbb{Z}^{s+1}.$$

Definition 1.2. Let \mathcal{P} be a polytope. For $i = 1, \dots, p$, let $w_i: K^s \rightarrow K$, $a \mapsto w_i(a)$, be a weight such that $w_i(\mathbb{Z}^s) \subset \mathbb{Z}$, and let $w: K^s \rightarrow K^p$ be the function $w = (w_1, \dots, w_p)$. We now define weighted Ehrhart functions using the w_i 's and new auxiliary variables q_1, \dots, q_p :

- (1) The *q-weighted multivariate Ehrhart function* and *q-weighted multivariate Ehrhart series* of \mathcal{P} relative to w , denoted $E_{\mathcal{P}}^{q,w}$ and $F_{\mathcal{P}}^{q,w}$ respectively, are given by

$$E_{\mathcal{P}}^{q,w}(q, t, n) := \sum_{(a,n) \in \text{cone}(\mathcal{P}) \cap \mathbb{Z}^{s+1}} q^{w(a)} t^a, \quad F_{\mathcal{P}}^{q,w}(q, t, x) := \sum_{n=0}^{\infty} E_{\mathcal{P}}^{q,w}(q, t, n) x^n,$$

where $q^{w(a)} = q_1^{w_1(a)} \cdots q_p^{w_p(a)}$.

- (2) The *r-weighted multivariate Ehrhart function* and *r-weighted multivariate Ehrhart series* of \mathcal{P} relative to w , denoted $E_{\mathcal{P}}^{r,w}$ and $F_{\mathcal{P}}^{r,w}$ respectively, are given by

$$E_{\mathcal{P}}^{r,w}(q, t, n) := \sum_{\substack{(a,n) \in \text{cone}(\mathcal{P}) \cap \mathbb{Z}^{s+1} \\ 0 \leq b_i \leq w_i(a) \ \forall i}} t^a q_1^{b_1} \cdots q_p^{b_p}, \quad F_{\mathcal{P}}^{r,w}(q, t, x) := \sum_{n=0}^{\infty} E_{\mathcal{P}}^{r,w}(q, t, n) x^n.$$

Finally, when there is only one single weight function $w: K^s \rightarrow K$, $a \mapsto w(a)$. We define the last kind of weighting using w directly:

- (3) The *s-weighted Ehrhart function* and the *s-weighted multivariate Ehrhart series* of \mathcal{P} relative to w , denoted $E_{\mathcal{P}}^{s,w}$ and $F_{\mathcal{P}}^{s,w}$ respectively, are given by

$$E_{\mathcal{P}}^{s,w}(t, n) := \sum_{(a,n) \in \text{cone}(\mathcal{P}) \cap \mathbb{Z}^{s+1}} w(a) t^a, \quad \text{and} \quad F_{\mathcal{P}}^{s,w}(t, x) := \sum_{n=0}^{\infty} E_{\mathcal{P}}^{s,w}(t, n) x^n,$$

Remark 1.3. Of course, the above definitions are very general definitions, but note we recover old versions directly via substitutions and specializations. For the q -weighted multivariate Ehrhart function and series when we set $t_j = 1$:

$$E_{\mathcal{P}}^{q,w}(q, 1, n) = \sum_{(a,n) \in \text{cone}(\mathcal{P}) \cap \mathbb{Z}^{s+1}} q^{w(a)}, \quad F_{\mathcal{P}}^{q,w}(q, 1, x) = \sum_{n=0}^{\infty} E_{\mathcal{P}}^{q,w}(q, 1, n) x^n.$$

This type of weighting was investigated [19] (but only for w_i linear and $p = 1$). Note further that when we set $q_j = 1$, we recover the traditional Ehrhart series. Regarding the s -weighting definition, if we set $t_j = 1$, we recover the univariate versions introduced in [12]. $E_{\mathcal{P}}^{s,w}(n) := \sum_{(a,n) \in \text{cone}(\mathcal{P}) \cap \mathbb{Z}^{s+1}} w(a)$, and $F_{\mathcal{P}}^{s,w}(x) := \sum_{n=0}^{\infty} E_{\mathcal{P}}^{s,w}(n) x^n$. The definition of r -weighting is new, but as we will see, it connects the other two definitions.

The questions we discuss in this paper are: *When are the weighted Ehrhart series rational functions? What classical properties are preserved? And, when are these abstract formal power series the Hilbert series of Ehrhart rings of some polytopes?*

Prior work on weighted Ehrhart theory: The theory of weighted Ehrhart functions and weighted Ehrhart series has been developed in several papers since at least the 1990's. See [5, 6, 12, 16, 17, 19] and references therein, as well as the very recent papers [2, 28]. Brion and Vergne [12] presented in 1997 a generalization of Ehrhart's theorem in the context of Euler-Maclaurin formulas where the points are counted with “ s -weights” given by a polynomial function w , i.e., $E_{\mathcal{P}}^{s,w}(n) = \sum_{a \in n\mathcal{P} \cap \mathbb{Z}^s} w(a)$. Since then, it is known that when w is a polynomial with integer coefficients and \mathcal{P} is a lattice polytope, the s -weighted Ehrhart function is a polynomial and the s -weighted Ehrhart series is the power series expansion of a rational function. See [12] and Corollary 5.2. Here, the s -weighted Ehrhart functions will be studied in Section 5.

To the best of our knowledge, research on q -weightings is less expansive, and the first paper on the subject comes from Chapoton, who introduced the q -weighted Ehrhart functions [19] in one variable thinking of q -analogues of the Ehrhart counting methods. Chapoton only looked at the q -weights of degree 1, thus when w is linear. The recent paper of Reiner and Rhoades [28] also looks at q -analogues of the Ehrhart series that come from deformations of the ring of polynomials modulo the ideal of polynomials that vanish at the lattice points of a polytope. Like us, they care about constructing rings whose Hilbert series is exactly their new Ehrhart series. We seem to be the first to introduce r -weightings.

In later sections, we revisit the classical results in Ehrhart theory and see some that extend to weighted multivariate Ehrhart series (Theorems 2.5 and 2.9).

Ehrhart rings and \mathbf{h}^* -vectors. There is a rich relationship between Ehrhart functions and commutative algebra, which we briefly recall here (see [28, 31, 35, 37] and references therein for all details). Let \mathcal{P} be a rational polytope in \mathbb{R}^s of dimension d and let $B_{\mathcal{P}}$ be as in Eq. (1.1). We are interested in the affine semigroup $\mathbb{R}_+ B_{\mathcal{P}} \cap \mathbb{Z}^{s+1}$, where $\mathbb{R}_+ B_{\mathcal{P}}$ denotes the polyhedral cone generated by $B_{\mathcal{P}}$. We assume \mathcal{P} is always a rational polytope to be sure this semigroup is finitely generated. The cone $\mathbb{R}_+ B_{\mathcal{P}}$ is called the *cone over \mathcal{P}* and is denoted by $\text{cone}(\mathcal{P})$. The resulting monomial algebras are often called *toric algebras* and are core to the theory of toric varieties [13, 20, 36]; their generating sets come from Hilbert bases, and their Hilbert functions are investigated by several researchers [13, 14, 31, 35].

On the algebraic side, when \mathcal{P} is a lattice polytope (i.e., its vertices are lattice points), the *Ehrhart ring* of \mathcal{P} , denoted $A(\mathcal{P})$, is the monomial ring of $S[z]$ given by

$$(1.2) \quad A(\mathcal{P}) := K[t^a z^n \mid a \in n\mathcal{P} \cap \mathbb{Z}^s] = K[t^a z^n \mid (a, n) \in \mathbb{R}_+ B_{\mathcal{P}} \cap \mathbb{Z}^{s+1}] \subset S[z],$$

where z is a *grading* variable and $n\mathcal{P}$ denotes the n -th dilation of \mathcal{P} . The Ehrhart ring $A(\mathcal{P})$ has a natural \mathbb{Z} -grading by

$$A(\mathcal{P}) = \bigoplus_{n=0}^{\infty} A(\mathcal{P})_n,$$

where $t^a z^n \in A(\mathcal{P})_n$ if and only if $a \in n\mathcal{P} \cap \mathbb{Z}^s$, which is exactly the n -level of $\text{cone}(\mathcal{P})$.

We homogenize the set of vertices $\{v_i\}_{i=1}^m$ of \mathcal{P} with $\mathcal{B} := \{(v_i, 1)\}_{i=1}^m$. According to [37, Theorem 9.3.6], one has

$$A(\mathcal{P}) = K[\mathbb{Z}^{s+1} \cap \mathbb{R}_+ \mathcal{B}] = K[t^a z^n \mid (a, n) \in \mathbb{Z}^{s+1} \cap \mathbb{R}_+ \mathcal{B}],$$

where $\mathbb{R}_+ \mathcal{B}$ is the cone generated by \mathcal{B} . Let $\mathcal{H}_{\mathcal{P}} = \{(c_1, d_1), \dots, (c_r, d_r)\}$, $c_i \in \mathbb{N}^s$, $d_i \in \mathbb{N}$, be a Hilbert basis of $\mathbb{R}_+ \mathcal{B}$, that is, $\mathcal{H}_{\mathcal{P}}$ is a finite subset of \mathbb{N}^{s+1} and $\mathbb{R}_+ \mathcal{B} \cap \mathbb{Z}^{s+1} = \mathbb{N}\mathcal{H}_{\mathcal{P}}$, where $\mathbb{N}\mathcal{H}_{\mathcal{P}}$ is the semigroup spanned by $\mathcal{H}_{\mathcal{P}}$. If $\mathcal{H}_{\mathcal{P}}$ is minimal (with respect to inclusion), then $\mathcal{H}_{\mathcal{P}}$ is unique [29, Proposition 1]. The program *Normaliz* [15] can be used to compute the Hilbert

basis of $\mathbb{R}_+\mathcal{B}$. The Ehrhart ring $A(\mathcal{P})$ of \mathcal{P} is equal to the monomial subring

$$K[\mathbb{N}\mathcal{H}_{\mathcal{P}}] = K[t^{c_1}z^{d_1}, \dots, t^{c_r}z^{d_r}],$$

that is, $A(\mathcal{P})$ is the *semigroup ring* of the affine semigroup $\mathbb{N}\mathcal{H}_{\mathcal{P}}$.

It is well-known that the *Ehrhart function* $E_{\mathcal{P}}: \mathbb{N} \rightarrow \mathbb{N}$, $n \mapsto |n\mathcal{P} \cap \mathbb{Z}^s|$, of \mathcal{P} is a *quasipolynomial* of degree $d = \dim(\mathcal{P})$ whose generating function $F_{\mathcal{P}}$ is a rational function, and a nice *reciprocity law* is satisfied between the Ehrhart function of \mathcal{P} and that of its relative interior. When in addition \mathcal{P} is a lattice polytope, the quasipolynomial becomes, in fact, a polynomial whose leading coefficient is the relative volume $\text{vol}(\mathcal{P})$ of \mathcal{P} [11, 22, 23], and the generating function $F_{\mathcal{P}}$ of $E_{\mathcal{P}}$ is a rational function,

$$F_{\mathcal{P}}(x) := \sum_{n=0}^{\infty} E_{\mathcal{P}}(n)x^n = \frac{h^*(x)}{(1-x)^{d+1}},$$

where its numerator $h^*(x)$ is a polynomial with nonnegative integer coefficients of degree at most d [32, Theorem 2.1] and is called the *h^* -polynomial* of \mathcal{P} . The vector formed with the coefficients of $h^*(x)$ is called the *h^* -vector* of \mathcal{P} and is denoted by $h^*(\mathcal{P})$. See [11, 13, 22, 24, 35, 37] and references therein for all details on these facts.

Our Contributions: Our contributions have two themes: first, the combinatorics of the generating functions and the weighted Ehrhart series, and second, the construction of graded rings whose Hilbert functions are precisely the weighted Ehrhart functions.

Positivity of h^* -coefficients for q -weighted Ehrhart series. Two famous results in Ehrhart theory due to Stanley are the *positivity* theorem showing that the numerator of the rational function representing the Ehrhart series of a lattice polytope is a polynomial $h^*(x)$ with nonnegative coefficients [32], and the *monotonicity* theorem for h^* -vectors showing that given lattice polytopes \mathcal{P}_1 and \mathcal{P}_2 with $\mathcal{P}_1 \subset \mathcal{P}_2$, then their h^* -vectors satisfy $h^*(\mathcal{P}_1) \leq h^*(\mathcal{P}_2)$ component-wise [34]. Could it be that the weighted Ehrhart series we consider are rational functions and that Stanley's results extend to the weighted Ehrhart series? We give answers to these questions.

It was shown recently that Stanley's positivity and monotonicity theorems [32, 34] continue to hold for s -weighted Ehrhart series of rational polytopes where lattice points are counted with homogeneous polynomial weights that are sums of products of linear forms which are nonnegative on the polytope [3].

Inspired by the fact that the numerator of the Ehrhart series always has nonnegative coefficients, Chapoton investigated in [19] the numerator of univariate q -weighted Ehrhart series and noticed that such nonnegativity result does not hold anymore. In Section 2, we take a deeper look and give a sufficient, but not necessary, geometric condition to show when the nonnegativity result holds.

Definition 1.4. For a lattice polytope $\mathcal{P} \subset \mathbb{R}^s$ and p linear functions $w_i: \mathbb{R}^s \rightarrow \mathbb{R}$ ($i = 1, \dots, p$), we say a triangulation \mathcal{T} of the polytope \mathcal{P} is (w_1, \dots, w_p) -compatible if every simplex $\Delta \in \mathcal{T}$ satisfies that the multiset of weight vectors on the vertices of Δ is identical.

We come to one of our positivity results.

Theorem 2.4. *If w_i 's are linear weights and a lattice polytope \mathcal{P} has a (w_1, \dots, w_p) -compatible triangulation \mathcal{T} , then the numerator of the rational form of its q -weighted Ehrhart series has positive coefficients.*

The condition of Theorem 2.4 is only a sufficient condition for positivity.

Example 1.5. Consider the polytope $\mathcal{P} = \text{conv}((0, 0, 0), (1, 0, 0), (1, 1, 0), (1, 1, 1), (2, 1, 1))$ and the linear function $w(y_1, y_2, y_3) = y_1 + y_2 + y_3$, we find that its q -weighted Ehrhart series is:

$$\frac{1 + q^2x}{(1-x)(1-qx)(1-q^3x)(1-q^4x)},$$

and the polytope \mathcal{P} does not have any w -compatible triangulation.

Reciprocity for weighted Ehrhart series. Another famous result in Ehrhart theory, also due to Stanley, is the *multivariate law of reciprocity*, an elegant functional relation showing that when inverting each variable in the integer point transform of a rational cone [9], [11, p. 60], one obtains (up to sign) the integer point transform for the interior of the cone [30]. The classical Ehrhart-Macdonald law of reciprocity relating the Ehrhart function and the interior Ehrhart function follows as a specialization [11, p. 84]. Inspired by this and Chapoton's reciprocity result for q -weighted Ehrhart series for only one weight, in Section 2, we investigate the reciprocity properties for other weighted Ehrhart series we considered.

Definition 1.6. For a polytope \mathcal{P} , we define:

- (1) the *interior q -weighted multivariate Ehrhart series* as

$$F_{\mathcal{P}^\circ}^{q, w_1, \dots, w_p}(q, t, x) := \sum_{(a, n) \in \text{cone}(\mathcal{P})^\circ \cap \mathbb{Z}^{s+1}} q_1^{-w_1(-a)} \dots q_p^{-w_p(-a)} t^a x^n,$$

- (2) the *interior q -weighted Ehrhart series* as

$$F_{\mathcal{P}^\circ}^{q, w_1, \dots, w_p}(q, 1, x) := \sum_{(a, n) \in \text{cone}(\mathcal{P})^\circ \cap \mathbb{Z}^{s+1}} q_1^{-w_1(-a)} \dots q_p^{-w_p(-a)} x^n, \text{ and}$$

- (3) the *interior s -weighted Ehrhart series* as

$$F_{\mathcal{P}^\circ}^{s, w}(t, x) := \sum_{(a, n) \in \text{cone}(\mathcal{P})^\circ \cap \mathbb{Z}^{s+1}} w(-a) t^a x^n.$$

We show the following reciprocity law for the Ehrhart series in the case of the multivariate q -weighting of the Ehrhart series.

Theorem 2.5. *If w_1, \dots, w_p are linear weights and \mathcal{P} is a polytope whose cone has dimension $s + 1$, then the q -weighted multivariate Ehrhart series satisfies the reciprocity property, i.e.,*

$$F_{\mathcal{P}}^{q, w_1, \dots, w_p}(q^{-1}, t^{-1}, x^{-1}) = (-1)^{s+1} F_{\mathcal{P}^\circ}^{q, w_1, \dots, w_p}(q, t, x).$$

As a consequence, $F_{\mathcal{P}}^{q, w_1, \dots, w_p}(q^{-1}, t^{-1}, x^{-1})$ can be represented by interior q -weighted multivariate Ehrhart series (Corollary 2.6).

We also show the following reciprocity law in the case of the s -weighting of Ehrhart functions.

Theorem 2.9. *Let $\mathcal{P} \subset \mathbb{R}^s$ be a rational polytope and $h(a) = \prod_{i=1}^{s+1} \sum_{j=1}^{k_i} P_{ij}(a_i) \gamma_{ij}^{a_i}$, where the P_{ij} 's are polynomials and γ_{ij} are nonzero complex numbers. Then*

- (1) $F_{\mathcal{P}}^{s, h}(t, x)$ and $F_{\mathcal{P}^\circ}^{s, h}(t, x)$ are rational s -weighted multivariate Ehrhart series, and
 (2) they satisfy the reciprocity relation,

$$F_{\mathcal{P}}^{s, h}(t^{-1}, x^{-1}) = (-1)^{s+1} F_{\mathcal{P}^\circ}^{s, h}(t, x).$$

Weighted Ehrhart rings.

Definition 1.7. Let \mathcal{P} be a lattice polytope in \mathbb{R}^s and let $w_i: \mathbb{R}^s \rightarrow \mathbb{R}$ ($i = 1, \dots, p$) be p weight functions such that $w_i(e_j) \in \mathbb{N}$ for $j = 1, \dots, s$. We define *weighted Ehrhart-rings*:

- (1) The q -weighted Ehrhart ring of \mathcal{P} , denoted $A_q^w(\mathcal{P})$, is a monomial subring given by

$$A_q^w(\mathcal{P}) := K[t^a q^{w(a)} z^n \mid (a, n) \in \text{cone}(\mathcal{P}) \cap \mathbb{Z}^{s+1}] \subset S[q, z],$$

where $w = (w_1, \dots, w_p)$ and $q^{w(a)} = q_1^{w_1(a)} \dots q_p^{w_p(a)}$. We allow the case $t_j = 1$.

- (2) The r -weighted Ehrhart ring of \mathcal{P} , denoted $A_r^w(\mathcal{P})$, is a monomial subring given by

$$A_r^w(\mathcal{P}) := K[t^a q_1^{b_1} \dots q_p^{b_p} z^n \mid (a, n) \in \text{cone}(\mathcal{P}) \cap \mathbb{Z}^{s+1}, 0 \leq b_i \leq w_i(a) \text{ for all } i].$$

If w_1, \dots, w_p are linear functions, then the weighted Ehrhart rings become finitely generated graded K -algebras. The graded structures of $A_r^w(\mathcal{P})$ and $A_q^w(\mathcal{P})$ are given by considering all monomials containing z^n , $n \geq 0$ (Eq. (1.4), Theorem 1.8(b)). Let $\mathcal{P} = \text{conv}(v_1, \dots, v_m)$ be a lattice polytope. Note that by making $w_i = 0$ for all i , $A_q^w(\mathcal{P})$ and $A_r^w(\mathcal{P})$ are equal to the classical Ehrhart ring $A(\mathcal{P})$ of \mathcal{P} .

Now, we describe the q - and r -weighted Ehrhart rings using classical Ehrhart rings. The strategy is to construct new polytopes \mathcal{P}^w and \mathcal{P}_w (see Eqs. (1.3) and (1.6)) in a higher dimension such that the q - and r -weighted Ehrhart rings and the classic Ehrhart rings are correspondent, that is, $A_q^w(\mathcal{P}) = A(\mathcal{P}^w)$ and $A_r^w(\mathcal{P}) = A(\mathcal{P}_w)$. We will call them *q- and r-weight lifting polytopes*, respectively. One can also denote \mathcal{P}^w and \mathcal{P}_w by $\mathcal{P}^{q,w}$ and $\mathcal{P}^{r,w}$, respectively, to match the left hand side of the two equalities. The same type of r -weight lifting polytopes are studied in [21, Theorem 1.1]. The difference in representing the r -weight lifting polytope is that they use linear inequalities, and we use vertices.

q-weight lifting polytopes and q-weighted Ehrhart rings. Let $w_i: \mathbb{R}^s \rightarrow \mathbb{R}$, $i = 1, \dots, p$, be linear functions such that $w_i(e_j) \in \mathbb{N}$ for all i, j , let $w: \mathbb{R}^s \rightarrow \mathbb{R}^p$ be the linear function $w = (w_1, \dots, w_p)$, and let $\mathcal{P} = \text{conv}(\{v_i\}_{i=1}^m)$ be a lattice polytope in \mathbb{R}_+^s . In Section 3, we prove that $A_q^w(\mathcal{P})$ is the Ehrhart ring of the *q-weight lifting polytope*:

$$(1.3) \quad \mathcal{P}^w := \text{conv}(\{(v_i, w(v_i))\}_{i=1}^m) \subset \mathbb{R}^{s+p},$$

that is, $A_q^w(\mathcal{P}) = A(\mathcal{P}^w)$ and $E_{\mathcal{P}}(n) = |n\mathcal{P} \cap \mathbb{Z}^s| = |n\mathcal{P}^w \cap \mathbb{Z}^{s+p}| = E_{\mathcal{P}^w}(n)$. We have the following theorem.

Theorem 1.8. *Let $C_{\mathcal{P}}$ be the cone over $\mathcal{P} = \text{conv}(\{v_i\}_{i=1}^m)$. The following hold.*

- (a) $A_q^w(\mathcal{P}) = K[t^a q^{w(a)} z^n \mid (a, n) \in C_{\mathcal{P}} \cap \mathbb{Z}^{s+1}] = K[t^a q^{w(a)} z^n \mid a \in n\mathcal{P} \cap \mathbb{Z}^s]$.
 (b) $A_q^w(\mathcal{P}) = \bigoplus_{n=0}^{\infty} A_q^w(\mathcal{P})_n$ is a graded K -algebra whose n -th graded component is

$$A_q^w(\mathcal{P})_n := \sum_{a \in n\mathcal{P} \cap \mathbb{Z}^s} K t^a q^{w(a)} z^n \text{ and } \dim_K(A_q^w(\mathcal{P})_n) = E_{\mathcal{P}}(n), \forall n \geq 0.$$

- (c) $A(\mathcal{P}^w) = A_q^w(\mathcal{P})$ and $E_{\mathcal{P}}(n) = E_{\mathcal{P}^w}(n)$ for $n \geq 1$.

- (d) There exist $\mathcal{H}_{\mathcal{P}} = \{(c_i, d_i)\}_{i=1}^r$, $c_i \in \mathbb{Z}^s$, $d_i \in \mathbb{Z}$ such that $C_{\mathcal{P}} \cap \mathbb{Z}^{s+1} = \mathbb{N}\mathcal{H}_{\mathcal{P}}$,

$$A(\mathcal{P}) = K[\mathbb{N}\mathcal{H}_{\mathcal{P}}], \text{ and } A(\mathcal{P}^w) = K[t^a q^{w(a)} z^n \mid a \in n\mathcal{P} \cap \mathbb{Z}^s] = K[\{t^{c_i} q^{w(c_i)} z^{d_i}\}_{i=1}^r].$$

- (e) The q -weighted multivariate Ehrhart series $F_{\mathcal{P}}^{q,w}(t, q, x)$ of \mathcal{P} is a rational function.

From this, we get that $A_q^w(\mathcal{P})$ is finitely generated as a K -algebra and that the dimension of \mathcal{P}^w is the dimension of \mathcal{P} . The rationality of $F_{\mathcal{P}}^{q,w}$ will follow noticing that the integer-point transform of $\text{cone}(\mathcal{P}^w)$, in the sense of [9, p. 60], is equal to $F_{\mathcal{P}}^{q,w}$ and applying [11, Corollary 3.7].

As we will see in Proposition 2.2, the hypothesis that the w_i 's are linear is essential to prove that $A_q^w(\mathcal{P})$ is finitely generated.

r-weight lifting polytopes and r-weighted Ehrhart rings. In Section 4, we show that the r -weighted Ehrhart ring $A_r^w(\mathcal{P})$ of \mathcal{P} relative to w is the Ehrhart ring of the r -weight lifting lattice polytope \mathcal{P}_w defined below in Eq. (1.6) (Theorem 4.6).

The r -weighted Ehrhart ring of \mathcal{P} has the same graded ring structure as $A(\mathcal{P})$. The ring $A_r^w(\mathcal{P}) = \bigoplus_{n=0}^{\infty} A_r^w(\mathcal{P})_n$ is a graded K -algebra whose n -th graded component is

$$(1.4) \quad A_r^w(\mathcal{P})_n = \sum_{\substack{a \in n\mathcal{P} \cap \mathbb{Z}^s \\ 0 \leq b_i \leq w_i(a)}} K t^a q_1^{b_1} \cdots q_p^{b_p} z^n$$

and, by Lemma 4.5, the dimension of $A_r^w(\mathcal{P})_n$ as a K -vector space is

$$\dim_K(A_r^w(\mathcal{P})_n) = E_{\mathcal{P}}^{s, \prod_{i=1}^p (w_i+1)}(n),$$

where $E_{\mathcal{P}}^{s,f}$ is the s -weighted Ehrhart function of $f = \prod_{i=1}^p (w_i + 1)$.

As before, we need to construct from \mathcal{P} and the weights a new polytope \mathcal{P}_w . Denote by

$$(1.5) \quad \mathcal{G} := \{(v_i, 0) + w_{j_1}(v_i)e_{s+j_1} + \cdots + w_{j_k}(v_i)e_{s+j_k} \mid 1 \leq i \leq m, 1 \leq j_1 < \cdots < j_k \leq p\},$$

we allow $k = 0$ by setting $w_{j_1}(v_i)e_{s+j_1} + \cdots + w_{j_k}(v_i)e_{s+j_k} := 0$ if $k = 0$,

that is, if $k = 0$, $(v_i, 0) \in \mathcal{G}$ for $i = 1, \dots, m$,

$$(1.6) \quad \mathcal{P}_w := \text{conv}(\mathcal{G}), \quad \Gamma := B_{\mathcal{G}} = \{(\alpha, 1) \mid \alpha \in \mathcal{G}\},$$

$$(1.7) \quad \mathcal{H}_{\mathcal{P}_w} := \{(c_i, b, d_i) \mid b = (b_1, \dots, b_p) \in \mathbb{N}^p, 0 \leq b_j \leq w_j(c_i) \forall j, c_i \in d_i\mathcal{P}, 1 \leq i \leq r\}.$$

If w is a monomial of S , the following result shows that $A_r^w(\mathcal{P})$ is the Ehrhart ring $A(\mathcal{P}_w)$ of \mathcal{P}_w . Ehrhart rings are normal [37] and then, by a classical theorem of Hochster [26], the ring $A(\mathcal{P}_w)$ is Cohen–Macaulay. We come to one of our main results about r -weighted Ehrhart rings:

Theorem 4.6. *If w_1, \dots, w_p are linear functions and $w_i(e_j) \in \mathbb{N}$ for all i, j , then*

- (a) $\mathbb{R}_+\Gamma \cap \mathbb{Z}^{s+p+1} = \mathbb{N}\mathcal{H}_{\mathcal{P}_w}$, that is, $\mathcal{H}_{\mathcal{P}_w}$ is a Hilbert basis of $\mathbb{R}_+\Gamma$,
- (b) $A_r^w(\mathcal{P}) = K[\mathbb{N}\mathcal{H}_{\mathcal{P}_w}]$, $w = w_1 \cdots w_p$,
- (c) $A(\mathcal{P}_w) = A_r^w(\mathcal{P})$, and
- (d) $E_{\mathcal{P}_w}(n) = E_{\mathcal{P}}^{s, \prod_{i=1}^p (w_i+1)}(n)$ for $n \geq 0$.
- (e) The r -weighted multivariate Ehrhart series $F_{\mathcal{P}}^{r,w}(t, q, x)$ of \mathcal{P} is a rational function.

In Section 4, we also relate the Ehrhart function of \mathcal{P}_w with s -weighted Ehrhart functions when the weight w is a product $w_1 \cdots w_p$ of linear functions. As an application of Theorem 4.6, we determine the dimension of \mathcal{P}_w when w is a monomial. Let $w = w_1 \cdots w_p$ be any monomial of S of degree p with $w_i \in \{t_1, \dots, t_s\}$ for $i = 1, \dots, p$. If $w_i \not\equiv 0$ on \mathcal{P} for $i \leq r$ and $w_i \equiv 0$ on \mathcal{P} for $i > r$, then

$$\dim(\mathcal{P}_w) = \deg(E_{\mathcal{P}_w}) = \dim(\mathcal{P}) + r,$$

see Lemma 4.8. As a consequence if \mathcal{P} is full-dimensional, that is $\dim(\mathcal{P}) = s$, then the degree of $E_{\mathcal{P}_w}$ is $\dim(\mathcal{P}) + p$.

We conclude this introduction illustrating Theorem 1.8 and Theorem 4.6, as well as its application to s -weightings:

Example 1.9. Let $\mathcal{P} = [0, 1] \times [0, 1]$ be the unit square in \mathbb{R}^2 and let w be the linear weight $w(y_1, y_2) = y_1 + y_2$. The weight lifting polytopes \mathcal{P}^w and \mathcal{P}_w of \mathcal{P} relative to w are given by

$$\mathcal{P}^w = \text{conv}((0, 0, 0), (1, 0, 1), (0, 1, 1), (1, 1, 2)),$$

$$\mathcal{P}_w = \text{conv}((0, 0, 0), (1, 0, 0), (0, 1, 0), (1, 1, 0), (1, 0, 1), (0, 1, 1), (1, 1, 2)).$$

The polytope \mathcal{P}_w is depicted in Figure 1. Note that the ‘‘upper facet’’ of \mathcal{P}_w is precisely \mathcal{P}^w and \mathcal{P}_w contains not only \mathcal{P}^w but also all points below \mathcal{P}^w . By, Corollary 5.4, one has

$$\sum_{a \in n\mathcal{P} \cap \mathbb{Z}^2} w(a) = E_{\mathcal{P}_w}(n) - E_{\mathcal{P}^w}(n) \text{ for } n \geq 0.$$




FIGURE 1. The weight lifting polytope \mathcal{P}_w of the unit square $\mathcal{P} = [0, 1]^2$ relative to $w = y_1 + y_2$. The weight lifting polytope \mathcal{P}^w appears as the upper facet.

s-weighted Ehrhart functions and Ehrhart series. In Section 5, we give other applications of Theorem 4.6 and recover some results from the literature. If $w_i \neq 0$ on \mathcal{P} for $i = 1, \dots, p$, then $E_{\mathcal{P}}^{s,w}$ is a polynomial whose leading coefficient is $\text{vol}(\mathcal{P}_w)$ and

$$\deg(E_{\mathcal{P}}^{s,w}) = \dim(\mathcal{P}) + p,$$

see Proposition 5.1. If f is a polynomial, writing f as a K -linear combination of monomials, this result allows us to find $E_{\mathcal{P}}^{s,f}$ and $F_{\mathcal{P}}^{s,f}$ using polynomial interpolation (Corollary 5.2(a)).

We say that $\mathcal{P} = \text{conv}(v_1, \dots, v_m)$ is *non-degenerate* if for each $1 \leq i \leq s$, there is v_j such that the i -th entry of v_j is non-zero.

Corollary 5.2. *Let f be a non-zero polynomial of $K[t_1, \dots, t_s]$ of degree p and let d be the dimension of \mathcal{P} . The following holds:*

- (a) $E_{\mathcal{P}}^{s,f}$ is a polynomial of n of degree at most $d + p$ and $F_{\mathcal{P}}^{s,f}(x)$ is a rational function.
- (b) If \mathcal{P} is non-degenerate and f is a monomial, then $\deg(E_{\mathcal{P}}^{s,f}) = d + p$.
- (c) $E_{\mathcal{P}}^{s,f}$ is a K -linear combination of Ehrhart polynomials.
- (d) [12, Proposition 4.1] If the interior \mathcal{P}° of \mathcal{P} is nonempty, $K = \mathbb{R}$, and f is homogeneous and $f \geq 0$ on \mathcal{P} , then $E_{\mathcal{P}}^{s,f}$ is a polynomial of degree $s + p$.

It is known that in part (d), the leading coefficient of $E_{\mathcal{P}}^{s,f}$ is equal to $\int_{\mathcal{P}} f$. This fact appears in [6, p. 437] and [17, Proposition 5]. Integrals of the type $\int_{\mathcal{P}} f$ with f a polynomial and \mathcal{P} a rational polytope were studied in [5, 7, 8, 17, 27]. Algorithms and implementation to compute this integral were developed independently in *NmzIntegrate* [18] and *LattE integrale* [4].

The following inequalities compare $E_{\mathcal{P}}^{s,f}(n)$ with $E_{\mathcal{P}}(n)$. If f is a homogeneous polynomial of degree p in S and $K = \mathbb{R}$, then there are $c_1, c_2 \in \mathbb{R}$, $c_1 \leq c_2$, such that

$$c_1 n^p E_{\mathcal{P}}(n) \leq E_{\mathcal{P}}^{s,f}(n) \leq c_2 n^p E_{\mathcal{P}}(n) \text{ for } n \geq 0 \quad (\text{Proposition 5.3}).$$

We use Theorem 4.6 linking classical and weighted Ehrhart theories, together with Stanley's positivity and monotonicity theorems for h^* -vectors of polytopes [32, 34] to prove the following corollary.

Corollary 5.4. *Let v_1, \dots, v_m be points in \mathbb{N}^s , let $\mathcal{P} = \text{conv}(v_1, \dots, v_m)$, and let $w: \mathbb{R}^s \rightarrow \mathbb{R}$ be a non-zero linear function such that $w(e_i) \in \mathbb{N}$ for $i = 1, \dots, s$. The following holds:*

(a) $E_{\mathcal{P}}^{s,w}(n) := \sum_{a \in n\mathcal{P} \cap \mathbb{Z}^s} w(a) = E_{\mathcal{P}_w}(n) - E_{\mathcal{P}}(n)$ for $n \geq 0$,

where $\mathcal{P}_w = \text{conv}((v_1, 0), \dots, (v_m, 0), (v_1, w(v_1)), \dots, (v_m, w(v_m)))$.

(b) If $w = \sum_{i=1}^s \eta_i t_i$, $t_i \neq 0$ on \mathcal{P} whenever $\eta_i > 0$, and $d = \dim(\mathcal{P})$, then $E_{\mathcal{P}}^{s,w}$ is a polynomial of degree $d + 1$ and the generating function $F_{\mathcal{P}}^{s,w}$ of $E_{\mathcal{P}}^{s,w}$ is equal to

$$F_{\mathcal{P}}^{s,w}(x) := \sum_{n=0}^{\infty} E_{\mathcal{P}}^{s,w}(n)x^n = \frac{h(x)}{(1-x)^{d+2}},$$

where $h(x)$ is a polynomial of degree at most $d + 1$ with nonnegative integer coefficients.

For unexplained terminology and additional information, we refer to [11, 13] for Ehrhart theory and [14, 31, 33, 35] for commutative algebra and enumerative combinatorics.

2. WEIGHTED EHRHART SERIES

We start with some useful general observations. In the introduction, we saw that from a polytope \mathcal{P} , we can assign a weight(s) to the lattice points; the weights are either a number, a monomial, a polynomial, etc., depending on the method of weighting. From it, we saw in Definitions 1.2 that the method of weighting gives different formal power series, but all weighted multivariate Ehrhart series can be computed from a single one.

Proposition 2.1. *Both r -weighted multivariate Ehrhart series and s -weighted Ehrhart series can be recovered from q -weighted multivariate Ehrhart series.*

Proof. Note that one can rewrite the r -weighted multivariate Ehrhart function as follows:

$$E_{\mathcal{P}}^{r,w}(q, t, n) := \sum_{\substack{(a,n) \in \text{cone}(\mathcal{P}) \cap \mathbb{Z}^{s+1} \\ 0 \leq b_i \leq w_i(a) \forall i}} t^a q_1^{b_1} \cdots q_p^{b_p} = \sum_{a \in n\mathcal{P} \cap \mathbb{Z}^s} \left[\sum_{0 \leq b_i \leq w_i(a) \forall i} q_1^{b_1} \cdots q_p^{b_p} \right] t^a = \sum_{a \in n\mathcal{P} \cap \mathbb{Z}^s} \left[\prod_{i=1}^p \left(\sum_{j=0}^{w_i(a)} q_i^j \right) \right] t^a.$$

Next, note that

$$\prod_{i=1}^p \left(\sum_{j=0}^{w_i(a)} q_i^j \right) t^a x^n = \prod_{i=1}^p \left(\frac{1 - q_i^{w_i(a)+1}}{1 - q_i} \right) t^a x^n = \sum_{I \subseteq [p]} \left(\frac{\prod_{j \in I} (-q_j)}{\prod_{i=1}^p (1 - q_i)} \right) \left(\prod_{j \in I} q_j^{w_j(a)} \right) t^a x^n,$$

where we allow the case $I = \emptyset$ and by convention, the product over an empty set is 1. For any fixed index set I , one has

$$\left(\prod_{j \in I} q_j^{w_j(a)} \right) t^a x^n = q_1^{w_1(a)} \cdots q_p^{w_p(a)} t^a x^n \Big|_{q_k=1, k \notin I}.$$

Therefore, $F_{\mathcal{P}}^{r, w_1, \dots, w_p}(q, t, x) = \sum_{I \subseteq [p]} \left(\frac{\prod_{j \in I} (-q_j)}{\prod_{i=1}^p (1 - q_i)} \right) \left(F_{\mathcal{P}}^{q, w_1, \dots, w_p}(q, t, x) \Big|_{q_k=1, k \notin I} \right)$, that is,

$$F_{\mathcal{P}}^{r, w_1, \dots, w_p}(q, t, x) = \frac{1}{\prod_{i=1}^p (1 - q_i)} \sum_{k=0}^p \left(\sum_{1 \leq j_1 < \dots < j_k \leq p} (-q_{j_1}) \cdots (-q_{j_k}) F_{\mathcal{P}}^{q, w_{j_1}, \dots, w_{j_k}}(q, t, x) \right).$$

The first term in the summation, i.e., the term corresponding to $k = 0$ is

$$F_{\mathcal{P}}^{q, 0}(q, t, x) = \sum_{n=0}^{\infty} E_{\mathcal{P}}^{q, 0}(q, t, n) x^n = \sum_{n=0}^{\infty} \left(\sum_{a \in n\mathcal{P} \cap \mathbb{Z}^s} t^a \right) x^n.$$

Note that

$$\left[q \cdot \frac{\partial}{\partial q} \left(q^{w(a)} t^a x^n \right) \right] \Big|_{q=1} = w(a) q^{w(a)} t^a x^n \Big|_{q=1} = w(a) t^a x^n,$$

therefore, $F_{\mathcal{P}}^{s, w}(t, x) = \left[q \cdot \frac{\partial}{\partial q} \left(F_{\mathcal{P}}^{q, w}(q, t, x) \right) \right] \Big|_{q=1}$. \square

As the following result shows, the hypothesis in Theorem 1.8 that w_i is linear for $i = 1, \dots, p$ is essential to prove that the q -weighted graded monomial algebra is finitely generated.

Proposition 2.2. *Let $\mathcal{P} = \{1\}$ and $w(a) = a^2$, then the ring*

$$A_q^w(\mathcal{P}) = K[q^{a^2} x^a \mid a \in \mathbb{N}]$$

is not Noetherian. In particular, $A_q^w(\mathcal{P})$ is not finitely generated as a K -algebra.

Proof. A monomial $q^m x^n$ is in $A_q^w(\mathcal{P})$ if and only if there is a partition λ of n , denoted $\lambda \vdash n$, such that the sum of the squares $\lambda_1^2 + \lambda_2^2 + \dots$ is equal to m . Consider the following ideals of the ring $A_q^w(\mathcal{P})$

$$I_k = (q^{i^2} x^i \mid 1 \leq i \leq k), \quad k \geq 1.$$

It suffices to show that $I_{k-1} \subsetneq I_k$ for $k \geq 2$. We claim that $q^{k^2} x^k \in I_k \setminus I_{k-1}$. We argue by contradiction assuming that $q^{k^2} x^k \in I_{k-1}$. Then,

$$q^{k^2} x^k = (q^m x^n)(q^{i^2} x^i)$$

for some $q^m x^n \in A_q^w(\mathcal{P})$ and $1 \leq i \leq k-1$. Hence, there is a partition (n_1, \dots, n_ℓ) of n such that

$$\begin{aligned} n &= n_1 + \dots + n_\ell, \quad m = n_1^2 + \dots + n_\ell^2, \quad n_i \in \mathbb{N}, \\ k &= n + i = n_1 + \dots + n_\ell + i, \quad k^2 = m + i^2 = n_1^2 + \dots + n_\ell^2 + i^2, \quad \dots \\ (n_1 + \dots + n_\ell + i)^2 &= n_1^2 + \dots + n_\ell^2 + i^2. \end{aligned}$$

Hence, from the last equality, we get $n_i = 0$ for $i = 1, \dots, \ell$, $n = 0, m = 0$, and consequently $q^{k^2} x^k = q^{i^2} x^i$ for some $1 \leq i \leq k-1$, a contradiction. \square

When weights are defined by linear functions, then we are able to conclude that all the weighted Ehrhart series we considered are rational functions, just as in the traditional case.

Proposition 2.3. *If $w_i(a) = v_i^\top a + b_i$'s are linear weights, then the q -weighted multivariate Ehrhart series, the q -weighted Ehrhart series, the r -weighted multivariate Ehrhart series, the r -weighted Ehrhart series and the s -weighted multivariate Ehrhart series are all rational functions.*

Proof. The multivariate Ehrhart series of a polytope \mathcal{P} is

$$\sum_{(a,n) \in \text{cone}(\mathcal{P}) \cap \mathbb{Z}^{s+1}} t^a x^n.$$

Recall that t^a is the abbreviation of $t_1^{a_1} \cdots t_s^{a_s}$. We can apply (and in fact effectively compute) the following monomial substitutions: $t_1 \mapsto q_1^{v_1,1} \cdots q_p^{v_p,1} t_1$, \dots , $t_s \mapsto q_1^{v_1,s} \cdots q_p^{v_p,s} t_s$. Then, by the linearity of weights w_i 's,

$$t^a x^n \mapsto q_1^{v_1^\top a} \cdots q_p^{v_p^\top a} t^a x^n.$$

Lastly, we can just multiply the series by $q_1^{b_1} \cdots q_p^{b_p}$. The initial multivariate Ehrhart series has a rational form, then the monomial substitutions gives another rational form for the q -weighted multivariate Ehrhart series. Hence, by Proposition 2.1, the rest of the weighted Ehrhart series are also rational functions. \square

2.1. Positivity of h^* -vector. We begin this section by giving a sufficient condition for the positivity of h^* -coefficients. Then, we give extensions of Ehrhart's reciprocity law.

Theorem 2.4. *If w_i 's are linear weights and a lattice polytope \mathcal{P} has a (w_1, \dots, w_p) -compatible triangulation \mathcal{T} , then the numerator of the rational form of its q -weighted Ehrhart series has positive coefficients. Nevertheless, this is only a sufficient condition for positivity.*

Proof. We can construct a disjoint partition of \mathcal{P} using the triangulation \mathcal{T} , $\mathcal{P} = \sqcup_{\Delta \in \mathcal{T}} \Delta^*$ with Δ^* being obtained possibly by removing several facets of Δ . Note that the q -weighted Ehrhart function is additive with respect to disjoint union. Therefore,

$$E_{\mathcal{P}}^{q,w_1,\dots,w_p}(q, 1, n) = \sum_{\Delta \in \mathcal{T}} E_{\Delta^*}^{q,w_1,\dots,w_p}(q, 1, n),$$

and similarly,

$$F_{\mathcal{P}}^{q,w_1,\dots,w_p}(q, 1, n) = \sum_{\Delta \in \mathcal{T}} F_{\Delta^*}^{q,w_1,\dots,w_p}(q, 1, n).$$

Since Δ^* is a simplex with several facets possibly removed, using [11, Theorem 3.5], it is easy to see that its q -weighted Ehrhart series has the following rational form

$$F_{\Delta^*}^{q,w_1,\dots,w_p}(q, 1, n) = \frac{h_{\Delta^*}(q, x)}{\prod_{i=1}^{d+1} (1 - q_1^{w_1(v_i)} \cdots q_p^{w_p(v_i)} x)},$$

with v_i being the vertices of Δ^* and $h_{\Delta^*}(q, x) \in \mathbb{N}[q, x]$.

Since \mathcal{T} is (w_1, \dots, w_p) -compatible, the denominator is same for every $\Delta \in \mathcal{T}$, without loss of generality, we can denote it as $\prod_{i=1}^{d+1} (1 - q^{\alpha_i} x)$,

$$F_{\mathcal{P}}^{q,w_1,\dots,w_p}(q, 1, n) = \frac{\sum_{\Delta \in \mathcal{T}} h_{\Delta^*}(q, x)}{\prod_{i=1}^{d+1} (1 - q^{\alpha_i} x)}.$$

To see this rational form is reduced, we can simply degenerate $q = 1$ and use the classical Ehrhart theory. \square

2.2. Reciprocity on weighted Ehrhart series.

Theorem 2.5. *If w_1, \dots, w_p are linear weights and \mathcal{P} is a polytope whose cone has dimension $s + 1$, then the q -weighted multivariate Ehrhart series satisfies the reciprocity property, i.e.,*

$$F_{\mathcal{P}}^{q, w_1, \dots, w_p}(q^{-1}, t^{-1}, x^{-1}) = (-1)^{s+1} F_{\mathcal{P}^\circ}^{q, w_1, \dots, w_p}(q, t, x).$$

Proof. By assumption $\text{cone}(\mathcal{P})$ has dimension $s + 1$. Apply Stanley's reciprocity theorem for rational cones [11, Theorem 4.3] to $\text{cone}(\mathcal{P})$, which states that

$$\sum_{(a, n) \in \text{cone}(\mathcal{P}) \cap \mathbb{Z}^{s+1}} (t_1^{-1})^{a_1} \dots (t_s^{-1})^{a_s} (x^{-1})^n = (-1)^{s+1} \sum_{(a, n) \in \text{cone}(\mathcal{P})^\circ \cap \mathbb{Z}^{s+1}} t_1^{a_1} \dots t_s^{a_s} x^n,$$

apply the appropriate monomial substitutions: $t_i \mapsto q_1^{w_1(e_i)} \dots q_p^{w_p(e_i)} t_i$ for $i = 1, \dots, s$. For $(a, n) \in \text{cone}(\mathcal{P}) \cap \mathbb{Z}^{s+1}$, $(a, n) = (a_1, \dots, a_s, n)$, by the linearity of weights w_i 's, we can see

$$(t_1^{-1})^{a_1} \dots (t_s^{-1})^{a_s} (x^{-1})^n \mapsto (q_1^{-1})^{w_1(a)} \dots (q_p^{-1})^{w_p(a)} (t_1^{-1})^{a_1} \dots (t_s^{-1})^{a_s} (x^{-1})^n.$$

Similarly for $(a, n) \in \text{cone}(\mathcal{P})^\circ \cap \mathbb{Z}^{s+1}$,

$$t_1^{a_1} \dots t_s^{a_s} x^n \mapsto q_1^{w_1(a)} \dots q_p^{w_p(a)} t_1^{a_1} \dots t_s^{a_s} x^n,$$

and the asserted equality follows readily. \square

Corollary 2.6. *$F_{\mathcal{P}}^{r, w_1, \dots, w_p}(q^{-1}, t^{-1}, x^{-1})$ can be represented by interior q -weighted multivariate Ehrhart series.*

Proof. This follows from the equalities

$$\begin{aligned} F_{\mathcal{P}}^{r, w_1, \dots, w_p}(q^{-1}, t^{-1}, x^{-1}) &= \sum_{I \subseteq [p]} \left(\frac{\prod_{j \in I} (-q_j^{-1})}{\prod_{i=1}^p (1 - q_i^{-1})} \right) \left(F_{\mathcal{P}}^{q, w_1, \dots, w_p}(q^{-1}, t^{-1}, x^{-1}) \Big|_{q_k=1, k \notin I} \right) \\ &= (-1)^{s+1} \sum_{I \subseteq [p]} \left(\frac{\prod_{j \in I} (-q_j^{-1})}{\prod_{i=1}^p (1 - q_i^{-1})} \right) \left(F_{\mathcal{P}^\circ}^{q, w_1, \dots, w_p}(q, t, x) \Big|_{q_k=1, k \notin I} \right). \square \end{aligned}$$

Now we look at reciprocity in the case of the s -weighting of the Ehrhart series.

Let \mathcal{P} be a simplex in \mathbb{R}^s with vertices v_1, \dots, v_{s+1} . We simplify notation for the rest of this section, we identify v_i with $(v_i, 1)$. With this identification v_1, \dots, v_{s+1} is a basis of \mathbb{R}^{s+1} . We denote by ϕ , the projection $\mathbb{R}^{s+1} \mapsto \mathbb{R}$, $a = (a_i) \mapsto a_{s+1}$. Thus, from $\phi(z)$, we recover the dilation level of z .

Lemma 2.7 (Naive Weighted Reciprocity Lemma). *Assume the following conditions.*

- The \mathcal{P} is a simplex, i.e., the vertices of \mathcal{P} , $\{v_i\}$, forms a basis of \mathbb{R}^{s+1} .
- $h(a)$ is separable and multiplicative with respect to the basis $\{v_i\}$, i.e., there exist univariate functions h_i such that $h(a) = h_1(\alpha_1) \dots h_{s+1}(\alpha_{s+1})$ with $a = \alpha_1 v_1 + \dots + \alpha_{s+1} v_{s+1}$.
- $h_i(x) = \sum_{j=1}^{k_i} P_{ij}(x) \gamma_{ij}^x$, where the P_{ij} 's are polynomials and $\gamma_{ij} \in \mathbb{C} \setminus \{0\}$.

Then, $F_{\mathcal{P}}^{s, h}(t, x)$ and $F_{\mathcal{P}^\circ}^{s, h}(t, x)$ are rational s -weighted multivariate Ehrhart series, and they satisfy the reciprocity relation,

$$F_{\mathcal{P}}^{s, h}(t^{-1}, x^{-1}) = (-1)^{s+1} F_{\mathcal{P}^\circ}^{s, h}(t, x).$$

Proof. Denote $\diamond = \{\sum c_i v_i : 0 \leq c_i < 1\}$ as the fundamental parallelepiped generated by $\{v_i\}$. For any $a \in \text{cone}(\mathcal{P}) \cap \mathbb{Z}^{s+1}$, we can represent $a = \alpha_1 v_1 + \cdots + \alpha_{s+1} v_{s+1}$. Decompose it into integer parts and fractional parts, we have

$$a = \underbrace{[\alpha_1]v_1 + \cdots + [\alpha_{s+1}]v_{s+1}}_{\text{integral parts}} + \underbrace{\{\alpha_1\}v_1 + \cdots + \{\alpha_{s+1}\}v_{s+1}}_{\text{fractional parts}}.$$

For simplicity, we denote the fractional parts as $r \in \diamond$. Then

$$\begin{aligned} F_{\mathcal{P}}^{s,h}(t,x) &= \sum_{a \in \text{cone}(\mathcal{P}) \cap \mathbb{Z}^{s+1}} h(a) t^a x^{\phi(a)} \\ &= \sum_{\alpha_1 v_1 + \cdots + \alpha_{s+1} v_{s+1} \in \text{cone}(\mathcal{P}) \cap \mathbb{Z}^{s+1}} \prod_{i=1}^{s+1} \left(h_i(\alpha_i) t^{[\alpha_i]v_i} x^{\phi([\alpha_i]v_i)} \right) \cdot t^r x^{\phi(r)} \\ &= \sum_{r \in \diamond} \sum_{n_1=0}^{\infty} \cdots \sum_{n_{s+1}=0}^{\infty} \prod_{i=1}^{s+1} \left(h_i(n_i + \{\alpha_i\}) t^{n_i v_i} x^{\phi(n_i v_i)} \right) \cdot t^r x^{\phi(r)} \\ &= \sum_{\beta \in \diamond} \prod_{i=1}^{s+1} \left(\sum_{n_i=0}^{\infty} h_i(n_i + \{\alpha_i\}) t^{n_i v_i} x^{\phi(n_i v_i)} \right) \cdot t^r x^{\phi(r)}. \end{aligned}$$

Similarly, for any $a \in \text{cone}(\mathcal{P}^\circ) \cap \mathbb{Z}^{s+1}$, we can decompose it into

$$a = \underbrace{[\alpha_1]v_1 + \cdots + [\alpha_{s+1}]v_{s+1}}_{\text{integral parts}} - \underbrace{(\{-\alpha_1\}v_1 + \cdots + \{-\alpha_{s+1}\}v_{s+1})}_{\text{fractional parts}}.$$

Denote the fractional part as $r \in \diamond$. Then

$$\begin{aligned} F_{\mathcal{P}^\circ}^{s,h}(t,x) &= \sum_{a \in \text{cone}(\mathcal{P}^\circ) \cap \mathbb{Z}^{s+1}} h(-a) t^a x^{\phi(a)} \\ &= \sum_{\alpha_1 v_1 + \cdots + \alpha_{s+1} v_{s+1} \in \text{cone}(\mathcal{P}^\circ) \cap \mathbb{Z}^{s+1}} \prod_{i=1}^{s+1} \left(h_i(-\alpha_i) t^{[\alpha_i]v_i} x^{\phi([\alpha_i]v_i)} \right) \cdot t^{-r} x^{\phi(-r)} \\ &= \sum_{r \in \diamond} \sum_{n_1=1}^{\infty} \cdots \sum_{n_{s+1}=1}^{\infty} \prod_{i=1}^{s+1} \left(h_i(-n_i + \{-\alpha_i\}) t^{n_i v_i} x^{\phi(n_i v_i)} \right) \cdot t^{-r} x^{\phi(-r)} \\ &= \sum_{r \in \diamond} \prod_{i=1}^{s+1} \left(\sum_{n_i=1}^{\infty} h_i(-n_i + \{-\alpha_i\}) t^{n_i v_i} x^{\phi(n_i v_i)} \right) \cdot t^{-r} x^{\phi(-r)}. \end{aligned}$$

According to the assumption and the equivalent characterizations of rational power series. For each $r \in \diamond$, generating functions

$$G_i(t,x) := \sum_{n_i=0}^{\infty} h_i(n_i + \{\alpha_i\}) t^{n_i v_i} x^{\phi(n_i v_i)} \quad \text{and} \quad \overline{G}_i(t,x) := \sum_{n_i=1}^{\infty} h_i(-n_i + \{-\alpha_i\}) t^{n_i v_i} x^{\phi(n_i v_i)}$$

are rational. In particular, they satisfy the reciprocity property, i.e.,

$$G_i(t^{-1}, x^{-1}) = (-1) \overline{G}_i(t, x).$$

Therefore,

$$\begin{aligned} F_{\mathcal{P}}^{s,h}(t^{-1}, x^{-1}) &= \sum_{r \in \diamond} \prod_{i=1}^{s+1} (G_i(t^{-1}, x^{-1})) \cdot (t^{-1})^r (x^{-1})^{\phi(-r)} \\ &= \sum_{r \in \diamond} \prod_{i=1}^{s+1} ((-1)\overline{G}_i(t, x)) \cdot t^{-r} x^{\phi(-r)} = (-1)^{s+1} F_{\mathcal{P}^\circ}^{s,h}(t, x). \quad \square \end{aligned}$$

Lemma 2.8. *Assume $h(a) = \prod_{i=1}^{s+1} \sum_{j=1}^{k_i} P_{ij}(a_i) \gamma_{ij}^{a_i}$, where the P_{ij} 's are polynomials and γ_{ij} are nonzero complex numbers, then $h(a)$ can be decomposed into finite terms where each term is separable and multiplicative with respect to any basis $\{v_i\}$ of \mathbb{R}^{s+1} .*

Proof. Only need to prove when $h(a) = a_1^{m_1} \cdots a_{s+1}^{m_{s+1}} \cdot \gamma_1^{a_1} \cdots \gamma_{s+1}^{a_{s+1}}$. Suppose $a = \alpha_1 v_1 + \cdots + \alpha_{s+1} v_{s+1}$, then we can simply replace a_i by $\alpha_1 v_{1,i} + \cdots + \alpha_{s+1} v_{s+1,i}$ to get

$$\begin{aligned} h(a) &= \prod_{i=1}^{s+1} (\alpha_1 v_{1,i} + \cdots + \alpha_{s+1} v_{s+1,i})^{m_i} \cdot \prod_{i=1}^{s+1} \gamma_i^{\alpha_1 v_{1,i} + \cdots + \alpha_{s+1} v_{s+1,i}} \\ &= \sum_{\mathbf{k}} c_{\mathbf{k}} \alpha^{\mathbf{k}} \prod_{i=1}^{s+1} (\gamma_i^{v_{1,1}} \cdots \gamma_d^{v_{i,s+1}})^{\alpha_i} = \sum_{\mathbf{k}} c_{\mathbf{k}} \alpha^{\mathbf{k}} \prod_{i=1}^{s+1} \hat{\gamma}_i^{\alpha_i}. \quad \square \end{aligned}$$

Theorem 2.9. *Let $\mathcal{P} \subset \mathbb{R}^s$ be a rational polytope and $h(a) = \prod_{i=1}^{s+1} \sum_{j=1}^{k_i} P_{ij}(a_i) \gamma_{ij}^{a_i}$, where the P_{ij} 's are polynomials and γ_{ij} are nonzero complex numbers. Then, $F_{\mathcal{P}}^{s,h}(t, x)$ and $F_{\mathcal{P}^\circ}^{s,h}(t, x)$ are rational s -weighted multivariate Ehrhart series, and they satisfy the reciprocity relation,*

$$F_{\mathcal{P}}^{s,h}(t^{-1}, x^{-1}) = (-1)^{s+1} F_{\mathcal{P}^\circ}^{s,h}(t, x).$$

Proof. Using the fact that every rational polyhedral cone can be triangulated into simplicial cones and the inclusion-exclusion principle, we can assume that \mathcal{P} is simplicial. Then, applying Lemma 2.8, we can assume $h(a)$ is separable and multiplicative with respect to a basis. Finally, we can apply Naive Weighted Reciprocity Lemma 2.7. \square

3. q -WEIGHTED EHRHART RINGS

Let $S = K[t_1, \dots, t_s]$ be a polynomial ring over a field K containing \mathbb{R} as a subfield, let v_1, \dots, v_m be points in \mathbb{N}^s , let $\mathcal{P} = \text{conv}(v_1, \dots, v_m)$, let $w_i: \mathbb{R}^s \rightarrow \mathbb{R}$, $i = 1, \dots, p$, be linear functions such that $w_i(e_j) \in \mathbb{N}$ for all i, j , and let $w: \mathbb{R}^s \rightarrow \mathbb{R}^p$ be the linear map $w = (w_1, \dots, w_p)$.

Consider the q -weight lifting lattice polytope $\mathcal{P}^w = \text{conv}(\{(v_i, w(v_i))\}_{i=1}^m) \subset \mathbb{R}^{s+p}$. The linear map $\psi: \mathbb{R}^s \rightarrow \mathbb{R}^{s+p}$, $a \mapsto (a, w(a))$ induces a bijective map

$$n\mathcal{P} \cap \mathbb{Z}^s \xrightarrow{\psi} n\mathcal{P}^w \cap \mathbb{Z}^{s+p}, \text{ for } n \in \mathbb{N},$$

and consequently $E_{\mathcal{P}}(n) = E_{\mathcal{P}^w}(n)$ for $n \geq 1$. We will give a different explanation of this equality by showing that $A_q^w(\mathcal{P})$ is the Ehrhart ring of \mathcal{P}^w . Note that $\text{cone}(\mathcal{P})$, the cone over \mathcal{P} , is the cone $C_{\mathcal{P}} := \mathbb{R}_+ \{(v_i, 1)\}_{i=1}^m$.

Proof of Theorem 1.8. (a) Note that $C_{\mathcal{P}}$ is the cone generated by $\{(v_i, 1)\}_{i=1}^m$. The equality follows noticing that $(a, n) \in C_{\mathcal{P}} \cap \mathbb{Z}^{s+1}$ if and only if $a \in n\mathcal{P} \cap \mathbb{Z}^s$.

(b) The inclusion $A_q^w(\mathcal{P})_k A_q^w(\mathcal{P})_n \subset A_q^w(\mathcal{P})_{k+n}$, $k, n \in \mathbb{N}$, follows from the linearity of $w = (w_1, \dots, w_p)$ and the convexity of \mathcal{P} . As the set $\{t^{a q w(a)} z^n \mid a \in n\mathcal{P} \cap \mathbb{Z}^s\}$ has $|n\mathcal{P} \cap \mathbb{Z}^s|$ elements and is linearly independent over K , one has $\dim_K(A_q^w(\mathcal{P})_n) = E_{\mathcal{P}}(n)$.

(c) To show that $A(\mathcal{P}^w) \subset A_q^w(\mathcal{P})$ take $t^a q^b z^n \in A(\mathcal{P}^w)$. Then, $(a, b, n) \in C_{\mathcal{P}^w} \cap \mathbb{Z}^{s+p+1}$,

$$(a, b, n) = \sum_{i=1}^m \lambda_i(v_i, w(v_i), 1), \lambda_i \geq 0, (a, n) \in C_{\mathcal{P}} \cap \mathbb{Z}^{s+1},$$

and $b = w(a)$. Thus, $t^a q^b z^n = t^a q^{w(a)} z^n$ and $(a, n) \in C_{\mathcal{P}} \cap \mathbb{Z}^s$, that is, $t^a q^b z^n \in A_q^w(\mathcal{P})$. To show that $A(\mathcal{P}^w) \supset A_q^w(\mathcal{P})$ take $t^a q^{w(a)} z^n \in A_q^w(\mathcal{P})$. Then, $(a, n) \in C_{\mathcal{P}} \cap \mathbb{Z}^{s+1}$,

$$(a, n) = \sum_{i=1}^m \lambda_i(v_i, 1), \lambda_i \geq 0, \text{ and } w(a) = \sum_{i=1}^m \lambda_i w(v_i).$$

Then, $(a, w(a), n) = \sum_{i=1}^m \lambda_i(v_i, w(v_i), 1) \in C_{\mathcal{P}^w} \cap \mathbb{Z}^{s+p+1}$, and $t^a q^{w(a)} z^n \in A(\mathcal{P}^w)$. Finally, the equality $E_{\mathcal{P}}(n) = E_{\mathcal{P}^w}(n)$ for $n \geq 0$ follows from the equality $A(\mathcal{P}^w) = A_q^w(\mathcal{P})$ and part (b).

(d) The existence of $\mathcal{H}_{\mathcal{P}}$ satisfying $C_{\mathcal{P}} \cap \mathbb{Z}^{s+1} = \mathbb{N}\mathcal{H}_{\mathcal{P}}$ and $A(\mathcal{P}) = K[\mathbb{N}\mathcal{H}_{\mathcal{P}}]$ follows from [37, Theorem 9.3.6]. The other expressions for $A(\mathcal{P}^w)$ follow from parts (a) and (c).

(e) By the proof of part (c), $(a, b, n) \in C_{\mathcal{P}^w} \cap \mathbb{Z}^{s+p+1}$ if and only if $(a, n) \in C_{\mathcal{P}} \cap \mathbb{Z}^{s+1}$ and $b = w(a)$. Then, by [11, Corollary 3.7], the integer-point transform of the cone $C_{\mathcal{P}^w}$ over \mathcal{P}^w is a rational function given by

$$\frac{G(t, q, x)}{\prod_{i=1}^{\ell} (1 - t^{\alpha_i} q^{\beta_i} x^{n_i})} = \sum_{(a, b, n) \in C_{\mathcal{P}^w} \cap \mathbb{Z}^{s+p+1}} t^a q^b x^n = \sum_{(a, n) \in C_{\mathcal{P}} \cap \mathbb{Z}^{s+1}} t^a q^{w(a)} x^n = F_{\mathcal{P}}^{q, w}(t, q, x).$$

Thus, $F_{\mathcal{P}}^{q, w}(t, q, x)$ is a rational function. \square

4. r -WEIGHTED EHRHART RINGS

In this part, we prove one of our main results and show applications to weighted Ehrhart theory. To avoid repetitions, we continue to employ the notations and definitions used in Section 1.

The following lemmas are used to show that the r -weighted Ehrhart ring of a lattice polytope is the Ehrhart ring of some weight lifting lattice polytope when the weight function is a product of linear functions (Theorem 4.6).

Lemma 4.1. *Let $\delta, \eta_1, \dots, \eta_q$ be a sequence of nonnegative integers. If $\eta_1 + \dots + \eta_q \geq \delta$, then there are integers $\alpha_1, \dots, \alpha_q$ such that $\alpha_1 + \dots + \alpha_q = \delta$ and $0 \leq \alpha_i \leq \eta_i$ for all i .*

Proof. We proceed by induction on $\delta \geq 0$. If $\delta = 0$, we set $\alpha_i = 0$ for $i = 1, \dots, q$. Assume that $\delta \geq 1$ and $\eta_1 + \dots + \eta_q \geq \delta$. Then, $\eta_j \geq 1$ for some j and

$$\eta_1 + \dots + \eta_{j-1} + (\eta_j - 1) + \eta_{j+1} + \dots + \eta_q \geq \delta - 1.$$

Then, by induction, there are integers $\alpha'_1, \dots, \alpha'_q$ such that $\delta - 1 = \alpha'_1 + \dots + \alpha'_q$, $0 \leq \alpha'_i \leq \eta_i$ for $i \neq j$, and $\alpha'_j \leq \eta_j - 1$. Setting $\alpha_i = \alpha'_i$ for $i \neq j$ and $\alpha_j = \alpha'_j + 1$, we obtain $\delta = \alpha_1 + \dots + \alpha_q$ and $0 \leq \alpha_i \leq \eta_i$ for $i = 1, \dots, q$. \square

Lemma 4.2. *Let δ, ϵ, μ be positive integers. If $\delta \leq \epsilon\mu$, then there are positive integers ϵ_i, μ_i , $i = 1, \dots, r$, such that $\delta = \sum_{i=1}^r \epsilon_i \mu_i$, $\epsilon \geq \sum_{i=1}^r \epsilon_i$, and $1 \leq \mu_i \leq \mu$ for $i = 1, \dots, r$.*

Proof. We argue by induction on $\epsilon \geq 1$. If $\epsilon = 1$, we set $\epsilon_1 = 1$, $\mu_1 = \delta$, and $r = 1$. Assume that $\epsilon > 1$ and $\delta \leq \epsilon\mu$. If $\delta \leq \mu$, we set $\epsilon_1 = 1$, $\mu_1 = \delta$, and $r = 1$. Thus, we may assume that $\delta > \mu$. Then, $1 \leq \delta - \mu \leq (\epsilon - 1)\mu$ and, by induction, there are positive integers ϵ_i, μ_i , $i = 1, \dots, q$, such that $\delta - \mu = \sum_{i=1}^q \epsilon_i \mu_i$, $\epsilon - 1 \geq \sum_{i=1}^q \epsilon_i$, and $1 \leq \mu_i \leq \mu$ for $i = 1, \dots, q$. To complete the induction process, we set $r = q + 1$, $\epsilon_r = 1$ and $\mu_r = \mu$. \square

Lemma 4.3. *Let r be a positive integer and let $n, \delta, \epsilon_1, \dots, \epsilon_r, w_1, \dots, w_r$ be positive real (resp. rational) numbers. If $\delta \leq \sum_{i=1}^r n\epsilon_i w_i$, then there are nonnegative real (resp. rational) numbers μ_1, \dots, μ_r such that $\delta = \sum_{i=1}^r n\epsilon_i \mu_i$ and $0 \leq \mu_i \leq w_i$ for $i = 1, \dots, r$.*

Proof. We proceed by induction on r . If $r = 1$, setting $\mu_1 = \delta/(n\epsilon_1)$, one has $\delta = n\epsilon_1 \mu_1$ and $\mu_1 = \delta/(n\epsilon_1) \leq w_1$. Assume that $r > 1$ and $\delta \leq \sum_{i=1}^r n\epsilon_i w_i$. If $\delta \leq n\epsilon_1 w_1$, setting $\mu_1 = \delta/(n\epsilon_1)$ and $\mu_i = 0$ for $i = 2, \dots, r$, one has $\delta = \sum_{i=1}^r n\epsilon_i \mu_i$ and $0 \leq \mu_i \leq w_i$ for $i = 1, \dots, r$. If $\delta > n\epsilon_1 w_1$, then $0 < \delta - n\epsilon_1 w_1 \leq \sum_{i=2}^r n\epsilon_i w_i$. Hence, by induction, then there are nonnegative numbers μ_2, \dots, μ_r such that $\delta - n\epsilon_1 w_1 = \sum_{i=2}^r n\epsilon_i \mu_i$ and $0 \leq \mu_i \leq w_i$ for $i = 2, \dots, r$. Then, the result follows by setting $\mu_1 = w_1$. \square

Lemma 4.4. *Let w_1, \dots, w_d be nonnegative real numbers and let $T: \mathbb{R}^d \mapsto \mathbb{R}^d$ be the linear map such that $T(e_i) = w_i e_i$ for $i = 1, \dots, d$. Then, $T([0, 1]^d) = [0, w_1] \times \dots \times [0, w_d]$.*

Proof. To show the inclusion “ \subset ” take $\alpha \in T([0, 1]^d)$. Then, $\alpha = T(x_1, \dots, x_d)$, $1 \leq x_i \leq 1$ for all i , and consequently $\alpha = x_1 w_1 e_1 + \dots + x_d w_d e_d$. Thus, α is in $[0, w_1] \times \dots \times [0, w_d]$ because $0 \leq x_i w_i \leq w_i$ for all i . Now, we show the inclusion “ \supset ”. Take $\gamma \in [0, w_1] \times \dots \times [0, w_d]$. Then, $\gamma = \gamma_1 e_1 + \dots + \gamma_d e_d$, $0 \leq \gamma_i \leq w_i$ for all i . Consider the vector $\beta = (\beta_1, \dots, \beta_d)$, given by $\beta_i = 0$ if $w_i = 0$ and $\beta_i = \gamma_i/w_i$ if $w_i > 0$. Then, $\beta \in [0, 1]^d$ and $\gamma = T(\beta)$, that is, $\gamma \in T([0, 1]^d)$. \square

Lemma 4.5. *Let $w_i: K^s \rightarrow K$, $i = 1, \dots, p$, be linear functions such that $w_i(e_j) \in \mathbb{N}$ for all i, j . If $w = w_1 \cdots w_p$ and $A_r^w(\mathcal{P}) = \bigoplus_{n=0}^{\infty} A_r^w(\mathcal{P})_n$ is the r -weighted Ehrhart ring of \mathcal{P} , then*

$$\dim_K(A_r^w(\mathcal{P})_n) = E_{\mathcal{P}}(n) + \left(\sum_{i=1}^p E_{\mathcal{P}}^{s, w_i}(n) \right) + \left(\sum_{1 \leq j_1 < j_2 \leq p} E_{\mathcal{P}}^{s, w_{j_1} w_{j_2}}(n) \right) + \dots + E_{\mathcal{P}}^{s, w_1 \cdots w_p}(n).$$

Proof. The n -th graded component of $A_r^w(\mathcal{P})$ is given by

$$A_r^w(\mathcal{P})_n = \sum_{\substack{a \in n\mathcal{P} \cap \mathbb{Z}^s \\ 0 \leq b_i \leq w_i(a)}} K t^a q_1^{b_1} \cdots q_p^{b_p} z^n.$$

To determine the dimension of $A_r^w(\mathcal{P})_n$, note that for each $a \in n\mathcal{P} \cap \mathbb{Z}^s$, the set

$$\{t^a q^b z^n \mid b = (b_1, \dots, b_p) \in ([0, w_1(a)] \times \dots \times [0, w_p(a)]) \cap \mathbb{Z}^p\}$$

has $\prod_{i=1}^p (w_i(a) + 1)$ elements. Then

$$\begin{aligned} \dim_K(A_r^w(\mathcal{P})_n) &= \sum_{a \in n\mathcal{P} \cap \mathbb{Z}^s} \prod_{i=1}^p (w_i(a) + 1) = E_{\mathcal{P}}^{s, \prod_{i=1}^p (w_i+1)}(n) \\ &= \sum_{k=0}^p \left(\sum_{1 \leq j_1 < \dots < j_k \leq p} E_{\mathcal{P}}^{s, w_{j_1} \cdots w_{j_k}}(n) \right) \\ &= E_{\mathcal{P}}(n) + \left(\sum_{i=1}^p E_{\mathcal{P}}^{s, w_i}(n) \right) + \left(\sum_{1 \leq j_1 < j_2 \leq p} E_{\mathcal{P}}^{s, w_{j_1} w_{j_2}}(n) \right) + \dots + E_{\mathcal{P}}^{s, w_1 \cdots w_p}(n), \end{aligned}$$

and the proof is complete. \square

Theorem 4.6. *Let w_1, \dots, w_p be linear functions such that $w_i(e_j) \in \mathbb{N}$ for all i, j and let $\mathcal{P}_w, \mathcal{H}_{\mathcal{P}_w}, \Gamma$, be as in Eqs. (1.5)-(1.7). Then*

- (a) $\mathbb{R}_+ \Gamma \cap \mathbb{Z}^{s+p+1} = \mathbb{N} \mathcal{H}_{\mathcal{P}_w}$, that is, $\mathcal{H}_{\mathcal{P}_w}$ is a Hilbert basis of $\mathbb{R}_+ \Gamma$,
- (b) $A_r^w(\mathcal{P}) = K[\mathbb{N} \mathcal{H}_{\mathcal{P}_w}]$, $w = w_1 \cdots w_p$,

- (c) $A(\mathcal{P}_w) = A_r^w(\mathcal{P})$, and
- (d) $E_{\mathcal{P}_w}(n) = E_{\mathcal{P}}^{s, \prod_{i=1}^p (w_i+1)}(n)$ for $n \geq 0$.
- (e) The r -weighted multivariate Ehrhart series $mF_{\mathcal{P}}^{r,w}(t, q, x)$ of \mathcal{P} is a rational function.

Proof. (a) To show the inclusion “ \subset ” we proceed by induction on $p \geq 1$. Assume that $p = 1$. Take $(a, b, n) \in \mathbb{R}_+\Gamma \cap \mathbb{Z}^{s+2}$, $b \in \mathbb{N}_+$. Then

$$(4.1) \quad (a, b, n) = \sum_{i=1}^m \mu_{0,i}(v_i, 0, 1) + \sum_{i=1}^m \mu_{1,i}(v_i, w_1(v_i), 1),$$

where $\mu_{i,j} \geq 0$, and consequently we obtain the following equalities

$$(4.2) \quad a = \mu_{0,1}v_1 + \cdots + \mu_{0,m}v_m + \mu_{1,1}v_1 + \cdots + \mu_{1,m}v_m,$$

$$(4.3) \quad b = \mu_{1,1}w_1(v_1) + \cdots + \mu_{1,m}w_1(v_m),$$

$$n = \mu_{0,1} + \cdots + \mu_{0,m} + \mu_{1,1} + \cdots + \mu_{1,m},$$

$$(4.4) \quad (a, n) = \sum_{i=1}^m \mu_{0,i}(v_i, 1) + \sum_{i=1}^m \mu_{1,i}(v_i, 1) = \eta_1(c_1, d_1) + \cdots + \eta_r(c_r, d_r),$$

for some $\eta_1, \dots, \eta_r \in \mathbb{N}$. The last equality follows recalling that $\mathbb{R}_+\mathcal{B} \cap \mathbb{Z}^{s+1} = \mathbb{N}\mathcal{H}_{\mathcal{P}}$ and noticing that $(a, n) \in \mathbb{R}_+\mathcal{B} \cap \mathbb{Z}^{s+1}$. From Eqs. (4.2) and (4.3), one has

$$b = \sum_{i=1}^m \mu_{1,i}w_1(v_i) \leq \sum_{i=1}^m \mu_{0,i}w_1(v_i) + \sum_{i=1}^m \mu_{1,i}w_1(v_i) = w_1(a),$$

that is, $b \leq w_1(a)$. Hence, from Eq. (4.4), we get

$$(4.5) \quad b \leq w_1(a) = \eta_1 w_1(c_1) + \cdots + \eta_r w_1(c_r).$$

As $\eta_i, w_1(c_i) \in \mathbb{N}$ for $i = 1, \dots, r$, by Lemma 4.1, we can write $b = b_{1,1} + \cdots + b_{1,r}$ for some $b_{1,i}$'s in \mathbb{N} such that $b_{1,i} \leq \eta_i w_1(c_i)$ for $i = 1, \dots, r$. If $b_{1,i} = 0$, we set $k_{1,i} = 1$ and $\eta_{1,i} = u_{1,i} = 0$. Then, applying Lemma 4.2 to each inequality $b_{1,i} \leq \eta_i w_1(c_i)$ with $b_{1,i} > 0$, there are nonnegative integers $\eta_{1,i}^{(j)}, u_{1,i}^{(j)}$ such that

$$(4.6) \quad b_{1,i} = \sum_{j=1}^{k_{1,i}} \eta_{1,i}^{(j)} u_{1,i}^{(j)}, \quad \eta_i \geq \sum_{j=1}^{k_{1,i}} \eta_{1,i}^{(j)}, \quad u_{1,i}^{(j)} \leq w_1(c_i),$$

for $i = 1, \dots, r$. Writing $\eta_i = (\sum_{j=1}^{k_{1,i}} \eta_{1,i}^{(j)}) + (\eta_i - \sum_{j=1}^{k_{1,i}} \eta_{1,i}^{(j)})$, by Eqs. (4.4)–(4.6), we get

$$\begin{aligned} (a, b, n) &= \sum_{i=1}^r \eta_i(c_i, 0, d_i) + \left(\sum_{i=1}^r b_{1,i} \right) e_{s+1} \\ &= \sum_{i=1}^r \left(\sum_{j=1}^{k_{1,i}} \eta_{1,i}^{(j)} \right) (c_i, 0, d_i) + \left(\sum_{i=1}^r \left(\sum_{j=1}^{k_{1,i}} \eta_{1,i}^{(j)} u_{1,i}^{(j)} \right) \right) e_{s+1} + \sum_{i=1}^r \left(\eta_i - \sum_{j=1}^{k_{1,i}} \eta_{1,i}^{(j)} \right) (c_i, 0, d_i) \\ &= \sum_{i=1}^r \left(\sum_{j=1}^{k_{1,i}} \eta_{1,i}^{(j)} (c_i, 0, d_i) + \sum_{j=1}^{k_{1,i}} \eta_{1,i}^{(j)} u_{1,i}^{(j)} e_{s+1} \right) + \sum_{i=1}^r \left(\eta_i - \sum_{j=1}^{k_{1,i}} \eta_{1,i}^{(j)} \right) (c_i, 0, d_i) \\ &= \sum_{i=1}^r \left(\sum_{j=1}^{k_{1,i}} \eta_{1,i}^{(j)} (c_i, u_{1,i}^{(j)}, d_i) \right) + \sum_{i=1}^r \left(\eta_i - \sum_{j=1}^{k_{1,i}} \eta_{1,i}^{(j)} \right) (c_i, 0, d_i), \end{aligned}$$

that is, $(a, b, n) \in \mathbb{N}\mathcal{H}_{\mathcal{P}_w}$. Assume that $p > 1$. We set

$$\begin{aligned} \mathcal{G}' &:= \{(v_i, 0) + w_{j_1}(v_i)e_{s+j_1} + \cdots + w_{j_k}(v_i)e_{s+j_k} \mid 1 \leq i \leq m, 1 \leq j_1 < \cdots < j_k < p\}, \\ \mathcal{P}' &:= \text{conv}(\mathcal{G}'), \quad \Gamma' := \{(\alpha', 1) \mid \alpha' \in \mathcal{G}'\}, \\ (4.7) \quad \mathcal{H}_{\mathcal{P}'} &:= \{(c_i, b', d_i) \mid b' = (b_1, \dots, b_{p-1}) \in \mathbb{N}^{p-1}, 0 \leq b_j \leq w_j(c_i) \forall j, c_i \in d_i \mathcal{P}, 1 \leq i \leq r\}. \end{aligned}$$

Take $(a, b, n) \in \mathbb{R}_+\Gamma \cap \mathbb{Z}^{s+p+1}$, $b = (b_1, \dots, b_p)$. Then, one has

$$(4.8) \quad (a, b, n) = \sum_{\substack{1 \leq i \leq m \\ 1 \leq j_1 < \cdots < j_k \leq p}} \mu_{i, j_1, \dots, j_k} ((v_i, 0) + w_{j_1}(v_i)e_{s+j_1} + \cdots + w_{j_k}(v_i)e_{s+j_k} + e_{s+p+1}),$$

where $\mu_{i, j_1, \dots, j_k} \geq 0$. Consequently, noticing that b_i is the $(i+s)$ -entry of (a, b, n) , we obtain the following equalities

$$(4.9) \quad a = \sum \mu_{i, j_1, \dots, j_k} v_i,$$

$$(4.10) \quad b_1 e_{s+1} + \cdots + b_p e_{s+p} = \sum \mu_{i, j_1, \dots, j_k} (w_{j_1}(v_i)e_{s+j_1} + \cdots + w_{j_k}(v_i)e_{s+j_k}),$$

$$(4.11) \quad n = \sum \mu_{i, j_1, \dots, j_k},$$

$$(4.12) \quad (a, n) = \sum \mu_{i, j_1, \dots, j_k} (v_i + e_{s+p+1}).$$

Consider the following Cartesian products

$$(4.13) \quad ([0, w_1(c_1)] \times \cdots \times [0, w_{p-1}(c_1)]) \cap \mathbb{Z}^{p-1} = \{\beta_{1,1}, \dots, \beta_{1,q_1}\},$$

$$\vdots$$

$$(4.14) \quad ([0, w_1(c_r)] \times \cdots \times [0, w_{p-1}(c_r)]) \cap \mathbb{Z}^{p-1} = \{\beta_{r,1}, \dots, \beta_{r,q_r}\}.$$

Setting $b' = (b_1, \dots, b_{p-1})$, one has $(a, b', n) \in \mathbb{R}_+\Gamma' \cap \mathbb{Z}^{s+p}$. Then, by induction, we get $(a, b', n) \in \mathbb{N}\mathcal{H}_{\mathcal{P}'}$. Therefore, we can write

$$(4.15) \quad (a, b', n) = \sum_{i=1}^r \sum_{j=1}^{q_i} \eta_{i,j} (c_i, \beta_{i,j}, d_i),$$

where $\eta_{i,j} \in \mathbb{N}$. From Eqs. (4.9) and (4.10), one has

$$b_p = \sum_{j_k=p} \mu_{i, j_1, \dots, j_k} w_p(v_i) \leq \sum \mu_{i, j_1, \dots, j_k} w_p(v_i) = w_p(a),$$

that is, $b_p \leq w_p(a)$. Hence, from Eq. (4.15), we get

$$b_p \leq w_p(a) = \sum_{i=1}^r \sum_{j=1}^{q_i} \eta_{i,j} w_p(c_i).$$

As $\eta_{i,j} w_p(c_i) \in \mathbb{N}$, by Lemma 4.1, we can write

$$b_p = \sum_{i=1}^r \sum_{j=1}^{q_i} \alpha_{i,j},$$

where $\alpha_{i,j} \in \mathbb{N}$ and $\alpha_{i,j} \leq \eta_{i,j} w_p(c_i)$. Then, applying Lemma 4.2 to each inequality, there are nonnegative integers $\eta_{i,j}^{(\ell)}, u_{i,j}^{(\ell)}$ such that

$$b_p = \sum_{i=1}^r \sum_{j=1}^{q_i} \sum_{\ell=1}^{k_{i,j}} \eta_{i,j}^{(\ell)} u_{i,j}^{(\ell)},$$

$u_{i,j}^{(\ell)} \leq w_p(c_i)$ and $\eta_{i,j} \geq \sum_{\ell=1}^{k_{i,j}} \eta_{i,j}^{(\ell)}$. As $(a, b, n) = (a, b', n) + b_p e_{s+p}$, by Eq. (4.15), we get

$$\begin{aligned} (a, b, n) &= \sum_{i=1}^r \sum_{j=1}^{q_i} \eta_{i,j}(c_i, \beta_{i,j}, 0, d_i) + b_p e_{s+p} = \sum_{i=1}^r \sum_{j=1}^{q_i} \left(\sum_{\ell=1}^{k_{i,j}} \eta_{i,j}^{(\ell)} \right) (c_i, \beta_{i,j}, 0, d_i) + b_p e_{s+p} \\ &+ \sum_{i=1}^r \sum_{j=1}^{q_i} \left(\eta_{i,j} - \sum_{\ell=1}^{k_{i,j}} \eta_{i,j}^{(\ell)} \right) (c_i, \beta_{i,j}, 0, d_i) \\ &= \sum_{i=1}^r \sum_{j=1}^{q_i} \sum_{\ell=1}^{k_{i,j}} \eta_{i,j}^{(\ell)} (c_i, \beta_{i,j}, 0, d_i) + \left(\sum_{i=1}^r \sum_{j=1}^{q_i} \sum_{\ell=1}^{k_{i,j}} \eta_{i,j}^{(\ell)} u_{i,j}^{(\ell)} \right) e_{s+p} \\ &+ \sum_{i=1}^r \sum_{j=1}^{q_i} \left(\eta_{i,j} - \sum_{\ell=1}^{k_{i,j}} \eta_{i,j}^{(\ell)} \right) (c_i, \beta_{i,j}, 0, d_i) \\ &= \sum_{i=1}^r \sum_{j=1}^{q_i} \sum_{\ell=1}^{k_{i,j}} \eta_{i,j}^{(\ell)} (c_i, \beta_{i,j}, u_{i,j}^{(\ell)}, d_i) + \sum_{i=1}^r \sum_{j=1}^{q_i} \left(\eta_{i,j} - \sum_{\ell=1}^{k_{i,j}} \eta_{i,j}^{(\ell)} \right) (c_i, \beta_{i,j}, 0, d_i). \end{aligned}$$

Therefore, $(a, b, n) \in \mathbb{N}\mathcal{H}_{\mathcal{P}_w}$, and the induction process is complete.

To show the inclusion “ \supset ” take $(c_i, b, d_i) \in \mathcal{H}_{\mathcal{P}_w}$, $b = (b_1, \dots, b_p)$. It suffices to show that $(c_i, b, d_i) \in \mathbb{R}_+\Gamma$. As $c_i \in d_i\mathcal{P}$ and $0 \leq b_j \leq w_j(c_i)$ for $j = 1, \dots, p$, one has

$$\begin{aligned} c_i &= d_i \sum_{\ell=1}^m \epsilon_\ell v_\ell, \quad \epsilon_\ell \geq 0, \quad \sum_{\ell=1}^m \epsilon_\ell = 1, \\ 0 \leq b_j &\leq w_j(c_i) = d_i \sum_{\ell=1}^m \epsilon_\ell w_j(v_\ell), \quad j = 1, \dots, p. \end{aligned}$$

Hence, noticing that b_i is the $(s+i)$ -entry of (c_i, b, d_i) and using Lemma 4.3, we obtain

$$\begin{aligned} b_j &= d_i \sum_{\ell=1}^m \epsilon_\ell \mu_{j,\ell}, \quad 0 \leq \mu_{j,\ell} \leq w_j(v_\ell), \quad \ell = 1, \dots, m, \quad j = 1, \dots, p, \\ (c_i, b, d_i) &= d_i \left[\sum_{\ell=1}^m \epsilon_\ell (v_\ell, 0) + \sum_{j=1}^p \left(\sum_{\ell=1}^m \epsilon_\ell \mu_{j,\ell} \right) e_{s+j} + \left(\sum_{\ell=1}^m \epsilon_\ell \right) e_{s+p+1} \right] \\ &= d_i \left[\sum_{\ell=1}^m \epsilon_\ell \left((v_\ell, 0) + \left(\sum_{j=1}^p \mu_{j,\ell} e_{s+j} \right) + e_{s+p+1} \right) \right]. \end{aligned}$$

Thus, setting $\gamma_\ell := (v_\ell, 0) + \left(\sum_{j=1}^p \mu_{j,\ell} e_{s+j} \right) + e_{s+p+1}$, we need only show that $\gamma_\ell \in \mathbb{R}_+\Gamma$ for $\ell = 1, \dots, m$. Recall that

$$[0, 1]^p = \text{conv}(\{0\} \cup \{e_{j_1} + \dots + e_{j_k} \mid 1 \leq j_1 < \dots < j_k \leq p\}).$$

Fix $1 \leq \ell \leq m$ and consider the linear map $T_\ell: \mathbb{R}^p \rightarrow \mathbb{R}^p$, $e_i \mapsto w_i(v_\ell)e_i$, $i = 1, \dots, p$. Using that T_ℓ is linear and commutes with taking convex hulls, by Lemma 4.4, we get

$$\begin{aligned} [0, w_1(v_\ell)] \times \cdots \times [0, w_p(v_\ell)] &= T_\ell([0, 1]^p) \\ &= T_\ell(\text{conv}(\{0\} \cup \{e_{j_1} + \cdots + e_{j_k} \mid 1 \leq j_1 < \cdots < j_k \leq p\})) \\ &= \text{conv}(\{0\} \cup \{w_{j_1}(v_\ell)e_{j_1} + \cdots + w_{j_k}(v_\ell)e_{j_k} \mid 1 \leq j_1 < \cdots < j_k \leq p\}). \end{aligned}$$

Hence, as $0 \leq \mu_{j,\ell} \leq w_j(v_\ell)$ for $j = 1, \dots, p$, one obtains

$$\begin{aligned} \sum_{j=1}^p \mu_{j,\ell} e_j &\in \text{conv}(\{0\} \cup \{w_{j_1}(v_\ell)e_{j_1} + \cdots + w_{j_k}(v_\ell)e_{j_k} \mid 1 \leq j_1 < \cdots < j_k \leq p\}), \\ \sum_{j=1}^p \mu_{j,\ell} e_j &= \lambda_1 \cdot 0 + \sum \lambda_{j_1, \dots, j_k} w_{j_1}(v_\ell)e_{j_1} + \cdots + w_{j_k}(v_\ell)e_{j_k}, \\ \lambda_1 &\geq 0, \lambda_{j_1, \dots, j_k} \geq 0, \lambda_1 + \sum \lambda_{j_1, \dots, j_k} = 1, \\ (v_\ell, 0) + \sum_{j=1}^p \mu_{j,\ell} e_{s+j} &= (\lambda_1 + \sum \lambda_{j_1, \dots, j_k})(v_\ell, 0) + \lambda_1 \cdot 0 \\ &\quad + \sum \lambda_{j_1, \dots, j_k} w_{j_1}(v_\ell)e_{s+j_1} + \cdots + w_{j_k}(v_\ell)e_{s+j_k} \\ &= \lambda_1(v_\ell, 0) + \sum \lambda_{j_1, \dots, j_k} ((v_\ell, 0) + w_{j_1}(v_\ell)e_{s+j_1} + \cdots + w_{j_k}(v_\ell)e_{s+j_k}). \end{aligned}$$

Then, adding $e_{s+p+1} = (\lambda_1 + \sum \lambda_{j_1, \dots, j_k})e_{s+p+1}$ to the last two equalities, we get

$$\begin{aligned} \gamma_\ell &= (v_\ell, 0) + \left(\sum_{j=1}^p \mu_{j,\ell} e_{s+j} \right) + e_{s+p+1} = \lambda_1((v_\ell, 0) + e_{s+p+1}) \\ &\quad + \sum \lambda_{j_1, \dots, j_k} ((v_\ell, 0) + w_{j_1}(v_\ell)e_{s+j_1} + \cdots + w_{j_k}(v_\ell)e_{s+j_k} + e_{s+p+1}). \end{aligned}$$

Thus, $\gamma_\ell \in \mathbb{R}_+\Gamma$ for $\ell = 1, \dots, m$.

(b) The semigroup ring $K[\mathbb{N}\mathcal{H}_{\mathcal{P}_w}]$ is equal to the monomial subring

$$K[t^a q^b z^n \mid (a, b, n) \in \mathbb{N}\mathcal{H}_{\mathcal{P}_w}],$$

and the inclusion $A_r^w(\mathcal{P}) \supset K[\mathbb{N}\mathcal{H}_{\mathcal{P}_w}]$ follows noticing that $t^{c_i} q^b z^{d_i}$ is in $A_r^w(\mathcal{P})$ for all (c_i, b, d_i) in $\mathcal{H}_{\mathcal{P}_w}$. The inclusion $A_r^w(\mathcal{P}) \subset K[\mathbb{N}\mathcal{H}_{\mathcal{P}_w}]$ follows from the proof of part (a). For clarity, we briefly explain the main steps. We proceed by induction on $p \geq 1$. Assume that $p = 1$. Take $t^a q_1^b z^n \in A_r^w(\mathcal{P})$. Then, $a \in n\mathcal{P} \cap \mathbb{Z}^s$, $b \in \mathbb{N}$, and $0 \leq b \leq w_1(a)$. Thus, $t^a z^n \in A(\mathcal{P})$ and consequently, by Eq. (1.2), we get

$$\begin{aligned} t^a z^n &= (t^{c_1} z^{d_1})^{\eta_1} \cdots (t^{c_r} z^{d_r})^{\eta_r}, \quad \eta_i \in \mathbb{N}, \\ (a, n) &= \eta_1(c_1, d_1) + \cdots + \eta_r(c_r, d_r), \\ b &\leq w_1(a) = \eta_1 w_1(c_1) + \cdots + \eta_r w_1(c_r). \end{aligned}$$

We can now proceed as in the case $p = 1$ of part (a), starting with Eq. (4.5), to obtain that $(a, b, n) \in \mathbb{N}\mathcal{H}_{\mathcal{P}_w}$, that is, $t^a q_1^b z^n \in K[\mathbb{N}\mathcal{H}_{\mathcal{P}_w}]$. Assume that $p > 1$. Take $t^a q^b z^n \in A_r^w(\mathcal{P})$, $b = (b_1, \dots, b_p)$. In particular one has $0 \leq b_i \leq w_i(a)$ for $i = 1, \dots, p$. With the notation of Eq. (4.7), by induction we get

$$t^a q_1^{b_1} \cdots q_{p-1}^{b_{p-1}} z^n \in K[\mathbb{N}\mathcal{H}_{\mathcal{P}'}],$$

that is, setting $b' = (b_1, \dots, b_{p-1})$, with the notation of Eqs. (4.13)–(4.14), one has

$$(a, b', n) = \sum_{j=1}^{q_1} \eta_{1,j}(c_1, \beta_{1,j}, d_1) + \dots + \sum_{j=1}^{q_r} \eta_{r,j}(c_r, \beta_{r,j}, d_r).$$

As $0 \leq b_p \leq w_p(a)$, starting with Eq. (4.15), we can proceed as in the case $p > 1$ of part (a) to obtain that $(a, b, n) \in \mathbb{N}\mathcal{H}_{\mathcal{P}_w}$.

(c) As $\mathcal{P}_w = \text{conv}(\mathcal{G})$ and $\Gamma = \{(\alpha, 1) \mid \alpha \in \mathcal{G}\}$, by [37, Theorem 9.3.6(b)], one has

$$A(\mathcal{P}_w) = K[t^a q^b z^n \mid (a, b, n) \in \mathbb{R}_+ \Gamma \cap \mathbb{Z}^{s+p+1}] = K[\mathbb{R}_+ \Gamma \cap \mathbb{Z}^{s+p+1}].$$

Then, by parts (a) and (b), $A(\mathcal{P}_w) = K[\mathbb{N}\mathcal{H}_{\mathcal{P}_w}] = A_r^w(\mathcal{P})$.

(d) By part (c), $A(\mathcal{P}_w) = A_r^w(\mathcal{P})$. Thus, by Lemma 4.5, we get

$$E_{\mathcal{P}_w}(n) = \dim_K(A(\mathcal{P}_w)_n) = \dim_K(A_r^w(\mathcal{P})_n) = E_{\mathcal{P}}^{s, \prod_{i=1}^p (w_i+1)}(n),$$

and the proof of (d) is complete.

(e) By part (a), $(a, b, n) \in \mathbb{R}_+ \Gamma \cap \mathbb{Z}^{s+p+1}$ if and only if $(a, b, n) \in \mathbb{N}\mathcal{H}_{\mathcal{P}_w}$ and, by part (b), $(a, b, n) \in \mathbb{N}\mathcal{H}_{\mathcal{P}_w}$ if and only if $(a, n) \in \text{cone}(\mathcal{P}) \cap \mathbb{Z}^{s+1}$ and $0 \leq b_i \leq w_i(a)$ for all i , where $b = (b_1, \dots, b_p)$. Then, by [11, Corollary 3.7], the integer-point transform of the cone $\mathbb{R}_+ \Gamma$ is a rational function given by

$$\sum_{(a,b,n) \in \mathbb{R}_+ \Gamma \cap \mathbb{Z}^{s+p+1}} t^a q^b x^n = \sum_{\substack{(a,n) \in \text{cone}(\mathcal{P}) \cap \mathbb{Z}^{s+1} \\ 0 \leq b_i \leq w_i(a) \ \forall i}} t^a q^b x^n = F_{\mathcal{P}}^{r,w}(t, q, x).$$

Thus, $F_{\mathcal{P}}^{r,w}(t, q, x)$ is a rational function. \square

Remark 4.7. From Theorem 4.6(c) it follows that $t^a q^b z^n \in A_r^w(\mathcal{P})_n$ if and only if $(a, b) \in n\mathcal{P}_w$.

The following result says in particular that if \mathcal{P} is full-dimensional, that is, $\dim(\mathcal{P}) = s$ and w is a monomial of degree p , then the degree of $E_{\mathcal{P}_w}$ is $\dim(\mathcal{P}) + p$.

Lemma 4.8. *Let $w = w_1 \cdots w_p$ be any monomial of S of degree p with $w_i \in \{t_1, \dots, t_s\}$ for $i = 1, \dots, p$. If $w_i \not\equiv 0$ on \mathcal{P} for $i \leq r$ and $w_i \equiv 0$ on \mathcal{P} for $i > r$, then*

$$\dim(\mathcal{P}_w) = \deg(E_{\mathcal{P}_w}) = \dim(\mathcal{P}) + r.$$

Proof. The equality $\dim(\mathcal{P}_w) = \deg(E_{\mathcal{P}_w})$ is well known, see [11], [37, p. 383]. Let D be the following matrix of size $(m \sum_{k=0}^p \binom{p}{k}) \times (s+p+1)$ whose rows correspond to the vectors in Γ

$$D = \begin{bmatrix} v_1 & 0 & \cdots & 0 & 1 \\ \vdots & & & & \vdots \\ v_m & 0 & \cdots & 0 & 1 \\ v_i & \sum_{\ell=1}^k w_{j_\ell}(v_i) e_{s+j_\ell} & & 1 & \end{bmatrix}_{1 \leq i \leq m, 1 \leq j_1 < \cdots < j_k \leq p}$$

By using elementary row operations, D is equivalent to

$$D_1 = \begin{bmatrix} v_1 & 0 & \cdots & 0 & 1 \\ \vdots & & & & \vdots \\ v_m & 0 & \cdots & 0 & 1 \\ 0 & \sum_{\ell=1}^k w_{j_\ell}(v_i) e_{s+j_\ell} & & 0 & \end{bmatrix}_{1 \leq i \leq m, 1 \leq j_1 < \cdots < j_k \leq p}$$

Pick $v_{\ell_1}, \dots, v_{\ell_r}$ in $\{v_1, \dots, v_m\}$ such that $w_1(v_{\ell_1}) \neq 0, \dots, w_r(v_{\ell_r}) \neq 0$. The matrix

$$D_2 = \begin{bmatrix} v_1 & 0 & \cdots & 0 & 1 \\ \vdots & & & & \vdots \\ v_m & 0 & \cdots & 0 & 1 \end{bmatrix}$$

has rank $\dim(\mathcal{P}) + 1$ (see the proof of [37, Proposition 9.3.1]) whereas the matrix

$$D_3 = \left[0 \quad \sum_{\ell=1}^k w_{j_\ell}(v_i) e_{s+j_\ell} \quad 0 \right]_{1 \leq i \leq m, 1 \leq j_1 < \cdots < j_k \leq p}$$

has rank at most r because $w_i = 0$ on \mathcal{P} for $i > r$. To show that the rank of D_3 is r notice that $w_1(v_{\ell_1})e_{s+1}, \dots, w_r(v_{\ell_r})e_{s+r}$ are linearly independent rows of D_3 . Therefore

$$\begin{aligned} \dim(\mathcal{P}_w) + 1 &= \dim(A(\mathcal{P}_w)) = \text{rank}(D) \\ &= \text{rank}(D_1) = \text{rank}(D_2) + \text{rank}(D_3) = \dim(\mathcal{P}) + 1 + r, \end{aligned}$$

where the first two equalities follow from [37, Theorem 9.3.6(b), Corollary 8.2.21]. Hence $\dim(\mathcal{P}_w)$ is equal to $\dim(\mathcal{P}) + r$ and the proof is complete. \square

We end with an example that illustrates the results of this section:

Example 4.9. First, we give an illustration of the equality $E_{\mathcal{P}_w} = E_{\mathcal{P}}^{s, \prod_{i=1}^p (w_i+1)}$ of Theorem 4.6. Let $\mathcal{P} = \text{conv}(v_1, v_2, v_3, v_4)$, $v_1 = (0, 0)$, $v_2 = (1, 0)$, $v_3 = (0, 1)$, $v_4 = (1, 1)$, be the unit square in \mathbb{R}^2 and let w_i , $i = 1, 2, 3$, be the linear functions $w_1 = t_1 + t_2$, $w_2 = 2t_1 + 3t_2$ and $w_3 = t_1$. Using *Macaulay2* [25], we obtain

$$\begin{aligned} E_{\mathcal{P}}(n) &= (n+1)^2, \\ E_{\mathcal{P}}^{s, w_1}(n) &= (n^3 + 2n^2 + n), \\ E_{\mathcal{P}}^{s, w_2}(n) &= (5/2)n^3 + 5n^2 + (5/2)n, \\ E_{\mathcal{P}}^{s, w_3}(n) &= (1/2)n^3 + n^2 + (1/2)n, \\ E_{\mathcal{P}}^{s, w_1 w_2}(n) &= (35/12)n^4 + (20/3)n^3 + (55/12)n^2 + (5/6)n, \\ E_{\mathcal{P}}^{s, w_1 w_3}(n) &= (17/12)n^4 + (19/6)n^3 + (25/12)n^2 + (1/3)n, \\ E_{\mathcal{P}}^{s, w_2 w_3}(n) &= (7/12)n^4 + (4/3)n^3 + (11/12)n^2 + (1/6)n, \\ E_{\mathcal{P}}^{s, w_1 w_2 w_3}(n) &= (11/6)n^5 + (29/6)n^4 + (25/6)n^3 + (7/6)n^2. \end{aligned}$$

Therefore, by Theorem 4.6, we get

$$\begin{aligned} E_{\mathcal{P}_w}(n) &= E_{\mathcal{P}}^{s, \prod_{i=1}^p (w_i+1)}(n) = E_{\mathcal{P}}(n) + E_{\mathcal{P}}^{s, w_1}(n) + E_{\mathcal{P}}^{s, w_2}(n) + E_{\mathcal{P}}^{s, w_3}(n) + \\ &\quad E_{\mathcal{P}}^{s, w_1 w_2}(n) + E_{\mathcal{P}}^{s, w_1 w_3}(n) + E_{\mathcal{P}}^{s, w_2 w_3}(n) + E_{\mathcal{P}}^{s, w_1 w_2 w_3}(n) \\ &= (11/6)n^5 + (39/4)n^4 + (58/3)n^3 + (71/4)n^2 + (22/3)n + 1, \end{aligned}$$

where $\mathcal{P}_w = \text{conv}(\mathcal{G}) \subset \mathbb{R}^5$ and \mathcal{G} is the following multiset of vectors:

$$\begin{array}{cccc}
(v_1, & 0, & 0, & 0), & (v_1, & w_1(v_1), & w_2(v_1), & 0), \\
(v_2, & 0, & 0, & 0), & (v_2, & w_1(v_2), & w_2(v_2), & 0), \\
(v_3, & 0, & 0, & 0), & (v_3, & w_1(v_3), & w_2(v_3), & 0), \\
(v_4, & 0, & 0, & 0), & (v_4, & w_1(v_4), & w_2(v_4), & 0), \\
(v_1, & w_1(v_1), & 0, & 0), & (v_1, & w_1(v_1), & 0, & w_3(v_1)), \\
(v_2, & w_1(v_2), & 0, & 0), & (v_2, & w_1(v_2), & 0, & w_3(v_2)), \\
(v_3, & w_1(v_3), & 0, & 0), & (v_3, & w_1(v_3), & 0, & w_3(v_3)), \\
(v_4, & w_1(v_4), & 0, & 0), & (v_4, & w_1(v_4), & 0, & w_3(v_4)), \\
(v_1, & 0, & w_2(v_1), & 0), & (v_1, & 0, & w_2(v_1), & w_3(v_1)), \\
(v_2, & 0, & w_2(v_2), & 0), & (v_2, & 0, & w_2(v_2), & w_3(v_2)), \\
(v_3, & 0, & w_2(v_3), & 0), & (v_3, & 0, & w_2(v_3), & w_3(v_3)), \\
(v_4, & 0, & w_2(v_4), & 0), & (v_4, & 0, & w_2(v_4), & w_3(v_4)), \\
(v_1, & 0, & 0, & w_3(v_1)), & (v_1, & w_1(v_1), & w_2(v_1), & w_3(v_1)), \\
(v_2, & 0, & 0, & w_3(v_2)), & (v_2, & w_1(v_2), & w_2(v_2), & w_3(v_2)), \\
(v_3, & 0, & 0, & w_3(v_3)), & (v_3, & w_1(v_3), & w_2(v_3), & w_3(v_3)), \\
(v_4, & 0, & 0, & w_3(v_4)), & (v_4, & w_1(v_4), & w_2(v_4), & w_3(v_4)).
\end{array}$$

Computing the values of $w_i(v_j)$ for all i, j , this list of vectors becomes

$$\begin{array}{cc}
(0, 0, 0, 0, 0), & (0, 0, 0, 0, 0), \\
(1, 0, 0, 0, 0), & (1, 0, 1, 2, 0), \\
(0, 1, 0, 0, 0), & (0, 1, 1, 3, 0), \\
(1, 1, 0, 0, 0), & (1, 1, 2, 5, 0), \\
(0, 0, 0, 0, 0), & (0, 0, 0, 0, 0), \\
(1, 0, 1, 0, 0), & (1, 0, 1, 0, 1), \\
(0, 1, 1, 0, 0), & (0, 1, 1, 0, 0), \\
(1, 1, 2, 0, 0), & (1, 1, 2, 0, 1), \\
(0, 0, 0, 0, 0), & (0, 0, 0, 0, 0), \\
(1, 0, 0, 2, 0), & (1, 0, 0, 2, 1), \\
(0, 1, 0, 3, 0), & (0, 1, 0, 3, 0), \\
(1, 1, 0, 5, 0), & (1, 1, 0, 5, 1), \\
(0, 0, 0, 0, 0), & (0, 0, 0, 0, 0), \\
(1, 0, 0, 0, 1), & (1, 0, 1, 2, 1), \\
(0, 1, 0, 0, 0), & (0, 1, 1, 3, 0), \\
(1, 1, 0, 0, 1), & (1, 1, 2, 5, 1).
\end{array}$$

The Ehrhart series of \mathcal{P} and \mathcal{P}_w are

$$F_{\mathcal{P}}(x) = \frac{1+x}{(1-x)^3} \quad \text{and} \quad F_{\mathcal{P}_w}^{s,w}(x) = F_{\mathcal{P}_w}(x) = \frac{1+51x+129x^2+39x^3}{(1-x)^6}, \quad \text{respectively.}$$

5. s -WEIGHTED EHRHART FUNCTIONS AND EHRHART SERIES

In this section, we show applications to s -weighted Ehrhart theory. To avoid repetitions, we continue to employ the notations and definitions used in Section 4.

Proposition 5.1. *Let $w = w_1 \cdots w_p$ be any monomial of S of degree p with $w_i \in \{t_1, \dots, t_s\}$ for $i = 1, \dots, p$. If $w_i \not\equiv 0$ on \mathcal{P} for $i = 1, \dots, p$, then $E_{\mathcal{P}}^{s,w}$ is a polynomial whose leading coefficient is the relative volume $\text{vol}(\mathcal{P}_w)$ of \mathcal{P}_w and*

$$\deg(E_{\mathcal{P}}^{s,w}) = \dim(\mathcal{P}) + p.$$

Proof. We argue by induction on $p \geq 1$. By Theorem 4.6, we obtain

$$(5.1) \quad E_{\mathcal{P}_w}(n) = E_{\mathcal{P}}^{s, w_1 \cdots w_p}(n) + \cdots + \left(\sum_{1 \leq j_1 < \cdots < j_k \leq p} E_{\mathcal{P}}^{s, w_{j_1} \cdots w_{j_k}}(n) \right) + \cdots + E_{\mathcal{P}}(n).$$

Assume that $p = 1$. Then, by Eq. (5.1), we obtain

$$(5.2) \quad E_{\mathcal{P}_w}(n) = E_{\mathcal{P}}^{s, w}(n) + E_{\mathcal{P}}(n) \text{ for } n \geq 0.$$

Since $E_{\mathcal{P}_w}$ and $E_{\mathcal{P}}$ are Ehrhart functions of lattice polytopes, one has $\deg(E_{\mathcal{P}_w}) = \dim(\mathcal{P}_w)$, $\deg(E_{\mathcal{P}}) = \dim(\mathcal{P})$, the leading coefficient of $E_{\mathcal{P}_w}$ is $\text{vol}(\mathcal{P}_w)$ and, by Lemma 4.8, we get that $\deg(E_{\mathcal{P}_w}) = \dim(\mathcal{P}) + 1$. Hence, by Eq. (5.2), $E_{\mathcal{P}}^{s, w}$ is a polynomial of degree $\dim(\mathcal{P}) + 1$ whose leading coefficient is $\text{vol}(\mathcal{P}_w)$. Assume that $p > 1$. By induction, using Eq. (5.1), we obtain

$$E_{\mathcal{P}_w}(n) = E_{\mathcal{P}}^{s, w}(n) + g(n) \text{ for } n \geq 0,$$

where g is a polynomial of degree $\dim(\mathcal{P}) + p - 1$. Thus, by Lemma 4.8, $E_{\mathcal{P}}^{s, w} = E_{\mathcal{P}_w} - g$ is a polynomial of degree $\dim(\mathcal{P}) + p$ whose leading coefficient is $\text{vol}(\mathcal{P}_w)$. \square

A function $E: \mathbb{N} \rightarrow K$ is called a *polynomial* if there is a polynomial $g(x) \in K[x]$ such that $E(n) = g(n)$ for all $n \in \mathbb{N}$. If $E(n) = g(n)$ for $n \gg 0$, we say that E is a *polynomial function*.

The Ehrhart function $E_{\mathcal{P}}$ of a lattice polytope \mathcal{P} is a polynomial in n and so is the weighted Ehrhart function $E_{\mathcal{P}}^{s, f}$, $f \in S$.

Corollary 5.2. *Let f be a non-zero polynomial of $S = K[t_1, \dots, t_s]$ of degree p , $\mathbb{R} \subset K$, and let d be the dimension of the lattice polytope \mathcal{P} . The following hold.*

- (a) $E_{\mathcal{P}}^{s, f}$ is a polynomial of n of degree at most $d + p$ and $F_{\mathcal{P}}^{s, f}(x)$ is a rational function.
- (b) If \mathcal{P} is non-degenerate and f is a monomial, then $\deg(E_{\mathcal{P}}^{s, f}) = d + p$.
- (c) $E_{\mathcal{P}}^{s, f}$ is a K -linear combination of Ehrhart polynomials.
- (d) [12, Proposition 4.1] If the interior \mathcal{P}° of \mathcal{P} is nonempty, $K = \mathbb{R}$, and f is homogeneous and $f \geq 0$ on \mathcal{P} , then $E_{\mathcal{P}}^{s, f}$ is a polynomial of degree $s + p$.

Proof. We can write $f = \sum_b \mu_b t^b$, $\mu_b \in K \setminus \{0\}$. Then, for $n \geq 0$, one has

$$(5.3) \quad E_{\mathcal{P}}^{s, f}(n) = \sum_{a \in n\mathcal{P} \cap \mathbb{Z}^s} f(a) = \sum_{a \in n\mathcal{P} \cap \mathbb{Z}^s} \left(\sum_b \mu_b t^b(a) \right) = \sum_b \left(\mu_b \sum_{a \in n\mathcal{P} \cap \mathbb{Z}^s} t^b(a) \right) = \sum_b \mu_b E_{\mathcal{P}}^{s, t^b}(n).$$

(a) By Proposition 5.1, it follows readily that $E_{\mathcal{P}}^{s, t^b}$ is a polynomial of n of degree at most $d + p$ for all b . Thus, by Eq. (5.3), $E_{\mathcal{P}}^{s, f}$ is a polynomial of n of degree at most $d + p$. Then, by [33, Corollary 4.3.1], $F_{\mathcal{P}}^{s, f}(x)$ is a rational function [33, Corollary 4.3.1].

(b) Writing $f = w_1 \cdots w_p$ with $w_i \in \{t_1, \dots, t_s\}$ for all i and noticing that \mathcal{P} is non-degenerate if and only if $t_i \not\equiv 0$ on \mathcal{P} for $i = 1, \dots, s$, by Proposition 5.1, we get $\deg(E_{\mathcal{P}}^{s, f}) = d + p$.

(c) By Eq. (5.3), it suffices to show that $E_{\mathcal{P}}^{s, t^b}$ is a K -linear combination of Ehrhart polynomials, and this follows using induction on $\deg(t^b)$ and Theorem 4.6(d).

(d) As $\mathcal{P}^\circ \neq \emptyset$, the dimension of \mathcal{P} is equal to s , and consequently $t_i \not\equiv 0$ on \mathcal{P} for $i = 1, \dots, s$. Then, by Proposition 5.1, $E_{\mathcal{P}}^{s, t^b}$ is a polynomial of degree $s + p$ for all t^b . Let u_b be the leading coefficient of $E_{\mathcal{P}}^{s, t^b}$. By Eq. (5.3), it suffices to show that $\sum_b \mu_b u_b \neq 0$. Setting $h = t^b$ and

noticing that $h(a) = n^p h(\frac{a}{n})$ for all $a \in \mathbb{R}^s$ and $n \geq 1$, one has

$$u_b = \lim_{n \rightarrow \infty} \frac{E_{\mathcal{P}}^{s,h}(n)}{n^{s+p}} = \lim_{n \rightarrow \infty} \sum_{a \in n\mathcal{P} \cap \mathbb{Z}^s} \frac{h(\frac{a}{n})}{n^s} = \lim_{n \rightarrow \infty} \sum_{\frac{a}{n} \in \mathcal{P} \cap (\frac{1}{n}\mathbb{Z}^s)} \frac{h(\frac{a}{n})}{n^s} = \lim_{n \rightarrow \infty} \sum_{c \in \mathcal{P} \cap (\frac{1}{n}\mathbb{Z}^s)} \frac{h(c)}{n^s} = \int_{\mathcal{P}} h,$$

where the last equality is followed by elementary integration theory. Thus, $u_b = \int_{\mathcal{P}} t^b$. The fact that the leading coefficient of $E_{\mathcal{P}}^{s,h}$ is equal to $\int_{\mathcal{P}} h$ appears in [6, p. 437], and more recently in [17, Proposition 5]. Therefore

$$(5.4) \quad \sum_b \mu_b u_b = \sum_b \mu_b \int_{\mathcal{P}} t^b = \int_{\mathcal{P}} \sum_b \mu_b t^b = \int_{\mathcal{P}} f.$$

As $\mathcal{P}^\circ \neq \emptyset$, there is a box $B = B_1 \times \cdots \times B_s$ contained in an open subset of \mathcal{P} , where B_i is a closed interval of \mathbb{R} with $|B_i| = \infty$ for all i . Then, by Alon's combinatorial Nullstellensatz [1], [37, Theorem 8.4.11], there is $\beta \in B$ with $f(\beta) > 0$. Therefore, $\int_{\mathcal{P}} f > 0$ because f is a continuous function and $f \geq 0$ on \mathcal{P} . Hence, by Eq. (5.4), $\sum_b \mu_b u_b > 0$. \square

Proposition 5.3. *Let f be a homogeneous polynomial of degree p in $K[t_1, \dots, t_s]$, $K = \mathbb{R}$, and let \mathcal{P} be a lattice polytope in \mathbb{R}^s . Then, there are $c_1, c_2 \in \mathbb{R}$, $c_1 \leq c_2$, such that*

$$c_1 n^p E_{\mathcal{P}}(n) \leq E_{\mathcal{P}}^{s,f}(n) \leq c_2 n^p E_{\mathcal{P}}(n) \text{ for } n \geq 0.$$

Proof. For $a \in n\mathcal{P} \cap \mathbb{Z}^s$ and $n \geq 1$, we can write $a = nb$, with $b \in \mathcal{P}$. Hence, $f(a) = n^p f(b)$ because f is homogeneous of degree p . We set

$$c_1 = \inf\{f(x) \mid x \in \mathcal{P}\} \text{ and } c_2 = \sup\{f(x) \mid x \in \mathcal{P}\}.$$

Since f is a continuous function and \mathcal{P} is a compact subset of \mathbb{R}^s , c_1 and c_2 are real numbers and $f(\mathcal{P}) \subset [c_1, c_2]$. Thus

$$\begin{aligned} n^p c_1 &\leq f(a) = n^p f(b) \leq n^p c_2 \quad \therefore \\ c_1 n^p E_{\mathcal{P}}(n) &\leq E_{\mathcal{P}}^{s,f}(n) = \sum_{a \in n\mathcal{P} \cap \mathbb{Z}^s} f(a) \leq c_2 n^p E_{\mathcal{P}}(n), \end{aligned}$$

and the proof is complete. \square

Corollary 5.4. *Let v_1, \dots, v_m be points in \mathbb{N}^s , let $\mathcal{P} = \text{conv}(v_1, \dots, v_m)$, and let $w: \mathbb{R}^s \rightarrow \mathbb{R}$ be a non-zero linear function such that $w(e_i) \in \mathbb{N}$ for $i = 1, \dots, s$. The following hold.*

$$(a) \quad E_{\mathcal{P}}^{s,w}(n) := \sum_{a \in n\mathcal{P} \cap \mathbb{Z}^s} w(a) = E_{\mathcal{P}_w}(n) - E_{\mathcal{P}}(n) \text{ for } n \geq 0,$$

where $\mathcal{P}_w = \text{conv}((v_1, 0), \dots, (v_m, 0), (v_1, w(v_1)), \dots, (v_m, w(v_m)))$.

(b) *If $w = \sum_{i=1}^s \eta_i t_i$, $t_i \not\equiv 0$ on \mathcal{P} whenever $\eta_i > 0$, and $d = \dim(\mathcal{P})$, then $E_{\mathcal{P}}^{s,w}$ is a polynomial of degree $d + 1$ and the generating function $F_{\mathcal{P}}^{s,w}$ of $E_{\mathcal{P}}^{s,w}$ is equal to*

$$F_{\mathcal{P}}^{s,w}(x) := \sum_{n=0}^{\infty} E_{\mathcal{P}}^{s,w}(n) x^n = \frac{h(x)}{(1-x)^{d+2}},$$

where $h(x)$ is a polynomial of degree at most $d + 1$ with nonnegative integer coefficients.

Proof. (a) From Theorem 4.6(d), we get $E_{\mathcal{P}_w}(n) = E_{\mathcal{P}}^{s,w+1}(n) = E_{\mathcal{P}}^{s,w}(n) + E_{\mathcal{P}}(n)$ for $n \geq 0$.

(b) If $\eta_i > 0$, by Proposition 5.1, $\eta_i E_{\mathcal{P}}^{s,t_i}$ is a polynomial of degree $\dim(\mathcal{P}) + 1$ whose leading coefficient is positive. As $E_{\mathcal{P}}^{s,w} = \sum_{i=1}^s \eta_i E_{\mathcal{P}}^{s,t_i}$, $E_{\mathcal{P}}^{s,w}$ is a polynomial of degree $\dim(\mathcal{P}) + 1$. By

part (a), Lemma 4.8, and Stanley's result that h^* -vectors of lattice polytopes have nonnegative integer coefficients [32, Theorem 2.1], one has

$$F_{\mathcal{P}}^{s,w}(x) = F_{\mathcal{P}_w}(x) - F_{\mathcal{P}}(x) = \frac{f(x)}{(1-x)^{d+2}} - \frac{(1-x)g(x)}{(1-x)^{d+2}} = \frac{h(x)}{(1-x)^{d+2}},$$

where $f(x)$ and $g(x)$ are polynomials with non-negative integer coefficients of degrees at most $d+1$ and d , respectively. Note that $\mathcal{P}' := \text{conv}((v_1, 0), \dots, (v_m, 0)) \subset \mathcal{P}_w$, the Ehrhart series of \mathcal{P}' and \mathcal{P} are equal, and $f(x)$ and $g(x)$ are the h^* -polynomials of \mathcal{P}_w and \mathcal{P} , respectively. Then, by Stanley's monotonicity property of h^* -vectors [34, Theorem 3.3] (cf. [10, Theorem 3.3]), we get that $h^*(\mathcal{P}_w) \geq h^*(\mathcal{P}') = h^*(\mathcal{P})$ componentwise, and consequently $f(x) \geq g(x) \geq 0$ coefficientwise. Therefore, from the equalities

$$h(x) = f(x) - (1-x)g(x) = f(x) - g(x) + xg(x),$$

we obtain that $h(x)$ has nonnegative integer coefficients. \square

Corollary 5.5. *Let v_1, \dots, v_m be points in \mathbb{N}^s , let $\mathcal{P} = \text{conv}(v_1, \dots, v_m)$, and let $w: \mathbb{R}^s \rightarrow \mathbb{R}$ be a non-zero linear function such that $w(e_i) \in \mathbb{N}$ for all i . The following hold.*

- (a) *For each $n \geq 1$, $E_{\mathcal{P}}^{s,w}(n) = \lambda_n n E_{\mathcal{P}}(n)$ for some $\lambda_n \in \text{conv}(\{w(v_i)\}_{i=1}^m)$.*
- (b) *If $t_i \not\equiv 0$ on \mathcal{P} whenever $w(e_i) > 0$, then $\dim(\mathcal{P}_w) = \dim(\mathcal{P}) + 1$ and $\min\{w(v_i)\} \leq \text{vol}(\mathcal{P}_w)/\text{vol}(\mathcal{P}) \leq \max\{w(v_i)\}$.*
- (c) *If $w(v_i) = k$ for $i = 1, \dots, m$, then $E_{\mathcal{P}}^{s,w}(n) = kn E_{\mathcal{P}}(n)$ and $\text{vol}(\mathcal{P}_w) = k \text{vol}(\mathcal{P})$.*

Proof. Setting $u_i = w(v_i)$ for $i = 1, \dots, m$, we may assume $u_1 \leq \dots \leq u_m$. Note that

$$(5.5) \quad n[u_1, u_m] \cap \mathbb{Z} = [nu_1, nu_m] \cap \mathbb{Z} = \{nu_1, nu_1 + 1, nu_1 + 2, \dots, nu_m\},$$

and $p_n := |n[u_1, u_m] \cap \mathbb{Z}| = (u_m - u_1)n + 1$. We set $d = \dim(\mathcal{P})$.

- (a) For $a \in n\mathcal{P} \cap \mathbb{Z}^s$, one has $w(a) \in [nu_1, nu_m] \cap \mathbb{Z}$. Hence, by Eq. (5.5), we can write

$$E_{\mathcal{P}}^{s,w}(n) = c_1(nu_1) + c_2(nu_1 + 1) + c_3(nu_1 + 2) + \dots + c_{p_n}(nu_m),$$

where $c_i \in \mathbb{N}$ and $\sum_{i=1}^{p_n} c_i = E_{\mathcal{P}}(n)$. Then, $\lambda_n := E_{\mathcal{P}}^{s,w}(n)/nE_{\mathcal{P}}(n) \in \text{conv}(\{w(v_i)\}_{i=1}^m)$, that is, λ_n is a rational number in $[u_1, u_m]$. This completes the proof.

- (b) By part (a), we get $u_1 n E_{\mathcal{P}}(n) \leq E_{\mathcal{P}}^{s,w}(n) \leq u_m n E_{\mathcal{P}}(n)$ for $n \geq 1$. By Corollary 5.4 and [11, pp. 64 and 72], we get that $E_{\mathcal{P}}^{s,w}(n)$ is a polynomial in n of degree $d+1 = \dim(\mathcal{P}_w)$ whose leading coefficient is $\text{vol}(\mathcal{P}_w)$. As $nE_{\mathcal{P}}(n)$ is a polynomial in n of degree $\dim(\mathcal{P}) + 1$ it follows readily that $u_1 \leq \text{vol}(\mathcal{P}_w)/\text{vol}(\mathcal{P}) \leq u_m$.

- (c) This follows from parts (a) and (b) since $w(v_i) = k$ for all i . \square

ACKNOWLEDGMENTS

We used *Normaliz* [15] and *Macaulay2* [25] to compute weighted Ehrhart polynomials and series and volumes of lattice polytopes.

On behalf of all authors, the corresponding author states that there is no conflict of interest.

Data sharing is not applicable to this article as no datasets were generated or analyzed during the current study. 0

REFERENCES

- [1] N. Alon, Combinatorial Nullstellensatz, *Combin. Probab. Comput.* **8** (1999), no. 1-2, 7–29.
- [2] E. Bajo, M. Beck and A. R. Vindas-Meléndez, q -Chromatic polynomials. Preprint, 2024, [arXiv:2403.19573](https://arxiv.org/abs/2403.19573).
- [3] E. Bajo, R. Davis, J. A. De Loera, A. Garber, S. Garzón Mora, K. Jochemko and J. Yu, Weighted Ehrhart theory: extending Stanley’s nonnegativity theorem, *Adv. Math.* **444** (2024), Paper No. 109627, 30 pp.
- [4] V. Baldoni, N. Berline, J. A. De Loera, B. Dutra, M. Köppe, S. Moreinis, G. Pinto, M. Vergne, J. Wu, *A User’s Guide for LattE integrale v1.7.2*, 2013, software package LattE is available at <http://www.math.ucdavis.edu/latte/>.
- [5] V. Baldoni, N. Berline, J. A. De Loera, M. Köppe and M. Vergne, How to integrate a polynomial over a simplex, *Math. Comp.* **80** (2011), no. 273, 297–325.
- [6] V. Baldoni, N. Berline, J. A. De Loera, M. Köppe and M. Vergne, Computation of the highest coefficients of weighted Ehrhart quasi-polynomials of rational polyhedra, *Found. Comput. Math.* **12** (2012), no. 4, 435–469.
- [7] A.I. Barvinok, Calculation of exponential integrals. (Russian) *Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)* **192** (1991), Teor. Slozhn. Vychisl. 5, 149–162, 175–176; translation in *J. Math. Sci.* **70** (1994), no. 4, 1934–1943.
- [8] A.I. Barvinok, Partition functions in optimization and computational problems, *St. Petersburg. Math. J.* **4** (1992), no. 1, 1–49; translation from *Algebra Anal.* **4** (1992), no. 1, 3–53.
- [9] M. Beck, Stanley’s major contributions to Ehrhart theory, *The mathematical legacy of Richard P. Stanley*, 53–63. American Mathematical Society, Providence, RI, 2016.
- [10] M. Beck, B. Braun and A. R. Vindas-Meléndez, Decompositions of Ehrhart h^* -polynomials for rational polytopes, *Discrete Comput. Geom.* **68** (2022), no. 1, 50–71.
- [11] M. Beck and S. Robins, *Computing the continuous discretely*, Second edition, Undergraduate Texts in Mathematics, Springer, New York, 2017.
- [12] M. Brion and M. Vergne, Lattice points in simple polytopes, *J. Amer. Math. Soc.* **10** (1997), no. 2, 371–392.
- [13] W. Bruns and J. Gubeladze, *Polytopes, Rings, and K-Theory*, Springer Monographs in Mathematics, Springer, Dordrecht, 2009.
- [14] W. Bruns and J. Herzog, *Cohen-Macaulay Rings*, Cambridge University Press, Cambridge, Revised Edition, 1998.
- [15] W. Bruns, B. Ichim, T. Römer, R. Sieg and C. Söger: Normaliz. Algorithms for rational cones and affine monoids. Available at <https://normaliz.uos.de>.
- [16] W. Bruns, B. Ichim and C. Söger, Computations of volumes and Ehrhart series in four candidates elections, *Ann. Oper. Res.* **280** (2019), no. 1-2, 241–265.
- [17] W. Bruns and C. Söger, The computation of generalized Ehrhart series in Normaliz, *J. Symbolic Comput.* **68** (2015), part 2, 75–86.
- [18] W. Bruns and C. Söger, *NmzIntegrate*. Available from <http://www.mathematik.uni-osnabrueck.de/normaliz/Normaliz2.11/NmzIntegrate1.2.pdf>
- [19] F. Chapoton, q -analogues of Ehrhart polynomials, *Proc. Edinb. Math. Soc. (2)* **59** (2016), no. 2, 339–358.
- [20] D. Cox, J. Little and H. Schenck, *Toric Varieties*, Graduate Studies in Mathematics **124**, American Mathematical Society, Providence, RI, 2011.
- [21] J. A. De Loera, L. Escobar, N. Kaplan and C. Wang, Sums of Weighted Lattice Points of Polytopes, *Séminaire Lotharingien de Combinatoire 89B* (2023), Proceedings of the 35th Conference on Formal Power Series and Algebraic Combinatorics (Davis), Article no. 17, 12 pp.
- [22] E. Ehrhart, *Polynômes arithmétiques et méthode des polyèdres en combinatoire*, International Series of Numerical Mathematics, Vol. 35, Birkhäuser Verlag, Basel-Stuttgart, 1977.
- [23] C. Escobar, J. Martínez-Bernal and R. H. Villarreal, Relative volumes and minors in monomial subrings, *Linear Algebra Appl.* **374** (2003), 275–290.
- [24] I. Gitler and R. H. Villarreal, *Graphs, Rings and Polyhedra*, Aportaciones Mat. Textos, **35**, Soc. Mat. Mexicana, México, 2011.
- [25] D. Grayson and M. Stillman, *Macaulay2*, 1996. Available at <http://www.math.uiuc.edu/Macaulay2/>.
- [26] M. Hochster, Rings of invariants of tori, Cohen–Macaulay rings generated by monomials, and polytopes, *Ann. of Math.* **96** (1972), 318–337.
- [27] J. B. Lasserre, Integration on a convex polytope, *Proc. Amer. Math. Soc.* **126** (1998), no. 8, 2433–2441.
- [28] V. Reiner and B. Rhoades, Harmonics and graded Ehrhart theory. Preprint, 2024, [arXiv:2407.06511](https://arxiv.org/abs/2407.06511).
- [29] A. Schrijver, On total dual integrality, *Linear Algebra Appl.* **38** (1981), 27–32.
- [30] R. P. Stanley, Combinatorial reciprocity theorems, *Adv. Math.* **14** (1974), no. 2, 194–253.
- [31] R. P. Stanley, Hilbert functions of graded algebras, *Adv. Math.* **28** (1978), 57–83.

- [32] R. P. Stanley, Decompositions of rational convex polytopes, *Ann. Discrete Math.* **6** (1980), 333–342.
- [33] R. P. Stanley, *Enumerative Combinatorics I*, Wadsworth-Brooks/Cole, Monterey, California, 1986.
- [34] R. P. Stanley, A monotonicity property of h -vectors and h^* -vectors, *European J. Combin.* **14** (1993), no. 3, 251–258.
- [35] R. P. Stanley, *Combinatorics and Commutative Algebra*, Birkhäuser Boston, 2nd ed., 1996.
- [36] B. Sturmfels, *Gröbner Bases and Convex Polytopes*, University Lecture Series **8**, American Mathematical Society, Rhode Island, 1996.
- [37] R. H. Villarreal, *Monomial Algebras*, Second edition, Monographs and Research Notes in Mathematics, Chapman and Hall/CRC, Boca Raton, FL, 2015.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, DAVIS.

Email address: `deloera@math.ucdavis.edu`

DEPARTAMENTO DE MATEMÁTICAS, CINVESTAV, Av. IPN 2508, 07360, CDMX, MÉXICO.

Email address: `cvalencia@math.cinvestav.edu.mx`

DEPARTAMENTO DE MATEMÁTICAS, CINVESTAV, Av. IPN 2508, 07360, CDMX, MÉXICO.

Email address: `rvillarreal@cinvestav.mx`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, DAVIS.

Email address: `cywang@ucdavis.edu`