## arXiv:2412.17785v1 [hep-th] 23 Dec 2024

## Krylov spread complexity as holographic complexity beyond JT gravity

Michal P. Heller,<sup>1,\*</sup> Jacopo Papalini,<sup>1,†</sup> and Tim Schuhmann<sup>1,‡</sup>

<sup>1</sup>Department of Physics and Astronomy, Ghent University, 9000 Ghent, Belgium

One of the important open problems in quantum black hole physics is a dual interpretation of holographic complexity proposals. To date the only quantitative match is the equality between the Krylov spread complexity in triple-scaled SYK at infinite temperature and the complexity = volume proposal in classical JT gravity. Our work utilizes the recent connection between double-scaled SYK and sine-dilaton gravity to show that the quantitative relation between Krylov spread complexity and complexity = volume extends to finite temperatures and to full quantum regime on the gravity side at disk level. From the latter we isolate the first quantum correction to the complexity = volume proposal and propose to view it as a complexity of quantum fields in the bulk. Finally, we comment on the switchback effect, whose presence would make the Krylov spread complexity a fully fledged holographic complexity at least in sine-dilaton gravity.

Introduction. – Understanding black hole interiors has been an important theme of research in quantum gravity ever since the foundational papers on black holes appeared. A major modern incarnation of this research front has been studies of holographic complexity over the course of the past decade [1, 2]. This research direction stems from a heuristic yet convincing observation in [3] that connected perpetual growth of the black hole interiors in spatial volume with the expected (and proven in random quantum circuits [4] behaviour of quantum circuit complexity in chaotic systems. In the realm of holography this property of black hole interiors is now formalized within the complexity = anything paradigm [5, 6], which provides an infinite set of boundary-anchored bulk geometric objects which all grow linearly with the boundary time at late enough times. Arguably the simplest complexity = anything proposal is the one that was considered first: the complexity = volume proposal dealing with the spatial volume of extremal time slices of the bulk [7].

The key open question in the studies of holographic complexity is that of its interpretation in the microscopic description on the boundary. To date, despite a decade of research on the topic there has been only one fully quantitative match discovered<sup>1</sup>. In 2023 [11] (see also 2022 [12]) showed that complexity = volume in the simplest holographic gravity theory (JT gravity) is given by the Krylov spread complexity [13] in the dual description in terms of a particular (triple-scaling) limit of the SYK model at infinite temperature. While this may come as a surprise, since Krylov complexity is a priori a distinct quantity from quantum circuit complexity originally proposed in this context, both quantities realize Susskind's expectation about linear growth behavior in chaotic quantum systems [4, 13, 14].

The so far isolated nature of [11] raises the crucial question: does the relation relation between holographic complexity and microscopic Krylov spread complexity extends also to other gravity setups? The aim of our letter is to present that the answer is affirmative within a freshly discovered duality between two-dimensional sine-dilaton gravity and a double-scaled SYK model. From the point of view of the parameter space of the double-scaled SYK model, the result of [11] lies at a single point and our letter extends it to a two-dimensional plane spanned by the interaction strength (nonlocality of the model), in which case we make use of the quantum generalization of the complexity = volume proposal put forward in [15], and inverse temperature  $\beta$ . In the outlook we speculate on the implications of our study for the broader field of holographic complexity.

**Setup.**– The SYK model [16, 17] describes a system of N Majorana fermions, with dynamics governed by a p-body interaction of the form:

$$H_{\text{SYK}} = i^{p/2} \sum_{1 \le i_1 < \dots < i_p \le N} J_{i_1 \cdots i_p} \psi_{i_1} \cdots \psi_{i_p}, \qquad (1)$$

where the couplings  $J_{i_1\cdots i_p}$  are typically assumed to be Gaussian random variables. We will be focusing on the double-scaled SYK model (DSSYK), where both N and pare sent to infinity, while keeping the ratio  $|\log q| = \frac{p^2}{N}$ finite. A striking feature of this regime is that all amplitudes in DSSYK are computable in a exact way, by evaluating the so called chord diagrams, an intermediate combinatorial tool to solve the system [18, 19]. Specifically, this is done by introducing an auxiliary quantum mechanical system governed by an effective Hamiltonian, the transfer matrix  $\hat{T}$ , which acts on a Hilbert state spanned by chord states  $|n\rangle$ . See for instance [20] for a recent comprehensive review on the subject.

Our gravitational setup is sine-dilaton gravity, a recently proposed gravity dual to double-scaled SYK [21, 22]. The main object of interest for us is the length, which is the extremal volume of the complexity = volume proposal in two dimensions, in this gravity theory. The gravita-

<sup>&</sup>lt;sup>1</sup> Note other efforts, including [8–10], that reported a quantitative match in the context of a broad complexity research, but never directly with any holographic complexity proposal.

tional path integral of sine-dilaton gravity takes the form

$$\int \mathcal{D}g\mathcal{D}\Phi \,\exp\left(\frac{1}{2}\int \mathrm{d}^2x\sqrt{g}\left(\Phi R + \frac{\sin(2|\log q|\Phi)}{|\log q|}\right)\right),\quad(2)$$

where we suppressed the boundary terms. In (2) the potential for dilaton  $\Phi$  exhibits a sine profile and thus corresponds to a deformation of the linear dilaton potential of JT gravity, see e.g. [23] for a review of the latter. Specifically, to prove the complexity=volume proposal in this theory, we will be interested in the gravitational twopoint function in sine-dilaton gravity, corresponding the the boundary to boundary propagator of a non-minimally coupled scalar field in the bulk [21]. The effective geometry this matter probe will experience is given by

$$ds_{\rm eff}^2 = ds_{\rm sine}^2 e^{-2i|\log q|\Phi} = -\left(\rho^2 - \sin(\theta)^2\right) dt_S^2 + \frac{d\rho^2}{\rho^2 - \sin(\theta)^2},$$
(3)

which represents the Schwarzschild patch of an AdS<sub>2</sub> black hole with Hawking temperature  $\beta_{BH} = 2\pi/\sin(\theta)$ , which reduces to the JT black hole when  $\theta \ll 1$ . The complexity = volume in the Kruskal extension of the geometry (3), is, after holographic renormalization, proportional to

$$L = 2\log\left(\cosh\left(t\sin(\theta)/2\right)\right) - 2\log(\sin(\theta)) \qquad (4)$$

in terms of the two-sided boundary Lorentzian time t. The expression (4) represents the semiclassical limit of the full quantum expectation value of the length of the thermofield double state [24] in sine dilaton gravity as a function of t, which we will compute in (16) and match with Krylov complexity in DSSYK. By introducing the canonical conjugate P of the length L and performing canonical quantization, the gravitational Hamiltonian  $\hat{H}_{\text{grav}}$  of sine-dilaton gravity reads

$$\hat{H}_{\rm grav} = \frac{1}{\sqrt{2 |\log q|} \sqrt{1 - q^2}} \left[ -\cos\left(\hat{P}\right) + \frac{1}{2} e^{i\hat{P}} e^{-\hat{L}} \right], \quad (5)$$

which precisely matches the DSSYK transfer matrix  $\hat{T}$  [12, 18, 19] (see supplemental material for details), making the duality manifest. Moreover, as discussed in detail in [25], the invariance of the Hamiltonian (5) under periodic shifts  $P \rightarrow P + 2\pi$  of the momentum instructs us to consider this as a redundancy and to treat P as a compact variable. Upon doing so, the theory is projected onto discretized lenghts. We point out how the discretization of chord number is obvious from the point of view of DSSYK [12, 18, 19], but the quantization of the length is far from obvious from the gravity side. Upon projection, the physical Hilbert space  $|L\rangle$  can then be identified with the chord Hilbert space of DSSYK, spanned by the chord number states  $|n\rangle$  [12, 18, 19], via the holographic dictionary

$$L = 2 \left| \log q \right| n. \tag{6}$$

Indeed, gauging  $P \rightarrow P + 2\pi$  is also responsible for selecting  $|L = 0\rangle$  as the true gravitational vacuum state of the theory, since all negative lengths states are removed from the physical Hilbert space [25] . This reduces the Euclidean periodicity  $\beta_{\rm BH}=2\pi/\sin(\theta)$  of the black hole background (3) to

$$\beta_{\text{DSSYK}} = \frac{2\pi - 4\theta}{\sin(\theta)},\tag{7}$$

reproducing the true microscopic temperature of DSSYK<sup>2</sup>. The global slice Wheeler-DeWitt wavefunctions of sine dilaton gravity are the solutions of the Schroedinger equation associated with (5). One finds the following spectrum

$$E(\theta) = 2\cos(\theta)/\sqrt{2|\log q|(1-q^2)},\tag{8}$$

and the eigenvectors in the L-basis are given by q-Hermite polynomials, i.e.

$$\psi_{\theta}(L) = \langle \theta | L \rangle = H_n \left( \cos(\theta) | q^2 \right), \tag{9}$$

with right eigenvectors given instead by  $\langle L|\theta\rangle = H_n\left(\cos(\theta)|q^2\right)/\left(q^2;q^2\right)_n$  [22]. With the Hamiltonian now diagonalized, one can determine the two-point function of a massive probe in sine-dilaton gravity, which is captured by the insertion of the bilocal operator  $e^{-\Delta \hat{L}}$  of conformal weight  $\Delta$  [21]. The structure of this expectation value takes a similar form to the JT gravity one [27, 28], i.e.

$$\langle e^{-\Delta \hat{L}} \rangle = Z_{\beta}^{-1} \langle L = 0 | e^{-\tau \hat{H}_{\text{grav}}} e^{-\Delta \hat{L}} e^{-(\beta - \tau) \hat{H}_{\text{grav}}} | L = 0 \rangle,$$
(10)

where the disk has been divided in two parts of length  $\tau$ and  $\beta - \tau$  corresponding to the Euclidean boundary times to which the semiclassical geodesics is anchored. Inserting two completeness relation in the  $\theta$  basis, we obtain

$$\langle e^{-\Delta \hat{L}} \rangle = Z_{\beta}^{-1} \int_{0}^{\pi} \mathrm{d}\theta_{1} \,\rho(\theta_{1}) \int_{0}^{\pi} \mathrm{d}\theta_{2} \,\rho(\theta_{2}) \exp\left(-\tau E(\theta_{1})\right)$$
$$\exp\left(-(\beta - \tau)E(\theta_{2})\right) \langle\theta_{1}| \,e^{-\Delta \hat{L}} \,|\theta_{2}\rangle \,, \tag{11}$$

where one has [18, 22]:

$$\langle \theta_1 | e^{-\Delta \hat{L}} | \theta_2 \rangle = \frac{(q^{4\Delta}; q^2)_\infty}{(q^{2\Delta} e^{\pm 2i\theta_1 \pm 2i\theta_2}; q^2)_\infty}$$
$$= \sum_{n=0}^\infty \frac{q^{2n\Delta}}{(q^2; q^2)_n} H_n \left( \cos \theta_1 | q^2 \right) H_n \left( \cos \theta_2 | q^2 \right)$$
(12)

and  $\rho(\theta) = (q^2, e^{\pm 2i\theta}; q^2)_{\infty}$  is the spectral density associated with the wavefunctions (9). Analytically continuing as  $\tau = \frac{\beta}{2} + it$  in (11), we finally obtain the Lorentzian two-point function in sine dilaton gravity. We finally point out the latter is given by the expectation value of the bilocal operator in the state

$$|\psi(t)\rangle = Z_{\beta}^{-\frac{1}{2}} e^{-it\hat{H}_{\text{grav}}} |\psi(\beta)\rangle , \qquad (13)$$

<sup>&</sup>lt;sup>2</sup> The Euclidean time periodicity of the solution (3) corresponds to the so called "fake temperature"  $\beta_{\rm BH} = \beta_{\rm fake} = \frac{2}{\frac{2\pi}{n(\theta)}}$  [26].

which corresponds to the Lorentzian time evolution of the Hartle-Hawking state  $|\psi(\beta)\rangle = Z_{\beta}^{-\frac{1}{2}} e^{-\beta \hat{H}_{\text{grav}}/2} |L=0\rangle$  obtained by the Eucliden half-disk preparation amplitude.

**Deriving Krylov complexity from length.** – The Krylov spread complexity is defined as the expectation value of the position of a state  $|\psi(t)\rangle$  spreading over the Krylov basis  $|K_n\rangle$  [13]

$$C_K(t) = \sum_{n=0}^{\infty} n \left| \langle K_n | \psi(t) \rangle \right|^2, \quad |\psi(t)\rangle = e^{-iHt} |R\rangle.$$
 (14)

The state  $|\psi(t)\rangle$  follows Hamiltonian time evolution from the reference state  $|R\rangle$  and the Krylov basis  $|K_n\rangle$  is the unique basis that arises from Gram-Schmidt orthonormalization of the set of states constructed via repeated application of the Hamiltonian to the reference state  $H^n|R\rangle$ , a procedure commonly referred to as the Lanczos algorithm.

On the gravity side, the results of [11] motivate us to study the Lorentzian evolution of the expectation value of the length in sine dilaton gravity (2). Specifically, we will extract the length from the two point function of scaling dimension  $\Delta$  operators (10) as

$$\langle \hat{L} \rangle = \left[ -\partial_{\Delta} \langle e^{-\Delta \hat{L}} \rangle \right]_{\Delta=0}.$$
 (15)

Note that this starting point coincides with the way the authors of [15] lay out their study of the black hole interior at very late times in JT gravity.

The key observation we make is that by leveraging the relation (15) to derive  $\langle \theta_1 | \hat{L} | \theta_2 \rangle$  from (12), one obtains

$$\begin{aligned} \langle \hat{L} \rangle &= \frac{\left| \log q^2 \right|}{Z_\beta} \int_0^\pi \mathrm{d}\theta_1 \,\rho(\theta_1) \int_0^\pi \mathrm{d}\theta_2 \,\rho(\theta_2) e^{-\left(\frac{\beta}{2} + it\right)E(\theta_1)} \\ e^{-\left(\frac{\beta}{2} - it\right)E(\theta_2)} \sum_{n=0}^\infty \frac{n}{(q^2; q^2)_n} H_n\left(\cos\theta_1 | q^2\right) H_n\left(\cos\theta_2 | q^2\right) , \end{aligned}$$
(16)

where, importantly, a sum over n appears. As we will now show, the result can be brought to the form of a Krylov spread complexity (14). On the way, we will crucially identify the state  $|\psi(t)\rangle$  that this Krylov spread complexity is associated to.

By construction, we can identify the components appearing in (16) as eigenvalues  $E(\theta)$ , given in (8), and eigenfunctions (in the chord basis)  $\langle n|\theta\rangle$ , given in (9), of the transfer matrix  $\hat{T}$ , the effective Hamiltonian of the chord Hilbert space of DSSYK [12, 18, 19]<sup>3</sup>. This is because the gravitational Hamiltonian  $\hat{H}_{\text{grav}}$  of sine dilaton gravity, as discussed in the previous section, is constructed

3

to precisely match  $\hat{T}$ . Moreover, we leverage the fact that the Krylov basis  $|K_n\rangle$  coincides with the chord basis  $|n\rangle$ in DSSYK, as discussed in [11]. While recent arguments suggest that this identification holds only for times shorter than those exponential in the entropy [29], we argue that any potential corrections to this correspondence lie beyoud the strict  $N \to \infty$  limit of DSSYK. In this regime, we are probing time scales smaller than the saturation time. This perspective aligns with the expectation that late-time deviations arise from higher-genus contributions to the gravitational path integral of sine-dilaton gravity. Since such contributions are not considered in our analysis, the identification of the Krylov and chord basis remains robust within the scope of our work. These combined observations let us identify the length, evolving in Lorentzian time t at nonzero DSSYK inverse temperature  $\beta$ , with the Krylov complexity  $C_K(t)_{\beta}$ 

$$\langle \hat{L} \rangle = \frac{\left|\log q^{2}\right|}{Z_{\beta}} \sum_{n=0}^{\infty} n \left| \int_{0}^{\pi} \mathrm{d}\theta \,\rho(\theta) e^{-\left(\frac{\beta}{2}+it\right)E(\theta)} \langle n|\theta \rangle \langle \theta|0 \rangle \right|^{2}$$

$$= \left|\log q^{2}\right| \sum_{n} n \left| \langle n| \frac{e^{-iT(t-i\beta/2)}}{\sqrt{Z_{\beta}}} |0 \rangle \right|^{2} = \left|\log q^{2}\right| C_{K}(t)_{\beta} \quad (17)$$

of the (normalized) state  $|\psi(t)\rangle_{\beta} = Z_{\beta}^{-\frac{1}{2}} e^{-iT(t-i\beta/2)}|0\rangle$ that arises by mixed Lorentzian plus Euclidean transfer matrix evolution from the zero chord state  $|0\rangle$  as reference state. The matching

$$2\left|\log q\right|C_{K}(t)_{\beta} = \langle \hat{L} \rangle = \left[-\partial_{\Delta} \langle e^{-\Delta \hat{L}} \rangle\right]_{\Delta=0}$$
(18)

is one of the main results of this letter. Note that this is a match on the quantum level that is valid for all allowed values of  $q \in (0, 1)$  in DSSYK, as well as for all values of  $\beta$ 

**Two limits.** – Although our identification (17) holds at the full quantum level, it is important to check the correct semiclassical limit, as  $q \rightarrow 1$  limit, is recovered. Indeed, as one can observe in Fig. 1, the numerical evaluation of (16) is very well interpolated by the semiclassical effective length in sine-dilaton gravity, presented in (4). The same semiclassical limit was also checked for the exact two-point function in [30], leading, by extrapolation through (15), to the same result (4). Moreover, the relation (7) between inverse temperature  $\beta$  and  $\theta$  is also matched with the saddle point approximation of the exact DSSYK partition function [30]. We can also verify that the correct JT limit is recovered. This corresponds to small values of  $\theta$ , where the spectral edge of the DSSYK spectrum is probed. In this regime, the semiclassical length simplifies to

$$L_{\rm JT} = 2\log\left(\cosh\left(t\theta/2\right)\right) - 2\log(\theta) \tag{19}$$

which corresponds to the length of the ERB bridge in the JT black hole background [31], at temperature  $\beta_{\rm JT} = 2\pi/\theta$ . On the other hand, we can also investigate the regime where  $\beta \to 0$  which, according to (7), corresponds to  $\theta = \frac{\pi}{2}$ , the midpoint of the DSSYK spectral support. In this regime, the length becomes:

$$L_{\beta=0} = 2\log\left(\cosh\left(t/2\right)\right) \tag{20}$$

<sup>&</sup>lt;sup>3</sup> Without loss of generality, we set the DSSYK coupling J that appears as a multiplicative factor in  $E(\theta)$  to J = 1 for the rest of this letter. Hence, all times are understood to be measured in units of J.

In [11] this limit, corresponding to a Krylov complexity at  $\beta = 0$ , was connected again to a JT classical length by performing a particular combination of limits. We clarify that taking only the  $\beta \to 0$  limit does not lead to JT gravity. This does not contradict the results of [11] because their combination of limits is different from the pure  $\beta \to 0$  limit that we take. At the level of sine-dilaton gravity, the  $\beta \to 0$  regime is different from the standard JT limit, as it corresponds to an expansion of the dilaton potential around  $\Phi = \pi/2$ . In this case, as shown in [25], the gravity theory reduces to a regularization of flat-space JT gravity, reproducing the results of [32] associated to the high-temperature limit of DSSYK.

Finite temperature lessons. – There are two main lessons to be drawn from the result (18). Firstly, the calculation shows that gravity demands to assign complexity to both Lorentzian evolution as well as Euclidean state preparation in order to match Krylov complexity to the Lorentzian evolution of the volume  $\langle \hat{L} \rangle$  at nonzero inverse temperature  $\beta$ . Importantly, this notion of complexity is different from starting at reference state  $e^{-\beta/2T} |0\rangle$  and assigning complexity only to the Lorentzian evolution. This is because the different choice of reference state leads to a different Krylov basis, and hence a different Krylov complexity.

$$C_K \text{ for } e^{-iT(t-i\beta/2)} |0\rangle \neq C_K \text{ for } e^{-iTt} e^{-\beta T/2} |0\rangle |R\rangle$$
.

The second lesson concerns the dependence of the Krylov complexity  $C_K(t)_\beta$  on the temperature. We observe two important features. A visualization of  $C_K(t)_{\beta}$  where both features are clearly visible is given in Fig. 1. The complexity (solid line) acquires a positive onset that increases with increasing  $\beta$ . This is the complexity associated to Euclidean preparation of the Hartle-Hawking state  $e^{-\beta T/2}|0\rangle$ . The other distinct feature is that finite temperature slows down the complexity growth. This agrees with the finite temperature observations in [33] and, as discussed there, is consistent with what has been confirmed for other instances of the SYK model. When contrasted with the analytic result for the classical length (dashed line) given in eq. (4), we observe correlation but not a precise match for a generic choice of q and  $\beta$ . We will now show in the following section that this is expected and can be resolved by taking into account quantum corrections to the classical length.

The role of quantum corrections to the length.-In [11, 33] showing that certain Krylov complexities, calculated in specific DSSYK setups, are dual to JT gravity results, was achieved by explicit calculation of both and observation of coinciding analytic structure in the semiclassical approximation, combined with matching the parameters. This is different from what is reported in



FIG. 1. Numerical values for the finite temperature Krylov spread complexity  $C_K(t)_\beta$  at various temperatures  $\beta$  and q = 0.6 (solid). It is visible that for larger  $\beta$ , the initial onset at t = 0 that accounts for higher complexity of the Euclidean state preparation is larger and the complexity growth slows down. The complexities are contrasted with the expectation value of the classical length in sine dilaton gravity at temperature  $\beta$ (dashed), normalized by 2  $|\log q|$ .

this letter. Here, we can explicitly derive the Krylov complexity from the length in sine-dilaton gravity directly.

The strength of our result lies in the fact that we leverage a full gravitational dual to DSSYK, extending beyond the triple-scaling limit where JT gravity emerges. Consequently, we uncover a fully quantum mechanical bulk manifestation of complexity within the dual gravity theory.

Let us focus on the region of q close to but not exactly one. In this region, the full quantum answer of the length in sine dilaton gravity is well approximated by the classical answer  $\langle \hat{L} \rangle_{\rm cl.}$ , given in eq. (4), modified by the first correction in  $|\log q|$ . The analytical expression for this correction has been very recently derived by one of us and his collaborators in [34] and reads

$$\langle \hat{L} \rangle_{q\text{-corr.}} = -\frac{|\log q|}{4((\pi - 2\theta)\cot(\theta) + 2)} \\ \left[ \left( -t^2 \sin^2(\theta) \left( \cot^2(\theta) + \tanh^2\left(\frac{1}{2}t\sin(\theta)\right) \right) \right) \\ + \frac{4\left(t\sin(\theta)\tanh\left(\frac{1}{2}t\sin(\theta)\right) - 2\right)}{(\pi - 2\theta)\cot(\theta) + 2} \\ - \left( (\pi - 2\theta)\cot(\theta) + 2 \right)^2 \operatorname{sech}^2\left(\frac{1}{2}t\sin(\theta)\right) \\ + \left( t\tanh\left(\frac{1}{2}t\sin(\theta)\right) - 2\csc(\theta)\right)^2 + 4 \right].$$
(21)

Numerically, we can now compare the Krylov spread complexity  $C_K(t)_\beta$  we derived in (17), which is the full quantum answer at generic value of q, and check whether it matches with  $\langle \hat{L} \rangle_{\text{cl.}}$  in (4) plus  $\langle \hat{L} \rangle_{q\text{-corr.}}$  in (21) for values of q close to q = 1. This comparison is visualized in Fig. 2. We find excellent agreement around q = 1. As



FIG. 2. Numerically evaluated Krylov complexity  $C_K(t)_\beta$ (solid), normalized by the classical length  $\langle \hat{L} \rangle_{\rm cl.}$  (and a factor of  $2 |\log q|$  that is omitted in the label) for  $\beta = 1$  and various q (color coding) vs. the leading order quantum correction  $\langle \hat{L} \rangle_{q\text{-corr.}}$  (dot-dashed), also normalized by  $\langle \hat{L} \rangle_{\rm cl.}$ , for the same values of q and  $\beta$ . For q close to one, the curves fall on top of each other, signaling perfect agreement; For q further away from q = 1, the deviation systematically increases because of higher order corrections that were not taken into account.

expected, the agreement gets worse for lower q; this is because here, higher order corrections to the volume that are not known analytically become relevant.

**Complexity equals volume: operator statement.**– Let us highlight that Krylov complexity equals volume in sine dilaton gravity dual to finite temperature DSSYK can also be made as an operator statement. Krylov spread complexity can be rewritten as the expectation value

$$C_K(t) = \langle \psi(t) | \hat{K} | \psi(t) \rangle, \quad \hat{K} = \sum_{n=0}^{\infty} n | K_n \rangle \langle K_n | \quad (22)$$

of the Krylov operator  $\hat{K}$  in the state  $|\psi(t)\rangle$  that arises via transfer matrix evolution from the zero chord state as reference state. The Krylov basis  $|K_n\rangle$  coincides with the chord basis  $|n\rangle$  in DSSYK [11] for times smaller than exponential in the entropy which is the natural restriction of the presented argument. Through the holographic dictionary  $L = 2 |\log q| n$ , established in [25] and discussed in the setup, it is also known that the chord number operator  $\hat{n} = \sum_{n=0}^{\infty} n |n\rangle \langle n|$  is the length operator  $\hat{L}$  in sine dilaton gravity. Combining these two statements implies that the Krylov operator coincides with the length operator

$$2\left|\log q\right|\hat{K} = \hat{L}\tag{23}$$

in sine dilaton gravity dual to finite temperature DSSYK. This is an operator manifestation of the complexity equals volume conjecture in the setup of this letter. By looking at the expectation value of this expression in the state  $|\psi(t)\rangle$ , we find

$$2\left|\log q\right| C_K(t) = \langle \psi(t) | \hat{L} | \psi(t) \rangle \tag{24}$$

which precisely agrees with the result we have derived by explicit calculation in eq. (17).

Summary. – We presented a precise duality between Krylov spread complexity in DSSYK at finite temperature and length in sine dilaton gravity. This is accomplished by identifying the analytic expression for the length in sine dilaton gravity as a Krylov spread complexity. In particular, to establish this duality, gravity demands to assign complexity to the Euclidean preparation of the Hartle-Hawking state. The derived duality holds on the quantum level, meaning for all DSSYK parameters  $q \in (0, 1)$ . This is exemplified by a numerical comparison of the derived Krylov spread complexity with the first quantum correction to the length. For q close to one where the full quantum answer is well approximated by the first quantum correction, we observe perfect agreement. The duality studied in this paper is also made manifest as an operator statement.

**Outlook.** – Krylov complexity is expected to saturate at times exponential in the entropy. Ab inito, this seems to conflict with the linear growth that is observed in [11] and this letter as the asymptotic behaviour of the studied Krylov spread complexities. The resolution of this puzzle lies in taking into account non-perturbative corrections. From the gravitational perspective, the two-point function is expected to receive contributions from higher topologies in the gravitational path integral of sine-dilaton gravity. In particular, the saturation of the two-point function—manifesting as a late-time plateau that halts the linear growth—is likely driven by non-perturbative effects in the topological expansion. This mirrors the behavior observed in JT gravity, where the dual matrix model [35] provided a framework to explain such non-perturbative late-time phenomena [15]. Remarkable progress in this direction was very recently also reported in [29]. We conjecture that similar mechanisms should govern sinedilaton gravity, whose dual matrix model is expected to correspond to the ETH matrix model for q-deformed JT gravity [36]. On the boundary side, any potential incarnation of Krylov complexity is thus expected to go beyond the double-scaled SYK model, which corresponds to the strict  $N \to \infty$  limit.

Moreover, it is important to state that for Krylov spread complexity, a proof of the presence of the switchback effect - i.e. scrambling time delay of the linear growth associated with the presence of shockwaves in the bulk [7] - would complete its understanding as a genuine holographic complexity. There are strong reasons to believe that this characteristic can be made manifest in the context of the presented model. We expect that insights from JT gravity, such as the observation of Shapiro time delay in the two-point function of a heavy bilocal operator via a boundary clock analysis [23] and the connection of the crossed four-point function with shockwave scattering [37], translate to sine dilaton gravity. These characteristics are very closely tied to what would be interpreted as the switchback effect from the boundary complexity viewpoint, justifying the conjecture of existence of the switchback effect in DSSYK Krylov complexity. Recently, important boundary insights connecting to this line of argument have been reported for the triple-scaled DSSYK model in [33, 38]. A finite  $\beta$  adaptation of these methods, together with relaxation of the triple-scaling limit, could facilitate the proposed study of the switchback effect in Krylov complexity in the general case that we elaborate on in this letter.

Furthermore, the analysis of the quantum correction to the length and its connection to Krylov complexity leads us to two important questions to be investigated. Firstly, it is important to gain a geometric understanding of the the first quantum correction to the length in sine dilaton gravity. This carries the potential to obtain a refined understanding of the complexity equals volume conjecture in the full quantum regime where quantum corrections to the volume have to be taken into account. Secondly, the functional form of the quantum correction to the length suggests a particular interesting interpretation: In the early time limit, the correction itself behaves quadratically, supplemented by an onset in the case of finite temperature, while in the late time limit it exhibits linear growth. This coincides with the expected behaviour of a Krylov spread complexity as studied in this letter. Hence, reminiscent of the case of holographic entanglement entropy where the first quantum correction is given by entanglement between bulk quantum fields [39], we conjecture that the length quantum correction has a dual interpretation as Krylov spread complexity of a bulk quantum field.

Finally, the most important question is whether the paradigm presented in this letter extends to higher dimensions. An interesting lesson seems to be that a crucial ingredient in the match is to account for the Euclidean segment of time evolution.

We would like to thank Thomas G. Mertens, Andreas Belaey and Thomas Tappeiner for inspiring discussions that helped to spark this project and Mario Flory, Aranya Bhattacharya and Emiliano Rizza for collaborations on related questions. MPH acknowledges welcoming hospitality of Jagiellonian University and discussions with Romuald Janik during the completion of this project. TS is supported by the Research Foundation - Flanders (FWO) doctoral fellowship 11I5425N. JP acknowledge financial support from the European Research Council (grant BHHQG-101040024). Funded by the European Union. Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or the European Research Council. Neither the European Union nor the granting authority can be held responsible for them.

- \* michal.p.heller@ugent.be
- <sup>†</sup> jacopo.papalini@ugent.be
- <sup>‡</sup> tim.schuhmann@ugent.be
- L. Susskind, Three Lectures on Complexity and Black Holes, SpringerBriefs in Physics, Springer, 10, 2018. arXiv:1810.11563.
- [2] S. Chapman and G. Policastro, Quantum computational complexity from quantum information to black holes and back, Eur. Phys. J. C 82 (2022), no. 2 128 [arXiv:2110.14672].
- [3] L. Susskind, Computational Complexity and Black Hole Horizons, Fortsch. Phys. 64 (2016) 24–43
   [arXiv:1403.5695]. [Addendum: Fortsch.Phys. 64, 44–48 (2016)].
- [4] J. Haferkamp, P. Faist, N. B. T. Kothakonda, J. Eisert and N. Y. Halpern, *Linear growth of quantum circuit complexity*, Nature Phys. 18 (2022), no. 5 528–532 [arXiv:2106.05305].
- [5] A. Belin, R. C. Myers, S.-M. Ruan, G. Sárosi and A. J. Speranza, *Does Complexity Equal Anything?*, Phys. Rev. Lett. **128** (2022), no. 8 081602 [arXiv:2111.02429].
- [6] A. Belin, R. C. Myers, S.-M. Ruan, G. Sárosi and A. J. Speranza, *Complexity equals anything II*, JHEP **01** (2023) 154 [arXiv:2210.09647].
- [7] D. Stanford and L. Susskind, Complexity and Shock Wave Geometries, Phys. Rev. D 90 (2014), no. 12 126007 [arXiv:1406.2678].
- [8] N. Chagnet, S. Chapman, J. de Boer and C. Zukowski, Complexity for Conformal Field Theories in General Dimensions, Phys. Rev. Lett. **128** (2022), no. 5 051601 [arXiv:2103.06920].
- [9] J. Erdmenger, A.-L. Weigel, M. Gerbershagen and M. P. Heller, From complexity geometry to holographic spacetime, Phys. Rev. D 108 (2023), no. 10 106020 [arXiv:2212.00043].
- [10] P. Caputa, B. Chen, R. W. McDonald, J. Simón and B. Strittmatter, Spread Complexity Rate as Proper Momentum, arXiv:2410.23334.
- [11] E. Rabinovici, A. Sánchez-Garrido, R. Shir and J. Sonner, A bulk manifestation of Krylov complexity, JHEP 08 (2023) 213 [arXiv:2305.04355].
- [12] H. W. Lin, The bulk Hilbert space of double scaled SYK, JHEP **11** (2022) 060 [arXiv:2208.07032].
- [13] V. Balasubramanian, P. Caputa, J. M. Magan and Q. Wu, Quantum chaos and the complexity of spread of states, Phys. Rev. D 106 (2022), no. 4 046007 [arXiv:2202.06957].
- [14] V. Balasubramanian, M. Decross, A. Kar and O. Parrikar, *Quantum Complexity of Time Evolution* with Chaotic Hamiltonians, JHEP **01** (2020) 134 [arXiv:1905.05765].
- [15] L. V. Iliesiu, M. Mezei and G. Sárosi, The volume of the black hole interior at late times, JHEP 07 (2022) 073 [arXiv:2107.06286].
- [16] S. Sachdev and J. Ye, Gapless spin fluid ground state in a random, quantum Heisenberg magnet, Phys. Rev. Lett. 70 (1993) 3339 [arXiv:cond-mat/9212030].

- [17] A. Kitaev, "A simple model of quantum holography." http://online.kitp.ucsb.edu/online/entangled15/ kitaev/ and http://online.kitp.ucsb.edu/online/ entangled15/kitaev2/, 2015.
- [18] M. Berkooz, M. Isachenkov, V. Narovlansky and G. Torrents, *Towards a full solution of the large N* double-scaled SYK model, JHEP **03** (2019) 079 [arXiv:1811.02584].
- [19] M. Berkooz, P. Narayan and J. Simon, Chord diagrams, exact correlators in spin glasses and black hole bulk reconstruction, JHEP 08 (2018) 192 [arXiv:1806.04380].
- [20] M. Berkooz and O. Mamroud, A Cordial Introduction to Double Scaled SYK, arXiv:2407.09396.
- [21] A. Blommaert, T. G. Mertens and J. Papalini, *The dilaton gravity hologram of double-scaled SYK*, arXiv:2404.03535.
- [22] A. Blommaert, T. G. Mertens and S. Yao, Dynamical actions and q-representation theory for double-scaled SYK, JHEP 02 (2024) 067 [arXiv:2306.00941].
- [23] T. G. Mertens and G. J. Turiaci, Solvable models of quantum black holes: a review on Jackiw-Teitelboim gravity, Living Rev. Rel. 26 (2023), no. 1 4 [arXiv:2210.10846].
- [24] J. M. Maldacena, Eternal black holes in anti-de Sitter, JHEP 04 (2003) 021 [arXiv:hep-th/0106112].
- [25] A. Blommaert, A. Levine, T. G. Mertens, J. Papalini and K. Parmentier, An entropic puzzle in periodic dilaton gravity and DSSYK, arXiv:2411.16922.
- [26] H. W. Lin and D. Stanford, A symmetry algebra in double-scaled SYK, SciPost Phys. 15 (2023), no. 6 234 [arXiv:2307.15725].
- [27] A. Blommaert, T. G. Mertens and H. Verschelde, *The Schwarzian Theory A Wilson Line Perspective*, JHEP 12 (2018) 022 [arXiv:1806.07765].
- [28] L. V. Iliesiu, S. S. Pufu, H. Verlinde and Y. Wang, An exact quantization of Jackiw-Teitelboim gravity, JHEP 11

(2019) 091 [arXiv:1905.02726].

- [29] V. Balasubramanian, J. M. Magan, P. Nandi and Q. Wu, Spread complexity and the saturation of wormhole size, arXiv:2412.02038.
- [30] A. Goel, V. Narovlansky and H. Verlinde, Semiclassical geometry in double-scaled SYK, JHEP 11 (2023) 093 [arXiv:2301.05732].
- [31] D. Harlow and D. Jafferis, The Factorization Problem in Jackiw-Teitelboim Gravity, JHEP 02 (2020) 177 [arXiv:1804.01081].
- [32] A. Almheiri, A. Goel and X.-Y. Hu, Quantum gravity of the Heisenberg algebra, JHEP 08 (2024) 098 [arXiv:2403.18333].
- [33] J. Xu, On Chord Dynamics and Complexity Growth in Double-Scaled SYK, arXiv:2411.04251.
- [34] L. Bossi, L. Griguolo, J. Papalini, L. Russo and D. Seminara, Sine-dilaton gravity vs double-scaled SYK: exploring one-loop quantum corrections, arXiv:2411.15957.
- [35] P. Saad, S. H. Shenker and D. Stanford, JT gravity as a matrix integral, arXiv:1903.11115.
- [36] D. L. Jafferis, D. K. Kolchmeyer, B. Mukhametzhanov and J. Sonner, Jackiw-Teitelboim gravity with matter, generalized eigenstate thermalization hypothesis, and random matrices, Phys. Rev. D 108 (2023), no. 6 066015 [arXiv:2209.02131].
- [37] H. T. Lam, T. G. Mertens, G. J. Turiaci and H. Verlinde, Shockwave S-matrix from Schwarzian Quantum Mechanics, JHEP 11 (2018) 182 [arXiv:1804.09834].
- [38] M. Ambrosini, E. Rabinovici, A. Sánchez-Garrido, R. Shir and J. Sonner, Operator K-complexity in DSSYK: Krylov complexity equals bulk length, arXiv:2412.15318.
- [39] T. Faulkner, A. Lewkowycz and J. Maldacena, *Quantum corrections to holographic entanglement entropy*, JHEP 11 (2013) 074 [arXiv:1307.2892].

## Supplemental material

More details on sine dilaton gravity and the duality. – Let us provide further details about the duality between sine-dilaton gravity, described by the action in eq. (2), and double-scaled SYK. The ADM energy of sine-dilaton gravity is given by [21]:

$$E_{\text{ADM}} = H_{\text{grav}} = -\frac{\cos(\theta)}{2\left|\log q\right|} \,. \tag{25}$$

We now aim to perform a canonical transformation in phase space to express the gravitational Hamiltonian in more convenient variables, following an approach analogous to the one used for JT gravity in [31]. In eq. (4), we computed the length of the thermofield double (TFD) state in the black hole geometry (3), expressed in terms of the two-sided Lorentzian time evolution t, which is canonically conjugate to the ADM energy (25). To proceed with canonical quantization, we derive the canonical conjugate variable P to the TFD length L (given in (4)) by imposing that the symplectic measure on phase space adopts the canonical form in terms of L and P:

$$\omega = \mathrm{d}t \wedge \mathrm{d}H_{\mathrm{grav}} = \frac{1}{2\left|\log q\right|} \mathrm{d}L \wedge \mathrm{d}P \,. \tag{26}$$

This is achieved through

$$P = i \log (i \sin(\theta) \tanh(\sin(\theta)t/2) + \cos(\theta)). \qquad (27)$$

By inverting  $\theta$  and t in terms of L and P, we express the gravitational Hamiltonian  $H_{\text{grav}}$  in the new variables:

$$H_{\rm grav}(L,P) = -\frac{\cos(P)}{2\left|\log q\right|} + \frac{1}{4\left|\log q\right|}e^{iP}e^{-L}.$$
 (28)

Upon quantization of this Hamiltonian, operator ordering ambiguities may arise. A particular ordering choice, replacing  $e^{i\hat{P}} \rightarrow e^{\alpha(q)\hat{L}}e^{i\hat{P}}e^{-\alpha(q)\hat{L}}$  for  $\alpha(q) = \frac{1}{4|\log q|}\log\left(\frac{2|\log q|}{1-q^2}\right)$  (we thank the authors of [34] for a discussion about this point), yields the following quantum Hamiltonian

$$\hat{H}_{\rm grav}(\hat{L},\hat{P}) = \frac{1}{\sqrt{2|\log q|}\sqrt{1-q^2}} \left[ -\cos\left(\hat{P}\right) + \frac{1}{2}e^{i\hat{P}}e^{-\hat{L}} \right].$$
(29)

This exactly corresponds to the DSSYK transfer matrix  $\hat{T}$ , in accordance with the holographic dictionary (6). The mapping

$$a_{\text{norm}}, a_{\text{norm}}^{\dagger} = -e^{\pm i\vec{P}}$$
 (30)

relates to the normalized raising and lowering operators in the chord basis [12, 20]. In the gravitational framework, the symplectic structure (26) induces the standard commutation relation  $[\hat{L}, \hat{P}] = 2i |\log q|$ , implying that  $e^{i\hat{P}}$  acts as a shift operator by  $2 |\log q|$ . Consequently, the Schrödinger equation associated with (29) takes the form of a difference equation in the *L*-basis:

$$2E(\theta)\psi_{\theta}(L) = \psi_{\theta}(L+2|\log q|) + (1-e^{-L})\psi_{\theta}(L-2|\log q|).$$
(31)

This equation can be solved for a general continuous variable L, but it reduces to the DSSYK wavefunctions (9) after discretization [25]. Furthermore, the orthogonality of these wavefunctions yields the DSSYK spectral density via:

$$\begin{aligned} \langle \theta | \theta' \rangle &= \sum_{n=0}^{+\infty} \frac{1}{(q^2; q^2)_n} H_n(\cos(\theta) | q^2) H_n(\cos(\theta') | q^2) \\ &= \frac{\delta(\theta - \theta')}{(e^{\pm 2i\theta}; q^2)_\infty}. \end{aligned}$$
(32)

Numerical evaluation of finite temperature Krylov spread complexity. – For transparency, let us lay out briefly how we obtain data points in our numerical evaluation of Krylov spread complexity at finite  $\beta$  and generic q. Numerical evaluation is facilitated by introduction of two cutoffs  $n_{\text{cut}}$  and  $k_{\text{cut}}$  (chosen sufficiently large) in the relevant infinite sums. We numerically evaluate

$$C_K(t)_{\beta} = Z_{\beta}^{-1} \sum_{n=1}^{n_{\text{cut}}} \frac{n}{(q;q)_n} |\phi_n(t - i\beta/2)|^2 \qquad (33)$$

where

$$\phi_{0}(t) = \sum_{k=-k_{\text{cut}}}^{k_{\text{cut}}} q^{\binom{k}{2}} J_{2k} \left( \frac{2t/\sqrt{1-q}}{\sqrt{-\log(q)}} \right), \qquad (34)$$

$$\phi_{n}(t) = (-i)^{n} \sum_{k=0}^{k_{\text{cut}}} \frac{q^{\binom{k}{2}} \left(1-q^{2k+n}\right) (q;q)_{k+n} J_{2k+n} \left(\frac{2t/\sqrt{1-q}}{\sqrt{-\log(q)}}\right)}{(1-q^{k+n}) (q;q)_{k}} \qquad (35)$$

We owe these analytic expressions for the wave functions  $\phi_n(t)$ , which are key for the numerical implementation, to the detailed analysis in [33]. The infinite sum in k converges sufficiently fast (for thorough analysis see [33]) such that cutting off the sum at  $k_{\rm cut}$  large enough is justified. We have chosen  $k_{\rm cut} = 50$  in our implementation. A sufficiently large cutoff  $n_{\rm cut}$  can now be identified by requiring the norm of the state

$$1 \stackrel{!}{=} Z_{\beta}^{-1} \sum_{n=0}^{n_{\text{cut}}} \frac{1}{(q;q)_n} |\phi_n(t - i\beta/2)|^2$$
(36)

to agree with one within a desired accuracy. Here,  $Z_\beta$  also depends on  $n_{\rm cut}$  and  $k_{\rm cut}$  as

$$Z_{\beta} = \sum_{n=0}^{n_{\text{cut}}} \frac{1}{(q;q)_n} |\phi_n(-i\beta/2)|^2.$$
 (37)